

- [85762] mutated in adult-onset type II citrullinaemia encodes a putative mitochondrial
- [85763] carrier protein. *Nature Genet.* 22: 159–163, 1999.
- [85764] 15035.
- [85765] Saheki, T.; Kobayashi, K.: Mitochondrial aspartate glutamate carrier
- [85766] (citrin) deficiency as the cause of adult-onset type II citrullinemia
- [85767] (CTLN2) and idiopathic neonatal hepatitis (NICCD). *J. Hum. Genet.* 47:
- [85768] 333–341, 2002.
- [85769] 15036.
- [85770] Hara, Y.; Wakamori, M.; Ishii, M.; Maeno, E.; Nishida, M.; Yoshida,
- [85771] T.; Yamada, H.; Shimizu, S.; Mori, E.; Kudoh, J.; Shimizu, S.; Kurose,
- [85772] H.; Okada, Y.; Imoto, K.; Mori, Y.: LTRPC2
- Ca(2+)-permeable channel
- [85773] activated by changes in redox status confers susceptibility to cell
- [85774] death. *Molec. Cell* 9: 163–173, 2002.

[85775] 15037.

[85776] Harteneck, C.; Plant, T. D.; Schultz, G.: From worm to man: three

[85777] subfamilies of TRP channels. Trends Neurosci. 23: 159–166, 2000.

[85778] 15038.

[85779] Kudoh, J.; Nagamine, K.; Asakawa, S.; Abe, I.; Kawasaki, K.; Maeda,

[85780] H.; Tsujimoto, S.; Minoshima, S.; Ito, F.; Shimizu, N.: Localization

[85781] of 16 exons to a 450-kb region involved in the autoimmune polyglandular

[85782] disease type I (APECED) on human chromosome 21q22.3. DNA Res. 4:

[85783] 45–52, 1997.

[85784] 15039.

[85785] Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.; Asakawa,

[85786] S.; Ito, F.; Shimizu, N.: Molecular cloning of a novel putative Ca(2+)

[85787] channel protein (TRPC7) highly expressed in brain. Genomics 54:

- [85788] 124–131, 1998.
- [85789] 15040.
- [85790] Perraud, A.–L.; Fleig, A.; Dunn, C. A.; Bagley, L. A.; Launay,
- [85791] P.; Schmitz, C.; Stokes, A. J.; Zhu, Q.; Bessman, M. J.; Penner, R.;
- [85792] Kinet, J.–P.; Scharenberg, A. M.: ADP–ribose gating of the calcium–permeable
- [85793] LTRPC2 channel revealed by Nudix motif homology. Nature 411: 595–599,
- [85794] 2001.
- [85795] 15041.
- [85796] Sano, Y.; Inamura, K.; Miyake, A.; Mochizuki, S.; Yokoi, H.; Matsushime,
- [85797] H.; Furuichi, K.: Immunocyte Ca(2+) influx system mediated by LTRPC2. Science 293:
- [85798] 1327–1330, 2001.
- [85799] 15042.
- [85800] Koyama, N.; Ishibashi, K.; Kuwahara, M.; Inase, N.; Ichioka, M.;
- [85801] Sasaki, S.; Marumo, F.: Cloning and functional expression of human

- [85802] aquaporin8 cDNA and analysis of its gene. Genomics 54: 169–172, 1998.
- [85803] 1998.
- [85804] 15043.
- [85805] Viggiano, L.; Rocchi, M.; Svelto, M.; Calamita, G.: Assignment of the aquaporin–8 water channel gene (AQP8) to human chromosome 16p12. Cytogenet. Cell Genet. 84: 208–210, 1999.
- [85806] 15044.
- [85807] Sperandio, M. P.; Borsani, G.; Incerti, B.; Zollo, M.; Rossi, E.;
- [85808] Zuffardi, O.; Castaldo, P.; Taglialatela, M.; Andria, G.; Sebastio, G.: The gene encoding a cationic amino acid transporter (SLC7A4) maps to the region deleted in the velocardiofacial syndrome. Genomics 49: 230–236, 1998.
- [85809] 15045.
- [85810] Koegl, M.; Hoppe, T.; Schlenker, S.; Ulrich, H. D.; Mayer, T.

U.;

[85816] Jentsch, S.: A novel ubiquitination factor, E4, is involved in multiubiquitin

[85817] chain assembly. Cell 96: 635–644, 1999.

[85818] 15046.

[85819] Nonaka, S.; Tanaka, Y.; Okada, Y.; Takeda, S.; Harada, A.; Kanai,

[85820] Y.; Kido, M.; Hirokawa, N.: Randomization of left–right asymmetry

[85821] due to loss of nodal cilia generating leftward flow of extraembryonic

[85822] fluid in mice lacking KIF3B motor protein. Cell 95: 829–837, 1998.

[85823] 15047.

[85824] Yamazaki, H.; Nakata, T.; Okada, Y.; Hirokawa, N.: KIF3A/B: a

[85825] heterodimeric kinesin superfamily protein that works as a microtubule

[85826] plus end–directed motor for membrane organelle transport. J. Cell

[85827] Biol. 130: 1387–1399, 1995.

[85828] 15048.

- [85829] Suzuki, N.; Zara, J.; Sato, T.; Ong, E.; Bakhiet, N.; Oshima, R.
- [85830] G.; Watson, K. L.; Fukuda, M. N.: A cytoplasmic protein, bystin,
- [85831] interacts with trophinin, tastin, and cytokeratin and may be involved
- [85832] in trophinin-mediated cell adhesion between trophoblast and endometrial
- [85833] epithelial cells. Proc. Nat. Acad. Sci. 95: 5027-5032, 1998.
- [85834] 15049.
- [85835] Hurskainen, T. L.; Hirohata, S.; Seldin, M. F.; Apte, S. S.: ADAM-TS5,
- [85836] ADAM-TS6, and ADAM-TS7, novel members of a new family of zinc metalloproteases:
- [85837] general features and genomic distribution of the ADAM-TS family. J.
- [85838] Biol. Chem. 274: 25555-25563, 1999.
- [85839] 15050.
- [85840] Tang, B. L.; Hong, W.: ADAMTS: a novel family of proteases with
- [85841] an ADAM protease domain and thrombospondin 1 re-

peats. FEBS Lett. 445:

[85842] 223–225, 1999.

[85843] 15051.

[85844] Tortorella, M. D.; Burn, T. C.; Pratta, M. A.; Abbaszade, I.;
Hollis,

[85845] J. M.; Liu, R.; Rosenfeld, S. A.; Copeland, R. A.; Decicco, C.
P.;

[85846] Wynn, R.; Rockwell, A.; Yang, F.; and 16 others: Purifica-
tion and

[85847] cloning of aggrecanase-1: a member of the ADAMTS fam-
ily of proteins. Science 284:

[85848] 1664–1666, 1999.

[85849] 15052.

[85850] Shiratsuchi, T.; Oda, K.; Nishimori, H.; Suzuki, M.; Taka-
hashi,

[85851] E.; Tokino, T.; Nakamura, Y.: Cloning and characterization
of BAP3

[85852] (BAI-associated protein 3), a C2 domain-containing pro-
tein that interacts

[85853] with BAI1. Biochem. Biophys. Res. Commun. 251:
158–165, 1998.

[85854] 15053.

- [85855] Almeida, R.; Amado, M.; David, L.; Levery, S. B.; Holmes, E. H.;
- [85856] Merckx, G.; Geurts van Kessel, A.; Rygaard, E.; Hassan, H.; Bennett,
- [85857] E.; Clausen, H.: A family of human beta-4-galactosyltransferases:
- [85858] cloning and expression of two novel UDP-galactose:beta-N-acetylglucosamine
- [85859] beta-1,4-galactosyltransferases, beta-4-Gal-T2 and beta-4-Gal-T3. J.
- [85860] Biol. Chem. 272: 31979-31991, 1997.
- [85861] 15054.
- [85862] Sato, T.; Furukawa, K.; Bakker, H.; Van den Eijnden, D. H.; Van
- [85863] Die, I.: Molecular cloning of a human cDNA encoding beta-1,4-galactosyltransferase
- [85864] with 37% identity to mammalian UDP-Gal:GlcNAc beta-1,4-galactosyltransferase. Proc.
- [85865] Nat. Acad. Sci. 95: 472-477, 1998.
- [85866] 15055.
- [85867] Shaper, N.: Personal Communication. Baltimore, Md. 11/2/1999.

[85868] 15056.

[85869] Gundelfinger, E.: Personal Communication. Madgeburg, FRG. 1/8/1999.

[85870] 15057.

[85871] Hashida, H.; Goto, J.; Zhao, N.; Takahashi, N.; Hirai, M.; Kanazawa,

[85872] I.; Sakaki, Y.: Cloning and mapping of ZNF231, a novel brain-specific

[85873] gene encoding neuronal double zinc finger protein whose expression

[85874] is enhanced in a neurodegenerative disorder, multiple system atrophy

[85875] (MSA). Genomics 54: 50-58, 1998.

[85876] 15058.

[85877] Sparks, A. B.; Peterson, S. N.; Bell, C.; Loftus, B. J.; Hocking,

[85878] L.; Cahill, D. P.; Frassica, F. J.; Streeten, E. A.; Levine, M. A.;

[85879] Fraser, C. M.; Adams, M. D.; Broder, S.; Venter, J. C.; Vogelstein,

[85880] B.; Ralston, S. H.: Mutation screening of the TNFRSF11A gene encoding

- [85881] receptor activator of NF- κ B (RANK) in familial and sporadic Paget's
- [85882] disease of bone and osteosarcoma. *Calcif. Tissue Int.* 68: 151–155,
- [85883] 2001.
- [85884] 15059.
- [85885] Santorelli, F. M.; Patrono, C.; Fortini, D.; Tessa, A.; Comanducci,
- [85886] G.; Bertini, E.; Pierallini, A.; Amabile, G. A.; Casali, C.: Intrafamilial
- [85887] variability in hereditary spastic paraplegia associated with an SPG4
- [85888] gene mutation. *Neurology* 55: 702–705, 2000.
- [85889] 15060.
- [85890] Sauter, S.; Mitterski, B.; Klimpe, S.; Bonsch, D.; Schols, L.;
- [85891] Visbeck, A.; Papke, T.; Hopf, H. C.; Engel, W.; Deufel, T.; Epplen,
- [85892] J. T.; Neesen, J.: Mutation analysis of the spastin gene (SPG4) in
- [85893] patients in Germany with autosomal dominant hereditary spastic paraplegia. *Hum.*

- [85894] *Mutat.* 20: 127–132, 2002.
- [85895] 15061.
- [85896] Svenson, I. K.; Ashley–Koch, A. E.; Pericak–Vance, M. A.;
Marchuk,
- [85897] D. A.: A second leaky splice–site mutation in the spastin
gene. (Letter) *Am.*
- [85898] *J. Hum. Genet.* 69: 1407–1409, 2001.
- [85899] 15062.
- [85900] Flannery, C. R.; Hughes, C. E.; Schumacher, B. L.; Tudor,
D.; Aydelotte,
- [85901] M. B.; Kuettner, K. E.; Caterson, B.: Articular cartilage su-
perficial
- [85902] zone protein (SZP) is homologous to megakaryocyte stim-
ulating factor
- [85903] precursor and is a multifunctional proteoglycan with po-
tential growth–promoting,
- [85904] cytoprotective, and lubricating properties in cartilage
metabolism. *Biochem.*
- [85905] *Biophys. Res. Commun.* 254: 535–541, 1999.
- [85906] 15063.
- [85907] Ikegawa, S.; Sano, M.; Koshizuka, Y.; Nakamura, Y.: Isola–

tion,

[85908] characterization and mapping of the mouse and human PRG4 (proteoglycan

[85909] 4) genes. *Cytogenet. Cell Genet.* 90: 291–297, 2000.

[85910] 15064.

[85911] Merberg, D. M.; Fitz, L. J.; Temple, P.; Giannotti, J.; Murtha,

[85912] P.; Fitzgerald, M.; Scaltreto, H.; Kelleher, K.; Preissner, K.; Kriz,

[85913] R.; Jacobs, K.; Turner, K.: In: Preissner, K. T.; Rosenblatt, S.;

[85914] Kost, C.; Wegerhoff, J.; Mosher, D. F. (eds.): *Biology of Vitronectins*

[85915] and Their Receptors. Elsevier Science, B.V. 1993. Pp. 45–53.

[85916] 15065.

[85917] Schumacher, B. L.; Block, J. A.; Schmid, T. M.; Aydelotte, M. B.;

[85918] Kuettner, K. E.: A novel proteoglycan synthesized and secreted by

[85919] chondrocytes of the superficial zone of articular cartilage. *Arch.*

[85920] *Biochem. Biophys.* 15: 144–152, 1994.

[85921] 15066.

[85922] Pfeiffer, R.; et al; et al: Amino acid transport of
y(0,+)-L-type

[85923] by heterodimers of 4F2hc/CD98 and members of the gly-
coprotein-associated

[85924] amino acid transporter family. EMBO J. 18: 49-57, 1999.

[85925] 15067.

[85926] Pineda, M.; et al.; et al: Identification of a membrane pro-
tein

[85927] (LAT-2) that co-expresses with 4F2hc an L type amino
acid transport

[85928] activity with broad specificity for small and large zwitteri-
onic amino

[85929] acids. J. Biol. Chem. 274: 19738-19744, 1999.

[85930] 15068.

[85931] Sato, H.; Tamba, M.; Ishii, T.; Bannai, S.: Cloning and ex-
pression

[85932] of a plasma membrane cystine/glutamate exchange
transporter composed

[85933] of two distinct proteins. J. Biol. Chem. 274: 11455-11458,
1999.

[85934] 15069.

[85935] Torrents, D.; et al.; et al: Identification and characterization

[85936] of a membrane protein (y+L amino acid transporter-1) that associates

[85937] with 4F2hc to encode the amino acid transport activity y+L: a candidate

[85938] gene for lysinuric protein intolerance. J. Biol. Chem. 273: 32437-32445,

[85939] 1998.

[85940] 15070.

[85941] Dominguez, A.; Ramos-Morales, F.; Romero F.; Rios R. M.; Dreyfus,

[85942] F.; Tortolero, M.; Pintor-Toro, J. A.: hpttg, a human homologue of

[85943] rat pttg, is overexpressed in hematopoietic neoplasms: evidence for

[85944] a transcriptional activation function of hPTTG. Oncogene 17: 2187-2193,

[85945] 1998.

[85946] 15071.

[85947] Ishikawa, H.; Heaney, A. P.; Yu, R.; Horwitz, G. A.; Melmed, S.

[85948] : Human pituitary tumor-transforming gene induces angiogenesis. J.

[85949] Clin. Endocr. Metab. 86: 867-874, 2001.

[85950] 15072.

[85951] Jallepalli, P. V.; Waizenegger, I. C.; Bunz, F.; Langer, S.; Speicher,

[85952] M. R.; Peters, J.-M.; Kinzler, K. W.; Vogelstein, B.; Lengauer, C.

[85953] : Securin is required for chromosomal stability in human cells. Cell 105:

[85954] 445-457, 2001.

[85955] 15073.

[85956] Kakar, S. S.: Assignment of the human tumor transforming gene

[85957] TUTR1 to chromosome band 5q35.1 by fluorescence in situ hybridization. Cytogenet.

[85958] Cell Genet. 83: 93-95, 1998.

[85959] 15074.

[85960] Pei, L.: Pituitary tumor-transforming gene protein asso-

ciates

[85961] with ribosomal protein S10 and a novel human homologue of DnaJ in

[85962] testicular cells. J. Biol. Chem. 274: 3151–3158, 1999.

[85963] 15075.

[85964] Wang, Z.; Yu, R.; Melmed, S.: Mice lacking pituitary tumor transforming

[85965] gene show testicular and splenic hypoplasia, thymic hyperplasia, thrombocytopenia,

[85966] aberrant cell cycle progression, and premature centromere division. Molec.

[85967] Endocr. 15: 1870–1879, 2001. 1. Dominguez, A.; Ramos-Morales, F.;

[85968] Romero, F.; Rios, R. M.; Dreyfus,

[85969] 15076.

[85970] Yu, R.; Ren, S.-G.; Horwitz, G. A.; Wang, Z.; Melmed, S.: Pituitary

[85971] tumor transforming gene (PTTG) regulates placental JEG-3 cell division

[85972] and survival: evidence from live cell imaging. Molec. Endocr. 14:

[85973] 1137–1146, 2000.

[85974] 15077.

[85975] Zhang, X.; Horwitz, G. A.; Heaney, A. P.; Nakashima, M.;
Prezant,

[85976] T. R.; Bronstein, M. D.; Melmed, S.: Pituitary tumor trans-
forming

[85977] gene (PTTG) expression in pituitary adenomas. J. Clin. En-
docr. Metab. 84:

[85978] 761–767, 1999.

[85979] 15078.

[85980] Zhang, X.; Horwitz, G. A.; Prezant, T. R.; Valentini, A.;
Nakashima,

[85981] M.; Bronstein, M. D.; Melmed, S.: Structure, expression,
and function

[85982] of human pituitary tumor-transforming gene (PTTG).
Molec. Endocr. 13:

[85983] 156–166, 1999.

[85984] 15079.

[85985] Pennica, D.; Swanson, T. A.; Welsh, J. W.; Roy, M. A.;
Lawrence,

[85986] D. A.; Lee, J.; Brush, J.; Taneyhill, L. A.; Deuel, B.; Lew, M.;
Watanabe,

- [85987] C.; Cohen, R. L.; Melhem, M. F.; Finley, G. G.; Quirke, P.;
Goddard,
[85988] A. D.; Hillan, K. J.; Gurney, A. L.; Botstein, D.; Levine, A. J.:
[85989] WISP genes are members of the connective tissue growth
factor family
[85990] that are up-regulated in Wnt-1-transformed cells and
aberrantly expressed
[85991] in human colon tumors. Proc. Nat. Acad. Sci. 95:
14717-14722, 1998.
[85992] 15080.
[85993] Tanaka, S.; Sugimachi, K.; Saeki, H.; Kinoshita, J.; Ohga,
T.;
[85994] Shimada, M.; Maehara, Y.; Sugimachi, K.: A novel variant
of WISP1
[85995] lacking a von Willebrand type C module overexpressed in
scirrhous
[85996] gastric carcinoma. Oncogene 20: 5525-5532, 2001.
[85997] 15081.
[85998] Feng, L.; Novak, E. K.; Hartnell, L. M.; Bonifacino, J. S.;
Collinson,
[85999] L. M.; Swank, R. T.: The Hermansky-Pudlak syndrome 1
(HPS1) and HPS2

- [86000] genes independently contribute to the production and function of platelet
- [86001] dense granules, melanosomes, and lysosomes. Blood 99: 1651–1658,
- [86002] 2002.
- [86003] 15082.
- [86004] Vanhooren, J. C. T.; Marynen, P.; Mannaerts, G. P.; Van Veldhoven,
- [86005] P. P.: Evidence for the existence of a pristanoyl-CoA oxidase gene
- [86006] in man. Biochem. J. 325: 593–599, 1997.
- [86007] 15083.
- [86008] Gee, S.; Krauss, S. W.; Miller, E.; Aoyagi, K.; Arenas, J.; Conboy,
- [86009] J. G.: Cloning of mDEAH9, a putative RNA helicase and mammalian homologue
- [86010] of *Saccharomyces cerevisiae* splicing factor Prp43. Proc. Nat. Acad.
- [86011] Sci. 94: 11803–11807, 1997.
- [86012] 15084.
- [86013] Imamura, O.; Sugawara, M.; Furuichi, Y.: Cloning and

characterization

[86014] of a putative human RNA helicase gene of the DEAH-box protein family. *Biochem.*

[86015] *Biophys. Res. Commun.* 240: 335–340, 1997.

[86016] 15085.

[86017] Del Mastro, R. G.; Wang, L.; Simmons, A. D.; Gallardo, T. D.; Clines,

[86018] G. A.; Ashley, J. A.; Hilliard, C. J.; Wasmuth, J. J.; McPherson,

[86019] J. D.; Lovett, M.: Human chromosome-specific cDNA libraries: new

[86020] tools for gene identification and genome annotation.

Genome Res. 5:

[86021] 185–194, 1995.

[86022] 15086.

[86023] Sano, K.; Tanihara, H.; Heimark, R. L.; Obata, S.; Davidson, M.;

[86024] St. John, T.; Taketani, S.; Suzuki, S.: Protocadherins: a large family

[86025] of cadherin-related molecules in central nervous system. *EMBO J.* 12:

[86026] 2249–2256, 1993.

- [86027] 15087.
- [86028] Frigerio, J.-M.; Berthezene, P.; Garrido, P.; Ortiz, E.;
Barthelley,
- [86029] S.; Vasseur, S.; Sastre, B.; Seleznieff, I.; Dagorn, J. C.; Iovanna,
vanna,
- [86030] J. L.: Analysis of 2166 clones from a human colorectal
cancer cDNA
- [86031] library by partial sequencing. Hum. Molec. Genet. 4:
37-43, 1995.
- [86032] 15088.
- [86033] Frigerio, J.-M.; Dagorn, J.-C.; Iovanna, J. L.: Cloning, se-
quencing
- [86034] and expression of the L5, L21, L27a, L28, S5, S9, S10 and
S29 human
- [86035] ribosomal protein mRNAs. Biochim. Biophys. Acta 1262:
64-68, 1995.
- [86036] 15089.
- [86037] Dominguez, O.; Ashhab, Y.; Sabater, L.; Belloso, E.; Caro,
P.;
- [86038] Pujol-Borrell, R.: Cloning of ARE-containing genes by AU-
motif-directed
- [86039] display. Genomics 54: 278-286, 1998.

- [86040] 15090.
- [86041] Kostrub, C. F.; Knudsen, K.; Subramani, S.; Enoch, T.:
Hus1p,
[86042] a conserved fission yeast checkpoint protein, interacts
with Rad1p
[86043] and is phosphorylated in response to DNA damage. EMBO
J. 17: 2055–2066,
[86044] 1998.
- [86045] 15091.
- [86046] Lieberman, H. B.; Hopkins, K. M.; Nass, M.; Demetrick, D.;
Davey,
[86047] S.: A human homolog of the Schizosaccharomyces pombe
rad9+ checkpoint
[86048] control gene. Proc. Nat. Acad. Sci. 93: 13890–13895,
1996.
- [86049] 15092.
- [86050] Stocker, M.; Kerschensteiner, D.: Cloning and tissue dis-
tribution
[86051] of two new potassium channel alpha-subunits from rat
brain. Biochem.
- [86052] Biophys. Res. Commun. 248: 927–934, 1998.

[86053] 15093.

[86054] Seki, N.; Hattori, A.; Hayashi, A.; Kozuma, S.; Hori, T.;
Saito,

[86055] T.: The human regulator of G-protein signaling protein 6
gene (RGS6)

[86056] maps between markers WI-5202 and D14S277 on chro-
mosome 14q24.3. J.

[86057] Hum. Genet. 44: 138-140, 1999.

[86058] 15094.

[86059] Leegwater, P. A. J.; Konst, A. A. M.; Kuyt, B.; Sandkuijl, L.
A.;

[86060] Naidu, S.; Oudejans, C. B. M.; Schutgens, R. B. H.; Pronk, J.
C.;

[86061] van der Knaap, M. S.: The gene for leukoencephalopathy
with vanishing

[86062] white matter is located on chromosome 3q27. Am. J.
Hum. Genet. 65:

[86063] 728-734, 1999.

[86064] 15095.

[86065] tom Dieck, S.; Sanmarti-Vila, L.; Langnaese, K.; Richter, K.;
Kindler,

- [86066] S.; Soyke, A.; Wex, H.; Smalla, K.-H.; Kampf, U.; Franzer, J.-T.;
- [86067] Stumm, M.; Garner, C. C.; Gundelfinger, E. D.: Bassoon, a novel zinc-finger
- [86068] CAG/glutamine-repeat protein selectively localized at the active zone
- [86069] of presynaptic nerve terminals. J. Cell Biol. 142: 499-509, 1998.
- [86070] 15096.
- [86071] George, A. L., Jr.; Komisarof, J.; Kallen, R. G.; Barchi, R. L.
- [86072] : Primary structure of the adult human skeletal muscle voltage-dependent
- [86073] sodium channel. Ann. Neurol. 31: 131-137, 1992.
- [86074] 15097.
- [86075] Heine, R.; Pika, U.; Lehmann-Horn, F.: A novel SCN4A mutation
- [86076] causing myotonia aggravated by cold and potassium. Hum. Molec. Genet. 2:
- [86077] 1349-1353, 1993.
- [86078] 15098.
- [86079] Meyer-Kleine, C.; Otto, M.; Zoll, B.; Koch, M. C.: Molecular

- [86080] and genetic characterization of German families with paramyotonia
- [86081] congenita and demonstration of founder effect in the Ravensberg families. Hum.
- [86082] Genet. 93: 707–710, 1994.
- [86083] 15099.
- [86084] Orrell, R. W.; Jurkat-Rott, K.; Lehmann-Horn, F.; Lane, R. J. M.: Familial cramp due to potassium-aggravated myotonia. J. Neurol.
- [86086] Neurosurg. Psychiat. 65: 569–572, 1998.
- [86087] 15100.
- [86088] Rojas, C. V.: Personal Communication. Pittsburgh, Pa. 2/6/1992.
- [86089] 15101.
- [86090] Rojas, C. V.; Wang, J.; Schwartz, L. S.; Hoffman, E. P.; Powell, B. R.; Brown, R. H., Jr.: A met-to-val mutation in the skeletal muscle Na⁺ channel alpha-subunit in hyperkalaemic periodic paralysis. Nature 354: 387–389, 1991.

[86094] 15102.

[86095] Winter, c.; tom Dieck, S.; Boeckers, T. M.; Bockmann, J.;
Kampf,

[86096] U.; Sanmarti-Vila, L.; Langnaese, K.; Altrock, W.; Stumm,
M.; Soyke,

[86097] A.; Wieacker, P.; Garner, C. C.; Gundelfinger, E. D.: The
presynaptic

[86098] cytomatrix protein Bassoon: sequence and chromosomal
localization

[86099] of the human BSN gene. Genomics 57: 389–397, 1999.

[86100] 15103.

[86101] Bui, T. D.; Levy, E. R.; Subramaniam, V. N.; Lowe, S. L.;
Hong,

[86102] W.: cDNA characterization and chromosomal mapping of
human Golgi

[86103] SNARE GS27 and GS28 to chromosome 17. Genomics 57:
285–288, 1999.

[86104] 15104.

[86105] Lowe, S. L.; Peter, F.; Subramaniam, V. N.; Wong, S. H.;
Hong,

[86106] W.: A SNARE involved in protein transport through the
Golgi apparatus. Nature 389:

- [86107] 881–884, 1997.
- [86108] 15105.
- [86109] Nagahama, M.; Orci, L.; Ravazzola, M.; Amherdt, M.; La-comis, L.;
- [86110] Tempst, P.; Rothman, J. E.; Sollner, T. H.: A v-SNARE im-plicated
- [86111] in intra-Golgi transport. J. Cell Biol. 133: 507–516, 1996.
- [86112] 15106.
- [86113] Subramaniam, V. N.; Peter, F.; Philp, R.; Wong, S. H.; Hong, W.
- [86114] : GS28, a 28-kilodalton Golgi SNARE that participates in ER-Golgi
- [86115] transport. Science 272: 1161–1163, 1996.
- [86116] 15107.
- [86117] Hay, J. C.; Chao, D. S.; Kuo, C. S.; Scheller, R. H.: Protein
- [86118] interactions regulating vesicle transport between the en-doplasmic
- [86119] reticulum and Golgi apparatus in mammalian cells. Cell 89: 149–158,
- [86120] 1997.

- [86121] 15108.
- [86122] Li, J.; Ding, S.-F.; Habib, N. A.; Fermor, B. F.; Wood, C. B.;
- [86123] Gilmour, R. S.: Partial characterization of a cDNA for human stearyl-CoA
- [86124] desaturase and changes in its mRNA expression in some normal and malignant
- [86125] tissues. *Int. J. Cancer* 57: 348-352, 1994.
- [86126] 15109.
- [86127] Ntambi, J. M.; Miyazaki, M.; Stoeckl, J. P.; Lan, H.; Kendziora,
- [86128] C. M.; Yandell, B. S.; Song, Y.; Cohen, P.; Friedman, J. M.; Attie,
- [86129] A. D.: Loss of stearyl-CoA desaturase-1 function protects mice against
- [86130] adiposity. *Proc. Nat. Acad. Sci.* 99: 11482-11486, 2002.
- [86131] 15110.
- [86132] Thiede, M. A.; Ozols, J.; Strittmatter, P.: Construction and sequence
- [86133] of cDNA for rat liver stearyl coenzyme A desaturase. *J. Biol. Chem.* 261:
- [86134] 13230-13235, 1986.

- [86135] 15111.
- [86136] Zhang, L.; Ge, L.; Parimoo, S.; Stenn, K.; Prouty, S. M.: Human
man
- [86137] stearyl-CoA desaturase: alternative transcripts generated
from a
- [86138] single gene by usage of tandem polyadenylation sites.
Biochem. J. 340:
- [86139] 255–264, 1999.
- [86140] 15112.
- [86141] Zheng, Y.; Eilertsen, K. J.; Ge, L.; Zhang, L.; Sundberg, J. P.;
- [86142] Prouty, S. M.; Stenn, K. S.; Parimoo, S.: Scd1 is expressed
in sebaceous
- [86143] glands and is disrupted in the asebia mouse. (Letter) Na-
ture Genet. 23:
- [86144] 268–270, 1999.
- [86145] 15113.
- [86146] Harding, H. P.; Zeng, H.; Zhang, Y.; Jungries, R.; Chung, P.;
Plesken,
- [86147] H.; Sabatini, D. D.; Ron, D.: Diabetes mellitus and exocrine
pancreatic
- [86148] dysfunction in Perk $-/-$ mice reveals a role for transla-

tional control

[86149] in secretory cell survival. *Molec. Cell* 7: 1153–1163, 2001.

[86150] 15114.

[86151] Harding, H. P.; Zhang, Y.; Bertolotti, A.; Zeng, H.; Ron, D.:

[86152] Perk is essential for translational regulation and cell survival during

[86153] the unfolded protein response. *Molec. Cell* 5: 897–904, 2000.

[86154] 15115.

[86155] Duh, F.-M.; Latif, F.; Weng, Y.; Geil, L.; Modi, W.; Stackhouse,

[86156] T.; Matsumura, F.; Duan, D. R.; Linehan, W. M.; Lerman, M. I.; Gnarr,

[86157] J. R.: cDNA cloning and expression of the human homolog of the sea

[86158] urchin fascin and *Drosophila* singed genes which encodes an actin-bundling

[86159] protein. *DNA Cell Biol.* 13: 821–827, 1994.

[86160] 15116.

[86161] Mosialos, G.; Birkenbach, M.; Ayehunie, S.; Matsumura, F.; Pinkus,

- [86162] G. S.; Kieff, E.; Langhoff, E.: Circulating human dendritic cells
- [86163] differentially express high levels of a 55-kd actin-bundling protein. Am. J. Path. 148: 593-600, 1996.
- [86164] 15117.
- [86165] Ono, S.; Yamakita, Y.; Yamashiro, S.; Matsudaira, P. T.; Gnarra, J. R.; Obinata, T.; Matsumura, F.: Identification of an actin binding region and a protein kinase C phosphorylation site on human fascin. J. Biol. Chem. 272: 2527-2533, 1997.
- [86166] 15118.
- [86167] Pinkus, G. S.; Pinkus, J. L.; Langhoff, E.; Matsumura, F.; Yamashiro, S.; Mosialos, G.; Said, J. W.: Fascin, a sensitive new marker for Reed-Sternberg cells of Hodgkin's disease: evidence for a dendritic or B cell derivation? Am. J. Path. 150: 543-562, 1997.

- [86175] 15119.
- [86176] Sonderbye, L.; Magerstadt, R.; Blatman, R. N.; Preffer, F. I.;
- [86177] Langhoff, E.: Selective expression of human fascin (p55)
by dendritic
- [86178] leukocytes. Adv. Exp. Med. Biol. 471: 41–46, 1997.
- [86179] 15120.
- [86180] Yamakita, Y.; Ono, S.; Matsumura, F.; Yamashiro, S.: Phos-
phorylation
- [86181] of human fascin inhibits its actin binding and bundling
activities. J.
- [86182] Biol. Chem. 271: 12632–12638, 1996.
- [86183] 15121.
- [86184] Yamashiro–Matsumura, S.; Matsumura, F.: Intracellular lo-
calization
- [86185] of the 55–kD actin–bundling protein in cultured cells:
spatial relationships
- [86186] with actin, alpha–actinin, tropomyosin, and fimbrin. J. Cell
Biol. 103:
- [86187] 631–640, 1986.
- [86188] 15122.

- [86189] Yamashiro–Matsumura, S.; Matsumura, F.: Purification and characterization
- [86190] of an F–actin–bundling 55–kilodalton protein from HeLa cells. J.
- [86191] Biol. Chem. 260: 5087–5097, 1985.
- [86192] 15123.
- [86193] Gibbs, P. E. M.; McGregor, W. G.; Maher, V. M.; Nisson, P.; Lawrence,
- [86194] C. W.: A human homolog of the *Saccharomyces cerevisiae* REV3 gene,
- [86195] which encodes the catalytic subunit of DNA polymerase zeta. Proc.
- [86196] Nat. Acad. Sci. 95: 6876–6880, 1998.
- [86197] 15124.
- [86198] Morelli, C.; Mungall, A. J.; Negrini, M.; Barbanti–Brodano, G.;
- [86199] Croce, C. M.: Alternative splicing, genomic structure, and fine chromosome
- [86200] localization of REV3L. Cytogenet. Cell Genet. 83: 18–20, 1998.
- [86201] 15125.

- [86202] Xiao, W.; Lechler, T.; Chow, B. L.; Fontanie, T.; Agustus, M.;
- [86203] Carter, K. C.; Wei, Y.-F.: Identification, chromosomal mapping and
- [86204] tissue-specific expression of hREV3 encoding a putative human DNA
- [86205] polymerase zeta. *Carcinogenesis* 19: 945-949, 1998.
- [86206] 15126.
- [86207] Kahn, M. L.; Zheng, Y.-W.; Huang, W.; Bigornia, V.; Zeng, D.; Moff,
- [86208] S.; Farese, R. V., Jr.; Tam, C.; Coughlin, S. R.: A dual thrombin
- [86209] receptor system for platelet activation. *Nature* 394: 690-694, 1998.
- [86210] 15127.
- [86211] Xu, W.-F.; Andersen, H.; Whitmore, T. E.; Presnell, S. R.; Yee,
- [86212] D. P.; Ching, A.; Gilbert, T.; Davie, E. W.; Foster, D. C.: Cloning
- [86213] and characterization of human protease-activated receptor 4. *Proc.*
- [86214] *Nat. Acad. Sci.* 95: 6642-6646, 1998.

[86215] 15128.

[86216] Santoro, B.; Grant, S. G. N.; Bartsch, D.; Kandel, E. R.: In-
teractive

[86217] cloning with the SH3 domain of N-src identifies a new
brain specific

[86218] ion channel protein, with homology to Eag and cyclic nu-
cleotide-gated

[86219] channels. Proc. Nat. Acad. Sci. 94: 14815-14820, 1997.

[86220] 15129.

[86221] Santoro, B.; Liu, D. T.; Yao, H.; Bartsch, D.; Kandel, E. R.;
Siegelbaum,

[86222] S. A.; Tibbs, G. R.: Identification of a gene encoding a hy-
perpolarization-activated

[86223] pacemaker channel of brain. Cell 93: 717-729, 1998.

[86224] 15130.

[86225] Wainger, B. J.; DeGennaro, M.; Santoro, B.; Siegelbaum, S.
A.;

[86226] Tibbs, G. R.: Molecular mechanism of cAMP modulation of
HCN pacemaker

[86227] channels. Nature 411: 805-810, 2001.

[86228] 15131.

- [86229] Ludwig, A.; Zong, X.; Stieber, J.; Hullin, R.; Hofmann, F.; Biel,
- [86230] M.: Two pacemaker channels from human heart with profoundly different
- [86231] activation kinetics. EMBO J. 18: 2323–2329, 1999.
- [86232] 15132.
- [86233] Casari, G.; De Fusco, M.; Ciarmatori, S.; Zeviani, M.; Mora, M.;
- [86234] Fernandez, P.; De Michele, G.; Filla, A.; Coccozza, S.; Marconi, R.;
- [86235] Durr, A.; Fontaine, B.; Ballabio, A.: Spastic paraplegia and OXPHOS
- [86236] impairment caused by mutations in paraplegin, a nuclear-encoded mitochondrial
- [86237] metalloprotease. Cell 93: 973–983, 1998.
- [86238] 15133.
- [86239] De Michele, G.; De Fusco, M.; Cavalcanti, F.; Filla, A.; Marconi,
- [86240] R.; Volpe, G.; Monticelli, A.; Ballabio, A.; Casari, G.; Coccozza,
- [86241] S.: A new locus for autosomal recessive hereditary spastic paraplegia

- [86242] maps to chromosome 16q24.3. *Am. J. Hum. Genet.* 63: 135–139, 1998.
- [86243] 15134.
- [86244] Settasatian, C.; Whitmore, S. A.; Crawford, J.; Bilton, R. L.;
- [86245] Cleton-Jansen, A.-M.; Sutherland, G. R.; Callen, D. F.: Ge-
nomic structure
- [86246] and expression analysis of the spastic paraplegia gene,
SPG7. *Hum.*
- [86247] *Genet.* 105: 139–144, 1999.
- [86248] 15135.
- [86249] Nykjaer, A.; Dragun, D.; Walther, D.; Vorum, H.; Jacobsen,
C.;
- [86250] Herz, J.; Melsen, F.; Christensen, E. I.; Willnow, T. E.: An
endocytic
- [86251] pathway essential for renal uptake and activation of the
steroid 25-(OH)
- [86252] vitamin D3. *Cell* 96: 507–515, 1999.
- [86253] 15136.
- [86254] Quelle, D. E.; Ashmun, R. A.; Hannon, G. J.; Rehberger, P.
A.;
- [86255] Trono, D.; Richter, K. H.; Walker, C.; Beach, D.; Sherr, C. J.;

Serrano,

[86256] M.: Cloning and characterization of murine p16(INK4a) and p15(INK4b)

[86257] genes. *Oncogene* 11: 635–645, 1995.

[86258] 15137.

[86259] Quelle, D. E.; Cheng, M.; Ashmun, R. A.; Sherr, C. J.: Cancer-associated

[86260] mutations at the INK4a locus cancel cell cycle arrest by p16(INK4a)

[86261] but not by the alternative reading frame protein p19(ARF). *Proc.*

[86262] *Nat. Acad. Sci.* 94: 669–673, 1997.

[86263] 15138.

[86264] Quelle, D. E.; Zindy, F.; Ashmun, R. A.; Sherr, C. J.: Alternative

[86265] reading frames of the INK4a tumor suppressor gene encode two unrelated

[86266] proteins capable of inducing cell cycle arrest. *Cell* 83: 993–1000,

[86267] 1995.

[86268] 15139.

- [86269] Ranade, K.; Hussussian, C. J.; Sikorski, R. S.; Varmus, H. E.;
- [86270] Goldstein, A. M.; Tucker, M. A.; Serrano, M.; Hannon, G. J.;
Beach,
- [86271] D.; Dracopoli, N. C.: Mutations associated with familial
melanoma
- [86272] impair p16(INK4) function. *Nature Genet.* 10: 114–116,
1995.
- [86273] 15140.
- [86274] Schmitt, C. A.; Fridman, J. S.; Yang, M.; Lee, S.; Baranov,
E.;
- [86275] Hoffman, R. M.; Lowe, S. W.: A senescence program con-
trolled by p53
- [86276] and p16–INK4a contributes to the outcome of cancer
therapy. *Cell* 109:
- [86277] 335–346, 2002.
- [86278] 15141.
- [86279] Serrano, J.; Goebel, S. U.; Peghini, P. L.; Lubensky, I. A.;
- Gibril,
- [86280] F.; Jensen, R. T.: Alterations in the p16INK4a/CDKN2A tu-
mor suppressor
- [86281] gene in gastrinomas. *J. Clin. Endocr. Metab.* 85:
4146–4156, 2000.

[86282] 15142.

[86283] Serrano, M.; Hannon, G. J.; Beach, D.: A new regulatory motif

[86284] in cell-cycle control causing specific inhibition of cyclin D/CDK4. Nature 366:

[86285] 704–707, 1993.

[86286] 15143.

[86287] Serrano, M.; Lee, H.-W.; Chin, L.; Cordon-Cardo, C.; Beach, D.;

[86288] DePinho, R. A.: Role of the INK4a locus in tumor suppression and

[86289] cell mortality. Cell 85: 27–37, 1996.

[86290] 15144.

[86291] Sharpless, N. E.; Bardeesy, N.; Lee, K.-H.; Carrasco, D.; Castrillon,

[86292] D. H.; Aguirre, A. J.; Wu, E. A.; Horner, J. W.; DePinho, R. A.:

[86293] Loss of p16(Ink4a) with retention of p19(Arf) predisposes mice to

[86294] tumorigenesis. Nature 413: 86–91, 2001.

[86295] 15145.

- [86296] Smith–Sorensen, B.; Hovig, E.: CDKN2A (p16–INK4A) so-
matic and
- [86297] germline mutations. Hum. Mutat. 7: 294–303, 1996.
- [86298] 15146.
- [86299] Stone, S.; Jiang, P.; Dayananth, P.; Tavtigian, S. W.;
Katcher,
- [86300] H.; Parry, D.; Peters, G.; Kamb, A.: Complex structure and
regulation
- [86301] of the p16(MTS1) locus. Cancer Res. 55: 2988–2994,
1995.
- [86302] 15147.
- [86303] Stott, F. J.; Bates, S.; James, M. C.; McConnell, B. B.; Star-
borg,
- [86304] M.; Brookes, S.; Palmero, I.; Ryan, K.; Hara, E.; Vousden, K.
H.;
- [86305] Peters, G.: The alternative product from the human
CDKN2A locus,
- [86306] p14(ARF), participates in a regulatory feedback loop with
p53 and
- [86307] MDM2. EMBO J. 17: 5001–5014, 1998.
- [86308] 15148.

- [86309] Tsutsumi, M.; Tsai, Y. C.; Gonzalgo, M. L.; Nichols, P. W.; Jones,
[86310] P. A.: Early acquisition of homozygous deletions of
p16/p19 during
[86311] squamous cell carcinogenesis and genetic mosaicism in
bladder cancer. *Oncogene* 17:
[86312] 3021–3027, 1998.
[86313] 15149.
[86314] van der Velden, P. A.; Sandkuijl, L. A.; Bergman, W.; Hille,
E.
[86315] T. M.; Frants, R. R.; Gruis, N. A.: A locus linked to p16
modifies
[86316] melanoma risk in Dutch familial atypical multiple mole
melanoma (FAMMM)
[86317] syndrome families. *Genome Res.* 9: 575–580, 1999.
[86318] 15150.
[86319] Vasen, H. F. A.; Gruis, N. A.; Frants, R. R.; van der Velden,
[86320] P. A.; Hille, E. T. M.; Bergman, W.: Risk of developing pan-
creatic
[86321] cancer in families with familial atypical multiple mole
melanoma associated

- [86322] with a specific 19 deletion of p16(p16-Leiden). *Int. J. Cancer* 87:
809–811, 2000.
- [86323] 15151.
- [86324] Wainwright, B.: Familial melanoma and p16—a hung jury.
Nature Genet. 8: 3–5, 1994.
- [86325] 15152.
- [86326] Walker, G. J.; Hussussian, C. J.; Flores, J. F.; Glendening, J.
M.; Haluska, F. G.; Dracopoli, N. C.; Hayward, N. K.; Fountain, J.
W.: Mutations of the CDKN2/p16(INK4) gene in Australian
melanoma kindreds. *Hum. Mol. Genet.* 4: 1845–1852, 1995.
- [86327] 15153.
- [86328] Wiest, J. S.; Franklin, W. A.; Otstot, J. T.; Forbey, K.;
Varella-Garcia,
M.; Rao, K.; Drabkin, H.; Gemmill, R.; Ahrent, S.; Sidransky, D.;
Saccomanno, G.; Fountain, J. W.; Anderson, M. W.: Identification

- [86336] of a novel region of homozygous deletion on chromosome 9p in squamous
- [86337] cell carcinoma of the lung: the location of a putative tumor suppressor
- [86338] gene. *Cancer Res.* 57: 1–6, 1997.
- [86339] 15154.
- [86340] Williamson, M. P.; Elder, P. A.; Shaw, M. E.; Devlin, J.; Knowles,
- [86341] M. A.: p16 (CDKN2) is a major deletion target at 9p21 in bladder
- [86342] cancer. *Hum. Molec. Genet.* 4: 1569–1577, 1995.
- [86343] 15155.
- [86344] Woloschak, M.; Yu, A.; Xiao, J.; Post, K. D.: Frequent loss of
- [86345] the P16(INK4a) gene product in human pituitary tumors. *Cancer Res.* 56:
- [86346] 2493–2496, 1996.
- [86347] 15156.
- [86348] Zhang, Y.; Xiong, Y.: Mutations in human ARF exon 2 disrupt its
- [86349] nucleolar localization and impair its ability to block nu-

clear export

[86350] of MDM2 and p53. *Molec. Cell* 3: 579–591, 1999.

[86351] 15157.

[86352] Zhang, Y.; Xiong, Y.; Yarbrough, W. G.: ARF promotes
MDM2 degradation

[86353] and stabilizes p53: ARF–INK4a locus deletion impairs both
the Rb and

[86354] p53 tumor suppression pathways. *Cell* 92: 725–734,
1998.

[86355] 15158.

[86356] Zhu, G.; Duffy, D. L.; Eldridge, A.; Grace, M.; Mayne, C.;
O'Gorman,

[86357] L.; Aitken, J. F.; Neale, M. C.; Hayward, N. K.; Green, A. C.;
Martin,

[86358] N. G.: A major quantitative–trait locus for mole density is
linked

[86359] to the familial melanoma gene CDKN2A: a maximum–like–
lihood combined

[86360] linkage and association analysis in twins and their sibs.
Am. J.

[86361] *Hum. Genet.* 65: 483–492, 1999.

[86362] 15159.

- [86363] Pennica, D.; Swanson, T. A.; Shaw, K. J.; Kuang, W.-J.; Gray, C.
- [86364] L.; Beatty, B. G.; Wood, W. I.: Human cardiotrophin-1: protein and
- [86365] gene structure, biological and binding activities, and chromosomal
- [86366] localization. *Cytokine* 8: 183-189, 1996.
- [86367] 15160.
- [86368] Bonnefont, J.-P.; Taroni, F.; Cavadini, P.; Cepanec, C.; Brivet,
- [86369] M.; Saudubray, J.-M.; Leroux, J.-P.; Demaugre, F.: Molecular analysis
- [86370] of carnitine palmitoyltransferase II deficiency with hepatocardiomyopathy
- [86371] expression. *Am. J. Hum. Genet.* 58: 971-978, 1996.
- [86372] 15161.
- [86373] Candia, A. F.; Hu, J.; Crosby, J.; Lalley, P. A.; Noden, D.; Nadeau,
- [86374] J. H.; Wright, C. V. E.: Mox-1 and Mox-2 define a novel homeobox
- [86375] gene subfamily and are differentially expressed during early mesodermal

- [86376] patterning in mouse embryos. Development 116: 1123–1136, 1992.
- [86377] 15162.
- [86378] Gorski, D. H.; LePage, D. F.; Patel, C. V.; Copeland, N. G.; Jenkins, N. A.; Walsh, K.: Molecular cloning of a diverged homeobox gene that is rapidly down-regulated during the G0/G1 transition in vascular smooth muscle cells. Molec. Cell Biol. 13: 3722–3733, 1993.
- [86382] 15163.
- [86383] Grigoriou, M.; Kastrinaki, M.–C.; Modi, W. S.; Theodorakis, K.;
- [86384] Mankoo, B.; Pachnis, V.; Karagogeos, D.: Isolation of the human MOX2 homeobox gene and localization to chromosome 7p22.1–p21.3. Genomics 26: 550–555, 1995.
- [86387] 15164.
- [86388] LePage, D. F.; Altomare, D. A.; Testa, J. R.; Walsh, K.: Molecular

[86389] cloning and localization of the human GAX gene to 7p21.

Genomics 24:

[86390] 535–540, 1994.

[86391] 15165.

[86392] Mankoo, B. S.; Collins, N. S.; Ashby, P.; Grigorieva, E.;

Pevny,

[86393] L. H.; Candia, A.; Wright, C. V. E.; Rigby, P. W. J.; Pachnis,

V.

[86394] : Mox2 is a component of the genetic hierarchy controlling
limb muscle

[86395] development. Nature 400: 69–73, 1999.

[86396] 15166.

[86397] Hayashi, Y. K.; Chou, F.-L.; Engvall, E.; Ogawa, M.; Mat-

suda, C.;

[86398] Hirabayashi, S.; Yokochi, K.; Ziober, B. L.; Kramer, R. H.;

Kaufman,

[86399] S. J.; Ozawa, E.; Goto, Y.; Nonaka, I.; Tsukahara, T.; Wang,

J.; Hoffman,

[86400] E. P.; Arahata, K.: Mutations in the integrin alpha-7 gene
cause

[86401] congenital myopathy. Nature Genet. 19: 94–97, 1998.

[86402] 15167.

[86403] Mayer, U.; Saher, G.; Fassler, R.; Bornemann, A.; Echter-
meyer,

[86404] F.; von der Mark, H.; Miosge, N.; Poschl, E.; von der Mark,
K.: Absence

[86405] of integrin alpha-7 causes a novel form of muscular dys-
trophy. Nature

[86406] Genet. 17: 318-323, 1997.

[86407] 15168.

[86408] Wang, W.; Wu, W.; Desai, T.; Ward, D. C.; Kaufman, S. J.:
Localization

[86409] of the alpha-7 integrin gene (ITGA7) on human chromo-
some 12q13: clustering

[86410] of integrin and Hox genes implies parallel evolution of
these gene

[86411] families. Genomics 26: 563-570, 1995.

[86412] 15169.

[86413] Lindor, N. M.; Furuichi, Y.; Kitao, S.; Shimamoto, A.; Arndt,
C.;

[86414] Jalal, S.: Rothmund-Thomson syndrome due to RECQ4 he-
licase mutations:

- [86415] report and clinical and molecular comparisons with Bloom syndrome
- [86416] and Werner syndrome. *Am. J. Med. Genet.* 90: 223–228, 2000.
- [86417] 15170.
- [86418] Puranam, K. L.; Blackshear, P. J.: Cloning and characterization
- [86419] of RECQL, a potential human homologue of the *Escherichia coli* DNA
- [86420] helicase RecQ. *J. Biol. Chem.* 269: 29838–29845, 1994.
- [86421] 15171.
- [86422] Puranam, K. L.; Kennington, E.; J.-Sait, S. N.; Shows, T. B.; Rochelle,
- [86423] J. M.; Seldin, M. F.; Blackshear, P. J.: Chromosomal localization
- [86424] of the gene encoding the human DNA helicase RECQL and its mouse homologue. *Genomics* 26:
- [86425] 595–598, 1995.
- [86426] 15172.
- [86427] Hagen, G.; Muller, S.; Beato, M.; Suske, G.: Cloning by recognition

- [86428] site screening of two novel GT box binding proteins: a family of Sp1
- [86429] related genes. *Nucleic Acids Res.* 20: 5519–5525, 1992.
- [86430] 15173.
- [86431] Lamour, V.; Lecluse, Y.; Desmaze, C.; Spector, M.; Bode-scot, M.;
- [86432] Aurias, A.; Osley, M. A.; Lipinski, M.: A human homolog of the S.
- [86433] cerevisiae HIR1 and HIR2 transcriptional repressors cloned from the
- [86434] DiGeorge syndrome critical region. *Hum. Molec. Genet.* 4: 791–799,
- [86435] 1995.
- [86436] 15174.
- [86437] Llevadot, R.; Scambler, P.; Estivill, X.; Pritchard, M.: Ge-nomic
- [86438] organization of TUPLE1/HIRA: a gene implicated in DiGe-orge syndrome. *Mammalian*
- [86439] *Genome* 7: 911–914, 1996.
- [86440] 15175.
- [86441] Lorain, S.; Quivy, J.-P.; Monier-Gavelle, F.; Scamps, C.;

Lecluse,

[86442] Y.; Almouzni, G.; Lipinski, M.: Core histones and HIRIP3, a novel

[86443] histone-binding protein, directly interact with WD repeat protein

[86444] HIRA. *Molec. Cell. Biol.* 18: 5546–5556, 1998.

[86445] 15176.

[86446] Magnaghi, P.; Roberts, C.; Lorain, S.; Lipinski, M.; Scambler,

[86447] P. J.: HIRA, a mammalian homologue of *Saccharomyces cerevisiae* transcriptional

[86448] co-repressors, interacts with Pax3. *Nature Genet.* 20: 74–77, 1998.

[86449] 15177.

[86450] Mattei, M.-G.; Halford, S.; Scambler, P. J.: Mapping of the TUPLE1

[86451] gene to mouse chromosome 16A-B1. *Genomics* 23: 717–718, 1994.

[86452] 15178.

[86453] Roberts, C.; Daw, S. C. M.; Halford, S.; Scambler, P. J.: Cloning

[86454] and developmental expression analysis of chick Hira

(Chira), a candidate

[86455] gene for DiGeorge syndrome. Hum. Molec. Genet. 6:
237-245, 1997.

[86456] 15179.

[86457] Wilming, L. G.; Snoeren, C. A. S.; van Rijswijk, A.;
Grosveld,

[86458] F.; Meijers, C.: The murine homologue of HIRA, a DiGe-
orge syndrome

[86459] candidate gene, is expressed in embryonic structures af-
fected in human

[86460] CATCH22 patients. Hum. Molec. Genet. 6: 247-258, 1997.

[86461] 15180.

[86462] Akao, Y.; Matsuda, Y.: Identification and chromosome
mapping of

[86463] the mouse homologue of the human gene (DDX6) that en-
codes a putative

[86464] RNA helicase of the DEAD box protein family. Cytogenet.
Cell Genet. 75:

[86465] 38-44, 1996.

[86466] 15181.

[86467] Akao, Y.; Seto, M.; Takahashi, T.; Kubonishi, I.; Miyoshi, I.;

- [86468] Nakazawa, S.; Tsujimoto, Y.; Croce, C. M.; Ueda, R.: Molecular cloning
[86469] of the chromosomal breakpoint of a B-cell lymphoma with
the t(11;14)(q23;q32)
[86470] chromosome translocation. Cancer Res. 51: 1574–1576,
1991.
[86471] 15182.
- [86472] Akao, Y.; Seto, M.; Yamamoto, K.; Iida, S.; Nakazawa, S.;
Inazawa,
[86473] J.; Abe, T.; Takahashi, T.; Ueda, R.: The RCK gene associated with
[86474] t(11;14) translocation is distinct from the MLL/ALL-1 gene
with t(4;11)
[86475] and t(11;19) translocations. Cancer Res. 52: 6083–6087,
1992.
[86476] 15183.
- [86477] Akao, Y.; Tsujimoto, Y.; Finan, J.; Nowell, P. C.; Croce, C.
M.
[86478] : Molecular characterization of a t(11;14)(q23;q32) chromosome translocation
[86479] in a B-cell lymphoma. Cancer Res. 50: 4856–4859, 1990.
[86480] 15184.

- [86481] Lu, D.; Yunis, J. J.: Cloning, expression and localization of
- [86482] an RNA helicase gene from a human lymphoid cell line with chromosomal
- [86483] breakpoint 11q23.3. *Nucleic Acids Res.* 20: 1967–1972, 1992.
- [86484] 15185.
- [86485] Seto, M.; Yamamoto, K.; Takahashi, T.; Ueda, R.: Cloning and expression
- [86486] of a murine cDNA homologous to the human RCK/P54, a lymphoma-linked
- [86487] chromosomal translocation junction gene on 11q23. *Gene* 166: 293–296,
- [86488] 1995.
- [86489] 15186.
- [86490] Tunnacliffe, A.; Perry, H.; Radice, P.; Budarf, M. L.; Emanuel,
- [86491] B. S.: A panel of sequence tagged sites for chromosome band 11q23. *Genomics* 17:
- [86492] 744–747, 1993.
- [86493] 15187.
- [86494] Ricker, K.; Moxley, R. T.; Rohkamm, R.: Rippling muscle

disease. Arch.

[86495] Neurol. 46: 405–408, 1989.

[86496] 15188.

[86497] Dumon, K. R.; Ishii, H.; Fong, L. Y. Y.; Zanesi, N.; Fidanza, V.;

[86498] Mancini, R.; Vecchione, A.; Baffa, R.; Trapasso, F.; During, M. J.;

[86499] Huebner, K.; Croce, C. M.: FHIT gene therapy prevents tumor development

[86500] in Fhit-deficient mice. Proc. Nat. Acad. Sci. 98: 3346–3351, 2001.

[86501] 15189.

[86502] Ikeda, A.; Zheng, Q. Y.; Zuberi, A. R.; Johnson, K. R.; Nagert,

[86503] J. K.; Nishina, P. M.: Microtubule-associated protein 1A is a modifier

[86504] of tubby hearing (moth1). Nature Genet. 30: 401–405, 2002.

[86505] 15190.

[86506] Roy, S. K.; Hu, J.; Meng, Q.; Xia, Y.; Shapiro, P. S.; Reddy, S.

[86507] P. M.; Platanius, L. C.; Lindner, D. J.; Johnson, P. F.;

Pritchard,

[86508] C.; Pages, G.; Pouyssegur, J.; Kalvakolanu, D. V.: MEKK1 plays a

[86509] critical role in activating the transcription factor C/EBP-beta-dependent

[86510] gene expression in response to IFN-gamma. Proc. Nat. Acad. Sci. 99:

[86511] 7945-7950, 2002.

[86512] 15191.

[86513] Lindsten, T.; Ross, A. J.; King, A.; Zong, W.-X.; Rathmell, J.

[86514] C.; Shiels, H. A.; Ulrich, E.; Waymire, K. G.; Mahar, P.; Frauwirth,

[86515] K.; Chen, Y.; Wei, M.; and 9 others: The combined functions of

[86516] proapoptotic Bcl-2 family members Bak and Bax are essential for normal

[86517] development of multiple tissues. Molec. Cell 6: 1389-1399, 2000.

[86518] 15192.

[86519] Matikainen, T.; Perez, G. I.; Jurisicova, A.; Pru, J. K.; Schlezinger,

[86520] J. J.; Ryu, H.-Y.; Laine, J.; Sakai, T.; Korsmeyer, S. J.;

Casper,

[86521] R. F.; Sherr, D. H.; Tilly, J. L.: Aromatic hydrocarbon receptor-driven

[86522] Bax gene expression is required for premature ovarian failure caused

[86523] by biohazardous environmental chemicals. *Nature Genet.* 28: 355–360,

[86524] 2001.

[86525] 15193.

[86526] Wei, M. C.; Zong, W.-X.; Cheng, E. H.-Y.; Lindsten, T.; Panoutsakopoulou,

[86527] V.; Ross, A. J.; Roth, K. A.; MacGregor, G. R.; Thompson, C. B.; Korsmeyer,

[86528] S. J.: Proapoptotic BAX or BAK: a requisite gateway to mitochondrial

[86529] dysfunction and death. *Science* 292: 727–730, 2001.

[86530] 15194.

[86531] Dasari, V. R.; Sandhu, A. K.; Mills, D. C. B.; Athwal, R. S.; Kunapuli,

[86532] S. P.: Mapping of the P2U purinergic receptor gene to human chromosome

[86533] 11q13.5–14.1. *Somat. Cell Molec. Genet.* 22: 75–79,

1996.

[86534] 15195.

[86535] Katzur, A. C.; Koshimizu, T.-A.; Tomic, M.; Schultze-Mosgau, A.;

[86536] Ortmann, O.; Stojilkovic, S. S.: Expression and responsiveness of

[86537] P2Y2 receptors in human endometrial cancer cell lines. J. Clin. Endocr.

[86538] Metab. 84: 4085-4091, 1999.

[86539] 15196.

[86540] Parr, C. E.; Sullivan, D. M.; Paradiso, A. M.; Lazarowski, E. R.;

[86541] Burch, L. H.; Olsen, J. C.; Erb, L.; Weisman, G. A.; Boucher, R. C.;

[86542] Turner, J. T.: Cloning and expression of a human P(2U) nucleotide

[86543] receptor, a target for cystic fibrosis pharmacotherapy. Proc. Nat.

[86544] Acad. Sci. 91: 3275-3279, 1994.

[86545] 15197.

[86546] Somers, G. R.; Hammet, F.; Woollatt, E.; Richards, R. I.;

Southey,

[86547] M. C.; Venter, D. J.: Chromosomal localization of the human P2Y(6)

[86548] purinoceptor gene and phylogenetic analysis of the P2y purinoceptor

[86549] family. Genomics 44: 127–130, 1997.

[86550] 15198.

[86551] Tai, C.-J.; Kang, S. K.; Cheng, K. W.; Choi, K.-C.; Nathwani, P.

[86552] S.; Leung, P. C. K.: Expression and regulation of P2U–purinergic

[86553] receptor in human granulosa–luteal cells. J. Clin. Endocr. Metab. 85:

[86554] 1591–1597, 2000.

[86555] 15199.

[86556] Santamarina–Fojo, S.; Peterson, K.; Knapper, C.; Qiu, Y.; Freeman,

[86557] L.; Cheng, J.-F.; Osorio, J.; Remaley, A.; Yang, X.-P.; Hauschild,

[86558] C.; Prades, C.; Chimini, G.; Blackmon, E.; Francois, T.; Duverger,

[86559] N.; Rubin, E. M.; Rosier, M.; Deneffe, P.; Fredrickson, D. S.;

Brewer,

- [86560] H. B., Jr.: Complete genomic sequence of the human ABCA1 gene: analysis
[86561] of the human and mouse ATP-binding cassette A promoter. Proc. Nat.
[86562] Acad. Sci. 97: 7987–7992, 2000. Note: Erratum: Proc. Nat.
Acad. Sci.
[86563] 99: 1098 only, 2002.
[86564] 15200.
- [86565] Szakacs, G.; Langmann, T.; Ozvegy, C.; Orso, E.; Schmitz, G.;
[86566] Varadi, A.; Sarkadi, B.: Characterization of the ATPase cycle of
[86567] human ABCA1: implications for its function as a regulator rather than
[86568] an active transporter. Biochem. Biophys. Res. Commun. 288: 1258–1264,
[86569] 2001.
[86570] 15201.
- [86571] Utech, M.; Hobbel, G.; Rust, S.; Reinecke, H.; Assmann, G.; Walter,
[86572] M.: Accumulation of RhoA, RhoB, RhoG, and Rac1 in fi-

broblasts from

[86573] Tangier disease subjects suggests a regulatory role of Rho family

[86574] proteins in cholesterol efflux. Biochem. Biophys. Res. Commun. 280:

[86575] 229–236, 2001.

[86576] 15202.

[86577] Zhao, L.-X.; Zhou, C.-J.; Tanaka, A.; Nakata, M.; Hirabayashi,

[86578] T.; Amachi, T.; Shioda, S.; Ueda, K.; Inagaki, N.: Cloning, characterization

[86579] and tissue distribution of the rat ATP-binding cassette (ABC) transporter

[86580] ABC2/ABCA2. Biochem J. 350: 865–872, 2000.

[86581] 15203.

[86582] Zwarts, K. Y.; Clee, S. M.; Zwinderman, A. H.; Engert, J. C.;

[86583] Singaraja, R.; Loubser, O.; James, E.; Roomp, K.; Hudson, T. J.; Jukema,

[86584] J. W.; Kastelein, J. J. P.; Hayden, M. R.: ABCA1 regulatory variants

[86585] influence coronary artery disease independent of effects

on plasma

[86586] lipid levels. Clin. Genet. 61: 115–125, 2002.

[86587] 15204.

[86588] Ing, Y. L.; Leung, I. W. L.; Heng, H. H. Q.; Tsui, L.-C.; Las-sam,

[86589] N. J.: MLK-3: identification of a widely-expressed protein kinase

[86590] bearing an SH3 domain and a leucine zipper-basic region domain. Oncogene 9:

[86591] 1745–1750, 1994.

[86592] 15205.

[86593] Bernard, O. A.; Mauchauffe, M.; Mecucci, C.; Van Den Berghe, H.;

[86594] Berger, R.: A novel gene, AF-1p, fused to HRX in t(1;11)(p32;q23),

[86595] is not related to AF-4, AF-9 nor ENL. Oncogene 9: 1039–1045, 1994.

[86596] 15206.

[86597] Fazioli, F.; Minichiello, L.; Matoska, V.; Castagnino, P.; Miki,

[86598] T.; Wong, W. T.; Di Fiore, P. P.: Eps8, a substrate for the

epidermal

[86599] growth factor receptor kinase, enhances EGF-dependent mitogenic signals.

[86600] EMBO J. 12: 3799–3808, 1993.

[86601] 15207.

[86602] Fazioli, F.; Minichiello, L.; Matoskova, B.; Wong, W. T.; Di Fiore,

[86603] P. P.: eps 15, a novel tyrosine kinase substrate, exhibits transforming

[86604] activity. Molec. Cell. Biol. 13: 5814–5828, 1993.

[86605] 15208.

[86606] Fazioli, F.; Wong, W. T.; Ullrich, S. J.; Sakaguchi, K.; Appella,

[86607] E.; Di Fiore, P. P.: The ezrin-like family of tyrosine kinase substrates:

[86608] receptor-specific pattern of tyrosine phosphorylation and relationship

[86609] to malignant transformation. Oncogene 8: 1335–1345, 1993.

[86610] 15209.

[86611] Wong, W. T.; Kraus, M. H.; Carlomagno, F.; Zelano, A.;

Druck, T.;

[86612] Croce, C. M.; Huebner, K.; Di Fiore, P. P.: The human
eps15 gene,

[86613] encoding a tyrosine kinase substrate, is conserved in evolution and

[86614] maps to 1p31-p32. *Oncogene* 9: 1591-1597, 1994.

[86615] 15210.

[86616] Frydman, J.; Nimmesgern, E.; Erdjument-Bromage, H.;
Wall, J. S.;

[86617] Tempst, P.; Hartl, F.-U.: Function in protein folding of
TRiC, a

[86618] cytosolic ring complex containing TCP-1 and structurally
related subunits. *EMBO*

[86619] *J.* 11: 4767-4778, 1992.

[86620] 15211.

[86621] Joly, E. C.; Seigny, G.; Todorov, I. T.; Bibor-Hardy, V.:
cDNA

[86622] encoding a novel TCP1-related protein. *Biochim. Biophys.*
Acta 1217:

[86623] 224-226, 1994.

[86624] 15212.

- [86625] Sevigny, G.; Joly, E. C.; Bibor-Hardy, V.; Lemieux, N.: Assignment
- [86626] of the human homologue of the mTRiC-P5 gene (TRiC5) to band 1q23 by
- [86627] fluorescence in situ hybridization. Genomics 22: 634-636, 1994.
- [86628] 15213.
- [86629] Sevigny, G.; Lemieux, N.; Steyaert, A.; Bibor-Hardy, V.: Structure
- [86630] of the gene coding for the mouse TRiC-P5 subunit of the cytosolic

- [86631] chaperonin TRiC. Genomics 31: 107–110, 1996.
- [86632] 15214.
- [86633] Walkley, N. A.; Demaine, A. G.; Malik, A. N.: Cloning,
structure
- [86634] and mRNA expression of human Cctg, which encodes the
chaperonin subunit
- [86635] CCT-gamma. Biochem. J. 313: 381–389, 1996.
- [86636] 15215.
- [86637] Hoenicka, J.; Vidal, L.; Morales, B.; Ampuero, I.; Jimenez-
Jimenez,
- [86638] F. J.; Berciano, J.; del Ser, T.; Jimenez, A.; Ruiz, P. G.; de
Yebeles,
- [86639] J. G.: Molecular findings in familial Parkinson disease in
Spain. Arch.
- [86640] Neurol. 59: 966–970, 2002.
- [86641] 15216.
- [86642] Kitada, T.; Asakawa, S.; Hattori, N.; Matsumine, H.; Yama-
mura,
- [86643] Y.; Minoshima, S.; Yokochi, M.; Mizuno, Y.; Shimizu, N.:
Mutations
- [86644] in the parkin gene cause autosomal recessive juvenile

parkinsonism. Nature 392:

[86645] 605–608, 1998.

[86646] 15217.

[86647] Matsumine, H.; Saito, M.; Shimoda–Matsubayashi, S.;
Tanaka, H.;

[86648] Ishikawa, A.; Nakagawa–Hattori, Y.; Yokochi, M.;
Kobayashi, T.; Igarashi,

[86649] S.; Takano, H.; Sanpei, K.; Koike, R.; Mori, H.; Kondo, T.;
Mizutani,

[86650] Y.; Schaffer, A. A.; Yamamura, Y.; Nakamura, S.; Kuzuhura,
S.; Tsuji,

[86651] S.; Mizuno, Y.: Localization of a gene for an autosomal re-
cessive

[86652] form of juvenile parkinsonism to chromosome 6q25.2–27.
Am. J. Hum.

[86653] Genet. 60: 588–596, 1997.

[86654] 15218.

[86655] Tassin, J.; Durr, A.; de Broucker, T.; Abbas, N.; Bonifati, V.;

[86656] De Michele, G.; Bonnet, A.–M.; Broussolle, E.; Pollak, P.;
Vidailhet,

[86657] M.; De Mari, M.; Marconi, R.; Medjbeur, S.; Filla, A.; Meco,

G.; Agid,

[86658] Y.; Brice, A.; French Parkinson's Disease Genetics Study Group;

[86659] European Consortium on Genetic Susceptibility in Parkinson's Disease

[86660] : Chromosome 6-linked autosomal recessive early-onset parkinsonism:

[86661] linkage in European and Algerian families, extension of the clinical

[86662] spectrum, and evidence of a small homozygous deletion in one family. Am.

[86663] J. Hum. Genet. 63: 88-94, 1998.

[86664] 15219.

[86665] Kozu, T.; Henrich, B.; Schafer, K. P.: Structure and expression

[86666] of the gene (HNRPA2B1) encoding the human hnRNP protein A2/B1. Genomics 25:

[86667] 365-371, 1995.

[86668] 15220.

[86669] Kordeli, E.; Lambert, S.; Bennett, V.: Ankyrin-G: a new ankyrin

[86670] gene with neural-specific isoforms localized at the axonal

initial

[86671] segment and node of Ranvier. J. Biol. Chem. 270:
2352–2359, 1995.

[86672] 15221.

[86673] Hsu, L. C.; Chang, W.-C.; Yoshida, A.: Human aldehyde
dehydrogenase

[86674] genes, ALDH7 and ALDH8: genomic organization and
gene structure comparison. Gene 189:

[86675] 89–94, 1997.

[86676] 15222.

[86677] Hsu, L. C.; Chang, W.-C.; Yoshida, A.: Cloning of a cDNA
encoding

[86678] human ALDH7, a new member of the aldehyde dehydro-
genase family. Gene 151:

[86679] 285–289, 1994.

[86680] 15223.

[86681] Aruga, J.; Yokota, N.; Hashimoto, M.; Furuichi, T.; Fukuda,
M.;

[86682] Mikoshiba, K.: A novel zinc finger protein, Zic, is involved
in neurogenesis,

[86683] especially in the cell lineage of cerebellar granule cells. J.

Neurochem. 63:

[86684] 1880–1890, 1994.

[86685] 15224.

[86686] Salero, E.; Perez–Sen, R.; Aruga, J.; Gimenez, C.; Zafra, F.:

[86687] Transcription factors Zic1 and Zic2 bind and transactivate
the apolipoprotein

[86688] E gene promoter. J. Biol. Chem. 276: 1881–1888, 2001.

[86689] 15225.

[86690] Yokota, N.; Aruga, J.; Takai, S.; Yamada, K.; Hamazaki, M.;
Iwase,

[86691] T.; Sugimura, H.; Mikoshiba, K.: Predominant expression
of human

[86692] Zic in cerebellar granule cell lineage and medulloblas-
toma. Cancer

[86693] Res. 56: 377–383, 1996.

[86694] 15226.

[86695] Bevins, C. L.; Jones, D. E.; Dutra, A.; Schaffzin, J.; Muenke,

[86696] M.: Human enteric defensin genes: chromosomal map po-
sition and a

[86697] model for possible evolutionary relationships. Genomics

31: 95–106,

[86698] 1996.

[86699] 15227.

[86700] Jones, D. E.; Bevins, C. L.: Defensin-6 mRNA in human Paneth cells:

[86701] implications for antimicrobial peptides in host defense of the human

[86702] bowel. FEBS Lett. 315: 187–192, 1993.

[86703] 15228.

[86704] Mallow, E. B.; Harris, A.; Salzman, N.; Russell, J. P.; DeBerardinis,

[86705] R. J.; Ruchelli, E.; Bevins, C. L.: Human enteric defensins: gene

[86706] structure and developmental expression. J. Biol. Chem. 271: 4038–4045,

[86707] 1996.

[86708] 15229.

[86709] Jones, D. E.; Bevins, C. L.: Paneth cells of the human small intestine

[86710] express an antimicrobial peptide gene. J. Biol. Chem. 267: 23216–23225,

- [86711] 1992.
- [86712] 15230.
- [86713] Porter, E. M.; Liu, L.; Oren, A.; Anton, P. A.; Ganz, T.: Localization
- [86714] of human intestinal defensin 5 in Paneth cell granules. Infect. Immun. 65:
- [86715] 2389–2395, 1997.
- [86716] 15231.
- [86717] Quayle, A. J.; Porter, E. M.; Nussbaum, A. A.; Wang, Y. M.; Brabec,
- [86718] C.; Yip, K.-P.; Mok, S. C.: Gene expression, immunolocalization,
- [86719] and secretion of human defensin-5 in human female reproductive tract. Am.
- [86720] J. Path. 152: 1247–1258, 1998.
- [86721] 15232.
- [86722] Matsuzono, Y.; Kinoshita, N.; Tamura, S.; Shimozawa, N.; Hamasaki,
- [86723] M.; Ghaedi, K.; Wanders, R. J. A.; Suzuki, Y.; Kondo, N.; Fujiki,
- [86724] Y.: Human PEX19: cDNA cloning by functional comple-

mentation, mutation

[86725] analysis in a patient with Zellweger syndrome, and potential role

[86726] in peroxisomal membrane assembly. Proc. Nat. Acad. Sci. 96: 2116–2121,

[86727] 1999.

[86728] 15233.

[86729] Kinoshita, N.; Ghaedi, K.; Shimozawa, N.; Wanders, R. J. A.; Matsuzono,

[86730] Y.; Imanaka, T.; Okumoto, K.; Suzuki, Y.; Kondo, N.; Fujiki, Y.:

[86731] Newly identified Chinese hamster ovary cell mutants are defective

[86732] in biogenesis of peroxisomal membrane vesicles (peroxisomal ghosts),

[86733] representing a novel complementation group in mammals J. Biol. Chem. 273:

[86734] 24122–24130, 1998.

[86735] 15234.

[86736] Chen, K.-S.; Gunaratne, P. H.; Hoheisel, J. D.; Young, I. G.; Gabor

[86737] Miklos, G. L.; Greenberg, F.; Shaffer, L. G.; Campbell, H.

D.; Lupski,

[86738] J. R.: The human homologue of the *Drosophila melanogaster* flightless-I

[86739] gene (flil) maps within the Smith-Magenis microdeletion critical region

[86740] in 17p11.2. *Am. J. Hum. Genet.* 56: 175–182, 1995.

[86741] 15235.

[86742] Heisterkamp, N.; Kaartinen, V.; van Soest, S.; Bokoch, G. M.; Groffen,

[86743] J.: Human ABR encodes a protein with GAP-rac activity and homology

[86744] to the DBL nucleotide exchange factor domain. *J. Biol. Chem.* 268:

[86745] 16903–16906, 1993.

[86746] 15236.

[86747] Heisterkamp, N.; Morris, C.; Groffen, J.: ABR, an active BCR-related

[86748] gene. *Nucleic Acids Res.* 17: 8821–8831, 1989.

[86749] 15237.

[86750] McDonald, J. D.; Daneshvar, L.; Willert, J. R.; Matsumura, K.;

- [86751] Waldman, F.; Cogen, P. H.: Physical mapping of chromosome 17p13.3
- [86752] in the region of a putative tumor suppressor gene important in medulloblastoma. *Genomics* 23:
- [86753] 229–232, 1994.
- [86754] 15238.
- [86755] Shimomura, H.; Sanke, T.; Hanabusa, T.; Tsunoda, K.; Furuta, H.;
- [86756] Nanjo, K.: Nonsense mutation of islet-1 gene (Q310X) found in a type
- [86757] 2 diabetic patient with a strong family history. *Diabetes* 49: 1597–1600,
- [86758] 2000.
- [86759] 15239.
- [86760] Tanizawa, Y.; Riggs, A. C.; Dagogo-Jack, S.; Vaxillaire, M.; Froguel,
- [86761] P.; Liu, L.; Donis-Keller, H.; Permutt, M. A.: Isolation of the human
- [86762] LIM/homeodomain gene islet-1 and identification of a simple sequence
- [86763] repeat 1. *Diabetes* 43: 935–941, 1994.

- [86764] 15240.
- [86765] Corti, O.; Finocchiaro, G.; Rossi, E.; Zuffardi, O.; DiDonato,
[86766] S.: Molecular cloning of cDNAs encoding human carnitine
acetyltransferase
[86767] and mapping of the corresponding gene to chromosome
9q34.1. Genomics 23:
[86768] 94–99, 1994.
- [86769] 15241.
- [86770] Kalaria, R. N.; Harik, S. I.: Carnitine acetyltransferase ac-
tivity
[86771] in the human brain and its microvessels is decreased in
Alzheimer's
[86772] disease. Ann. Neurol. 32: 583–586, 1992.
- [86773] 15242.
- [86774] van der Leij, F. R.; Huijkman, N. C. A.; Boomsma, C.;
Kuipers,
[86775] J. R. G.; Bartelds, B.: Genomics of the human carnitine
acyltransferase
[86776] genes. Molec. Genet. Metab. 71: 139–153, 2000.
- [86777] 15243.

- [86778] Barkardottir, R. B.; Sarantaus, L.; Arason, A.; Vehmanen, P.; Bendahl,
- [86779] P.-O.; Kainu, T.; Syrjakoski, K.; Krahe, R.; Huusko, P.; Pyrhonen,
- [86780] S.; Holli, K.; Kallioniemi, O.-P.; Egilsson, V.; Kere, J.; Nevanlinna,
- [86781] H.: Haplotype analysis in Icelandic and Finnish BRCA2 999del5 breast
- [86782] cancer families. *Europ. J. Hum. Genet.* 9: 773-779, 2001.
- [86783] 15244.
- [86784] Breast Cancer Linkage Consortium: Cancer risks in BRCA2 mutation
- [86785] carriers. *J. Nat. Cancer Inst.* 91: 1310-1316, 1999.
- [86786] 15245.
- [86787] Connor, F.; Smith, A.; Wooster, R.; Stratton, M.; Dixon, A.; Campbell,
- [86788] E.; Tait, T.-M.; Freeman, T.; Ashworth, A.: Cloning, chromosomal
- [86789] mapping and expression pattern of the mouse Brca2 gene. *Hum. Molec.*
- [86790] *Genet.* 6: 291-300, 1997.

- [86791] 15246.
- [86792] Couch, F. J.; Rommens, J. M.; Neuhausen, S. L.; Belanger, C.; Dumont, M.; Abel, K.; Bell, R.; Berry, S.; Bogden, R.; Cannon-Albright, L.; Farid, L.; Frye, C.; and 30 others: Generation of an integrated transcription map of the BRCA2 region on chromosome 13q12-q13. *Genomics* 36: 86-99, 1996.
- [86797] 15247.
- [86798] Davies, A. A.; Masson, J.-Y.; McIlwraith, M. J.; Stasiak, A. Z.; Stasiak, A.; Venkitaraman, A. R.; West, S. C.: Role of BRCA2 in control of the RAD51 recombination and DNA repair protein. *Molec. Cell* 7: 273-282, 2001.
- [86802] 15248.
- [86803] Easton, D. F.; Ford, D.; Bishop, D. T.; Breast Cancer Linkage

- [86804] Consortium: Breast and ovarian cancer incidence in
BRCA1-mutation
- [86805] carriers. Am. J. Hum. Genet. 56: 265-271, 1995.
- [86806] 15249.
- [86807] Easton, D. F.; Steele, L.; Fields, P.; Ormiston, W.; Averill,
[86808] D.; Daly, P. A.; McManus, R.; Neuhausen, S. L.; Ford, D.;
Wooster,
[86809] R.; Cannon-Albright, L. A.; Stratton, M. R.; Goldgar, D. E.:
Cancer
[86810] risks in two large breast cancer families linked to BRCA2
on chromosome
[86811] 13q12-13. Am. J. Hum. Genet. 61: 120-128, 1997.
- [86812] 15250.
- [86813] Fackenthal, J. D.; Cartegni, L.; Krainer, A. R.; Olopade, O. I.
[86814] : BRCA2 T2722R is a deleterious allele that causes exon
skipping. Am.
[86815] J. Hum. Genet. 71: 625-631, 2002.
- [86816] 15251.
- [86817] Garcia-Marco, J. A.; Caldas, C.; Price, C. M.; Wiedemann, L.
M.;

- [86818] Ashworth, A.; Catovsky, D.: Frequent somatic deletion of the 13q12.3
- [86819] locus encompassing BRCA2 in chronic lymphocytic leukemia. Blood 88:
- [86820] 1568–1575, 1996.
- [86821] 15252.
- [86822] Gayther, S. A.; Mangion, J.; Russell, P.; Seal, S.; Barfoot, R.;
- [86823] Ponder, B. A. J.; Stratton, M. R.; Easton, D.: Variation in risks
- [86824] of breast and ovarian cancer associated with different germline mutations
- [86825] of the BRCA2 gene. Nature Genet. 15: 103–105, 1997.
- [86826] 15253.
- [86827] Gronberg, H.; Ahman, A.-K.; Emanuelsson, M.; Bergh, A.; Damber,
- [86828] J.-E.; Borg, A.: BRCA2 mutation in a family with hereditary prostate
- [86829] cancer. Genes Chromosomes Cancer 30: 299–301, 2001.
- [86830] 15254.
- [86831] Gudmundsson, J.; Johannesdottir, G.; Arason, A.; Bergthorsson,

- [86832] J. T.; Ingvarsson, S.; Egilsson, V.; Barkardottir, R. B.: Frequent
occurrence of BRCA2 linkage in Icelandic breast cancer families and
segregation of a common BRCA2 haplotype. Am. J. Hum. Genet. 58:
749–756, 1996.
- [86836] 15255.
- [86837] Healey, C. S.; Dunning, A. M.; Teare, M. D.; Chase, D.; Parker,
L.; Burn, J.; Chang–Claude, J.; Mannermaa, A.; Kataja, V.; Huntsman,
D. G.; Pharoah, P. D. P.; Luben, R. N.; Easton, D. F.; Ponder, B.
A. J.: A common variant in BRCA2 is associated with both breast cancer
risk and prenatal viability. Nature Genet. 26: 362–364, 2000.
- [86842] 15256.
- [86843] Gengyo–Ando, K.; Kitayama, H.; Mukaida, M.; Ikawa, Y.: A murine
neural–specific homolog corrects cholinergic defects in

Caenorhabditis

- [86845] *elegans* unc-18 mutants. J. Neurosci. 16: 6695–6702, 1996.
- [86846] 15257.
- [86847] Pevsner, J.; Hsu, S.-C.; Scheller, R. H.: n-Sec1: a neural-specific
- [86848] syntaxin-binding protein. Proc. Nat. Acad. Sci. 91: 1445–1449, 1994.
- [86849] 15258.
- [86850] Swanson, D. A.; Steel, J. M.; Valle, D.: Identification and characterization
- [86851] of the human ortholog of rat STXBP1, a protein implicated in vesicle
- [86852] trafficking and neurotransmitter release. Genomics 48: 373–376,
- [86853] 1998.
- [86854] 15259.
- [86855] Verhage, M.; Mala, A. S.; Plomp, J. J.; Brussaard, A. B.; Heeroma,
- [86856] J. H.; Vermeer, H.; Toonen, R. F.; Hammer, R. E.; van den Berg, T.
- [86857] K.; Missler, M.; Geuze, H. J.; Sudhof, T. C.: Synaptic as-

sembly of

[86858] the brain in the absence of neurotransmitter secretion.

Science 287:

[86859] 864–869, 2000.

[86860] 15260.

[86861] Monaco, C.; Negrini, M.; Sozzi, G.; Veronese, M. L.; Vorechovsky,

[86862] I.; Godwin, A. K.; Croce, C. M.: Molecular cloning and characterization

[86863] of LOH11CR2A, a new gene within a refined minimal region of LOH at

[86864] 11q23. Genomics 46: 217–222, 1997.

[86865] 15261.

[86866] Topper, J. N.; Cai, J.; Qiu, Y.; Anderson, K. R.; Xu, Y.–Y.; Deeds,

[86867] J. D.; Feeley, R.; Gimeno, C. J.; Woolf, E. A.; Tayber, O.; Mays,

[86868] G. G.; Sampson, B. A.; Schoen, F. J.; Gimbrone, M. A., Jr.; Falb,

[86869] D.: Vascular MADs: two novel MAD-related genes selectively inducible

[86870] by flow in human vascular endothelium. Proc. Nat. Acad.

Sci. 94:

[86871] 9314–9319, 1997.

[86872] 15262.

[86873] Festing, M. F. W. (ed.): Inbred strains in biomedical research. New

[86874] York: Oxford University Press , 1979. p. 255.

[86875] 15263.

[86876] Illa, I.; Serrano–Munuera, C.; Gallardo, E.; Lasa, A.; Rojas–Garcia,

[86877] R.; Palmer, J.; Gallano, P.; Baiget, M.; Matsuda, C.; Brown, R. H.

[86878] : Distal anterior compartment myopathy: a dysferlin mutation causing

[86879] a new muscular dystrophy phenotype. Ann. Neurol. 49: 130–134, 2001.

[86880] 15264.

[86881] Illarioshkin, S. N.; Ivanova–Smolenskaya, I. A.; Greenberg, C.

[86882] R.; Nylen, E.; Sukhorukov, V. S.; Poleshchuk, V. V.; Markova, E. D.;

[86883] Wrogemann, K.: Identical dysferlin mutation in limb–girdle

muscular

- [86884] dystrophy type 2B and distal myopathy. *Neurology* 55: 1931–1933, 2000.
- [86885] 2000.
- [86886] 15265.
- [86887] Liu, J.; Wu, C.; Bossie, K.; Bejaoui, K.; Hosler, B. A.; Gingrich, J. C.; Ben Hamida, M.; Hentati, F.; Schurr, E.; de Jong, P. J.; Brown, R. H., Jr.: Generation of 3–Mb PAC contig spanning the Miyoshi myopathy/limb girdle muscular dystrophy (MM/LGMD2B) locus on chromosome 2p13. *Genomics* 49: 23–29, 1998.
- [86889] 2000.
- [86890] 15266.
- [86891] Mahjneh, I.; Vannelli, G.; Bushby, K.; Marconi, G. P.: A large inbred Palestinian family with two forms of muscular dystrophy. *Neuromusc. Disord.* 2: 277–283, 1992.
- [86892] 15267.

- [86897] Weiler, T.; Bashir, R.; Anderson, L. V. B.; Davison, K.; Moss, J. A.; Britton, S.; Nylen, E.; Keers, S.; Vafiadaki, E.; Greenberg, C. R.; Bushby, K. M. D.; Wrogemann, K.: Identical mutation in patients with limb girdle muscular dystrophy type 2B or Miyoshi myopathy suggests a role for modifier gene(s). Hum. Molec. Genet. 8: 871–877, 1999.
- [86902] 15268.
- [86903] Hannon, G. J.; Beach, D.: p15(INK4B) is a potential effector of TGF-beta-induced cell cycle arrest. Nature 371: 257–261, 1994.
- [86905] 15269.
- [86906] Stone, S.; Dayananth, P.; Jiang, P.; Weaver-Feldhaus, J. M.; Tavtigian, S. V.; Cannon-Albright, L.; Kamb, A.: Genomic structure, expression and mutational analysis of the P15 (MTS2) gene. Oncogene 11: 987–991, 1995.
- [86909] 1995.

- [86910] 15270.
- [86911] Pennica, D.; Arce, V.; Swanson, T. A.; Vejsada, R.; Pollock, R.
- [86912] A.; Armanini, M.; Dudley, K.; Phillips, H. S.; Rosenthal, A.; Kato, A. C.; Henderson, C. E.: Cardiotrophin-1, a cytokine present in embryonic muscle, supports long-term survival of spinal motoneurons. *Neuron* 17: 63-74, 1996.
- [86913] 15271.
- [86914] Pennica, D.; King, K. L.; Shaw, K. J.; Luis, E.; Rullamas, J.; Luoh, S.-M.; Darbonne, W. C.; Knutzon, D. S.; Yen, R.; Chien, K. R.; Baker, J. B.; Wood, W. I.: Expression cloning of cardiotrophin 1, a cytokine that induces cardiac myocyte hypertrophy. *Proc. Nat. Acad. Sci.* 92: 1142-1146, 1995.
- [86915] 15272.
- [86916] Ghosh, A.: Learning more about NMDA receptor regula-

tion. Science 295:

[86924] 449–451, 2002.

[86925] 15273.

[86926] Grunwald, I. C.; Korte, M.; Wolfer, D.; Wilkinson, G. A.; Unsicker,

[86927] K.; Lipp, H.-P.; Bonhoeffer, T.; Klein, R.: Kinase-independent requirement

[86928] of EphB2 receptors in hippocampal synaptic plasticity.

Neuron 32:

[86929] 1027–1040, 2001.

[86930] 15274.

[86931] Henderson, J. T.; Georgiou, J.; Jia, Z.; Robertson, J.; Elowe,

[86932] S.; Roder, J. C.; Pawson, T.: The receptor tyrosine kinase EphB2

[86933] regulates NMDA-dependent synaptic function. Neuron 32: 1041–1056,

[86934] 2001.

[86935] 15275.

[86936] Himanen, J.-P.; Rajashankar, K. R.; Lackmann, M.; Cowan, C. A.;

[86937] Henkemeyer, M.; Nikolov, D. B.: Crystal structure of an

Eph receptor-ephrin

[86938] complex. Nature 414: 933–938, 2001.

[86939] 15276.

[86940] Sakano, S.; Serizawa, R.; Inada, T.; Iwama, A.; Itoh, A.; Kato,

[86941] C.; Shimizu, Y.; Shinkai, F.; Shimizu, R.; Kondo, S.; Ohno, M.; Suda,

[86942] T.: Characterization of a ligand for receptor protein-tyrosine kinase

[86943] HTK expressed in immature hematopoietic cells. Oncogene 13: 813–822,

[86944] 1996.

[86945] 15277.

[86946] Takasu, M. A.; Dalva, M. B.; Zigmond, R. E.; Greenberg, M. E.

[86947] : Modulation of NMDA receptor-dependent calcium influx and gene expression

[86948] through EphB receptors. Science 295: 491–495, 2002.

[86949] 15278.

[86950] Wang, H. U.; Chen, Z.-F.; Anderson, D. J.: Molecular distinction

- [86951] and angiogenic interaction between embryonic arteries and veins revealed
- [86952] by ephrin-B2 and its receptor Eph-B4. *Cell* 93: 741-753, 1998.
- [86953] 15279.
- [86954] Slama, A.; Brivet, M.; Boutron, A.; Legrand, A.; Saudubray, J.-M.;
- [86955] Demaugre, F.: Complementation analysis of carnitine palmitoyltransferase
- [86956] I and II defects. *Pediat. Res.* 40: 542-546, 1996.
- [86957] 15280.
- [86958] Doudney, K.; Murdoch, J. N.; Paternotte, C.; Bentley, L.; Gregory,
- [86959] S.; Copp, A. J.; Stanier, P.: Comparative physical and transcript
- [86960] maps of approximately 1 Mb around loop-tail, a gene for severe neural
- [86961] tube defects on distal mouse chromosome 1 and human chromosome 1q22-q23. *Genomics* 72:
- [86962] 180-192, 2001.
- [86963] 15281.

- [86964] Kibar, Z.; Vogan, K. J.; Groulx, N.; Justice, M. J.; Underhill,
[86965] D. A.; Gros, P.: Ltap, a mammalian homolog of *Drosophila*
Strabismus/Van
[86966] Gogh, is altered in the mouse neural tube mutant loop-
tail. *Nature*
[86967] *Genet.* 28: 251–255, 2001.
[86968] 15282.
[86969] Mullick, A.; Trasler, D.; Gros, P.: High-resolution linkage
map
[86970] in the vicinity of the Lp locus. *Genomics* 26: 479–488,
1995.
[86971] 15283.
[86972] Murdoch, J. N.; Doudney, K.; Paternotte, C.; Copp, A. J.;
Stanier,
[86973] P.: Severe neural tube defects in the loop-tail mouse re-
sult from
[86974] mutation of *Lpp1*, a novel gene involved in floor plate
specification. *Hum.*
[86975] *Molec. Genet.* 10: 2593–2601, 2001.
[86976] 15284.
[86977] Stanier, P.; Henson, J. N.; Eddleston, J.; Moore, G. E.; Copp,

- [86978] A. J.: Genetic basis of neural tube defects: the mouse gene loop-tail
- [86979] maps to a region of chromosome 1 syntenic with human 1q21-q23. Genomics 26:
- [86980] 473-478, 1995.
- [86981] 15285.
- [86982] Yu, J. X.; Chao, L.; Chao, J.: Protasin is a novel human serine
- [86983] proteinase from seminal fluid: purification, tissue distribution,
- [86984] and localization in prostate gland. J. Biol. Chem. 269: 18843-18848,
- [86985] 1994.
- [86986] 15286.
- [86987] Yu, J. X.; Chao, L.; Chao, J.: Molecular cloning, tissue-specific
- [86988] expression, and cellular localization of human protasin mRNA. J.
- [86989] Biol. Chem. 270: 13483-13489, 1995.
- [86990] 15287.
- [86991] Yu, J. X.; Chao, L.; Ward, D. C.; Chao, J.: Structure and

chromosomal

[86992] localization of the human prostatic (PRSS8) gene. Genomics 32: 334–340,

[86993] 1996.

[86994] 15288.

[86995] Carlberg, C.; Hooft van Huijsduijnen, R.; Staple, J. K.; DeLamar, T.

[86996] J. F.; Becker-Andre, M.: RZR, a new family of retinoid-related orphan

[86997] receptors that function as both monomers and homodimers. Mol. Endocrinol. 8:

[86998] 757–770, 1994.

[86999] 15289.

[87000] Ueda, H. R.; Chen, W.; Adachi, A.; Wakamatsu, H.; Hayashi, S.;

[87001] Takasugi, T.; Nagano, M.; Nakahama, K.; Suzuki, Y.; Sugano, S.; Iino,

[87002] M.; Shigemoto, Y.; Hashimoto, S.: A transcription factor response

[87003] element for gene expression during circadian night. Nature 418:

[87004] 534–539, 2002.

[87005] 15290.

[87006] Masternak, K.; Barras, E.; Zufferey, M.; Conrad, B.;
Corthals,

[87007] G.; Aebersold, R.; Sanchez, J.-C.; Hochstrasser, D. F.;
Mach, B.;

[87008] Reith, W.: A gene encoding a novel RFX-associated trans-
activator

[87009] is mutated in the majority of MHC class II deficiency pa-
tients. Nature

[87010] Genet. 20: 273-277, 1998.

[87011] 15291.

[87012] Prange, C. K.; Pennacchio, L. A.; Lieuallen, K.; Fan, W.;
Lennon,

[87013] G. G.: Characterization of the human neurocan gene,
CSPG3. Gene 221:

[87014] 199-205, 1998.

[87015] 15292.

[87016] Rauch, U.; Grimpe, B.; Kulbe, G.; Arnold-Ammer, I.; Beier,
D. R.;

[87017] Fassler, R.: Structure and chromosomal localization of the
mouse

- [87018] neurocan gene. Genomics 28: 405–410, 1995.
- [87019] 15293.
- [87020] Rauch, U.; Karthikeyan, L.; Maurel, P.; Margolis, R. U.;
Margolis,
- [87021] R. K.: Cloning and primary structure of neurocan, a devel-
opmentally
- [87022] regulated, aggregating chondroitin sulfate proteoglycan of
brain. J.
- [87023] Biol. Chem. 267: 19536–19547, 1992.
- [87024] 15294.
- [87025] Drummond, J. T.; Genschel, J.; Wolf, E.; Modrich, P.: DHFR/
MSH3
- [87026] amplification in methotrexate-resistant cells alters the
hMutS-alpha/hMutS-beta
- [87027] ratio and reduces the efficiency of base-base mismatch
repair. Proc.
- [87028] Nat. Acad. Sci. 94: 10144–10149, 1997.
- [87029] 15295.
- [87030] Fujii, H.; Shimada, T.: Isolation and characterization of
cDNA
- [87031] clones derived from the divergently transcribed gene in

the region

[87032] upstream from the human dihydrofolate reductase gene. J.

Biol. Chem. 264:

[87033] 10057–10064, 1989.

[87034] 15296.

[87035] Inokuchi, K.; Ikejima, M.; Watanabe, A.; Nakajima, E.;

Orimo, H.;

[87036] Nomura, T.; Shimada, T.: Loss of expression of the human

MSH3 gene

[87037] in hematological malignancies. Biochem. Biophys. Res.

Commun. 214:

[87038] 171–179, 1995.

[87039] 15297.

[87040] Linton, J. P.; Yen, J. Y.-J.; Selby, E.; Chen, Z.; Chinsky, J.

[87041] M.; Liu, K.; Kellems, R. E.; Crouse, G. F.: Dual bidirectional

promoters

[87042] at the mouse dhfr locus: cloning and characterization of

two mRNA

[87043] classes of the divergently transcribed Rep-1 gene. Molec.

Cell. Biol. 9:

[87044] 3058–3072, 1989.

- [87045] 15298.
- [87046] Marra, G.; Iaccarino, I.; Lettieri, T.; Roscilli, G.; Delmastro,
[87047] P.; Jiricny, J.: Mismatch repair deficiency associated with
overexpression
[87048] of the MSH3 gene. Proc. Nat. Acad. Sci. 95: 8568–8573,
1998.
[87049] 15299.
- [87050] Nakajima, E.; Orimo, H.; Ikejima, M.; Shimada, T.: Nine-bp
repeat
[87051] polymorphism in exon 1 of the hMSH3 gene. Jpn. J. Hum.
Genet. 40:
[87052] 343–345, 1995.
[87053] 15300.
- [87054] Risinger, J. I.; Umar, A.; Boyd, J.; Berchuck, A.; Kunkel, T.
A.;
- [87055] Barrett, J. C.: Mutation of MSH3 in endometrial cancer and
evidence
[87056] for its functional role in heteroduplex repair. Nature
Genet. 14:
[87057] 102–109, 1996.
[87058] 15301.

- [87059] Smith, M. L.; Mitchell, P. J.; Crouse, G. F.: Analysis of the
[87060] mouse Dhfr/Rep-3 major promoter region by using
linker-scanning and
[87061] internal deletion mutations and DNase I footprinting.
Molec. Cell.
[87062] Biol. 10: 6003-6012, 1990.
[87063] 15302.
[87064] Watanabe, A.; Ikejima, M.; Suzuki, N.; Shimada, T.: Ge-
nomic organization
[87065] and expression of the human MSH3 gene. Genomics 31:
311-318, 1996.
[87066] 15303.
[87067] Yin, J.; Kong, D.; Wang, S.; Zou, T.-T. Souza, R. F.; Smolin-
ski,
[87068] K. N.; Lynch, P. M.; Hamilton, S. R.; Sugimura, H.; Powell,
S. M.;
[87069] Young, J.; Abraham, J. M.; Meltzer, S. J.: Mutation of
hMSH3 and
[87070] hMSH6 mismatch repair genes in genetically unstable hu-
man colorectal
[87071] and gastric carcinomas. Hum. Mutat. 10: 474-478, 1997.

[87072] 15304.

[87073] Burger, B.; Uhlhaas, S.; Mangold, E.; Propping, P.; Friedl, W.;

[87074] Jenne, D.; Dockter, G.; Back, W.: Novel de novo mutation of MADH4/SMAD4

[87075] in a patient with juvenile polyposis. (Letter) Am. J. Med. Genet. 110:

[87076] 289–291, 2002.

[87077] 15305.

[87078] Sakamoto, A.; Ono, K.; Abe, M.; Jasmin, G.; Eki, T.; Murakami,

[87079] Y.; Masaki, T.; Toyo-oka, T.; Hanaoka, F.: Both hypertrophic and

[87080] dilated cardiomyopathies are caused by mutation of the same gene,

[87081] delta-sarcoglycan, in hamster: an animal model of disrupted dystrophin-associated

[87082] glycoprotein complex. Proc. Nat. Acad. Sci. 94: 13873–13878, 1997.

[87083] 15306.

[87084] Kurima, K.; Peters. L. M.; Yang, Y.; Riazuddin, S.; Ahmed, Z. M.;

- [87085] Naz, S.; Arnaud, D.; Drury, S.; Mo, J.; Makishima, T.; Ghosh, M.;
- [87086] Menon, P. S. N.; and 13 others: Dominant and recessive deafness
- [87087] caused by mutations of a novel gene, TMC1, required for cochlear hair-cell
- [87088] function. *Nature Genet.* 30: 277–284, 2002.
- [87089] 15307.
- [87090] Bryant, P. J.; Huettner, B.; Held, L. I., Jr.; Ryerse, J.; Szidonya,
- [87091] J.: Mutations at the fat locus interfere with cell proliferation
- [87092] control and epithelial morphogenesis in *Drosophila*. *Dev. Biol.* 129:
- [87093] 541–554, 1988.
- [87094] 15308.
- [87095] Dunne, J.; Hanby, A. M.; Poulsom, R.; Jones, T. A.; Sheer, D.;
- [87096] Chin, W. G.; Da, S. M.; Zhao, Q.; Beverley, P. C. L.; Owen, M. J.
- [87097] : Molecular cloning and tissue expression of FAT, the human homologue

[87098] of the *Drosophila* fat gene that is located on chromosome 4q34–q35

[87099] and encodes a putative adhesion molecule. *Genomics* 30: 207–223,

[87100] 1995.

[87101] 15309.

[87102] Hortsch, M.; Goodman, C. S.: Cell and substrate adhesion molecules

[87103] in *Drosophila*. *Ann. Rev. Cell. Biol.* 7: 505–557, 1991.

[87104] 15310.

[87105] Mahoney, P. A.; Weber, U.; Onofrechuk, P.; Biessmann, H.; Bryant,

[87106] P. J.; Goodman, C. S.: The fat tumor suppressor gene in *Drosophila*

[87107] encodes a novel member of the cadherin gene superfamily. *Cell* 67:

[87108] 853–868, 1991.

[87109] 15311.

[87110] Browning, J. L.; Ngam-ek, A.; Lawton, P.; DeMarinis, J.; Tizard,

[87111] R.; Chow, E. P.; Hession, C.; O'Brine-Greco, B.; Foley, S. F.;

Ware,

[87112] C. F.: Lymphotoxin beta, a novel member of the TNF family that forms

[87113] a heteromeric complex with lymphotoxin on the cell surface. Cell 72:

[87114] 847-856, 1993.

[87115] 15312.

[87116] Crowe, P. D.; VanArsdale, T. L.; Walter, B. N.; Ware, C. F.; Hession,

[87117] C.; Ehrenfels, B.; Browning, J. L.; Din, W. S.; Goodwin, R. G.; Smith,

[87118] C. A.: A lymphotoxin-beta-specific receptor. Science 264: 707-710,

[87119] 1994.

[87120] 15313.

[87121] Nakamura, T.; Tashiro, K.; Nazarea, M.; Nakano, T.; Sasayama, S.;

[87122] Honjo, T.: The murine lymphotoxin-beta receptor cDNA: isolation by

[87123] the signal sequence trap and chromosomal mapping. Genomics 30: 312-319,

[87124] 1995.

- [87125] 15314.
- [87126] Warzocha, K.; Renard, N.; Charlot, C.; Bienvenu, J.;
Coiffier,
[87127] B.; Salles, G.: Identification of two lymphotoxin beta iso-
forms expressed
[87128] in human lymphoid cell lines and non-Hodgkin's lym-
phomas. Biochem.
[87129] Biophys. Res. Commun. 238: 273-276, 1997.
- [87130] 15315.
- [87131] Tashiro, K.; Tada, H.; Heilker, R.; Shirozu, M.; Nakano, T.;
Honjo,
[87132] T.: Signal sequence trap: a cloning strategy for secreted
proteins
[87133] and type I membrane proteins. Science 261: 600-603,
1993.
- [87134] 15316.
- [87135] George, A.; Sabsay, B.; Simonian, P. A.; Veis, A.: Charac-
terization
[87136] of a novel dentin matrix acidic phosphoprotein: implica-
tions for induction
[87137] of biomineralization . J. Biol. Chem. 268: 12624-12630,
1993.

[87138] 15317.

[87139] Hirst, K. L.; Simmons, D.; Feng, J.; Aplin, H.; Dixon, M. J.;
MacDougall,

[87140] M.: Elucidation of the sequence and the genomic organi-
zation of the

[87141] human dentin matrix acidic phosphoprotein 1 (DMP1)
gene: exclusion

[87142] of the locus from a causative role in the pathogenesis of
dentinogenesis

[87143] imperfecta type II. Genomics 42: 38-45, 1997.

[87144] 15318.

[87145] MacDougall, M.; DuPont, B. R.; Simmons, D.; Leach, R. J.:
Assignment

[87146] of DMP1 to human chromosome 4 band q21 by in situ hy-
bridization. Cytogenet.

[87147] Cell. Genet. 74: 189 only, 1996.

[87148] 15319.

[87149] Lu, Z.; Xu, S.; Joazeiro, C.; Cobb, M. H.; Hunter, T.: The
PHD

[87150] domain of MEKK1 acts as an E3 ubiquitin ligase and medi-
ates ubiquitination

- [87151] and degradation of ERK1/2. *Molec. Cell* 9: 945–956, 2002.
- [87152] 15320.
- [87153] Xu, G. M.; Sikaneta, T.; Sullivan, B. M.; Zhang, Q.; Andreucci,
M.; Stehle, T.; Drummond, I.; Arnaout, M. A.: Polycystin-1
interacts
with intermediate filaments. *J. Biol. Chem.* 276:
46544–46552, 2001.
- [87156] 15321.
- [87157] Zerres, K.; Rudnik-Schoneborn, S.; Deget, F.; Beetz, R.;
Brodehl,
J.; Kuwertz-Broking, E.; Muller-Wiefel, D.; Rascher, W.;
Ruder, H.;
Scharer, K.; Weber, H. P.: Childhood onset autosomal
dominant polycystic
kidney disease in sibs: clinical picture and recurrence risk.
J.
Med. Genet. 30: 583–588, 1993.
- [87162] 15322.
- [87163] Barnard, R. C.; Pascall, J. C.; Brown, K. D.; McKay, I. A.;
Williams,

- [87164] N. S.; Bustin, S. A.: Coding sequence of ERF-1, the human homologue
[87165] of Tis11b/cMG1, members of the Tis11 family of early re-
[87166] sponse genes. Nucleic
[87167] Acids Res. 21: 3580 only, 1993.
[87167] 15323.
- [87168] Bustin, S. A.; Xiao-Feng, N.; Barnard, R. C.; Kumar, V.;
Pascall,
[87169] J. C.; Brown, K. D.; Leigh, I. M.; Williams, N. S.; McKay, I.
A.:
[87170] Cloning and characterisation of ERF1, a human member of
the Tis11
[87171] family of early-response genes. DNA Cell Biol. 13:
449-459, 1994.
[87172] 15324.
- [87173] Maclean, K. N.; See, C. G.; McKay, I. A.; Bustin, S. A.: The
human
[87174] immediate early gene BRF1 maps to chromosome
14q22-q24. Genomics 30:
[87175] 89-90, 1995.
[87176] 15325.

- [87177] Ning, Z.-Q.; Norton, J. D.; Li, J.; Murphy, J. J.: Distinct mechanisms
- [87178] for rescue from apoptosis in Ramos human B cells by signaling through
- [87179] CD40 and interleukin-4 receptor: a role for inhibition of an early
- [87180] response gene, Berg36. *Europ. J. Immun.* 26: 2356-2363, 1996.
- [87181] 15326.
- [87182] Maas, S.; Kim, Y.-G.; Rich, A.: Genomic clustering of tRNA-specific
- [87183] adenosine deaminase ADAT1 and two tRNA synthetases. *Mammalian Genome* 12:
- [87184] 387-393, 2001.
- [87185] 15327.
- [87186] Bolz, H.; von Brederlow, B.; Ramirez, A.; Bryda, E. C.; Kutsche,
- [87187] K.; Nothwang, H. G.; Seeliger, M.; Cabrera, M. C.-S.; Vila, M. C.;
- [87188] Molina, O. P.; Gal, A.; Kubisch, C.: Mutation of CDH23, encoding
- [87189] a new member of the cadherin gene family, causes Usher

syndrome type

[87190] 1D. Nature Genet. 27: 108–112, 2001.

[87191] 15328.

[87192] Bork, J. M.; Peters, L. M.; Riazuddin, S.; Bernstein, S. L.;
Ahmed,

[87193] Z. M.; Ness, S. L.; Polomeno, R.; Ramesh, A.; Schloss, M.;
Srisailpathy,

[87194] C. R. S.; Wayne, S.; Bellman, S.; and 16 others: Usher syn-
drome

[87195] 1D and nonsyndromic autosomal recessive deafness
DFNB12 are caused

[87196] by allelic mutations of the novel cadherin-like gene
CDH23. Am. J.

[87197] Hum. Genet. 68: 26–37, 2001.

[87198] 15329.

[87199] Derynck, R.; Gelbart, W. M.; Harland, R. M.; Heldin, C.–H.;
Kern,

[87200] S. E.; Massague, J.; Melton, D. A.; Mlodzik, M.; Padgett, R.
W.; Roberts,

[87201] A. B.; Smith, J.; Thomsen, G. H.; Vogelstein, B.; Wang,
X.–F.: Nomenclature:

[87202] vertebrate mediators of TGF-beta family signals. (Letter)

Cell 87:

[87203] 173 only, 1996.

[87204] 15330.

[87205] Friedl, W.; Kruse, R.; Uhlhaas, S.; Stolte, M.; Schartmann, B.;

[87206] Keller, K. M.; Jungck, M.; Stern, M.; Loff, S.; Back, W.;
Propping,

[87207] P.; Jenne, D. E.: Frequent 4-bp deletion in exon 9 of the
SMAD4/MADH4

[87208] gene in familial juvenile polyposis patients. Genes Chromosomes Cancer 25:

[87209] 403–406, 1999.

[87210] 15331.

[87211] Hahn, S. A.; Schutte, M.; Hoque, T. M. S.; Moskaluk, C. A.;
da

[87212] Costa, L. T.; Rozenblum, E.; Weinstein, C. L.; Fischer, A.;
Yeo, C.

[87213] J.; Hruban, R. H.; Kern, S. E.: DPC4, a candidate tumor
suppressor

[87214] gene at human chromosome 18q21.1. Science 271:
350–354, 1996.

[87215] 15332.

- [87216] Howe, J. R.; Shellnut, J.; Wagner, B.; Ringold, J. C.; Sayed,
[87217] M. G.; Ahmed, A. F.; Lynch, P. M.; Amos, C. I.; Sistonen, P.;
Aaltonen,
[87218] L. A.: Common deletion of SMAD4 in juvenile polyposis is
a mutational
[87219] hotspot. Am. J. Hum. Genet. 70: 1357–1362, 2002.
[87220] 15333.
[87221] Inman, G. J.; Nicolas, F. J.; Hill, C. S.: Nucleocytoplasmic
[87222] shuttling of Smads 2, 3, and 4 permits sensing of TGF-
beta receptor
[87223] activity. Molec. Cell 10: 283–294, 2002.
[87224] 15334.
[87225] Kim, S. K.; Fan, Y.; Papadimitrakopoulou, V.; Clayman, G.;
Hittelman,
[87226] W. N.; Hong, W. K.; Lotan, R.; Mao, L.: DPC4, a candidate
tumor suppressor
[87227] gene, is altered infrequently in head and neck squamous
cell carcinoma. Cancer
[87228] Res. 56: 2519–2521, 1996.
[87229] 15335.

- [87230] Kinzler, K. W.; Vogelstein, B.: Landscaping the cancer terrain. *Science* 280:
- [87231] 1036–1037, 1998.
- [87232] 15336.
- [87233] MacGrogan, D.; et al.; et al: Comparative mutational analysis
- [87234] of DPC4 (Smad4) in prostatic and colorectal carcinomas. *Oncogene* 15:
- [87235] 1111–1114, 1997.
- [87236] 15337.
- [87237] Roth, S.; Johansson, M.; Loukola, A.; Peltomaki, P.; Jarvinen,
- [87238] H.; Mecklin, J.-P.; Aaltonen, L. A.: Mutation analysis of SMAD2,
- [87239] SMAD3, and SMAD4 genes in hereditary non-polyposis colorectal cancer. *J.*
- [87240] *Med. Genet.* 37: 298–300, 2000.
- [87241] 15338.
- [87242] Schutte, M.; Hruban, R. H.; Hedrick, L.; Cho, K. R.; Nadasdy,
- [87243] G. M.; Weinstein, C. L.; Bova, G. S.; Isaacs, W. B.; Cairns, P.;

Nawroz,

[87244] H.; Sidransky, D.; Casero, R. A., Jr.; Meltzer, P. S.; Hahn, S. A.;

[87245] Kern, S. E.: DPC4 gene in various tumor types. Cancer Res. 56:

[87246] 2527–2530, 1996.

[87247] 15339.

[87248] Shioda, T.; Lechleider, R. J.; Dunwoodie, S. L.; Li, H.; Yahata,

[87249] T.; de Caestecker, M. P.; Fenner, M. H.; Roberts, A. B.; Is-selbacher,

[87250] K. J.: Transcriptional activating activity of Smad4: roles of SMAD

[87251] hetero-oligomerization and enhancement by an associating transactivator. Proc.

[87252] Nat. Acad. Sci. 95: 9785–9790, 1998.

[87253] 15340.

[87254] Sirard, C.; de la Pompa, J. L.; Elia, A.; Itie, A.; Mirtsos, C.;

[87255] Cheung, A.; Hahn, S.; Wakeham, A.; Schwartz, L.; Kern, S. E.; Rossant,

[87256] J.; Mak, T. W.: The tumor suppressor gene Smad4/Dpc4 is

required

[87257] for gastrulation and later for anterior development of the mouse embryo. Genes

[87258] Dev. 12: 107–119, 1998.

[87259] 15341.

[87260] Tagaki, Y.; et al.; et al: Somatic alterations of the DPC4 gene

[87261] in human colorectal cancers in vivo. Gastroenterology 111: 1369–1372,

[87262] 1996.

[87263] 15342.

[87264] Takaku, K.; Oshima, M.; Miyoshi, H.; Matsui, M.; Seldin, M. F.;

[87265] Taketo, M. M.: Intestinal tumorigenesis in compound mutant mice of

[87266] both Dpc4 (Smad4) and Apc genes. Cell 92: 645–656, 1998.

[87267] 15343.

[87268] Thiagalingam, S.; Lengauer, C.; Leach, F. S.; Schutte, M.; Hahn,

[87269] S. A.; Overhauser, J.; Willson, J. K. V.; Markowitz, S.;

Hamilton,

- [87270] S. R.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Evaluation of
- [87271] candidate tumour suppressor genes on chromosome 18 in colorectal cancers. Nature
- [87272] Genet. 13: 343–346, 1996.
- [87273] 15344.
- [87274] Zawel, L.; Dai, J. L.; Buckhaults, P.; Zhou, S.; Kinzler, K. W.;
- [87275] Vogelstein, B.; Kern, S. E.: Human Smad3 and Smad4 are sequence-specific
- [87276] transcription activators. Molec. Cell 1: 611–617, 1998.
- [87277] 15345.
- [87278] Zhou, S.; Buckhaults, P.; Zawel, L.; Bunz, F.; Riggins, G.; Le
- [87279] Dai, J.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Targeted deletion
- [87280] of Smad4 shows it is required for transforming growth factor beta
- [87281] and activin signaling in colorectal cancer cells. Proc. Nat. Acad.
- [87282] Sci. 95: 2412–2416, 1998.
- [87283] 15346.

- [87284] Abbs, S.; Roberts, R. G.; Mathew, C. G.; Bentley, D. R.; Bo-
brow,
- [87285] M.: Accurate assessment of intragenic recombination fre-
quency within
- [87286] the Duchenne muscular dystrophy gene. *Genomics* 7:
602-606, 1990.
- [87287] 15347.
- [87288] Ahn, A. H.; Kunkel, L. M.: The structural and functional di-
versity
- [87289] of dystrophin. *Nature Genet.* 3: 283-291, 1993.
- [87290] 15348.
- [87291] Alwine, J. C.; Kemp, D. J.; Stark, G. R.: Method for detec-
tion
- [87292] of specific RNAs in agarose gels by transfer to diazoben-
zyloxymethyl-paper
- [87293] and by hybridization with DNA probes. *Proc. Nat. Acad.*
Sci. 74:
- [87294] 5350-5354, 1977.
- [87295] 15349.
- [87296] Angelini, C.; Beggs, A. H.; Hoffman, E. P.; Fanin, M.;
Kunkel,

- [87297] L. M.: Enormous dystrophin in a patient with Becker muscular dystrophy. *Neurology* 40: 808–812, 1990.
- [87298] 15350.
- [87299] 15350.
- [87300] Badorff, C.; Berkely, N.; Mehrotra, S.; Talhouk, J. W.; Rhoads, R. E.; Knowlton, K. U.: Enteroviral protease 2A directly cleaves dystrophin and is inhibited by a dystrophin-based substrate analogue. *J. Biol. Chem.* 275: 11191–11197, 2000.
- [87301] 15351.
- [87302] Badorff, C.; Lee, G.-H.; Lamphear, B. J.; Martone, M. E.; Campbell, K. P.; Rhoads, R. E.; Knowlton, K. U.: Enteroviral protease 2A cleaves dystrophin: evidence of cytoskeletal disruption in an acquired cardiomyopathy. *Nature Med.* 5: 320–326, 1999.
- [87303] 15352.
- [87304] Bakker, E.; Pearson, P. L.: Mutation of the Duchenne mus-

cular

- [87311] dystrophy gene associated with meiotic recombination.
(Letter) Clin.
- [87312] Genet. 30: 347–349, 1986.
- [87313] 15353.
- [87314] Bakker, E.; Hofker, M. H.; Goor, N.; Mandel, J. L.; Wroge-
mann,
- [87315] K.; Davies, K. E.; Kunkel, L. M.; Willard, H. F.; Fenton, W.
A.; Sandkuyl,
- [87316] L.; Majoor-Krakauer, D.; van Essen, A. J.; Jahoda, M. G. J.;
Sachs,
- [87317] E. S.; van Ommen, G. J. B.; Pearson, P. L.: Prenatal diagno-
sis and
- [87318] carrier detection of Duchenne muscular dystrophy with
closely linked
- [87319] RFLPs. Lancet I: 655–658, 1985.
- [87320] 15354.
- [87321] Bakker, E.; Van Broeckhoven, C.; Bonten, E. J.; van de
Vooren,
- [87322] M. J.; Veenema, H.; Van Hul, W.; Van Ommen, G. J. B.;
Vandenberghe,
- [87323] A.; Pearson, P. L.: Germline mosaicism and Duchenne

muscular dystrophy

[87324] mutations. Nature 329: 554–556, 1987.

[87325] 15355.

[87326] Bar, S.; Barnea, E.; Levy, Z.; Neuman, S.; Yaffe, D.; Nudel, U.

[87327] : A novel product of the Duchenne muscular dystrophy gene which greatly

[87328] differs from the known isoforms in its structure and tissue distribution. Biochem.

[87329] J. 272: 557–560, 1990.

[87330] 15356.

[87331] Barbieri, A. M.; Soriani, N.; Tubiello, G. M.; Ferrari, M.; Carrera,

[87332] P.: A nonsense mutation (gln-673-term) in exon 17 of the human dystrophin

[87333] gene detected by heteroduplex analysis. Hum. Genet. 96: 343–344,

[87334] 1995.

[87335] 15357.

[87336] Bartlett, R. J.; Pericak-Vance, M. A.; Koh, J.; Yamaoka, L. H.;

- [87337] Chen, J. C.; Hung, W.-Y.; Speer, M. C.; Wapenaar, M. C.;
Van Ommen,
- [87338] G. J. B.; Bakker, E.; Pearson, P. L.; Kandt, R. S.; Siddique,
T.;
- [87339] Gilbert, J. R.; Lee, J. E.; Sirotkin-Roses, M. J.; Roses, A. D.:
- [87340] Duchenne muscular dystrophy: high frequency of dele-
tions. *Neurology* 38:
- [87341] 1-4, 1988.
- [87342] 15358.
- [87343] Barton-Davis, E. R.; Cordier, L.; Shoturma, D. I.; Leland, S.
- [87344] E.; Sweeney, H. L.: Aminoglycoside antibiotics restore dys-
trophin
- [87345] function to skeletal muscles of mdx mice. *J. Clin. Invest.*
104:
- [87346] 375-381, 1999.
- [87347] 15359.
- [87348] Bastianutto, C.; Bestard, J. A.; Lahnakoski, K.; Broere, D.;
- De
- [87349] Visser, M.; Zaccolo, M.; Pozzan, T.; Ferlini, A.; Muntoni, F.;
- Patarnello,
- [87350] T.; Klamut, H. J.: Dystrophin muscle enhancer 1 is impli-

cated in

[87351] the activation of non-muscle isoforms in the skeletal muscle of patients

[87352] with X-linked dilated cardiomyopathy. Hum. Molec. Genet. 10: 2627-2635,

[87353] 2001.

[87354] 15360.

[87355] Baumbach, L. L.; Chamberlain, J. S.; Ward, P. A.; Farwell, N.

[87356] J.; Caskey, C. T.: Molecular and clinical correlation of deletion

[87357] leading to Duchenne and Becker muscular dystrophies. Neurology 39:

[87358] 465-474, 1989.

[87359] 15361.

[87360] Baumbach, L. L.; Ward, P. A.; Fenwick, R.; Caskey, C. T.: Analysis

[87361] of mutations at the Duchenne muscular dystrophy locus provides no

[87362] evidence for illegitimate recombination in deletion formation. (Abstract) Am.

[87363] J. Hum. Genet. 45 (suppl.): A173, 1989.

[87364] 15362.

[87365] Beggs, A. H.; Koenig, M.; Boyce, F. M.; Kunkel, L. M.: De-
tection

[87366] of 98% of DMD/BMD gene deletions by polymerase chain
reaction. Hum.

[87367] Genet. 86: 45–48, 1990.

[87368] 15363.

[87369] Berko, B. A.; Swift, M.: X-linked dilated cardiomyopathy.
New

[87370] Eng. J. Med. 316: 1186–1191, 1987.

[87371] 15364.

[87372] Bettecken, T.; Muller, C. R.: Identification of a 220-kb in-
sertion

[87373] into the Duchenne gene in a family with an atypical course
of muscular

[87374] dystrophy. Genomics 4: 592–596, 1989.

[87375] 15365.

[87376] Bies, R. D.: X-linked dilated cardiomyopathy. (Letter) New
Eng.

[87377] J. Med. 330: 368–369, 1994.

- [87378] 15366.
- [87379] Bies, R. D.; Caskey, C. T.; Fenwick, R.: An intact cysteine-rich
domain is required for dystrophin function. J. Clin. Invest.
90:
666-672, 1992.
- [87382] 15367.
- [87383] Bittner, R. E.; Streubel, B.; Shorny, S.; Schaden, G.; Voit, T.;
Hoger, H.: Coisogenic all-plus-one immunization: a
model for identifying
missing proteins in null-mutant conditions. Antibodies to
dystrophin
in mdx mouse after transplantation of muscle from normal coisogenic
donor. Neuropediatrics 25: 176-182, 1994.
- [87388] 15368.
- [87389] Bleyl, S. B.; Mumford, B. R.; Brown-Harrison, M.-C.;
Pagotto, L.
- [87390] T.; Carey, J. C.; Pysher, T. J.; Ward, K.; Chin, T. K.:
Xq28-linked
noncompaction of the left ventricular myocardium: prena-

tal diagnosis

[87392] and pathologic analysis of affected individuals. Am. J. Med. Genet. 72:

[87393] 257–265, 1997.

[87394] 15369.

[87395] Bleyl, S. B.; Mumford, B. R.; Thompson, V.; Carey, J. C.; Pysker,

[87396] T. J.; Chin, T. K.; Ward, K.: Neonatal, lethal noncompaction of the

[87397] left ventricular myocardium is allelic with Barth syndrome. Am. J.

[87398] Hum. Genet. 61: 868–872, 1997.

[87399] 15370.

[87400] Mohapatra, B.; Verma, S.; Shankar, S.; Suri, A.: Molecular cloning

[87401] of human testis mRNA specifically expressed in haploid germ cells,

[87402] having structural homology with the A-kinase anchoring proteins. Biochem.

[87403] Biophys. Res. Commun. 244: 540–545, 1998.

[87404] 15371.

- [87405] Turner, R. M. O.; Johnson, L. R.; Haig-Ladewig, L.; Gerton, G.
- [87406] L.; Moss, S. B.: An X-linked gene encodes a major human sperm fibrous
- [87407] sheath protein, hAKAP82: genomic organization, protein kinase A-RII
- [87408] binding, and distribution of the precursor in the sperm tail. J.
- [87409] Biol. Chem. 273: 32135-32141, 1998.
- [87410] 15372.
- [87411] Dever, T. E.; Wei, C.-L.; Benkowski, L. A.; Browning, K.; Merrick,
- [87412] W. C.; Hershey, J. W. B.: Determination of the amino acid sequence
- [87413] of rabbit, human, and wheat germ protein synthesis factor eIF-4C by
- [87414] cloning and chemical sequencing. J. Biol. Chem. 269: 3212-3218,
- [87415] 1994.
- [87416] 15373.
- [87417] Knebelmann, B.; Breillat, C.; Forestier, L.; Arrondel, C.; Jaccassier,

- [87418] D.; Giatras, I.; Drouot, L; Deschenes, G.; Grunfeld, J.-P.; Broyer,
- [87419] M.; Gubler, M.-C.; Antignac, C.: Spectrum of mutations in the COL4A5
- [87420] collagen gene in X-linked Alport syndrome. Am. J. Hum. Genet. 59:
- [87421] 1221-1232, 1996.
- [87422] 15374.
- [87423] Perkoff, G. T.; Stephens, F. E.; Dolowitz, D. A.; Tyler, F. H.
- [87424] : A clinical study of hereditary interstitial pyelonephritis. Arch.
- [87425] Intern. Med. 88: 191-200, 1951.
- [87426] 15375.
- [87427] Lee, S.-H.; Kim, W.-H.; Kim, H.-K.; Woo, K.-M.; Nam, H.-S.; Kim,
- [87428] H.-S.; Kim, J.-G.; Cho, M.-H.: Altered expression of the fragile
- [87429] histidine triad gene in primary gastric adenocarcinomas. Biochem.
- [87430] Biophys. Res. Commun. 284: 850-855, 2001.
- [87431] 15376.

- [87432] Markkanen, A.; Heinonen, K.; Knuutila, S.; de la Chapelle, A.
- [87433] : Methotrexate-induced increase in gap formation in human chromosome
- [87434] band 3p14. Hereditas 96: 317–319, 1982.
- [87435] 15377.
- [87436] Markkanen, A.; Knuutila, S.; de la Chapelle, A.: Inducible fragile
- [87437] site on chromosome 3. (Letter) Hum. Genet. 65: 217 only, 1983.
- [87438] 15378.
- [87439] Bell, M. V.; Hirst, M. C.; Nakahori, Y.; MacKinnon, R. N.; Roche,
- [87440] A.; Flint, T. J.; Jacobs, P. A.; Tommerup, N.; Tranebjaerg, L.; Froster-Iskenius,
- [87441] U.; Kerr, B.; Turner, G.; Lindenbaum, R. H.; Winter, R.; Pembrey,
- [87442] M.; Thibodeau, S.; Davies, K. E.: Physical mapping across the fragile
- [87443] X: hypermethylation and clinical expression of the fragile X syndrome. Cell 64:
- [87444] 861–866, 1991.

- [87445] 15379.
- [87446] Matsumoto, T.; Imamura, O.; Yamabe, Y.; Kuromitsu, J.;
Tokutake,
- [87447] Y.; Shimamoto, A.; Suzuki, N.; Satoh, M.; Kitao, S.;
Ichikawa, K.;
- [87448] Kataoka, H.; Sugawara, K.; Thomas, W.; Mason, B.; Tsuchi-
hashi, Z.;
- [87449] Drayna, D.; Sugawara, M.; Sugimoto, M.; Furuichi, Y.;
Goto, M.: Mutation
- [87450] and haplotype analyses of the Werner's syndrome gene
based on its
- [87451] genomic structure: genetic epidemiology in the Japanese
population. Hum.
- [87452] Genet. 100: 123–130, 1997.
- [87453] 15380.
- [87454] Filmus, J.; Shi, W.; Wong, Z. M.; Wong, M. J.: Identification
- [87455] of a new membrane-bound heparan sulphate proteogly-
can. Biochem. J. 311:
- [87456] 561–565, 1995.
- [87457] 15381.
- [87458] Zhou, J.; Hertz, J. M.; Tryggvason, K.: Mutation in the al-

pha-5(IV)

[87459] collagen chain in juvenile-onset Alport syndrome without hearing loss

[87460] or ocular lesions: detection by denaturing gradient gel electrophoresis

[87461] of a PCR product. *Am. J. Hum. Genet.* 50: 1291-1300, 1992.

[87462] 15382.

[87463] Bohler, M. C.; Seger, R. A.; Mouy, R.; Vilmer, E.; Fischer, A.;

[87464] Griscelli, C.: A study of 25 patients with chronic granulomatous

[87465] disease: a new classification by correlating respiratory burst, cytochrome

[87466] b and flavoprotein. *J. Clin. Immun.* 6: 136-145, 1986.

[87467] 15383.

[87468] Basler, E.; Grompe, M.; Parenti, G.; Caskey, C. T.; Ballabio,

[87469] A.: Identification of point mutations in three patients with steroid

[87470] sulfatase deficiency. (Abstract) *Am. J. Hum. Genet.* 47 (suppl.):

[87471] A150, 1990.

[87472] 15384.

[87473] Arranz, V.; Harper, F.; Florentin, Y.; Puvion, E.; Kress, M.;
Ernoul-Lange,

[87474] M.: Human and mouse MOK2 proteins are associated with
nuclear ribonucleoprotein

[87475] components and bind specifically to RNA and DNA
through their zinc

[87476] finger domains. *Molec. Cell. Biol.* 17: 2116–2126, 1997.

[87477] 15385.

[87478] Arranz, V.; Kress, M.; Ernoul-Lange, M.: Localization of
zinc

[87479] finger Mok2 gene to mouse chromosome 6, a new region
of homology with

[87480] human chromosome 19. *Mammalian Genome* 7: 77–78,
1996.

[87481] 15386.

[87482] Ernoul-Lange, M.; Arranz, V.; Le Conait, M.; Berger, R.;
Kress,

[87483] M.: Human and mouse Kruppel-like (MOK2) orthologue
genes encode two

[87484] different zinc finger proteins. *J. Molec. Evol.* 41: 784–794,
1995.

[87485] 15387.

[87486] Yasunaga, S.; Grati, M.; Cohen-Salmon, M.; El-Amraoui, A.; Mustapha,

[87487] M.; Salem, N.; El-Zir, E.; Loiselet, J.; Petit, C.: A mutation in

[87488] OTOF, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic

[87489] form of deafness. *Nature Genet.* 21: 363-369, 1999.

[87490] 15388.

[87491] Berry, A.; Scott, H. S.; Kudoh, J.; Talior, I.; Korostishevsky,

[87492] M.; Wattenhofer, M.; Guipponi, M.; Barras, C.; Rossier, C.; Shibuya,

[87493] K.; Wang, J.; Kawasaki, K.; Asakawa, S.; Minoshima, S.; Shimizu, N.;

[87494] Antonarakis, S.; Bonne-Tamir, B.: Refined localization of autosomal

[87495] recessive nonsyndromic deafness DFNB10 locus using 34 novel microsatellite

[87496] markers, genomic structure, and exclusion of six known genes in the

[87497] region. *Genomics* 68: 22-29, 2000.

[87498] 15389.

- [87499] Scott, H. S.; Kudoh, J.; Wattenhofer, M.; Shibuya, K.; Berry, A.;
- [87500] Chrast, R.; Guipponi, M.; Wang, J.; Kawasaki, K.; Asakawa, S.; Minoshima,
- [87501] S.; Younus, F.; and 10 others: Insertion of beta-satellite repeats
- [87502] identifies a transmembrane protease causing both congenital and childhood
- [87503] onset autosomal recessive deafness. *Nature Genet.* 27: 59–63, 2001.
- [87504] 15390.
- [87505] Good, P. J.; Chen, Q.; Warner, S. J.; Herring, D. C.: A family
- [87506] of human RNA-binding proteins related to the *Drosophila* Bruno transcriptional
- [87507] regulator. *J. Biol. Chem.* 275: 28583–28592, 2000.
- [87508] 15391.
- [87509] Roberts, R.; Timchenko, N. A.; Miller, J. W.; Reddy, S.; Caskey,
- [87510] C. T.; Swanson, M. S.; Timchenko, L. T.: Altered phosphorylation
- [87511] and intracellular distribution of a (CUG)_n triplet repeat RNA-binding

- [87512] protein in patients with myotonic dystrophy and in myotonic protein
- [87513] kinase knockout mice. *Proc. Nat. Acad. Sci.* 94: 13221–13226, 1997.
- [87514] 15392.
- [87515] Li, Z.; Jiang, H.; Xie, W.; Zhang, Z.; Smrcka, A. V.; Wu, D.:
- [87516] Roles of PLC- β 2 and - β 3 and PI3K- γ in chemoattractant-mediated
- [87517] signal transduction. *Science* 287: 1046–1049, 2000.
- [87518] 15393.
- [87519] Makita, N.; Sloan-Brown, K.; Weghuis, D. O.; Ropers, H. H.; George,
- [87520] A. L., Jr.: Genomic organization and chromosomal assignment of the
- [87521] human voltage-gated Na(+) channel β -1 subunit gene (SCN1B). *Genomics* 23:
- [87522] 628–634, 1994.
- [87523] 15394.
- [87524] Scheffer, I. E.; Berkovic, S. F.: Generalized epilepsy with febrile
- [87525] seizures plus: a genetic disorder with heterogeneous clin-

ical phenotypes. Brain 120:

[87526] 479–490, 1997.

[87527] 15395.

[87528] Wallace, R. H.; Wang, D. W.; Singh, R.; Scheffer, I. E.;
George,

[87529] A. L., Jr.; Phillips, H. A.; Saar, K.; Reis, A.; Johnson, E. W.;
Sutherland,

[87530] G. R.; Berkovic, S. F.; Mulley, J. C.: Febrile seizures and
generalized

[87531] epilepsy associated with a mutation in the Na(+)-channel
beta-1 subunit

[87532] gene SCN1B. Nature Genet. 19: 366–370, 1998.

[87533] 15396.

[87534] Gassmann, M.; Casagrande, F.; Orioli, D.; Simon, H.; Lai,
C.; Klein,

[87535] R.; Lemke, G.: Aberrant neural and cardiac development in
mice lacking

[87536] the ErbB4 neuregulin receptor. Nature 378: 390–394,
1995.

[87537] 15397.

[87538] Golding, J. P.; Trainor, P.; Krumlauf, R.; Gassmann, M.: De-

fects

[87539] in pathfinding by cranial neural crest cells in mice lacking the neuregulin

[87540] receptor ErbB4. *Nature Cell Biol.* 2: 103–109, 2000.

[87541] 15398.

[87542] Zimonjic, D. B.; Alimandi, M.; Miki, T.; Popescu, N. C.; Kraus,

[87543] M. H.: Localization of the human HER4/erbB-4 gene to chromosome 2. *Oncogene* 10:

[87544] 1235–1237, 1995.

[87545] 15399.

[87546] Adibi, S. A.: The oligopeptide transporter (Pept-1) in human intestine:

[87547] biology and function. *Gastroenterology* 113: 332–340, 1997.

[87548] 15400.

[87549] Fei, Y. -J.; Kanai, Y.; Nussberger, S.; Ganapathy, V.; Leibach,

[87550] F. H.; Romero, M. F.; Singh, S. K.; Boron, W. F.; Hediger, M. A.:

[87551] Expression cloning of a mammalian proton-coupled

oligopeptide transporter. Nature 368:

[87552] 563–566, 1994.

[87553] 15401.

[87554] Liang, R.; Fei, Y.-J.; Prasad, P. D.; Ramamoorthy, S.; Han, H.;

[87555] Yang–Feng, T. L.; Hediger, M. A.; Ganapathy, V.; Leibach, F. H.:

[87556] Human intestinal H(+)/peptide cotransporter: cloning, functional expression,

[87557] and chromosomal localization. J. Biol. Chem. 270: 6456–6463, 1995.

[87558] 15402.

[87559] Perry, S.; Harries, H.; Scholfield, C.; Lock, T.; King, L.; Gibson,

[87560] G.; Goldfarb, P.: Molecular cloning and expression of a cDNA for

[87561] human kidney cysteine conjugate beta-lyase. FEBS Lett. 360: 277–280,

[87562] 1995.

[87563] 15403.

[87564] Kaul, S. C.; Wadhwa, R.; Matsuda, Y.; Hensler, P. J.;

Pereira-Smith,

[87565] O. M.; Komatsu, Y.; Mitsui, Y.: Mouse and human chromosomal assignments

[87566] of mortalin, a novel member of the murine hsp70 family of proteins. FEBS

[87567] Lett. 361: 269-272, 1995.

[87568] 15404.

[87569] Heiber, M.; Docherty, J. M.; Shah, G.; Nguyen, T.; Cheng, R.; Heng,

[87570] H. H. Q.; Marchese, A.; Tsui, L.-C.; Shi, X.; George, S. R.; O'Dowd,

[87571] B. F.: Isolation of three novel human genes encoding G protein-coupled

[87572] receptors. DNA Cell Biol. 14: 25-35, 1995.

[87573] 15405.

[87574] Mahadevan, M. S.; Baird, S.; Bailly, J. E.; Shutler, G. G.; Sabourin,

[87575] L. A.; Tsilfidis, C.; Neville, C. E.; Narang, M.; Korneluk, R. G.

[87576] : Isolation of a novel G protein-coupled receptor (GPR4) localized

[87577] to chromosome 19q13.3. Genomics 30: 84-88, 1995.

[87578] 15406.

[87579] Ye, R. D.; Prossnitz, E. R.; Zou, A.; Cochrane, C. G.: Characterization

[87580] of a human cDNA that encodes a functional receptor for platelet activating

[87581] factor. Biochem. Biophys. Res. Commun. 180: 105–111, 1991.

[87582] 15407.

[87583] Hossain, A.; Saunders, G. F.: The human sex-determining gene

[87584] SRY is a direct target of WT1. J. Biol. Chem. 276: 16817–16823,

[87585] 2001.

[87586] 15408.

[87587] Iida, T.; Nakahori, Y.; Komaki, R.; Mori, E.; Hayashi, N.; Tsutsumi,

[87588] O.; Taketani, Y.; Nakagome, Y.: A novel missense mutation in the

[87589] HMG box of the SRY gene in a patient with XY sex reversal. Hum. Molec.

[87590] Genet. 3: 1437–1438, 1994.

- [87591] 15409.
- [87592] Jager, R. J.; Harley, V. R.; Pfeiffer, R. A.; Goodfellow, P. N.;
- [87593] Scherer, G.: A familial mutation in the testis-determining
gene SRY
- [87594] shared by both sexes. Hum. Genet. 90: 350–355, 1992.
- [87595] 15410.
- [87596] Kawakura, K.; Miyake, Y.-I.; Murakami, R.-K.; Kondoh, S.;
- Hirata,
- [87597] T.-I.; Kaneda, Y.: Deletion of the SRY region on the Y
chromosome
- [87598] detected in bovine gonadal hypoplasia (XY female) by PCR.
Cytogenet.
- [87599] Cell Genet. 72: 183–184, 1996.
- [87600] 15411.
- [87601] Lahr, G.; Maxson, S. C.; Mayer, A.; Just, W.; Pilgrim, C.;
- Reisert,
- [87602] I.: Transcription of the Y chromosomal gene, Sry, in adult
mouse
- [87603] brain. Molec. Brain Res. 33: 179–182, 1995.
- [87604] 15412.

- [87605] Li, B.; Zhang, W.; Chan, G.; Jancso-Radek, A.; Liu, S.; Weiss,
Weiss,
- [87606] M. A.: Human sex reversal due to impaired nuclear localization of
- [87607] SRY: a clinical correlation. J. Biol. Chem. 276:
46480-46484, 2001.
- [87608] 15413.
- [87609] Lovell-Badge, R.; Robertson, E.: XY female mice resulting from
- [87610] a heritable mutation in the primary testis-determining gene, Tdy. Development 109:
- [87611] 635-646, 1990.
- [87612] 15414.
- [87613] Margarit, E.; Coll, M. D.; Oliva, R.; Gomez, D.; Soler, A.; Ballesta,
- [87614] F.: SRY gene transferred to the long arm of the X chromosome in a
- [87615] Y-positive XX true hermaphrodite. Am. J. Med. Genet. 90:
25-28,
- [87616] 2000.
- [87617] 15415.

- [87618] Mittwoch, U.: Sex determination and sex reversal: genotype, phenotype,
[87619] dogma and semantics. Hum. Genet. 89: 467–479, 1992.
[87620] 15416.
- [87621] Muller, J.; Schwartz, M.; Skakkebaek, N. E.: Analysis of the
[87622] sex-determining region of the Y-chromosome (SRY) in
sex reversed patients:
[87623] point-mutation in SRY causing sex-reversion in a 46,XY
female. J.
[87624] Clin. Endocr. Metab. 75: 331–333, 1992.
[87625] 15417.
- [87626] Peters, R.; King, C.-Y.; Ukiyama, E.; Falsafi, S.; Donahoe, P.
[87627] K.; Weiss, M. A.: An SRY mutation causing human sex re-
versal resolves
[87628] a general mechanism of structure-specific DNA recogni-
tion: application
[87629] to the four-way DNA junction. Biochemistry 34:
4569–4576, 1995.
[87630] 15418.
- [87631] Schaffler, A.; Barth, N.; Winkler, K.; Zietz, B.; Rummele, P.;

- [87632] Knuchel, R.; Scholmerich, J.; Palitzsch, K.-D.: Identification of
- [87633] a new missense mutation (gly95glu) in a highly conserved codon within
- [87634] the high-mobility group box of the sex-determining region Y gene:
- [87635] report on a 46,XY female with gonadal dysgenesis and yolk-sac tumor. J.
- [87636] Clin. Endocr. Metab. 85: 2287-2292, 2000.
- [87637] 15419.
- [87638] Schmitt-Ney, M.; Thiele, H.; Kaltwasser, P.; Bardoni, B.; Cisternino,
- [87639] M.; Scherer, G.: Two novel SRY missense mutations reducing DNA binding
- [87640] identified in XY females and their mosaic fathers. Am. J. Hum. Genet. 56:
- [87641] 862-869, 1995.
- [87642] 15420.
- [87643] Sinclair, A. H.; Berta, P.; Palmer, M. S.; Hawkins, R.; Griffiths,
- [87644] B. L.; Smith, M.; Foster, J. W.; Frischauf, A.-M.; Lovell-Badge, R.;

- [87645] Goodfellow, P. N.: A gene from the human sex-determining region encodes
- [87646] a protein with homology to a conserved DNA-binding motif.. Nature 346:
- [87647] 240-245, 1990.
- [87648] 15421.
- [87649] Su, H.; Lau, Y.-F. C.: Identification of the transcriptional

- [87650] unit, structural organization, and promoter sequence of the human
- [87651] sex-determining region Y (SRY) gene, using a reverse genetic approach. Am.
- [87652] J. Hum. Genet. 52: 24–38, 1993.
- [87653] 15422.
- [87654] Tucker, P. K.; Lundrigan, B. L.: Rapid evolution of the sex determining
- [87655] locus in Old World mice and rats. Nature 364: 715–717, 1993.
- [87656] 15423.
- [87657] Uehara, S.; Funato, T.; Yaegashi, N.; Suzuki, H.; Sato, J.; Sasaki,
- [87658] T.; Yajima, A.: SRY mutation and tumor formation on the gonads of
- [87659] XY pure gonadal dysgenesis patients. Cancer Genet. Cytogenet. 113:
- [87660] 78–84, 1999.
- [87661] 15424.
- [87662] Uehara, S.; Hashiyada, M.; Sato, K.; Nata, M.; Funato, T.; Okamura,

- [87663] K.: Complete XY gonadal dysgenesis and aspects of the SRY genotype
- [87664] and gonadal tumor formation. J. Hum. Genet. 47: 279–284, 2002.
- [87665] 15425.
- [87666] Veitia, R.; Ion, A.; Barbaux, S.; Jobling, M. A.; Souleyreau,
- [87667] N.; Ennis, K.; Ostrer, H.; Tosi, M.; Meo, T.; Chibani, J.; Fellous,
- [87668] M.; McElreavey, K.: Mutations and sequence variants in the testis-determining
- [87669] region of the Y chromosome in individuals with a 46,XY female phenotype. Hum.
- [87670] Genet. 99: 648–652, 1997.
- [87671] 15426.
- [87672] Werner, M. H.; Huth, J. R.; Gronenborn, A. M.; Clore, G. M.:
- [87673] Molecular basis of human 46X,Y (sic) sex reversal revealed from the
- [87674] three-dimensional solution structure of the human SRY-DNA complex. Cell 81:
- [87675] 705–714, 1995.
- [87676] 15427.

- [87677] Whitfield, L. S.; Lovell-Badge, R.; Goodfellow, P. N.: Rapid
[87678] sequence evolution of the mammalian sex-determining
gene SRY. *Nature* 364:
[87679] 713–715, 1993.
[87680] 15428.
- [87681] Zeng, Y.; Ren, Z.; Zhang, M.; Huang, Y.; Zeng, F.; Huang,
S.:
[87682] A new de novo mutation (A113T) in HMG box of the SRY
gene leads to
[87683] XY gonadal dysgenesis. *J. Med. Genet.* 30: 655–657,
1993.
[87684] 15429.
- [87685] Zenteno, J. C.; Lopez, M.; Vera, C.; Mendez, J. P.; Kofman-
Alfaro,
[87686] S.: Two SRY-negative XX male brothers without genital
ambiguity. *Hum.*
[87687] *Genet.* 100: 606–610, 1997.
[87688] 15430.
- [87689] Henkemeyer, M.; Orioli, D.; Henderson, J. T.; Saxton, T.
M.; Roder,
[87690] J.; Pawson, T.; Klein, R.: Nuk controls pathfinding of com-

missural

- [87691] axons in the mammalian central nervous system. Cell 86: 35–46, 1996.
- [87692] 15431.
- [87693] Ikegaki, N.; Tang, X. X.; Liu, X.-G.; Biegel, J. A.; Allen, C.;
- [87694] Yoshioka, A.; Sulman, E. P.; Brodeur, G. M.; Pleasure, D. E.: Molecular
- [87695] characterization and chromosomal localization of DRT (EPHT3): a developmentally
- [87696] regulated human protein-tyrosine kinase gene of the EPH family. Hum.
- [87697] Molec. Genet. 4: 2033–2045, 1995.
- [87698] 15432.
- [87699] Saito, T.; Seki, N.; Matsuda, Y.; Kitahara, M.; Murata, M.; Kanda,
- [87700] N.; Nomura, N.; Yamamoto, T.; Hori, T.: Identification of the human
- [87701] ERK gene as a putative receptor tyrosine kinase and its chromosomal
- [87702] localization to 1p36.1: a comparative mapping of human, mouse, and
- [87703] rat chromosomes. Genomics 26: 382–384, 1995.

- [87704] 15433.
- [87705] Wybenga-Groot, L. E.; Baskin, B.; Ong, S. H.; Tong, J.; Pawson,
- [87706] T.; Sicheri, F.: Structural basis for autoinhibition of the EphB2
- [87707] receptor tyrosine kinase by the unphosphorylated jux-
- tamembrane region. Cell 106:
- [87708] 745-757, 2001.
- [87709] 15434.
- [87710] Elpeleg, O. N.; Hammerman, C.; Saada, A.; Shaag, A.; Golzand, E.;
- [87711] Hochner-Celnikier, D.; Berger, I.; Nadjari, M.: Antenatal
- presentation
- [87712] of carnitine palmitoyltransferase II deficiency. Am. J. Med.
- Genet. 102:
- [87713] 183-187, 2001.
- [87714] 15435.
- [87715] Finocchiaro, G.; Taroni, F.; Rocchi, M.; Liras Martin, A.; Colombo,
- [87716] I.; Tarelli, G. T.; DiDonato, S.: cDNA cloning, sequence
- analysis,

- [87717] and chromosomal localization of the gene for human carnitine palmitoyltransferase. Proc.
- [87718] Nat. Acad. Sci. 88: 661–665, 1991.
- [87719] 15436.
- [87720] Haap, M.; Thamer, C.; Machann, J.; Tschritter, O.; Loblein, K.;
- [87721] Kellerer, M.; Schick, F.; Jacob, S.; Haring, H.–U.; Stumvoll, M.:
- [87722] Metabolic characterization of a woman homozygous for the ser113leu
- [87723] missense mutation in carnitine palmitoyl transferase II. J. Clin.
- [87724] Endocr. Metab. 87: 2139–2143, 2002.
- [87725] 15437.
- [87726] Handig, I.; Dams, E.; Taroni, F.; Van Laere, S.; de Barse, T.;
- [87727] Willems, P. J.: Inheritance of the S113L mutation within an inbred
- [87728] family with carnitine palmitoyltransferase enzyme deficiency. Hum.
- [87729] Genet. 97: 291–293, 1996.
- [87730] 15438.

- [87731] Martin, M. A.; Rubio, J. C.; De Bustos, F.; Del Hoyo, P.; Campos,
- [87732] Y.; Garcia, A.; Bornstein, B.; Cabello, A.; Arenas, J.: Molecular
- [87733] analysis in Spanish patients with muscle carnitine palmitoyltransferase
- [87734] deficiency. *Muscle Nerve* 22: 941–943, 1999.
- [87735] 15439.
- [87736] Minoletti, F.; Colombo, I.; Liras Martin, A.; DiDonato, S.; Taroni,
- [87737] F.; Finocchiaro, G.; Pandolfo, M.: Localization of the human gene
- [87738] for carnitine palmitoyltransferase to 1p13–p11 by nonradioactive in
- [87739] situ hybridization. *Genomics* 13: 1372–1374, 1992.
- [87740] 15440.
- [87741] Montermini, L.; Wang, H.; Verderio, E.; Taroni, F.; DiDonato,
- [87742] S.; Finocchiaro, G: Identification of 5–prime regulatory regions
- [87743] of the human carnitine palmitoyltransferase II gene. *Biochim. Biophys.*

- [87744] Acta 1219: 237–240, 1994.
- [87745] 15441.
- [87746] Taroni, F.; Verderio, E.; Fiorucci, S.; Cavadini, P.; Finoc-
chiaro,
- [87747] G.; Uziel, G.; Lamantea, E.; Gellera, C.; DiDonato, S.:
Molecular
- [87748] characterization of inherited carnitine palmitoyltrans-
ferase II deficiency. Proc.
- [87749] Nat. Acad. Sci. 89: 8429–8433, 1992.
- [87750] 15442.
- [87751] Verderio, E.; Cavadini, P.; Montermini, L.; Wang, H.;
Lamantea,
- [87752] E.; Finocchiaro, G.; DiDonato, S.; Gellera, C.; Taroni, F.:
Carnitine
- [87753] palmitoyltransferase II deficiency: structure of the gene
and characterization
- [87754] of two novel disease-causing mutations. Hum. Molec.
Genet. 4: 19–29,
- [87755] 1995.
- [87756] 15443.
- [87757] Vladutiu, G. D.; Bennett, M. J.; Smail, D.; Wong, L.-J.; Tag-

gart,

[87758] R. T.; Lindsley, H. B.: A variable myopathy associated with heterozygosity

[87759] for the R503C mutation in the carnitine palmitoyltransferase II gene. Molec.

[87760] Genet. Metab. 70: 134-141, 2000.

[87761] 15444.

[87762] Vladutiu, G. D.; Taggart, R. T.; Smail, D.; Lindsley, H. B.; Hogan,

[87763] K.: A carnitine palmitoyl transferase II (CPT2) arg503cys mutation

[87764] confers malignant hyperthermia and variable myopathy. (Abstract) Am.

[87765] J. Hum. Genet. 63 (suppl.): A5 only, 1998.

[87766] 15445.

[87767] Yamamoto, S.; Abe, H.; Kohgo, T.; Ogawa, A.; Ohtake, A.; Hayashibe,

[87768] H.; Sakuraba, H.; Suzuki, Y.; Aramaki, S.; Takayanagi, M.; Hasegawa,

[87769] S.; Niimi, H.: Two novel gene mutations (glu174-to-lys, phe383-to-tyr)

[87770] causing the 'hepatic' form of carnitine palmitoyltrans-

ferase II deficiency. Hum.

[87771] Genet. 98: 116–118, 1996.

[87772] 15446.

[87773] Johnson, D. F.; Hamon, M.; Fischel-Ghodsian, N.: Characterization

[87774] of the human mitochondrial ribosomal S12 gene. Genomics 52: 363–368,

[87775] 1998.

[87776] 15447.

[87777] Shah, Z. H.; Migliosi, V.; Miller, S. C. M.; Wang, A.; Friedman,

[87778] T. B.; Jacobs, H. T.: Chromosomal locations of three human nuclear

[87779] genes (RPSM12, TUFM, and AFG3L1) specifying putative components of

[87780] the mitochondrial gene expression apparatus. Genomics 48: 384–388,

[87781] 1998.

[87782] 15448.

[87783] Zhu, K.; Baudhuin, L. M.; Hong, G.; Williams, F. S.; Cristina,

[87784] K. L.; Kabarowski, J. H. S.; Witte, O. N.; Xu, Y.: Sphingo-

sylphosphorylcholine

[87785] and lysophosphatidylcholine are ligands for the G protein-coupled

[87786] receptor GPR4. J. Biol. Chem. 276: 41325–41335, 2001.

[87787] 15449.

[87788] Vinik, B. S.; Kay, E. S.; Fiedorek, F. T., Jr.: Mapping of the

[87789] MEK kinase gene (Mekk) to mouse chromosome 13 and human chromosome

[87790] 5. Mammalian Genome 6: 782–783, 1995.

[87791] 15450.

[87792] Yujiri, T.; Sather, S.; Fanger, G. R.; Johnson, G. L.: Role of

[87793] MEKK1 in cell survival and activation of JNK and ERK pathways defined

[87794] by targeted gene disruption. Science 282: 1911–1914, 1998.

[87795] 15451.

[87796] Arriza, J. L.; Weinberger, C.; Cerelli, G.; Glaser, T. M.; Handelin,

[87797] B. L.; Housman, D. E.; Evans, R. M.: Cloning of human mineralocorticoid

[87798] receptor complementary DNA: structural and functional

kinship with

[87799] the glucocorticoid receptor. *Science* 237: 268–275, 1987.

[87800] 15452.

[87801] Berger, S.; Bleich, M.; Schmid, W.; Cole, T. J.; Peters, J.;
Watanabe,

[87802] H.; Kriz, W.; Warth, R.; Greger, R.; Schutz, G.: Mineralocor-
ticoid

[87803] receptor knockout mice: pathophysiology of Na+
metabolism. *Proc.*

[87804] *Nat. Acad. Sci.* 95: 9424–9429, 1998.

[87805] 15453.

[87806] Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[87807] M.; Nowak, N. J.; Shows, T. B.: The human mineralocorti-
coid receptor

[87808] gene (MLR) is located on chromosome 4 at q31.2. *Cyto-*
genet. Cell Genet. 52:

[87809] 83–84, 1989.

[87810] 15454.

[87811] Geller, D. S.; Farhi, A.; Pinkerton, N.; Fradley, M.; Moritz,
M.;

[87812] Spitzer, A.; Meinke, G.; Tsai, F. T. F.; Sigler, P. B.; Lifton, R.

- [87813] P.: Activating mineralocorticoid receptor mutation in hypertension
- [87814] exacerbated by pregnancy. Science 289: 119–123, 2000.
- [87815] 15455.
- [87816] Geller, D. S.; Rodriguez–Soriano, J.; Vallo Boado, A.; Schifter,
- [87817] S.; Bayer, M.; Chang, S. S.; Lifton, R. P.: Mutations in the mineralocorticoid
- [87818] receptor gene cause autosomal dominant pseudohypoaldosteronism type
- [87819] I. Nature Genet. 19: 279–281, 1998.
- [87820] 15456.
- [87821] Hellal–Levy, C.; Fagart, J.; Souque, A.; Wurtz, J.–M.; Moras, D.;
- [87822] Rafestin–Oblin, M.–E.: Crucial role of the H11–H12 loop in stabilizing
- [87823] the active conformation of the human mineralocorticoid receptor. Molec.
- [87824] Endocr. 14: 1210–1221, 2000.
- [87825] 15457.
- [87826] Le Menuet, D.; Isnard, R.; Bichara, M.; Viengchareun, S.;

Muffat-Joly,

- [87827] M.; Walker, F.; Zennaro, M.-C.; Lombes, M.: Alteration of cardiac
- [87828] and renal functions in transgenic mice overexpressing human mineralocorticoid
- [87829] receptor. J. Biol. Chem. 276: 38911–38920, 2001.
- [87830] 15458.
- [87831] Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor, J.
- [87832] M.: Regional chromosomal assignment of the human mineralocorticoid
- [87833] receptor gene to 4q31.1. Hum. Genet. 85: 130–132, 1990.
- [87834] 15459.
- [87835] Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor,
- [87836] J. M.: Regional chromosomal assignment of the human mineralocorticoid
- [87837] receptor gene to 4q31.1. (Abstract) Cytogenet. Cell Genet. 51: 1048,
- [87838] 1989.
- [87839] 15460.
- [87840] Bristow, J.; Tee, M. K.; Gitelman, S. E.; Mellon, S. H.; Miller,

- [87841] W. L.: Tenascin-X: a novel extracellular matrix protein encoded by
- [87842] the human XB gene overlapping P450c21B. *J. Cell Biol.* 122: 265–278,
- [87843] 1993.
- [87844] 15461.
- [87845] Jay, P.; Goze, C.; Marsollier, C.; Taviaux, S.; Hardelin, J.-P.;
- [87846] Koopman, P.; Berta, P.: The human SOX11 gene: cloning, chromosomal
- [87847] assignment and tissue expression. *Genomics* 29: 541–545, 1995.
- [87848] 15462.
- [87849] Laudet, V.; Stehelin, D.; Clevers, H.: Ancestry and diversity
- [87850] of the HMG box superfamily. *Nucleic Acids Res.* 21: 2493–2501, 1993.
- [87851] 15463.
- [87852] Weigt, C.; Gaertner, A.; Wegner, A.; Korte, H.; Meyer, H. E.:
- [87853] Occurrence of an actin-inserting domain in tensin. *J. Molec. Biol.* 227:
- [87854] 593–595, 1992.

- [87855] 15464.
- [87856] Charest, A.; Wagner, J.; Muise, E. S.; Heng, H. H. Q.; Tremblay,
- [87857] M. L.: Structure of the murine MPTP-PEST gene: genomic organization
- [87858] and chromosomal mapping. *Genomics* 28: 501-507, 1995.
- [87859] 15465.
- [87860] Takekawa, M.; Itoh, F.; Hinoda, Y.; Adachi, M.; Ariyama, T.; Inazawa,
- [87861] J.; Imai, K.; Yachi, A.: Chromosomal localization of the protein
- [87862] tyrosine phosphatase G1 gene and characterization of the aberrant
- [87863] transcripts in human colon cancer cells. *FEBS Lett.* 339: 222-228,
- [87864] 1994.
- [87865] 15466.
- [87866] Takekawa, M.; Itoh, F.; Hinoda, Y.; Arimura, Y.; Toyota, M.; Sekiya,
- [87867] M.; Adachi, M.; Imai, K.; Yachi, A.: Cloning and characterization

- [87868] of a human cDNA encoding a novel putative cytoplasmic protein-tyrosine-phosphatase. *Biochem.*
- [87869] *Biophys. Res. Commun.* 189: 1223-1230, 1992.
- [87870] 15467.
- [87871] Yang, Q.; Co, D.; Sommercorn, J.; Tonks, N. K.: Cloning and expression
- [87872] of PTP-PEST: a novel, human, nontransmembrane protein tyrosine phosphatase. *J.*
- [87873] *Biol. Chem.* 268: 6622-6628, 1993.
- [87874] 15468.
- [87875] Coopman, P. J. P.; Do, M. T. H.; Barth, M.; Bowden, E. T.; Hayes,
- [87876] A. J.; Basyuk, E.; Blancato, J. K.; Vezza, P. R.; McLeskey, S. W.;
- [87877] Mangeat, P. H.; Mueller, S. C.: The Syk tyrosine kinase suppresses
- [87878] malignant growth of human breast cancer cells. *Nature* 406: 742-747,
- [87879] 2000.
- [87880] 15469.
- [87881] Inatome, R.; Yanagi, S.; Takano, T.; Yamamura, H.: A criti-

cal

[87882] role for Syk in endothelial cell proliferation and migration.

Biochem.

[87883] Biophys. Res. Commun. 286: 195–199, 2001.

[87884] 15470.

[87885] Mocsai, A.; Zhou, M.; Meng, F.; Tybulewicz, V. L.; Lowell,

C. A.

[87886] : Syk is required for integrin signaling in neutrophils. Im-

munity 16:

[87887] 547–558, 2002.

[87888] 15471.

[87889] Muller, B.; Cooper, L.; Terhorst, C.: Molecular cloning of

the

[87890] human homologue to the pig protein-tyrosine kinase syk.

Immunogenetics 39:

[87891] 359–362, 1994.

[87892] 15472.

[87893] Taniguchi, T.; Kobayashi, T.; Kondo, J.; Takahashi, K.;

Nakamura,

[87894] H.; Suzuki, J.; Nagai, K.; Yamada, T.; Nakamura, S.; Yama-

mura, H.

- [87895] : Molecular cloning of a porcine gene syk that encodes a 72-kDa protein-tyrosine
- [87896] kinase showing high susceptibility to proteolysis. J. Biol. Chem. 266:
- [87897] 15790–15796, 1991.
- [87898] 15473.
- [87899] Zioncheck, T. F.; Harrison, M. L.; Isaacson, C. C.; Geahlen, R.
- [87900] L.: Generation of an active protein-tyrosine kinase from lymphocytes
- [87901] by proteolysis. J. Biol. Chem. 263: 19195–19202, 1988.
- [87902] 15474.
- [87903] Satre, M. A.; Zgombic-Knight, M.; Duester, G.: The complete structure
- [87904] of human class IV alcohol dehydrogenase (retinol dehydrogenase) determined
- [87905] from the ADH7 gene. J. Biol. Chem. 269: 15606–15612, 1994.
- [87906] 15475.
- [87907] Yokoyama, H.; Baraona, E.; Lieber, C. S.: Molecular cloning and

- [87908] chromosomal localization of the ADH7 gene encoding human class IV
- [87909] (sigma) ADH. Genomics 31: 243–245, 1996.
- [87910] 15476.
- [87911] Zgombic-Knight, M.; Foglio, M. H.; Duester, G.: Genomic structure
- [87912] and expression of the ADH7 gene encoding human class IV alcohol dehydrogenase,
- [87913] the form most efficient for retinol metabolism in vitro. J. Biol. Chem. 270: 4305–4311, 1995.
- [87914] 15477.
- [87915] Bashir, M. M.; Abrams, W. R.; Tucker, T.; Sellinger, B.; Budarf, M.; Emanuel, B.; Rosenbloom, J.: Molecular cloning and characterization
- [87916] of the bovine and human tuftelin genes. Connect. Tissue Res. 39: 13–24, 1998.
- [87917] 15478.
- [87918] Deutsch, D.: Structure and function of enamel gene prod-

ucts. Anat.

[87922] Rec. 224: 189–210, 1989.

[87923] 15479.

[87924] Beguin, P.; Nagashima, K.; Gonoï, T.; Shibasaki, T.; Takahashi,

[87925] K.; Kashima, Y.; Ozaki, N.; Geering, K.; Iwanaga, T.; Seino, S.:

[87926] Regulation of Ca(2+) channel expression at the cell surface by the

[87927] small G-protein kir/Gem. Nature 411: 701–706, 2001.

[87928] 15480.

[87929] Maguire, J.; Santoro, T.; Jensen, P.; Siebenlist, U.; Yewdell,

[87930] J.; Kelly, K.: GEM: an induced, immediate early protein belonging

[87931] to the Ras family. Science 265: 241–244, 1994.

[87932] 15481.

[87933] Santoro, T.; Maguire, J.; McBride, O. W.; Avraham, K. B.; Copeland,

[87934] N. G.; Jenkins, N. A.; Kelly, K.: Chromosomal organization and transcriptional

[87935] regulation of human GEM and localization of the human

and mouse Gem

[87936] loci encoding an inducible Ras-like protein. Genomics 30:
558-564,

[87937] 1995.

[87938] 15482.

[87939] Inoue, I.; Taniuchi, I.; Kitamura, D.; Jenkins, N. A.; Gilbert,

[87940] D. J.; Copeland, N. G.; Watanabe, T.: Characteristics of the
mouse

[87941] genomic histamine H1 receptor gene. Genomics 36:
178-181, 1996.

[87942] 15483.

[87943] Le Coniat, M.; Traiffort, E.; Ruat, M.; Arrang, J.-M.; Berger,

[87944] R.: Chromosomal localization of the human histamine
H1-receptor gene. Hum.

[87945] Genet. 94: 186-188, 1994.

[87946] 15484.

[87947] Ma, R. Z.; Gao, J.; Meeker, N. D.; Fillmore, P. D.; Tung, K.
S.

[87948] K.; Watanabe, T.; Zachary, J. F.; Offner, H.; Blankenhorn, E.
P.;

[87949] Teuscher, C.: Identification of Bphs, an autoimmune dis-

ease locus,

[87950] as histamine receptor H-1. Science 297: 620-623, 2002.

[87951] 15485.

[87952] Yamashita, M.; Fukui, H.; Sugama, K.; Horio, Y.; Ito, S.; Mizuguchi,

[87953] H.; Wada, H.: Expression cloning of a cDNA encoding the bovine histamine

[87954] H1 receptor. Proc. Nat. Acad. Sci. 88: 11515-11519, 1991.

[87955] 15486.

[87956] Gaudino, G.; Follenzi, A.; Naldini, L.; Collesi, C.; Santoro, M.;

[87957] Gallo, K. A.; Godowski, P. J.; Comoglio, P. M.: RON is a heterodimeric

[87958] tyrosine kinase receptor activated by the HGF homologue MSP. EMBO

[87959] J. 13: 3524-3532, 1994.

[87960] 15487.

[87961] Muraoka, R. S.; Sun, W. Y.; Colbert, M. C.; Waltz, S. E.; Witte,

[87962] D. P.; Degen, J. L.; Degen, S. J. F.: The Ron/STK receptor

tyrosine

[87963] kinase is essential for peri-implantation development in the mouse. J.

[87964] Clin. Invest. 103: 1277-1285, 1999.

[87965] 15488.

[87966] Ronsin, C.; Muscatelli, F.; Mattei, M.-G.; Breathnach, R.: A novel

[87967] putative receptor protein tyrosine kinase of the met family. Oncogene 8:

[87968] 1195-1202, 1993.

[87969] 15489.

[87970] Santoro, M. M.; Collesi, C.; Grisendi, S.; Gaudino, G.; Comoglio,

[87971] P. M.: Constitutive activation of the RON gene promotes invasive

[87972] growth but not transformation. Molec. Cell. Biol. 16: 7072-7083,

[87973] 1996.

[87974] 15490.

[87975] Wang, M.-H.; Ronsin, C.; Gesnel, M.-C.; Coupey, L.; Skeel, A.;

- [87976] Leonard, E. J.; Breathnach, R.: Identification of the RON gene product
- [87977] as the receptor for the human macrophage stimulating protein. Science 266:
- [87978] 117–119, 1994.
- [87979] 15491.
- [87980] Bauer, S.; Groh, V.; Wu, J.; Steinle, A.; Phillips, J. H.; Lanier, L. L.; Spies, T.: Activation of NK cells and T cells by NKG2D, a
- [87982] receptor for stress-inducible MICA. Science 285: 727–729, 1999.
- [87983] 15492.
- [87984] Diefenbach, A.; Jensen, E. R.; Jamieson, A. M.; Raulet, D. H.:
- [87985] Rae1 and H60 ligands of the NKG2D receptor stimulate tumour immunity. Nature 413:
- [87986] 165–171, 2001.
- [87987] 15493.
- [87988] Girardi, M.; Oppenheim, D. E.; Steele, C. R.; Lewis, J. M.; Glusac,
- [87989] E.; Filler, R.; Hobby, P.; Sutton, B.; Tigelaar, R. E.; Hayday,

A.

[87990] C.: Regulation of cutaneous malignancy by gamma-delta T cells. Science 294:

[87991] 605–609, 2001.

[87992] 15494.

[87993] Groh, V.; Rhinehart, R.; Randolph-Habecker, J.; Topp, M. S.; Riddell,

[87994] S. R.; Spies, T.: Costimulation of CD8-alpha-beta T cells by NKG2D

[87995] via engagement by MIC induced on virus-infected cells. Nature Immun. 2:

[87996] 255–260, 2001.

[87997] 15495.

[87998] Groh, V.; Steinle, A.; Bauer, S.; Spies, T.: Recognition of stress-induced

[87999] MHC molecules by intestinal epithelial gamma-delta T cells. Science 279:

[88000] 1737–1740, 1998.

[88001] 15496.

[88002] Groh, V.; Wu, J.; Yee, C.; Spies, T.: Tumour-derived soluble MIC

[88003] ligands impair expression of NKG2D and T-cell activation.

Nature 419:

[88004] 734–738, 2002.

[88005] 15497.

[88006] Li, P.; Morris, D. L.; Willcox, B. E.; Steinle, A.; Spies, T.;

[88007] Strong, R. K.: Complex structure of the activating immunoreceptor

[88008] NKG2D and its MHC class I-like ligand MICA. Nature Immun. 2: 443–451,

[88009] 2001.

[88010] 15498.

[88011] Nakashima, T.; Sekiguchi, T.; Kuraoka, A.; Fukushima, K.; Shibata,

[88012] Y.; Komiyama, S.; Nishimoto, T.: Molecular cloning of a human cDNA

[88013] encoding a novel protein, DAD1, whose defect causes apoptotic cell

[88014] death in hamster BHK21 cells. Molec. Cell. Biol. 13: 6367–6374,

[88015] 1993.

[88016] 15499.

[88017] Ylug, I. G.; See, C. G.; Fisher, E. M. C.: The DAD1 protein,
whose

[88018] defect causes apoptotic cell death, maps to human chro-
mosome 14. Genomics 26:

[88019] 433–435, 1995. Note: Erratum: Genomics: 28: 129 only,
1995.

[88020] 15500.

[88021] Yulug, I. G.; See, C. G.; Fisher, E. M. C.: The DAD1 protein,

[88022] whose defect causes apoptotic cell death, maps to human
chromosome

[88023] 14. Genomics 26: 433–435, 1995.

[88024] 15501.

[88025] Jepsen, K. J.; Wu, F.; Peragallo, J. H.; Paul, J.; Roberts, L.;

[88026] Ezura, Y.; Oldberg, A.; Birk, D. E.; Chakravarti, S.: A syn-
drome

[88027] of joint laxity and impaired tendon integrity in lumican-
and fibromodulin-deficient

[88028] mice. J. Biol. Chem. 277: 35532–35540, 2002.

[88029] 15502.

[88030] Sztrolovics, R.; Chen, X.-N.; Grover, J.; Roughley, P. J.; Ko-
renberg,

- [88031] J. R.: Localization of the human fibromodulin gene (FMOD) to chromosome
- [88032] 1q32 and completion of the cDNA sequence. *Genomics* 23: 715–717,
- [88033] 1994.
- [88034] 15503.
- [88035] Dalton, S.; Treisman, R.: Characterization of SAP–1, a protein
- [88036] recruited by serum response factor to the c–fos serum response element. *Cell* 68:
- [88037] 597–612, 1992.
- [88038] 15504.
- [88039] Mo, Y.; Vaessen, B.; Johnston, K.; Marmorstein, R.: Structures
- [88040] of SAP–1 bound to DNA targets from the E74 and c–fos promoters: insights
- [88041] into DNA sequence discrimination by Ets proteins. *Molec. Cell* 2:
- [88042] 201–212, 1998.
- [88043] 15505.
- [88044] Shipley, J.; Sheer, D.; Dalton, S.; Treisman, R.; Patel, K.:

Mapping

- [88045] of the human SAP1 (SRF accessory protein 1) gene and SAP2, a gene
- [88046] encoding a related protein, to chromosomal bands 1q32 and 12q23, respectively. Genomics 23:
- [88047] 710–711, 1994.
- [88048] 15506.
- [88049] Ikeuchi, T.; Asaka, T.; Saito, M.; Tanaka, H.; Higuchi, S.; Tanaka,
- [88050] K.; Saida, K.; Uyama, E.; Mizusawa, H.; Fukuhara, N.; Nonaka, I.;
- [88051] Takamori, M.; Tsuji, S.: Gene locus for autosomal recessive distal
- [88052] myopathy with rimmed vacuoles maps to chromosome 9. Ann. Neurol. 41:
- [88053] 432–437, 1997.
- [88054] 15507.
- [88055] Erickson, J. D.; Varoqui, H.; Schafer, M. K.–H.; Modi, W.; Diebler,
- [88056] M.–F.; Weihe, E.; Rand, J.; Eiden, L. E.; Bonner, T. I.; Usdin, T.
- [88057] B.: Functional identification of a vesicular acetylcholine

transporter

[88058] and its expression from a 'cholinergic' gene locus. J. Biol. Chem. 269:

[88059] 21929–21932, 1994.

[88060] 15508.

[88061] Probst, M. R.; Beer, M.; Beer, D.; Jenö, P.; Meyer, U. A.; Gasser,

[88062] R.: Human liver arylacetamide deacetylase: molecular cloning of a

[88063] novel esterase involved in the metabolic activation of arylamine carcinogens

[88064] with high sequence similarity to hormone-sensitive lipase. J. Biol.

[88065] Chem. 269: 21650–21656, 1994.

[88066] 15509.

[88067] Yamazaki, K.; Kusano, K.; Tadano, K.; Tanaka, I.: Radiation hybrid

[88068] mapping of human arylacetamide deacetylase (AADAC) locus to chromosome 15510.

[88069] Worman, H. J.; Yuan, J.; Blobel, G.; Georgatos, S. D.: A lamin

[88070] B receptor in the nuclear envelope. Proc. Nat. Acad. Sci.

85: 8531–8534,

[88071] 1988.

[88072] 15511.

[88073] Tanahashi, H.; Tabira, T.: Alzheimer's disease-associated
presenilin

[88074] 2 interacts with DRAL, an LIM-domain protein. Hum.
Molec. Genet. 9:

[88075] 2281–2289, 2000.

[88076] 15512.

[88077] Erkinheimo, T.-L.; Saukkonen, K.; Narko, K.; Jalkanen, J.;
Ylikorkala,

[88078] O.; Ristimäki, A.: Expression of cyclooxygenase-2 and
prostanoid

[88079] receptors by human myometrium. J. Clin. Endocr. Metab.
85: 3468–3475,

[88080] 2000.

[88081] 15513.

[88082] Hla, T.; Neilson, K.: Human cyclooxygenase-2 cDNA. Proc.
Nat.

[88083] Acad. Sci. 89: 7384–7388, 1992.

[88084] 15514.

- [88085] Jones, D. A.; Carlton, D. P.; McIntyre, T. M.; Zimmerman, G. A.;
- [88086] Prescott, S. M.: Molecular cloning of human prostaglandin endoperoxide
- [88087] synthase type II and demonstration of expression in response to cytokines. J.
- [88088] Biol. Chem. 268: 9049–9054, 1993.
- [88089] 15515.
- [88090] Kraemer, S. A.; Meade, E. A.; DeWitt, D. L.: Prostaglandin endoperoxide
- [88091] synthase gene structure: identification of the transcriptional start
- [88092] site and 5-prime-flanking regulatory sequences. Arch. Biochem. Biophys. 293:
- [88093] 391–400, 1992.
- [88094] 15516.
- [88095] Lassus, P.; Wolff, H.; Andersson, S.: Cyclooxygenase-2 in human
- [88096] perinatal lung. Pediat. Res. 47: 602–605, 2000.
- [88097] 15517.
- [88098] Lim, H.; Paria, B. C.; Das, S. K.; Dinchuk, J. E.; Langenbach,

- [88099] R.; Trzaskos, J. M.; Dey, S. K.: Multiple female reproductive failures
- [88100] in cyclooxygenase 2-deficient mice. Cell 91: 197–208, 1997.
- [88101] 15518.
- [88102] Morrow, J. A.; Collie, I. T.; Dunbar, D. R.; Walker, G. B.; Shahid,
- [88103] M.; Hill, D. R.: Molecular cloning and functional expression of the
- [88104] human glycine transporter GlyT2 and chromosomal localisation of the
- [88105] gene in the human genome. FEBS Lett. 439: 334–340, 1998.
- [88106] 15519.
- [88107] Iacobazzi, V.; Palmieri, F.; Runswick, M. J.; Walker, J. E.: Sequences
- [88108] of the human and bovine genes for the mitochondrial 2-oxoglutarate
- [88109] carrier. DNA Seq. 3: 79–88, 1992.
- [88110] 15520.
- [88111] Piccininni, S.; Iacobazzi, V.; Lauria, G.; Rocchi, M.; Palmieri,

- [88112] F.: Assignment of the oxoglutarate carrier gene (SLC20A4) to human
- [88113] chromosome 17p13.3. Cytogenet. Cell Genet. 83: 256–257, 1998.
- [88114] 15521.
- [88115] Harvey, T. J.; Hooper, J. D.; Myers, S. A.; Stephenson, S.-A.;
- [88116] Ashworth, L. K.; Clements, J. A.: Tissue-specific expression patterns
- [88117] and fine mapping of the human kallikrein (KLK) locus on proximal 19q13.4. J.
- [88118] Biol. Chem. 275: 37397–37406, 2000.
- [88119] 15522.
- [88120] Chaudhry, F. A.; Reimer, R. J.; Krizaj, D.; Barber, D.; Storm-Mathisen,
- [88121] J.; Copenhagen, D. R.; Edwards, R. H.: Molecular analysis of system
- [88122] N suggests novel physiological roles in nitrogen metabolism and synaptic
- [88123] transmission. Cell 99: 769–780, 1999.
- [88124] 15523.

[88125] Yamada, K.; Nishida, K.; Hibi, M.; Hirano, T.; Matsuda, Y.:
Comparative

[88126] FISH mapping of Gab1 and Gab2 genes in human, mouse
and rat. Cytogenet.

[88127] Cell Genet. 94: 39–42, 2001.

[88128] 15524.

[88129] Afonina, E.; Stauber, R.; Pavlakis, G. N.: The human
poly(A)–binding

[88130] protein 1 shuttles between the nucleus and the cytoplasm.
J. Biol.

[88131] Chem. 273: 13015–13021, 1998.

[88132] 15525.

[88133] Deo, R. C.; Bonanno, J. B.; Sonenberg, N.; Burley, S. K.:
Recognition

[88134] of polyadenylate RNA by the poly(A)–binding protein. Cell
98: 835–845,

[88135] 1999.

[88136] 15526.

[88137] Gorlach, M.; Burd, C. G.; Dreyfuss, G.: The mRNA
poly(A)–binding

[88138] protein: localization, abundance, and RNA–binding speci–

ficity. Exp.

[88139] Cell Res. 211: 400–407, 1994.

[88140] 15527.

[88141] Grange, T.; Martins de Sa, C.; Oddos, J.; Pictet, R.: Human mRNA

[88142] polyadenylate binding protein: evolutionary conservation of a nucleic

[88143] acid binding motif. Nucleic Acids Res. 15: 4771–4787, 1987.

[88144] 15528.

[88145] Feral, C.; Guellaen, G.; Pawlak, A.: Human testis expresses a

[88146] specific poly(A)–binding protein. Nucleic Acids Res. 29: 1872–1883,

[88147] 2001.

[88148] 15529.

[88149] Danpure, C. J.; Purdue, P. E.; Fryer, P.; Griffiths, S.; Allsop,

[88150] J.; Lumb, M. J.; Guttridge, K. M.; Jennings, P. R.; Scheinman, J.

[88151] I.; Mauer, S. M.; Davidson, N. O.: Enzymological and mutational analysis

- [88152] of a complex primary hyperoxaluria type I phenotype involving alanine:glyoxylate
- [88153] aminotransferase peroxisome-to-mitochondrion mistargeting and intraperoxisomal
- [88154] aggregation. *Am. J. Hum. Genet.* 53: 417-432, 1993.
- [88155] 15530.
- [88156] Mori, M.; Oda, T.; Nishiyama, K.; Serikawa, T.; Yamada, J.; Ichiyama,
- [88157] A.: A single serine:pyruvate aminotransferase gene on rat chromosome
- [88158] 9q34-q36. *Genomics* 13: 686-689, 1992.
- [88159] 15531.
- [88160] Purdue, P. E.; Allsop, J.; Isaya, G.; Rosenberg, L. E.; Danpure,
- [88161] C. J.: Mistargeting of peroxisomal L-alanine:glyoxylate aminotransferase
- [88162] to mitochondria in primary hyperoxaluria patients depends upon activation
- [88163] of a cryptic mitochondrial targeting sequence by a point mutation. *Proc.*
- [88164] *Nat. Acad. Sci.* 88: 10900-10904, 1991.

- [88165] 15532.
- [88166] Purdue, P. E.; Lumb, M. J.; Allsop, J.; Minatogawa, Y.; Danpure,
- [88167] C. J.: A glycine-to-glutamate substitution abolishes alanine:glyoxylate
- [88168] aminotransferase catalytic activity in a subset of patients with primary
- [88169] hyperoxaluria type 1. *Genomics* 13: 215-218, 1992.
- [88170] 15533.
- [88171] Purdue, P. E.; Takada, Y.; Danpure, C. J.: Identification of
- [88172] mutations associated with peroxisome-to-mitochondrion mistargeting
- [88173] of alanine/glyoxylate aminotransferase in primary hyperoxaluria type
- [88174] 1. *J. Cell Biol.* 111: 2341-2351, 1990.
- [88175] 15534.
- [88176] Takada, Y.; Kaneko, N.; Esumi, H.; Purdue, P. E.; Danpure, C.
- [88177] J.: Human peroxisomal L-alanine:glyoxylate aminotransferase: evolutionary
- [88178] loss of a mitochondrial targeting signal by point mutation

of the

[88179] initiation codon. Biochem. J. 268: 517–520, 1990.

[88180] 15535.

[88181] von Schnakenburg, C.; Rumsby, G.: Primary hyperoxaluria
type

[88182] 1: a cluster of new mutations in exon 7 of the AGXT gene.
J. Med.

[88183] Genet. 34: 489–492, 1997.

[88184] 15536.

[88185] Kobayashi, H.; Hino, M.; Shimodahira, M.; Iwakura, T.;
Ishihara,

[88186] T.; Ikekubo, K.; Ogawa, Y.; Nakao, K.; Kurahachi, H.: Mis-
sense mutation

[88187] of TRPS1 in a family of tricho–rhino–phalangeal syndrome
type III. Am.

[88188] J. Med. Genet. 107: 26–29, 2002.

[88189] 15537.

[88190] Yoshida, A.; Minowa, M. T.; Takamatsu, S.; Hara, T.; Ike-
naga, H.;

[88191] Takeuchi, M.: A novel second isoenzyme of the human
UDP–N–acetylglucosamine:alpha–1,3–D–mannoside

- [88192] beta-1,4-N-acetylglucosaminyltransferase family: cDNA cloning, expression,
[88193] and chromosomal assignment. Glycoconj. J. 15:
1115-1123, 1998.
[88194] 15538.
- [88195] Jakobsson, P.-J.; Mancini, J. A.; Riendeau, D.; Ford-
Hutchinson,
[88196] A. W.: Identification and characterization of a novel micro-
somal
[88197] enzyme with glutathione-dependent transferase and per-
oxidase activities. J.
[88198] Biol. Chem. 272: 22934-22939, 1997.
[88199] 15539.
- [88200] Poliak, S.; Gollan, L.; Martinez, R.; Custer, A.; Einheber, S.;
[88201] Salzer, J. L.; Trimmer, J. S.; Shrager, P.; Peles, E.: Caspr2, a
[88202] new member of the neurexin superfamily, is localized at
the juxtaparanodes
[88203] of myelinated axons and associates with K⁺ channels.
Neuron 24:
[88204] 1037-1047, 1999.
[88205] 15540.

- [88206] Hata, M.; Ohtsuka, K.: Characterization of HSE sequences in human
- [88207] Hsp40 gene: structural and promoter analysis. *Biochim. Biophys. Acta* 1397:
- [88208] 43–55, 1998.
- [88209] 15541.
- [88210] Hata, M.; Okumura, K.; Seto, M.; Ohtsuka, K.: Genomic cloning
- [88211] of a human heat shock protein 40 (Hsp40) gene (HSPF1) and its chromosomal
- [88212] localization to 19p13.2. *Genomics* 38: 446–449, 1996.
- [88213] 15542.
- [88214] Ohtsuka, K.: Cloning of a cDNA for heat-shock protein hsp40, a
- [88215] human homologue of bacterial DnaJ. *Biochem. Biophys. Res. Commun.* 197:
- [88216] 235–240, 1993.
- [88217] 15543.
- [88218] Sultz, B. M.: Genetic control of leucocyte responses to endotoxin. *Nature* 219:
- [88219] 1253–1254, 1968.

- [88220] 15544.
- [88221] Donger, C.; Krejci, E.; Serradell, A. P.; Eymard, B.; Bon, S.;
- [88222] Nicole, S.; Chateau, D.; Gary, F.; Fardeau, M.; Massoulie, J.;
Guicheney,
- [88223] P.: Mutation in the human acetylcholinesterase-associated
collagen
- [88224] gene, COLQ, is responsible for congenital myasthenic
syndrome with
- [88225] end-plate acetylcholinesterase deficiency (type Ic). *Am. J.*
Hum.
- [88226] *Genet.* 63: 967–975, 1998.
- [88227] 15545.
- [88228] Krejci, E.; Coussen, F.; Duval, N.; Chatel, J.-M.; Legay, C.;
- Puype,
- [88229] M.; Vandekerckhove, J.; Cartaud, J.; Bon, S.; Massoulie, J.:
- Primary
- [88230] structure of a collagenic tail peptide of torpedo acetyl-
cholinesterase:
- [88231] co-expression with catalytic subunit induces the produc-
tion of collagen-tailed
- [88232] forms in transfected cells. *EMBO J.* 10: 1285–1293, 1991.

- [88233] 15546.
- [88234] Krejci, E.; Thomine, S.; Boschetti, N.; Legay, C.; Sketelj, J.;
- [88235] Massoulie, J.: The mammalian gene of acetyl-
cholinesterase-associated
- [88236] collagen. J. Biol. Chem. 272: 22840-22847, 1997.
- [88237] 15547.
- [88238] Ohno, K.; Brengman, J. M.; Felice, K. J.; Cornblath, D. R.;
Engel,
- [88239] A. G.: Congenital end-plate acetylcholinesterase defi-
ciency caused
- [88240] by a nonsense mutation and an A-to-G splice-donor-site
mutation at
- [88241] position +3 of the collagenlike-tail-subunit gene (COLQ):
how does
- [88242] G at position +3 result in aberrant splicing? Am. J. Hum.
Genet. 65:
- [88243] 635-644, 1999.
- [88244] 15548.
- [88245] Muravenko, O. V.; Gizatullin, R. Z.; Protopopov, A. I.;
Kashuba,
- [88246] V. I.; Zabarovsky, E. R.; Zelenin, A. V.: Assignment of

CDK5R2 coding

- [88247] for the cyclin-dependent kinase 5, regulatory subunit 2 (NCK5A1 protein)
- [88248] to human chromosome band 2q35 by fluorescent in situ hybridization. Cytogenet.
- [88249] Cell Genet. 89: 160-161, 2000.
- [88250] 15549.
- [88251] Nilden, F.; Backstrom, A.; Bark, C.: Molecular cloning and characterisation
- [88252] of a mouse gene encoding an isoform of the neuronal cyclin-dependent
- [88253] kinase 5 (CDK5) activator. Biochim. Biophys. Acta 1398: 371-376,
- [88254] 1998.
- [88255] 15550.
- [88256] Tang, D.; Yeung, J.; Lee, K-Y.; Matsushita, M.; Matsui, H.; Tomizawa,
- [88257] K.; Hatase, O.; Wang, J. H.: An isoform of the neuronal cyclin-dependent
- [88258] kinase 5 (Cdk5) activator. J. Biol. Chem. 270: 26897-26903, 1995.
- [88259] 15551.

- [88260] Fujioka, M.; Takahashi, N.; Odai, H.; Araki, S.; Ichikawa, K.;
- [88261] Feng, J.; Nakamura, M.; Kaibuchi, K.; Hartshorne, D. J.;
Nakano, T.;
- [88262] Ito, M.: A new isoform of human myosin phosphatase tar-
geting/regulatory
- [88263] subunit (MYPT2): cDNA cloning, tissue expression, and
chromosomal
- [88264] mapping. *Genomics* 49: 59–68, 1998.
- [88265] 15552.
- [88266] Hallas, C.; Pekarsky, Y.; Itoyama, T.; Varnum, J.; Bichi, R.;
Rothstein,
- [88267] J. L.; Croce, C. M.: Genomic analysis of human and mouse
TCL1 loci
- [88268] reveals a complex of tightly clustered genes. *Proc. Nat.*
Acad. Sci. 96:
- [88269] 14418–14423, 1999.
- [88270] 15553.
- [88271] Pekarsky, Y.; Hallas, C.; Isobe, M.; Russo, G.; Croce, C. M.:
- [88272] Abnormalities at 14q32.1 in T cell malignancies involve
two oncogenes. *Proc.*
- [88273] *Nat. Acad. Sci.* 96: 2949–2951, 1999.

- [88274] 15554.
- [88275] Koga, Y.; Fabrizi, G. M.; Mita, S.; Arnaudo, E.; Lomax, M. I.;
- [88276] Aqua, M. S.; Grossman, L. I.; Schon, E. A.: Sequence of a
cDNA specifying
- [88277] subunit VIIc of human cytochrome c oxidase. *Nucleic
Acids Res.* 18:
- [88278] 684 only, 1990.
- [88279] 15555.
- [88280] Harju, K.; Glumoff, V.; Hallman, M.: Ontogeny of Toll-like
receptors
- [88281] Tlr2 and Tlr4 in mice. *Pediat. Res.* 49: 81–83, 2001.
- [88282] 15556.
- [88283] Qu, L. H.; Nicoloso, M.; Michot, B.; Azum, M. C.; Caizer-
gues-Ferrer,
- [88284] M.; Renalier, M. H.; Bachellerie, J. P.: U21, a novel small
nucleolar
- [88285] RNA with a 13 nt. complementarity to 28S rRNA, is en-
coded in an intron
- [88286] of ribosomal protein L5 gene in chicken and mammals.
Nucleic Acids
- [88287] *Res.* 22: 4073–4081, 1994.

[88288] 15557.

[88289] Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.;
Grandy, D.;

[88290] Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an
11q22.3-9p22

[88291] translocation with affective disorder: proximity of the
dopamine D2

[88292] receptor gene relative to the translocation breakpoint.
(Abstract) Am.

[88293] J. Hum. Genet. 45 (suppl.): A220 only, 1989.

[88294] 15558.

[88295] Abuladze, N.; Lee, I.; Newman, D.; Hwang, J.; Boorer, K.;
Pushkin,

[88296] A.; Kurtz, I.: Molecular cloning, chromosomal localization,
tissue

[88297] distribution, and functional expression of the human pan-
creatic sodium

[88298] bicarbonate cotransporter. J. Biol. Chem. 273:
17689-17695, 1998.

[88299] 15559.

[88300] Burnham, C. E.; Amlal, H.; Wang, Z.; Shull, G. E.; Soleimani,
M.

- [88301] : Cloning and functional expression of a human kidney Na⁺:HCO₃⁻ cotransporter. J.
- [88302] Biol. Chem. 272: 19111–19114, 1997.
- [88303] 15560.
- [88304] Choi, I.; Romero, M. F.; Khandoudi, N.; Bril, A.; Boron, W. F.
- [88305] : Cloning and characterization of a human electrogenic Na⁽⁺⁾-HCO₃⁻
- [88306] cotransporter isoform (hhNBC). Am. J. Physiol. 276: C576–C584, 1999.
- [88307] 15561.
- [88308] Igarashi, T.; Inatomi, J.; Sekine, T.; Cha, S. H.; Kanai, Y.; Kunimi,
- [88309] M.; Tsukamoto, K.; Satoh, H.; Shimadzu, M.; Tozawa, F.; Mori, T.;
- [88310] Shiobara, M.; Seki, G.; Endou, H.: Mutations in SLC4A4 cause permanent
- [88311] isolated proximal renal tubular acidosis with ocular abnormalities.
- [88312] (Letter) Nature Genet. 23: 264–265, 1999.
- [88313] 15562.

- [88314] Romero, M. F.; Boron, W. F.: Electrogenic Na(+)/HCO(3-) cotransporters:
[88315] cloning and physiology. *Annu. Rev. Physiol.* 61: 699–723, 1999.
[88316] 15563.
- [88317] Soleimani, M.; Burnham, C. E.: Physiologic and molecular aspects
[88318] of the Na(+):HCO(3-) cotransporter in health and disease processes. *Kidney Int.* 57: 371–384, 2000.
[88319] 15564.
- [88320] Usui, T.; et al.; et al. :*Pflugers Arch.* 438: 458–462, 1999.
[88321] 15565.
- [88322] Hayes, S. E.; Conner, L. J.; Stramm, L. E.; Shi, Y.: Assignment
[88323] of pancreatic eIF-2a kinase (EIF2AK3) to human chromosome band 2p12
[88324] by radiation hybrid mapping and in situ hybridization. *Cytogenet. Cell Genet.* 86: 327–328, 1999.
[88325] 15566.

- [88328] Avraham, K. B.; Cho, B. C.; Gilbert, D.; Fujii, H.; Okamoto, K.;
- [88329] Shimazaki, T.; Ito, T.; Shoji, H.; Wakamatsu, Y.; Kondoh, H.; Takahashi,
- [88330] N.; Muramatsu, M.; Hamada, H.; Copeland, N. G.; Jenkins, N. A.: Murine
- [88331] chromosomal location of four class III POU transcription factors. *Genomics* 18:
- [88332] 131–133, 1993.
- [88333] 15567.
- [88334] Egan, S.; Herbrick, J.-A.; Tsui, L.-C.; Cohen, B.; Flock, G.; Beatty,
- [88335] B.; Scherer, S. W.: Mapping of the human lunatic fringe (LFNG) gene
- [88336] to 7p22 and manic fringe (MFNG) to 22q12. *Genomics* 54: 576–577,
- [88337] 1998.
- [88338] 15568.
- [88339] Johnston, S. H.; Rauskolb, C.; Wilson, R.; Prabhakaran, B.; Irvine,
- [88340] K. D.; Vogt, T. F.: A family of mammalian Fringe genes implicated

[88341] in boundary determination and the Notch pathway. Development 124:

[88342] 2245–2254, 1997.

[88343] 15569.

[88344] Moran, J. L.; Johnston, S. H.; Rauskolb, C.; Bhalerao, J.; Bowcock,

[88345] A. M.; Vogt, T. F.: Genomic structure, mapping, and expression analysis

[88346] of the mammalian lunatic, manic, and radical fringe genes. Mammalian

[88347] Genome 10: 535–541, 1999.

[88348] 15570.

[88349] Van Tine, B. A.; Knops, J.; Shaw, G. M.; May, W. A.: Assignment

[88350] of human MFNG, manic fringe Drosophila homolog, to 22q13.1 using tyramide

[88351] fluorescence in situ hybridization (T-FISH). Cytogenet. Cell Genet. 87:

[88352] 132–133, 1999.

[88353] 15571.

[88354] Laufer, E.; Dahn, R.; Orozco, O. E.; Yeo, C.-Y.; Pisenti, J.;

Henrique,

[88355] D.; Abbott, U. K.; Fallon, J. F.; Tabin, C.: Expression of
Radical

[88356] fringe in limb-bud ectoderm regulates apical ectodermal
ridge formation. Nature 386:

[88357] 366–373, 1997.

[88358] 15572.

[88359] Rodriguez-Esteban, C.; Schwabe, J. W. R.; De La Pena, J.;
Foys,

[88360] B.; Eshelman, B.; Izpisua Belmonte, J. C.: Radical fringe
positions

[88361] the apical ectodermal ridge at the dorsoventral boundary
of the vertebrate

[88362] limb. Nature 386: 360–366, 1997.

[88363] 15573.

[88364] Wiley, S. R.; Cassiano, L.; Lofton, T.; Davis-Smith, T.; Win-
kles,

[88365] J. A.; Lindner, V.; Liu, H.; Daniel, T. O.; Smith, C. A.;
Fanslow,

[88366] W. C.: A novel TNF receptor family member binds TWEAK
and is implicated

[88367] in angiogenesis. Immunity 15: 837–846, 2001.

[88368] 15574.

[88369] Hoey, T.; Sun, Y.-L.; Williamson, K.; Xu, X.: Isolation of two

[88370] new members of the NF-AT gene family and functional
characterization

[88371] of the NF-AT proteins. *Immunity* 2: 461-472, 1995.

[88372] 15575.

[88373] Arany, Z.; Newsome, D.; Oldread, E.; Livingston, D. M.;
Eckner,

[88374] R.: A family of transcriptional adaptor proteins targeted by
the

[88375] E1A oncoprotein. *Nature* 374: 81-84, 1995.

[88376] 15576.

[88377] Dong, C.; Zhu, S.; Wang, T.; Yoon, W.; Li, Z.; Alvarez, R. J.;

[88378] ten Dijke, P.; White, B.; Wigley, F. M.; Goldschmidt-Cler-
mont, P.

[88379] J.: Deficient Smad7 expression: a putative molecular de-
fect in scleroderma. *Proc.*

[88380] *Nat. Acad. Sci.* 99: 3908-3913, 2002.

[88381] 15577.

[88382] Kavsak, P.; Rasmussen, R. K.; Causing, C. G.; Bonni, S.;

Zhu, H.;

[88383] Thomsen, G. H.; Wrana, J. L.: Smad7 binds to Smurf2 to form an E3

[88384] ubiquitin ligase that targets the TGF-beta receptor for degradation. Molec.

[88385] Cell 6: 1365-1375, 2000.

[88386] 15578.

[88387] Lallemand, F.; Mazars, A.; Prunier, C.; Bertrand, F.; Korn-prost,

[88388] M.; Gallea, S.; Roman-Roman, S.; Cherqui, G.; Atfi, A.: Smad7 inhibits

[88389] the survival nuclear factor kappa-B and potentiates apoptosis in epithelial

[88390] cells. Oncogene 20: 879-884, 2001.

[88391] 15579.

[88392] Roijer, E.; Moren, A.; ten Dijke, P.; Stenman, G.: Assignment

[88393] of the Smad7 gene (MADH7) to human chromosome 18q21.1 by fluorescence

[88394] in situ hybridization. Cytogenet. Cell Genet. 81: 189-190, 1998.

[88395] 15580.

- [88396] Cravatt, B. F.; Giang, D. K.; Mayfield, S. P.; Boger, D. L.; Lerner, R. A.; Gilula, N. B.: Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature* 384: 83–87, 1996.
- [88399] 15581.
- [88400] Giang, D. K.; Cravatt, B. F.: Molecular characterization of human and mouse fatty acid amide hydrolases. *Proc. Nat. Acad. Sci.* 94: 2238–2242, 1997.
- [88403] 15582.
- [88404] Sipe, J. C.; Chiang, K.; Gerber, A. L.; Beutler, E.; Cravatt, B. F.: A missense mutation in human fatty acid amide hydrolase associated with problem drug use. *Proc. Nat. Acad. Sci.* 99: 8394–8399, 2002.
- [88407] 15583.
- [88408] Wan, M.; Cravatt, B. F.; Ring, H. Z.; Zhang, X.; Francke, U.: Conserved chromosomal location and genomic structure

of human and

[88410] mouse fatty-acid amide hydrolase genes and evaluation of
clasper as

[88411] a candidate neurological mutation. Genomics 54:
408–414, 1998.

[88412] 15584.

[88413] Baier, M.; Bannert, N.; Werner, A.; Lang, K.; Kurth, R.:
Molecular

[88414] cloning, sequence, expression, and processing of the in-
terleukin 16

[88415] precursor. Proc. Nat. Acad. Sci. 94: 5273–5277, 1997.

[88416] 15585.

[88417] Roh, M. H.; Makarova, O.; Liu, C.-J.; Shin, K.; Lee, S.; Lau-
rinec,

[88418] S.; Goyal, M.; Wiggins, R.; Margolis, B.: The Maguk protein,
Pals1,

[88419] functions as an adapter, linking mammalian homologues
of Crumbs and

[88420] Discs Lost. J. Cell Biol. 157: 161–172, 2002.

[88421] 15586.

[88422] Bannert, N.; Baier, M.; Werner, A.; Kurth, R.: Interleukin-16

- [88423] or not? Nature 381: 30 only, 1996.
- [88424] 15587.
- [88425] Bannert, N.; Kurth, R.; Baier, M.: The gene encoding mouse interleukin-16
- [88426] consists of seven exons and maps to Chromosome 7 D2-D3. Immunogenetics 49:
- [88427] 704-706, 1999.
- [88428] 15588.
- [88429] Cruikshank, W.; Center, D. M.: Modulation of lymphocyte migration
- [88430] by human lymphokines. II. Purification of a lymphotactic factor (LCF). J.
- [88431] Immun. 128: 2569-2574, 1982.
- [88432] 15589.
- [88433] Cruikshank, W. W.; Center, D. M.; Nisar, N.; Wu, M.; Natke, B.;
- [88434] Theodore, A. C.; Kornfeld, H.: Molecular and functional analysis
- [88435] of a lymphocyte chemoattractant factor: association of biologic function
- [88436] with CD4 expression. Proc. Nat. Acad. Sci. 91:

5109–5113, 1994.

[88437] 15590.

[88438] Drwinga, H. L.; Toji, L. H.; Kim, C. H.; Greene, A. E.;
Mulivor,

[88439] R. A.: NIGMS human/rodent somatic cell hybrid mapping
panels 1 and

[88440] 2. Genomics 16: 311–314, 1993.

[88441] 15591.

[88442] Hudson, T. J.; Stein, L. D.; Gerety, S. S.; Ma, J.; Castle, A.

[88443] B.; Silva, J.; Slonim, D. K.; Baptista, R.; Kruglyak, L.; Xu, S.
H.;

[88444] Hu, X.; Colbert, A. M. E.; and 39 others: An STS-based
map of the

[88445] human genome. Science 270: 1945–1954, 1995.

[88446] 15592.

[88447] Keane, J.; Nicoll, J.; Kim, S.; Wu, D. M. H.; Cruikshank, W.
W.;

[88448] Brazer, W.; Natke, B.; Zhang, Y.; Center, D. M.; Kornfeld,
H.: Conservation

[88449] of structure and function between human and murine IL–
16. J. Immun. 160:

[88450] 5945–5954, 1998.

[88451] 15593.

[88452] Tarnasky, H.; Gill, D.; Murthy, S.; Shao, X.; Demetrick, D. J.;

[88453] van der Hoorn, F. A.: A novel testis-specific gene, SPAG4, whose

[88454] product interacts specifically with outer dense fiber protein ODF27,

[88455] maps to human chromosome 20q11.2. *Cytogenet. Cell Genet.* 81: 65–67,

[88456] 1998.

[88457] 15594.

[88458] Hurlin, P. J.; Queva, C.; Eisenman, R. N.: Mnt, a novel Max-interacting

[88459] protein is coexpressed with Myc in proliferating cells and mediates

[88460] repression at Myc binding sites. *Genes Dev.* 11: 44–58, 1997.

[88461] 15595.

[88462] Lo Nigro, C.; Venesio, T.; Reymond, A.; Meroni, G.; Alberici, P.;

- [88463] Cainarca, S.; Enrico, F.; Stack, M.; Ledbetter, D. H.; Liscia, D.
- [88464] S.; Ballabio, A.; Carrozzo, R.: The human ROX gene: genomic structure
- [88465] and mutation analysis in human breast tumors. *Genomics* 49: 275–282,
- [88466] 1998.
- [88467] 15596.
- [88468] Meroni, G.; Reymond, A.; Alcalay, M.; Borsani, G.; Tanigami, A.;
- [88469] Tonlorenzi, R.; Lo Nigro, C.; Messali, S.; Zollo, M.; Ledbetter, D.
- [88470] H.; Brent, R.; Ballabio, A.; Carrozzo, R.: Rox, a novel bHLHZip protein
- [88471] expressed in quiescent cells that heterodimerizes with Max, binds
- [88472] a non-canonical E box and acts as a transcriptional repressor. *EMBO*
- [88473] *J.* 16: 2892–2906, 1997.
- [88474] 15597.
- [88475] Lennon-Dumenil, A.-M.; Barbouche, M.-R.; Vedrenne, J.; Prod'Homme,

- [88476] T.; Bejaoui, M.; Ghariani, S.; Charron, D.; Fellous, M.; Del-
lagi,
- [88477] K.; Alcaide-Loridan, C.: Uncoordinated HLA-D gene ex-
pression in a
- [88478] RFXANK-defective patient with MHC class II deficiency. J.
Immun. 166:
- [88479] 5681-5687, 2001.
- [88480] 15598.
- [88481] Lisowska-Groszpiere, B.; Fondaneche, M. C.; Rols, M. P.;
Griscelli,
- [88482] C.; Fischer, A.: Two complementation groups account for
most cases
- [88483] of inherited MHC class II deficiency. Hum. Molec. Genet. 3:
953-958,
- [88484] 1994.
- [88485] 15599.
- [88486] Gerloff, T.; Stieger, B.; Hagenbuch, B.; Madon, J.; Land-
mann, L.;
- [88487] Roth, J.; Hofmann, A. F.; Meier, P. J.: The sister of P-
glycoprotein
- [88488] represents the canalicular bile salt export pump of mam-
malian liver. J.

- [88489] Biol. Chem. 273: 10046–10050, 1998.
- [88490] 15600.
- [88491] Wang, R.; Salem, M.; Yousef, I. M.; Tuchweber, B.; Lam, P.; Childs,
- [88492] S. J.; Helgason, C. D.; Ackerley, C.; Phillips, M. J.; Ling, V.:
- [88493] Targeted inactivation of sister of P-glycoprotein gene (spgp) in mice
- [88494] results in nonprogressive but persistent intrahepatic cholestasis. Proc.
- [88495] Nat. Acad. Sci. 98: 2011–2016, 2001.
- [88496] 15601.
- [88497] Wiszniewski, W.; Fondaneche, M.-C.; Lambert, N.; Master-nak, K.;
- [88498] Picard, C.; Notarangelo, L.; Schwartz, K.; Bal, J.; Reith, W.; Alcaide,
- [88499] C.; de Saint Basile, G.; Fischer, A.; Lisowska-Grospierre, B.: Founder
- [88500] effect for a 26-bp deletion in the RFXANK gene in North African major
- [88501] histocompatibility complex class II-deficient patients belonging to

- [88502] complementation group B. Immunogenetics 51: 261–267, 2000.
- [88503] 15602.
- [88504] Stec, I.; Wright, T. J.; van Ommen, G.–J. B.; de Boer, P. A. J.;
- [88505] van Haeringen, A.; Moorman, A. F. M.; Altherr, M. R.; den Dunnen,
- [88506] J. T.: WHSC1, a 90 kb SET domain–containing gene, expressed in early
- [88507] development and homologous to a Drosophila dysmorphia gene maps in
- [88508] the Wolf–Hirschhorn syndrome critical region and is fused to IgH in
- [88509] t(4;14) multiple myeloma. Hum. Molec. Genet. 7: 1071–1082, 1998.
- [88510] 15603.
- [88511] Buanne, P.; Incerti, B.; Guardavaccaro, D.; Avvantaggiato, V.;
- [88512] Simeone, A.; Tirone, F.: Cloning of the human interferon–related
- [88513] developmental regulator (IFRD1) gene coding for the PC4 protein, a
- [88514] member of a novel family of developmentally regulated

genes. Genomics 51:

[88515] 233–242, 1998.

[88516] 15604.

[88517] Chen, D.; Lathrop, W.; Dong, Y.: Molecular cloning of mouse Lrp7(Lr3)

[88518] cDNA and chromosomal mapping of orthologous genes in mouse and human. Genomics 55:

[88519] 314–321, 1999.

[88520] 15605.

[88521] Dong, Y.; Lathrop, W.; Weaver, D.; Qiu, Q.; Cini, J.; Bertolini,

[88522] D.; Chen, D.: Molecular cloning and characterization of LR3, a novel

[88523] LDL receptor family protein with mitogenic activity. Biochem. Biophys.

[88524] Res. Commun. 251: 784–790, 1998.

[88525] 15606.

[88526] Mao, J.; Wang, J.; Liu, B.; Pan, W.; Farr, G. H., III; Flynn, C.;

[88527] Yuan, H.; Takada, S.; Kimelman, D.; Li, L.; Wu, D.: Low-density lipoprotein

[88528] receptor-related protein-5 binds to axin and regulates the

canonical

[88529] Wnt signaling pathway. *Molec. Cell* 7: 801–809, 2001.

[88530] 15607.

[88531] Mao, B.; Wu, W.; Li, Y.; Hoppe, D.; Stannek, P.; Glinka, A.; Niehrs,

[88532] C.: LDL-receptor-related protein 6 is a receptor for Dickkopf proteins. *Nature* 411:

[88533] 321–325, 2001.

[88534] 15608.

[88535] Vinkemeier, U.; Obermann, W.; Weber, K.; Furst, D. O.: The globular

[88536] head domain of titin extends into the center of the sarcomeric M band:

[88537] cDNA cloning, epitope mapping and immunoelectron microscopy of two

[88538] titin-associated proteins. *J. Cell Sci.* 106: 319–330, 1993.

[88539] 15609.

[88540] van der Ven, P. F. M.; Speel, E. J. M.; Albrechts, J. C. M.; Ramaekers,

[88541] F. C. S.; Hopman, A. H. N.; Furst, D. O.: Assignment of the human

- [88542] gene for endosarcomeric cytoskeletal M-protein (MYOM2) to 8p23.3. Genomics 55: 253–255, 1999.
- [88543] 15610.
- [88544] 15610.
- [88545] Murakami, M.; Mori, M.: Identification of immunogenic regions in human thyrotropin receptor for immunoglobulin G of patients with Graves' disease. Biochem. Biophys. Res. Commun. 171: 512–518, 1990.
- [88546] 15611.
- [88547] Nagayama, Y.; Kaufman, K. D.; Seto, P.; Rapoport, B.: Molecular cloning, sequence and functional expression of the cDNA for the human thyrotropin receptor. Biochem. Biophys. Res. Commun. 165: 1184–1190, 1989.
- [88548] 15612.
- [88549] Hirohata, S.; Seldin, M. F.; Apte, S. S.: Chromosomal assignment

- [88555] of two ADAM genes, TACE (ADAM17) and MLTNB (ADAM19), to human chromosomes
- [88556] 2 and 5, respectively, and of Mltnb to mouse chromosome 11. Genomics 54:
- [88557] 178–179, 1998.
- [88558] 15613.
- [88559] Inoue, D.; Reid, M.; Lum, L.; Kratzschmar, J.; Weskamp, G.; Myung,
- [88560] Y. M.; Baron, R.; Blobel, C. P.: Cloning and initial characterization
- [88561] of mouse meltrin beta and analysis of the expression of four metalloprotease–disintegrins
- [88562] in bone cells. J. Biol. Chem. 273: 4180–4187, 1998.
- [88563] 15614.
- [88564] Petruzzella, V.; Tiranti, V.; Fernandez, P.; Ianna, P.; Carrozzo,
- [88565] R.; Zeviani, M.: Identification and characterization of human cDNAs
- [88566] specific to BCS1, PET112, SCO1, COX15, and COX11, five genes involved
- [88567] in the formation and function of the mitochondrial respiratory chain. Genomics 54:

[88568] 494–504, 1998.

[88569] 15615.

[88570] Sakuma, H.; Ikeda, A.; Oka, S.; Kozutsumi, Y.; Zanetta, J.-P.;

[88571] Kawasaki, T.: Molecular cloning and functional expression of a cDNA

[88572] encoding a new member of mixed lineage protein kinase from human brain. J.

[88573] Biol. Chem. 272: 28622–28629, 1997.

[88574] 15616.

[88575] Iyer, N. V.; Leung, S. W.; Semenza, G. L.: The human hypoxia-inducible

[88576] factor 1- α gene: HIF1A structure and evolutionary conservation. Genomics 52:

[88577] 159–165, 1998.

[88578] 15617.

[88579] Kline, D. D.; Peng, Y.-J.; Manalo, D. J.; Semenza, G. L.; Prabhakar,

[88580] N. R.: Defective carotid body function and impaired ventilatory responses

[88581] to chronic hypoxia in mice partially deficient for hypoxia-

inducible

[88582] factor 1- α . Proc. Nat. Acad. Sci. 99: 821-826, 2002.

[88583] 15618.

[88584] Lando, D.; Peet, D. J.; Whelan, D. A.; Gorman, J. J.;
Whitelaw,

[88585] M. L.: Asparagine hydroxylation of the HIF transactivation
domain:

[88586] a hypoxic switch. Science 295: 858-861, 2002.

[88587] 15619.

[88588] Marti, H. H.; Katschinski, D. M.; Wagner, K. F.; Schaffer, L.;

[88589] Stier, B.; Wenger, R. H.: Isoform-specific expression of hy-
poxia-inducible

[88590] factor-1- α during the late stages of mouse spermi-
genesis. Molec.

[88591] Endocr. 16: 234-243, 2002.

[88592] 15620.

[88593] Maxwell, P. H.; Wiesener, M. S.; Chang, G.-W.; Clifford, S.
C.;

[88594] Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.;
Maher, E.

[88595] R.; Ratcliffe, P. J.: The tumour suppressor protein VHL tar-

gets hypoxia-inducible

[88596] factors for oxygen-dependent proteolysis. *Nature* 399: 271–275, 1999.

[88597] 15621.

[88598] Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W. G., Jr.;

[88599] Pavletich, N. P.: Structure of an HIF-1- α -pVHL complex: hydroxyproline

[88600] recognition in signaling. *Science* 296: 1886–1889, 2002.

[88601] 15622.

[88602] Semenza, G. L.: HIF-1 and human disease: one highly involved

[88603] factor. *Genes Dev.* 14: 1983–1991, 2000.

[88604] 15623.

[88605] Semenza, G. L.; Rue, E. A.; Iyer, N. V.; Pang, M. G.; Kearns,

[88606] W. G.: Assignment of the hypoxia-inducible factor 1- α gene to

[88607] a region of conserved synteny on mouse chromosome 12 and human chromosome

[88608] 14q. *Genomics* 34: 437–439, 1996.

[88609] 15624.

- [88610] Sutter, C. H.; Laughner, E.; Semenza, G. L.: Hypoxia-inducible
- [88611] factor 1- α protein expression is controlled by oxygen-regulated
- [88612] ubiquitination that is disrupted by deletions and missense mutations. Proc.
- [88613] Nat. Acad. Sci. 97: 4748-4753, 2000.
- [88614] 15625.
- [88615] Wenger, R. H.; Rolfs, A.; Kvietikova, I.; Spielmann, P.; Zimmermann,
- [88616] D. R.; Gassmann, M.: The mouse gene for hypoxia-inducible factor-1- α --genomic
- [88617] organization, expression and characterization of an alternative first
- [88618] exon and 5-prime flanking sequence. Europ. J. Biochem. 246: 155-165,
- [88619] 1997.
- [88620] 15626.
- [88621] Wenger, R. H.; Rolfs, A.; Marti, H. H.; Guenet, J.-L.; Gassmann,
- [88622] M.: Nucleotide sequence, chromosomal assignment and mRNA expression

- [88623] of mouse hypoxia-inducible factor-1-alpha. Biochem. Biophys. Res.
- [88624] Commun. 223: 54-59, 1996.
- [88625] 15627.
- [88626] Wenger, R. H.; Rolfs, A.; Spielmann, P.; Zimmermann, D. R.; Gassmann,
- [88627] M.: Mouse hypoxia-inducible factor-1-alpha is encoded by two different
- [88628] mRNA isoforms: expression from a tissue-specific and a housekeeping-type
- [88629] promoter. Blood 91: 3471-3480, 1998.
- [88630] 15628.
- [88631] Hovanessian, A. G.; Laurent, A. G.; Chebath, J.; Galabru, J.; Robert,
- [88632] N.; Svab, J.: Identification of 69-kd and 100-kd forms of 2-5A synthetase
- [88633] in interferon-treated human cells by specific monoclonal antibodies. EMBO
- [88634] J. 6: 1273-1280, 1987.
- [88635] 15629.
- [88636] Marie, I.; Galabru, J.; Svab, J.; Hovanessian, A. G.: Prepara-

tion

[88637] and characterization of polyclonal antibodies specific for the 69

[88638] and 100 k-dalton forms of human 2-5A synthetase.

Biochem. Biophys.

[88639] Res. Commun. 160: 580-587, 1989.

[88640] 15630.

[88641] Marie, I.; Hovanessian, A. G.: The 69-kDa 2-5A synthetase is composed

[88642] of two homologous and adjacent functional domains. J. Biol. Chem. 267:

[88643] 9933-9939, 1992.

[88644] 15631.

[88645] Kools, P.; Van Imschoot, G.; van Roy, F.: Characterization of

[88646] three novel human cadherin genes (CDH7, CDH19, and CDH20) clustered

[88647] on chromosome 18q22-q23 and with high homology to chicken cadherin-7. Genomics 68:

[88648] 283-295, 2000.

[88649] 15632.

- [88650] Amado, M.; Almeida, R.; Carneiro, F.; Lavery, S. B.; Holmes, E.
- [88651] H.; Nomoto, M.; Hollingsworth, M. A.; Hassan, H.; Schwientek, T.;
- [88652] Nielsen, P. A.; Bennett, E. P.; Clausen, H.: A family of human beta-3-galactosyltransferases:
- [88653] characterization of four members of a UDP-galactose:beta-N-acetyl-glucosamine/beta-N-acetyl-galactosamine
- [88654] beta-1,3-galactosyltransferase family. J. Biol. Chem. 273: 12770-12778,
- [88655] 1998.
- [88656] 15633.
- [88657] Hennet, T.; Dinter, A.; Kuhnert, P.; Mattu, T. S.; Rudd, P. M.;
- [88658] Berger, E. G.: Genomic cloning and expression of three murine UDP-galactose:beta-N-acetylglucosamine
- [88659] beta-1,3-galactosyltransferase genes. J. Biol. Chem. 273: 58-65,
- [88660] 1998.
- [88661] 15634.
- [88662] Kolbinger, F.; Streiff, M. B.; Katopodis, A. G.: Cloning of a

[88663] human UDP-galactose:2-acetamido-2-deoxy-D-glucose
3-beta-galactosyltransferase

[88664] catalyzing the formation of type 1 chains. J. Biol. Chem.
273: 433-440,

[88665] 1998.

[88666] 15635.

[88667] Hsu, D. R.; Economides, A. N.; Wang, X.; Eimon, P. M.;
Harland,

[88668] R. M.: The Xenopus dorsalizing factor gremlin identifies a
novel

[88669] family of secreted proteins that antagonize BMP activities.

Molec.

[88670] Cell 1: 673–683, 1998.

[88671] 15636.

[88672] Topol, L. Z.; Modi, W. S.; Koochekpour, S.; Blair, D. G.:

DRM–Gremlin

[88673] (CKTSF1B1) maps to human chromosome 15 and is highly expressed in

[88674] adult and fetal brain. Cytogenet. Cell Genet. 89: 79–84, 2000.

[88675] 15637.

[88676] Beaconsfield, P.; Rainsbury, R.; Kalton, G.: Glucose–6-phosphate

[88677] dehydrogenase deficiency and the incidence of cancer. Oncologia 19:

[88678] 11–19, 1965.

[88679] 15638.

[88680] Ben–Bassat, J.; Ben–Ishay, D.: Hereditary hemolytic anemia associated

[88681] with glucose–6–phosphate dehydrogenase deficiency (Mediterranean type). Israel

[88682] J. Med. Sci. 5: 1053–1059, 1969.

[88683] 15639.

[88684] Benabadji, M.; Merad, F.; Benmoussa, M.; Trabuchet, G.;
Junien,

[88685] C.; Dreyfus, J. C.; Kaplan, J. C.: Heterogeneity of glucose–
6–phosphate

[88686] dehydrogenase deficiency in Algeria. Hum. Genet. 40:
177–184, 1978.

[88687] 15640.

[88688] Benohr, H. C.; Klumpp, F.; Waller, H. D.: Glucose–
6–phosphat–Dehydrogenase

[88689] Typ Schwaben. Dtsch. Med. Wschr. 96: 1029–1032, 1971.

[88690] 15641.

[88691] Benohr, H. C.; Waller, H. D.: Eigenschaften der Glucose–
6–p–dehydrogenase,

[88692] Typ Tubingen. Klin. Wschr. 48: 71–74, 1970.

[88693] 15642.

[88694] Benohr, H. C.; Waller, H. D.; Arnold, H.; Blume, K. G.; Lohr,

[88695] G. W.: Glucose–6–P–Dehydrogenase Typ Bodensee (eine
neue Enzymvariante). Klin.

[88696] Wschr. 49: 1058–1062, 1971.

[88697] 15643.

[88698] Beutler, E.: Glucose 6–phosphate dehydrogenase deficiency, a

[88699] new Indian variant, G6PD Jammu. In: Sen, N. N.; Basu, A. K.: Trends

[88700] in Haematology. Calcutta: Chatterjea Memorial Committee (pub.)

[88701] 1975. Pp. 279–283.

[88702] 15644.

[88703] Beutler, E.: Personal Communication. La Jolla, Calif. 11/12/1990.

[88704] 15645.

[88705] Beutler, E.: G6PD deficiency. Blood 84: 3613–3636, 1994.

[88706] 15646.

[88707] Beutler, E.: Glucose–6–phosphate dehydrogenase deficiency. New

[88708] Eng. J. Med. 324: 169–174, 1991.

[88709] 15647.

[88710] Beutler, E.: Selectivity of proteases as a basis for tissue

distribution

[88711] of enzymes in hereditary deficiencies. Proc. Nat. Acad. Sci.
80:

[88712] 3767–3768, 1983.

[88713] 15648.

[88714] Beutler, E.: Glucose–6–phosphate dehydrogenase deficiency. In:

[88715] Wintrobe, M. M.: Red Cell Metabolism in Hemolytic Anemia. New York:

[88716] Plenum Press (pub.) 1978.

[88717] 15649.

[88718] Beutler, E.: The hemolytic effect of primaquine and related compounds:

[88719] a review. Blood 14: 103–139, 1959.

[88720] 15650.

[88721] Beutler, E.; Grooms, A. M.; Morgan, S. K.; Trinidad, F.: Chronic

[88722] severe hemolytic anemia due to G–6–PD Charleston: a new deficiency

[88723] variant. J. Pediat. 80: 1005–1009, 1972.

[88724] 15651.

- [88725] Beutler, E.; Hartman, K.; Gelbart, T.; Forman, L.: G-6-PD
Walter
- [88726] Reed: possible insight into 'structural' NADP in G-6-PD.
Am. J. Hemat. 23:
- [88727] 25-30, 1986.
- [88728] 15652.
- [88729] Beutler, E.; Keller, J. W.; Matsumoto, F.: A new glucose
6-phosphate
- [88730] dehydrogenase (G6PD) variant associated with nonsphero-
cytic hemolytic
- [88731] anemia: G6PD Atlanta. I.R.C.S. 4: 479, 1976.
- [88732] 15653.
- [88733] Beutler, E.; Kuhl, W.: The NT 1311 polymorphism of G6PD:
G6PD
- [88734] Mediterranean mutation may have originated indepen-
dently in Europe
- [88735] and Asia. Am. J. Hum. Genet. 47: 1008-1012, 1990.
- [88736] 15654.
- [88737] Beutler, E.; Kuhl, W.: Linkage between a PvuII restriction
fragment
- [88738] length polymorphism and G6PD A- (202A/376G): evi-

dence for a single

[88739] origin of the common G6PD A- mutation. Hum. Genet. 85: 9-11, 1990.

[88740] 15655.

[88741] Beutler, E.; Kuhl, W.; Gelbart, T.; Forman, L.: DNA sequence

[88742] abnormalities of human glucose-6-phosphate dehydrogenase variants. J.

[88743] Biol. Chem. 266: 4145-4150, 1991.

[88744] 15656.

[88745] Beutler, E.; Kuhl, W.; Ramirez, E.; Lisker, R.: Some Mexican

[88746] glucose-6-phosphate dehydrogenase variants revisited. Hum. Genet. 86:

[88747] 371-374, 1991.

[88748] 15657.

[88749] Beutler, E.; Kuhl, W.; Saenz, G. F.; Rodriguez, W.: Mutation

[88750] analysis of glucose-6-phosphate dehydrogenase (G6PD) variants in Costa

[88751] Rica. Hum. Genet. 87: 462-464, 1991.

[88752] 15658.

- [88753] Beutler, E.; Mathai, C. K.; Smith, J. E.: Biochemical variants
[88754] of glucose-6-phosphate dehydrogenase giving rise to
congenital nonspherocytic
[88755] hemolytic disease. Blood 31: 131-150, 1968.
[88756] 15659.
- [88757] Beutler, E.; Matsumoto, F.: A new glucose 6-phosphate
dehydrogenase
[88758] variant: G6PD (-) Los Angeles. I.R.C.S. 5: 89, 1977.
[88759] 15660.
- [88760] Beutler, E.; Matsumoto, F.; Daiber, A.: Nonspherocytic
hemolytic
[88761] anemia due to G-6-PD Panama. I.R.C.S. 2: 1389, 1974.
[88762] 15661.
- [88763] Beutler, E.; Rosen, R.: Nonspherocytic congenital
hemolytic anemia
[88764] due to a new G-6-PD variant: G-6-PD Alhambra. Pediatrics 45: 230-235,
[88765] 1970.
[88766] 15662.
- [88767] Beutler, E.; Westwood, B.; Prchal, J. T.; Vaca, G.; Bartsocas,

- [88768] C. S.; Baronciani, L.: New glucose-6-phosphate dehydrogenase mutations
- [88769] from various ethnic groups. Blood 80: 255-256, 1992.
- [88770] 15663.
- [88771] Beutler, E.; Yoshida, A.: Human glucose-6-phosphate dehydrogenase
- [88772] variants: a supplementary tabulation. Ann. Hum. Genet. 37: 151-156,
- [88773] 1973.
- [88774] 15664.
- [88775] Boivin, P.; Galand, C.: Nouvelles variantes de la glucose-6-phosphate
- [88776] dehydrogenase erythrocytaire. Rev. Franc. Etud. Clin. Biol. 13:
- [88777] 30-39, 1968.
- [88778] 15665.
- [88779] Botha, M. C.; Dern, R. J.; Mitchell, M.; West, C.; Beutler, E.
- [88780] : G6PD Capetown, a variant of glucose-6-phosphate dehydrogenase. Am.
- [88781] J. Hum. Genet. 21: 547-551, 1969.

- [88782] 15666.
- [88783] Boyer, S. H.; Graham, J. B.: Linkage between the X chromosome
- [88784] loci for glucose-6-phosphate dehydrogenase electrophoretic variation
- [88785] and hemophilia A. Am. J. Hum. Genet. 17: 320-324, 1965.
- [88786] 15667.
- [88787] Boyer, S. H.; Porter, I. H.; Weilbaecher, R. G.: Electrophoretic
- [88788] heterogeneity of glucose-6-phosphate dehydrogenase and its relationship
- [88789] to enzyme deficiency in man. Proc. Nat. Acad. Sci. 48: 1868-1876,
- [88790] 1962.
- [88791] 15668.
- [88792] Busch, D.; Bote, K.: Glucose-6-phosphate-dehydrogenase-Defect
- [88793] in Deutschland. II. Eigenschaften des Enzyms (Typ Freiburg). Klin.
- [88794] Wschr. 48: 74-78, 1970.
- [88795] 15669.

- [88796] Calabro, V.; Giacobbe, A.; Vallone, D.; Montanaro, V.; Cascone,
[88797] A.; Filosa, S.; Battistuzzi, G.: Genetic heterogeneity at the
glucose-6-phosphate
[88798] dehydrogenase locus in southern Italy: a study on a population from
[88799] the Matera district. Hum. Genet. 86: 49-53, 1990.
[88800] 15670.
- [88801] Cappadoro, M.; Giribaldi, G.; O'Brien, E.; Turrini, F.;
Mannu,
[88802] F.; Ulliers, D.; Simula, G.; Luzzatto, L.; Arese, P.: Early
phagocytosis
[88803] of glucose-6-phosphate dehydrogenase (G6PD)-deficient
erythrocytes
[88804] parasitized by Plasmodium falciparum may explain
malaria protection
[88805] in G6PD deficiency. Blood 92: 2527-2534, 1998.
[88806] 15671.
- [88807] Cappellini, M. D.; Sampietro, M.; Toniolo, D.; Carandina,
G.;
[88808] Pittalis, S.; Martinez di Montemuros, F.; Tavazzi, D.;
Fiorelli, G.

- [88809] : Biochemical and molecular characterization of a new sporadic glucose-6-phosphate
- [88810] dehydrogenase variant described in Italy: G6PD Modena. Brit. J. Haemat. 87:
- [88811] 209-211, 1994.
- [88812] 15672.
- [88813] Carandina, G.; Moretto, E.; Zecchi, G.; Conighi, C.: Glucose
- [88814] 6-phosphate dehydrogenase Ferrara. A new variant of G6PD identified
- [88815] in Northern Italy. Acta Haemat. 56: 116-122, 1976.
- [88816] 15673.
- [88817] Carson, P. E.; Flanagan, C. L.; Ickes, C. E.; Alving, A. S.:
- [88818] Enzymatic deficiency in primaquine-sensitive erythrocytes. Science 124:
- [88819] 484-485, 1956.
- [88820] 15674.
- [88821] Poy, F.; Yaffe, M. B.; Sayos, J.; Saxena, K.; Morra, M.; Sumegi,
- [88822] J.; Cantley, L. C.; Terhorst, C.; Eck, M. J.: Crystal structures
- [88823] of the XLP protein SAP reveal a class of SH2 domains with

extended,

[88824] phosphotyrosine-independent sequence recognition.

Molec. Cell 4:

[88825] 555-561, 1999.

[88826] 15675.

[88827] Provisor, A. J.; Iaccone, J. J.; Chilcote, R. R.; Neiburger, R.

[88828] G.; Crussi, F. G.; Baehner, R. L.: Acquired agammaglobulinemia after

[88829] a life-threatening illness with clinical and laboratory features of

[88830] infectious mononucleosis in three related male children.
New Eng.

[88831] J. Med. 293: 62-65, 1975.

[88832] 15676.

[88833] Purtilo, D. T.: Pathogenesis and phenotypes of an X-linked
recessive

[88834] lymphoproliferative syndrome. Lancet II: 882-885, 1976.

[88835] 15677.

[88836] Purtilo, D. T.: X-linked lymphoproliferative syndrome: an
immunodeficiency

[88837] disorder with acquired agammaglobulinemia, fatal infec-

tious mononucleosis,

[88838] or malignant lymphoma. Arch. Path. Lab. Med. 105:
119-121, 1981.

[88839] 15678.

[88840] Purtilo, D. T.; Bhawan, J.; Hutt, L. M.; De Nicola, L.; Szy-
manski,

[88841] I.; Yang, J. P. S.; Boto, W.; Naier, R.; Thorley-Lawson, D.:
Epstein-Barr

[88842] virus in the X-linked recessive lymphoproliferative syn-
drome. Lancet I:

[88843] 798-801, 1978.

[88844] 15679.

[88845] Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.: Fatal infectious

[88846] mononucleosis in familial lymphohistiocytosis. (Letter)
New Eng.

[88847] J. Med. 201: 736 only, 1974.

[88848] 15680.

[88849] Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.; Harper, R.;
Stephenson,

[88850] S. R.; Landing, B. H.; Vewter, G. F.: X-linked recessive pro-
gressive

[88851] combined variable immunodeficiency (Duncan's disease).

Lancet I:

[88852] 935–941, 1975.

[88853] 15681.

[88854] Purtilo, D. T.; DeFlorio, D., Jr.; Hutt, L. M.; Bhawan, J.;

Yang,

[88855] J. P. S.; Otto, R. L.; Edwards, W.: Variable phenotypic expression

[88856] of an X-linked recessive lymphoproliferative syndrome.

New Eng. J.

[88857] Med. 297: 1077–1081, 1977.

[88858] 15682.

[88859] Purtilo, D. T.; Grierson, H. L.: Methods of detection of new

[88860] families with X-linked lymphoproliferative disease. Cancer Genet.

[88861] Cytogenet. 51: 143–153, 1991.

[88862] 15683.

[88863] Purtilo, D. T.; Sakamoto, K.; Barnabei, V.; Seeley, J.; Bechtold,

[88864] T.; Rogers, G.; Yetz, J.; Harada, S.; the XLP collaborators: Epstein–Barr

- [88865] virus-induced diseases in boys with the X-linked lymphoproliferative
- [88866] syndrome (XLP): update on studies of the registry. *Am. J. Med.* 73:
- [88867] 49-56, 1982.
- [88868] 15684.
- [88869] Purtilo, D. T.; Yang, J. P. S.; Allegra, S.; DeFlorio, D.; Hutt,
- [88870] L. M.; Soltani, M.; Vawter, G. F.: Hematopathology and pathogenesis
- [88871] of the X-linked recessive lymphoproliferative syndrome. *Am. J. Med.* 62:
- [88872] 225-233, 1977.
- [88873] 15685.
- [88874] Sanger, W. G.; Grierson, H. L.; Skare, J.; Wyandt, H.; Pirruccello,
- [88875] S.; Fordyce, R.; Purtilo, D. T.: Partial Xq25 deletion in a family
- [88876] with the X-linked lymphoproliferative disease (XLP). *Cancer Genet.*
- [88877] *Cytogenet.* 47: 163-169, 1990.
- [88878] 15686.

- [88879] Sayos, J.; Wu, C.; Morra, M.; Wang, N.; Zhang, X.; Allen, D.;
- [88880] van Schaik, S.; Notarangelo, L.; Gehat, R.; Roncarolo, M. G.; Oettgen,
- [88881] H.; De Vries, J. E.; Aversall, G.; Terhorst, C.: The X-linked lymphoproliferative-disease
- [88882] gene product SAP regulates signals induced through the co-receptor
- [88883] SLAM. Nature 395: 462-469, 1998.
- [88884] 15687.
- [88885] Scher, I.: The CBA/N mouse strain: an experimental model illustrating
- [88886] the influence of the X-chromosome on immunity. Adv. Immun. 33: 1-71,
- [88887] 1982.
- [88888] 15688.
- [88889] Schuster, V.; Kreth, H. W.: X-linked lymphoproliferative disease.In:
- [88890] Ochs, H. D.; Smith, C. I. E.; Puck, J. M. (eds.): Primary Immunodeficiency
- [88891] Diseases: A Molecular and Genetic Approach. New York: Oxford University

[88892] Press 1999. Pp. 222–232.

[88893] 15689.

[88894] Seemayer, T. A.; Gross, T. G.; Egeler, R. M.; Pirruccello, S.

[88895] J.; Davis, D. J.; Kelly, C. M.; Okano, M.; Lanyi, A.; Sumegi, J.:

[88896] X-linked lymphoproliferative disease: twenty-five years after the

[88897] discovery. *Pediat. Res.* 38: 471–478, 1995.

[88898] 15690.

[88899] Skare, J.; Grierson, H.; Wyandt, H.; Sanger, W.; Milunsky, J.;

[88900] Purtilo, D.; Sullivan, J.; Milunsky, A.: Genetics of the X-linked

[88901] lymphoproliferative syndrome. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.):

[88902] A161 only, 1989.

[88903] 15691.

[88904] Skare, J.; Madan, S.; Glaser, J.; Purtilo, D.; Nitowsky, H.; Pulijaal,

[88905] V.; Milunsky, A.: First prenatal diagnosis of X-linked lymphoproliferative

- [88906] disease. Am. J. Med. Genet. 44: 79–81, 1992.
- [88907] 15692.
- [88908] Skare, J.; Milunsky, A.; Byron, K.; Sullivan, J.: The mutation
- [88909] causing X-linked lymphoproliferative syndrome lies in Xq26. (Abstract) Am.
- [88910] J. Hum. Genet. 41: A185 only, 1987.
- [88911] 15693.
- [88912] Skare, J.; Wu, B.-L.; Madan, S.; Pulijaal, V.; Purtilo, D.;
- Haber,
- [88913] D.; Nelson, D.; Sylla, B.; Grierson, H.; Nitowsky, H.; Glaser, J.;
- [88914] Wissink, J.; White, B.; Holden, J.; Housman, D.; Lenoir, G.; Wyandt,
- [88915] H.; Milunsky, A.: Characterization of three overlapping deletions
- [88916] causing X-linked lymphoproliferative disease. Genomics 16: 254–255,
- [88917] 1993.
- [88918] 15694.
- [88919] Skare, J. C.; Grierson, H. L.; Sullivan, J. L.; Nussbaum, R. L.;

- [88920] Purtilo, D. T.; Sylla, B. S.; Lenoir, G. M.; Reilly, D. S.; White,
[88921] B. N.; Milunsky, A.: Linkage analysis of seven kindreds
with the
[88922] X-linked lymphoproliferative syndrome (XLP) confirms
that the XLP
[88923] locus is near DXS42 and DXS37. Hum. Genet. 82:
354-358, 1989.
[88924] 15695.
- [88925] Skare, J. C.; Milunsky, A.; Byron, K. S.; Sullivan, J. L.: Map-
ping
[88926] the X-linked lymphoproliferative syndrome. Proc. Nat.
Acad. Sci. 84:
[88927] 2015-2018, 1987.
[88928] 15696.
- [88929] Skare, J. C.; Sullivan, J. L.; Milunsky, A.: Mapping the mu-
tation
[88930] causing the X-linked lymphoproliferative syndrome in re-
lation to restriction
[88931] fragment length polymorphisms on Xq. Hum. Genet. 82:
349-353, 1989.
[88932] 15697.

- [88933] Steinherz, R.; Levy, Y.; Litwin, A.; Nitzan, M.; Friedman, E.;
- [88934] Levin, S.: X-linked lymphoproliferative syndrome: a new
kindred with
- [88935] variable phenotypic expression. *Am. J. Dis. Child.* 139:
191–193,
- [88936] 1985.
- [88937] 15698.
- [88938] Nehlin, J. O.; Hara, E.; Kuo, W.-L.; Collins, C.; Campisi, J.:
- [88939] Genomic organization, sequence, and chromosomal local-
ization of the
- [88940] human helix-loop-helix Id1 gene. *Biochem. Biophys. Res.*
Commun. 231:
- [88941] 628–634, 1997.
- [88942] 15699.
- [88943] Inoue, S.; Orimo, A.; Matsuda, Y.; Inazawa, J.; Emi, M.;
Nakamura,
- [88944] Y.; Hori, T.; Muramatsu, M.: Chromosome mapping of hu-
man (ZNF147)
- [88945] and mouse genes for estrogen-responsive finger protein
(efp), a member
- [88946] of the RING finger family. *Genomics* 25: 581–583, 1995.

[88947] 15700.

[88948] Law, D. J.; Prasad, M. A.; King, S. E.; Spranger, K. D.; Lee, Y.

[88949] H.; Fox, R. E.; Collins, E. E.; Gebuhr, T. C.; Miller, D. E.; Petty,

[88950] E. M.: Localization of the human estrogen-responsive finger protein

[88951] (EFP) gene (ZNF147) within a YAC contig containing the myeloperoxidase

[88952] (MPO) gene. Genomics 28: 361–363, 1995.

[88953] 15701.

[88954] Fujii, H.; Chen, S.-H.; Akatsuka, J.; Miwa, S.; Yoshida, A.: Use

[88955] of cultured lymphoblastoid cells for the study of abnormal enzymes:

[88956] molecular abnormality of a phosphoglycerate kinase variant associated

[88957] with hemolytic anemia. Proc. Nat. Acad. Sci. 78: 2587–2590, 1981.

[88958] 15702.

[88959] Lindenbaum, R. H.; Clarke, G.; Patel, C.; Moncrieff, M.; Hughes,

[88960] J. T.: Muscular dystrophy in an X;1 translocation female suggests
[88961] that Duchenne locus is on X chromosome short arm. J.
Med. Genet. 16:
[88962] 389–392, 1979.
[88963] 15703.
[88964] Loftin, C. D.; Trivedi, D. B.; Tiano, H. F.; Clark, J. A.; Lee,
[88965] C. A.; Epstein, J. A.; Morham, S. G.; Breyer, M. D.; Nguyen,
M.; Hawkins,
[88966] B. M.; Goulet, J. L.; Smithies, O.; Koller, B. H.; Langenbach,
R.
[88967] : Failure of ductus arteriosus closure and remodeling in
neonatal
[88968] mice deficient in cyclooxygenase–1 and cyclooxygenase–2.
Proc. Nat.
[88969] Acad. Sci. 98: 1059–1064, 2001.
[88970] 15704.
[88971] Macchia, L.; Di Paola, R.; Guerrese, M.–C.; Chiechi, L. M.;
Tursi,
[88972] A.; Caiaffa, M. F.; Haeggstrom, J. Z.: Expression of
prostaglandin

- [88973] endoperoxide H synthase 1 and 2 in human placenta at term. Biochem.
- [88974] Biophys. Res. Commun. 233: 496–501, 1997.
- [88975] 15705.
- [88976] Morham, S. G.; Langenbach, R.; Loftin, C. D.; Tiano, H. F.; Vouloumanos,
- [88977] N.; Jennette, J. C.; Mahler, J. F.; Kluckman, K. D.; Ledford, A.;
- [88978] Lee, C. A.; Smithies, O.: Prostaglandin synthase 2 gene disruption
- [88979] causes severe renal pathology in the mouse. Cell 83: 473–482, 1995.
- [88980] 15706.
- [88981] Neufang, G.; Furstenberger, G.; Heidt, M.; Marks, F.; Muller-Decker,
- [88982] K.: Abnormal differentiation of epidermis in transgenic mice constitutively
- [88983] expressing cyclooxygenase–2 in skin. Proc. Nat. Acad. Sci. 98: 7629–7634,
- [88984] 2001.
- [88985] 15707.

[88986] O'Banion, M. K.; Sadowski, H. B.; Winn, V.; Young, D. A.: A serum-
[88987] and glucocorticoid-regulated 4-kilobase mRNA encodes a
cyclooxygenase-related
[88988] protein. J. Biol. Chem. 266: 23261-23267, 1991.
[88989] 15708.
[88990] O'Banion, M. K.; Winn, V. D.; Young, D. A.: cDNA cloning
and
[88991] functional activity of a glucocorticoid-regulated inflam-
matory cyclooxygenase. Proc.
[88992] Nat. Acad. Sci. 89: 4888-4892, 1992.
[88993] 15709.
[88994] Oshima, M.; Dinchuk, J. E.; Kargman, S. L.; Oshima, H.;
Hancock,
[88995] B.; Kwong, E.; Trzaskos, J. M.; Evans, J. F.; Taketo, M. M.:
Suppression
[88996] of intestinal polyposis in Apc(delta-716) knockout mice by
inhibition
[88997] of cyclooxygenase 2 (COX-2). Cell 87: 803-809, 1996.
[88998] 15710.
[88999] Salmenkivi, K.; Haglund, C.; Ristimaki, A.; Arola, J.;

Heikkila,

[89000] P.: Increased expression of cyclooxygenase-2 in malignant pheochromocytomas. J.

[89001] Clin. Endocr. Metab. 86: 5615-5619, 2001.

[89002] 15711.

[89003] Tay, A.; Squire, J. A.; Goldberg, H.; Skorecki, K.: Assignment

[89004] of the human prostaglandin-endoperoxide synthase 2 (PTGS2) gene to

[89005] 1q25 by fluorescence in situ hybridization. Genomics 23: 718-719,

[89006] 1994.

[89007] 15712.

[89008] Swan, S. K.; Rudy, D. W.; Lasseter, K. C.; Ryan, C. F.; Buechel,

[89009] K. L.; Lambrecht, L. J.; Pinto, M. B.; Dilzer, S. C.; Obrda, O.; Sundblad,

[89010] K. J.; Gumbs, C. P.; Ebel, D. L.; Quan, H.; Larson, P. J.; Schwartz,

[89011] J. I.; Musliner, T. A.; Gertz, B. J.; Brater, D. C.; Yao, S.-L.:

[89012] Effect of cyclooxygenase-2 inhibition on renal function in

elderly

[89013] persons receiving a low-salt diet: a randomized controlled trial. *Ann.*

[89014] *Intern. Med.* 133: 1-9, 2000.

[89015] 15713.

[89016] Tazawa, R.; Xu, X.-M.; Wu, K. K.; Wang, L.-H.: Characterization

[89017] of the genomic structure, chromosomal location and promoter of human

[89018] prostaglandin H synthase-2 gene. *Biochem. Biophys. Res. Commun.* 203:

[89019] 190-199, 1994.

[89020] 15714.

[89021] Tsujii, M.; DuBois, R. N.: Alterations in cellular adhesion and

[89022] apoptosis in epithelial cells overexpressing prostaglandin endoperoxide

[89023] synthase 2. *Cell* 83: 493-501, 1995.

[89024] 15715.

[89025] Zhou, X.-L.; Lei, Z. M.; Rao, C. V.: Treatment of human endometrial

- [89026] gland epithelial cells with chorionic gonadotropin/luteinizing hormone
- [89027] increases the expression of the cyclooxygenase-2 gene. J. Clin. Endocr.
- [89028] Metab. 84: 3364–3377, 1999.
- [89029] 15716.
- [89030] Inoue, S.; Orimo, A.; Hosoi, T.; Kondo, S.; Toyoshima, H.; Kondo,
- [89031] T.; Ikegami, A.; Ouchi, Y.; Orimo, H.; Muramatsu, M.: Genomic binding-site
- [89032] cloning reveals an estrogen-responsive gene that encodes a RING finger
- [89033] protein. Proc. Nat. Acad. Sci. 90: 11117–11121, 1993.
- [89034] 15717.
- [89035] Maho, A.; Bensimon, A.; Vassart, G.; Parmentier, M.: Mapping of
- [89036] the CCXCR1, CX3CR1, CCBP2 and CCR9 genes to the CCR cluster within
- [89037] the 3p21.3 region of the human genome. Cytogenet. Cell Genet. 87:
- [89038] 265–268, 1999.

[89039] 15718.

[89040] Anderson, D. M.; Johnson, L.; Glaccum, M. B.; Copeland, N. G.;

[89041] Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Kirstein, M. N.; Shapiro,

[89042] D. N.; Morris, S. W.; Grabstein, K.; Cosman, D.: Chromosomal assignment

[89043] and genomic structure of IL15. Genomics 25: 701-706, 1995.

[89044] 15719.

[89045] Grabstein, K. H.; Eisenman, J.; Shanebeck, K.; Rauch, C.; Srinivasan,

[89046] S.; Fung, V.; Beers, C.; Richardson, J.; Schoenborn, M. A.; Ahdieh,

[89047] M.; Johnson, L.; Alderson, M. R.; Watson, J. D.; Anderson, D. M.;

[89048] Giri, J. G.: Cloning of a T cell growth factor that interacts with

[89049] the beta chain of the interleukin-2 receptor. Science 264: 965-968,

[89050] 1994.

[89051] 15720.

- [89052] Roberts, A. I.; Lee, L.; Schwarz, E.; Groh, V.; Spies, T.; Ebert,
E. C.; Jabri, B.: Cutting edge: NKG2D receptors induced by IL-15
costimulate CD28-negative effector CTL in the tissue microenvironment. J.
Immun. 167: 5527-5530, 2001.
- [89056] 15721.
- [89057] Robinson, P.; Okhuysen, P. C.; Chappell, C. L.; Lewis, D. E.; Shahab,
I.; Lahoti, S.; White, A. C., Jr.: Expression of IL-15 and IL-4
in
IFN-gamma-independent control of experimental human
Cryptosporidium
parvum infection. Cytokine 15: 39-46, 2001.
- [89061] 15722.
- [89062] Chen, X.; Vinkemeier, U.; Zhao, Y.; Jeruzalmi, D.; Darnell,
J.
E., Jr.; Kuriyan, J.: Crystal structure of a tyrosine phosphorylated
STAT-1 dimer bound to DNA. Cell 93: 827-839, 1998.

- [89065] 15723.
- [89066] Copeland, N. G.; Gilbert, D. J.; Schindler, C.; Zhong, Z.;
Wen,
- [89067] Z.; Darnell, J. E., Jr.; Mui, A. L.-F.; Miyajima, A.; Quelle, F.
W.;
- [89068] Ihle, J. N.; Jenkins, N. A.: Distribution of the mammalian
Stat gene
- [89069] family in mouse chromosomes. Genomics 29: 225–228,
1995.
- [89070] 15724.
- [89071] Darnell, J. E., Jr.; Kerr, I. M.; Stark, G. M.: Jak–STAT path-
ways
- [89072] and transcriptional activation in response to IFNs and
other extracellular
- [89073] signaling proteins. Science 264: 1415–1421, 1994.
- [89074] 15725.
- [89075] Durbin, J. E.; Hackenmiller, R.; Simon, M. C.; Levy, D. E.:
Targeted
- [89076] disruption of the mouse Stat1 gene results in compro-
mised innate immunity
- [89077] to viral disease. Cell 84: 443–450, 1996.

- [89078] 15726.
- [89079] Haddad, B.; Pabon-Pena, C. R.; Young, H.; Sun, W. H.: Assignment
[89080] of STAT1 to human chromosome 2q32 by FISH and radiation hybrids. Cytogenet.
[89081] Cell Genet. 83: 58-59, 1998.
- [89082] 15727.
- [89083] Ihle, J. N.: STATs: signal transducers and activators of transcription. Cell 84:
[89084] 331-334, 1996.
- [89085] 15728.
- [89086] Meraz, M. A.; White, J. M.; Sheehan, K. C. F.; Bach, E. A.; Rodig,
[89087] S. J.; Dighe, A. S.; Kaplan, D. H.; Riley, J. K.; Greenlund, A. C.
- [89088] Campbell, D.; Carver-Moore, K.; DuBois, R. N.; Clark, R.; Aguet, M.;
- [89089] Schreiber, R. D.: Targeted disruption of the Stat1 gene in mice reveals
[89090] unexpected physiologic specificity in the JAK-STAT signaling pathway. Cell 84:

- [89091] 431–442, 1996.
- [89092] 15729.
- [89093] Mowen, K. A.; Tang, J.; Zhu, W.; Schurter, B. T.; Shuai, K.; Herschman,
- [89094] H. R.; David, M.: Arginine methylation of STAT1 modulates IFN- α /beta-induced
- [89095] transcription. *Cell* 104: 731–741, 2001.
- [89096] 15730.
- [89097] Ramana, C. V.; Chatterjee-Kishore, M.; Nguyen, H.; Stark, G. R.
- [89098] : Complex roles of Stat1 in regulating gene expression. *Oncogene* 19:
- [89099] 2619–2627, 2000.
- [89100] 15731.
- [89101] Yamamoto, K.; Kobayashi, H.; Arai, A.; Miura, O.; Hiro-sawa, S.;
- [89102] Miyasaka, N.: cDNA cloning, expression and chromosome mapping of
- [89103] the human STAT4 gene: both STAT4 and STAT1 genes are mapped to 2q32.2–q32.3. *Cytogenet.*
- [89104] *Cell Genet.* 77: 207–210, 1997.

- [89105] 15732.
- [89106] Fukuda, M.: Biogenesis of the lysosomal membrane. Sub-cell. Biochem. 22:
- [89107] 199–230, 1994.
- [89108] 15733.
- [89109] Fukuda, M.; Viitala, J.; Matteson, J.; Carlsson, S. R.: Cloning
- [89110] of the cDNAs encoding human lysosomal membrane glycoproteins, h-lamp-1
- [89111] and h-lamp-2: comparison of their deduced amino acid sequences. J.
- [89112] Biol. Chem. 263: 18920–18928, 1988.
- [89113] 15734.
- [89114] Konecki, D. S.; Foetisch, K.; Schlotter, M.; Lichter-Konecki, U.
- [89115] : Complete cDNA sequence of human lysosome-associated membrane protein-2. Biochem.
- [89116] Biophys. Res. Commun. 205: 1–5, 1994.
- [89117] 15735.
- [89118] Lindsay, S.; Bird, A. P.: Use of restriction enzymes to detect

- [89119] potential gene sequences in mammalian DNA. *Nature* 327: 336–338,
- [89120] 1987.
- [89121] 15736.
- [89122] Manoni, M.; Tribioli, C.; Lazzari, B.; DeBellis, G.; Patrosso,
- [89123] C.; Pergolizzi, R.; Pellegrini, M.; Maestrini, E.; Rivella, S.; Vezzoni,
- [89124] P.; Toniolo, D.: The nucleotide sequence of a CpG island demonstrates
- [89125] the presence of the first exon of the gene encoding the human lysosomal
- [89126] membrane protein LAMP2 and assigns the gene to Xq24. *Genomics* 9: 551–554, 1991.
- [89127] 15737.
- [89128] 15737.
- [89129] Takahashi, M.; Yamamoto, A.; Takano, K.; Sudo, A.; Wada, T.; Goto,
- [89130] Y. I.; Nishino, I.; Saitoh, S.: Germline mosaicism of a novel mutation
- [89131] in lysosome-associated membrane protein-2 deficiency (Danon disease). *Ann.*

- [89132] Neurol. 52: 122–125, 2002.
- [89133] 15738.
- [89134] Tanaka, Y.; Guhe, G.; Suter, A.; Eskelinen, E.–L.; Hartmann, D.;
- [89135] Lullmann–Rauch, R.; Janssen, P. M. L.; Blanz, J.; von Figura, K.;
- [89136] Saftig, P.: Accumulation of autophagic vacuoles and cardiomyopathy
- [89137] in LAMP–2–deficient mice. Nature 406: 902–906, 2000.
- [89138] 15739.
- [89139] Falls, H. F.: The role of the sex chromosome in hereditary ocular
- [89140] pathology. Trans. Am. Ophthal. Soc. 50: 421–467, 1952.
- [89141] 15740.
- [89142] Rosse, W. F.: Paroxysmal nocturnal hemoglobinuria: the biochemical
- [89143] defects and the clinical syndrome. Blood Rev. 3: 192–200, 1989.
- [89144] 15741.
- [89145] Rotoli, B.; Luzzatto, L.: Paroxysmal nocturnal haemoglobinuria. Clin.

- [89146] Haemat. 2: 113–138, 1989.
- [89147] 15742.
- [89148] Sauer, B.; Henderson, N.: Site-specific DNA recombination
in
- [89149] mammalian cells by the Cre recombinase of bacteriophage
P1. Proc.
- [89150] Nat. Acad. Sci. 85: 5166–5170, 1988.
- [89151] 15743.
- [89152] Savoia, A.; Ianzano, L.; Lunardi, C.; De Sandre, G.;
Carotenuto,
- [89153] M.; Musto, P.; Zelante, L.: Identification of three novel mu-
tations
- [89154] in the PIG-A gene in paroxysmal nocturnal haemoglobin-
uria (PNH) patients. Hum.
- [89155] Genet. 97: 45–48, 1996.
- [89156] 15744.
- [89157] Socie, G.; Mary, J.-Y.; de Gramont, A.; Rio, B.; Leporrier,
M.;
- [89158] Rose, C.; Heudier, P.; Rochant, H.; Cahn, J.-Y.; Gluckman,
E.: Paroxysmal
- [89159] nocturnal haemoglobinuria: long-term follow-up and

prognostic factors. Lancet 348:

[89160] 573–577, 1996.

[89161] 15745.

[89162] Stevens, V. L.; Raetz, C. R. H.: Defective glycosyl phosphatidylinositol

[89163] biosynthesis in extracts of three thy-1 negative lymphoma cell mutants. J.

[89164] Biol. Chem. 266: 10039–10042, 1991.

[89165] 15746.

[89166] Sugiyama, E.; DeGasperi, R.; Urakaze, M.; Chang, H.-M.; Thomas,

[89167] L. J.; Hyman, R.; Warren, C. D.; Yeh, E. T. H.: Identification of

[89168] defects in glycosylphosphatidylinositol anchor biosynthesis in the

[89169] thy-1 expression mutants. J. Biol. Chem. 266: 12119–12122, 1991.

[89170] 15747.

[89171] Takahashi, M.; Takeda, J.; Hirose, S.; Hyman, R.; Inoue, N.; Miyata,

[89172] T.; Ueda, E.; Kitani, T.; Medof, M. E.; Kinoshita, T.: Defi-

cient

- [89173] biosynthesis of N-acetylglucosaminyl-phosphatidylinositol, the first intermediate of glycosyl phosphatidylinositol anchor biosynthesis,
- [89174] in cell lines established from patients with paroxysmal nocturnal hemoglobinuria. J. Exp. Med. 177: 517-521, 1993.
- [89175] 15748.
- [89176] Takeda, J.; Miyata, T.; Kawagoe, K.; Iida, Y.; Endo, Y.; Fujita, T.; Takahashi, M.; Kitani, T.; Kinoshita, T.: Deficiency of the GPI anchor caused by a somatic mutation of the PIG-A gene in paroxysmal nocturnal hemoglobinuria. Cell 73: 703-711, 1993.
- [89177] 15749.
- [89178] Ueda, E.; Nishimura, J.; Kitani, T.; Nasu, K.; Kageyama, T.; Kim, Y. U.; Takeda, J.; Kinoshita, T.: Deficient surface expression of glycosylphosphatidylinositol-anchored proteins in B cell

lines established

[89186] from patients with paroxysmal nocturnal hemoglobinuria.

Int. Immun. 4:

[89187] 1263–1271, 1992.

[89188] 15750.

[89189] Ware, R. E.; Howard, T. A.; Kamitani, T.; Chang, H.–M.;

Yeh, E.

[89190] T. H.; Seldin, M. F.: Chromosomal assignment of genes involved in

[89191] glycosylphosphatidylinositol anchor biosynthesis: implications for

[89192] the pathogenesis of paroxysmal nocturnal hemoglobinuria. Blood 83:

[89193] 3753–3757, 1994.

[89194] 15751.

[89195] Ware, R. E.; Rosse, W. F.; Howard, T. A.: Mutations within the

[89196] Piga gene in patients with paroxysmal nocturnal hemoglobinuria. Blood 83:

[89197] 2418–2422, 1994.

[89198] 15752.

- [89199] Watanabe, R.; Inoue, N.; Westfall, B.; Taron, C. H.; Orlean, P.;
- [89200] Takeda, J.; Kinoshita, T.: The first step of glycosylphosphatidylinositol
- [89201] biosynthesis is mediated by a complex of PIG-A, PIG-H, PIG-C and GPI1. EMBO
- [89202] J. 17: 877-885, 1998.
- [89203] 15753.
- [89204] Watanabe, R.; Kinoshita, T.; Masaki, R.; Yamamoto, A.; Takeda,
- [89205] J.; Inoue, N.: PIG-A and PIG-H, which participate in glycosylphosphatidylinositol
- [89206] anchor biosynthesis, form a protein complex in the endoplasmic reticulum. J.
- [89207] Biol. Chem. 271: 26868-26875, 1996.
- [89208] 15754.
- [89209] Yeh, E. T. H.; Rosse, W. F.: Paroxysmal nocturnal hemoglobinuria
- [89210] and the glycosylphosphatidylinositol anchor. J. Clin. Invest. 93:
- [89211] 2305-2310, 1994.

[89212] 15755.

[89213] Yu, J.; Nagarajan, S.; Ueda, E.; Knez, J. J.; Petersen, R. B.;

[89214] Medof, M. E.: Characterization of alternatively spliced PIG-A transcripts

[89215] in normal and paroxysmal nocturnal hemoglobinuria cells. Brazilian

[89216] J. Med. Biol. Res. 27: 195–201, 1994.

[89217] 15756.

[89218] Chen, S.-H.; Malcolm, L. A.; Yoshida, A.; Giblett, E. R.: Phosphoglycerate

[89219] kinase: an X-linked polymorphism in man. Am. J. Hum. Genet. 23:

[89220] 87–91, 1971.

[89221] 15757.

[89222] Cooper, D. W.; Johnston, P. G.; Murtagh, C. E.; Sharman, G. B.;

[89223] Vandeberg, J. L.; Poole, W. E.: Sex-linked isozymes and sex-chromosome

[89224] evolution and inactivation in kangaroos. In: Markert, C. L.: Isozymes.

[89225] Developmental Biology. New York: Academic Press (pub.)

III: 1975.

[89226] Pp. 559–573.

[89227] 15758.

[89228] Cooper, D. W.; Vandeberg, J. L.; Sharman, G. B.; Poole, W.
E.:

[89229] Phosphoglycerate kinase polymorphism in kangaroos provides further

[89230] evidence for paternal inactivation. Nature N.B. 230:
155–157, 1971.

[89231] 15759.

[89232] Deys, B. F.; Grzeschik, K.–H.; Grzeschik, A.; Jaffe, E. R.;
Siniscalco,

[89233] M.: Human phosphoglycerate kinase and inactivation of
the X chromosome. Science 175:

[89234] 1002–1003, 1972.

[89235] 15760.

[89236] DiMauro, S.; Dalakas, M.; Miranda, A. S.: Phosphoglycerate
kinase

[89237] deficiency: another cause of recurrent myoglobinuria.
Ann. Neurol. 13:

[89238] 11–19, 1983.

- [89239] 15761.
- [89240] Echard, G.; Gillois, M.: G6PD--alpha-GAL--PGK--HPRT
synteny in
- [89241] the rabbit, *Oryctolagus cuniculus*. (Abstract) Cytogenet.
Cell Genet. 25:
- [89242] 148-149, 1979.
- [89243] 15762.
- [89244] Patil, S. R.; Bartley, J. A.; Murray, J. C.; Ionasescu, V. V.;
- [89245] Pearson, P. L.: X-linked glycerol kinase, adrenal hypopla-
sia and
- [89246] myopathy maps at Xp21. (Abstract) Cytogenet. Cell Genet.
40: 720-721,
- [89247] 1985.
- [89248] 15763.
- [89249] Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.; Goldstein, L. S.:
- [89250] Axonal transport of amyloid precursor protein is mediated
by direct
- [89251] binding to the kinesin light chain subunit of kinesin-I.
Neuron 28:
- [89252] 449-459, 2000.

[89253] 15764.

[89254] Ahn, A. H.; Yoshida, M.; Anderson, M. S.; Feener, C. A.;
Selig,

[89255] S.; Hagiwara, Y.; Ozawa, E.; Kunkel, L. M.: Cloning of hu-
man basic

[89256] A1, a distinct 59-kDa dystrophin-associated protein en-
coded on chromosome

[89257] 8q23-24. Proc. Nat. Acad. Sci. 91: 4446-4450, 1994.

[89258] 15765.

[89259] Yang, B.; Ibraghimov-Beskrovnaya, O.; Moomaw, C. R.;
Slaughter,

[89260] C. A.; Campbell, K. P.: Heterogeneity of the 59-kDa dys-
trophin-associated

[89261] protein revealed by cDNA cloning and expression. J. Biol.
Chem. 269:

[89262] 6040-6044, 1994.

[89263] 15766.

[89264] Merendino, J. J., Jr.; Spiegel, A. M.; Crawford, J. D.;
O'Carroll,

[89265] A.-M.; Brownstein, M. J.; Lolait, S. J.: A mutation in the va-
sopressin

- [89266] V2-receptor gene in a kindred with X-linked nephrogenic diabetes insipidus. New
- [89267] Eng. J. Med. 328: 1538-1541, 1993.
- [89268] 15767.
- [89269] Moses, A. M.; Sangani, G.; Miller, J. L.: Proposed cause of marked
- [89270] vasopressin resistance in a female with an X-linked recessive V2 receptor
- [89271] abnormality. J. Clin. Endocr. Metab. 80: 1184-1186, 1995.
- [89272] 15768.
- [89273] Nakano, K. K.: Familial nephrogenic diabetes insipidus. Hawaii
- [89274] Med. J. 28: 205-208, 1969.
- [89275] 15769.
- [89276] Nomura, Y.; Onigata, K.; Nagashima, T.; Yutani, S.; Mochizuki,
- [89277] H.; Nagashima, K.; Morikawa, A.: Detection of skewed X-inactivation
- [89278] in two female carriers of vasopressin type 2 receptor gene mutation. J.
- [89279] Clin. Endocr. Metab. 82: 3434-3437, 1997.

- [89280] 15770.
- [89281] Pan, Y.; Metzenberg, A.; Das, S.; Jing, B.; Gitschier, J.: Mutations
[89282] in the V2 vasopressin receptor gene are associated with
X-linked nephrogenic
[89283] diabetes insipidus. Nature Genet. 2: 103–106, 1992.
- [89284] 15771.
- [89285] Pasel, K.; Schulz, A.; Timmermann, K.; Linnemann, K.;
Hoeltzenbein,
[89286] M.; Jaaskelainen, J.; Gruters, A.; Filler, G.; Schoneberg, T.:
Functional
[89287] characterization of the molecular defects causing nephro-
genic diabetes
[89288] insipidus in eight families. J. Clin. Endocr. Metab. 85:
1703–1710,
[89289] 2000.
- [89290] 15772.
- [89291] Rosenthal, W.; Seibold, A.; Antaramian, A.; Lonergan, M.;
Arthus,
[89292] M.-F.; Hendy, G. N.; Birnbaumer, M.; Bichet, D. G.: Molec-
ular identification

- [89293] of the gene responsible for congenital nephrogenic diabetes insipidus. *Nature* 359:
- [89294] 233–235, 1992.
- [89295] 15773.
- [89296] Sadeghi, H.; Robertson, G. L.; Bichet, D. G.; Innamorati, G.;
- [89297] Birnbaumer, M.: Biochemical basis of partial nephrogenic diabetes
- [89298] insipidus phenotypes. *Molec. Endocr.* 11: 1806–1813, 1997.
- [89299] 15774.
- [89300] Sadeghi, H. M.; Innamorati, G.; Birnbaumer, M.: An X-linked NDI
- [89301] mutation reveals a requirement for cell surface V2R expression. *Molec.*
- [89302] *Endocr.* 11: 706–713, 1997.
- [89303] 15775.
- [89304] Sawin, C. T.: Endocrine classic. *Endocrinologist* 8: 143–148,
- [89305] 1998.
- [89306] 15776.

- [89307] Schoneberg, T.; Sandig, V.; Wess, J.; Gudermann, T.; Schultz,
G.: Reconstitution of mutant V2 vasopressin receptors by
adenovirus-mediated
gene transfer: molecular basis and clinical implication. J.
Clin. Invest. 100: 1547-1556, 1997.
15777.
- [89312] Schoneberg, T.; Schulz, A.; Biebermann, H.; Gruters, A.;
Grimm,
T.; Hubschmann, K.; Filler, G.; Gudermann, T.; Schultz, G.:
V2 vasopressin
receptor dysfunction in nephrogenic diabetes insipidus
caused by different
molecular mechanisms. Hum. Mutat. 12: 196-205, 1998.
15778.
- [89317] Nathan, D. G.; Baehner, R. L.; Weaver, D. K.: Failure of ni-
tro
blue tetrazolium reduction in the phagocytic vacuoles of
leukocytes
in chronic granulomatous disease. J. Clin. Invest. 48:
1895-1904,

- [89320] 1969.
- [89321] 15779.
- [89322] Nelson, C. E.; Dahl, M. V.; Goltz, R. W.: Arcuate dermal erythema
- [89323] in a carrier of chronic granulomatous disease. Arch. Derm. 113:
- [89324] 789–800, 1977.
- [89325] 15780.
- [89326] Newburger, P. E.; Cohen, H. J.; Rothchild, S. B.; Hobbins, J.
- [89327] C.; Malawista, S. E.; Mahoney, M. J.: Prenatal diagnosis of chronic
- [89328] granulomatous disease. New Eng. J. Med. 300: 178–181, 1979.
- [89329] 15781.
- [89330] Noack, D.; Heyworth, P. G.; Kyono, W.; Cross, A. R.: A second
- [89331] case of somatic triple mosaicism in the CYBB gene causing chronic
- [89332] granulomatous disease. Hum. Genet. 109: 234–238, 2001.
- [89333] 15782.

- [89334] Okamura, N.; Malawista, S. E.; Roberts, R. L.; Rosen, H.; Ochs,
- [89335] H. D.; Babior, B. M.; Curnutte, J. T.: Phosphorylation of the oxidase-related
- [89336] 48K phosphoprotein family in the unusual cytochrome-negative and X-linked
- [89337] cytochrome-positive types of chronic granulomatous disease. Blood 72:
- [89338] 811-816, 1988.
- [89339] 15783.
- [89340] Orkin, S. H.: Stem cell alchemy. Nature Med. 6:
- [89341] 1212-1213,
- [89342] 2000.
- [89343] 15784.
- [89344] Orkin, S. H.: X-linked chronic granulomatous disease: from chromosomal
- [89345] position to the in vivo gene product. Trends Genet. 3:
- [89346] 149-151,
- [89347] 1987.
- [89348] 15785.
- [89349] Orkin, S. H.: Molecular genetics of chronic granulomatous

disease. Annu.

[89348] Rev. Immun. 7: 277–307, 1989.

[89349] 15786.

[89350] Patino, P. J.; Perez, J. E.; Lopez, J. A.; Condino-Neto, A.; Grumach,

[89351] A. S.; Botero, J. H.; Curnutte, J. T.; Garcia de Olarte, D.:
Molecular

[89352] analysis of chronic granulomatous disease caused by defects in gp91-phox. Hum.

[89353] Mutat. 13: 29–37, 1999.

[89354] 15787.

[89355] Pollock, J. D.; Williams, D. A.; Gifford, M. A. C.; Li, L. L.;

[89356] Du, X.; Fisherman, J.; Orkin, S. H.; Doerschuk, C. M.; Din-
auer, M.

[89357] C.: Mouse model of X-linked chronic granulomatous disease, an inherited

[89358] defect in phagocyte superoxide production. Nature Genet.
9: 202–209,

[89359] 1995.

[89360] 15788.

[89361] Porter, C. D.; Parkar, M. H.; Levinsky, R. J.; Collins, M. K.

[89362] L.; Kinnon, C.: X-linked chronic granulomatous disease: correction

[89363] of NADPH oxidase defect by retrovirus-mediated expression of gp91-phox. Blood 82:

[89364] 2196-2202, 1993.

[89365] 15789.

[89366] Quie, P. G.; White, J. G.; Holmes, B.; Good, R. A.: In vitro

[89367] bactericidal capacity of human polymorphonuclear leukocytes: diminished

[89368] activity in chronic granulomatous disease of childhood. J. Clin.

[89369] Invest. 46: 668-679, 1967.

[89370] 15790.

[89371] Rae, J.; Newburger, P. E.; Dinanuer, M. C.; Noack, D.; Hopkins,

[89372] P. J.; Kuruto, R.; Curnutte, J. T.: X-linked chronic granulomatous

[89373] disease: mutations in the CYBB gene encoding the gp91-phox component

[89374] of respiratory-burst oxidase. Am. J. Hum. Genet. 62: 1320-1331,

[89375] 1998.

[89376] 15791.

[89377] Roos, D.: The genetic basis of chronic granulomatous disease. Immun.

[89378] Rev. 138: 121–157, 1994.

[89379] 15792.

[89380] Roos, D.; de Boer, M.; Kuribayashi, F.; Meischl, C.; Weening,

[89381] R. S.; Segal, A. W.; Ahlin, A.; Nemet, K.; Hossle, J. P.; Bernatowska-Matuszkiewicz,

[89382] E.; Middleton-Price, H.: Mutations in the X-linked and autosomal

[89383] recessive forms of chronic granulomatous disease. Blood 87: 1663–1681,

[89384] 1996.

[89385] 15793.

[89386] Royer-Pokora, B.; Kunkel, L. M.; Monaco, A. P.; Goff, S. C.; Newburger,

[89387] P. E.; Baehner, R. L.; Cole, F. S.; Curnutte, J. T.; Orkin, S. H.

[89388] : Cloning the gene for an inherited human disorder—chronic granulomatous

- [89389] disease--on the basis of its chromosomal location. *Nature* 322: 32-38,
- [89390] 1986.
- [89391] 15794.
- [89392] Schmalzer, E. A.; Miller, D. R.: Chronic granulomatous disease. *Prog. Med. Genet.* 1: 145-184, 1976.
- [89393] 15795.
- [89394] Segal, A. W.: Cytochrome b(-245) and its involvement in the molecular
- [89395] pathology of chronic granulomatous disease. *Hemat. Oncol. Clin. North Am.* 2: 213-223, 1988.
- [89396] 100. Segal, A. W.; Cross, A. R.; Garcia, R. C.; Borregaard, N.; Valerius,
- [89397] N. H.; Soothill, J. F.; Jones, O. T. G.: Absence of cytochrome b(-245)
- [89398] in chronic granulomatous disease: a multicenter European evaluation
- [89399] of its incidence and relevance. *New Eng. J. Med.* 308:

245–251, 1983.

[89403]

[89404] 101. Segal, A. W.; Peters, T. J.: Characterisation of the enzyme

[89405] defect in chronic granulomatous disease. Lancet I: 1363–1365, 1976.

[89406]

[89407] 102. Segal, A. W.; Webster, D.; Jones, O. T. G.; Allison, A. C.:

[89408] Absence of a newly described cytochrome b from neutrophils of patients

[89409] with chronic granulomatous disease. Lancet II: 446–449, 1978.

[89410]

[89411] 103. Soothill, J. F.: Inheritance of chronic granulomatous disease.

[89412] (Letter) Lancet II: 543–544, 1969.

[89413]

[89414] 104. Southwick, F. S.; van der Meer, J. W. M.: Recurrent cystitis

[89415] and bladder mass in two adults with chronic granuloma-

tous disease. Ann.

[89416] Intern. Med. 109: 118–121, 1988.

[89417]

[89418] 105. Stasia, M. J.; Lardy, B.; Maturana, A.; Rousseau, P.;
Martel,

[89419] C.; Bordigoni, P.; Demaurex, N.; Morel, F.: Molecular and
functional

[89420] characterization of a new X-linked chronic granulomatous
disease variant

[89421] (X91+) case with a double missense mutation in the cy-
tosolic gp91phox

[89422] C-terminal tail. Biochim. Biophys. Acta 1586: 316–330,
2002.

[89423]

[89424] 106. Taswell, H. F.; Lewis, J. C.; Marsh, W. L.; Wimer, B. M.;
Pineda,

[89425] A. A.; Brzica, S. M., Jr.: Erythrocyte morphology in genetic
defects

[89426] of the Rh and Kell blood group systems. Mayo Clin. Proc.
52: 157–159,

[89427] 1977.

[89428]

- [89429] 107. Tauber, A. I.; Borregaard, N.; Simons, E.; Wright, J.:
Chronic
[89430] granulomatous disease: a syndrome of phagocyte oxidase
deficiencies. Medicine 62:
[89431] 286–309, 1983.
[89432]
[89433] 108. Teahan, C.; Rowe, P.; Parker, P.; Totty, N.; Segal, A.
W.: The
[89434] X-linked chronic granulomatous disease gene codes for
the beta-chain
[89435] of cytochrome b(–245). Nature 327: 720–721, 1987.
[89436]
[89437] 109. Thompson, E. N.; Chandra, R. K.; Cope, W. A.;
Soothill, J. F.
[89438] : Leukocyte abnormality in both parents of a patient with
chronic
[89439] granulomatous disease. Lancet I: 799–800, 1969.
[89440]
[89441] 110. Thompson, E. N.; Soothill, J. F.: Chronic granuloma-
tous disease:
[89442] quantitative clinicopathological relationships. Arch. Dis.

Child. 45:

[89443] 24–32, 1970.

[89444]

[89445] 111. Tsuda, M.; Kaneda, M.; Sakiyama, T.; Inana, I.;
Owada, M.; Kiryu,

[89446] C.; Shiraishi, T.; Kakinuma, K.: A novel mutation at a
probable heme-binding

[89447] ligand in neutrophil cytochrome b(558) in atypical X-
linked chronic

[89448] granulomatous disease. Hum. Genet. 103: 377–381,
1998.

[89449]

[89450] 112. Windhorst, D. B.: Inheritance of chronic granuloma-
tous disease.

[89451] (Letter) Lancet II: 543–544, 1969.

[89452]

[89453] 113. Windhorst, D. B.; Holmes, B.; Good, R. A.: A newly
defined X-linked

[89454] trait in man with demonstration of the Lyon effect in car-
rier females. Lancet I:

[89455] 737–739, 1967.

[89456]

[89457] 114. Wolff, G.; Muller, C. R.; Jobke, A.: Linkage of genes
for chronic

[89458] granulomatous disease and Xg. Hum. Genet. 54: 269–271,
1980.

[89459] 15796.

[89460] Yen, P. H.; Ellison, J.; Salido, E. C.; Mohandas, T.; Shapiro,

[89461] L.: Isolation of a new gene from the distal short arm of the
human

[89462] X chromosome that escapes X-inactivation. Hum. Molec.
Genet. 1:

[89463] 47–52, 1992.

[89464] 15797.

[89465] Abrams, E. S.; Murdaugh, S. E.; Lerman, L. S.: Comprehen-
sive detection

[89466] of single base changes in human genomic DNA using de-
naturing gradient

[89467] gel electrophoresis and a GC clamp. Genomics 7:
463–475, 1990.

[89468] 15798.

[89469] Acquila, M.; Caprino, D.; Pecorara, M.; Baudo, F.; Morfini,

M.;

[89470] Mori, P. G.: Two novel mutations at 373 codon of FVIII gene detected

[89471] by DGGE. Thromb. Haemost. 69: 392–393, 1993.

[89472] 15799.

[89473] Alexander, B.; Goldstein, R.: Dual hemostatic defect in pseudohemophilia.

[89474] (Abstract) J. Clin. Invest. 32: 551 only, 1953.

[89475] 15800.

[89476] Aly, A. M.; Arai, M.; Hoyer, L. W.: Cysteamine enhances the procoagulant

[89477] activity of factor VIII–East Hartford, a dysfunctional protein due

[89478] to a light chain thrombin cleavage site mutation (arginine–1689 to

[89479] cysteine). J. Clin. Invest. 89: 1375–1381, 1992.

[89480] 15801.

[89481] Muller, C. R.; Migl, B.; Ropers, H.–H.; Happle, R.: Heterozygote

[89482] detection in steroid sulphatase deficiency. (Letter) Lancet I: 546–547,

- [89483] 1980.
- [89484] 15802.
- [89485] Muller, C. R.; Wahlstrom, J.; Ropers, H.-H.: Further evidence
- [89486] for the assignment of the steroid sulfatase X-linked ichthyosis locus
- [89487] to the telomer of Xp. (Letter) Hum. Genet. 58: 446, 1981.
- [89488] 15803.
- [89489] Munke, M.; Kruse, K.; Goos, M.; Ropers, H. H.; Tolksdorf, M.:
- [89490] Genetic heterogeneity of the ichthyosis, hypogonadism, mental retardation,
- [89491] and epilepsy syndrome: clinical and biochemical investigations on
- [89492] two patients with Rud syndrome and review of the literature. Europ.
- [89493] J. Pediat. 141: 8-13, 1983.
- [89494] 15804.
- [89495] Nomura, K.; Nakano, H.; Umeki, K.; Harada, K.; Kon, A.; Tamai,
- [89496] K.; Sawamura, D.; Hashimoto, I.: A study of the steroid

sulfatase

[89497] gene in families with X-linked ichthyosis using polymerase chain reaction. *Acta*

[89498] *Derm. Venereol.* 75: 340-342, 1995.

[89499] 15805.

[89500] Okano, M.; Kitano, Y.; Nakamura, T.; Matsuzawa, Y.: Detection

[89501] of heterozygotes of X-linked ichthyosis by measuring steroid sulphatase

[89502] activity of lymphocytes: mode of inheritance in three families. *Brit.*

[89503] *J. Derm.* 113: 645-649, 1985.

[89504] 15806.

[89505] Orel, H.: Die Vererbung der Ichthyosis congenita und der Ichthyosis

[89506] vulgaris. *Z. Kinderheilk.* 47: 312-340, 1929.

[89507] 15807.

[89508] Passarge, E.; Post, B.; Schopf, E.: Possible genetic heterogeneity

[89509] of X-linked ichthyosis. *Birth Defects Orig. Art. Ser.* VII(8): 46-49,

[89510] 1971.

[89511] 15808.

[89512] Pike, M. G.; Hammerton, M.; Edge, J.; Atherton, D. J.;
Grant,

[89513] D. B.: A family with X-linked ichthyosis and hypogonadism. Europ.

[89514] J. Pediat. 148: 442-444, 1989.

[89515] 15809.

[89516] Ropers, H.-H.; Migl, B.; Zimmer, J.; Fraccaro, M.; Maraschio,

[89517] P. P.; Westerveld, A.: Activity of steroid sulfatase in fibroblasts

[89518] with numerical and structural X chromosome aberrations. Hum. Genet. 57:

[89519] 354-356, 1981.

[89520] 15810.

[89521] Ropers, H.-H.; Wiberg, U.: Evidence for X-linkage and non-inactivation

[89522] of steroid sulphatase locus in wood lemming. Nature 296: 766-767,

[89523] 1982.

[89524] 15811.

[89525] Ross, J. B.; Allderdice, P. W.; Shapiro, L. J.; Aveling, J.;
Eales,

[89526] B. A.; Simms, D., Jr.: Familial X-linked ichthyosis, steroid
sulfatase

[89527] deficiency, mental retardation, and nullisomy for
Xp223-pter. Arch.

[89528] Derm. 121: 1524-1528, 1985.

[89529] 15812.

[89530] Salido, E. C.; Li, X. M.; Yen, P. H.; Martin, N.; Mohandas, T.

[89531] K.; Shapiro, L. J.: Cloning and expression of the mouse
pseudoautosomal

[89532] steroid sulphatase gene (Sts). Nature Genet. 13: 83-86,
1996.

[89533] 15813.

[89534] Schlamadinger, J.; Meyer, J. C.; Vajda, I.; Szabo, G.: X-
linked

[89535] recessive ichthyosis: reinvestigation of a family first de-
scribed

[89536] in 1928. Dermatologica 175: 217-223, 1987.

[89537] 15814.

[89538] Schmickel, R. D.: Chromosomal deletions and enzyme deficiencies. J.

[89539] *Pediat.* 108: 244–246, 1986.

[89540]

[89541] 100. Schnyder, U. W.: Inherited ichthyoses. *Arch. Derm.* 102: 240–252,

[89542] 1970.

[89543]

[89544] 101. Sever, R. J.; Frost, P.; Weinstein, G.: Eye changes in ichthyosis. *J.A.M.A.* 206:

[89545] 2283–2286, 1968.

[89546]

[89547] 102. Shapiro, L. J.: Personal Communication. Los Angeles, Calif.

[89548] 3/1987.

[89549]

[89550] 103. Shapiro, L. J.: Personal Communication. San Francisco, Calif.

[89551] 5/20/1997.

[89552]

[89553] 104. Shapiro, L. J.: Personal Communication. Torrance, Calif.

[89554] 1977.

[89555]

[89556] 105. Shapiro, L. J.; Mohandas, T.; Weiss, R.; Romeo, G.:
Non-activation

[89557] of a X-chromosome locus in man. Science 204:
1224-1226, 1979.

[89558]

[89559] 106. Shapiro, L. J.; Weiss, R.: Diminished cholesterol sul-
fatase

[89560] activity in fibroblasts of placental sulfatase deficiency pa-
tients.

[89561] (Abstract) Vth Int. Conf. on Birth Defects, Montreal ,
8/1977.

[89562]

[89563] 107. Shapiro, L. J.; Weiss, R.; Buxman, M. M.; Vidgoff, J.;
Dimond,

[89564] R. L.: Enzymatic basis of typical X-linked ichthyosis.
Lancet II:

[89565] 756-757, 1978.

[89566]

[89567] 108. Shapiro, L. J.; Weiss, R.; Webster, D.; France, J. T.: X-linked

[89568] ichthyosis due to steroid-sulphatase deficiency. Lancet I: 70-72,

[89569] 1978.

[89570]

[89571] 109. Shapiro, L. J.; Yen, P.; Pomerantz, D.; Martin, E.; Rolewic,

[89572] L.; Mohandas, T.: Molecular studies of deletions at the human steroid

[89573] sulfatase locus. Proc. Nat. Acad. Sci. 86: 8477-8481, 1989.

[89574]

[89575] 110. Shapiro, L. J.; Yen, P. H.: Reply to letter of Andria et al.,

[89576] 1987. (Letter) Ann. Neurol. 22: 99, 1987.

[89577]

[89578] 111. Shapiro, L. J.; Yen, P. H.; Marsh, B.; Mohandas, T.: Frequent

[89579] deletions at the steroid sulfatase (STS) locus. (Abstract)

Am. J.

[89580] Hum. Genet. 41: A238, 1987.

[89581]

[89582] 112. Solomon, I. L.; Schoen, E. J.: Sex-linked ichthyosis in XO gonadal

[89583] dysgenesis. (Letter) Lancet I: 1304-1305, 1971.

[89584]

[89585] 113. Stein, C.; Hille, A.; Seidel, J.; Rijnbout, S.; Waheed, A.; Schmidt,

[89586] B.; Geuze, H.; von Figura, K.: Cloning and expression of human steroid-sulfatase:

[89587] membrane topology, glycosylation, and subcellular distribution in

[89588] BHK-21 cells. J. Biol. Chem. 264: 13865-13872, 1989.

[89589]

[89590] 114. Steinmann, B.; Mieth, D.; Gitzelmann, R.: A newly recognized

[89591] cause of low urinary estriol in pregnancy: multiple sulfatase deficiency

[89592] of the fetus. Gynec. Obstet. Invest. 12: 107-109, 1981.

[89593]

- [89594] 115. Stoll, C.; Grosshans, E.; Binder, P.; Roth, M.: Hyper-trophic
- [89595] pyloric stenosis associated with X-linked ichthyosis in two brothers. Clin.
- [89596] Exp. Derm. 8: 61–64, 1983.
- [89597]
- [89598] 116. Sunohara, N.; Sakuragawa, N.; Satoyoshi, E.; Tanae, A.; Shapiro,
- [89599] L. J.: A new syndrome of anosmia, ichthyosis, hypogonadism, and various
- [89600] neurological manifestations with deficiency of steroid sulphatase
- [89601] and arylsulphatase C. Ann. Neurol. 19: 174–181, 1986.
- [89602]
- [89603] 117. Tabei, F.; Heinrichs, W. L.: Diagnosis of placental sulphatase
- [89604] deficiency. Am. J. Obstet. Gynec. 124: 409–414, 1976.
- [89605]
- [89606] 118. Tiepolo, L.; Zuffardi, O.; Fraccaro, M.; di Natale, D.; Gargantini,
- [89607] L.; Muller, C. R.; Ropers, H.–H.: Assignment by deletion

mapping

[89608] of the steroid sulfatase X-linked ichthyosis locus to
Xp223. Hum.

[89609] Genet. 54: 205–206, 1980.

[89610]

[89611] 119. Toder, R.; Rappold, G. A.; Schiebel, K.; Schempp, W.:
ANT3 and

[89612] STS are autosomal in prosimian lemurs: implications for
the evolution

[89613] of the pseudoautosomal region. Hum. Genet. 95: 22–28,
1995.

[89614]

[89615] 120. Traupe, H.; Happle, R.: Clinical spectrum of steroid
sulfatase

[89616] deficiency: X-linked recessive ichthyosis, birth complica-
tions, and

[89617] cryptorchidism. Europ. J. Pediat. 140: 19–21, 1983.

[89618]

[89619] 121. Traupe, H.; Muller-Migl, C. R.; Kolde, G.; Happle, R.;
Kovary,

[89620] P. M.; Hameister, H.; Ropers, H. H.: Ichthyosis vulgaris

with hypogenitalism

[89621] and hypogonadism: evidence for different genotypes by lipoprotein

[89622] electrophoresis and steroid sulfatase testing. Clin. Genet. 25:

[89623] 42-51, 1984.

[89624]

[89625] 122. Traupe, H.; Ropers, H.-H.: Cryptorchidism and hypogenitalism

[89626] in X-linked recessive ichthyosis vulgaris. (Letter) Hum. Genet. 60:

[89627] 206, 1982.

[89628]

[89629] 123. Turpin, R.; Desvignes, P.; Demassieux, J.-L.: Sur une variete

[89630] d'ichthyose hereditaire avec alteration du fond d'oeil. Nouveau syndrome

[89631] ectoblastique?. Sem. Hop. Paris 21: 343, 1945.

[89632]

[89633] 124. Valdes-Flores, M.; Kofman-Alfaro, S. H.; Jimenez-Vaca, A. L.;

- [89634] Cuevas–Covarrubias, S. A.: Carrier identification by FISH analysis
- [89635] in isolated cases of X-linked ichthyosis. Am. J. Med. Genet. 102:
- [89636] 146–148, 2001.
- [89637]
- [89638] 125. Valdes–Flores, M.; Kofman–Alfaro, S. H.; Jimenez Vaca, A. L.;
- [89639] Cuevas–Covarrubias, S. A.: A novel partial deletion of exons 2–10
- [89640] of the STS gene in recessive X-linked ichthyosis. J. Invest. Derm. 114:
- [89641] 591–593, 2000.
- [89642]
- [89643] 126. Vogel, W.; Grompe, M.; Storz, R.; Pentz, S.: A comparative study
- [89644] on steroid sulfatase and arylsulfatase C in fibroblast clones from
- [89645] 45,X/47,XXX and 69,XXY. Hum. Genet. 66: 367–369, 1984.
- [89646]

[89647] 127. Wells, R. S.; Jennings, M. C.: X-linked ichthyosis and ichthyosis

[89648] vulgaris: clinical and genetic distinctions in a second series of

[89649] families. J.A.M.A. 202: 485–488, 1967.

[89650]

[89651] 128. Went, L. N.; DeGroot, W. P.; Sanger, R.; Tippet, P.; Gavin,

[89652] J.: X-linked ichthyosis: linkage relationship with the Xg blood groups

[89653] and other studies in a large Dutch kindred. Ann. Hum. Genet. 32:

[89654] 333–346, 1969.

[89655]

[89656] 129. Wieacker, P.; Davies, K. E.; Mevorah, B.; Ropers, H. H.: Linkage

[89657] studies in a family with X-linked recessive ichthyosis employing a

[89658] cloned DNA sequence from the distal short arm of the X chromosome. Hum.

[89659] Genet. 63: 113–116, 1983.

[89660]

[89661] 130. Willard, H. F.; Holmes, M. T.: A sensitive and dependable assay

[89662] for distinguishing hamster and human X-linked steroid sulfatase activity

[89663] in somatic cell hybrids. Hum. Genet. 66: 272-275, 1984.

[89664]

[89665] 131. Wirth, B.; Herrmann, F. H.; Neugebauer, M.; Gillard, E. F.; Wulff,

[89666] K.; Stein, C.; von Figura, K.; Ferguson-Smith, M. A.; Gal, A.: Linkage

[89667] analysis in X-linked ichthyosis (steroid sulfatase deficiency). Hum.

[89668] Genet. 80: 191-192, 1988.

[89669]

[89670] 132. Yen, P. H.; Allen, E.; Marsh, B.; Mohandas, T.; Wang, N.; Taggart,

[89671] R. T.; Shapiro, L. J.: Cloning and expression of steroid sulfatase

[89672] cDNA and the frequent occurrence of deletions in STS deficiency: implications

[89673] for X-Y interchange. Cell 49: 443-454, 1987.

[89674]

[89675] 133. Yen, P. H.; Marsh, B.; Allen, E.; Tsai, S. P.; Ellison, J.; Connolly,

[89676] L.; Neiswanger, K.; Shapiro, L. J.: The human X-linked steroid sulfatase

[89677] gene and a Y-encoded pseudogene: evidence for an inversion of the

[89678] Y chromosome during primate evolution. Cell 55: 1123-1135, 1988.

[89679]

[89680] 134. Zettersten, E.; Man, M.-Q.; Sato, J.; Denda, M.; Farrell, A.;

[89681] Ghadially, R.; Williams, M. L.; Feingold, K. R.; Elias, P. M. Recessive

[89682] X-linked ichthyosis: role of cholesterol-sulfate accumulation in the

[89683] barrier abnormality. J. Invest. Derm. 111: 784-790, 1998.

[89684] 15815.

[89685] Holinski-Feder, E.; Chahrokh-Zadeh, S.; Jedelev, K. B.; Meindl,

[89686] A.; Steinbach, P.; Wohrle, D.: FRAXE testing. (Letter) Am. J. Hum.

[89687] Genet. 59: 1168–1169, 1996.

- [89688] 15816.
- [89689] Knight, S. J. L.; Flannery, A. V.; Hirst, M. C.; Campbell, L.;
- [89690] Christodoulou, Z.; Phelps, S. R.; Middleton-Price, H. R.;
- Bamicoat,
- [89691] A.; Pembrey, M. E.; Holland, J.; Bobrow, M.; Oostra, B. A.;
- Davies,
- [89692] K. E.: FRAXE mental retardation associated with trinucleotide repeat
- [89693] expansion and hypermethylation of a CpG island in Xq28. (Abstract) Am.
- [89694] J. Hum. Genet. 53 (suppl.): A79, 1993.
- [89695] 15817.
- [89696] Knight, S. J. L.; Flannery, A. V.; Hirst, M. C.; Campbell, L.;
- [89697] Christodoulou, Z.; Phelps, S. R.; Pointon, J.; Middleton-Price, H.
- [89698] R.; Barnicoat, A.; Pembrey, M. E.; Holland, J.; Oostra, B. A.;
- Bobrow,
- [89699] M.; Davies, K. E.: Trinucleotide repeat amplification and hypermethylation
- [89700] of a CpG island in FRAXE mental retardation. Cell 74: 127-134, 1993.

- [89701] 15818.
- [89702] Knight, S. J. L.; Ritchie, R. J.; Chakrabarti, L.; Cross, G.;
- [89703] Taylor, G. R.; Mueller, R. F.; Hurst, J.; Paterson, J.; Yates, J.
- [89704] R. W.; Dow, D. J.; Davies, K. E.: A study of FRAXE in men-
tally retarded
- [89705] individuals referred for fragile X syndrome (FRAXA) testing
in the
- [89706] United Kingdom. Am. J. Hum. Genet. 58: 906–913, 1996.
- [89707] 15819.
- [89708] Knight, S. J. L.; Voelckel, M. A.; Hirst, M. C.; Flannery, A.
- [89709] V.; Moncla, A.; Davies, K. E.: Triplet repeat expansion at
the FRAXE
- [89710] locus and X-linked mild mental handicap. Am. J. Hum.
Genet. 55:
- [89711] 81–86, 1994.
- [89712] 15820.
- [89713] Mila, M.; Sanchez, A.; Badenas, C.; Brun, C.; Jimenez, D.;
- Villa,
- [89714] M. P.; Castellvi-Bel, S.; Estivill, X.: Screening for FMR1 and
FMR2

- [89715] mutations in 222 individuals from Spanish special schools: identification
- [89716] of a case of FRAXE-associated mental retardation. Hum. Genet. 100:
- [89717] 503–507, 1997.
- [89718] 15821.
- [89719] Mulley, J. C.; Yu, S.; Loesch, D. Z.; Hay, D. A.; Donnelly, A.;
- [89720] Gedeon, A. K.; Carbonell, P.; Lopez, I.; Glover, G.; Gabar-
ron, I.;
- [89721] Yu, P. W. L.; Baker, E.; Haan, E. A.; Hockey, A.; Knight, S. J.
L.;
- [89722] Davies, K. E.; Richards, R. I.; Sutherland, G. R.: FRAXE and
mental
- [89723] retardation. J. Med. Genet. 32: 162–169, 1995.
- [89724] 15822.
- [89725] Murray, A.; Webb, J.; Dennis, N.; Conway, G.; Morton, N.:
Microdeletions
- [89726] in FMR2 may be a significant cause of premature ovarian
failure. J.
- [89727] Med. Genet. 36: 767–770, 1999.
- [89728] 15823.

- [89729] Russo, S.; Selicorni, A.; Bedeschi, M. F.; Natacci, F.; Viziello, P.; Fortuna, R.; Pagani, G.; Dalpra, L.; Larizza, L.: Molecular characterization of FRAXE-positive subjects with mental impairment in two unrelated Italian families. Am. J. Med. Genet. 75: 304–308, 1998.
- [89732] 15824.
- [89733] Sutherland, G. R.: Personal Communication. Adelaide, Australia 11/17/1993.
- [89734] 15825.
- [89735] Sutherland, G. R.; Baker, E.: Characterisation of a new rare fragile site easily confused with the fragile X. Hum. Molec. Genet. 1: 111–113, 1992.
- [89736] 15826.
- [89737] Abitbol, M.; Menini, C.; Delezoide, A.–L.; Rhyner, T.; Veke-mans, M.; Mallet, J.: Nucleus basalis magnocellularis and hippocampus are

- [89743] the major sites of FMR-1 expression in the human fetal brain. *Nature*
- [89744] *Genet.* 4: 147-153, 1993.
- [89745] 15827.
- [89746] Abrams, M. T.; Kaufmann, W. E.; Rousseau, F.; Oostra, B. A.; Wolozin,
- [89747] B.; Taylor, C. V.; Lishaa, N.; Morel, M.-L.; Hoogeveen, A.; Reiss,
- [89748] A. L.: FMR1 gene expression in olfactory neuroblasts from two males
- [89749] with fragile X syndrome. *Am. J. Med. Genet.* 82: 25-30, 1999.
- [89750] 15828.
- [89751] Allingham-Hawkins, D. J.; Babul-Hirji, R.; Chitayat, D.; Holden,
- [89752] J. J. A.; Yang, K. T.; Lee, C.; Hudson, R.; Gorwill, H.; Nolin, S.
- [89753] L.; Glicksman, A.; Jenkins, E. C.; Brown, W. T.; and 27 others:
- [89754] Fragile X premutation is a significant risk factor for premature ovarian
- [89755] failure: the international collaborative POF in fragile X

study--preliminary

[89756] data. Am. J. Med. Genet. 83: 322-325, 1999.

[89757] 15829.

[89758] Ashley, C. T.; Sutcliffe, J. S.; Kunst, C. B.; Leiner, H. A.;
Eichler,

[89759] E. E.; Nelson, D. L.; Warren, S. T.: Human and murine FMR-
1: alternative

[89760] splicing and translational initiation downstream of the
CGG-repeat. Nature

[89761] Genet. 4: 244-251, 1993.

[89762] 15830.

[89763] Ashley, C. T., Jr.; Wilkinson, K. D.; Reines, D.; Warren, S. T.

[89764] : FMR1 protein: conserved RNP family domains and selec-
tive RNA binding. Science 262:

[89765] 563-566, 1993.

[89766] 15831.

[89767] Bachner, D.; Steinbach, P.; Wohrle, D.; Just, W.; Vogel, W.;
Hameister,

[89768] H.; Manca, A.; Poustka, A.: Enhanced Fmr-1 expression in
testis.

[89769] (Letter) Nature Genet. 4: 115-116, 1993.

[89770] 15832.

[89771] Backes, M.; Genc, B.; Schreck, J.; Doerfler, W.; Lehmkuhl, G.;

[89772] von Gontard, A.: Cognitive and behavioral profile of fragile X boys:

[89773] correlations to molecular data. Am. J. Med. Genet. 95: 150–156,

[89774] 2000.

[89775] 15833.

[89776] Maglott, D. R.; Durkin, A. S.; Lane, S. A.; Callen, D. F.; Feldblyum,

[89777] T. V.; Nierman, W. C.: The gene for membrane protein E16 (D16S469E)

[89778] maps to human chromosome 16q24.3 and is expressed in human brain,

[89779] thymus, and retina. Genomics 23: 303–304, 1994.

[89780] 15834.

[89781] Kastury, K.; Druck, T.; Huebner, K.; Barletta, C.; Acampora, D.;

[89782] Simeone, A.; Faiella, A.; Boncinelli, E.: Chromosome locations of

- [89783] human EMX and OTX genes. Genomics 22: 41–45, 1994.
- [89784] 15835.
- [89785] Simeone, A.; Acampora, D.; Gulisano, M.; Stornaiuolo, A.; Boncinelli, E.: Nested expression domains of four homeobox genes in developing rostral brain. Nature 358: 687–690, 1992.
- [89787] 15836.
- [89788] Bishop, K. M.; Goudreau, G.; O'Leary, D. D. M.: Regulation of area identity in the mammalian neocortex by Emx2 and Pax6. Science 288: 344–349, 2000.
- [89789] 15837.
- [89790] Boncinelli, E.; Gulisano, M.; Spada, F.; Broccoli, V.: Emx and Otx gene expression in the developing mouse brain. Ciba Found. Symp. 193: 100–116, 1995.
- [89791] 15838.

- [89797] Bosetti, A.; Faiella, A.; Boncinelli, E.; Consalez, G. G.: Linkage
- [89798] mapping of Emx2 to mouse chromosome 19. *Mammalian Genome* 8: 71-72,
- [89799] 1997.
- [89800] 15839.
- [89801] Noonan, F. C.; Mutch, D. G.; Mallon, M. A.; Goodfellow, P. J.:
- [89802] Characterization of the homeodomain gene EMX2: sequence conservation,
- [89803] expression analysis, and a search for mutations in endometrial cancers. *Genomics* 76:
- [89804] 37-44, 2001.
- [89805] 15840.
- [89806] Acampora, D.; Mazan, S.; Avantaggiato, V.; Barone, P.; Tuorto,
- [89807] F.; Lallemand, Y.; Brulet, P.; Simeone, A.: Epilepsy and brain abnormalities
- [89808] in mice lacking the Otx1 gene. *Nature Genet.* 14: 218-222, 1996.
- [89809] 15841.

- [89810] Boncinelli, E.; Gulisano, M.; Broccoli, V.: Emx and Otx homeobox
genes in the developing mouse brain. J. Neurobiol. 24:
1356–1366,
1993.
15842.
- [89814] Frantz, G. D.; Weimann, J. M.; Levin, M. E.; McConnell, S.
K.:
Otx1 and Otx2 define layers and regions in developing
cerebral cortex
and cerebellum. J. Neurosci. 14: 5725–5740, 1994.
15843.
- [89818] Avraham, S.; Jiang, S.; Ota, S.; Fu, Y.; Deng, B.; Dowler, L.
L.;
White, R. A.; Avraham, H.: Structural and functional studies of the
intracellular tyrosine kinase MATK gene and its translated
product. J.
Biol. Chem. 270: 1833–1842, 1995.
15844.
- [89823] Bennett, B. D.; Cowley, S.; Jiang, S.; London, R.; Deng, B.;

Grabarek,

[89824] J.; Groopman, J. E.; Goeddel, D. V.; Avraham, H.: Identification

[89825] and characterization of a novel tyrosine kinase from megakaryocytes. J.

[89826] Biol. Chem. 269: 1068–1074, 1994.

[89827] 15845.

[89828] Klages, S.; Adam, D.; Class, K.; Fargnoli, J.; Bolen, J. B.; Penhallow,

[89829] R. C.: Ctk: a protein–tyrosine kinase related to Csk that defines

[89830] an enzyme family. Proc. Nat. Acad. Sci. 91: 2597–2601, 1994.

[89831] 15846.

[89832] Sakano, S.; Iwama, A.; Inazawa, J.; Ariyama, T.; Ohno, M.; Suda,

[89833] T.: Molecular cloning of a novel non–receptor tyrosine kinase, HYL

[89834] (hematopoietic consensus tyrosine–lacking kinase). Oncogene 9: 1155–1161,

[89835] 1994.

[89836] 15847.

- [89837] Zrihan-Licht, S.; Lim, J.; Keydar, I.; Sliwkowski, M. X.; Groopman, J. E.; Avraham, H.: Association of Csk-homologous kinase (CHK) (formerly MATK) with HER-2/ErbB-2 in breast cancer cells. J. Biol. Chem. 272: 1856-1863, 1997.
- [89840] 15848.
- [89841] Shimizu, S.; Narita, M.; Tsujimoto, Y.: Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC. Nature 399: 483-487, 1999.
- [89844] 15849.
- [89845] Nichols, A. F.: Personal Communication. Berkeley, Calif. 10/4/1995.
- [89846] 15850.
- [89847] Campbell, H. D.; Schimansky, T.; Claudianos, C.; Ozsarac, N.; Kasprzak, A. B.; Cotsell, J. N.; Young, I. G.; de Couet, H. G.; Gabor Miklos,

- [89850] G. L.: The *Drosophila melanogaster* flightless-I gene involved in
- [89851] gastrulation and muscle degeneration encodes gelsolin-like and leucine-rich
- [89852] repeat domains and is conserved in *Caenorhabditis elegans* and humans. *Proc.*
- [89853] *Nat. Acad. Sci.* 90: 11386–11390, 1993.
- [89854] 15851.
- [89855] Harmon, D. L.; Gardner-Medwin, D.; Stirling, J. L.: Two new mutations
- [89856] in a late infantile Tay-Sachs patient are both in exon 1 of the beta-hexosaminidase
- [89857] alpha subunit gene. *J. Med. Genet.* 30: 123–128, 1993.
- [89858] 15852.
- [89859] Buckle, V.; Mondello, C.; Darling, S.; Craig, I. W.; Goodfellow,
- [89860] P. N.: Homologous expressed genes in the human sex chromosome pairing
- [89861] region. *Nature* 317: 739–741, 1985.
- [89862] 15853.
- [89863] Slaugenhaupt, S. A.; Roca, A. L.; Liebert, C. B.; Altherr, M.

R.;

[89864] Gusella, J. F.; Reppert, S. M.: Mapping of the gene for the Mel1a-melatonin

[89865] receptor to human chromosome 4 (MTNR1A) and mouse chromosome 8 (Mtnr1a). Genomics 27:

[89866] 355-357, 1995.

[89867] 15854.

[89868] Hulsebos, T. J. M.; Cerosaletti, K. M.; Fournier, R. E. K.; Sinke,

[89869] R. J.; Rocchi, M.; Marzella, R.; Jenkins, N. A.; Gilbert, D. J.; Copeland,

[89870] N. G.: Identification of the human beta-A2 crystallin gene (CRYBA2):

[89871] localization of the gene on human chromosome 2 and of the homologous

[89872] gene on mouse chromosome 1. Genomics 28: 543-548, 1995.

[89873] 15855.

[89874] Grimsby, J.; Toth, M.; Chen, K.; Kumazawa, T.; Klaidman, L.; Adams,

[89875] J. D.; Karoum, F.; Gal, J.; Shih, J. C.: Increased stress response

- [89876] and beta-phenylethylamine in MAOB-deficient mice. *Nature Genet.* 17:
[89877] 206-210, 1997.
- [89878] 15856.
- [89879] Lenders, J. W. M.; Eisenhofer, G.; Abeling, N. G. G. M.;
Berger,
[89880] W.; Murphy, D. L.; Konings, C. H.; Bleeker Wagemakers, L.
M.; Kopin,
[89881] I. J.; Karoum, F.; van Gennip, A. H.; Brunner, H. G.: Specific
genetic
[89882] deficiencies of the A and B isoenzymes of monoamine ox-
idase are characterized
[89883] by distinct neurochemical and clinical phenotypes. *J. Clin.*
Invest. 97:
[89884] 1010-1019, 1996.
- [89885] 15857.
- [89886] Rice, J.; McGuffin, P.; Goldin, L. R.; Shaskan, E. G.; Ger-
shon,
[89887] E. S.: Platelet monoamine oxidase (MAO) activity: evidence
for a
[89888] single major locus. *Am. J. Hum. Genet.* 36: 36-43, 1984.

[89889] 15858.

[89890] Weinshilboum, R. M.: Biochemical genetics of catecholamines in

[89891] humans. Mayo Clin. Proc. 58: 319–330, 1983.

[89892] 15859.

[89893] Wyatt, R. J.; Murphy, D. L.; Belmaker, R.; Cohen, S.; Donnelly,

[89894] C. H.; Pollin, W.: Reduced monoamine oxidase activity in platelets:

[89895] a possible genetic marker for vulnerability to schizophrenia. Science 179:

[89896] 916–918, 1973.

[89897] 15860.

[89898] Hattori, Y.; Yamashiro, Y.; Ohba, Y.; Miyaji, T.; Morishita, M.;

[89899] Yamamoto, K.; Yamamoto, K.; Narai, S.; Kimura, A.: A new beta-thalassemia

[89900] mutation (initiation codon ATG-to-GTG) found in the Japanese population. Hemoglobin 15:

[89901] 317–325, 1991.

[89902] 15861.

- [89903] Holmes, L. B.: Norrie's disease--an X-linked syndrome of retinal
malformation, mental retardation and deafness. New Eng.
J. Med. 284:
367-368, 1971.
- [89906] 15862.
- [89907] Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Doi, N.;
Nakagawa,
M.; Ozawa, M.; Kuroda, N.: Novel mutation at the initiation
codon
in the Norrie disease gene in two Japanese families. Hum.
Genet. 95:
105-108, 1995.
- [89911] 15863.
- [89912] Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Hotta, Y.;
Hayakawa,
M.; Fujiki, K.; Tanabe, U.: Mutations in the Norrie disease
gene:
a new mutation in a Japanese family. (Letter) Brit. J. Oph-
thal. 79:
703-708, 1995.

- [89916] 15864.
- [89917] Johnson, K.; Mintz-Hittner, H. A.; Conley, Y. P.; Ferrell, R.
- [89918] E.: X-linked exudative vitreoretinopathy caused by an
arginine to
- [89919] leucine substitution (R121L) in the Norrie disease protein.
Clin.
- [89920] Genet. 50: 113–115, 1996.
- [89921] 15865.
- [89922] Johnston, S. S.; Hanna, J. E.; Nevin, N. C.; Bryars, J. H.:
Norrie's
- [89923] disease. Birth Defects Orig. Art. Ser. 18(6): 729–738,
1982.
- [89924] 15866.
- [89925] Katayama, S.; Wohlferd, M.; Golbus, M. S.: First demon-
stration
- [89926] of recombination between the gene for Norrie disease and
probe L1.28. Am.
- [89927] J. Med. Genet. 30: 967–970, 1988.
- [89928] 15867.
- [89929] Kivlin, J. D.; Sanborn, G. E.; Wright, E.; Cannon, L.; Carey,

- [89930] J.: Further linkage data on Norrie disease. *Am. J. Med. Genet.* 26:
- [89931] 733–736, 1987.
- [89932] 15868.
- [89933] Lindsay, S.; Thiselton, D. L.; Bateman, J. B.; Ngo, J. T.; Sparkes,
- [89934] R. S.; Coleman, M.; Davies, K. E.; Bhattacharya, S. S.: Localisation
- [89935] of the gene for Norrie disease to between DXS7 and DXS426 on Xp. *Hum. Genet.* 88: 349–350, 1992.
- [89936] 15869.
- [89937] Meindl, A.; Berger, W.; Meitinger, T.; van de Pol, D.; Achatz,
- [89938] H.; Dorner, C.; Haasemann, M.; Hellebrand, H.; Gal, A.; Cremers, F.;
- [89939] Ropers, H.-H.: Norrie disease is caused by mutations in an extracellular
- [89940] protein resembling C-terminal globular domain of mucins. *Nature Genet.* 2:
- [89941] 139–143, 1992.

[89943] 15870.

[89944] Meindl, A.; Lorenz, B.; Achatz, H.; Hellebrand, H.;
Schmitz-Valckenberg,

[89945] P.; Meitinger, T.: Missense mutations in the NDP gene in
patients

[89946] with a less severe course of Norrie disease. Hum. Molec.
Genet. 4:

[89947] 489–490, 1995.

[89948] 15871.

[89949] Meitinger, T.; Meindl, A.; Bork, P.; Rost, B.; Sander, C.;
Haasemann,

[89950] M.; Murken, J.: Molecular modeling of the Norrie disease
protein

[89951] predicts a cystine knot growth factor tertiary structure.
Nature

[89952] Genet. 5: 376–380, 1993.

[89953] 15872.

[89954] Moreira-Filho, C. A.; Neustein, I.: A presumptive new vari-
ant

[89955] of Norrie's disease. J. Med. Genet. 16: 125–128, 1979.

[89956] 15873.

- [89957] Nance, W. E.; Hara, S.; Hansen, A.; Elliott, J.; Lewis, M.; Chown,
[89958] B.: Genetic linkage studies in a Negro kindred with Norrie's disease. Am.
[89959] J. Hum. Genet. 21: 423-429, 1969.
- [89960] 15874.
- [89961] Ngo, J.; Spence, M. A.; Cortessis, V.; Bateman, J. B.; Sparkes,
[89962] R. S.: Duplicate report crossing over in Norrie disease family. (Letter) Am.
[89963] J. Med. Genet. 33: 286, 1989.
- [89964] 15875.
- [89965] Ngo, J. T.; Bateman, J. B.; Cortessis, V.; Sparkes, R. S.; Mohandas,
[89966] T.; Inana, G.; Spence, M. A.: Norrie disease: linkage analysis using
[89967] a 4.2-kb RFLP detected by a human ornithine aminotransferase cDNA
[89968] probe. Genomics 4: 539-545, 1989.
- [89969] 15876.
- [89970] Ngo, J. T.; Spence, M. A.; Cortessis, V.; Sparkes, R. S.;

Bateman,

[89971] J. B.: Recombinational event between Norrie disease and DXS7 loci. Clin.

[89972] Genet. 34: 43–47, 1988.

[89973] 15877.

[89974] Norrie, G.: Nogle Blindhedsaarsager: en oversigt. Hospitalstidende 76:

[89975] 141–147, 1933.

[89976] 15878.

[89977] Ohba, N.; Yamashita, T.: Primary vitreoretinal dysplasia resembling

[89978] Norrie's disease in a female: association with X autosome chromosomal

[89979] translocation. Brit. J. Ophthal. 70: 64–71, 1986.

[89980] 15879.

[89981] Phillips, C. I.; Newton, M.; Duvall, J.; Holloway, S.; Levy, A.

[89982] M.: Probably Norrie's disease due to mutation: two sporadic sibships

[89983] of two males each, a necropsy of one case, and, given Norrie's disease,

[89984] a calculation of the gene mutation frequency. Brit. J. Oph–

thal. 70:

[89985] 305–313, 1986.

[89986] 15880.

[89987] Rehm, H. L.; Gutierrez–Espeleta, G. A.; Garcia, R.; Jimenez, G.;

[89988] Khetarpal, U.; Priest, J. M.; Sims, K. B.; Keats, B. J. B.; Morton,

[89989] C. C.: Norrie disease gene mutation in a large Costa Rican kindred

[89990] with a novel phenotype including venous insufficiency. Hum. Mutat.

[89991] 9: 402–408, 1997.

[89992] 15881.

[89993] Schuback, D. E.; Chen, Z. Y.; Craig, I. W.; Breakefield, X. O.;

[89994] Sims, K. B.: Mutations in the Norrie disease gene. Hum. Mutat. 5:

[89995] 285–292, 1995.

[89996] 15882.

[89997] Shastry, B. S.: Identification of a recurrent missense mutation

- [89998] in the Norrie disease gene associated with a simplex case of exudative
- [89999] vitreoretinopathy. *Biochem. Biophys. Res. Commun.* 246: 35–38, 1998.
- [90000] 15883.
- [90001] Shastry, B. S.; Hejtmancik, J. F.; Plager, D. A.; Hartzler, M.
- [90002] K.; Trese, M. T.: Linkage and candidate gene analysis in X-linked
- [90003] familial exudative vitreoretinopathy. *Genomics* 27: 341–344, 1995.
- [90004] 15884.
- [90005] Shastry, B. S.; Hejtmancik, J. F.; Trese, M. T.: Identification
- [90006] of novel missense mutations in the Norrie disease gene associated
- [90007] with one X-linked and four sporadic cases of familial exudative vitreoretinopathy. *Hum.*
- [90008] *Mutat.* 9: 396–401, 1997.
- [90009] 15885.
- [90010] Sims, K. B.; Lebo, R. V.; Benson, G.; Shalish, C.; Schuback, D.;
- [90011] Chen, Z. Y.; Bruns, G.; Craig, I. W.; Golbus, M. S.; Breake-

field,

[90012] X. O.: The Norrie disease gene maps to a 150 kb region on chromosome

[90013] Xp11.3. Hum. Molec. Genet. 1: 83-89, 1992.

[90014] 15886.

[90015] Sims, K. B.; Ozelius, L.; Corey, T.; Rinehart, W. B.; Libera-

[90016] R.; Haines, J.; Chen, W. J.; Norio, R.; Sankila, E.; de la Chapelle,

[90017] A.; Murphy, D. L.; Gusella, J.; Breakefield, X. O.: Norrie disease

[90018] gene is distinct from the monoamine oxidase genes. Am. J. Hum. Genet. 45:

[90019] 424-434, 1989.

[90020] 15887.

[90021] Taylor, P. J.; Coates, T.; Newhouse, M. L.: Episkopi blindness:

[90022] hereditary blindness in a Greek Cypriot family. Brit. J. Ophthalmol. 43:

[90023] 340-344, 1959.

[90024] 15888.

- [90025] Torrente, I.; Mangino, M.; Gennarelli, M.; Novelli, G.; Giannotti,
[90026] A.; Vadala, P.; Dallapiccola, B.: Two new missense mutations (A105T
[90027] and C110G) in the norrin gene in two Italian families with
Norrie
[90028] disease and familial exudative vitreoretinopathy. (Letter)
Am. J.
[90029] Med. Genet. 72: 242-244, 1997.
[90030] 15889.
[90031] Burgoyne, P. S.: Genetic homology and crossing over in
the X and
[90032] Y chromosomes of mammals. Hum. Genet. 61: 85-90,
1982.
[90033] 15890.
[90034] Cooke, H. J.; Brown, W. R. A.; Rappold, G. A.: Hypervariable telomeric
[90035] sequences from the human sex chromosomes are pseudoautosomal. Nature 317:
[90036] 687-692, 1985.
[90037] 15891.

- [90038] Darling, S. M.; Banting, G. S.; Pym, B.; Wolfe, J.; Goodfellow,
low,
- [90039] P. N.: Cloning an expressed gene shared by the human
sex chromosomes. *Proc.*
- [90040] *Nat. Acad. Sci.* 83: 135–139, 1986.
- [90041] 15892.
- [90042] Dracopoli, N. C.; Rettig, W. J.; Albino, A. P.; Esposito, D.;
Archidiacono,
- [90043] N.; Rocchi, M.; Siniscalco, M.; Old, L. J.: Genes controlling
gp25/30
- [90044] cell-surface molecules map to chromosomes X and Y and
escape X-inactivation. *Am.*
- [90045] *J. Hum. Genet.* 37: 199–207, 1985.
- [90046] 15893.
- [90047] Gelin, C.; Aubrit, F.; Phalipon, A.; Raynal, B.; Cole, S.; Kaczorek,
zorek,
- [90048] M.; Bernard, A.: The E2 antigen, a 32 kD glycoprotein involved in
- [90049] T-cell adhesion processes, is the MIC2 gene product.
EMBO J. 8:
- [90050] 3253–3259, 1989.

[90051] 15894.

[90052] Geller, R. L.; Shapiro, L. J.; Mohandas, T. K.: Fine mapping
of

[90053] the distal short arm of the human X chromosome using X/
Y translocations. Am.

[90054] J. Hum. Genet. 38: 884–890, 1986.

[90055] 15895.

[90056] Goodfellow, P.; Banting, G.; Sheer, D.; Ropers, H. H.;
Caine,

[90057] A.; Ferguson-Smith, M. A.; Povey, S.; Voss, R.: Genetic evi-
dence

[90058] that a Y-linked gene in man is homologous to a gene on
the X chromosome. Nature 302:

[90059] 346–349, 1983.

[90060] 15896.

[90061] Goodfellow, P.; Pym, B.; Mohandas, T.; Shapiro, L. J.: The
cell

[90062] surface antigen locus, MIC2X, escapes X-inactivation. Am.
J. Hum.

[90063] Genet. 36: 777–782, 1984.

[90064] 15897.

- [90065] Goodfellow, P. J.; Darling, S. M.; Thomas, N. S.; Goodfellow,
low,
- [90066] P. N.: A pseudoautosomal gene in man. Science 234:
740–743, 1986.
- [90067] 15898.
- [90068] Goodfellow, P. N.; Tippet, P.: A human quantitative polymorphism
morphism
- [90069] related to Xg blood groups. Nature 289: 404–405, 1981.
- [90070] 15899.
- [90071] Levy, R.; Dilley, J.; Fox, R. I.; Warnke, R.: A human thymus-leukemia
leukemia
- [90072] antigen defined by hybridoma monoclonal antibodies.
Proc. Nat. Acad.
- [90073] Sci. 76: 6552–6556, 1979.
- [90074] 15900.
- [90075] Pettersen, R. D.; Bernard, G.; Olafsen, M. K.; Pourtein, M.;
Lie,
- [90076] S. O.: CD99 signals caspase-independent T cell death. J.
Immun. 166:
- [90077] 4931–4942, 2001.
- [90078] 15901.

- [90079] Ropers, H. H.; Zimmer, J.; Strobl, G.; Goodfellow, P.: The MIC2X
- [90080] (12E7) locus maps distally from STS on Xp. (Abstract) Cytogenet.
- [90081] Cell Genet. 40: 736 only, 1985.
- [90082] 15902.
- [90083] Simmler, M.-C.; Rouyer, F.; Vergnaud, G.; Nystrom-Lahti, M.; Ngo,
- [90084] K. Y.; de la Chapelle, A.; Weissenbach, J.: Pseudoautosomal DNA sequences
- [90085] in the pairing region of the human sex chromosomes. Nature 317:
- [90086] 692-697, 1985.
- [90087] 15903.
- [90088] Tippett, P.; Shaw, M.-A.; Green, C. A.; Daniels, G. L.: The 12E7
- [90089] red cell quantitative polymorphism: control by the Y-borne locus,
- [90090] Yg. Ann. Hum. Genet. 50: 339-347, 1986.
- [90091] 15904.
- [90092] Janz, R.; Sudhof, T. C.; Hammer, R. E.; Unni, V.; Siegel-

baum, S.

- [90093] A.; Bolshakov, V. Y.: Essential roles in synaptic plasticity for
- [90094] synaptogyrin I and synaptophysin I. *Neuron* 24: 687–700, 1999.
- [90095] 15905.
- [90096] McMahon, H. T.; Bolshakov, V. Y.; Janz, R.; Hammer, R. E.; Siegelbaum,
- [90097] S. A.; Sudhof, T. C.: Synaptophysin, a major synaptic vesicle protein,
- [90098] is not essential for neurotransmitter release. *Proc. Nat. Acad. Sci.* 93:
- [90099] 4760–4764, 1996.
- [90100] 15906.
- [90101] Brown, C. J.; Sekiguchi, T.; Nishimoto, T.; Willard, H. F.: Regional
- [90102] localization of CCG1 gene which complements hamster cell cycle mutation
- [90103] BN462 to Xq11–Xq13. *Somat. Cell Molec. Genet.* 15: 93–96, 1989.
- [90104] 15907.
- [90105] Derry, J. M. J.; Barnard, P. J.: Localization of the Ccg1 gene

- [90106] on the mouse X chromosome. (Abstract) Cytogenet. Cell Genet. 51:
- [90107] 988, 1989.
- [90108] 15908.
- [90109] Dikstein, R.; Ruppert, S.; Tjian, R.: TAFII250 is a bipartite
- [90110] protein kinase that phosphorylates the base transcription factor RAP74. Cell 84:
- [90111] 781–790, 1996.
- [90112] 15909.
- [90113] Giles, R. E.; Ruddle, F. H.: X-linkage of a human genetic locus
- [90114] that corrects the DNA synthesis lesion in tsC1AGOH mouse cells. Genetics 93:
- [90115] 975–996, 1979.
- [90116] 15910.
- [90117] Hisatake, K.; Hasegawa, S.; Takada, R.; Nakatani, Y.; Horikoshi,
- [90118] M.; Roeder, R. G.: The p250 subunit of native TATA box-binding factor
- [90119] TFIID is the cell-cycle regulatory protein CCG1. Nature 362: 179–181,

[90120] 1993.

[90121] 15911.

[90122] Jacobson, R. H.; Ladurner, A. G.; King, D. S.; Tjian, R.:
Structure

[90123] and function of a human TAFII250 double bromodomain
module. Science 288:

[90124] 1422–1425, 2000.

[90125] 15912.

[90126] Jha, K. K.; Ozer, H. L.: Genetic studies with a mutant
mouse cell,

[90127] ts-2 Balb-3T3, with a temperature-sensitive defect in
DNA synthesis.

[90128] (Abstract) Genetics 86: s32–s33, 1977.

[90129] 15913.

[90130] Jha, K. K.; Siniscalco, M.; Ozer, H. L.: Temperature-sensi-
tive

[90131] mutants of Balb-3T3 cells. III. Hybrids between ts2 and
other mouse

[90132] mutant cells affected in DNA synthesis and correction of
ts2 defect

[90133] by human X chromosome. Somat. Cell Genet. 6: 603–614,

1980.

[90134] 15914.

[90135] Allen, F. H.; Krabbe, S. M. R.; Corcoran, P. A.: A new phenotype

[90136] (McLeod) in the Kell blood-group system. Vox Sang. 6: 555-560, 1961.

[90137] 15915.

[90138] Bertelson, C. J.; Pogo, A. O.; Chaudhuri, A.; Marsh, W. L.; Redman,

[90139] C. M.; Banerjee, D.; Symmans, W. A.; Simon, T.; Frey, D.; Kunkel,

[90140] L. M.: Localization of the McLeod locus (XK) within Xp21 by deletion

[90141] analysis. Am. J. Hum. Genet. 42: 703-711, 1988.

[90142] 15916.

[90143] Danek, A.; Rubio, J. P.; Rampoldi, L.; Ho, M.; Dobson-Stone, C.;

[90144] Tison, F.; Symmans, W. A.; Oechsner, M.; Kalckreuth, W.; Watt, J.

[90145] M.; Corbett, A. J.; Hamdalla, H. H. M.; Marshall, A. G.; Sutton, I.;

[90146] Dotti, M. T.; Malandrini, A.; Walker, R. H.; Daniels, G.;

Monaco,

[90147] A. P.: McLeod neuroacanthocytosis: genotype and phenotype. Ann.

[90148] Neurol. 50: 755-764, 2001.

[90149] 15917.

[90150] Danek, A.; Witt, T. N.; Stockmann, H. B. A. C.; Weiss, B. J.; Schotland,

[90151] D. L.; Fischbeck, K. H.: Normal dystrophin in McLeod myopathy. Ann.

[90152] Neurol. 28: 720-722, 1990.

[90153] 15918.

[90154] Densen, P.; Wilkinson-Kroovand, S.; Mandell, G. L.; Sullivan, G.;

[90155] Oyen, R.; Marsh, W. L.: Kx: its relationship to chronic granulomatous

[90156] disease and genetic linkage with Xg. Blood 58: 34-37, 1981.

[90157] 15919.

[90158] de Saint-Basile, G.; Bohler, M. C.; Fischer, A.; Cartron, J.; Dufier,

[90159] J. L.; Griscelli, C.; Orkin, S. H.: Xp21 DNA microdeletion in

a patient

[90160] with chronic granulomatous disease, retinitis pigmentosa,
and McLeod

[90161] phenotype. Hum. Genet. 80: 85-89, 1988.

[90162] 15920.

[90163] Faillace, R. T.; Kingston, W. J.; Nanda, N. C.; Griggs, R. C.:

[90164] Cardiomyopathy associated with the syndrome of amy-
otrophic chorea

[90165] and acanthocytosis. Ann. Intern. Med. 96: 616-617, 1982.

[90166] 15921.

[90167] Frey, D.; Machler, M.; Seger, R.; Schmid, W.; Orkin, S. H.:
Gene

[90168] deletion in a patient with chronic granulomatous disease
and McLeod

[90169] syndrome: fine mapping of the Xk gene locus. Blood 71:
252-255,

[90170] 1988.

[90171] 15922.

[90172] Giblett, E. R.; Klebanoff, S. J.; Pincus, S. H.; Swanson, J.;

[90173] Park, B. H.; McCullough, J.: Kell phenotypes in chronic

granulomatous

[90174] disease: a potential transfusion hazard. Lancet I:
1235–1236, 1971.

[90175] 15923.

[90176] Hanaoka, N.; Yoshida, K.; Nakamura, A.; Furihata, K.; Seo,
T.;

[90177] Tani, Y.; Takahashi, J.; Ikeda, S.; Hanyu, N.: A novel
frameshift

[90178] mutation in the McLeod syndrome gene in a Japanese
family. J. Neurol.

[90179] Sci. 165: 6–9, 1999.

[90180] 15924.

[90181] Ho, M.; Chelly, J.; Carter, N.; Danek, A.; Crocker, P.;
Monaco,

[90182] A. P.: Isolation of the gene for McLeod syndrome that en-
codes a novel

[90183] membrane transport protein. Cell 77: 869–880, 1994.

[90184] 15925.

[90185] Almind, K.; Ahlgren, M. G.; Hansen, T.; Urhammer, S. A.;
Clausen,

[90186] J. O.; Pedersen, O.: Discovery of a Met300Val variant in

Shc and

- [90187] studies of its relationship to birth weight and length, impaired insulin
- [90188] secretion, insulin resistance, and type 2 diabetes mellitus. J. Clin.
- [90189] Endocr. Metab. 84: 2241–2244, 1999.
- [90190] 15926.
- [90191] Harun, R. B.; Smith, K. K.; Leek, J. P.; Markham, A. F.; Norris,
- [90192] A.; Morrison, J. F. J.: Characterization of human SHC p66 cDNA and
- [90193] its processed pseudogene mapping to Xq12–q13.1. Genomics 42: 349–352,
- [90194] 1997.
- [90195] 15927.
- [90196] McGlade, J.; Cheng, A.; Pelicci, G.; Pelicci, P. G.; Pawson, T.
- [90197] : Shc proteins are phosphorylated and regulated by the v-src and v-fps
- [90198] protein-tyrosine-kinases. Proc. Nat. Acad. Sci. 89: 8869–8873, 1992.
- [90199] 15928.

- [90200] Migliaccio, E.; Giorgio, M.; Mele, S.; Pelicci, G.; Reboldi, P.;
- [90201] Pandolfi, P. P.; Lanfranccone, L.; Pelicci, P. G.: The p66(shc)
adaptor
- [90202] protein controls oxidative stress response and life span in
mammals. Nature 402:
- [90203] 309–313, 1999.
- [90204] 15929.
- [90205] Nemoto, S.; Finkel, T.: Redox regulation of forkhead pro-
teins
- [90206] through a p66shc-dependent signaling pathway. Science
295: 2450–2452,
- [90207] 2002.
- [90208] 15930.
- [90209] Pelicci, G.; Lanfranccone, L.; Grignani, F.; McGlade, J.; Cav-
allo,
- [90210] F.; Forni, G.; Nicoletti, I.; Grignani, F.; Pawson, T.; Pelicci,
P.
- [90211] G.: A novel transforming protein (SHC) with an SH2 do-
main is implicated
- [90212] in mitogenic signal transduction. Cell 70: 93–104, 1992.
- [90213] 15931.

- [90214] Yulug, I. G.; Egan, S. E.; See, C. G.; Fisher, E. M. C.: A human
- [90215] SHC-related sequence maps to chromosome 17, the SHC gene maps to chromosome
- [90216] 1. Hum. Genet. 96: 245-248, 1995.
- [90217] 15932.
- [90218] Zhang, L.; Camerini, V.; Bender, T. P.; Ravichandran, K. S.: A
- [90219] nonredundant role for the adapter protein Shc in thymic T cell development. Nature
- [90220] Immun. 3: 749-755, 2002.
- [90221] 15933.
- [90222] Selig, S.; Bruno, S.; Scharf, J. M.; Wang, C. H.; Vitale, E.; Gilliam,
- [90223] T. C.; Kunkel, L. M.: Expressed cadherin pseudogenes are localized
- [90224] to the critical region of the spinal muscular atrophy gene. Proc.
- [90225] Nat. Acad. Sci. 92: 3702-3706, 1995.
- [90226] 15934.
- [90227] Thompson, T. G.; DiDonato, C. J.; Simard, L. R.; Ingraham,

S. E.;

[90228] Burghes, A. H. M.; Crawford, T. O.; Rochette, C.; Mendell, J. R.;

[90229] Wasmuth, J. J.: A novel cDNA detects homozygous microdeletions in

[90230] greater than 50% of type I spinal muscular atrophy patients.. Nature

[90231] Genet. 9: 56-62, 1995.

[90232] 15935.

[90233] Abramovitz, M.; Boie, Y.; Nguyen, T.; Rushmore, T. H.; Bayne, M.

[90234] A.; Metters, K. M.; Slipetz, D. M.; Grygorczyk, R.: Cloning and expression

[90235] of a cDNA for the human prostanoid FP receptor. J. Biol. Chem. 269:

[90236] 2632-2636, 1994.

[90237] 15936.

[90238] Betz, R.; Lagercrantz, J.; Kedra, D.; Dumanski, J. P.; Nordenskjold,

[90239] A.: Genomic structure, 5-prime flanking sequences, and precise localization

[90240] in 1p31.1 of the human prostaglandin F receptor gene.

Biochem. Biophys.

[90241] Res. Commun. 254: 413–416, 1999.

[90242] 15937.

[90243] Sugimoto, Y.; Yamasaki, A.; Segi, E.; Tsuboi, K.; Aze, Y.;
Nishimura,

[90244] T.; Oida, H.; Yoshida, N.; Tanaka, T.; Katsuyama, M.; Ha-
sumoto, K.;

[90245] Murata, T.; Hirata, M.; Ushikubi, F.; Negishi, M.; Ichikawa,
A.; Narumiya,

[90246] S.: Failure of parturition in mice lacking the prostaglandin
F receptor. Science 277:

[90247] 681–683, 1997.

[90248] 15938.

[90249] Jean, L.; Smih, F.; Olivier, E.; Soury, E.; Simon–Chazottes,
D.;

[90250] Guenet, J. L.; Mattei, M. G.; Salier, J. P.: Comparative as-
signments

[90251] of the genes of the inter- α -inhibitor family in human
and mouse:

[90252] ITIH4 is close to ITIH1 and ITIH3, on HSA 3 and MMU 14.
Genomics 41:

[90253] 139–140, 1997.

[90254] 15939.

[90255] Nishimura, H.; Kakizaki, I.; Muta, T.; Sasaki, N.; Pu, P. X.; Yamashita,

[90256] T.; Nagasawa, S.: cDNA and deduced amino acid sequence of human PK-120,

[90257] a plasma kallikrein-sensitive glycoprotein. FEBS Lett. 357: 207-211,

[90258] 1995.

[90259] 15940.

[90260] Saguchi, K.; Tobe, T.; Hashimoto, K.; Sano, Y.; Nakano, Y.; Miura,

[90261] N.-H.; Tomita, M.: Cloning and characterization of cDNA for inter-alpha-trypsin

[90262] inhibitor family heavy chain-related protein (IHRP), a novel human

[90263] plasma glycoprotein. J. Biochem. 117: 14-18, 1995.

[90264] 15941.

[90265] Tobe, T.; Saguchi, K.; Hashimoto, K.; Miura, N.-H.; Tomita, M.;

[90266] Li, F.; Wang, Y.; Minoshima, S.; Shimizu, N.: Mapping of human inter-alpha-trypsin

- [90267] inhibitor family heavy chain-related protein gene (ITIH1)
to human
- [90268] chromosome 3p21-p14. Cytogenet. Cell. Genet. 71:
296-298, 1995.
- [90269] 15942.
- [90270] Geppert, M.; Khvotchev, M.; Krasnoperov, V.; Goda, Y.;
Missler,
- [90271] M.; Hammer, R. E.; Ichtchenko, K.; Petrenko, A. G.; Sudhof,
T. C.
- [90272] : Neurexin I-alpha is a major alpha-latrotoxin receptor
that cooperates
- [90273] in alpha-latrotoxin action. J. Biol. Chem. 273: 1705-1710,
1998.
- [90274] 15943.
- [90275] Ichtchenko, K.; Hata, Y.; Nguyen, T.; Ullrich, B.; Missler,
M.;
- [90276] Moomaw, C.; Sudhof, T. C.: Neuroligin 1: a splice site-
specific ligand
- [90277] for beta-neurexins. Cell 81: 435-443, 1995.
- [90278] 15944.
- [90279] Kleiderlein, J. J.; Nisson, P. E.; Jessee, J.; Li, W.-B.; Becker,

- [90280] K. G.; Derby, M. L.; Ross, C. A.; Margolis, R. L.: CCG repeats in
- [90281] cDNAs from human brain. Hum. Genet. 103: 666–673, 1998. Note: Erratum:
- [90282] Hum. Genet. 104: 113 only, 1999.
- [90283] 15945.
- [90284] Schoneberg, T.; Yun, J.; Wenkert, D.; Wess, J.: Functional rescue
- [90285] of mutant V2 vasopressin receptors causing nephrogenic diabetes insipidus
- [90286] by a co-expressed receptor polypeptide. EMBO J. 15: 1283–1291, 1996.
- [90287] 15946.
- [90288] Aly, A. M.; Higuchi, M.; Kasper, C. K.; Kazazian, H. H., Jr.; Antonarakis,
- [90289] S. E.; Hoyer, L. W.: Hemophilia A due to mutations that create new
- [90290] N-glycosylation sites. Proc. Nat. Acad. Sci. 89: 4933–4937, 1992.
- [90291] 15947.
- [90292] Greenberg, J. H.; Turner, C. G., II; Zegura, S. L.: The settlement

[90293] of the Americas: a comparison of the linguistic, dental,
and genetic

[90294] evidence. *Curr. Anthropol.* 27: 477–497, 1986.

[90295] 15948.

[90296] Margolis, R.: Personal Communication. Baltimore, Md.
3/29/2000.

[90297] 15949.

[90298] Missler, M.; Sudhof, T. C.: Neurexins: three genes and
1001 products. *Trends*

[90299] *Genet.* 14: 20–26, 1998.

[90300] 15950.

[90301] Ikeshima, H.; Imai, S.; Shimoda, K.; Hata, J.; Takano, T.:
Expression

[90302] of a MADS box gene, MEF2D, in neurons of the mouse
central nervous

[90303] system: implication of its binary function in myogenic and
neurogenic

[90304] cell lineages. *Neurosci. Lett.* 200: 117–120, 1995.

[90305] 15951.

[90306] Delhase, M.; Hayakawa, M.; Chen, Y.; Karin, M.: Positive
and negative

- [90307] regulation of I-kappa-B kinase activity through IKK-beta subunit phosphorylation. Science 284:
- [90308] 309-312, 1999.
- [90309] 15952.
- [90310] DiDonato, J. A.; Hayakawa, M.; Rothwarf, D. M.; Zandi, E.; Karin,
- [90311] M.: A cytokine-responsive IkappaB kinase that activates the transcription
- [90312] factor NF-kappaB. Nature 388: 548-554, 1997.
- [90313] 15953.
- [90314] Mercurio, F.; Zhu, H.; Murray, B. W.; Shevchenko, A.; Bennett,
- [90315] B. L.; Li, J.; Young, D. B.; Barbosa, M.; Mann, M.; Manning, A.; Rao,
- [90316] A.: IKK-1 and IKK-2: cytokine-activated I-kappa-B kinases essential
- [90317] for NF-kappa-B activation. Science 278: 860-866, 1997.
- [90318] 15954.
- [90319] Matsushima, A.; Kaisho, T.; Rennert, P. D.; Nakano, H.; Kurosawa,
- [90320] K.; Uchida, D.; Takeda, K.; Akira, S.; Matsumoto, M.: Es-

sential role

[90321] of nuclear factor (NF)-kappa-B-inducing kinase and inhibitor of kappa-B

[90322] (I-kappa-B) kinase alpha in NF-kappa-B activation through lymphotoxin

[90323] beta receptor, but not through tumor necrosis factor receptor I. J.

[90324] Exp. Med. 193: 631-636, 2001.

[90325] 15955.

[90326] Brzezinski, A.: Melatonin in humans. New Eng. J. Med. 336: 186-195,

[90327] 1997.

[90328] 15956.

[90329] Liu, R.-Y.; Zhou, J.-N.; van Heerikhuize, J.; Hofman, M. A.; Swaab,

[90330] D. F.: Decreased melatonin levels in postmortem cerebrospinal fluid

[90331] in relation to aging, Alzheimer's disease, and apolipoprotein E-epsilon-4/4

[90332] genotype. J. Clin. Endocr. Metab. 84: 323-327, 1999.

[90333] 15957.

- [90334] Nelson, C. S.; Ikeda, M.; Gompf, H. S.; Robinson, M. L.; Fuchs,
- [90335] N. K.; Yoshioka, T.; Neve, K. A.; Allen, C. N.: Regulation of melatonin
- [90336] 1a receptor signaling and trafficking by asparagine-124. Molec. Endocr. 15:
- [90337] 1306-1317, 2001.
- [90338] 15958.
- [90339] Reppert, S. M.; Weaver, D. R.: Melatonin madness. Cell 83: 1059-1062,
- [90340] 1995.
- [90341] 15959.
- [90342] Reppert, S. M.; Weaver, D. R.; Ebisawa, T.: Cloning and characterization
- [90343] of a mammalian melatonin receptor that mediates reproductive and circadian
- [90344] responses. Neuron 13: 1177-1185, 1994.
- [90345] 15960.
- [90346] Deng, Z.; Centola, M.; Chen, X.; Sood, R.; Vedula, A.; Fischel-Ghodsian,
- [90347] N.; Kastner, D. L.: Identification of two Kruppel-related

zinc finger

- [90348] genes (ZNF200 and ZNF210) from human chromosome 16p13.3. *Genomics* 53: 97–103, 1998.
- [90349] 15961.
- [90350] 15961.
- [90351] Muller-Pillasch, F.; Zimmerhackl, F.; Lacher, U.; Schultz, N.;
- [90352] Hameister, H.; Varga, G.; Friess, H.; Buchler, M.; Adler, G.; Gress,
- [90353] T. M.: Cloning of novel transcripts of the human guanine-nucleotide-exchange
- [90354] factor Mss4: in situ chromosomal mapping and expression in pancreatic
- [90355] cancer. *Genomics* 46: 389–396, 1997.
- [90356] 15962.
- [90357] Yu, H.; Schreiber, S. L.: Cloning, Zn(2+) binding, and structural
- [90358] characterization of the guanine nucleotide exchange factor human Mss4. *Biochemistry* 34: 9103–9110, 1995.
- [90359] 9103–9110, 1995.
- [90360] 15963.

- [90361] Kordes, E.; Savelyeva, L.; Schwab, M.; Rommelaere, J.; Jau-
niaux,
- [90362] J.-C.; Cziepluch, C.: Isolation and characterization of hu-
man SGT
- [90363] and identification of homologues in *Saccharomyces cere-*
visiae and *Caenorhabditis*
- [90364] *elegans*. *Genomics* 52: 90–94, 1998.
- [90365] 15964.
- [90366] Yabe, D.; Nakamura, T.; Kanazawa, N.; Tashiro, K.; Honjo,
T.:
- [90367] Calumenin, a Ca(2+)-binding protein retained in the en-
doplasmic reticulum
- [90368] with a novel carboxyl-terminal sequence, *HDEF. J. Biol.*
Chem. 272:
- [90369] 18232–18239, 1997.
- [90370] 15965.
- [90371] Yabe, D.; Taniwaki, M.; Nakamura, T.; Kanazawa, N.;
Tashiro, K.;
- [90372] Honjo, T.: Human calumenin gene (CALU): cDNA isolation
and chromosomal
- [90373] mapping to 7q32. *Genomics* 49: 331–333, 1998.

[90374] 15966.

[90375] Angelin-Duclos, C.; Cattoretti, G.; Lin, K.-I.; Calame, K.:
Commitment

[90376] of B lymphocytes to a plasma cell fate is associated with
Blimp-1

[90377] expression in vivo. J. Immun. 165: 5462-5471, 2000.

[90378] 15967.

[90379] Chang, D. H.; Angelin-Duclos, C.; Calame, K.: BLIMP-1:
trigger

[90380] for differentiation of myeloid lineage. Nature Immun. 1:
169-176,

[90381] 2000.

[90382] 15968.

[90383] Huang, S.: Blimp-1 is the murine homolog of the human
transcriptional

[90384] repressor PRDI-BF1. (Letter) Cell 78: 9 only, 1994.

[90385] 15969.

[90386] Keller, A. D.; Maniatis, T.: Identification and characteriza-
tion

[90387] of a novel repressor of beta-interferon gene expression.
Genes Dev. 5:

- [90388] 868–879, 1991.
- [90389] 15970.
- [90390] Mock, B. A.; Liu, L.; Le Paslier, D.; Huang, S.: The B-lymphocyte
- [90391] maturation promoting transcription factor BLIMP1/PRDI-BF1 maps to
- [90392] D6S447 on human chromosome 6q21–q22.1 and the syntenic region of mouse
- [90393] chromosome 10. Genomics 37: 24–28, 1996.
- [90394] 15971.
- [90395] Ito, T.; Yang, M.; May, W. S.: RAX, a cellular activator for double-stranded
- [90396] RNA-dependent protein kinase during stress signaling. J. Biol. Chem. 274:
- [90397] 15427–15432, 1999.
- [90398] 15972.
- [90399] Patel, R. C.; Sen, G. C.: PACT, a protein activator of the interferon-induced
- [90400] protein kinase, PKR. EMBO J. 17: 4379–4390, 1998.
- [90401] 15973.

- [90402] Scott, A. F.: Personal Communication. Baltimore, Md.
2001.
- [90403] 15974.
- [90404] Shearman, L. P.; Zylka, M. J.; Weaver, D. R.; Kolakowski, L.
F.,
[90405] Jr.; Reppert, S. M.: Two period homologs: circadian ex-
pression and
[90406] photic regulation in the suprachiasmatic nuclei. Neuron
19: 1261–1269,
[90407] 1997.
- [90408] 15975.
- [90409] Toh, K. L.; Jones, C. R.; He, Y.; Eide, E. J.; Hinz, W. A.; Vir-
shup,
[90410] D. M.; Ptacek, L. J.; Fu, Y.-H.: An hPer2 phosphorylation
site mutation
[90411] in familial advanced sleep phase syndrome. Science 291:
1040–1043,
[90412] 2001.
- [90413] 15976.
- [90414] Engelender, S.; Wanner, T.; Kleiderlein, J. J.; Wakabayashi,
K.;

- [90415] Tsuji, S.; Takahashi, H.; Ashworth, R.; Margolis, R. L.; Ross, C.
- [90416] A.: Organization of the human synphilin-1 gene, a candidate for Parkinson's
- [90417] disease. *Mammalian Genome* 11: 763-766, 2000.
- [90418] 15977.
- [90419] Kitao, S.; Ohsugi, I.; Ichikawa, K.; Goto, M.; Furuichi, Y.; Shimamoto,
- [90420] A.: Cloning of two new human helicase genes of the RecQ family: biological
- [90421] significance of multiple species in higher eukaryotes. *Genomics* 54:
- [90422] 443-452, 1998.
- [90423] 15978.
- [90424] Mohaghegh, P.; Hickson, I. D.: DNA helicase deficiencies associated
- [90425] with cancer predisposition and premature ageing disorders. *Hum. Molec.*
- [90426] *Genet.* 10: 741-746, 2001.
- [90427] 15979.
- [90428] Chien, W.; Pei, L.: A novel binding factor facilitates nuclear

- [90429] translocation and transcriptional activation function of the pituitary
- [90430] tumor-transforming gene product. J. Biol. Chem. 275: 19422–19427,
- [90431] 2000.
- [90432] 15980.
- [90433] Yaspo, M.–L.; Aaltonen, J.; Horelli-Kuitunen, N.; Peltonen, L.;
- [90434] Lehrach, H.: Cloning of a novel human putative type Ia integral membrane
- [90435] protein mapping to 21q22.3. Genomics 49: 133–136, 1998.
- [90436] 15981.
- [90437] Yaspo, M.–L.; Gellen, L.; Mott, R.; Korn, B.; Nizetic, D.; Poustka,
- [90438] A. M.; Lehrach, H.: Model for a transcript map of human chromosome
- [90439] 21: isolation of new coding sequences from exon and enriched cDNA
- [90440] libraries. Hum. Molec. Genet. 4: 1291–1304, 1995.
- [90441] 15982.

- [90442] Su, K.; Kyaw, H.; Fan, P.; Zeng, Z.; Shell, B. K.; Carter, K. C.;
- [90443] Li, Y.: Isolation, characterization, and mapping of two human potassium
- [90444] channels. *Biochem. Biophys. Res. Commun.* 241: 675–681, 1997.
- [90445] 15983.
- [90446] Ghiso, J. A.; Holton, J.; Miravalle, L.; Calero, M.; Lashley, T.;
- [90447] Vidal, R.; Houlden, H.; Wood, N.; Neubert, T. A.; Rostagno, A.; Plant,
- [90448] G.; Revesz, T.; Frangione, B.: Systemic amyloid deposits in familial
- [90449] British dementia. *J. Biol. Chem.* 276: 43909–43914, 2001.
- [90450] 15984.
- [90451] Stromgren, E.; Dalby, A.; Dalby, M.; Ranheim, B. :*Acta Neurol.*
- [90452] *Scand.* 46 (suppl. 43): 97–98, 1970.
- [90453] 15985.
- [90454] Gruber, A. D.; Schreur, K. D.; Ji, H.–L.; Fuller, C. M.; Pauli,
- [90455] B. U.: Molecular cloning and transmembrane structure of

hCLCA2 from

[90456] human lung, trachea, and mammary gland. *Am. J. Physiol.* 276: C1261–C1270,

[90457] 1999.

[90458] 15986.

[90459] Claustres, M.; Gerrard, B.; White, M. B.; Desgeorges, M.; Kjellberg,

[90460] P.; Rollin, B.; Dean, M.: A new mutation (1078delT) in exon 7 of

[90461] the CFTR gene in a southern French adult with cystic fibrosis. *Genomics* 13:

[90462] 907–908, 1992.

[90463] 15987.

[90464] Claustres, M.; Laussel, M.; Desgeorges, M.; Giansily, M.; Culard,

[90465] J.-F.; Razakatsara, G.; Demaille, J.: Analysis of the 27 exons and

[90466] flanking regions of the cystic fibrosis gene: 40 different mutations

[90467] account for 91.2% of the mutant alleles in Southern France. *Hum.*

[90468] *Molec. Genet.* 2: 1209–1213, 1993.

[90469] 15988.

[90470] Costes, B.; Girodon, E.; Ghanem, N.; Flori, E.; Jardin, A.; Soufir,

[90471] J. C.; Goossens, M.: Frequent occurrence of the CFTR intron 8 (TG)_n

[90472] 5T allele in men with congenital bilateral absence of the vas deferens. *Europ.*

[90473] *J. Hum. Genet.* 3: 285–293, 1995.

[90474] 15989.

[90475] Cromwell, O.; Walport, M. J.; Morris, H. R.; Taylor, G. W.; Hodson,

[90476] M. E.; Batten, J.; Kay, A. B.: Identification of leukotrienes D and

[90477] B in sputum from cystic fibrosis patients. *Lancet* II: 164–165, 1981.

[90478] 15990.

[90479] Cuppens, H.; Lin, W.; Jaspers, M.; Costes, B.; Teng, H.; Vankeerberghen,

[90480] A.; Jorissen, M.; Droogmans, G.; Reynaert, I.; Goossens, M.; Nilius,

[90481] B.; Cassiman, J.–J.: Polyvariant mutant cystic fibrosis transmembrane

- [90482] conductance regulator genes: the polymorphic (TG)_n locus explains
- [90483] the partial penetrance of the T5 polymorphism as a disease mutation. J.
- [90484] Clin. Invest. 101: 487–496, 1998.
- [90485] 15991.
- [90486] Cuppens, H.; Marynen, P.; De Boeck, C.; De Baets, F.; Eggermont,
- [90487] E.; Van den Berghe, H.; Cassiman, J. J.: A child, homozygous for
- [90488] a stop codon in exon 11, shows milder cystic fibrosis symptoms than
- [90489] her heterozygous nephew. J. Med. Genet. 27: 717–719, 1990.
- [90490] 15992.
- [90491] Cutting, G. R.; Kasch, L. M.; Rosenstein, B. J.; Tsui, L.-C.;
- [90492] Kazazian, H. H., Jr.; Antonarakis, S. E.: Two patients with cystic
- [90493] fibrosis, nonsense mutations in each cystic fibrosis gene, and mild
- [90494] pulmonary disease. New Eng. J. Med. 323: 1685–1689, 1990.

- [90495] 15993.
- [90496] Cutting, G. R.; Kasch, L. M.; Rosenstein, B. J.; Zielenski, J.;
- [90497] Tsui, L.-C.; Antonarakis, S. E.; Kazazian, H. H., Jr.: A cluster
ter
- [90498] of cystic fibrosis mutations in the first nucleotide-binding
fold
- [90499] of the cystic fibrosis conductance regulator protein. Nature 346:
ture 346:
- [90500] 366-369, 1990.
- [90501] 15994.
- [90502] Daigneault, J.; Aubin, G.; Simard, F.; De Braekeleer, M.:
Genetic
- [90503] epidemiology of cystic fibrosis in Saguenay-Lac-St-Jean
(Quebec, Canada). Clin.
- [90504] Genet. 40: 298-303, 1991.
- [90505] 15995.
- [90506] Dean, M.; White, M. B.; Amos, J.; Gerrard, B.; Stewart, C.;
Khaw,
- [90507] K.-T.; Leppert, M.: Multiple mutations in highly conserved
residues
- [90508] are found in mildly affected cystic fibrosis patients. Cell

61:

[90509] 863–870, 1990.

[90510] 15996.

[90511] Dean, M.; White, M. B.; Gerrard, B.; Amos, J.; Milunsky, A.:

[90512] A 22–bp deletion in the coding region of the cystic fibrosis gene. *Genomics* 13:

[90513] 235–236, 1992.

[90514] 15997.

[90515] Delaney, S. J.; Alton, E. W. F. W.; Smith, S. N.; Lunn, D. P.;

[90516] Farley, R.; Lovelock, P. K.; Thomson, S. A.; Hume, D. A.; Lamb, D.;

[90517] Porteous, D. J.; Dorin, J. R.; Wainwright, B. J.: Cystic fibrosis

[90518] mice carrying the missense mutation G551D replicate human genotype–phenotype

[90519] correlations. *EMBO J.* 15: 955–963, 1996.

[90520] 15998.

[90521] de Meeus, A.; Guittard, C.; Desgeorges, M.; Carles, S.; Demaille,

[90522] J.; Claustres, M.: Linkage disequilibrium between the

M470V variant

[90523] and the IVS8 polyT alleles of the CFTR gene in CBAVD. J.

Med. Genet. 35:

[90524] 594–596, 1998.

[90525] 15999.

[90526] Denning, G. M.; Anderson, M. P.; Amara, J. F.; Marshall, J.;

Smith,

[90527] A. E.; Welsh, M. J.: Processing of mutant cystic fibrosis

transmembrane

[90528] conductance regulator is temperature-sensitive. Nature

358: 761–764,

[90529] 1992.

[90530] 16000.

[90531] Devor, D. C.; Schultz, B. D.: Ibuprofen inhibits cystic fibrosis

[90532] transmembrane conductance regulator-mediated Cl(–) secretion. J.

[90533] Clin. Invest. 102: 679–687, 1998.

[90534] 16001.

[90535] Dickinson, P.; Smith, S. N.; Webb, S.; Kilanowski, F. M.;
Campbell,

- [90536] I. J.; Taylor, M. S.; Porteous, D. J.; Willemsen, R.; de Jonge, H.
- [90537] R.; Farley, R.; Alton, E. W. F. W.; Dorin, J. R.: The severe G480C
- [90538] cystic fibrosis mutation, when replicated in the mouse, demonstrates
- [90539] mistrafficking, normal survival and organ-specific bio-electrics. Hum.
- [90540] Molec. Genet. 11: 243-251, 2002.
- [90541] 16002.
- [90542] Nicolino, P. M.; Dupin, H.; Macebeo, V.; Treppoz, S.; Chatelain,
- [90543] P. G.; Wolcott-Rallison syndrome (diabetes mellitus and spondyloepiphyseal
- [90544] dysplasia): a plausible existence of a gene(s) important for the
- [90545] mutation of neonatal pancreatic beta cell function. (Abstract) Hormone
- [90546] Res. 50 (suppl. 3): 77 only, 1998.
- [90547] 16003.
- [90548] Shi, Y.; An, J.; Liang, J.; Hayes, S. E.; Sandusky, G. E.; Stramm,

- [90549] L. E.; Yang, N. N.: Characterization of a mutant pancreatic eIF-2-alpha
- [90550] kinase, PEK, and co-localization with somatostatin in islet delta
- [90551] cells. J. Biol. Chem. 274: 5723-5730, 1999.
- [90552] 16004.
- [90553] Shi, Y.; Vатtem, K. M.; Sood, R.; An, J.; Liang, J.; Stramm, L.;
- [90554] Wek, R. C.: Identification and characterization of pancreatic eukaryotic
- [90555] initiation factor 2 alpha-subunit kinase, PEK, involved in translational
- [90556] control. Molec. Cell. Biol. 18: 7499-7509, 1998.
- [90557] 16005.
- [90558] Csoka, A. B.; Scherer, S. W.; Stern, R.: Expression analysis of
- [90559] six paralogous human hyaluronidase genes clustered on chromosomes
- [90560] 3p21 and 7q31. Genomics 60: 356-361, 1999.
- [90561] 16006.
- [90562] Dolganov, G. M.; Maser, R. S.; Novikov, A.; Tosto, L.;

Chong, S.;

[90563] Bressan, D. A.; Petrini, J. H. J.: Human Rad50 is physically associated

[90564] with human Mre11: identification of a conserved multi-protein complex

[90565] implicated in recombinational DNA repair. *Molec. Cell Biol.* 16:

[90566] 4832–4841, 1996.

[90567] 16007.

[90568] Hopfner, K.-P.; Craig, L.; Moncalian, G.; Zinkel, R. A.; Usui,

[90569] T.; Owen, B. A. L.; Karcher, A.; Henderson, B.; Bodmer, J.-L.; McMurray,

[90570] C. T.; Carney, J. P.; Petrini, J. H. J.; Tainer, J. A.: The Rad50

[90571] zinc-hook is a structure joining Mre11 complexes in DNA recombination

[90572] and repair. *Nature* 418: 562–566, 2002.

[90573] 16008.

[90574] Luo, G.; Yao, M. S.; Bender, C. F.; Mills, M.; Bladl, A. R.; Bradley,

[90575] A.; Petrini, J. H. J.: Disruption of mRad50 causes embryonic stem

- [90576] cell lethality, abnormal embryonic development, and sensitivity to
- [90577] ionizing radiation. *Proc. Nat. Acad. Sci.* 96: 7376–7381, 1999.
- [90578] 16009.
- [90579] Trujillo, K. M.; Yuan, S.-S. F.; Lee, E. Y.-H. P.; Sung, P.: Nuclease
- [90580] activities in a complex of human recombination and DNA repair factors
- [90581] Rad50, Mre11, and p95. *J. Biol. Chem.* 273: 21447–21450, 1998.
- [90582] 16010.
- [90583] Clark, J.; Lu, Y. J.; Sidhar, S. K.; Parker, C.; Gill, S.; Smedley,
- [90584] D.; Hamoudi, R.; Linehan, W.; Shipley, J.; Cooper, C.: Fusion of
- [90585] splicing factor genes PSF and NonO (p54nrb) to the TFE3 gene in papillary
- [90586] renal cell carcinoma. *Oncogene* 15: 2233–2239, 1997.
- [90587] 16011.
- [90588] Heimann, P.; El Housni, H.; Ogur, G.; Weterman, M. A. J.; Petty,

- [90589] E. M.; Vassart, G.: Fusion of a novel gene, RCC17, to the TFE3 gene
- [90590] in t(X;17)(p11.2;q25.3)-bearing papillary renal cell carcinomas. Cancer
- [90591] Res. 61: 4130-4135, 2001.
- [90592] 16012.
- [90593] Henthorn, P. S.; Stewart, C. C.; Kadesch, T.; Puck, J. M.: The
- [90594] gene encoding human TFE3, a transcription factor that binds the immunoglobulin
- [90595] heavy-chain enhancer, maps to Xp11.22. Genomics 11: 374-378, 1991.
- [90596] 16013.
- [90597] Joyama, S.; Ueda, T.; Shimizu, K.; Kudawara, I.; Mano, M.; Funai,
- [90598] H.; Takemura, K.; Yoshikawa, H.: Chromosome rearrangement at 17q25
- [90599] and Xp11.2 in alveolar soft-part sarcoma: a case report and review
- [90600] of the literature. Cancer 86: 1246-1250, 1999.
- [90601] 16014.

- [90602] Ladanyi, M.; Lui, M. Y.; Antonescu, C. R.; Krause-Boehm, A.; Meindl,
- [90603] A.; Argani, P.; Healey, J. H.; Ueda, T.; Yoshikawa, H.; Meloni-Ehrig,
- [90604] A.; Sorensen, P. H. B.; Mertens, F.; Mandahl, N.; van den Berghe,
- [90605] H.; Sciot, R.; Dal Cin, P.; Bridge, J.: The
der(17)t(X;17)(p11;q25)
- [90606] of human alveolar soft part sarcoma fuses the TFE3 transcription factor
- [90607] gene to ASPL, a novel gene at 17q25. *Oncogene* 20: 48–57, 2001.
- [90608] 16015.
- [90609] Macchi, P.; Notarangelo, L.; Giliani, S.; Strina, D.; Repetto,
- [90610] M.; Sacco, M. G.; Vezzoni, P.; Villa, A.: The genomic organization
- [90611] of the human transcription factor 3 (TFE3) gene. *Genomics* 28: 491–494,
- [90612] 1995.
- [90613] 16016.
- [90614] Shipley, J. M.; Birdsall, S.; Clark, J.; Crew, J.; Gill, S.; Linehan,

- [90615] M.; Gnarr, J.; Gisher, S.; Craig, I. W.; Cooper, C. S.: Mapping
[90616] the X chromosome breakpoint in two papillary renal cell
carcinoma
[90617] cell lines with a t(X;1)(p11.2;q21.2) and the first report of
a female
[90618] case. Cytogenet. Cell. Genet. 71: 280–284, 1995.
[90619] 16017.
[90620] Sidhar, S. K.; Clark, J.; Gill, S.; Hamoudi, R.; Crew, A. J.;
[90621] Gwilliam, R.; Ross, M.; Linehan, W. M.; Birdsall, S.; Shipley,
J.;
[90622] Cooper, C. S.: The t(X;1)(p11.2;q21.2) translocation in
papillary
[90623] renal cell carcinoma fuses a novel gene PRCC to the TFE3
transcription
[90624] factor gene. Hum. Molec. Genet. 5: 1333–1338, 1996.
[90625] 16018.
[90626] Larola, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.: Control
[90627] of mRNA decay by heat shock–ubiquitin–proteasome
pathway. Science 284:
[90628] 499–502, 1999.

[90629] 16019.

[90630] Deutsch, D.; Palmon, A.; Dafni, L.; Fisher, L.; Termine, J. D.;

[90631] Young, M.: Cloning, sequencing, and characterization of tuftelin:

[90632] a novel acidic enamel protein. Connect. Tissue Res. 27: 121 only,

[90633] 1992.

[90634] 16020.

[90635] Genomics 44: 248-250, 1997.

[90636] 16021.

[90637] Deiss, L. P.; Feinstein, E.; Berissi, H.; Cohen, O.; Kimchi, A.

[90638] : Identification of a novel serine/threonine kinase and a novel 15-kD

[90639] protein as potential mediators of the gamma interferon-induced cell

[90640] death. Genes Dev. 9: 15-30, 1995.

[90641] 16022.

[90642] Feinstein, E.; Druck, T.; Kastury, K.; Berissi, H.; Goodart, S.

[90643] A.; Overhauser, J.; Kimchi, A.; Huebner, K.: Assignment of

DAP1 and

[90644] DAPK: genes that positively mediate programmed cell death triggered

[90645] by IFN- γ --to chromosome regions 5p12.2 (sic) and 9q34.1, respectively. Genomics 29:

[90646] 305-307, 1995.

[90647] 16023.

[90648] Raveh, T.; Droguett, G.; Horwitz, M. S.; DePinho, R. A.; Kimchi,

[90649] A.: DAP kinase activates a p19-ARF/p53-mediated apoptotic checkpoint

[90650] to suppress oncogenic transformation. Nature Cell Biol. 3: 1-7,

[90651] 2001.

[90652] 16024.

[90653] Simpson, D. J.; Clayton, R. N.; Farrell, W. E.: Preferential loss

[90654] of death associated protein kinase expression in invasive pituitary

[90655] tumours is associated with either CpG island methylation or homozygous

[90656] deletion. Oncogene 21: 1217-1224, 2002.

- [90657] 16025.
- [90658] Hughes, K. A.; Hurlstone, A. F. L.; Tobias, E. S.; McFarlane, R.;
- [90659] Black, D. M.: Absence of ST7 mutations in tumor-derived cell lines
- [90660] and tumors. *Nature Genet.* 29: 380–381, 2001.
- [90661] 16026.
- [90662] Thomas, N. A.; Choong, D. Y. H.; Jokubaitis, V. J.; Neville, P.
- [90663] J.; Campbell, I. G.: Mutation of the ST7 tumor suppressor gene on
- [90664] 7q31.1 is rare in breast, ovarian and colorectal cancers. *Nature*
- [90665] *Genet.* 29: 379–380, 2001.
- [90666] 16027.
- [90667] Zenklusen, J. C.; Conti, C. J.; Green, E. D.: Mutational and functional
- [90668] analyses reveal that ST7 is a highly conserved tumor-suppressor gene
- [90669] on human chromosome 7q31. *Nature Genet.* 27: 392–398, 2001.

[90670] 16028.

[90671] Zenklusen, J. C.; Rodriguez, L. V.; LaCava, M.; Wang, Z.;
Goldstein,

[90672] L. S.; Conti, C. J.: Novel susceptibility locus for mouse
hepatomas:

[90673] evidence for a conserved tumor suppressor gene. Genome
Res. 6: 1070–1076,

[90674] 1996.

[90675] 16029.

[90676] Zenklusen, J. C.; Weitzel, J. N.; Ball, H. G.; Conti, C. J.: Al-
lelic

[90677] loss at 7q31.1 in human primary ovarian carcinomas sug-
gests the existence

[90678] of a tumor suppressor gene. Oncogene 11: 359–363,
1995.

[90679] 16030.

[90680] Lee, P. L.; Gelbart, T.; West, C.; Adams, M.; Blackstone, R.;
Beutler,

[90681] E.: Three genes encoding zinc finger proteins on human
chromosome

[90682] 6p21.3: members of a new subclass of the Kruppel gene
family containing

- [90683] the conserved SCAN box domain. Genomics 43: 191–201, 1997.
- [90684] 16031.
- [90685] Deutsch, D.; Palmon, A.; Fisher, L. W.; Kolodny, N.; Ter-
mine, J.
- [90686] D.; Young, M. F.: Sequencing of bovine enamelin (tuftelin),
a novel
- [90687] acidic enamel protein. J. Biol. Chem. 266: 16021–16028,
1991.
- [90688] 16032.
- [90689] Deutsch, D.; Palmon, A.; Young, M. F.; Selig, S.; Kearns, W.
G.;
- [90690] Fisher, L. W.: Mapping of the human tuftelin (TUFT1) gene
to chromosome
- [90691] 1 by fluorescence in situ hybridization. (Abstract) Mam-
malian Genome 5:
- [90692] 461–462, 1994.
- [90693] 16033.
- [90694] MacDougall, M.; Simmons, D.; Dodds, A.; Knight, C.; Luan,
X.; Zeichner–David,
- [90695] M.; Zhang, C.; Ryu, O. H.; Qian, Q.; Simmer, J. P.; Hu,
C.–C.: Cloning,

[90696] characterization, and tissue expression pattern of mouse tuftelin

[90697] cDNA. J. Dent. Res. 77: 1970–1978, 1998.

[90698] 16034.

[90699] Fujii, H.; Kanno, H.; Hirono, A.; Shiomura, T.; Miwa, S.: A single

[90700] amino acid substitution (157gly-to-val) in a phosphoglycerate kinase

[90701] variant (PGK Shizuoka) associated with chronic hemolysis and myoglobinuria. Blood 79:

[90702] 1582–1585, 1992.

[90703] 16035.

[90704] Fujii, H.; Krietsch, W. K. G.; Yoshida, A.: A single amino acid

[90705] substitution (asp-to-asn) in a phosphoglycerate kinase variant (PGK

[90706] Munchen) associated with enzyme deficiency. J. Biol. Chem. 255:

- [90707] 6421–6423, 1980.
- [90708] 16036.
- [90709] Fujii, H.; Yoshida, A.: Molecular abnormality of phospho-glycerate
- [90710] kinase–Uppsala associated with chronic nonspherocytic hemolytic anemia. Proc.
- [90711] Nat. Acad. Sci. 77: 5461–5465, 1980.
- [90712] 16037.
- [90713] Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors–Cayla, M.
- [90714] C.; Gillois, M.: Pig gene mapping: syntenry, attempt to assign the
- [90715] histocompatibility complex (SLA). (Abstract) Cytogenet. Cell Genet. 25:
- [90716] 159, 1979.
- [90717] 16038.
- [90718] Grzeschik, K.–H.; Allderdice, P. W.; Grzeschik, A.; Opitz, J.
- [90719] M.; Miller, O. J.; Siniscalco, M.: Cytological mapping of human X–linked
- [90720] genes by use of somatic cell hybrids involving an X–

autosome translocation. Proc.

[90721] Nat. Acad. Sci. 69: 69–73, 1972.

[90722] 16039.

[90723] Guis, M. S.; Karadsheh, N.; Mentzer, W. C.: Phosphoglycerate

[90724] kinase San Francisco: a new variant associated with hemolytic anemia

[90725] but not with neuromuscular manifestations. Am. J. Hemat. 25: 175–182,

[90726] 1987.

[90727] 16040.

[90728] Huijing, F.; Eicher, E. M.; Coleman, D. L.: Location of phosphorylase

[90729] kinase (Phk) in the mouse X-chromosome. Biochem. Genet. 9: 193–196,

[90730] 1973.

[90731] 16041.

[90732] Hutz, M. H.; Michelson, A. M.; Antonarakis, S. E.; Orkin, S. H.;

[90733] Kazazian, H. H., Jr.: Restriction site polymorphism in the phosphoglycerate

- [90734] kinase gene on the X chromosome. Hum. Genet. 66: 217–219, 1984.
- [90735] 16042.
- [90736] Konrad, P. N. J.; McCarthy, D. J.; Mauer, A. M.; Valentine, W.
- [90737] N.; Paglia, D. E.: Erythrocyte and leukocyte phosphoglycerate kinase
- [90738] deficiency with neurologic disease. J. Pediat. 82: 456–460, 1973.
- [90739] 16043.
- [90740] Kozak, L. P.; McLean, G. K.; Eicher, E. M.: X-linkage of phosphoglycerate
- [90741] kinase in the mouse. Biochem. Genet. 2: 41–47, 1974.
- [90742] 16044.
- [90743] Kraus, A. P.; Langston, M. F., Jr.; Lynch, B. L.: Red cell phosphoglycerate
- [90744] kinase deficiency: a new cause of non-spherocytic hemolytic anemia. Biochem.
- [90745] Biophys. Res. Commun. 30: 173–177, 1968.
- [90746] 16045.
- [90747] Krietsch, W. K.; Krietsch, H.; Kaiser, W.; Dunnwald, M.;

Kuntz,

[90748] G. W.; Duhn, J.; Bucher, T.: Hereditary deficiency of phosphoglycerate

[90749] kinase: a new variant in erythrocytes and leucocytes, not associated

[90750] with hemolytic anemia. *Europ. J. Clin. Invest.* 7: 427–435, 1977.

[90751] 16046.

[90752] Krietsch, W. K. G.; Eber, S. W.; Haas, B.; Rubbelt, W.; Kuntz,

[90753] G. W. K.: Characterization of a phosphoglycerate kinase deficiency

[90754] variant not associated with hemolytic anemia. *Am. J. Hum. Genet.* 32:

[90755] 364–373, 1980.

[90756] 16047.

[90757] Lay, A. J.; Jiang, X.-M.; Kisker, O.; Flynn, E.; Underwood, A.;

[90758] Condrón, R.; Hogg, P. J.: Phosphoglycerate kinase acts in tumour

[90759] angiogenesis as a disulphide reductase. *Nature* 408: 869–873, 2000.

- [90760] 16048.
- [90761] Maeda, M.; Bawle, E. V.; Kulkarni, R.; Beutler, E.; Yoshida, A.
- [90762] : Molecular abnormalities of a phosphoglycerate kinase variant generated
- [90763] by spontaneous mutation. Blood 79: 2759–2762, 1992.
- [90764] 16049.
- [90765] Maeda, M.; Yoshida, A.: Molecular defect of a phosphoglycerate
- [90766] kinase variant (PGK–Matsue) associated with hemolytic anemia: leu-to-pro
- [90767] substitution caused by T/A-to-C/G transition in exon 3. Blood 77:
- [90768] 1348–1352, 1991.
- [90769] 16050.
- [90770] Meera Khan, P.; Westerveld, A.; Grzeschik, K.–H.; Deys, B. F.;
- [90771] Garson, O. M.; Siniscalco, M.: X-linkage of human phosphoglycerate
- [90772] kinase confirmed in man–mouse and man–Chinese hamster somatic cell

- [90773] hybrids. *Am. J. Hum. Genet.* 23: 614–623, 1971.
- [90774] 16051.
- [90775] Michelson, A. M.; Blake, C. C. F.; Evans, S. T.; Orkin, S. H.
- [90776] : Structure of the human phosphoglycerate kinase gene
and the intron-mediated
- [90777] evolution and dispersal of the nucleotide-binding domain.
Proc. Nat.
- [90778] *Acad. Sci.* 82: 6965–6969, 1985.
- [90779] 16052.
- [90780] Michelson, A. M.; Markham, A. F.; Orkin, S. H.: Isolation
and
- [90781] DNA sequence of a full-length cDNA clone for human X
chromosome-encoded
- [90782] phosphoglycerate kinase. *Proc. Nat. Acad. Sci.* 80:
472–476, 1983.
- [90783] 16053.
- [90784] Miwa, S.; Nakawhima, K.; Oda, S.; Ogawa, H.; Nagafuji, H.;
Arima,
- [90785] M.; Okuna, T.; Nakashima, T.: Phosphoglycerate kinase
deficiency
- [90786] hereditary nonspherocytic hemolytic anemia: report of a

case found

[90787] in a Japanese family. *Acta Haemat. Jpn.* 35: 571–574, 1972.

[90788] 16054.

[90789] Ricciuti, F. C.; Ruddle, F. H.: Assignment of three gene loci
[90790] (PGK, HGPRT, and G6PD) to the long arm of the human X-
chromosome by

[90791] somatic cell genetics. *Genetics* 74: 661–678, 1973.

[90792] 16055.

[90793] Rosa, R.; George, C.; Fardeau, M.; Calvin, M. C.; Rapin, M.;
Rosa,

[90794] J.: A new case of phosphoglycerate kinase deficiency: PGK
Creteil

[90795] associated with rhabdomyolysis and lacking hemolytic
anemia. *Blood* 60:

[90796] 84–91, 1982.

[90797] 16056.

[90798] Saidi, N.; Hors-Cayla, M. C.; Van Cong, N.; Benne, F.:
Sheep

[90799] gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

- [90800] Cell Genet. 25: 200, 1979.
- [90801] 16057.
- [90802] Schwab, A. J.; Krietsch, W. K. G.: Linkage between phosphoglycerate
- [90803] kinase and Xg in a large German kindred. Hum. Genet. 38: 217-221,
- [90804] 1977.
- [90805] 16058.
- [90806] Shows, T. B.; Brown, J. A.: Human X-linked genes regionally mapped
- [90807] utilizing X-autosome translocations and somatic cell hybrids. Proc.
- [90808] Nat. Acad. Sci. 72: 2125-2129, 1975.
- [90809] 16059.
- [90810] Sugie, H.; Sugie, Y.; Ito, M.; Fukuda, T.: A novel missense mutation
- [90811] (837T-C) in the phosphoglycerate kinase gene of a patient with a myopathic
- [90812] form of phosphoglycerate kinase deficiency. J. Child Neurol. 13:
- [90813] 95-97, 1998.

- [90814] 16060.
- [90815] Buj-Bello, A.; Biancalana, V.; Moutou, C.; Laporte, J.; Mandel,
- [90816] J.-L.: Identification of novel mutations in the MTM1 gene causing
- [90817] severe and mild forms of X-linked myotubular myopathy. Hum. Mutat. 14:
- [90818] 320-325, 1999.
- [90819] 16061.
- [90820] Cui, X.; De Vivo, I.; Slany, R.; Miyamoto, A.; Firestein, R.;
- [90821] Cleary, M. L.: Association of SET domain and myotubularin-related
- [90822] proteins modulates growth control. Nature Genet. 18: 331-337, 1998.
- [90823] 16062.
- [90824] Dickeson, S. K.; Helmkamp, G. M., Jr.; Yarbrough, L. R.: Sequence
- [90825] of a human cDNA encoding phosphatidylinositol transfer protein and
- [90826] occurrence of a related sequence in widely divergent eukaryotes. Gene 142:

- [90827] 301–305, 1994.
- [90828] 16063.
- [90829] Dickeson, S. K.; Lim, C. N.; Schuyler, G. T.; Dalton, T. P.;
Helmkamp,
- [90830] G. M., Jr.; Yarbrough, L. R.: Isolation and sequence of
cDNA clones
- [90831] encoding rat phosphatidylinositol transfer protein. J. Biol.
Chem. 264:
- [90832] 16557–16564, 1989.
- [90833] 16064.
- [90834] Hay, J. C.; Martin, T. F. J.: Phosphatidylinositol transfer
protein
- [90835] required for ATP–dependent priming of $\text{Ca}(2+)$ –activated
secretion. Nature 366:
- [90836] 572–575, 1993.
- [90837] 16065.
- [90838] Vihtelic, T. S.; Goebel, M.; Milligan, S.; O'Tousa, J. E.; Hyde,
- [90839] D. R.: Localization of Drosophila retinal degeneration B, a
membrane–associated
- [90840] phosphatidylinositol transfer protein. J. Cell Biol. 122:
1013–1022,

- [90841] 1993.
- [90842] 16066.
- [90843] Wirtz, K. W. A.: Phospholipid transfer proteins. *Annu. Rev. Biochem.* 60:
- [90844] 73–99, 1991.
- [90845] 16067.
- [90846] Fitzgibbon, J.; Pilz, A.; Gayther, S.; Appukuttan, B.; Dulai, K.
- [90847] S.; Delhanty, J. D. A.; Helmkamp, G. M., Jr.; Yarbrough, L. R.; Hunt, D. M.: Localization of the gene encoding human phosphatidylinositol
- [90848] transfer protein (PITPN) to 17p13.3: a gene showing homology to the
- [90849] *Drosophila* retinal degeneration B gene (rdgB). *Cytogenet. Cell Genet.* 67:
- [90850] 205–207, 1994.
- [90851] 16068.
- [90852] Hamilton, B. A.; Smith, D. J.; Mueller, K. L.; Kerrebrock, A. W.;
- [90853] Bronson, R. T.; van Berkel, V.; Daly, M. J.; Kruglyak, L.;

Reeve,

[90855] M. P.; Nemhauser, J. L.; Hawkins, T. L.; Rubin, E. M.; Lander, E.

[90856] S.: The vibrator mutation causes neurodegeneration via reduced expression

[90857] of P1TP- α : positional complementation cloning and extragenic suppression. *Neuron* 18:

[90858] 711-722, 1997.

[90859] 16069.

[90860] Ishida, N.; Miura, N.; Yoshioka, S.; Kawakita, M.: Molecular cloning

[90861] and characterization of a novel isoform of the human UDP-galactose

[90862] transporter, and of related complementary DNAs belonging to the nucleotide-sugar

[90863] transporter gene family. *J. Biochem.* 120: 1074-1078, 1996.

[90864] 16070.

[90865] Sugie, H.; Sugie, Y.; Nishida, M.; Ito, M.; Tsurui, S.; Suzuki,

[90866] M.; Miyamoto, R.; Igarashi, Y.: Recurrent myoglobinuria in a child

[90867] with mental retardation: phosphoglycerate kinase defi-

ciency. J.

[90868] Child Neurol. 4: 95–99, 1989.

[90869] 16071.

[90870] Sugie, H.; Sugie, Y.; Tsurui, S.; Ito, M.: Phosphoglycerate kinase

[90871] deficiency. (Letter) Neurology 44: 1364–1365, 1994.

[90872] 16072.

[90873] Tani, K.; Takizawa, T.; Yoshida, A.: Normal mRNA content in a

[90874] phosphoglycerate kinase variant with severe enzyme deficiency. Am.

[90875] J. Hum. Genet. 37: 931–937, 1985.

[90876] 16073.

[90877] Tonin, P.; Shanske, S.; Miranda, A. F.; Brownell, A. K.; Wyse,

[90878] J. P.; Tsujino, S.; Di Mauro, S.: Phosphoglycerate kinase deficiency:

[90879] biochemical and molecular genetic studies in a new myopathic variant. Neurology 43:

[90880] 387–391, 1993.

[90881] 16074.

- [90882] Valentin, C.; Birgens, H.; Craescu, C. T.; Brodum-Nielsen, K.;
- [90883] Cohen-Solal, M.: A phosphoglycerate kinase mutant (PGK Herlev; D285V)
- [90884] in a Danish patient with isolated chronic hemolytic anemia: mechanism
- [90885] of mutation and structure-function relationships. Hum. Mutat. 12:
- [90886] 280-287, 1998.
- [90887] 16075.
- [90888] Valentine, W. N.; Hsieh, H.-S.; Paglia, D. E.; Anderson, H. M.;
- [90889] Baughan, M. A.; Jaffe, E. R.; Garson, O. M.: Hereditary hemolytic
- [90890] anemia associated with phosphoglycerate kinase deficiency in erythrocytes
- [90891] and leukocytes: a probable X-chromosome-linked syndrome. New Eng.
- [90892] J. Med. 280: 528-534, 1969.
- [90893] 16076.
- [90894] Yoshida, A.; Miwa, S.: Characterization of a phosphoglycerate

- [90895] kinase variant associated with hemolytic anemia. Am. J. Hum. Genet. 26: 378–384, 1974.
- [90896] 16077.
- [90897] Yoshida, A.; Twele, T. W.; Dave, V.; Beutler, E.: Molecular abnormality of a phosphoglycerate kinase variant (PGK–Alabama). Blood Cells Mol. Dis. 21: 179–181, 1995.
- [90900] 16078.
- [90901] Yoshida, A.; Watanabe, S.; Chen, S.–H.; Giblett, E. R.; Malcolm, L. A.: Human phosphoglycerate kinase II: structure of a variant enzyme. J. Biol. Chem. 247: 446–449, 1972.
- [90902] 16079.
- [90903] Becker, M. A.; Kostel, P. J.; Meyer, L. J.; Seegmiller, J. E.: Human phosphoribosylpyrophosphate synthetase: increased enzyme specific activity in a family with gout and excessive purine synthesis. Proc.

- [90909] Nat. Acad. Sci. 70: 2749–2752, 1973.
- [90910] 16080.
- [90911] Becker, M. A.; Losman, M. J.; Rosenberg, A. L.; Mehlman, I.; Levinson, D. J.; Holmes, E. W.: Phosphoribosylpyrophosphate synthetase superactivity: a study of five patients with catalytic defects in the enzyme. Arthritis Rheum. 29: 880–888, 1986.
- [90912] 16081.
- [90913] Becker, M. A.; Losman, M. J.; Wilson, J.; Simmonds, H. A.: Superactivity of human phosphoribosyl pyrophosphate synthetase due to altered regulation by nucleotide inhibitors and inorganic phosphate. Biochim. Biophys. Acta 882: 168–176, 1986.
- [90914] 16082.
- [90915] Becker, M. A.; Meyer, L. J.; Seegmiller, J. E.: Gout with purine overproduction due to increased phosphoribosylphos-

phate synthetase

[90923] activity. Am. J. Med. 55: 232-242, 1973.

[90924] 16083.

[90925] Becker, M. A.; Meyer, L. J.; Wood, A. W.; Seegmiller, J. E.:
Purine

[90926] overproduction in man associated with increased phosphoribosylpyrophosphate

[90927] synthetase activity. Science 179: 1123-1126, 1973.

[90928] 16084.

[90929] Becker, M. A.; Puig, J. G.; Mateos, F. A.; Jimenez, M. L.;
Kim,

[90930] M.; Simmonds, H. A.: Inherited superactivity of phosphoribosylpyrophosphate

[90931] synthetase: association of uric acid overproduction and
sensorineural

[90932] deafness. Am. J. Med. 85: 383-390, 1988.

[90933] 16085.

[90934] Becker, M. A.; Raivio, K. O.; Bakay, B.; Adams, W. B.; Nyhan, W.

[90935] L.: Variant human phosphoribosylpyrophosphate synthetase altered

[90936] in regulatory and catalytic functions. J. Clin. Invest. 65:
109–120,

[90937] 1980.

[90938] 16086.

[90939] Becker, M. A.; Smith, P. R.; Taylor, W.; Mustafi, R.; Switzer,

[90940] R. L.: The genetic and functional basis of purine nucleotide feedback-resistant

[90941] phosphoribosylpyrophosphate synthetase superactivity. J. Clin. Invest. 96:

[90942] 2133–2141, 1995.

[90943] 16087.

[90944] Becker, M. A.; Taylor, W.; Smith, P. R.; Ahmed, M.: Overexpression

[90945] of the normal phosphoribosylpyrophosphate synthetase 1 isoform underlies

[90946] catalytic superactivity of human phosphoribosylpyrophosphate synthetase. J.

[90947] Biol. Chem. 271: 19894–19899, 1996.

[90948] 16088.

[90949] Becker, M. A.; Yen, R. C. K.; Goss, S. J.; Seegmiller, J. E.;

- [90950] Itkin, P.; Lazar, C.; Adams, W. B.: Localization of the structural
- [90951] gene for human phosphoribosylpyrophosphate synthetase on the X-chromosome.
- [90952] (Abstract) Clin. Res. 26: 500A, 1978.
- [90953] 16089.
- [90954] Becker, M. A.; Yen, R. C. K.; Itkin, P.; Goss, S. J.; Seeg-
miller,
- [90955] J. E.; Bakay, B.: Regional localization of the gene for hu-
man phosphoribosylpyrophosphate
- [90956] synthetase on the X-chromosome. Science 203:
1016-1019, 1979.
- [90957] 16090.
- [90958] Christen, H.-J.; Hanefeld, F.; Duley, J. A.; Simmonds, H. A.:
- [90959] Distinct neurological syndrome in two brothers with hype-
ruricaemia.
- [90960] (Letter) Lancet 340: 1167-1168, 1992.
- [90961] 16091.
- [90962] De Vries, A.; Sperling, O.: Familial gouty malignant uric
acid
- [90963] lithiasis due to mutant phosphoribosylpyrophosphatase

synthetase. Der

[90964] Urologe 12: 153–157, 1973.

[90965] 16092.

[90966] Iinuma, K.; Wada, Y.; Onuma, A.; Tanabu, M.: Electroencephalographic

[90967] study of an infant with phosphoribosylpyrophosphate synthetase deficiency. Tohoku

[90968] J. Exp. Med. 116: 53–55, 1975.

[90969] 16093.

[90970] Lebo, R. V.; Martin, D. W., Jr.: Electrophoretic heterogeneity

[90971] of 5–phosphoribosyl–1–pyrophosphate synthetase within and among humans. Biochem.

[90972] Genet. 16: 905–916, 1978.

[90973] 16094.

[90974] Nyhan, W. L.; James, J. A.; Teberg, A. J.; Sweetman, L.; Nelson,

[90975] L. G.: A new disorder of purine metabolism with behavioral manifestations. J.

[90976] Pediat. 74: 20–27, 1969.

[90977] 16095.

- [90978] Riccardi, V. M.: Personal Communication. Denver, Colo. 1974.
- [90979] 16096.
- [90980] Roessler, B. J.; Bell, G.; Heidler, S.; Seino, S.; Becker, M.;
- [90981] Palella, T. D.: Cloning of two distinct copies of human phosphoribosylpyrophosphate
- [90982] synthetase cDNA. Nucleic Acids Res. 18: 193 only, 1990.
- [90983] 16097.
- [90984] Roessler, B. J.; Nosal, J. M.; Smith, P. R.; Heidler, S. A.; Palella,
- [90985] T. D.; Switzer, R. L.; Becker, M. A.: Human X-linked phosphoribosylpyrophosphate
- [90986] synthetase superactivity is associated with distinct point mutations
- [90987] in the PRPS1 gene. J. Biol. Chem. 268: 26476-26481, 1993.
- [90988] 16098.
- [90989] Lund, A.; Udd, B.; Juvonen, V.; Andersen, P. M.; Ced-erquist, K.;
- [90990] Davis, M.; Gellera, C.; Kolmel, C.; Ronnevi, L.-O.; Sperfeld, A.-D.;

- [90991] Sorensen, S.-A.; Tranebjaerg, L.; Van Maldergem, L.; Watanabe, M.;
- [90992] Weber, M.; Yeung, L.; Savontaus, M.-L.: Multiple founder effects
- [90993] in spinal and bulbar muscular atrophy (SBMA, Kennedy disease) around
- [90994] the world. *Europ. J. Hum. Genet.* 9: 431–436, 2001.
- [90995] 16099.
- [90996] Edelhoff, S.; Ayer, D. E.; Zervos, A. S.; Steingrimsson, E.; Jenkins,
- [90997] N. A.; Copeland, N. G.; Eisenman, R. N.; Brent, R.; Distech, C. M.
- [90998] : Mapping of two genes encoding members of a distinct subfamily of
- [90999] MAX interacting proteins: MAD to human chromosome 2 and mouse chromosome
- [91000] 6, and MXI1 to chromosome 10 and mouse chromosome 19. *Oncogene* 9:
- [91001] 665–668, 1994.
- [91002] 16100.
- [91003] Ferguson, G. D.; Anagnostaras, S. G.; Silva, A. J.; Herschman,

- [91004] H. R.: Deficits in memory and motor performance in synaptotagmin
- [91005] IV mutant mice. *Proc. Nat. Acad. Sci.* 97: 5598–5603, 2000.
- [91006] 16101.
- [91007] Ferguson, G. D.; Chen, X.-N.; Korenberg, J. R.; Herschman, H. R.
- [91008] : The human synaptotagmin IV gene defines an evolutionary break point
- [91009] between syntenic mouse and human chromosome regions but retains ligand
- [91010] inducibility and tissue specificity. *J. Biol. Chem.* 275: 36920–36926,
- [91011] 2000.
- [91012] 16102.
- [91013] Holterhus, P.-M.; Wiebel, J.; Sinnecker, G. H. G.; Bruggenwirth,
- [91014] H. T.; Sippell, W. G.; Brinkmann, A. O.; Kruse, K.; Hiort, O.: Clinical
- [91015] and molecular spectrum of somatic mosaicism in androgen insensitivity
- [91016] syndrome. *Pediat. Res.* 46: 684–690, 1999.

- [91017] 16103.
- [91018] Foresta, C.; Bertella, A.; Moro, E.; Roverato, A.; Merico, M.;
- [91019] Ferlin, A.: Sertoli cell function in infertile patients with and
- [91020] without microdeletions of the azoospermia factors on the
Y chromosome
- [91021] long arm. J. Clin. Endocr. Metab. 86: 2414–2419, 2001.
- [91022] 16104.
- [91023] Kobayashi, K.; Mizuno, K.; Hida, A.; Komaki, R.; Tomita,
K.; Matsushita,
- [91024] I.; Namiki, M.; Iwamoto, T.; Tamura, S.; Minowada, S.;
- Nakahori, Y.;
- [91025] Nakagome, Y.: PCR analysis of the Y chromosome long
arm in azoospermic
- [91026] patients: evidence for a second locus required for sper-
matogenesis. Hum.
- [91027] Molec. Genet. 3: 1965–1967, 1994.
- [91028] 16105.
- [91029] Agulnik, A. I.; Zharkikh, A.; Boettger–Tong, H.; Bourgeron,
T.;
- [91030] McElreavey, K.; Bishop, C. E.: Evolution of the DAZ gene
family suggests

- [91031] that Y-linked DAZ plays little, or a limited, role in spermatogenesis
- [91032] but underlines a recent African origin for human populations. Hum.
- [91033] Molec. Genet. 7: 1371–1377, 1998.
- [91034] 16106.
- [91035] Anonymous: A missing piece on the Y. (Editorial) Nature Genet. 10:
- [91036] 367–368, 1995.
- [91037] 16107.
- [91038] Cooke, H. J.; Lee, M.; Kerr, S.; Ruggiu, M.: A murine homologue
- [91039] of the human DAZ gene is autosomal and expressed only in male and
- [91040] female gonads. Hum. Molec. Genet. 5: 513–516, 1996.
- [91041] 16108.
- [91042] Foresta, C.; Moro, E.; Garolla, A.; Onisto, M.; Ferlin, A.: Y
- [91043] chromosome microdeletions in cryptorchidism and idiopathic infertility. J.
- [91044] Clin. Endocr. Metab. 84: 3660–3665, 1999.

- [91045] 16109.
- [91046] Makova, K. D.; Li, W.-H.: Strong male-driven evolution of
DNA
[91047] sequences in humans and apes. *Nature* 416: 624–626,
2002.
- [91048] 16110.
- [91049] Menke, D. B.; Mutter, G. L.; Page, D. C.: Expression of
DAZ, an
[91050] azoospermia factor candidate, in human spermatogonia.
(Letter) *Am.*
- [91051] *J. Hum. Genet.* 60: 237–241, 1997.
- [91052] 16111.
- [91053] Moro, E.; Ferlin, A.; Yen, P. H.; Franchi, P. G.; Palka, G.;
Foresta,
[91054] C.: Male infertility caused by a de novo partial deletion of
the
[91055] DAZ cluster on the Y chromosome. *J. Clin. Endocr. Metab.*
85: 4069–4073,
[91056] 2000.
- [91057] 16112.
- [91058] Reijo, R.; Lee, T.-Y.; Salo, P.; Alagappan, R.; Brown, L. G.;

- [91059] Rosenberg, M.; Rozen, S.; Jaffe, T.; Straus, D.; Hovatta, O.;
de la
- [91060] Chapelle, A.; Silber, S.; Page, D. C.: Diverse spermatogenic
defects
- [91061] in humans caused by Y chromosome deletions encom-
passing a novel RNA-binding
- [91062] protein gene. *Nature Genet.* 10: 383–393, 1995.
- [91063] 16113.
- [91064] Reijo, R.; Seligman, J.; Dinulos, M. B.; Jaffe, T.; Brown, L.
- [91065] G.; Disteche, C. M.; Page, D. C.: Mouse autosomal ho-
molog of DAZ,
- [91066] a candidate male sterility gene in humans, is expressed in
male germ
- [91067] cells before and after puberty. *Genomics* 35: 346–352,
1996.
- [91068] 16114.
- [91069] Saxena, R.; Brown, L. G.; Hawkins, T.; Alagappan, R. K.;
Skaletsky,
- [91070] H.; Reeve, M. P.; Reijo, R.; Rozen, S.; Dinulos, M. B.; Dis-
teche,
- [91071] C. M.; Page, D. C.: The DAZ gene cluster on the human Y
chromosome

- [91072] arose from an autosomal gene that was transposed, repeatedly amplified
- [91073] and pruned. *Nature Genet.* 14: 292–299, 1996.
- [91074] 16115.
- [91075] Najmabadi, H.; Huang, V.; Yen, P.; Subbarao, M. N.; Bhasin, D;
- [91076] Banaag, L.; Naseeruddin, S.; de Kretser, D. M.; Baker, H. W. G.; McLachlan,
- [91077] R. I.; Loveland, K. A.; Bhasin, S.: Substantial prevalence of microdeletions
- [91078] of the Y-chromosome in infertile men with idiopathic azoospermia and
- [91079] oligozoospermia detected using a sequence-tagged site-based mapping
- [91080] strategy. *J. Clin. Endocr. Metab.* 81: 1347–1352, 1996.
- [91081] 16116.
- [91082] Wang, G.; Sawai, N.; Kotliarova, S.; Kanazawa, I.; Nukina, N.:
- [91083] Ataxin-3, the MJD1 gene product, interacts with the two human homologs
- [91084] of yeast DNA repair protein RAD23, HHR23A and HHR23B. *Hum. Molec.*

- [91085] Genet. 9: 1795–1803, 2000.
- [91086] 16117.
- [91087] Ng, J. M. Y.; Vrieling, H.; Sugasawa, K.; Ooms, M. P.;
Grootegoed,
- [91088] J. A.; Vreeburg, J. T. M.; Visser, P.; Beems, R. B.; Gorgels,
T. G.
- [91089] M. F.; Hanaoka, F.; Hoeijmakers, J. H. J.; van der Horst, G.
T. J.
- [91090] : Developmental defects and male sterility in mice lacking
the ubiquitin-like
- [91091] DNA repair gene mHR23B. Molec. Cell. Biol. 22:
1233–1245, 2002.
- [91092] 16118.
- [91093] Volker, M.; Mone, M. J.; Karmakar, P.; van Hoffen, A.;
Schul, W.;
- [91094] Vermeulen, W.; Hoeijmakers, J. H. J.; van Driel, R.; van
Zeeland,
- [91095] A. A.; Mullenders, L. H. F.: Sequential assembly of the nu-
cleotide
- [91096] excision repair factors in vivo. Molec. Cell 8: 213–224,
2001.
- [91097] 16119.

- [91098] Jernstrom, H.; Lerman, C.; Ghadirian, P.; Lynch, H. T.; Weber,
B.; Garber, J.; Daly, M.; Olopade, O. I.; Foulkes, W. D.;
Warner,
E.; Brunet, J.-S.; Narod, S. A.: Pregnancy and risk of early
breast
cancer in carriers of BRCA1 and BRCA2. Lancet 354:
1846–1850, 1999.
16120.
- [91103] Khittoo, G.; Manning, A.; Mustun, H.; Appadoo, J.;
Venkatasamy,
S.; Fagoonee, I.; Ghadirian, P.; Tonin, P. N.: Mutation anal-
ysis
of a Mauritian hereditary breast cancer family reveals the
BRCA2 6503delTT
mutation previously found to recur in different ethnic
populations. Hum.
Hered. 52: 55–58, 2001.
16121.
- [91109] Kinzler, K. W.; Vogelstein, B.: Gatekeepers and caretakers.
Nature 386:
761–763, 1997.

- [91111] 16122.
- [91112] Carrie, A.; Piccolo, F.; Leturcq, F.; de Toma, C.; Azibi, K.; Beldjord, C.; Vallat, J.-M.; Merlini, L.; Voit, T.; Sewry, C.; Urtizberea, J.
- [91114] A.; Romero, N.; Tome, F. M. S.; Fardeau, M.; Sunada, Y.; Campbell, K. P.; Kaplan, J.-C.; Jeanpierre, M.: Mutational diversity and hot spots in the alpha-sarcoglycan gene in autosomal recessive muscular dystrophy (LGMD2D). J. Med. Genet. 34: 470-475, 1997.
- [91118] 16123.
- [91119] Coral-Vazquez, R.; Cohn, R. D.; Moore, S. A.; Hill, J. A.; Weiss, R. M.; Davisson, R. L.; Straub, V.; Barresi, R.; Bansal, D.; Hrstka, R. F.; Williamson, R.; Campbell, K. P.: Disruption of the sarcoglycan-sarcospan complex in vascular smooth muscle: a novel mechanism for cardiomyopathy and muscular dystrophy. Cell 98: 465-474, 1999.

[91124] 16124.

[91125] Duggan, D. J.; Gorospe, J. R.; Fanin, M.; Hoffman, E. P.;
Angelini,

[91126] C.: Mutations in the sarcoglycan genes in patients with
myopathy. New

[91127] Eng. J. Med. 336: 618–624, 1997.

[91128] 16125.

[91129] Fadic, R.; Sunada, Y.; Waclawik, A. J.; Buck, S.; Lewandoski,
P.

[91130] J.; Campbell, K. P.; Lotz, B. P.: Deficiency of a dystrophin–
associated

[91131] glycoprotein (adhalin) in a patient with muscular dystro-
phy and cardiomyopathy. New

[91132] Eng. J. Med. 334: 362–366, 1996.

[91133] 16126.

[91134] Kawai, H.; Akaike, M.; Endo, T.; Adachi, K.; Inui, T.; Mitsui,

[91135] T.; Kashiwagi, S.; Fujiwara, T.; Okuno, S.; Shin, S.; Miyoshi,
K.;

[91136] Campbell, K. P.; Yamada, H.; Shimizu, T.; Matsumura, K.;
Saito, S.

[91137] : Adhalin gene mutations in patients with autosomal re-

cessive childhood

[91138] onset muscular dystrophy with adhalin deficiency. J. Clin. Invest. 96:

[91139] 1202–1207, 1995.

[91140] 16127.

[91141] Ljunggren, A.; Duggan, D.; McNally, E.; Boylan, K. B.; Gama, C.

[91142] H.; Kunkel, L. M.; Hoffman, E. P.: Primary adhalin deficiency as

[91143] a cause of muscular dystrophy in patients with normal dystrophin. Ann.

[91144] Neurol. 38: 367–372, 1995.

[91145] 16128.

[91146] McNally, E. M.; Yoshida, M.; Mizuno, Y.; Ozawa, E.; Kunkel, L.

[91147] M.: Human adhalin is alternatively spliced and the gene is located

[91148] on chromosome 17q21. Proc. Nat. Acad. Sci. 91: 9690–9694, 1994.

[91149] 16129.

[91150] Passos–Bueno, M. R.; Oliveira, J. R.; Bakker, E.; Anderson,

R.

[91151] D.; Marie, S. K.; Vainzof, M.; Roberds, S. L.; Campbell, K. P.; Zatz,

[91152] M.: Genetic heterogeneity for Duchenne-like muscular dystrophy (DLMD)

[91153] based on linkage and 50 DAG analysis. Hum. Molec. Genet. 2: 1945–1947,

[91154] 1993.

[91155] 16130.

[91156] Passos Bueno, M. R.; Moreira, E. S.; Vainzof, M.; Chamberlain,

[91157] J.; Marie, S. K.; Pereira, L.; Akiyama, J.; Roberds, S. L.; Campbell,

[91158] K. P.; Zatz, M.: A common missense mutation in the adhalin gene in

[91159] three unrelated Brazilian families with a relatively mild form of

[91160] autosomal recessive limb-girdle muscular dystrophy. Hum. Molec. Genet. 4:

[91161] 1163–1167, 1995.

[91162] 16131.

[91163] Piccolo, F.; Roberds, S. L.; Jeanpierre, M.; Leturcq, F.; Az-

ibi,

[91164] K.; Beldjord, C.; Carrie, A.; Recan, D.; Chaouch, M.; Reghis, A.;

[91165] El Kerch, F.; Sefiani, A.; Voit, T.; Merlini, L.; Collin, H.; Ey-
mard,

[91166] B.; Beckmann, J. S.; Romero, N. B.; Tome, F. M. S.; Fardeau, M.; Campbell,

[91167] K. P.; Kaplan, J.-C.: Primary adhalinopathy: a common
cause of autosomal

[91168] recessive muscular dystrophy of variable severity. *Nature*
Genet. 10:

[91169] 243–245, 1995.

[91170] 16132.

[91171] Roberds, S. L.; Anderson, R. D.; Ibraghimov-Beskrovnaya, O.; Campbell,

[91172] K. P.: Primary structure and muscle-specific expression on
the 50-kDa

[91173] dystrophin-associated glycoprotein (adhalin). *J. Biol.*
Chem. 268:

[91174] 23739–23742, 1993.

[91175] 16133.

[91176] Roberds, S. L.; Leturcq, F.; Allamand, V.; Piccolo, F.; Jean-

pierre,

[91177] M.; Anderson, R. D.; Lim, L. E.; Lee, J. C.; Tome, F. M. S.; Romero,

[91178] N. B.; Fardeau, M.; Beckmann, J. S.; Kaplan, J.-C.; Campbell, K. P.

[91179] : Missense mutations in the adhalin gene linked to autosomal recessive

[91180] muscular dystrophy. Cell 78: 625–633, 1994.

[91181] 16134.

[91182] Romero, N. B.; Tome, F. M. S.; Leturcq, F.; El Kerch, F.; Azibi,

[91183] K.; Bachner, L.; Anderson, R. D.; Roberds, S. L.; Campbell, K. P.;

[91184] Fardeau, M.; Kaplan, J.-C.: Genetic heterogeneity of severe childhood

[91185] autosomal recessive muscular dystrophy with adhalin (50 kDa dystrophin-associated

[91186] glycoprotein) deficiency. Comp. Rend. Acad. Sci. (Paris) 317: 70–76,

[91187] 1994.

[91188] 16135.

[91189] Schara, U.; Gencik, M.; Mortier, J.; Langen, M.; Gencikova,

A.;

[91190] Epplen, J. T.; Mortier, W.: Alpha-sarcoglycanopathy previously misdiagnosed

[91191] as Duchenne muscular dystrophy: implications for current diagnostics

[91192] and patient care. *Europ. J. Pediat.* 160: 452–453, 2001.

[91193] 16136.

[91194] Perelman, B.; Dafni, N.; Naiman, T.; Eli, D.; Yaakov, M.; Yang

[91195] Feng, T. L.; Sinha, S.; Weber, G.; Khodaei, S.; Sancar, A.; Dotan,

[91196] I.; Canaani, D.: Molecular cloning of a novel human gene encoding

[91197] a 63-kDa protein and its sublocalization within the 11q13 locus. *Genomics* 41:

[91198] 397–405, 1997.

[91199] 16137.

[91200] Teitz, T.; Penner, M.; Eli, D.; Stark, M.; Bakhanashvili, M.; Naiman,

[91201] T.; Canaani, D.: Isolation by polymerase chain reaction of a cDNA

[91202] whose product partially complements the ultraviolet sen–

sitivity of

[91203] xeroderma pigmentosum group C cells. Gene 87:
295–298, 1990.

[91204] 16138.

[91205] Kramps, T.; Peter, O.; Brunner, E.; Nellen, D.; Froesch, B.;
Chatterjee,

[91206] S.; Murone, M.; Zullig, S.; Basler, K.: Wnt/Wingless signal-
ing requires

[91207] BCL9/legless-mediated recruitment of pygopus to the nu-
clear beta-catenin-TCF

[91208] complex. Cell 109: 47–60, 2002.

[91209] 16139.

[91210] Willis, T. G.; Zalcborg, I. R.; Coignet, L. J. A.; Wlodarska,
M.;

[91211] Stul, D. M.; Jadayel, D. M.; Bastard, C.; Treleaven, J. G.;
Catovsky,

[91212] D.; Silva, M. L. M.; Dyer, M. J. S.: Molecular cloning of
translocation

[91213] t(1;14)(q21;q32) defines a novel gene (BCL9) at chromo-
some 1q21. Blood 91:

[91214] 1873–1881, 1998.

[91215] 16140.

- [91216] Davidson, R. G.; Nitowsky, H. M.; Childs, B.: Demonstration of
- [91217] two populations of cells in the human female heterozygous for glucose-6-phosphate
- [91218] dehydrogenase variants. *Proc. Nat. Acad. Sci.* 50: 481-485, 1963.
- [91219] 16141.
- [91220] De Flora, A.; Morelli, A.; Benatti, U.; Giuntini, P.; Ferraris, A. M.; Galiano, S.; Ravazzolo, R.; Gaetani, G. F.: G6PD
- [91221] Napoli and
- [91222] Ferrara II: two new glucose-6-phosphate dehydrogenase variants having
- [91223] similar characteristics but different intracellular lability and specific
- [91224] activity. *Brit. J. Haemat.* 48: 417-423, 1981.
- [91225] 16142.
- [91226] Dern, R. J.; McCurdy, P. R.; Yoshida, A.: A new structural variant
- [91227] of glucose-6-phosphate dehydrogenase with a high production rate (G6PD
- [91228] Hektoen). *J. Lab. Clin. Med.* 73: 283-290, 1969.

- [91229] 16143.
- [91230] De Vita, G.; Alcalay, M.; Sampietro, M.; Cappelini, M. D.;
Fiorelli,
- [91231] G.; Toniolo, D.: Two point mutations are responsible for
G6PD polymorphism
- [91232] in Sardinia. Am. J. Hum. Genet. 44: 233–240, 1989.
- [91233] 16144.
- [91234] Du, C.; Xu, Y.; Hua, X.; Wu, Q.; Liu, L.; Wu, M.: Studies on
- [91235] erythrocyte glucose–6–phosphate dehydrogenase variants
in Chinese.
- [91236] III: Gd(–) Miaozi–Baisha. Acta Genet. Sinica 11(2):
153–158, 1984.
- [91237] 16145.
- [91238] Du, C.–S.; Hua, X.–Y.; Wu, Q.–L.; Li, C.–Q.; Zheng, J.–F.; Li,
- [91239] H.–L.: Studies on erythrocyte glucose–6–phosphate dehy-
drogenase variants
- [91240] in Chinese. IV. Gd(–) Gaohe associated with paroxysmal
nocturnal hemoglobinuria. Chinese
- [91241] J. Pathophysiol. 1: 12–15, 1985.
- [91242] 16146.

- [91243] Du, C. S.; Xu, Y. K.; Hua, X. Y.; Wu, Q. L.; Liu, L. B.: Glucose-6-phosphate
- [91244] dehydrogenase variants and their frequency in Guangdong, China. Hum.
- [91245] Genet. 80: 385-388, 1988.
- [91246] 16147.
- [91247] Eber, S. W.; Gahr, M.; Schroter, W.: Glucose-6-phosphate dehydrogenase
- [91248] (G6PD) Iserlohn and G6PD Regensburg: two new severe enzyme defects
- [91249] in German families. Blut 51: 109-115, 1985.
- [91250] 16148.
- [91251] Echard, G.; Gillois, M.: G6PD--alpha-GAL-PGK--HPRT syntenry in
- [91252] the rabbit, *Oryctolagus cuniculus*. (Abstract) Cytogenet. Cell Genet. 25:
- [91253] 148-149, 1979.
- [91254] 16149.
- [91255] Elizondo, J.; Saenz, G. F.; Paez, C. A.; Ramon, M.; Garcia, M.;
- [91256] Gutierrez, A.; Estrada, M.: G6PD Puerto Limon: a new defi-

cient variant

[91257] of glucose-6-phosphate dehydrogenase associated with congenital nonspherocytic

[91258] hemolytic anemia. Hum. Genet. 62: 110-112, 1982.

[91259] 16150.

[91260] Engstrom, P. F.; Beutler, E.: G-6-PD Tripler: a unique variant

[91261] associated with chronic hemolytic disease. Blood 36: 10-13, 1970.

[91262] 16151.

[91263] Epstein, C. J.: Mammalian oocytes: X-chromosome activity. Science 163:

[91264] 1078-1079, 1969.

[91265] 16152.

[91266] Ermakov, N. V.; Chernyak, N. B.; Tokarev, Y. N.: Properties of

[91267] new variant of glucose-6-phosphate dehydrogenase (Regar variant):

[91268] glucose metabolism in erythrocytes containing abnormal enzyme. Biokhimiia 48:

[91269] 577-583, 1983.

- [91270] 16153.
- [91271] Estrada, M.; Garcia, M.; Gutierrez, A.; Quintero, I.; Gonzalez,
lez,
- [91272] R.: G6PD Varadero. Vox Sang. 43: 102–104, 1982.
- [91273] 16154.
- [91274] Fairbanks, V. F.; Nepo, A. G.; Beutler, E.; Dickson, E. R.;
Honig,
- [91275] G.: Glucose–6–phosphate dehydrogenase variants: reex-
amination of
- [91276] G6PD Chicago and Cornell and a new variant (G6PD Pea
Ridge) resembling
- [91277] G6PD Chicago. Blood 55: 216–220, 1980.
- [91278] 16155.
- [91279] Faust, C. J.; Levinson, B.; Gitschier, J.; Herman, G. E.: Ex-
tension
- [91280] of the physical map in the region of the mouse X chromo-
some homologous
- [91281] to human Xq28 and identification of an exception to con-
served linkage. Genomics 13:
- [91282] 1289–1295, 1992.
- [91283] 16156.

- [91284] Feldman, R.; Gromisch, D. S.; Luhby, A. L.; Beutler, E.:
Congenital
- [91285] nonspherocytic hemolytic anemia due to glucose-
6-phosphate dehydrogenase
- [91286] East Harlem: a new deficient variant. J. Pediat. 90: 89-91,
1977.
- [91287] 16157.
- [91288] Fernandez, M.; Fairbanks, V. F.: Glucose-6-phosphate de-
hydrogenase
- [91289] deficiency in the Philippines: report of a new variant-
-G6PD Panay. Mayo
- [91290] Clin. Proc. 43: 645-660, 1968.
- [91291] 16158.
- [91292] Hostikka, S. L.; Eddy, R. L.; Byers, M. G.; Hoyhtya, M.;
Shows,
- [91293] T. B.; Tryggvason, K.: Identification of a distinct type IV
collagen
- [91294] alpha chain with restricted kidney distribution and assign-
ment of
- [91295] its gene to the locus of X chromosome-linked Alport syn-
drome. Proc.
- [91296] Nat. Acad. Sci. 87: 1606-1610, 1990.

[91297] 16159.

[91298] Bland, J. H. (General Chairman): Proceedings of seminars
on the

[91299] Lesch-Nyhan syndrome. Fed. Proc. 27: 1017-1112, 1968.

[91300] 16160.

[91301] Ferraris, A. M.; Broccia, G.; Meloni, T.; Forteleoni, G.; Gaetan-
tani,

[91302] G. F.: Glucose-6-phosphate dehydrogenase deficiency
and incidence

[91303] of hematologic malignancy. Am. J. Hum. Genet. 42:
516-520, 1988.

[91304]

[91305] 100. Filosa, E.: Personal Communication. La Jolla, Calif.
1989.

[91306]

[91307] 101. Filosa, S.; Calabro, V.; Vallone, D.; Poggi, V.; Mason,
P.; Pagnini,

[91308] D.; Alfinito, F.; Rotoli, B.; Martini, G.; Luzzatto, L.; Battis-
tuzzi,

[91309] G.: Molecular basis of chronic non-spherocytic haemolytic
anaemia:

[91310] a new G6PD variant (393arg-to-his) with abnormal K(m)
G6P and marked

[91311] in vivo instability. *Brit. J. Haemat.* 80: 111–116, 1992.

[91312]

[91313] 102. Filosa, S.; Giacometti, N.; Wangwei, C.; De Mattia, D.;
Pagnini,

[91314] D.; Alfinito, F.; Schettini, F.; Luzzatto, L.; Martini, G.: So-
matic-cell

[91315] selection is a major determinant of the blood-cell pheno-
type in heterozygotes

[91316] for glucose-6-phosphate dehydrogenase mutations caus-
ing severe enzyme

[91317] deficiency. *Am. J. Hum. Genet.* 59: 887–895, 1996.

[91318]

[91319] 103. Fiorelli, G.; Manoussakis, C.; Sampietro, M.; Pittalis,
S.; Guglielmino,

[91320] C. R.; Cappellini, M. D.: Different polymorphic variants of
glucose-6-phosphate

[91321] dehydrogenase (G6PD) in Italy. *Ann. Hum. Genet.* 53:
229–236, 1989.

[91322]

[91323] 104. Fite, E.; Morell, F.; Zuazu, J.; Julia, A.; Morera, J.: Leucocyte

[91324] glucose-6-phosphate dehydrogenase deficiency and necrotizing pneumonia. Europ.

[91325] J. Resp. Dis. 64: 150-154, 1983.

[91326]

[91327] 105. Francke, U.; Bakay, B.; Connor, J. D.; Coldwell, J. G.; Nyhan,

[91328] W. L.: Linkage relationships of X-linked enzymes glucose-6-phosphate

[91329] dehydrogenase and hypoxanthine guanine phosphoribosyltransferase. Am.

[91330] J. Hum. Genet. 26: 512-522, 1974.

[91331]

[91332] 106. Friedman, M. J.; Trager, W.: The biochemistry of resistance

[91333] to malaria. Sci. Am. 244(3): 154-164, 1981.

[91334]

[91335] 107. Frigerio, R.; Sole, G.; Olla, N.; Lovicu, M.; Passiu, G.; Carcassi,

[91336] U.: Cagliari II: a new G-6-PD variant. Haematologica 72:

241–243,

[91337] 1987.

[91338]

[91339] 108. Fujii, H.; Miwa, S.; Takegawa, S.; Takahashi, K.; Hirono, A.;

[91340] Takizawa, T.; Morisaki, T.; Kanno, H.; Taguchi, T.; Okamura, J.:

[91341] Gd(–) Gifu and Gd(–) Fukuoka: two new variants of glucose-6-phosphate

[91342] dehydrogenase found in Japan. Hum. Genet. 66: 276–278, 1984.

[91343]

[91344] 109. Fujii, H.; Miwa, S.; Tani, K.; Takegawa, S.; Fujinami, N.; Takahashi,

[91345] K.; Nakayama, S.; Konno, M.; Sato, T.: Glucose 6-phosphate dehydrogenase

[91346] variants: a unique variant (G6PD Kobe) showed an extremely increased

[91347] affinity for galactose 6-phosphate and a new variant (G6PD Sapporo)

[91348] resembling G6PD Pea Ridge. Hum. Genet. 58: 405–407, 1981.

[91349]

[91350] 110. Gaetani, G. F.; Galiano, S.; Melani, C.; Miglino, M.;
Forni,

[91351] G. L.; Napoli, G.; Perrone, L.; Ferraris, A. M.: A new glu-
cose-6-phosphate

[91352] dehydrogenase variant with congenital nonspherocytic
hemolytic anemia

[91353] (G6PD Genova): biochemical characterization and mo-
saicism expression

[91354] in the heterozygote. Hum. Genet. 84: 337-340, 1990.

[91355]

[91356] 111. Gahr, M.; Bornhalm, D.; Schroeter, W.: Biochemische
Eigenschaften

[91357] einer neuen Variante des Glucose-
6-phosphadehydrogenase (G6PD) Mangels

[91358] mit Favismus: G6PD Bielefeld. Klin. Wschr. 55: 379-384,
1977.

[91359]

[91360] 112. Gahr, M.; Bornhalm, D.; Schroeter, W.: Haemolytic
anemia due

[91361] to glucose 6-phosphate dehydrogenase (G6PD) deficiency:
demonstration

[91362] of two new biochemical variants, G6PD Hamm and G6PD Tarsus. Brit.

[91363] J. Haemat. 33: 363–370, 1976.

[91364]

[91365] 113. Gahr, M.; Schroeter, W.: Glucose 6–phosphate dehydrogenase (G6PD)

[91366] Hamburg, a new variant with chronic nonspherocytic hemolytic anemia. Europ.

[91367] J. Clin. Invest. 4: 187–191, 1974.

[91368]

[91369] 114. Gahr, M.; Schroeter, W.; Sturzenegger, M.; Bornhalm, D.; Marti,

[91370] H. R.: Glucose 6–phosphate dehydrogenase (G6PD) deficiency in Switzerland. Helv.

[91371] Paediat. Acta 31: 159–166, 1976.

[91372]

[91373] 115. Geerdink, R. A.; Horst, R.; Staal, G. E.: An Iraqi Jewish family

[91374] with a new red cell glucose 6–phosphate dehydrogenase variant (Gd–Bagdad)

[91375] and kernicterus. Israel J. Med. Sci. 9: 1040–1043, 1973.

[91376]

[91377] 116. Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors-Cayla, M.

[91378] C.; Gillois, M.: Pig gene mapping: synteny, attempt to assign the

[91379] histocompatibility complex (SLA). (Abstract) Cytogenet. Cell Genet. 25:

[91380] 159, 1979.

[91381]

[91382] 117. Gomez-Gallego, F.; Garrido-Pertierra, A.; Bautista, J. M.: Structural

[91383] defects underlying protein dysfunction in human glucose-6-phosphate

[91384] dehydrogenase A- deficiency. J. Biol. Chem. 275: 9256-9262, 2000.

[91385]

[91386] 118. Gonzalez, R.; Estrada, M.; Garcia, M.; Gutierrez, A.: G6PD Ciudad

[91387] de la Habana: a new slow variant with deficiency found in a Cuban

[91388] family. Hum. Genet. 55: 133-135, 1980.

[91389]

[91390] 119. Gonzalez, R.; Wade, M.; Estrada, M.; Svarch, E.;
Colombo, B.

[91391] : G6PD Pinar del Rio: a new variant discovered in a Cuban
family. Biochem.

[91392] Genet. 15: 909–913, 1977.

[91393]

[91394] 120. Goss, S. J.; Harris, H.: Gene transfer by means of cell
fusion.

[91395] I. Statistical mapping of the human X–chromosome by
analysis of radiation

[91396] induced gene segregation. J. Cell Sci. 25: 17–37, 1977.

[91397]

[91398] 121. Gourdin, D.; Vergnes, H.; Bouloux, C.; Ruffie, J.;
Gherardi,

[91399] M.: Polymorphism of erythrocyte G6PD in the baboon. Am.
J. Phys.

[91400] Anthropol. 37: 281–288, 1972.

[91401]

[91402] 122. Gray, G. R.; Stamatoyannopoulos, G.; Naiman, S. C.;
Kliman, M.

- [91403] R.; Klebanoff, S. J.; Austin, T.; Yoshida, A.; Robinson, G. C. G.
- [91404] : Neutrophil dysfunction, chronic granulomatous disease, and non-spherocytic
- [91405] haemolytic anaemia caused by complete deficiency of glucose-6-phosphate
- [91406] dehydrogenase. *Lancet* II: 530-534, 1973.
- [91407]
- [91408] 123. Grossman, A.; Ramanathan, K.; Justice, P.; Gordon, J.; Shahidi,
- [91409] N. T.; Hsia, D. Y. Y.: Congenital nonspherocytic hemolytic anemia
- [91410] associated with erythrocyte G-6-PD deficiency in a Negro family. *Pediatrics* 37:
- [91411] 624-629, 1966.
- [91412]
- [91413] 124. Gutierrez, A.; Garcia, M.; Estrada, M.; Quintero, I.; Gonzalez,
- [91414] R.: Glucose-6-phosphate dehydrogenase (G6PD) Guantano and G6PD
- [91415] Caujeri: two new glucose-6-phosphate dehydrogenase-deficient variants

[91416] found in Cuba. Biochem. Genet. 25: 231–238, 1987.

[91417]

[91418] 125. Hall, K.; Schreeder, M. T.; Prchal, J. T.: G6PD
Huntsville:

[91419] a new glucose–6–phosphate dehydrogenase associated
with chronic hemolytic

[91420] anemia. Hum. Genet. 79: 90–91, 1988.

[91421]

[91422] 126. Harkonen, M.; Vuopio, P.: Red cell glucose–
6–phosphate dehydrogenase

[91423] deficiency in Finland. Ann. Clin. Res. 6: 187–197, 1974.

[91424]

[91425] 127. Harley, J. D.; Agar, N. S.; Yoshida, A.: Glucose
6–phosphate

[91426] dehydrogenase variant Gd(+) Alexandra associated with
neonatal jaundice

[91427] and Gd(–) Camperdown in a young man with lamellar
cataracts. J. Lab.

[91428] Clin. Med. 91: 295–300, 1978.

[91429]

- [91430] 128. Helge, H.; Borner, K.: Kongenitale nichtsphaerozytare haemolytische
- [91431] Anaemie, Katarakt und Glucose-6-phosphat-dehydrogenase-mangel. Dtsch.
- [91432] Med. Wschr. 91: 1584-1589, 1966.
- [91433]
- [91434] 129. Hirono, A.; Beutler, E.: Alternative splicing of human glucose-6-phosphate
- [91435] dehydrogenase messenger RNA in different tissues. J. Clin. Invest. 83:
- [91436] 343-346, 1989.
- [91437]
- [91438] 130. Hirono, A.; Beutler, E.: Molecular cloning and nucleotide sequence
- [91439] of cDNA for human glucose-6-phosphate dehydrogenase variant A(-). Proc.
- [91440] Nat. Acad. Sci. 85: 3951-3954, 1988.
- [91441]
- [91442] 131. Hirono, A.; Fujii, H.; Shima, M.; Miwa, S.: G6PD Nara: a new
- [91443] class 1 glucose-6-phosphate dehydrogenase variant with

an eight amino

[91444] acid deletion. Blood 82: 3250–3252, 1998.

[91445]

[91446] 132. Hirono, A.; Kawate, K.; Honda, A.; Fujii, H.; Miwa, S.:
A single

[91447] mutation 202G→A in the human glucose-6-phosphate de-
hydrogenase gene

[91448] (G6PD) can cause acute hemolysis by itself. (Letter) Blood
99: 1498

[91449] only, 2002.

[91450]

[91451] 133. Hirono, A.; Kuhl, W.; Gelbart, T.; Forman, L.; Fair-
banks, V.

[91452] F.; Beutler, E.: Identification of the binding domain for
NADP(+)

[91453] of human glucose-6-phosphate dehydrogenase by se-
quence analysis of

[91454] mutants. Proc. Nat. Acad. Sci. 86: 10015–10017, 1989.

[91455]

[91456] 134. Hitzeroth, H. W.; Bender, K.: Age-dependency of so-
matic selection

[91457] in South African Negro G-6-PD heterozygotes. Hum.
Genet. 58: 338-343,

[91458] 1981.

[91459]

[91460] 135. Honig, G. R.; Habacon, E.; Vida, L. N.; Matsumoto, F.;
Beutler,

[91461] E.: Three new variants of glucose-6-phosphate dehydro-
genase associated

[91462] with chronic nonspherocytic hemolytic anemia: G6PD Lin-
coln Park, G6PD

[91463] Arlington Heights, and G6PD West Town. Am. J. Hemat. 6:
353-360,

[91464] 1979.

[91465]

[91466] 136. Hook, E. B.; Stamatoyannopoulos, G.; Yoshida, A.;
Motulsky, A.

[91467] G.: Glucose-6-phosphate dehydrogenase Madrona: a slow
electrophoretic

[91468] glucose-6-phosphate dehydrogenase variant with kinetic
characteristics

[91469] similar to those of normal type. J. Lab. Clin. Med. 72:
404-409,

[91470] 1968.

[91471]

[91472] 137. Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Benne, F.: Cattle

[91473] gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

[91474] Cell Genet. 25: 165-166, 1979.

[91475]

[91476] 138. Howell, E. B.; Nelson, A. J.; Jones, O. W.: A new G-6-PD variant

[91477] associated with chronic non-spherocytic haemolytic
anaemia in a Negro

[91478] family. J. Med. Genet. 9: 160-164, 1972.

[91479]

[91480] 139. Hutz, M. H.; Yoshida, A.; Salzano, F. M.: Three rare
G-6-PD

[91481] variants from Porto Alegre, Brazil. Hum. Genet. 39:
191-197, 1977.

[91482]

[91483] 140. Ishwad, C. S.; Naik, S. N.: A new glucose-6-phosphate dehydrogenase

[91484] variant (G-6-PD Kalyan) found in a Koli family. Hum.

Genet. 66:

[91485] 171-175, 1984.

[91486]

[91487] 141. Johnson, G. J.; Kaplan, M. E.; Beutler, E.: G6PD Long
Prairie:

[91488] a new mutant exhibiting normal sensitivity to inhibition by
NADPH

[91489] and accompanied by nonspherocytic hemolytic anemia.
Blood 49: 247-251,

[91490] 1977.

[91491]

[91492] 142. Johnston, P. G.; VandeBerg, J. L.; Sharman, G. B.: In-
heritance

[91493] of erythrocyte glucose 6-phosphate dehydrogenase in the
red-necked

[91494] wallaby, *Macropus rufogriseus* (Desmarest) consistent
with paternal

[91495] X inactivation. Biochem. Genet. 13: 235-242, 1975.

[91496]

[91497] 143. Junien, C.; Kaplan, J.-C.; Meienhofer, M. C.; Maigret,

P.; Sender,

[91498] A.: G6PD Baudelocque: a new unstable variant characterized in cultured

[91499] fibroblasts. *Enzyme* 18: 48–59, 1974.

[91500]

[91501] 144. Kaeda, J. S.; Chhotray, G. P.; Ranjit, M. R.; Bautista, J. M.;

[91502] Reddy, P. H.; Stevens, D.; Naidu, J. M.; Britt, R. P.; Vulliamy, T.

[91503] J.; Luzzatto, L.; Mason, P. J.: A new glucose-6-phosphate dehydrogenase

[91504] variant, G6PD Orissa (44 ala-gly), is the major polymorphic variant

[91505] in tribal populations in India. *Am. J. Hum. Genet.* 57: 1335–1341,

[91506] 1995.

[91507]

[91508] 145. Kageoka, T.; Satoh, C.; Goriki, K.; Fujita, M.; Neriishi, S.;

[91509] Yamamura, K.; Kaneko, J.; Masunari, N.: Electrophoretic variants

[91510] of blood proteins in Japanese. IV. Prevalence and enzyme-

logic characteristics

[91511] of G6PD variants in Hiroshima and Nagasaki. Hum. Genet.
70: 101–108,

[91512] 1985.

[91513]

[91514] 146. Kahn, A.; Bernard, J.-F.; Cottreau, D.; Mazie, J.;
Boivin, P.

[91515] : Gd(–) Abrami, a deficient G6PD variant with hemizygous
expression

[91516] in blood cells of a woman with primary myelofibrosis. Hu-
mangenetik 30:

[91517] 41–46, 1975.

[91518]

[91519] 147. Kahn, A.; Boivin, P.; Hakim, J.; Lagneau, J.: Hetero-
geneite

[91520] des glucose-6-phosphate dehydrogenase erythrocytaire
deficitaires

[91521] dans la race noire: etude cinetique et description de deux
nouvelles

[91522] variants Gd(–) Dakar et Gd(–) Mali. Nouv. Rev. Franc.
Hemat. 11:

[91523] 741–758, 1971.

[91524]

[91525] 148. Kahn, A.; Boivin, P.; Lagneau, J.: Phenotypes de la glucose-6-phosphate

[91526] dehydrogenase erythrocytaire dans la race noire. Human-genetik 18:

[91527] 261-270, 1973.

[91528]

[91529] 149. Kahn, A.; Boulard, M.; Hakim, J.; Schaison, G.; Boivin, P.; Bernard,

[91530] J.: Anemie hemolytique congenitale non spherocytaire par deficit

[91531] en glucose 6-phosphate dehydrogenase erythrocytaire: description de

[91532] deux nouvelles variants: Gd(-) Saint Louis (Paris) et Gd(-) Hayem. Nouv.

[91533] Rev. Franc. Hemat. 14: 587-600, 1974.

[91534]

[91535] 150. Kahn, A.; Dao, C.; Cottreau, D.; Bilski-Pasquier, G.: 'Gd(-)

[91536] Hotel Dieu': a new G-6PD variant with chronic hemolysis in a Negro

[91537] patient from Senegal. Hum. Genet. 39: 353–358, 1977.

[91538]

[91539] 151. Kahn, A.; Exters, A.; Habedank, M.: Gd(–) Aachen, a new variant

[91540] of deficient glucose–6–phosphate dehydrogenase. Humangenetik 32:

[91541] 171–180, 1976.

[91542]

[91543] 152. Kahn, A.; Hakim, J.; Cottreau, D.; Boivin, P.: Gd (–) Matam.

[91544] An African glucose 6–phosphate dehydrogenase variant with enzyme deficiency.

[91545] Biochemical and immunological properties in various hemopoietic tissues. Clin.

[91546] Chim. Acta 59: 183–190, 1975.

[91547]

[91548] 153. Kahn, A.; North, M. L.; Cottreau, D.; Giron, G.; Lang, J. M.

[91549] : G6PD Vientiane: a new glucose–6–phosphate dehydrogenase variant

[91550] with increased stability. Hum. Genet. 43: 85–89, 1978.

[91551]

[91552] 154. Kahn, A.; North, M. L.; Messer, J.; Boivin, P.: G-6PD
'Ankara':

[91553] a new G-6PD variant with deficiency found in a Turkish
family. Humangenetik 27:

[91554] 247-250, 1975.

[91555]

[91556] 155. Kanno, H.; Huang, I.-Y.; Kan, Y. W.; Yoshida, A.: Two
structural

[91557] genes on different chromosomes are required for encod-
ing the major

[91558] subunit of human red cell glucose-6-phosphate dehydro-
genase. Cell 58:

[91559] 595-606, 1989.

[91560]

[91561] 156. Kanno, H.; Takano, T.; Fujii, H.; Tani, K.; Morisaki, T.;
Hirono,

[91562] A.; Kumakawa, T.; Ogura, H.; Takahashi, K.; Tsutsumi, H.;
Miwa, S.

[91563] : A new glucose-6-phosphate variant (G6PD Iwate) associ-
ated with congenital

[91564] non-spherocytic hemolytic anemia. Acta Haemat. Jpn. 51:
715-719,

[91565] 1987.

[91566]

[91567] 157. Kaplan, J. C.; Hanzlickova-Leroux, A.; Nicholas, A.
M.; Rosa,

[91568] R.; Weiler, C.; Lepercq, G.: A new glucose-6-phosphate
dehydrogenase

[91569] variant (G6PD Port-Royal). Enzyme 12: 25-32, 1971.

[91570]

[91571] 158. Kaplan, J. C.; Rosa, R.; Seringe, P.; Hoeffel, J. C.: Le
polymorphisme

[91572] genetique de la glucose-6-phosphate deshydrogenase
erythrocytaire

[91573] chez l'homme. II. Etude d'une nouvelle variete a activite
diminuee:

[91574] le type 'Kabyle.'. Enzym. Biol. Clin. 8: 332-340, 1967.

[91575]

[91576] 159. Kaplan, M.; Hammerman, C.; Vreman, H. J.; Steven-
son, D. K.; Beutler,

[91577] E.: Acute hemolysis and severe neonatal hyperbilirubine-

mia in glucose-6-phosphate

[91578] dehydrogenase deficient heterozygotes. J. Pediat. 139:
137-140,

[91579] 2001.

[91580]

[91581] 160. Kaplan, M.; Renbaum, P.; Levy-Lahad, E.; Hammer-
man, C.; Lahad,

[91582] A. Beutler, E.: Gilbert syndrome and glucose-6-phosphate
dehydrogenase

[91583] deficiency: a dose-dependent genetic interaction crucial
to neonatal

[91584] hyperbilirubinemia. Proc. Nat. Acad. Sci. 94:
12128-12132, 1997.

[91585]

[91586] 161. Kappas, A.; Drummond, G. S.; Valaes, T.: A single
dose of Sn-mesoporphyrin

[91587] prevents development of severe hyperbilirubinemia in
glucose-6-phosphate

[91588] dehydrogenase-deficient newborns. Pediatrics 108:
25-30, 2001.

[91589]

[91590] 162. Karadsheh, N. S.; Awidi, A. S.; Tarawneh, M. S.: Two

new glucose-6

[91591] phosphate dehydrogenase (G6PD) variants associated with hemolytic

[91592] anemia: G6PD Amman-1 and G6PD Amman-2. Am. J. Hemat. 22: 185-192,

[91593] 1986.

[91594]

[91595] 163. Kay, A. C.; Kuhl, W.; Prchal, J.; Beutler, E.: The origin of

[91596] glucose-6-phosphate-dehydrogenase (G6PD) polymorphisms in African-Americans. Am.

[91597] J. Hum. Genet. 50: 394-398, 1992.

[91598]

[91599] 164. Kirkman, H. N.; Kidson, C.; Kennedy, M.: Variants of human glucose-6-phosphate

[91600] dehydrogenase. Studies of samples from New Guinea. In: Beutler, E.

[91601] : Hereditary Disorders of Erythrocyte Metabolism. New York: Grune

[91602] and Stratton (pub.) 1968. Pp. 126-145.

[91603]

- [91604] 165. Kirkman, H. N.; Lie-Injo, L. E.: Variants of glucose 6-phosphate
- [91605] dehydrogenase in Indonesia. *Nature* 221: 959, 1969.
- [91606]
- [91607] 166. Kirkman, H. N.; McCurdy, P. R.; Naiman, J. L.: Functionally
- [91608] abnormal glucose-6-phosphate dehydrogenases. *Cold Spring Harbor Symp.*
- [91609] *Quant. Biol.* 29: 391-398, 1964.
- [91610]
- [91611] 167. Kirkman, H. N.; Ramot, B.; Lee, J. T.: Altered aggregational
- [91612] properties in a genetic variant of human glucose-6-phosphate dehydrogenase. *Biochem.*
- [91613] *Genet.* 3: 137-150, 1969.
- [91614]
- [91615] 168. Kirkman, H. N.; Riley, H. D., Jr.: Congenital nonspherocytic
- [91616] hemolytic anemia. *Am. J. Dis. Child.* 102: 313-320, 1961.
- [91617]

[91618] 169. Kirkman, H. N.; Rosenthal, I. M.; Simon, E. B.; Carson, P. E.;

[91619] Brinson, A. G.: 'Chicago I' variant of glucose-6-phosphate dehydrogenase

[91620] in congenital hemolytic disease. J. Lab. Clin. Med. 63: 715-725,

[91621] 1964.

[91622]

[91623] 170. Kirkman, H. N.; Schettini, E.; Pickard, B. M.: Mediter-ranean

[91624] variant of glucose-6-phosphate dehydrogenase. J. Lab. Clin. Med. 63:

[91625] 726-735, 1964.

[91626]

[91627] 171. Kirkman, H. N.; Simon, E. R.; Pickard, B. M.: Seattle variant

[91628] of glucose-6-phosphate dehydrogenase. J. Lab. Clin. Med. 66: 834-840,

[91629] 1965.

[91630]

[91631] 172. Kissin, C.; Cotte, J.: Etude d'un variant de glucose-

6-phosphate

[91632] deshydrogenase: I B type Constantine. Enzyme 11: 277-284, 1970.

[91633]

[91634] 173. Kitao, T.; Ito, K.; Hattori, K.; Matsuki, T.; Yoneyama, Y.:

[91635] G6PD Kanazawa: a new variant of glucose-6-phosphate dehydrogenase

[91636] associated with congenital nonspherocytic hemolytic anemia. Acta

[91637] Haemat. 68: 131-135, 1982.

[91638]

[91639] 174. Kojima, H.: Congenital nonspherocytic hemolytic disease (CNHD)

[91640] due to a G-6-PD variant: G-6-PD Kyoto. Acta Haemat. Jpn. 35: 32-38,

[91641] 1972.

[91642]

[91643] 175. Koliakos, G.; Kalomenopoulou, M.; Grammatikos, P.; Dimitriadou,

[91644] A.; Kouzi-Koliakos, K.; Zacharaki, R.; Skaragas, G.; Kokka,

A.; Trakatellis,

[91645] A.: A new glucose 6-phosphate dehydrogenase variant
(G6PD Thessaloniki)

[91646] in a patient with idiopathic myelofibrosis. Hum. Hered. 39:
141–149,

[91647] 1989.

[91648]

[91649] 176. Krasnopol'skaya, K. D.; Bochkov, N. P.: Genetic het-
erogeneity

[91650] of hereditary enzymopathies. Vestn. Akad. Med. Nauk.
SSSR 9: 56–64,

[91651] 1982.

[91652]

[91653] 177. Krasnopol'skaya, K. D.; Shatskaya, T. L.; Filippov, I.
K.; Annenkov,

[91654] G. A.; Zakharova, T. V.; Mekhtiev, N. K.; Movsum-Zade, K.
M.: Genetic

[91655] heterogeneity of G6PD deficiency: study of mutant alleles
in Shekii

[91656] district of Azerbaijan. Genetika 13: 1455–1461, 1977.

[91657]

- [91658] 178. Kumakawa, T.; Suzuki, S.; Fujii, H.; Miwa, S.: Frequency of
[91659] glucose 6-phosphate dehydrogenase (G6PD) deficiency in
Tokyo and a
[91660] new variant: G6PD Musashino. *Acta Haemat. Jpn.* 50:
25-28, 1987.
[91661]
- [91662] 179. Kurdi-Haidar, B.; Mason, P. J.; Berrebi, A.; Ankra-
Badu, G.;
[91663] Al-Ali, A.; Oppenheim, A.; Luzzatto, L.: Origin and spread
of the
[91664] glucose-6-phosphate dehydrogenase variant
(G6PD-Mediterranean) in
[91665] the Middle East. *Am. J. Hum. Genet.* 47: 1013-1019,
1990.
[91666]
- [91667] 180. Kwiatkowska, J.; Kacprzak-Bergman, I.: New erythro-
cyte glucose
[91668] 6-phosphate dehydrogenase variant. *Acta Haemat.* 46:
188-192, 1971.
[91669]
- [91670] 181. Kwok, C. J.; Martin, A. C. R.; Au, S. W. N.; Lam, V. M.

S.:

- [91671] G6PDdb, an integrated database of glucose-6-phosphate dehydrogenase
- [91672] (G6PD) mutations. Hum. Mutat. 19: 217-224, 2002.
- [91673]
- [91674] 182. Lee, K. T.; Thomas, W. A.; Janakidevi, K.; Kroms, M.; Reiner, J. M.; Borg, K. Y.: Mosaicism in female hybrid hares heterozygous for glucose-6-phosphate dehydrogenase (G-6-PD). I. General properties
- [91677] of a hybrid hare model with special reference to atherogenesis. Exp.
- [91678] Molec. Path. 34: 191-201, 1981.
- [91679]
- [91680] 183. Lenzerini, L.; Meera Khan, P.; Filippi, G.; Rattazzi, M. C.;
- [91681] Rat, A. K.: Characterization of glucose-6-phosphate dehydrogenase
- [91682] variants. I. Occurrence of a G6PD Seattle-like variant in Sardinia
- [91683] and its interaction with G6PD Mediterranean variant. Am.

J. Hum.

[91684] Genet. 21: 142–153, 1969.

[91685]

[91686] 184. Lisker, R.; Linares, C.; Motulsky, A. G.: Glucose-6-phosphate

[91687] dehydrogenase Mexico, a new variant with enzyme deficiency, abnormal

[91688] mobility and absence of hemolysis. J. Lab. Clin. Med. 29: 788–793,

[91689] 1972.

[91690]

[91691] 185. Lisker, R.; Perez-Briceno, R.; Beutler, E.: A new glucose-6-phosphate

[91692] dehydrogenase variant, Gd(–) Tepic, characterized by moderate enzyme

[91693] deficiency and mild episodes of hemolytic anemia. Hum. Genet. 69:

[91694] 19–21, 1985.

[91695]

[91696] 186. Lisker, R.; Perez-Briceno, R.; Rave, V.; Yoshida, A.: Glucosa-6-fosfato

- [91697] deshidrogenasa Gd(-) Distrito Federal: nueva variante asociada a deficiencia
- [91698] enzimatica moderada y anemia hemolitica ocasional. Rev. Invest. Clin. 33:
- [91699] 209-211, 1981.
- [91700]
- [91701] 187. Lisker, R.; Perez Briceno, R.; Agrilar, L.; Yoshida, A.: A variant
- [91702] glucose-6-phosphate dehydrogenase Gd(-) Chiapas associated with moderate
- [91703] enzyme deficiency and occasional hemolytic anemia. Hum. Genet. 43:
- [91704] 81-84, 1978.
- [91705]
- [91706] 188. Lisker, R.; Perez Briceno, R.; Zavala, C.; Navarrette, J. I.;
- [91707] Wessels, M.; Yoshida, A.: A glucose 6-phosphate dehydrogenase Gd(-)
- [91708] Castilla variant characterized by mild deficiency associated with
- [91709] drug-induced hemolytic anemia. J. Lab. Clin. Med. 90: 754-759, 1977.

[91710]

[91711] 189. Liu, Y.; Phelan, J.; Go, R. C. P.; Prchal, J. F.; Prchal, J.

[91712] T.: Rapid determination of clonality by detection of two closely-linked

[91713] X chromosome exonic polymorphisms using allele-specific PCR. J. Clin.

[91714] Invest. 99: 1984-1990, 1997.

[91715]

[91716] 190. Long, W. K.; Kirkman, H. N.; Sutton, H. H.: Electrophoretically

[91717] slow variants of glucose-6-phosphate dehydrogenase from red cells

[91718] of Negroes. J. Lab. Clin. Med. 65: 81-87, 1965.

[91719]

[91720] 191. Longo, L.; Vanegas, O. C.; Patel, M.; Rosti, V.; Li, H.; Waka,

[91721] J.; Merghoub, T.; Pandolfi, P. P.; Notaro, R.; Manova, K.; Luzzatto,

[91722] L.: Maternally transmitted severe glucose 6-phosphate dehydrogenase

[91723] deficiency is an embryonic lethal. EMBO J. 21: 4229-4239,

2002.

[91724]

[91725] 192. Luzzatto, L.: Genetic heterogeneity and pathophysiology of G6PD

[91726] deficiency. Brit. J. Haemat. 28: 151–156, 1974.

[91727]

[91728] 193. Luzzatto, L.: Personal Communication. London, England 4/1990.

[91729]

[91730] 194. Luzzatto, L.; Afolayam, A.: Enzyme properties of different types

[91731] of human erythrocyte glucose–6–phosphate dehydrogenase with characterization

[91732] of two new genetic variants. J. Clin. Invest. 47: 1833–1842, 1968.

[91733]

[91734] 195. Luzzatto, L.; Martini, G.: X-Linked Wiskott–Aldrich syndrome

[91735] in a girl. (Letter) New Eng. J. Med. 338: 1850–1851, 1998.

[91736]

[91737] 196. Luzzatto, L.; Usanga, E. A.; Bienzle, U.; Esan, G. F. J.; Fusuan,

[91738] F. A.: Imbalance in X–chromosome expression: evidence for a human

[91739] X-linked gene affecting growth of hemopoietic cells. Sci–

ence 205:

[91740] 1418–1420, 1979.

[91741]

[91742] 197. Luzzatto, L.; Usanga, E. A.; Reddy, S.: Glucose-6-phosphate

[91743] dehydrogenase deficient red cells: resistance to infection by malarial

[91744] parasites. Science 164: 839–842, 1969.

[91745]

[91746] 198. MacDonald, D.; Town, M.; Mason, P.; Vulliamy, T.; Luzzatto, L.;

[91747] Goff, D. K.: Deficiency in red blood cells. (Letter) Nature 350:

[91748] 115, 1991.

[91749]

[91750] 199. Maeda, M.; Constantoulakis, P.; Chen, C.–S.; Stamatoyannopoulos,

[91751] G.; Yoshida, A.: Molecular abnormalities of a human glucose-6-phosphate

[91752] dehydrogenase variant associated with undetectable enzyme activity

[91753] and immunologically cross-reacting material. Am. J. Hum. Genet. 51:

[91754] 386-395, 1992.

[91755]

[91756] 200. Mallouh, A. A.; Abu-Osba, Y. K.: Bacterial infections in children

[91757] with glucose-6-phosphate dehydrogenase deficiency. J. Pediat. 111:

[91758] 850-852, 1987.

[91759]

[91760] 201. Mamlok, R. J.; Mamlok, V.; Mills, G. C.; Daeschner, C. W., III;

[91761] Schmalstieg, F. C.; Anderson, D. C.: Glucose-6-phosphate dehydrogenase

[91762] deficiency, neutrophil dysfunction and Chromobacterium violaceum sepsis. J.

[91763] Pediat. 111: 852-854, 1987.

[91764]

[91765] 202. Mamlok, R. J.; Mills, G. C.; Goldblum, R. M.; Daeschner, C. W.

[91766] : Glucose-6-phosphate dehydrogenase Beaumont: a new

variant with severe

[91767] enzyme deficiency and chronic nonspherocytic hemolytic anemia. Enzyme 34:

[91768] 15–21, 1985.

[91769]

[91770] 203. Mandelli, F.; Amadori, S.; DeLaurenzi, A.; Kahn, A.; Isacchi,

[91771] G.; Papa, G.: Glucose–6–phosphate dehydrogenase Vellettri: a new variant

[91772] with reduced activity in a patient with congenital nonspherocytic

[91773] haemolytic anemia. Acta Haemat. 57: 121–126, 1977.

[91774]

[91775] 204. Marks, P. A.; Banks, J.; Gross, R.: Genetic heterogeneity of

[91776] glucose–6–phosphate dehydrogenase deficiency. Nature 194: 454–456,

[91777] 1962.

[91778]

[91779] 205. Martin, S. K.; Miller, L. H.; Alling, D.; Okoye, V. C.; Esan,

- [91780] G. J. F.; Osunkoya, B. O.; Deane, M.: Severe malaria and glucose-6-phosphate-dehydrogenase
- [91781] deficiency: a reappraisal of the malaria-G6PD hypothesis. Lancet I:
- [91782] 524-526, 1979.
- [91783]
- [91784] 206. Martini, G.; Toniolo, D.; Vulliamy, T.; Luzzatto, L.;
- Dono, R.;
- [91785] Viglietto, G.; Paonessa, G.; D'Urso, M.; Persico, M. G.: Structural
- [91786] analysis of the X-linked gene encoding human glucose 6-phosphate dehydrogenase. EMBO
- [91787] J. 5: 1849-1855, 1986.
- [91788]
- [91789] 207. Mason, P. J.: New insights into G6PD deficiency. (Annotation) Brit.
- [91790] J. Haemat. 94: 585-591, 1996.
- [91791]
- [91792] 208. Mason, P. J.; Stevens, D. J.; Luzzatto, L.; Brenner, S.;
- Aparicio,
- [91793] S.: Genomic structure and sequence of the Fugu rubripes

glucose-6-phosphate

[91794] dehydrogenase gene (G6PD). Genomics 26: 587-591, 1995.

[91795]

[91796] 209. McCann, S. R.; Smithwick, A. M.; Temperley, I. J.; Tipton, K.

[91797] : G6PD (Dublin): chronic non-spherocytic haemolytic anaemia resulting

[91798] from glucose-6-phosphate dehydrogenase deficiency in an Irish kindred. J.

[91799] Med. Genet. 17: 191-193, 1980.

[91800]

[91801] 210. McCurdy, P. R.: Use of genetic linkage for the detection of

[91802] female carriers of hemophilia. New Eng. J. Med. 285: 218-219, 1971.

[91803]

[91804] 211. McCurdy, P. R.; Blackwell, R. Q.; Todd, D.; Tso, S. C.; Tuchinda,

[91805] S.: Further studies on glucose-6-phosphate dehydrogenase deficiency

[91806] in Chinese subjects. J. Lab. Clin. Med. 75: 788-797, 1970.

[91807]

[91808] 212. McCurdy, P. R.; Kamel, K.; Selim, O.: Heterogeneity of red cell

[91809] glucose 6-phosphate dehydrogenase (G6PD) deficiency in Egypt. J.

[91810] Lab. Clin. Med. 84: 673-680, 1974.

[91811]

[91812] 213. McCurdy, P. R.; Kirkman, H. N.; Naiman, J. L.; Jim, R. T. S.;

[91813] Pickard, B. M.: A Chinese variant of glucose-6-phosphate dehydrogenase. J.

[91814] Lab. Clin. Med. 67: 374-385, 1966.

[91815]

[91816] 214. McCurdy, P. R.; Mahmood, L.: Red cell glucose-6-phosphate dehydrogenase

[91817] deficiency in Pakistan. J. Lab. Clin. Med. 76: 943-948, 1970.

[91818]

[91819] 215. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.: Variants of

[91820] glucose-6-phosphate dehydrogenase (G-6-PD) associated

with G-6-PD

[91821] deficiency in Puerto Ricans. J. Lab. Clin. Med. 82:
432-437, 1973.

[91822]

[91823] 216. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.; Conrad, M. E.

[91824] : Variants of glucose-6-phosphate dehydrogenase
(G-6-PD) associated

[91825] with G-6-PD deficiency in Puerto Ricans. J. Lab. Clin. Med.
82:

[91826] 432-437, 1973.

[91827]

[91828] 217. Meloni, T.; Carta, F.; Forteleoni, G.; Carta, A.; Ena, F.;
Meloni,

[91829] G. F.: Glucose 6-phosphate dehydrogenase deficiency and
cataract

[91830] of patients in northern Sardinia. Am. J. Ophthal. 110:
661-664,

[91831] 1990.

[91832]

[91833] 218. Mentzer, W. C., Jr.; Warner, R.; Addiego, J.; Smith, B.;

Walter,

- [91834] T.: G6PD San Francisco: a new variant of glucose-6-phosphate dehydrogenase
- [91835] associated with congenital nonspherocytic hemolytic anemia. Blood 55:
- [91836] 195–198, 1980.
- [91837]
- [91838] 219. Mesbah–Namin, S. A.; Sanati, M. H.; Mowjoodi, A.; Mason, P. J.;
- [91839] Vulliamy, T. J.; Noori–Dalooi, M. R.: Three major glucose-6-phosphate
- [91840] dehydrogenase–deficient polymorphic variants identified in Mazandaran
- [91841] state of Iran. Brit. J. Haemat. 117: 763–764, 2002.
- [91842]
- [91843] 220. Miller, D. R.; Wollman, M. R.: A new variant of glucose 6–phosphate
- [91844] dehydrogenase deficiency hereditary hemolytic anemia, G6PD Cornell:
- [91845] erythrocyte, leukocyte and platelet studies. Blood 44: 277–284,
- [91846] 1974.

[91847]

[91848] 221. Mills, G. C.; Alperin, J. B.; Trimmer, K. B.: Studies on variant

[91849] glucose-6-phosphate dehydrogenase: G6PD Fort Worth. Biochem. Med. 13:

[91850] 264-275, 1975.

[91851]

[91852] 222. Milner, G.; Delamore, I. W.; Yoshida, A.: G-6-PD Manchester:

[91853] a new variant associated with chronic nonspherocytic hemolytic anemia. Blood 43:

[91854] 271-276, 1974.

[91855]

[91856] 223. Miwa, S.; Fujii, H.: Molecular basis of erythroenzymopathies

[91857] associated with hereditary hemolytic anemia: tabulation of mutant

[91858] enzymes. Am. J. Hemat. 51: 122-132, 1996.

[91859]

[91860] 224. Miwa, S.; Fujii, H.; Nakashima, K.; Miura, Y.; Yamada, K.; Hagiwara,

- [91861] T.; Fukuda, M.: Three new electrophoretically normal glucose-6-phosphate
- [91862] dehydrogenase variants associated with congenital nonspherocytic hemolytic
- [91863] anemia found in Japan: G6PD Ogikubo, Yokohama, and Akita. Hum. Genet. 45:
- [91864] 11-17, 1978.
- [91865]
- [91866] 225. Miwa, S.; Fujii, H.; Nakatsuji, T.; Ishida, Y.; Oda, E.; Kaneto,
- [91867] A.; Motokawa, M.; Ariga, Y.; Fukuchi, S.; Sasai, S.; Hiraoka, K.;
- [91868] Kashii, H.; Kodama, T.: Four new electrophoretically slow-moving
- [91869] glucose-6-phosphate dehydrogenase variants associated with congenital
- [91870] nonspherocytic hemolytic anemia found in Japan: Gd(-) Kurume, Gd(-)
- [91871] Fukushima, Gd(-) Yamaguchi, and Gd(-) Wakayama. Am. J. Hemat. 5:
- [91872] 131-138, 1978.
- [91873]

- [91874] 226. Miwa, S.; Nakashima, K.; Ono, J.; Fujii, H.; Suzuki, E.:
Three
[91875] glucose 6-phosphate dehydrogenase variants found in
Japan. Hum. Genet. 36:
[91876] 327-334, 1977.
[91877]
- [91878] 227. Miwa, S.; Ono, J.; Nakashima, K.; Abe, S.; Kageoka,
T.; Shinohara,
[91879] K.; Isobe, J.; Yamaguchi, H.: Two new glucose
6-phosphate dehydrogenase
[91880] variants associated with congenital nonspherocytic
hemolytic anemia
[91881] found in Japan: Gd(-) Tokushima and Gd(-) Tokyo. Am. J.
Hemat. 1:
[91882] 433-442, 1976.
[91883]
- [91884] 228. Modiano, G.; Battistuzzi, G.; Esan, G. J. F.; Testa, U.;
Luzzatto,
[91885] L.: Genetic heterogeneity of 'normal' human erythrocyte
glucose-6-phosphate
[91886] dehydrogenase: an isoelectrophoretic polymorphism.
Proc. Nat. Acad.

[91887] Sci. 76: 852–856, 1979.

[91888]

[91889] 229. Mohrenweiser, H. W.; Neel, J. V.: Frequency of thermostability

[91890] variants: estimation of total 'rare' variant frequency in human populations. Proc.

[91891] Nat. Acad. Sci. 78: 5729–5733, 1981.

[91892]

[91893] 230. Morelli, A.; Benatti, U.; Guida, L.; De Flora, A.: G6PD Cagliari:

[91894] a new low activity glucose 6–phosphate dehydrogenase variant characterized

[91895] by enhanced intracellular lability. Hum. Genet. 66: 62–65, 1984.

[91896]

[91897] 231. Morisaki, T.; Fujii, H.; Takegawa, S.; Tani, K.; Hirono, A.;

[91898] Takizawa, T.; Takahashi, K.; Shinogi, M.; Teshirogi, T.; Miwa, S.

[91899] : G6PD Sendagi: a new glucose–6–phosphate dehydrogenase variant associated

- [91900] with congenital hemolytic anemia. Hum. Genet. 65: 214–215, 1983.
- [91901]
- [91902] 232. Nafa, K.; Reghis, A.; Osmani, N.; Baghli, L.; Ben-abadji, M.;
- [91903] Kaplan, J.-C.; Vulliamy, T. J.; Luzzatto, L.: G6PD Aures: a new mutation
- [91904] (48 ile-to-thr) causing mild G6PD deficiency is associated with favism. Hum.
- [91905] Molec. Genet. 2: 81–82, 1993.
- [91906]
- [91907] 233. Nagel, R. L.; Ranney, H. M.: Genetic epidemiology of structural
- [91908] mutations of the beta-globin gene. Semin. Hemat. 27: 342–359, 1990.
- [91909]
- [91910] 234. Nakai, T.; Yoshida, A.: G6PD Heian. A glucose-6-phosphate dehydrogenase
- [91911] variant associated with hemolytic anemia found in Japan. Clin. Chim.
- [91912] Acta 51: 199–203, 1974.

[91913]

[91914] 235. Nakashima, K.; Ono, J.; Abe, S.; Miwa, S.; Yoshida, A.:
G6PD

[91915] Ube: a glucose 6-phosphate dehydrogenase variant found
in four unrelated

[91916] Japanese families. Am. J. Hum. Genet. 29: 24-30, 1977.

[91917]

[91918] 236. Nakatsuji, T.; Miwa, S.: Incidence and characteristics
of glucose-6-phosphate

[91919] dehydrogenase variants in Japan. Hum. Genet. 51:
297-305, 1979.

[91920]

[91921] 237. Nance, W. E.: Turner's syndrome, twinning, and an
unusual variant

[91922] of glucose-6-phosphate dehydrogenase. Am. J. Hum.
Genet. 16: 380-392,

[91923] 1964.

[91924]

[91925] 238. Necheles, T. F.; Snyder, L. M.; Strauss, W.: Glucose-
6-phosphate

[91926] dehydrogenase Boston. A new variant associated with

congenital nonspherocytic

[91927] hemolytic disease. Humangenetik 13: 218–221, 1971.

[91928]

[91929] 239. Niazi, G.; Adeyokunu, A.; Westwood, B.; Beutler, E.:
G6PD Aures:

[91930] a rare mutant of G6PD in Saudi Arabia: molecular and
clinical presentations. Saudi

[91931] Med. J. 17: 311–314, 1996.

[91932]

[91933] 240. Ninfali, P.; Baronciani, L.; Bardoni, A.; Bresolin, N.:
Muscle

[91934] expression of glucose–6–phosphate dehydrogenase defi-
ciency in different

[91935] variants. Clin. Genet. 48: 232–237, 1995.

[91936]

[91937] 241. Notaro, R.; Afolayan, A.; Luzzatto, L.: Human muta-
tions in glucose

[91938] 6–phosphate dehydrogenase reflect evolutionary history.
FASEB J. 14:

[91939] 485–494, 2000.

[91940]

- [91941] 242. Nowicki, L.; Strobel, S.; Martin, H.; Koschwitz, U.: Ueber eine
ber eine
- [91942] neue erythrocytaere glucose 6-phosphatdehydrogenase
Variante, Typ
- [91943] Frankfurt. Klin. Wschr. 52: 478-484, 1974.
- [91944]
- [91945] 243. Nsouly, G. M.; Prchal, J. T.: Characterization of a new
G6PD
- [91946] variant and its associated oxidative damage. Clin. Res. 29:
829,
- [91947] 1981.
- [91948]
- [91949] 244. O'Brien, S. J.: The extent and character of biochemical genetic
cal genetic
- [91950] variation in the domestic cat. J. Hered. 71: 2-8, 1980.
- [91951]
- [91952] 245. Ogura, H.; Morisaki, T.; Tani, K.; Kanno, H.; Tsutsumi, H.; Takahashi,
sumi, H.; Takahashi,
- [91953] K.; Miyamori, T.; Fujii, H.; Miwa, S.: A new glucose-
6-phosphate
- [91954] dehydrogenase variant (G6PD Tsukui) associated with

congenital hemolytic

[91955] anemia. Hum. Genet. 78: 369–371, 1988.

[91956]

[91957] 246. Ohno, S.: Sex Chromosomes and Sex-linked Genes.
Berlin and

[91958] New York: Springer (pub.) 1967.

[91959]

[91960] 247. Orzalesi, N.; Sorcinelli, R.; Guiso, G.: Increased incidence

[91961] of cataracts in male subjects deficient in glucose-
6-phosphate dehydrogenase. Arch.

[91962] Ophthalmol. 99: 69–70, 1981.

[91963]

[91964] 248. Othieno-Obel, A.: East African variant of glucose-
6-phosphate

[91965] dehydrogenase. East Afr. Med. J. 49: 230–234, 1972.

[91966]

[91967] 249. Pai, G. S.; Sprenkle, J. A.; Do, T. T.; Maren, C. E.; Mi-
geon,

[91968] B. R.: Localization of loci for hypoxanthine phosphoribo-

sytransferase

[91969] **and glucose-6-phosphate dehydrogenase and biochemical evidence of**

[91970] **nonrandom X chromosome expression from studies of a human X-autosome**

[91971] **translocation. Proc. Nat. Acad. Sci. 77: 2810-2813, 1980.**

[91972]

[91973] **250. Panich, V.: G6PD variants in Laotians. Hum. Hered. 24: 285-290,**

[91974] **1974.**

[91975]

[91976] **251. Panich, V.: G6PD Intanon, a new glucose 6-phosphate dehydrogenase**

[91977] **variant. Humangenetik 21: 203-205, 1974.**

[91978]

[91979] **252. Panich, V.: G6PD characterization in Thailand. Genetics 74**

[91980] **(suppl.): s208 only, 1973.**

[91981]

[91982] **253. Panich, V.: Glucose-6-phosphate dehydrogenase in**

Thailand. Hum.

[91983] Genet. 53: 227–228, 1980.

[91984]

[91985] 254. Panich, V.; Bumrungtrakul, P.; Jitjai, C.; Kamolmatayakul, S.;

[91986] Khoprasert, B.; Klaisuvan, C.; Kongmuang, U.; Maneechai, P.; Pornpatkul,

[91987] M.; Ruengrairatanaroje, P.; Surapruk, P.; Viriyayudhakorn, S.: Glucose–6–phosphate

[91988] dehydrogenase deficiency in South Vietnamese. Hum. Hered. 30: 361–364,

[91989] 1980.

[91990]

[91991] 255. Panich, V.; Na–Nakorn, S.: G6PD variants in Thailand. J. Med.

[91992] Assoc. Thai. 63: 537–543, 1980.

[91993]

[91994] 256. Panich, V.; Sungnate, T.: Characterization of glucose 6–phosphate

[91995] dehydrogenase in Thailand: the occurrence of 6 variants among 50 G6PD

[91996] deficient Thai. Humangenetik 18: 39–46, 1973.

[91997]

[91998] 257. Panich, V.; Sungnate, T.; Na-Nakorn, S.: Acute intravascular

[91999] hemolysis and renal failure in a new glucose 6-phosphate dehydrogenase

[92000] variant: G6PD Siriraj. J. Med. Assoc. Thai. 55: 726–731, 1972.

[92001]

[92002] 258. Panich, V.; Sungnate, T.; Wasi, P.; Na-Nakorn, S.: G-6-PD Mahidol:

[92003] the most common glucose-6-phosphate dehydrogenase variant in Thailand. J.

[92004] Med. Assoc. Thai. 55: 576–585, 1972.

[92005]

[92006] 259. Pawlak, A. L.; Mazurkiewicz, C. A.; Ordynski, J.; Ruzynkowa,

[92007] D.; Horst, A.: G6PD Poznan, variant with severe enzyme deficiency. Humangenetik 28:

[92008] 163–165, 1975.

[92009]

[92010] 260. Pawlak, A. L.; Zagorski, Z.; Rozyrkowa, D.; Horst, A.:
Polish

[92011] variant of glucose-6-phosphate dehydrogenase (G-6-PD
Lublin). Humangenetik 10:

[92012] 340-343, 1970.

[92013]

[92014] 261. Pekrun, A.; Eber, S. W.; Schroter, W.: G6PD Avenches
and G6PD

[92015] Moosburg: biochemical and erythrocyte membrane char-
acterization. Blut 58:

[92016] 11-14, 1989.

[92017]

[92018] 262. Perroni, L.; Tassara, P.; Baldi, M.; Reali, R.;
Scartezzini,

[92019] P.: G6PD variants detected in Genoa area. In: Weatherall, D.
J.; Fiorelli,

[92020] G.; Gorini, S.: Advances in Red Blood Cell Biology. New
York: Raven

[92021] Press (pub.) 1982. Pp. 409-416.

[92022]

[92023] 263. Persico, M. G.; Toniolo, D.; Nobile, C.; D'Urso, M.;

Luzzatto,

[92024] L.: cDNA sequences of human glucose 6-phosphate dehydrogenase cloned

[92025] in pBR322. Nature 294: 778-780, 1981.

[92026]

[92027] 264. Persico, M. G.; Viglietto, G.; Martini, G.; Toniolo, D.; Paonessa,

[92028] G.; Moscatelli, C.; Dono, R.; Vulliamy, T.; Luzzatto, L.; D'Urso,

[92029] M.: Isolation of human glucose-6-phosphate dehydrogenase (G6PD) cDNA

[92030] clones: primary structure of the protein and unusual 5-prime non-coding

[92031] region. Nucleic Acids Res. 14: 2511-2522, 1986.

[92032]

[92033] 265. Picat, C.; Etienne, J.; Boivin, P.; Le Prise, P.-Y.: Gd(-)

[92034] Rennes, a new deficient variant of glucose-6-phosphate dehydrogenase

[92035] associated with congenital nonspherocytic hemolytic anemia found in

[92036] France. Hum. Genet. 55: 125-127, 1980.

[92037]

[92038] 266. Pinto, P. V. C.; Newton, W. A., Jr.; Richardson, K. E.:
Evidence

[92039] for four types of erythrocyte glucose-6-phosphate dehydrogenase from

[92040] G-6-PD deficient human subjects. J. Clin. Invest. 45:
823-831, 1966.

[92041]

[92042] 267. Poggi, V.: Personal Communication. London, England
1989.

[92043]

[92044] 268. Poon, M.-C.; Hall, K.; Scott, C. W.; Prchal, J. T.: G6PD
Viangchan:

[92045] a new glucose 6-phosphate dehydrogenase variant from
Laos. Hum. Genet. 78:

[92046] 98-99, 1988.

[92047]

[92048] 269. Porter, I. H.; Boyer, S. H.; Watson-Williams, E. J.;
Adam, A.;

[92049] Szeinberg, A.; Siniscalco, M.: Variation of glucose-
6-phosphate dehydrogenase

[92050] in different populations. Lancet I: 895–899, 1964.

[92051]

[92052] 270. Porter, I. H.; Schulze, J.; McKusick, V. A.: Genetical linkage

[92053] between the loci for glucose–6–phosphate dehydrogenase deficiency

[92054] and colour–blindness in American Negroes. Ann. Hum. Genet. 26: 107–122,

[92055] 1962.

[92056]

[92057] 271. Prchal, J.; Moreno, H.; Conrad, M.; Vitek, A.: G–6–PD Dothan:

[92058] a new variant associated with chronic hemolytic anemia. I.R.C.S. 7:

[92059] 348, 1979.

[92060]

[92061] 272. Prchal, J. T.: Personal Communication. Birmingham, Ala. 1985.

[92062]

[92063] 273. Prchal, J. T.; Crist, W. M.; Malluh, A.; Vitek, A.; Tauxe, W.

- [92064] N.; Carroll, A. J.: A new glucose-6-phosphate dehydrogenase deficient
- [92065] variant in a patient with Chediak-Higashi syndrome. Blood 56: 476-480,
- [92066] 1980.
- [92067]
- [92068] 274. Prchal, J. T.; Hall, K.; Csepregy, M.; Lilly, M.; Berkow, R.;
- [92069] Scott, C. W.: Two apparent glucose-6-phosphate dehydrogenase variants
- [92070] in normal XY males: G6PD Alabama. Am. J. Med. 84: 517-523, 1988.
- [92071]
- [92072] 275. Pretsch, W.; Charles, D. J.; Merkle, S.: X-linked glucose-6-phosphate-dehydrogenase
- [92073] deficiency in *Mus musculus*. Biochem. Genet. 26: 89-103, 1988.
- [92074]
- [92075] 276. Puck, J. M.; Willard, H. F.: X inactivation in females with
- [92076] X-linked disease. New Eng. J. Med. 338: 325-327, 1998.

[92077]

[92078] 277. Ramot, B.; Ben-Bassat, I.; Shchory, M.: New glucose-6-phosphate

[92079] dehydrogenase variants observed in Israel and their association with

[92080] congenital nonspherocytic hemolytic disease. J. Lab. Clin. Med. 74:

[92081] 895-901, 1969.

[92082]

[92083] 278. Ramot, B.; Brok, F.: A new glucose-6-phosphate dehydrogenase

[92084] mutant (Tel-Hashomer mutant). Ann. Hum. Genet. 28: 167-172, 1964.

[92085]

[92086] 279. Rattazzi, M. C.; Corash, L. M.; Van Zzanen, G. E.; Jaffe, E.

[92087] R.; Piomelli, S.: G6PD deficiency and chronic hemolysis: four new

[92088] mutants--relationships between clinical syndrome and enzyme kinetics. Blood 38:

[92089] 205-218, 1971.

[92090]

[92091] 280. Rattazzi, M. C.; Lenzerini, L.; Meera Khan, P.; Luzzatto, L.

[92092] : Characterization of glucose-6-phosphate dehydrogenase variants.

[92093] II. G6PD Kephalaria, G6PD Attica, and G6PD 'Seattle-like' found in

[92094] Greece. Am. J. Hum. Genet. 21: 154-167, 1969.

[92095]

[92096] 281. Ravindranath, Y.; Beutler, E.: Two new variants of glucose-6-phosphate

[92097] dehydrogenase associated with hereditary non-spherocytic hemolytic

[92098] anemia: G6PD Wayne and G6PD Huron. Am. J. Hemat. 24: 357-363, 1987.

[92099]

[92100] 282. Reys, L.; Manso, C.; Stamatoyannopoulos, G.: Genetic studies

[92101] on Southeastern Bantu of Mozambique. I. Variants of glucose-6-phosphate

[92102] dehydrogenase. Am. J. Hum. Genet. 22: 203-215, 1970.

[92103]

[92104] 283. Rinaldi, A.; Filippi, G.; Siniscalco, M.: Variability of
red

[92105] cell phenotypes between and within individuals in an un-
biased sample

[92106] of 77 heterozygotes for G6PD deficiency in Sardinia. Am.
J. Hum.

[92107] Genet. 28: 496–505, 1976.

[92108]

[92109] 284. Rosenstrauss, M.; Chasin, L. A.: Isolation of mam-
malian cell

[92110] mutants deficient in glucose-6-phosphate dehydrogenase
activity: linkage

[92111] to hypoxanthine phosphoribosyl transferase. Proc. Nat.
Acad. Sci. 72:

[92112] 493–497, 1975.

[92113]

[92114] 285. Roth, E. F., Jr.; Raventos-Suarez, C.; Rinaldi, A.;
Nagel, R.

[92115] L.: Glucose-6-phosphate dehydrogenase deficiency in-
hibits in vitro

- [92116] growth of *Plasmodium falciparum*. Proc. Nat. Acad. Sci.
80: 298–299,
[92117] 1983.
- [92118]
- [92119] 286. Roychoudhury, A. K.; Nei, M.: Human Polymorphic
Genes: World
[92120] Distribution. New York: Oxford Univ. Press (pub.) 1988.
- [92121]
- [92122] 287. Ruwando, C.; Khea, S. C.; Snow, R. W.; Yates, S. N. R.;
Kwiatkoweld,
[92123] D.; Gupta, S.; Warn, P.; Alisopp, G. E. M.; Gilbert, S. C.;
Peschu,
[92124] N.; Newbold, C. I.; Greenwood, S. M.; Marsh, K.; Hill, A. V.
S.:
[92125] Natural selection of hemi- and heterozygotes for G6PD
deficiency in
[92126] Africa by resistance to severe malaria. Nature 376:
246–249, 1995.
- [92127]
- [92128] 288. Saenz, G. F.; Chaves, M.; Berrantes, A.; Elizondo, J.;
Montero,

[92129] A. G.; Yoshida, A.: A glucose-6-phosphate dehydrogenase variant,

[92130] Gd(-) Santamaria found in Costa Rica. *Acta Haemat.* 72: 37-40, 1984.

[92131]

[92132] 289. Saidi, N.; Hors-Cayla, M. C.; Van Cong, N.; Benne, F.: Sheep

[92133] gene mapping by somatic cell hybridization. (Abstract) *Cytogenet.*

[92134] *Cell Genet.* 25: 200, 1979.

[92135]

[92136] 290. Samuel, A. P. W.; Saha, N.; Omer, A.; Hoffbrand, A. V.: Quantitative

[92137] expression of G6PD activity of different phenotypes of G6PD and haemoglobin

[92138] in a Sudanese population. *Hum. Hered.* 31: 110-115, 1981.

[92139]

[92140] 291. Sansone, G.; Perroni, L.; Testa, U.; Mareni, C.; Luz-zatto, L.

[92141] : New genetic variants of glucose 6-phosphate dehydrogenase (G6PD)

[92142] in Italy. Ann. Hum. Genet. 45: 97–104, 1981.

[92143]

[92144] 292. Sansone, G.; Perroni, L.; Yoshida, A.: Glucose–
6–phosphate dehydrogenase

[92145] variants from Italian subjects associated with severe
neonatal jaundice. Brit.

[92146] J. Haemat. 31: 159–165, 1975.

[92147]

[92148] 293. Sansone, G.; Perroni, L.; Yoshida, A.; Dave, V.: A new
glucose

[92149] 6–phosphate dehydrogenase variant (Gd Trinacria) in two
unrelated

[92150] families of Sicilian ancestry. Ital. J. Biochem. 26: 44–50,
1977.

[92151]

[92152] 294. Shatskaya, T. L.; Krasnopolskaya, K. D.; Annenkov, G.
A.: A

[92153] description of new mutant forms of erythrocyte glucose–
6–phosphate

[92154] dehydrogenase isolated at the territory of the Soviet
Union. Genetika 11:

[92155] 116–122, 1975.

[92156]

[92157] 295. Shatskaya, T. L.; Krasnopolskaya, K. D.; Idelson, L. J.:
The

[92158] new form of glucose 6–phosphate dehydrogenase (G6PD
'Kaluga') from

[92159] erythrocytes of a patient with chronic non–spherocytic
hemolytic anemia. Vopr.

[92160] Med. Khim. 22: 764–768, 1976.

[92161]

[92162] 296. Shatskaya, T. L.; Krasnopolskaya, K. D.; Idelson, L. J.:
Mutant

[92163] forms of erythrocyte glucose 6–phosphate dehydrogenase
in Ashkenazi:

[92164] description of two new variants, G6PD Kirovograd and
G6PD Zhitomir. Humangenetik 33:

[92165] 175–178, 1976.

[92166]

[92167] 297. Shatskaya, T. L.; Krasnopolskaya, K. D.; Tzoneva, M.;
Mavrudieva,

[92168] M.; Toncheva, D.: Variants of erythrocyte glucose–

6-phosphate dehydrogenase

[92169] (G6PD) in Bulgarian populations. Hum. Genet. 54: 115–117, 1980.

[92170]

[92171] 298. Shatskaya, T. L.; Krasnopolskaya, K. D.; Zakharova, T. V.: Regularities

[92172] of distribution of Gd-alleles in Azerbaijan. II. Identification of

[92173] G6PD mutant forms. Genetika 16: 2217–2225, 1980.

[92174]

[92175] 299. Shows, T. B.; Brown, J. A.: Human X-linked genes regionally

[92176] mapped utilizing X-autosome translocations and somatic cell hybrids. Proc.

[92177] Nat. Acad. Sci. 72: 2125–2129, 1975.

[92178]

[92179] 300. Shows, T. B.; Brown, J. A.; Chapman, V. M.: Comparative gene

[92180] mapping of HPRT, G6PD and PGK in man, mouse, and Muntjac deer. Birth

[92181] Defects Orig. Art. Ser. XII(7): 436–439, 1976.

[92182]

[92183] 301. Shows, T. B.; Tashian, R. E.; Brewer, G. J.: Erythrocyte glucose-6-phosphate

[92184] dehydrogenase in Caucasians: new inherited variant. Science 145:

[92185] 1056-1057, 1964.

[92186]

[92187] 302. Sidi, Y.; Aderka, D.; Brok-Simoni, F.; Benjamin, D.; Ramot, B.;

[92188] Pinkhas, J.: Viral hepatitis with extreme hyperbilirubinaemia, massive

[92189] hemolysis and encephalopathy in a patient with a new G6PD variant. Israel

[92190] J. Med. Sci. 16: 130-133, 1980.

[92191]

[92192] 303. Siegel, N. H.; Beutler, E.: Hemolytic anemia caused by G-6-PD

[92193] Carswell, a new variant. Ann. Intern. Med. 75: 437-439, 1971.

[92194]

[92195] 304. Smith, J. E.; Ryer, K.; Wallace, L.: Glucose-

6-phosphate dehydrogenase

[92196] deficiency in a dog. *Enzyme* 21: 379–382, 1976.

[92197]

[92198] 305. Snyder, L. M.; Necheles, T. F.; Reddy, W. J.: G-6-PD
Worcester:

[92199] a new variant, associated with X-linked optic atrophy. *Am.
J. Med.* 49:

[92200] 125–132, 1970.

[92201]

[92202] 306. Stamatoyannopoulos, G.; Kotsakis, P.; Voigtlander,
V.; Akrivakis,

[92203] A.; Motulsky, A. G.: Electrophoretic diversity of glucose-
6-phosphate

[92204] dehydrogenase among Greeks. *Am. J. Hum. Genet.* 22:
587–596, 1970.

[92205]

[92206] 307. Stamatoyannopoulos, G.; Voigtlander, V.; Akrivakis,
A.: Thessaly

[92207] variant of glucose-6-phosphate dehydrogenase. *Human-
genetik* 9: 23–25,

[92208] 1970.

[92209]

[92210] 308. Stamatoyannopoulos, G.; Voightlander, V.; Kotsakis, P.; Akrivakis,

[92211] A.: Genetic diversity of the 'Mediterranean' glucose-6-phosphate

[92212] dehydrogenase deficiency phenotype. J. Clin. Invest. 50: 1253-1261,

[92213] 1971.

[92214]

[92215] 309. Stamatoyannopoulos, G.; Yoshida, A.; Bacopoulos, C.; Motulsky,

[92216] A. G.: Athens variant of glucose-6-phosphate dehydrogenase. Science 157:

[92217] 831-833, 1967.

[92218]

[92219] 310. Stevens, D. J.; Wanachiwanawin, W.; Mason, P. J.; Vulliamy, T.

[92220] J.; Luzzatto, L.: G6PD Canton a common deficient variant in South

[92221] East Asia caused by a 459 arg-to-leu mutation. Nucleic Acids Res. 18:

[92222] 7190, 1990.

[92223]

[92224] 311. Stocco dos Santos, R. C.; Barretto, O. C. O.;
Nonoyama, K.; Castro,

[92225] N. H. C.; Ferraz, O. P.; Walter-Moura, J.; Vescio, C. C. S.;
Becak,

[92226] W.: X-linked syndrome: mental retardation, hip luxation,
and G6PD

[92227] variant (Gd(+)) Butantan). Am. J. Med. Genet. 39: 133-136,
1991.

[92228]

[92229] 312. Stockham, S. L.; Harvey, J. W.; Kinden, D. A.: Equine
glucose-6-phosphate

[92230] dehydrogenase deficiency. Vet. Path. 31: 518-527, 1994.

[92231]

[92232] 313. Streiff, F.; Vigneron, C.: Anémie hémolytique
chronique par

[92233] déficit en glucose 6-phosphate déshydrogénase dans une
famille d'origine

[92234] Lorraine. Nouv. Rev. Franc. Hemat. 11: 279-290, 1971.

[92235]

- [92236] 314. Takahashi, K.; Fujii, H.; Takegawa, S.; Tani, K.; Hirono, A.;
- [92237] Takizawa, T.; Kawakatsu, T.; Miwa, S.: A new glucose-6-phosphate
- [92238] dehydrogenase variant (G6PD Nagano) associated with congenital hemolytic
- [92239] anemia. Hum. Genet. 62: 368-370, 1982.
- [92240]
- [92241] 315. Takizawa, T.; Fujii, H.; Takegawa, S.; Takahashi, K.; Hirono,
- [92242] A.; Morisaki, T.; Kanno, H.; Oka, R.; Yoshioka, H.; Miwa, S.: A unique
- [92243] electrophoretic slow-moving glucose 6-phosphate dehydrogenase variant
- [92244] (G6PD Asahikawa) with a markedly acidic pH optimum. Hum. Genet. 68:
- [92245] 70-72, 1984.
- [92246]
- [92247] 316. Takizawa, T.; Huang, I.-Y.; Ikuta, T.; Yoshida, A.: Human glucose-6-phosphate
- [92248] dehydrogenase: primary structure and cDNA cloning. Proc. Nat. Acad.

[92249] Sci. 83: 4157-4161, 1986.

[92250]

[92251] 317. Takizawa, T.; Yoneyama, Y.; Miwa, S.; Yoshida, A.: A single

[92252] nucleotide base transition is the basis of the common human glucose-6-phosphate

[92253] dehydrogenase variant A(+). Genomics 1: 228-231, 1987.

[92254]

[92255] 318. Takizawa, T.; Yoshida, A.: Molecular abnormality of the common

[92256] glucose-6-phosphate dehydrogenase variant, G6PD A(+), and restriction-fragment-length

[92257] polymorphism. (Abstract) Am. J. Hum. Genet. 41: A241, 1987.

[92258]

[92259] 319. Talalak, P.; Beutler, E.: G-6-PD Bangkok: a new variant found

[92260] in congenital nonspherocytic hemolytic disease (CNHD). Blood 33:

[92261] 772-776, 1969.

[92262]

- [92263] 320. Tanaka, K. R.; Beutler, E.: Hereditary hemolytic anemia due
[92264] to glucose-6-phosphate dehydrogenase Torrance: a new
variant. J.
[92265] Lab. Clin. Med. 73: 657-667, 1969.
[92266]
[92267] 321. Tang, T. K.; Huang, C.-S.; Huang, M. J.; Tam, K.-B.;
Yeh, C.-H.;
[92268] Tang, C.-J. C.: Diverse point mutations result in glucose-
6-phosphate
[92269] dehydrogenase (G6PD) polymorphism in Taiwan. Blood
79: 2135-2140,
[92270] 1992.
[92271]
[92272] 322. Testa, U.; Meloni, T.; Lania, A.; Battistuzzi, G.; Cuttillo, S.;
[92273] Luzzatto, L.: Genetic heterogeneity of glucose-
6-phosphate dehydrogenase
[92274] deficiency in Sardinia. Hum. Genet. 56: 99-105, 1980.
[92275]
[92276] 323. Thigpen, J. T.; Steinberg, M. H.; Beutler, E.; Gillespie,

G.

[92277] T., Jr.; Dreiling, B. J.; Morrison, F. S.: Glucose-6-phosphate dehydrogenase

[92278] Jackson, a new variant associated with hemolytic anemia. Acta Haemat. 51:

[92279] 310-314, 1974.

[92280]

[92281] 324. Tishkoff, S. A.; Varkonyi, R.; Cahinhinan, N.; Abbes, S.; Argyropoulos,

[92282] G.; Destro-Bisol, G.; Drousiotou, A.; Dangerfield, B.; Lefranc, G.;

[92283] Loiselet, J.; Piro, A.; Stoneking, M.; Tagarelli, A.; Tagarelli, G.;

[92284] Touma, E. H.; Williams, S. M.; Clark, A. G.: Haplotype diversity

[92285] and linkage disequilibrium at human G6PD: recent origin of alleles

[92286] that confer malarial resistance. Science 293: 455-462, 2001.

[92287]

[92288] 325. Tokarev, Y. N.; Chernyak, N. B.; Batischev, A. I.; Lanzina, N.

- [92289] V.; Alexeyev, G. A.: Etude des proprietes electrophoretiques et cinetiques
- [92290] de la glucose-6-phosphate deshydrogenase (Gd)
- [92291] d'erythrocytes dans
- [92291] les deficits hereditaires de l'enzyme: description d'une nouvelle
- [92292] variante de glucose-6-phosphate deshydrogenase: la Gd Kremenchug. Nouv.
- [92293] Rev. Franc. Hemat. 20: 557-564, 1978.
- [92294]
- [92295] 326. Toncheva, D.: Variants of glucose-6-phosphate dehydrogenase
- [92296] in a Vietnamese population. Hum. Hered. 36: 348-351, 1986.
- [92297]
- [92298] 327. Toncheva, D.; Tzoneva, M.: Genetic polymorphism of G6PD in a
- [92299] Bulgarian population. Hum. Genet. 67: 340-342, 1984.
- [92300]
- [92301] 328. Toniolo, D.; Martini, G.; Migeon, B. R.; Dono, R.: Expression

[92302] of the G6PD locus on the human X chromosome is associated with demethylation

[92303] of three CpG islands within 100 kb of DNA. EMBO J. 7: 401–406, 1988.

[92304]

[92305] 329. Town, M.; Athanasiou–Metaxa, M.; Luzzatto, L.: In–tragenic interspecific

[92306] complementation of glucose 6–phosphate dehydrogenase in human–hamster

[92307] cell hybrids. Somat. Cell Molec. Genet. 16: 97–108, 1990.

[92308]

[92309] 330. Town, M.; Bautista, J. M.; Mason, P. J.; Luzzatto, L.: Both

[92310] mutations in G6PD A– are necessary to produce the G6PD deficient phenotype. Hum.

[92311] Molec. Genet. 1: 171–174, 1992.

[92312]

[92313] 331. Usanga, E. A.; Bienzle, U.; Cancedda, K.; Fasuan, F. A.; Ajayi,

[92314] O.; Luzzatto, L.: Genetic variants of human erythrocyte glucose 6–phosphate

[92315] dehydrogenase: new variants in West Africa characterized by column

[92316] chromatography. Ann. Hum. Genet. 40: 279–286, 1977.

[92317]

[92318] 332. Vaca, G.; Ibarra, B.; Garcia Cruz, D.; Medina, C.; Romero, F.;

[92319] Cantu, J. M.; Beutler, E.: G-6-PD Jalisco and G-6-PD Morelia: two

[92320] new Mexican variants. Hum. Genet. 71: 82–85, 1985.

[92321]

[92322] 333. Vaca, G.; Ibarra, B.; Romero, F.; Olivares, N.; Cantu, J. M.;

[92323] Beutler, E.: G-6-PD Guadalajara: a new mutant associated with chronic

[92324] nonspherocytic hemolytic anemia. Hum. Genet. 61: 175–176, 1982.

[92325]

[92326] 334. Ventura, A.; Panizon, F.; Soranzo, M. R.; Veneziano, G.; Sansone,

[92327] G.; Testa, U.; Luzzatto, L.: Congenital dyserythropoietic anaemia

[92328] type II associated with a new type of G6PD deficiency (G6PD Gabrovizza). Acta

[92329] Haemat. 71: 227-234, 1984.

[92330]

[92331] 335. Vergnes, H.; Gherardi, M.; Bouloux, C.: Erythrocyte glucose-6-phosphate

[92332] dehydrogenase in the Niokolonko (Malinke of the Niokolo) of the Eastern

[92333] Senegal: identification of a slow variant with normal activity (Tacoma-like). Hum.

[92334] Hered. 25: 80-87, 1975.

[92335]

[92336] 336. Vergnes, H.; Gherardi, M.; Quilici, J. C.; Yoshida, A.; Giacardy,

[92337] R.: G6PD Luz-Saint-Sauveur: a new variant with abnormal electrophoretic

[92338] mobility, mild enzyme deficiency and absence of hemolytic disorders. I.R.C.S. 7:

[92339] 14, 1973.

[92340]

[92341] 337. Vergnes, H.; Gherardi, M.; Yoshida, A.: G6PD Lozere

and Trinacria-like:

[92342] segregation of two non-hemolytic variants in a French family. Hum.

[92343] Genet. 34: 293-298, 1976.

[92344]

[92345] 338. Vergnes, H.; Ribet, A.; Bommelaer, G.; Amadiou, J.; Brun, H.

[92346] : GD(-) Muret and GD(-) Colomiers, two new variants of glucose-6-phosphate

[92347] dehydrogenase associated with favism. Hum. Genet. 57: 332-334, 1981.

[92348]

[92349] 339. Vergnes, H.; Yoshida, A.; Gourdin, D.; Gherardi, M.; Bierme,

[92350] R.; Ruffie, J.: Glucose 6-phosphate dehydrogenase Toulouse: a new

[92351] variant with marked instability and severe deficiency discovered in

[92352] a family of Mediterranean ancestry. Acta Haemat. 51: 240-249, 1974.

[92353]

[92354] 340. Viglietto, G.; Montanaro, V.; Calabro, V.; Vallone, D.;

D'Urso,

[92355] M.; Persico, M. G.; Battistuzzi, G.: Common glucose-6-phosphate dehydrogenase

[92356] (G6PD) variants from the Italian population: biochemical and molecular

[92357] characterization. Ann. Hum. Genet. 54: 1-15, 1990.

[92358]

[92359] 341. Vives-Corrons, J.-L.; Kuhl, W.; Pujades, M. A.; Beutler, E.:

[92360] Molecular genetics of the glucose-6-phosphate dehydrogenase (G6PD)

[92361] Mediterranean variant and description of a new G6PD mutant, G6PD Andalus(1361A). Am.

[92362] J. Hum. Genet. 47: 575-579, 1990.

[92363]

[92364] 342. Vives-Corrons, J. L.; Feliu, E.; Pujades, M. A.; Cardellach,

[92365] F.; Rozman, C.; Carreras, A.; Jou, J. M.; Vallespi, M. T.; Zuazu,

[92366] F. J.: Severe glucose-6-phosphate dehydrogenase (G6PD) deficiency

[92367] associated with chronic hemolytic anemia, granulocyte

dysfunction,

[92368] and increased susceptibility to infections: description of a new molecular

[92369] variant (G6PD Barcelona). Blood 59: 428-434, 1982.

[92370]

[92371] 343. Vives-Corrons, J. L.; Pujades, A.: Heterogeneity of 'Mediterranean

[92372] type' glucose-6-phosphate dehydrogenase (G6PD) deficiency in Spain

[92373] and description of two new variants associated with favism. Hum.

[92374] Genet. 60: 216-221, 1982.

[92375]

[92376] 344. Vives-Corrons, J. L.; Pujades, A.; Curia, M. D.: Caracterizacion

[92377] molecular de la glucosa-6-fosfato deshidrogenasa (G6PD) en 24 casos

[92378] de deficit enzimatico y descripcion de una nueva variante (G6PD-Betica). Sangre 25:

[92379] 1049-1064, 1980.

[92380]

[92381] 345. Vives-Corrons, J. L.; Pujades, M. A.; Petit, J.; Colomer, D.;

[92382] Corbella, M.; Aguilar i Bascompte, J. L.; Merino, A.:
Chronic nonspherocytic

[92383] hemolytic anemia (CNSHA) and glucose 6 phosphate de-
hydrogenase (G6PD)

[92384] deficiency in a patient with familial amyloidotic polyneu-
ropathy (FAP):

[92385] molecular study of a new variant (G6PD Clinic) with
markedly acidic

[92386] pH optimum. Hum. Genet. 81: 161-164, 1989.

[92387]

[92388] 346. Vulliamy, T.; Beutler, E.; Luzzatto, L.: Variants of glu-
cose-6-phosphate

[92389] dehydrogenase are due to missense mutations spread
throughout the

[92390] coding region of the gene. Hum. Mutat. 2: 159-167,
1993.

[92391]

[92392] 347. Vulliamy, T.; Mason, P.; Luzzatto, L.: The molecular
basis of

[92393] glucose-6-phosphate dehydrogenase deficiency. Trends

Genet. 8: 138–143,

[92394] 1992.

[92395]

[92396] 348. Vulliamy, T.; Rovira, A.; Yusoff, N.; Colomer, D.; Luzzatto,

[92397] L.; Vives-Corrons, J.-L.: Independent origin of single and double

[92398] mutations in the human glucose 6-phosphate dehydrogenase gene. Hum.

[92399] Mutat. 8: 311–318, 1996.

[92400]

[92401] 349. Vulliamy, T. J.: Personal Communication. London, England

[92402] 1989.

[92403]

[92404] 350. Vulliamy, T. J.; D'Urso, M.; Battistuzzi, G.; Estrada, M.; Foulkes,

[92405] N. S.; Martini, G.; Calabro, V.; Poggi, V.; Giordano, R.; Town, M.;

[92406] Luzzatto, L.; Persico, M. G.: Diverse point mutations in the human

- [92407] glucose-6-phosphate dehydrogenase gene cause enzyme deficiency and
- [92408] mild or severe hemolytic anemia. Proc. Nat. Acad. Sci. 85: 5171-5175,
- [92409] 1988.
- [92410]
- [92411] 351. Vulliamy, T. J.; Kaeda, J. S.; Ait-Chafa, D.; Mangerini, R.;
- [92412] Roper, D.; Barbot, J.; Mehta, A. B.; Athanassiou-Metaxa, M.; Luzzatto,
- [92413] L.; Mason, P. J.: Clinical and haematological consequences of recurrent
- [92414] G6PD mutations and a single new mutation causing chronic nonspherocytic
- [92415] haemolytic anaemia. Brit. J. Haemat. 101: 670-675, 1998.
- [92416]
- [92417] 352. Vulliamy, T. J.; Othman, A.; Town, M.; Nathwani, A.; Falusi,
- [92418] A. G.; Mason, P. J.; Luzzatto, L.: Polymorphic sites in the African
- [92419] population detected by sequence analysis of the glucose-6-phosphate

[92420] dehydrogenase gene outline the evolution of the variants A and A-. Proc.

[92421] Nat. Acad. Sci. 88: 8568-8571, 1991.

[92422]

[92423] 353. Vulliamy, T. J.; Wanachiwanawin, W.; Mason, P. J.; Luzzatto,

[92424] L.: G6PD Mahidol, a common deficient variant in South East Asia is

[92425] caused by a (163)glycine-to-serine mutation. Nucleic Acids Res. 17:

[92426] 5868, 1989.

[92427]

[92428] 354. Vuopio, P.; Harkonen, M.; Helske, T.; Naeveri, H.: Red cell

[92429] glucose-6-phosphate dehydrogenase deficiency in Finland: characterization

[92430] of a new variant with severe enzyme deficiency. Scand. J. Haemat. 15:

[92431] 145-152, 1975.

[92432]

[92433] 355. Vuopio, P.; Harkonen, M.; Johnsson, P.; Nuutinen, M.:

Red cell

[92434] glucose-phosphate dehydrogenase deficiency in Finland.
Ann. Clin.

[92435] Res. 5: 168-178, 1973.

[92436]

[92437] 356. Waitz, R.; Boivin, P.; Oberling, F.; Casenave, J. P.;
North,

[92438] M. L.; Mayer, S.: Variante Gd(-) Strasbourg de la glucose-
6-phosphate

[92439] dehydrogenase. Nouv. Rev. Franc. Hemat. 10: 312-314,
1970.

[92440]

[92441] 357. Wang, Y. M.; Patterson, J. H.; Van Eys, J.: The poten-
tial use

[92442] of xylitol in glucose-6-phosphate dehydrogenase defi-
ciency anemia. J.

[92443] Clin. Invest. 50: 1421-1428, 1971.

[92444]

[92445] 358. Weimer, T. A.; Salzano, F. M.; Hutz, M. H.: Erythro-
cyte isozymes

[92446] and hemoglobin types in a southern Brazilian population.

J. Hum.

[92447] *Evol.* 10: 319–328, 1981.

[92448]

[92449] 359. Weimer, T. A.; Schuler, L.; Beutler, E.; Salzano, F. M.:
Gd(+)

[92450] Laguna, a new rare glucose-6-phosphate dehydrogenase
variant from

[92451] Brazil. *Hum. Genet.* 65: 402–404, 1984.

[92452]

[92453] 360. Weinreich, J.; Busch, D.; Gottstein, U.; Schaefer, J.;
Rohr,

[92454] J.: Ueber zwei neue Faelle von hereditaerer nichtsphaero-
cytaerer

[92455] haemolytischer Anaemie bei glucose-
6-phosphat-dehydrogenase-Defekt

[92456] in einer Nord Deutschen Familie. *Klin. Wschr.* 46:
146–149, 1968.

[92457]

[92458] 361. Welch, S. G.; McGregor, I. A.; Williams, K.: A new
variant of

[92459] human erythrocyte G6PD occurring at a high frequency

amongst the population

[92460] of two villages in The Gambia, West Africa. Hum. Genet.
40: 305–309,

[92461] 1978.

[92462]

[92463] 362. Westring, D. W.; Pisciotto, A. V.: Anemia, cataracts,
and seizures

[92464] in patients with glucose–6–phosphate dehydrogenase de-
ficiency. Arch.

[92465] Intern. Med. 118: 385–390, 1966.

[92466]

[92467] 363. WHO: Nomenclature of glucose–6–phosphate dehy-
drogenase in

[92468] man. Bull. WHO 36: 319–322, 1967. Note: See Also:
Canad. Med. Assoc.

[92469] J. 97: 422–424, 1967.

[92470]

[92471] 364. WHO: Scientific group on the standardization of pro-
cedures

[92472] for the study of glucose–6–phosphate dehydrogenase.
WHO Techn. Rep.

[92473] (pub.) Ser. No. 366: 1967.

[92474]

[92475] 365. Wilson, W. W.: Congenital hemolytic anemia due to a deficiency

[92476] of glucose 6-phosphate dehydrogenase. Rocky Mt. Med. J. 73: 160-162,

[92477] 1976.

[92478]

[92479] 366. Witt, I.; Yoshioka, S.: Biochemical characterization of a glucose-6-phosphate

[92480] dehydrogenase variant with favism: G-6-PD Zaehringen. Klin. Wschr. 50:

[92481] 205-209, 1972.

[92482]

[92483] 367. Wong, P. W. K.; Shih, L.-Y.; Hsia, D. Y. Y.: Characterization

[92484] of glucose-6-phosphate dehydrogenase among Chinese. Nature 208:

[92485] 1323-1324, 1965.

[92486]

- [92487] 368. Yermakov, N.; Tokarev, J.; Chernjak, N.; Schoenian, G.; Grieger, M.; Guckler, G.; Jacobasch, G.; Mahmudova, M.; Bahramov, S.: New stable mutant Gd(-) variants: G6PD Tashkent and G6PD Nucus: molecular basis of hereditary enzyme deficiency. Acta Biol. Med. Ger. 40: 559–562, 1981.
- [92492]
- [92493] 369. Yoshida, A.: Personal Communication. Duarte, Calif. 2/26/1996.
- [92494]
- [92495] 370. Yoshida, A.: Amino acid substitution (histidine to tyrosine) in a glucose-6-phosphatate dehydrogenase variant (G6PD Hektoen) associated with over-production. J. Mol. Biol. 52: 483–490, 1970.
- [92498]
- [92499] 371. Yoshida, A.: A single amino acid substitution (asparagine to

- [92500] aspartic acid) between normal (B plus) and the common Negro variant
- [92501] (A plus) of human glucose-6-phosphate dehydrogenase. Proc. Nat. Acad.
- [92502] Sci. 57: 835-840, 1967.
- [92503]
- [92504] 372. Yoshida, A.: Human glucose-6-phosphate dehydrogenase: purification
- [92505] and characterization of Negro type variant (A+) and comparison with
- [92506] normal enzyme (B+). Biochem. Genet. 1: 81-99, 1967.
- [92507]
- [92508] 373. Yoshida, A.; Baur, E. W.; Motulsky, A. G.: A Philippino glucose-6-phosphate
- [92509] dehydrogenase variant (G6PD Union) with enzyme deficiency and altered
- [92510] substrate specificity. Blood 35: 506-513, 1970.
- [92511]
- [92512] 374. Yoshida, A.; Beutler, E.: Human glucose-6-phosphate dehydrogenase
- [92513] variants: a supplementary tabulation. Ann. Hum. Genet.

41: 347-355,

[92514] 1978.

[92515]

[92516] 375. Yoshida, A.; Beutler, E.; Motulsky, A. G.: Table of human glucose-6-phosphate

[92517] dehydrogenase variants. Bull. WHO 45: 243-253, 1971.

[92518]

[92519] 376. Yoshida, A.; Stamatoyannopoulos, G.; Motulsky, A. G.: Negro

[92520] variant of glucose-6-phosphate dehydrogenase deficiency (A-) in man. Science 155:

[92521] 97-99, 1967.

[92522]

[92523] 377. Yoshida, A.; Takizawa, T.: The same extra FokI cleavage site

[92524] exists in glucose-6-phosphate dehydrogenase variants A(+) and A(-). Am.

[92525] J. Hum. Genet. 43: 131-133, 1988.

[92526]

[92527] 378. Yoshida, A.; Takizawa, T.; Prchal, J. T.: RFLP of the X

chromosome-linked

[92528] glucose-6-phosphate dehydrogenase locus in blacks. Am. J. Hum. Genet. 42:

[92529] 872-876, 1988.

[92530]

[92531] 379. Zinkham, W. H.: A deficiency of glucose-6-phosphate dehydrogenase

[92532] activity in lens from individuals with primaquine-sensitive erythrocytes. Bull.

[92533] Johns Hopkins Hosp. 109: 206-216, 1961.

[92534]

[92535] 380. Zuo, L.; Chen, E.; Du, C. S.; Chang, C. N.; Chiu, D. T. Y.:

[92536] Genetic study of Chinese G6PD variants by direct PCR sequencing. (Abstract) Blood 76

[92537] (suppl. 1): 51A, 1990.

[92538] 16161.

[92539] Antignac, C.; Gros, F.; Geffriaud, C.; Kashtan, C.; Proesmans,

[92540] W.; Tryggvason, K.; Gubler, M. C.: Alport syndrome (AS) and diffuse

- [92541] esophageal leiomyomatosis (DL): deletions in the 5-prime end and upstream
- [92542] region of the COL4A5 collagen gene. (Abstract) Am. J. Hum. Genet. 51
- [92543] (suppl.): A7, 1992.
- [92544] 16162.
- [92545] Antignac, C.; Knebelmann, B.; Drouot, L.; Gros, F.; Deschenes,
- [92546] G.; Hors-Cayla, M.-C.; Zhou, J.; Tryggvason, K.; Grunfeld, J.-P.;
- [92547] Broyer, M.; Gubler, M.-C.: Deletions in the COL4A5 collagen gene
- [92548] in X-linked Alport syndrome: characterization of the pathological
- [92549] transcripts in nonrenal cells and correlation with disease expression. J.
- [92550] Clin. Invest. 93: 1195-1207, 1994.
- [92551] 16163.
- [92552] Antignac, C.; Zhou, J.; Sanak, M.; Cochat, P.; Roussel, B.; Deschenes,
- [92553] G.; Gros, F.; Knebelmann, B.; Hors-Cayla, M.-C.; Tryggvason, K.; Gubler,

[92554] M.-C.: Alport syndrome and diffuse leiomyomatosis: deletions in the

[92555] 5-prime end of the COL4A5 collagen gene. *Kidney Int.* 42: 1178-1183,

[92556] 1992.

[92557] 16164.

[92558] Barker, D. F.; Denison, J. C.; Atkin, C. L.; Gregory, M. C.: Common

[92559] ancestry of three Ashkenazi-American families with Alport syndrome

[92560] and COL4A5 R1677Q. *Hum. Genet.* 99: 681-684, 1997.

[92561] 16165.

[92562] Barker, D. F.; Pruchno, C. J.; Jiang, X.; Atkin, C. L.; Stone,

[92563] E. M.; Denison, J. C.; Fain, P. R.; Gregory, M. C.: A mutation causing

[92564] Alport syndrome with tardive hearing loss is common in the western

[92565] United States. *Am. J. Hum. Genet.* 58: 1157-1165, 1996.

[92566] 16166.

[92567] Boye, E.; Vetrie, D.; Flinter, F.; Buckle, B.; Pihlajaniemi, T.;

- [92568] Hamalainen, E.-R.; Myers, J. C.; Bobrow, M.; Harris, A.:
Major rearrangements
- [92569] in the alpha-5-(IV) collagen gene in three patients with
Alport syndrome. Genomics 11:
- [92570] 1125-1132, 1991.
- [92571] 16167.
- [92572] Guo, C.; Van Damme, B.; Vanrenterghem, Y.; Devriendt,
K.; Cassiman,
- [92573] J.-J.; Marynen, P.: Severe Alport phenotype in a woman
with two missense
- [92574] mutations in the same COL4A5 gene and preponderant
inactivation of
- [92575] the X chromosome carrying the normal allele. J. Clin. In-
vest. 95:
- [92576] 1832-1837, 1995.
- [92577] 16168.
- [92578] Campuzano, V.; Montermini, L.; Molto, M. D.; Pianese, L.;
Cossee,
- [92579] M.; Cavalcanti, F.; Monros, E.; Rodius, F.; Duclos, F.; Mon-
ticelli,
- [92580] A.; Zara, F.; Canizares, J.; Koutnikova, H.; Bidichandani, S.
I.;

- [92581] Gellera, C.; Brice, A.; Trouillas, P.; De Michele, G.; Filla, A.;
- [92582] De Frutos, R.; Palau, F.; Patel, P. I.; Di Donato, S.; Mandel, J.
- [92583] –L.; Coccozza, S.; Koenig, M.; Pandolfo, M.: Friedreich's ataxia:
- [92584] autosomal recessive disease caused by an intronic GAA triplet repeat
- [92585] expansion. *Science* 271: 1423–1427, 1996.
- [92586] 16169.
- [92587] Boyd, M.; Lanyon, W. G.; Connor, J. M.: Screening for molecular
- [92588] pathologies in Lesch–Nyhan syndrome. *Hum. Mutat.* 2: 127–130, 1993.
- [92589] 16170.
- [92590] Brennand, J.; Chinault, A. C.; Konecki, D. S.; Melton, D. W.; Caskey,
- [92591] C. T.: Cloned cDNA sequences of the hypoxanthine–guanine phosphoribosyltransferase
- [92592] gene from a mouse neuroblastoma cell line found to have amplified
- [92593] genomic sequences. *Proc. Nat. Acad. Sci.* 79: 1950–1954, 1982.

[92594] 16171.

[92595] Cariello, N. F.; Scott, J. K.; Kat, A. G.; Thilly, W. G.; Keo-
havong,

[92596] P.: Resolution of a missense mutant in human genomic
DNA by denaturing

[92597] gradient gel electrophoresis and direct sequencing using
in vitro

[92598] DNA amplification: HPRT(Munich). Am. J. Hum. Genet. 42:
726-734,

[92599] 1988.

[92600] 16172.

[92601] Caskey, C. T.; Kruh, G. D.: The HPRT locus: review. Cell
16:

[92602] 1-9, 1979.

[92603] 16173.

[92604] Chinault, A. C.; Caskey, C. T.: The hypoxanthine phospho-
ribosyltransferase

[92605] gene: a model for the study of mutation in mammalian
cells. Prog.

[92606] Nucleic Acids Res. Molec. Biol. 31: 317-465, 1984.

[92607] 16174.

- [92608] Cianfriglia, M.; Miggiano, V. C.; Meo, T.; Muller, H. J.;
Muller,
- [92609] E.; Battistuzzi, G.: Evidence for syntenic between the rabbit
gene
- [92610] loci coding for HPRT, PGK and G6PD in mouse-rabbit so-
matic cell hybrids.
- [92611] (Abstract) Cytogenet. Cell Genet. 25: 142 only, 1979.
- [92612] 16175.
- [92613] Colgin, L. M.; Hackmann, A. F. M.; Emond, M. J.; Monnat,
R. J.,
- [92614] Jr.: The unexpected landscape of in vivo somatic mutation
in a human
- [92615] epithelial cell lineage. Proc. Nat. Acad. Sci. 99:
1437-1442, 2002.
- [92616] 16176.
- [92617] Cox, R. P.; Krauss, M. R.; Balis, M. E.; Dancis, J.: Evidence
- [92618] for transfer of enzyme product as the basis of metabolic
cooperation
- [92619] between tissue culture fibroblasts of Lesch-Nyhan disease
and normal
- [92620] cells. Proc. Nat. Acad. Sci. 67: 1573-1579, 1970.

[92621] 16177.

[92622] Dancis, J.; Yip, L. C.; Cox, R. P.; Piomelli, S.; Balis, M. E.

[92623] : Disparate enzyme activity in erythrocytes and leukocytes:
a variant

[92624] of hypoxanthine phosphoribosyltransferase deficiency
with an unstable

[92625] enzyme. J. Clin. Invest. 52: 2068–2074, 1973.

[92626] 16178.

[92627] Davidson, B. L.: Personal Communication. Ann Arbor,
Mich.

[92628] 1990.

[92629] 16179.

[92630] Davidson, B. L.; Chin, S.-J.; Wilson, J. M.; Kelley, W. N.;
Palella,

[92631] T. D.: Hypoxanthine–guanine phosphoribosyltransferase:
genetic evidence

[92632] for identical mutations in two partially deficient subjects.
J. Clin.

[92633] Invest. 82: 2164–2167, 1988.

[92634] 16180.

[92635] Davidson, B. L.; Pashmforoush, M.; Kelley, W. N.; Palella, T.

[92636] D.: Genetic basis of hypoxanthine–guanine phosphoribosyltransferase

[92637] deficiency in a patient with the Lesch–Nyhan syndrome (HPRT–Flint). Gene 63:

[92638] 331–336, 1988.

[92639] 16181.

[92640] Davidson, B. L.; Pashmforoush, M.; Kelley, W. N.; Palella, T.

[92641] D.: Human hypoxanthine–guanine phosphoribosyltransferase deficiency:

[92642] the molecular defect in a patient with gout (HPRT–Ashville). J. Biol.

[92643] Chem. 264: 520–525, 1989.

[92644] 16182.

[92645] Davidson, B. L.; Tarle, S. A.; Palella, T. D.; Kelley, W. N.:

[92646] Molecular basis of hypoxanthine–guanine phosphoribosyltransferase

[92647] deficiency in 10 subjects determined by direct sequencing of amplified

[92648] transcripts. J. Clin. Invest. 84: 342–346, 1989.

- [92649] 16183.
- [92650] Davidson, B. L.; Tarle, S. A.; Van Antwerp, M.; Gibbs, D. A.;
- [92651] Watts, R. W. E.; Kelley, W. N.; Palella, T. D.: Identification
of
- [92652] 17 independent mutations responsible for human hypox-
anthine-guanine
- [92653] phosphoribosyltransferase (HPRT) deficiency. Am. J. Hum.
Genet. 48:
- [92654] 951-958, 1991.
- [92655] 16184.
- [92656] De Gregorio, L.; Nyhan, W. L.; Serafin, E.; Chamoles, N. A.:
- [92657] An unexpected affected female patient in a classical
Lesch-Nyhan family. Molec.
- [92658] Genet. Metab. 69: 263-268, 2000.
- [92659] 16185.
- [92660] Demars, R. I.; Sarto, G. E.; Felix, J. S.; Benke, P.: Lesch-
Nyhan
- [92661] mutation: prenatal detection with amniotic fluid cells. Sci-
ence 164:
- [92662] 1303-1305, 1969.

[92663] 16186.

[92664] Dempsey, J. L.; Morley, A. A.; Seshadri, R. S.; Emmerson, B. T.;

[92665] Gordon, R.; Bhagat, C. I.: Detection of the carrier state for an

[92666] X-linked disorder, the Lesch-Nyhan syndrome, by the use of lymphocyte

[92667] cloning. Hum. Genet. 64: 288-290, 1983.

[92668] 16187.

[92669] Dobrovic, A.; Gareau, P.; Seifert, A.-M.; Messing, K.; Bradley,

[92670] W. E. C.: A HindIII RFLP for the HPRT pseudogene on chromosome 3

[92671] (HPRT1). Nucleic Acids Res. 15: 1346, 1987.

[92672] 16188.

[92673] Doetschman, T.; Gregg, R. G.; Maeda, N.; Hooper, M. L.; Melton,

[92674] D. W.; Thompson, S.; Smithies, O.: Targetted correction of a mutant

[92675] HPRT gene in mouse embryonic stem cells. Nature 330: 576-578, 1987.

[92676] 16189.

[92677] Echard, G.; Gillois, M.: G6PD-PGK-GAL-HPRT syntenry in the rabbit,

[92678] *Oryctolagus cuniculus*. (Abstract) Cytogenet. Cell Genet. 25: 148-149,

[92679] 1979.

[92680] 16190.

[92681] Edwards, A.; Caskey, C. T.: Personal Communication. Houston,

[92682] Tex. 8/1990.

[92683] 16191.

[92684] Edwards, A.; Voss, H.; Rice, P.; Civitello, A.; Stegemann, J.;

[92685] Schwager, C.; Zimmermann, J.; Erfle, H.; Caskey, C. T.; An-sorge, W.

[92686] : Automated DNA sequencing of the human HPRT locus. Genomics 6:

[92687] 593-608, 1990.

[92688] 16192.

[92689] Emmerson, B. T.; Thompson, C. J.; Wallace, D. C.: Partial deficiency

- [92690] hypoxanthine-guanine phosphoribosyltransferase: intermediate enzyme
- [92691] deficiency in heterozygote red cells. *Ann. Intern. Med.* 76: 285-288,
- [92692] 1972.
- [92693] 16193.
- [92694] Engle, S. J.; Womer, D. E.; Davies, P. M.; Boivin, G.; Sahota, A.; Simmonds, H. A.; Stambrook, P. J.; Tischfield, J. A.: HPRT-APRT-deficient
- [92696] mice are not a model for Lesch-Nyhan syndrome. *Hum. Mol. Genet.* 5:
- [92697] 1607-1610, 1996.
- [92698] 16194.
- [92699] Epstein, C. J.: Expression of the mammalian X chromosome before
- [92700] and after fertilization. *Science* 175: 1467-1468, 1972.
- [92701] 16195.
- [92702] Ernst, M.; Zametkin, A. J.; Matochik, J. A.; Pascualvaca, D.;
- [92703] Jons, P. H.; Hardy, K.; Hankerson, J. G.; Doudet, D. J.; Cohen, R.

- [92704] M.: Presynaptic dopaminergic deficits in Lesch–Nyhan disease. New
- [92705] Eng. J. Med. 334: 1568–1572, 1996.
- [92706] 16196.
- [92707] Fenwick, R. G., Jr.: Reversion of a mutation affecting the molecular
- [92708] weight of HGPRT: intragenic suppression and localization of X-linked
- [92709] genes. Somat. Cell Genet. 6: 477–494, 1980.
- [92710] 16197.
- [92711] Fox, I. H.; Dwosh, I. L.; Marchant, P. J.; Lacroix, S.; Moore,
- [92712] M. R.; Omura, S.; Wyhofsky, V.: Hypoxanthine–guanine phosphoribosyltransferase:
- [92713] characterization of a mutant in a patient with gout. J. Clin. Invest. 56:
- [92714] 1239–1249, 1975.
- [92715] 16198.
- [92716] Francke, U.; Bakay, B.; Connor, J. D.; Coldwell, J. G.; Nyhan,
- [92717] W. L.: Linkage relationships of X-linked enzymes glucose–6-phosphate

- [92718] dehydrogenase and hypoxanthine guanine phosphoribosyltransferase. Am.
- [92719] J. Hum. Genet. 26: 512–522, 1974.
- [92720] 16199.
- [92721] Francke, U.; Felsenstein, J.; Gartler, S. M.; Migeon, B. R.; Dancis,
- [92722] J.; Seegmiller, J. E.; Bakay, B.; Nyhan, W. L.: The occurrence of
- [92723] new mutants in the X-linked recessive Lesch–Nyhan disease. Am. J.
- [92724] Hum. Genet. 28: 123–137, 1976.
- [92725] 16200.
- [92726] Francke, U.; Felsenstein, J.; Gartler, S. M.; Nyhan, W. L.; Seegmiller,
- [92727] J. E.: Answer to criticism of Morton and Lalouel. (Letter) Am. J.
- [92728] Hum. Genet. 29: 307–310, 1977.
- [92729] 16201.
- [92730] Gartler, S. M.; Francke, U.: Half–chromatid mutation: transmission
- [92731] in humans? Am. J. Hum. Genet. 27: 218–223, 1975.

- [92732] 16202.
- [92733] Gibbs, R. A.; McFadyen, I. R.; Crawford, M. d'A.; de Muinck Keizer, E. E.; Headhouse-Benson, C. M.; Wilson, T. M.; Farrant, P. H.: First-trimester diagnosis of Lesch-Nyhan syndrome. *Lancet* II: 1180-1183, 1984.
- [92734] 16203.
- [92735] Graham, G. W.; Aitken, D. A.; Connor, J. M.: Prenatal diagnosis by enzyme analysis in 15 pregnancies at risk for the Lesch-Nyhan syndrome. *Prenatal Diag.* 16: 647-651, 1996.
- [92736] 16204.
- [92737] Greene, M. L.; Nyhan, W. L.; Seegmiller, J. E.: Hypoxanthine-guanine phosphoribosyltransferase deficiency and Xg blood group. *Am. J. Hum. Genet.* 22: 50-54, 1970.
- [92738] 16205.

- [92745] Henderson, J. F.; Kelley, W. N.; Rosenbloom, F. M.; Seeg-
miller,
[92746] J. E.: Inheritance of purine phosphoribosyltransferases in
man. Am.
[92747] J. Hum. Genet. 21: 61-70, 1969.
[92748] 16206.
[92749] Hoefnagel, D.; Andrew, E. D.; Mireault, N. G.; Berndt, W.
O.:
[92750] Hereditary choreoathetosis, self-mutilation and hyper-
uricemia in young
[92751] males. New Eng. J. Med. 273: 130-135, 1965.
[92752] 16207.
[92753] Lloyd, K. G.; Hornykiewicz, O.; Davidson, L.; Shannak, K.;
Farley,
[92754] I.; Goldstein, M.; Shibuya, M.; Kelley, W. N.; Fox, I. H.: Bio-
chemical
[92755] evidence of dysfunction of brain neurotransmitters in the
Lesch-Nyhan
[92756] syndrome. New Eng. J. Med. 305: 1106-1111, 1981.
[92757] 16208.
[92758] Migeon, B. R.; Der Kaloustian, V. M.; Nyhan, W. L.; Young,

W.

- [92759] J.; Childs, B.: X-linked hypoxanthine-guanine phosphoribosyl transferase deficiency: heterozygote has two clonal populations. *Science* 160: 425-427, 1968.
- [92761] 16209.
- [92762] 16209.
- [92763] Morton, N. E.; Lalouel, J. M.: Genetic epidemiology of Lesch-Nyhan disease. (Letter) *Am. J. Hum. Genet.* 29: 304-307, 1977.
- [92764] 16210.
- [92765] 16210.
- [92766] Nabholz, M.; Miggiano, V.; Bodmer, W.: Genetic analysis with human-mouse somatic cell hybrids. *Nature* 223: 358-363, 1969.
- [92767] 16211.
- [92768] 16211.
- [92769] Nyhan, W. L.; Bakay, B.; Connor, J. D.; Marks, J. F.; Keele, D. K.: Hemizygous expression of glucose-6-phosphate dehydrogenase in erythrocytes of heterozygotes for the Lesch-Nyhan syn-

drome. Proc.

[92772] Nat. Acad. Sci. 65: 214–218, 1970.

[92773] 16212.

[92774] Nyhan, W. L.; Olivier, W. J.; Lesch, M.: A familial disorder

[92775] of uric acid metabolism and central nervous system function. J. Pediat. 67:

[92776] 257–263, 1965.

[92777] 16213.

[92778] Nyhan, W. L.; Wong, D. F.: New approaches to understanding Lesch–Nyhan

[92779] disease. (Editorial) New Eng. J. Med. 334: 1602–1604, 1996.

[92780] 16214.

[92781] Burwinkel, B.; Amat, L.; Gray, R. G. F.; Matsuo, N.; Muroya, K.;

[92782] Narisawa, K.; Sokol, R. J.; Vilaseca, M. A.; Kilimann, M. W.: Variability

[92783] of biochemical and clinical phenotype in X-linked liver glycogenosis

[92784] with mutations in the phosphorylase kinase PHKA2 gene. Hum. Genet. 102:

[92785] 423–429, 1998.

[92786] 16215.

[92787] Burwinkel, B.; Shin, Y. S.; Bakker, H. D.; Deutsch, J.;
Lozano,

[92788] M. J.; Maire, I.; Kilimann, M. W.: Mutation hotspots in the
PHKA2

[92789] gene in X-linked liver glycogenosis due to phosphorylase
kinase deficiency

[92790] with atypical activity in blood cells (XLG2). Hum. Molec.
Genet. 5:

[92791] 653–658, 1996.

[92792] 16216.

[92793] Davidson, J. J.; Ozcelik, T.; Hamacher, C.; Willems, P. J.;
Francke,

[92794] U.; Kilimann, M. W.: cDNA cloning of a liver isoform of the
phosphorylase

[92795] kinase alpha subunit and mapping of the gene to
Xp22.2–p22.1, the

[92796] region of human X-linked liver glycogenosis. Proc. Nat.
Acad. Sci. 89:

[92797] 2096–2100, 1992.

- [92798] 16217.
- [92799] Davisson, M. T.: X-linked genetic homologies between mouse and
- [92800] man. *Genomics* 1: 213–227, 1987.
- [92801] 16218.
- [92802] Garibaldi, L. R.; Borrone, C.; De Martini, I.; Battistini, E.:
- [92803] Dextrothyroxine treatment of phosphorylase-kinase deficiency glycogenosis
- [92804] in four boys. *Helv. Paediat. Acta* 33: 435–444, 1978.
- [92805] 16219.
- [92806] Goji, K.; Morishita, Y.; Kodama, S.; Takahashi, T.; Matsuo, T.
- [92807] : Lymphocyte phosphorylase kinase activities in the sex-linked form
- [92808] of liver phosphorylase kinase deficiency. *Europ. J. Pediat.* 143:
- [92809] 179–182, 1985.
- [92810] 16220.
- [92811] Hendrickx, J.; Bosshard, N. U.; Willems, P.; Gitzelmann, R.: Clinical,

- [92812] biochemical and molecular findings in a patient with X-linked liver
- [92813] glycogenosis followed for 40 years. *Europ. J. Pediat.* 157: 919–923,
- [92814] 1998.
- [92815] 16221.
- [92816] Hendrickx, J.; Coucke, P.; Bossuyt, P.; Wauters, J.; Raeymaekers,
- [92817] P.; Marchau, F.; Smit, G. P. A.; Stolte, I.; Sardharwalla, I. B.;
- [92818] Berthelot, J.; Van den Bergh, I.; Berger, R.; Van Broeckhoven, C.;
- [92819] Baussan, C.; Wapenaar, M.; Fernandes, J.; Willems, P. J.: X-linked
- [92820] liver glycogenosis: localization and isolation of a candidate gene. *Hum.*
- [92821] *Molec. Genet.* 2: 583–589, 1993.
- [92822] 16222.
- [92823] Hendrickx, J.; Coucke, P.; Dams, E.; Lee, P.; Odievre, M.; Corbeel,
- [92824] L.; Fernandes, J. F.; Willems, P. J.: Mutations in the phosphorylase

- [92825] kinase gene PHKA2 are responsible for X-linked liver glycogen storage
- [92826] disease. Hum. Molec. Genet. 4: 77-83, 1995.
- [92827] 16223.
- [92828] Hendrickx, J.; Coucke, P.; Hors-Cayla, M.-C.; Smit, G. P. A.;
- [92829] Shin, Y. S.; Deutsch, J.; Smeitink, J.; Berger, R.; Lee, P.; Fernandes,
- [92830] J.; Willems, P. J.: Localization of a new type of X-linked liver
- [92831] glycogenosis to the chromosomal region Xp22 containing the liver alpha-subunit
- [92832] of phosphorylase kinase (PHKA2). Genomics 21: 620-625, 1994.
- [92833] 16224.
- [92834] Hendrickx, J.; Coucke, P.; Raeymaekers, P.; Willems, P. J.: X-linked
- [92835] liver glycogenosis: localization and isolation of a strong candidate
- [92836] gene. (Abstract) Am. J. Hum. Genet. 51 (suppl.): A190 only, 1992.
- [92837] 16225.

- [92838] Hendrickx, J.; Dams, E.; Coucke, P.; Lee, P.; Fernandes, J.; Willems,
- [92839] P. J.: X-linked liver glycogenosis type II (XLG II) is caused by
- [92840] mutations in PHKA2, the gene encoding the liver alpha subunit of phosphorylase
- [92841] kinase. Hum. Molec. Genet. 5: 649-652, 1996.
- [92842] 16226.
- [92843] Hendrickx, J.; Lee, P.; Keating, J. P.; Carton, D.; Sardharwalla,
- [92844] I. B.; Tuchman, M.; Baussan, C.; Willems, P. J.: Complete genomic
- [92845] structure and mutational spectrum of PHKA2 in patients with X-linked
- [92846] liver glycogenosis type I and II. Am. J. Hum. Genet. 64: 1541-1549,
- [92847] 1999.
- [92848] 16227.
- [92849] Hers, H.-G.; Van Hoof, F.; de Barse, T.: Glycogen storage diseases. In:
- [92850] Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D.: The Metabolic

[92851] Basis of Inherited Disease. New York: McGraw-Hill (pub.)
(6th

[92852] ed.) I: 1989. Pp. 425-452.

[92853] 16228.

[92854] Hug, G.: Personal Communication. Cincinnati, Ohio 1974.

[92855] 16229.

[92856] Hug, G.; Schubert, W. K.; Chuck, G.: Deficient activity of
dephosphophosphorylase

[92857] kinase and accumulation of glycogen in the liver. J. Clin.
Invest. 48:

[92858] 704-715, 1969.

[92859] 16230.

[92860] Huijing, F.: Phosphorylase kinase deficiency. Biochem.
Genet. 4:

[92861] 187-194, 1970.

[92862] 16231.

[92863] Huijing, F.: Phosphorylase kinase in leucocytes of normal
subjects

[92864] and of patients with glycogen-storage disease. Biochim.
Biophys.

- [92865] Acta 148: 601–603, 1967.
- [92866] 16232.
- [92867] Huijing, F.: Glycogen–storage disease type VIa: low phosphorylase
- [92868] kinase activity caused by a low enzyme–substrate affinity. Biochim.
- [92869] Biophys. Acta 206: 199–201, 1970.
- [92870] 16233.
- [92871] Huijing, F.; Fernandez, J.: Liver glycogenosis and phosphorylase
- [92872] kinase deficiency. (Letter) Am. J. Hum. Genet. 22: 484–485, 1970.
- [92873] 16234.
- [92874] Heidet, L.; Dahan, K.; Zhou, J.; Xu, Z.; Cochat, P.; Gould, J.
- [92875] D. M.; Leppig, K. A.; Proesmans, W.; Guyot, C.; Guillot, M.; Roussel,
- [92876] B.; Tryggvason, K.; Grunfeld, J.–P.; Gubler, M.–C.; Antignac, C.:
- [92877] Deletions of both alpha–5(IV) and alpha–6(IV) collagen genes in Alport
- [92878] syndrome and in Alport syndrome associated with smooth

muscle tumours. Hum.

[92879] Molec. Genet. 4: 99–108, 1995.

[92880] 16235.

[92881] Morton, C. C.; Nussenzweig, M. C.; Sousa, R.; Sorenson, G. D.;

[92882] Pettengill, O. S.; Shows, T. B.: Mapping and characterization of

[92883] an X-linked processed gene related to MYCL1. Genomics 4: 367–375,

[92884] 1989.

[92885] 16236.

[92886] Ambler, M. W.; Neave, C.; Singer, D. B.: X-linked recessive myotubular

[92887] myopathy. II. Muscle morphology and human myogenesis. Hum. Path. 15:

[92888] 1107–1120, 1984.

[92889] 16237.

[92890] Ambler, M. W.; Neave, C.; Tutschka, B. G.; Pueschel, S. M.; Orson,

[92891] J. M.; Singer, D. B.: X-linked recessive myotubular myopathy. I.

- [92892] Clinical and pathologic findings in a family. Hum. Path.
15: 566–574,
[92893] 1984.
- [92894] 16238.
- [92895] Askanas, V.; Engel, W. K.; Reddy, N. B.; Barth, P. G.; Beth-
lem,
[92896] J.; Krauss, D. R.; Hibberd, M. E.; Lawrence, J. V.; Carter, L.
S.
- [92897] : X-linked recessive congenital muscle fiber hypotrophy
with central
[92898] nuclei: abnormalities of growth and adenylate cyclase in
muscle tissue
[92899] cultures. Arch. Neurol. 36: 604–609, 1979.
- [92900] 16239.
- [92901] Barth, P. G.; Dubowitz, V.: X-linked myotubular myopa-
thy--a long-term
[92902] follow-up study. Europ. J. Paediat. Neurol. 2: 49–56,
1998.
- [92903] 16240.
- [92904] Ilyin, G. P.; Rialland, M.; Pigeon, C.; Guguen-Guillouzo, C.:
[92905] cDNA cloning and expression analysis of new members of

the mammalian

[92906] F-box protein family. Genomics 67: 40–47, 2000.

[92907] 16241.

[92908] Chan, A. M.–L.; Miki, T.; Meyers, K. A.; Aaronson, S. A.: A human

[92909] oncogene of the RAS superfamily unmasked by expression cDNA cloning. Proc.

[92910] Nat. Acad. Sci. 91: 7558–7562, 1994.

[92911] 16242.

[92912] Drivas, G. T.; Shih, A.; Coutavas, E.; Rush, M. G.; D'Eustachio,

[92913] P.: Characterization of four novel ras-like genes expressed in a

[92914] human teratocarcinoma cell line. Molec. Cell. Biol. 10: 1793–1798,

[92915] 1990.

[92916] 16243.

[92917] Brugnera, E.; Georgiev, O.; Radtke, F.; Heuchel, R.; Baker, E.;

[92918] Sutherland, G. R.; Schaffner, W.: Cloning, chromosomal mapping and

- [92919] characterization of the human metal-regulatory transcription factor
- [92920] MTF-1. Nucleic Acids Res. 22: 3167-3173, 1994.
- [92921] 16244.
- [92922] Gunes, C.; Heuchel, R.; Georgiev, O.; Muller, K.-H.; Lichtlen,
- [92923] P.; Bluthmann, H.; Marino, S.; Aguzzi, A.; Schaffner, W.: Embryonic
- [92924] lethality and liver degeneration in mice lacking the metal-responsive
- [92925] transcriptional activator MTF-1. EMBO J. 17: 2846-2854, 1998.
- [92926] 16245.
- [92927] Radtke, F.; Heuchel, R.; Georgiev, O.; Hergersberg, M.; Gariglio,
- [92928] M.; Dembic, Z.; Schaffner, W.: Cloned transcription factor MTF-1
- [92929] activates the mouse metallothionein I promoter. EMBO J. 12: 1355-1362,
- [92930] 1993.
- [92931] 16246.

- [92932] Westin, W.; Schaffner, W.: A zinc-responsive factor interacts
- [92933] with a metal-regulated enhancer element (MRE) of the mouse metallothionein-I
- [92934] gene. EMBO J. 7: 3763-3770, 1988.
- [92935] 16247.
- [92936] Andersson, P.; McGuire, J.; Rubio, C.; Gradin, K.; Whitelaw, M.
- [92937] L.; Pettersson, S.; Hanberg, A.; Poellinger, L.: A constitutively
- [92938] active dioxin/aryl hydrocarbon receptor induces stomach tumors. Proc.
- [92939] Nat. Acad. Sci. 99: 9990-9995, 2002.
- [92940] 16248.
- [92941] Ema, M.; Matsushita, N.; Sogawa, K.; Ariyama, T.; Inazawa, J.;
- [92942] Nemoto, T.; Ota, M.; Oshimura, M.; Fujii-Kuriyama, Y.: Human arylhydrocarbon
- [92943] receptor: functional expression and chromosomal assignment to 7p21. J.
- [92944] Biochem. 116: 845-851, 1994.

[92945] 16249.

[92946] Le Beau, M. M.; Carver, L. A.; Espinosa, R., III; Schmidt, J. V.;

[92947] Bradfield, C. A.: Chromosomal localization of the human AHR locus

[92948] encoding the structural gene for the Ah receptor to 7p21-p15. Cytogenet.

[92949] Cell Genet. 66: 172-176, 1994.

[92950] 16250.

[92951] Micka, J.; Milatovich, A.; Menon, A.; Grabowski, G. A.; Puga, A.;

[92952] Nebert, D. W.: Human Ah receptor (AHR) gene: localization to 7p15

[92953] and suggestive correlation of polymorphism with CYP1A1 inducibility. Pharmacogenetics 7:

[92954] 95-101, 1997.

[92955] 16251.

[92956] Shimizu, Y.; Nakatsuru, Y.; Ichinose, M.; Takahashi, Y.; Kume,

[92957] H.; Mimura, J.; Fujii-Kuriyama, Y.; Ishikawa, T.: Benzo[a]pyrene

- [92958] carcinogenicity is lost in mice lacking the aryl hydrocarbon receptor. Proc.
- [92959] Nat. Acad. Sci. 97: 779–782, 2000.
- [92960] 16252.
- [92961] Chen, P.; Hao, W.; Rife, L.; Wang, X. P.; Shen, D.; Chen, J.; Ogden,
- [92962] T.; Van Boemel, G. B.; Wu, L.; Yang, M.; Fong, H. K. W.: A photic
- [92963] visual cycle of rhodopsin regeneration is dependent on Rgr. Nature
- [92964] Genet. 28: 256–260, 2001.
- [92965] 16253.
- [92966] Chen, X.-N.; Korenberg, J. R.; Jiang, M.; Shen, D.; Fong, H. K.
- [92967] W.: Localization of the human RGR opsin gene to chromosome 10q23. Hum.
- [92968] Genet. 97: 720–722, 1996.
- [92969] 16254.
- [92970] Jiang, M.; Pandey, S.; Fong, H. K. W.: An opsin homologue in the
- [92971] retina and pigment epithelium. Invest. Ophthalm. Visual Sci.

34:

[92972] 3669–3678, 1993.

[92973] 16255.

[92974] Morimura, H.; Saindelle–Ribeaudau, F.; Berson, E. L.;
Dryja, T.

[92975] P.: Mutations in RGR, encoding a light-sensitive opsin ho-
mologue,

[92976] in patients with retinitis pigmentosa. (Letter) Nature
Genet. 23:

[92977] 393–394, 1999.

[92978] 16256.

[92979] Shen, D.; Jiang, M.; Hao, W.; Tao, L.; Salazar, M.; Fong, H.
K.

[92980] W.: A human opsin-related gene that encodes a retinalde-
hyde-binding

[92981] protein. Biochemistry 33: 13117–13125, 1994.

[92982] 16257.

[92983] Sasada, R.; Ono, Y.; Taniyama, Y.; Shing, Y.; Folkman, J.;
Igarashi,

[92984] K.: Cloning and expression of cDNA encoding human be-
tacellulin, a

- [92985] new member of the EGF family. *Biochem. Biophys. Res. Commun.* 190: 1173–1179, 1993.
- [92986] 16258.
- [92987] Watanabe, T.; Shintani, A.; Nakata, M.; Shing, Y.; Folkman, J.;
- [92988] Igarashi, K.; Sasada, R.: Recombinant human betacellulin: molecular structure, biological activities, and receptor interaction. *J. Biol. Chem.* 269: 9966–9973, 1994.
- [92989] 16259.
- [92990] Asano, H.; Ishida, A.; Hasegawa, M.; Ono, T.; Yoshida, M. C.; Taniguchi, M.; Kanno, M.: The mouse Mel-18 'RING-finger' gene: genomic organization, promoter analysis and chromosomal assignment. *DNA Seq.* 3: 369–377, 1993.
- [92991] 16260.
- [92992] Ishida, A.; Asano, H.; Hasegawa, M.; Koseki, H.; Ono, T.;

Yoshida,

[92999] M. C.; Taniguchi, M.; Kanno, M.: Cloning and chromosome mapping of

[93000] the human Mel-18 gene which encodes a DNA-binding protein with a new

[93001] 'RING-finger' motif. Gene 129: 249-255, 1993.

[93002] 16261.

[93003] Tagawa, M.; Sakamoto, T.; Shigemoto, K.; Matsubara, H.; Tamura,

[93004] Y.; Ito, T.; Nakamura, I.; Okitsu, A.; Imai, K.; Taniguchi, M.: Expression

[93005] of novel DNA-binding protein with zinc finger structure in various

[93006] tumor cells. J. Biol. Chem. 265: 20021-20026, 1990.

[93007] 16262.

[93008] Hara, E.; Yamaguchi, T.; Nojima, H.; Ide, T.; Campisi, J.; Okayama,

[93009] H.; Oda, K.: Id-related genes encoding helix-loop-helix proteins

[93010] are required for G1 progression and are repressed in senescent human

[93011] fibroblasts. J. Biol. Chem. 269: 2139-2145, 1994.

- [93012] 16263.
- [93013] Mathew, S.; Chen, W.; Murty, V. V. V. S.; Benezra, R.; Chaganti,
- [93014] R. S. K.: Chromosomal assignment of human ID1 and ID2 genes. *Genomics* 30:
- [93015] 385–387, 1995.
- [93016] 16264.
- [93017] Chen, M. X.; McPartlin, A. E.; Brown, L.; Chen, Y. H.; Barker,
- [93018] H. M.; Cohen, P. T. W.: A novel human protein serine/threonine phosphatase,
- [93019] which possesses four tetratricopeptide repeat motifs and localizes
- [93020] to the nucleus. *EMBO J.* 13: 4278–4290, 1994.
- [93021] 16265.
- [93022] Xu, X.; Lagercrantz, J.; Zickert, P.; Bajalica-Lagercrantz, S.;
- [93023] Zetterberg, A.: Chromosomal localization and 5-prime sequence of
- [93024] the human protein serine/threonine phosphatase 5-prime gene. *Biochem.*
- [93025] *Biophys. Res. Commun.* 218: 514–517, 1996.

- [93026] 16266.
- [93027] Yong, W. H.; Ueki, K.; Chou, D.; Reeves, S. A.; von Deimling, A.;
- [93028] Gusella, J. F.; Mohrenweiser, H. W.; Buckler, A. J.; Louis, D. N.
- [93029] : Cloning of a highly conserved human protein serine-threonine phosphatase
- [93030] gene from the glioma candidate region on chromosome 19q13.3. Genomics 29:
- [93031] 533-536, 1995.
- [93032] 16267.
- [93033] Breitbart, R. E.; Liang, C.; Smoot, L. B.; Laheru, D. A.; Mahdavi,
- [93034] V.; Nadal-Ginard, B.: A fourth human MEF2 transcription factor, hMEF2D,
- [93035] is an early marker of the myogenic lineage. Development 118: 1095-1106,
- [93036] 1993.
- [93037] 16268.
- [93038] Hobson, G. M.; Krahe, R.; Garcia, E.; Siciliano, M. J.; Fumanage,

- [93039] V. L.: Regional chromosomal assignments for four members of the MADS
- [93040] domain transcription enhancer factor 2 (MEF2) gene family to human
- [93041] chromosomes 15q26, 19p12, 5q14, and 1q12-q23. Genomics 29: 704-711,
- [93042] 1995.
- [93043] 16269.
- [93044] Mao, Z.; Bonni, A.; Xia, F.; Nadal-Vicans, M.; Greenberg, M. E.
- [93045] : Neuronal activity-dependent cell survival mediated by transcription
- [93046] factor MEF2. Science 286: 785-790, 1999.
- [93047] 16270.
- [93048] Martin, J. F.; Miano, J. M.; Hustad, C. M.; Copeland, N. G.; Jenkins,
- [93049] N. A.; Olson, E. N.: A Mef2 gene that generates a muscle-specific
- [93050] isoform via alternative mRNA splicing. Molec. Cell. Biol. 14: 1647-1656,
- [93051] 1994.

- [93052] 16271.
- [93053] Molkenstin, J. D.; Black, B. L.; Martin, J. F.; Olson, E. N.: Co-operative
activation of muscle gene expression by MEF2 and myogenic bHLH proteins. Cell 83:
1125–1136, 1995.
- [93055] 16272.
- [93056] Naya, F. J.; Black, B. L.; Wu, H.; Bassel-Duby, R.; Richardson,
J. A.; Hill, J. A.; Olson, E. N.: Mitochondrial deficiency and cardiac
sudden death in mice lacking the MEF2A transcription factor. Nature
Med. 15Oct, 2002. Note: Advance Electronic Publication.
- [93060] 16273.
- [93061] Pollock, R.; Treisman, R.: Human SRF-related proteins: DNA-binding
properties and potential regulatory targets. Genes Dev. 5:
2327–2341,
1991.
- [93064] 16274.

- [93066] Suzuki, E.; Lowry, J.; Sonoda, G.; Testa, J. R.; Walsh, K.:
Structures
- [93067] and chromosome locations of the human MEF2A gene and
a pseudogene
- [93068] MEF2AP. Cytogenet. Cell Genet. 73: 244–249, 1996.
- [93069] 16275.
- [93070] Yu, Y.-T.; Breitbart, R. E.; Smoot, L. B.; Lee, Y.; Mahdavi,
V.;
- [93071] Nadal-Ginard, B.: Human myocyte-specific enhancer fac-
tor 2 comprises
- [93072] a group of tissue-restricted MADS box transcription fac-
tors. Genes
- [93073] Dev. 6: 1783–1798, 1992.
- [93074] 16276.
- [93075] Youn, H.-D.; Sun, L.; Prywes, R.; Liu, J. O.: Apoptosis of T
cells
- [93076] mediated by Ca(2+)-induced release of the transcription
factor MEF2. Science 286:
- [93077] 790–793, 1999.
- [93078] 16277.
- [93079] Krainc, D.; Haas, M.; Ward, D. C.; Lipton, S. A.; Bruns, G.;

Leifer,

[93080] D.: Assignment of human myocyte-specific enhancer binding factor

[93081] 2C (hMEF2C) to human chromosome 5q14 and evidence that MEF2C is evolutionarily

[93082] conserved. Genomics 29: 809–811, 1995.

[93083] 16278.

[93084] Leifer, D.; Krainc, D.; Yu, Y.-T.; McDermott, J.; Breitbart, R.

[93085] E.; Heng, J.; Neve, R. L.; Kosofsky, B.; Nadal-Ginard, B.; Lipton,

[93086] S. A.: MEF2C, a MADS/MEF2-family transcription factor expressed in

[93087] a laminar distribution in cerebral cortex. Proc. Nat. Acad. Sci. 90:

[93088] 1546–1550, 1993.

[93089] 16279.

[93090] Holmbeck, K.; Bianco, P.; Caterina, J.; Yamada, S.; Kromer, M.;

[93091] Kuznetsov, S. A.; Mankani, M.; Robey, P. G.; Poole, A. R.; Pidoux,

[93092] I.; Ward, J. M.; Birkedal-Hansen, H.: MT1-MMP-deficient

mice develop

[93093] dwarfism, osteopenia, arthritis, and connective tissue disease due

[93094] to inadequate collagen turnover. Cell 99: 81–92, 1999.

[93095] 16280.

[93096] Sato, H.; Takino, T.; Okada, Y.; Cao, J.; Shinagawa, A.; Yamamoto,

[93097] E.; Seiki, M.: A matrix metalloproteinase expressed on the surface

[93098] of invasive tumor cells. Nature 370: 61–65, 1994.

[93099] 16281.

[93100] Takino, T.; Sato, H.; Yamamoto, E.; Seiki, M.: Cloning of a human

[93101] gene potentially encoding a novel matrix metalloproteinase having

[93102] a C-terminal transmembrane domain. Gene 155: 293–298, 1995.

[93103] 16282.

[93104] McCright, B.; Rivers, A. M.; Audlin, S.; Virshup, D. M.: The B56

[93105] family of protein phosphatase 2A (PP2A) regulatory sub-

units encodes

[93106] differentiation-induced phosphoproteins that target PP2A to both nucleus

[93107] and cytoplasm. J. Biol. Chem. 271: 22081-22089, 1996.

[93108] 16283.

[93109] Van Hoof, C.; Aly, M. S.; Garcia, A.; Cayla, X.; Cassiman, J. J.;

[93110] Merlevede, W.; Goris, J.: Structure and chromosomal localization

[93111] of the human gene of the phosphotyrosyl phosphatase activator (PTPA)

[93112] of protein phosphatase 2A. Genomics 28: 261-272, 1995.

[93113] 16284.

[93114] Lory, P.; Ophoff, R. A.; Nahmias, J.: Towards a unified nomenclature

[93115] describing voltage-gated calcium channel genes. Hum. Genet. 100:

[93116] 149-150, 1997.

[93117] 16285.

[93118] Rosenfeld, M. R.; Wong, E.; Dalmau, J.; Manley, G.; Posner, J.

- [93119] B.; Sher, E.; Furneaux, H. M.: Cloning and characterization of a
- [93120] Lambert-Eaton myasthenic syndrome antigen. *Ann. Neurol.* 33: 113-120,
- [93121] 1993.
- [93122] 16286.
- [93123] Taviaux, S.; Williams, M. E.; Harpold, M. M.; Nargeot, J.; Lory,
- [93124] P.: Assignment of human genes for beta-2 and beta-4 subunits of voltage-dependent
- [93125] Ca(2+) channels to chromosomes 10p12 and 2q22-q23. *Hum. Genet.* 100:
- [93126] 151-154, 1997.
- [93127] 16287.
- [93128] Bontron, S.; Steimle, V.; Ucla, C.; Eibl, M. M.; Mach, B.: Two
- [93129] novel mutations in the MHC class II transactivator CIITA in a second
- [93130] patient from MHC class II deficiency complementation group A. *Hum.*
- [93131] *Genet.* 99: 541-546, 1997.
- [93132] 16288.

- [93133] Dziembowska, M.; Fondaneche, M.-C.; Vedrenne, J.; Barbieri, G.;
- [93134] Wiszniewski, W.; Picard, C.; Cant, A. J.; Steimle, V.; Charon, D.;
- [93135] Alca-Loridan, C.; Fischer, A.; Lisowska-Grosppierre, B.:
Three novel
mutations of the CIITA gene in MHC class II-deficient patients with
a severe immunodeficiency. Immunogenetics 53:
821-829, 2002.
- [93138] 16289.
- [93139] Harton, J. A.; Cressman, D. E.; Chin, K.-C.; Der, C. J.; Ting,
J. P.-Y.: GTP binding by class II transactivator: role in nuclear
import. Science 285: 1402-1405, 1999.
- [93142] 16290.
- [93143] Harton, J. A.; Ting, J. P.-Y.: Class II transactivator: mastering
the art of major histocompatibility complex expression.
Molec. Cell.
- [93145] Biol. 20: 6185-6194, 2000.

- [93146] 16291.
- [93147] Mach, B.; Steimle, V.; Martinez-Soria, E.; Reith, W.: Regulation
[93148] of MHC class II genes: lessons from a disease. *Annu. Rev. Immun.* 14:
[93149] 301–331, 1996.
- [93150] 16292.
- [93151] Mahanta, S. K.; Scholl, T.; Yang, F.-C.; Strominger, J. L.: Transactivation
[93152] by CIITA, the type II bare lymphocyte syndrome-associated factor,
[93153] requires participation of multiple regions of the TATA box binding
[93154] protein. *Proc. Nat. Acad. Sci.* 94: 6324–6329, 1997.
- [93155] 16293.
- [93156] Raval, A.; Howcroft, T. K.; Weissman, J. D.; Kirshner, S.; Zhu,
[93157] X.-S.; Yokoyama, K.; Ting, J.; Singer, D. S.: Transcriptional coactivator,
[93158] CIITA, is an acetyltransferase that bypasses a promoter requirement

[93159] for TAFII250. Molec. Cell 7: 105–115, 2001.

[93160] 16294.

[93161] Reith, W.: Personal Communication. Geneva, Switzerland
5/30/1997.

[93162] 16295.

[93163] Scholl, T.; Mahanta, S. K.; Strominger, J. L.: Specific complex

[93164] formation between the type II bare lymphocyte syndrome-associated

[93165] transactivators CIITA and RFX5. Proc. Nat. Acad. Sci. 94:
6330–6334,

[93166] 1997.

[93167] 16296.

[93168] Steimle, V.; Otten, L. A.; Zufferey, M.; Mach, B.: Complementa-

[93169] tion cloning of an MHC class II transactivator mutated in
hereditary MHC

[93170] class II deficiency (or bare lymphocyte syndrome). Cell 75:
135–146,

[93171] 1993.

[93172] 16297.

- [93173] Zhu, X.-S.; Linhoff, M. W.; Li, G.; Chin, K.-C.; Maity, S. N.;
- [93174] Ting, J. P.-Y.: Transcriptional scaffold: CIITA interacts with NF- κ B, RFX, and CREB to cause stereospecific regulation of the class II major histocompatibility complex promoter. *Molec. Cell. Biol.* 20: 6051-6061, 2000.
- [93175] 16298.
- [93176] Emery, P.; Durand, B.; Mach, B.; Reith, W.: RFX proteins, a novel family of DNA binding proteins conserved in the eukaryotic kingdom. *Nucleic Acids Res.* 24: 803-807, 1996.
- [93177] 16299.
- [93178] Pugliatti, L.; Reith, W.; Fey, S.; Mach, B.: Mapping the RF-X gene, encoding a DNA-binding protein controlling HLA class II gene expression, to 19p13.(Abstract) *Cytogenet. Cell Genet.* 51: 1061 only, 1989.

- [93186] 16300.
- [93187] Mizuta, M.; Inagaki, N.; Nemoto, Y.; Matsukura, S.; Takahashi,
- [93188] M.; Seino, S.: Synaptotagmin III is a novel isoform of rat synaptotagmin
- [93189] expressed in endocrine and neuronal cells. J. Biol. Chem. 269: 11675–11678,
- [93190] 1994.
- [93191] 16301.
- [93192] Li, X.-J.; Wang, D.-Y.; Zhu, Y.; Guo, R.-J.; Wang, X.-D.; Lubomir,
- [93193] K.; Mukai, K.; Sasaki, H.; Yoshida, H.; Oka, T.; Machinami, R.; Shinmura,
- [93194] K.; Tanaka, M.; Sugimura, H.: Mxi1 mutations in human neurofibrosarcomas. Jpn.
- [93195] J. Cancer Res. 90: 740–746, 1999.
- [93196] 16302.
- [93197] Prochownik, E. V.; Grove, L. E.; Deubler, D.; Zhu, X. L.; Stephenson,
- [93198] R. A.; Rohr, L. R.; Yin, X.; Brothman, A. R.: Commonly occurring

[93199] loss and mutation of the MXI1 gene in prostate cancer.

Genes Chromosomes

[93200] *Cancer* 22: 295–304, 1998.

[93201] 16303.

[93202] Schreiber–Agus, N.; Meng, Y.; Hoang, T.; Hou, H., Jr.;
Chen, K.;

[93203] Greenberg, R.; Cordon–Cardo, C.; Lee, H.–W.; DePinho, R.
A.: Role

[93204] of Mxi1 in ageing organ systems and the regulation of
normal and neoplastic

[93205] growth. *Nature* 393: 483–487, 1998.

[93206] 16304.

[93207] Shapiro, D. N.; Valentine, V.; Eagle, L.; Yin, X.; Morris, S.
W.;

[93208] Prochownik, E. V.: Assignment of the human MAD and
MXI1 genes to

[93209] chromosomes 2p12–p13 and 10q24–q25. *Genomics* 23:
282–285, 1994.

[93210] 16305.

[93211] Wechsler, D. S.; Hawkins, A. L.; Li, X.; Jabs, E. W.; Griffin,

[93212] C. A.; Dang, C. V.: Localization of the human Mxi1 tran–

scription

[93213] factor gene (MXI1) to chromosome 10q24-q25. Genomics 21: 669-672,

[93214] 1994.

[93215] 16306.

[93216] Wechsler, D. S.; Shelly, C. A.; Dang, C. V.: Genomic organization

[93217] of human MXI1, a putative tumor suppressor gene. Genomics 32: 466-470,

[93218] 1996.

[93219] 16307.

[93220] Zervos, A. S.; Gyuris, J.; Brent, R.: Mxi1, a protein that specifically

[93221] interacts with Max to bind Myc-Max recognition sites. Cell 72: 223-232,

[93222] 1993. Note: Erratum: Cell 79: 389 only, 1994.

[93223] 16308.

[93224] Boie, Y.; Rushmore, T. H.; Darmon-Goodwin, A.; Grygorczyk, R.;

[93225] Slipetz, D. M.; Metters, K. M.; Abramovitz, M.: Cloning and expression

- [93226] of a cDNA for the human prostanoid IP receptor. J. Biol. Chem. 269:
- [93227] 12173–12178, 1994.
- [93228] 16309.
- [93229] Ishikawa, T.; Tamai, Y.; Rochelle, J. M.; Hirata, M.; Namba, T.;
- [93230] Sugimoto, Y.; Ichikawa, A.; Narumiya, S.; Taketo, M. M.; Seldin, M.
- [93231] F.: Mapping of the genes encoding mouse prostaglandin D, E, and F
- [93232] and prostacyclin receptors. Genomics 32: 285–288, 1996.
- [93233] 16310.
- [93234] Murata, T.; Ushikubi, F.; Matsuoka, T.; Hirata, M.; Yamasaki, A.;
- [93235] Sugimoto, Y.; Ichikawa, A.; Aze, Y.; Tanaka, T.; Yoshida, N.; Ueno,
- [93236] A.; Ohishi, S.; Narumiya, S.: Altered pain perception and inflammatory
- [93237] response in mice lacking prostacyclin receptor. Nature 388: 678–682,
- [93238] 1997.

- [93239] 16311.
- [93240] Ogawa, Y.; Tanaka, I.; Inoue, M.; Yoshitake, Y.; Isse, N.; Nakagawa,
- [93241] O.; Usui, T.; Itoh, H.; Yoshimasa, T.; Narumiya, S.; Nakao, K.: Structural
- [93242] organization and chromosomal assignment of the human prostacyclin
- [93243] receptor gene. *Genomics* 27: 142–148, 1995.
- [93244] 16312.
- [93245] Hoffman, I.; Balling, R.: Chromosomal localization of the murine
- [93246] cadherin-11. *Mammalian Genome* 6: 304 only, 1995.
- [93247] 16313.
- [93248] Okazaki, M.; Takeshita, S.; Kawai, S.; Kikuno, R.; Tsujimura, A.;
- [93249] Kudo, A.; Amann, E.: Molecular cloning and characterization of OB-cadherin,
- [93250] a new member of cadherin family expressed in osteoblasts. *J. Biol.*
- [93251] *Chem.* 269: 12092–12098, 1994.
- [93252] 16314.

- [93253] Tanihara, H.; Sano, K.; Heimark, R. L.; St. John, T.; Suzuki, S.
- [93254] : Cloning of five human cadherins clarifies characteristic features
- [93255] of cadherin extracellular domain and provides further evidence for
- [93256] two structurally different types of cadherin. Cell Adhes. Commun. 2:
- [93257] 15–26, 1994.
- [93258] 16315.
- [93259] Courvalin, J.-C.; Lassoued, K.; Worman, H. J.; Blobel, G.: Identification
- [93260] and characterization of autoantibodies against the nuclear envelope
- [93261] lamin B receptor from patients with primary biliary cirrhosis. J.
- [93262] Exp. Med. 172: 961–967, 1990.
- [93263] 16316.
- [93264] Karl, H.: Die besondere Haufung von Merkmalsträgern der Pelger–Huet–Kernanomalie
- [93265] der Leukozyten in Gelenau im Erzgebirge. Thesis: Humboldt Univ. Berlin,

- [93266] Germany , 1967.
- [93267] 16317.
- [93268] Nachtsheim, H.: The Pelger-anomaly in man and rabbit. J. Hered. 41:
- [93269] 131-137, 1950.
- [93270] 16318.
- [93271] Schuler, E.; Lin, F.; Worman, H. J.: Characterization of the human
- [93272] gene encoding LBR, an integral protein of the nuclear envelope inner
- [93273] membrane. J. Biol. Chem. 269: 11312-11317, 1994.
- [93274] 16319.
- [93275] Christie, P. T.; Curley, A.; Nesbit, M. A.; Chapman, C.; Genet,
- [93276] S.; Harper, P. S.; Keeling, S. L.; Wilkie, A. O. M.; Winter, R. M.;
- [93277] Thakker, R. V.: Mutational analysis in X-linked spondyloepiphyseal
- [93278] dysplasia tarda. J. Clin. Endocr. Metab. 86: 3233-3236, 2001.
- [93279] 16320.

- [93280] Fiedler, J.; Bittner, M.; Puhl, W.; Brenner, R. E.: Mutations
[93281] in the X-linked spondyloepiphyseal dysplasia tarda (SEDL)
coding sequence
[93282] are not a common cause of early primary osteoarthritis in
men. (Letter) Clin.
[93283] Genet. 62: 94–95, 2002.
[93284] 16321.
[93285] Gecz, J.; Hillman, M. A.; Gedeon, A. K.; Cox, T. C.; Baker,
E.;
[93286] Mulley, J. C.: Gene structure and expression study of the
SEDL gene
[93287] for spondyloepiphyseal dysplasia tarda. Genomics 69:
242–251, 2000.
[93288] 16322.
[93289] Gedeon, A. K.; Colley, A.; Jamieson, R.; Thompson, E. M.;
Rogers,
[93290] J.; Sillence, D.; Tiller, G. E.; Mulley, J. C.; Gecz, J.: Identifi-
cation
[93291] of the gene (SEDL) causing X-linked spondyloepiphyseal
dysplasia tarda. Nature
[93292] Genet. 22: 400–404, 1999.

[93293] 16323.

[93294] Gedeon, A. K.; Tiller, G. E.; Le Merrer, M.; Heuertz, S.;
Tranebjaerg,

[93295] L.; Chitayat, D.; Robertson, S.; Glass, I. A.; Savarirayan, R.;
Cole,

[93296] W. G.; Rimoin, D. L.; Kousseff, B. G.; Ohashi, H.; Zabel, B.;
Munnich,

[93297] A.; Gecz, J.; Mulley, J. C.: The molecular basis of X-linked
spondyloepiphyseal

[93298] dysplasia tarda. Am. J. Hum. Genet. 68: 1386–1397,
2001.

[93299] 16324.

[93300] Grunebaum, E.; Arpaia, E.; MacKenzie, J. J.; Fitzpatrick, J.;
Ray,

[93301] P. N.; Roifman, C. M.: A missense mutation in the SEDL
gene results

[93302] in delayed onset of X linked spondyloepiphyseal dysplasia
in a large

[93303] pedigree. (Letter) J. Med. Genet. 38: 409–411, 2001.

[93304] 16325.

[93305] Mumm, S.; Christie, P. T.; Finnegan, P.; Jones, J.; Dixon, P.
H.;

- [93306] Pannett, A. A. J.; Harding, B.; Gottesman, G. S.; Thakker, R. V.;
- [93307] Whyte, M. P.: A five-base pair deletion in the sedlin gene causes
- [93308] spondyloepiphyseal dysplasia tarda in a six-generation Arkansas kindred. J.
- [93309] Clin. Endocr. Metab. 85: 3343–3347, 2000.
- [93310] 16326.
- [93311] Mumm, S.; Zhang, X.; Vacca, M.; D'Esposito, M.; Whyte, M. P.:
- [93312] The sedlin gene for spondyloepiphyseal dysplasia tarda escapes X-inactivation
- [93313] and contains a non-canonical splice site. Gene 273: 285–293, 2001.
- [93314] 16327.
- [93315] Takahashi, T.; Takahashi, I.; Tsuchida, S.; Oyama, K.; Ko-matsu,
- [93316] M.; Saito, H.; Takada, G.: An SEDL gene mutation in a Japanese kindred
- [93317] of X-linked spondyloepiphyseal dysplasia tarda. (Letter) Clin. Genet. 61:
- [93318] 319–320, 2002.

- [93319] 16328.
- [93320] Tiller, G. E.; Hannig, V. L.; Dozier, D.; Carrel, L.; Trevarthen,
- [93321] K. C.; Wilcox, W. R.; Mundlos, S.; Haines, J. L.; Gedeon, A. K.; Gecz,
- [93322] J.: A recurrent RNA-splicing mutation in the SEDL gene causes X-linked
- [93323] spondyloepiphyseal dysplasia tarda. *Am. J. Hum. Genet.* 68: 1398-1407,
- [93324] 2001.
- [93325] 16329.
- [93326] Whyte, M. P.; Gottesman, G. S.; Eddy, M. C.; McAlister, W. H.
- [93327] : X-linked recessive spondyloepiphyseal dysplasia tarda: clinical
- [93328] and radiographic evolution in a 6-generation kindred and review of
- [93329] the literature. *Medicine* 78: 9-25, 1999.
- [93330] 16330.
- [93331] Montini, E.; Andolfi, G.; Caruso, A.; Buchner, G.; Walpole, S.

- [93332] M.; Mariani, M.; Consalez, G.; Trump, D.; Ballabio, A.; Franco, B.
- [93333] : Identification and characterization of a novel serine-threonine
- [93334] kinase gene from the Xp22 region. *Genomics* 51: 427-433, 1998.
- [93335] 16331.
- [93336] Becker, M. A.; Heidler, S. A.; Bell, G. I.; Seino, S.; Le Beau, M. M.; Westbrook, C. A.; Neuman, W.; Shapiro, L. J.; Mohandas, T.
- [93338] K.; Roessler, B. J.; Palella, T. D.: Cloning of cDNAs for human phosphoribosylpyrophosphate
- [93339] synthetases 1 and 2 and X chromosome localization of PRPS1 and PRPS2
- [93340] genes. *Genomics* 8: 555-561, 1990.
- [93341] 16332.
- [93342] Braverman, N.; Lin, P.; Moebius, F. F.; Obie, C.; Moser, A.; Glossmann,
- [93343] H.; Wilcox, W. R.; Rimoin, D. L.; Smith, M.; Kratz, L.; Kelley, R.
- [93344] I.; Valle, D.: Mutations in the gene encoding 3-beta-hydroxysteroid-delta(8),delta(7)-isomerase

- [93345] cause X-linked dominant Conradi-Hunermann syndrome.
Nature Genet. 22:
[93346] 291-294, 1999.
- [93347] 16333.
- [93348] Cho, S. Y.; Kim, J. H.; Paik, Y. K.: Cholesterol biosynthesis
[93349] from lanosterol: differential inhibition of sterol delta
8-isomerase
[93350] and other lanosterol-converting enzymes by tamoxifen.
Molec. Cells 8:
[93351] 233-239, 1998.
- [93352] 16334.
- [93353] Clayton, P. T.; Kalter, D. C.; Atherton, D. J.; Besley, G. T.;
[93354] Broadhead, D. M.: Peroxisomal enzyme deficiency in X-
linked dominant
[93355] Conradi-Hunermann syndrome. J. Inherit. Metab. Dis. 12:
358-360,
[93356] 1989.
- [93357] 16335.
- [93358] Derry, J. M. J.; Gormally, E.; Means, G. D.; Zhao, W.;
Meindl,

- [93359] A.; Kelley, R. I.; Boyd, Y.; Herman, G. E.: Mutations in a delta(8)-delta(7)
- [93360] sterol isomerase in the tattered mouse and X-linked dominant chondrodysplasia
- [93361] punctata. Nature Genet. 22: 286-290, 1999.
- [93362] 16336.
- [93363] Grange, D. K.; Kratz, L. E.; Braverman, N. E.; Kelley, R. I.:
- [93364] CHILD syndrome caused by deficiency of 3-beta-hydroxysteroid-delta-8,delta-7-isomerase. Am.
- [93365] J. Med. Genet. 90: 328-335, 2000.
- [93366] 16337.
- [93367] Hanner, M.; Moebius, F. F.; Weber, F.; Grabner, M.; Striessnig,
- [93368] J.; Glossmann, H.: Phenylalkylamine Ca(2+) antagonist binding protein:
- [93369] molecular cloning, tissue distribution, and heterologous expression. J.
- [93370] Biol. Chem. 270: 7551-7557, 1995.
- [93371] 16338.
- [93372] Has, C.; Bruckner-Tuderman, L.; Muller, D.; Floeth, M.; Folkers,

- [93373] E.; Donnai, D.; Traupe, H.: The Conradi–Hunermann–Happle syndrome
- [93374] (CDPX2) and emopamil binding protein: novel mutations, and somatic
- [93375] and gonadal mosaicism. Hum. Molec. Genet. 9: 1951–1955, 2000.
- [93376] 16339.
- [93377] Holmes, R. D.; Wilson, G. N.; Hajra, A. K.: Peroxisomal enzyme
- [93378] deficiency in the Conradi–Hunerman (sic) form of chondrodysplasia
- [93379] punctata. New Eng. J. Med. (Letter) 316: 1608 only, 1987.
- [93380] 16340.
- [93381] Ikegawa, S.; Ohashi, H.; Ogata, T.; Honda, A.; Tsukahara, M.; Kubo,
- [93382] T.; Kimizuka, M.; Shimode, M.; Hasegawa, T.; Nishimura, G.; Nakamura,
- [93383] Y.: Novel and recurrent EBP mutations in X-linked dominant chondrodysplasia
- [93384] punctata. Am. J. Med. Genet. 94: 300–305, 2000.
- [93385] 16341.

- [93386] Kelley, R. I.; Wilcox, W. G.; Smith, M.; Kratz, L. E.; Moser, A.; Rimoin, D. S.: Abnormal sterol metabolism in patients with Conradi-Hunermann-Happle syndrome and sporadic lethal chondrodysplasia punctata. Am. J. Med. Genet. 83: 213-219, 1999.
- [93387] 16342.
- [93388] Rasmussen, U. B.; Wolf, C.; Mattei, M.-G.; Chenard, M.-P.; Bellocq, J.-P.; Chambon, P.; Rio, M.-C.; Basset, P.: Identification of a new interferon- α -inducible gene (p27) on human chromosome 14q32 and its expression in breast carcinoma. Cancer Res. 53: 4096-4101, 1993.
- [93389] 16343.
- [93390] Lan, M. S.; Li, Q.; Lu, J.; Modi, W. S.; Notkins, A. L.: Genomic organization, 5-prime-upstream sequence, and chromosomal localization of an insulinoma-associated intronless gene, IA-1. J. Biol. Chem. 269:

[93399] 14170–14174, 1994.

[93400] 16344.

[93401] Bennett, B. D.; Wang, Z.; Kuang, W.-J.; Wang, A.; Groopman, J.

[93402] E.; Goeddel, D. V.; Scadden, D. T.: Cloning and characterization

[93403] of HTK, a novel transmembrane tyrosine kinase of the EPH subfamily. J.

[93404] Biol. Chem. 269: 14211–14218, 1994.

[93405] 16345.

[93406] Berclaz, G.; Andres, A.-C.; Albrecht, D.; Dreher, E.; Ziemiecki,

[93407] A.; Gusterson, B. A.; Crompton, M. R.: Expression of the receptor

[93408] protein tyrosine kinase myk-1/htk in normal and malignant mammary

[93409] epithelium. Biochem. Biophys. Res. Commun. 226: 869–875, 1996.

[93410] 16346.

[93411] Gerety, S. S.; Wang, H. U.; Chen, Z.-F.; Anderson, D. J.: Symmetrical

- [93412] mutant phenotypes of the receptor EphB4 and its specific transmembrane
- [93413] ligand ephrin-B2 in cardiovascular development. *Molec. Cell* 4: 403-414,
- [93414] 1999.
- [93415] 16347.
- [93416] Moynihan, T. P.; Ardley, H. C.; Leek, J. P.; Thompson, J.; Brindle,
- [93417] N. S.; Markham, A. F.; Robinson, P. A.: Characterization of a human
- [93418] ubiquitin-conjugating enzyme gene UBE2L3. *Mammalian Genome* 7: 520-525,
- [93419] 1996.
- [93420] 16348.
- [93421] Flanagan, J. R.; Becker, K. G.; Ennist, D. L.; Gleason, S. L.;
- [93422] Driggers, P. H.; Levi, B.-Z.; Appella, E.; Ozato, K.: Cloning of
- [93423] a negative transcription factor that binds to the upstream conserved
- [93424] region of Moloney murine leukemia virus. *Molec. Cell. Biol.* 12:

- [93425] 38–44, 1992.
- [93426] 16349.
- [93427] Hariharan, N.; Kelley, D. E.; Perry, R. P.: Delta, a transcription
- [93428] factor that binds to downstream elements in several polymerase II
- [93429] promoters, is a functionally versatile zinc finger protein. Proc.
- [93430] Nat. Acad. Sci. 88: 9799–9803, 1991.
- [93431] 16350.
- [93432] Oei, S. L.; Shi, Y.: Transcription factor Yin Yang 1 stimulates
- [93433] poly(ADP-ribosyl)ation and DNA repair. Biochem. Biophys. Res. Commun. 284:
- [93434] 450–454, 2001.
- [93435] 16351.
- [93436] Park, K.; Atchison, M. L.: Isolation of a candidate repressor/activator,
- [93437] NF-E1 (YY-1, delta), that binds to the immunoglobulin kappa 3-prime
- [93438] enhancer and the immunoglobulin heavy-chain micro-E1

site.. Proc.

[93439] Nat. Acad. Sci. 88: 9804–9808, 1991.

[93440] 16352.

[93441] Shi, Y.; Seto, E.; Chang, L.-S.; Shenk, T.: Transcriptional repression

[93442] by YY1, a human GLI-Kruppel-related protein, and relief of repression

[93443] by adenovirus E1A protein. Cell 67: 377–388, 1991.

[93444] 16353.

[93445] Yao, Y.-L.; Dupont, B. R.; Ghosh, S.; Fang, Y.; Leach, R. J.; Seto,

[93446] E.: Cloning, chromosomal localization and promoter analysis of the

[93447] human transcription factor YY1. Nucleic Acids Res. 26: 3776–3783,

[93448] 1998.

[93449] 16354.

[93450] Zhu, W.; Lossie, A. C.; Camper, S. A.; Gumucio, D. L.: Chromosomal

[93451] localization of the transcription factor YY1 in the mouse and human. Mammalian

- [93452] Genome 5: 234–236, 1994.
- [93453] 16355.
- [93454] Hakimi, M.–A.; Bochar, D. A.; Schmiesing, J. A.; Dong, Y.; Barak,
- [93455] O. G.; Speicher, D. W.; Yokomori, K.; Shiekhattar, R.: A chromatin
- [93456] remodelling complex that loads cohesin onto human chromosomes. Nature 418:
- [93457] 994–998, 2002.
- [93458] 16356.
- [93459] Berglund, E. O.; Murai, K. K.; Fredette, B.; Sekerkova, G.; Marturano,
- [93460] B.; Weber, L.; Mugnaini, E.; Ranscht, B.: Ataxia and abnormal cerebellar
- [93461] microorganization in mice with ablated contactin gene expression. Neuron 24:
- [93462] 739–750, 1999.
- [93463] 16357.
- [93464] Nagasaki, K.; Manabe, T.; Hanzawa, H.; Maass, N.; Tsukada, T.;
- [93465] Yamaguchi, K.: Identification of a novel gene, LDOC1,

down-regulated

[93466] in cancer cell lines. Cancer Lett. 140: 227-234, 1999.

[93467] 16358.

[93468] Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.;
Kawarabayasi,

[93469] Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.:
Prediction

[93470] of the coding sequences of unidentified human genes. I.
The coding

[93471] sequences of 40 new genes (KIAA0001-KIAA0040) de-
duced by analysis

[93472] of randomly samples cDNA clones from human immature
myeloid cell line

[93473] KG-1. DNA Res. 1: 27-35, 1994.

[93474] 16359.

[93475] Kikuno, R.; Nagase, T.; Ishikawa, K.; Hirose, M.; Miya-
jima, N.;

[93476] Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction
of the

[93477] coding sequences of unidentified human genes. XIV. The
complete sequences

[93478] of 100 new cDNA clones from brain which code for large

proteins in

[93479] *in vitro*. DNA Res. 6: 197–205, 1999.

[93480] 16360.

[93481] Polsky, D.; Young, A. Z.; Busam, K. J.; Alani, R. M.: The transcriptional

[93482] repressor of p16/Ink4a, Id1, is up-regulated in early melanomas. Cancer

[93483] Res. 61: 6008–6011, 2001.

[93484] 16361.

[93485] Dahl, N.; Hu, L. J.; Chery, M.; Fardeau, M.; Gilgenkrantz, S.;

[93486] Nivelon-Chevallier, A.; Sidaner-Noisette, I.; Mugneret, F.; Gouyon,

[93487] J. B.; Gal, A.; Kioschis, P.; d'Urso, M.; Mandel, J.-L.: Myotubular

[93488] myopathy in a girl with a deletion at Xq27–q28 and unbalanced X inactivation

[93489] assigns the MTM1 gene to a 600–kb region. Am. J. Hum. Genet. 56:

[93490] 1108–1115, 1995.

[93491] 16362.

[93492] Dahl, N.; Samson, F.; Thomas, N. S. T.; Hu, L. J.; Gong, W.;

Herman,

[93493] G.; Laporte, J.; Kioschis, P.; Poustka, A.; Mandel, J. L.: X linked

[93494] myotubular myopathy (MTM1) maps between DXS304 and DXS305, closely

[93495] linked to the DXS455 VNTR and a new, highly informative microsatellite

[93496] marker (DXS1684). J. Med. Genet. 31: 922–924, 1994.

[93497] 16363.

[93498] Darnfors, C.; Borje Larsson, H. E.; Oldfors, A.; Kyllerman, M.;

[93499] Gustavson, K.–H.; Bjursell, G.; Wahlstrom, J.: X-linked myotubular

[93500] myopathy: a linkage study. Clin. Genet. 37: 335–340, 1990.

[93501] 16364.

[93502] Darnfors, C.; Borje Larsson, H. E.; Oldfors, A.; Kyllerman, M.;

[93503] Gustavson, K.–H.; Bjursell, G.; Wahlstrom, J.: X-linked myotubular

[93504] myopathy: a linkage study.(Abstract) Cytogenet. Cell Genet. 51:

- [93505] 983, 1989.
- [93506] 16365.
- [93507] de Gouyon, B.; Chatterjee, A.; Monaco, A.; Quaderi, N.; Brown,
- [93508] S. D. M.; Herman, G. E.: Comparative mapping on the mouse X chromosome
- [93509] defines a myotubular myopathy equivalent region. *Mammalian Genome* 7:
- [93510] 575–579, 1996.
- [93511] 16366.
- [93512] de Gouyon, B. M.; Zhao, W.; Laporte, J.; Mandel, J.-L.; Metzenberg,
- [93513] A.; Herman, G. E.: Characterization of mutations in the myotubularin
- [93514] gene in twenty six patients with X-linked myotubular myopathy. *Hum. Molec. Genet.* 6: 1499–1504, 1997.
- [93515] 16367.
- [93517] Engel, W. K.; Gold, G. N.; Karpatis, B.: Type I fiber hypotrophy
- [93518] and central nuclei. *Arch. Neurol.* 18: 435–444, 1968.

- [93519] 16368.
- [93520] Fidzianska, A.; Warlo, I.; Goebel, H. H.: Neonatal centronuclear
- [93521] myopathy with N-Cam decorated myotubes. Neuropediatrics 25: 158-161,
- [93522] 1994.
- [93523] 16369.
- [93524] Guiraud-Chaumeil, C.; Vincent, M. C.; Laporte, J.; Fardeau, M.;
- [93525] Samson, F.; Mandel, J.-L.: A mutation in the MTM1 gene invalidates
- [93526] a previous suggestion of nonallelic heterogeneity in X-linked myotubular
- [93527] myopathy. (Letter) Am. J. Hum. Genet. 60: 1542-1544, 1997.
- [93528] 16370.
- [93529] Hammans, S. R.; Robinson, D. O.; Moutou, C.; Kennedy, C. R.; Dennis,
- [93530] N. R.; Hughes, P. J.; Ellison, D. W.: A clinical and genetic study
- [93531] of a manifesting heterozygote with X-linked myotubular myopathy. Neuromusc.

[93532] Disord. 10: 133–137, 2000.

[93533] 16371.

[93534] Hane, B. G.; Rogers, R. C.; Schwartz, C. E.: Germline mosaicism

[93535] in X-linked myotubular myopathy. Clin. Genet. 56: 77–81, 1999.

[93536] 16372.

[93537] Heckmatt, J. Z.; Sewry, C. A.; Hodes, D.; Dubowitz, V.: Congenital

[93538] centronuclear (myotubular) myopathy: a clinical, pathological and

[93539] genetic study in eight children. Brain 108: 941–964, 1985.

[93540] 16373.

[93541] Herman, G. E.; Finegold, M.; de Gouyon, B.; Metzenberg, A.: Medical

[93542] complications in long-term survivors with X-linked myotubular myopathy. J.

[93543] Pediat. 134: 206–214, 1999.

[93544] 16374.

[93545] Herman, G. E.; Kopacz, K.; Zhao, W.; Mills, P. L.; Metzenberg,

- [93546] A.; Das, S.: Characterization of mutations in fifty North American
- [93547] patients with X-linked myotubular myopathy. Hum. Mutat. 19: 114–121,
- [93548] 2002.
- [93549] 16375.
- [93550] Janssen, E. A. M.; Hensels, G. W.; van Oost, B. A.; Hamel, B.
- [93551] C. J.; Kemp, S.; Baas, F.; Weber, J. W.; Barth, P. G.; Bolhuis, P.
- [93552] A.: The gene for X-linked myotubular myopathy is located in an 8
- [93553] Mb region at the border of Xq27.3 and Xq28. Neuromusc. Disord. 4:
- [93554] 455–461, 1994.
- [93555] 16376.
- [93556] Joseph, M.; Pai, G. S.; Holden, K. R.; Herman, G.: X-linked myotubular
- [93557] myopathy: clinical observations in ten additional cases. Am. J. Med.
- [93558] Genet. 59: 168–173, 1995.
- [93559] 16377.

- [93560] Keppen, L. D.; Husain, M. M.; Woody, R. C.: X-linked myotubular
- [93561] myopathy: intrafamilial variability and normal muscle biopsy in a
- [93562] heterozygous female. Clin. Genet. 32: 95-99, 1987.
- [93563] 16378.
- [93564] Kioschis, P.; Rogner, U. C.; Pick, E.; Klauck, S. M.; Heiss, N.;
- [93565] Siebenhaar, R.; Korn, B.; Coy, J. F.; Laporte, J.; Liechti-Gallati,
- [93566] S.; Poustka, A.: A 900-kb cosmid contig and 10 new transcripts within
- [93567] the candidate region for myotubular myopathy (MTM1). Genomics 33:
- [93568] 365-373, 1996.
- [93569] 16379.
- [93570] Laporte, J.; Biancalana, V.; Tanner, S. M.; Kress, W.; Schneider,
- [93571] V.; Wallgren-Pettersson, C.; Herger, F.; Buj-Bello, A.; Blondeau,
- [93572] F.; Liechti-Gallati, S.; Mandel, J.-L.: MTM1 mutations in X-linked

- [93573] myotubular myopathy. Hum. Mutat. 15: 393–409, 2000.
- [93574] 16380.
- [93575] Laporte, J.; Blondeau, F.; Buj-Bello, A.; Mandel, J.-L.: The
- [93576] myotubularin family: from genetic disease to phospho-
inositide metabolism. Trends
- [93577] Genet. 17: 221–228, 2001.
- [93578] 16381.
- [93579] Laporte, J.; Blondeau, F.; Buj-Bello, A.; Tentler, D.; Kretz,
- [93580] C.; Dahl, N.; Mandel, J.-L.: Characterization of the my-
otubularin
- [93581] dual specificity phosphatase gene family from yeast to
human. Hum.
- [93582] Molec. Genet. 7: 1703–1712, 1998.
- [93583] 16382.
- [93584] Laporte, J.; Guiraud-Chaumeil, C.; Tanner, S. M.; Blon-
deau, F.;
- [93585] Hu, L.-J.; Vicaire, S.; Liechti-Gallati, S.; Mandel, J.-L.: Ge-
nomic
- [93586] organization of the MTM1 gene implicated in X-linked
myotubular myopathy. Europ.

- [93587] J. Hum. Genet. 6: 325–330, 1998.
- [93588] 16383.
- [93589] Laporte, J.; Guiraud–Chaumeil, C.; Vincent, M.–C.; Mandel, J.–L.;
- [93590] Tanner, S. M.; Liechti–Gallati, S.; Wallgren–Pettersson, C.; Dahl,
- [93591] N.; Kress, W.; Bolhuis, P. A.; Fardeau, M.; Samson, F.; Bertini, E.;
- [93592] members of the ENMC International Consortium on Myotubular Myopathy
- [93593] : Mutations in the MTM1 gene implicated in X-linked myotubular myopathy. Hum.
- [93594] Molec. Genet. 6: 1505–1511, 1997.
- [93595] 16384.
- [93596] Lehesjoki, A.–E.; Sankila, E.–M.; Miao, J.; Somer, M.; Salonen,
- [93597] R.; Rapola, J.; de la Chapelle, A.: X linked neonatal myotubular
- [93598] myopathy: one recombination detected with four polymorphic DNA markers
- [93599] from Xq28. J. Med. Genet. 27: 288–291, 1990.

- [93600] 16385.
- [93601] Liechti-Gallati, S.; Muller, B.; Grimm, T.; Kress, W.; Muller, C.; Boltshauser, E.; Moser, H.; Braga, S.: X-linked centronuclear myopathy: mapping the gene to Xq28. Neuromusc. Disord. 1: 239-245, 1991.
- [93604] 16386.
- [93605] 16386.
- [93606] Meyers, K. R.; Golomb, H. M.; Hansen, J. L.; McKusick, V. A.: Familial neuromuscular disease with 'myotubes.'. Clin. Genet. 5: 327-337, 1974.
- [93608] 16387.
- [93610] Walker, J. L.; Dixon, J.; Fenton, C. R.; Hungerford, J.; Lynch, S. A.; Stenhouses, S. A. R.; Christian, A.; Craig, I. W.: Two new mutations in exon 3 of the NDP gene: S73X and S101F associated with severe and less severe ocular phenotype, respectively.

Hum. Mutat. 9:

[93614] 53–56, 1997.

[93615] 16388.

[93616] Lee, D. K.; Horikoshi, M.; Roeder, R. G.: Interaction of
TFIID

[93617] in the minor groove of the TATA element. Cell 67:
1241–1250, 1991.

[93618] 16389.

[93619] Ho, M. F.; Chalmers, R. M.; Davis, M. B.; Harding, A. E.;
Monaco,

[93620] A. P.: A novel point mutation in the McLeod syndrome
gene in neuroacanthocytosis. Ann.

[93621] Neurol. 39: 672–675, 1996.

[93622] 16390.

[93623] Wilson, R. B.; Kiledjian, M.; Shen, C.–P.; Benezra, R.;
Zwollo,

[93624] P.; Dymecki, S. M.; Desiderio, S. V.; Kadesch, T.: Repres-
sion of

[93625] immunoglobulin enhancers by the helix–loop–helix pro-
tein Id: implications

[93626] for B–lymphoid–cell development. Molec. Cell. Biol. 11:

6185-6191,

[93627] 1991.

[93628] 16391.

[93629] Warburg, M.: Norrie's disease: a new hereditary bilateral
pseudotumour

[93630] of the retina. Acta Ophthal. (Copenh) 39: 757-772, 1961.

[93631] 16392.

[93632] Warburg, M.: Norrie's disease (atrofia bulborum hereditaria). Acta

[93633] Ophthal. 41: 134-146, 1963.

[93634] 16393.

[93635] Warburg, M.: Norrie's disease, a congenital progressive
oculo-acoustico-cerebral

[93636] degeneration. Acta Ophthal. 89 (suppl.): 1-147, 1966.

[93637] 16394.

[93638] Warburg, M.; Hauge, M.; Sanger, R.: Norrie's disease and
the

[93639] Xg blood group system: linkage data. Acta Genet. Statist.
Med. 15:

[93640] 103-115, 1965.

- [93641] 16395.
- [93642] Whitnall, S. E.; Norman, R. M.: Microphthalmia and the visual
[93643] pathways: a case associated with blindness and imbecility,
and sex-linked. Brit.
[93644] J. Ophthal. 24: 229-244, 1940.
- [93645] 16396.
- [93646] Wilson, W. M. G.: Congenital blindness (pseudoglioma) oc-
curring
[93647] as a sex-linked developmental anomaly. Canad. Med. As-
soc. J. 60:
[93648] 580-584, 1949.
- [93649] 16397.
- [93650] Wolff, G.; Mayerova, A.; Wienker, T. F.; Atalianis, P.; Ioan-
nou,
[93651] P.; Warburg, M.: Clinical reinvestigation and linkage analy-
sis in
[93652] the family with Episkopi blindness (Norrie disease). J. Med.
Genet. 29:
[93653] 816-819, 1992.
- [93654] 16398.

- [93655] Wong, F.; Goldberg, M. F.; Hao, Y.: Identification of a non-sense
- [93656] mutation at codon 128 of the Norrie's disease gene in a male infant. Arch.
- [93657] Ophthal. 111: 1553-1557, 1993.
- [93658] 16399.
- [93659] Woodruff, G.; Newbury-Ecob, R.; Plaha, D. S.; Young, I. D.: Manifesting
- [93660] heterozygosity in Norrie's disease?. Brit. J. Ophthal. 77: 813-814,
- [93661] 1993.
- [93662] 16400.
- [93663] Zhu, D.; Antonarakis, S. E.; Schmeckpeper, B. J.; Diergaarde,
- [93664] P. J.; Greb, A. E.; Maumenee, I. H.: Microdeletion in the X-chromosome
- [93665] and prenatal diagnosis in a family with Norrie disease. Am. J. Med.
- [93666] Genet. 33: 485-488, 1989.
- [93667] 16401.
- [93668] Avner, P.; Bucan, M.; Arnaud, D.; Lehrach, H.; Rapp, U.: A-

raf

- [93669] oncogene localizes on mouse X chromosome to region
some 10–17 centimorgans
- [93670] proximal to hypoxanthine phosphoribosyltransferase
gene. Somat. Cell
- [93671] Molec. Genet. 13: 267–272, 1987.
- [93672] 16402.
- [93673] Beck, T. W.; Huleihel, M.; Gunnell, M.; Bonner, T. I.; Rapp,
U.
- [93674] R.: The complete coding sequence of the human A-raf–1
oncogene and
- [93675] transforming activity of a human A-raf carrying retrovirus.
Nucleic
- [93676] Acids Res. 15: 595–609, 1987.
- [93677] 16403.
- [93678] Huebner, K.; ar–Rushdi, A.; Griffin, C. A.; Isobe, M.; Kozak,
C.;
- [93679] Emanuel, B. S.; Nagarajan, L.; Cleveland, J. L.; Bonner, T. I.;
Goldsborough,
- [93680] M. D.; Croce, C. M.; Rapp, U.: Actively transcribed genes in
the
- [93681] raf oncogene group, located on the X chromosome in

mouse and human. Proc.

[93682] Nat. Acad. Sci. 83: 3934–3938, 1986.

[93683] 16404.

[93684] Lee, J.-E.; Beck, T. W.; Brennscheidt, U.; DeGennaro, L. J.;
Rapp,

[93685] U. R.: The complete sequence and promoter activity of the
human A-raf-1

[93686] gene (ARAF1). Genomics 20: 43–55, 1994.

[93687] 16405.

[93688] Mark, G. E.; Seeley, T. W.; Shows, T. B.; Mountz, J. D.: Pks,

[93689] a raf-related sequence in humans. Proc. Nat. Acad. Sci.
83: 6312–6316,

[93690] 1986.

[93691] 16406.

[93692] Popescu, N. C.; Mark, G. E.: Localization of the pKs gene, a
raf

[93693] related sequence on human chromosomes X and 7. Onco-
gene 4: 517–519,

[93694] 1989.

[93695] 16407.

- [93696] Dowdy, S. F.; Fasching, C. L.; Araujo, D.; Lai, K.-M.; Li-
vanos,
[93697] E.; Weissman, B. E.; Stanbridge, E. J.: Suppression of tu-
morigenicity
[93698] in Wilms' tumor by the p15.5-p14 region of chromosome
11. Science 254:
[93699] 293-295, 1991.
[93700] 16408.
- [93701] Dowdy, S. F.; Lai, K.-M.; Weissman, B. E.; Matsui, Y.;
Hogan, B.
[93702] L. M.; Stanbridge, E. J.: The isolation and characterization
of a
[93703] novel cDNA demonstrating an altered mRNA level in non-
tumorigenic Wilms'
[93704] microcell hybrid cells. Nucleic Acids Res. 19: 5763-5769,
1991.
[93705] 16409.
- [93706] Farmer, A. A.; Loftus, T. M.; Mills, A. A.; Sato, K. Y.; Neill,
[93707] J. D.; Tron, T.; Yang, M.; Trumpower, B. L.; Stanbridge, E.
J.: Extreme
[93708] evolutionary conservation of QM, a novel c-Jun associated
transcription

- [93709] factor. Hum. Molec. Genet. 3: 723–728, 1994.
- [93710] 16410.
- [93711] Kaneko, K.; Kobayashi, H.; Onodera, O.; Miyatake, T.;
Tsuji, S.
- [93712] : Genomic organization of a cDNA (QM) demonstrating an
altered mRNA
- [93713] level in nontumorigenic Wilms' microcell hybrid cells and
its localization
- [93714] to Xq28. Hum. Molec. Genet. 1: 529–533, 1992.
- [93715] 16411.
- [93716] Korn, B.; Sedlacek, Z.; Manca, A.; Kioschis, P.; Konecki, D.;
Lehrach,
- [93717] H.; Poustka, A.: A strategy for the selection of transcribed
sequences
- [93718] in the Xq28 region. Hum. Molec. Genet. 1: 235–242,
1992.
- [93719] 16412.
- [93720] van den Ouweland, A. M. W.; Verdijk, M.; Mannens, M. M.
A. M.;
- [93721] van Oost, B. A.: The QM gene is X-linked and therefore
not involved

- [93722] in suppression of tumorigenesis in Wilms' tumor. Hum. Molec. Genet. 90: 144-146, 1992.
- [93723] 16413.
- [93724] Weissman, B. E.; Saxon, P. J.; Pasquale, S. R.; Jones, G. R.; Geiser, A. G.; Stanbridge, E. J.: Introduction of a normal human chromosome 11 into a Wilms' tumor cell line controls its tumorigenic expression. Science 236: 175-180, 1987.
- [93725] 16414.
- [93726] Nishimoto, T.; Sekiguchi, T.; Kai, R.; Yamashita, K.; Takahashi, T.; Sekiguchi, M.: Large-scale selection and analysis of temperature-sensitive mutants for cell reproduction from BHK cells. Somat. Cell Genet. 8: 811-812, 1982.
- [93727] 16415.
- [93728] O'Brien, T.; Tjian, R.: Functional analysis of the human

TAFII250

[93736] N-terminal kinase domain. Molec. Cell 1: 905–911, 1998.

[93737] 16416.

[93738] Ruppert, S.; Wang, E. H.; Tjian, R.: Cloning and expression of

[93739] human TAF(II)250: a TBP-associated factor implicated in cell-cycle

[93740] regulation. Nature 362: 175–179, 1993.

[93741] 16417.

[93742] Schwartz, H. E.; Holmes, S.; Meiss, H. K.: Assignment of temperature-sensitive

[93743] mutations of BHK cells to the X-chromosome. (Abstract) J. Cell Biol. 75:

[93744] 393A, 1977.

[93745] 16418.

[93746] Schwartz, H. E.; Moser, G. C.; Holmes, S.; Meiss, H. K.: Assignment

[93747] of temperature-sensitive mutations of BHK cells to the X-chromosome. Somat.

[93748] Cell Genet. 5: 217–224, 1979.

[93749] 16419.

[93750] Sekiguchi, T.; Miyata, T.; Nishimoto, T.: Molecular cloning of
[93751] the cDNA of human X chromosomal gene (CCG1) which complements the
[93752] temperature-sensitive G(1) mutants, tsBN462 and ts13, of the BHK cell
[93753] line. EMBO J. 7: 1683–1687, 1988.
[93754] 16420.
[93755] Sekiguchi, T.; Nohiro, Y.; Nakamura, Y.; Hisamoto, N.; Nishimoto,
[93756] T.: The human CCG1 gene, essential for progression of the G1 phase,
[93757] encodes a 210-kilodalton nuclear DNA-binding protein. Molec. Cell.
[93758] Biol. 11: 3317–3325, 1991.
[93759] 16421.
[93760] Sekiguchi, T.; Yoshida, M. C.; Sekiguchi, M.; Nishimoto, T.:
[93761] Isolation of a human X chromosome-linked gene essential for progression
[93762] from G1 to S phase of the cell cycle. Exp. Cell Res. 169: 395–407,

[93763] 1987.

[93764] 16422.

[93765] Simchen, G.: Cell cycle mutants. Annu. Rev. Genet. 12:
161–191,

[93766] 1978.

[93767] 16423.

[93768] Slater, M. L.; Ozer, H. L.: Temperature-sensitive mutants
of

[93769] Balb–3T3 cells: description of a mutant affected in cellular
and polyoma

[93770] virus DNA synthesis. Cell 7: 289–295, 1976.

[93771] 16424.

[93772] Starr, D. B.; Hawley, D. K.: TFIID binds in the minor groove
[93773] of the TATA box. Cell 67: 1231–1240, 1991.

[93774] 16425.

[93775] Abel, A.; Walcott, J.; Woods, J.; Duda, J.; Merry, D. E.: Ex-
pression

[93776] of expanded repeat androgen receptor produces neuro-
logic disease in

[93777] transgenic mice. Hum. Molec. Genet. 10: 107–116, 2001.

[93778] 16426.

[93779] Bailey, C. K.; Andriola, I. F. M.; Kampinga, H. H.; Merry, D. E.

[93780] : Molecular chaperones enhance the degradation of expanded polyglutamine

[93781] repeat androgen receptor in a cellular model of spinal and bulbar

[93782] muscular atrophy. Hum. Molec. Genet. 11: 515–523, 2002.

[93783] 16427.

[93784] Batch, J. A.; Williams, D. M.; Davies, H. R.; Brown, B. D.; Evans,

[93785] B. A. J.; Hughes, I. A.; Patterson, M. N.: Androgen receptor gene

[93786] mutations identified by SSCP in fourteen subjects with androgen insensitivity

[93787] syndrome. Hum. Molec. Genet. 1: 497–503, 1992.

[93788] 16428.

[93789] Beitel, L. K.; Kazemi-Esfarjani, P.; Kaufman, M.; Lumbroso, R.;

[93790] DiGeorge, A. M.; Killinger, D. W.; Trifiro, M. A.; Pinsky, L.: Substitution

[93791] of arginine-839 by cysteine or histidine in the androgen

receptor

[93792] causes different receptor phenotypes in cultured cells and coordinate

[93793] degrees of clinical androgen resistance. J. Clin. Invest. 94: 546–554,

[93794] 1994.

[93795] 16429.

[93796] Belsham, D. D.; Pereira, F.; Greenberg, C. R.; Liao, S.; Wro-
gemann,

[93797] K.: Leu676-to-pro mutation of the androgen receptor
causes complete

[93798] androgen insensitivity syndrome in a large Hutterite kin-
dred. Hum.

[93799] Mutat. 5: 28–33, 1995.

[93800] 16430.

[93801] Bevan, C. L.; Brown, B. B.; Davies, H. R.; Evans, B. A. J.;
Hughes,

[93802] I. A.; Patterson, M. N.: Functional analysis of six androgen
receptor

[93803] mutations identified in patients with partial androgen in-
sensitivity

[93804] syndrome. Hum. Molec. Genet. 5: 265–273, 1996.

- [93805] 16431.
- [93806] Biancalana, V.; Serville, F.; Pommier, J.; Julien, J.; Hanauer, A.; Mandel, J. L.: Moderate instability of the trinucleotide repeat
- [93808] in spinobulbar muscular atrophy. Hum. Molec. Genet. 1: 255–258,
- [93809] 1992.
- [93810] 16432.
- [93811] Boehmer, A. L. M.; Brinkmann, A. O.; Niermeijer, M. F.; Bakker, L.; Halley, D. J. J.; Drop, S. L. S.: Germ–line and somatic mosaicism
- [93813] in the androgen insensitivity syndrome: implications for genetic counseling.
- [93814] (Letter) Am. J. Hum. Genet. 60: 1003–1006, 1997.
- [93815] 16433.
- [93816] Boehmer, A. L. M.; Brinkmann, A. O.; Nijman, R. M.; Verleun–Mooijman, M. C. T.; de Ruiter, P.; Niermeijer, M. F.; Drop, S. L. S.: Phenotypic
- [93818] variation in a family with partial androgen insensitivity

syndrome

[93819] explained by differences in 5- α dihydrotestosterone availability. J.

[93820] Clin. Endocr. Metab. 86: 1240–1246, 2001.

[93821] 16434.

[93822] Bouvattier, C.; Carel, J.-C.; Lecointre, C.; David, A.; Sultan,

[93823] C.; Bertrand, A.-M., Morel, Y.; Chaussain, J.-L.: Postnatal changes

[93824] of T, LH, and FSH in 46,XY infants with mutations in the AR gene. J.

[93825] Clin. Endocr. Metab. 87: 29–32, 2002.

[93826] 16435.

[93827] Brown, C. J.; Goss, S. J.; Lubahn, D. B.; Joseph, D. R.; Wilson,

[93828] E. M.; French, F. S.; Willard, H. F.: Androgen receptor locus on

[93829] the human X chromosome: regional localization to Xq11–12 and description

[93830] of a DNA polymorphism. Am. J. Hum. Genet. 44: 264–269, 1989.

[93831] 16436.

- [93832] Brown, T. R.; Lubahn, D. B.; Wilson, E. M.; Joseph, D. R.; French,
- [93833] F. S.; Migeon, C. J.: Deletion of the steroid-binding domain of the
- [93834] human androgen receptor gene in one family with complete androgen
- [93835] insensitivity syndrome: evidence for further genetic heterogeneity
- [93836] in this syndrome. *Proc. Nat. Acad. Sci.* 85: 8151-8155, 1988.
- [93837] 16437.
- [93838] Bruggenwirth, H. T.; Boehmer, A. L. M.; Ramnarain, S.; Verleun-Mooijman,
- [93839] M. C. T.; Satijn, D. P. E.; Trapman, J.; Grootegoed, J. A.; Brinkmann,
- [93840] A. O.: Molecular analysis of the androgen-receptor gene in a family
- [93841] with receptor-positive partial androgen insensitivity: an unusual
- [93842] type of intronic mutation. *Am. J. Hum. Genet.* 61: 1067-1077, 1997.
- [93843] 16438.

- [93844] Buchanan, G.; Yang, M.; Harris, J. M.; Nahm, H. S.; Han, G.; Moore,
- [93845] N.; Bentel, J. M.; Matusik, R. J.; Horsfall, D. J.; Marshall, V. R.;
- [93846] Greenberg, N. M.; Tilley, W. D.: Mutations at the boundary of the
- [93847] hinge and ligand binding domain of the androgen receptor confer increased
- [93848] transactivation function. *Molec. Endocr.* 15: 46–56, 2001.
- [93849] 16439.
- [93850] Butler, R.; Leigh, P. N.; McPhaul, M. J.; Gallo, J.–M.: Truncated
- [93851] forms of the androgen receptor are associated with polyglutamine expansion
- [93852] in X-linked spinal and bulbar muscular atrophy. *Hum. Molec. Genet.* 7:
- [93853] 121–127, 1998.
- [93854] 16440.
- [93855] Ho, M. F.; Monaco, A. P.; Blonden, L. A. J.; van Ommen, G. J.
- [93856] B.; Affara, N. A.; Ferguson-Smith, M. A.; Lehrach, H.: Fine mapping

- [93857] of the McLeod locus (XK) to a 150–380–kb region in Xp21.
Am. J. Hum.
- [93858] Genet. 50: 317–330, 1992.
- [93859] 16441.
- [93860] Jung, H. H.; Hergersberg, M.; Kneifel, S.; Alkadhi, H.;
Schiess,
- [93861] R.; Weigell–Weber, M.; Daniels, G.; Kollias, S.; Hess, K.:
McLeod
- [93862] syndrome: a novel mutation, predominant psychiatric
manifestations,
- [93863] and distinct striatal imaging findings. Ann. Neurol. 49:
384–392,
- [93864] 2001.
- [93865] 16442.
- [93866] Malandrini, A.; Fabrizi, G. M.; Truschi, F.; Di Pietro, G.;
Moschini,
- [93867] F.; Bartalucci, P.; Berti, G.; Salvadori, C.; Bucalossi, A.;
Guazzi,
- [93868] G.: Atypical McLeod syndrome manifested as X–linked
chorea–acanthocytosis,
- [93869] neuromyopathy and dilated cardiomyopathy: report of a
family. J.

- [93870] Neurol. Sci. 124: 89–94, 1994.
- [93871] 16443.
- [93872] Marsh, W. L.: Personal Communication. New York, N. Y.
11/13/1979.
- [93873] 16444.
- [93874] Marsh, W. L.: The Kell blood groups and their relationship
to
- [93875] chronic granulomatous disease. In: Antigens and Disease.
Am. Assoc.
- [93876] Blood Banks Symposium. (pub.) 1977. Pp. 52–66.
- [93877] 16445.
- [93878] Marsh, W. L.: Linkage relationship of the Xg and Xk loci.
Cytogenet.
- [93879] Cell Genet. 22: 531–533, 1978.
- [93880] 16446.
- [93881] Marsh, W. L.: Chronic granulomatous disease, the McLeod
syndrome,
- [93882] and the Kell blood groups. Birth Defects Orig. Art. Ser.
XIV(6A):
- [93883] 9–25, 1978.

- [93884] 16447.
- [93885] Marsh, W. L.: Chronic granulomatous disease, Kx antigen and the
- [93886] Kell blood groups. In: Brewer, G. J.: Progress in Clinical and Biological
- [93887] Research: The Red Cell. New York: Alan R. Liss (pub.) 1978. Pp.
- [93888] 493–507.
- [93889] 16448.
- [93890] Marsh, W. L.; Marsh, N. J.; Moore, A.; Symmans, W. A.; Johnson,
- [93891] C. L.; Redman, C. M.: Elevated serum creatine phosphokinase in subjects
- [93892] with McLeod syndrome. Vox Sang. 40: 403–411, 1981.
- [93893] 16449.
- [93894] Marsh, W. L.; Oyen, R.; Nichols, M. E.: Kx antigen, the McLeod
- [93895] phenotype, and chronic granulomatous disease: further studies. Vox
- [93896] Sang. 31: 356–362, 1976.
- [93897] 16450.

- [93898] Marsh, W. L.; Oyen, R.; Nichols, M. E.; Allen, F. H., Jr.:
Chronic
- [93899] granulomatous disease and the Kell blood groups. Brit. J.
Haemat. 29:
- [93900] 247–262, 1975.
- [93901] 16451.
- [93902] Schwartz, S. A.; Marsh, W. L.; Symmans, A.; et al: 'New'
clinical
- [93903] features of McLeod syndrome. (Abstract) Transfusion 22:
404 only,
- [93904] 1982.
- [93905] 16452.
- [93906] Stanfield, G. M.; Horvitz, H. R.: The ced-8 gene controls
the
- [93907] timing of programmed cell deaths in *C. elegans*. Molec.
Cell 5: 423–433,
- [93908] 2000.
- [93909] 16453.
- [93910] Supple, S. G.; Iland, H. J.; Barnett, M. H.; Pollard, J. D.:
- [93911] A spontaneous novel XK gene mutation in a patient with
McLeod syndrome. Brit.

- [93912] J. Haemat. 115: 369–372, 2001.
- [93913] 16454.
- [93914] Swash, M.; Schwartz, M. S.; Carter, N. D.; Heath, R.; Leak, M.;
- [93915] Rogers, K. L.: Benign X-linked myopathy with acanthocytes (McLeod
- [93916] syndrome): its relationship to X-linked muscular dystrophy. Brain 106:
- [93917] 717–733, 1983.
- [93918] 16455.
- [93919] Symmans, W. A.; Sheperd, C. S.; Marsh, W. L.; Oyen, R.; Shohet,
- [93920] S. B.; Linehan, B. J.: Hereditary acanthocytosis associated with
- [93921] the McLeod phenotype of the Kell blood group system. Brit. J. Haemat. 42:
- [93922] 575–583, 1979.
- [93923] 16456.
- [93924] Tippet, P.: Personal Communication. London, England 7/1981.
- [93925] 16457.

- [93926] Wimer, B. M.; Marsh, W. L.; Taswell, H. F.: Clinical characteristics
- [93927] of the McLeod blood group phenotype. (Abstract) Am. Soc. Hemat.,
- [93928] Boston , 12/1976.
- [93929] 16458.
- [93930] Wimer, B. M.; Marsh, W. L.; Taswell, H. F.; Galey, W. R.: Haematological
- [93931] changes associated with the McLeod phenotype of the Kell blood group
- [93932] system. Brit. J. Haemat. 36: 219-224, 1977.
- [93933] 16459.
- [93934] Adler, D. A.; Bressler, S. L.; Chapman, V. M.; Page, D. C.; Disteche,
- [93935] C. M.: Inactivation of the Zfx gene on the mouse X chromosome. Proc.
- [93936] Nat. Acad. Sci. 88: 4592-4595, 1991.
- [93937] 16460.
- [93938] Chong, S. S.; Kristjansson, K.; Cota, J.; Handyside, A. H.; Hughes,
- [93939] M. R.: Preimplantation prevention of X-linked disease: re-

liable and

[93940] rapid sex determination of single human cells by restriction analysis

[93941] of simultaneously amplified ZFX and ZFY sequences. Hum. Molec. Genet. 2:

[93942] 1187-1191, 1993.

[93943] 16461.

[93944] Luoh, S.-W.; Jegalian, K.; Lee, A.; Chen, E. Y.; Ridley, A.; Page,

[93945] D. C.: CpG islands in human ZFX and ZFY and mouse Zfx genes: sequence

[93946] similarities and methylation differences. Genomics 29: 353-363,

[93947] 1995.

[93948] 16462.

[93949] Muller, G.; Schempp, W.: Mapping the human ZFX locus to Xp21.3

[93950] by in situ hybridization. Hum. Genet. 82: 82-84, 1989.

[93951] 16463.

[93952] Page, D. C.; Disteché, C. M.; Simpson, E. M.; de la Chapelle, A.;

- [93953] Andersson, M.; Alitalo, T.; Brown, L. G.; Green, P.; Akots, G.: Chromosomal
- [93954] localization of ZFX--a human gene that escapes X inactivation--and
- [93955] its murine homologs. *Genomics* 7: 37-46, 1990.
- [93956] 16464.
- [93957] Palmer, M. S.; Berta, P.; Sinclair, A. H.; Pym, B.; Goodfellow,
- [93958] P. N.: Comparison of human ZFY and ZFX transcripts. *Proc. Nat.*
- [93959] *Acad. Sci.* 87: 1681-1685, 1990.
- [93960] 16465.
- [93961] Palmer, M. S.; Sinclair, A. H.; Berta, P.; Ellis, N. A.; Goodfellow,
- [93962] P. N.; Abbas, N. E.; Fellous, M.: Genetic evidence that ZFY is not
- [93963] the testis-determining factor. *Nature* 342: 937-939, 1989.
- [93964] 16466.
- [93965] Scherer, G.; Schempp, W.; Baccichetti, C.; Lenzini, E.; Dagna Bricarelli,

- [93966] F.; Carbone, L. D. L.; Wolf, U.: Duplication of an Xp segment that
- [93967] includes the ZFX locus causes sex inversion in man. Hum. Genet. 81:
- [93968] 291–294, 1989.
- [93969] 16467.
- [93970] Schneider–Gadicke, A.; Beer–Romero, P.; Brown, L. G.; Nussbaum,
- [93971] R.; Page, D. C.: ZFX has a gene structure similar to ZFY, the putative
- [93972] human sex determinant, and escapes X inactivation. Cell 57: 1247–1258,
- [93973] 1989.
- [93974] 16468.
- [93975] Seibold, A.; Brabet, P.; Rosenthal, W.; Birnbaumer, M.: Structure
- [93976] and chromosomal localization of the human antidiuretic hormone receptor
- [93977] gene. Am. J. Hum. Genet. 51: 1078–1083, 1992.
- [93978] 16469.
- [93979] Trousseau, A.: Polydipsia. Lecture LXV. In: Clinique Medi–

cale

[93980] de L'Hotel-Dieu de Paris. Vol. 2. Paris: J.-B. Balliere 1862.

[93981] P. 607.

[93982] 16470.

[93983] Tsukaguchi, H.; Matsubara, H.; Taketani, S.; Mori, Y.;
Seido,

[93984] T.; Inada, M.: Binding-, intracellular transport-, and
biosynthesis-defective

[93985] mutants of vasopressin type 2 receptor in patients with X-
linked nephrogenic

[93986] diabetes insipidus. J. Clin. Invest. 96: 2043-2050, 1995.

[93987] 16471.

[93988] Uttley, W. S.; Thistlethwaite, D.: Failure to detect the car-
rier

[93989] in congenital nephrogenic diabetes insipidus. Arch. Dis.
Child. 47:

[93990] 137-138, 1972.

[93991] 16472.

[93992] van den Ouweland, A.; Markslag, P.; Knoers, N.; Ropers, H.
H.;

[93993] Poustka, A.; Jans, D.; Fahrenholz, F.; van Oost, B.: Colocal-

ization

- [93994] of the locus for human nephrogenic diabetes insipidus and the vasopressin
- [93995] type-2 receptor gene in the Xq28 region. (Series) Miami Short
- [93996] Reports. Advances in Gene Technology: The Molecular Biology of Human
- [93997] Genetic Disease. New York: IRL Press (pub.) 1: 1991. Pp. 20
- [93998] only.
- [93999] 16473.
- [94000] van den Ouweland, A. M. W.; Dreesen, J. C. F. M.; Verdijk, M.;
- [94001] Knoers, N. V. A. M.; Monnens, L. A. H.; Rocchi, M.; van Oost, B. A.
- [94002] : Mutations in the vasopressin type 2 receptor gene (AVPR2) associated
- [94003] with nephrogenic diabetes insipidus. Nature Genet. 2: 99-102, 1992.
- [94004] 16474.
- [94005] van den Ouweland, A. M. W.; Knoop, M. T.; Knoers, V. V. A. M.;

- [94006] Markslag, P. W. B.; Rocchi, M.; Warren, S. T.; Ropers, H. H.;
Fahrenholz,
- [94007] F.; Monnens, L. A.; van Oost, B. A. H.: Colocalization of the
gene
- [94008] for nephrogenic diabetes insipidus (DIR) and the vaso-
pressin type
- [94009] 2 receptor gene (AVPR2) in the Xq28 region. Genomics 13:
1350–1352,
- [94010] 1992.
- [94011] 16475.
- [94012] van Lieburg, A. F.; Knoers, N. V. A. M.; Monnens, L. A. H.:
Clinical
- [94013] presentation and follow-up of 30 patients with congenital
nephrogenic
- [94014] diabetes insipidus. J. Am. Soc. Nephrol. 10: 1958–1964,
1999.
- [94015] 16476.
- [94016] van Lieburg, A. F.; Verdijk, M. A. J.; Schoute, F.; Ligten-
berg,
- [94017] M. J. L.; van Oost, B. A.; Waldhauser, F.; Dobner, M.; Mon-
nens, L.
- [94018] A. H.; Knoers, N. V. A. M.: Clinical phenotype of nephro-

genic diabetes

[94019] insipidus in females heterozygous for a vasopressin type 2 receptor

[94020] mutation. Hum. Genet. 96: 70–78, 1995.

[94021] 16477.

[94022] Wildin, R. S.; Antush, M. J.; Bennett, R. L.; Schoof, J. M.; Scott,

[94023] C. R.: Heterogeneous AVPR2 gene mutations in congenital nephrogenic

[94024] diabetes insipidus. Am. J. Hum. Genet. 55: 266–277, 1994.

[94025] 16478.

[94026] Hata, A.; Matsuura, T.; Setoyama, C.; Shimada, K.; Yokoi, T.;

[94027] Akaboshi, I.; Matsuda, I.: A novel missense mutation in exon 8 of

[94028] the ornithine transcarbamylase gene in two unrelated male patients

[94029] with mild ornithine transcarbamylase deficiency. Hum. Genet. 87:

[94030] 28–32, 1991.

[94031] 16479.

- [94032] Masutani, C.; Sugasawa, K.; Yanagisawa, J.; Sonoyama, T.;
Ui, M.;
- [94033] Enomoto, T.; Takio, K.; Tanaka, K.; van der Spek, P. J.;
Bootsma,
- [94034] D.; Hoeijmakers, J. H. J.; Hanaoka, F.: Purification and
cloning
- [94035] of a nucleotide excision repair complex involving the xe-
roderma pigmentosum
- [94036] group C protein and a human homologue of yeast RAD23.
EMBO J. 13:
- [94037] 1831-1843, 1994.
- [94038] 16480.
- [94039] Angelini, C.; Fanin, M.; Menegazzo, E.; Freda, M. P.; Dug-
gan, D.
- [94040] J.; Hoffman, E. P.: Homozygous alpha-sarcoglycan muta-
tion in two
- [94041] siblings: one asymptomatic and one steroid-responsive
mild limb-girdle
- [94042] muscular dystrophy patient. Muscle Nerve 21: 769-775,
1998.
- [94043] 16481.
- [94044] Derst, C.; Konrad, M.; Kockerling, A.; Karolyi, L.; Desch-

enes,

[94045] G.; Daut, J.; Karschin, A.; Seyberth, H. W.: Mutations in the ROMK

[94046] gene in antenatal Bartter syndrome are associated with impaired K(+)

[94047] channel function. Biochem. Biophys. Res. Commun. 203: 641–645, 1997.

[94048] 16482.

[94049] Derst, C.; Wischmeyer, E.; Preisig–Muller, R.; Spauschus, A.; Konrad,

[94050] M.; Hensen, P.; Jeck, N.; Seyberth, H. W.; Daut, J.; Karschin, A.

[94051] : A hyperprostaglandin E syndrome mutation in Kir1.1 (renal outer

[94052] medullary potassium) channels reveals a crucial residue for channel

[94053] function in Kir1.3 channels. J. Biol. Chem. 273: 23884–23891, 1998.

[94054] 16483.

[94055] Ho, K.; Nichols, C. G.; Lederer, W. J.; Lytton, J.; Vassilev, P.

[94056] M.; Kanazirska, M. V.; Hebert, S. C.: Cloning and expression of an

[94057] inwardly rectifying ATP-regulated potassium channel. Nature 362:

[94058] 31-38, 1993.

[94059] 16484.

[94060] International Collaborative Study Group for Bartter-like Syndromes

[94061] : Mutations in the gene encoding the inwardly-rectifying renal potassium

[94062] channel, ROMK, cause the antenatal variant of Bartter syndrome: evidence

[94063] for genetic heterogeneity. Hum. Molec. Genet. 6: 17-26, 1997.

[94064] 16485.

[94065] Krishnan, S. N.; Desai, T.; Ward, D. C.; Haddad, G. G.: Isolation

[94066] and chromosomal localization of a human ATP-regulated potassium channel. Hum.

[94067] Genet. 96: 155-160, 1995.

[94068] 16486.

[94069] Shuck, M. E.; Bock, J. H.; Benjamin, C. W.; Tsai, T.-D.; Lee, K.

- [94070] S.; Slightom, J. L.; Bienkowski, M. J.: Cloning and characterization
of multiple forms of the human kidney ROM-K potassium
channel. J.
- [94072] Biol. Chem. 269: 24261–24270, 1994.
- [94073] 16487.
- [94074] Simon, D. B.; Karet, F. E.; Rodriguez-Soriano, J.; Hamdan,
J. H.;
- [94075] DiPietro, A.; Trachtman, H.; Sanjad, S. A.; Lifton, R. P.: Ge-
netic
heterogeneity of Bartter's syndrome revealed by mutations
in the K(+)
channel, ROMK. Nature Genet. 14: 152–156, 1996.
- [94078] 16488.
- [94079] Yano, H.; Philipson, L. H.; Kugler, J. L.; Tokuyama, Y.;
Davis,
E. M.; Le Beau, M. M.; Nelson, D. J.; Bell, G. I.; Takeda, J.:
Alternative
splicing of human inwardly rectifying K(+) channel ROMK1
mRNA. Molec.
- [94082] Pharm. 45: 854–860, 1994.

- [94083] 16489.
- [94084] Campbell, H. D.; Fountain, S.; McLennan, I. S.; Berven, L. A.;
- [94085] Crouch, M. F.; Davy, D. A.; Hooper, J. A.; Waterford, K.; Chen, K.-S.;
- [94086] Lupski, J. R.; Ledermann, B.; Young, I. G.; Matthaei, K. I.: Fliih,
- [94087] a gelsolin-related cytoskeletal regulator essential for early mammalian
- [94088] embryonic development. *Molec. Cell. Biol.* 22: 3518–3526, 2002.
- [94089] 16490.
- [94090] Campbell, H. D.; Fountain, S.; Young, I. G.; Claudianos, C.; Hoheisel,
- [94091] J. D.; Chen, K.-S.; Lupski, J. R.: Genomic structure, evolution,
- [94092] and expression of human FLII, a gelsolin and leucine-rich-repeat family
- [94093] member: overlap with LLGL. *Genomics* 42: 46–54, 1997.
- [94094] 16491.
- [94095] Connelly, M. A.; Zhang, H.; Kieleczawa, J.; Anderson, C. W.: The

- [94096] promoters for human DNA-PK(cs) (PRKDC) and MCM4: divergently transcribed
- [94097] genes located at chromosome 8 band q11. Genomics 47: 71–83, 1998.
- [94098] 16492.
- [94099] Aly, A. M.; Hoyer, L. W.: Factor VIII–East Hartford (arginine1689
- [94100] to cysteine) has procoagulant activity when separated from von Willebrand
- [94101] factor. J. Clin. Invest. 89: 1382–1387, 1992.
- [94102] 16493.
- [94103] Antonarakis, S. E.; Copeland, K. L.; Carpenter, R. J., Jr.; Carta,
- [94104] C. A.; Hoyer, L. W.; Caskey, C. T.; Toole, J. J.; Kazazian, H. H.,
- [94105] Jr.: Prenatal diagnosis of haemophilia A by factor VIII gene analysis. Lancet I:
- [94106] 1407–1409, 1985.
- [94107] 16494.
- [94108] Antonarakis, S. E.; Kazazian, H. H.; Tuddenham, G. D.: Molecular

- [94109] etiology of factor VIII deficiency in hemophilia A. Hum. Mutat. 5: 1-22, 1995.
- [94110] 16495.
- [94111] Antonarakis, S. E.; Rossiter, J. P.; Young, M.; Horst, J.; de Moerloose, P.; Sommer, S. S.; Ketterling, R. P.; Kazazian, H. H., Jr.; Ne-grier, C.; Vinciguerra, C.; Gitschier, J.; Goossens, M.; and 54 others : Factor VIII inversions in severe hemophilia A: results from an international consortium. Blood 86: 2206-2212, 1995.
- [94112] 16496.
- [94113] Antonarakis, S. E.; Waber, P. G.; Kittur, S. D.; Patel, A. S.; Kazazian, H. H., Jr.; Mellis, M. A.; Counts, R. B.; Stamatoy-annopoulos, G.; Bowie, E. J. W.; Fass, D. N.; Pittman, D. D.; Wozney, J. M.; Toole, J. J.: Hemophilia A: detection of molecular defects and of carriers

- [94122] by DNA analysis. New Eng. J. Med. 313: 842–848, 1985.
- [94123] 16497.
- [94124] Antonarakis, S. E.; Youssoufian, H.; Kazazian, H. H.:
Molecular
- [94125] genetics of hemophilia–A in man (factor VIII deficiency).
Molec.
- [94126] Biol. Med. 4: 81–95, 1987.
- [94127] 16498.
- [94128] Arai, M.; Higuchi, M.; Antonarakis, S. E.; Kazazian, H. H.,
Jr.;
- [94129] Phillips, J. A., III; Janco, R. L.; Hoyer, L. W.: Characteriza-
tion
- [94130] of a thrombin cleavage site mutation (arg1689–to–cys) in
the factor
- [94131] VIII gene of two unrelated patients with cross–reacting
material–positive
- [94132] hemophilia A. Blood 75: 384–389, 1990.
- [94133] 16499.
- [94134] Arai, M.; Inaba, H.; Higuchi, M.; Antonarakis, S. E.;
Kazazian,
- [94135] H. H., Jr.; Fujimaki, M.; Hoyer, L. W.: Direct characteriza–

tion of

[94136] factor VIII in plasma: detection of a mutation altering a
thrombin

[94137] cleavage site (arginine372-to-histidine). Proc. Nat. Acad.
Sci. 86:

[94138] 4277-4281, 1989.

[94139] 16500.

[94140] Arrants, J. E.; Jordan, P. H., Jr.; Newcomb, T. F.: Von Wille-
brand's

[94141] disease: a cause for massive postoperative bleeding-
report of a case. Ann.

[94142] Surg. 156: 845-851, 1962.

[94143] 16501.

[94144] Arveiler, B.; Vincent, A.; Mandel, J.-L.: Toward a physical
map

[94145] of the Xq28 region in man: linking color vision, G6PD, and
coagulation

[94146] factor VIII genes to an X-Y homology region. Genomics 4:
460-471,

[94147] 1989.

[94148] 16502.

- [94149] Bardoni, B.; Sampietro, M.; Romano, M.; Crapanzano, M.; Mannucci,
[94150] P. M.; Camerino, G.: Characterization of a partial deletion
of the
[94151] factor VIII gene in a haemophiliac with inhibitor. Hum.
Genet. 79:
[94152] 86–88, 1988.
[94153] 16503.
[94154] Barker, D.; Schafer, M.; White, R.: Restriction sites con-
taining
[94155] CpG show a higher frequency of polymorphism in human
DNA. Cell 36:
[94156] 131–138, 1984.
[94157] 16504.
[94158] Barrai, I.; Cann, H. M.; Cavalli-Sforza, L. L.: Segregation
analysis
[94159] of hemophilia A and B. (Letter) Am. J. Hum. Genet. 31:
226–227,
[94160] 1979.
[94161] 16505.
[94162] Barrai, I.; Cann, H. M.; Cavalli-Sforza, L. L.; de Nicola, P.

- [94163] : The effect of parental age on rates of mutation for hemophilia and
- [94164] evidence for differing mutation rates for hemophilia A and B. Am.
- [94165] J. Hum. Genet. 20: 175–196, 1968.
- [94166] 16506.
- [94167] Barrow, E. M. S.; Graham, J. B.: Factor VIII. (Letter) Lancet I:
- [94168] 1312–1313, 1973.
- [94169] 16507.
- [94170] Baty, B. J.; Drayna, D.; Leonard, C. O.; White, R.: Prenatal
- [94171] diagnosis of factor VIII deficiency to help with the management of
- [94172] pregnancy and delivery. (Letter) Lancet I: 207 only, 1986.
- [94173] 16508.
- [94174] Becker, J.; Schwaab, R.; Moller-Taube, A.; Schwaab, U.; Schmidt,
- [94175] W.; Brackmann, H. H.; Grimm, T.; Olek, K.; Oldenburg, J.: Characterization
- [94176] of the factor VIII defect in 147 patients with sporadic hemophilia

- [94177] A: family studies indicate a mutation type-dependent sex ratio of
- [94178] mutation frequencies. *Am. J. Hum. Genet.* 58: 657-670, 1996.
- [94179] 16509.
- [94180] Bennett, B.; Ratnoff, O. D.: Deletion of the carrier state for
- [94181] classic hemophilia. *New Eng. J. Med.* 7: 342-345, 1974.
- [94182] 16510.
- [94183] Bennett, E.; Huehns, E. R.: Immunological differentiation of
- [94184] three types of hemophilia and identification of some female carriers. *Lancet* II:
- [94185] 956-958, 1970.
- [94186] 16511.
- [94187] Berg, L.-P.; Wieland, K.; Millar, D. S.; Schlosser, M.; Wagner,
- [94188] M.; Kakkar, V. V.; Reiss, J.; Cooper, D. N.: Detection of a novel
- [94189] point mutation causing haemophilia A by PCR/direct sequencing of ectopically-transcribed
- [94190] factor VIII mRNA. *Hum. Genet.* 85: 655-658, 1990.

- [94191] 16512.
- [94192] Bernardi, F.; Marchetti, G.; Bertagnolo, V.; Faggioli, L.;
Volinia,
- [94193] S.; Patracchini, P.; Bartolai, S.; Vannini, F.; Felloni, L.;
Rossi,
- [94194] L.; Panicucci, F.; Conconi, F.: RFLP analysis in families with
sporadic
- [94195] hemophilia A: estimate of the mutation ratio in male and
female gametes. Hum.
- [94196] Genet. 76: 253–256, 1987.
- [94197] 16513.
- [94198] Bernardi, F.; Volinia, S.; Patracchini, P.; Gemmati, D.;
Boninsegna,
- [94199] S.; Schwienbacher, C.; Marchetti, G.: A recurrent missense
mutation
- [94200] (arg-to-gln) and a partial deletion in factor VIII gene caus-
ing severe
- [94201] haemophilia A. Brit. J. Haemat. 71: 271–276, 1989.
- [94202] 16514.
- [94203] Bidichandani, S. I.; Shiach, C. R.; Lanyon, W. G.; Connor, J.
- [94204] M.: A novel splice donor mutation affecting position +3 in

intron

[94205] 6 of the factor VIII gene. Hum. Molec. Genet. 3: 651–653, 1994.

[94206] 16515.

[94207] Biggs, R.; Rizza, C. R.: The sporadic case of haemophilia A. Lancet II:

[94208] 431–433, 1976.

[94209] 16516.

[94210] Bird, A. P.: DNA methylation and the frequency of CpG in animal

[94211] DNA. Nucleic Acids Res. 8: 1499–1504, 1980.

[94212] 16517.

[94213] Bloom, A. L.; Peake, I. R.: Molecular genetics of factor VIII

[94214] and its disorders. Am. J. Path. 88: 319–340, 1977.

[94215] 16518.

[94216] Bond, T. P.; Levin, W. C.; Celandier, D. R.; Guest, M. M.: 'Mild

[94217] hemophilia' affecting both males and females. New Eng. J. Med. 266:

[94218] 220–223, 1962.

- [94219] 16519.
- [94220] Boyer, S. H.; Siggers, D. C.; Krueger, L. J.: A caveat to protein
- [94221] replacement therapy for genetic disease. Immunologic implications
- [94222] of accurate molecular diagnosis. Lancet II: 654-659, 1973.
- [94223] 16520.
- [94224] Briet, E.; Bakker, B.; Brocker-Vriends, A.: Somatic and germinal
- [94225] mosaicism for a deletion in the factor VIII gene. (Abstract) Brit.
- [94226] J. Haemat. 71 (suppl. 1): 1 only, 1989.
- [94227] 16521.
- [94228] Greenough, W. T.; Klintsova, A. Y.; Irwin, S. A.; Galvez, R.;
- [94229] Bates, K. E.; Weiler, I. J.: Synaptic regulation of protein synthesis
- [94230] and the fragile X protein. Proc. Nat. Acad. Sci. 98: 7101-7106,
- [94231] 2001.
- [94232] 16522.

- [94233] Griffiths, M. J.; Strachan, M. C.: A single lymphocyte culture
- [94234] for fragile X induction and prometaphase chromosome analysis. J.
- [94235] Med. Genet. 28: 837–839, 1991.
- [94236] 16523.
- [94237] Haataja, R.; Vaisanen, M.–L.; Li, M.; Ryyananen, M.; Leisti, J.
- [94238] : The fragile X syndrome in Finland: demonstration of a founder effect
- [94239] by analysis of microsatellite haplotypes. Hum. Genet. 94: 479–483,
- [94240] 1994.
- [94241] 16524.
- [94242] Hagerman, R. J.; Leehey, M.; Heinrichs, W.; Tassone, F.; Wilson,
- [94243] R.; Hills, J.; Grigsby, J.; Gage, B.; Hagerman, P. J.: Intention
- [94244] tremor, parkinsonism, and generalized brain atrophy in male carriers
- [94245] of fragile X. Neurology 57: 127–130, 2001.
- [94246] 16525.

- [94247] Hagerman, R. J.; Synhorst, D. P.: Mitral valve prolapse and aortic
- [94248] dilatation in the fragile X syndrome. *Am. J. Med. Genet.* 17: 123–131,
- [94249] 1984.
- [94250] 16526.
- [94251] Hagerman, R. J.; Van Housen, K.; Smith, A. C. M.; McGavran, L.
- [94252] : Consideration of connective tissue dysfunction in the fragile X
- [94253] syndrome. *Am. J. Med. Genet.* 17: 111–121, 1984.
- [94254] 16527.
- [94255] Hanzlik, A. J.; Osemlak-Hanzlik, M. M.; Hauser, M. A.; Kurnit,
- [94256] D. M.: A recombination-based assay demonstrates that the fragile
- [94257] X sequence is transcribed widely during development. *Nature Genet.* 3:
- [94258] 44–48, 1993.
- [94259] 16528.
- [94260] Harvey, J.; Judge, C.; Wiener, S.: Familial X-linked mental

retardation

[94261] with an X chromosome abnormality. J. Med. Genet. 14: 46-50, 1977.

[94262] 16529.

[94263] Hecht, F.; Cannizzaro, L. A.; Hecht, B. K.: Gene symbols for

[94264] fragile sites: a proposal. (Letter) Hum. Genet. 82: 394, 1989.

[94265] 16530.

[94266] Hecht, F.; Glover, T. W.: Antibiotics containing trimethoprim

[94267] and the fragile X chromosome. (Letter) New Eng. J. Med. 308: 285-286,

[94268] 1983.

[94269] 16531.

[94270] Hecht, F.; Jacky, P. B.; Sutherland, G. R.: The fragile X chromosome:

[94271] current methods. Am. J. Med. Genet. 11: 489-495, 1982.

[94272] 16532.

[94273] Hinds, H. L.; Ashley, C. T.; Sutcliffe, J. S.; Nelson, D. L.;

[94274] Warren, S. T.; Housman, D. E.; Schalling, M.: Tissue spe-

cific expression

[94275] of FMR-1 provides evidence for a functional role in fragile X syndrome. *Nature*

[94276] *Genet.* 3: 36-43, 1993.

[94277] 16533.

[94278] Hirst, M.; Grewal, P.; Flannery, A.; Slatter, R.; Maher, E.; Barton,

[94279] D.; Fryns, J.-P.; Davies, K.: Two new cases of FMR1 deletion associated

[94280] with mental impairment. *Am. J. Hum. Genet.* 56: 67-74, 1995.

[94281] 16534.

[94282] Hirst, M. C.; Knight, S. J. L.; Christodoulou, Z.; Grewal, P.

[94283] K.; Fryns, J. P.; Davies, K. E.: Origins of the fragile X syndrome

[94284] mutation. *J. Med. Genet.* 30: 647-650, 1993.

[94285] 16535.

[94286] Hofker, M. H.; Bergen, A. A. B.; Skraastad, M. I.; Carpenter,

[94287] N. J.; Veenema, H.; Connor, J. M.; Bakker, E.; van Ommen, G. J. B.;

[94288] Pearson, P. L.: Efficient isolation of X chromosome-spe-

cific single

[94289] copy probes from a cosmid library of a human X/hamster hybrid-cell

[94290] line: mapping of new probes close to the locus for X-linked mental

[94291] retardation. Am. J. Hum. Genet. 40: 312-328, 1987.

[94292] 16536.

[94293] Howard-Peebles, P. N.; Stoddard, G. R.: Familial X-linked mental

[94294] retardation with a marker X chromosome and its relationship to macroorchidism. Clin.

[94295] Genet. 17: 125-128, 1980.

[94296] 16537.

[94297] Hundscheid, R. D. L.; Sistermans, E. A.; Thomas, C. M. G.; Braat,

[94298] D. D. M.; Straatman, H.; Kiemeney, L. A. L. M.; Oostra, B. A.; Smits,

[94299] A. P. T.: Imprinting effect in premature ovarian failure confined

[94300] to paternally inherited fragile X premutations. Am. J. Hum. Genet. 66:

[94301] 413-418, 2000.

[94302] 16538.

[94303] Hundscheid, R. D. L.; Thomas, C. M. G.; Braat, D. D. M.;
Oostra,

[94304] B. A.; Smits, A. P. T.: Reply to the letters from Murray et al.
and

[94305] Vianna-Morgante and Costa. (Letter) Am. J. Hum. Genet.
67: 256-258,

[94306] 2000.

[94307] 16539.

[94308] ISCN: An International System for Human Cytogenetic
Nomenclature

[94309] (1978). Cytogenet. Cell Genet. 21: 309-404, 1978.

[94310] 16540.

[94311] Jacky, P. B.; Dill, F. J.: Expression in fibroblast culture of

[94312] the satellited-X chromosome associated with familial sex-
linked mental

[94313] retardation. Hum. Genet. 53: 267-269, 1980.

[94314] 16541.

[94315] Jacobs, P. A.: Personal Communication. Honolulu, Hawaii
1982.

- [94316] 16542.
- [94317] Jacobs, P. A.: The fragile X syndrome. (Editorial) J. Med. Genet. 28:
[94318] 809–810, 1991.
- [94319] 16543.
- [94320] Jacobs, P. A.; Glover, T. W.; Mayer, M.; Fox, P.; Gerrard, J.
[94321] W.; Dunn, H. G.; Herbst, D. S.: X-linked mental retardation: a study
[94322] of 7 families. Am. J. Med. Genet. 7: 471–479, 1980.
- [94323] 16544.
- [94324] Jacobs, P. A.; Hunt, P. A.; Mayer, M.; Wang, J.-C.; Boss, G.
R.;
- [94325] Erbe, R. W.: Expression of the marker(X)(q28) in lymphoblastoid cell
[94326] lines. Am. J. Hum. Genet. 34: 552–557, 1982.
- [94327] 16545.
- [94328] Jakala, P.; Hanninen, T.; Ryyanen, M.; Laakso, M.; Partanen,
[94329] K.; Mannermaa, A.; Soininen, H.: Fragile-X: neuropsychological test

[94330] performance, CGG triplet repeat lengths, and hippocampal volumes. J.

[94331] Clin. Invest. 100: 331–338, 1997.

[94332] 16546.

[94333] Jenkins, E. C.; Brown, W. T.; Brooks, J.; Duncan, C. J.; Rudelli,

[94334] R. D.; Wisniewski, H. M.: Experience with prenatal fragile X detection. Am.

[94335] J. Med. Genet. 17: 215–239, 1984.

[94336] 16547.

[94337] Jenkins, E. C.; Brown, W. T.; Duncan, C. J.; Brooks, J.: Demonstration

[94338] of the fragile X chromosome in amniotic fluid cells. (Abstract) Clin.

[94339] Res. 30: 292A, 1982.

[94340] 16548.

[94341] Jin, P.; Warren, S. T.: Understanding the molecular basis of

[94342] fragile X syndrome. Hum. Molec. Genet. 9: 901–908, 2000.

[94343]

[94344] 100. Johnson, V. P.; Carpenter, N. J.; Skorey, P. A.: Martin-Bell

[94345] syndrome segregating in a large kindred with normal transmitting males:

[94346] clinical, cytogenetic, and linkage study. Am. J. Med. Genet. 38:

[94347] 275-282, 1991.

[94348]

[94349] 101. Kaiser-McCaw, B.; Hecht, F.; Cadien, J. D.; Moore, B. C.: Fragile

[94350] X-linked mental retardation. Am. J. Hum. Genet. 7: 503-505, 1980.

[94351]

[94352] 102. Kenneson, A.; Zhang, F.; Hagedorn, C. H.; Warren, S. T.: Reduced

[94353] FMRP and increased FMR1 transcription is proportionally associated

[94354] with CGG repeat number in intermediate-length and pre-mutation carriers. Hum.

[94355] Molec. Genet. 10: 1449-1454, 2001.

[94356]

- [94357] 103. Khalifa, M. M.; Reiss, A. L.; Migeon, B. R.: Methylation status
[94358] of genes flanking the fragile site in males with the fragile-X syndrome:
[94359] a test of the imprinting hypothesis. Am. J. Hum. Genet. 46: 744-753,
[94360] 1990.
[94361]
[94362] 104. Khandjian, E. W.; Corbin, F.; Woerly, S.; Rousseau, F.: The
[94363] fragile X mental retardation protein is associated with ribosomes. Nature
[94364] Genet. 12: 91-93, 1996.
[94365]
[94366] 105. Kinnell, H. G.: Fragile-X disorder associated with antisocial
[94367] personality. (Letter) Lancet II: 1104, 1982.
[94368]
[94369] 106. Kirchgessner, C. U.; Warren, S. T.; Willard, H. F.: X in-activation
[94370] if the FMR1 fragile X mental retardation gene. J. Med.

Genet. 32:

[94371] 925–929, 1995.

[94372]

[94373] 107. Kirkilionis, A. J.; Chudley, A. E.; Greenberg, C. R.; Yan, D.

[94374] L.; McGillivray, B.; Hamerton, J. L.: Transmission of the fra(X)

[94375] haplotype from three nonpenetrant brothers to their affected grandsons. Am.

[94376] J. Med. Genet. 43: 588–591, 1992.

[94377]

[94378] 108. Klauck, S. M.; Munstermann, E.; Bieber–Martig, B.; Ruhl, D.;

[94379] Lisch, S.; Schmotzer, G.; Poustka, A.; Poustka, F.: Molecular genetic

[94380] analysis of the FRM–1 gene in a large collection of autistic patients. Hum.

[94381] Genet. 100: 224–229, 1997.

[94382]

[94383] 109. Knight, S. J. L.; Flannery, A. V.; Hirst, M. C.; Campbell, L.;

- [94384] Christodoulou, Z.; Phelps, S. R.; Pointon, J.; Middleton-Price, H.
- [94385] R.; Barnicoat, A.; Pembrey, M. E.; Holland, J.; Oostra, B. A.; Bobrow,
- [94386] M.; Davies, K. E.: Trinucleotide repeat amplification and hypermethylation
- [94387] of a CpG island in FRAXE mental retardation. Cell 74: 127-134, 1993.
- [94388]
- [94389] 110. Kooy, R. F.; Reyniers, E.; Verhoye, M.; Sijbers, J.; Bakker,
- [94390] C. E.; Oostra, B. A.; Willems, P. J.; Van Der Linden, A.: Neuroanatomy
- [94391] of the fragile X knockout mouse brain studied using in vivo high resolution
- [94392] magnetic resonance imaging. Europ. J. Hum. Genet. 7: 526-532, 1999.
- [94393]
- [94394] 111. Krawczun, M. S.; Jenkins, E. C.; Brown, W. T.: Analysis of the
- [94395] fragile-X chromosome: localization and detection of the fragile site

[94396] in high resolution preparations. Hum. Genet. 69:
209–211, 1985.

[94397]

[94398] 112. Kruyer, H.; Mila, M.; Glover, G.; Carbonell, P.;
Ballesta, F.;

[94399] Estivill, X.: Fragile X syndrome and the (CGG)_n mutation:
two families

[94400] with discordant MZ twins. Am. J. Hum. Genet. 54:
437–442, 1994.

[94401]

[94402] 113. Kunst, C. B.; Leeflang, E. P.; Iber, J. C.; Arnheim, N.;
Warren,

[94403] S. T.: The effect of FMR1 CGG repeat interruptions on mu-
tation frequency

[94404] as measured by sperm typing. J. Med. Genet. 34:
627–631, 1997.

[94405]

[94406] 114. Kunst, C. B.; Zerylnick, C.; Karickhoff, L.; Eichler, E.;
Bullard,

[94407] J.; Chalifoux, M.; Holden, J. J. A.; Torroni, A.; Nelson, D. L.;
Warren,

[94408] S. T.: FMR1 in global populations. Am. J. Hum. Genet. 58:

513–522,

[94409] 1996.

[94410]

[94411] 115. Lachiewicz, A. M.; Dawson, D. V.; Spiridigliozzi, G.
A.: Physical

[94412] characteristics of young boys with fragile X syndrome:
reasons for

[94413] difficulties in making a diagnosis in young males. Am. J.
Med. Genet. 92:

[94414] 229–236, 2000.

[94415]

[94416] 116. Lagerbauer, B.; Ostareck, D.; Keidel, E.–M.;
Ostareck–Lederer,

[94417] A.; Fischer, U.: Evidence that fragile X mental retardation
protein

[94418] is a negative regulator of translation. Hum. Molec. Genet.
10: 329–338,

[94419] 2001.

[94420]

[94421] 117. Laird, C. D.: Possible erasure of the imprint on a
fragile X

[94422] chromosome when transmitted by a male. Am. J. Med.
Genet. 38: 391–395,

[94423] 1991.

[94424]

[94425] 118. Laird, C. D.: Proposed mechanism of inheritance and
expression

[94426] of the human fragile X syndrome of mental retardation.
Genetics 117:

[94427] 587–599, 1987.

[94428]

[94429] 119. Laird, C. D.; Jaffe, E.; Karpen, G.; Lamb, M.; Nelson,
R.: Fragile

[94430] sites in human chromosomes as regions of late-
replicating DNA. Trends

[94431] Genet. 3: 274–281, 1987.

[94432]

[94433] 120. Laird, C. D.; Lamb, M. M.; Thorne, J. L.: Two progeni-
tor cells

[94434] for human oogonia inferred from pedigree data and the
X-inactivation

[94435] imprinting model of the fragile-X syndrome. Am. J. Hum.

Genet. 46:

[94436] 696–719, 1990.

[94437]

[94438] 121. Langenbeck, U.; Schmidtke, J.; Bartels, I.; Hansmann, I.; Knuppel,

[94439] H.: Mean corpuscular hemoglobin is increased in Martin–Bell syndrome. Hum.

[94440] Genet. 66: 365–366, 1984.

[94441]

[94442] 122. Larsen, L. A.; Armstrong, J. S. M.; Gronskov, K.; Hjalgrim, H.;

[94443] Brondum–Nielsen, K.; Hasholt, L.; Norgaard–Pedersen, B.; Vuust, J.

[94444] : Analysis of FMR1 (CGG)*n* alleles and FRAXA microsatellite haplotypes

[94445] in the population of Greenland: implications for the population of

[94446] the New World from Asia. Europ. J. Hum. Genet. 7: 771–777, 1999.

[94447]

[94448] 123. Larsen, L. A.; Vuust, J.; Nystad, M.; Evseeva, I.; Van

Ghelue,

- [94449] M.; Tranebjaerg, L.: Analysis of FMR1 (CGG)_n alleles and DXS548-FRAXAC1
- [94450] haplotypes in three European circumpolar populations: traces of genetic
- [94451] relationship with Asia. *Europ. J. Hum. Genet.* 9: 724-727, 2001.
- [94452]
- [94453] 124. Laval, S. H.; Blair, H. J.; Hirst, M. C.; Davies, K. E.; Boyd,
- [94454] Y.: Mapping of FMR1, the gene implicated in fragile X-linked mental
- [94455] retardation, on the mouse X chromosome. *Genomics* 12: 818-821, 1992.
- [94456]
- [94457] 125. Lejeune, J.; Legrand, N.; Lafourcade, J.; Rethore, M.-O.; Raoul,
- [94458] O.; Maunoury, C.: Fragilite du chromosome X et effets de la trimethoprimine. *Ann.*
- [94459] *Genet.* 25: 149-151, 1982.
- [94460]
- [94461] 126. Lewis, H. A.; Musunuru, K.; Jensen, K. B.; Edo, C.;

Chen, H.;

[94462] Darnell, R. B.; Burley, S. K.: Sequence-specific RNA binding by a

[94463] Nova KH domain: implications for paraneoplastic disease and the fragile

[94464] X syndrome. Cell 100: 323–332, 2000.

[94465]

[94466] 127. Limprasert, P.; Jaruratanasirikul, S.; Vasiknanonte, P.: Unilateral

[94467] macroorchidism in fragile X syndrome. (Letter) Am. J. Med. Genet. 95:

[94468] 516–517, 2000.

[94469]

[94470] 128. Linden, M. G.; Tassone, F.; Gane, L. W.; Hills, J. L.; Hagerman,

[94471] R. J.; Taylor, A. K.: Compound heterozygous female with fragile X

[94472] syndrome. Am. J. Med. Genet. 83: 318–321, 1999.

[94473]

[94474] 129. Loesch, D. Z.; Hay, D. A.: Clinical features and reproductive

[94475] patterns in fragile X female heterozygotes. J. Med. Genet.
25: 407–414,

[94476] 1988.

[94477]

[94478] 130. Lubs, H.; Travers, H.; Lujan, E.; Carroll, A.: A large
kindred

[94479] with X-linked mental retardation, marker X and
macroorchidism. Am.

[94480] J. Med. Genet. 17: 145–157, 1984.

[94481]

[94482] 131. Lubs, H. A.; Watson, M.; Breg, R.; Lujan, E.: Restudy
of the

[94483] original marker X family. Am. J. Med. Genet. 17: 133–144,
1984.

[94484]

[94485] 132. Lubs, H. A., Jr.: A marker X chromosome. Am. J.
Hum. Genet. 21:

[94486] 231–244, 1969.

[94487]

[94488] 133. Lugenbeel, K. A.; Peier, A. M.; Carson, N. L.; Chudley,
A. E.;

- [94489] Nelson, D. L.: Intragenic loss of function mutations demonstrate
- [94490] the primary role of FMR1 in fragile X syndrome. *Nature Genet.* 10:
- [94491] 483–485, 1995.
- [94492]
- [94493] 134. Malter, H. E.; Iber, J. C.; Willemsen, R.; de Graaff, E.; Tarleton,
- [94494] J. C.; Leisti, J.; Warren, S. T.; Oostra, B. A.: Characterization
- [94495] of the full fragile X syndrome mutation in fetal gametes. *Nature Genet.* 15: 165–169, 1997.
- [94496]
- [94497]
- [94498] 135. Mandel, J.–L.; Hagerman, R.; Froster, U.; Brown, W. T.; Jenkins,
- [94499] E. C.; Jacobs, P.; Turner, G.; Lubs, H.; Neri, G.: Fifth International
- [94500] Workshop on the Fragile X and X-Linked Mental Retardation. *Am. J. Med. Genet.* 43: 5–27, 1992.
- [94501]
- [94502]

[94503] 136. Martin, J. P.; Bell, J.: A pedigree of mental defect showing

[94504] sex-linkage. J. Neurol. Psychiat. 6: 154–157, 1943.

[94505]

[94506] 137. Martin, R. H.; Lin, C. C.; Mathies, B. J.; Lowry, R. B.: X-linked

[94507] mental retardation with macro-orchidism and marker-X chromosomes. Am.

[94508] J. Med. Genet. 7: 433–441, 1980.

[94509]

[94510] 138. Mattei, J. F.; Mattei, M. G.; Aumeras, C.; Auger, M.; Giraud,

[94511] F.: X-linked mental retardation with the fragile X: a study of 15

[94512] families. Hum. Genet. 59: 281–289, 1981.

[94513]

[94514] 139. McCabe, E. R. B.; de la Cruz, F.; Clapp, K.: Workshop on Fragile

[94515] X: future research directions. Am. J. Med. Genet. 85: 317–322, 1999.

[94516]

- [94517] 140. McConkie-Rosell, A.; Lachiewicz, A. M.;
Spiridigliozzi, G. A.;
- [94518] Tarleton, J.; Schoenwald, S.; Phelan, M. C.; Goonewardena,
P.; Ding,
- [94519] X.; Brown, W. T.: Evidence that methylation of the FMR-1
locus is
- [94520] responsible for variable phenotypic expression of the
fragile X syndrome. Am.
- [94521] J. Hum. Genet. 53: 800-809, 1993.
- [94522]
- [94523] 141. Meijer, H.; de Graaff, E.; Merckx, D. M. L.; Jongbloed,
R. J.
- [94524] E.; de Die-Smulders, C. E. M.; Engelen, J. J. M.; Fryns, J.-P.;
- Curfs,
- [94525] P. M. G.; Oostra, B. A.: A deletion of 1.6 kb proximal to
the CGG
- [94526] repeat of the FMR1 gene causes the clinical phenotype of
the fragile
- [94527] X syndrome. Hum. Molec. Genet. 3: 615-620, 1994.
- [94528]
- [94529] 142. Meryash, D. L.; Cronk, C. E.; Sachs, B.; Gerald, P. S.:
An anthropometric

[94530] study of males with the fragile-X syndrome. Am. J. Med.
Genet. 17:

[94531] 159-174, 1984.

[94532]

[94533] 143. Mila, M.; Castellvi-Bel, S.; Gine, R.; Vazquez, C.;
Badenas,

[94534] C.; Sanchez, A.; Estivill, X.: A female compound heterozy-
gote (pre-

[94535] and full mutation) for the CGG FMR1 expansion. Hum.
Genet. 98: 419-421,

[94536] 1996.

[94537]

[94538] 144. Mila, M.; Sanchez, A.; Badenas, C.; Brun, C.; Jimenez,
D.; Villa,

[94539] M. P.; Castellvi-Bel, S.; Estivill, X.: Screening for FMR1 and
FMR2

[94540] mutations in 222 individuals from Spanish special
schools: identification

[94541] of a case of FRAXE-associated mental retardation. Hum.
Genet. 100:

[94542] 503-507, 1997.

[94543]

[94544] 145. Mingroni-Netto, R. C.; Angeli, C. B.; Auricchio, M. T. B. M.;

[94545] Leal-Mesquita, E. R.; Ribeiro-dos-Santos, A. K. C.; Ferrari, I.; Hutz,

[94546] M. H.; Salzano, F. M.; Hill, K.; Hurtado, A. M.; Vianna-Morgante,

[94547] A. M.: Distribution of CGG repeats and FRAXAC1/DXS548 alleles in

[94548] South American populations. Am. J. Med. Genet. 111: 243-252, 2002.

[94549]

[94550] 146. Morales, J.; Hiesinger, P. R.; Schroeder, A. J.; Kume, K.; Verstreken,

[94551] P.; Jackson, F. R.; Nelson, D. L.; Hassan, B. A.: Drosophila fragile

[94552] X protein, DFXR, regulates neuronal morphology and function in the

[94553] brain. Neuron 34: 961-972, 2002.

[94554]

[94555] 147. Morton, J. E.; Bunday, S.; Webb, T. P.; MacDonald, F.; Rindl,

[94556] P. M.; Bullock, S.: Fragile X syndrome is less common than previously

[94557] estimated. J. Med. Genet. 34: 1–5, 1997.

[94558]

[94559] 148. Morton, N. E.; Macpherson, J. N.: Population genetics of the

[94560] fragile–X syndrome: multiallelic model for the FMR1 locus. Proc.

[94561] Nat. Acad. Sci. 89: 4215–4217, 1992.

[94562]

[94563] 149. Mulley, J. C.; Yu, S.; Gedeon, A. K.; Donnelly, A.; Turner, G.;

[94564] Loesch, D.; Chapman, C. J.; Gardner, R. J. M.; Richards, R. I.; Sutherland,

[94565] G. R.: Experience with direct molecular diagnosis of fragile X. J.

[94566] Med. Genet. 29: 368–374, 1992.

[94567]

[94568] 150. Mulligan, L. M.; Phillips, M. A.; Forster–Gibson, C. J.; Beckett,

[94569] J.; Partington, M. W.; Simpson, N. E.; Holden, J. J. A.; White,

B.

[94570] N.: Genetic mapping of DNA segments relative to the locus for the

[94571] fragile-X syndrome at Xq27.3. *Am. J. Hum. Genet.* 37: 463-472, 1985.

[94572]

[94573] 151. Murray, A.; Ennis, S.; Morton, N.: No evidence for parent of

[94574] origin influencing premature ovarian failure in fragile X premutation

[94575] carriers. (Letter) *Am. J. Hum. Genet.* 67: 253-254, 2000.

[94576]

[94577] 152. Murray, A.; Webb, J.; Grimley, S.; Conway, G.; Jacobs, P.: Studies

[94578] of FRAXA and FRAXE in women with premature ovarian failure. *J. Med.*

[94579] *Genet.* 35: 637-640, 1998.

[94580]

[94581] 153. Nolin, S. L.; Lewis, F. A., III; Ye, L. L.; Houck, G. E., Jr.;

[94582] Glicksman, A. E.; Limprasert, P.; Li, S. Y.; Zhong, N.; Ashley, A.

[94583] E.; Feingold, E.; Sherman, S. L.; Brown, W. T.: Familial transmission

[94584] of the FMR1 GCC repeat. *Am. J. Hum. Genet.* 59: 1252–1261, 1996.

[94585]

[94586] 154. Nussbaum, R. L.; Ledbetter, D. H.: Fragile X syndrome: a unique

[94587] mutation in man. *Annu. Rev. Genet.* 20: 109–145, 1986.

[94588]

[94589] 155. Oberle, I.; Rousseau, F.; Heitz, D.; Kretz, C.; Devys, D.; Hanauer,

[94590] A.; Boue, J.; Bertheas, M. F.; Mandel, J. L.: Instability of a 550–base

[94591] pair DNA segment and abnormal methylation in fragile X syndrome. *Science* 252:

[94592] 1097–1102, 1991.

[94593]

[94594] 156. Oostra, B. A.; Chiurazzi, P.: The fragile X gene and its function. *Clin.*

[94595] *Genet.* 60: 399–408, 2001.

[94596]

- [94597] 157. Opitz, J. M.; Sutherland, G. R.: International work-shop on the
[94598] fragile X and X-linked mental retardation. Am. J. Med.
Genet. 17:
[94599] 5-94, 1984.
- [94600]
- [94601] 158. Opitz, J. M.; Westphal, J. M.; Daniel, A.: Discovery of a
connective
[94602] tissue dysplasia in the Martin-Bell syndrome. Am. J. Med.
Genet. 17:
[94603] 101-109, 1984.
- [94604]
- [94605] 159. Oudet, C.; Mornet, E.; Serre, J. L.; Thomas, F.;
Lentes-Zengerling,
[94606] S.; Kretz, C.; Deluchat, C.; Tejada, I.; Boue, J.; Boue, A.;
Mandel,
[94607] J. L.: Linkage disequilibrium between the fragile X muta-
tion and
[94608] two closely linked CA repeats suggests that fragile X chro-
mosomes
[94609] are derived from a small number of founder chromo-
somes. Am. J. Hum.

[94610] Genet. 52: 297–304, 1993.

[94611]

[94612] 160. Park, V.; Howard–Peebles, P.; Sherman, S.; Taylor, A.; Wulfsberg,

[94613] E.: Policy statement: American College of Medical Genetics. Fragile

[94614] X syndrome: diagnostic and carrier testing. Am. J. Med. Genet. 53:

[94615] 380–381, 1994.

[94616]

[94617] 161. Parrish, J. E.; Oostra, B. A.; Verkerk, A. J. M. H.; Richards,

[94618] C. S.; Reynolds, J.; Spikes, A. S.; Shaffer, L. G.; Nelson, D. L.

[94619] : Isolation of a GCC repeat showing expansion in FRAXF, a fragile

[94620] site distal to FRAXA and FRAXE. Nature Genet. 8: 229–235, 1994.

[94621]

[94622] 162. Peier, A. M.; McIlwain, K. L.; Kenneson, A.; Warren, S. T.; Paylor,

[94623] R.; Nelson, D. L.: (Over)correction of FMR1 deficiency with YAC transgenics:

[94624] behavioral and physical features. Hum. Molec. Genet. 9: 1145–1159,

[94625] 2000.

[94626]

[94627] 163. Pellissier, M.–C.; Voelckel, M.–A.; Piquet, C.; Mattei, M.–G.;

[94628] Mattei, J.–F.: Transmission of mental retardation with fragile X

[94629] site by two normal transmitter brothers. Am. J. Med. Genet. 38:

[94630] 367–369, 1991.

[94631]

[94632] 164. Pembrey, M. E.; Winter, R. M.; Davies, K. E.: A premutation

[94633] that generates a defect at crossing over explains the inheritance

[94634] of fragile X mental retardation. Am. J. Med. Genet. 21: 709–717,

[94635] 1985.

[94636]

[94637] 165. Pieretti, M.; Zhang, F.; Fu, Y.-H.; Warren, S. T.; Oostra, B.

[94638] A.; Caskey, C. T.; Nelson, D. L.: Absence of expression of the FMR-1

[94639] gene in fragile X syndrome. Cell 66: 817-822, 1991.

[94640]

[94641] 166. Poustka, A.; Dietrich, A.; Langenstein, G.; Toniolo, D.; Warren,

[94642] S. T.; Lehrach, H.: Physical map of human Xq27-qter: localizing the

[94643] region of the fragile X mutation. Proc. Nat. Acad. Sci. 88: 8302-8306,

[94644] 1991.

[94645]

[94646] 167. Purrello, M.; Alhadeff, B.; Esposito, D.; Szabo, P.; Rocchi,

[94647] M.; Truett, M.; Masiarz, F.; Siniscalco, M.: The human genes for

[94648] hemophilia A and hemophilia B flank the X chromosome fragile site

[94649] at Xq27.3. EMBO J. 4: 725–729, 1985.

[94650]

[94651] 168. Pyeritz, R. E.; Stamberg, J.; Thomas, G. H.; Bell, B. B.; Zahka,

[94652] K. G.; Bernhardt, B. A.: The marker Xq28 syndrome (fragile X syndrome)

[94653] in a retarded man with mitral valve prolapse. Johns Hopkins Med.

[94654] J. 151: 231–237, 1982.

[94655]

[94656] 169. Quan, F.; Grompe, M.; Jakobs, P.; Popovich, B. W.: Spontaneous

[94657] deletion in the FMR1 gene in a patient with fragile X syndrome and

[94658] cherubism. Hum. Molec. Genet. 4: 1681–1684, 1995.

[94659]

[94660] 170. Quan, F.; Zonana, J.; Gunter, K.; Peterson, K. L.; Magenis, R.

[94661] E.; Popovich, B. W.: An atypical case of fragile X syndrome caused

[94662] by a deletion that includes the FMR1 gene. Am. J. Hum.

Genet. 56:

[94663] 1042-1051, 1995.

[94664]

[94665] 171. Ramos, F. J.; Emanuel, B. S.; Spinner, N. B.: Frequency of the

[94666] common fragile site at Xq27.2 under conditions of thymidylate stress:

[94667] implications for cytogenetic diagnosis of the fragile-X syndrome. Am.

[94668] J. Med. Genet. 42: 835-838, 1992.

[94669]

[94670] 172. Reiss, A. L.; Aylward, E.; Freund, L. S.; Joshi, P. K.; Bryan,

[94671] R. N.: Neuroanatomy of fragile X syndrome: the posterior fossa. Ann.

[94672] Neurol. 29: 26-32, 1991.

[94673]

[94674] 173. Reiss, A. L.; Freund, L.; Tseng, J. E.; Joshi, P. K.: Neuroanatomy

[94675] in fragile X females: the posterior fossa. Am. J. Hum. Genet. 49:

[94676] 279–288, 1991.

[94677]

[94678] 174. Reiss, A. L.; Lee, J.; Freund, L.: Neuroanatomy of fragile X

[94679] syndrome: the temporal lobe. *Neurology* 44: 1317–1324, 1994.

[94680]

[94681] 175. Reyniers, E.; Vits, L.; De Boulle, K.; Van Roy, B.; Van Velzen,

[94682] D.; de Graaff, E.; Verkerk, A. J. M. H.; Jorens, H. Z. J.; Darby,

[94683] J. K.; Oostra, B.; Willems, P. J.: The full mutation in the FMR-1

[94684] gene of male fragile X patients is absent in their sperm. *Nature*

[94685] *Genet.* 4: 143–148, 1993.

[94686]

[94687] 176. Richards, B. W.; Sylvester, P. E.; Brooker, C.: Fragile X-linked

[94688] mental retardation: the Martin–Bell syndrome. *J. Ment. Defic. Res.* 25:

[94689] 253–256, 1981.

[94690]

[94691] 177. Richards, R. I.; Holman, K.; Friend, K.; Kremer, E.; Hillen,

[94692] D.; Staples, A.; Brown, W. T.; Goonewardena, P.; Tarleton, J.; Schwartz,

[94693] C.; Sutherland, G. R.: Evidence of founder chromosomes in fragile

[94694] X syndrome. *Nature Genet.* 1: 257–260, 1992.

[94695]

[94696] 178. Richards, R. I.; Holman, K.; Kozman, H.; Kremer, E.; Lynch, M.;

[94697] Pritchard, M.; Yu, S.; Mulley, J.; Sutherland, G. R.: Fragile X syndrome:

[94698] genetic localisation by linkage mapping of two microsatellite repeats

[94699] FRAXAC1 and FRAXAC2 which immediately flank the fragile site. *J.*

[94700] *Med. Genet.* 28: 818–823, 1991.

[94701]

[94702] 179. Richards, R. I.; Holman, K.; Kozman, H.; Kremer, E.;

Lynch, M.;

[94703] Pritchard, M.; Yu, S.; Mulley, J.; Sutherland, G. R.: Fragile X syndrome:

[94704] genetic localisation by linkage mapping of two microsatellite repeats

[94705] FRAXAC1 and FRAXAC2 which immediately flank the fragile site. J.

[94706] Med. Genet. 28: 818–823, 1991.

[94707]

[94708] 180. Richards, R. I.; Sutherland, G. R.: Dynamic mutations: a new

[94709] class of mutations causing human disease. Cell 70: 709–712, 1992.

[94710]

[94711] 181. Rocchi, M.; Archidiacono, N.; Rinaldi, A.; Filippi, G.; Bartolucci,

[94712] G.; Fancello, G. S.; Siniscalco, M.: Mental retardation in heterozygotes

[94713] for the fragile-X mutation: evidence in favor of an X inactivation-dependent

[94714] effect. Am. J. Hum. Genet. 46: 738–743, 1990.

[94715]

[94716] 182. Rodewald, L.; Miller, D. C.; Sciorra, L.; Barabas, G.;
Lee, M.-L.

[94717] : Central nervous system neoplasm in a young man with
Martin-Bell

[94718] syndrome--fra(X)-XLMR. Am. J. Med. Genet. 26: 7-12,
1987.

[94719]

[94720] 183. Rousseau, F.; Heitz, D.; Biancalana, V.; Blumenfeld,
S.; Kretz,

[94721] C.; Boue, J.; Tommerup, N.; Van Der Hagen, C.; DeLozier-
Blanchet,

[94722] C.; Croquette, M.-F.; Gilgenkrantz, S.; Jalbert, P.; Voelckel,
M.-A.;

[94723] Oberle, I.; Mandel, J.-L.: Direct diagnosis by DNA analysis
of the

[94724] fragile X syndrome of mental retardation. New Eng. J. Med.
325:

[94725] 1673-1681, 1991.

[94726]

[94727] 184. Rousseau, F.; Heitz, D.; Mandel, J.-L.: The unstable
and methylatable

[94728] mutations causing the fragile X syndrome. Hum. Mutat. 1:

91–96,

[94729] 1992.

[94730]

[94731] 185. Rousseau, F.; Heitz, D.; Oberle, I.; Mandel, J.-L.: Se-
lection

[94732] in blood cells from female carriers of the fragile X syn-
drome: inverse

[94733] correlation between age and proportion of active X chro-
mosomes carrying

[94734] the full mutation. J. Med. Genet. 28: 830–836, 1991.

[94735]

[94736] 186. Rousseau, F.; Heitz, D.; Tarleton, J.; MacPherson, J.;
Malmgren,

[94737] H.; Dahl, N.; Barnicoat, A.; Mathew, C.; Mornet, E.; Tejada,
I.; Maddalena,

[94738] A.; Spiegel, R.; Schinzel, A.; Marcos, J. A. G.; Schorderet,
D. F.;

[94739] Schaap, T.; Maccioni, L.; Russo, S.; Jacobs, P. A.; Schwartz,
C.;

[94740] Mandel, J. L.: A multicenter study on genotype–phenotype
correlations

[94741] in the fragile X syndrome, using direct diagnosis with

probe StB12.3:

[94742] the first 2,253 cases. Am. J. Hum. Genet. 55: 225–237, 1994.

[94743]

[94744] 187. Russo, S.; Briscioli, V.; Cogliati, F.; Macchi, M.; Lalatta,

[94745] F.; Larizza, L.: An unusual fragile X sibship: female compound heterozygote

[94746] and male with a partially methylated full mutation. Clin. Genet. 54:

[94747] 309–314, 1998.

[94748]

[94749] 188. Ruvalcaba, R. H. A.; Myhre, S. A.; Roosen-Runge, E. C.; Beckwith,

[94750] J. B.: X-linked mental deficiency megalotestes syndrome. J.A.M.A. 238:

[94751] 1646–1650, 1977.

[94752]

[94753] 189. Schrandt-Stumpel, C.; Gerver, W.-J.; Meyer, H.; Engelen, J.;

[94754] Mulder, H.; Fryns, J.-P.: Prader-Willi-like phenotype in

fragile

[94755] X syndrome. Clin. Genet. 45: 175–180, 1994.

[94756]

[94757] 190. Sherman, S. L.: Premature ovarian failure among fragile X premutation

[94758] carriers: parent-of-origin effect? (Editorial) Am. J. Hum. Genet. 67:

[94759] 11–13, 2000.

[94760]

[94761] 191. Sherman, S. L.; Jacobs, P. A.; Morton, N. E.; Froster-Iskenius,

[94762] U.; Howard-Peebles, P. N.; Nielsen, K. B.; Partington, M. W.; Sutherland,

[94763] G. R.; Turner, G.; Watson, M.: Further segregation analysis of the

[94764] fragile X syndrome with special reference to transmitting males. Hum.

[94765] Genet. 69: 289–299, 1985.

[94766]

[94767] 192. Siomi, H.; Choi, M.; Siomi, M. C.; Nussbaum, R. L.; Dreyfuss,

- [94768] G.: Essential role for KH domains in RNA binding: impaired RNA binding
- [94769] by a mutation in the KH domain of FMR1 that causes fragile X syndrome. Cell 77:
- [94770] 33–39, 1994.
- [94771]
- [94772] 193. Siomi, H.; Siomi, M. C.; Nussbaum, R. L.; Dreyfuss, G.: The
- [94773] protein product of the fragile X gene, FMR1, has characteristics of
- [94774] an RNA-binding protein. Cell 74: 291–298, 1993.
- [94775]
- [94776] 194. Sittler, A.; Devys, D.; Weber, C.; Mandel, J.-L.: Alternative
- [94777] splicing of exon 14 determines nuclear or cytoplasmic localisation
- [94778] of fmr1 protein isoforms. Hum. Molec. Genet. 5: 95–102, 1996.
- [94779]
- [94780] 195. Smeets, H. J. M.; Smits, A. P. T.; Verheij, C. E.; Theelen, J.

[94781] P. G.; Willemsen, R.; van de Burgt, I.; Hoogeveen, A. T.;
Oosterwijk,

[94782] J. C.; Oostra, B. A.: Normal phenotype in two brothers with
a full

[94783] FMR1 mutation. Hum. Molec. Genet. 4: 2103–2108, 1995.

[94784]

[94785] 196. Smits, A. P. T.; Dreesen, J. C. F. M.; Post, J. G.;
Smeets, D.

[94786] F. C. M.; de Die-Smulders, C.; Spaans-van der Bijl, T.; Go-
vaerts,

[94787] L. C. P.; Warren, S. T.; Oostra, B. A.; van Oost, B. A.: The
fragile

[94788] X syndrome: no evidence for any recent mutations. J. Med.
Genet. 30:

[94789] 94–96, 1993.

[94790]

[94791] 197. Snow, K.; Doud, L. K.; Hagerman, R.; Pergolizzi, R. G.;
Erster,

[94792] S. H.; Thibodeau, S. N.: Analysis of a CGG sequence at the
FMR–1

[94793] locus in fragile X families and in the general population.
Am. J.

[94794] Hum. Genet. 53: 1217–1228, 1993.

[94795]

[94796] 198. Snyder, F. F.; Lin, C. C.; Harasym, C. A.; Jamro, H. K.;

Kushnig,

[94797] M. L.; Lowe, J. K.; O'Brien, S. I.: Evidence for close association

[94798] between the fragile X(q27-28) chromosome site and glucose-6-phosphate

[94799] dehydrogenase (G6PD) but not with hypoxanthine-guanine phosphoribosyltransferase

[94800] (HPRT). (Abstract) Cytogenet. Cell Genet. 37: 587, 1984.

[94801]

[94802] 199. Soysa, P.; Senanayahe, M.; Mikkelsen, M.; Poulsen, H.: Martin-Bell

[94803] syndrome fra(X)(q28) in a Sri Lankan family. J. Ment. Defic. Res. 26:

[94804] 251-257, 1982.

[94805]

[94806] 200. Steinbach, P.: Mental impairment in Martin-Bell syndrome is

[94807] probably determined by interaction of several genes: simple explanation

[94808] of phenotypic differences between unaffected and affected males with

[94809] the same X chromosome. Hum. Genet. 72: 248-252,

1986.

[94810]

[94811] 201. Stoll, C.: Problems in the diagnosis of fragile X syndrome in

[94812] young children are still present. Am. J. Med. Genet. 100: 110–115,

[94813] 2001.

[94814]

[94815] 202. Storm, K.; Handig, I.; Reyniers, E.; Oostra, B. A.; Kooy, R.

[94816] F.; Willems, P. J.: Incomplete EcoRI digestion may lead to false

[94817] diagnosis of fragile X syndrome. Hum. Genet. 102: 54–56, 1998.

[94818]

[94819] 203. Subramanian, P. S.; Nelson, D. L.; Chinault, A. C.: Large domains

[94820] of apparent delayed replication timing associated with triplet repeat

[94821] expansion at FRAXA and FRAXE. Am. J. Hum. Genet. 59: 407–416, 1996.

[94822]

- [94823] 204. Sun, Y. J.; Baumer, A.: Nonrandom X inactivation and selection
[94824] of fragile X full mutation in fetal fibroblasts. Am. J. Med. Genet. 86:
[94825] 162–164, 1999.
[94826]
- [94827] 205. Sutherland, G. R.: Personal Communication. Adelaide, Australia
[94828] 11/3/1993.
[94829]
- [94830] 206. Sutherland, G. R.: Fragile sites on human chromosomes: demonstration
[94831] of their dependence on the type of tissue culture medium. Science 197:
[94832] 265–266, 1977.
[94833]
- [94834] 207. Sutherland, G. R.: Heritable sites on human chromosomes. VII.
[94835] Children homozygous for the BrdU–requiring fra10q25 are phenotypically
[94836] normal. Am. J. Hum. Genet. 33: 946–949, 1981.

[94837]

[94838] 208. Sutherland, G. R.: Heritable fragile sites on human chromosomes.

[94839] II. Distribution, phenotypic effects, and cytogenetics. Am. J. Hum.

[94840] Genet. 31: 136–148, 1979.

[94841]

[94842] 209. Sutherland, G. R.: Personal Communication. Adelaide, Australia

[94843] 6/1989.

[94844]

[94845] 210. Sutherland, G. R.; Ashford, P. L. C.: X-linked mental retardation

[94846] with macro-orchidism and the fragile site at Xq27 or 28. Hum. Genet. 48:

[94847] 117–120, 1979.

[94848]

[94849] 211. Sutherland, G. R.; Baker, E.: Characterisation of a new rare

[94850] fragile site easily confused with the fragile X. Hum. Molec. Genet. 1:

[94851] 111–113, 1992.

[94852]

[94853] 212. Sutherland, G. R.; Gedeon, A.; Kornman, L.; Donnelly, A.; Byard,

[94854] R. W.; Mulley, J. C.; Kremer, E.; Lynch, M.; Pritchard, M.; Yu, S.;

[94855] Richards, R. I.: Prenatal diagnosis of fragile X syndrome by direct

[94856] detection of the unstable DNA sequence. New Eng. J. Med. 325: 1720–1722,

[94857] 1991.

[94858]

[94859] 213. Sved, J. A.; Laird, C. D.: Population genetic consequences of

[94860] the fragile X syndrome, based on the X-inactivation imprinting model. Am.

[94861] J. Hum. Genet. 46: 443–451, 1990.

[94862]

[94863] 214. Sved, J. A.; Laird, C. D.: The X-inactivation imprinting model

[94864] can explain the incidence of the fragile-X syndrome of

mental retardation

[94865] in mother-offspring pairs. Brain Dysfunction 1: 245-254, 1988.

[94866]

[94867] 215. Szabo, P.; Purrello, M.; Rocchi, M.; Archidiacono, N.; Alhadeff,

[94868] B.; Filippi, G.; Toniolo, D.; Martini, G.; Luzzatto, L.; Siniscalco,

[94869] M.: Cytological mapping of the human glucose-6-phosphate dehydrogenase

[94870] gene distal to the fragile-X site suggests a high rate of meiotic

[94871] recombination across this site. Proc. Nat. Acad. Sci. 81: 7855-7859,

[94872] 1984.

[94873]

[94874] 216. Tamanini, F.; Bontekoe, C.; Bakker, C. E.; van Unen, L.; Anar,

[94875] B.; Willemsen, R.; Yoshida, M.; Galjaard, H.; Oostra, B. A.; Hoogeveen,

[94876] A. T.: Different targets for the fragile X-related proteins revealed

[94877] by their distinct nuclear localizations. Hum. Molec. Genet.
8: 863–869,

[94878] 1999.

[94879]

[94880] 217. Tarleton, J.; Richie, R.; Schwartz, C.; Rao, K.;
Aylsworth, A.

[94881] S.; Lachiewicz, A.: An extensive de novo deletion removing
FMR1 in

[94882] a patient with mental retardation and the fragile X syn-
drome phenotype. Hum.

[94883] Molec. Genet. 2: 1973–1974, 1993.

[94884]

[94885] 218. Thibodeau, S. N.; Dorkins, H. R.; Faulk, K. R.; Berry,
R.; Smith,

[94886] A. C. M.; Hagerman, R.; King, A.; Davies, K. E.: Linkage
analysis

[94887] using multiple DNA polymorphic markers in normal fami-
lies and in families

[94888] with fragile X syndrome. Hum. Genet. 79: 219–227, 1988.

[94889]

[94890] 219. Thode, A.; Laing, S.; Partington, M. W.; Turner, G.: Is

there

[94891] a fragile(X) negative Martin-Bell syndrome?. Am. J. Med. Genet. 30:

[94892] 459-471, 1988.

[94893]

[94894] 220. Toledano-Alhadeff, H.; Basel-Vanagaite, L.; Magal, N.; Davidov,

[94895] B.; Ehrlich, S.; Drasinover, V.; Taub, E.; Halpern, G. J.; Ginott,

[94896] N.; Shohat, M.: Fragile X carrier screening and the prevalence of

[94897] premutation and full-mutation carriers in Israel. Am. J. Hum. Genet. 69:

[94898] 351-360, 2001.

[94899]

[94900] 221. Tommerup, N.; Poulsen, H.; Brondum-Nielsen, K.: 5-Fluoro-2-prime-deoxyuridine

[94901] induction of the fragile site on Xq28 associated with X-linked mental

[94902] retardation. J. Med. Genet. 33: 234-242, 1981.

[94903]

- [94904] 222. Turner, G.; Brookwell, R.; Daniel, A.; Selikowitz, M.; Zilibowitz,
- [94905] M.: Heterozygous expression of X-linked mental retardation and X-chromosome
- [94906] marker fra(X)(q27). New Eng. J. Med. 303: 662-664, 1980.
- [94907]
- [94908] 223. Turner, G.; Daniel, A.; Frost, M.: X-linked mental retardation,
- [94909] macro-orchidism, and the Xq27 fragile site. J. Pediat. 96: 837-841,
- [94910] 1980.
- [94911]
- [94912] 224. Turner, G.; Eastman, C.; Casey, J.; McLeay, A.; Procopis, P.;
- [94913] Turner, B.: X-linked mental retardation associated with macro-orchidism. J.
- [94914] Med. Genet. 12: 367-371, 1975.
- [94915]
- [94916] 225. Turner, G.; Till, R.; Daniel, A.: Marker X chromosomes, mental
- [94917] retardation and macro-orchidism. (Letter) New Eng. J.

Med. 299:

[94918] 1472, 1978.

[94919]

[94920] 226. Turner, G.; Webb, T.; Wake, S.; Robinson, H.: Prevalence of

[94921] fragile X syndrome. Am. J. Med. Genet. 64: 196–197, 1996.

[94922]

[94923] 227. Uchida, I. A.; Joyce, E. M.: Activity of the fragile X in heterozygous

[94924] carriers. Am. J. Hum. Genet. 34: 286–293, 1982.

[94925]

[94926] 228. van den Ouweland, A. M. W.; Deelen, W. H.; Kunst, C. B.; Uzielli,

[94927] M.–L. G.; Nelson, D. L.; Warren, S. T.; Oostra, B. A.; Halley, D.

[94928] J. J.: Loss of mutation at the FMR1 locus through multiple exchanges

[94929] between maternal X chromosomes. Hum. Molec. Genet. 3: 1823–1827,

[94930] 1994.

[94931]

[94932] 229. Van Roy, B. C.; De Smedt, M. C.; Raes, R. A.; Dumon, J. E.; Leroy,

[94933] J. G.: Fragile X trait in a large kindred: transmission also through

[94934] normal males. J. Med. Genet. 20: 286–289, 1983.

[94935]

[94936] 230. Verheij, C.; Bakker, C. E.; de Graaff, E.; Keulemans, J.; Willemsen,

[94937] R.; Verkerk, A. J. M. H.; Galjaard, H.; Reuser, A. J. J.; Hoogeveen,

[94938] A. T.; Oostra, B. A.: Characterization and localization of the FMR-1

[94939] gene product associated with fragile X syndrome. Nature 363: 722–724,

[94940] 1993.

[94941]

[94942] 231. Verheij, C.; de Graaff, E.; Bakker, C. E.; Willemsen, R.; Willems,

[94943] P. J.; Meijer, N.; Galjaard, H.; Reuser, A. J. J.; Oostra, B. A.;

[94944] Hoogeveen, A. T.: Characterization of FMR1 proteins iso-

lated from

[94945] different tissues. Hum. Molec. Genet. 4: 895–901, 1995.

[94946]

[94947] 232. Verkerk, A. J. M. H.; Pieretti, M.; Sutcliffe, J. S.; Fu, Y.-H.;

[94948] Kuhl, D. P. A.; Pizzuti, A.; Reiner, O.; Richards, S.; Victoria, M.

[94949] F.; Zhang, F.; Eussen, B. E.; van Ommen, G.-J. B.; Blonden, L. A.

[94950] J.; Riggins, G. J.; Chastain, J. L.; Kunst, C. B.; Galjaard, H.; Caskey,

[94951] C. T.; Nelson, D. L.; Oostra, B. A.; Warren, S. T.: Identification

[94952] of a gene (FMR-1) containing a CCG repeat coincident with a breakpoint

[94953] cluster region exhibiting length variation in fragile X syndrome. Cell 65:

[94954] 905–914, 1991.

[94955]

[94956] 233. Vianna-Morgante, A. M.; Costa, S. S.: Premature ovarian failure

[94957] is associated with maternally and paternally inherited pre-

mutation

- [94958] in Brazilian families with fragile X. (Letter) Am. J. Hum. Genet. 67: 254–255, 2000.
- [94959]
- [94960]
- [94961] 234. Vincent, J. B.; Gurling, H. M. D.: Point mutation in intron 10 of FMR1 is unlikely to be a cause of fragile X syndrome. (Letter) Hum. Mutat. 12: 431 only, 1998.
- [94964]
- [94965] 235. Voelckel, M. A.; Mattei, M. G.; N'Guyen, C.; Philip, N.; Birg, F.; Mattei, J. F.: Dissociation between mental retardation and fragile site expression in a family with fragile X-linked mental retardation. Hum. Genet. 80: 375–378, 1988.
- [94969]
- [94970] 236. Vogel, F.; Crusio, W. E.; Kovac, C.; Fryns, J.-P.; Freund, M.

[94971] : Selective advantage of fra (X) heterozygotes. Hum.

Genet. 86:

[94972] 25–32, 1990.

[94973]

[94974] 237. Wallace, D. C.; Torroni, A.: American Indian prehistory as written

[94975] in the mitochondrial DNA: a review. Hum. Biol. 64: 403–416, 1992.

[94976]

[94977] 238. Wang, Y.-C.; Li, S.-Y.: Response to Vincent and Gurling. (Letter) Hum.

[94978] Mutat. 12: 432 only, 1998.

[94979]

[94980] 239. Wang, Y.-C.; Lin, M.-L.; Lin, S. J.; Li, Y.-C.; Li, S.-Y.: Novel

[94981] point mutation within intron 10 of FMR-1 gene causing fragile X syndrome. Hum.

[94982] Mutat. 10: 393–399, 1997.

[94983]

[94984] 240. Warren, S. T.; Glover, T. W.; Davidson, R. L.; Jagadeeswaran,

[94985] P.: Linkage and recombination between fragile X-linked mental retardation

[94986] and the factor IX gene. (Letter) Hum. Genet. 69: 44–46, 1985.

[94987]

[94988] 241. Warren, S. T.; Knight, S. J. L.; Peters, J. F.; Stayton, C. L.;

[94989] Consalez, G. G.; Zhang, F. P.: Isolation of the human chromosomal

[94990] band Xq28 within somatic cell hybrids by fragile X site breakage. Proc.

[94991] Nat. Acad. Sci. 87: 3856–3860, 1990.

[94992]

[94993] 242. Warren, S. T.; Zhang, F.; Licameli, G. R.; Peters, J. F.: The

[94994] fragile X site in somatic cell hybrids: an approach for molecular

[94995] cloning of fragile sites. Science 237: 420–423, 1987.

[94996]

[94997] 243. Warren, S. T.; Zhang, F. P.; Sutcliffe, J. S.; Peters, J. F.

[94998] : Strategy for molecular cloning of the fragile X site DNA.

Am. J.

[94999] Med. Genet. 30: 613–623, 1988.

[95000]

[95001] 244. Weaver, D. D.; Sherman, S. L.: A counseling guide to the Martin–Bell

[95002] syndrome. (Letter) Am. J. Med. Genet. 26: 39–44, 1987.

[95003]

[95004] 245. Webb, T.; Jacobs, P. A.: Fragile Xq27.3 in female heterozygotes

[95005] for the Martin–Bell syndrome. J. Med. Genet. 27: 627–631, 1990.

[95006]

[95007] 246. Webb, T. P.; Bunday, S.; Thake, A.; Todd, J.: The frequency

[95008] of the fragile X chromosome among schoolchildren in Coventry. J.

[95009] Med. Genet. 23: 396–399, 1986.

[95010]

[95011] 247. Wheeler, R. F.; Roberts, S. H.: An improved lymphocyte culture

[95012] technique: deoxycytidine release of a thymidine block and

use of a

[95013] constant humidity chamber for slide making. J. Med.

Genet. 24: 113–115,

[95014] 1987.

[95015]

[95016] 248. Willemsen, R.; Anar, B.; De Diego Otero, Y.; de Vries,

B. B.

[95017] A.; Hilhorst–Hofstee, Y.; Smits, A.; van Looveren, E.;

Willems, P.

[95018] J.; Galjaard, H.; Oostra, B. A.: Noninvasive test for fragile X syndrome,

[95019] using hair root analysis. Am. J. Hum. Genet. 65: 98–103, 1999.

[95020]

[95021] 249. Willemsen, R.; Mohkamsing, S.; De Vries, B.; Devys, D.; van den

[95022] Ouweland, A.; Mandel, J. L.; Galjaard, H.; Oostra, B.: Rapid antibody

[95023] test for fragile X syndrome. Lancet 345: 1147–1148, 1995.

[95024]

[95025] 250. Willemsen, R.; Smits, A.; Mohkamsing, S.; van

Beerendonk, H.;

[95026] de Haan, A.; de Vries, B.; van den Ouweland, A.; Sistermans, A.; Galjaard,

[95027] H.; Oostra, B. A.: Rapid antibody test for diagnosing fragile X syndrome:

[95028] a validation of the technique. Hum. Genet. 99: 308–311, 1997.

[95029]

[95030] 251. Winter, R. M.; Pembrey, M. E.: Analysis of linkage relationships

[95031] between genetic markers around the fragile X locus with special reference

[95032] to the daughters of normal transmitting males. Hum. Genet. 74: 93–97,

[95033] 1986.

[95034]

[95035] 252. Wohrle, D.; Hennig, I.; Vogel, W.; Steinbach, P.: Mitotic stability

[95036] of fragile X mutations in differentiated cells indicates early post-conceptual

[95037] trinucleotide repeat expansion. Nature Genet. 4: 140–142, 1993.

[95038]

[95039] 253. Wohrle, D.; Kotzot, D.; Hirst, M. C.; Manca, A.; Korn, B.; Schmidt,

[95040] A.; Barbi, G.; Rott, H.-D.; Poustka, A.; Davies, K. E.; Steinbach,

[95041] P.: A microdeletion of less than 250 kb, including the proximal part

[95042] of the FMR-1 gene and the fragile-X site, in a male with the clinical

[95043] phenotype of fragile-X syndrome. Am. J. Hum. Genet. 51: 299-306,

[95044] 1992.

[95045]

[95046] 254. Wohrle, D.; Salat, U.; Hameister, H.; Vogel, W.; Steinbach, P.

[95047] : Demethylation, reactivation, and destabilization of human fragile

[95048] X full-mutation alleles in mouse embryocarcinoma cells. Am. J. Hum.

[95049] Genet. 69: 504-515, 2001.

[95050]

- [95051] 255. Yamauchi, M.; Nagata, S.; Seki, N.; Toyama, Y.; Harada, N.; Niikawa,
[95052] N.; Masuno, I.; Kajii, T.; Hori, T.: Prenatal diagnosis of
fragile
[95053] X syndrome by direct detection of the dynamic mutation
due to an unstable
[95054] DNA sequence. Clin. Genet. 44: 169–172, 1993.
[95055]
[95056] 256. Yeshaya, J.; Shalgi, R.; Shohat, M.; Avivi, L.: Replica-
tion
[95057] timing of the various FMR1 alleles detected by FISH: infer-
ences regarding
[95058] their transcriptional status. Hum. Genet. 102: 6–14, 1998.
[95059]
[95060] 257. Yu, S.; Mulley, J.; Loesch, D.; Turner, G.; Donnelly, A.;
Gedeon,
[95061] A.; Hillen, D.; Kremer, E.; Lynch, M.; Pritchard, M.; Suther-
land,
[95062] G. R.; Richards, R. I.: Fragile-X syndrome: unique genetics
of the
[95063] heritable unstable element. Am. J. Hum. Genet. 50:
968–980, 1992.

[95064]

[95065] 258. Yu, S.; Pritchard, M.; Kremer, E.; Lynch, M.; Nancarrow, J.;

[95066] Baker, E.; Holman, K.; Mulley, J. C.; Warren, S. T.; Schlessinger,

[95067] D.; Sutherland, G. R.; Richards, R. I.: Fragile X genotype characterized

[95068] by an unstable region of DNA. Science 252: 1179–1181, 1991.

[95069]

[95070] 259. Zhang, Y. Q.; Bailey, A. M.; Matthies, H. J. G.; Renden, R. B.;

[95071] Smith, M. A.; Speese, S. D.; Rubin, G. M.; Broadie, K.: Drosophila

[95072] fragile X-related gene regulates the MAP1B homolog Futsch to control

[95073] synaptic structure and function. Cell 107: 591–603, 2001.

[95074]

[95075] 260. Zhong, N.; Dobkin, C.; Brown, W. T.: A complex mutable polymorphism

[95076] located within the fragile X gene. Nature Genet. 5: 248–253, 1993.

[95077]

[95078] 261. Zoll, B.; Arnemann, J.; Krawczak, M.; Cooper, D. N.;
Pescia,

[95079] G.; Wahli, W.; Steinbach, P.; Schmidtke, J.: Evidence against
close

[95080] linkage of the loci for fraXq of Martin-Bell syndrome and
for factor

[95081] IX. Hum. Genet. 71: 122-126, 1985.

[95082] 16549.

[95083] Bartley, T. D.; Bogenberger, J.; Hunt, P.; Li, Y.-S.; Lu, H. S.;

[95084] Martin, F.; Chang, M.-S.; Samal, B.; Nichol, J. L.; Swift, S.;
Johnson,

[95085] M. J.; Hsu, R.-Y.; and 41 others: Identification and cloning
of

[95086] a megakaryocyte growth and development factor that is a
ligand for

[95087] the cytokine receptor Mpl. Cell 77: 1117-1124, 1994.

[95088] 16550.

[95089] Chang, M.; McNinch, J.; Basu, R.; Shutter, J.; Hsu, R.;
Perkins,

[95090] C.; Mar, V.; Suggs, S.; Welcher, A.; Li, L.; Lu, H.; Bartley, T.;

- [95091] Hunt, P.; Martin, F.; Samal, B.; Bogenberger, J.: Cloning and characterization
- [95092] of the human megakaryocyte growth and development factor (MGDF) gene. J.
- [95093] Biol. Chem. 270: 511–514, 1995.
- [95094] 16551.
- [95095] Chang, M.-S.; Hsu, R.-Y.; McNinch, J.; Copeland, N. G.; Jenkins,
- [95096] N. A.: The gene for murine megakaryocyte growth and development factor
- [95097] (thrombopoietin, Thpo) is located on mouse chromosome 16. Genomics 26:
- [95098] 636–637, 1995.
- [95099] 16552.
- [95100] de Sauvage, F. J.; Hass, P. E.; Spencer, S. D.; Malloy, B. E.;
- [95101] Gurney, A. L.; Spencer, S. A.; Darbonne, W. C.; Henzel, W. J.; Wong,
- [95102] S. C.; Kuang, W.-J.; Oles, K. J.; Hultgren, B.; Solberg, L. A., Jr.;
- [95103] Goeddel, D. V.; Eaton, D. L.: Stimulation of megakaryocytopoiesis

- [95104] and thrombopoiesis by the c-Mpl ligand. *Nature* 369: 533–538, 1994.
- [95105] 16553.
- [95106] Farese, A. M.; Hunt, P.; Grab, L. B.; MacVittie, T. J.: Combined
- [95107] administration of recombinant human megakaryocyte growth and development
- [95108] factor and granulocyte colony-stimulating factor enhances multilineage
- [95109] hematopoietic reconstitution in nonhuman primates after radiation-induced
- [95110] marrow aplasia. *J. Clin. Invest.* 97: 2145–2151, 1996.
- [95111] 16554.
- [95112] Foster, D. C.; Sprecher, C. A.; Grant, F. J.; Kramer, J. M.; Kuijper,
- [95113] J. L.; Holly, R. D.; Whitmore, T. E.; Heipel, M. D.; Bell, L. A.;
- [95114] Ching, A. F. T.; McGrane, V.; Hart, C.; O'Hara, P. J.; Lok, S.: Human
- [95115] thrombopoietin: gene structure, cDNA sequence, expression, and chromosomal
- [95116] localization. *Proc. Nat. Acad. Sci.* 91: 13023–13027, 1994.

- [95117] 16555.
- [95118] Ghilardi, N.; Wiestner, A.; Kikuchi, M.; Ohsaka, A.; Skoda, R.
- [95119] C.: Hereditary thrombocythaemia in a Japanese family is caused by
- [95120] a novel point mutation in the thrombopoietin gene. *Brit. J. Haemat.* 107:
- [95121] 310–316, 1999.
- [95122] 16556.
- [95123] Kaushansky, K.; Lok, S.; Holly, R. D.; Broudy, V. C.; Lin, N.;
- [95124] Bailey, M. C.; Forstrom, J. W.; Buddle, M. M.; Oort, P. J.; Hagen,
- [95125] F. S.; Roth, G. J.; Papayannopoulou, T.; Foster, D. C.: Promotion
- [95126] of megakaryocyte progenitor expansion and differentiation by the c-Mpl
- [95127] ligand thrombopoietin. *Nature* 369: 568–571, 1994.
- [95128] 16557.
- [95129] Kondo, T.; Okabe, M.; Sanada, M.; Kurosawa, M.; Suzuki, S.; Kobayashi,
- [95130] M.; Hosokawa, M.; Asaka, M.: Familial essential thrombo-

cythemia associated

- [95131] with one-base deletion in the 5-prime-untranslated region of the thrombopoietin
- [95132] gene. Blood 92: 1091-1096, 1998.
- [95133] 16558.
- [95134] Li, J.; Yang, C.; Xia, Y.; Bertino, A.; Glaspy, J.; Roberts, M.;
- [95135] Kuter, D. J.: Thrombocytopenia caused by the development of antibodies
- [95136] to thrombopoietin. Blood 98: 3241-3248, 2001.
- [95137] 16559.
- [95138] Lok, S.; Kaushansky, K.; Holly, R. D.; Kuijper, J. L.; Lofton-Day,
- [95139] C. E.; Oort, P. J.; Grant, F. J.; Heipel, M. D.; Burkhead, S. K.;
- [95140] Kramer, J. M.; Bell, L. A.; Sprecher, C. A.; Blumberg, H.; Johnson,
- [95141] R.; Prunkard, D.; Ching, A. F. T.; Mathewes, S. L.; Bailey, M. C.;
- [95142] Forstrom, J. W.; Buddle, M. M.; Osborn, S. G.; Evans, S. J.; Sheppard,
- [95143] P. O.; Presnell, S. R.; O'Hara, P. J.; Hagen, F. S.; Roth, G. J.;
- [95144] Foster, D. C.: Cloning and expression of murine throm-

bopoietin cDNA

[95145] and stimulation of platelet production in vivo. Nature 369:
565–568,

[95146] 1994.

[95147] 16560.

[95148] Metcalf, D.: Thrombopoietin: at last. Nature 369:
519–520,

[95149] 1994.

[95150] 16561.

[95151] Methia, N.; Louache, F.; Vainchenker, W.; Wendling, F.:
Oligodeoxynucleotides

[95152] antisense to the proto-oncogene c-mpl specifically inhibit
in vitro

[95153] megakaryocytopoiesis. Blood 82: 1395–1401, 1993.

[95154] 16562.

[95155] Ratajczak, M. Z.; Ratajczak, J.; Marlicz, W.; Pletcher, C. H.,

[95156] Jr.; Machalinski, B.; Moore, J.; Hung, H.; Gewirtz, A. M.:

Recombinant

[95157] human thrombopoietin (TPO) stimulates erythropoiesis by
inhibiting

[95158] erythroid progenitor cell apoptosis. Brit. J. Haemat. 98:

8-17,

[95159] 1997.

[95160] 16563.

[95161] Sohma, Y.; Akahori, H.; Seki, N.; Hori, T.; Ogami, K.; Kato, T.;

[95162] Shimada, Y.; Kawamura, K.; Miyazaki, H.: Molecular cloning and chromosomal

[95163] localization of the human thrombopoietin gene. FEBS Lett. 353: 57-61,

[95164] 1994.

[95165] 16564.

[95166] Wendling, F.; Maraskovsky, E.; Debili, N.; Florindo, C.; Teepe,

[95167] M.; Titeux, M.; Methia, N.; Breton-Gorius, J.; Cosman, D.; Vainchenker,

[95168] W.: c-Mpl ligand is a humoral regulator of megakaryocytopoiesis. Nature 369:

[95169] 571-574, 1994.

[95170] 16565.

[95171] Dualan, R.; Brody, T.; Keeney, S.; Nichols, A. F.; Admon, A.; Linn,

- [95172] S.: Chromosomal localization and cDNA cloning of the genes (DDB1
- [95173] and DDB2) for the p127 and p48 subunits of a human damage-specific
- [95174] DNA binding protein. *Genomics* 29: 62–69, 1995.
- [95175] 16566.
- [95176] Koken, M. H. M.; Reynolds, P.; Jaspers–Dekker, I.; Prakash, L.;
- [95177] Prakash, S.; Bootsma, D.; Hoeijmakers, J. H. J.: Structural and functional
- [95178] conservation of two human homologs of the yeast DNA repair gene RAD6. *Proc.*
- [95179] *Nat. Acad. Sci.* 88: 8865–8869, 1991.
- [95180] 16567.
- [95181] Knoll, A.; Schunkert, H.; Reichwald, K.; Danser, A. H. J.; Bauer,
- [95182] D.; Platzer, M.; Stein, G.; Rosenthal, A.: Human renin binding protein:
- [95183] complete genomic sequence and association of an intronic T/C polymorphism
- [95184] with the prorenin level in males. *Hum. Molec. Genet.* 6: 1527–1534,

- [95185] 1997.
- [95186] 16568.
- [95187] Laan, M.; Paabo, S.: Mapping genes by drift-generated linkage disequilibrium. (Letter) Am. J. Hum. Genet. 63: 654-656, 1998.
- [95189] 16569.
- [95190] Takahashi, S.; Inoue, H.; Miyake, Y.: The human gene for renin-binding protein. J. Biol. Chem. 267: 13007-13013, 1992.
- [95192] 16570.
- [95193] Terwilliger, J. D.; Zollner, S.; Laan, M.; Paabo, S.: Mapping genes through the use of linkage disequilibrium generated by genetic drift: 'drift mapping' in small populations with no demographic expansion. Hum. Hered. 48: 138-154, 1998.
- [95197] 16571.
- [95198] van den Ouweland, A. M. W.; Verdijk, M.; Kiochis, P.; Poustka,

- [95199] A.; van Oost, B. A.: The human renin-binding protein gene (RENBP)
- [95200] maps in Xq28. *Genomics* 21: 279–281, 1994.
- [95201] 16572.
- [95202] Warren, S. T.; Knight, S. J. L.; Peters, J. F.; Stayton, C. L.;
- [95203] Consalez, G. G.; Zhang, F.: Isolation of the human chromosomal band
- [95204] Xq28 within somatic cell hybrids by fragile X site breakage. *Proc.*
- [95205] *Nat. Acad. Sci.* 87: 3856–3860, 1990.
- [95206] 16573.
- [95207] Calvo, R. M.; Asuncion, M.; Sancho, J.; San Millan, J. L.; Escobar-Morreale,
- [95208] H. F.: The role of the CAG repeat polymorphism in the androgen receptor
- [95209] gene and of skewed X-chromosome inactivation, in the pathogenesis
- [95210] of hirsutism. *J. Clin. Endocr. Metab.* 85: 1735–1740, 2000.
- [95211] 16574.
- [95212] Sinclair, A. H.; Foster, J. W.; Spencer, J. A.; Page, D. C.; Palmer,

- [95213] M.; Goodfellow, P. N.; Marshall Graves, J. A.: Sequences homologous
- [95214] to ZFY, a candidate human sex-determining gene, are autosomal in marsupials.
- [95215] Nature 336: 780–783, 1988.
- [95216] 16575.
- [95217] Gao, X.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Gridley,
- [95218] T.: Assignment of the murine Notch2 and Notch3 genes to chromosomes
- [95219] 3 and 17. Genomics 49: 160–161, 1998.
- [95220] 16576.
- [95221] Larsson, C.; Lardelli, M.; White, I.; Lendahl, U.: The human NOTCH1,
- [95222] 2, and 3 genes are located at chromosome positions 9q34, 1p13–p11,
- [95223] and 19p13.2–p13.1 in regions of neoplasia-associated translocation. Genomics 24:
- [95224] 253–258, 1994.
- [95225] 16577.
- [95226] Dichgans, M.; Herzog, J.; Gasser, T.: NOTCH3 mutation in–

volving

[95227] three cysteine residues in a family with typical CADASIL.

Neurology 57:

[95228] 1714–1717, 2001.

[95229] 16578.

[95230] Joutel, A.; Corpechot, C.; Ducros, A.; Vahedi, K.; Chabriat, H.;

[95231] Mouton, P.; Alamowitch, S.; Domenga, V.; Cecillion, M.; Marechal,

[95232] E.; Maciazek, J.; Vayssiere, C.; Cruaud, C.; Cabanis, E.–A.; Ruchoux,

[95233] M. M.; Weissenbach, J.; Bach, J. F.; Bousser, M. G.; Tournier–Lasserre,

[95234] E.: Notch3 mutations in CADASIL, a hereditary adult-onset condition

[95235] causing stroke and dementia. Nature 383: 707–710, 1996.

[95236] 16579.

[95237] Joutel, A.; Dodick, D. D.; Parisi, J. E.; Cecillon, M.; Tournier–Lasserre,

[95238] E.; Bousser, M. G.: De novo mutation in the Notch3 gene causing CADASIL. Ann.

- [95239] *Neurol.* 47: 388–391, 2000.
- [95240] 16580.
- [95241] Joutel, A.; Vahedi, K.; Corpechot, C.; Troesch, A.; Chabriat, H.;
- [95242] Vayssiere, C.; Cruaud, C.; Maciazek, J.; Weissenbach, J.; Bousser, M.-G.; Bach, J.-F.; Tournier-Lasserre, E.: Strong clustering and stereotyped nature of Notch3 mutations in CADASIL patients. *Lancet* 350: 1511–1515, 1997.
- [95243] 16581.
- [95244] Rebay, I.; Fehon, R. G.; Artavanis-Tsakonas, S.: Specific truncations of *Drosophila* Notch define dominant activated and dominant negative forms of the receptor. *Cell* 74: 319–329, 1993.
- [95245] 16582.
- [95246] Deed, R. W.; Hirose, T.; Mitchell, E. L. D.; Santibanez-Koref, M. F.; Norton, J. D.: Structural organisation and chromo-

somal mapping

[95253] of the human Id-3 gene. Gene 151: 309-314, 1994.

[95254] 16583.

[95255] Ellmeier, W.; Aguzzi, A.; Kleiner, E.; Kurzbauer, R.; Weith, A.

[95256] : Mutually exclusive expression of a helix-loop-helix gene and N-myc

[95257] in human neuroblastomas and in normal development. EMBO J. 11: 2563-2571,

[95258] 1992.

[95259] 16584.

[95260] Kee, B. L.; Rivera, R. R.; Murre, C.: Id3 inhibits B lymphocyte

[95261] progenitor growth and survival in response to TGF-beta. Nature Immun. 2:

[95262] 242-247, 2001.

[95263] 16585.

[95264] Lyden, D.; Young, A. Z.; Zagzag, D.; Yan, W.; Gerald, W.; O'Reilly,

[95265] R.; Bader, B. L.; Hynes, R. O.; Zhuang, Y.; Manova, K.; Benezra, R.

- [95266] : Id1 and Id3 are required for neurogenesis, angiogenesis and vascularization
- [95267] of tumour xenografts. *Nature* 401: 670–677, 1999.
- [95268] 16586.
- [95269] Pan, L.; Sato, S.; Frederick, J. P.; Sun, X.–H.; Zhuang, Y.: Impaired
- [95270] immune responses and B–cell proliferation in mice lacking the Id3
- [95271] gene. *Molec. Cell. Biol.* 19: 5969–5980, 1999.
- [95272] 16587.
- [95273] Yeh, K.; Lim, R. W.: Genomic organization and promoter analysis
- [95274] of the murine Id3 gene. *Gene* 254: 163–171, 2000.
- [95275] 16588.
- [95276] Rubinfeld, B.; Munemitsu, S.; Clark, R.; Conroy, L.; Watt, K.;
- [95277] Crosier, W. J.; McCormick, F.; Polakis, P.: Molecular cloning of
- [95278] a GTPase activating protein specific for the Krev–1 protein p21–rap1. *Cell* 65:
- [95279] 1033–1042, 1991.

[95280] 16589.

[95281] Weiss, J.; Rubinfeld, B.; Polakis, P. G.; McCormick, F.;
Cavenee,

[95282] W. K.; Arden, K. C.: The RAP1GA1 locus for human
Rap1-GTPase activating

[95283] protein 1 maps to chromosome 1p36.1-p35. Cytogenet.
Cell Genet. 66:

[95284] 18-21, 1994.

[95285] 16590.

[95286] Braun, A.; Kammerer, S.; Weissenhorn, W.; Weiss, E. H.;
Cleve,

[95287] H.: Sequence of a putative human housekeeping gene
(HK33) localized

[95288] on chromosome 1. Gene 146: 291-295, 1994.

[95289] 16591.

[95290] Gotte, K.; Girzalsky, W.; Linkert, M.; Baumgart, E.; Kam-
merer,

[95291] S.; Kunau, W.-H.; Erdmann, R.: Pex19p, a farnesylated
protein essential

[95292] for peroxisome biogenesis. Molec. Cell. Biol. 18: 616-628,
1998.

- [95293] 16592.
- [95294] James, G. L.; Goldstein, J. L.; Pathak, R. K.; Anderson, R. G.
- [95295] W.; Brown, M. S.: PxF, a prenylated protein of peroxisomes
J. Biol.
- [95296] Chem. 269: 14182–14190, 1994.
- [95297] 16593.
- [95298] Kammerer, S.; Arnold, N.; Gutensohn, W.; Mewes, H.–W.;
Kunau, W.–H.;
- [95299] Hofler, G.; Roscher, A. A.; Braun, A.: Genomic organization
and molecular
- [95300] characterization of a gene encoding HsPXF, a human per-
oxisomal farnesylated
- [95301] protein. Genomics 45: 200–210, 1997.
- [95302] 16594.
- [95303] Caskey, C. T.; Pizzuti, A.; Fu, Y.–H.; Fenwick, R. G., Jr.;
Nelson,
- [95304] D. L.: Triplet repeat mutations in human disease. Science
256:
- [95305] 784–789, 1992.
- [95306] 16595.

- [95307] Chang, B.; Zheng, S. L.; Hawkins, G. A.; Isaacs, S. D.; Wiley, K. E.; Turner, A.; Carpten, J. D.; Bleecker, E. R.; Walsh, P. C.;
- [95309] Trent, J. M.; Meyers, D. A.; Isaacs, W. B.; Xu, J.: Polymorphic GGC repeats in the androgen receptor gene are associated with hereditary and sporadic prostate cancer risk. Hum. Genet. 110: 122–129, 2002.
- [95312] 16596.
- [95313] Chang, C.; Kokontis, J.; Liao, S.: Molecular cloning of human and rat complementary DNA encoding androgen receptors. Science 240: 324–326, 1988.
- [95316] 16597.
- [95317] Choong, C. S.; Kempainen, J. A.; Zhou, Z.-X.; Wilson, E. M.:
- [95318] Reduced androgen receptor gene expression with first exon CAG repeat expansion. Molec. Endocr. 10: 1527–1535, 1996.

- [95320] 16598.
- [95321] Choong, C. S.; Quigley, C. A.; French, F. S.; Wilson, E. M.:
- [95322] A novel missense mutation in the amino-terminal domain
of the human
- [95323] androgen receptor gene in a family with partial androgen
insensitivity
- [95324] syndrome causes reduced efficiency of protein translation.
J. Clin.
- [95325] Invest. 98: 1423-1431, 1996.
- [95326] 16599.
- [95327] Chu, J.; Zhang, R.; Zhao, Z.; Zou, W.; Han, Y.; Qi, Q.;
Zhang,
- [95328] H.; Wang, J.-C.; Tao, S.; Liu, X.; Luo, Z.: Male fertility is
compatible
- [95329] with an Arg840Cys substitution in the AR in a large Chi-
nese family
- [95330] with divergent phenotypes of AR insensitivity syndrome. J.
Clin.
- [95331] Endocr. Metab. 87: 347-351, 2002.
- [95332] 16600.
- [95333] Coetzee, G. A.; Ross, R. K.: Re: Prostate cancer and the

androgen

[95334] receptor. (Letter) J. Nat. Cancer Inst. 86: 872–873, 1994.

[95335] 16601.

[95336] Correa–Cerro, L.; Wohr, G.; Haussler, J.; Berthon, P.;
Drelon,

[95337] E.; Mangin, P.; Fournier, G.; Cussenot, O.; Kraus, P.; Just,
W.; Paiss,

[95338] T.; Cantu, J. M.; Vogel, W.: (CAG)nCAA and GGN repeats in
the human

[95339] androgen receptor gene are not associated with prostate
cancer in

[95340] a French–German population. Europ. J. Hum. Genet. 7:
347–362, 1999.

[95341] 16602.

[95342] Dowsing, A. T.; Yong, E. L.; Clark, M.; McLachlan, R. I.; de
Kretser,

[95343] D. M.; Trounson, A. O.: Linkage between male infertility
and trinucleotide

[95344] repeat expansion in the androgen–receptor gene. Lancet
354: 640–643,

[95345] 1999.

[95346] 16603.

- [95347] Elo, J. P.; Kvist, L.; Leinonen, K.; Isomaa, V.; Hentuu, P.;
Lukkarinen,
- [95348] O.; Vihko, P.: Mutated human androgen receptor gene de-
tected in a
- [95349] prostatic cancer patient is also activated by estradiol. J.
Clin.
- [95350] Endocr. Metab. 80: 3494–3500, 1995.
- [95351] 16604.
- [95352] Gaddipati, J. P.; McLeod, D. G.; Heidenberg, H. B.; Sester-
henn,
- [95353] I. A.; Finger, M. J.; Moul, J. W.; Srivastava, S.: Frequent de-
tection
- [95354] of codon 877 mutation in the androgen receptor gene in
advanced prostate
- [95355] cancers. Cancer Res. 54: 2861–2864, 1994.
- [95356] 16605.
- [95357] Gehring, U.; Tomkins, G. M.: Characterization of a hor-
mone receptor
- [95358] defect in the androgen–insensitivity mutant. Cell 3:
59–64, 1974.
- [95359] 16606.

- [95360] Gingrich, J. R.; Barrios, R. J.; Kattan, M. W.; Nahm, H. S.; Finegold,
[95361] M. J.; Greenberg, N. M.: Androgen-independent prostate
cancer progression
[95362] in the TRAMP model. *Cancer Res.* 57: 4687–4691, 1997.
[95363] 16607.
- [95364] Gingrich, J. R.; Greenberg, N. M.: A transgenic mouse
prostate
[95365] cancer model. *Toxicol. Path.* 24: 502–504, 1996.
[95366] 16608.
- [95367] Giovannucci, E.; Stampfer, M. J.; Krithivas, K.; Brown, M.; Brufsky,
[95368] A.; Talcott, J.; Hennekens, C. H.; Kantoff, P. W.: The CAG
repeat
[95369] within the androgen receptor gene and its relationship to
prostate
[95370] cancer. *Proc. Nat. Acad. Sci.* 94: 3320–3323, 1997.
[95371] 16609.
- [95372] Gottlieb, B.; Beitel, L. K.; Trifiro, M. A.: Variable expressiv-
ity
[95373] and mutation databases: the androgen receptor gene mu-

tations database. Hum.

[95374] **Mutat. 17: 382–388, 2001.**

[95375] **16610.**

[95376] **Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Pinsky, L.: The an-
drogen**

[95377] **receptor gene mutations database. Nucleic Acids Res. 25:
158–162,**

[95378] **1997.**

[95379] **16611.**

[95380] **Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Vasiliou, D. M.; Pin-
sky,**

[95381] **L.: The androgen receptor gene mutations database. Nu-
cleic Acids**

[95382] **Res. 24: 151–154, 1996.**

[95383] **16612.**

[95384] **Griffin, J. E.: Androgen resistance--the clinical and molec-
ular**

[95385] **spectrum. New Eng. J. Med. 326: 611–618, 1992.**

[95386] **16613.**

[95387] **Grino, P. B.; Isidro-Gutierrez, R. F.; Griffin, J. E.; Wilson,**

- [95388] J. D.: Androgen resistance associated with a qualitative abnormality
- [95389] of the androgen receptor and responsive to high dose androgen therapy. J.
- [95390] Clin. Endocr. Metab. 68: 578–584, 1989.
- [95391] 16614.
- [95392] Hardy, D. O.; Scher, H. I.; Bogenreider, T.; Sabbatini, P.; Zhang,
- [95393] Z.-F.; Nanus, D. M.; Catterall, J. F.: Androgen receptor CAG repeat
- [95394] lengths in prostate cancer: correlation with age of onset. J. Clin.
- [95395] Endocr. Metab. 81: 4400–4405, 1996.
- [95396] 16615.
- [95397] Hellwinkel, O. J.-C.; Holterhus, P.-M.; Struve, D.; Marschke,
- [95398] C.; Homburg, N.; Hiort, O.: A unique exonic splicing mutation in
- [95399] the human androgen receptor gene indicates a physiologic relevance
- [95400] of regular androgen receptor transcript variants. J. Clin. Endocr.

- [95401] Metab. 86: 2569–2575, 2001.
- [95402] 16616.
- [95403] Hickey, T.; Chandy, A.; Norman, R. J.: The androgen receptor
- [95404] CAG repeat polymorphism and X-chromosome inactivation in Australian
- [95405] Caucasian women with infertility related to polycystic ovary syndrome. J.
- [95406] Clin. Endocr. Metab. 87: 161–165, 2002.
- [95407] 16617.
- [95408] Hiort, O.; Sinnecker, G. H. G.; Holterhus, P.–M.; Nitsche, E.
- [95409] M.; Kruse, K.: Inherited and de novo androgen receptor gene mutations:
- [95410] investigation of single-case families. J. Pediat. 132: 939–943,
- [95411] 1998.
- [95412] 16618.
- [95413] Holterhus, P.–M.; Bruggenwirth, H. T.; Hiort, O.; Kleinkauf–Houcken,
- [95414] A.; Kruse, K.; Sinnecker, G. H. G.; Brinkmann, A. O.: Mo-
saicism due

- [95415] to a somatic mutation of the androgen receptor gene determines phenotype
- [95416] in androgen insensitivity syndrome. *J. Clin. Endocr. Metab.* 82:
- [95417] 3584–3589, 1997.
- [95418] 16619.
- [95419] Shibuya, H.; Yamaguchi, K.; Shirakabe, K.; Tonegawa, A.; Gotoh,
- [95420] Y.; Ueno, N.; Irie, K.; Nishida, E.; Matsumoto, K.: TAB1: an activator
- [95421] of the TAK1 MAPKKK in TGF- β signal transduction. *Science* 272:
- [95422] 1179–1182, 1996.
- [95423] 16620.
- [95424] Chadwick, B. P.; Obermayr, F.; Frischauf, A.-M.: FKHL15, a new
- [95425] human member of the forkhead gene family located on chromosome 9q22. *Genomics* 41:
- [95426] 390–396, 1997.
- [95427] 16621.
- [95428] De Felice, M.; Ovitt, C.; Biffali, E.; Rodriguez-Mallon, A.;

Arra,

[95429] C.; Anastassiadis, K.; Macchia, P. E.; Mattei, M.-G.; Mariano, A.;

[95430] Scholer, H.; Macchia, V.; Di Lauro, R.: A mouse model for hereditary

[95431] thyroid dysgenesis and cleft palate. *Nature Genet.* 19: 395–398,

[95432] 1998.

[95433] 16622.

[95434] Kaufmann, E.; Knochel, W.: Five years on the wings of fork head. *Mech.*

[95435] *Dev.* 57: 3–20, 1996.

[95436] 16623.

[95437] Manley, N. R.; Capecchi, M. R.: Hox group 3 paralogs regulate

[95438] the development and migration of the thymus, thyroid, and parathyroid

[95439] gland. *Dev. Biol.* 195: 1–15, 1998.

[95440] 16624.

[95441] Toubanc, J. E.: Comparison of epidemiological data on congenital

- [95442] hypothyroidism in Europe with those of other parts in the world. Horm.
- [95443] Res. 38: 230–235, 1992.
- [95444] 16625.
- [95445] Zannini, M.; Avantaggiato, V.; Biffali, E.; Arnone, M. I.; Sato,
- [95446] K.; Pischetola, M.; Taylor, B. A.; Phillips, S. J.; Simeone, A.; Di
- [95447] Lauro, R.: TTF–2, a new forkhead protein, shows a temporal expression
- [95448] in the developing thyroid which is consistent with a role in controlling
- [95449] the onset of differentiation. EMBO J. 16: 3185–3197, 1997.
- [95450] 16626.
- [95451] Boyd, J. M.; Subramanian, T.; Schaeper, U.; La Regina, M.; Bayley,
- [95452] S.; Chinnadurai, G.: A region in the C–terminus of adenovirus 2/5
- [95453] E1a protein is required for association with a cellular phosphoprotein
- [95454] and important for the negative modulation of T24–ras

mediated transformation,

[95455] tumorigenesis and metastasis. EMBO J. 469–478, 1993.

[95456] 16627.

[95457] Furusawa, T.; Moribe, H.; Kondoh, H.; Higashi, Y.: Identification

[95458] of CtBP1 and CtBP2 as corepressors of zinc finger-homeodomain factor

[95459] delta-EF1. Molec. Cell. Biol. 19: 8581–8590, 1999.

[95460] 16628.

[95461] Katsanis, N.; Fisher, E. M. C.: A novel C-terminal binding protein

[95462] (CTBP2) is closely related to CTBP1, an adenovirus E1A-binding protein,

[95463] and maps to human chromosome 21q21.3. Genomics 47: 294–299, 1998.

[95464] 16629.

[95465] Schaeper, U.; Boyd, J. M.; Verma, S.; Uhlmann, E.; Subramanian,

[95466] T.; Chinnadurai, G.: Molecular cloning and characterization of a

[95467] cellular phosphoprotein that interacts with a conserved C-

terminal

- [95468] domain of adenovirus E1A involved in negative modulation of oncogenic transformation. Proc. Nat. Acad. Sci. 92: 10467-10471, 1995.
- [95469] 16630.
- [95470] Sewalt, R. G. A. B.; Gunster, M. J.; van der Vlag, J.; Satijn, D. P. E.; Otte, A. P.: C-terminal binding protein is a transcriptional repressor that interacts with a specific class of vertebrate polycomb proteins. Molec. Cell. Biol. 19: 777-787, 1999.
- [95471] 16631.
- [95472] Zhang, Q.; Piston, D. W.; Goodman, R. H.: Regulation of corepressor function by nuclear NADH. Science 295: 1895-1897, 2002.
- [95473] 16632.
- [95474] Aya, K.; Tanaka, H.; Seino, Y.: Novel mutation in the nephrin gene of a Japanese patient with congenital nephrotic syn-

drome of the

[95481] Finnish type. *Kidney Int.* 57: 401–404, 2000.

[95482] 16633.

[95483] Beltcheva, O.; Martin, P.; Lenkkeri, U.; Tryggvason, K.: Mutation

[95484] spectrum in the nephrin gene (NPHS1) in congenital nephrotic syndrome. *Hum.*

[95485] *Mutat.* 17: 368–373, 2001.

[95486] 16634.

[95487] Guez, S.; Giani, M.; Melzi, M. L.; Antignac, C.; Assael, B. M.

[95488] : Adequate clinical control of congenital nephrotic syndrome by enalapril. *Pediat.*

[95489] *Nephrol.* 12: 130–132, 1998.

[95490] 16635.

[95491] Koziell, A.; Grech, V.; Hussain, S.; Lee, G.; Lenkkeri, U.; Tryggvason,

[95492] K.; Scambler, P.: Genotype/phenotype correlations of NPHS1 and NPHS2

[95493] mutations in nephrotic syndrome advocate a functional inter-relationship

[95494] in glomerular filtration. *Hum. Molec. Genet.* 11: 379–388,

2002.

[95495] 16636.

[95496] Lenkkeri, U.; Mannikko, M.; McCready, P.; Lamerdin, J.;
Gribouval,

[95497] O.; Niaudet, P.; Antignac, C.; Kashtan, C. E.; Holmberg, C.;
Olsen,

[95498] A.; Kestila, M.; Tryggvason, K.: Structure of the gene for
congenital

[95499] nephrotic syndrome of the Finnish type (NPHS1) and char-
acterization

[95500] of mutations. Am. J. Hum. Genet. 64: 51-61, 1999.

[95501] 16637.

[95502] Liu, L.; Done, S. C.; Khoshnoodi, J.; Bertorello, A.; Wartio-
vaara,

[95503] J.; Berggren, P.-O.; Tryggvason, K.: Defective nephrin traf-
ficking

[95504] caused by missense mutations in the NPHS1 gene: insight
into the mechanisms

[95505] of congenital nephrotic syndrome. Hum. Molec. Genet. 10:
2637-2644,

[95506] 2001.

[95507] 16638.

- [95508] Putaala, H.; Soininen, R.; Kilpelainen, P.; Wartiovaara, J.; Tryggvason,
K.: The murine nephrin gene is specifically expressed in
kidney,
brain and pancreas: inactivation of the gene leads to mas-
sive proteinuria
and neonatal death. Hum. Molec. Genet. 10: 1-8, 2001.
16639.
- [95513] Ruotsalainen, V.; Ljungberg, P.; Wartiovaara, J.; Lenkkeri,
U.;
Kestila, M.; Jalanko, H.; Holmberg, C.; Tryggvason, K.:
Nephrin is
specifically located at the slit diaphragm of glomerular
podocytes. Proc.
Nat. Acad. Sci. 96: 7962-7967, 1999.
16640.
- [95518] Watanabe, T.; Inoue, S.; Hiroi, H.; Orimo, A.; Kawashima,
H.; Muramatsu,
M.: Isolation of estrogen-responsive genes with a CpG is-
land library. Molec.
Cell. Biol. 18: 442-449, 1998.

- [95521] 16641.
- [95522] Adra, C. N.; Kobayashi, H.; Rowley, J. D.; Lim, B.: Assign-
ment
- [95523] of the human GDID4 gene, a GDP/GTP-exchange regula-
tor, to chromosome
- [95524] 12p12.3. Genomics 24: 188-190, 1994.
- [95525] 16642.
- [95526] Leffers, H.; Nielsen, M. S.; Andersen, A. H.; Honore, B.;
Madsen,
- [95527] P.; Vandekerckhove, J.; Celis, J. E.: Identification of two
human
- [95528] Rho GDP dissociation inhibitor proteins whose overex-
pression leads
- [95529] to disruption of the actin cytoskeleton. Exp. Cell Res. 209:
165-174,
- [95530] 1993.
- [95531] 16643.
- [95532] Lelias, J.-M.; Adra, C. N.; Wulf, G. M.; Guillemot, J.-C.;
Khagad,
- [95533] M.; Caput, D.; Lim, B.: cDNA cloning of a human mRNA
preferentially

- [95534] expressed in hematopoietic cells and with homology to a GDP-dissociation
- [95535] inhibitor for the rho GTP-binding proteins. Proc. Nat. Acad. Sci. 90:
- [95536] 1479-1483, 1993.
- [95537] 16644.
- [95538] Scherle, P.; Behrens, T.; Staudt, L. M.: Ly-GDI, a GDP-dissociation
- [95539] inhibitor of the RhoA GTP-binding protein, is expressed preferentially
- [95540] in lymphocytes. Proc. Nat. Acad. Sci. 90: 7568-7572, 1993.
- [95541] 16645.
- [95542] Telford, E. A. R.; Wightman, P.; Leek, J.; Markham, A. F.; Lench,
- [95543] N. J.; Bonthron, D. T.: cDNA cloning, genomic organization, and chromosomal
- [95544] localization of a novel human gene that encodes a kinesin-related
- [95545] protein highly similar to mouse Kif3C. Biochem. Biophys. Res. Commun. 242:
- [95546] 407-412, 1998.

- [95547] 16646.
- [95548] Kalchman, M. A.; Graham, R. K.; Xia, G.; Koide, H. B.;
Hodgson,
- [95549] J. G.; Graham, K. C.; Goldberg, Y. P.; Gietz, R. D.; Pickart,
C. M.;
- [95550] Hayden, M. R.: Huntingtin is ubiquitinated and interacts
with a specific
- [95551] ubiquitin-conjugating enzyme. J. Biol. Chem. 271:
19385-19394, 1996.
- [95552] 16647.
- [95553] Chiariotti, L.; Benvenuto, G.; Fedele, M.; Santoro, M.;
Simeone,
- [95554] A.; Fusco, A.; Bruni, C. B.: Identification and characteriza-
tion
- [95555] of a novel RING-finger gene (RNF4) mapping at 4p16.3.
Genomics 47:
- [95556] 258-265, 1998.
- [95557] 16648.
- [95558] Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.;
Tanaka, A.;
- [95559] Kotani, H.; Nomura, N.; Ohara, O.: Prediction of coding
sequences

- [95560] of unidentified human genes. X. The complete sequences of 100 new
- [95561] cDNA clones from brain which can code for large proteins in vitro. DNA
- [95562] Res. 5: 169–176, 1998.
- [95563] 16649.
- [95564] Waldegger, S.; Erdel, M.; Nagl, U. O.; Barth, P.; Raber, G.; Steuer,
- [95565] S.; Utermann, G.; Paulmichl, M.; Lang, F.: Genomic organization and
- [95566] chromosomal localization of the human SGK protein kinase gene. Genomics 51:
- [95567] 299–302, 1998.
- [95568] 16650.
- [95569] Webster, M. K.; Goya, L.; Ge, Y.; Maiyar, A. C.; Firestone, G.
- [95570] L.: Characterization of sgk, a novel member of the serine/threonine
- [95571] protein kinase gene family which is transcriptionally induced by glucocorticoids
- [95572] and serum. Molec. Cell. Biol. 13: 2031–2040, 1993.

- [95573] 16651.
- [95574] Aoki, K.; Ishida, R.; Kasai, M.: Isolation and characterization
of a cDNA encoding a translin-like protein, TRAX. FEBS
Lett. 401:
109–112, 1997.
- [95577] 16652.
- [95578] Meng, G.; Aoki, K.; Tokura, K.; Nakahara, K.; Inazawa, J.;
Kasai,
M.: Genomic structure and chromosomal localization of
the gene encoding
TRAX, a translin-associated factor X. J. Hum. Genet. 45:
305–308,
2000.
- [95582] 16653.
- [95583] Collins, C.; Rommens, J. M.; Kowbel, D.; Godfrey, T.; Tan-
ner, M.;
- [95584] Hwang, S.; Polikoff, D.; Nonet, G.; Cochran, J.; Myambo,
K.; Jay,
K. E.; Froula, J.; Cloutier, T.; Kuo, W.-L.; Yaswen, P.; Dair-
kee,

- [95586] S.; Giovanola, J.; Hutchinson, G. B.; Isola, J.; Kallioniemi, O.-P.;
- [95587] Palazzolo, M.; Martin, C.; Ericsson, C.; Pinkel, D.; Albertson, D.;
- [95588] Li, W.-B.; Gray, J. W.: Positional cloning of ZNF217 and NABC1: genes
- [95589] amplified at 20q13.2 and overexpressed in breast carcinoma. Proc.
- [95590] Nat. Acad. Sci. 95: 8703-8708, 1998.
- [95591] 16654.
- [95592] Eudy, J. D.; Yao, S.; Weston, M. D.; Ma-Edmonds, M.; Talmadge,
- [95593] C. B.; Cheng, J. J.; Kimberling, W. J.; Sumegi, J.: Isolation of
- [95594] a gene encoding a novel member of the nuclear receptor superfamily
- [95595] from the critical region of Usher syndrome type IIa at 1q41. Genomics 50:
- [95596] 382-384, 1998.
- [95597] 16655.
- [95598] Greschik, H.; Wurtz, J.-M.; Sanglier, S.; Bourguet, W.; van Dorsselaer,

- [95599] A.; Moras, D.; Renaud, J.-P.: Structural and functional evidence
- [95600] for ligand-independent transcriptional activation by the estrogen-related
- [95601] receptor 3. *Molec. Cell* 9: 303-313, 2002.
- [95602] 16656.
- [95603] Matsuyoshi, N.; Imamura, S.: Multiple cadherins are expressed
- [95604] in human fibroblasts. *Biochem. Biophys. Res. Commun.* 235: 355-358,
- [95605] 1997.
- [95606] 16657.
- [95607] Bertilsson, G.; Heidrich, J.; Svensson, K.; Asman, M.; Jendeberg,
- [95608] L.; Sydow-Backman, M.; Ohlsson, R.; Postlind, H.; Blomquist, P.; Berkenstam,
- [95609] A.: Identification of a human nuclear receptor defines a new signaling
- [95610] pathway for CYP3A induction. *Proc. Nat. Acad. Sci.* 95: 12208-12213,
- [95611] 1998.

- [95612] 16658.
- [95613] Blumberg, B.; Sabbagh, W., Jr.; Juguilon, H.; Bolado, J., Jr.;
- [95614] van Meter, C. M.; Ong, E. S.; Evans, R. M.: SXR, a novel
steroid
- [95615] and xenobiotic-sensing nuclear receptor. *Genes Dev.* 12:
3195-3205,
- [95616] 1998.
- [95617] 16659.
- [95618] Watkins, R. E.; Wisely, G. B.; Moore, L. B.; Collins, J. L.;
- Lambert,
- [95619] M. H.; Williams, S. P.; Willson, T. M.; Kliewer, S. A.; Red-
inbo, M.
- [95620] R.: The human nuclear xenobiotic receptor PXR: structural
determinants
- [95621] of directed promiscuity. *Science* 292: 2329-2333, 2001.
- [95622] 16660.
- [95623] Nemes, J. P.; Benzow, K. A.; Koob, M. D.: The SCA8 tran-
script
- [95624] is an antisense RNA to a brain-specific transcript encod-
ing a novel
- [95625] actin-binding protein (KLHL1). *Hum. Molec. Genet.* 9:

1543–1551,

[95626] 2000.

[95627] 16661.

[95628] Adato, A.; Raskin, L.; Petit, C.; Bonne–Tamir, B.: Deafness heterogeneity

[95629] in a Druze isolate from the Middle East: novel OTOF and PDS mutations,

[95630] low prevalence of GJB2 35delG mutation and indication for a new DFNB

[95631] locus. *Europ. J. Hum. Genet.* 8: 437–442, 2000.

[95632] 16662.

[95633] Yasunaga, S.; Grati, M.; Chardenoux, S.; Smith, T. N.; Friedman,

[95634] T. B.; Lalwani, A. K.; Wilcox, E. R.; Petit, C.: OTOF encodes multiple

[95635] long and short isoforms: genetic evidence that the long ones underlie

[95636] recessive deafness DFNB9. *Am. J. Hum. Genet.* 67: 591–600, 2000.

[95637] 16663.

[95638] Yasunaga, S.; Petit, C.: Physical map of the region sur–

rounding

[95639] the OTOFERLIN locus on chromosome 2p22–p23. Genomics 66: 110–112,

[95640] 2000.

[95641] 16664.

[95642] Boffa, M. B.; Reid, S.; Joo, E.; Nesheim, M. E.; Koschinsky, M.

[95643] L.: Characterization of the gene encoding human TAFI (thrombin–activable

[95644] fibrinolysis inhibitor; plasma procarboxypeptidase B). Biochemistry 38:

[95645] 6547–6558, 1999.

[95646] 16665.

[95647] Eaton, D. L.; Malloy, B. E.; Tsai, S. P.; Henzel, W.; Drayna, D.

[95648] : Isolation, molecular cloning, and partial characterization of a

[95649] novel carboxypeptidase B from human plasma. J. Biol. Chem. 266:

[95650] 21833–21838, 1991.

[95651] 16666.

- [95652] Koschinsky, M. L.; Boffa, M. B.; Nesheim, M. E.; Zinman, B.; Hanley,
- [95653] A. J. G.; Harris, S. B.; Cao, H.; Hegele, R. A.: Association of a
- [95654] single nucleotide polymorphism in CPB2 encoding the thrombin-activable
- [95655] fibrinolysis inhibitor (TAFI) with blood pressure. Clin. Genet. 60:
- [95656] 345-349, 2001.
- [95657] 16667.
- [95658] Tregouet, D.-A.; Aubert, H.; Henry, M.; Morange, P.; Visvikis,
- [95659] S.; Juhan-Vague, I.; Tired, L.: Combined segregation-linkage analysis
- [95660] of plasma thrombin-activatable fibrinolysis inhibitor (TAFI) antigen
- [95661] levels with TAFI gene polymorphisms. Hum. Genet. 109: 191-197, 2001.
- [95662] 16668.
- [95663] Tsai, S. P.; Drayna, D.: The gene encoding human plasma carboxypeptidase
- [95664] B (CPB2) resides on chromosome 13. Genomics 14:

549–550, 1992.

[95665] 16669.

[95666] Vanhoof, G.; Wauters, J.; Schatteman, K.; Hendriks, D.;
Goossens,

[95667] F.; Bossuyt, P.; Scharpe, S.: The gene for human car-
boxypeptidase

[95668] U (CPU)--a proposed novel regulator of plasminogen acti-
vation--maps

[95669] to 13q14.11. Genomics 38: 454–455, 1996.

[95670] 16670.

[95671] Zhao, L.; Morser, J.; Bajzar, L.; Nesheim, M.; Nagashima,
N.:

[95672] Identification and characterization of two thrombin-acti-
vatable fibrinolysis

[95673] inhibitor isoforms. Thromb. Haemost. 80: 949–955, 1998.

[95674] 16671.

[95675] McGwire, G. B.; Tan, F.; Michel, B.; Rehli, M.; Skidgel, R. A.

[95676] : Identification of a membrane-bound carboxypeptidase
as the mammalian

[95677] homolog of duck gp180, a hepatitis B virus-binding pro-
tein. Life

[95678] Sci. 60: 715–724, 1997.

[95679] 16672.

[95680] Riley, D. A.; Tan, F.; Miletich, D. J.; Skidgel, R. A.: Chromosomal

[95681] localization of the genes for human carboxypeptidase D (CPD) and the

[95682] active 50–kilodalton subunit of human carboxypeptidase N (CPN1). Genomics 50:

[95683] 105–108, 1998.

[95684] 16673.

[95685] Tan, F.; Rehli, M.; Krause, S. W.; Skidgel, R. A.: Sequence of

[95686] human carboxypeptidase D reveals it to be a member of the regulatory

[95687] carboxypeptidase family with three tandem active site domains. Biochem.

[95688] J. 327: 81–87, 1997.

[95689] 16674.

[95690] Rajadhyaksha, A. Riviere, M.; Van Vooren, P.; Szpirer, J.; Szpirer,

[95691] C.; Babin, J.; Bina, M.: Assignment of AR1, transcription

factor

[95692] 20 (TCF20), to human chromosome 22q13.3 with somatic cell hybrids

[95693] and in situ hybridization. Cytogenet. Cell Genet. 81: 176–177, 1998.

[95694] 16675.

[95695] Sanz, L.; Moscat, J.; Diaz-Meco, M. T.: Molecular characterization

[95696] of a novel transcription factor that controls stromelysin expression. Molec.

[95697] Cell. Biol. 15: 3164–3170, 1995.

[95698] 16676.

[95699] Ding, H.; Descheemaeker, K.; Marynen, P.; Nelles, L.; Carvalho,

[95700] T.; Carmo-Fonseca, M.; Collen, D.; Belayew, A.: Characterization

[95701] of a helicase-like transcription factor involved in the expression

[95702] of the human plasminogen activator inhibitor-1 gene. DNA Cell Biol. 15:

[95703] 429–442, 1996.

[95704] 16677.

- [95705] Lin, Y.; Sheridan, P. L.; Jones, K. A.; Evans, G. A.: The
HIP116
- [95706] SNF2/SWI2-related transcription factor gene (SNF2L3) is
located on
- [95707] human chromosome 3q25.1-q26.1 Genomics 27:
381-382, 1995.
- [95708] 16678.
- [95709] Moinova, H. R.; Chen, W.-D.; Shen, L.; Smiraglia, D.;
Olechnowicz,
- [95710] J.; Ravi, L.; Kasturi, L.; Myeroff, L.; Plass, C.; Parsons, R.;
Minna,
- [95711] J.; Willson, J. K. V.; Green, S. B.; Issa, J.-P.; Markowitz, S. D.
- [95712] : HLTF gene silencing in human colon cancer. Proc. Nat.
Acad. Sci. 99:
- [95713] 4562-4567, 2002.
- [95714] 16679.
- [95715] Sheridan, P. L.; Schorpp, M.; Voz, M. L.; Jones, K. A.:
Cloning
- [95716] of an SNF2/SWI2-related protein that binds specifically to
the SPH
- [95717] motifs of the SV40 enhancer and to the HIV-1 promoter. J.
Biol. Chem. 270:

- [95718] 4575–4587, 1995.
- [95719] 16680.
- [95720] Ambros, P. F.; Schmid, J.; Rumpler, S.; Binder, B. R.; de
Martin,
- [95721] R.: Localization of the human I-kappa-B kinase-beta
(IKBKB) to chromosome
- [95722] 8p11.2 by fluorescence in situ hybridization and radiation
hybrid
- [95723] mapping. Genomics 54: 575–576, 1998.
- [95724] 16681.
- [95725] Pasparakis, M.; Courtois, G.; Hafner, M.; Schmidt-Sup-
prian, M.;
- [95726] Nenci, A.; Toksoy, A.; Krampert, M.; Goebeler, M.;
Gillitzer, R.;
- [95727] Israel, A.; Krieg, T.; Rajewsky, K.; Haase, I.: TNF-mediated
inflammatory
- [95728] skin disease in mice with epidermis-specific deletion of
IKK2. Nature 417:
- [95729] 861–866, 2002.
- [95730] 16682.
- [95731] Rossi, A.; Kapahi, P.; Natoli, G.; Takahashi, T.; Chen, Y.;

Karin,

[95732] M.; Santoro, M. G.: Anti-inflammatory cyclopentenone
prostaglandins

[95733] are direct inhibitors of I-kappa-B kinase. Nature 403:
103-108,

[95734] 2000.

[95735] 16683.

[95736] Shindo, M.; Nakano, H.; Sakon, S.; Yagita, H.; Mihara, M.;
Okumura,

[95737] K.: Assignment of I-kappa-B kinase beta (IKBKB) to human
chromosome

[95738] band 8p12-p11 by in situ hybridization. Cytogenet. Cell
Genet. 82:

[95739] 32-33, 1998.

[95740] 16684.

[95741] Tang, G.; Yang, J.; Minemoto, Y.; Lin, A.: Blocking cas-
pase-3-mediated

[95742] proteolysis of IKK-beta suppresses TNF-alpha-induced
apoptosis. Molec.

[95743] Cell 8: 1005-1016, 2001.

[95744] 16685.

[95745] Woronicz, J. D.; Gao, X.; Cao, Z.; Rothe, M.; Goeddel, D. V.:

[95746] IkappaB kinase-beta: NF-kappa-B activation and complex formation with

[95747] IkappaB kinase-alpha and NIK. Science 278: 866-869, 1997.

[95748] 16686.

[95749] Yin, M.-J.; Yamamoto, Y.; Gaynor, R. B.: The anti-inflammatory

[95750] agents aspirin and salicylate inhibit the activity of I-kappa-B kinase-beta. Nature 396:

[95751] 77-80, 1998.

[95752] 16687.

[95753] Zandi, E.; Rothwarf, D. M.; Delhase, M.; Hayakawa, M.; Karin,

[95754] M.: The IkappaB kinase complex (IKK) contains two kinase subunits,

[95755] IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappa-B

[95756] activation. Cell 91: 243-252, 1997.

[95757] 16688.

[95758] Aoki, M.; Hamada, F.; Sugimoto, T.; Sumida, S.; Akiyama,

T.; Toyoshima,

[95759] K.: The human cot proto-oncogene encodes two protein serine/threonine

[95760] kinases with different transforming activities by alternative initiation

[95761] of translation. J. Biol. Chem. 268: 22723–22732, 1993.

[95762] 16689.

[95763] Bataille, V.; Hiles, R.; Bishop, J. A. N.: Retinoblastoma, melanoma

[95764] and the atypical mole syndrome. Brit. J. Derm. 132: 134–138, 1995.

[95765] 16690.

[95766] Chilosi, M.; Doglioni, C.; Yan, Z.; Lestani, M.; Menestrina, F.;

[95767] Sorio, C.; Benedetti, A.; Vinante, F.; Pizzolo, G.; Inghirami, G.

[95768] : Differential expression of cyclin-dependent kinase 6 in cortical

[95769] thymocytes and T-cell lymphoblastic lymphoma/leukemia. Am. J. Pathol. 152:

[95770] 209–217, 1998.

[95771] 16691.

- [95772] Costello, J. F.; Plass, C.; Arap, W.; Chapman, V. M.; Held, W.
- [95773] A.; Berger, M. S.; Su Huang, H. J.; Cavenee, W. K.: Cyclin-dependent
- [95774] kinase 6 (cdk6) amplification in human gliomas identified using two-dimensional
- [95775] separation of genomic DNA. *Cancer Res.* 57: 1250–1254, 1997.
- [95776] 16692.
- [95777] Guan, K.-L.; Jenkins, C. W.; Li, Y.; Nichols, M. A.; Wu, X.; O'Keefe,
- [95778] C. L.; Matera, A. G.; Xiong, Y.: Growth suppression by p18, a p16(INK4/MTS1)–
- [95779] and p14(INK4B/MTS2)–related CDK6 inhibitor, correlates with wild-type
- [95780] pRb function. *Genes Dev.* 8: 2939–2952, 1994.
- [95781] 16693.
- [95782] Lien, H.-C.; Lin, C.-W.; Huang, P.-H.; Chang, M.-L.; Hsu, S.-M.
- [95783] : Expression of cyclin-dependent kinase 6 (cdk6) and frequent loss
- [95784] of CD44 in nasal–nasopharyngeal NK/T-cell lymphomas:

comparison with

[95785] CD56-negative peripheral T-cell lymphomas. Lab. Invest.
80: 893-900,

[95786] 2000.

[95787] 16694.

[95788] Meyerson, M.; Harlow, E.: Identification of G1 kinase activ-
ity

[95789] for cdk6, a novel cyclin D partner. Molec. Cell. Biol. 14:
2077-2086,

[95790] 1994.

[95791] 16695.

[95792] Shennan, M. G.; Badin, A.-C.; Walsh, S.; Summers, A.;
From, L.;

[95793] McKenzie, M.; Goldstein, A. M.; Tucker, M. A.; Hogg, D.;
Lassam, N.

[95794] : Lack of germline CDK6 mutations in familial melanoma.
Oncogene 19:

[95795] 1849-1852, 2000.

[95796] 16696.

[95797] Traboulsi, E. I.; Zimmerman, L. E.; Manz, H. J.: Cutaneous
malignant

[95798] melanoma in survivors of heritable retinoblastoma. Arch.
Ophthal. 106:

[95799] 1059–1061, 1988.

[95800] 16697.

[95801] Blais, A.; Labrie, Y.; Pouliot, F.; Lachance, Y.; Labrie, C.:

- [95802] Structure of the gene encoding the human cyclin-dependent kinase inhibitor
- [95803] p18 and mutational analysis in breast cancer. Biochem. Biophys. Res.
- [95804] Commun. 247: 146–153, 1998.
- [95805] 16698.
- [95806] Lapointe, J.; Lachance, Y.; Labrie, Y.; Labrie, C.: A p18 mutant
- [95807] defective in CDK6 binding in human breast cancer cells. Cancer Res. 56:
- [95808] 4586–4589, 1996.
- [95809] 16699.
- [95810] Bontemps, Y.; Maquart, F.-X.; , Wegrowski, Y.: Human UDP-glucose
- [95811] dehydrogenase gene: complete cloning and transcription start mapping. Biochem.
- [95812] Biophys. Res. Commun. 275: 981–985, 2000.
- [95813] 16700.
- [95814] Hempel, J.; Perozich, J.; Romovacek, H.; Hinich, A.; Kuo, I.; Feingold,
- [95815] D. S.: UDP-glucose dehydrogenase from bovine liver: pri-

mary structure

[95816] and relationship to other dehydrogenases. Protein Sci. 3: 1074–1080,

[95817] 1994.

[95818] 16701.

[95819] Marcu, O.; Stathakis, D. G.; Marsh, J. L.: Assignment of the UGDH

[95820] locus encoding UDP–glucose dehydrogenase to human chromosome band

[95821] 4p15.1 by radiation hybrid mapping. Cytogenet. Cell Genet. 86: 244–245,

[95822] 1999.

[95823] 16702.

[95824] Spicer, A. P.; Kaback, L. A.; Smith, T. J.; Seldin, M. F.: Molecular

[95825] cloning and characterization of the human and mouse UDP–glucose dehydrogenase

[95826] genes. J. Biol. Chem. 273: 25117–25124, 1998.

[95827] 16703.

[95828] Walsh, E. C.; Stainier, D. Y. R.: UDP–glucose dehydrogenase required

- [95829] for cardiac valve formation in zebrafish. Science 293:
1670–1674,
- [95830] 2001.
- [95831] 16704.
- [95832] Akamizu, T.; Ikuyama, S.; Saji, M.; Kosugi, S.; Kozak, C.;
McBride,
- [95833] O. W.; Kohn, L. D.: Cloning, chromosomal assignment, and
regulation
- [95834] of the rat thyrotropin receptor: expression of the gene is
regulated
- [95835] by thyrotropin, agents that increase cAMP levels, and thy-
roid autoantibodies. Proc.
- [95836] Nat. Acad. Sci. 87: 5677–5681, 1990.
- [95837] 16705.
- [95838] Reyes, R.; Duprat, F.; Lesage, F.; Fink, M.; Salinas, M.; Far-
man,
- [95839] N.; Lazdunski, M.: Cloning and expression of a novel pH-
sensitive
- [95840] two pore domain K⁺ channel from human kidney. J. Biol.
Chem. 273:
- [95841] 30863–30869, 1998.

- [95842] 16706.
- [95843] Becker, W.; Weber, Y.; Wetzel, K.; Eirmbter, K.; Tejedor, F. J.;
- [95844] Joost, H.-G.: Sequence characteristics, subcellular localization,
- [95845] and substrate specificity of DYRK-related kinases, a novel family
- [95846] of dual specificity protein kinases. J. Biol. Chem. 273: 25893-25902,
- [95847] 1998.
- [95848] 16707.
- [95849] Good, D.; Busfield, F.; Duffy, D.; Lovelock, P. K.; Kesting, J.
- [95850] B.; Cameron, D. P.; Shaw, J. T. E.: Familial Paget's disease of bone:
- [95851] nonlinkage to the PDB1 and PDB2 loci on chromosomes 6p and 18q in
- [95852] a large pedigree. J. Bone Miner. Res. 16: 33-38, 2001.
- [95853] 16708.
- [95854] Hocking, L.; Slee, F.; Haslam, S. I.; Cundy, T.; Nicholson, G.;
- [95855] van Hul, W.; Ralston, S. H.: Familial Paget's disease of

bone: patterns

[95856] of inheritance and frequency of linkage to chromosome
18q. Bone 26:

[95857] 577–580, 2000.

[95858] 16709.

[95859] Marco–Mingot, M.; San–Millan, J. L.; Wuyts, W.; Bachiller–
Corral,

[95860] J.; Van Hul, W.; Morales–Piga, A. A.: Lack of mutations in
the RANK

[95861] gene in Spanish patients with Paget disease of bone.
(Letter) Clin.

[95862] Genet. 60: 86–88, 2001.

[95863] 16710.

[95864] Li, J.; Sarosi, I.; Yan, X.–Q.; Morony, S.; Capparelli, C.; Tan,

[95865] H.–L.; McCabe, S.; Elliott, R.; Scully, S.; Van, G.; Kaufman,
S.;

[95866] Juan, S.–C.; Sun, Y.; Tarpley, J.; Martin, L.; Christensen, K.;
McCabe,

[95867] J.; Kostenuik, P.; Hsu, H.; Fletcher, F.; Dunstan, C. R.;
Lacey, D.

[95868] L.; Boyle, W. J.: RANK is the intrinsic hematopoietic cell

surface

[95869] receptor that controls osteoclastogenesis and regulation of bone mass

[95870] and calcium metabolism. Proc. Nat. Acad. Sci. 97: 1566–1571, 2000.

[95871] 16711.

[95872] Nakagawa, N.; Kinosaki, M.; Yamaguchi, K.; Shima, N.; Yasuda, H.;

[95873] Yano, K.; Morinaga, T.; Higashio, K.: RANK is the essential signaling

[95874] receptor for osteoclast differentiation factor in osteoclastogenesis. Biochem.

[95875] Biophys. Res. Commun. 253: 395–400, 1998.

[95876] 16712.

[95877] Wuyts, W.; Van Wesenbeeck, L.; Morales-Piga, A.; Ralston, S.;

[95878] Hocking, L.; Vanhoenacker, F.; Westhovens, R.; Verbruggen, L.; Anderson,

[95879] D.; Hughes, A.; Van Hul, W.: Evaluation of the role of RANK and OPG

[95880] genes in Paget's disease of bone. Bone 28: 104–107, 2001.

- [95881] 16713.
- [95882] Devilder, M.-C.; Cadoret, E.; Cherel, M.; Moreau, I.; Ron-
deau,
- [95883] G.; Bezieau, S.; Moisan, J.-P.: cDNA cloning, gene charac-
terization
- [95884] and 13q14.3 chromosomal assignment of CHC1-L, a
chromosome condensation
- [95885] regulator-like guanine nucleotide exchange factor. Ge-
nomics 54:
- [95886] 99-106, 1998.
- [95887] 16714.
- [95888] Phillips, N. J.; Zeigler, M. R.; Deaven, L. L.: A cDNA from
the
- [95889] ovarian cancer critical region of deletion on chromosome
17p13.3. Cancer
- [95890] Lett. 102: 85-90, 1996.
- [95891] 16715.
- [95892] Schultz, D. C.; Vanderveer, L.; Berman, D. B.; Hamilton, T.
C.;
- [95893] Wong, A. J.; Godwin, A. K.: Identification of two candidate
tumor

- [95894] suppressor genes on chromosome 17p13.3. Cancer Res. 56: 1997–2002,
- [95895] 1996.
- [95896] 16716.
- [95897] Peyrard, M.; Parveneh, S.; Lagercrantz, S.; Ekman, M.; Fransson,
- [95898] I.; Sahlen, S.; Dumanski, J. P.: Cloning, expression pattern, and
- [95899] chromosomal assignment to 16q23 of the human gamma-adaptin gene (ADTG). Genomics 50:
- [95900] 275–280, 1998.
- [95901] 16717.
- [95902] Adamou, J. E.; Aiyar, N.; Van Horn, S.; Elshourbagy, N. A.: Cloning
- [95903] and functional characterization of the human vasoactive intestinal
- [95904] peptide (VIP)–2 receptor. Biochem. Biophys. Res. Commun. 209: 385–392,
- [95905] 1995.
- [95906] 16718.
- [95907] Harmar, A. J.; Marston, H. M.; Shen, S.; Spratt, C.; West, K.

M.;

[95908] Sheward, W. J.; Morrison, C. F.; Dorin, J. R.; Piggins, H. D.;
Reubi,

[95909] J.-C.; Kelly, J. S.; Maywood, E. S.; Hastings, M. H.: The
VPAC2 receptor

[95910] is essential for circadian function in the mouse suprachiasmatic nuclei. Cell 109:

[95911] 497–508, 2002.

[95912] 16719.

[95913] Mackay, M.; Fantes, J.; Scherer, S.; Boyle, S.; West, K.; Tsui,

[95914] L.-C.; Belloni, E.; Lutz, E.; Van Heyningen, V.; Harmar, A.
J.: Chromosomal

[95915] localization in mouse and human of the vasoactive intestinal peptide

[95916] receptor type 2 gene: a possible contributor to the holoprosencephaly

[95917] 3 phenotype. Genomics 37: 345–353, 1996.

[95918] 16720.

[95919] Svoboda, M.; Tastenoy, M.; Van Rampelbergh, J.;
Goossens, J.-F.;

[95920] De Neef, P.; Waelbroeck, M.; Robberecht, P.: Molecular

cloning and

[95921] functional characterization of a human VIP receptor from SUP-T1 lymphoblasts. *Biochem.*

[95922] *Biophys. Res. Commun.* 205: 1617-1624, 1994.

[95923] 16721.

[95924] Andre, E.; Conquet, F.; Steinmayr, M.; Stratton, S. C.; Porciatti,

[95925] V.; Becker-Andre, M.: Disruption of retinoid-related orphan receptor

[95926] beta changes circadian behavior, causes retinal degeneration and leads

[95927] to vacillans phenotype in mice. *EMBO J.* 17: 3867-3877, 1998.

[95928] 16722.

[95929] Paravicini, G.; Steinmayr, M.; Andre, E.; Becker-Andre, M.: The

[95930] metastasis suppressor candidate nucleotide diphosphate kinase NM23

[95931] specifically interacts with members of the ROR/RZR nuclear orphan

[95932] receptor subfamily. *Biochem. Biophys. Res. Commun.* 227: 82-87, 1996.

[95933] 16723.

[95934] Sirlin, J. L.: Vacillans, a neurological mutant in the house mouse

[95935] linked with brown. J. Genet. 54: 42–48, 1956.

[95936] 16724.

[95937] Bourdi, M.; Demady, D.; Martin, J. L.; Jabbour, S. K.; Martin,

[95938] B. M.; George, J. W.; Pohl, L. R.: cDNA cloning and baculovirus expression

[95939] of the human liver endoplasmic reticulum P58: characterization as

[95940] a protein disulfide isomerase isoform, but not as a protease or a

[95941] carnitine acyltransferase. Arch. Biochem. Biophys. 323: 397–403,

[95942] 1995.

[95943] 16725.

[95944] Briquet-Laugier, V.; Xia, Y.-R.; Rooke, K.; Mehrabian, M.; Lusi,

[95945] A. J.; Doolittle, M. H.: Mapping of three members of the mouse protein

- [95946] disulfide isomerase family. Mammalian Genome 9: 176–177, 1998.
- [95947] 16726.
- [95948] Hirano, N.; Shibasaki, F.; Sakai, R.; Tanaka, T.; Nishida, J.;
- [95949] Yazaki, Y.; Takenawa, T.; Hirai, H.: Molecular cloning of the human
- [95950] glucose-regulated protein ERp57/GRP58, a thiol-dependent reductase:
- [95951] identification of its secretory form and inducible expression by the
- [95952] oncogenic transformation. Europ. J. Biochem. 234: 336–342, 1995.
- [95953] 16727.
- [95954] Koivunen, P.; Helaakoski, T.; Annunen, P.; Veijola, J.; Raisanen,
- [95955] S.; Pihlajaniemi, T.; Kivirikko, K. I.: ERp60 does not substitute
- [95956] for protein disulphide isomerase as the beta-subunit of prolyl 4-hydroxylase. Biochem.
- [95957] J. 316: 599–605, 1996.
- [95958] 16728.

- [95959] Koivunen, P.; Horelli-Kuitunen, N.; Helaakoski, T.; Karvonen, P.;
- [95960] Jaakkola, M.; Palotie, A.; Kivirikko, K. I.: Structures of the human
- [95961] gene for the protein disulfide isomerase-related polypeptide ERp60
- [95962] and a processed gene and assignment of these genes to 15q15 and 1q21. Genomics 42:
- [95963] 397-404, 1997.
- [95964] 16729.
- [95965] Oliver, J. D.; van der Wal, F. J.; Bulleid, N. J.; High, S.: Interaction
- [95966] of the thiol-dependent reductase ERp57 with nascent glycoproteins. Science 275:
- [95967] 86-88, 1997.
- [95968] 16730.
- [95969] Berthelsen, J.; Viggiano, L.; Schulz, H.; Ferretti, E.; Con-
salez,
- [95970] G. G.; Rocchi, M.; Blasi, F.: PKNOX1, a gene encoding PREP1, a new
- [95971] regulator of Pbx activity, maps on human chromosome 21q22.3 and murine

- [95972] chromosome 17B/C. *Genomics* 47: 323–324, 1998.
- [95973] 16731.
- [95974] Chen, H.; Rossier, C.; Nakamura, Y.; Lynn, A.; Chakravarti, A.;
- [95975] Antonarakis, S. E.: Cloning of a novel homeobox–containing gene,
- [95976] PKNOX1, and mapping to human chromosome 21q22.3. *Genomics* 41: 193–200,
- [95977] 1997.
- [95978] 16732.
- [95979] Morita, K.; Furuse, M.; Fujimoto, K.; Tsukita, S.: Claudin multigene
- [95980] family encoding four–transmembrane domain protein components of tight
- [95981] junction strands. *Proc. Nat. Acad. Sci.* 96: 511–516, 1999.
- [95982] 16733.
- [95983] Chiang, P.–W.; Fogel, E.; Jackson, C. L.; Lieuallen, K.; Lennon,
- [95984] G.; Qu, X.; Wang, S.–Q.; Kurnit, D. M.: Isolation, sequencing, and
- [95985] mapping of the human homologue of the yeast transcrip–

tion factor,

[95986] SPT5. Genomics 38: 421–424, 1996.

[95987] 16734.

[95988] Chiang, P.-W.; Stubbs, L.; Zhang, L.; Kurnit, D. M.: Isolation

[95989] of murine SPT5 homologue: completion of the isolation and characterization

[95990] of human and murine homologues of yeast chromatin structural protein

[95991] complex SPT4, SPT5, and SPT6. Genomics 47: 426–428, 1998.

[95992] 16735.

[95993] Iwaki, A.; Nagano, T.; Nakagawa, M.; Iwaki, T.; Fukumaki, Y.:

[95994] Identification and characterization of the gene encoding a new member

[95995] of the alpha-crystallin/Small hsp family, closely linked to the alpha-B-crystallin

[95996] gene in a head-to-head manner. Genomics 45: 386–394, 1997.

[95997] 16736.

[95998] de Coo, R. F. M.; Buddiger, P.; Smeets, H. J. M.; van Oost,

B.

- [95999] A.: Molecular cloning and characterization of the human mitochondrial
- [96000] NADH:oxidoreductase 10-kDa gene (NDUFV3). Genomics 45: 434–437,
- [96001] 1997.
- [96002] 16737.
- [96003] Canu, N.; Possenti, R.; Ricco, A. S.; Rocchi, M.; Levi, A.: Cloning,
- [96004] structural organization analysis, and chromosomal assignment of the
- [96005] human gene for the neurosecretory protein VGF. Genomics 45: 443–446,
- [96006] 1997.
- [96007] 16738.
- [96008] Ferri, G.-L.; Possenti, R.: vgf: a neurotrophin-inducible gene
- [96009] expressed in neuroendocrine tissues. Trends Endocr. Metab. 7: 233–239,
- [96010] 1996.
- [96011] 16739.

[96012] Levi, A.; Eldridge, J. D.; Paterson, B. M.: Molecular cloning
[96013] of a gene sequence regulated by nerve growth factor. Sci-
ence 229:
[96014] 393–395, 1985.
[96015] 16740.
[96016] Salton, S. R. J.; Fischberg, D. J.; Dong, K. W.: Structure of
[96017] the gene encoding VGF, a nervous system-specific mRNA
that is rapidly
[96018] and selectively induced by nerve growth factor in PC12
cells. Molec.
[96019] Cell. Biol. 11: 2335–2349, 1991.
[96020] 16741.
[96021] Gautvik, K. M.; de Lecea, L.; Gautvik, V. T.; Danielson, P.
E.;
[96022] Tranque, P.; Dopazo, A.; Bloom, F. E.; Sutcliffe, J. G.:
Overview
[96023] of the most prevalent hypothalamus-specific mRNAs as
identified by
[96024] directional tag PCR subtraction. Proc. Nat. Acad. Sci. 93:
8733–8738,
[96025] 1996.

[96026] 16742.

[96027] Green, E. D.; Olson, M. V.: Chromosomal region of the
cystic

[96028] fibrosis gene in yeast artificial chromosomes: a model for
human genome

[96029] mapping. Science 250: 94–98, 1990.

[96030] 16743.

[96031] Fritzler, M. J.; Lung, C.-C.; Hamel, J. C.; Griffith, K. J.;
Chan,

[96032] E. K. L.: Molecular characterization of golgin-245, a novel
Golgi

[96033] complex protein containing a granin signature. J. Biol.
Chem. 270:

[96034] 31262–31268, 1995.

[96035] 16744.

[96036] Gencik, M.; Dahmen, N.; Wieczorek, S.; Kasten, M.; Bier-
brauer,

[96037] J.; Anghelescu, I.; Szegedi, A.; Menezes Saecker, A. M.;
Epplen, J.

[96038] T.: A prepro-orexin gene polymorphism is associated with
narcolepsy. Neurology 56:

- [96039] 115–117, 2001.
- [96040] 16745.
- [96041] Hagan, J. J.; Leslie, R. A.; Patel, S.; Evans, M. L.; Wattam, T.
- [96042] A.; Holmes, S.; Benham, C. D.; Taylor, S. G.; Routledge, C.; Hemmati,
- [96043] P.; Munton, R. P.; Ashmeade, T. E.; Shah, A. S.; Hatcher, J. P.; Hatcher,
- [96044] P. D.; Jones, D. N. C.; Smith, M. I.; Piper, D. C.; Hunter, A. J.;
- [96045] Porter, R. A.; Upton, N.: Orexin A activates locus coeruleus cell
- [96046] firing and increases arousal in the rat. Proc. Nat. Acad. Sci. 96:
- [96047] 10911–10916, 1999.
- [96048] 16746.
- [96049] Hara, J.; Beuckmann, C. T.; Nambu, T.; Willie, J. T.; Chemelli,
- [96050] R. M.; Sinton, C. M.; Sugiyama, F.; Yagami, K.; Goto, K.; Yanagisawa,
- [96051] M.; Sakurai, T.: Genetic ablation of orexin neurons in mice results

- [96052] in narcolepsy, hypophagia, and obesity. *Neuron* 30: 345–354, 2001.
- [96053] 16747.
- [96054] Hungs, M.; Lin, L.; Okun, M.; Mignot, E.: Polymorphisms in the
- [96055] vicinity of the hypocretin/orexin are not associated with human narcolepsy. *Neurology* 57:
- [96056] 1893–1895, 2001.
- [96057] 16748.
- [96058] Nishino, S.; Ripley, B.; Overeem, S.; Lammers, G. J.; Mignot, E.
- [96059] : Hypocretin (orexin) deficiency in human narcolepsy. *Lancet* 355:
- [96060] 39–40, 2000.
- [96061] 16749.
- [96062] Chen, L.-S. K.; Lo, C. F.; Numann, R.; Cuddy, M.: Characterization
- [96063] of the human and rat phospholemmann (PLM) cDNAs and localization of
- [96064] the human PLM gene to chromosome 19q13.1. *Genomics* 41: 435–443,

[96065] 1997.

[96066] 16750.

[96067] Sakurai, T.; Amemiya, A.; Ishii, M.; Matsuzaki, I.; Chemelli,

[96068] R. M.; Tanaka, H.; Williams, S. C.; Richardson, J. A.; Ko-
złowski,

[96069] G. P.; Wilson, S.; Arch, J. R. S.; Buckingham, R. E.; Haynes,
A. C.;

[96070] Carr, S. A.; Annan, R. S.; McNulty, D. E.; Liu, W.-S.; Terrett,
J.

[96071] A.; Elshourbagy, N. A.; Bergsma, D. J.; Yanagisawa, M.:
Orexins and

[96072] orexin receptors: a family of hypothalamic neuropeptides
and G protein-coupled

[96073] receptors that regulate feeding behavior. Cell 92:
573-585, 1998.

[96074] 16751.

[96075] Heard, D. J.; Norby, P. L.; Holloway, J.; Vissing, H.: Human
ERR-gamma,

[96076] a third member of the estrogen receptor-related receptor
(ERR) subfamily

[96077] of orphan nuclear receptors: tissue-specific isoforms are
expressed

- [96078] during development in the adult. *Molec. Endocr.* 14: 382–392, 2000.
- [96079] 16752.
- [96080] Hong, H.; Yang, L.; Stallcup, M. R.: Hormone-independent transcriptional
- [96081] activation and coactivator binding by novel orphan nuclear receptor
- [96082] ERR3. *J. Biol. Chem.* 274: 22618–22626, 1999.
- [96083] 16753.
- [96084] Billin, A. N.; Eilers, A. L.; Queva, C.; Ayer, D. E.: Mlx, a novel
- [96085] Max-like BHLHZip protein that interacts with the Max network of transcription
- [96086] factors. *J. Biol. Chem.* 274: 36344–36350, 1999.
- [96087] 16754.
- [96088] Bjerknes, M.; Cheng, H.: TCFL4: a gene at 17q21.1 encoding a putative
- [96089] basic helix-loop-helix leucine-zipper transcription factor. *Gene* 181:
- [96090] 7–11, 1996.
- [96091] 16755.

- [96092] Cairo, S.; Merla, G.; Ubinati, F.; Ballabio, A.; Reymond, A.:
[96093] WBSCR14, a gene mapping to the Williams–Beuren syn-
drome deleted region,
[96094] is a new member of the Mlx transcription factor network.
Hum. Molec.
[96095] Genet. 10: 617–627, 2001.
[96096] 16756.
[96097] Fukuoka, S.–I.; Freedman, S. D.; Scheele, G. A.: A single
gene
[96098] encodes membrane–bound and free forms of GP–2, the
major glycoprotein
[96099] in pancreatic secretory (zymogen) granule membranes.
Proc. Nat. Acad.
[96100] Sci. 88: 2898–2902, 1991.
[96101] 16757.
[96102] Fukuoka, S.–I.; Freedman, S. D.; Yu, H.; Sukhatme, V. P.;
Scheele,
[96103] G. A.: GP–2/THP gene family encodes self–binding glyco-
sylphosphatidylinositol–anchored
[96104] proteins in apical secretory compartments of pancreas
and kidney. Proc.

- [96105] Nat. Acad. Sci. 89: 1189–1193, 1992.
- [96106] 16758.
- [96107] Fukuoka, S.-I.; Suzuki, M.; Okabayashi, K.; Takahashi, E.:
Assignment
[96108] of pancreatic zymogen granule membrane protein GP2
(GP2) to human
[96109] chromosome band 9q21.11 to q21.2 by in situ hybridiza-
tion. Cytogenet.
[96110] Cell Genet. 79: 231–232, 1997.
- [96111] 16759.
- [96112] Gregory, R. J.; Rich, D. P.; Cheng, S. H.; Souza, D. W.; Paul,
[96113] S.; Manavalan, P.; Anderson, M. P.; Welsh, M. J.; Smith, A.
E.: Maturation
[96114] and function of cystic fibrosis transmembrane conduc-
tance regulator
[96115] variants bearing mutations in putative nucleotide-binding
domains
[96116] 1 and 2. Molec. Cell. Biol. 11: 3886–3893, 1991.
- [96117] 16760.
- [96118] Greil, I.; Wagner, K.; Rosenkranz, W.: A new missense mu-
tation

- [96119] G1249E in exon 20 of the cystic fibrosis transmembrane conductance
- [96120] regulator (CFTR) gene. Hum. Hered. 44: 238–240, 1994.
- [96121] 16761.
- [96122] Guillermit, H.; Fanen, P.; Ferec, C.: A 3–prime splice site consensus
- [96123] sequence mutation in the cystic fibrosis gene. Hum. Genet. 85: 450–453,
- [96124] 1990.
- [96125] 16762.
- [96126] Guillermit, H.; Jehanne, M.; Quere, I.; Audrezet, M. P.; Mercier,
- [96127] B.; Ferec, C.: A novel mutation in exon 3 of the CFTR gene. Hum.
- [96128] Genet. 91: 233–235, 1993.
- [96129] 16763.
- [96130] Haardt, M.; Benharouga, M.; Lechardeur, D.; Kartner, N.; Lukacs,
- [96131] G. L.: C–terminal truncations destabilize the cystic fibrosis transmembrane
- [96132] conductance regulator without impairing its biogenesis: a

novel class

[96133] of mutation. J. Biol. Chem. 274: 21873–21877, 1999.

[96134] 16764.

[96135] Hamosh, A.; King, T. M.; Rosenstein, B. J.; Corey, M.; Levi-son,

[96136] H.; Durie, P.; Tsui, L.-C.; McIntosh, I.; Keston, M.; Brock, D. J.

[96137] H.; Macek, M., Jr.; Zemkova, D.; Krasnicanova, H.; Vavrova, V.; Macek,

[96138] M., Sr.; Golder, N.; Schwarz, M. J.; Super, M.; Watson, E. K.; Williams,

[96139] C.; Bush, A.; O'Mahoney, S. M.; Humphries, P.; DeArce, M. A.; Reis,

[96140] A.; Burger, J.; Stuhmann, M.; Schmidtke, J.; Wulbrand, U.; Dork,

[96141] T.; Tummeler, B.; Cutting, G. R.: Cystic fibrosis patients bearing

[96142] both the common missense mutation gly-to-aspartate at codon 551 and the

[96143] delta-F508 mutation are clinically indistinguishable from delta-F508

[96144] homozygotes, except for decreased risk of meconium

ileus. Am. J.

[96145] Hum. Genet. 51: 245–250, 1992.

[96146] 16765.

[96147] Hamosh, A.; Trapnell, B. C.; Zeitlin, P. L.; Montrose-Rafizadeh,

[96148] C.; Rosenstein, B. J.; Crystal, R. G.; Cutting, G. R.: Severe deficiency

[96149] of cystic fibrosis transmembrane conductance regulator messenger RNA

[96150] carrying nonsense mutations R553X and W1316X in respiratory epithelial

[96151] cells of patients with cystic fibrosis. J. Clin. Invest. 88: 1880–1885,

[96152] 1991.

[96153] 16766.

[96154] Hefferon, T. W.; Broackes-Carter, F. C.; Harris, A.; Cutting,

[96155] G. R.: Atypical 5-prime splice sites cause CFTR exon 9 to be vulnerable

[96156] to skipping. Am. J. Hum. Genet. 71: 294–303, 2002.

[96157] 16767.

[96158] Highsmith, W. E.; Burch, L. H.; Boat, T. F.; Boucher, R. C.;

Silverman,

- [96159] L. M.; Knowles, M. R.: Identification of a homozygous point mutation
- [96160] in intron 19 in an inbred CF patient with mild disease and normal
- [96161] sweat chloride: creation of an alternative splice site resulting in
- [96162] base-sequence insertion in CFTR coding region between exons 19 and
- [96163] 20. (Abstract) *Pediat. Pulmonol. Suppl.* 6: 22 only, 1991.
- [96164] 16768.
- [96165] Highsmith, W. E., Jr.: Personal Communication. Chapel Hill,
- [96166] N.C. 1991.
- [96167] 16769.
- [96168] Iannuzzi, M. C.; Stern, R. C.; Collins, F. S.; Tom Hon, C.; Hidaka,
- [96169] N.; Strong, T.; Becker, L.; Drumm, M. L.; White, M. B.; Gerard, B.;
- [96170] Dean, M.: Two frameshift mutations in the cystic fibrosis gene. *Am.*
- [96171] *J. Hum. Genet.* 48: 227-231, 1991.

[96172] 16770.

[96173] Ivaschenko, T. E.; White, M. B.; Dean, M.; Baranov, V. S.: A

[96174] deletion of two nucleotides in exon 10 of the CFTR gene
in a Soviet

[96175] family with cystic fibrosis causing early infant death. Ge-
nomics 10:

[96176] 298-299, 1991.

[96177] 16771.

[96178] Jensen, T. J.; Loo, M. A.; Pind, S.; Williams, D. B.; Goldberg,

[96179] A. L.; Riordan, J. R.: Multiple proteolytic systems, including
the

[96180] proteasome, contribute to CFTR processing. Cell 83:
129-135, 1995.

[96181] 16772.

[96182] Jiang, Q.; Engelhardt, J. F.: Cellular heterogeneity of CFTR

[96183] expression and function in the lung: implications for gene
therapy

[96184] of cystic fibrosis. Europ. J. Hum. Genet. 6: 12-31, 1998.

[96185] 16773.

[96186] Jones, C. T.; McIntosh, I.; Keston, M.; Ferguson, A.; Brock,

D.

[96187] J. H.: Three novel mutations in the cystic fibrosis gene detected

[96188] by chemical cleavage: analysis of variant splicing and a nonsense

[96189] mutation. Hum. Molec. Genet. 1: 11–17, 1992.

[96190]

[96191] 100. Kabra, M; Kabra, S. K.; Ghosh, M.; Khanna, A.; Arora, S.; Menon,

[96192] P. S. N.; Verma, I. C.: Is the spectrum of mutations in Indian patients

[96193] with cystic fibrosis different? (Letter) Am. J. Med. Genet. 93:

[96194] 161–163, 2000.

[96195]

[96196] 101. Kalin, N.; Claass, A.; Sommer, M.; Puchelle, E.; Tummeler, B.

[96197] : Delta-F508 CFTR protein expression in tissues from patients with

[96198] cystic fibrosis. J. Clin. Invest. 103: 1379–1389, 1999.

[96199]

[96200] 102. Kelley, K. A.; Stamm, S.; Kozak, C. A.: Expression and chromosome

[96201] localization of the murine cystic fibrosis transmembrane conductance

[96202] regulator. *Genomics* 13: 381–388, 1992.

[96203]

[96204] 103. Kerem, B.; Buchanan, J. A.; Durie, P.; Corey, M. L.; Levison,

[96205] H.; Rommens, J. M.; Buchwald, M.; Tsui, L.-C.: DNA marker haplotype

[96206] association with pancreatic sufficiency in cystic fibrosis. *Am. J.*

[96207] *Hum. Genet.* 44: 827–834, 1989.

[96208]

[96209] 104. Kerem, B.; Rommens, J. M.; Buchanan, J. A.; Markiewicz, D.; Cox,

[96210] T. K.; Chakravarti, A.; Buchwald, M.; Tsui, L.-C.: Identification

[96211] of the cystic fibrosis gene: genetic analysis. *Science* 245: 1073–1080,

[96212] 1989.

[96213]

[96214] 105. Kerem, B.; Zielenski, J.; Markiewicz, D.; Bozon, D.;
Gazit, E.;

[96215] Yahav, J.; Kennedy, D.; Riordan, J. R.; Collins, F. S.; Rom-
mens, J.

[96216] M.; Tsui, L.-C.: Identification of mutations in regions cor-
responding

[96217] to the 2 putative nucleotide (ATP)-binding folds of the
cystic fibrosis

[96218] gene. Proc. Nat. Acad. Sci. 87: 8447-8451, 1990.

[96219]

[96220] 106. Kerem, E.; Corey, M.; Kerem, B.; Rommens, J.;
Markiewicz, D.;

[96221] Levison, H.; Tsui, L.-C.; Durie, P.: The relation between
genotype

[96222] and phenotype in cystic fibrosis--analysis of the most
common mutation

[96223] (delta-F508). New Eng. J. Med. 323: 1517-1522, 1990.

[96224]

[96225] 107. Kerem, E.; Kalman, Y. M.; Yahav, Y.; Shoshani, T.;
Abeliovich,

- [96226] D.; Szeinberg, A.; Rivlin, J.; Blau, H.; Tal, A.; Ben-Tur, L.; Springer,
- [96227] C.; Augarten, A.; Godfrey, S.; Lerer, I.; Branski, D.; Friedman, M.;
- [96228] Kerem, B.: Highly variable incidence of cystic fibrosis and different
- [96229] mutation distribution among different Jewish ethnic groups in Israel. Hum.
- [96230] Genet. 96: 193–197, 1995.
- [96231]
- [96232] 108. Kiesewetter, S.; Macek, M., Jr.; Davis, C.; Curristin, S. M.;
- [96233] Chu, C.-S.; Graham, C.; Shrimpton, A. E.; Cashman, S. M.; Tsui, L.-C.;
- [96234] Mickle, J.; Amos, J.; Highsmith, W. E.; Shuber, A.; Witt, D. R.; Crystal,
- [96235] R. G.; Cutting, G. R.: A mutation in CFTR produces different phenotypes
- [96236] depending on chromosomal background. Nature Genet. 5: 274–278, 1993.
- [96237]
- [96238] 109. Klinger, K.; Horn, G. T.; Stanislovitis, P.; Schwartz, R.

H.;

[96239] Fujiwara, T. M.; Morgan, K.: Cystic fibrosis mutations in the Hutterite

[96240] brethren. Am. J. Hum. Genet. 46: 983–987, 1990.

[96241]

[96242] 110. Kobayashi, K.; Knowles, M. R.; Boucher, R. C.; O'Brien, W. E.;

[96243] Beaudet, A. L.: Benign missense variations in the cystic fibrosis

[96244] gene. Am. J. Hum. Genet. 47: 611–615, 1990.

[96245]

[96246] 111. Kocher, O.; Comella, N.; Tognazzi, K.; Brown, L. F.: Identification

[96247] and partial characterization of PDZK1: a novel protein containing

[96248] PDZ interaction domains. Lab. Invest. 78: 117–125, 1998.

[96249]

[96250] 112. Konstan, M. W.; Byard, P. J.; Hoppel, C. L.; Davis, P. B.: Effect

[96251] of high-dose ibuprofen in patients with cystic fibrosis. New Eng.

[96252] J. Med. 332: 848–854, 1995.

[96253]

[96254] 113. Laroche, D.; Travert, G.: Abnormal frequency of
delta-F(508)

[96255] mutation in neonatal transitory hypertrypsinaemia. (Letter)
Lancet 337:

[96256] 55 only, 1991.

[96257]

[96258] 114. Latham, T.; Grabowski, G. A.; Theophilus, B. D. M.;
Smith, F.

[96259] I.: Complex alleles of the acid beta-glucosidase gene in
Gaucher

[96260] disease. Am. J. Hum. Genet. 47: 79–86, 1990.

[96261]

[96262] 115. Leoni, G.; Pitzalis, S.; Podda, R.; Zanda, M.; Silvetti,
M.;

[96263] Caocci, L.; Cao, A.; Rosatelli, M. C.: A specific cystic fibro-
sis

[96264] mutation (T338I) associated with the phenotype of iso-
lated hypotonic

[96265] dehydration. J. Pediat. 127: 281–283, 1995.

[96266]

[96267] 116. Lerer, I.; Sagi, M.; Cutting, G. R.; Abeliovich, D.: Cystic

[96268] fibrosis mutations delta-F508 and G542X in Jewish patients. J. Med.

[96269] Genet. 29: 131–133, 1992.

[96270]

[96271] 117. Logan, J.; Hiestand, D.; Daram, P.; Huang, Z.; Muccio, D. D.;

[96272] Hartman, J.; Haley, B.; Cook, W. J.; Sorscher, E. J.: Cystic fibrosis

[96273] transmembrane conductance regulator mutations that disrupt nucleotide

[96274] binding. J. Clin. Invest. 94: 228–236, 1994.

[96275]

[96276] 118. Loirat, F.; Hazout, S.; Lucotte, G.: G542X as a probable Phoenician

[96277] cystic fibrosis mutation. Hum. Biol. 69: 419–425, 1997.

[96278]

[96279] 119. Marino, C. R.; Matovcik, L. M.; Gorelick, F. S.; Cohn, J. A.

- [96280] : Localization of the cystic fibrosis transmembrane conductance regulator
- [96281] in pancreas. *J. Clin. Invest.* 88: 712–716, 1991.
- [96282]
- [96283] 120. Marshal, R. D.; Koontz, J.; Sklar, J.: Detection of mutations
- [96284] by cleavage of DNA heteroduplexes with bacteriophage resolvases. *Nature*
- [96285] *Genet.* 9: 177–183, 1995.
- [96286]
- [96287] 121. McCombie, W. R.; Adams, M. D.; Kelley, J. M.; FitzGerald, M.
- [96288] G.; Utterback, T. R.; Khan, M.; Dubnick, M.; Kerlavage, A. R.; Venter,
- [96289] J. C.; Fields, C.: *Caenorhabditis elegans* expressed sequence tags
- [96290] identify gene families and potential disease gene homologues. *Nature*
- [96291] *Genet.* 1: 124–131, 1992.
- [96292]
- [96293] 122. Mickle, J. E.; Macek, M., Jr.; Fulmer-Smentek, S. B.;

Egan, M.

[96294] M.; Schwiebert, E.; Guggino, W.; Moss, R.; Cutting, G. R.: A mutation

[96295] in the cystic fibrosis transmembrane conductance regulator gene associated

[96296] with elevated sweat chloride concentrations in the absence of cystic

[96297] fibrosis. Hum. Molec. Genet. 7: 729–735, 1998.

[96298]

[96299] 123. Mickle, J. E.; Milewski, M. I.; Macek, M., Jr.; Cutting, G. R.

[96300] : Effects of cystic fibrosis and congenital bilateral absence of the

[96301] vas deferens-associated mutations on cystic fibrosis transmembrane

[96302] conductance regulator-mediated regulation of separate channels. Am.

[96303] J. Hum. Genet. 66: 1485–1495, 2000.

[96304]

[96305] 124. Morral, N.; Nunes, V.; Casals, T.; Estivill, X.: CA/GT microsatellite

[96306] alleles within the cystic fibrosis transmembrane conduc-

tance regulator

[96307] (CFTR) gene are not generated by unequal crossing-over.

Genomics 10:

[96308] 692–698, 1991.

[96309]

[96310] 125. Moyer, B. D.; Duhaime, M.; Shaw, C.; Denton, J.;
Reynolds, D.;

[96311] Karlson, K. H.; Pfeiffer, J.; Wang, S.; Mickle, J. E.; Milewski,
M.;

[96312] Cutting, G. R.; Guggino, W. B.; Li, M.; Stanton, B. A.: The
PDZ-interacting

[96313] domain of cystic fibrosis transmembrane conductance
regulator is required

[96314] for functional expression in the apical plasma membrane.
J. Biol.

[96315] Chem. 275: 27069–27074, 2000.

[96316]

[96317] 126. Nelson, P. V.; Carey, W. F.; Morris, C. P.: Identifica-
tion of

[96318] a cystic fibrosis mutation: deletion of isoleucine-506.
Hum. Genet. 86:

[96319] 391–393, 1991.

[96320]

[96321] 127. Nunes, V.; Bonizzato, A.; Gaona, A.; Dognini, M.;
Chillon, M.;

[96322] Casals, T.; Pignatti, P. F.; Novelli, G.; Estivill, X.; Gasparini,
[96323] P.: A frameshift mutation (2869insG) in the second trans-
membrane

[96324] domain of the CFTR gene: identification, regional distribu-
tion, and

[96325] clinical presentation. (Letter) Am. J. Hum. Genet. 50:
1140–1142,

[96326] 1992.

[96327]

[96328] 128. Nunes, V.; Casals, T.; Gaona, A.; Antinolo, G.; Ferrer-
Calvete,

[96329] J.; Perez-Frias, J.; Tardio, E.; Molano, J.; Estivill, X.: Cystic

[96330] fibrosis patients with mutation 1949del84 in exon 13 of
the CFTR gene

[96331] have a similar clinical severity as delF508 homozygotes.
Hum. Mutat. 1:

[96332] 375–379, 1992.

[96333]

- [96334] 129. Nunes, V.; Chillon, M.; Dork, T.; Tummler, B.; Casals, T.; Estivill, X.: A new missense mutation (E92K) in the first trans-membrane domain of the CFTR gene causes a benign cystic fibrosis phenotype. Hum. Molec. Genet. 2: 79–80, 1993.
- [96338]
- [96339] 130. Nunes, V.; Gasparini, P.; Novelli, G.; Gaona, A.; Bonizzato, A.; Sangiuolo, F.; Balassopoulou, A.; Gimenez, F. J.; Dognini, M.; Ravnik-Glavac, M.; Cikuli, M.; Mokini, V.; Komel, R.; Dal-lapiccola, B.; Pignatti, P. F.; Loukopoulos, D.; Casals, T.; Estivill, X.: Analysis of 14 cystic fibrosis mutations in five south European populations. Hum. Genet. 87: 737–738, 1991.
- [96345]
- [96346] 131. Orita, M.; Iwahana, H.; Kanazawa, H.; Hayashi, K.; Sekiya, T.

[96347] : Detection of polymorphisms of human DNA by gel electrophoresis as

[96348] single-strand conformation polymorphisms. Proc. Nat. Acad. Sci. 86:

[96349] 2766–2770, 1989.

[96350]

[96351] 132. Orita, M.; Suzuki, Y.; Sekiya, T.; Hayashi, K.: Rapid and sensitive

[96352] detection of point mutations and DNA polymorphisms using the polymerase

[96353] chain reaction. Genomics 5: 874–879, 1989.

[96354]

[96355] 133. Orozco, L.; Friedman, K.; Chavez, M.; Lezana, J. L.; Villarreal,

[96356] M. T.; Carnevale, A.: Identification of the I507 deletion by site-directed

[96357] mutagenesis. Am. J. Med. Genet. 51: 137–139, 1994.

[96358]

[96359] 134. Osborne, L.; Knight, R.; Santis, G.; Hodson, M.: A mutation

[96360] in the second nucleotide binding fold of the cystic fibrosis

gene. Am.

[96361] J. Hum. Genet. 48: 608–612, 1991.

[96362]

[96363] 135. Osborne, L.; Santis, G.; Schwarz, M.; Klinger, K.;
Dork, T.;

[96364] McIntosh, I.; Schwartz, M.; Nunes, V.; Macek, M., Jr.; Reiss,
J.;

[96365] 46 others: Incidence and expression of the N1303K muta-
tion of the

[96366] cystic fibrosis (CFTR) gene. Hum. Genet. 89: 653–658,
1992.

[96367]

[96368] 136. Pier, G. B.; Grout, M.; Zaidi, T.; Meluleni, G.;
Mueschenborn,

[96369] S. S.; Banting, G.; Ratcliff, R.; Evans, M. J.; Colledge, W. H.:

[96370] Salmonella typhi uses CFTR to enter intestinal epithelial
cells. Nature 393:

[96371] 79–82, 1998.

[96372]

[96373] 137. Pier, G. B.; Grout, M.; Zaldi, T. S.; Olsen, J. C.; John-
son,

- [96374] L. G.; Yankaskas, J. R.; Goldberg, J. B.: Role of mutant CFTR in
- [96375] hypersusceptibility of cystic fibrosis patients to lung infections. Science 271:
- [96376] 63–67, 1996.
- [96377]
- [96378] 138. Pignatti, P. F.; Bombieri, C.; Benetazzo, M.; Casartelli, A.;
- [96379] Trabetti, E.; Gile, L. S.; Martinati, L. C.; Boner, A. L.; Luisetti,
- [96380] M.: CFTR gene variant IVS8–5T in disseminated bronchiectasis. (Letter) Am.
- [96381] J. Hum. Genet. 58: 889–892, 1996.
- [96382]
- [96383] 139. Pind, S.; Riordan, J. R.; Williams, D. B.: Participation of
- [96384] the endoplasmic reticulum chaperone calnexin (p88,IP90) in the biogenesis
- [96385] of the cystic fibrosis transmembrane conductance regulator. J. Biol.
- [96386] Chem. 269: 12784–12788, 1994.

[96387]

[96388] 140. Ramjeesingh, M.; Li, C.; Garami, E.; Huan, L.-J.; Galley, K.;

[96389] Wang, Y.; Bear, C. E.: Walker mutations reveal loose relationship

[96390] between catalytic and channel-gating activities of purified CFTR (cystic

[96391] fibrosis transmembrane conductance regulator). Biochemistry 38:

[96392] 1463-1468, 1999.

[96393]

[96394] 141. Reddy, M. M.; Light, M. J.; Quinton, P. M.: Activation of the

[96395] epithelial Na(+) channel (ENaC) requires CFTR Cl(-) channel function. Nature 402:

[96396] 301-304, 1999.

[96397]

[96398] 142. Rich, D. P.; Anderson, M. P.; Gregory, R. J.; Cheng, S. H.; Paul,

[96399] S.; Jefferson, D. M.; McCann, J. D.; Klinger, K. W.; Smith, A. E.;

- [96400] Welsh, M. J.: Expression of cystic fibrosis transmembrane conductance
- [96401] regulator corrects defective chloride channel regulation in cystic
- [96402] fibrosis airway epithelial cells. Nature 347: 358–363, 1990.
- [96403]
- [96404] 143. Riordan, J. R.; Rommens, J. M.; Kerem, B.; Alon, N.; Rozmahel,
- [96405] R.; Grzelczak, Z.; Zielenski, J.; Lok, S.; Plavsic, N.; Chou, J. L.;
- [96406] Drumm, M. L.; Iannuzzi, M. C.; Collins, F. S.; Tsui, L.-C.: Identification
- [96407] of the cystic fibrosis gene: cloning and characterization of complementary
- [96408] DNA. Science 245: 1066–1073, 1989.
- [96409]
- [96410] 144. Romey, M.-C.; Aguilar-Martinez, P.; Demaille, J.; Claustres,
- [96411] M.: Rapid detection of single nucleotide deletions: application to
- [96412] the beta-6 (-A) mutation of the beta-globin gene and to

cystic fibrosis. Hum.

[96413] Genet. 92: 627–628, 1993.

[96414]

[96415] 145. Romey, M.–C.; Desgeorges, M.; Laussel, M.; Durand, M.–F.; Demaille,

[96416] J.; Claustres, M.: Two novel rare frameshift mutations (2423 del

[96417] G in exon 13 and 1215 del G in exon 7) and one novel rare sequence

[96418] variation (3271 + 18 C or T) identified in a patient with cystic fibrosis. Hum.

[96419] Molec. Genet. 3: 1003–1004, 1994.

[96420]

[96421] 146. Romey, M.–C.; Guittard, C.; Carles, S.; Demaille, J.; Claustres,

[96422] M.; Ramsay, M.: First putative sequence alterations in the minimal

[96423] CFTR promoter region. (Letter) J. Med. Genet. 36: 263–264, 1999.

[96424]

[96425] 147. Romey, M.–C.; Guittard, C.; Chazalotte, J.–P.;

Frossard, P.;

[96426] Dawson, K. P.; Patton, M. A.; Casals, T.; Bazarbachi, T.;
Girodon,

[96427] E.; Rault, G.; Bozon, D.; Seguret, F.; Demaille, J.; Claustres,
M.

[96428] : Complex allele (-102T-to-A+S549R(T-to-G) is associ-
ated with milder

[96429] forms of cystic fibrosis than allele S549R(T-to-G) alone.
Hum. Genet. 105:

[96430] 145-150, 1999.

[96431]

[96432] 148. Romey, M.-C.; Pallares-Ruiz, N.; Mange, A.; Mettling,
C.; Peytavi,

[96433] R.; Demaille, J.; Claustres, M.: A naturally occurring se-
quence variation

[96434] that creates a YY1 element is associated with increased
cystic fibrosis

[96435] transmembrane conductance regulator gene expression. J.
Biol. Chem. 275:

[96436] 3561-3567, 2000.

[96437]

[96438] 149. Rommens, J. M.; Iannuzzi, M. C.; Kerem, B.; Drumm,

M. L.; Melmer,

[96439] G.; Dean, M.; Rozmahel, R.; Cole, J. L.; Kennedy, D.; Hidakaka, N.;

[96440] Zsiga, M.; Buchwald, M.; Riordan, J. R.; Tsui, L.-C.; Collins, F.

[96441] S.: Identification of the cystic fibrosis gene: chromosome walking

[96442] and jumping. Science 245: 1059–1065, 1989.

[96443]

[96444] 150. Ronchetto, P.; Telleria Orriols, J. J.; Fanen, P.; Cremonesi,

[96445] L.; Ferrari, M.; Magnani, C.; Seia, M.; Goossens, M.; Romeo, G.; Devoto,

[96446] M.: A nonsense mutation (R1158X) and a splicing mutation (3849 +

[96447] 4A-to-G) in exon 19 of the cystic fibrosis transmembrane conductance

[96448] regulator gene. Genomics 12: 417–418, 1992.

[96449]

[96450] 151. Rosenfeld, M. A.; Yoshimura, K.; Trapnell, B. C.; Yoneyama, K.;

[96451] Rosenthal, E. R.; Dalemans, W.; Fukayama, M.; Bargon, J.;

Stier, L.

[96452] E.; Stratford-Perricaudet, L.; Perricaudet, M.; Guggino, W. B.; Pavirani,

[96453] A.; Lecocq, J.-P.; Crystal, R. G.: In vivo transfer of the human

[96454] cystic fibrosis transmembrane conductance regulator gene to the airway

[96455] epithelium. Cell 68: 143-155, 1992.

[96456]

[96457] 152. Rowntree, R. K.; Vassaux, G.; McDowell, T. L.; Howe, S.; McGuigan,

[96458] A.; Phylactides, M.; Huxley, C.; Harris, A.: An element in intron

[96459] 1 of the CFTR gene augments intestinal expression in vivo. Hum. Molec.

[96460] Genet. 10: 1455-1464, 2001.

[96461]

[96462] 153. Rozen, R.; De Braekeleer, M.; Daigneault, J.; Ferreira-Rajabi,

[96463] L.; Gerdes, M.; Lamoureux, L.; Aubin, G.; Simard, F.; Fujiwara, T.

[96464] M.; Morgan, K.: Cystic fibrosis mutations in French Cana-

dians: three

[96465] CFTR mutations are relatively frequent in a Quebec population with

[96466] an elevated incidence of cystic fibrosis. Am. J. Med. Genet. 42:

[96467] 360–364, 1992.

[96468]

[96469] 154. Rozen, R.; Ferreira–Rajabi, L.; Robb, L.; Colman, N.: L206W

[96470] mutation of the cystic fibrosis gene, relatively frequent in French

[96471] Canadians, is associated with atypical presentations of cystic fibrosis. Am.

[96472] J. Med. Genet. 57: 437–439, 1995.

[96473]

[96474] 155. Rozen, R.; Schwartz, R. H.; Hilman, B. C.; Stanislovis, P.;

[96475] Horn, G. T.; Klinger, K.; Daigneault, J.; De Braekeleer, M.; Kerem,

[96476] B.; Tsui, L.–C.; Fujiwara, T. M.; Morgan, K.: Cystic fibrosis mutations

[96477] in North American populations of French ancestry: analy–

sis of Quebec

[96478] French-Canadian and Louisiana Acadian families. Am. J. Hum. Genet. 47:

[96479] 606-610, 1990.

[96480]

[96481] 156. Russo, M. P.; Romeo, G.; Devoto, M.; Barbujani, G.; Cabrini,

[96482] G.; Giunta, A.; D'Alcamo, E.; Leoni, G.; Sangiuolo, F.; Magnani, C.;

[96483] Cremonesi, L.; Ferrari, M.: Analysis of linkage disequilibrium between

[96484] different cystic fibrosis mutations and three intragenic microsatellites

[96485] in the Italian population. Hum. Mutat. 5: 23-27, 1995.

[96486]

[96487] 157. Sangiuolo, F.; Cicero, S. L.; Maceratesi, P.; Quattrucci, S.;

[96488] Novelli, G.; Dallapiccola, B.: Molecular characterization of a frameshift

[96489] mutation in exon 19 of the CFTR gene. Hum. Mutat. 2: 422-424, 1993.

[96490]

[96491] 158. Sangiuolo, F.; Novelli, G.; Murru, S.; Dallapiccola, B.:
A serine-to-arginine

[96492] (AGT-to-CGT) mutation in codon 549 of the CFTR gene in
an Italian

[96493] patient with severe cystic fibrosis. Genomics 9: 788–789,
1991.

[96494]

[96495] 159. Sato, S.; Ward, C. L.; Kopito, R. R.: Cotranslational
ubiquitination

[96496] of cystic fibrosis transmembrane conductance regulator in
vitro. J.

[96497] Biol. Chem. 273: 7189–7192, 1998.

[96498]

[96499] 160. Sato, S.; Ward, C. L.; Krouse, M. E.; Wine, J. J.; Kopito,
R.

[96500] R.: Glycerol reverses the misfolding phenotype of the
most common

[96501] cystic fibrosis mutation. J. Biol. Chem. 271: 635–638,
1996.

[96502]

[96503] 161. Schaedel, C.; Kristoffersson, A.-C.; Kornfalt, R.;
Holmberg,

- [96504] L.: A novel cystic fibrosis mutation, Y109C, in the first transmembrane
- [96505] domain of CFTR. Hum. Molec. Genet. 3: 1001–1002, 1994.
- [96506]
- [96507] 162. Schwartz, M.; Anvret, M.; Claustres, M.; Geir Eiken, H.; Eiklid,
- [96508] K.; Schaedel, C.; Stolpe, L.; Tranebjaerg, L.: 394delTT: a Nordic
- [96509] cystic fibrosis mutation. Hum. Genet. 93: 157–161, 1994.
- [96510]
- [96511] 163. Schwiebert, E. M.; Egan, M. E.; Hwang, T.–H.; Fulmer, S. B.;
- [96512] Allen, S. S.; Cutting, G. R.; Guggino, W. B.: CFTR regulates outwardly
- [96513] rectifying chloride channels through an autocrine mechanism involving
- [96514] ATP. Cell 81: 1063–1073, 1995.
- [96515]
- [96516] 164. Scotet, V.; De Braekeleer, M.; Audrezet, M.–P.; Lode, L.; Verlingue,

- [96517] C.; Quere, I.; Mercier, B.; Dugueperoux, I.; Codet, J.-P.;
Moineau,
[96518] M.-P.; Parent, P.; Ferec, C.: Prevalence of CFTR mutations
in hypertrypsinaemia
[96519] detected through neonatal screening for cystic fibrosis.
Clin. Genet. 59:
[96520] 42–47, 2001.
[96521]
[96522] 165. Sharer, N.; Schwarz, M.; Malone, G.; Howarth, A.;
Painter, J.;
[96523] Super, M.; Braganza, J.: Mutations of the cystic fibrosis
gene in
[96524] patients with chronic pancreatitis. New Eng. J. Med. 339:
645–652,
[96525] 1998.
[96526]
[96527] 166. Sheppard, D. N.; Rich, D. P.; Ostedgaard, L. S.; Gre-
gory, R.
[96528] J.; Smith, A. E.; Welsh, M. J.: Mutations in CFTR associated
with
[96529] mild–disease form CI– channels with altered pore proper-
ties. Nature 362:

[96530] 160–164, 1993.

[96531]

[96532] 167. Shoshani, T.; Augarten, A.; Gazit, E.; Bashan, N.; Yahav, Y.;

[96533] Rivlin, Y.; Tal, A.; Seret, H.; Yaar, L.; Kerem, E.; Kerem, B.:
Association

[96534] of a nonsense mutation (W1282X), the most common mutation in the Ashkenazi

[96535] Jewish cystic fibrosis patients in Israel, with presentation
of severe

[96536] disease. Am. J. Hum. Genet. 50: 222–228, 1992.

[96537]

[96538] 168. Shoshani, T.; Augarten, A.; Yahav, J.; Gazit, E.;
Kerem, B.:

[96539] Two novel mutations in the CFTR gene: W1089X in exon
17B and 4010delTATT

[96540] in exon 21. Hum. Molec. Genet. 3: 657–658, 1994.

[96541]

[96542] 169. Shoshani, T.; Berkun, Y.; Yahav, Y.; Augarten, A.;
Bashan, N.;

[96543] Rivlin, Y.; Gazit, E.; Sereth, H.; Kerem, E.; Kerem, B.: A new

mutation

[96544] in the CFTR gene, composed of two adjacent DNA alterations, is a common

[96545] cause of cystic fibrosis among Georgian Jews. Genomics 15: 236–237,

[96546] 1993.

[96547]

[96548] 170. Shoshani, T.; Kerem, E.; Szeinberg, A.; Augarten, A.; Yahav,

[96549] Y.; Cohen, D.; Rivlin, J.; Tal, A.; Kerem, B. S.: Similar levels

[96550] of mRNA from the W1282X and the delta-F508 cystic fibrosis alleles,

[96551] in nasal epithelial cells. J. Clin. Invest. 93: 1502–1507, 1994.

[96552]

[96553] 171. Siegel, D.; Irving, N. G.; Friedman, J. M.; Wainwright, B. J.

[96554] : Localization of the cystic fibrosis transmembrane conductance regulator

[96555] (Cftr) to mouse chromosome 6. Cytogenet. Cell Genet. 61: 184–185,

[96556] 1992.

[96557]

[96558] 172. Smit, L. S.; Nasr, S. Z.; Iannuzzi, M. C.; Collins, F. S.:

An

[96559] African-American cystic fibrosis patient homozygous for a novel frameshift

[96560] mutation associated with reduced CFTR mRNA levels.

Hum. Mutat. 2:

[96561] 148–151, 1993.

[96562]

[96563] 173. Smit, L. S.; Strong, T. V.; Wilkinson, D. J.; Macek, M., Jr.;

[96564] Mansoura, M. K.; Wood, D. L.; Cole, J. L.; Cutting, G. R.; Cohn, J.

[96565] A.; Dawson, D. C.; Collins, F. S.: Missense mutation (G480C) in the

[96566] CFTR gene associated with protein mislocalization but normal chloride

[96567] channel activity. Hum. Molec. Genet. 4: 269–273, 1995.

[96568]

[96569] 174. Snouwaert, J. N.; Brigman, K. K.; Latour, A. M.; Malouf, N. N.;

[96570] Boucher, R. C.; Smithies, O.; Koller, B. H.: An animal model
for

[96571] cystic fibrosis made by gene targeting. Science 257:
1083–1088,

[96572] 1992.

[96573]

[96574] 175. Stern, R. C.: The diagnosis of cystic fibrosis. New
Eng. J.

[96575] Med. 336: 487–491, 1997.

[96576]

[96577] 176. Strong, T. V.; Smit, L. S.; Nasr, S.; Wood, D. L.; Cole,
J. L.;

[96578] Iannuzzi, M. C.; Stern, R. C.; Collins, F. S.: Characterization
of

[96579] an intron 12 splice donor mutation in the cystic fibrosis
transmembrane

[96580] conductance regulator (CFTR) gene. Hum. Mutat. 1:
380–387, 1992.

[96581]

[96582] 177. Strong, T. V.; Smit, L. S.; Turpin, S. V.; Cole, J. L.;
Tom Hon,

- [96583] C.; Markiewicz, D.; Petty, T. L.; Craig, M. W.; Rosenow, E. C., III;
- [96584] Tsui, L.-C.; Iannuzzi, M. C.; Knowles, M. R.; Collins, F. S.:
Cystic
fibrosis gene mutation in two sisters with mild disease
and normal
sweat electrolyte levels. New Eng. J. Med. 325:
1630-1634, 1991.
- [96587]
- [96588] 178. Tata, F.; Stanier, P.; Wicking, C.; Halford, S.; Kruyer, H.;
- [96589] Lench, N. J.; Scambler, P. J.; Hansen, C.; Braman, J. C.;
Williamson,
R.; Wainwright, B. J.: Cloning the mouse homolog of the
human cystic
fibrosis transmembrane conductance regulator gene. Ge-
nomics 10:
301-307, 1991.
- [96593]
- [96594] 179. Teng, H.; Jorissen, M.; Van Poppel, H.; Legius, E.;
Cassiman,
J.-J.; Cuppens, H.: Increased proportion of exon 9 alterna-

tively

[96596] spliced CFTR transcripts in vas deferens compared with nasal epithelial

[96597] cells. Hum. Molec. Genet. 6: 85–90, 1997.

[96598]

[96599] 180. The Cystic Fibrosis Genotype–Phenotype Consortium: Correlation

[96600] between genotype and phenotype in patients with cystic fibrosis. New

[96601] Eng. J. Med. 329: 1308–1313, 1993.

[96602]

[96603] 181. Trapnell, B. C.; Chu, C.-S.; Paakko, P. K.; Banks, T. C.; Yoshimura,

[96604] K.; Ferrans, V. J.; Chernick, M. S.; Crystal, R. G.: Expression of

[96605] the cystic fibrosis transmembrane conductance regulator gene in the

[96606] respiratory tract of normal individuals and individuals with cystic

[96607] fibrosis. Proc. Nat. Acad. Sci. 88: 6565–6569, 1991.

[96608]

- [96609] 182. Trezise, A. E. O.; Szpirer, C.; Buchwald, M.: Localization of
- [96610] the gene encoding the cystic fibrosis transmembrane conductance regulator
- [96611] (CFTR) in the rat to chromosome 4 and implications for the evolution
- [96612] of mammalian chromosomes. *Genomics* 14: 869–874, 1992.
- [96613]
- [96614] 183. Van Doorninck, J. H.; French, P. J.; Verbeek, E.; Peters, R.
- [96615] H. P. C.; Morreau, H.; Bijman, J.; Scholte, B. J.: A mouse model
- [96616] for the cystic fibrosis delta-F508 mutation. *EMBO J.* 14: 4403–4411,
- [96617] 1995.
- [96618]
- [96619] 184. Varon, R.; Magdorf, K.; Staab, D.; Wahn, H.–U.; Krawczak, M.;
- [96620] Sperling, K.; Reis, A.: Recurrent nasal polyps as a monosymptomatic
- [96621] form of cystic fibrosis associated with a novel in-frame

deletion

[96622] (591del18) in the CFTR gene. Hum. Molec. Genet. 4: 1463–1464, 1995.

[96623]

[96624] 185. Varon, R.; Stuhmann, M.; Macek, M., Jr.; Kufardjieva, A.; Angelicheva,

[96625] D.; Magdorf, K.; Jordanova, A.; Savov, A.; Wahn, U.; Macek, M.; Lalov,

[96626] V.; Ivanova, T.; Ellemunter, H.; Vavrova, V.; Ferak, V.; Kayserova,

[96627] H.; Reis, A.; Kalaydjieva, L.: Pancreatic insufficiency and pulmonary

[96628] disease in German and Slavic cystic fibrosis patients with the R347P

[96629] mutation. Hum. Mutat. 6: 219–225, 1995.

[96630]

[96631] 186. Verlingue, C.; Mercier, B.; Lecoq, I.; Audrezet, M. P.; Laroche,

[96632] D.; Travert, G.; Ferec, C.: Retrospective study of the cystic fibrosis

[96633] transmembrane conductance regulator (CFTR) gene mutations in Guthrie

- [96634] cards from a large cohort of neonatal screening for cystic fibrosis. Hum.
- [96635] Genet. 93: 429–434, 1994.
- [96636]
- [96637] 187. Vidaud, M.; Fanen, P.; Martin, J.; Ghanem, N.; Nicolas, S.; Goossens, M.: Three point mutations in the CFTR gene in French cystic fibrosis patients: identification by denaturing gradient gel electrophoresis. Hum.
- [96639] Genet. 85: 446–449, 1990.
- [96640]
- [96641]
- [96642] 188. Vidaud, M.; Gattoni, R.; Stevenin, J.; Vidaud, D.; Amselem, S.; Chibani, J.; Rosa, J.; Goossens, M.: A 5-prime splice-region G-to-C mutation in exon 1 of the human beta-globin gene inhibits pre-mRNA splicing: a mechanism for beta(+)-thalassemia. Proc. Nat. Acad. Sci. 86: 1041–1045, 1989.
- [96646]

[96647]

[96648] 189. Wang, J.; Bowman, M. C.; Hsu, E.; Wertz, K.; Wong, L.-J. C.:

[96649] A novel mutation in the CFTR gene correlates with severe clinical

[96650] phenotype in seven Hispanic patients. J. Med. Genet. 37: 215-218,

[96651] 2000.

[96652]

[96653] 190. Wang, S.; Yue, H.; Derin, R. B.; Guggino, W. B.; Li, M.: Accessory

[96654] protein facilitated CFTR-CFTR interaction, a molecular mechanism to

[96655] potentiate the chloride channel activity. Cell 103: 169-179, 2000.

[96656]

[96657] 191. Waterston, R.; Martin, C.; Craxton, M.; Huynh, C.; Coulson, A.;

[96658] Hillier, L.; Durbin, R.; Green, P.; Shownkeen, R.; Halloran, N.; Metzstein,

[96659] M.; Hawkins, T.; Wilson, R.; Berks, M.; Du, Z.; Thomas, K.; Thierry-Mieg,

[96660] J.; Sulston, J.: A survey of expressed genes in *Caenorhabditis elegans*. *Nature*

[96661] *Genet.* 1: 114–123, 1992.

[96662]

[96663] 192. Wauters, J. G.; Hendrickx, J.; Coucke, P.; Vits, L.; Stuer, K.;

[96664] van Schil, L.; van der Auwera, B. J.; Van Elsen, A.; Dumon, J.; Willems,

[96665] P. J.: Frequency of the phenylalanine deletion (delta-F508) in the

[96666] CF gene of Belgian cystic fibrosis patients. *Clin. Genet.* 39: 89–92,

[96667] 1991.

[96668]

[96669] 193. Wei, L.; Vankeerberghen, A.; Cuppens, H.; Droogmans, G.; Cassiman,

[96670] J.-J.; Nilius, B.: Phosphorylation site independent single R-domain

[96671] mutations affect CFTR channel activity. *FEBS Lett.* 439: 121–126,

[96672] 1998.

[96673]

[96674] 194. Weixel, K. M.; Bradbury, N. A.: The carboxyl terminus of the

[96675] cystic fibrosis transmembrane conductance regulator binds to AP-2

[96676] clathrin adaptors. J. Biol. Chem. 275: 3655-3660, 2000.

[96677]

[96678] 195. White, M. B.; Amos, J.; Hsu, J. M. C.; Gerrard, B.; Finn, P.;

[96679] Dean, M.: A frame shift mutation in the cystic fibrosis gene. Nature 344:

[96680] 655-667, 1990.

[96681]

[96682] 196. White, M. B.; Leppert, M.; Nielsen, D.; Zielenski, J.; Gerrard,

[96683] B.; Stewart, C.; Dean, M.: A de novo cystic fibrosis mutation: CGA

[96684] (arg) to TGA (stop) at codon 851 of the CFTR gene. Genomics 11:

[96685] 778-779, 1991.

[96686]

- [96687] 197. White, S. M.; Lucassen, A.; Norbury, G.: Cystic fibrosis: a
[96688] further case of an asymptomatic compound heterozygote.
(Letter) Am.
[96689] J. Med. Genet. 103: 342–343, 2001.
[96690]
[96691] 198. Will, K.; Dork, T.; Stuhrmann, M.; Meitinger, T.;
Bertele-Harms,
[96692] R.; Tummler, B.; Schmidtke, J.: A novel exon in the cystic
fibrosis
[96693] transmembrane conductance regulator gene activated by
the nonsense
[96694] mutation E92X in airway epithelial cells of patients with
cystic fibrosis. J.
[96695] Clin. Invest. 93: 1852–1859, 1994.
[96696]
[96697] 199. Yang, Y.; Devor, D. C.; Engelhardt, J. F.; Ernst, S. A.;
Strong,
[96698] T. V.; Collins, F. S.; Cohn, J. A.; Frizzell, R. A.; Wilson, J. M.
[96699] : Molecular basis of defective anion transport in L cells
expressing

- [96700] recombinant forms of CFTR. Hum. Molec. Genet. 2: 1253–1261, 1993.
- [96701]
- [96702] 200. Yang, Y.; Janich, S.; Cohn, J. A.; Wilson, J. M.: The common
- [96703] variant of cystic fibrosis transmembrane conductance regulator is
- [96704] recognized by hsp70 and degraded in a pre-Golgi nonlysosomal compartment. Proc.
- [96705] Nat. Acad. Sci. 90: 9480–9484, 1993.
- [96706]
- [96707] 201. Yang, Y.; Raper, S. E.; Cohn, J. A.; Engelhardt, J. F.; Wilson,
- [96708] J. M.: An approach for treating the hepatobiliary disease of cystic
- [96709] fibrosis by somatic gene transfer. Proc. Nat. Acad. Sci. 90: 4601–4605,
- [96710] 1993.
- [96711]
- [96712] 202. Youil, R.; Kemper, B. W.; Cotton, R. G. H.: Screening for mutations

- [96713] by enzyme mismatch cleavage with T4 endonuclease VII.
Proc. Nat.
- [96714] Acad. Sci. 92: 87–91, 1995.
- [96715]
- [96716] 203. Zeiher, B. G.; Eichwald, E.; Zabner, J.; Smith, J. J.;
Puga,
- [96717] A. P.; McCray, P. B., Jr.; Capecchi, M. R.; Welsh, M. J.;
Thomas,
- [96718] K. R.: A mouse model for the delta-F508 allele of cystic fi-
brosis. J.
- [96719] Clin. Invest. 96: 2051–2064, 1995.
- [96720]
- [96721] 204. Zeitlin, P. L.; Crawford, I.; Lu, L.; Woel, S.; Cohen, M.
E.;
- [96722] Donowitz, M.; Montrose, M. H.; Hamosh, A.; Cutting, G. R.;
Gruenert,
- [96723] D.; Haganir, R.; Maloney, P.; Guggino, W. B.: CFTR protein
expression
- [96724] in primary and cultured epithelia. Proc. Nat. Acad. Sci. 89:
344–347,
- [96725] 1992.

[96726]

[96727] 205. Zhang, F.; Kartner, N.; Lukacs, G. L.: Limited proteolysis as

[96728] a probe for arrested conformational maturation of deltaF508 CFTR. Nature

[96729] Struct. Biol. 5: 180–183, 1998.

[96730]

[96731] 206. Zielenski, J.; Bozon, D.; Kerem, B.; Markiewicz, D.; Durie, P.;

[96732] Rommens, J. M.; Tsui, L.-C.: Identification of mutations in exons

[96733] 1 through 8 of the cystic fibrosis transmembrane conductance regulator

[96734] (CFTR) gene. Genomics 10: 229–235, 1991.

[96735]

[96736] 207. Zielenski, J.; Markiewicz, D.; Rininsland, F.; Rommens, J.; Tsui,

[96737] L.-C.: A cluster of highly polymorphic dinucleotide repeats in intron

[96738] 17b of the cystic fibrosis transmembrane conductance regulator (CFTR)

[96739] gene. Am. J. Hum. Genet. 49: 1256–1262, 1991.

[96740]

[96741] 208. Zielenski, J.; Patrizio, P.; Corey, M.; Handelin, B.;
Markiewicz,

[96742] D.; Asch, R.; Tsui, L.-C.: CFTR gene variant for patients
with congenital

[96743] absence of vas deferens. (Letter) Am. J. Hum. Genet. 57:
958–960,

[96744] 1995.

[96745] 16774.

[96746] Martin, F.; Schaller, A.; Eglite, S.; Schumperli, D.; Muller, B.

[96747] : The gene for histone RNA hairpin binding protein is lo-
cated on human

[96748] chromosome 4 and encodes a novel type of RNA binding
protein. EMBO

[96749] J. 16: 769–778, 1997.

[96750] 16775.

[96751] Wang, Z.-F.; Whitfield, M. L.; Ingledue, T. C., III; Dominski,

[96752] Z.; Marzluff, W. F.: The protein that binds the 3–prime end
of histone

- [96753] mRNA: a novel RNA-binding protein required for histone pre-mRNA processing. *Genes Dev.* 10: 3028-3040, 1996.
- [96754] 16776.
- [96755] Kooy, J.; Toh, B.-H.; Pettitt, J. M.; Erlich, R.; Gleeson, P. A.
- [96757] : Human autoantibodies as reagents to conserved Golgi components:
- [96758] characterization of a peripheral, 230-kDa compartment-specific Golgi
- [96759] protein. *J. Biol. Chem.* 267: 20255-20263, 1992.
- [96760] 16777.
- [96761] Marneros, A. G.; Mehenni, H.; Reichenberger, E.; Antonarakis, S.
- [96762] E.; Krieg, T.; Olsen, B. R.: Gene for the human transmembrane-type
- [96763] protein tyrosine phosphatase H (PTPRH): genomic structure, fine-mapping
- [96764] and its exclusion as a candidate for Peutz-Jeghers syndrome. *Cytogenet.*
- [96765] *Cell Genet.* 92: 213-216, 2001.
- [96766] 16778.

- [96767] Matozaki, T.; Suzuki, T.; Uchida, T.; Inazawa, J.; Ariyama, T.;
- [96768] Matsuda, K.; Horita, K.; Noguchi, H.; Mizuno, H.; Sakamoto, C.; Kasuga,
- [96769] M.: Molecular cloning of a human transmembrane-type protein tyrosine
- [96770] phosphatase and its expression in gastrointestinal cancers. J. Biol.
- [96771] Chem. 269: 2075–2081, 1994.
- [96772] 16779.
- [96773] Seo, Y.; Matozaki, T.; Tsuda, M.; Hayashi, Y.; Itoh, H.; Kasuga,
- [96774] M.: Overexpression of SAP-1, a transmembrane-type protein tyrosine
- [96775] phosphatase, in human colorectal cancers. Biochem. Biophys. Res.
- [96776] Commun. 231: 705–711, 1997.
- [96777] 16780.
- [96778] Snow, B. E.; Antonio, L.; Suggs, S.; Gutstein, H. B.; Siderovski,
- [96779] D. P.: Molecular cloning and expression analysis of rat Rgs12 and

- [96780] Rgs14. Biochem. Biophys. Res. Commun. 233: 770–777, 1997.
- [96781] 16781.
- [96782] Kimple, R. J.; Kimple, M. E.; Betts, L.; Sondek, J.; Siderovski,
- [96783] D. P.: Structural determinants for GoLoco-induced inhibition of nucleotide
- [96784] release by G- α subunits. Nature 416: 878–881, 2002.
- [96785] 16782.
- [96786] Takesono, A.; et al; et al: Receptor-independent activators of
- [96787] heterotrimeric G-protein signalling pathways. J. Biol. Chem. 274:
- [96788] 33202–33205, 1999.
- [96789] 16783.
- [96790] Traver, S.; Bidot, C.; Spassky, N.; Baltauss, T.; de Tand, M.-F.;
- [96791] Thomas, J.-L.; Zalc, B.; Janoueix-Lerosey, I.; de Gunzburg, J.: RGS14
- [96792] is a novel Rap effector that preferentially regulates the GTPase activity

[96793] of G- α -0. Biochem. J. 350: 19-29, 2000.

[96794] 16784.

[96795] Zheng, B.; Chen, D.; Farquhar, M. G.: MIR16, a putative membrane

[96796] glycerophosphodiester phosphodiesterase, interacts with RGS16. Proc.

[96797] Nat. Acad. Sci. 97: 3999-4004, 2000.

[96798] 16785.

[96799] Schmitz, F.; Konigstorfer, A.; Sudhof, T. C.: RIBEYE, a component

[96800] of synaptic ribbons: a protein's journey through evolution provides

[96801] insight into synaptic ribbon function. Neuron 28: 857-872, 2000.

[96802] 16786.

[96803] Dale, M.; Hammond, D. W.; Cox, A.; Nicklin, M. J. H.: The human

[96804] gene encoding the interleukin-1 receptor accessory protein (IL1RAP)

[96805] maps to chromosome 3q28 by fluorescence in situ hybridization and

- [96806] radiation hybrid mapping. Genomics 47: 325–326, 1998.
- [96807] 16787.
- [96808] Huang, J.; Gao, X.; Li, S.; Cao, Z.: Recruitment of IRAK to the
- [96809] interleukin 1 receptor complex requires interleukin 1 re-
ceptor accessory
- [96810] protein. Proc. Nat. Acad. Sci. 94: 12829–12832, 1997.
- [96811] 16788.
- [96812] Wesche, H.; Korherr, C.; Kracht, M.; Falk, W.; Resch, K.;
Martin,
- [96813] M. U.: The interleukin–1 receptor accessory protein
(IL–1RAcP) is
- [96814] essential for IL–1–induced activation of interleukin–1 re-
ceptor–associated
- [96815] kinase (IRAK) and stress–activated protein kinases (SAP ki-
nases). J.
- [96816] Biol. Chem. 272: 7727–7731, 1997.
- [96817] 16789.
- [96818] Saha, P.; Chen, J.; Thome, K. C.; Lawlis, S. J.; Hou, Z.–H.;
Hendricks,
- [96819] M.; Parvin, J. D.; Dutta, A.: Human CDC6/Cdc18 associates

with Orc1

[96820] and cyclin-cdk and is selectively eliminated from the nucleus at the

- [96821] onset of S phase. *Molec. Cell. Biol.* 18: 2758–2767, 1998.
- [96822] 16790.
- [96823] Williams, R. S.; Shohet, R. V.; Stillman, B.: A human protein
- [96824] related to yeast Cdc6p. *Proc. Nat. Acad. Sci.* 94: 142–147, 1997.
- [96825] 16791.
- [96826] Yan, Z.; DeGregori, J.; Shohet, R.; Leone, G.; Stillman, B.; Nevins,
- [96827] J. R.; Williams, R. S.: Cdc6 is regulated by E2F and is essential
- [96828] for DNA replication in mammalian cells. *Proc. Nat. Acad. Sci.* 95:
- [96829] 3603–3608, 1998.
- [96830] 16792.
- [96831] Brais, B.; Bouchard, J.-P.; Xie, Y.-G.; Rochefort, D. L.; Chretien,
- [96832] N.; Tome, F. M. S.; Lafreniere, R. G.; Rommens, J. M.; Uyama, E.;
- [96833] Nohira, O.; Blumen, S.; Korczyn, A. D.; Heutink, P.; Mathieu, J.;
- [96834] Duranceau, A.; Codere, F.; Fardeau, M.; Rouleau, G. A.:

Short GCG

- [96835] expansions in the PABP2 gene cause oculopharyngeal muscular dystrophy. Nature
- [96836] Genet. 18: 164–167, 1998. Note: Erratum: Nature Genet. 19: 404 only,
- [96837] 1998.
- [96838] 16793.
- [96839] Calado, A.; Tome, F. M. S.; Brais, B.; Rouleau, G. A.; Kuhn, U.;
- [96840] Wahle, E.; Carmo-Fonseca, M.: Nuclear inclusions in oculopharyngeal
- [96841] muscular dystrophy consist of poly(A) binding protein 2 aggregates
- [96842] which sequester poly(A) RNA. Hum. Molec. Genet. 9: 2321–2328, 2000.
- [96843] 16794.
- [96844] Fan, X.; Dion, P.; Laganier, J.; Brais, B.; Rouleau, G. A.: Oligomerization
- [96845] of polyalanine expanded PABPN1 facilitates nuclear protein aggregation
- [96846] that is associated with cell death. Hum. Molec. Genet. 10: 2341–2351,

- [96847] 2001.
- [96848] 16795.
- [96849] Ikeda, S.; Shiva, N.; Ikeda, A.; Smith, R. S.; Nusinowitz, S.;
- [96850] Yan, G.; Lin, T. R.; Chu, S.; Heckenlively, J. R.; North, M. A.;
Naggert,
- [96851] J. K.; Nishina, P. M.; Duyao, M. P.: Retinal degeneration but
not
- [96852] obesity is observed in null mutants of the tubby-like pro-
tein 1 gene. Hum.
- [96853] Molec. Genet. 9: 155–163, 2000.
- [96854] 16796.
- [96855] North, M. A.; Naggert, J. K.; Yan, Y.; Noben-Trauth, K.;
- Nishina,
- [96856] P. M.: Molecular characterization of TUB, TULP1, and
TULP2, members
- [96857] of the novel tubby gene family and their possible relation
to ocular
- [96858] diseases. Proc. Nat. Acad. Sci. 94: 3128–3133, 1997.
- [96859] 16797.
- [96860] Paloma, E.; Hjelmqvist, L.; Bayes, M.; Garcia-Sandoval, B.;
- Ayuso,

- [96861] C.; Balcells, S.; Gonzalez-Duarte, R.: Novel mutations in the TULP1
- [96862] gene causing autosomal recessive retinitis pigmentosa. Invest. Ophthalm.
- [96863] Vis. Sci. 41: 656–659, 2000.
- [96864] 16798.
- [96865] Aoki, N.; Ishii, T.; Ohira, S.; Yamaguchi, Y.; Negi, M.; Adachi,
- [96866] T.; Nakamura, R.; Matsuda, T.: Stage specific expression of milk
- [96867] fat globule membrane glycoproteins in mouse mammary gland: comparison
- [96868] of MFG-E8, butyrophilin, and CD36 with a major milk protein, beta-casein. Biochim.
- [96869] Biophys. Acta 1334: 182–190, 1997.
- [96870] 16799.
- [96871] Collins, C.; Nehlin, J. O.; Stubbs, J. D.; Kowbel, D.; Kuo, W.-L.;
- [96872] Parry, G.: Mapping of a newly discovered human gene homologous to
- [96873] the apoptosis associated-murine mammary protein, MFG-E8, to chromosome

- [96874] 15q25. Genomics 39: 117–118, 1997.
- [96875] 16800.
- [96876] Haggqvist, B.; Naslund, J.; Sletten, K.; Westermark, G. T.;
Mucchiano,
- [96877] G.; Tjernberg, L. O.; Nordstedt, C.; Engstrom, U.; Wester-
mark, P.
- [96878] : Medin: an integral fragment of aortic smooth muscle
cell-produced
- [96879] lactadherin forms the most common human amyloid.
Proc. Nat. Acad.
- [96880] Sci. 96: 8669–8674, 1999.
- [96881] 16801.
- [96882] Hanayama, R.; Tanaka, M.; Miwa, K.; Shinohara, A.; Iwa-
matsu, A.;
- [96883] Nagata, S.: Identification of a factor that links apoptotic
cells
- [96884] to phagocytes. Nature 417: 182–187, 2002.
- [96885] 16802.
- [96886] Larocca, D.; Peterson, J. A.; Urrea, R.; Kuniyoshi, J.; Bis-
train,
- [96887] A. M.; Ceriani, R. L.: A M(r) 46,000 human milk fat globule

protein

[96888] that is highly expressed in human breast tumors contains factor VIII-like

[96889] domains. *Cancer Res.* 51: 4994–4998, 1991.

[96890] 16803.

[96891] Stubbs, J. D.; Lekutis, C.; Singer, K. L.; Bui, A.; Yuzuki, D.;

[96892] Srinivasan, U.; Parry, G.: cDNA cloning of a mouse mammary epithelial

[96893] cell surface protein reveals the existence of epidermal growth factor-like

[96894] domains linked to factor VIII-like sequences. *Proc. Nat. Acad. Sci.* 87:

[96895] 8417–8421, 1990.

[96896] 16804.

[96897] Zhang, J.; Kuehl, P.; Green, E. D.; Touchman, J. W.; Watkins, P.

[96898] B.; Daly, A.; Hall, S. D.; Maurel, P.; Relling, M.; Brimer, C.; Yasuda,

[96899] K.; Wrighton, S. A.; Hancock, M.; Kim, R. B.; Strom, S.; Thummel,

[96900] K.; Russell, C. G.; Hudson, J. R., Jr.; Schuetz, E. G.; Bo-

guski, M.

- [96901] S.: The human pregnane X receptor: genomic structure and identification
- [96902] and functional characterization of natural allelic variants. Pharmacogenetics 11:
- [96903] 555–572, 2001.
- [96904] 16805.
- [96905] Berrueta, L.; Kraeft, S.-K.; Tirnauer, J. S.; Schuyler, S. C.;
- [96906] Chen, L. B.; Hill, D. E.; Pellman, D.; Bierer, B. E.: The adenomatous
- [96907] polyposis coli-binding protein EB1 is associated with cytoplasmic
- [96908] and spindle microtubules. Proc. Nat. Acad. Sci. 95: 10596–10601,
- [96909] 1998.
- [96910] 16806.
- [96911] Berrueta, L.; Tirnauer, J. S.; Schuyler, S. C.; Pellman, D.; Bierer,
- [96912] B. E.: The APC-associated protein EB1 associates with components
- [96913] of the dynactin complex and cytoplasmic dynein interme-

diate chain. Curr.

[96914] Biol. 9: 425–428, 1999.

[96915] 16807.

[96916] Su, L.-K.; Burrell, M.; Hill, D. E.; Gyuris, J.; Brent, R.; Wiltshire,

[96917] R.; Trent, J.; Vogelstein, B.; Kinzler, K. W.: APC binds to the novel

[96918] protein EB1. Cancer Res. 55: 2972–2977, 1995.

[96919] 16808.

[96920] Su, L.-K.; Qi, Y.: Characterization of human MAPRE genes and their

[96921] proteins. Genomics 71: 143–149, 2001.

[96922] 16809.

[96923] Tirnauer, J. S.; Bierer, B. E.: EB1 proteins regulate microtubule

[96924] dynamics, cell polarity, and chromosome stability. J. Cell Biol. 149:

[96925] 761–766, 2000.

[96926] 16810.

[96927] Arai, T.; Akiyama, Y.; Okabe, S.; Ando, M.; Endo, M.;

Yuasa, Y.

[96928] : Genomic structure of the human Smad3 gene and its infrequent alterations

[96929] in colorectal cancers. Cancer Lett. 122: 157–163, 1998.

[96930] 16811.

[96931] Zhang, Y.; Feng, X.-H.; Wu, R.-Y.; Derynck, R.: Receptor-associated

[96932] Mad homologues synergize as effectors of the TGF- β response. Nature 383:

[96933] 168–172, 1996.

[96934] 16812.

[96935] Zhu, Y.; Richardson, J. A.; Parada, L. F.; Graff, J. M.: Smad3

[96936] mutant mice develop metastatic colorectal cancer. Cell 94: 703–714,

[96937] 1998.

[96938] 16813.

[96939] Wicki, R.; Marenholz, I.; Mischke, D.; Schafer, B. W.; Heizmann,

[96940] C. W.: Characterization of the human S100A12 (calgranulin C, p6,

[96941] CAAF1, CGRP) gene, a new member of the S100 gene

cluster on chromosome

[96942] 1q21. Cell Calcium 20: 459–464, 1996.

[96943] 16814.

[96944] Tan, K. B.; Harrop, J.; Reddy, M.; Young, P.; Terrett, J.;
Emery,

[96945] J.; Moore, G.; Truneh, A.: Characterization of a novel TNF-
like ligand

[96946] and recently described TNF ligand and TNF receptor su-
perfamily genes

[96947] and their constitutive and inducible expression in
hematopoietic and

[96948] non-hematopoietic cells. Gene 204: 35–46, 1997.

[96949] 16815.

[96950] Gruter, P.; Tabernero, C.; von Kobbe, C.; Schmitt, C.;
Saavedra,

[96951] C.; Bachi, A.; Wilm, M.; Felber, B. K.; Izaurralde, E.: TAP,
the

[96952] human homolog of Mex67p, mediates CTE-dependent
RNA export from the

[96953] nucleus. Molec. Cell 1: 649–659, 1998.

[96954] 16816.

- [96955] Segref, A.; Sharma, K.; Doye, V.; Hellwig, A.; Huber, J.;
Luhrmann,
[96956] R.; Hurt, E.: Mex67p, a novel factor for nuclear mRNA ex-
port, binds
[96957] to both poly(A)+ RNA and nuclear pores. EMBO J. 16:
3256–3271, 1997.
[96958] 16817.
- [96959] Yoon, D.-W.; Lee, H.; Seol, W.; DeMaria, M.; Rosenzweig,
M.; Jung,
[96960] J. U.: Tap: a novel cellular protein that interacts with Tip of
herpesvirus
[96961] saimiri and induces lymphocyte aggregation. Immunity 6:
571–582,
[96962] 1997.
[96963] 16818.
- [96964] Bonini, J. A.; Martin, S. K.; Dralyuk, F.; Roe, M. W.; Philip-
son,
[96965] L. H.; Steiner, D. F.: Cloning, expression, and chromoso-
mal mapping
[96966] of a novel human CC-chemokine receptor (CCR10) that
displays high-affinity
[96967] binding for MCP-1 and MCP-3. DNA Cell Biol. 16:

1249–1256, 1997.

[96968] 16819.

[96969] Nibbs, R. J. B.; Wylie, S. M.; Yang, J.; Landau, N. R.; Graham,

[96970] G. J.: Cloning and characterization of a novel promiscuous human

[96971] beta-chemokine receptor D6. *J. Biol. Chem.* 272: 32078–32083, 1997.

[96972] 16820.

[96973] Karet, F. E.; Finberg, K. E.; Nayir, A.; Bakkaloglu, A.; Ozen,

[96974] S.; Hulton, S. A.; Sanjad, S. A.; Al-Sabban, E. A.; Medina, J. F.;

[96975] Lifton, R. P.: Localization of a gene for autosomal recessive distal

[96976] renal tubular acidosis with normal hearing (rdRTA2) to 7q33–34. *Am.*

[96977] *J. Hum. Genet.* 65: 1656–1665, 1999.

[96978] 16821.

[96979] Smith, A. N.; Skaug, J.; Choate, K. A.; Nayir, A.; Bakkaloglu,

[96980] A.; Ozen, S.; Hulton, S. A.; Sanjad, S. A.; Al-Sabban, E. A.;

Lifton,

- [96981] R. P.; Scherer, S. W.; Karet, F. E.: Mutations in ATP6N1B, encoding
[96982] a new kidney vacuolar proton pump 116-kD subunit, cause recessive
[96983] distal renal tubular acidosis with preserved hearing. Nature Genet. 26:
[96984] 71–75, 2000.
[96985] 16822.
- [96986] McKie, J. M.; Sutherland, H. F.; Harvey, E.; Kim, U.-J.; Scambler,
[96987] P. J.: A human gene similar to Drosophila melanogaster peanut maps
[96988] to the DiGeorge syndrome region of 22q11. Hum. Genet. 101: 6–12,
[96989] 1997.
[96990] 16823.
- [96991] Yagi, M.; Zieger, B.; Roth, G. J.; Ware, J.: Structure and expression
[96992] of the human septin gene HCDCREL-1. Gene 212: 229–236, 1998.
[96993] 16824.

- [96994] Zieger, B.; Tran, H.; Hainmann, I.; Wunderle, D.; Zgaga-Griesz,
[96995] A.; Blaser, S.; Ware, J.: Characterization and expression
analysis
[96996] of two human septin genes, PNUTL1 and PNUTL2. Gene
261: 197–203,
[96997] 2000.
[96998] 16825.
- [96999] Latif, F.; Duh, F.–M.; Bader, S.; Sekido, Y.; Li, H.; Geil, L.;
[97000] Zbar, B.; Minna, J. D.; Lerman, M. I.: The human homolog
of the rodent
[97001] immediate early response genes, PC4 and TIS7, resides in
the lung
[97002] cancer tumor suppressor gene region on chromosome
3p21. Hum. Genet. 99:
[97003] 334–341, 1997.
[97004] 16826.
- [97005] Brandt, S.; Jentsch, T. J.: CIC–6 and CIC–7 are two novel
broadly
[97006] expressed members of the CLC chloride channel family.
FEBS Lett. 377:

[97007] 15–20, 1995.

[97008] 16827.

[97009] Eggermont, J.; Buyse, G.; Voets, T.; Tytgat, J.; De Smedt, H.;

[97010] Droogmans, G.: Alternative splicing of ClC-6 (a member of the ClC

[97011] chloride-channel family) transcripts generates three truncated isoforms

[97012] one of which, ClC-6c, is kidney-specific. *Biochem. J.* 325: 269–276,

[97013] 1997.

[97014] 16828.

[97015] Cleiren, E.; Benichou, O.; Van Hul, E.; Gram, J.; Bollerslav, J.;

[97016] Singer, F. R.; Beaverson, K.; Aledo, A.; Whyte, M. P.; Yoneyama, T.;

[97017] deVernejou, M.-C.; Van Hul, W.: Albers-Schonberg disease (autosomal

[97018] dominant osteopetrosis, type II) results from mutations in the CLCN7

[97019] chloride channel gene. *Hum. Molec. Genet.* 10: 2861–2867, 2001.

- [97020] 16829.
- [97021] Hedblom, E.; Kirkness, E. F.: A novel class of GABA-A receptor
subunit in tissues of the reproductive system. J. Biol.
Chem. 272:
15346-15350, 1997.
- [97022] 16830.
- [97023] Caloca, M. J.; Garcia-Bermejo, M. L.; Blumberg, P. M.;
Lewin, N.
- [97024] E.; Kremmer, E.; Mischak, H.; Wang, S.; Nacro, K.; Bienfait,
B.; Marquez,
- [97025] V. E.; Kazanietz, M. G.: Beta-2-chimaerin is a novel target
for diacylglycerol:
binding properties and changes in subcellular localization
mediated
by ligand binding to its C1 domain. Proc. Nat. Acad. Sci.
96: 11854-11859,
1999.
- [97026] 16831.
- [97027] Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Cerebellar beta-
2-chimaerin,

- [97033] a GTPase-activating protein for p21 Ras-related Rac is specifically
- [97034] expressed in granule cells and has a unique N-terminal SH2 domain J.
- [97035] Biol. Chem. 269: 12888–12892, 1994.
- [97036] 16832.
- [97037] Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Germ cell beta-chimaerin,
- [97038] a new GTPase-activating protein for p21rac, is specifically expressed
- [97039] during the acrosomal assembly stage in rat testis. J. Biol. Chem. 268:
- [97040] 3813–3816, 1993.
- [97041] 16833.
- [97042] Yuan, S.; Miller, D. W.; Barnett, G. H.; Hahn, J. F.; Williams,
- [97043] B. R.: Identification and characterization of human beta 2-chimaerin:
- [97044] association with malignant transformation in astrocytoma. Cancer
- [97045] Res. 55: 3456–3461, 1995.
- [97046] 16834.

- [97047] Holland, P. C.; Dillon, M. J.; Pincott, J.; Simmonds, H. A.; Barratt,
T. M.: Hypoxanthine guanine phosphoribosyl transferase deficiency
presenting with gout and renal failure in infancy. Arch. Dis. Child. 58:
831–833, 1983.
- [97051] 16835.
- [97052] Jolly, D. J.; Esty, A. C.; Bernard, H. U.; Friedmann, T.: Isolation
of a genomic clone partially encoding human hypoxanthine phosphoribosyltransferase. Proc. Nat. Acad. Sci. 79: 5038–5041, 1982.
- [97055] 16836.
- [97056] Jolly, D. J.; Okayama, H.; Berg, P.; Esty, A. C.; Filpula, D.; Bohlen, P.; Johnson, G. G.; Shively, J. E.; Hunkapiller, T.; Friedmann, T.: Isolation and characterization of a full-length expressible cDNA
for human hypoxanthine phosphoribosyltransferase. Proc. Nat. Acad.

- [97060] Sci. 80: 477–481, 1983.
- [97061] 16837.
- [97062] Chung, E.; Hanukoglu, A.; Rees, M.; Thompson, R.; Dillon, M.; Hanukoglu, I.; Bistritzer, T.; Kuhnle, U.; Seckl, J.; Gardiner, R. M.: Exclusion of the locus for autosomal recessive pseudohypoaldosteronism type 1 from the mineralocorticoid receptor gene region on human chromosome 4q by linkage analysis. J. Clin. Endocr. Metab. 80: 3341–3345, 1995.
- [97063] 16838.
- [97064] Schwienbacher, C.; Sabbioni, S.; Campi, M.; Veronese, A.; Bernardi, G.; Menegatti, A.; Hatada, I.; Mukai, T.; Ohashi, H.; Barbanti-Brodano, G.; Croce, C. M.; Negrini, M.: Transcriptional map of 170-kb region at chromosome 11p15.5: identification and mutational analysis of the BWR1A gene reveals the presence of mutations in tumor

samples. Proc.

[97073] Nat. Acad. Sci. 95: 3873–3878, 1998.

[97074] 16839.

[97075] Bartsch, O.; Kress, W.; Wagner, A.; Seemanova, E.: The novel contiguous

[97076] gene syndrome of myotubular myopathy (MTM1), male hypogenitalism and

[97077] deletion in Xq28: report of the first familial case. Cytogenet. Cell

[97078] Genet. 85: 310–314, 1999.

[97079] 16840.

[97080] Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hiro-sawa, M.;

[97081] Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction

[97082] of the coding sequences of unidentified human genes. XIII. The complete

[97083] sequences of 100 new cDNA clones from brain which code for large proteins

[97084] in vitro. DNA Res. 6: 63–70, 1999.

[97085] 16841.

- [97086] Philibert, R. A.; Winfield, S. L.; Sandhu, H. K.; Martin, B. M.;
- [97087] Ginns, E. I.: The structure and expression of the human
neurologin-3
- [97088] gene. *Gene* 246: 303-310, 2000.
- [97089] 16842.
- [97090] Zonana, J.; Clarke, A.; Sarfarazi, M.; Thomas, N. S. T.;
- Roberts,
- [97091] K.; Marymee, K.; Harper, P. S.: X-linked hypohidrotic ecto-
dermal
- [97092] dysplasia: localization within the region Xq11-21.1 by
linkage analysis
- [97093] and implications for carrier detection and prenatal diag-
nosis. *Am.*
- [97094] *J. Hum. Genet.* 43: 75-85, 1988.
- [97095] 16843.
- [97096] Arico, M.; Imashuku, S.; Clementi, R.; Hibi, S.; Teramura,
T.;
- [97097] Danesino, C.; Haber, D. A.; Nichols, K. E.: Hemophagocytic
lymphohistiocytosis
- [97098] due to germline mutations in SH2D1A, the X-linked lym-
phoproliferative

- [97099] disease gene. *Blood* 97: 1131–1133, 2001.
- [97100] 16844.
- [97101] Arkwright, P. D.; Makin, G.; Will, A. M.; Ayres, M.; Gokhale, D.
- [97102] A.; Fergusson, W. D.; Taylor, G. M.: X linked lymphoproliferative
- [97103] disease in a United Kingdom family. *Arch. Dis. Child.* 79: 52–55,
- [97104] 1998.
- [97105] 16845.
- [97106] Bar, R. S.; DeLor, C. J.; Clausen, K. P.; Hurtubise, P.; Henle, W.; Hewetson, J. F.: Fatal infectious mononucleosis in a family. *New*
- [97107] Eng. J. Med. 290: 363–367, 1974.
- [97108] 16846.
- [97109] Benoit, L.; Wang, X.; Pabst, H. F.; Dutz, J.; Tan, R.: Cutting
- [97110] edge: defective NK cell activation in X-linked lymphoproliferative
- [97111] disease. *J. Immun.* 165: 3549–3553, 2000.
- [97112] 16847.
- [97113]

- [97114] Brandau, O.; Schuster, V.; Weiss, M.; Hellebrand, H.; Fink, F.
- [97115] M.; Kreczy, A.; Friedrich, W.; Strahm, B.; Niemeyer, C.; Be-lohradsky,
- [97116] B. H.; Meindl, A.: Epstein-Barr virus-negative boys with non-Hodgkin
- [97117] lymphoma are mutated in the SH2D1A gene, as are patients with X-linked
- [97118] lymphoproliferative disease (XLP). Hum. Molec. Genet. 8: 2407-2413,
- [97119] 1999.
- [97120] 16848.
- [97121] Coffey, A. J.; Brooksbank, R. A.; Brandau, O.; Oohashi, T.; Howell,
- [97122] G. R.; Bye, J. M.; Cahn, A. P.; Durham, J.; Heath, P.; Wray, P.; Pavitt,
- [97123] R.; Wilkinson, J.; and 31 others: Host response to EBV infection
- [97124] in X-linked lymphoproliferative disease results from mutations in
- [97125] an SH2-domain encoding gene. Nature Genet. 20: 129-135, 1998.

- [97126] 16849.
- [97127] Czar, M. J.; Kersh, E. N.; Mijares, L. A.; Lanier, G.; Lewis, J.;
- [97128] Yap, G.; Chen, A.; Sher, A.; Duckett, C. S.; Ahmed, R.;
Schwartzberg,
- [97129] P. L.: Altered lymphocyte responses and cytokine produc-
tion in mice
- [97130] deficient in the X-linked lymphoproliferative disease gene
SH2D1A/DSHP/SAP. Proc.
- [97131] Nat. Acad. Sci. 98: 7449–7454, 2001.
- [97132] 16850.
- [97133] Dutz, J. P.; Benoit, L.; Wang, X.; Demetrick, D. J.; Junker,
A.;
- [97134] de Sa, D.; Tan, R.: Lymphocytic vasculitis in X-linked lym-
phoproliferative
- [97135] disease. Blood 97: 95–100, 2001.
- [97136] 16851.
- [97137] Grierson, H. L.; Skare, J.; Church, J.; Silberman, T.; Davis, J.
- [97138] R.; Kобрinsky, N.; McGregor, R.; Israels, S.; McCarty, J.;
Andrews,
- [97139] L. G.; Blecha, T.; Erdman, S.; Obringer, A.; Scharnhorst, D.;
Purtilo,

- [97140] D. T.: Evaluation of families wherein a single male manifests a phenotype
- [97141] of X-linked lymphoproliferative disease (XLP). Am. J. Med. Genet. 47:
- [97142] 458–463, 1993.
- [97143] 16852.
- [97144] Hambleton, G.; Cottom, D. G.: Familial lymphoma. Proc. Roy.
- [97145] Soc. Med. 62: 1095 only, 1969.
- [97146] 16853.
- [97147] Hamilton, J. K.; Paquin, L. A.; Sullivan, J. L.; Maurer, H. S.;
- [97148] Cruzei, F. G.; Provisor, A. J.; Steuber, C. P.; Hawkins, E.; Yawn,
- [97149] D.; Cornet, J.; Clausen, K.; Finkelstein, G. Z.; Landing, B.; Grunnet,
- [97150] M.; Purtilo, D. T.: X-linked lymphoproliferative syndrome registry
- [97151] report. J. Pediat. 96: 669–673, 1980.
- [97152] 16854.
- [97153] Harris, A.; Docherty, Z.: X-linked lymphoproliferative disease:

[97154] a karyotype analysis. Cytogenet. Cell Genet. 47: 92–94, 1988.

[97155] 16855.

[97156] Harris, A.; Lenoir, G. M.; Lankester, S. A.: X-linked lymphoproliferative

[97157] disease: linkage studies using DNA probes. Clin. Genet. 33: 162–168,

[97158] 1988.

[97159] 16856.

[97160] Hayoz, D.; Lenoir, G. M.; Nicole, A.; Pugin, P.; Regamey, C.:

[97161] X-linked lymphoproliferative syndrome: identification of a large family

[97162] in Switzerland. Am. J. Med. 84: 529–534, 1988.

[97163] 16857.

[97164] Klein, G.; Klein, E.: Sinking surveillance's flagship. Nature 395:

[97165] 441–445, 1998.

[97166] 16858.

[97167] Levine, P. H.; Kamaraju, L. S.; Connelly, R. R.; Berard, C. W.;

- [97168] Dorfman, R. F.; Magrath, I.; Easton, J. M.: The American Burkitt's
- [97169] Lymphoma Registry: eight years' experience. Cancer 49: 1016–1022,
- [97170] 1982.
- [97171] 16859.
- [97172] Loeffel, S.; Chang, C.-H.; Heyn, R.; Harada, S.; Lipscomb, H.;
- [97173] Sinangil, F.; Volsky, D. J.; McClain, K.; Ochs, H.; Purtilo, D. T.
- [97174] : Necrotizing lymphoid vasculitis in X-linked lymphoproliferative
- [97175] syndrome. Arch. Path. Lab. Med. 109: 546–550, 1985.
- [97176] 16860.
- [97177] Lyon, M. F.; Loutit, J. F.: X-linked factor in acquired immunodeficiency
- [97178] syndrome?. (Letter) Lancet I: 768 only, 1983.
- [97179] 16861.
- [97180] Mulley, J. C.; Turner, A. M.; Gedeon, A. K.; Berdoukas, V. A.;
- [97181] Huang, T. H. M.; Ledbetter, D. H.; Grierson, H.; Purtilo, D.

T.:

[97182] X-linked lymphoproliferative disease: prenatal detection
of an unaffected

[97183] histocompatible male. Clin. Genet. 42: 76-79, 1992.

[97184] 16862.

[97185] Eva, A.; Aaronson, S. A.: Isolation of a new human onco-
gene from

[97186] a diffuse B-cell lymphoma. Nature 316: 273-275, 1985.

[97187] 16863.

[97188] Anderson, S. R.; Warburg, M.: Norrie's disease. Arch. Oph-
thal. 66:

[97189] 614-618, 1961.

[97190] 16864.

[97191] Bergen, A. A. B.; Wapenaar, M. C.; Schuurman, E. J. M.;
Diergaarde,

[97192] P. J.; Lerach, H.; Monaco, A. P.; Bakker, E.; Bleeker-
Wagemakers,

[97193] E. M.; van Ommen, G. J. B.: Detection of a new submicro-
scopic Norrie

[97194] disease deletion interval with a novel DNA probe isolated
by differential

- [97195] Alu PCR fingerprint cloning. *Cytogenet. Cell Genet.* 62: 231–235,
- [97196] 1993.
- [97197] 16865.
- [97198] Berger, W.; Meindl, A.; van de Pol, T. J. R.; Cremers, F. P. M.;
- [97199] Ropers, H. H.; Doerner, C.; Monaco, A.; Bergen, A. A. B.; Lebo, R.;
- [97200] Warburg, M.; Zergollern, L.; Lorenz, B.; Gal, A.; Bleeker-Wagemakers,
- [97201] E. M.; Meitinger, T.: Isolation of a candidate gene for Norrie disease
- [97202] by positional cloning. *Nature Genet.* 1: 199–203, 1992.
- [97203] 16866.
- [97204] Berger, W.; van de Pol, D.; Bachner, D.; Oerlemans, F.; Winkens,
- [97205] H.; Hameister, H.; Wieringa, B.; Hendriks, W.; Ropers, H.–H.: An
- [97206] animal model for Norrie disease (ND): gene targeting of the mouse
- [97207] ND gene. *Hum. Molec. Genet.* 5: 51–59, 1996.

- [97208] 16867.
- [97209] Eva, A.; Vecchio, G.; Rao, C. D.; Tronick, S. R.; Aaronson, S.
- [97210] A.: The predicted DBL oncogene product defines a distinct class of
- [97211] transforming proteins. Proc. Nat. Acad. Sci. 85: 2061–2065, 1988.
- [97212] 16868.
- [97213] Galland, F.; Stefanova, M.; Lafage, M.; Birnbaum, D.: Localization
- [97214] of the 5-prime end of the MCF2 oncogene to human chromosome 15q15–q23.
- [97215] Cytogenet. Cell Genet. 60: 114–116, 1992.
- [97216] 16869.
- [97217] Grant, S. G.; Mattei, M.-G.; Galland, F.; Stephenson, D. A.; Keitz,
- [97218] B. T.; Birnbaum, D.; Chapman, V. M.: Localization of the mouse Mcf-2
- [97219] (Dbl) protooncogene within a conserved linkage group on the mouse
- [97220] X chromosome. Cytogenet. Cell Genet. 54: 175–181, 1990.

- [97221] 16870.
- [97222] Nguyen, C.; Pontarotti, P.; Birnbaum, D.; Chimini, G.; Rey, J.
- [97223] A.; Mattei, J.-F.; Jordan, B. R.: Large scale physical mapping in
- [97224] the q27 region of the human X chromosome: the coagulation factor IX
- [97225] gene and the mcf.2 transforming sequence are separated by at most
- [97226] 270 kilobase pairs and are surrounded by several 'HTF islands.' EMBO
- [97227] J. 6: 3285-3289, 1987.
- [97228] 16871.
- [97229] Nguyen, C.; Poustka, A.-M.; Djabali, M.; Roux, D.; Mattei, J.-F.;
- [97230] Lehrach, H.; Jordan, B. R.: Large-scale mapping and chromosome jumping
- [97231] in the q27 region of the human X chromosome. Genomics 5: 298-303,
- [97232] 1989.
- [97233] 16872.

- [97234] Noguchi, T.; Mattei, M.-G.; Oberle, I.; Planche, J.; Imbert, J.;
- [97235] Pelassy, C.; Birg, F.; Birnbaum, D.: Localization of the mcf.2 transforming
- [97236] sequence to the X chromosome. EMBO J. 6: 1301-1307, 1987.
- [97237] 16873.
- [97238] Ron, D.; Tronick, S. R.; Aaronson, S. A.; Eva, A.: Molecular cloning
- [97239] and characterization of the human DBL proto-oncogene: evidence that
- [97240] its overexpression is sufficient to transform NIH/3T3 cells. EMBO
- [97241] J. 7: 2465-2473, 1988.
- [97242] 16874.
- [97243] Srivastava, S. K.; Wheelock, R. H. P.; Aaronson, S. A.; Eva, A.
- [97244] : Identification of the protein encoded by the human dif-fuse B-cell
- [97245] lymphoma (dbl) oncogene. Proc. Nat. Acad. Sci. 83: 8868-8872, 1986.
- [97246] 16875.

- [97247] Tronick, S. R.; McBride, O. W.; Popescu, N. C.; Eva, A.:
Chromosomal
- [97248] localization of DBL oncogene sequences. Genomics 5:
546–553, 1989.
- [97249] 16876.
- [97250] Giovane, A.; Sobieszczuk, P.; Mignon, C.; Mattei, M.–G.;
Wasylyk,
- [97251] B.: Locations of the ets subfamily members net, elk1, and
sap1 (ELK3,
- [97252] ELK1, and ELK4) on three homologous regions of the
mouse and human
- [97253] genomes. Genomics 29: 769–772, 1995.
- [97254] 16877.
- [97255] Acland, G. M.; Blanton, S. H.; Hershfield, B.; Aguirre, G. D.:
- [97256] XLPRa: a canine retinal degeneration inherited as an X–
linked trait. Am.
- [97257] J. Med. Genet. 52: 27–33, 1994.
- [97258] 16878.
- [97259] Aldred, M. A.; Teague, P. W.; Jay, M.; Bunday, S.; Redmond,
R.
- [97260] M.; Jay, B.; Bird, A. C.; Bhattacharya, S. S.; Wright, A. F.:

Retinitis

- [97261] pigmentosa families showing apparent X linked inheritance but unlinked
- [97262] to the RP2 or RP3 loci. J. Med. Genet. 31: 848-852, 1994.
- [97263] 16879.
- [97264] Allan, W.: Eugenic significance of retinitis pigmentosa. Arch.
- [97265] Ophthal. 18: 938-947, 1937.
- [97266] 16880.
- [97267] Bhattacharya, S. S.; Clayton, J. F.; Harper, P. S.; Hoare, G. W.;
- [97268] Jay, M. R.; Lyness, A. L.; Wright, A. F.: A genetic linkage study
- [97269] of a kindred with X-linked retinitis pigmentosa. Brit. J. Ophthal. 69:
- [97270] 340-347, 1985.
- [97271] 16881.
- [97272] Bhattacharya, S. S.; Wright, A. F.; Clayton, J. F.; Price, W. H.;
- [97273] Phillips, C. I.; McKeown, C. M. E.; Jay, M.; Bird, A. C.; Pearson,

- [97274] P. L.; Southern, E. M.; Evans, H. J.: Close genetic linkage between
- [97275] X-linked retinitis pigmentosa and a restriction fragment length polymorphism
- [97276] identified by recombinant DNA probe L1.28. Nature 309: 253–255,
- [97277] 1984.
- [97278] 16882.
- [97279] Bird, A. C.: X-linked retinitis pigmentosa. Brit. J. Ophthal. 59:
- [97280] 177–199, 1975.
- [97281] 16883.
- [97282] Bunday, S.; Crews, S. J.: A study of retinitis pigmentosa in the
- [97283] City of Birmingham. (Letter) J. Med. Genet. 23: 188–191, 1986.
- [97284] 16884.
- [97285] Chapple, J. P.; Hardcastle, A. J.; Grayson, C.; Spackman, L. A.;
- [97286] Willison, K. R.; Cheetham, M. E.: Mutations in the N-terminus of

- [97287] the X-linked retinitis pigmentosa protein RP2 interfere with the normal
- [97288] targeting of the protein to the plasma membrane. Hum. Molec. Genet. 9:
- [97289] 1919–1926, 2000.
- [97290] 16885.
- [97291] Chen, J.-D.; Halliday, F.; Keith, G.; Sheffield, L.; Dickinson, P.; Gray, R.; Constable, I.; Denton, M.: Linkage heterogeneity between
- [97292] X-linked retinitis pigmentosa and a map of 10 RFLP loci. Am. J. Hum.
- [97293] Genet. 45: 401–411, 1989.
- [97294] 16886.
- [97295] Chen, J. D.; Denton, M. J.; Donald, J.; Serravalle, S.; Colley, P.; Lam Po Tang, P. R. L.; Halliday, F. B.: Analysis of linkage relationships
- [97296] of X-linked retinitis pigmentosa with the following Xp loci: DXS7,
- [97297] OTC, DXS84, XJ-1, DXS164, and DXS28. (Abstract) Cytogenet. Cell Genet. 46:
- [97300] 592 only, 1987.

- [97301] 16887.
- [97302] Chen, J. D.; Halliday, F.; Keith, G.; Sheffield, L.; Fraser, N.;
- [97303] Dickinson, P.; Gray, R.; Constable, I.; Denton, M. J.: Two
loci for
- [97304] X-linked retinitis pigmentosa suggested by multipoint
linkage analysis
- [97305] in nine families. (Abstract) Cytogenet. Cell Genet. 51: 976
only,
- [97306] 1989.
- [97307] 16888.
- [97308] Clayton, J. F.; Wright, A. F.; Jay, M.; McKeown, C. M. E.;
Dempster,
- [97309] M.; Jay, B. S.; Bird, A. C.; Bhattacharya, S. S.: Genetic link-
age
- [97310] between X-linked retinitis pigmentosa and DNA probe
DXS7 [L1.28]:
- [97311] further linkage data, heterogeneity testing, and risk esti-
mation. Hum.
- [97312] Genet. 74: 168-171, 1986.
- [97313] 16889.
- [97314] Coleman, M.; Bhattacharya, S.; Lindsay, S.; Wright, A.; Jay,

M.;

- [97315] Litt, M.; Craig, I.; Davies, K.: Localization of the mi-
crosatellite
- [97316] probe DXS426 between DXS7 and DXS255 on Xp and link-
age to X-linked
- [97317] retinitis pigmentosa. Am. J. Hum. Genet. 47: 935-940,
1990.
- [97318] 16890.
- [97319] Ernst, W.; Clover, G.; Faulkner, D. J.: X-linked retinitis pig-
mentosa:
- [97320] reduced rod flicker sensitivity in heterozygous females.
Invest.
- [97321] Ophthal. Vis. Sci. 6: 812-816, 1981.
- [97322] 16891.
- [97323] Farrar, G. J.; Geraghty, M. T.; Moloney, J. M. B.; McConnell,
- [97324] D. J.; Humphries, P.: Linkage analysis of X linked retinitis
pigmentosa
- [97325] in the Irish population. J. Med. Genet. 25: 222-226, 1988.
- [97326] 16892.
- [97327] Fishman, G. A.; Farber, M. D.; Derlacki, D. J.: X-linked re-
tinitis

- [97328] pigmentosa: profile of clinical findings. Arch. Ophthalm.
106: 369–375,
[97329] 1988.
[97330] 16893.
[97331] Friedrich, U.; Warburg, M.; Jorgensen, A. L.: X-inactivation
[97332] pattern in carriers of X-linked retinitis pigmentosa: a
valuable means
[97333] of prognostic evaluation?. Hum. Genet. 92: 359–363,
1993.
[97334] 16894.
[97335] Friedrich, U.; Warburg, M.; Kruse, T. A.; Andreasson, S.: X-
linked
[97336] retinitis pigmentosa: new map studies of XLRP2, and a
possible human
[97337] centromere effect. Hum. Genet. 88: 683–687, 1992.
[97338] 16895.
[97339] Friedrich, U.; Warburg, M.; Wieacker, P.; Wienker, T. F.;
Gal,
[97340] A.; Ropers, H.–H.: X-linked retinitis pigmentosa: linkage
with the
[97341] centromere and a cloned DNA sequence from the proxi-

mal short arm of

[97342] the X chromosome. Hum. Genet. 71: 93–99, 1985.

[97343] 16896.

[97344] Gieser, D. K.; Fishman, G. A.; Cunha-Vaz, J.: X-linked recessive

[97345] retinitis pigmentosa and vitreous fluorophotometry: a study of female

[97346] heterozygotes. Arch. Ophthal. 98: 307–310, 1980.

[97347] 16897.

[97348] Grutzner, P.; Sanger, R.; Spivey, B. E.: Linkage studies in X-linked

[97349] retinitis pigmentosa. Humangenetik 14: 155–158, 1972.

[97350] 16898.

[97351] Heck, A. F.: Presumptive X-linked intermediate transmission of

[97352] retinal degenerations: variations and coincidental occurrence with

[97353] ataxia in a large family. Arch. Ophthal. 70: 143–149, 1963.

[97354] 16899.

[97355] Hoare, G. W.: Choroido-retinal dystrophy. Brit. J. Ophthal.

49:

[97356] 449–459, 1965.

[97357] 16900.

[97358] Bermingham, J. R., Jr.; Scherer, S. S.; O'Connell, S.; Arroyo,

[97359] E.; Kalla, K. A.; Powell, F. L.; Rosenfeld, M. G.: Tst–

1/Oct–6/SCIP

[97360] regulates a unique step in peripheral myelination and is
required

[97361] for normal respiration. Genes Dev. 10: 1751–1762, 1996.

[97362] 16901.

[97363] Jaegle, M.; Mandemakers, W.; Broos, L.; Zwart, R.; Karis,
A.; Visser,

[97364] P.; Grosveld, F.; Meijer, D.: The POU factor Oct–6 and
Schwann cell

[97365] differentiation. Science 273: 507–510, 1996.

[97366] 16902.

[97367] Sumiyama, K.; Washio–Watanabe, K.; Ono, T.; Yoshida, M.
C.; Hayakawa,

[97368] T.; Ueda, S.: Human class III POU genes, POU3F1 and
POU3F3, map to

[97369] chromosomes 1p34.1 and 3p14.1. Mammalian Genome 9:

180-181, 1998.

[97370] 16903.

[97371] Costache, M.; Apoil, P.-A.; Cailleau, A.; Elmgren, A.; Larson,

[97372] G.; Henry, S.; Blancher, A.; Iordachescu, D.; Oriol, R.; Mollicone,

[97373] R.: Evolution of fucosyltransferase genes in vertebrates. J. Biol.

[97374] Chem. 272: 29721-29728, 1997.

[97375] 16904.

[97376] Yanagidani, S.; Uozumi, N.; Ihara, Y.; Miyoshi, E.; Yamaguchi,

[97377] N.; Taniguchi, N.: Purification and cDNA cloning of GDP-L-Fuc:N-acetyl-beta-D-glucosaminide:alpha-1-6

[97378] fucosyltransferase (alpha-1-6 FucT) from human gastric cancer MKN45

[97379] cells. J. Biochem. 121: 626-632, 1997.

[97380] 16905.

[97381] Bekri, S.; Adelaide, J.; Merscher, S.; Grosgeorge, J.; Caroli-Bosc,

[97382] F.; Perucca-Lostanlen, D.; Kelley, P. M.; Pebusque, M.-J.;

Theillet,

- [97383] C.; Birnbaum, D.; Gaudray, P.: Detailed map of a region commonly
[97384] amplified at 11q13-q14 in human breast carcinoma. Cy-
togenet. Cell
[97385] Genet. 79: 125-131, 1997.
[97386] 16906.
- [97387] Goodwin, R. G.; Alderson, M. R.; Smith, C. A.; Armitage, R.
J.;
- [97388] VandenBos, T.; Jerzy, R.; Tough, T. W.; Schoenborn, M. A.;
Davis-Smith,
[97389] T.; Hennen, K.; Falk, B.; Cosman, D.; Baker, E.; Sutherland,
G. R.;
- [97390] Grabstein, K. H.; Farrah, T.; Giri, J. G.; Beckmann, M. P.:
Molecular
[97391] and biological characterization of a ligand for CD27 de-
fines a new
[97392] family of cytokines with homology to tumor necrosis fac-
tor. Cell 73:
[97393] 447-456, 1993.
[97394] 16907.
- [97395] Kawabata, M.; Saeki, K.: Multiple alternative transcripts of

the

[97396] human homologue of the mouse TRAD/R51H3/RAD51D gene, a member of the

[97397] rec A/RAD51 gene family. Biochem. Biophys. Res. Commun. 257: 156–162,

[97398] 1999.

[97399] 16908.

[97400] Pittman, D. L.; Weinberg, L. R.; Schimenti, J. C.: Identification,

[97401] characterization, and genetic mapping of Rad51d, a new mouse and human

[97402] RAD51/RecA-related gene. Genomics 49: 103–111, 1998.

[97403] 16909.

[97404] Bell, B.; Scheer, E.; Tora, L.: Identification of hTAFII80-delta

[97405] links apoptotic signaling pathways to transcription factor TFIID function. Molec.

[97406] Cell 8: 591–600, 2001.

[97407] 16910.

[97408] Hisatake, K.; Ohta, T.; Takada, R.; Guermah, M.; Horikoshi, M.;

- [97409] Nakatani, Y.; Roeder, R. G.: Evolutionary conservation of human TATA-binding-polypeptide-associated
- [97410] factors TAFII31 and TAFII80 and interactions of TAFII80 with other
- [97411] TAFs and with general transcription factors. Proc. Nat. Acad. Sci. 92:
- [97412] 8195–8199, 1995.
- [97413] 16911.
- [97414] Weinzierl, R. O. J.; Ruppert, S.; Dynlacht, B. D.; Tanese, N.;
- [97415] Tjian, R.: Cloning and expression of Drosophila TAFII60 and human
- [97416] TAFII70 reveal conserved interactions with other subunits of TFIID. EMBO
- [97417] J. 12: 5303–5309, 1993.
- [97418] 16912.
- [97419] de Winter, J. P.; Waisfisz, Q.; Rooimans, M. A.; van Berkel, C.
- [97420] G. M.; Bosnoyan–Collins, L.; Alon, N.; Carreau, M.; Bender, O.; Demuth,
- [97421] I.; Schindler, D.; Pronk, J. C.; Arwert, F.; Hoehn, H.; Digweed, M.;

- [97422] Buchwald, M.; Joenje, H.: The Fanconi anaemia group G gene FANCG
- [97423] is identical with XRCC9. Nature Genet. 20: 281–283, 1998.
- [97424] 16913.
- [97425] Demuth, I.; Wlodarski, M.; Tipping, A. J.; Morgan, N. V.; de Winter,
- [97426] J. P.; Thiel, M.; Grasl, S.; Schindler, D.; D'Andrea, A. D.; Altay,
- [97427] C.; Kayserili, H.; Zatterdale, A.; Kunze, J.; Ebell, W.; Mathew, C.
- [97428] G.; Joenje, H.; Sperling, K.; Digweed, M.: Spectrum of mutations
- [97429] in the Fanconi anaemia group G gene, FANCG/XRCC9. Europ. J. Hum.
- [97430] Genet. 8: 861–868, 2000.
- [97431] 16914.
- [97432] Futaki, M.; Watanabe, S.; Kajigaya, S.; Liu, J. M.: Fanconi anemia
- [97433] protein, FANCG, is a phosphoprotein and is upregulated with FANCA
- [97434] after TNF- α treatment. Biochem. Biophys. Res. Com-

mun. 281: 347–351,

[97435] 2001.

[97436] 16915.

[97437] Koomen, M.; Cheng, N. C.; van de Vrugt, H. J.; Godthelp, B. C.;

[97438] van der Valk, M. A.; Oostra, A. B.; Zdzienicka, M. Z.; Joenje, H.;

[97439] Arwert, F.: Reduced fertility and hypersensitivity to mitomycin C

[97440] characterize Fancg/Xrcc9 null mice. Hum. Molec. Genet. 11: 273–281,

[97441] 2002.

[97442] 16916.

[97443] Kuang, Y.; Garcia-Higuera, I.; Moran, A.; Mondoux, M.; Digweed,

[97444] M.; D'Andrea, A. D.: Carboxy terminal region of the Fanconi anemia

[97445] protein, FANCG/XRCC9, is required for functional activity. Blood 96:

[97446] 1625–1632, 2000.

[97447] 16917.

- [97448] Liu, N.; Lamerdin, J. E.; Tucker, J. D.; Zhou, Z.-Q.; Walter, C.
- [97449] A.; Albala, J. S.; Busch, D. B.; Thompson, L. H.: The human XRCC9
- [97450] gene corrects chromosomal instability and mutagen sensitivities in
- [97451] CHO UV40 cells. *Proc. Nat. Acad. Sci.* 94: 9232–9237, 1997.
- [97452] 16918.
- [97453] Saar, K.; Schindler, D.; Wegner, R. D.; Reis, A.; Wienker, T. F.;
- [97454] Hoehn, H.; Joenje, H.; Sperling, K.; Digweed, M.: Localisation of
- [97455] a Fanconi anaemia gene to chromosome 9p. *Europ. J. Hum. Genet.* 6:
- [97456] 501–508, 1998.
- [97457] 16919.
- [97458] Yamada, T.; Tachibana, A.; Shimizu, T.; Mugishima, H.; Okubo, M.;
- [97459] Sasaki, M. S.: Novel mutations of the FANCG gene causing alternative
- [97460] splicing in Japanese Fanconi anemia. *J. Hum. Genet.* 45:

159–166,

[97461] 2000.

[97462] 16920.

[97463] Yang, Y.; Kuang, Y.; Montes De Oca, R.; Hays, T.; Moreau, L.;

[97464] Lu, N.; Seed, B.; D'Andrea, A. D.: Targeted disruption of the murine

[97465] Fanconi anemia gene, Fancg/Xrcc9. Blood 98: 3435–3440, 2001.

[97466] 16921.

[97467] Tsai, K. J.; Chen, S. K.; Ma, Y. L.; Hsu, W. L.; Lee, E. H. Y.

[97468] : sgk, a primary glucocorticoid-induced gene, facilitates memory consolidation

[97469] of spatial learning in rats. Proc. Nat. Acad. Sci. 99: 3990–3995,

[97470] 2002.

[97471] 16922.

[97472] Waldegger, S.; Barth, P.; Raber, G.; Lang, F.: Cloning and characterization

[97473] of a putative human serine/threonine protein kinase transcriptionally

- [97474] modified during anisotonic and isotonic alterations of cell volume. Proc.
- [97475] Nat. Acad. Sci. 94: 4440–4445, 1997.
- [97476] 16923.
- [97477] Lynch, R. A.; Piper, M.; Bankier, A.; Bhugra, B.; Surti, U.; Liu,
- [97478] J.; Buckler, A.; Dear, P. H.; Menon, A. G.: Genomic and functional
- [97479] map of the chromosome 14 t(12;14) breakpoint cluster region in uterine
- [97480] leiomyoma. Genomics 52: 17–26, 1998.
- [97481] 16924.
- [97482] Adam–Klages, S.; Adam, D.; Wiegmann, K.; Struve, S.; Kolanus, W.;
- [97483] Schneider–Mergener, J.; Kronke, M.: FAN, a novel WD–repeat protein,
- [97484] couples the p55 TNF–receptor to neutral sphingomyelinase. Cell 86:
- [97485] 937–947, 1996.
- [97486] 16925.
- [97487] Zhang, Y.; Kreder, D.; Schwandner, R.; Krut, O.; Scherer,

- G.; Adam-Klages,
[97488] S.; Siebert, R.; Kronke, M.; Schlegelberger, B.: Assignment
of the
[97489] human FAN protein gene (NSMAF) to human chromosome
region 8q12-q13
[97490] by in situ hybridization. Cytogenet. Cell Genet. 87:
115-116, 1999.
[97491] 16926.
- [97492] Ide, H.; Katoh, M.; Sasaki, H.; Yoshida, T.; Aoki, K.; Nawa,
Y.;
[97493] Osada, Y.; Sugimura, T.; Terada, M.: Cloning of human
bone morphogenetic
[97494] protein type 1B receptor (BMPR-1B) and its expression in
prostate
[97495] cancer in comparison with other BMPRs. Oncogene 14:
1377-1382, 1997.
[97496] 16927.
- [97497] ten Dijke, P.; Yamashita, H.; Ichijo, H.; Franzen, P.; Laiho,
M.;
[97498] Miyazono, K.; Heldin, C.-H.: Characterization of type I re-
ceptors
[97499] for transforming growth factor-beta and activin. Science

264: 101–104,

[97500] 1994.

[97501] 16928.

[97502] Mitsuhashi, N.; Miki, T.; Senbongi, H.; Yokoi, N.; Yano, H.; Miyazaki,

[97503] M.; Nakajima, N.; Iwanaga, T.; Yokoyama, Y.; Shibata, T.; Seino, S.

[97504] : MTABC3, a novel mitochondrial ATP-binding cassette protein involved

[97505] in iron homeostasis. J. Biol. Chem. 275: 17536–17540, 2000.

[97506] 16929.

[97507] Bai, C.; Connolly, B.; Metzker, M. L.; Hilliard, C. A.; Liu, X.;

[97508] Sandig, V.; Soderman, A.; Galloway, S. M.; Liu, Q.; Austin, C. P.;

[97509] Caskey, C. T.: Overexpression of M68/DCR3 in human gastrointestinal

[97510] tract tumors independent of gene amplification and its location in

[97511] a four-gene cluster. Proc. Nat. Acad. Sci. 97: 1230–1235, 2000.

[97512] 16930.

- [97513] Pitti, R. M.; Marsters, S. A.; Lawrence, D. A.; Roy, M.; Kischkel,
- [97514] F. C.; Dowd, P.; Huang, A.; Donahue, C. J.; Sherwood, S. W.; Baldwin,
- [97515] D. T.; Godowski, P. J.; Wood, W. I.; Gurney, A. L.; Hillan, K. J.;
- [97516] Cohen, R. L.; Goddard, A. D.; Botstein, D.; Ashkenazi, A.: Genomic
- [97517] amplification of a decoy receptor for Fas ligand in lung and colon
- [97518] cancer. *Nature* 396: 699–703, 1998.
- [97519] 16931.
- [97520] Ringstad, N.; Nemoto, Y.; De Camilli, P.: The SH3p4/Sh3p8/SH3p13
- [97521] protein family: binding partners for synaptojanin and dynamin via
- [97522] a Grb2-like Src homology 3 domain. *Proc. Nat. Acad. Sci.* 94: 8569–8574,
- [97523] 1997.
- [97524] 16932.
- [97525] Sittler, A.; Walter, S.; Wedemeyer, N.; Hasenbank, R.; Scherzinger,

- [97526] E.; Eickhoff, H.; Bates, G. P.; Lehrach, H.; Wanker, E. E.:
SH3GL3
- [97527] associates with the huntington exon 1 protein and pro-
motes the formation
- [97528] of polyglIn-containing protein aggregates. Molec. Cell 2:
427-436,
- [97529] 1998.
- [97530] 16933.
- [97531] Sparks, A. B.; Hoffman, N. G.; McConnell, S. J.; Fowlkes, D.
M.;
- [97532] Kay, B. K.: Cloning of ligand targets: systematic isolation
of SH3
- [97533] domain-containing proteins. Nature Biotech. 14:
741-744, 1996.
- [97534] 16934.
- [97535] Fracchiolla, N. S.; Colombo, G.; Finelli, P.; Maiolo, A. T.;
- Neri,
- [97536] A.: EHT, a new member of the MTG8/ETO gene family,
maps on 20q11
- [97537] region and is deleted in acute myeloid leukemias. (Letter)
Blood 92:
- [97538] 3481-3484, 1998.

- [97539] 16935.
- [97540] Kitabayashi, I.; Ida, K.; Morohoshi, F.; Yokoyama, A.; Mit-
suhashi,
- [97541] N.; Shimizu, K.; Nomura, N.; Hayashi, Y.; Ohki, M.: The
AML1-MTG8
- [97542] leukemic fusion protein forms a complex with a novel
member of the
- [97543] MTG8(ETO/CDR) family, MTGR1. Molec. Cell. Biol. 18:
846-858, 1998.
- [97544] 16936.
- [97545] Jensen, D. E.; Proctor, M.; Marquis, S. T.; Gardner, H. P.;
Ha,
- [97546] S. I.; Chodosh, L. A.; Ishov, A. M.; Tommerup, N.; Vissing,
H.; Sekido,
- [97547] Y.; Minna, J.; Borodovsky, A.; Schultz, D. C.; Wilkinson, K.
D.; Maul,
- [97548] G. G.; Barlev, N.; Berger, S. L.; Prendergast, G. C.;
Rauscher, F.
- [97549] J., III: BAP1: a novel ubiquitin hydrolase which binds to the
BRCA1
- [97550] RING finger and enhances BRCA1-mediated cell growth
suppression. Oncogene 16:

[97551] 1097–1112, 1998.

[97552] 16937.

[97553] Nelson, H.; Mandiyan, S.; Noumi, T.; Moriyama, Y.; Miedel, M. C.;

[97554] Nelson, N.: Molecular cloning of cDNA encoding the C subunit of H(+)-ATPase

[97555] from bovine chromaffin granules. J. Biol. Chem. 265: 20390–20393,

[97556] 1990.

[97557] 16938.

[97558] van Hille, B.; Vanek, M.; Richener, H.; Green, J. R.; Bilbe, G.

[97559] : Cloning and tissue distribution of subunits C, D, and E of the human

[97560] vacuolar H(+)-ATPase. Biochem. Biophys. Res. Commun. 197: 15–21,

[97561] 1993.

[97562] 16939.

[97563] Yasuda, T.; Yamaguchi, N.; Kobayashi, K.; Nishi, I.; Hori-nouchi,

[97564] H.; Jalil, M. A.; Li, M. X.; Ushikai, M.; Iijima, M.; Kondo, I.; Saheki,

- [97565] T.: Identification of two novel mutations in the SLC25A13 gene and
- [97566] detection of seven mutations in 102 patients with adult-onset type
- [97567] II citrullinemia. Hum. Genet. 107: 537–545, 2000.
- [97568] 16940.
- [97569] Kondoh, N.; Schweinfest, C. W.; Henderson, K. W.; Papas, T. S.
- [97570] : Differential expression of S19 ribosomal protein, laminin-binding
- [97571] protein, and human lymphocyte antigen class I messenger RNAs associated
- [97572] with colon carcinoma progression and differentiation. Cancer Res. 52:
- [97573] 791–796, 1992.
- [97574] 16941.
- [97575] Matsson, H.; Klar, J.; Draptchinskaia, N.; Gustavsson, P.; Carlsson,
- [97576] B.; Bowers, D.; de Bont, E.; Dahl, N.: Truncating ribosomal protein
- [97577] S19 mutations and variable clinical expression in Diamond-Blackfan

- [97578] anemia. Hum. Genet. 105: 496–500, 1999.
- [97579] 16942.
- [97580] Tentler, D.; Gustavsson, P.; Elinder, G.; Eklof, O.; Gordon, L.;
- [97581] Mandel, A.; Dahl, N.: A microdeletion in 19q13.2 associated with
- [97582] mental retardation, skeletal malformations, and Diamond–Blackfan anaemia
- [97583] suggests a novel contiguous gene syndrome. J. Med. Genet. 37: 128–131,
- [97584] 2000.
- [97585] 16943.
- [97586] Hoornaert, I.; Marynen, P.; Baens, M.: CREBL2, a novel transcript
- [97587] from the chromosome 12 region flanked by ETV6 and CDKN1B. Genomics 51:
- [97588] 154–157, 1998.
- [97589] 16944.
- [97590] Nishiu, J.; Tanaka, T.; Nakamura, Y.: Identification of a novel
- [97591] gene (ECM2) encoding a putative extracellular matrix pro-

tein expressed

[97592] predominantly in adipose and female-specific tissues and its chromosomal

[97593] localization to 9q22.3. Genomics 52: 378-381, 1998.

[97594] 16945.

[97595] Anderson, P.; Nagler-Anderson, C.; O'Brien, C.; Levine, H.; Watkins,

[97596] S.; Slayter, H. S.; Blue, M. L.; Schlossman, S. F.: A monoclonal

[97597] antibody reactive with a 15-kDa cytoplasmic granule-associated protein

[97598] defines a subpopulation of CD8+ T lymphocytes. J. Immun. 144: 574-582,

[97599] 1990.

[97600] 16946.

[97601] Forch, P.; Puig, O.; Kedersha, N.; Martinez, C.; Granneman, S.;

[97602] Seraphin, B.; Anderson, P.; Valcarcel, J.: The apoptosis-promoting

[97603] factor TIA-1 is a regulator of alternative pre-mRNA splicing. Molec.

[97604] Cell 6: 1089-1098, 2000.

- [97605] 16947.
- [97606] Kawakami, A.; Tian, Q.; Streuli, M.; Poe, M.; Edelhoff, S.;
Disteche,
- [97607] C. M.; Anderson, P.: Intron-exon organization and chro-
mosomal localization
- [97608] of the human TIA-1 gene. J. Immun. 152: 4937-4945,
1994.
- [97609] 16948.
- [97610] Tian, Q.; Streuli, M.; Saito, H.; Schlossman, S. F.; Ander-
son,
- [97611] P.: A polyadenylate binding protein localized to the gran-
ules of
- [97612] cytolytic lymphocytes induces DNA fragmentation in tar-
get cells. Cell 67:
- [97613] 629-639, 1991.
- [97614] 16949.
- [97615] Neubauer, G.; King, A.; Rappsilber, J.; Calvio, C.; Watson,
M.;
- [97616] Ajuh, P.; Sleeman, J.; Lamond, A.; Mann, M.: Mass spec-
trometry and
- [97617] EST-database searching allows characterisation of the
multi-protein

- [97618] spliceosome complex. *Nature Genet.* 20: 46–50, 1998.
- [97619] 16950.
- [97620] Talbot, K.; Miguel–Aliaga, I.; Mohaghegh, P.; Ponting, C. P.; Davies, P.; Davies, K. E.: Characterization of a gene encoding survival motor neuron (SMN)–related protein, a constituent of the spliceosome complex. *Hum. Molec. Genet.* 7: 2149–2156, 1998.
- [97624] 16951.
- [97625] Sillekens, P. T. G.; Beijer, R. P.; Habets, W. J.; van Venrooij, W. J.: Molecular cloning of the cDNA for the human U2 snRNA–specific A–prime protein. *Nucleic Acids Res.* 17: 1893–1906, 1989.
- [97628] 16952.
- [97629] Zheng, B.; Albrecht, U.; Kaasik, K.; Sage, M.; Lu, W.; Vaishnav, S.; Li, Q.; Sun, Z. S.; Eichele, G.; Bradley, A.; Lee, C. C.: Nonredundant roles of the mPer1 and mPer2 genes in the mammalian

circadian clock. Cell 105:

[97632] 683–694, 2001.

[97633] 16953.

[97634] Mattei, M.–G.; Roeckel, N.; Olsen, B. R.; Apte, S. S.: Genes
of

[97635] the membrane–type matrix metalloproteinase (MT–MMP)
gene family, MMP14,

[97636] MMP15, and MMP16, localize to human chromosomes 14,
16, and 8, respectively. Genomics 40:

[97637] 168–169, 1997.

[97638] 16954.

[97639] Sato, H.; Tanaka, M.; Takino, T.; Inoue, M.; Seiki, M.: As-
signment

[97640] of the human genes for membrane–type–1, –2, and –3
matrix metalloproteinases

[97641] (MMP14, MMP15, and MMP16) to 14q12.2, 16q12.2–q21,
and 8q21, respectively,

[97642] by in situ hybridization. Genomics 39: 412–413, 1997.

[97643] 16955.

[97644] Takino, T.; Sato, H.; Shinagawa, A.; Seiki, M.: Identification

[97645] of the second membrane–type matrix metalloproteinase

(MT-MMP-2) gene

[97646] from a human placenta cDNA library: MT-MMPs form a unique membrane-type

[97647] subclass in the MMP family. *J. Biol. Chem.* 270: 23013-23020, 1995.

[97648] 16956.

[97649] Will, H.; Hinzmann, B.: cDNA sequence and mRNA tissue distribution

[97650] of a novel human matrix metalloproteinase with a potential transmembrane

[97651] segment. *Europ. J. Biochem.* 231: 602-608, 1995.

[97652] 16957.

[97653] Alliel, P. M.; Perin, J.-P.; Jolles, P.; Bonnet, F. J.: Testican,

[97654] a multidomain testicular proteoglycan resembling modulators of cell

[97655] social behaviour. *Europ. J. Biochem.* 214: 347-350, 1993.

[97656] 16958.

[97657] Charbonnier, F.; Perin, J.-P.; Mattei, M.-G.; Camuzat, A.; Bonnet,

[97658] F.; Gressin, L.; Alliel, P. M.: Genomic organization of the human

- [97659] SPOCK gene and its chromosomal localization to 5q31.
Genomics 48:
[97660] 377–380, 1998.
- [97661] 16959.
- [97662] Cheng, J.-F.; Boyartchuk, V.; Zhu, Y.: Isolation and mapping of
[97663] human chromosome 21 cDNA: progress in constructing a
chromosome 21
[97664] expression map. Genomics 23: 75–84, 1994.
- [97665] 16960.
- [97666] Sirotkin, H.; O'Donnell, H.; DasGupta, R.; Halford, S.; St.
Jore,
[97667] B.; Puech, A.; Parimoo, S.; Morrow, B.; Skoultchi, A.;
Weissman, S.
[97668] M.; Scambler, P.; Kucherlapati, R.: Identification of a new
human
[97669] catenin gene family member (ARVCF) from the region
deleted in velo–cardio–facial
[97670] syndrome. Genomics 41: 75–83, 1997.
- [97671] 16961.
- [97672] Andrews, N. C.: Mining copper transport genes. Proc. Nat.

Acad.

[97673] Sci. 98: 6543–6545, 2001.

[97674] 16962.

[97675] Boultonwood, J.; Strickson, A. J.; Jabs, E. W.; Cheng, J.-F.; Fidler,

[97676] C.; Wainscoat, J. S.: Physical mapping of the human ATX1
homologue

[97677] (HAH1) to the critical region of the 5q- syndrome within
5q32, and

[97678] immediately adjacent to the SPARC gene. Hum. Genet.
106: 127–129,

[97679] 2000.

[97680] 16963.

[97681] Hamza, I.; Faisst, A.; Prohaska, J.; Chen, J.; Gruss, P.;
Gitlin,

[97682] J. D.: The metallochaperone Atox1 plays a critical role in
perinatal

[97683] copper homeostasis. Proc. Nat. Acad. Sci. 98: 6848–6852,
2001.

[97684] 16964.

[97685] Klomp, L. W. J.; Lin, S.-J.; Yuan, D. S.; Klausner, R. D.; Cu-

lotta,

[97686] V. C.; Gitlin, J. D.: Identification and functional expression of

[97687] HAH1, a novel human gene involved in copper homeostasis. J. Biol.

[97688] Chem. 272: 9221–9226, 1997.

[97689] 16965.

[97690] Castro, A. M.; Snyder, L. M.: G6PD San Jose: a new variant characterized

[97691] by NADPH inhibition studies. Humangenetik 21: 361–363, 1974.

[97692] 16966.

[97693] Cayanis, E.; Gomperts, E. D.; Balinsky, D.; Disler, P.; Myers,

[97694] A.: G6PD Hillbrow: a new variant of glucose–6–phosphate dehydrogenase

[97695] associated with drug–induced haemolytic anaemia. Brit. J. Haemat. 30:

[97696] 343–350, 1975.

[97697] 16967.

[97698] Cayanis, E.; Lane, A. B.; Jenkins, T.; Nurse, G. T.; Balinsky,

[97699] D.: Glucose–6–phosphate dehydrogenase Porbandar: a

new slow variant

[97700] with slightly reduced activity in a South African family of Indian

[97701] descent. *Biochem. Genet.* 15: 765–773, 1977.

[97702] 16968.

[97703] Cederbaum, A. I.; Beutler, E.: Nonspherocytic hemolytic anemia

[97704] due to G6PD Grand Prairie. *I.R.C.S.* 3: 579, 1975.

[97705] 16969.

[97706] Chan, T. K.; Todd, D.: Characteristics and distribution of glucose–6–phosphate

[97707] dehydrogenase–deficient variants in South China. *Am. J. Hum. Genet.* 24:

[97708] 475–484, 1972.

[97709] 16970.

[97710] Chan, T. K.; Todd, D.; Lai, M. C. S.: Glucose 6–phosphate dehydrogenase:

[97711] identity of erythrocyte and leukocyte enzyme with report of a new

[97712] variant in Chinese. *Biochem. Genet.* 6: 119–124, 1972.

[97713] 16971.

- [97714] Chang, J.-G.; Chiou, S.-S.; Perng, L.-I.; Chen, T.-C.; Liu, T.-C.;
- [97715] Lee, L.-S.; Chen, P.-H.; Tang, T. K.: Molecular characterization
- [97716] of glucose-6-phosphate dehydrogenase (G6PD) deficiency by natural
- [97717] and amplification created restriction sites: five mutations account
- [97718] for most G6PD deficiency cases in Taiwan. Blood 80: 1079-1082, 1992.
- [97719] 16972.
- [97720] Chen, E. Y.; Cheng, A.; Lee, A.; Kuang, W.-J.; Hillier, L.; Green,
- [97721] P.; Schlessinger, D.; Ciccodicola, A.; D'Urso, M.: Sequence of human
- [97722] glucose-6-phosphate dehydrogenase cloned in plasmids and a yeast artificial
- [97723] chromosome. Genomics 10: 792-800, 1991.
- [97724] 16973.
- [97725] Chen, H.-L.; Huang, M.-J.; Huang, C.-S.; Tang, T. K.: G6PD NanKang
- [97726] (517 T-to-C; 173 phe-to-leu): a new Chinese G6PD vari-

ant associated

[97727] with neonatal jaundice. Hum. Hered. 46: 201–204, 1996.

[97728] 16974.

[97729] Chernyak, N. B.; Batischev, A. I.; Lanzina, N. V.; Tokarev, Y.

[97730] N.; Alexeyev, G. A.: Electrophoretic and kinetic properties of glucose-6-phosphate

[97731] dehydrogenase from erythrocytes or patients with hemolytic anemia,

[97732] related to deficiency of the enzyme activity. Vopr. Med. Khim. 23:

[97733] 166–171, 1977.

[97734] 16975.

[97735] Childs, B.; Zinkham, W.; Browne, E. A.; Kimbro, E. L.; Torbert,

[97736] J. V.: A genetic study of a defect in glutathione metabolism of the

[97737] erythrocytes. Bull. Johns Hopkins Hosp. 102: 21–37, 1958.

[97738] 16976.

[97739] Chiu, D. T. Y.; Zuo, L.; Chao, L.; Chen, E.; Louie, E.; Lubin,

- [97740] B.; Liu, T. Z.; Du, C. S.: Molecular characterization of glucose-6-phosphate
- [97741] dehydrogenase (G6PD) deficiency in patients of Chinese descent and
- [97742] identification of new base substitutions in the human G6PD gene. Blood 81:
- [97743] 2150-2154, 1993.
- [97744] 16977.
- [97745] Chockkalingam, K.; Board, P. G.: Further evidence for heterogeneity
- [97746] of glucose-6-phosphate dehydrogenase deficiency in Papua New Guinea. Hum.
- [97747] Genet. 56: 209-212, 1980.
- [97748] 16978.
- [97749] Chockkalingam, K.; Board, P. G.; Breguet, G.: Glucose-6-phosphate
- [97750] dehydrogenase variants of Bali Island (Indonesia). Hum. Genet. 60:
- [97751] 60-62, 1982.
- [97752] 16979.
- [97753] Chockkalingam, K.; Board, P. G.; Nurse, G. T.: Glucose-

6-phosphate

- [97754] dehydrogenase deficiency in Papua New Guinea: the description of 13
- [97755] new variants. Hum. Genet. 60: 189-192, 1982.
- [97756] 16980.
- [97757] Chuanshu, D.; Yankang, X.; Lin, W. Q. R.; Xiaoyun, H.: Studies
- [97758] on erythrocyte glucose-6-phosphate dehydrogenase variants in Chinese.
- [97759] I. Gd(-) Lizu-Baisha. Acta Acad. Med. Zhong Shan 2: 649-658, 1981.
- [97760] 16981.
- [97761] Cocco, P.; Todde, P.; Fornera, S.; Manca, M. B.; Manca, P. Sias,
- [97762] A. R.: Mortality in a cohort of men expressing the glucose-6-phosphate
- [97763] dehydrogenase deficiency. Blood 91: 706-709, 1998.
- [97764] 16982.
- [97765] Cohn, J.; Carter, N.; Warburg, M.: Glucose-6-phosphate dehydrogenase
- [97766] deficiency in a native Danish family: a new variant. Scand.

J. Haemat. 23:

[97767] 403–406, 1979.

[97768] 16983.

[97769] Colonna–Romano, S.; Iolascon, A.; Lippo, S.; Pinto, L.; Cutillo,

[97770] S.; Battistuzzi, G.: Genetic heterogeneity at the glucose–6-phosphate

[97771] dehydrogenase locus in southern Italy: a study on the population of

[97772] Naples. Hum. Genet. 69: 228–232, 1985.

[97773] 16984.

[97774] Cooper, D. W.; Johnston, P. G.; Murtagh, C. E.; Sharman, G. B.;

[97775] Vandeberg, J. L.; Poole, W. E.: Sex–linked isozymes and sex–chromosome

[97776] evolution and inactivation in kangaroos. In: Markeit, C. L.: Isozymes.

[97777] Developmental Biology. New York: Academic Press (pub.) III: 1975.

[97778] Pp. 559–573.

[97779] 16985.

- [97780] Cooper, M. R.; Dechatelet, L. R.; McCall, C. E.; Lavia, M. F.;
- [97781] Spurr, C. L.; Baehner, R. L.: Complete deficiency of leukocyte glucose-6-phosphate
- [97782] dehydrogenase with defective bactericidal activity. J. Clin. Invest. 51:
- [97783] 769-778, 1972.
- [97784] 16986.
- [97785] Corash, L.; Spielberg, S.; Bartsocas, C.; Boxer, L.; Steiner,
- [97786] R.; Sheetz, M.; Egan, M.; Schlesselman, J.; Schulman, J. D.: Reduced
- [97787] chronic hemolysis during high-dose vitamin E administration in Mediterranean-type
- [97788] glucose-6-phosphate dehydrogenase deficiency. New Eng. J. Med. 303:
- [97789] 416-420, 1980.
- [97790] 16987.
- [97791] Corcoran, C. M.; Calabro, V.; Tamagnini, G.; Town, M.; Haidar,
- [97792] B.; Vulliamy, T. J.; Mason, P. J.; Luzzatto, L.: Molecular heterogeneity

- [97793] underlying the G6PD Mediterranean phenotype. Hum. Genet. 88: 688–690, 1992.
- [97794] 1992.
- [97795] 16988.
- [97796] Costa, E.; Cabeda, J. M.; Vieira, E.; Pinto, R.; Pereira, S. A.; Ferraz, L.; Santos, R.; Barbot, J.: Glucose-6-phosphate dehydrogenase
- [97798] Aveiro: a de novo mutation associated with chronic non-spherocytic
- [97799] hemolytic anemia. Blood 95: 1499–1501, 2000.
- [97800] 16989.
- [97801] Crookston, J. H.; Yoshida, A.; Lin, M.; Booser, D. J.: G6PD Toronto. Biochem. Genet. 8: 259–265, 1973.
- [97802] 16990.
- [97803] 16990.
- [97804] Csepregy, M.; Hall, M. K.; Berkow, R. L.; Jackson, S.; Prchal,
- [97805] J. T.: Characterization of a new G6PD variant: G6PD Titusville. Am. J. Med. Sci. 297: 114–117, 1989.

- [97807] 16991.
- [97808] Csepregy, M.; Yeilding, A.; Lilly, M.; Hall, K.; Scott, C. W.;
- [97809] Prchal, J. T.: Characterization of a new glucose-
6-phosphate dehydrogenase
- [97810] variant: G6PD Central City. Am. J. Hemat. 28: 61-62,
1988.
- [97811] 16992.
- [97812] D'Urso, M.; Luzzatto, L.; Perroni, L.; Ciccodicola, A.; Gen-
tile,
- [97813] G.; Peluso, I.; Persico, M. G.; Pizzella, T.; Toniolo, D.; Vul-
liamy,
- [97814] T. J.: An extensive search for RFLP in the human glucose-
6-phosphate
- [97815] dehydrogenase locus has revealed a silent mutation in the
coding sequence. Am.
- [97816] J. Hum. Genet. 42: 735-741, 1988.
- [97817] 16993.
- [97818] Roessler, B. J.; Palella, T. D.; Heidler, S.; Becker, M. A.:
- [97819] Identification of distinct PRPS1 mutations in two patients
with X-linked
- [97820] phosphoribosylpyrophosphate synthetase superactivity.

(Abstract) Clin.

[97821] Res. 39: 267A, 1991.

[97822] 16994.

[97823] Moreno Fuenmayor, H.; Roldan-Paris, L.; Bermudez, H.:
Ectodermal

[97824] dysplasia in females and inversion of chromosome 9. J.
Med. Genet. 18:

[97825] 214-217, 1981.

[97826] 16995.

[97827] Nakata, M.; Koshiba, H.; Eto, K.; Nance, W. E.: A genetic
study

[97828] of anodontia in X-linked hypohidrotic ectodermal dysplasia. Am. J.

[97829] Hum. Genet. 32: 908-919, 1980.

[97830] 16996.

[97831] Passarge, E.; Fries, E.: X-chromosome inactivation in X-linked

[97832] hypohidrotic ectodermal dysplasia. Nature N.B. 245:
58-59, 1973.

[97833] 16997.

[97834] Pinheiro, M.; Freire-Maia, N.: Christ-Siemens-Touraine

syndrome--a

[97835] clinical and genetic analysis of a large Brazilian kindred. I.

Affected

[97836] females. II. Affected males. III. Carrier detection. Am. J.

Med.

[97837] Genet. 4: 113-134, 1979.

[97838] 16998.

[97839] Pinheiro, M.; Ideriha, M. T.; Chautard-Freire-Maia, E. A.;

Freire-Maia,

- [97840] N.; Primo-Parmo, S. L.: Christ-Siemens-Touraine syndrome: investigations
[97841] on two large Brazilian kindreds with a new estimate of the
manifestation
[97842] rate among carriers. Hum. Genet. 57: 428-431, 1981.
[97843] 16999.
- [97844] Reed, W. B.; Lopez, D. A.; Landing, B. H.: Clinical spectrum
[97845] of anhidrotic ectodermal dysplasia. Arch. Derm. 102:
134-143, 1970.
[97846] 17000.
- [97847] Richards, W.; Kaplan, M.: Anhidrotic ectodermal dysplasia:
an
[97848] unusual case of pyrexia in the newborn. Am. J. Dis. Child.
117:
[97849] 597-598, 1969.
[97850] 17001.
- [97851] Roberts, E.: The inheritance of anhidrosis associated with
anodontia. J.A.M.A. 93:
[97852] 277-279, 1929.
[97853] 17002.

- [97854] Saksena, S. S.; Bixler, D.: Facial morphometrics in the identification
- [97855] of gene carriers of X-linked hypohidrotic ectodermal dysplasia. Am.
- [97856] J. Med. Genet. 35: 105–114, 1990.
- [97857] 17003.
- [97858] Simpson, J. L.; Allen, F. H., Jr.; New, M.; German, J.: Absence
- [97859] of close linkage between the locus for Xg and the locus for anhidrotic
- [97860] ectodermal dysplasia. Vox Sang. 17: 465–467, 1969.
- [97861] 17004.
- [97862] Singh, A.; Jolly, S. S.; Kaur, S.: Hereditary ectodermal dysplasia. Brit.
- [97863] J. Derm. 74: 34–37, 1962.
- [97864] 17005.
- [97865] Soderholm, A.-L.; Kaitila, I.: Expression of X-linked hypohidrotic
- [97866] ectodermal dysplasia in six males and in their mothers. Clin. Genet. 28:
- [97867] 136–144, 1985.

- [97868] 17006.
- [97869] Srivastava, A. K.; Durmowicz, M. C.; Hartung, A. J.; Hudson, J.;
- [97870] Ouzts, L. V.; Donovan, D. M.; Cui, C.-Y.; Schlessinger, D.:
Ectodysplasin-A1
- [97871] is sufficient to rescue both hair growth and sweat glands
in Tabby
- [97872] mice. Hum. Molec. Genet. 10: 2973-2981, 2001.
- [97873] 17007.
- [97874] Srivastava, A. K.; Montonen, O.; Saarialho-Kere, U.; Chen,
E.;
- [97875] Baybayan, P.; Pispá, J.; Limon, J.; Schlessinger, D.; Kere, J.:
Fine
- [97876] mapping of the EDA gene: a translocation breakpoint is
associated
- [97877] with a CpG island that is transcribed. Am. J. Hum. Genet.
58: 126-132,
- [97878] 1996.
- [97879] 17008.
- [97880] Srivastava, A. K.; Pispá, J.; Hartung, A. J.; Du, Y.; Ezer, S.;
- [97881] Jenks, T.; Shimada, T.; Pekkanen, M.; Mikkola, M. L.; Ko,

M. S. H.;

- [97882] Thesleff, I.; Kere, J.; Schlessinger, D.: The Tabby phenotype is
[97883] caused by mutation in a mouse homologue of the EDA
gene that reveals
[97884] novel mouse and human exons and encodes a protein
(ectodysplasin-A)
[97885] with collagenous domains. Proc. Nat. Acad. Sci. 94:
13069–13074,
[97886] 1997.
[97887] 17009.
- [97888] Thadani, K. I.: The toothless men of Sind. J. Hered. 25:
483–484,
[97889] 1934.
[97890] 17010.
- [97891] Thurnam, J.: Two cases in which the skin, hair and teeth
were
[97892] very imperfectly developed. Proc. R. M. Chir. Soc. London
31: 71–82,
[97893] 1848.
[97894] 17011.

- [97895] Turleau, C.; Niaudet, P.; Cabanis, M.-O.; Plessis, G.; Cau, D.;
- [97896] de Grouchy, J.: X-linked hypohidrotic ectodermal dysplasia and t(X;12)
- [97897] in a female. Clin. Genet. 35: 462–466, 1989.
- [97898] 17012.
- [97899] Vargas, G. A.; Fantino, E.; George-Nascimento, C.; Gargus, J.
- [97900] J.; Haigler, H. T.: Reduced epidermal growth factor receptor expression
- [97901] in hypohidrotic ectodermal dysplasia and Tabby mice. J. Clin. Invest. 97:
- [97902] 2426–2432, 1996.
- [97903] 17013.
- [97904] Vincent, M. C.; Biancalana, V.; Ginisty, D.; Mandel, J. L.; Calvas,
- [97905] P.: Mutational spectrum of the ED1 gene in X-linked hypohidrotic
- [97906] ectodermal dysplasia. Europ. J. Hum. Genet. 9: 355–363, 2001.
- [97907] 17014.

- [97908] WPA, Federal Writers' Project: Mississippi: A Guide to the
[97909] Magnolia State. New York: Hastings House (pub.) 1938.
Pp. 359
[97910] only.
[97911] 17015.
[97912] Yotsumoto, S.; Fukumaru, S.; Matsushita, S.; Oku, T.;
Kobayashi,
[97913] K.; Saheki, T.; Kanzaki, T.: A novel point mutation of the
EDA gene
[97914] in a Japanese family with anhidrotic ectodermal dysplasia.
(Letter) J.
[97915] Invest. Derm. 111: 1246–1247, 1998.
[97916] 17016.
[97917] Ballabio, A.: Personal Communication. Houston, Texas
3/11/1993.
[97918] 17017.
[97919] Bardin, C. W.; Ross, G. T.; Rifkind, A. B.; Cargille, C. M.;
Lipsett,
[97920] M. B.: Studies of the pituitary Leydig cell axis in young
men with
[97921] hypogonadotropic hypogonadism and hyposmia: compar-

ison with normal

[97922] men, prepubertal boys, and hypo-pituitary patients. J.

Clin. Invest. 48:

[97923] 2046-2056, 1969.

[97924] 17018.

[97925] Bick, D.; Franco, B.; Sherins, R. J.; Heye, B.; Pike, L.; Crawford,

[97926] J.; Maddalena, A.; Incerti, B.; Pragliola, A.; Meitinger, T.; Ballabio,

[97927] A.: Brief report: intragenic deletion of the KALIG-1 gene in Kallmann's

[97928] syndrome. New Eng. J. Med. 326: 1752-1755, 1992.

[97929] 17019.

[97930] Bick, D. P.; Schwanzel-Fukuda, M.; Pfaff, D. W.; Schorderet, D.

[97931] F.; Price, P. A.; Campbell, L.; Huff, R. W.; Moore, C. M.: Prenatal

[97932] diagnosis and investigation of a fetus with chondrodysplasia punctata,

[97933] ichthyosis and Kallmann syndrome due to an Xp deletion: evidence for

[97934] a neuronal migration defect in Kallmann syndrome.

(Abstract) Am.

[97935] J. Hum. Genet. 45 (suppl.): A252, 1989.

[97936] 17020.

[97937] Birnbacher, R.; Wandl-Vergesslich, K.; Frisch, H.: Diagnosis of

[97938] X-recessive Kallmann syndrome in early infancy: evidence of hypoplastic

[97939] rhinencephalon. Europ. J. Pediat. 153: 245-247, 1994.

[97940] 17021.

[97941] Bulow, H. E.; Berry, K. L.; Topper, L. H.; Peles, E.; Hobert, O.

[97942] : Heparan sulfate proteoglycan-dependent induction of axon branching

[97943] and axon misrouting by the Kallmann syndrome gene kal-1. Proc. Nat.

[97944] Acad. Sci. 99: 6346-6351, 2002.

[97945] 17022.

[97946] del Castillo, I.; Cohen-Salmon, M.; Blanchard, S.; Lutfalla, G.;

[97947] Petit, C.: Structure of the X-linked Kallmann syndrome gene and its

- [97948] homologous pseudogene on the Y chromosome. *Nature Genet.* 2: 305–310, 1992.
- [97949] 17023.
- [97950] 17023.
- [97951] De Morsier, G.: Etudes sur les dysraphies cranio-encephaliques.
- [97952] I. Agenesie des lobes olfactifs (telencephaloschizis lateral) et des
- [97953] commissures calleuse et anterieure (telencephaloschizis median): la
- [97954] dysplasie olfacto-genitale. *Schweiz. Arch. Neurol. Psychiat.* 74: 309–361, 1954.
- [97955] 17024.
- [97956] 17024.
- [97957] Basse, F.; Stout, J. G.; Sims, P. J.; Wiedmer, T.: Isolation of
- [97958] an erythrocyte membrane protein that mediates Ca^{2+} -dependent transbilayer
- [97959] movement of phospholipid. *J. Biol. Chem.* 271: 17205–17210, 1996.
- [97960] 17025.
- [97961] Garrett, R. M.; Johnson, J. L.; Graf, T. N.; Feigenbaum, A.;

Rajagopalan,

- [97962] K. V.: Human sulfite oxidase R160Q: identification of the mutation
[97963] in a sulfite oxidase-deficient patient and expression and characterization
[97964] of the mutant enzyme. Proc. Nat. Acad. Sci. 95: 6394-6398, 1998.
[97965] 17026.
- [97966] Kisker, C.; Schindelin, H.; Pacheco, A.; Wehbi, W. A.; Garrett,
[97967] R. M.; Rajagopalan, K. V.; Enemark, J. H.; Rees, D. C.: Molecular
[97968] basis of sulfite oxidase deficiency from the structure of sulfite
[97969] oxidase. Cell 91: 973-983, 1997.
[97970] 17027.
- [97971] Naylor, J.; Brinke, A.; Hassock, S.; Green, P. M.; Giannelli,
[97972] F.: Characteristic mRNA abnormality found in half the patients with
[97973] severe haemophilia A is due to large DNA inversions. Hum. Molec.
[97974] Genet. 2: 1773-1778, 1993.

- [97975] 17028.
- [97976] Renieri, A.; Seri, M.; Myers, J. C.; Pihlajaniemi, T.; Massella,
[97977] L.; Rizzoni, G.; De Marchi, M.: De novo mutation in the
COL4A5 gene
[97978] converting glycine 325 to glutamic acid in Alport syn-
drome. Hum.
[97979] Molec. Genet. 1: 127-129, 1992.
- [97980] 17029.
- [97981] Corral, J.; Forster, A.; Thompson, S.; Lampert, F.; Kaneko,
Y.;
- [97982] Slater, R.; Kroes, W. G.; van der Schoot, C. E.; Ludwig,
W.-D.; Karpas,
[97983] A.; Pocock, C.; Cotter, F.; Rabbitts, T. H.: Acute leukemias
of different
[97984] lineages have similar MLL gene fusions encoding related
chimeric proteins
[97985] resulting from chromosomal translocation. Proc. Nat.
Acad. Sci. 90:
[97986] 8538-8542, 1993.
- [97987] 17030.
- [97988] Knebelmann, B.; Deschenes, G.; Gros, F.; Hors, M.-C.;

Grunfeld,

- [97989] J.-P.; Tryggvason, K.; Gubler, M.-C.; Antignac, C.: Substitution
[97990] of arginine for glycine 325 in the collagen alpha-5 (IV)
chain associated
[97991] with X-linked Alport syndrome: characterization of the
mutation by
[97992] direct sequencing of PCR-amplified lymphoblast cDNA
fragments. Am.
[97993] J. Hum. Genet. 51: 135-142, 1992.
[97994] 17031.
[97995] Renieri, A.; Seri, M.; Myers, J. C.; Pihlajaniemi, T.; Sessa,
[97996] A.; Rizzoni, G.; De Marchi, M.: Alport syndrome caused by
a 5-prime
[97997] deletion within the COL4A5 gene. Hum. Genet. 89:
120-121, 1992.
[97998] 17032.
[97999] Smeets, H. J. M.; Melenhorst, J. J.; Lemmink, H. H.;
Schroder,
[98000] C. H.; Nelen, M. R.; Zhou, J.; Hostikka, S. L.; Tryggvason,
K.; Ropers,
[98001] H.-H.; Jansweijer, M. C. E.; Monnens, L. A. H.; Brunner, H.

G.; van

[98002] Oost, B. A.: Different mutations in the COL4A5 collagen gene in two

[98003] patients with different features of Alport syndrome. Kidney Int. 42:

[98004] 83–88, 1992.

[98005] 17033.

[98006] Thorner, P. S.; Zheng, K.; Kalluri, R.; Jacobs, R.; Hudson, B.

[98007] G.: Coordinate gene expression of the alpha-3, alpha-4, and alpha-5

[98008] chains of collagen type IV. J. Biol. Chem. 271: 13821–13828, 1996.

[98009] 17034.

[98010] Turco, A. E.; Rossetti, S.; Biasi, M. O.; Rizzoni, G.; Massella,

[98011] L.; Saarinen, N. H.; Renieri, A.; Pignatti, P. F.; De Marchi, M.:

[98012] A novel missense mutation in exon 3 of the COL4A5 gene associated

[98013] with late-onset Alport syndrome. Clin. Genet. 48: 261–263, 1995.

[98014] 17035.

- [98015] Vetrie, D.; Boye, E.; Flinter, F.; Bobrow, M.; Harris, A.: DNA
[98016] rearrangements in the alpha-5(IV) collagen gene (COL4A5)
of individuals
[98017] with Alport syndrome: further refinement using pulsed-
field gel electrophoresis. Genomics 14:
[98018] 624-633, 1992.
[98019] 17036.
[98020] Vetrie, D.; Flinter, F.; Bobrow, M.; Harris, A.: Long-range
mapping
[98021] of the gene for the human alpha-5(IV) collagen chain at
Xq22-q23. Genomics 12:
[98022] 130-138, 1992.
[98023] 17037.
[98024] Zheng, K.; Thorner, P. S.; Marrano, P.; Baumal, R.;
McInnes, R.
[98025] R.: Canine X chromosome-linked hereditary nephritis: a
genetic model
[98026] for human X-linked hereditary nephritis resulting from a
single base
[98027] mutation in the gene encoding the alpha-5 chain of colla-
gen type IV. Proc.

- [98028] Nat. Acad. Sci. 91: 3989–3993, 1994.
- [98029] 17038.
- [98030] Zhou, J.; Barker, D. F.; Hostikka, S. L.; Gregory, M. C.;
Atkin,
- [98031] C. L.; Tryggvason, K.: Single base mutation in alpha-5(IV)
collagen
- [98032] chain gene converting a conserved cysteine to serine in
Alport syndrome. Genomics 9:
- [98033] 10–18, 1991.
- [98034] 17039.
- [98035] Zhou, J.; Hertz, J. M.; Leinonen, A.; Tryggvason, K.: Com-
plete
- [98036] amino acid sequence of the human alpha-5(IV) collagen
chain and identification
- [98037] of a single-base mutation in exon 23 converting glycine
521 in the
- [98038] collagenous domain to cysteine in an Alport syndrome
patient. J.
- [98039] Biol. Chem. 267: 12475–12481, 1992.
- [98040] 17040.
- [98041] Zhou, J.; Hostikka, S. L.; Chow, L. T.; Tryggvason, K.:

Characterization

[98042] of the 3-prime half of the human type IV collagen alpha-5 gene that

[98043] is affected in the Alport syndrome. Genomics 9: 1-9, 1991.

[98044] 17041.

[98045] Oohashi, T.; Sugimoto, M.; Mattei, M.-G.; Ninomiya, Y.: Identification

[98046] of a new collagen IV chain, alpha-6(IV), by cDNA isolation and assignment

[98047] of the gene to chromosome Xq22, which is the same locus for COL4A5. J.

[98048] Biol. Chem. 269: 7520-7526, 1994.

[98049] 17042.

[98050] Oohashi, T.; Ueki, Y.; Sugimoto, M.; Ninomiya, Y.: Isolation and

[98051] structure of the COL4A6 gene encoding the human alpha-6(IV) collagen

[98052] chain and comparison with other type IV collagen genes. J. Biol.

[98053] Chem. 270: 26863-26867, 1995.

[98054] 17043.

- [98055] Sugimoto, M.; Oohashi, T.; Ninomiya, Y.: The genes COL4A5 and
- [98056] COL4A6, coding for basement membrane collagen chains alpha-5(IV) and
- [98057] alpha-6(IV), are located head-to-head in close proximity on human
- [98058] chromosome Xq22 and COL4A6 is transcribed from two alternative promoters. Proc.
- [98059] Nat. Acad. Sci. 91: 11679-11683, 1994.
- [98060] 17044.
- [98061] Parry, P.; Wei, Y.; Evans, G.: Cloning and characterization of
- [98062] the t(X;11) breakpoint from a leukemic cell line identify a new member
- [98063] of the forkhead gene family. Genes Chromosomes Cancer 11: 79-84,
- [98064] 1994.
- [98065] 17045.
- [98066] Peters, U.; Haberhausen, G.; Kostrzewa, M.; Nolte, D.; Muller,
- [98067] U.: AFX1 and p54(nrb): fine mapping, genomic structure, and exclusion

- [98068] as candidate genes of X-linked dystonia parkinsonism.
Hum. Genet. 100:
- [98069] 569–572, 1997.
- [98070] 17046.
- [98071] Bohme, B.; VandenBos, T.; Cerretti, D. P.; Park, L. S.;
Holtrich,
- [98072] U.; Rubsamen-Waigmann, H.; Strebhardt, K.: Cell-cell ad-
hesion mediated
- [98073] by binding of membrane-anchored ligand LERK-2 to the
EPH-related receptor
- [98074] human embryonal kinase 2 promotes tyrosine kinase ac-
tivity. J. Biol.
- [98075] Chem. 271: 24747–24752, 1996.
- [98076] 17047.
- [98077] Fletcher, F. A.; Huebner, K.; Shaffer, L. G.; Fairweather, N.
D.;
- [98078] Monaco, A. P.; Muller, U.; Druck, T.; Simoneaux, D. K.;
Chelly, J.;
- [98079] Belmont, J. W.; Beckmann, M. P.; Lyman, S. D.: Assignment
of the
- [98080] gene (EPLG2) encoding a high-affinity binding protein for
the receptor

[98081] tyrosine kinase Elk to a 200–kilobasepair region in human chromosome

[98082] Xq12. Genomics 25: 334–335, 1995.

[98083] 17048.

[98084] Fletcher, F. A.; Renshaw, B.; Hollingsworth, T.; Baum, P.; Lyman,

[98085] S. D.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Davison, B.

[98086] L.: Genomic organization and chromosomal localization of mouse Eplg2,

[98087] a gene encoding a binding protein for the receptor tyrosine kinase

[98088] Elk. Genomics 24: 127–132, 1994.

[98089] 17049.

[98090] Palmer, A.; Zimmer, M.; Erdmann, K. S.; Eulenburg, V.; Porthin,

[98091] A.; Heumann, R.; Deutsch, U.; Klein, R.: EphrinB phosphorylation

[98092] and reverse signaling: regulation by Src kinases and PTP-BL phosphatase. Molec.

[98093] Cell 9: 725–737, 2002.

- [98094] 17050.
- [98095] Gregor, P.; Nash, S. R.; Caron, M. G.; Seldin, M. F.; Warren, S.
- [98096] T.: Assignment of the creatine transporter gene (SLC6A8) to human
- [98097] chromosome Xq28 telomeric to G6PD. *Genomics* 25: 332–333, 1995.
- [98098] 17051.
- [98099] Hahn, K. A.; Salomons, G. S.; Tackels–Horne, D.; Wood, T. C.; Taylor,
- [98100] H. A.; Schroer, R. J.; Lubs, H. A.; Jakobs, C.; Olson, R. L.; Holden,
- [98101] K. R.; Stevenson, R. E.; Schwartz, C. E.: X-linked mental retardation
- [98102] with seizures and carrier manifestations is caused by a mutation in
- [98103] the creatine–transporter gene (SLC6A8) located in Xq28. *Am. J. Hum. Genet.* 70: 1349–1356, 2002.
- [98104] 17052.
- [98105] Item, C. B.; Stockler–Ipsiroglu, S.; Stromberger, C.; Muhl, A.;

- [98107] Alessandri, M. G.; Bianchi, M. C.; Tosetti, M.; Fornai, F.; Cioni,
[98108] G.: Arginine:glycine amidinotransferase deficiency: the third inborn
[98109] error of creatine metabolism in humans. *Am. J. Hum. Genet.* 69: 1127–33,
[98110] 2001.
[98111] 17053.
[98112] Iyer, G. S.; Krahe, R.; Goodwin, L. A.; Doggett, N. A.; Siciliano,
[98113] M. J.; Funanage, V. L.; Proujansky, R.: Identification of a testis-expressed
[98114] creatine transporter gene at 16p11.2 and confirmation of the X-linked
[98115] locus to Xq28. *Genomics* 34: 143–146, 1996.
[98116] 17054.
[98117] Nash, S. R.; Giros, B.; Kingsmore, S. F.; Rochelle, J. M.; Suter,
[98118] S. T.; Gregor, P.; Seldin, M. F.; Caron, M. G.: Cloning, pharmacological
[98119] characterization, and genomic localization of the human creatine transporter. *Receptors*

- [98120] Channels 2: 165–174, 1994.
- [98121] 17055.
- [98122] Salomons, G. S.; van Dooren, S. J. M.; Verhoeven, N. M.; Cecil,
- [98123] K. M.; Ball, W. S.; Degrauw, T. J.; Jakobs, C.: X-linked creatine-transporter
- [98124] gene (SLC6A8) defect: a new creatine-deficiency syndrome. Am. J. Hum. Genet. 68: 1497–1500, 2001.
- [98125] 17056.
- [98126] Sandoval, N.; Bauer, D.; Brenner, V.; Coy, J. F.; Drescher, B.;
- [98127] Kioschis, P.; Korn, B.; Nyakatura, G.; Poustka, A.; Reichwald, K.;
- [98128] Rosenthal, A.; Platzer, M.: The genomic organization of a human creatine
- [98129] transporter (CRTR) gene located in Xq28. Genomics 35: 383–385, 1996.
- [98130] 17057.
- [98131] Stockler, S.; Hanefeld, F.; Frahm, J.: Creatine replacement therapy

- [98133] in guanidinoacetate methyltransferase deficiency, a novel inborn error
- [98134] of metabolism. *Lancet* 348: 789–90, 1996.
- [98135] 17058.
- [98136] Brzustowicz, L. M.; Farrell, S.; Khan, M. B.; Weksberg, R.: Mapping
- [98137] of a new SGBS locus to chromosome Xp22 in a family with a severe form
- [98138] of Simpson–Golabi–Behmel syndrome. *Am. J. Hum. Genet.* 65: 779–783,
- [98139] 1999.
- [98140] 17059.
- [98141] Filmus, J.; Church, J. G.; Buick, R. N.: Isolation of a cDNA corresponding
- [98142] to a developmentally regulated transcript in rat intestine. *Molec. Cell Biol.* 4243–4249, 1988.
- [98143] 17060.
- [98144] 17060.
- [98145] Gelb, B. D.; Shi, G.–P.; Chapman, H. A.; Desnick, R. J.: Pycnodysostosis,
- [98146] a lysosomal disease caused by cathepsin K deficiency. *Sci-*

ence 273:

[98147] 1236–1238, 1996.

[98148] 17061.

[98149] Gelb, B. D.; Willner, J. P.; Dunn, T. M.; Kardon, N. B.; Verloes,

[98150] A.; Poncin, J.; Desnick, R. J.: Paternal uniparental disomy for chromosome

[98151] 1 revealed by molecular analysis of a patient with pycnodysostosis. Am.

[98152] J. Hum. Genet. 62: 848–854, 1998.

[98153] 17062.

[98154] Polymeropoulos, M. H.; Ortiz De Luna, R. I.; Ide, S. E.; Torres,

[98155] R.; Rubenstein, J.; Francomano, C. A.: The gene for pycnodysostosis

[98156] maps to human chromosome 1cen–q21. Nature Genet. 10: 238–239, 1995.

[98157] 17063.

[98158] Bourke, E.; Frindt, G.; Flynn, P.; Schreiner, G. E.: Primary hyperoxaluria

[98159] with normal alpha–ketoglutarate:glyoxylate carboligase

activity: treatment

[98160] with isocarboxazid. Ann. Intern. Med. 76: 279–284, 1972.

[98161] 17064.

[98162] Danpure, C. J.: Personal Communication. Middlesex, England

[98163] 6/16/1988.

[98164] 17065.

[98165] Danpure, C. J.: Primary hyperoxaluria type 1 and peroxisome-to-mitochondrion

[98166] mistargeting of alanine:glyoxylate aminotransferase. Biochimie 75:

[98167] 309–315, 1993.

[98168] 17066.

[98169] Danpure, C. J.; Jennings, P. R.: Peroxisomal alanine:glyoxylate

[98170] aminotransferase deficiency in primary hyperoxaluria type I. FEBS

[98171] Lett. 201: 20–24, 1986.

[98172] 17067.

[98173] Danpure, C. J.; Jennings, P. R.; Watts, R. W. E.: Enzymolog–

ical

[98174] diagnosis of primary hyperoxaluria type 1 by measurement of hepatic

[98175] alanine:glyoxylate aminotransferase activity. Lancet I: 289-291,

[98176] 1987.

[98177] 17068.

[98178] Brunelli, S.; Faiella, A.; Capra, V.; Nigro, V.; Simeone, A.; Cama,

[98179] A.; Boncinelli, E.: Germline mutation in the homeobox gene EMX2 in

[98180] patients with severe schizencephaly. Nature Genet. 12: 94-96, 1996.

[98181] 17069.

[98182] Burg, J.; Conzelmann, E.; Sandhoff, K.; Solomon, E.; Swallow, D.

[98183] M.: Mapping of the gene coding for the human GM2 activator protein

[98184] to chromosome 5. Ann. Hum. Genet. 49: 41-45, 1985.

[98185] 17070.

[98186] Chen, B.; Rigat, B.; Curry, C.; Mahuran, D. J.: Structure of

the

- [98187] **GM2A gene: identification of an exon 2 nonsense mutation and a naturally occurring transcript with an in-frame deletion of exon 2. Am. J.**
- [98189] **Hum. Genet. 65: 77–87, 1999.**
- [98190] **17071.**
- [98191] **Conzelmann, E.; Sandhoff, K.: AB variant of infantile Gm2–gangliosidosis:**
- [98192] **deficiency of a factor necessary for stimulation of hexosaminidase**
- [98193] **A–catalyzed degradation of ganglioside Gm2 and glycolipid Ga2. Proc.**
- [98194] **Nat. Acad. Sci. 75: 3979–3983, 1978.**
- [98195] **17072.**
- [98196] **Hechtman, P.; Gordon, B. A.; Ng Ying Kin, N. M. K.: Deficiency**
- [98197] **of the hexosaminidase A activator protein in a case of GM2 gangliosidosis;**
- [98198] **variant AB. Pediat. Res. 16: 217–222, 1982.**
- [98199] **17073.**

- [98200] Heng, H. H. Q.; Xie, B.; Shi, X.-M.; Tsui, L.-C.; Mahuran, D. J.
- [98201] : Refined mapping of the GM2 activator protein (GM2A) locus to 5q31.3-q33.1,
- [98202] distal to the spinal muscular atrophy locus. *Genomics* 18: 429-431,
- [98203] 1993.
- [98204] 17074.
- [98205] Kleyn, P. W.; Brzustowicz, L. M.; Wilhelmsen, K. C.; Freimer, N.
- [98206] B.; Miller, J. M.; Munsat, T. L.; Gilliam, T. C.: Spinal muscular
- [98207] atrophy is not the result of mutations at the beta-hexosaminidase
- [98208] or GM(2)-activator locus. *Neurology* 41: 1418-1422, 1991.
- [98209] 17075.
- [98210] Li, S. C.; Hirabayashi, Y.; Li, Y. T.: A new variant of type-AB
- [98211] GM2-gangliosidosis. *Biochem. Biophys. Res. Commun.* 101: 479-485,
- [98212] 1981.

[98213] 17076.

[98214] Barton, D. E.; Yang-Feng, T. L.; Francke, U.: The human tyrosine

[98215] aminotransferase gene mapped to the long arm of chromosome 16 (region

[98216] 16q22-q24) by somatic cell hybrid analysis and in situ hybridization. Hum.

[98217] Genet. 72: 221-224, 1986.

[98218] 17077.

[98219] Beinfang, D. C.; Kuwabara, T.; Pueschel, S. M.: The Richner-Hanhart

[98220] syndrome: report of a case with associated tyrosinemia. Arch. Ophthal. 94:

[98221] 1133-1137, 1976.

[98222] 17078.

[98223] Bohnert, A.; Anton-Lamprecht, I.: Richner-Hanhart's syndrome:

[98224] ultrastructural abnormalities of epidermal keratinization indicating

[98225] a causal relationship to high intracellular tyrosine levels. J. Invest.

[98226] Derm. 79: 68–74, 1982.

[98227] 17079.

[98228] Buist, N.: Phenylketonuria and related problems. In: Nyhan, W.

[98229] L.: Amino Acid Metabolism and Genetic Variation. New York: McGraw-Hill

[98230] (pub.) 1967. Pp. 117 only.

[98231] 17080.

[98232] Chitayat, D.; Balbul, A.; Hani, V.; Mamer, O. A.; Clow, C.; Sriver,

[98233] C. R.: Hereditary tyrosinaemia type II in a consanguineous Ashkenazi

[98234] Jewish family: intrafamilial variation in phenotype; absence of parental

[98235] phenotype effects on the fetus. J. Inherit. Metab. Dis. 15: 198–203,

[98236] 1992.

[98237] 17081.

[98238] Crovato, F.; Desirello, G.; Gatti, R.; Babbini, N.; Rebora, A.

[98239] : Richner–Hanhart syndrome spares a plantar autograft. Arch. Derm. 121:

[98240] 539–540, 1985.

[98241] 17082.

[98242] Fellman, J. H.; Vanbellinghen, P. J.; Jones, R. T.; Koler, R. D.

[98243] : Soluble and mitochondrial forms of tyrosine aminotransferase: relationship

[98244] to human tyrosinemia. *Biochemistry* 8: 615–622, 1969.

[98245] 17083.

[98246] Garibaldi, L. R.; Siliato, F.; De Martini, I.; Scarsi, M. R.; Romano,

[98247] C.: Oculocutaneous tyrosinosis: report of two cases in the same family. *Helv.*

[98248] *Paediat. Acta* 32: 173–180, 1977.

[98249] 17084.

[98250] Goldsmith, L. A.; Kang, E. S.; Bienfang, D. C.; Jimbow, K.; Gerald,

[98251] P. S.; Baden, H. P.: Tyrosinemia with plantar and palmar keratosis

[98252] and keratitis. *J. Pediat.* 83: 798–805, 1973.

[98253] 17085.

- [98254] Hanhart, E.: Neue Sonderformen von Keratosis palmo-plantaris,
- [98255] u.a. eine regelmaessig-dominante mit systematisierten Lipomen, ferner
- [98256] 2 einfach-rezessive mit Schwachsinn und z.T. mit Hornhautveraenderungen
- [98257] des Auges (Ektodermatosyndrom). Dermatologica 94: 286-308, 1947.
- [98258] 17086.
- [98259] Hunziker, N.: Richner-Hanhart syndrome and tyrosinemia type II. Dermatologica 160:
- [98260] 180-189, 1980.
- [98261] 17087.
- [98262] Kennaway, N. G.; Buist, N. R. M.: Metabolic studies in a patient
- [98263] with hepatic cytosol tyrosine aminotransferase deficiency. Pediat.
- [98264] Res. 5: 287-297, 1971.
- [98265] 17088.
- [98266] Muller, G.; Scherer, G.; Zentgraf, H.; Ruppert, S.; Hermann,

- [98267] B.; Lehrach, H.; Schutz, G.: Isolation, characterization and chromosomal
- [98268] mapping of the mouse tyrosine aminotransferase gene. J. Molec. Biol. 184:
- [98269] 367–373, 1985.
- [98270] 17089.
- [98271] Natt, E.; Kao, F.-T.; Rettenmeier, R.; Scherer, G.: Assignment
- [98272] of the human tyrosine aminotransferase gene to chromosome 16. Hum.
- [98273] Genet. 72: 225–228, 1986.
- [98274] 17090.
- [98275] Kim, E. K.; Yoo, O. J.; Song, K. Y.; Yoo, H. W.; Choi, S. Y.;
- [98276] Cho, S. W.; Hahn, S. H.: Identification of three novel mutations
- [98277] and a high frequency of the arg778-to-leu mutation in Korean patients
- [98278] with Wilson disease. Hum. Mutat. 11: 275–278, 1998.
- [98279] 17091.
- [98280] Kuo, Y.-M.; Gitschier, J.; Packman, S.: Developmental expression

- [98281] of the mouse mottled and toxic milk genes suggests distinct functions
- [98282] for the Menkes and Wilson disease copper transporters. Hum. Molec.
- [98283] Genet. 6: 1043–1049, 1997.
- [98284] 17092.
- [98285] Loudianos, G.; Dessi, V.; Lovicu, M.; Angius, A.; Figus, A.; Lilliu,
- [98286] F.; De Virgiliis, S.; Nurchi, A. M.; Deplano, A.; Moi, P.; Pirastu,
- [98287] M.; Cao, A.: Molecular characterization of Wilson disease in the
- [98288] Sardinian population—evidence of a founder effect. Hum. Mutat. 14:
- [98289] 294–303, 1999.
- [98290] 17093.
- [98291] Loudianos, G.; Dessi, V.; Lovicu, M.; Angius, A.; Nurchi, A.;
- [98292] Sturniolo, G. C.; Marcellini, M.; Zancan, L.; Bragetti, P.; Akar,
- [98293] N.; Yagci, R.; Vegnente, A.; Cao, A.; Pirastu, M.: Further delineation

- [98294] of the molecular pathology of Wilson disease in the
Mediterranean
- [98295] population. Hum. Mutat. 12: 89–94, 1998.
- [98296] 17094.
- [98297] Okada, T.; Shiono, Y.; Hayashi, H.; Satoh, H.; Sawada, T.;
Suzuki,
- [98298] A.; Takeda, Y.; Yano, M.; Michitaka, K.; Onji, M.; Mabuchi,
H.: Mutational
- [98299] analysis of ATP7B and genotype–phenotype correlation in
Japanese with
- [98300] Wilson's disease. Hum. Mutat. 15: 454–462, 2000.
- [98301] 17095.
- [98302] Sasaki, N.; Hayashizaki, Y.; Muramatsu, M.; Matsuda, Y.;
Ando,
- [98303] Y.; Kuramoto, T.; Serikawa, T.; Azuma, T.; Naito, A.; Agui,
T.; Yamashita,
- [98304] T.; Miyoshi, I.; Takeichi, N.; Kasai, N.: The gene responsi-
ble for
- [98305] LEC hepatitis, located on rat chromosome 16, is the ho-
molog to the
- [98306] human Wilson disease gene. Biochem. Biophys. Res. Com-
mun. 202: 512–518,

- [98307] 1994.
- [98308] 17096.
- [98309] McKusick, V. A.: On the X Chromosome of Man. Washington: Am.
- [98310] Inst. Biol. Sci. (pub.) 1964.
- [98311] 17097.
- [98312] Ichinose, H.; Ohye, T.; Suzuki, T.; Sumi-Ichinose, C.; Nomura,
- [98313] T.; Hagino, Y.; Nagatsu, T.: Molecular cloning of the human Nurr1
- [98314] gene: characterization of the human gene and cDNAs. Gene 230: 233-239,
- [98315] 1999.
- [98316] 17098.
- [98317] Law, S. W.; Conneely, O. M.; DeMayo, F. J.; O'Malley, B. W.: Identification
- [98318] of a new brain-specific transcription factor, NURR1. Molec. Endocr. 6:
- [98319] 2129-2135, 1992.
- [98320] 17099.

- [98321] Mages, H. W.; Rilke, O.; Bravo, R.; Senger, G.; Krocze, R. A.
- [98322] : NOT, a human immediate-early response gene closely related to the
- [98323] steroid/thyroid hormone receptor NAK1/TR3. *Molec. Endocr.* 8: 1583-1591,
- [98324] 1994.
- [98325] 17100.
- [98326] McEvoy, A. N.; Murphy, E. A.; Ponnio, T.; Conneely, O. M.; Bresnihan,
- [98327] B.; FitzGerald, O.; Murphy, E. P.: Activation of nuclear orphan receptor
- [98328] NURR1 transcription by NF-kappaB and cyclic adenosine 5-prime-monophosphate
- [98329] response element-binding protein in rheumatoid arthritis synovial
- [98330] tissue. *J. Immun.* 168: 2979-2987, 2002.
- [98331] 17101.
- [98332] Okabe, T.; Takayanagi, R.; Imasaki, K.; Haji, M.; Nawata, H.; Watanabe,
- [98333] T.: cDNA cloning of a NGFI-B/nur77-related transcription factor from

- [98334] an apoptotic human T cell line. *J. Immun.* 154: 3871–3879, 1995.
- [98335] 17102.
- [98336] Torii, T.; Kawai, T.; Nakamura, S.; Kawakami, H.: Organization
- [98337] of the human orphan nuclear receptor Nurr1 gene. *Gene* 230: 225–232,
- [98338] 1999.
- [98339] 17103.
- [98340] Xu, P.-Y.; Liang, R.; Jankovic, J.; Hunter, C.; Zeng, Y.-X.; Ashizawa,
- [98341] T.; Lai, D.; Le, W.-D.: Association of homozygous 7048G7049 variant
- [98342] in the intron six of Nurr1 gene with Parkinson's disease. *Neurology* 58:
- [98343] 881–884, 2002.
- [98344] 17104.
- [98345] Hanahan, D.: Signaling vascular morphogenesis and maintenance. *Science* 277:
- [98346] 48–50, 1997.
- [98347] 17105.

- [98348] Wong, S. M. E.; Lowe, A. W.: Sequence of the cDNA encoding human
- [98349] GP-2, the major membrane protein in the secretory granule of the exocrine
- [98350] pancreas. *Gene* 171: 311–312, 1996.
- [98351] 17106.
- [98352] Miyoshi, J.; Higashi, T.; Mukai, H.; Ohuchi, T.; Kakunaga, T.:
- [98353] Structure and transforming potential of the human cot oncogene encoding
- [98354] a putative protein kinase. *Molec. Cell. Biol.* 11: 4088–4096, 1991.
- [98355] 17107.
- [98356] Ohara, R.; Miyoshi, J.; Aoki, M.; Toyoshima, K.: The murine cot
- [98357] proto-oncogene: genome structure and tissue-specific expression. *Jpn.*
- [98358] *J. Cancer Res.* 84: 518–525, 1993.
- [98359] 17108.
- [98360] Ishihara, H.; Shibasaki, Y.; Kizuki, N.; Wada, T.; Yazaki, Y.;
- [98361] Asano, T.; Oka, Y.: Type I phosphatidylinositol–

4-phosphate 5-kinases:

- [98362] cloning of the third isoform and deletion/substitution analysis of
- [98363] members of this novel lipid kinase family. *J. Biol. Chem.* 273: 8741-8748,
- [98364] 1998.
- [98365] 17109.
- [98366] Law, D. J.; Tarle, S. A.; Merchant, J. L.: The human ZBP-89 homolog,
- [98367] located at chromosome 3q21, represses gastrin gene expression. *Mammalian*
- [98368] *Genome* 9: 165-167, 1998.
- [98369] 17110.
- [98370] Merchant, J. L.; Iyer, G. R.; Taylor, B. R.; Kitchen, J. R.; Mortensen,
- [98371] E. R.; Wang, Z.; Flintoft, R. J.; Michel, J. B.; Bassel-Duby, R.:
- [98372] ZBP-89, a Kruppel-like zinc finger protein, inhibits epidermal growth
- [98373] factor induction of the gastrin promoter. *Molec. Cell. Biol.* 16:
- [98374] 6644-6653, 1996.

[98375] 17111.

[98376] Wang, Y.; Kobori, J. A.; Hood, L.: The ht-beta gene encodes a

[98377] novel CACCC box-binding protein that regulates T-cell receptor gene

[98378] expression. Molec. Cell. Biol. 13: 5691-5701, 1993.

[98379] 17112.

[98380] Everett, R. D.; Meredith, M.; Orr, A.; Cross, A.; Kathoria, M.;

[98381] Parkinson, J.: A novel ubiquitin-specific protease is dynamically

[98382] associated with the PML nuclear domain and binds to a herpesvirus

[98383] regulatory protein. EMBO J. 16: 1519-1530, 1997. Note: Corrected

[98384] republication of EMBO J.:16: 566-577, 1997.

[98385] 17113.

[98386] Li, M.; Chen, D.; Shiloh, A.; Luo, J.; Nikolaev, A. Y.; Qin, J.;

[98387] Gu, W.: Deubiquitination of p53 by HAUSP is an important pathway

[98388] for p53 stabilization. Nature 416: 648-653, 2002.

- [98389] 17114.
- [98390] Robinson, P. A.; Lomonte, P.; Leek, J. P.; Markham, A. F.;
Everett,
- [98391] R. D.: Assignment of herpesvirus-associated ubiquitin-specific protease
- [98392] gene HAUSP to human chromosome band 16p13.3 by in situ hybridization. Cytogenet.
- [98393] Cell Genet. 83: 100 only, 1998.
- [98394] 17115.
- [98395] Zhou, G.; Bao, Z. Q.; Dixon, J. E.: Components of a new human
- [98396] protein kinase signal transduction pathway. J. Biol. Chem. 270:
- [98397] 12665-12669, 1995.
- [98398] 17116.
- [98399] Lee, J.-D.; Ulevitch, R. J.; Han, J.: Primary structure of BMK1:
- [98400] a new mammalian MAP kinase. Biochem. Biophys. Res. Commun. 213:
- [98401] 715-724, 1995.
- [98402] 17117.

- [98403] Purandare, S. M.; Lee, J.-D.; Patel, P. I.: Assignment of big
- [98404] MAP kinase (PRKM7) to human chromosome 17 band
p11.2 with somatic
- [98405] cell hybrids. Cytogenet. Cell Genet. 83: 258–259, 1998.
- [98406] 17118.
- [98407] Regan, C. P.; Li, W.; Boucher, D. M.; Spatz, S.; Su, M. S.;
Kuida,
- [98408] K.: Erk5 null mice display multiple extraembryonic vascu-
lar and embryonic
- [98409] cardiovascular defects. Proc. Nat. Acad. Sci. 99:
9248–9253, 2002.
- [98410] 17119.
- [98411] Brennan, T. M. H.; Landau, D.; Shalev, H.; Lamb, F.;
Schutte, B.
- [98412] C.; Walder, R. Y.; Mark, A. L.; Carmi, R.; Sheffield, V. C.:
Linkage
- [98413] of infantile Bartter syndrome with sensorineural deafness
to chromosome
- [98414] 1p. Am. J. Hum. Genet. 62: 355–361, 1998.
- [98415] 17120.
- [98416] Birkenhager, R.; Otto, E.; Schurmann, M. J.; Vollmer, M.;

Ruf,

[98417] E.-M.; Maier-Lutz, I.; Beekmann, F.; Fekete, A.; Omran, H.;
Feldmann,

[98418] D.; Milford, D. V.; Jeck, N.; Konrad, M.; Landau, D.; Knoers,
N. V.

[98419] A. M.; Antignac, C.; Sudbrak, R.; Kispert, A.; Hildebrandt,
F.: Mutation

[98420] of BSND causes Bartter syndrome with sensorineural deaf-
ness and kidney

[98421] failure. *Nature Genet.* 29: 310–314, 2001.

[98422] 17121.

[98423] Barlow, G. M.; Chen, X.-N.; Shi, Z. Y.; Lyons, G. E.; Kurnit,
D.

[98424] M.; Celle, L.; Spinner, N. B.; Zackai, E.; Pettenati, M. J.; Van
Riper,

[98425] A. J.; Vekemans, M. J.; Mjaatvedt, C. H.; Korenberg, J. R.:
Down

[98426] syndrome congenital heart disease: a narrowed region and
a candidate

[98427] gene. *Genet. Med.* 3: 91–101, 2001.

[98428] 17122.

[98429] Schmucker, D.; Clemens, J. C.; Shu, H.; Worby, C. A.; Xiao,

J.;

[98430] Muda, M.; Dixon, J. E.; Zipursky, S. L.: *Drosophila Dscam is an axon*

[98431] *guidance receptor exhibiting extraordinary molecular diversity. Cell* 101:

[98432] 671–684, 2000.

[98433] 17123.

[98434] Yamakawa, K.; Huo, Y.-K.; Haendel, M. A.; Hubert, R.; Chen, X.-N.;

[98435] Lyons, G. E.; Korenberg, J. R.: *DSCAM: a novel member of the immunoglobulin*

[98436] *superfamily maps in a Down syndrome region and is involved in the*

[98437] *development of the nervous system. Hum. Molec. Genet.* 7: 227–237,

[98438] 1998.

[98439] 17124.

[98440] Rowles, J.; Scherer, S. W.; Xi, T.; Majer, M.; Nickle, D. C.; Rommens,

[98441] J. M.; Popov, K. M.; Harris, R. A.; Riebow, N. L.; Xia, J.; Tsui,

[98442] L.-C.; Bogardus, C.; Prochazka, M.: *Cloning and charac-*

terization

[98443] of PDK4 on 7q21.3 encoding a fourth pyruvate dehydrogenase kinase

[98444] isoenzyme in human. J. Biol. Chem. 271: 22376–22382, 1996.

[98445] 17125.

[98446] Dode, C.; Weil, D.; Levilliers, J.; Crozet, F.; Chaib, H.; Levi-Acobas,

[98447] F.; Guilford, P.; Petit, C.: Sequence characterization of a newly

[98448] identified human alpha-tubulin gene (TUBA2). Genomics 47: 125–130,

[98449] 1998.

[98450] 17126.

[98451] Bertolino, E.; Reimund, B.; Wildt-Perinic, D.; Clerc, R. G.: A

[98452] novel homeobox protein which recognizes a TGT core and functionally

[98453] interferes with a retinoid-responsive motif. J. Biol. Chem. 270:

[98454] 31178–31188, 1995.

[98455] 17127.

- [98456] Bertolino, E.; Wildt, S.; Richards, G.; Clerc, R. G.: Expression
of a novel murine homeobox gene in the developing cerebellar external
granular layer during its proliferation. Dev. Dyn. 205:
410–420,
1996.
- [98459] 17128.
- [98460] 17128.
- [98461] Gripp, K. W.; Edwards, M. C.; Mowat, D.; Meinecke, P.;
Richieri-Costa,
A.; Zackai, E. H.; Elledge, S.; Muenke, M.: Mutations in the
transcription
factor TGIF in holoprosencephaly. (Abstract) Am. J. Hum.
Genet. 63:
A32, 1998.
- [98464] 17129.
- [98465] 17129.
- [98466] Sulik, K. K.; Dehart, D. B.; Rogers, J. M.; Chernoff, N.: Teratogenicity
of low doses of all-trans retinoic acid in presomite mouse
embryos. Teratology 51:
398–403, 1995.

- [98469] 17130.
- [98470] Wotton, D.; Lo, R. S.; Lee, S.; Massague, J.: A Smad transcriptional
- [98471] corepressor. Cell 97: 29–39, 1999.
- [98472] 17131.
- [98473] Zhu, J.; Petersen, S.; Tessarollo, L.; Nussenzweig, A.: Targeted
- [98474] disruption of the Nijmegen breakage syndrome gene NBS1 leads to early
- [98475] embryonic lethality in mice. Curr. Biol. 11: 105–109, 2001.
- [98476] 17132.
- [98477] Cooper, P. R.; Smilnich, N. J.; Day, C. D.; Nowak, N. J.; Reid,
- [98478] L. H.; Pearsall, R. S.; Reece, M.; Prawitt, D.; Landers, J.; Housman,
- [98479] D. E.; Winterpacht, A.; Zabel, B. U.; Pelletier, J.; Weissman, B.
- [98480] E.; Shows, T. B.; Higgins, M. J.: Divergently transcribed overlapping
- [98481] genes expressed in liver and kidney and located in the 11p15.5 imprinted

- [98482] domain. Genomics 49: 38–51, 1998.
- [98483] 17133.
- [98484] Kershaw, D. B.; Beck, S. G.; Wharram, B. L.; Wiggins, J. E.;
Goyal,
- [98485] M.; Thomas, P. E.; Wiggins, R. C.: Molecular cloning and
characterization
- [98486] of human podocalyxin-like protein: orthologous relation-
ship to rabbit
- [98487] PCLP1 and rat podocalyxin. J. Biol. Chem. 272:
15708–15714, 1997.
- [98488] 17134.
- [98489] Kershaw, D. B.; Wiggins, J. E.; Wharram, B. L.; Wiggins, R.
C.
- [98490] : Assignment of the human podocalyxin-like protein
(PODXL) gene to
- [98491] 7q32–q33. Genomics 45: 239–240, 1997.
- [98492] 17135.
- [98493] Chan, K. K.; Tsui, S. K. W.; Lee, S. M. Y.; Luk, S. C. W.;
Liew,
- [98494] C. C.; Fung, K. P.; Waye, M. M. Y.; Lee, C. Y.: Molecular
cloning

- [98495] and characterization of FHL2, a novel LIM domain protein preferentially
- [98496] expressed in human heart. *Gene* 210: 345–350, 1998.
- [98497] 17136.
- [98498] Genini, M.; Schwalbe, P.; Scholl, F. A.; Remppis, A.; Mattei, M.-G.;
- [98499] Schafer, B. W.: Subtractive cloning and characterization of DRAL,
- [98500] a novel LIM-domain protein down-regulated in rhabdomyosarcoma. *DNA*
- [98501] *Cell Biol.* 16: 433–442, 1997.
- [98502] 17137.
- [98503] Prols, F.; Liehr, T.; Rinke, R.; Rautenstrauss, B.: Assignment
- [98504] of the microvascular endothelial differentiation gene 1 (MDG1) to
- [98505] human chromosome band 14q24.2–q24.3 by fluorescence in situ hybridization. *Cytogenet.*
- [98506] *Cell Genet.* 79: 149–150, 1997.
- [98507] 17138.
- [98508] Claverie-Martin, F.; Wang, M.; Cohen, S. N.: ARD-1 cDNA

from human

[98509] cells encodes a site-specific single-strand endoribonuclease that

[98510] functionally resembles Escherichia coli RNase E. J. Biol. Chem. 272:

[98511] 13823-13828, 1997.

[98512] 17139.

[98513] Van Eynde, A.; Wera, S.; Beullens, M.; Torrekens, S.; Van Leuven,

[98514] F.; Stalmans, W.; Bollen, M.: Molecular cloning of NIPP-1, a nuclear

[98515] inhibitor of protein phosphatase-1, reveals homology with polypeptides

[98516] involved in RNA processing. J. Biol. Chem. 270: 28068-28074, 1995.

[98517] 17140.

[98518] Tekin, M.; Dogu, F.; Tacyildiz, N.; Akar, E.; Ikinciogullari,

[98519] A.; Ogur, G.; Yavuz, G.; Babacan, E.; Akar, N.: 657del5 mutation

[98520] in the NBS1 gene is associated with Nijmegen breakage syndrome in

[98521] a Turkish family. Clin. Genet. 62: 84-88, 2002.

- [98522] 17141.
- [98523] Varon, R.; Reis, A.; Henze, G.; Einsiedel, H. G.; Sperling, K.;
- [98524] Seeger, K.: Mutations in the Nijmegen breakage syndrome
gene (NBS1)
- [98525] in childhood acute lymphoblastic leukemia (ALL). Cancer
Res. 61:
- [98526] 3570–3572, 2001.
- [98527] 17142.
- [98528] Wilda, M.; Demuth, I.; Concannon, P.; Sperling, K.;
- Hameister,
- [98529] H.: Expression pattern of the Nijmegen breakage syn-
drome gene, Nbs1,
- [98530] during murine development. Hum. Molec. Genet. 9:
1739–1744, 2000.
- [98531] 17143.
- [98532] Wu, X.; Ranganathan, V.; Weisman, D. S.; Heine, W. F.; Cic-
cone,
- [98533] D. N.; O'Neill, T. B.; Crick, K. E.; Pierce, K. A.; Lane, W. S.;
- Rathbun,
- [98534] G.; Livingston, D. M.; Weaver, D. T.: ATM phosphorylation
of Nijmegen

- [98535] breakage syndrome protein is required in a DNA damage response. Nature 405:
- [98536] 477–482, 2000.
- [98537] 17144.
- [98538] Zhao, S.; Weng, Y.–C.; Yuan, S.–S. F.; Lin, Y.–T.; Hsu, H.–C.;
- [98539] Lin, S.–C. J.; Gerbino, E.; Song, M.; Zdzenicka, M. Z.; Gatti, R.
- [98540] A.; Shay, J. W.; Ziv, Y.; Shiloh, Y.; Lee, E. Y.–H. P.: Functional
- [98541] link between ataxia–telangiectasia and Nijmegen breakage syndrome
- [98542] gene products. Nature 405: 473–477, 2000.
- [98543] 17145.
- [98544] Mankodi, A.; Urbinati, C. R.; Yuan, Q.–P.; Moxley, R. T.; Sansone,
- [98545] V.; Krym, M.; Henderson, D.; Schalling, M.; Swanson, M. S.; Thornton,
- [98546] C. A.: Muscleblind localizes to nuclear foci of aberrant RNA in myotonic
- [98547] dystrophy types 1 and 2. Hum. Molec. Genet. 10: 2165–2170, 2001.

- [98548] 17146.
- [98549] Hittner, H. M.; Kretzer, F. L.; Antoszyk, J. H.; Ferrell, R. E.;
- [98550] Mehta, R. S.: Variable expressivity of autosomal dominant
anterior
- [98551] segment mesenchymal dysgenesis in six generations. Am.
J. Ophthal. 93:
- [98552] 57–70, 1982.
- [98553] 17147.
- [98554] Semina, E. V.; Ferrell, R. E.; Mintz–Hittner, H. A.; Bitoun, P.;
- [98555] Alward, W. L. M.; Reiter, R. S.; Funkhauser, C.; Daack–
Hirsch, S.;
- [98556] Murray, J. C.: A novel homeobox gene PITX3 is mutated in
families
- [98557] with autosomal–dominant cataracts and ASMD. Nature
Genet. 19: 167–170,
- [98558] 1998.
- [98559] 17148.
- [98560] Semina, E. V.; Murray, J. C.; Reiter, R.; Hrstka, R. F.; Graw,
[98561] J.: Deletion in the promoter region and altered expression
of Pitx3

- [98562] homeobox gene in aphakia mice. Hum. Molec. Genet. 9:
1575–1585,
[98563] 2000.
[98564] 17149.
- [98565] Semina, E. V.; Reiter, R. S.; Murray, J. C.: Isolation of a new
[98566] homeobox gene belonging to the Pitx/Rieg family: ex-
pression during
[98567] lens development and mapping to the aphakia region on
mouse chromosome
[98568] 19. Hum. Molec. Genet. 6: 2109–2116, 1997.
[98569] 17150.
- [98570] Lin, C.-S.; Shen, W.; Chen, Z. P.; Tu, Y.-H.; Matsudaira, P.:
[98571] Identification of I-plastin, a human fimbrin isoform ex-
pressed in
[98572] intestine and kidney. Molec. Cell. Biol. 14: 2457–2467,
1994.
[98573] 17151.
- [98574] Kent, J.; Lee, M.; Schedl, A.; Boyle, S.; Fantes, J.; Powell, M.;
[98575] Rushmere, N.; Abbott, C.; van Heyningen, V.; Bickmore, W.
A.: The

- [98576] reticulocalbin gene maps to the WAGR region in human and to the Small
- [98577] eye Harwell deletion in mouse. *Genomics* 42: 260–267, 1997.
- [98578] 17152.
- [98579] Ozawa, M.: Cloning of a human homologue of mouse reticulocalbin
- [98580] reveals conservation of structural domains in the novel endoplasmic
- [98581] reticulum resident Ca^{2+} -binding protein with multiple EF-hand motifs. *J.*
- [98582] *Biochem.* 117: 1113–1119, 1995.
- [98583] 17153.
- [98584] Ozawa, M.; Muramatsu, T.: Reticulocalbin, a novel endoplasmic
- [98585] reticulum resident Ca^{2+} -binding protein with multiple EF-hand motifs
- [98586] and a carboxyl-terminal HDEL sequence. *J. Biol. Chem.* 268: 699–705,
- [98587] 1993.
- [98588] 17154.

- [98589] Yan, W. L.; Lerner, T. J.; Haines, J. L.; Gusella, J. F.: Sequence
[98590] analysis and mapping of a novel human mitochondrial
ATP synthase subunit
[98591] 9 cDNA (ATP5G3). Genomics 24: 375–377, 1994.
[98592] 17155.
[98593] Aho, S.; McLean, W. H. I.; Li, K.; Uitto, J.: cDNA cloning,
mRNA
[98594] expression, and chromosomal mapping of human and
mouse periplakin
[98595] genes. Genomics 48: 242–247, 1998.
[98596] 17156.
[98597] Luo, G.; Leroy, E.; Kozak, C. A.; Polymeropoulos, M. H.;
Horowitz,
[98598] R.: Mapping of the gene (NRAP) encoding N-RAP in the
mouse and human
[98599] genomes. Genomics 45: 229–232, 1997.
[98600] 17157.
[98601] Luo, G.; Zhang, J. Q.; Nguyen, T.-P.; Herrera, A. H.; Pater-
son,
[98602] B.; Horowitz, R.: Complete cDNA sequence and tissue lo-

calization

[98603] of N-RAP, a novel nebulin-related protein of striated muscle. Cell

[98604] Motil. Cytoskeleton 38: 75-90, 1997.

[98605] 17158.

[98606] Ando-Akatsuka, Y.; Saitou, M.; Hirase, T.; Kishi, M.; Sakakibara,

[98607] A.; Itoh, M.; Yonemura, S.; Furuse, M.; Tsukita, S.: Inter-species

[98608] diversity of the occludin sequence: cDNA cloning of human, mouse,

[98609] dog, and rat-kangaroo homologues. J. Cell Biol. 133: 43-47, 1996.

[98610] 17159.

[98611] Strahm, B.; Rittweiler, K.; Duffner, U.; Brandau, O.; Orłowska-Volk,

[98612] M.; Karajannis, M. A.; zur Stadt, U.; Tiemann, M.; Reiter, A.; Brandis,

[98613] M.; Meindl, A.; Niemeyer, C. M.: Recurrent B-cell non-Hodgkin's lymphoma

[98614] in two brothers with X-linked lymphoproliferative disease without

- [98615] evidence for Epstein-Barr virus infection. *Brit. J. Haemat.* 108:
- [98616] 377-382, 2000.
- [98617] 17160.
- [98618] Sylla, B. S.; Murphy, K.; Cahir-McFarland, E.; Lane, W. S.; Mosialos,
- [98619] G.; Kieff, E.: The X-linked lymphoproliferative syndrome gene product
- [98620] SH2D1A associates with p62(dok) (Dok1) and activates NF-kappa-beta. *Proc.*
- [98621] *Nat. Acad. Sci.* 97: 7470-7475, 2000.
- [98622] 17161.
- [98623] Sylla, B. S.; Wang, Q.; Hayoz, D.; Lathrop, G. M.; Lenoir, G.
- [98624] M.: Multipoint linkage mapping of the Xq25-q26 region in a family
- [98625] affected by the X-linked lymphoproliferative syndrome. *Clin. Genet.* 36:
- [98626] 359-462, 1989.
- [98627] 17162.
- [98628] Tangye, S. G.; Lazetic, S.; Woollatt, E.; Sutherland, G. R.; Lanier,

- [98629] L. L.; Phillips, J. H.: Cutting edge: human 2B4, an activating NK
- [98630] cell receptor, recruits the protein tyrosine phosphatase SHP-2 and
- [98631] the adaptor signaling protein SAP. J. Immun. 162: 6981-6985, 1999.
- [98632] 17163.
- [98633] Tangye, S. G.; Phillips, J. H.; Lanier, L. L.; Nichols, K. E.
- [98634] : Cutting edge: functional requirement for SAP in 2B4-mediated activation
- [98635] of human natural killer cells as revealed by the X-linked lymphoproliferative
- [98636] syndrome. J. Immun. 165: 2932-2936, 2000.
- [98637] 17164.
- [98638] Thorley-Lawson, D. A.; Schooley, R. T.; Bhan, A. K.; Nadler, L.
- [98639] M.: Epstein-Barr virus superinduces a new human B cell differentiation
- [98640] antigen (B-LAST 1) expressed on transformed lymphoblasts. Cell 30:
- [98641] 415-425, 1982.

- [98642] 17165.
- [98643] Vowels, M. R.; Lam-Po-Tang, R.; Berdoukas, V.; Ford, D.;
Thierry,
- [98644] D.; Purtilo, D.; Gluckman, E.: Correction of X-linked lym-
phoproliferative
- [98645] disease by transplantation of cord-blood stem cells. New
Eng. J.
- [98646] Med. 329: 1623-1625, 1993.
- [98647] 17166.
- [98648] Williams, L. L.; Rooney, C. M.; Conley, M. E.; Brenner, M.
K.;
- [98649] Krance, R. A.; Heslop, H. E.: Correction of Duncan's syn-
drome by
- [98650] allogeneic bone marrow transplantation. Lancet 342:
587-588, 1993.
- [98651] 17167.
- [98652] Wu, C.; Nguyen, K. B.; Pien, G. C.; Wang, N.; Gullo, C.;
Duncan,
- [98653] H.; Sosa, M. R.; Edwards, M. J.; Borrow, P.; Satoskar, A. R.;
Sharpe,
- [98654] A. H.; Biron, C. A.; Terhorst, C.: SAP controls T cell re-
sponses

- [98655] to virus and terminal differentiation of T(H)2 cells. *Nature Immun.* 2:
410–414, 2001.
- [98656] 17168.
- [98657] Wyandt, H. E.; Skare, J. C.; Grierson, H. L.; Purtilo, D. T.;
- [98658] Milunsky, A.: Detection of a chromosomal deletion of Xq25 in an affected male with X-linked lymphoproliferative disease. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A108 only, 1989.
- [98659] 17169.
- [98660] Yin, L.; Ferrand, V.; Lavoue, M.-F.; Hayoz, D.; Philippe, N.;
- [98661] Souillet, G.; Seri, M.; Giacchino, R.; Castagnola, E.; Hodgson, S.;
- [98662] Sylla, B. S.; Romeo, G.: SH2D1A mutation analysis for diagnosis of XLP in typical and atypical patients. *Hum. Genet.* 105: 501–505, 1999.
- [98663] 17170.

[98669] Yin, L.; Tocco, T.; Pauly, S.; Lenoir, G. M.; Romeo, G.: Absence

[98670] of SH2D1A point mutation in 62 Burkitts lymphoma cell lines. Am.

[98671] J. Hum. Genet. 65 (suppl. 1868): A331 only, 1999.

[98672] 17171.

[98673] Berglund, E. O.; Ranscht, B.: Molecular cloning and in situ localization

[98674] of the human contactin gene (CNTN1) on chromosome 12q11-q12. Genomics 21:

[98675] 571-582, 1994.

[98676] 17172.

[98677] Laxminarayan, K. M.; Chan, B. K.; Tetaz, T.; Bird, P. I.; Mitchell,

[98678] C. A.: Characterization of a cDNA encoding the 43-kDa membrane-associated

[98679] inositol-polyphosphate 5-phosphatase. J. Biol. Chem. 269: 17305-17310,

[98680] 1994.

[98681] 17173.

[98682] Fink, J. K.; Jones, S. M.; Esposito, C.; Wilkowski, J.: Human

- [98683] microtubule-associated protein 1a (MAP1A) gene: genomic organization,
- [98684] cDNA sequence, and developmental-and tissue-specific expression. *Genomics* 35:
- [98685] 577-585, 1996.
- [98686] 17174.
- [98687] Quinton, R.; Duke, V. M.; de Zoysa, P. A.; Platts, A. D.; Valentine,
- [98688] A.; Kendall, B.; Pickman, S.; Kirk, J. M. W.; Besser, G. M.; Jacobs,
- [98689] H. S.; Bouloux, P. M. G.: The neuroradiology of Kallmann's syndrome:
- [98690] a genotypic and phenotypic analysis. *J. Clin. Endocr. Metab.* 81:
- [98691] 3010-3017, 1996.
- [98692] 17175.
- [98693] Rogol, A.; Mittal, K.; White, B. J.; McGinnis, M. H.; Liebllich,
- [98694] J. M.; Rosen, S. W.: HLA-compatible paternity in two 'fertile eunuchs'
- [98695] with congenital hypogonadotrophic hypogonadism and anosmia (the Kallmann

- [98696] syndrome). J. Clin. Endocr. Metab. 51: 275–279, 1980.
- [98697] 17176.
- [98698] Befort, K.; Filliol, D.; Decaillot, F. M.; Gaveriaux–Ruff, C.;
- [98699] Hoehe, M. R; Kieffer, B. L.: A single nucleotide polymorphic mutation
- [98700] in the human mu–opioid receptor severely impairs receptor signaling. J.
- [98701] Biol. Chem. 276: 3130–3137, 2001.
- [98702] 17177.
- [98703] Bond, C.; LaForge, K. S.; Tian, M.; Melia, D.; Zhang, S.; Borg,
- [98704] L.; Gong, J.; Schluger, J.; Strong, J. A.; Leal, S. M.; Tischfield,
- [98705] J. A.; Kreek, M. J.; Yu, L.: Single–nucleotide polymorphism in the
- [98706] human mu opioid receptor gene alters beta–endorphin binding and activity:
- [98707] possible implications for opiate addiction. Proc. Nat. Acad. Sci. 95:
- [98708] 9608–9613, 1998.
- [98709] 17178.

- [98710] Chen, Y.; Mestek, A.; Liu, J.; Hurley, J. A.; Yu, L.: Molecular
[98711] cloning and functional expression of a mu-opioid recep-
tor from rat
[98712] brain. *Molec. Pharm.* 44: 8–12, 1993.
[98713] 17179.
- [98714] He, L.; Fong, J.; von Zastrow, M.; Whistler, J. L.: Regulation
[98715] of opioid receptor trafficking and morphine tolerance by
receptor
[98716] oligomerization. *Cell* 108: 271–282, 2002.
[98717] 17180.
- [98718] Hoehe, M. R.; Kopke, K.; Wendel, B.; Rohde, K.; Flach-
meier, C.;
[98719] Kidd, K. K.; Berrettini, W. H.; Church, G. M.: Sequence vari-
ability
[98720] and candidate gene analysis in complex disease: associa-
tion of mu
[98721] opioid receptor gene variation with substance depen-
dence. *Hum. Molec.*
[98722] *Genet.* 9: 2895–2908, 2000.
[98723] 17181.

- [98724] Holtt, V.: A polymorphism (A118G) in the mu-opioid receptor gene
- [98725] affects the response to morphine-6-glucuronide in humans. Pharmacogenetics 12:
- [98726] 1-2, 2002.
- [98727] 17182.
- [98728] LaForge, K. S.; Yuferov, V.; Kreek, M. J.: Opioid receptor and
- [98729] peptide gene polymorphisms: potential implications for addictions. Europ.
- [98730] J. Pharm. 410: 249-268, 2000.
- [98731] 17183.
- [98732] Lotsch, J.; Skarke, C.; Grosch, S.; Darimont, J.; Schmidt, H.;
- [98733] Geisslinger, G.: The polymorphism A118G of the human mu-opioid receptor
- [98734] gene decreases the pupil constrictory effect of morphine-6-glucuronide
- [98735] but not that of morphine. Pharmacogenetics 12: 3-9, 2002.
- [98736] 17184.
- [98737] Mansour, A.; Fox, C. A.; Burke, S.; Akil, H.; Watson, S. J.:

- [98738] Immunohistochemical localization of the clone mu opioid receptor in
- [98739] the rat CNS. J. Chem. Neuroanat. 8: 283–305, 1995.
- [98740] 17185.
- [98741] Meyer, U. A.: Pharmacogenetics and adverse drug reactions. Lancet 356:
- [98742] 1667–1671, 2000.
- [98743] 17186.
- [98744] Mogil, J. S.: The genetic mediation of individual differences
- [98745] in sensitivity to pain and its inhibition. Proc. Nat. Acad. Sci. 96:
- [98746] 7744–7751, 1999.
- [98747] 17187.
- [98748] Uhl, G. R.; Sora, I.; Wang, Z.: The mu opiate receptor as a candidate
- [98749] gene for pain: polymorphisms, variations in expression, nociception,
- [98750] and opiate responses. Proc. Nat. Acad. Sci. 96:
- [98751] 7752–7755, 1999.
- [98751] 17188.

- [98752] Wang, J.-B.; Johnson, P. S.; Persico, A. M.; Hawkins, A. L.; Griffin,
[98753] C. A.; Uhl, G. R.: Human mu opiate receptor: cDNA and
genomic clones,
[98754] pharmacologic characterization and chromosomal assign-
ment. FEBS Lett. 338:
[98755] 217-222, 1994.
[98756] 17189.
- [98757] Wu, D.; Kang, Y.-S.; Bickel, U.; Pardridge, W. M.: Blood-
brain
[98758] barrier permeability to morphine-6-glucuronide is
markedly reduced
[98759] compared with morphine. Drug Metab. Dispos. 25:
768-771, 1997.
[98760] 17190.
- [98761] Zubietta, J.-K.; Smith, Y. R.; Bueller, J. A.; Xu, Y.; Kilbourn,
[98762] M. R.; Jewett, D. M.; Meyer, C. R.; Koeppe, R. A.; Stohler, C.
S.
[98763] : Regional mu opioid receptor regulation of sensory and
affective
[98764] dimensions of pain. Science 293: 311-315, 2001.

[98765] 17191.

[98766] McBride, O. W.; Yi, H. F.; Srivastava, M.: The human cytochrome

[98767] b561 gene (CYB561) is located at 17q11-qter. Genomics 21: 662-663,

[98768] 1994.

[98769] 17192.

[98770] Srivastava, M.: Genomic structure and expression of the human

[98771] gene encoding cytochrome b(561), an integral protein of the cromaffin

[98772] granule membrane. J. Biol. Chem. 270: 22714-22720, 1995.

[98773] 17193.

[98774] Albarosa, R.; DiDonato, S.; Finocchiaro, G.: Redefinition of the

[98775] coding sequence of the MXI1 gene and identification of a polymorphic

[98776] repeat in the 3-prime non-coding region that allows the detection

[98777] of loss of heterozygosity of chromosome 10q25 in glioblastomas. Hum.

- [98778] Genet. 95: 709–711, 1995.
- [98779] 17194.
- [98780] Mitchell, C. A.; Speed, C. J.; Nicholl, J.; Sutherland, G. R.:
- [98781] Chromosomal mapping of the gene (INPP5A) encoding the
43-kDa membrane-associated
- [98782] inositol polyphosphate 5-phosphatase to 10q26.3 by flu-
orescence in
- [98783] situ hybridization. Genomics 31: 139–140, 1996.
- [98784] 17195.
- [98785] Phillips, D. R.; Agin, R. P.: Platelet membrane defects in
Glanzmann's
- [98786] thrombasthenia: evidence for decreased amounts of two
major glycoproteins. J.
- [98787] Clin. Invest. 60: 535–545, 1977.
- [98788] 17196.
- [98789] Pittman, M. A., Jr.; Graham, J. B.: Glanzmann's thrombopa-
thy:
- [98790] an autosomal recessive trait in one family. Am. J. Med. Sci.
247:
- [98791] 293–303, 1964.

- [98792] 17197.
- [98793] Poncz, M.; Rifat, S.; Coller, B. S.; Newman, P. J.; Shattil, S. J.; Parrella, T.; Fortina, P.; Bennett, J. S.: Glanzmann thrombasthenia secondary to a gly273-to-aspartate mutation adjacent to the first calcium-binding domain of platelet glycoprotein IIb. J. Clin. Invest. 93: 172-179, 1994.
- [98797] 17198.
- [98798] 17198.
- [98799] Prandini, M. H.; Denarier, E.; Frachet, P.; Uzan, G.; Marguerie, G.: Isolation of the human platelet glycoprotein IIb gene and characterization of the 5-prime flanking region. Biochem. Biophys. Res. Commun. 156: 595-601, 1988.
- [98802] 595-601, 1988.
- [98803] 17199.
- [98804] Pytela, R.; Pierschbacher, M. D.; Ginsberg, M. H.; Plow, E. F.;
- [98805] Ruoslahti, E.: Platelet membrane glycoprotein IIb/IIIa:

member of

[98806] a family of arg-gly-asp-specific adhesion receptors. Science 231:

[98807] 1559-1562, 1986.

[98808] 17200.

[98809] Ruan, J.; Peyruchaud, O.; Alberio, L.; Valles, G.; Clemetson,

[98810] K.; Bourre, F.; Nurden, A. T.: Double heterozygosity of the GPIIb

[98811] gene in a Swiss patient with Glanzmann's thrombasthenia. Brit. J.

[98812] Haemat. 102: 918-925, 1998.

[98813] 17201.

[98814] Russell, M. E.; Seligsohn, U.; Coller, B. S.; Ginsberg, M. H.;

[98815] Skoglund, P.; Quertermous, T.: Structural integrity of the glycoprotein

[98816] IIb and IIIa genes in Glanzmann thrombasthenia patients from Israel. Blood 72:

[98817] 1833-1836, 1988.

[98818] 17202.

[98819] Schlegel, N.; Gayet, O.; Morel-Kopp, M.-C.; Wyler, B.; Hur-

taud-Roux,

- [98820] M.-F.; Kaplan, C.; McGregor, J.: The molecular genetic basis of Glanzmann's
- [98821] thrombasthenia in a gypsy population in France: identification of
- [98822] a new mutation on the alpha(IIB) gene. Blood 86: 977-982, 1995.
- [98823] 17203.
- [98824] Seligsohn, U.; Mibashan, R. S.; Rodeck, C. H.; Nicolaides, K.
- [98825] H.; Millar, D. S.; Coller, B. S.: Prenatal diagnosis of Glanzmann's
- [98826] thrombasthenia. (Letter) Lancet II: 1419 only, 1985.
- [98827] 17204.
- [98828] Sosnoski, D.; Emanuel, B. S.; Hawkins, A. L.; van Tuinen, P.;
- [98829] Ledbetter, D. H.; Nussbaum, R. L.; Kaos, F.-T.; Schwartz, E.; Phillips,
- [98830] D.; Bennett, J. S.; Fitzgerald, L. A.; Poncz, M.: Chromosomal localization
- [98831] of the genes for the vitronectin and fibronectin receptors alpha-subunits

- [98832] and for platelet glycoproteins IIb and IIIa. J. Clin. Invest. 81:
- [98833] 1993–1998, 1988.
- [98834] 17205.
- [98835] Tongio, M. M.; Lutz, P.; Hauptmann, G.; Rodier, L.; Levy, J.-M.;
- [98836] Mayer, S.; Cazenave, J.-P.: Type I Glanzmann's thrombasthenia segregates
- [98837] independently of Ss and Duffy systems and the A, B, C, factor B, C2
- [98838] and C4 loci of the HLA complex. Tissue Antigens 20: 22–27, 1982.
- [98839] 17206.
- [98840] Uzan, G.; Van Cong, N.; Gross, M. S.; Jegou-Foubert, C.; Frachet,
- [98841] P.; Boucheix, C.; Margerie, G.; Frezal, J.: Assignment of a gene
- [98842] coding for human platelet GPIIb (GP2B) to 17q21.1–q21.3 by in situ
- [98843] hybridization. (Abstract) Cytogenet. Cell Genet. 46: 707 only, 1987.
- [98844] 17207.

- [98845] Van Cong, N.; Uzan, G.; Gross, M. S.; Jegou-Foubert, C.; Frachet,
- [98846] P.; Boucheix, C.; Marguerie, G.; Frezal, J.: Assignment of human
- [98847] platelet GP2B (GPIIb) gene to chromosome 17, region q21.1–q21.3. Hum.
- [98848] Genet. 80: 389–392, 1988.
- [98849] 17208.
- [98850] von dem Borne, A. E. G. K.; von Riesz, E.; Verheugt, F. W. A.;
- [98851] ten Cate, J. W.; Koppe, J. G.; Engelfriet, C. P.; Nijenhuis, L. E.
- [98852] : Bak(a), a new platelet-specific antigen involved in neonatal allo-immune
- [98853] thrombocytopenia. Vox Sang. 39: 113–120, 1980.
- [98854] 17209.
- [98855] Waller, H. D.; Gross, R.: Genetische Enzymdefecte als Ursache
- [98856] von Thrombocytopathien. Verh. Dtsch. Ges. Inn. Med. 70: 476–494,
- [98857] 1964.

[98858] 17210.

- [98859] Wilcox, D. A.; Gill, J.; Newman, P. J.: Glanzmann thrombasthenia
- [98860] resulting from a single amino acid substitution flanking the fibrinogen
- [98861] gamma-chain dodecapeptide binding domain on GPIIb. (Abstract) Blood 82:
- [98862] 210a only, 1993.
- [98863] 17211.
- [98864] Wilcox, D. A.; Paddock, C. M.; Lyman, S.; Gill, J. C.; Newman,
- [98865] P. J.: Glanzmann thrombasthenia resulting from a single amino acid
- [98866] substitution between the second and third calcium-binding domains
- [98867] of GPIIb: role of the GPIIb amino terminus in integrin subunit association. J.
- [98868] Clin. Invest. 95: 1553 only, 1995.
- [98869] 17212.
- [98870] Wilcox, D. A.; Wautier, J. L.; Pidard, D.; Newman, P. J.: A single
- [98871] amino acid substitution flanking the fourth calcium binding domain

- [98872] of alpha-IIb prevents maturation of the alpha-IIb-beta-3 integrin
- [98873] complex. J. Biol. Chem. 269: 4450-4457, 1994.
- [98874] 17213.
- [98875] Zaizov, R.; Cohen, I.; Matoth, Y.: Thrombasthenia: a study of
- [98876] two siblings. Acta Paediat. Scand. 57: 522-526, 1968.
- [98877] 17214.
- [98878] Dreyer, B.; Tranebjaerg, L.; Rosenberg, T.; Weston, M. D.; Kimberling,
- [98879] W. J.; Nilssen, O.: Identification of novel USH2A mutations: implications
- [98880] for the structure of USH2A protein. Europ. J. Hum. Genet. 8: 500-506,
- [98881] 2000.
- [98882] 17215.
- [98883] Eudy, J. D.; Weston, M. D.; Yao, S.; Hoover, D. M.; Rehm, H. L.;
- [98884] Ma-Edmonds, M.; Yan, D.; Ahmad, I.; Cheng, J. J.; Ayuso, C.; Cremers,
- [98885] C.; Davenport, S.; Moller, C.; Talmadge, C. B.; Beisel, K. W.;

Tamayo,

[98886] M.; Morton, C. C.; Swaroop, A.; Kimberling, W. J.; Sumegi, J.: Mutation

[98887] of a gene encoding a protein with extracellular matrix motifs in Usher

[98888] syndrome type IIa. Science 280: 1753–1757, 1998.

[98889] 17216.

[98890] Huang, D.; Eudy, J. D.; Uzvolgyi, E.; Davis, J. R.; Talmadge, C.

[98891] B.; Pretto, D.; Weston, M. D.; Lehman, J. E.; Zhou, M.; Seemayer,

[98892] T. A.; Ahmad, I.; Kimberling, W. J.; Sumegi, J.: Identification of

[98893] the mouse and rat orthologs of the gene mutated in Usher syndrome

[98894] type IIA and the cellular source of USH2A mRNA in retina, a target

[98895] tissue of the disease. Genomics 80: 195–203, 2002.

[98896] 17217.

[98897] Kimberling, W. J.; Weston, M. D.; Moller, C.; Davenport, S. L.

[98898] H.; Shugart, Y. Y.; Priluck, I. A.; Martini, A.; Milani, M.;

Smith,

[98899] R. J.: Localization of Usher syndrome type II to chromosome 1q. Genomics 7:

[98900] 245–249, 1990.

[98901] 17218.

[98902] Kimberling, W. J.; Weston, M. D.; Moller, C.; van Aarem, A.; Cremers,

[98903] C. W. R. J.; Sumegi, J.; Ing, P. S.; Connolly, C.; Martini, A.; Milani,

[98904] M.; Tamayo, M. L.; Bernal, J.; Greenberg, J.; Ayuso, C.: Gene mapping

[98905] of Usher syndrome type IIa: localization of the gene to a 2.1-cM segment

[98906] on chromosome 1q41. Am. J. Hum. Genet. 56: 216–223, 1995.

[98907] 17219.

[98908] Lewis, R. A.; Otterud, B.; Stauffer, D.; Lalouel, J.-M.; Lerpert,

[98909] M.: Mapping recessive ophthalmic diseases: linkage of the locus for

[98910] Usher syndrome type II to a DNA marker on chromosome 1q. Genomics 7:

- [98911] 250–256, 1990.
- [98912] 17220.
- [98913] Liu, X.–Z.; Hope, C.; Liang, C. Y.; Zou, J. M.; Xu, L. R.; Cole,
[98914] T.; Mueller, R. F.; Bunday, S.; Nance, W.; Steel, K. P.;
Brown, S.
[98915] D. M.: A mutation (2314delG) in the Usher syndrome type
IIA gene:
[98916] high prevalence and phenotypic variation. (Letter) Am. J.
Hum. Genet. 64:
[98917] 1221–1225, 1999.
- [98918] 17221.
- [98919] Pieke–Dahl, S.; van Aarem, A.; Dobin, A.; Cremers, C. W. R.
J.;
- [98920] Kimberling, W. J.: Genetic heterogeneity of Usher syn-
drome type II
[98921] in a Dutch population. J. Med. Genet. 33: 753–757, 1996.
- [98922] 17222.
- [98923] Pieke Dahl, S.; Weston, M. D.; Kimberling, W. J.; Gorin, M.
B.;
- [98924] Shugart, Y. Y.; Kenyon, J. B.: Possible genetic heterogene-
ity of

- [98925] Usher syndrome type 2: a family unlinked to chromosome 1q markers.
- [98926] (Abstract) Am. J. Hum. Genet. 49 (suppl.): 200 only, 1991.
- [98927] 17223.
- [98928] Rivolta, C.; Sweklo, E. A.; Berson, E. L.; Dryja, T. P.: Mis-sense
- [98929] mutation in the USH2A gene: association with recessive retinitis pigmentosa
- [98930] without hearing loss. Am. J. Hum. Genet. 66: 1975–1978, 2000.
- [98931] 17224.
- [98932] Robbins, J. H.; Scudiero, D. A.; Otsuka, F.; Tarone, R. E.; Brumback,
- [98933] R. A.; Wirtschafter, J. D.; Polinsky, R. J.; Barrett, S. F.; Moshell,
- [98934] A. N.; Scarpinato, R. G.; Ganges, M. B.; Nee, L. E.; Meyer, S. A.;
- [98935] Clatterbuck, B. E.: Hypersensitivity to DNA-damaging agents in cultured
- [98936] cells from patients with Usher's syndrome and Duchenne muscular dystrophy. J.
- [98937] Neurol. Neurosurg. Psychiat. 47: 391–398, 1984.

- [98938] 17225.
- [98939] Smith, R. J. H.; Lee, E. C.; Kimberling, W. J.; Daiger, S. P.;
- [98940] Pelias, M. Z.; Keats, B. J. B.; Jay, M.; Bird, A.; Reardon, W.;
Guest,
- [98941] M.; Ayyagari, R.; Hejtmancik, J. F.: Localization of two
genes for
- [98942] Usher syndrome type I to chromosome 11. *Genomics* 14:
995–1002, 1992.
- [98943] 17226.
- [98944] Weston, M. D.; Eudy, J. D.; Fujita, S.; Yao, S.-F.; Usami, S.;
- [98945] Cremers, C.; Greenburg, J.; Ramesar, R.; Martini, A.;
Moller, C.;
- [98946] Smith, R. J.; Sumegi, J.; Kimberling, W. J.: Genomic struc-
ture and
- [98947] identification of novel mutations in usherin, the gene re-
sponsible
- [98948] for Usher syndrome type IIa. *Am. J. Hum. Genet.* 66:
1199–1210, 2000.
- [98949] Note: Erratum: *Am. J. Hum. Genet.* 66: 2020 only, 2000.
- [98950] 17227.
- [98951] Gasparini, P.; De Fazio, A.; Croce, A. I.; Stanziale, P.; Ze-

lante,

[98952] L.: Usher syndrome type III (USH3) linked to chromosome 3q in an

[98953] Italian family. J. Med. Genet. 35: 666–667, 1998.

[98954] 17228.

[98955] Joensuu, T.; Hamalainen, R.; Yuan, B.; Johnson, C.; Tegelberg,

[98956] S.; Gasparini, P.; Zelante, L.; Pirvola, U.; Parakinen, L.; Lehesjoki,

[98957] A.–E.; de la Chapelle, A.; Sankila, E.–M.: Mutations in a novel gene

[98958] with transmembrane domains underlie Usher syndrome type 3. Am. J.

[98959] Hum. Genet. 69: 673–684, 2001. Note: Erratum: Am. J. Hum. Genet.

[98960] 69: 1160 only, 2001.

[98961] 17229.

[98962] Sankila, E.–M.; Pakarinen, L.; Kaariainen, H.; Aittomaki, K.; Karjalainen,

[98963] S.; Sistonen, P.; de la Chapelle, A.: Assignment of an Usher syndrome

[98964] type III (USH3) gene to chromosome 3q. Hum. Molec.

Genet. 4: 93–98,

[98965] 1995.

[98966] 17230.

[98967] Anantharaman, V.; Koonin, E. V.; Aravind, L.: Peptide–
N-glycanases

[98968] and DNA repair proteins, Xp-C/Rad4, are, respectively,
active and

[98969] inactivated enzymes sharing a common transglutaminase
fold. Hum.

[98970] Molec. Genet. 10: 1627–1630, 2001.

[98971] 17231.

[98972] Cheo, D. L.; Burns, D. K.; Meira, L. B.; Houle, J. F.; Fried–
berg,

[98973] E. C.: Mutational inactivation of the xeroderma pigmento–
sum group

[98974] C gene confers predisposition to
2-acetylaminofluorene-induced liver

[98975] and lung cancer and to spontaneous testicular cancer in
Trp53 –/–

[98976] mice. Cancer Res. 59: 771–775, 1999.

[98977] 17232.

- [98978] Cleaver, J. E.: DNA repair in human xeroderma pigmentosum group
- [98979] C cells involves a different distribution of damaged sites in confluent
- [98980] and growing cells. *Nucleic Acids Res.* 14: 8155–8165, 1986.
- [98981] 17233.
- [98982] Emmert, S.; Kobayashi, N.; Khan, S. G.; Kraemer, K. H.: The xeroderma
- [98983] pigmentosum group C gene leads to selective repair of cyclobutane
- [98984] pyrimidine dimers rather than 6–4 photoproducts. *Proc. Nat. Acad.*
- [98985] *Sci.* 97: 2151–2156, 2000.
- [98986] 17234.
- [98987] Halley, D. J. J.; Keijzer, W.; Jaspers, N. G. J.; Niermeijer, M.
- [98988] F.; Kleijer, W. J.; Boue, J.; Boue, A.; Bootsma, D.: Prenatal diagnosis
- [98989] of xeroderma pigmentosum (group C) using assays of unscheduled DNA
- [98990] synthesis and postreplication repair. *Clin. Genet.* 16: 137–146,

[98991] 1979.

[98992] 17235.

[98993] Hananian, J.; Cleaver, J. E.: Xeroderma pigmentosum ex-
hibiting

[98994] neurological disorders and systemic lupus erythematosus.
Clin. Genet. 17:

[98995] 39-45, 1980.

[98996] 17236.

[98997] Kaur, G. P.; Athwal, R. S.: Identification of a DNA repair
gene

[98998] on human chromosome 5 that complements xeroderma
pigmentosum cells

[98999] of group C. (Abstract) Am. J. Hum. Genet. 51 (suppl.):
A51, 1992.

[99000] 17237.

[99001] Khan, S. G.; Levy, H. L.; Legerski, R.; Quackenbush, E.;
Reardon,

[99002] J. T.; Emmert, S.; Sancar, A.; Li, L.; Schneider, T. D.;
Cleaver,

[99003] J. E.; Kraemer, K. H.: Xeroderma pigmentosum group C
splice mutation

- [99004] associated with autism and hypoglycinemia. J. Invest. Derm. 111:
- [99005] 791–796, 1998.
- [99006] 17238.
- [99007] Legerski, R.; Peterson, C.: Expression cloning of a human DNA
- [99008] repair gene involved in xeroderma pigmentosum group C. Nature 359:
- [99009] 70–73, 1992.
- [99010] 17239.
- [99011] Legerski, R. J.; Liu, P.; Li, L.; Peterson, C. A.; Zhao, Y.; Leach,
- [99012] R. J.; Naylor, S. L.; Siciliano, M. J.: Assignment of xeroderma pigmentosum
- [99013] group C (XPC) gene to chromosome 3p25. Genomics 21: 266–269, 1994.
- [99014] 17240.
- [99015] Li, L.; Bales, E. S.; Peterson, C. A.; Legerski, R. J.: Characterization
- [99016] of molecular defects in xeroderma pigmentosum group C. Nature Genet. 5:

- [99017] 413–417, 1993.
- [99018] 17241.
- [99019] Lynch, H. T.; Fusaro, R. M.; Johnson, J. A.: Xeroderma pigmentosum:
- [99020] complementation group C and malignant melanoma. Arch. Derm. 120:
- [99021] 175–179, 1984.
- [99022] 17242.
- [99023] Masutani, C.; Sugasawa, K.; Yanagisawa, J.; Sonoyama, T.; Ui,
- [99024] M.; Enomoto, T.; Takio, K.; Tanaka, K.; van der Spek, P.; Bootsma,
- [99025] D.; Hoeijmakers, J. H. J.; Hanaoka, F.: Purification and cloning
- [99026] of a nucleotide excision repair complex involving the xeroderma pigmentosum
- [99027] group C protein and a human homologue of yeast RAD23. EMBO J. 13:
- [99028] 1831–1843, 1994.
- [99029] 17243.
- [99030] Peterson, C.; Legerski, R.: High-frequency transformation

of

[99031] human repair-deficient cell lines by an Epstein-Barr virus-based cDNA

[99032] expression vector. *Gene* 107: 279-284, 1991.

[99033] 17244.

[99034] Sands, A. T.; Abuin, A.; Sanchez, A.; Conti, C. J.; Bradley, A.

[99035] : High susceptibility to ultraviolet-induced carcinogenesis in mice

[99036] lacking XPC. *Nature* 377: 162-165, 1995.

[99037] 17245.

[99038] Seguin, L. R.; Tarone, R. E.; Liao, K.; Robbins, J. H.: Ultraviolet

[99039] light-induced chromosomal aberrations in cultured cells from Cockayne

[99040] syndrome and complementation group C xeroderma pigmentosum patients:

[99041] lack of correlation with cancer susceptibility. *Am. J. Hum. Genet.* 42:

[99042] 468-475, 1988.

[99043] 17246.

- [99044] Slor, H.; Batko, S.; Khan, S. G.; Sobe, T.; Emmert, S.; Khadavi,
- [99045] A.; Frumkin, A.; Busch, D. B.; Albert, R. B.; Kraemer, K. H.: Clinical,
- [99046] cellular, and molecular features of an Israeli xeroderma pigmentosum
- [99047] family with a frameshift mutation in the XPC gene: sun protection
- [99048] prolongs life. J. Invest. Derm. 115: 974–980, 2000.
- [99049] 17247.
- [99050] Sugasawa, K.; Ng, J. M. Y.; Masutani, C.; Iwai, S.; van der Spek,
- [99051] P. J.; Eker, A. P. M.; Hanaoka, F.; Bootsma, D.; Hoeijmakers, J. H.
- [99052] J.: Xeroderma pigmentosum group C protein complex is the initiator
- [99053] of global genome nucleotide excision repair. Molec. Cell 2: 223–232,
- [99054] 1998.
- [99055] 17248.
- [99056] Suzuki, T.; Park, H.; Hollingsworth, N. M.; Sternglanz, R.; Lennarz,

- [99057] W. L.: PNG1, a yeast gene encoding a highly conserved peptide:N-glycanase. J.
- [99058] Cell Biol. 149: 1039–1052, 2000.
- [99059] 17249.
- [99060] Teitz, T.; Naiman, T.; Avissar, S. S.; Bar, S.; Okayama, H.; Canaani,
- [99061] D.: Complementation of the UV-sensitive phenotype of a xeroderma
- [99062] pigmentosum human cell line by transfection with a cDNA clone library. Proc.
- [99063] Nat. Acad. Sci. 84: 8801–8804, 1987.
- [99064] 17250.
- [99065] van der Spek, P. J.; Smit, E. M. E.; Beverloo, H. B.; Sugawara,
- [99066] K.; Masutani, C.; Hanaoka, F.; Hoeijmakers, J. H. J.; Hage-meijer,
- [99067] A.: Chromosomal localization of three repair genes: the xeroderma
- [99068] pigmentosum group C gene and two human homologs of yeast RAD23. Genomics 23:
- [99069] 651–658, 1994.

[99070] 17251.

[99071] van der Spek, P. J.; Visser, C. E.; Hanaoka, F.; Smit, B.;
Hagemeyer,

[99072] A.; Bootsma, D.; Hoeijmakers, J. H. J.: Cloning, comparative mapping,

[99073] and RNA expression of the mouse homologues of the *Saccharomyces cerevisiae*

[99074] nucleotide excision repair gene RAD23. *Genomics* 31:
20-27, 1996.

[99075] 17252.

[99076] Moser, H. W.: Personal Communication. Baltimore, Md.
6/29/1989.

[99077] 17253.

[99078] Moser, H. W.: Lorenzo's Oil. (Film Review). *Lancet* 341:
544,

[99079] 1993.

[99080] 17254.

[99081] Moser, H. W.: Personal Communication. Baltimore, Md.
1980.

[99082] 17255.

[99083] Moser, H. W.: Personal Communication. Baltimore, Md.

10/13/1987.

[99084] 17256.

[99085] Moser, H. W.; Kok, F.; Neumann, S.; Borel, J.; Bergin, A.;
Mostafa,

[99086] S. D.; Panoscha, R.; Davoli, C. T.; Shankroff, J.; Smith, K.
D.:

[99087] Adrenoleukodystrophy update: genetics and effect of
Lorenzo's oil

[99088] therapy in asymptomatic patients. Int. Pediat. 9: 196–204,
1994.

[99089] 17257.

[99090] Moser, H. W.; Moser, A. B.; Frayer, K. K.; Chen, W.; Schul-
man,

[99091] J. D.; O'Neill, B. P.; Kishimoto, Y.: Adrenoleukodystrophy:
increased

[99092] plasma content of saturated very long chain fatty acids.
Neurology 31:

[99093] 1241–1249, 1981.

[99094] 17258.

[99095] Moser, H. W.; Moser, A. B.; Kawamura, N.; Murphy, J.;
Suzuki,

[99096] K.; Schaumburg, H.; Kishimoto, Y.: Adrenoleukodystrophy:

elevated

[99097] C26 fatty acid in cultured skin fibroblasts. *Ann. Neurol.* 7:
542–549,

[99098] 1980.

[99099] 17259.

[99100] Moser, H. W.; Moser, A. B.; Naidu, S.; Bergin, A.: Clinical
aspects

[99101] of adrenoleukodystrophy and adrenomyeloneuropathy.
Dev. Neurosci. 13:

[99102] 254–261, 1991.

[99103] 17260.

[99104] Moser, H. W.; Moser, A. E.; Singh, I.; O'Neill, B. P.:
Adrenoleukodystrophy:

[99105] survey of 303 cases: biochemistry, diagnosis, and therapy.
Ann. Neurol. 16:

[99106] 628–641, 1984.

[99107] 17261.

[99108] Moser, H. W.; Smith, K. D.; Moser, A. B.: X-linked
adrenoleukodystrophy. In:

[99109] Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds): *The*

[99110] *Metabolic and Molecular Bases of Inherited Disease*. New

York: McGraw-Hill

[99111] 1995. Pp. 2325–2349.

[99112] 17262.

[99113] Moser, H. W.; Tutschka, P. J.; Brown, F. R., III; Moser, A. E.;

[99114] Yeager, A. M.; Singh, I.; Mark, S. A.; Kumar, A. A. J.; Mc-
Donnell,

[99115] J. M.; White, C. L., III; Maumenee, I. H.; Green, W. R.; Pow-
ers, J.

[99116] M.; Santos, G. W.: Bone marrow transplant in
adrenoleukodystrophy. *Neurology* 34:

[99117] 1410–1417, 1984.

[99118] 17263.

[99119] Mosser, J.; Douar, A.–M.; Sarde, C.–O.; Kioschis, P.; Feil,
R.;

[99120] Moser, H.; Poustka, A.–M.; Mandel, J.–L.; Aubourg, P.: Pu-
tative X-linked

[99121] adrenoleukodystrophy gene shares unexpected homology
with ABC transporters. *Nature* 361:

[99122] 726–730, 1993.

[99123] 17264.

[99124] Mosser, J.; Lutz, Y.; Stoeckel, M. E.; Sarde, C. O.; Kretz, C.;

- [99125] Douar, A. M.; Lopez, J.; Aubourg, P.; Mandel, J. L.: The gene responsible
- [99126] for adrenoleukodystrophy encodes a peroxisomal membrane protein. Hum.
- [99127] Molec. Genet. 3: 265–271, 1994.
- [99128] 17265.
- [99129] Naidu, S.; Bresnan, M. J.; Griffin, D.; O'Toole, S.; Moser, H.
- [99130] W.: Childhood adrenoleukodystrophy: failure of intensive immunosuppression
- [99131] to arrest neurologic progression. Arch. Neurol. 45: 846–848, 1988.
- [99132] 17266.
- [99133] Netik, A.; Forss-Petter, S.; Holzinger, A.; Molzer, B.; Unterrainer,
- [99134] G.; Berger, J.: Adrenoleukodystrophy-related protein can compensate
- [99135] functionally for adrenoleukodystrophy protein deficiency (X-ALD):
- [99136] implications for therapy. Hum. Molec. Genet. 8: 907–913, 1999.
- [99137] 17267.

- [99138] Noetzel, M. J.; Landau, W. M.; Moser, H. W.:
Adrenoleukodystrophy
- [99139] carrier state presenting as a chronic nonprogressive spinal
cord disorder. Arch.
- [99140] Neurol. 44: 566–567, 1987.
- [99141] 17268.
- [99142] O'Neill, B. P.; Marmion, L. C.; Feringa, E. R.: The
adrenoleukomyeloneuropathy
- [99143] complex: expression in four generations. Neurology 31:
151–156,
- [99144] 1981.
- [99145] 17269.
- [99146] O'Neill, B. P.; Moser, H. W.; Marmion, L. C.:
Adrenoleukodystrophy:
- [99147] elevated C26 fatty acid in cultured skin fibroblasts and
correlation
- [99148] with disease expression in three generations of a kindred.
Neurology 32:
- [99149] 540–542, 1982.
- [99150] 17270.
- [99151] O'Neill, B. P.; Moser, H. W.; Marmion, L. C.: The

adrenoleukomyeloneuropathy

[99152] (ALMN) complex: elevated C26 fatty acid in cultured skin fibroblasts

[99153] and correlation with disease expression in three generations of a

[99154] kindred. (Abstract) Neurology 30: 352 only, 1980.

[99155] 17271.

[99156] O'Neill, B. P.; Moser, H. W.; Saxena, K. M.: Familial X-linked

[99157] Addison disease as an expression of adrenoleukodystrophy (ALD): elevated

[99158] C26 fatty acid in cultured skin fibroblasts. Neurology 32: 543-547,

[99159] 1982.

[99160] 17272.

[99161] O'Neill, B. P.; Swanson, J. W.; Brown, F. R., III; Griffin, J.

[99162] W.; Moser, H. W.: Familial spastic paraparesis: an adrenoleukodystrophy

[99163] phenotype? Neurology 35: 1233-1235, 1985.

[99164] 17273.

[99165] Peckham, R. S.; Marshall, M. C., Jr.; Rosman, P. M.; Farag,

A.;

- [99166] Kabadi, U.; Wallace, E. Z.: A variant of adrenomyeloneu-
ropathy with
- [99167] hypothalamic-pituitary dysfunction and neurologic remis-
sion after
- [99168] glucocorticoid replacement therapy. Am. J. Med. 72:
173-176, 1982.
- [99169] 17274.
- [99170] Penman, R. W. B.: Addison's disease in association with
spastic
- [99171] paraplegia. Brit. Med. J. 1: 402, 1960.
- [99172] 17275.
- [99173] Poulos, A.; Gibson, R.; Sharp, P.; Beckman, K.; Grattan-
Smith,
- [99174] P.: Very long chain fatty acids in X-linked
adrenoleukodystrophy
- [99175] brain after treatment with Lorenzo's oil. Ann. Neurol. 36:
741-746,
- [99176] 1994.
- [99177]
- [99178] 100. Powers, J. M.; Schaumburg, H. H.: The testis in

adreno-leukodystrophy. Am.

[99179] J. Path. 102: 90-98, 1981.

[99180]

[99181] 101. Powers, J. M.; Schaumburg, H. H.: A fatal cause of sexual inadequacy

[99182] in men: adreno-leukodystrophy. J. Urol. 124: 583-585, 1980.

[99183]

[99184] 102. Powers, J. M.; Schaumburg, H. H.: Adreno-leukodystrophy (sex-linked

[99185] Schilder's disease): a pathogenic hypothesis based on ultrastructural

[99186] lesions in adrenal cortex, peripheral nerve and testis. Am. J. Path. 76:

[99187] 481-500, 1974.

[99188]

[99189] 103. Powers, J. M.; Schaumburg, H. H.; Gaffney, C. L.: Klu-ver-Bucy

[99190] syndrome caused by adreno-leukodystrophy. Neurology 30: 1131-1132,

[99191] 1980.

[99192]

[99193] 104. Probst, A.; Ulrich, J.; Heitz, P. U.; Herschkowitz, N.:
Adrenomyeloneuropathy:

[99194] a protracted, pseudosystematic variant of
adrenoleukodystrophy. Acta

[99195] Neuropath. 49: 105–115, 1980.

[99196]

[99197] 105. Restuccia, D.; Di Lazzaro, V.; Valeriani, M.; Oliviero,
A.; Le

[99198] Pera, D.; Barba, C.; Cappa, M.; Bertini, E.; Di Capua, M.;
Tonali,

[99199] P.: Neurophysiologic follow-up of long-term dietary treat-
ment in

[99200] adult-onset adrenoleukodystrophy. Neurology 52:
810–816, 1999.

[99201]

[99202] 106. Ropers, H.–H.; Burmeister, P.; von Petrykowski, W.;
Schindera,

[99203] F.: Leukodystrophy, skin hyperpigmentation, and adrenal
atrophy:

[99204] Siemerling–Creutzfeldt disease: transmission through
several generations

[99205] in two families. Am. J. Hum. Genet. 27: 547–553, 1975.

[99206]

[99207] 107. Ropers, H.–H.; Zimmermann, J.; Wienker, T.:

Adrenoleukodystrophy

[99208] (Siemerling–Creutzfeldt disease): heterozygote with two
clonal fibroblast

[99209] populations. Clin. Genet. 11: 114–118, 1977.

[99210]

[99211] 108. Rosen, N. L.; Lechtenberg, R.; Wisniewski, K.;

Pullarkat, R.;

[99212] Bennett, H. S.: Adrenoleukomyeloneuropathy with onset in
early childhood. Ann.

[99213] Neurol. 17: 311–312, 1985.

[99214]

[99215] 109. Ruiz, M.; Coll, M. J.; Pampols, T.; Giros, M.: X-linked
adrenoleukodystrophy:

[99216] phenotype distribution and expression of ALDP in Spanish
kindreds. Am.

[99217] J. Med. Genet. 76: 424–427, 1998.

[99218]

- [99219] 110. Sack, G. H., Jr.; Alpern, M.; Webster, T.; Feil, R. P.;
Morrell,
[99220] J. C.; Chen, G.; Chen, W.; Caskey, C. T.; Moser, H. W.:
Chromosomal
[99221] rearrangement segregating with adrenoleukodystrophy: a
molecular analysis. Proc.
[99222] Nat. Acad. Sci. 90: 9489–9493, 1993.
[99223]
[99224] 111. Sack, G. H., Jr.; Morrell, J. C.: Adrenoleukodystrophy:
overlapping
[99225] deletions point to a gene location in Xq28. Biochem. Bio-
phys. Res.
[99226] Commun. 191: 955–960, 1993.
[99227]
[99228] 112. Sack, G. H., Jr.; Raven, M. B.; Moser, H. W.: Color vi-
sion defects
[99229] in adrenomyeloneuropathy. Am. J. Hum. Genet. 44:
794–798, 1989.
[99230]
[99231] 113. Sadeghi-Nejad, A.; Senior, B.: Adrenomyeloneuropa-
thy presenting

[99232] as Addison's disease in childhood. New Eng. J. Med. 322:
13–16,

[99233] 1990.

[99234]

[99235] 114. Sarde, C.-O.; Mosser, J.; Kioschis, P.; Kretz, C.; Vi-
caire, S.;

[99236] Aubourg, P.; Poustka, A.; Mandel, J.-L.: Genomic organi-
zation of

[99237] the adrenoleukodystrophy gene. Genomics 22: 13–20,
1994.

[99238]

[99239] 115. Sarde, C.-O.; Thomas, J.; Sadoulet, H.; Garnier, J.-M.;
Mandel,

[99240] J.-L.: cDNA sequence of Aldgh, the mouse homolog of the
X-linked

[99241] adrenoleukodystrophy gene. Mammalian Genome 5:
810–813, 1994.

[99242]

[99243] 116. Schaumburg, H. H.; Powers, J. M.; Raine, C. S.;
Spencer, P. S.;

[99244] Griffin, J. W.; Prineas, J. W.; Boehme, D. M.: Adreno-
myeloneuropathy:

- [99245] a probable variant of adrenoleukodystrophy. II. General pathologic,
- [99246] neuropathologic, and biochemical aspects. Neurology 27: 1114-1119,
- [99247] 1977.
- [99248]
- [99249] 117. Schaumburg, H. H.; Powers, J. M.; Raine, C. S.; Suzuki, K.; Richardson,
- [99250] E. P., Jr.: Adrenoleukodystrophy: a clinical and pathological study
- [99251] of 17 cases. Arch. Neurol. 32: 577-591, 1975.
- [99252]
- [99253] 118. Schaumburg, H. H.; Richardson, E. P., Jr.; Johnson, P. C.; Cohen,
- [99254] R. B.; Powers, J. M.; Raine, C. S.: Schilder's disease: sex-linked
- [99255] recessive transmission with specific adrenal changes. Arch. Neurol. 27:
- [99256] 458-460, 1972.
- [99257]
- [99258] 119. Schilder, P.: Zur Frage der Encephalitis Periaxialis

Diffusa

[99259] (Sogenannte Diffuse Sklerose). Z. ges Neurol. Psychiat. 15: 359–376,

[99260] 1913.

[99261]

[99262] 120. Seneca, S.; Lissens, W.: DNA diagnosis of X-linked adrenoleukodystrophy. J.

[99263] Inherit. Metab. Dis. 18 (suppl. 1): 34–44, 1995.

[99264]

[99265] 121. Shani, N.; Watkins, P. A.; Valle, D.: PXA1, a possible *Saccharomyces*

[99266] *cerevisiae* ortholog of the human adrenoleukodystrophy gene. Proc.

[99267] Nat. Acad. Sci. 92: 6012–6016, 1995.

[99268]

[99269] 122. Siemerling, E.; Creutzfeldt, H. G.: Bronzekrankheit und sklerosierende

[99270] Encephalomyelitis. Arch. Psychiat. Nervkrankh. 68: 217–244, 1923.

[99271]

[99272] 123. Singh, I.; Lazo, O.; Contreras, M.; Hashmi, M.; Stan-

ley, W. S.

[99273] : Peroxisomal lignoceroyl-CoA ligase deficiency in X-linked adrenoleukodystrophy.

[99274] (Abstract) Am. J. Hum. Genet. 43: A16, 1988.

[99275]

[99276] 124. Sobue, G.; Ueno-Natsukari, I.; Okamoto, H.; Connell, T. A.; Aizawa,

[99277] I.; Mizoguchi, K.; Honma, M.; Ishikawa, G.; Mitsuma, T.; Natsukari,

[99278] N.: Phenotypic heterogeneity of an adult form of adrenoleukodystrophy

[99279] in monozygotic twins. Ann. Neurol. 36: 912-915, 1994.

[99280]

[99281] 125. Spira, T. J.; Adam, A.; Goodman, R. M.; Berger, A.: Recombination

[99282] between cerebral sclerosis--Addison's disease and the Xg blood-groups.

[99283] (Letter) Lancet II: 820-821, 1971.

[99284]

[99285] 126. Takano, H.; Koike, R.; Onodera, O.; Sasaki, R.; Tsuji, S.: Mutational

- [99286] analysis and genotype-phenotype correlation of 29 unrelated Japanese
- [99287] patients with X-linked adrenoleukodystrophy. Arch. Neurol. 56: 295-300,
- [99288] 1999.
- [99289]
- [99290] 127. Turkington, R. W.; Stempfel, R. S., Jr.: Adrenocortical atrophy
- [99291] and diffuse cerebral sclerosis (Addison-Schilder's disease). J. Pediat. 69:
- [99292] 406-412, 1966.
- [99293]
- [99294] 128. Ulrich, J.; Herschkowitz, N.; Heitz, P.; Sigrist, T. H.; Baerlocher,
- [99295] P.: Adrenoleukodystrophy: preliminary report of a congenital case;
- [99296] light- and electronmicroscopical, immunohistochemical and biochemical
- [99297] findings. Acta Neuropath. 43: 77-83, 1978.
- [99298]
- [99299] 129. Unterrainer, G.; Molzer, B.; Forss-Petter, S.; Berger,

J.: Co-expression

[99300] of mutated and normal adrenoleukodystrophy protein reduces protein

[99301] function: implications for gene therapy of X-linked adrenoleukodystrophy. Hum.

[99302] Molec. Genet. 9: 2609–2616, 2000.

[99303]

[99304] 130. Uyama, E.; Iwagoe, H.; Maeda, J.; Nakamura, M.; Terasaki, T.;

[99305] Ando, M.: Presenile-onset cerebral adrenoleukodystrophy presenting

[99306] as Balint's syndrome and dementia. Neurology 43: 1249–1251, 1993.

[99307]

[99308] 131. Valle, D.; Gartner, J.: Penetrating the peroxisome. Nature 361:

[99309] 682–683, 1993.

[99310]

[99311] 132. van den Bosch, H.; Schutgens, R. B. H.; Wanders, R. J. A.; Tager,

[99312] J. M.: Biochemistry of peroxisomes. Annu. Rev. Biochem.

61: 157-197,

[99313] 1992.

[99314]

[99315] 133. van Geel, B. M.; Assies, J.; Weverling, G. J.; Barth, P.
G.:

[99316] Predominance of the adrenomyeloneuropathy phenotype
of X-linked adrenoleukodystrophy

[99317] in the Netherlands: a survey of 30 kindreds. Neurology 44:
2343-2346,

[99318] 1994.

[99319]

[99320] 134. van Oost, B. A.; van Zandvoort, P.; Hoogeboom, A.;
Bakkeren,

[99321] J.; Hamel, B.; Brunner, H.; Knoers, N.; Ropers, H. H.: Tight
linkage

[99322] between adrenoleukodystrophy and DXS 52. (Abstract)
Cytogenet. Cell

[99323] Genet. 46: 708 only, 1987.

[99324]

[99325] 135. van Oost, B. A.; van Zandvoort, P. M.; Tunte, W.;
Brunner, H.

- [99326] G.; Hoogeboom, A. J. M.; Maaswinkel-Mooy, P. D.;
Bakkeren, J.; Hamel,
- [99327] B.; Ropers, H. H.: Linkage analysis in X-linked
adrenoleukodystrophy
- [99328] and application in post- and prenatal diagnosis. Hum.
Genet. 86:
- [99329] 404-407, 1991.
- [99330]
- [99331] 136. Walsh, P. J.: Adrenoleukodystrophy: report of two
cases with
- [99332] relapsing and remitting courses. Arch. Neurol. 37:
448-450, 1980.
- [99333]
- [99334] 137. Wanders, R. J. A.; van Roermund, C. W. T.; van Wij-
land, M. J.
- [99335] A.; Nijenhuis, A. A.; Tromp, A.; Schutgens, R. B. H.;
Brouwer-Kelder,
- [99336] E. M.; Schram, A. W.; Tager, J. M.; van den Bosch, H.;
Schalkwijk,
- [99337] C.: X-linked adrenoleukodystrophy: defective peroxisomal
oxidation
- [99338] of very long chain fatty acids but not of very long chain

fatty acyl-CoA

[99339] esters. Clin. Chim. Acta 165: 321-329, 1987.

[99340]

[99341] 138. Wanders, R. J. A.; van Roermund, C. W. T.; van Wijland, M. J.

[99342] A.; Schutgens, R. B. H.; Heikoop, J.; van den Bosch, H.; Schram, A.

[99343] W.; Tager, J. M.: Peroxisomal fatty acid beta-oxidation in relation

[99344] to the accumulation of very long chain fatty acids in cultured skin

[99345] fibroblasts from patients with Zellweger syndrome and other peroxisomal

[99346] disorders. J. Clin. Invest. 80: 1778-1783, 1987.

[99347]

[99348] 139. Wanders, R. J. A.; van Roermund, C. W. T.; van Wijland, M. J.

[99349] A.; Schutgens, R. B. H.; van den Bosch, H.; Schram, A. W.; Tager,

[99350] J. M.: Direct demonstration that the deficient oxidation of very

[99351] long chain fatty acids in X-linked adrenoleukodystrophy is

due to

[99352] an impaired ability of peroxisomes to activate very long chain fatty

[99353] acids. *Biochem. Biophys. Res. Commun.* 153: 618–624, 1988.

[99354]

[99355] 140. Watkiss, E.; Webb, T.; Bunday, S.: Is skewed X inactivation

[99356] responsible for symptoms in female carriers for adrenoleucodystrophy? *J.*

[99357] *Med. Genet.* 30: 651–654, 1993.

[99358]

[99359] 141. Wilichowski, E.; Ohlenbusch, A.; Korenke, G. C.; et al.: Identical

[99360] mitochondrial DNA in monozygotic twins with discordant adrenoleukodystrophy

[99361] phenotype. *Ann. Neurol.* 43: 835–836, 1998.

[99362]

[99363] 142. Willems, P. J.; Vits, L.; Wanders, R. J. A.; Coucke, P. J.; Van

[99364] der Auwera, B. J.; Van Elsen, A. F.; Raeymaekers, P.; Van

Broeckhoven,

[99365] C.; Schutgens, R. B. H.; Dacremont, G.; Leroy, J. G.; Martin, J.-J.;

[99366] Dumon, J. E.: Linkage of DNA markers at Xq28 to adrenoleukodystrophy

[99367] and adrenomyeloneuropathy present within the same family. Arch. Neurol. 47:

[99368] 665-669, 1990.

[99369]

[99370] 143. Wray, S. H.; Cogan, D. G.; Kuwabara, T.; Schaumburg, H. H.; Powers,

[99371] J. M.: Adrenoleukodystrophy with disease of the eye and optic nerve. Am.

[99372] J. Ophthalm. 82: 480-485, 1976.

[99373] 17276.

[99374] Dinulos, M. B.; Bassi, M. T.; Rugarli, E. I.; Chapman, V.; Ballabio,

[99375] A.; Disteche, C. M.: A new region of conservation is defined between

[99376] human and mouse X chromosomes. Genomics 35: 244-247, 1996.

[99377] 17277.

- [99378] Schiaffino, M. V.; Bassi, M. T.; Rugarli, E. I.; Renieri, A.; Galli,
L.; Ballabio, A.: Cloning of a human homologue of the
Xenopus laevis
APX gene from the ocular albinism type 1 critical region.
Hum. Molec. Genet. 4: 373–382, 1995.
17278.
- [99382] 17278.
- [99383] Bachner, D.; Sedlacek, Z.; Korn, B.; Hameister, H.; Poustka,
A.
: Expression patterns of two human genes coding for dif-
ferent rab
GDP-dissociation inhibitors (GDIs), extremely conserved
proteins involved
in cellular transport. Hum. Molec. Genet. 4: 701–708,
1995.
17279.
- [99387] 17279.
- [99388] Black, G. C. M.; Perveen, R.; Bonshek, R.; Cahill, M.; Clay-
ton-Smith,
J.; Lloyd, I. C.; McLeod, D.: Coats' disease of the retina
(unilateral
retinal telangiectasis) caused by somatic mutation in the

NDP gene:

- [99391] a role for norrin in retinal angiogenesis. Hum. Molec. Genet. 8: 2031–2035, 1999.
- [99392] 17280.
- [99393] Walpole, S. M.; Hiriyana, K. T.; Nicolaou, A.; Bingham, E. L.; Durham, J.; Vaudin, M.; Ross, M. T.; Yates, J. R. W.; Sieving, P.
- [99394] A.; Trump, D.: Identification and characterization of the human homologue (RAI2) of a mouse retinoic acid-induced gene in Xp22. Genomics 55: 275–283, 1999.
- [99395] 17281.
- [99396] Mao, M.; Fu, G.; Wu, J.-S.; Zhang, Q.-H.; Zhou, J.; Kan, L.-X.; Huang, Q.-H.; He, K.-L.; Gu, B.-W.; Han, Z.-G.; Shen, Y.; Gu, J.; Yu, Y.-P.; Xu, S.-H.; Wang, Y.-X.; Chen, S.-J.; Chen, Z.: Identification of genes expressed in human CD34+ hematopoietic
- [99402]
- [99403]

stem/progenitor cells

[99404] by expressed sequence tags and efficient full-length
cDNA cloning. Proc.

[99405] Nat. Acad. Sci. 95: 8175–8180, 1998.

[99406] 17282.

[99407] Pearce, W. G.; Sanger, R.; Race, R. R.: Ocular albinism and
Xg. Lancet I:

[99408] 1282–1283, 1968.

[99409] 17283.

[99410] Rosenberg, T.; Schwartz, M.: X-linked ocular albinism:
prevalence

[99411] and mutations—a national study. Europ. J. Hum. Genet. 6:
570–577,

[99412] 1998.

[99413] 17284.

[99414] Schiaffino, M. V.; Bassi, M. T.; Galli, L.; Renieri, A.; Bruttini,

[99415] M.; De Nigris, F.; Bergen, A. A. B.; Charles, S. J.; Yates, J. R.

[99416] W., Meindl, A.; Lewis, R. A.; King, R. A.; Ballabio, A.: Anal-
ysis

[99417] of the OA1 gene reveals mutations in only one-third of

patients with

[99418] X-linked ocular albinism. Hum. Molec. Genet. 4:
2319-2325, 1995.

[99419] 17285.

[99420] Schiaffino, M. V.; d'Addio, M.; Alloni, A.; Baschiroto, C.;
Valetti,

[99421] C.; Cortese, K.; Puri, C.; Bassi, M. T.; Colla, C.; De Luca,
M.; Tacchetti,

[99422] C.; Ballabio, A.: Ocular albinism: evidence for a defect in
an intracellular

[99423] signal transduction system. Nature Genet. 23: 108-112,
1999.

[99424] 17286.

[99425] Schnur, R. E.; Gao, M.; Wick, P. A.; Keller, M.; Benke, P. J.;

[99426] Edwards, M. J.; Grix, A. W.; Hockey, A.; Jung, J. H.; Kidd, K.
K.;

[99427] Kistenmacher, M.; Levin, A. V.; and 11 others: OA1 muta-
tions and

[99428] deletions in X-linked ocular albinism. Am. J. Hum. Genet.
62: 800-809,

[99429] 1998.

[99430] 17287.

- [99431] Schnur, R. E.; Trask, B. J.; van den Engh, G.; Punnett, H. H.;
- [99432] Kistenmacher, M.; Tomeo, M. A.; Naidu, R. E.; Nussbaum, R. L.: An
- [99433] Xp22 microdeletion associated with ocular albinism and ichthyosis:
- [99434] approximation of breakpoints and estimation of deletion size by using
- [99435] cloned DNA probes and flow cytometry. Am. J. Hum. Genet. 45: 706–720,
- [99436] 1989.
- [99437] 17288.
- [99438] Schnur, R. E.; Wick, P. A.; Bailey, C.; Rebbeck, T.; Weleber,
- [99439] R. G.; Wagstaff, J.; Grix, A. W.; Pagon, R. A.; Hockey, A.; Edwards,
- [99440] M. J.: Phenotypic variability in X-linked ocular albinism: relationship
- [99441] to linkage genotypes. Am. J. Hum. Genet. 55: 484–496, 1994.
- [99442] 17289.
- [99443] Vogt, A.: Die Iris: Albinismus solum bulbi. Atlas Spalt-Lampen-Mikroskopie

[99444] (pub.) (3rd ed.): 1942.

[99445] 17290.

[99446] Waardenburg, P. J.; Van den Bosch, J.: X-chromosomal ocular albinism

[99447] in Dutch family. Ann. Hum. Genet. 21: 101–122, 1956.

[99448] 17291.

[99449] Zhang, Y.; McMahon, R.; Charles, S. J.; Green, J. S.; Moore, A.

[99450] T.; Barton, D. E.; Yates, J. R. W.: Genetic mapping of the Kallmann

[99451] syndrome and X linked ocular albinism gene loci. J. Med. Genet. 30:

[99452] 923–925, 1993.

[99453] 17292.

[99454] Barth, P. G.; Van Wijngaarden, G. K.; Bethlem, J.: X-linked myotubular

[99455] myopathy with fatal neonatal asphyxia. Neurology 25: 531–536, 1975.

[99456] 17293.

[99457] Blondeau, F.; Laporte, J.; Bodin, S.; Superti-Furga, G.; Payastre,

- [99458] B.; Mandel, J.-L.: Myotubularin, a phosphatase deficient in myotubular
- [99459] myopathy, acts on phosphatidylinositol 3-kinase and phosphatidylinositol
- [99460] 3-phosphate pathway. Hum. Molec. Genet. 9: 2223-2229, 2000.
- [99461] 17294.
- [99462] Braga, S. E.; Gerber, A.; Meier, C.; Weiersmuller, A.; Zimmermann,
- [99463] A.; Herrmann, U.; Liechti, S.; Moser, H.: Severe neonatal asphyxia
- [99464] due to X-linked centronuclear myopathy. Europ. J. Pediat. 150: 132-135,
- [99465] 1990.
- [99466] 17295.
- [99467] Breningstall, G. N.; Grover, W. D.; Marks, H. G.: Maternal muscle
- [99468] biopsy in X-linked recessive centronuclear (myotubular) myopathy. Am.
- [99469] J. Med. Genet. 39: 13-18, 1991.
- [99470] 17296.

- [99471] Verga, V.; Hall, B. K.; Wang, S.; Johnson, S.; Higgins, J. V.;
- [99472] Glover, T. W.: Localization of the translocation breakpoint
in a
- [99473] female with Menkes syndrome to Xq13.2-q13.3 proximal
to PGK-1. Am.
- [99474] J. Hum. Genet. 48: 1133-1138, 1991.
- [99475] 17297.
- [99476] Seltzer, W. K.; Firminger, H.; Klein, J.; Pike, A.; Fennessey,
- [99477] P.; McCabe, E. R. B.: Adrenal dysfunction in glycerol kinase
deficiency. Biochem.
- [99478] Med. 33: 189-199, 1985.
- [99479] 17298.
- [99480] Cianfriglia, M.; Miggiano, V. C.; Meo, T.; Muller, H. J.;
- Muller,
- [99481] E.; Battistuzzi, G.: Evidence for syntenic between the rabbit
gene
- [99482] loci coding for HPRT, PGK and G6PD in mouse-rabbit so-
matic cell hybrids.
- [99483] (Abstract) Cytogenet. Cell Genet. 25: 142, 1979.
- [99484] 17299.

- [99485] Goss, S. J.; Harris, H.: Gene transfer by means of cell fusion.
- [99486] I. Statistical mapping of the human X-chromosome by analysis of radiation-induced
- [99487] gene segregation. J. Cell Sci. 25: 17–37, 1977.
- [99488] 17300.
- [99489] Gill, P.; Ivanov, P. L.; Kimpton, C.; Piercy, R.; Benson, N.;
- [99490] Tully, G.; Evett, I.; Hagelberg, E.; Sullivan, K.: Identification
- [99491] of the remains of the Romanov family by DNA analysis. Nature Genet. 6:
- [99492] 130–135, 1994.
- [99493] 17301.
- [99494] Gitschier, J.: Maternal duplication associated with gene deletion
- [99495] in sporadic hemophilia. Am. J. Hum. Genet. 43: 274–279, 1988.
- [99496] 17302.
- [99497] Gitschier, J.; Drayna, D.; Tuddenham, E. G. D.; White, R. L.;
- [99498] Lawn, R. M.: Genetic mapping and diagnosis of haemophilia A achieved

- [99499] through a BclI polymorphism in the factor VIII gene. Nature 314:
- [99500] 738–740, 1985.
- [99501] 17303.
- [99502] Gitschier, J.; Wood, W. I.; Shuman, M. A.; Lawn, R. M.: Identification
- [99503] of a missense mutation in the factor VIII gene of a mild hemophiliac. Science 232:
- [99504] 1415–1416, 1986.
- [99505] 17304.
- [99506] Gitschier, J.; Wood, W. I.; Tuddenham, E. G. D.; Shuman, M. A.;
- [99507] Goralka, T. M.; Chen, E. Y.; Lawn, R. M.: Detection and sequence
- [99508] of mutations in the factor VIII gene of haemophiliacs. Nature 315:
- [99509] 427–430, 1985.
- [99510] 17305.
- [99511] Graham, J. B.; Green, P. P.; McGraw, R. A.; Davis, L. M.: Application
- [99512] of molecular genetics to prenatal diagnosis and carrier

detection

- [99513] in the hemophilias: some limitations. Blood 66: 759-764, 1985.
- [99514] 17306.
- [99515] Graham, J. B.; McLendon, W. W.; Brinkhous, K. M.: Mild hemophilia:
- [99516] an allelic form of the disease. Am. J. Med. Sci. 225: 46-53, 1953.
- [99517] 17307.
- [99518] Green, P. M.; Montandon, A. J.; Bentley, D. R.; Giannelli, F.
- [99519] : Genetics and molecular biology of haemophilias A and B. Blood Coagul.
- [99520] Fibrinolysis 2: 539-565, 1991.
- [99521] 17308.
- [99522] Grover, H.; Phillips, M. A.; Lillicrap, D. P.; Giles, A. R.; Garvey,
- [99523] M. B.; Teitel, J.; Rivard, G.; Blanchette, V.; White, B. N.; Holden,
- [99524] J. J. A.: Carrier deletion of haemophilia A using DNA markers in
- [99525] families with an isolated affected male. Clin. Genet. 32:

- 10-19,
[99526] 1987.
- [99527] 17309.
- [99528] Grozdea, J.; Colombies, P.; Bierme, R.; Ducos, J.:
Myeloperoxidases
[99529] and genetics of haemophilia A. (Letter) Lancet II: 220 only,
1969.
- [99530] 17310.
- [99531] Haldane, J. B. S.; Smith, C. A. B.: A new estimate of the
linkage
[99532] between the genes for colour-blindness and haemophilia
in man. Ann.
[99533] Eugen. 14: 10-31, 1947.
- [99534] 17311.
- [99535] Harper, K.; Winter, R. M.; Pembrey, M. E.; Hartley, D.;
Davies,
[99536] K. E.; Tuddenham, E. G. D.: A clinically useful DNA probe
closely
[99537] linked to haemophilia A. Lancet II: 6-8, 1984.
- [99538] 17312.
- [99539] Hemker, H. C.; Muller, A. D.; Hermens, W. T.; Zwaal, R. F.

A.

- [99540] : Oral treatment of hemophilia A by gastrointestinal absorption of
- [99541] factor VIII entrapped in liposomes. Lancet I: 70–71, 1980.
- [99542] 17313.
- [99543] Hermann, J.: Der Einfluss des Zeugungsalters auf die Mutationen
- [99544] zu Haemophilie A. Humangenetik 3: 1–16, 1966.
- [99545] 17314.
- [99546] Higuchi, M.; Antonarakis, S. E.; Kasch, L.; Oldenburg, J.; Economou-Petersen,
- [99547] E.; Olek, K.; Arai, M.; Inaba, H.; Kazazian, H. H., Jr.: Molecular
- [99548] characterization of mild-to-moderate hemophilia A: detection of the
- [99549] mutation in 25 of 29 patients by denaturing gradient gel electrophoresis. Proc.
- [99550] Nat. Acad. Sci. 88: 8307–8311, 1991.
- [99551] 17315.
- [99552] Higuchi, M.; Kazazian, H. H., Jr.; Kasch, L.; Warren, T. C.; McGinniss,

- [99553] M. J.; Phillips, J. A., III; Kasper, C., III; Janco, R.; Antonarakis,
- [99554] S. E.: Molecular characterization of severe hemophilia A suggests
- [99555] that about half the mutations are not within the coding regions and
- [99556] splice junctions of the factor VIII gene. Proc. Nat. Acad. Sci. 88:
- [99557] 7405–7409, 1991.
- [99558] 17316.
- [99559] Higuchi, M.; Kochhan, L.; Olek, K.: A somatic mosaic for haemophilia
- [99560] A detected at the DNA level. Molec. Biol. Med. 5: 23–27, 1988.
- [99561] 17317.
- [99562] Higuchi, M.; Kochhan, L.; Schwaab, R.; Egli, H.; Brackmann, H.
- [99563] H.; Horst, J.; Olek, K.: Molecular defects in hemophilia A: identification
- [99564] and characterization of mutations in the factor VIII gene and family
- [99565] analysis. Blood 74: 1045–1051, 1989.

[99566] 17318.

[99567] Higuchi, M.; Wong, C.; Kochhan, L.; Olek, K.; Aronis, S.;
Kasper,

[99568] C. K.; Kazazian, H. H., Jr.; Antonarakis, S. E.: Characteriza-
tion

[99569] of mutations in the factor VIII gene by direct sequencing
of amplified

[99570] genomic DNA. Genomics 6: 65-71, 1990.

[99571] 17319.

[99572] Howard, P. L.; Hoag, J. B.; Bovill, E. G.; Heintz, N. H.: Spon-
taneous

[99573] mutation in the male gamete as a cause of hemophilia A:
clarification

[99574] of a case using DNA probes. Am. J. Hemat. 28: 167-169,
1988.

[99575] 17320.

[99576] Hoyer, L. W.; Breckenridge, R. T.: Two forms of
haemophilia?

[99577] (Letter) Lancet II: 457 only, 1968.

[99578] 17321.

[99579] Inaba, H.; Fujimaki, M.; Kazazian, H. H., Jr.; Antonarakis, S.

- [99580] E.: Mild hemophilia A resulting from arg-to-leu substitution in exon
- [99581] 26 of the factor VIII gene. Hum. Genet. 81: 335-338, 1989.
- [99582] 17322.
- [99583] Ivanov, P. L.; Wadhams, M. J.; Roby, R. K.; Holland, M. M.; Weedn,
- [99584] V. W.; Parsons, T. J.: Mitochondrial DNA sequence heteroplasmy in
- [99585] the Grand Duke of Russia Georgij Romanov establishes the authenticity
- [99586] of the remains of Tsar Nicholas II. Nature Genet. 12: 417-420, 1996.
- [99587] 17323.
- [99588] Jaffe, E. A.; Nachman, R. L.: Subunit structure of factor VIII
- [99589] antigen synthesized by cultured human endothelial cells. J. Clin.
- [99590] Invest. 56: 698-702, 1975.
- [99591] 17324.
- [99592] Janco, R. L.; Phillips, J. A., III; Orlando, P. J.; Woodard, M.
- [99593] J.; Wion, K. L.; Lawn, R. M.: Detection of hemophilia A car-

riers

[99594] using intragenic factor VIII:C DNA polymorphisms. Blood
69: 1539–1541,

[99595] 1987.

[99596] 17325.

[99597] Johnson, D. J.; Pemberton, S.; Acquila, M.; Mori, P. G.;
Tuddenham,

[99598] E. G.; O'Brien, D. P.: Factor VIII S373L: mutation at P1' site
confers

[99599] thrombin cleavage resistance, causing mild haemophilia
A. Thromb.

[99600] Haemost. 71: 428–433, 1994.

[99601] 17326.

[99602] Jonsdottir, S.; Diamond, C.; Levinson, B.; Magnusson, S.;
Jensson,

[99603] O.; Gitschier, J.: Missense mutations causing mild
hemophilia A in

[99604] Iceland detected by denaturing gradient gel electrophore-
sis. Hum.

[99605] Mutat. 1: 506–508, 1992.

[99606] 17327.

- [99607] Holterhus, P. M.; Sinnecker, G. H. G.; Wollmann, H. A.;
Struve,
[99608] D.; Homburg, N.; Kruse, K.; Hiort, O.: Expression of two
functionally
[99609] different androgen receptors in a patient with androgen
insensitivity. Europ.
[99610] J. Pediat. 158: 702–706, 1999.
[99611] 17328.
[99612] Shimmin, L. C.; Chang, B.–H.; Li, W.–H.: Male–driven evo–
lution
[99613] of DNA sequences. Nature 362: 745–747, 1993.
[99614] 17329.
[99615] Warburg, M.: Personal Communication. Copenhagen, Den–
mark 1966.
[99616] 17330.
[99617] Imasaki, K.; Okabe, T.; Murakami, H.; Tanaka, Y.; Haji, M.;
Takayanagi,
[99618] R.; Nawata, H.: Androgen insensitivity syndrome due to
new mutations
[99619] in the DNA–binding domain of the androgen receptor.
Molec. Cell.

- [99620] *Endocr.* 120: 15–24, 1996.
- [99621] 17331.
- [99622] Jakubiczka, S.; Nedel, S.; Werder, E. A.; Schleiermacher, E.;
- [99623] Theile, U.; Wolff, G.; Wieacker, P.: Mutations of the andro-
gen receptor
- [99624] gene in patients with complete androgen insensitivity.
Hum. Mutat. 9:
- [99625] 57–61, 1997.
- [99626] 17332.
- [99627] Kang, H.-Y.; Yeh, S.; Fujimoto, N.; Chang, C.: Cloning and
characterization
- [99628] of human prostate coactivator ARA54, a novel protein that
associates
- [99629] with the androgen receptor. *J. Biol. Chem.* 274:
8570–8576, 1999.
- [99630] 17333.
- [99631] Kazemi-Esfarjani, P.; Beitel, L. K.; Trifiro, M.; Kaufman, M.;
- [99632] Rennie, P.; Sheppard, P.; Matusik, R.; Pinsky, L.: Substitu-
tion of
- [99633] valine-865 by methionine or leucine in the human andro-
gen receptor

- [99634] causes complete or partial androgen insensitivity, respectively with
- [99635] distinct androgen receptor phenotypes. *Molec. Endocr.* 7: 37–46,
- [99636] 1993.
- [99637] 17334.
- [99638] Kittles, R. A.; Young, D.; Weinrich, S.; Hudson, J.; Argypoulos,
- [99639] G.; Ukoli, F.; Adams–Campbell, L.; Dunston, G. M.: Extent of linkage
- [99640] disequilibrium between the androgen receptor gene CAG and GGC repeats
- [99641] in human populations: implications for prostate cancer risk. *Hum.*
- [99642] *Genet.* 109: 253–261, 2001.
- [99643] 17335.
- [99644] Klocker, H.; Kaspar, F.; Eberle, J.; Uberreiter, S.; Radmayr, C.; Bartsch, G.: Point mutation in the DNA binding domain of the
- [99646] androgen receptor in two families with Reifenstein syndrome. *Am.*

[99647] J. Med. Genet. 50: 1318–1327, 1992.

[99648] 17336.

[99649] Knoke, I.; Allera, A.; Wieacker, P.: Significance of the CAG

[99650] repeat length in the androgen receptor gene (AR) for the
transactivation

[99651] function of an M780I mutant AR. Hum. Genet. 104:
257–261, 1999.

[99652] 17337.

[99653] Kobayashi, Y.; Miwa, S.; Merry, D. E.; Kume, A.; Mei, L.;
Doyu,

[99654] M.; Sobue, G.: Caspase–3 cleaves the expanded androgen
receptor protein

[99655] of spinal and bulbar muscular atrophy in a polyglutamine
repeat length–dependent

[99656] manner. Biochem. Biophys. Res. Commun. 252: 145–150,
1998.

[99657] 17338.

[99658] Koivisto, P. A.; Schleutker, J.; Helin, H.; Ehren–van Eekelen,

[99659] C.; Kallioniemi, O.–P.; Trapman, J.: Androgen receptor
gene alterations

[99660] and chromosomal gains and losses in prostate carcinomas

appearing

[99661] during finasteride treatment for benign prostatic hyperplasia. Clin.

[99662] Cancer Res. 5: 3578–3582, 1999.

[99663] 17339.

[99664] Kooy, R. F.; Reyniers, E.; Storm, K.; Vits, L.; van Velzen, D.;

[99665] de Ruiter, P. E.; Brinkmann, A. O.; de Paepe, A.; Willems, P. J.:

[99666] CAG repeat contraction in the androgen receptor gene in three brothers

[99667] with mental retardation. Am. J. Med. Genet. 85: 209–213, 1999.

[99668] 17340.

[99669] La Spada, A.; Fischbeck, K. H.: Androgen receptor gene defect

[99670] in X-linked spinal and bulbar muscular atrophy. (Abstract) Am. J.

[99671] Hum. Genet. 49 (suppl.): 20, 1991.

[99672] 17341.

[99673] Lewis, M.; Kaita, H.; Giblett, E. R.; Anderson, J.; Philipps,

[99674] S.; Steinberg, A. G.; McAlpine, P. J.: Multiplicity of genetic

polymorphisms

[99675] of blood in the Schmiedeleut Hutterites. Am. J. Med.

Genet. 22:

[99676] 477–485, 1985.

[99677] 17342.

[99678] Lim, H. N.; Chen, H.; McBride, S.; Dunning, A. M.; Nixon,

R. M.;

[99679] Hughes, I. A.; Hawkins, J. R.: Longer polyglutamine tracts

in the

[99680] androgen receptor are associated with moderate to severe

undermasculinized

[99681] genitalia in XY males. Hum. Molec. Genet. 9: 829–834,

2000.

[99682] 17343.

[99683] Lobaccaro, J.-M.; Lumbroso, S.; Belon, C.; Galtier-Dereure,

F.;

[99684] Bringer, J.; Lesimple, T.; Namer, M.; Cutuli, B. F.; Pujol, H.;

Sultan,

[99685] C.: Androgen receptor gene mutation in male breast can-

cer. Hum.

[99686] Molec. Genet. 2: 1799–1802, 1993.

[99687] 17344.

- [99688] Lubahn, D. B.; Brown, T. R.; Simental, J. A.; Higgs, H. N.; Migeon,
- [99689] C. J.; Wilson, E. M.; French, F. S.: Sequence of the intron/
exon
- [99690] junctions of the coding region of the human androgen re-
ceptor gene
- [99691] and identification of a point mutation in a family with
complete androgen
- [99692] insensitivity. Proc. Nat. Acad. Sci. 86: 9534–9538, 1989.
- [99693] 17345.
- [99694] Lubahn, D. B.; Joseph, D. R.; Sar, M.; Tan, J.; Higgs, H. N.;
- [99695] Larson, R. E.; French, F. S.; Wilson, E. M.: The human an-
drogen receptor:
- [99696] complementary deoxyribonucleic acid cloning, sequence
analysis and
- [99697] gene expression in prostate. Molec. Endocr. 2:
1265–1275, 1988.
- [99698] 17346.
- [99699] Lubahn, D. B.; Joseph, D. R.; Sullivan, P. M.; Willard, H. F.;
- [99700] French, F. S.; Wilson, E. M.: Cloning of human androgen
receptor

- [99701] complementary DNA and localization to the X chromosome. *Science* 240:
- [99702] 327–330, 1988.
- [99703] 17347.
- [99704] Lumbroso, S.; Lobaccaro, J. M.; Georget, V.; Leger, J.; Pouljol,
- [99705] N.; Terouanne, B.; Evain-Brion, D.; Czernichow, P.; Sultan, C.: A
- [99706] novel substitution (leu707-to-arg) in exon 4 of the androgen receptor
- [99707] gene causes complete androgen resistance. *J. Clin. Endocrinol. Metab.* 81:
- [99708] 1984–1988, 1996.
- [99709] 17348.
- [99710] Stuppia, L.; Calabrese, G.; Franchi, P. G.; Mingarelli, R.; Gatta,
- [99711] V.; Palka, G.; Dallapiccola, B.: Widening of a Y-chromosome interval–6
- [99712] deletion transmitted from a father to his infertile son accounts for
- [99713] an oligozoospermia critical region distal to the RBM1 and DAZ genes.

- [99714] (Letter) Am. J. Hum. Genet. 59: 1393–1395, 1996.
- [99715] 17349.
- [99716] Brown, G. M.; Furlong, R. A.; Sargent, C. A.; Erickson, R. P.;
- [99717] Longepied, G.; Mitchell, M.; Jones, M. H.; Hargreave, T. B.;
Cooke,
- [99718] H. J.; Affara, N. A.: Characterisation of the coding se-
quence and
- [99719] fine mapping of the human DFFRY gene and comparative
expression analysis
- [99720] and mapping to the Sxr-b interval of the mouse Y chro-
mosome of the
- [99721] Dffry gene. Hum. Molec. Genet. 7: 97–107, 1998.
- [99722] 17350.
- [99723] Ferlin, A.; Moro, E.; Garolla, A.; Foresta, C.: Human male
infertility
- [99724] and Y chromosome deletions: role of the AZF-candidate
genes DAZ, RBM
- [99725] and DFFRY. Hum. Reprod. 14: 1710–1716, 1999.
- [99726] 17351.
- [99727] Foresta, C.; Ferlin, A.; Moro, E.: Deletion and expression
analysis

- [99728] of AZFa genes on the human Y chromosome revealed a major role for
- [99729] DBY in male infertility. *Hum. Molec. Genet.* 9: 1161–1169, 2000.
- [99730] 17352.
- [99731] Sargent, C. A.; Boucher, C. A.; Kirsch, S.; Brown, G.; Weiss, B.; Trundley, A.; Burgoyne, P.; Saut, N.; Durand, C.; Levy, N.; Terriou, P.; Hargreave, T.; Cooke, H.; Mitchell, M.; Rappold, G. A.; Affara, N. A.: The critical region of overlap defining the AZFa male infertility interval of proximal Yq contains three transcribed sequences. *J. Med. Genet.* 36: 670–677, 1999.
- [99732] 17353.
- [99733] Shen, P.; Wang, F.; Underhill, P. A.; Franco, C.; Yang, W.-H.; Roxas, A.; Sung, R.; Lin, A. A.; Hyman, R. W.; Vollrath, D.; Davis, R. W.; Cavalli-Sforza, L. L.; Oefner, P. J.: Population genetic implications

- [99741] from sequence variation in four Y chromosome genes.
Proc. Nat. Acad.
- [99742] Sci. 97: 7354–7359, 2000.
- [99743] 17354.
- [99744] Sun, C.; Skaletsky, H.; Birren, B.; Devon, K.; Tang, Z.; Silber,
ber,
- [99745] S.; Oates, R.; Page, D. C.: An azoospermic man with a de
novo point
- [99746] mutation in the Y-chromosomal gene USP9Y. Nature
Genet. 23: 429–432,
- [99747] 1999.
- [99748] 17355.
- [99749] Thomson, R.; Pritchard, J. K.; Shen, P.; Oefner, P. J.; Feld-
man,
- [99750] M. W.: Recent common ancestry of human Y chromo-
somes: evidence from
- [99751] DNA sequence data. Proc. Nat. Acad. Sci. 97: 7360–7365,
2000.
- [99752] 17356.
- [99753] Schiebel, K.; Winkelmann, M.; Mertz, A.; Xu, X.; Page, D.
C.; Weil,

- [99754] D.; Petit, C.; Rappold,, G. A.: Abnormal XY interchange between a
- [99755] novel isolated protein kinase gene, PRKY, and its homologue, PRKX,
- [99756] accounts for one third of all (Y+)XX males and (Y-)XY females. Hum.
- [99757] Molec. Genet. 6: 1985-1989, 1997.
- [99758] 17357.
- [99759] Scott, D.; Addey, C.; Ellis, P; James, E.; Mitchell, M. J.; Saut,
- [99760] N.; Jurcevic, S.; Simpson, E.: Dendritic cells permit identification
- [99761] of genes encoding MHC class II-restricted epitopes of transplantation
- [99762] antigens. Immunity 12: 711-720, 2000.
- [99763] 17358.
- [99764] Wu, G.; Chai, J.; Suber, T. L.; Wu, J.-W.; Du, C.; Wang, X.; Shi,
- [99765] Y.: Structural basis of IAP recognition by Smac/DIABLO. Nature 408:
- [99766] 1008-1012, 2000.
- [99767] 17359.

- [99768] Shih, V. E.: Urea cycle disorders and other congenital hyperammonemic
- [99769] syndromes. In: Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D.
- [99770] S.: The Metabolic Basis of Inherited Disease. New York: McGraw-Hill
- [99771] (pub.) (4th ed.): 1978. Pp. 362–386.
- [99772]
- [99773] 100. Shih, V. E.; Berson, E. L.; Mandell, R.; Schmidt, S. Y.: Ornithine
- [99774] ketoacid transaminase deficiency in gyrate atrophy of the choroid
- [99775] and retina. Am. J. Hum. Genet. 30: 174–179, 1978.
- [99776]
- [99777] 101. Short, E. M.; Conn, H. O.; Snodgrass, P. J.; Campbell, A. G.
- [99778] M.; Rosenberg, L. E.: Evidence for X-linked dominant inheritance
- [99779] of ornithine transcarbamylase deficiency. New Eng. J. Med. 288:
- [99780] 7–12, 1973.

[99781]

[99782] 102. Snodgrass, P. J.; Wappner, R. S.; Brandt, I. K.: White cell

[99783] ornithine transcarbamylase activity cannot detect the liver enzyme

[99784] deficiency. (Letter) *Pediat. Res.* 12: 873, 1978.

[99785]

[99786] 103. Stoll, C.; Bieth, R.; Dreyfus, J.; Flori, E.; Lutz, P.; Levy,

[99787] J.-M.: Une nouvelle famille avec mutation du gene de structure de

[99788] l'ornithine carbamyltransferase humaine. *Arch. Franc. Pe-diat.* 35:

[99789] 512-518, 1978.

[99790]

[99791] 104. Strautnieks, S.; Rutland, P.; Malcolm, S.: Arginine 109 to glutamine

[99792] mutation in a girl with ornithine carbamoyl transferase deficiency. *J.*

[99793] *Med. Genet.* 28: 871-874, 1991.

[99794]

- [99795] 105. Suess, P. J.; Tsai, M. Y.; Holzknecht, R. A.; Horowitz, M.; Tuchman, M.: Screening for gene deletions and known mutations in 13 patients with ornithine transcarbamylase deficiency. *Biochem. Med. Metab.* Biol. 47: 250–259, 1992.
- [99799]
- [99800] 106. Sunshine, P.; Lindenbaum, J. E.; Levy, H. L.; Freeman, J. M. : Hyperammonemia due to a defect in hepatic ornithine transcarbamylase. *Pediatrics* 50: 100–111, 1972.
- [99803]
- [99804] 107. Thaler, M. M.; Hoogenraad, N. J.; Boswell, M.: Reye's syndrome due to a novel protein-tolerant variant of ornithine-transcarbamylase deficiency. *Lancet* II: 438–440, 1974.
- [99807]
- [99808] 108. Tripp, J. H.; Hargreaves, T.; Anthony, P. P.; Searle, J.

F.;

[99809] Miller, P.; Leonard, J. V.; Patrick, A. D.; Oberholzer, V. G.:

Sodium

[99810] valproate and ornithine carbamyl transferase deficiency.

(Letter) Lancet I:

[99811] 1165–1166, 1981.

[99812]

[99813] 109. Tuchman, M.: Mutations and polymorphisms in the human ornithine

[99814] transcarbamylase gene. Hum. Mutat. 2: 174–178, 1993.

[99815]

[99816] 110. Tuchman, M.; Holzknecht, R. A.; Gueron, A. B.; Berry, S. A.;

[99817] Tsai, M. Y.: Six new mutations in the ornithine transcarbamylase

[99818] gene detected by single-strand conformational polymorphism. Pediat.

[99819] Res. 32: 600–604, 1992.

[99820]

[99821] 111. Tuchman, M.; Jaleel, N.; Morizono, H.; Sheehy, L.; Lynch, M.

- [99822] G.: Mutations and polymorphisms in the human ornithine transcarbamylase
- [99823] gene. Hum. Mutat. 19: 93–107, 2002.
- [99824]
- [99825] 112. Tuchman, M.; Matsuda, I.; Munnich, A.; Malcolm, S.; Strautnieks,
- [99826] S.; Briede, T.: Proportions of spontaneous mutations in males and
- [99827] females with ornithine transcarbamylase deficiency. Am. J. Med. Genet. 55:
- [99828] 67–70, 1995.
- [99829]
- [99830] 113. Tuchman, M.; Morizono, H.; Reish, O.; Yuan, X.; Allewell, N.
- [99831] M.: The molecular basis of ornithine transcarbamylase deficiency:
- [99832] modelling the human enzyme and the effects of mutations. J. Med.
- [99833] Genet. 32: 680–688, 1995.
- [99834]
- [99835] 114. Tuchman, M.; Plante, R. J.; Garcia-Perez, M. A.; Ru-

bio, V.:

[99836] Relative frequency of mutations causing ornithine transcarbamylase

[99837] deficiency in 78 families. Hum. Genet. 97: 274–276, 1996.

[99838]

[99839] 115. Tuchman, M.; Plante, R. J.; McCann, M. T.; Qureshi, A. A.: Seven

[99840] new mutations in the human ornithine transcarbamylase gene. Hum.

[99841] Mutat. 4: 57–60, 1994.

[99842]

[99843] 116. Veres, G.; Gibbs, R. A.; Scherer, S. E.; Caskey, C. T.: The

[99844] molecular basis of the sparse fur mouse mutation. Science 237: 415–417,

[99845] 1987.

[99846]

[99847] 117. Wareham, K. A.; Lyon, M. F.; Glenister, P. H.; Williams, E. D.

[99848] : Age related reactivation of an X-linked gene. Nature 327: 725–727,

[99849] 1987.

[99850]

[99851] 118. Wettke-Schafer, R.; Kantner, G.: X-linked dominant inherited

[99852] diseases with lethality in hemizygous males. Hum. Genet. 64: 1-23,

[99853] 1983.

[99854]

[99855] 119. Wilson, C. J.; Lee, P. J.; Leonard, J. V.: Plasma glutamine

[99856] and ammonia concentrations in ornithine carbamoyltransferase deficiency

[99857] and citrullinaemia. J. Inherit. Metab. Dis. 24: 691-695, 2001.

[99858]

[99859] 120. Winter, R. M.; Donnai, D.; Burn, J.; Tuckers, S. M.: Fetal valproate

[99860] syndrome: is there a recognisable phenotype?. J. Med. Genet. 24:

[99861] 692-695, 1987.

[99862]

- [99863] 121. Yorifuji, T.; Muroi, J.; Uematsu, A.; Tanaka, K.; Kiwaki, K.;
- [99864] Endo, F.; Matsuda, I.; Nagasaka, H.; Furusho, K.: X-inactivation
- [99865] pattern in the liver of a manifesting female with ornithine transcarbamylase
- [99866] (OTC) deficiency. Clin. Genet. 54: 349–353, 1998.
- [99867]
- [99868] 122. Yudkoff, M.; Daikhin, Y.; Nissim, I.; Jawad, A.; Wilson, J.;
- [99869] Batshaw, M.: In vivo nitrogen metabolism in ornithine transcarbamylase
- [99870] deficiency. J. Clin. Invest. 98: 2167–2173, 1996.
- [99871]
- [99872] 123. Yudkoff, M.; Yang, W.; Snodgrass, P. J.; Segal, S.: Ornithine
- [99873] transcarbamylase deficiency in a boy with normal development. J.
- [99874] Pediat. 96: 441–443, 1980.
- [99875] 17360.
- [99876] Macke, J. P.; Hu, N.; Hu, S.; Bailey, M.; King, V. L.; Brown,

[99877] T.; Hamer, D.; Nathans, J.: Sequence variation in the androgen receptor

- [99878] gene is not a common determinant of male sexual orientation. Am.
- [99879] J. Hum. Genet. 53: 844–852, 1993.
- [99880] 17361.
- [99881] Nakamura, H.; Izumoto, Y.; Kambe, H.; Kuroda, T.; Mori, T.; Kawamura,
- [99882] K.; Yamamoto, H.; Kishimoto, T.: Molecular cloning of complementary
- [99883] DNA for a novel human hepatoma–derived growth factor: its homology
- [99884] with high mobility group–1 protein. J. Biol. Chem. 269: 25143–25149,
- [99885] 1994.
- [99886] 17362.
- [99887] Wanschura, S.; Schoenmakers, E. F. P. M.; Huysmans, C.; Bartnitzke,
- [99888] S.; Van de Ven, W. J. M.; Bullerdiek, J.: Mapping of the gene encoding
- [99889] the human hepatoma–derived growth factor (HDGF) with homology to the
- [99890] high–mobility group (HMG)–1 protein to Xq25. Genomics 32: 298–300,

- [99891] 1996.
- [99892] 17363.
- [99893] Gedeon, A. K.; Glass, I. A.; Connor, J. M.; Mulley, J. C.: Genetic
- [99894] localisation of MRX27 to Xq24-26 defines another discrete gene for
- [99895] non-specific X-linked mental retardation. Am. J. Med. Genet. 64:
- [99896] 121-124, 1996.
- [99897] 17364.
- [99898] Amir, J.; Alpert, G.; Statter, M.; Gutman, A.; Reisner, S. H.:
- [99899] Intracranial haemorrhage in siblings and ornithine transcarbamylase
- [99900] deficiency. Acta Paediat. Scand. 71: 671-673, 1982.
- [99901] 17365.
- [99902] Anadiotis, G.; Ierardi-Curto, L.; Kaplan, P. B.; Berry, G. T.:
- [99903] Ornithine transcarbamylase deficiency and pancreatitis. J. Pediat. 138:
- [99904] 123-124, 2001.
- [99905] 17366.

- [99906] Ardinger, H. H.; Atkin, J. F.; Blackston, R. D.; Elsas, L. J.;
- [99907] Clarren, S. K.; Livingstone, S.; Flannery, D. B.; Pellock, J. M.;
- [99908] Harrod, M. J.; Lammer, E. J.; Majewski, F.; Schinzel, A.; Toriello,
- [99909] H. V.; Hanson, J. W.: Verification of the fetal valproate syndrome
- [99910] phenotype. *Am. J. Med. Genet.* 29: 171–185, 1988.
- [99911] 17367.
- [99912] McMurtrie, E. B.; Barbosa, M. D. F. S.; Zerial, M.; Kingsmore,
- [99913] S. F.: Rab17 and Rab18, small GTPases with specificity for polarized
- [99914] epithelial cells: genetic mapping in the mouse. *Genomics* 45: 623–625,
- [99915] 1997.
- [99916] 17368.
- [99917] Schafer, U.; Seibold, S.; Schneider, A.; Neugebauer, E.: Isolation
- [99918] and characterisation of the human rab18 gene after stimulation of

- [99919] endothelial cells with histamine. FEBS Lett. 466: 148–154, 2000.
- [99920] 17369.
- [99921] Doupnik, C. A.; Davidson, N.; Lester, H. A.: The inward rectifier
- [99922] potassium channel family. Curr. Opin. Neurobiol. 5: 268–277, 1995.
- [99923] 17370.
- [99924] Tada, Y.; Horio, Y.; Takumi, T.; Terayama, M.; Tsuji, L.; Copeland,
- [99925] N. G.; Jenkins, N. A.; Kurachi, Y.: Assignment of the glial inwardly
- [99926] rectifying potassium channel K(AB)–2/Kir4.1 (Kcnj10) gene to the distal
- [99927] region of mouse chromosome 1. Genomics 45: 629–630, 1997.
- [99928] 17371.
- [99929] Takumi, T.; Ishii, T.; Horio, Y.; Morishige, K.–I.; Takahashi,
- [99930] N.; Yamada, M.; Yamashita, T.; Kiyama, H.; Sohmiya, K.; Nakanishi,
- [99931] S.; Kurachi, Y.: A novel ATP–dependent inward rectifier potassium

- [99932] channel expressed predominantly in glial cells. J. Biol. Chem. 270: 16339–16436, 1995.
- [99933] 17372.
- [99934] Schenker, T.; Trueb, B.: Assignment of the gene for a developmentally regulated GTP-binding protein (DRG2) to human chromosome bands 17p13–p12 by in situ hybridization. Cytogenet. Cell Genet. 79: 274–275, 1997.
- [99935] 17373.
- [99936] Heikkinen, J.; Risteli, M.; Wang, C.; Latvala, J.; Rossi, M.; Valtavaara, M.; Myllyla, R.: Lysyl hydroxylase 3 is a multifunctional protein possessing collagen glucosyltransferase activity. J. Biol. Chem. 275: 36158–36163, 2000.
- [99937] 17374.
- [99938] Passoja, K.; Rautavuoma, K.; Ala-Kokko, L.; Kosonen, T.; Kivirikko,

- [99945] K. I.: Cloning and characterization of a third human lysyl hydroxylase
- [99946] isoform. Proc. Nat. Acad. Sci. 95: 10482–10486, 1998.
- [99947] 17375.
- [99948] Valtavaara, M.; Szpirer, C.; Szpirer, J.; Myllyla, R.: Primary
- [99949] structure, tissue distribution, and chromosomal localization of a
- [99950] novel isoform of lysyl hydroxylase (lysyl hydroxylase 3). J. Biol.
- [99951] Chem. 273: 12881–12886, 1998.
- [99952] 17376.
- [99953] Ishibashi, T.; Bottaro, D. P.; Michieli, P.; Kelley, C. A.; Aaronson,
- [99954] S. A.: A novel dual specificity phosphatase induced by serum stimulation
- [99955] and heat shock. J. Biol. Chem. 269: 29897–29902, 1994.
- [99956] 17377.
- [99957] Kwak, S. P.; Dixon, J. E.: Multiple dual specificity protein tyrosine
- [99958] phosphatases are expressed and regulated differentially in liver cell

- [99959] lines. J. Biol. Chem. 270: 1156–1160, 1995.
- [99960] 17378.
- [99961] Kovalenko, O. V.; Golub, E. I.; Bray–Ward, P.; Ward, D. C.;
Radding,
- [99962] C. M.: A novel nucleic acid–binding protein that interacts
with human
- [99963] Rad51 recombinase. Nucleic Acids Res. 25: 4946–4953,
1997.
- [99964] 17379.
- [99965] Mizuta, R.; LaSalle, J. M.; Cheng, H.–L.; Shinohara, A.;
Ogawa,
- [99966] H.; Copeland, N.; Jenkins, N. A.; Lalande, M.; Alt, F. W.:
RAB22
- [99967] and RAB163/mouse BRCA2: proteins that specifically in–
teract with the
- [99968] RAD51 protein. Proc. Nat. Acad. Sci. 94: 6927–6932,
1997.
- [99969] 17380.
- [99970] Blasius, R.; Weber, R. G.; Lichter, P.; Ogilvie, A.: A novel
orphan
- [99971] G protein–coupled receptor primarily expressed in the
brain is localized

- [99972] on human chromosomal band 2q21. J. Neurochem. 70: 1357–1365, 1998.
- [99973] 17381.
- [99974] Raport, C. J.; Schweickart, V. L.; Chantry, D.; Eddy, R. L., Jr.;
- [99975] Shows, T. B.; Godiska, R.; Gray, P. W.: New members of the chemokine
- [99976] receptor gene family. J. Leukoc. Biol. 59: 18–23, 1996.
- [99977] 17382.
- [99978] Shindo, M.; Nakano, H.; Kuroyanagi, H.; Shirasawa, T.; Mi-hara,
- [99979] M.; Gilbert, D. J.; Jenkins, N. A.; Copeland, N. G.; Yagita, H.; Okumura,
- [99980] K.: cDNA cloning, expression, subcellular localization, and chromosomal
- [99981] assignment of mammalian aurora homologues, aurora-related kinase (ARK)
- [99982] 1 and 2. Biochem. Biophys. Res. Commun. 244: 285–292, 1998.
- [99983] 17383.
- [99984] Presky, D. H.; Yang, H.; Minetti, L. J.; Chua, A. O.; Nabavi, N.;

- [99985] Wu, C.-Y.; Gately, M. K.; Gubler, U.: A functional interleukin 12
- [99986] receptor complex is composed of two beta-type cytokine receptor subunits. Proc.
- [99987] Nat. Acad. Sci. 93: 14002-14007, 1996.
- [99988] 17384.
- [99989] Yamamoto, K.; Kobayashi, H.; Miura, O.; Hirose, S.; Miyasaka,
- [99990] N.: Assignment of IL12RB1 and IL12RB2, interleukin-12 receptor beta-1
- [99991] and beta-2 chains, to human chromosome 19 band p13.1 and chromosome
- [99992] 1 band p31.2, respectively, by in situ hybridization. Cytogenet.
- [99993] Cell Genet. 77: 257-258, 1997.
- [99994] 17385.
- [99995] Turelli, P.; Doucas, V.; Craig, E.; Mangeat, B.; Klages, N.; Evans,
- [99996] R.; Kalpana, G.; Trono, D.: Cytoplasmic recruitment of INI1 and PML
- [99997] on incoming HIV preintegration complexes: interference with early

[99998] steps of viral replication. *Molec. Cell* 7: 1245–1254, 2001.

[99999] 17386.

[100000] Lafreniere, R. G.; Rochefort, D. L.; Kibar, Z.; Fon, E. A.;
Han,

[100001] F.-Y.; Cochius, J.; Kang, X.; Baird, S.; Korneluk, R. G.; An-
dermann,

[100002] E.; Rommens, J. M.; Rouleau, G. A.: Isolation and charac-
terization

[100003] of GT335, a novel human gene conserved in *Escherichia*
coli and mapping

[100004] to 21q22.3. *Genomics* 38: 264–272, 1996.

[100005] 17387.

[100006] Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.;
Asakawa,

[100007] S.; Ito, F.; Shimizu, N.: Isolation of cDNA for a novel hu-
man protein

[100008] KNP-I that is homologous to the *E. coli* SCRP-27A protein
from the

[100009] autoimmune polyglandular disease type I (APECED) region
of chromosome

[100010] 21q22.3. *Biochem. Biophys. Res. Commun.* 225: 608–616,
1996.

[100011]17388.

[100012]Scott, H. S.; Chen, H.; Rossier, C.; Lalioti, M. D.; Antonarakis,

[100013]S. E.: Isolation of a human gene (HES1) with homology to an Escherichia

[100014]coli and a zebrafish protein that maps to chromosome 21q22.3. Hum.

[100015]Genet. 99: 616–623, 1997.

[100016]17389.

[100017]Bernier-Villamor, V.; Sampson, D. A.; Matunis, M. J.; Lima, C.

[100018]D.: Structural basis for E2-mediated SUMO conjugation revealed by

[100019]a complex between ubiquitin-conjugating enzyme Ubc9 and RanGAP1. Cell 108:

[100020]345–356, 2002.

[100021]17390.

[100022]Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association of

[100023]FHIT (fragile histidine triad), a candidate tumour suppressor gene,

- [100024]with the ubiquitin-conjugating enzyme hUBC9. *Biochem. J.* 352: 443-448,
- [100025]2000.
- [100026]17391.
- [100027]Tachibana, M.; Iwata, N.; Watanabe, A.; Nobukuni, Y.; Plo-
plis,
- [100028]B.; Kajigaya, S.: Assignment of the gene for a ubiquitin-
conjugating
- [100029]enzyme (UBE2I) to human chromosome band 16p13.3 by
in situ hybridization. *Cytogenet.*
- [100030]*Cell Genet.* 75: 222-223, 1996.
- [100031]17392.
- [100032]Wang, Z.-Y.; Qiu, Q.-Q.; Seufert, W.; Taguchi, T.; Testa, J.
R.;
- [100033]Whitmore, S. A.; Callen, D. F.; Welsh, D.; Shenk, T.; Deuel,
T. F.
- [100034]: Molecular cloning of the cDNA and chromosome local-
ization of the
- [100035]gene for human ubiquitin-conjugating enzyme 9. *J. Biol.*
Chem. 271:
- [100036]24811-24816, 1996.

[100037]17393.

[100038]Yasugi, T.; Howley, P. M.: Identification of the structural
and

[100039]functional human homolog of the yeast ubiquitin conjugating enzyme

[100040]UBC9. Nucleic Acids Res. 24: 2005–2010, 1996.

[100041]17394.

[100042]Bowen, M. A.; Patel, D. D.; Li, X.; Modrell, B.; Malacko, A.
R.;

[100043]Wang, W.-C.; Marquardt, H.; Neubauer, M.; Pesando, J. M.;
Francke,

[100044]U.; Haynes, B. F.; Aruffo, A.: Cloning, mapping, and characterization

[100045]of activated leukocyte–cell adhesion molecule (ALCAM), a
CD6 ligand. J.

[100046]Exp. Med. 181: 2213–2220, 1995.

[100047]17395.

[100048]Skonier, J. E.; Bowen, M. A.; Emswiler, J.; Aruffo, A.; Bajorath,

[100049]J.: Recognition of diverse proteins by members of the immunoglobulin

[100050]superfamily: delineation of the receptor binding site in the human

[100051]CD6 ligand ALCAM. Biochemistry 35: 12287–12291, 1996.

[100052]17396.

[100053]Bord, S.; Horner, A.; Beavan, S.; Compston, J.: Estrogen receptors

[100054]alpha and beta are differentially expressed in developing human bone. J.

[100055]Clin. Endocr. Metab. 86: 2309–2314, 2001.

[100056]17397.

[100057]Watanabe, T. K.; Fujiwara, T.; Kawai, A.; Shimizu, F.; Takami,

[100058]S.; Hirano, H.; Okuno, S.; Ozaki, K.; Takeda, S.; Shimada, Y.; Nagata,

[100059]M.; Takaichi, A.; Takahashi, E.; Nakamura, Y.; Shin, S.: Cloning,

[100060]expression, and mapping of UBE2I, a novel gene encoding a human homologue

[100061]of yeast ubiquitin-conjugating enzymes which are critical for regulating

[100062]the cell cycle. Cytogenet. Cell Genet. 72: 86–89, 1996.

[100063]17398.

[100064]Chaidarun, S. S.; Swearingen, B.; Alexander, J. M.: Differential

[100065]expression of estrogen receptor-beta (ER-beta) in human
pituitary

[100066]tumors: functional interactions with ER-alpha and a tumor-specific

[100067]splice variant. J. Clin. Endocr. Metab. 83: 3308-3315,
1998.

[100068]17399.

[100069]Chu, S.; Mamers, P.; Burger, H. G.; Fuller, P. J.: Estrogen
receptor

[100070]isoform gene expression in ovarian stromal and epithelial
tumors. J.

[100071]Clin. Endocr. Metab. 85: 1200-1205, 2000.

[100072]17400.

[100073]Enmark, E.; Peltö-Huikko, M.; Grandien, K.; Lagercrantz,
S.; Lagercrantz,

[100074]J.; Fried, G.; Nordenskjöld, M.; Gustafsson, J.-Å.: Human
estrogen

[100075]receptor beta-gene structure, chromosomal localization,
and expression

[100076]pattern. J. Clin. Endocr. Metab. 82: 4258–4265, 1997.

[100077]17401.

[100078]Esmaeli, B.; Harvey, J. T.; Hewlett, B.: Immunohistochemical evidence

[100079]for estrogen receptors in meibomian glands. Ophthalmology 107: 180–184,

[100080]2000.

[100081]17402.

[100082]Forsell, C.; Enmark, E.; Axelman, K.; Blomberg, M.; Wahlund, L.-O.;

[100083]Gustafsson, J.-A.; Lannfelt, L.: Investigations of a CA repeat in

[100084]the oestrogen receptor beta gene in patients with Alzheimer's disease. Europ.

[100085]J. Hum. Genet. 9: 802–804, 2001.

[100086]17403.

[100087]Korach, K. S.; Couse, J. F.; Curtis, S. W.; Washburn, T. F.; Lindzey,

[100088]J.; Kimbro, K. S.; Eddy, E. M.; Migliaccio, S.; Snedeker, S. M.; Lubahn,

[100089]D. B.; Schomberg, D. W.; Smith, E. P.: Estrogen receptor

gene disruption:

[100090]molecular characterization and experimental and clinical phenotypes. Recent

[100091]Prog. Horm. Res. 51: 159–188, 1996.

[100092]17404.

[100093]Krege, J. H.; Hodgin, J. B.; Couse, J. F.; Enmark, E.; Warner,

[100094]M.; Mahler, J. F.; Sar, M.; Korach, K. S.; Gustafsson, J.-A.; Smithies,

[100095]O.: Generation and reproductive phenotypes of mice lacking estrogen

[100096]receptor beta. Proc. Nat. Acad. Sci. 95: 15677–15682, 1998.

[100097]17405.

[100098]Krezel, W.; Dupont, S.; Krust, A.; Chambon, P.; Chapman, P. F.

[100099]: Increased anxiety and synaptic plasticity in estrogen receptor beta-deficient

[100100]mice. Proc. Nat. Acad. Sci. 98: 12278–12282, 2001.

[100101]17406.

[100102]Kuiper, G. G. J. M.; Enmark, E.; Pelto-Huikko, M.; Nilsson, S.;

[100103]Gustafsson, J.-A.: Cloning of a novel estrogen receptor expressed

[100104]in rat prostate and ovary. Proc. Nat. Acad. Sci. 93: 5925–5930,

[100105]1996.

[100106]17407.

[100107]Mosialos, G.; Birkenbach, M.; Yalamanchili, R.; VanArsdale, T.;

[100108]Ware, C.; Kieff, E.: The Epstein–Barr virus transforming protein

[100109]LMP1 engages signaling proteins for the tumor necrosis factor receptor

[100110]family. Cell 80: 389–399, 1995.

[100111]17408.

[100112]Rothe, M.; Wong, S. C.; Henzel, W. J.; Goeddel, D. V.: A novel

[100113]family of putative signal transducers associated with the cytoplasmic

[100114]domain of the 75 kDa tumor necrosis factor receptor. Cell 78: 681–692,

[100115]1994.

[100116]17409.

[100117]Sieminski, K.; Peters, N.; Scheurich, P.; Wajant, H.: Organization

[100118]of the human tumour necrosis factor receptor-associated factor 1 (TRAF1)

[100119]gene and mapping to chromosome 9q33-34. Gene 195: 35-39, 1997.

[100120]17410.

[100121]Tsitsikov, E. N.; Laouini, D.; Dunn, I. F.; Sannikova, T. Y.; Davidson,

[100122]L.; Alt, F. W.; Geha, R. S.: TRAF1 is a negative regulator of TNF

[100123]signaling: enhanced TNF signaling in TRAF1-deficient mice. Immunity 15:

[100124]647-657, 2001.

[100125]17411.

[100126]Miyamoto, K.; Kesterson, R. A.; Yamamoto, H.; Taketani, Y.; Nishiwaki,

[100127]E.; Tatsumi, S.; Inoue, Y.; Morita, K.; Takeda, E.; Pike, J. W.:

[100128]Structural organization of the human vitamin D receptor chromosomal

[100129]gene and its promoter. Molec. Endocr. 11: 1165–1179, 1997.

[100130]17412.

[100131]Pedeutour, F.; Merscher, S.; Durieux, E.; Montgomery, K.; Krauter,

[100132]K.; Clevy, J.-P.; Barcelo, G.; Kucherlapati, R.; Gaudray, P.; Turc-Carel,

[100133]C.: Mapping of the 12q12–q22 region with respect to tumor translocation

[100134]breakpoints. Genomics 22: 512–518, 1994.

[100135]17413.

[100136]Riggs, B. L.: Vitamin D-receptor genotypes and bone density.

[100137](Editorial) New Eng. J. Med. 337: 125–126, 1997.

[100138]17414.

[100139]Ritchie, H. H.; Hughes, M. R.; Thompson, E. T.; Malloy, P. J.;

[100140]Hochberg, Z.; Feldman, D.; Pike, J. W.; O'Malley, B. W.: An ochre

[100141]mutation in the vitamin D receptor gene causes hereditary 1,25-dihydroxyvitamin

[100142]D3-resistant rickets in three families. Proc. Nat. Acad. Sci.

86:

[100143]9783–9787, 1989.

[100144]17415.

[100145]Saijo, T.; Ito, M.; Takeda, E.; Mahbubul Huq, A. H. M.;

Naito,

[100146]E.; Yokota, I.; Sone, T.; Pike, J. W.; Kuroda, Y.: A unique

mutation

[100147]in the vitamin D receptor gene in three Japanese patients

with vitamin

[100148]D-dependent rickets type II: utility of single-strand con-

formation

[100149]polymorphism analysis for heterozygous carrier detection.

Am. J.

[100150]Hum. Genet. 49: 668–673, 1991.

[100151]17416.

[100152]Sainz, J.; Van Tornout, J. M.; Loro, L.; Sayre, J.; Roe, T. F.;

[100153]Gilsanz, V.: Vitamin D-receptor gene polymorphisms and

bone density

[100154]in prepubertal American girls of Mexican descent. New

Eng. J. Med. 337:

[100155]77–82, 1997.

[100156]17417.

[100157]Sone, T.; Marx, S. J.; Liberman, U. A.; Pike, J. W.: A unique

[100158]point mutation in the human vitamin D receptor chromo-
somal gene confers

[100159]hereditary resistance to 1,25-dihydroxyvitamin D(3).

Molec. Endocr. 4:

[100160]623–631, 1990.

[100161]17418.

[100162]Suarez, F.; Zeghoud, F.; Rossignol, C.; Walrant, O.;

Garabedian,

[100163]M.: Association between vitamin D receptor gene poly-
morphism and

[100164]sex-dependent growth during the first two years of life. J.
Clin.

[100165]Endocr. Metab. 82: 2966–2970, 1997.

[100166]17419.

[100167]Tao, C.; Yu, T.; Garnett, S.; Briody, J.; Knight, J.; Wood-
head,

[100168]H.; Cowell, C. T.: Vitamin D receptor alleles predict growth
and

[100169]bone density in girls. Arch. Dis. Child. 79: 488–494, 1998.

[100170]17420.

[100171]Uitterlinden, A. G.; Burger, H.; Huang, Q.; Odding, E.; van
Duijn,

[100172]C. M.; Hofman, A.; Birkenhager, J. C.; van Leeuwen, J. P. T.
M.; Pols,

[100173]H. A. P.: Vitamin D receptor genotype is associated with
radiographic

[100174]osteoarthritis at the knee. J. Clin. Invest. 100: 259–263,
1997.

[100175]17421.

[100176]Uitterlinden, A. G.; Weel, A. E. A. M.; Burger, H.; Fang, Y.;

[100177]Van Duijn, C. M.; Hofman, A.; Van Leeuwen, J. P. T. M.;
Pols, H. A.

[100178]P.: Interaction between the vitamin D receptor gene and
collagen

[100179]type I- α -1 gene in susceptibility for fracture. J. Bone
Miner.

[100180]Res. 16: 379–385, 2001.

[100181]17422.

[100182]Whitfield, G. K.; Selznick, S. H.; Haussler, C. A.; Hsieh,

J.-C.;

[100183]Galligan, M. A.; Jurutka, P. W.; Thompson, P. D.; Lee, S. M.;
Zerwekh,

[100184]J. E.; Haussler, M. R.: Vitamin D receptors from patients
with resistance

[100185]to 1,25-dihydroxyvitamin D(3): point mutations confer re-
duced transactivation

[100186]in response to ligand and impaired interaction with the
retinoid X

[100187]receptor heterodimeric partner. Molec. Endocr. 10:
1617-1631, 1996.

[100188]17423.

[100189]Yoshizawa, T.; Handa, Y.; Uematsu, Y.; Takeda, S.; Sekine,
K.;

[100190]Yoshihara, Y.; Kawakami, T.; Arioka, K.; Sato, H.;
Uchiyama, Y.; Masushige,

[100191]S.; Fukamizu, A.; Matsumoto, T.; Kato, S.: Mice lacking the
vitamin

[100192]D receptor exhibit impaired bone formation, uterine hy-
poplasia and

[100193]growth retardation after weaning. Nature Genet. 16:
391-396, 1997.

[100194]17424.

[100195]Zhu, W.; Malloy, P. J.; Delvin, E.; Chabot, G.; Feldman, D.:

[100196]Hereditary 1,25-dihydroxyvitamin D-resistant rickets due to an opal

[100197]mutation causing premature termination of the vitamin D receptor. J.

[100198]Bone Miner. Res. 13: 259–264, 1998.

[100199]17425.

[100200]Bejjani, B. A.; Lewis, R. A.; Tomey, K. F.; Anderson, K. L.; Dueker,

[100201]D. K.; Jabak, M.; Astle, W. F.; Otterud, B.; Leppert, M.; Lupski,

[100202]J. R.: Mutations in CYP1B1, the gene for cytochrome P4501B1, are

[100203]the predominant cause of primary congenital glaucoma in Saudi Arabia. Am.

[100204]J. Hum. Genet. 62: 325–333, 1998.

[100205]17426.

[100206]Bejjani, B. A.; Stockton, D. W.; Lewis, R. A.; Tomey, K. F.; Dueker,

[100207]D. K.; Jabak, M.; Astle, W. F.; Lupski, J. R.: Multiple CYP1B1 mutations

[100208]and incomplete penetrance in an inbred population segregating primary

[100209]congenital glaucoma suggest frequent de novo events and a dominant

[100210]modifier locus. Hum. Molec. Genet. 9: 367–374, 2000.

[100211]17427.

[100212]Lin, J.; Arnold, H. B.; Della-Fera, M. A.; Azain, M. J.; Hartzell,

[100213]D. L.; Baile, C. A.: Myostatin knockout in mice increases myogenesis

[100214]and decreases adipogenesis. Biochem. Biophys. Res. Commun. 291:

[100215]701–706, 2002.

[100216]17428.

[100217]McPherron, A. C.; Lawler, A. M.; Lee, S.-J.: Regulation of skeletal

[100218]muscle mass in mice by a new TGF- β superfamily member. Nature 387:

[100219]83–90, 1997.

[100220]17429.

[100221]Strautnieks, S. S.; Bull, L. N.; Knisely, A. S.; Kocoshis, S. A.;

[100222]Dahl, N.; Arnell, H.; Sokal, E.; Dahan, K.; Childs, S.; Ling, V.;

[100223]Tanner, M. S.; Kagalwalla, A. F.; Nemeth, A.; Pawlowska, J.; Baker,

[100224]A.; Mieli-Vergani, G.; Freimer, N. B.; Gardiner, R. M.; Thompson,

[100225]R. J.: A gene encoding a liver-specific ABC transporter is mutated

[100226]in progressive familial intrahepatic cholestasis. *Nature Genet.* 20:

[100227]233-238, 1998.

[100228]17430.

[100229]Roldan-Arjona, T.; Wei, Y.-F.; Carter, K. C.; Klungland, A.; Anselmino,

[100230]C.; Wang, R.-P.; Augustus, M.; Lindahl, T.: Molecular cloning and

[100231]functional expression of a human cDNA encoding the antitumor enzyme

[100232]8-hydroxyguanine-DNA glycosylase. *Proc. Nat. Acad. Sci.* 94: 8016-8020,

[100233]1997.

[100234]17431.

[100235]Goldman, M. J.; Anderson, G. M.; Stolzenberg, E. D.; Kari,
U. P.;

[100236]Zasloff, M.; Wilson, J. M.: Human beta-defensin-1 is a
salt-sensitive

[100237]antibiotic in lung that is inactivated in cystic fibrosis. Cell
88:

[100238]553-560, 1997.

[100239]17432.

[100240]Guo, Z.; Turner, C.; Castle, D.: Relocation of the t-SNARE
SNAP-23

[100241]from lamellipodia-like cell surface projections regulates
compound

[100242]exocytosis in mast cells. Cell 94: 537-548, 1998.

[100243]17433.

[100244]Lazo, P. A.; Nadal, M.; Ferrer, M.; Area, E.; Hernandez-Tor-
res,

[100245]J.; Nabokina, S. M.; Mollinedo, F.; Estivill, X.: Genomic or-
ganization,

[100246]chromosomal localization, alternative splicing, and iso-
forms of the

[100247]human synaptosome-associated protein-23 gene impli-
cated in vesicle-membrane

[100248]fusion processes. Hum. Genet. 108: 211–215, 2001.

[100249]17434.

[100250]Mollinedo, F.; Lazo, P. A.: Identification of two isoforms of

[100251]the vesicle–membrane fusion protein SNAP–23 in human
neutrophils and

[100252]HL–60 cells. Biochem. Biophys. Res. Commun. 231:
808–812, 1997.

[100253]17435.

[100254]Ravichandran, V.; Chawla, A.; Roche, P. A.: Identification
of

[100255]a novel syntaxin– and synaptobrevin/VAMP–binding pro-
tein, SNAP–23,

[100256]expressed in non–neuronal tissues. J. Biol. Chem. 271:
13300–13303,

[100257]1996.

[100258]17436.

[100259]Shukla, A.; Corydon, T. J.; Nielsen, S.; Hoffmann, H. J.;
Dahl,

[100260]R.: Identification of three new splice variants of the SNARE
protein

[100261]SNAP–23. Biochem. Biophys. Res. Commun. 285:

320–327, 2001.

[100262]17437.

[100263]Mugita, N.; Kumura, Y.; Ogawa, M.; Saya, H.; Nakao, M.:

Identification

[100264]of a novel, tissue-specific calpain htra-3; a human homologue of the

[100265]Caenorhabditis elegans sex determination gene. Biochem. Biophys.

[100266]Res. Commun. 239: 845–850, 1997.

[100267]17438.

[100268]Bohme, B.; Holtrich, U.; Wolf, G.; Luzius, H.; Grzeschik, K.-H.;

[100269]Strebhardt, K.; Rubsamen-Waigmann, H.: PCR mediated detection of

[100270]a new human receptor-tyrosine-kinase, HEK 2. Oncogene 8: 2857–2862,

[100271]1993.

[100272]17439.

[100273]Ruiz, J. C.; Conlon, F. L.; Robertson, E. J.: Identification of

[100274]novel protein kinases expressed in the myocardium of the developing

[100275]mouse heart. Mech. Dev. 48: 153–164, 1994.

[100276]17440.

[100277]Ashbourne, K. J.; Byth, B. C.; Meijers, J. C. M.; Cox, D. W.:

[100278]Polymorphism of the protein C inhibitor (PCI) gene on
chromosome 14. Hum.

[100279]Molec. Genet. 2: 92 only, 1993.

[100280]17441.

[100281]Sadler, J. E.: Combined factors V and VIII deficiency climbs
onto

[100282]the map. (Editorial) J. Clin. Invest. 99: 555–556, 1997.

[100283]17442.

[100284]Yasuda, T.; Nadano, D.; Iida, R.; Tanaka, Y.; Nakanaga, M.;
Kishi,

[100285]K.: Discovery of a genetic polymorphism of human plasma
protein C

[100286]inhibitor (PCI): genetic survey utilizing isoelectric focusing
followed

[100287]by immunoblotting, immunological and biochemical char-
acterization. Hum.

[100288]Genet. 89: 265–269, 1992.

[100289]17443.

[100290]Arturi, F.; Russo, D.; Schlumberger, M.; du Villard, J.-A.;
Caillou,

[100291]B.; Vigneri, P.; Wicker, R.; Chiefari, E.; Suarez, H. G.; Filetti,

[100292]S.: Iodide symporter gene expression in human thyroid
tumors. J.

[100293]Clin. Endocr. Metab. 83: 2493–2496, 1998.

[100294]17444.

[100295]Cho, J.-Y.; Leveille, R.; Kao, R.; Rousset, B.; Parlow, A. F.;

[100296]Burak, W. E., Jr.; Mazzaferri, E. L.; Jhiang, S. M.: Hormonal
regulation

[100297]of radioiodide uptake activity and Na⁺/I⁻ symporter ex-
pression in

[100298]mammary glands. J. Clin. Endocr. Metab. 85: 2936–2943,
2000.

[100299]17445.

[100300]Couch, R. M.; Dean, H. J.; Winter, J. S.: Congenital hy-
pothyroidism

[100301]caused by defective iodide transport. J. Pediat. 106:
950–953, 1985.

[100302]17446.

- [100303] Li, Z.; Matsuoka, S.; Hryshko, L. V.; Nicoll, D. A.; Bersohn, M.
- [100304] M.; Burke, E. P.; Lifton, R. P.; Philipson, K. D.: Cloning of the
- [100305] NCX2 isoform of the plasma membrane Na(+)-Ca(2+) exchanger. J. Biol.
- [100306] Chem. 269: 17434-17439, 1994.
- [100307] 17447.
- [100308] Detter, J. C.; Zhang, Q.; Mules, E. H.; Novak, E. K.; Mishra, V.
- [100309] S.; Li, W.; McMurtrie, E. B.; Tchernev, V. T.; Wallace, M. R.; Seabra,
- [100310] M. C.; Swank, R. T.; Kingsmore, S. F.: Rab geranylgeranyl transferase
- [100311] alpha mutation in the gunmetal mouse reduces Rab prenylation and platelet
- [100312] synthesis. Proc. Nat. Acad. Sci. 97: 4144-4149, 2000.
- [100313] 17448.
- [100314] Novak, E. K.; Reddington, M.; Zhen, L.; Stenberg, P. E.; Jackson,
- [100315] C. W.; McGarry, M. P.; Swank, R. T. :Blood 85: 1781-1789, 1995.

[100316]17449.

[100317]Swank, R. T.; Jiang, S. Y.; Reddington, M.; Conway, J.;
Stephenson,

[100318]D.; McGarry, M. P.; Novak, E. K. :Blood 81: 2626–2635,
1993.

[100319]17450.

[100320]Bui, T. D.; Rankin, J.; Smith, K.; Huguet, E. L.; Ruben, S.;
Strachan,

[100321]T.; Harris, A. L.; Lindsay, S.: A novel human Wnt gene,
WNT10B, maps

[100322]to 12q13 and is expressed in human breast carcinomas.
Oncogene 14:

[100323]1249–1253, 1997.

[100324]17451.

[100325]Hardiman, G.; Kastelein, R. A.; Bazan, J. F.: Isolation, char-
acterization

[100326]and chromosomal localization of human WNT10B. Cyto-
genet. Cell Genet. 77:

[100327]278–282, 1997.

[100328]17452.

[100329]Ross, S. E.; Hemati, N.; Longo, K. A.; Bennett, C. N.; Lucas,

P.

[100330]C.; Erickson, R. L.; MacDougald, O. A.: Inhibition of adipogenesis

[100331]by Wnt signaling. Science 289: 950–953, 2000.

[100332]17453.

[100333]Meyerhardt, J. A.; Look, A. T.; Bigner, S. H.; Fearon, E. R.:

[100334]Identification and characterization of neogenin, a DCC-related gene. Oncogene 14:

[100335]1129–1136, 1997.

[100336]17454.

[100337]Vielmetter, J.; Chen, X.-N.; Miskevich, F.; Lane, R. P.; Yamakawa,

[100338]K.; Korenberg, J. R.; Dreyer, W. J.: Molecular characterization of

[100339]human neogenin, a DCC-related protein, and the mapping of its gene

[100340](NEO1) to chromosomal position 15q22.3–q23. Genomics 41: 414–421,

[100341]1997.

[100342]17455.

[100343]Vielmetter, J.; Kayyem, J. F.; Roman, J. M.; Dreyer, W. J.:

Neogenin,

[100344]an avian cell surface protein expressed during terminal neuronal differentiation,

[100345]is closely related to the human tumor suppressor molecule deleted

[100346]in colorectal cancer. *J. Cell Biol.* 127: 2009–2020, 1994.

[100347]17456.

[100348]Rosenquist, T. A.; Zharkov, D. O.; Grollman, A. P.: Cloning and

[100349]characterization of a mammalian 8-oxoguanine DNA glycosylase. *Proc.*

[100350]*Nat. Acad. Sci.* 94: 7429–7434, 1997.

[100351]17457.

[100352]Tani, M.; Shinmura, K.; Kohno, T.; Shiroishi, T.; Wakama, S.;

[100353]Kim, S.-R.; Nohmi, T.; Kasai, H.; Takenoshita, S.; Nagamachi, Y.;

[100354]Yokota, J.: Genomic structure and chromosomal localization of the

[100355]mouse Ogg1 gene that is involved in the repair of 8-hydroxyguanine

[100356]in DNA damage. *Mammalian Genome* 9: 32–37, 1998.

[100357]17458.

[100358]Lim, H. N.; Hawkins, J. R.; Hughes, I. A.: Genetic evidence
to

[100359]exclude the androgen receptor co-factor, ARA70 (NCOA4)
as a candidate

[100360]gene for the causation of undermasculinised genitalia.
(Letter) Clin.

[100361]Genet. 59: 284–286, 2001.

[100362]17459.

[100363]Santoro, M.; Dathan, N. A.; Berlingieri, M. T.; Bongarzone,
I.;

[100364]Paulin, C.; Grieco, M.; Pierotti, M. A.; Vecchio, G.; Fusco,
A.:

[100365]Molecular characterization of RET/PTC3: a novel rear-
ranged version

[100366]of the RET proto-oncogene in a human thyroid papillary
carcinoma. Oncogene 9:

[100367]509–516, 1994.

[100368]17460.

[100369]Yeh, S.; Chang, C.: Cloning and characterization of a spe-
cific

[100370]coactivator, ARA-70, for the androgen receptor in human prostate cells. Proc.

[100371]Nat. Acad. Sci. 93: 5517-5521, 1996.

[100372]17461.

[100373]Huttner, K. M.; Kozak, C. A.; Bevins, C. L.: The mouse genome

[100374]encodes a single homolog of the antimicrobial peptide human beta-defensin

[100375]1. FEBS Lett. 413: 45-49, 1997.

[100376]17462.

[100377]Jia, H. P.; Starner, T.; Ackermann, M.; Kirby, P.; Tack, B. F.;

[100378]McCray, P. B., Jr.: Abundant human beta-defensin-1 expression in

[100379]milk and mammary gland epithelium. J. Pediat. 138: 109-112, 2001.

[100380]17463.

[100381]Morrison, G. M.; Davidson, D. J.; Kilanowski, F. M.; Borthwick,

[100382]D. W.; Crook, K.; Maxwell, A. I.; Govan, J. R. W.; Dorin, J. R.:

[100383]Mouse beta defensin-1 is a functional homolog of human

beta defensin-1. Mammalian

[100384]Genome 9: 453-457, 1998.

[100385]17464.

[100386]Singh, P. K.; Jia, H. P.; Wiles, K.; Hesselberth, J.; Liu, L.;

[100387]Conway, B.-A. D.; Greenberg, E. P.; Valore, E. V.; Welsh, M.

J.; Ganz,

[100388]T.; Tack, B. F.; McCray, P. B., Jr.: Production of beta-

defensins

[100389]by human airway epithelia. Proc. Nat. Acad. Sci. 95:

14961-14966,

[100390]1998.

[100391]17465.

[100392]Tunzi, C. R.; Harper, P. A.; Bar-Oz, B.; Valore, E. V.; Sem-

ple,

[100393]J. L.; Watson-MacDonell, J.; Ganz, T.; Ito, S.: Beta-defensin

expression

[100394]in human mammary gland epithelia. Pediat. Res. 48:

30-35, 2000.

[100395]17466.

[100396]Valore, E. V.; Park, C. H.; Quayle, A. J.; Wiles, K. R.; Mc-

Cray,

[100397]P. B., Jr.; Ganz, T.: Human beta-defensin-1: an antimicrobial peptide

[100398]of urogenital tissues. J. Clin. Invest. 101: 1633-1642, 1998.

[100399]17467.

[100400]Nagasawa, A.; Kubota, R.; Imamura, Y.; Nagamine, K.; Wang, Y.;

[100401]Asakawa, S.; Kudoh, J.; Minoshima, S.; Mashima, Y.; Oguchi, Y.; Shimizu,

[100402]N.: Cloning of the cDNA for a new member of the immunoglobulin superfamily

[100403](ISLR) containing leucine-rich repeat (LRR). Genomics 44: 273-279,

[100404]1997.

[100405]17468.

[100406]Paoloni-Giacobino, A.; Chen, H.; Peitsch, M. C.; Rossier, C.; Antonarakis,

[100407]S. E.: Cloning of the TMPRSS2 gene, which encodes a novel serine

[100408]protease with transmembrane, LDLRA, and SRCR domains and maps to 21q22.3. Genomics 44:

[100409]309-320, 1997. Note: Erratum: Genomics 44: 309-320,

1997.

[100410]17469.

[100411]McAllister, G.; Whiting, P.; Hammond, E. A.; Knowles, M.
R.; Atack,

[100412]J. R.; Bailey, F. J.; Maigetter, R.; Ragan, C. I.: cDNA cloning
of

[100413]human and rat brain myo-inositol monophosphatase: ex-
pression and characterization

[100414]of the human recombinant enzyme. Biochem. J. 284:
749-754, 1992.

[100415]17470.

[100416]Sjoholt, G.; Molven, A.; Lovlie, R.; Wilcox, A.; Sikela, J. M.;

[100417]Steen, V. M.: Genomic structure and chromosomal local-
ization of a

[100418]human myo-inositol monophosphatase gene (IMPA). Ge-
nomics 45: 113-122,

[100419]1997.

[100420]17471.

[100421]Steen, V. M.; Gulbrandsen, A. K.; Eiken, H. G.; Berle, J. O.:

[100422]Lack of genetic variation in the coding region of the myo-
inositol

[100423]monophosphatase gene in lithium-treated patients with
manic depressive

[100424]illness. Pharmacogenetics 6: 113–116, 1996.

[100425]17472.

[100426]McDonald, M. T.; Flejter, W.; Sheldon, S.; Putzi, M. J.;
Gorski,

[100427]J. L.: XY sex reversal and gonadal dysgenesis due to 9p24
monosomy. Am.

[100428]J. Med. Genet. 73: 321–326, 1997.

[100429]17473.

[100430]Muroya, K.; Okuyama, T.; Goishi, K.; Ogiso, Y.; Fukuda, S.;
Kameyama,

[100431]J.; Sato, H.; Suzuki, Y.; Terasaki, H.; Gomyo, H.; Wakui, K.;
Fukushima,

[100432]Y.; Ogata, T.: Sex-determining gene(s) on distal 9p: clinical and

[100433]molecular studies in six cases. J. Clin. Endocr. Metab. 85:
3094–3100,

[100434]2000.

[100435]17474.

[100436]Raymond, C. S.; Shamu, C. E.; Shen, M. M.; Seifert, K. J.;

Hirsch,

[100437]B.; Hodgkin, J.; Zarkower, D.: Evidence for evolutionary conservation

[100438]of sex-determining genes. Nature 391: 691-695, 1998.

[100439]17475.

[100440]Shan, Z.; Zabel, B.; Trautmann, U.; Hillig, U.; Ottolenghi, C.;

[100441]Wang, Y.; Haaf, T.: FISH mapping of the sex-reversal region on human

[100442]chromosome 9p in two XY females and in primates. Europ. J. Hum. Genet. 8:

[100443]167-173, 2000.

[100444]17476.

[100445]Shen, M. M.; Hodgkin, J.: mab-3, a gene required for sex-specific

[100446]yolk protein expression and a male-specific lineage in C. elegans. Cell 54:

[100447]1019-1031, 1988.

[100448]17477.

[100449]Smith, C. A.; McClive, P. J.; Western, P. S.; Reed, K. J.; Sinclair,

- [100450]A. H.: Conservation of a sex-determining gene. (Letter)
Nature 402:
[100451]601-602, 1999.
- [100452]17478.
- [100453]Veitia, R.; Nunes, M.; Brauner, R.; Doco-Fenzy, M.; Joanny-
Flinois,
[100454]O.; Jaubert, F.; Lortat-Jacob, S.; Fellous, M.; McElreavey,
K.: Deletions
[100455]of distal 9p associated with 46,XY male to female sex re-
versal: definition
[100456]of the breakpoints at 9p23.3-p24.1. Genomics 41:
271-274, 1997.
- [100457]17479.
- [100458]Takekawa, M.; Posas, F.; Saito, H.: A human homolog of
the yeast
[100459]Ssk2/Ssk22 MAP kinase kinase kinases, MTK1, mediates
stress-induced
[100460]activation of the p38 and JNK pathways. EMBO J. 16:
4973-4982, 1997.
- [100461]17480.
- [100462]Chapman, D. L.; Agulnik, I.; Hancock, S.; Silver, L. M.; Pa-
paioannou,

- [100463]V. E.: Tbx6, a mouse T-box gene implicated in paraxial mesoderm formation at gastrulation. Dev. Biol. 180: 534–542, 1996.
- [100465]17481.
- [100466]Chapman, D. L.; Papaioannou, V. E.: Three neural tubes in mouse embryos with mutations in the T-box gene Tbx6. Nature 391: 695–697, 1998.
- [100468]17482.
- [100469]17482.
- [100470]Papapetrou, C.; Putt, W.; Fox, M.; Edwards, Y. H.: The human TBX6 gene: cloning and assignment to chromosome 16p11.2. Genomics 55: 238–241, 1999.
- [100472]238–241, 1999.
- [100473]17483.
- [100474]Spicer, A. P.; Olson, J. S.; McDonald, J. A.: Molecular cloning and characterization of a cDNA encoding the third putative mammalian hyaluronan synthase. J. Biol. Chem. 272: 8957–8961,

1997.

[100477]17484.

[100478]Bagri, A.; Marin, O.; Plump, A. S.; Mak, J.; Pleasure, S. J.;
Rubenstein,

[100479]J. L. R.; Tessier-Lavigne, M.: Slit proteins prevent midline
crossing

[100480]and determine the dorsoventral position of major axonal
pathways in

[100481]the mammalian forebrain. Neuron 33: 233–248, 2002.

[100482]17485.

[100483]Dallol, A.; Forgacs, E.; Martinez, A.; Sekido, Y.; Walker, R.;

[100484]Kishida, T.; Rabbitts, P.; Maher, E. R.; Minna, J. D.; Latif, F.:

[100485]Tumour specific promoter region methylation of the hu-
man homologue

[100486]of the Drosophila Roundabout gene DUTT1 (ROBO1) in
human cancers. Oncogene 21:

[100487]3020–3028, 2002.

[100488]17486.

[100489]Kidd, T.; Brose, K.; Mitchell, K. J.; Fetter, R. D.; Tessier-
Lavigne,

[100490]M.; Goodman, C. S.; Tear, G.: Roundabout controls axon

crossing of

[100491]the CNS midline and defines a novel subfamily of evolutionarily conserved

[100492]guidance receptors. Cell 92: 205–215, 1998.

[100493]17487.

[100494]Sundaresan, V.; Chung, G.; Heppell–Parton, A.; Xiong, J.; Grundy,

[100495]C.; Roberts, I.; James, L.; Cahn, A.; Bench, A.; Douglas, J.; Minna,

[100496]J.; Sekido, Y.; Lerman, M.; Latif, F.; Bergh, J.; Li, H.; Lowe, N.;

[100497]Ogilvie, D.; Rabbitts, P.: Homozygous deletions at 3p12 in breast

[100498]and lung cancer. Oncogene 17: 1723–1729, 1998.

[100499]17488.

[100500]Zallen, J. A.; Yi, B. A.; Bargmann, C. I.: The conserved immunoglobulin

[100501]superfamily member SAX–3/Robo directs multiple aspects of axon guidance

[100502]in C. elegans. Cell 92: 217–227, 1998.

[100503]17489.

[100504]Faber, P. W.; Barnes, G. T.; Srinidhi, J.; Chen, J.; Gusella, J.
[100505]F.; MacDonald, M. E.: Huntingtin interacts with a family of
WW domain
[100506]proteins. Hum. Molec. Genet. 7: 1463–1474, 1998.
[100507]17490.
[100508]Hattula, K.; Peranen, J.: FIP-2, a coiled-coil protein, links
[100509]huntingtin to Rab8 and modulates cellular morphogene-
sis. Curr. Biol. 10:
[100510]1603–1606, 2000.
[100511]17491.
[100512]Li, Y.; Kang, J.; Horwitz, M. S.: Interaction of an adenovirus
[100513]E3 14.7-kilodalton protein with a novel tumor necrosis
factor alpha-inducible
[100514]cellular protein containing leucine zipper domains. Molec.
Cell.
[100515]Biol. 18: 1601–1610, 1998.
[100516]17492.
[100517]Fischer, U.; Heckel, D.; Michel, A.; Janka, M.; Hulsebos, T.;
Meese,
[100518]E.: Cloning of a novel transcription factor-like gene am-

plified in

[100519]human glioma including astrocytoma grade I. Hum. Molec.

Genet. 6:

[100520]1817–1822, 1997.

[100521]17493.

[100522]Gracia, E.; Fischer, U.; Elkahloun, A.; Trent, J. M.; Meese,

E.;

[100523]Meltzer, P. S.: Isolation of genes amplified in human can-

cers by

[100524]microdissection mediated cDNA capture. Hum. Molec.

Genet. 5: 595–600,

[100525]1996.

[100526]17494.

[100527]Gerard, M.; Hernandez, L.; Wevrick, R.; Stewart, C. L.: Dis-

ruption

[100528]of the mouse necdin gene results in early post-natal

lethality. Nature

[100529]Genet. 23: 199–202, 1999.

[100530]17495.

[100531]Hurst, L. D.; McVean, G.; Moore, T.: Imprinted genes have

few

[100532]and small introns. (Letter) Nature Genet. 12: 234–237,
1996.

[100533]17496.

[100534]Jay, P.; Rougeulle, C.; Massacrier, A.; Moncla, A.; Mattei,
M.–G.;

[100535]Malzac, P.; Roeckel, N.; Taviaux, S.; Lefranc, J.–L. B.; Cau,
P.;

[100536]Berta, P.; Lalande, M.; Muscatelli, F.: The human necdin
gene, NDN,

[100537]is maternally imprinted and located in the Prader–Willi
syndrome chromosomal

[100538]region. Nature Genet. 17: 357–360, 1997.

[100539]17497.

[100540]MacDonald, H. R.; Wevrick, R.: The necdin gene is deleted
in Prader–Willi

[100541]syndrome and is imprinted in human and mouse. Hum.
Molec. Genet. 6:

[100542]1873–1878, 1997.

[100543]17498.

[100544]Maruyama, K.; Usami, M.; Aizawa, T.; Yoshikawa, K.: A
novel brain–specific

[100545] mRNA encoding nuclear protein (necdin) expressed in neurally differentiated
[100546] embryonal carcinoma cells. Biochem. Biophys. Res. Comm. 178: 291-296,
[100547] 1991.
[100548] 17499.
[100549] Muscatelli, F.; Arous, D. N.; Massacrier, A.; Boccaccio, I.; Le
[100550] Moal, M.; Cau, P.; Cremer, H.: Disruption of the mouse Necdin gene
[100551] results in hypothalamic and behavioral alterations reminiscent of
[100552] the human Prader-Willi syndrome. Hum. Molec. Genet. 9: 3101-3110,
[100553] 2000.
[100554] 17500.
[100555] Nakada, Y.; Taniura, H.; Uetsuki, T.; Inazawa, J.; Yoshikawa, K.
[100556]: The human chromosomal gene for necdin, a neuronal growth suppressor,
[100557] in the Prader-Willi syndrome deletion region. Gene 213: 65-72, 1998.

[100558]17501.

[100559]Nicholls, R. D.: Incriminating gene suspects, Prader-Willi style. Nature

[100560]Genet. 23: 132-134, 1999.

[100561]17502.

[100562]Tsai, T.-F.; Armstrong, D.; Beaudet, A. L.: Necdin-deficient mice

[100563]do not show lethality or the obesity and infertility of Prader-Willi

[100564]syndrome. (Letter) Nature Genet. 22: 15-16, 1999.

[100565]17503.

[100566]Watrin, F.; Roeckel, N.; Lacroix, L.; Mignon, C.; Mattei, M.-G.;

[100567]Disteche, C.; Muscatelli, F.: The mouse necdin gene is expressed

[100568]from the paternal allele only and lies in the 7C region of the mouse

[100569]chromosome 7, a region of conserved synteny to the human Prader-Willi

[100570]syndrome region. Europ. J. Hum. Genet. 5: 324-332, 1997.

[100571]17504.

[100572]Woodage, T.; Basrai, M. A.; Baxevanis, A. D.; Hieter, P.;
Collins,

[100573]F. S.: Characterization of the CHD family of proteins. Proc.
Nat.

[100574]Acad. Sci. 94: 11472–11477, 1997.

[100575]17505.

[100576]Seelig, H. P.; Renz, M.; Targoff, I. N.; Ge, Q.; Frank, M. B.:

[100577]Two forms of the major antigenic protein of the dermato-
myositis-specific

[100578]Mi-2 autoantigen. (Letter) Arthritis Rheum. 39:
1769–1771, 1996.

[100579]17506.

[100580]Ernstsson, S.; Betz, R.; Lagercrantz, S.; Larsson, C.; Ericks-
son,

[100581]S.; Cederberg, A.; Carlsson, P.; Enerback, S.: Cloning and
characterization

[100582]of freac-9 (FKHL17), a novel kidney-expressed human
forkhead gene

[100583]that maps to chromosome 1p32–p34. Genomics 46:
78–85, 1997.

[100584]17507.

[100585]Ozkaynak, E.; Schnegelsberg, P. N. J.; Jin, D. F.; Clifford, G.

[100586]M.; Warren, F. D.; Drier, E. A.; Oppermann, H.: Osteogenic
protein-2:

[100587]a new member of the transforming growth factor-beta
superfamily expressed

[100588]in early embryogenesis. J. Biol. Chem. 267: 25220-25227,
1992.

[100589]17508.

[100590]Kinoh, H.; Hayashita, H.; Kajita, M.; Okada, A.; Seiki, M.:
Assignment

[100591]of the genes for membrane-type-4 matrix metallopro-
teinase (Mmp17,

[100592]MMP17) to mouse chromosome 5, human chromosome
band 12q24.3 and membrane-type-5

[100593]matrix metalloproteinase (Mmp24, MMP24) to mouse
chromosome 2 and

[100594]human chromosome band 20q11.2-q12, respectively, by
radiation hybrid

[100595]and in situ hybridization. Cytogenet. Cell Genet. 87:
97-98, 1999.

[100596]17509.

[100597]Bingle, C. D.; Gowan, S.: Molecular cloning of the forkhead transcription

[100598]factor HNF-3- α from a human pulmonary adenocarcinoma cell line. *Biochim.*

[100599]*Biophys. Acta* 1307: 17-20, 1996.

[100600]17510.

[100601]Hannigan, G. E.; Bayani, J.; Weksberg, R.; Beatty, B.; Pandita,

[100602]A.; Dedhar, S.; Squire, J.: Mapping of the gene encoding the integrin-linked

[100603]kinase, ILK, to human chromosome 11p15.5-p15.4. *Genomics* 42: 177-179,

[100604]1997.

[100605]17511.

[100606]Hannigan, G. E.; Leung-Hagesteijn, C.; Fitz-Gibbon, L.; Coppolino,

[100607]M. G.; Radeva, G.; Filmus, J.; Bell, J. C.; Dedhar, S.: Regulation

[100608]of cell adhesion and anchorage-dependent growth by a new beta-1-integrin-linked

[100609]protein kinase. *Nature* 379: 91-96, 1996.

[100610]17512.

[100611]Omeis, I. A.; Hsu, Y.-C.; Perin, M. S.: Mouse and human neuronal

[100612]pentraxin 1 (NPXT1): conservation, genomic structure, and chromosomal

[100613]localization. Genomics 36: 543–545, 1996.

[100614]17513.

[100615]Kubota, N.; Terauchi, Y.; Miki, H.; Tamemoto, H.; Yamauchi, T.;

[100616]Komeda, K.; Satoh, S.; Nakano, R.; Ishii, C.; Sugiyama, T.; Eto, K.;

[100617]Tsubamoto, Y.; and 17 others: PPAR- γ mediates high-fat diet-induced

[100618]adipocyte hypertrophy and insulin resistance. Molec. Cell 4: 597–609,

[100619]1999.

[100620]17514.

[100621]Lapsys, N. M.; Kriketos, A. D.; Lim-Fraser, M.; Poynten, A. M.;

[100622]Lowy, A.; Furler, S. M.; Chisholm, D. J.; Cooney, G. J.: Expression

[100623] of genes involved in lipid metabolism correlate with peroxisome proliferator-activated

[100624] receptor gamma expression in human skeletal muscle. J. Clin. Endocr.

[100625] Metab. 85: 4293–4297, 2000.

[100626] 17515.

[100627] Lehmann, J. M.; Moore, L. B.; Smith–Oliver, T. A.; Wilkison, W.

[100628] O.; Willson, T. M.; Kliewer, S. A.: An antidiabetic thiazolidinedione

[100629] is a high affinity ligand for peroxisome proliferator-activated receptor

[100630] gamma (PPAR gamma). J. Biol. Chem. 270: 12953–12956, 1995.

[100631] 17516.

[100632] Lowell, B. B.: PPAR-gamma: an essential regulator of adipogenesis

[100633] and modulator of fat cell function. Cell 99: 239–242, 1999.

[100634] 17517.

[100635] Martin, G.; Schoonjans, K.; Staels, B.; Auwerx, J.: PPAR-gamma

- [100636]activators improve glucose homeostasis by stimulating fatty acid uptake
- [100637]in the adipocytes. *Atherosclerosis* 137: S75–S80, 1998.
- [100638]17518.
- [100639]Meirhaeghe, A.; Fajas, L.; Helbecque, N.; Cottel, D.; Lebel, P.;
- [100640]Dallongeville, J.; Deeb, S.; Auwerx, J.; Amouyel, P.: A genetic polymorphism
- [100641]of the peroxisome proliferator–activated receptor gamma gene influences
- [100642]plasma leptin levels in obese tumors. *Hum. Molec. Genet.* 7: 435–440,
- [100643]1998.
- [100644]17519.
- [100645]Miles, P. D. G.; Barak, Y.; He, W.; Evans, R. M.; Olefsky, J.
- [100646]M.: Improved insulin–sensitivity in mice heterozygous for PPAR–gamma
- [100647]deficiency. *J. Clin. Invest.* 105: 287–292, 2000.
- [100648]17520.
- [100649]Mueller, E.; Sarraf, P.; Tontonoz, P.; Evans, R. M.; Martin, K.

[100650]J.; Zhang, M.; Fletcher, C.; Singer, S.; Spiegelman, B. M.:

Terminal

[100651]differentiation of human breast cancer through PPAR-
gamma. Molec.

[100652]Cell. 1: 465-470, 1998.

[100653]17521.

[100654]Mueller, E.; Smith, M.; Sarraf, P.; Kroll, T.; Aiyer, A.; Kauf-
man,

[100655]D. S.; Oh, W.; Demetri, G.; Figg, W. D.; Zhou, X.-P.; Eng,
C.; Spiegelman,

[100656]B. M.; Kantoff, P. W.: Effects of ligand activation of peroxi-
some

[100657]proliferator-activated receptor gamma in human prostate
cancer. Proc.

[100658]Nat. Acad. Sci. 97: 10990-10995, 2000.

[100659]17522.

[100660]Nagy, L.; Tontonoz, P.; Alvarez, J. G. A.; Chen, H.; Evans,
R.

[100661]M.: Oxidized LDL regulates macrophage gene expression
through ligand

[100662]activation of PPAR-gamma. Cell 93: 229-240, 1998.

[100663]17523.

[100664]Oh, E. Y.; Min, K. M.; Chung, J. H.; Min, Y.-K.; Lee, M.-S.;
Kim,

[100665]K.-W.; Lee, M.-K.: Significance of pro12ala mutation in
peroxisome

[100666]proliferator-activated receptor-gamma2 in Korean dia-
betic and obese

[100667]subjects. J. Clin. Endocr. Metab. 85: 1801-1804, 2000.

[100668]17524.

[100669]Pritchard, J. K.; Rosenberg, N. A.: Use of unlinked genetic
markers

[100670]to detect population stratification in association studies.
Am. J.

[100671]Hum. Genet. 65: 220-228, 1999.

[100672]17525.

[100673]Ristow, M.; Muller-Wieland, D.; Pfeiffer, A.; Krone, W.;
Kahn,

[100674]C. R.: Obesity associated with a mutation in a genetic reg-
ulator

[100675]of adipocyte differentiation. New Eng. J. Med. 339:
953-959, 1998.

[100676]17526.

[100677]Rocchi, S.; Picard, F.; Vamecq, J.; Gelman, L.; Potier, N.; Zeyer,

[100678]D.; Dubuquoy, L.; Bac, P.; Champy, M.-F.; Plunket, K. D.; Leesnitzer,

[100679]L. M.; Blanchard, S. G.; Desreumaux, P.; Moras, D.; Renaud, J.-P.;

[100680]Auwerx, J.: A unique PPAR- γ ligand with potent insulin-sensitizing

[100681]yet weak adipogenic activity. *Molec. Cell* 8: 737-747, 2001.

[100682]17527.

[100683]Rosen, E. D.; Sarraf, P.; Troy, A. E.; Bradwin, G.; Moore, K.;

[100684]Milstone, D. S.; Spiegelman, B. M.; Mortensen, R. M.: PPAR- γ

[100685]is required for the differentiation of adipose tissue in vivo and

[100686]in vitro. *Molec. Cell* 4: 611-617, 1999.

[100687]17528.

[100688]Ricote, M.; Huang, J.; Fajas, L.; Li, A.; Welch, J.; Najib, J.;

[100689]Witztum, J. L.; Auwerx, J.; Palinski, W.; Glass, C. K.: Ex-

pression

[100690]of the peroxisome proliferator-activated receptor gamma
(PPAR-gamma)

[100691]in human atherosclerosis and regulation in macrophages
by colony stimulating

[100692]factors and oxidized low density lipoprotein. Proc. Nat.
Acad. Sci. 95:

[100693]7614-7619, 1998.

[100694]17529.

[100695]Savage, D. B.; Agostini, M.; Barroso, I.; Gurnell, M.; Luan,
J.;

[100696]Meirhaeghe, A.; Harding, A.-H.; Ihrke, G.; Rajanayagam,
O.; Soos,

[100697]M. A.; George, S.; Berger, D.; and 9 others: Digenic inheri-
tance

[100698]of severe insulin resistance in a human pedigree. Nature
Genet. 31:

[100699]379-384, 2002. Note: Erratum: Nature Genet. 32: 211
only, 2002.

[100700]17530.

[100701]Smith, W. M.; Zhou, X.-P.; Kurose, K.; Gao, X.; Latif, F.;
Kroll,

[100702]T.; Sugano, K.; Cannistra, S. A.; Clinton, S. K.; Maher, E. R.;
Prior,
[100703]T. W.; Eng, C.: Opposite association of two PPARG variants
with cancer:
[100704]overrepresentation of H449H in endometrial carcinoma
cases and underrepresentation
[100705]of P12A in renal cell carcinoma cases. Hum. Genet. 109:
146–151,
[100706]2001.
[100707]17531.
[100708]Tarrade, A.; Schoonjans, K.; Pavan, L.; Auwerx, J.; Ro-
chette-Egly,
[100709]C.; Evain-Brion, D.; Fournier, T.: PPAR-gamma/RXR-alpha
heterodimers
[100710]control human trophoblast invasion. J. Clin. Endocr.
Metab. 86:
[100711]5017–5024, 2001.
[100712]17532.
[100713]Tontonoz, P.; Hu, E.; Devine, J.; Beale, E. G.; Spiegelman,
B.
[100714]M.: PPAR gamma 2 regulates adipose expression of the
phosphoenolpyruvate

[100715]carboxykinase gene. Molec. Cell. Biol. 15: 351–357, 1995.

[100716]17533.

[100717]Tontonoz, P.; Hu, E.; Graves, R. A.; Budavari, A. I.; Spiegelman,

[100718]B. M.: mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. Genes

[100719]Dev. 8: 1224–1234, 1994.

[100720]17534.

[100721]Tontonoz, P.; Nagy, L.; Alvarez, J. G. A.; Thomazy, V. A.; Evans,

[100722]R. M.: PPAR-gamma promotes monocyte/macrophage differentiation and

[100723]uptake of oxidized LDL. Cell 93: 241–252, 1998.

[100724]17535.

[100725]Wang, M.: Isolation and characterization of a human gene encoding

[100726]a single-strand-specific endoribonuclease. Ph.D. Thesis: Stanford

[100727]Univ. , 1995.

[100728]17536.

- [100729]Wang, M.; Cohen, S. N.: ard-1: a human gene that re-
verses the
- [100730]effects of temperature-sensitive and deletion mutations in
the Escherichia
- [100731]coli rne gene and encodes an activity producing RNase E-
like cleavages. Proc.
- [100732]Nat. Acad. Sci. 91: 10591-10595, 1994.
- [100733]17537.
- [100734]Amendt, B. A.; Sutherland, L. B.; Semina, E. V.; Russo, A.
F.:
- [100735]The molecular basis of Rieger syndrome: analysis of PITX2
homeodomain
- [100736]protein activities. J. Biol. Chem. 273: 20066-20072, 1998.
- [100737]17538.
- [100738]Arakawa, H.; Nakamura, T.; Zhadanov, A. B.; Fidanza, V.;
Yano,
- [100739]T.; Bullrich, F.; Shimizu, M.; Blechman, J.; Mazo, A.;
Canaani, E.;
- [100740]Croce, C. M.: Identification and characterization of the
ARP1 gene,
- [100741]a target for the human acute leukemia ALL1 gene. Proc.
Nat. Acad.

[100742]Sci. 95: 4573–4578, 1998.

[100743]17539.

[100744]Cushman, L. J.; Camper, S. A.: Molecular basis of pituitary dysfunction

[100745]in mouse and human. Mammalian Genome 12: 485–494, 2001.

[100746]17540.

[100747]Doward, W.; Perveen, R.; Lloyd, I. C.; Ridgway, A. E. A.; Wilson,

[100748]L.; Black, G. C. M.: A mutation in the RIEG1 gene associated with

[100749]Peters' anomaly. J. Med. Genet. 36: 152–155, 1999.

[100750]17541.

[100751]Flomen, R. H.; Vatcheva, R.; Gorman, P. A.; Baptista, P. R.; Groet,

[100752]J.; Barisic, I.; Ligutic, I.; Nizetic, D.: Construction and analysis

[100753]of a sequence-ready map in 4q25: Rieger syndrome can be caused by

[100754]haploinsufficiency of RIEG, but also by chromosome breaks approximately

[100755]90 kb upstream of this gene. Genomics 47: 409–413,
1998.

[100756]17542.

[100757]Gage, P. J.; Camper, S. A.: Pituitary homeobox 2, a novel
member

[100758]of the bicoid-related family of homeobox genes, is a po-
tential regulator

[100759]of anterior structure formation. Hum. Molec. Genet. 6:
457–464,

[100760]1997.

[100761]17543.

[100762]Kozak, M.: Interpreting cDNA sequences: some insights
from studies

[100763]on translation. Mammalian Genome 7: 563–574, 1996.

[100764]17544.

[100765]Kozlowski, K.; Walter, M. A.: Variation in residual PITX2
activity

[100766]underlies the phenotypic spectrum of anterior segment
developmental

[100767]disorders. Hum. Molec. Genet. 9: 2131–2139, 2000.

[100768]17545.

[100769]Lin, C. R.; Kioussi, C.; O'Connell, S.; Briata, P.; Szeto, D.;

[100770]Liu, F.; Izpisua-Belmonte, J. C.; Rosenfeld, M. G.: Pitx2
regulates

[100771]lung asymmetry, cardiac positioning and pituitary and
tooth morphogenesis. Nature 401:

[100772]279-282, 1999.

[100773]17546.

[100774]Logan, M.; Pagan-Westphal, S. M.; Smith, D. M.; Paganessi,
L.;

[100775]Tabin, C. J.: The transcription factor Pitx2 mediates situs-
specific

[100776]morphogenesis in response to left-right asymmetric sig-
nals. Cell 94:

[100777]307-317, 1998.

[100778]17547.

[100779]Lu, M.-F.; Pressman, C.; Dyer, R.; Johnson, R. L.; Martin, J.

[100780]F.: Function of Rieger syndrome gene in left-right asym-
metry and

[100781]craniofacial development. Nature 401: 276-278, 1999.

[100782]17548.

[100783]Martin, D. M.; Probst, F. J.; Fox, S. E.; Schimmenti, L. A.;
Semina,

[100784]E. V.; Hefner, M. A.; Belmont, J. W.; Camper, S. A.: Exclu-
sion of

[100785]PITX2 mutations as a major cause of CHARGE association.
Am. J. Med.

[100786]Genet. 111: 27–30, 2002.

[100787]17549.

[100788]Pellegrini-Bouiller, I.; Manrique, C.; Gunz, G.; Grino, M.;
Zamora,

[100789]A. J.; Figarella-Branger, D.; Grisoli, F.; Jaquet, P.; Enjalbert,

[100790]A.: Expression of the members of the Ptx family of tran-
scription

[100791]factors in human pituitary adenomas. J. Clin. Endocr.
Metab. 84:

[100792]2212–2220, 1999.

[100793]17550.

[100794]Piedra, M. E.; Icardo, J. M.; Albajar, M.; Rodriguez-Rey, J.
C.;

[100795]Ros, M. A.: Pitx2 participates in the late phase of the
pathway controlling

[100796]left-right asymmetry. Cell 94: 319–324, 1998.

[100797]17551.

[100798]Priston, M.; Kozlowski, K.; Gill, D.; Letwin, K.; Buys, Y.;
Levin,

[100799]A. V.; Walter, M. A.; Heon, E.: Functional analyses of two
newly

[100800]identified PITX2 mutants reveal a novel molecular mecha-
nism for Axenfeld-Rieger

[100801]syndrome. Hum. Molec. Genet. 10: 1631–1638, 2001.

[100802]17552.

[100803]Bayever, E.; Ladisch, S.; Philippart, M.; Brill, N.; Nuwer, M.;

[100804]Sparkes, R. S.; Feig, S. A.: Bone-marrow transplantation
for metachromatic

[100805]leucodystrophy. Lancet II: 471–473, 1985.

[100806]17553.

[100807]Beratis, N. G.; Danesino, C.; Hirschhorn, K.: Detection of
homozygotes

[100808]and heterozygotes for metachromatic leukodystrophy in
lymphoid cell

[100809]lines and peripheral leukocytes. Ann. Hum. Genet. 38:
485–493, 1975.

[100810]17554.

[100811]Berger, J.; Gmach, M.; Mayr, U.; Molzer, B.; Bernheimer, H.:

[100812]Coincidence of two novel arylsulfatase A alleles and mutation 459+1G→A

[100813]within a family with metachromatic leukodystrophy: molecular basis

[100814]of phenotypic heterogeneity. Hum. Mutat. 13: 61–68, 1999.

[100815]17555.

[100816]Betts, T. A.; Smith, W. T.; Kelly, R. E.: Adult metachromatic

[100817]leukodystrophy (sulphatide lipidosis) simulating acute schizophrenia:

[100818]report of a case. Neurology 18: 1140–1142, 1968.

[100819]17556.

[100820]Black, J. W.; Cumings, J. N.: Infantile metachromatic leukodystrophy. J.

[100821]Neurol. Neurosurg. Psychiat. 24: 233–239, 1961.

[100822]17557.

[100823]Bohne, W.; von Figura, K.; Gieselmann, V.: An 11-bp deletion

[100824]in the arylsulfatase A gene of a patient with late infantile metachromatic

[100825]leukodystrophy. Hum. Genet. 87: 155–158, 1991.

[100826]17558.

[100827]Bosch, E. P.; Hart, M. N.: Late adult-onset metachromatic leukodystrophy:

[100828]dementia and polyneuropathy in a 63-year-old man. Arch. Neurol. 35:

[100829]475–477, 1978.

[100830]17559.

[100831]Bruns, G. A. P.; Mintz, B. J.; Leary, A. C.; Regina, V. M.; Gerald,

[100832]P. S.: Expression of human arylsulfatase A in man-hamster somatic

[100833]cell hybrids. Cytogenet. Cell Genet. 22: 182–185, 1978.

[100834]17560.

[100835]Butterworth, J.; Broadhead, D. M.; Keay, A. J.: Low arylsulphatase

[100836]A activity in a family without metachromatic leukodystrophy. Clin.

[100837]Genet. 14: 213–218, 1978.

[100838]17561.

[100839]Caillaud, C.; Blanchor, C.; Akli, S.; Crosnier, J. M.; Puech,

[100840]J. P.; Kahn, A.; Poenaru, L.: Molecular basis of late infantile metachromatic

[100841]leukodystrophy in France. (Abstract) ESGLD Workshop Delphi, and

[100842]Greece, and Oct, 1993.

[100843]17562.

[100844]Chabas, A.; Castellvi, S.; Bayes, M.; Balcells, S.; Grinberg,

[100845]D.; Vilageliu, L.; Marfany, G.; Lissens, W.; Gonzalez-Duarte, R.:

[100846]Frequency of the arylsulphatase A pseudodeficiency allele in the Spanish

[100847]population. Clin. Genet. 44: 320-323, 1993.

[100848]17563.

[100849]Chang, P. L.; Davidson, R. G.: Pseudo arylsulphatase-A deficiency

[100850]in healthy individuals: genetic and biochemical relationship to metachromatic

[100851]leukodystrophy. Proc. Nat. Acad. Sci. 80: 7323-7327, 1983.

[100852]17564.

[100853]Chang, P. L.; Davidson, R. G.: Complementation of arylsul-
fatase

[100854]A in somatic hybrids of metachromatic leukodystrophy
and multiple

[100855]sulfatase deficiency disorder fibroblasts. Proc. Nat. Acad.
Sci. 77:

[100856]6166–6170, 1980.

[100857]17565.

[100858]Chang, P. L.; Rosa, N. E.; Davidson, R. G.: Somatic cell hy-
bridization

[100859]studies on the genetic regulation and allelic mutations in
metachromatic

[100860]leukodystrophy. Hum. Genet. 61: 231–235, 1982.

[100861]17566.

[100862]Coulter–Mackie, M. B.; Rip, J.; Ludman, M. D.; Beis, J.;
Cole,

[100863]D. E. C.: Metachromatic leucodystrophy (MLD) in a patient
with a

[100864]constitutional ring chromosome 22. J. Med. Genet. 32:
787–791, 1995.

[100865]17567.

[100866]Cravioto, H.; O'Brien, J.; Lockwood, R.; Kasten, F. H.;
Booker,

[100867]J.: Metachromatic leukodystrophy (sulfatide lipidoses) cul-
tured in

[100868]vitro. Science 156: 243-245, 1967.

[100869]17568.

[100870]DeLuca, C.; Brown, J. A.; Shows, T. B.: Lysosomal arylsul-
fatase

[100871]deficiencies in humans: chromosome assignment of aryl-
sulfatase A and

[100872]B. Proc. Nat. Acad. Sci. 76: 1957-1961, 1979.

[100873]17569.

[100874]DeLuca, C.; Champion, M. J.; Shows, T. B.: Arylsulfatase-A
(ARSA)

[100875]synteny with beta-glucuronidase (BGUS) indicates assign-
ment to human

[100876]chromosome 7 in man-Chinese hamster hybrids.
(Abstract) Winnipeg

[100877]Gene Mapping Conf. , 1977.

[100878]17570.

[100879] Draghia, R.; Letourneur, F.; Drugan, C.; Manicom, J.; Blanchot,

[100880] C.; Kahn, A.; Poenaru, L.; Caillaud, C.: Metachromic leukodystrophy:

[100881] identification of the first deletion in exon 1 and of nine novel point

[100882] mutations in the arylsulfatase gene. Hum. Mutat. 9: 234-242, 1997.

[100883] 17571.

[100884] Dubois, G.; Harzer, K.; Baumann, N.: Very low arylsulfatase A

[100885] and cerebroside sulfatase activities in leukocytes of healthy members

[100886] of metachromatic leukodystrophy family. Am. J. Hum. Genet. 29: 191-194,

[100887] 1977.

[100888] 17572.

[100889] Eto, Y.; Tahara, T.; Koda, N.; Yamaguchi, S.; Ito, F.; Okuno, A.: Prenatal diagnosis of metachromatic leukodystrophy: a diagnosis

[100891] by amniotic fluid and its confirmation. Arch. Neurol. 39: 29-32,

[100892]1982.

[100893]17573.

[100894]Farrell, D. F.: Heterozygote detection in MLD: allelic mutations

[100895]at the ARA locus. Hum. Genet. 59: 129–134, 1981.

[100896]17574.

[100897]Farrell, D. F.; MacMartin, M. P.; Clark, A. F.: Multiple
molecular
[100898]forms of arylsulfatase A in different forms of metachro-
matic leukodystrophy
[100899](MLD). Neurology 29: 16–20, 1979.
[100900]17575.

[100901]Felice, K. J.; Gomez Lira, M.; Natowicz, M.; Grunnet, M. L.;
Tsongalis,
[100902]G. J.; Sima, A. A. F.; Kaplan, R. F.: Adult-onset MLD: a
gene mutation
[100903]with isolated polyneuropathy. Neurology 55: 1036–1039,
2000.
[100904]17576.

[100905]Fluharty, A. L.; Fluharty, C. B.; Bohne, W.; von Figura, K.;
Gieselmann,
[100906]V.: Two new arylsulfatase A (ARSA) mutations in a juvenile
metachromatic
[100907]leukodystrophy (MLD) patient. Am. J. Hum. Genet. 49:
1340–1350,
[100908]1991.
[100909]17577.

[100910] Francke, U.; Tetri, P.; Taggart, R. T.; Oliver, N.: Conserved
[100911] autosomal syntenic group on mouse (MMU) chromosome
15 and human (HSA)
[100912] chromosome 22: assignment of a gene for arylsulfatase A
to MMU 15
[100913] and regional mapping of DIA1, ARSA, and ACO2 on
HSA22. Cytogenet.
[100914] Cell Genet. 31: 58–69, 1981.
[100915] 17578.
[100916] Geurts van Kessel, A. H. M.; Westerveld, A.; de Groot, P.
G.;
[100917] Meera Khan, P.; Hagemeijer, A.: Regional localization of
the genes
[100918] coding for human ACO2, ARSA, and NAGA on chromo-
some 22. Cytogenet.
[100919] Cell Genet. 28: 169–172, 1980.
[100920] 17579.
[100921] Gieselmann, V.: An assay for the rapid detection of the
arylsulfatase
[100922] A pseudodeficiency allele facilitates diagnosis and genetic
counseling

[100923]for metachromatic leukodystrophy. Hum. Genet. 86:
251–255, 1991.

[100924]17580.

[100925]Gieselmann, V.; Fluharty, A. L.; Tonnesen, T.; Von Figura, K.

[100926]: Mutations in the arylsulfatase A pseudodeficiency allele
causing

[100927]metachromatic leukodystrophy. Am. J. Hum. Genet. 49:
407–413, 1991.

[100928]17581.

[100929]Gieselmann, V.; Polten, A.; Kreysing, J.; von Figura, K.:
Arylsulfatase

[100930]A pseudodeficiency: loss of a polyadenylation (sic) signal
and N-glycosylation

[100931]site. Proc. Nat. Acad. Sci. 86: 9436–9440, 1989.

[100932]17582.

[100933]Gieselmann, V.; Zlotogora, J.; Harris, A.; Wenger, D. A.;
Morris,

[100934]C. P.: Molecular genetics of metachromatic leukodystro-
phy. Hum.

[100935]Mutat. 4: 233–242, 1994.

[100936]17583.

[100937]Goebel, H. H.; Pilz, H.; Argyrakis, A.: Adult metachromatic leukodystrophy.

[100938]II. Ultrastructural findings in peripheral nerve and skeletal muscle. Europ.

[100939]Neurol. 15: 308–317, 1977.

[100940]17584.

[100941]Beck, M.; Glossl, J.; Grubisic, A.; Spranger, J.: Heterogeneity

[100942]of Morquio disease. Clin. Genet. 29: 325–331, 1986.

[100943]17585.

[100944]Maroteaux, P.; Stanescu, V.; Stanescu, R.; Kresse, H.; Hors-Cayla,

[100945]M. C.: Heterogeneite des formes frustes de la maladie de Morquio. Arch.

[100946]Franc. Pediat. 39: 761–765, 1982.

[100947]17586.

[100948]Saunders, M.; Sweetman, L.; Robinson, B.; Roth, K.; Kohn, S.;

[100949]Sherwood, G.; Gravel, R.: Multiple carboxylase defects and complementation

[100950] studies in biotin responsive organic aciduria. (Abstract)
Am. J. Hum.

[100951] Genet. 31: 61A, 1979.

[100952] 17587.

[100953] Saunders, M. E.; Sherwood, W. G.; Duthie, M.; Surh, L.;
Gravel,

[100954] R. A.: Evidence for a defect of holocarboxylase synthetase
activity

[100955] in cultured lymphoblasts from a patient with biotin-
responsive multiple

[100956] carboxylase deficiency. Am. J. Hum. Genet. 34: 590-601,
1982.

[100957] 17588.

[100958] Suormala, T.; Fowler, B.; Duran, M.; Burtcher, A.;
Fuchshuber,

[100959] A.; Tratzmuller, R.; Lenze, M. J.; Raab, K.; Baur, B.; Wick,
H.; Baumgartner,

[100960] R.: Five patients with a biotin-responsive defect in holo-
carboxylase

[100961] formation: evaluation of responsiveness to biotin therapy
in vivo

[100962] and comparative biochemical studies in vitro. Pediat. Res.

41: 666-673,

[100963]1997.

[100964]17589.

[100965]Suzuki, Y.; Aoki, Y.; Ishida, Y.; Chiba, Y.; Iwamatsu, A.;
Kishino,

[100966]T.; Niikawa, N.; Matsubara, Y.; Narisawa, K.: Isolation and
characterization

[100967]of mutations in the human holocarboxylase synthetase
cDNA. Nature

[100968]Genet. 8: 122-128, 1994.

[100969]17590.

[100970]Sweetman, L.; Surh, L.; Baker, H.; Peterson, R. M.; Nyhan,
W.

[100971]L.: Clinical and metabolic abnormalities in a boy with di-
etary deficiency

[100972]of biotin. Pediatrics 68: 553-558, 1981.

[100973]17591.

[100974]Thoene, J.; Baker, H.; Yoshino, M.; Sweetman, L.: Biotin-
responsive

[100975]carboxylase deficiency associated with subnormal plasma
and urinary

[100976]biotin. New Eng. J. Med. 304: 817–820, 1981.

[100977]17592.

[100978]Thoene, J.; Sweetman, L.; Yoshino, M.: Biotin–responsive
multiple

[100979]carboxylase deficiency. (Abstract) Am. J. Hum. Genet. 31:
64A, 1979.

[100980]17593.

[100981]Wolf, B.; Grier, R. E.; Parker, W. D.; Goodman, S. I.; Allen,

[100982]R. J.: Deficient biotinidase activity in late–onset multiple
carboxylase

[100983]deficiency. (Letter) New Eng. J. Med. 308: 161, 1983.

[100984]17594.

[100985]Yang, X.; Aoki, Y.; Li, X.; Sakamoto, O.; Hiratsuka, M.; Gib-
son,

[100986]K. M.; Kure, S.; Narisawa, K.; Matsubara, Y.; Suzuki, Y.:
Haplotype

[100987]analysis suggests that the two predominant mutations in
Japanese patients

[100988]with holocarboxylase synthetase deficiency are founder
mutations. J.

[100989]Hum. Genet. 45: 358–362, 2000.

[100990]17595.

[100991]Yang, X.; Aoki, Y.; Li, X.; Sakamoto, O.; Hiratsuka, M.;
Kure,

[100992]S.; Taheri, S.; Christensen, E.; Inui, K.; Kubota, M.; Ohira,
M.;

[100993]Ohki, M.; and 10 others: Structure of human holocarboxy-
lase synthetase

[100994]gene and mutation spectrum of holocarboxylase syn-
thetase deficiency. Hum.

[100995]Genet. 109: 526-534, 2001.

[100996]17596.

[100997]Zhang, X. X.; Leon-Del-Rio, A.; Gravel, R. A.; Eydoux, P.:
Assignment

[100998]of holocarboxylase synthetase gene (HLCS) to human
chromosome band

[100999]21q22.1 and to mouse chromosome band 16C4 by in situ
hybridization. Cytogenet.

[101000]Cell Genet. 76: 179 only, 1997.

[101001]17597.

[101002]Mangos, J. A.; Opitz, J. M.; Lobeck, C. C.; Cookson, D. V.:
Familial

[101003]juvenile nephronophthisis: an unrecognized renal disease
in the United

[101004]States. Pediatrics 34: 337-345, 1964.

[101005]17598.

[101006]McKusick, V. A.: Mendelian Inheritance in Man. A Catalog
of Human

[101007]Genes and Genetic Disorders. (12th ed.) Baltimore: Johns
Hopkins

[101008]Univ. Press , 1998. Pp. I only. Note: Table 15.

[101009]17599.

[101010]Medhioub, M.; Cherif, D.; Benessy, F.; Silbermann, F.;
Gubler,

[101011]M. C.; Le Paslier, D.; Cohen, D.; Weissenbach, J.; Beck-
mann, J.; Antignac,

[101012]C.: Refined mapping of a gene (NPH1) causing familial ju-
venile nephronophthisis

[101013]and evidence for genetic heterogeneity. Genomics 22:
296-301, 1994.

[101014]17600.

[101015]Meier, D. A.; Hess, J. W.: Familial nephropathy with retini-
tis

[101016]pigmentosa: a new oculorenal syndrome in adults. Am. J. Med. 39:

[101017]58–69, 1965.

[101018]17601.

[101019]Mongeau, J. G.; Worthen, H. G.: Nephronophthisis and medullary

[101020]cystic disease. Am. J. Med. 43: 345–355, 1967.

[101021]17602.

[101022]Saunier, S.; Calado, J.; Benessy, F.; Silbermann, F.; Heilig,

[101023]R.; Weissenbach, J.; Antignac, C.: Characterization of the NPHP1

[101024]locus: mutational mechanism involved in deletions in familial juvenile

[101025]nephronophthisis. Am. J. Hum. Genet. 66: 778–789, 2000.

[101026]17603.

[101027]Saunier, S.; Calado, J.; Heilig, R.; Silbermann, F.; Benessy,

[101028]F.; Morin, G.; Konrad, M.; Broyer, M.; Gubler, M.–C.; Weissenbach,

[101029]J.; Antignac, C.: A novel gene that encodes a protein with a putative

[101030]src homology 3 domain is a candidate gene for familial juvenile nephronophthisis. Hum.

[101031]Molec. Genet. 6: 2317-2323, 1997.

[101032]17604.

[101033]Sherman, F. E.; Studnicki, F. M.; Fetterman, G. H.: Renal lesions

[101034]of familial juvenile nephronophthisis examined by microdissection. Am.

[101035]J. Clin. Path. 55: 391-400, 1971.

[101036]17605.

[101037]Simon, E. A.; Cook, S.; Davisson, M. T.; D'Eustachio, P.; Guay-Woodford,

[101038]L. M.: The mouse congenital polycystic kidney (cpk) locus maps within

[101039]1.3 cM of the chromosome 12 marker D12Nyu2. Genomics 21: 415-418,

[101040]1994.

[101041]17606.

[101042]Steele, B. T.; Lirenman, D. S.; Beattie, G. W.: Nephronophthisis. Am.

[101043]J. Med. 68: 531-538, 1980.

[101044]17607.

[101045]Strauss, M. B.; Sommers, S. C.: Medullary cystic disease
and

[101046]familial juvenile nephronophthisis: clinical and pathologi-
cal identity. New

[101047]Eng. J. Med. 277: 863-864, 1967.

[101048]17608.

[101049]Sworn, M. J.; Eisinger, A. J.: Medullary cystic disease and
juvenile

[101050]nephronophthisis in separate members of the same fam-
ily. Arch. Dis.

[101051]Child. 47: 278-281, 1972.

[101052]17609.

[101053]Von Sydow, G.; Ranstrom, S.: Familial juvenile
nephronophthisis. Acta

[101054]Paediat. 51: 561-574, 1962.

[101055]17610.

[101056]Waldherr, R.; Lennert, T.; Weber, H. P.; Fodisch, H. J.;
Scharer,

[101057]K.: The nephronophthisis complex: a clinicopathologic
study in children. Virchows

[101058]Arch. A Path. Anat. Histol. 394: 235–254, 1982.

[101059]17611.

[101060]Bolk, S.; Puffenberger, E. G.; Hudson, J.; Morton, D. H.;
Chakravarti,

[101061]A.: Elevated frequency and allelic heterogeneity of con-
genital nephrotic

[101062]syndrome, Finnish type, in the Old Order Mennonites.
(Letter) Am.

[101063]J. Hum. Genet. 65: 1785–1790, 1999.

[101064]17612.

[101065]Anikster, Y.; Kleta, R.; Shaag, A.; Gahl, W. A.; Elpeleg, O.:

[101066]Type III 3-methylglutaconic aciduria (optic atrophy plus
syndrome,

[101067]or Costeff optic atrophy syndrome): identification of the
OPA3 gene

[101068]and its founder mutation in Iraqi Jews. Am. J. Hum. Genet.
69: 1218–1224,

[101069]2001.

[101070]17613.

[101071]Bourdon, V.; Philippe, C.; Labrune, O.; Amsallem, D.;
Arnould,

[101072]C.; Jonveaux, P.: A detailed analysis of the MECP2 gene:
prevalence

[101073]of recurrent mutations and gross DNA rearrangements in
Rett syndrome

[101074]patients. Hum. Genet. 108: 43–50, 2001.

[101075]17614.

[101076]Brinke, A.; Green, P. M.; Giannelli, F.: Characterization of
the

[101077]gene (VBP1) and transcript for the von Hippel–Lindau
binding protein

[101078]and isolation of the highly conserved murine homologue.
Genomics 45:

[101079]105–112, 1997.

[101080]17615.

[101081]Brinke, A.; Tagliavacca, L.; Naylor, J.; Green, P.; Gian-
grande,

[101082]P.; Giannelli, F.: Two chimaeric transcription units result
from

[101083]an inversion breaking intron 1 of the factor VIII gene and a
region

[101084]reportedly affected by reciprocal translocations in T-cell
leukaemia. Hum.

- [101085]Molec. Genet. 5: 1945–1951, 1996.
- [101086]17616.
- [101087]Clifford, S. C.; Walsh, S.; Hewson, K.; Green, E. K.; Brinke, A.;
- [101088]Green, P. M.; Gianelli, F.; Eng, C.; Maher, E. R.: Genomic organization
- [101089]and chromosomal localization of the human CUL2 gene and the role of
- [101090]von Hippel–Lindau tumor suppressor–binding protein (CUL2 and VBP1)
- [101091]mutation and loss in renal–cell carcinoma development. Genes Chromosomes
- [101092]Cancer 26: 20–28, 1999.
- [101093]17617.
- [101094]Hemberger, M.; Himmelbauer, H.; Neumann, H. P. H.; Plate, K. H.;
- [101095]Schwarzkopf, G.; Fundele, R.: Expression of the von Hippel–Lindau–binding
- [101096]protein–1 (Vbp1) in fetal and adult mouse tissues. Hum. Molec. Genet. 8:
- [101097]229–236, 1999.

[101098]17618.

[101099]Tsuchiya, H.; Iseda, T.; Hino, O.: Identification of a novel protein

[101100](VBP-1) binding to the von Hippel-Lindau (VHL) tumor suppressor gene

[101101]product. Cancer Res. 56: 2881-2885, 1996.

[101102]17619.

[101103]Aoyama, T.; Tsushima, K.; Souri, M.; Kamijo, T.; Suzuki, Y.; Shimozawa,

[101104]N.; Orii, T.; Hashimoto, T.: Molecular cloning and functional expression

[101105]of a human peroxisomal acyl-coenzyme A oxidase. Biochem. Biophys.

[101106]Res. Commun. 198: 1113-1118, 1994.

[101107]17620.

[101108]Fan, C.-Y.; Pan, J.; Chu, R.; Lee, D.; Kluckman, K. D.; Usuda,

[101109]N.; Singh, I.; Yeldandi, A. V.; Rao, M. S.; Maeda, N.; Reddy, J. K.

[101110]: Hepatocellular and hepatic peroxisomal alterations in mice with

[101111]a disrupted peroxisomal fatty acyl-coenzyme A oxidase gene. J. Biol.

[101112]Chem. 271: 24698–24710, 1996.

[101113]17621.

[101114]Fournier, B.; Saudubray, J.-M.; Benichou, B.; Lyonnet, S.; Munnich,

[101115]A.; Clevers, H.; Poll-The, B. T.: Large deletion of the peroxisomal

[101116]acyl-CoA oxidase gene in pseudoneonatal adrenoleukodystrophy. J.

[101117]Clin. Invest. 94: 526–531, 1994.

[101118]17622.

[101119]Goldfischer, S. L.; Collins, J.; Rapin, I.; Neumann, P.; Neglia,

[101120]W.; Spiro, A. J.; Ishii, T.; Roels, F.; Vamecq, F.; Vamecq, J.; Van

[101121]Hoof, F.: Deficiencies in several peroxisomal oxidative activities. J.

[101122]Pediat. 108: 25–32, 1986.

[101123]17623.

[101124]Moghrabi, N. N.; Dawson, D. B.; Bennett, M. J.: Assignment

of

[101125]the human peroxisomal palmitoyl-CoA oxidase gene to
chromosome 17q23-qter

[101126]by PCR technique. J. Inherit. Metab. Dis. 18: 705-709,
1995.

[101127]17624.

[101128]Naidu, S.; Hoefler, G.; Watkins, P. A.; Chen, W. W.; Moser,
A.

[101129]B.; Hoefler, S.; Rance, N. E.; Powers, J. M.; Beard, M.;
Green, W.

[101130]R.; Hashimoto, T.; Moser, H. W.: Neonatal seizures and re-
tardation

[101131]in a female with biochemical features of X-linked
adrenoleukodystrophy:

[101132]a possible new peroxisomal disease entity. Neurology 38:
1100-1107,

[101133]1988.

[101134]17625.

[101135]Poll-The, B. T.; Roels, F.; Ogier, H.; Scotto, J.; Vamecq, J.;

[101136]Schutgens, R. B. H.; Wanders, R. J. A.; van Roermund, C. W.
T.; van

[101137]Wiljand, M. J. A.; Schram, A. W.; Tager, J. M.; Saudubray,

J.-M.:

[101138]A new peroxisomal disorder with enlarged peroxisomes and a specific

[101139]deficiency of acyl-CoA oxidase (pseudo-neonatal adrenoleukodystrophy). Am.

[101140]J. Hum. Genet. 42: 422-434, 1988.

[101141]17626.

[101142]Reddy, J. K.; Rao, M. S.: Peroxisome proliferators and cancer:

[101143]mechanisms and implications. Trends Pharm. Sci. 7: 438-443, 1986.

[101144]17627.

[101145]Varanasi, U.; Chu, R.; Chu, S.; Espinosa, R.; LeBeau, M. M.; Reddy,

[101146]J. K.: Isolation of the human peroxisomal acyl-CoA oxidase gene:

[101147]organization, promoter analysis, and chromosomal localization. Proc.

[101148]Nat. Acad. Sci. 91: 3107-3111, 1994.

[101149]17628.

[101150]Watkins, P. A.; McGuinness, M. C.; Raymond, G. V.; Hicks,

B. A.;

[101151]Sisk, J. M.; Moser, A. B.; Moser, H. W.: Distinction between
peroxisomal

[101152]bifunctional enzyme and acyl-CoA oxidase deficiencies.

Ann. Neurol. 38:

[101153]472-477, 1995.

[101154]17629.

[101155]Buyse, I. M.; Fang, P.; Hoon, K. T.; Amir, R. E.; Zoghbi, H.

Y.;

[101156]Roa, B. B.: Diagnostic testing for Rett syndrome by DHPLC
and direct

[101157]sequencing analysis of the MECP2 gene: identification of
several novel

[101158]mutations and polymorphisms. Am. J. Hum. Genet. 67:
1428-1436, 2000.

[101159]17630.

[101160]Cheadle, J. P.; Gill, H.; Fleming, N.; Maynard, J.; Kerr, A.;
Leonard,

[101161]H.; Krawczak, M.; Cooper, D. N.; Lynch, S.; Thomas, N.;
Hughes, H.;

[101162]Hulten, M.; Ravine, D.; Sampson, J. R.; Clarke, A.: Long-
read sequence

- [101163]analysis of the MECP2 gene in Rett syndrome patients:
correlation
- [101164]of disease severity with mutation type and location. Hum.
Molec.
- [101165]Genet. 9: 1119–1129, 2000.
- [101166]17631.
- [101167]Chen, R. Z.; Akbarian, S.; Tudor, M.; Jaenisch, R.: Defi-
ciency
- [101168]of methyl–CpG binding protein–2 in CNS neurons results
in a Rett–like
- [101169]phenotype in mice. Nature Genet. 27: 327–331, 2001.
- [101170]17632.
- [101171]Claes, S.; Devriendt, K.; D'Adamo, P.; Meireleire, J.; Raey-
maekers,
- [101172]P.; Toniolo, D.; Cassiman, J.–J.; Fryns, J.–P.: X–linked se-
vere mental
- [101173]retardation and a progressive neurological disorder in a
Belgian family:
- [101174]clinical and genetic studies. Clin. Genet. 52: 155–161,
1997.
- [101175]17633.

[101176]Clayton-Smith, J.; Watson, P.; Ramsden, S.; Black, G. C. M.:
[101177]Somatic mutation in MECP2 as a non-fatal neurodevelopmental disorder
[101178]in males. Lancet 356: 830-832, 2000.
[101179]17634.
[101180]Couvert, P.; Bienvenu, T.; Aquaviva, C.; Poirier, K.;
Moraine,
[101181]C.; Gendrot, C.; Verloes, A.; Andres, C.; Le Fevre, A. C.;
Souville,
[101182]J.; Steffann, J.; des Portes, V.; Ropers, H.-H.; Yntema, H.
G.; Fryns,
[101183]J.-P.; Briault, S.; Chelly, J.; Cherif, B.: MECP2 is highly mutated
[101184]in X-linked mental retardation. Hum. Molec. Genet. 10:
941-946,
[101185]2001.
[101186]17635.
[101187]D'Esposito, M.; Quaderi, N. A.; Ciccodicola, A.; Bruni, P.;
Esposito,
[101188]T.; D'Urso, M.; Brown, S. D. M.: Isolation, physical mapping, and

- [101189]Northern analysis of the X-linked human gene encoding methyl CpG-binding
- [101190]protein, MECP2. Mammalian Genome 7: 533–535, 1996.
- [101191]17636.
- [101192]De Bona, C.; Zappella, M.; Hayek, G.; Meloni, I.; Vitelli, F.;
- [101193]Bruttini, M.; Cusano, R.; Loffredo, P.; Longo, I.; Renieri, A.:
Preserved
- [101194]speech variant is allelic of classic Rett syndrome. Europ. J.
Hum.
- [101195]Genet. 8: 325–330, 2000.
- [101196]17637.
- [101197]Dragich, J.; Houwink–Manville, I.; Schanen, C.: Rett syn-
drome:
- [101198]a surprising result of mutation in MECP2. Hum. Molec.
Genet. 9:
- [101199]2365–2375, 2000.
- [101200]17638.
- [101201]Gendrot, C.; Ronce, N.; Raynaud, M.; Ayrault, A.–D.;
- Dourlens,
- [101202]J.; Castelnau, P.; Muh, J.–P.; Chelly, J.; Moraine, C.: X-
linked

[101203]nonspecific mental retardation (MRX16) mapping to distal
Xq28: linkage

[101204]study and neuropsychological data in a large family. Am.
J. Med.

[101205]Genet. 83: 411–418, 1999.

[101206]17639.

[101207]Guy, J.; Hendrich, B.; Holmes, M.; Martin, J. E.; Bird, A.: A

[101208]mouse Mecp2-null mutation causes neurological symptoms that mimic
toms that mimic

[101209]Rett syndrome. Nature Genet. 27: 322–326, 2001.

[101210]17640.

[101211]Hagberg, B.: Rett's syndrome: prevalence and impact on
progressive

[101212]severe mental retardation in girls. Acta Pediat. Scand. 74:
405–408,

[101213]1985.

[101214]17641.

[101215]Heilstedt, H. A.; Shahbazian, M. D.; Lee, B.: Infantile hypotonia
tonia

[101216]as a presentation of Rett syndrome. Am. J. Med. Genet.
111: 238–242,

[101217]2002.

[101218]17642.

[101219]Hoffbuhr, K.; Devaney, J. M.; LaFleur, B.; Sirianni, N.;
Scacheri,

[101220]C.; Giron, J.; Schuette, J.; Innis, J.; Marino, M.; Philippart,
M.;

[101221]Narayanan, V.; Umansky, R.; Kronn, D.; Hoffman, E. P.;
Naidu, S.:

[101222]MeCP2 mutations in children with and without the pheno-
type of Rett

[101223]syndrome. Neurology 56: 1486–1495, 2001.

[101224]17643.

[101225]Huppke, P.; Laccone, F.; Kramer, N.; Engel, W.; Hanefeld,
F.:

[101226]Rett syndrome: analysis of MECP2 and clinical characteri-
zation of

[101227]31 patients. Hum. Molec. Genet. 9: 1369–1375, 2000.

[101228]17644.

[101229]Imessaoudene, B.; Bonnefont, J.-P.; Royer, G.; Cormier-
Daire,

[101230]V.; Lyonnet, S.; Lyon, G.; Munnich, A.; Amiel, J.: MECP2

mutation

[101231]in non-fatal, non-progressive encephalopathy in a male. J.

Med. Genet. 38:

[101232]171-174, 2001.

[101233]17645.

[101234]Jones, P. L.; Veenstra, G. J. C.; Wade, P. A.; Vermaak, D.;

Kass,

[101235]S. U.; Landsberger, N.; Strouboulis, J.; Wolffe, A. P.:

Methylated

[101236]DNA and MeCP2 recruit histone deacetylase to repress

transcription. Nature

[101237]Genet. 19: 187-191, 1998.

[101238]17646.

[101239]Klauck, S. M.; Lindsay, S.; Beyer, K. S.; Splitt, M.; Burn, J.;

[101240]Poustka, A.: A mutation hot spot for nonspecific X-linked

mental

[101241]retardation in the MECP2 gene causes the PPM-X syn-

drome. Am. J. Hum.

[101242]Genet. 70: 1034-1037, 2002.

[101243]17647.

[101244]Kleefstra, T.; Yntema, H. G.; Oudakker, A. R.; Romein, T.;

Sisternans,

[101245]E.; Nillesen, W.; van Bokhoven, H.; de Vries, B. B. A.;
Hamel, B.

[101246]C. J.: De novo MECP2 frameshift mutation in a boy with
moderate mental

[101247]retardation, obesity and gynaecomastia. Clin. Genet. 61:
359–362,

[101248]2002.

[101249]17648.

[101250]LaSalle, J. M.; Goldstine, J.; Balmer, D.; Greco, C. M.:
Quantitative

[101251]localization of heterogeneous methyl–CpG–binding pro-
tein 2 (MeCP2)

[101252]expression phenotypes in normal and Rett syndrome brain
by laser scanning

[101253]cytometry. Hum. Molec. Genet. 10: 1729–1740, 2001.

[101254]17649.

[101255]Lewis, J. D.; Meehan, R. R.; Henzel, W. J.; Maurer–Fogy, I.;
Jeppesen,

[101256]P.; Klein, F.; Bird, A.: Purification, sequence, and cellular
localization

[101257]of a novel chromosomal protein that binds to methylated

DNA. Cell 69:

[101258]905–914, 1992.

[101259]17650.

[101260]Meloni, I.; Bruttini, M.; Longo, I.; Mari, F.; Rizzolio, F.;

D'Adamo,

[101261]P.; Denvriendt, K.; Fryns, J.-P.; Toniolo, D.; Renieri, A.: A

mutation

[101262]in the Rett syndrome gene, MECP2, causes X-linked men-

tal retardation

[101263]and progressive spasticity in males. Am. J. Hum. Genet.

67: 982–985,

[101264]2000. Note: Erratum: Am. J. Hum. Genet. 67: 1365 only,

2000.

[101265]17651.

[101266]Moncla, A.; Kpebe, A.; Missirian, C.; Mancini, J.; Villard, L.

[101267]: Polymorphisms in the C-terminal domain of MECP2 in

mentally handicapped

[101268]boys: implications for genetic counselling. Europ. J. Hum.

Genet. 10:

[101269]86–89, 2002.

[101270]17652.

[101271]Nan, X.; Campoy, F. J.; Bird, A.: MeCP2 is a transcriptional
[101272]repressor with abundant binding sites in genomic chromatin. Cell 88:
[101273]471–481, 1997.
[101274]17653.
[101275]Reuber, B. E.; Germain–Lee, E.; Collins, C. S.; Morrell, J. C.;
[101276]Ameritunga, R.; Moser, H. W.; Valle, D.; Gould, S. J.: Mutations
[101277]in PEX1 are the most common cause of peroxisome biogenesis disorders. Nature
[101278]Genet. 17: 445–448, 1997.
[101279]17654.
[101280]Corbaz, A.; ten Hove, T.; Herren, S.; Graber, P.; Schwartz–
[101281]B.; Belzer, I.; Harrison, J.; Plitz, T.; Kosco–Vilbois, M. H.; Kim,
[101282]S.–H.; Dinarello, C. A.; Novick, D.; van Deventer, S.; Chvatchko,
[101283]Y.: IL–18-binding protein expression by endothelial cells and macrophages
[101284]is up-regulated during active Crohn's disease. J. Immun.

168: 3608–3616,

[101285]2002.

[101286]17655.

[101287]Turner, N.; Mason, P. J.; Brown, R.; Fox, M.; Povey, S.;
Rees,

[101288]A.; Pusey, C. D.: Molecular cloning of the human Good-
pasture antigen

[101289]demonstrates it to be the alpha-3 chain of type IV colla-
gen. J. Clin.

[101290]Invest. 89: 592–601, 1992.

[101291]17656.

[101292]Alexander, R. W.: Hypertension and the pathogenesis of
atherosclerosis.

[101293]Oxidative stress and the mediation of arterial inflamma-
tory response:

[101294]a new perspective. Hypertension 25: 155–161, 1995.

[101295]17657.

[101296]Bu-Ghanim, H. N.; Casimir, C. M.; Povey, S.; Segal, A. W.:
The

[101297]alpha subunit of cytochrome b(-245) mapped to chromo-
some 16. Genomics 8:

[101298]568-570, 1990.

[101299]17658.

[101300]Clark, R. A.; Malech, H. L.; Gallin, J. I.; Nuno, H.; Volpp, B.

[101301]D.; Pearson, D. W.; Nauseef, W. M.; Curnutte, J. T.: Genetic variants

[101302]of chronic granulomatous disease: prevalence of deficiencies of two

[101303]cytosolic components of the NADPH oxidase system. New Eng. J. Med. 321:

[101304]647-652, 1989.

[101305]17659.

[101306]Abbott, M. H.; Folstein, S. E.; Abbey, H.; Pyeritz, R. E.: Psychiatric

[101307]manifestations of homocystinuria due to cystathionine beta-synthase

[101308]deficiency: prevalence, natural history, and relationship to neurologic

[101309]impairment and vitamin B(6)-responsiveness. Am. J. Med. Genet. 26:

[101310]959-969, 1987.

[101311]17660.

[101312]Almgren, B.; Eriksson, H.; Hemmingsson, A.; Hillerdal, G.;
Larsson,

[101313]E.; Aberg, H.: Abdominal aortic aneurysm in homocystin-
uria. Acta

[101314]Chir. Scand. 144: 545-546, 1978.

[101315]17661.

[101316]Aral, B.; Coude, M.; London, J.; Aupetit, J.; Chasse, J.-F.;
Zabot,

[101317]M.-T.; Chadeaux-Vekemans, B.; Kamoun, P.: Two novel
mutations (K384E

[101318]and L539S) in the C-terminal moiety of the cystathionine
beta-synthase

[101319]protein in two French pyridoxine-responsive homocystin-
uria patients. Hum.

[101320]Mutat. 9: 81-82, 1997.

[101321]17662.

[101322]Avramopoulos, D.; Cox, T.; Kraus, J. P.; Chakravarti, A.;
Antonarakis,

[101323]S. E.: Linkage mapping of the cystathionine beta-synthase
(CBS) gene

[101324]on human chromosome 21 using a DNA polymorphism in
the 3-prime untranslated

[101325]region. Hum. Genet. 90: 566–568, 1993.

[101326]17663.

[101327]Barber, G. W.; Spaeth, G. L.: Pyridoxine therapy in homocystinuria.

[101328](Letter) Lancet I: 337 only, 1967.

[101329]17664.

[101330]Bass, H. N.; LaGrave, D.; Mardach, R.; Cederbaum, S. D.; Fuster,

[101331]C. D.; Chetty, M.: Spontaneous pneumothorax in association with pyridoxine-responsive

[101332]homocystinuria. J. Inherit. Metab. Dis. 20: 831–832, 1997.

[101333]17665.

[101334]Boers, G. H. J.; Fowler, B.; Smals, A. G. H.; Trijbels, F. J. M.;

[101335]Leermakers, A. I.; Kleijer, W. J.; Kloppenborg, P. W. C.: Improved

[101336]identification of heterozygotes for homocystinuria due to cystathionine

[101337]synthase deficiency by the combination of methionine loading and enzyme

[101338]determination in cultured fibroblasts. Hum. Genet. 69: 164–169,

[101339]1985.

[101340]17666.

[101341]Boers, G. H. J.; Smals, A. G. H.; Trijbels, F. J. M.; Fowler, B.;

[101342]Bakkeren, J. A. J. M.; Schoonderwaldt, H. C.; Kleijer, W. J.;
Kloppenborg,

[101343]P. W. C.: Heterozygosity for homocystinuria in premature
peripheral

[101344]and cerebral occlusive arterial disease. New Eng. J. Med.
313: 709-715,

[101345]1985.

[101346]17667.

[101347]Bross, P.; Corydon, T. J.; Andresen, B. S.; Jorgensen, M. M.;
Bolund,

[101348]L.; Gregersen, N.: Protein misfolding and degradation in
genetic

[101349]diseases. Hum. Mutat. 14: 186-198, 1999.

[101350]17668.

[101351]Burke, J. P.; O'Keefe, M.; Bowell, R.; Naughten, E. R.: Ocular

[101352]complications in homocystinuria: early and late treated.
Brit. J.

[101353]Ophthalmol. 73: 427-431, 1989.

[101354]17669.

[101355]Carey, M. C.; Donovan, D. E.; Fitzgerald, O.; McAuley, F.
D.:

[101356]Homocystinuria: a clinical and pathological study of nine
subjects

[101357]in six families. Am. J. Med. 45: 7-25, 1968.

[101358]17670.

[101359]Carey, M. C.; Fennelly, J. J.; Fitzgerald, O.: Homocystinuria.

[101360]II. Subnormal serum folate levels, increased folate clear-
ance and

[101361]effects of folic acid therapy. Am. J. Med. 45: 26-31, 1968.

[101362]17671.

[101363]Carson, N. A. J.; Carre, I. J.: Treatment of homocystinuria
with

[101364]pyridoxine: a preliminary study. Arch. Dis. Child. 44:
387-392,

[101365]1969.

[101366]17672.

[101367]Carson, N. A. J.; Cusworth, D. C.; Dent, C. E.; Field, C. M.

B.;

[101368]Neill, D. W.; Westall, R. G.: Homocystinuria: a new inborn error

[101369]of metabolism associated with mental deficiency. Arch. Dis. Child. 38:

[101370]425-436, 1963.

[101371]17673.

[101372]Carson, N. A. J.; Neill, D. W.: Metabolic abnormalities detected

[101373]in a survey of mentally backward individuals in Northern Ireland. Arch.

[101374]Dis. Child. 37: 505-513, 1962.

[101375]17674.

[101376]Chadefaux, B.; Rethore, M. O.; Raoul, O.; Ceballos, I.; Poissonnier,

[101377]M.; Gilgenkrantz, S.; Allard, D.: Cystathionine beta synthase: gene

[101378]dosage effect in trisomy 21. Biochem. Biophys. Res. Commun. 128:

[101379]40-44, 1985.

[101380]17675.

[101381]Chao, C.-L.; Tsai, H.-H.; Lee, C.-M.; Hsu, S.-M.; Kao, J.-T.;

[101382]Chien, K.-L.; Sung, F.-C.; Lee, Y.-T.: The graded effect of
hyperhomocysteinemia

[101383]on the severity and extent of coronary atherosclerosis.
Atherosclerosis 147:

[101384]379-386, 1999.

[101385]17676.

[101386]Hoffmann, M. M.; Jacob, S.; Luft, D.; Schmulling, R.-M.;
Rett,

[101387]K.; Marz, W.; Haring, H.-U.; Matthaei, S.: Type I hyper-
lipoproteinemia

[101388]due to a novel loss of function mutation of lipoprotein li-
pase, cys239-trp,

[101389]associated with recurrent severe pancreatitis. J. Clin. En-
docr. Metab. 85:

[101390]4795-4798, 2000.

[101391]17677.

[101392]Holt, L. E., Jr.; Aylward, F. X.; Timbers, H. G.: Idiopathic

[101393]familial lipemia. Bull. Johns Hopkins Hosp. 64: 279-314,
1939.

[101394]17678.

[101395]Ishimura-Oka, K.; Faustinella, F.; Kihara, S.; Smith, L. C.; Oka, K.; Chan, L.: A missense mutation (trp86-to-arg) in exon 3 of the lipoprotein lipase gene: a cause of familial chylomicronemia. Am. J. Hum. Genet. 50: 1275-1280, 1992.

[101399]17679.

[101400]Kastelein, J. J. P.; Groenemeyer, B. E.; Hallman, D. M.; Henderson, H.; Reymer, P. W. A.; Gagne, S. E.; Jansen, H.; Seidell, J. C.; Kromhout, D.; Jukema, J. W.; Bruschke, A. V. G.; Boerwinkle, E.; Hayden, M. R.; The Regress Study Group: The asn9 variant of lipoprotein lipase is associated with the -93G promoter mutation and an increased risk of coronary artery disease. Clin. Genet. 53: 27-33, 1998.

[101406]17680.

[101407]Kirchgessner, T. G.; Chuat, J.-C.; Heinzmann, C.; Etienne, J.;

[101408]Guilhot, S.; Svenson, K.; Ameis, D.; Pilon, C.; d'Auriol, L.;
Andalibi,
[101409]A.; Schotz, M. C.; Galibert, F.; Lusi, A. J.: Organization of
the
[101410]human lipoprotein lipase gene and evolution of the lipase
gene family. Proc.
[101411]Nat. Acad. Sci. 86: 9647-9651, 1989.
[101412]17681.
[101413]Kirchgessner, T. G.; Svenson, K. L.; Lusi, A. J.; Schotz, M.
[101414]C.: The sequence of cDNA encoding lipoprotein lipase: a
member of
[101415]a lipase gene family. J. Biol. Chem. 262: 8463-8466,
1987.
[101416]17682.
[101417]Kobayashi, J.; Nishida, T.; Ameis, D.; Stahnke, G.; Schotz,
M.
[101418]C.; Hashimoto, H.; Fukamachi, I.; Shirai, K.; Saito, Y.;
Yoshida,
[101419]S.: A heterozygous mutation (the codon for ser477 to a
stop codon)
[101420]in lipoprotein lipase contributes to a defect in lipid inter-
face recognition

[101421]in a case with type I hyperlipidemia. Biochem. Biophys.

Res. Commun. 182:

[101422]70-77, 1992.

[101423]17683.

[101424]Kobayashi, J.; Sasaki, N.; Tashiro, J.; Inadera, H.; Saito, Y.;

[101425]Yoshida, S.: A missense mutation (ala334-to-thr) in exon
7 of the

[101426]lipoprotein lipase gene in a case with type I hyperlipi-
demia. Biochem.

[101427]Biophys. Res. Commun. 191: 1046-1054, 1993.

[101428]17684.

[101429]Langlois, S.; Deeb, S.; Brunzell, J.; Kastelein, J. J.; Hayden,

[101430]M. R.: A unique insertion accounts for a significant pro-
portion of

[101431]the mutations in the lipoprotein lipase (LPL)
gene.(Abstract) Am.

[101432]J. Hum. Genet. 43: A191, 1988.

[101433]17685.

[101434]Langlois, S.; Deeb, S.; Brunzell, J. D.; Kastelein, J. J.; Hay-
den,

[101435]M. R.: A major insertion accounts for a significant proportion of
[101436]mutations underlying human lipoprotein lipase deficiency.
Proc. Nat.
[101437]Acad. Sci. 86: 948–952, 1989.
[101438]17686.
[101439]Levak–Frank, S.; Radner, H.; Walsh, A.; Stollberger, R.;
Knipping,
[101440]G.; Hoefler, G.; Sattler, W.; Weinstock, P. H.; Breslow, J. L.;
Zechner,
[101441]R.: Muscle–specific overexpression of lipoprotein lipase
causes a
[101442]severe myopathy characterized by proliferation of mitochondria and
[101443]peroxisomes in transgenic mice. J. Clin. Invest. 96:
976–986, 1995.
[101444]17687.
[101445]Li, S.; Oka, K.; Galton, D.; Stocks, J.: Pvu–II RFLP at the human
[101446]lipoprotein lipase (LPL) gene locus. Nucleic Acids Res. 16:
2358,
[101447]1988.

[101448]17688.

[101449]Lo, J. Y.; Smith, L. C.; Chan, L.: Lipoprotein lipase: role of

[101450]intramolecular disulfide bonds in enzyme catalysis.

Biochem. Biophys.

[101451]Res. Commun. 206: 266-271, 1995.

[101452]17689.

[101453]Ma, Y.; Henderson, H. E.; Julien, P.; Roederer, G.; Brunzell,

[101454]J.; Hayden, M. R.: A missense mutation (pro-to-leu207) in

[101455]the human lipoprotein lipase gene is the major cause of
type I hyperlipoproteinemia

[101456]in French Canadians. (Series) Miami Short Reports. Advances

[101457]in Gene Technology: The Molecular Biology of Human Genetic Disease.

[101458]New York: IRL Press (pub.) 1: 1991. Pp. 34 only.

[101459]17690.

[101460]Ma, Y.; Henderson, H. E.; Ven Murthy, M. R.; Roederer, G.;
Monsalve,

[101461]M. V.; Clarke, L. A.; Normand, T.; Julien, P.; Gagne, C.;
Lambert,

[101462]M.; Davignon, J.; Lupien, P. J.; Brunzell, J.; Hayden, M. R.: A
mutation

[101463]in the human lipoprotein lipase gene as the most common
cause of familial

[101464]chylomicronemia in French Canadians. New Eng. J. Med.
324: 1761–1766,

[101465]1991.

[101466]17691.

[101467]Ma, Y.; Liu, M.–S.; Chitayat, D.; Bruin, T.; Beisiegel, U.;
Benlian,

[101468]P.; Foubert, L.; De Gennes, J. L.; Funke, H.; Forsythe, I.;
Blaichman,

[101469]S.; Papanikolaou, M.; Erkelens, D. W.; Kastelein, J.; Brun-
zell, J.

[101470]D.; Hayden, M. R.: Recurrent missense mutations at the
first and

[101471]second base of codon arg243 in human lipoprotein lipase
in patients

[101472]of different ancestries. Hum. Mutat. 3: 52–58, 1994.

[101473]17692.

[101474]Ma, Y.; Liu, M.–S.; Ginzinger, D.; Frohlich, J.; Brunzell, J.

[101475]D.; Hayden, M. R.: Gene-environment interaction in the
conversion
[101476]of a mild-to-severe phenotype in a patient homozygous
for a ser172-to-cys
[101477]mutation in the lipoprotein lipase gene. J. Clin. Invest. 91:
1953-1958,
[101478]1993.
[101479]17693.
[101480]Ma, Y.; Wilson, B. I.; Bijvoet, S.; Henderson, H. E.; Cramb,
E.;
[101481]Roederer, G.; Ven Murthy, M. R.; Julien, P.; Bakker, H. D.;
Kastelein,
[101482]J. J. P.; Brunzell, J. D.; Hayden, M. R.: A missense mutation
(asp250-to-asn)
[101483]in exon 6 of the human lipoprotein lipase gene causes
chylomicronemia
[101484]in patients of different ancestries. Genomics 13: 649-653,
1992.
[101485]17694.
[101486]Mattei, M. G.; Etienne, J.; Chuat, J. C.; Nguyen, V. C.;
Brault,
[101487]D.; Bernheim, A.; Galibert, F.: Assignment of the human

lipoprotein

[101488]lipase (LPL) gene to chromosome band 8p22. Cytogenet.

Cell Genet. 63:

[101489]45–46, 1993.

[101490]17695.

[101491]Miesenbock, G.; Holzl, B.; Foger, B.; Brandstatter, E.;

Paulweber,

[101492]B.; Sandhofer, F.; Patsch, J. R.: Heterozygous lipoprotein

lipase

[101493]deficiency due to a missense mutation as the cause of im-
paired triglyceride

[101494]tolerance with multiple lipoprotein abnormalities. J. Clin.

Invest. 91:

[101495]448–455, 1993.

[101496]17696.

[101497]Nevin, D. N.; Brunzell, J. D.; Deeb, S. S.: The LPL gene in
individuals

[101498]with familial combined hyperlipidemia and decreased LPL
activity. Arteriosclerosis

[101499]Thromb. 14: 869–873, 1994.

[101500]17697.

- [101501]Nevin, N. C.; Slack, J.: Hyperlipidaemic xanthomatosis II: mode
[101502]of inheritance in 55 families with essential hyperlipi-
daemia and xanthomatosis. J.
[101503]Med. Genet. 5: 9-28, 1968.
[101504]17698.
- [101505]Nickerson, D. A.; Taylor, S. L.; Weiss, K. M.; Clark, A. G.;
Hutchinson,
[101506]R. G.; Stengard, J.; Salomaa, V.; Vartiainen, E.; Boerwinkle,
E.;
[101507]Sing, C. F.: DNA sequence diversity in a 9.7-kb region of
the human
[101508]lipoprotein lipase gene. Nature Genet. 19: 233-240,
1998.
[101509]17699.
- [101510]Jacobson, J. H.; Stephens, G.: Hereditary choroidoretinal
degeneration:
[101511]study of a family including electroretinography and adap-
tometry. Arch.
[101512]Ophthalmol. 67: 321-335, 1962.
[101513]17700.

- [101514]Jay, M.: Personal Communication. London, England
6/18/1987.
- [101515]17701.
- [101516]Kaplan, J.; Bonneau, D.; Frezal, J.; Munnich, A.; Dufier, J. L.
- [101517]: Clinical and genetic heterogeneity in retinitis pigmentosa. Hum.
- [101518]Genet. 85: 635–642, 1990.
- [101519]17702.
- [101520]Kaplan, J.; Pelet, A.; Martin, C.; Delrieu, O.; Ayme, S.; Bonneau,
- [101521]D.; Briard, M. L.; Hanauer, A.; Larget-Piet, L.; Lefrancois, P.; Michel-Awad,
- [101522]A.; Plauchu, H.; Dufier, J. L.; Frezal, J.; Munnich, A.: Phenotype-genotype
- [101523]correlations in X linked retinitis pigmentosa. J. Med. Genet. 29:
- [101524]615–623, 1992.
- [101525]17703.
- [101526]Keats, B.; Ott, J.; Conneally, M.: Report of the committee
- [101527]on linkage and gene order. Cytogenet. Cell Genet. 51:

459–502, 1989.

[101528]17704.

[101529]Klein, D.; Franceschetti, A.; Hussels, I.; Race, R. R.; Sanger,

[101530]R.: X-linked retinitis pigmentosa and linkage studies with
the Xg

[101531]blood-groups. Lancet I: 974–975, 1967.

[101532]17705.

[101533]Litt, M.; Weleber, R. G.; Lovrien, E. W.; Litt, R. H.; Payne,

[101534]N.; Murphey, W.; Dubay, C.; Willard, H. F.: A family study
of X-linked

[101535]retinitis pigmentosa. (Abstract) Cytogenet. Cell Genet. 46:
651

[101536]only, 1987.

[101537]17706.

[101538]McQuarrie, M. D.: Two pedigrees of hereditary blindness
in man. J.

[101539]Genet. 30: 147–153, 1935.

[101540]17707.

[101541]Mears, A. J.; Gieser, L.; Yan, D.; Chen, C.; Fahrner, S.;
Hiriyanna,

[101542]S.; Fujita, R.; Jacobson, S. G.; Sieving, P. A.; Swaroop, A.:
Protein-truncation

[101543]mutations in the RP2 gene in a North American cohort of
families with

[101544]X-linked retinitis pigmentosa. (Letter) Am. J. Hum. Genet.
64: 897-900,

[101545]1999.

[101546]17708.

[101547]Meitinger, T.; Fraser, N. A.; Lorenz, B.; Zrenner, E.;
Murken,

[101548]J.; Craig, I. W.: Linkage of X-linked retinitis pigmentosa to
the

[101549]hypervariable DNA marker M27-beta (DXS255). Hum.
Genet. 81: 283-286,

[101550]1989.

[101551]17709.

[101552]Miano, M. G.; Testa, F.; Filippini, F.; Trujillo, M.; Conte, I.;

[101553]Lanzara, C.; Millan, J. M.; De Bernardo, C.; Grammatico, B.;
Mangino,

[101554]M.; Torrente, I.; Carrozzo, R.; Simonelli, F.; Rinaldi, E.;
Ventruto,

- [101555]V.; D'Urso, M.; Ayuso, C.; Ciccodicola, A.: Identification of novel
- [101556]RP2 mutations in a subset of X-linked retinitis pigmentosa families
- [101557]and prediction of new domains. Hum. Mutat. 18: 109–119, 2001.
- [101558]17710.
- [101559]Mukai, S.; Dryja, T. P.; Bruns, G. A. P.; Aldridge, J. F.; Berson,
- [101560]E. L.: Linkage between the X-linked retinitis pigmentosa locus and
- [101561]the L1.28 locus. Am. J. Ophthal. 100: 225–229, 1985.
- [101562]17711.
- [101563]Schwahn, U.; Lenzner, S.; Dong, J.; Feil, S.; Hinzmann, B.; van
- [101564]Duijnhoven, G.; Kirschner, R.; Hemberger, M.; Bergen, A. A. B.; Rosenberg,
- [101565]T.; Pinckers, A. J. L. G.; Fundele, R.; Rosenthal, A.; Cremers, F.
- [101566]P. M.; Ropers, H.-H.; Berger, W.: Positional cloning of the gene
- [101567]for X-linked retinitis pigmentosa 2. Nature Genet. 19:

327-332,

[101568]1998.

[101569]17712.

[101570]Schwahn, U.; Paland, N.; Techritz, S.; Lenzner, S.; Berger, W.

[101571]: Mutations in the X-linked RP2 gene cause intracellular misrouting

[101572]and loss of the protein. Hum. Molec. Genet. 10: 1177-1183, 2001.

[101573]17713.

[101574]Sharon, D.; Bruns, G. A. P.; McGee, T. L.; Sandberg, M. A.; Berson,

[101575]E. L.; Dryja, T. P.: X-linked retinitis pigmentosa: mutation spectrum

[101576]of the RPGR and RP2 genes and correlation with visual function. Invest.

[101577]Ophthal. Vis. Sci. 41: 2712-2721, 2000.

[101578]17714.

[101579]Spence, M. A.; Elston, R. C.; Cederbaum, S. D.: Pedigree analysis

[101580]to determine the mode of inheritance in a family with re-

tinitis pigmentosa. Clin.

[101581]Genet. 5: 338-343, 1974.

[101582]17715.

[101583]Thiselton, D. L.; Hampson, R. M.; Nayudu, M.; Van
Maldergem, L.;

[101584]Wolf, M. L.; Saha, B. K.; Bhattacharya, S. S.; Hardcastle, A.
J.:

[101585]Mapping the RP2 locus for X-linked retinitis pigmentosa
on proximal

[101586]Xp: a genetically defined 5-cM critical region and exclu-
sion of candidate

[101587]genes by physical mapping. Genome Res. 6: 1093-1102,
1996.

[101588]17716.

[101589]Usher, C. H.: Bowman lecture on a few hereditary eye af-
fections. Trans.

[101590]Ophthal. Soc. U.K. 55: 164-245, 1935.

[101591]17717.

[101592]Waardenburg, P. J.: Das menschliche Auge und seine Er-
banlagean.

[101593]S-Gravenhage: Martinus Nijhoff (pub.) 1932.

[101594]17718.

[101595]Waardenburg, P. J.; Franceschetti, A.; Klein, D.: Genetics
and

[101596]Ophthalmology. Oxford: Blackwell (pub.) 1: 1961. Pp. 799
only.

[101597]17719.

[101598]Warburg, M.; Simonsen, S. E.: Sex-linked recessive retinitis

[101599]pigmentosa: a preliminary study of the carriers. Acta Oph-
thal. 46:

[101600]494-499, 1968.

[101601]17720.

[101602]Wright, A. F.; Bhattacharya, S. S.; Aldred, M. A.; Jay, M.;
Carothers,

[101603]A. D.; Thomas, N. S. T.; Bird, A. C.; Jay, B.; Evans, H. J.: Ge-
netic

[101604]localisation of the RP2 type of X linked retinitis pigmen-
tosa in a

[101605]large kindred. J. Med. Genet. 28: 453-457, 1991.

[101606]17721.

[101607]Wright, A. F.; Bhattacharya, S. S.; Clayton, J. F.; Dempster,

[101608]M.; Tippett, P.; McKeown, C. M. E.; Jay, M.; Jay, B.; Bird, A. C.

[101609]; Linkage relationships between X-linked retinitis pigmentosa and

[101610]nine short-arm markers: exclusion of the disease locus from Xp21 and

[101611]localization to between DXS7 and DXS14. Am. J. Hum. Genet. 41: 635-644,

[101612]1987.

[101613]17722.

[101614]Wright, A. F.; Bhattacharya, S. S.; Clayton, J. F.; Jay, M.; Jay,

[101615]B.; Bird, A. C.; Evans, H. J.: Linkage relationships about X-linked

[101616]retinitis pigmentosa (RP2). (Abstract) Cytogenet. Cell Genet. 46:

[101617]720 only, 1987.

[101618]17723.

[101619]Lahn, B. T.; Page, D. C.: Retroposition of autosomal mRNA yielded

[101620]testis-specific gene family on human Y chromosome. Nature Genet. 21:

[101621]429–433, 1999.

[101622]17724.

[101623]Yen, P. H.: A long-range restriction map of deletion interval

[101624]6 of the human Y chromosome: a region frequently deleted in azoospermic

[101625]males. Genomics 54: 5–12, 1998.

[101626]17725.

[101627]MacLean, H. E.; Chu, S.; Warne, G. L.; Zajac, J. D.: Related

[101628]individuals with different androgen receptor gene deletions. J. Clin.

[101629]Invest. 91: 1123–1128, 1993.

[101630]17726.

[101631]Madgar, I.; Green, L.; Kent-First, M.; Weissenberg, R.; Gershoni-Baruch,

[101632]R.; Goldman, B.; Friedman, E.: Genotyping of Israeli infertile men

[101633]with idiopathic oligozoospermia. Clin. Genet. 62: 203–207, 2002.

[101634]17727.

[101635] Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Griffin, J. E.; Wilson,
son,
[101636] J. D.; McPhaul, M. J.: Definition of the human androgen re-
ceptor
[101637] gene structure permits the identification of mutations that
cause
[101638] androgen resistance: premature termination of the recep-
tor protein
[101639] at amino acid residue 588 causes complete androgen re-
sistance. Molec.
[101640] Endocr. 4: 1105-1116, 1990.
[101641] 17728.
[101642] Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Wilson, J. D.; Grif-
fin,
[101643] J. E.; McPhaul, M. J.: A single nucleotide substitution intro-
duces
[101644] a premature termination codon into the androgen recep-
tor gene of a
[101645] patient with receptor-negative androgen resistance. J.
Clin. Invest. 85:
[101646] 1522-1528, 1990.
[101647] 17729.

[101648]Marcelli, M.; Tilley, W. D.; Zoppi, S.; Griffin, J. E.; Wilson,
[101649]J. D.; McPhaul, M. J.: Androgen resistance associated with
a mutation
[101650]of the androgen receptor at amino acid 772 (arg-to-cys)
results from
[101651]a combination of decreased messenger ribonucleic acid
levels and impairment
[101652]of receptor function. J. Clin. Endocr. 73: 318–325, 1991.
[101653]17730.

[101654]McC Campbell, A.; Taylor, J. P.; Taye, A. A.; Robitschek, J.; Li,
[101655]M.; Walcott, J.; Merry, D.; Chai, Y.; Paulson, H.; Sobue, G.;
Fischbeck,
[101656]K. H.: CREB-binding protein sequestration by expanded
polyglutamine. Hum.
[101657]Molec. Genet. 9: 2197–2202, 2000.
[101658]17731.

[101659]McPhaul, M. J.; Griffin, J. E.: Male pseudohermaphroditism
caused
[101660]by mutations of the human androgen receptor. J. Clin. En-
docr. Metab. 84:
[101661]3435–3441, 1999.

[101662]17732.

[101663]McPhaul, M. J.; Marcelli, M.; Tilley, W. D.; Griffin, J. E.;
Isidro-Gutierrez,

[101664]R. F.; Wilson, J. D.: Molecular basis of androgen resistance
in a

[101665]family with a qualitative abnormality of the androgen re-
ceptor and

[101666]responsive to high-dose androgen therapy. J. Clin. Invest.
87: 1413-1421,

[101667]1991.

[101668]17733.

[101669]McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Griffin, J. E.; Wilson,

[101670]J. D.: Genetic basis of endocrine disease 4: the spectrum
of mutations

[101671]in the androgen receptor gene that causes androgen re-
sistance. J.

[101672]Clin. Endocr. Metab. 76: 17-23, 1993.

[101673]17734.

[101674]McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Wilson, C. M.; Grif-
fin,

[101675]J. E.; Wilson, J. D.: Mutations in the ligand-binding domain

of the

[101676] androgen receptor gene cluster in two regions of the
gene. J. Clin.

[101677] Invest. 90: 2097–2101, 1992.

[101678] 17735.

[101679] McPhaul, M. J.; Schweikert, H.-U.; Allman, D. R.: Assess-
ment

[101680] of androgen receptor function in genital skin fibroblasts
using a

[101681] recombinant adenovirus to deliver an androgen-respon-
sive reporter

[101682] gene. J. Clin. Endocr. Metab. 82: 1944–1948, 1997.

[101683] 17736.

[101684] Mifsud, A.; Ramirez, S.; Yong, E. L.: Androgen receptor
gene

[101685] CAG trinucleotide repeats in anovulatory infertility and
polycystic

[101686] ovaries. J. Clin. Endocr. Metab. 85: 3484–3488, 2000.

[101687] 17737.

[101688] Migeon, B. R.; Brown, T. R.; Axelman, J.; Migeon, C. J.:
Studies

[101689] of the locus for androgen receptor: localization on the human X and

[101690] evidence for homology with the Tfm locus in the mouse.
Proc. Nat.

[101691] Acad. Sci. 78: 6339–6343, 1981.

[101692] 17738.

[101693] Mongan, N. P.; Jaaskelainen, J.; Green, K.; Schwabe, J. W.;
Shimura,

[101694] N.; Dattani, M.; Hughes, I. A.: Two de novo mutations in
the AR gene

[101695] cause the complete androgen insensitivity syndrome in a
pair of monozygotic

[101696] twins. J. Clin. Endocr. Metab. 87: 1057–1061, 2002.

[101697] 17739.

[101698] Mononen, N.; Syrjakoski, K.; Matikainen, M.; Tammela, T.
L. J.;

[101699] Schleutker, J.; Kallioniemi, O.-P.; Trapman, J.; Koivisto, P.
A.:

[101700] Two percent of Finnish prostate cancer patients have a
germ-line mutation

[101701] in the hormone-binding domain of the androgen receptor
gene. Cancer

[101702]Res. 60: 6479–6481, 2000.

[101703]17740.

[101704]Muroño, K.; Mendonça, B. B.; Arnhold, I. J. P.; Rigon, A. C.
M.

[101705]M.; Migeon, C. J.; Brown, T. R.: Human androgen insensi-
tivity due

[101706]to point mutations encoding amino acid substitutions in
the androgen

[101707]receptor steroid-binding domain. Hum. Mutat. 6:
152–162, 1995.

[101708]17741.

[101709]Nakao, R.; Haji, M.; Yanase, T.; Ogo, A.; Takayanagi, R.;
Katsube,

[101710]T.; Fukumaki, Y.; Nawata, H.: A single amino acid substi-
tution (met786-to-val)

[101711]in the steroid-binding domain of human androgen recep-
tor leads to

[101712]complete androgen insensitivity syndrome. J. Clin. Endocr.
Metab. 74:

[101713]1152–1157, 1992.

[101714]17742.

- [101715]Newmark, J. R.; Hardy, D. O.; Tonb, D. C.; Carter, B. S.; Epstein,
stein,
[101716]J. I.; Isaacs, W. B.; Brown, T. R.; Barrack, E. R.: Androgen
receptor
[101717]gene mutations in human prostate cancer. Proc. Nat.
Acad. Sci. 89:
[101718]6319-6323, 1992.
[101719]17743.
[101720]Nguyen, D.; Steinberg, S. V.; Rouault, E.; Chagnon, S.;
Gottlieb,
[101721]B.; Pinsky, L.; Trifiro, M.; Mader, S.: A G577R mutation in
the human
[101722]AR P box results in selective decreases in DNA binding
and in partial
[101723]androgen insensitivity syndrome. Molec. Endocr. 15:
1790-1802, 2001.
[101724]17744.
[101725]Ohno, S.: The Y-linked antigen locus and the X-linked
Tfm locus
[101726]as major regulatory genes of the mammalian sex deter-
mining mechanism. J.
[101727]Steroid Biochem. 8: 585-592, 1977.

[101728]

[101729]100. Ohno, S.: Simplicity of mammalian regulatory systems inferred

[101730]by single gene determination of sex phenotypes. Nature
234: 134–137,

[101731]1971.

[101732]

[101733]101. Ong, Y. C.; Wong, H. B.; Adaikan, G.; Yong, E. L.: Directed

[101734]pharmacological therapy of ambiguous genitalia due to an
androgen

[101735]receptor gene mutation. (Letter) Lancet 354: 1444–1445,
1999.

[101736]

[101737]102. Patterson, M. N.; Hughes, I. A.; Gottlieb, B.; Pinsky,
L.: The

[101738]androgen receptor gene mutations database. Nucleic
Acids Res. 22:

[101739]3560–3562, 1994.

[101740]

[101741]103. Pinsky, L.; Kaufman, M.; Killinger, D. W.; Burko, B.;

Shatz,

[101742]D.; Volpe, R.: Human minimal androgen insensitivity with normal dihydrotestosterone-binding

[101743]capacity in cultured genital skin fibroblasts: evidence for an androgen-selective

[101744]qualitative abnormality of the receptor. Am. J. Hum. Genet. 36:

[101745]965-978, 1984.

[101746]

[101747]104. Pinsky, L.; Kaufman, M.; Levitsky, L. L.: Partial androgen resistance

[101748]due to a distinctive qualitative defect of the androgen receptor. Am.

[101749]J. Med. Genet. 27: 459-466, 1987.

[101750]

[101751]105. Pinsky, L.; Kaufman, M.; Summitt, R. L.: Congenital androgen

[101752]insensitivity due to a qualitatively abnormal androgen receptor. Am.

[101753]J. Med. Genet. 10: 91-99, 1981.

[101754]

[101755]106. Pinsky, L.; Trifiro, M.; Sebbaghian, N.; Kaufman, M.;
Chang,
[101756]C.; Trapman, J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris,
C. J.;
[101757]Brown, C. J.; Willard, H. F.; Sergovich, F.: A deletional al-
teration
[101758]of the androgen receptor (AR) gene in a sporadic patient
with complete
[101759]androgen insensitivity (CAI) who is mentally retarded.
(Abstract) Am.
[101760]J. Hum. Genet. 45 (suppl.): A212, 1989.
[101761]
[101762]107. Prior, L.; Bordet, S.; Trifiro, M. A.; Mhatre, A.; Kauf-
man, M.;
[101763]Pinsky, L.; Wrogeman, K.; Belsham, D. D.; Pereira, F.;
Greenberg,
[101764]C.; Trapman, J.; Brinkman, A. O.; Chang, C.; Liao, S.: Re-
placement
[101765]of arginine 773 by cysteine or histidine in the human an-
drogen receptor
[101766]causes complete androgen insensitivity with different re-
ceptor phenotypes. Am.

[101767]J. Hum. Genet. 51: 143–155, 1992.

[101768]

[101769]108. Quigley, C. A.; Friedman, K. J.; Johnson, A.; Lafreniere, R.

[101770]G.; Silverman, L. M.; Lubahn, D. B.; Brown, T. R.; Wilson, E. M.;

[101771]Willard, H. F.; French, F. S.: Complete deletion of the androgen

[101772]receptor gene: definition of the null phenotype of the androgen insensitivity

[101773]syndrome and determination of carrier status. J. Clin. Endocr. Metab. 74:

[101774]927–933, 1992.

[101775]

[101776]109. Ris-Stalpers, C.; Kuiper, G. G. J. M.; Faber, P. W.; Schweikert,

[101777]H. U.; van Rooij, H. C. J.; Zegers, N. D.; Hodgins, M. B.; Degenhart,

[101778]H. J.; Trapman, J.; Brinkmann, A. O.: Aberrant splicing of androgen

[101779]receptor mRNA results in synthesis of a nonfunctional receptor protein

[101780]in a patient with androgen insensitivity. Proc. Nat. Acad. Sci. 87:

[101781]7866–7870, 1990.

[101782]

[101783]110. Rodien, P.; Mebarki, F.; Mowszowicz, I.; Chaussain, J.-L.; Young,

[101784]J.; Morel, Y.; Schaison, G.: Different phenotypes in a family with

[101785]androgen insensitivity caused by the same M780I point mutation in

[101786]the androgen receptor gene. J. Clin. Endocr. Metab. 81: 2994–2998,

[101787]1996.

[101788]

[101789]111. Sai, T.; Seino, S.; Chang, C.; Trifiro, M.; Pinsky, L.; Mhatre,

[101790]A.; Kaufman, M.; Lambert, B.; Trapman, J.; Brinkmann, A. O.; Rosenfield,

[101791]R. L.; Liao, S.: An exonic point mutation of the androgen receptor

[101792]gene in a family with complete androgen insensitivity. Am. J. Hum.

[101793]Genet. 46: 1095–1100, 1990.

[101794]

[101795]112. Sammarco, I.; Grimaldi, P.; Rossi, P.; Cappa, M.;
Moretti, C.;

[101796]Frajese, G.; Geremia, R.: Novel point mutation in the splice
donor

[101797]site of exon–intron junction 6 of the androgen receptor
gene in a

[101798]patient with partial androgen insensitivity syndrome. J.
Clin. Endocr.

[101799]Metab. 85: 3256–3261, 2000.

[101800]

[101801]113. Schoenberg, M. P.; Hakimi, J. M.; Wang, S.; Bova, G.
S.; Epstein,

[101802]J. I.; Fischbeck, K. H.; Isaacs, W. B.; Walsh, P. C.; Barrack, E.

[101803]R.: Microsatellite mutation (CAG (24–to–18)) in the andro-
gen receptor

[101804]gene in human prostate cancer Biochem. Biophys. Res.
Commun. 198:

[101805]74–80, 1994.

[101806]

[101807]114. Shang, Y.; Myers, M.; Brown, M.: Formation of the androgen receptor

[101808]transcription complex. Molec. Cell 9: 601–610, 2002.

[101809]

[101810]115. Simeoni, S.; Mancini, M. A.; Stenoién, D. L.; Marcelli, M.; Weigel,

[101811]N. L.; Zanisi, M.; Martini, L.; Poletti, A.: Motoneuronal cell death

[101812]is not correlated with aggregate formation of androgen receptors containing

[101813]an elongated polyglutamine tract. Hum. Molec. Genet. 9: 133–144,

[101814]2000.

[101815]

[101816]116. Sullivan, D. A.; Sullivan, B. D.; Ullman, M. D.; Rocha, E. M.;

[101817]Krenzer, K. L.; Cermak, J. M.; Toda, I.; Doane, M. G.; Evans, J. E.;

[101818]Wickham, L. A.: Androgen influence on the meibomian gland. Invest.

[101819]Ophthal. Vis. Sci. 41: 3732–3742, 2000.

[101820]

[101821]117. Sultan, C.; Lumbroso, S.; Poujol, N.; Belon, C.;
Boudon, C.;

[101822]Lobaccaro, J.-M.: Mutations of androgen receptor gene in
androgen

[101823]insensitivity syndromes. J. Steroid Biochem. Molec. Biol.
46: 519-530,

[101824]1993.

[101825]

[101826]118. Sutherland, R. W.; Wiener, J. S.; Hicks, J. P.; Marcelli,
M.;

[101827]Gonzales, E. T.; Roth, D. R.; Lamb, D. J.: Androgen recep-
tor gene

[101828]mutations are rarely associated with isolated penile hy-
pospadias. J.

[101829]Urol. 156: 828-831, 1996.

[101830]

[101831]119. Taplin, M.-E.; Bubley, G. J.; Shuster, T. D.; Frantz, M.
E.;

[101832]Spooner, A. E.; Ogata, G. K.; Keer, H. N.; Balk, S. P.: Muta-
tion

[101833]of the androgen-receptor gene in metastatic androgen-independent prostate

[101834]cancer. New Eng. J. Med. 332: 1393-1398, 1995.

[101835]

[101836]120. Tilley, W. D.; Marcelli, M.; Wilson, J. D.; McPhaul, M. J.:

[101837]Characterization and expression of a cDNA encoding the human androgen

[101838]receptor. Proc. Nat. Acad. Sci. 86: 327-331, 1989.

[101839]

[101840]121. Trifiro, M.; Prior, L.; Pinsky, L.; Kaufman, M.; Chang, C.; Trapman,

[101841]J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris, C.: A single transition

[101842]at an exonic CpG site apparently abolishes androgen receptor (AR)-binding

[101843]activity in a family with complete androgen insensitivity (CAI). (Abstract) Am.

[101844]J. Hum. Genet. 45 (suppl.): A225, 1989.

[101845]

[101846]122. Trifiro, M.; Prior, R. L.; Sabbaghian, N.; Pinsky, L.;

Kaufman,

[101847]M.; Nylén, E. G.; Belsham, D. D.; Greenberg, C. R.; Wrogermann, K.

[101848]: Amber mutation creates a diagnostic Mael site in the androgen receptor

[101849]gene of a family with complete androgen insensitivity. Am. J. Med.

[101850]Genet. 40: 493–499, 1991.

[101851]

[101852]123. Tut, T. G.; Ghadessy, F. J.; Trifiro, M. A.; Pinsky, L.; Yong,

[101853]E. L.: Long polyglutamine tracts in the androgen receptor are associated

[101854]with reduced trans-activation, impaired sperm production, and male

[101855]infertility. J. Clin. Endocr. Metab. 82: 3777–3782, 1997.

[101856]

[101857]124. Visakorpi, T.; Hyytinen, E.; Koivisto, P.; Tanner, M.; Keinanen,

[101858]R.; Palmberg, C.; Palotie, A.; Tammela, T.; Isola, J.; Kallioniemi,

[101859]O.-P.: In vivo amplification of the androgen receptor gene

and progression

[101860] of human prostate cancer. *Nature Genet.* 9: 401–406,
1995.

[101861]

[101862] 125. Von Eckardstein, S.; Syska, A.; Gromoll, J.; Kamischke, A.; Simoni,

[101863] M.; Nieschlag, E.: Inverse correlation between sperm concentration

[101864] and number of androgen receptor CAG repeats in normal men. *J. Clin.*

[101865] *Endocr. Metab.* 86: 2585–2590, 2001.

[101866]

[101867] 126. Wang, Q.; Ghadessy, F. J.; Trounson, A.; de Kretser, D.; McLachlan,

[101868] R.; Ng, S. C.; Yong, E. L.: Azoospermia associated with a mutation

[101869] in the ligand-binding domain of an androgen receptor displaying normal

[101870] ligand binding, but defective trans-activation. *J. Clin. Endocr.*

[101871] *Metab.* 83: 4303–4309, 1998.

[101872]

[101873]127. Weidemann, W.; Peters, B.; Romalo, G.; Spindler, K.-D.; Schweikert, H.-U.: Response to androgen treatment in a patient with partial androgen insensitivity and a mutation in the deoxyribonucleic acid-binding domain of the androgen receptor. J. Clin. Endocr. Metab. 83: 1173-1176, 1998.

[101878]

[101879]128. Welch, W. J.; Diamond, M. I.: Glucocorticoid modulation of androgen receptor nuclear aggregation and cellular toxicity is associated with distinct forms of soluble expanded polyglutamine protein. Hum. Molec. Genet. 10: 3063-3074, 2001.

[101883]

[101884]129. Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.; Landen, M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Polymorphisms

[101886] of the androgen receptor gene and the estrogen receptor
beta gene

[101887] are associated with androgen levels in women. J. Clin. Endocr. Metab. 86:

[101888] 2562–2568, 2001.

[101889]

[101890] 130. Wieacker, P.; Breckwoldt, M.; Gal, A.: Testicular feminization:

[101891] diagnosis and search for closely linked restriction fragment length

[101892] polymorphism. Dis. Markers 3: 213–218, 1985.

[101893]

[101894] 131. Wieacker, P.; Griffin, J. E.; Wienker, T.; Lopez, J. M.; Wilson,

[101895] J. D.; Breckwoldt, M.: Linkage analysis with RFLPs in families with

[101896] androgen resistance syndromes: evidence for close linkage between

[101897] the androgen receptor locus and the DXS1 segment. Hum. Genet. 76:

[101898] 248–252, 1987.

[101899]

[101900]132. Wilson, C. M.; McPhaul, M. J.: A and B forms of the
androgen

[101901]receptor are present in human genital skin fibroblasts.
Proc. Nat.

[101902]Acad. Sci. 91: 1234–1238, 1994.

[101903]

[101904]133. Wilson, J. D.: The promiscuous receptor: prostate
cancer comes

[101905]of age. (Editorial) New Eng. J. Med. 332: 1440–1441,
1995.

[101906]

[101907]134. Wilson, J. D.; Carlson, B. R.; Weaver, D. D.; Kovacs, W.
J.;

[101908]Griffin, J. E.: Endocrine and genetic characterization of
cousins

[101909]with male pseudohermaphroditism: evidence that the Lubs
phenotype

[101910]can result from a mutation that alters the structure of the
androgen

[101911]receptor. Clin. Genet. 26: 363–370, 1984.

[101912]

[101913]135. Wooster, R.; Mangion, J.; Eeles, R.; Smith, S.;
Dowsett, M.;

[101914]Averill, D.; Barrett-Lee, P.; Easton, D. F.; Ponder, B. A. J.;
Stratton,

[101915]M. R.: A germline mutation in the androgen receptor gene
in two brothers

[101916]with breast cancer and Reifenstein syndrome. Nature
Genet. 2: 132–134,
[101917]1992.

[101918]

[101919]136. Zhang, L.; Leeflang, E. P.; Yu, J.; Arnheim, N.: Study-
ing human
[101920]mutations by sperm typing: instability of CAG trinucleotide repeats

[101921]in the human androgen receptor gene. Nature Genet. 7:
531–535, 1994.

[101922]

[101923]137. Zhu, Y.-S.; Cai, L.-Q.; Cordero, J. J.; Canovatchel, W.
J.; Katz,

[101924]M. D.; Imperato-McGinley, J.: A novel mutation in the CAG
triplet

[101925]region of exon 1 of androgen receptor gene causes complete androgen

[101926]insensitivity syndrome in a large kindred. J. Clin. Endocr.
Metab. 84:

[101927]1590–1594, 1999.

[101928]

[101929]138. Zoppi, S.; Wilson, C. M.; Harbison, M. D.; Griffin, J. E.;
Wilson,

[101930]J. D.; McPhaul, M. J.; Marcelli, M.: Complete testicular fem-
inization

[101931]caused by an amino-terminal truncation of the androgen
receptor with

[101932]downstream initiation. J. Clin. Invest. 91: 1105–1112,
1993.

[101933]17745.

[101934]Konig, A.; Happle, R.; Fink-Puches, R.; Soyer, H. P.; Born-
holdt,

[101935]D.; Engel, H.; Grzeschik, K.-H.: A novel missense mutation
of NSDHL

[101936]in an unusual case of CHILD syndrome showing bilateral,
almost symmetric

[101937]involvement. J. Am. Acad. Derm. 46: 594–596, 2002.

[101938]17746.

[101939]Labit-Le Bouteiller, C.; Jamme, M. F.; David, M.; Silve, S.;
Lanau,

[101940]C.; Dhers, C.; Picard, C.; Rahier, A.; Taton, M.; Loison, G.;
Caput,

[101941]D.; Ferrara, P.; Lupker, J.: Antiproliferative effects of

SR31747A

[101942]in animal cell lines are mediated by inhibition of cholesterol biosynthesis

[101943]at the sterol isomerase step. *Europ. J. Biochem.* 256: 342–349, 1998.

[101944]17747.

[101945]Liu, X. Y.; Dangel, A. W.; Kelley, R. I.; Zhao, W.; Denny, P.;

[101946]Botcherby, M.; Cattanach, B.; Peters, J.; Hunsicker, P. R.; Mallon,

[101947]A.–M.; Strivens, M. A.; Bate, R.; Miller, W.; Rhodes, M.; Brown, S.

[101948]D. M.; Herman, G. E.: The gene mutated in bare patches and striated

[101949]mice encodes a novel 3- β -hydroxysteroid dehydrogenase. *Nature*

[101950]*Genet.* 22: 182–187, 1999.

[101951]17748.

[101952]Schindelhauer, D.; Hellebrand, H.; Grimm, L.; Bader, I.; Meitinger,

[101953]T.; Wehnert, M.; Ross, M.; Meindl, A.: Long-range map of a 3.5-Mb

[101954]region in Xp11.23–22 with a sequence-ready map from a

1.1-Mb gene-rich

[101955]interval. *Genome Res.* 6: 1056–1069, 1996.

[101956]17749.

[101957]Silve, S.; Dupuy, P. H.; Labit-Lebouteiller, C.; Kaghad, M.;
Chalon,

[101958]P.; Rahier, A.; Taton, M.; Lupker, J.; Shire, D.; Loison, G.:
Emopamil-binding

[101959]protein, a mammalian protein that binds a series of struc-
turally diverse

[101960]neuroprotective agents, exhibits delta(8)-delta(7) sterol
isomerase

[101961]activity in yeast. *J. Biol. Chem.* 271: 22434–22440, 1996.

[101962]17750.

[101963]Traupe, H.; Muller, D.; Atherton, D.; Kalter, D. C.; Cremers,

[101964]F. P. M.; van Oost, B. A.; Ropers, H.-H.: Exclusion mapping
of the

[101965]X-linked dominant chondrodysplasia punctata/
ichthyosis/cataract/short

[101966]stature (Happle) syndrome: possible involvement of an
unstable pre-mutation. *Hum.*

[101967]*Genet.* 89: 659–665, 1992.

[101968]17751.

[101969]Jin, H.; Gardner, R. J.; Viswesvariah, R.; Muntoni, F.;
Roberts,

[101970]R. G.: Two novel members of the interleukin-1 receptor
gene family,

[101971]one deleted in Xp22.1-Xp21.3 mental retardation. *Europ.*
J. Hum. Genet. 8:

[101972]87-94, 2000.

[101973]17752.

[101974]Montini, E.; Buchner, G.; Spalluto, C.; Andolfi, G.; Caruso,
A.;

[101975]den Dunnen, J. T.; Trump, D.; Rocchi, M.; Ballabio, A.;
Franco, B.

[101976]: Identification of SCML2, a second human gene homolo-
gous to the *Drosophila*

[101977]sex comb on midleg (Scm): a new gene cluster on Xp22.
Genomics 58:

[101978]65-72, 1999.

[101979]17753.

[101980]Goto, M.; Imamura, O.; Kuromitsu, J.; Matsumoto, T.;
Yamabe, Y.;

- [101981]Tokutake, Y.; Suzuki, N.; Mason, B.; Drayna, D.; Sugawara, M.; Sugimoto, M.; Furuichi, Y.: Analysis of helicase gene mutations in Japanese Werner's syndrome patients. Hum. Genet. 99: 191–193, 1997.
- [101982]17754.
- [101983]Gray, M. D.; Shen, J.–C.; Kamath–Loeb, A. S.; Blank, A.; Sopher, B. L.; Martin, G. M.; Oshima, J.; Loeb, L. A.: The Werner syndrome protein is a DNA helicase. Nature Genet. 17: 100–103, 1997.
- [101984]17755.
- [101985]Imamura, O.; Ichikawa, K.; Yamabe, Y.; Goto, M.; Sugawara, M.; Furuichi, Y.: Cloning of a mouse homologue of the human Werner syndrome gene and assignment to 8A4 by fluorescence in situ hybridization. Genomics 41: 298–300, 1997.
- [101986]17756.

- [101994]Ishikawa, Y.; Sugano, H.; Matsumoto, T.; Furuichi, Y.;
Miller,
[101995]R. W.; Goto, M.: Unusual features of thyroid carcinomas in
Japanese
[101996]patients with Werner syndrome and possible genotype-
phenotype relations
[101997]to cell type and race. Cancer 85: 1345-1352, 1999.
[101998]17757.
- [101999]Marciniak, R. A.; Lombard, D. B.; Johnson, F. B.; Guarente,
L.
[102000]: Nucleolar localization of the Werner syndrome protein in
human cells. Proc.
[102001]Nat. Acad. Sci. 95: 6887-6892, 1998.
[102002]17758.
- [102003]Boyle, S.; Gilchrist, S.; Bridger, J. M.; Mahy, N. L.; Ellis, J.
[102004]A.; Bickmore, W. A.: The spatial organization of human
chromosomes
[102005]within the nuclei of normal and emerin-mutant cells.
Hum. Molec.
[102006]Genet. 10: 211-219, 2001.
[102007]17759.

- [102008]Clarke, A.: Hypohidrotic ectodermal dysplasia. J. Med. Genet. 24: 659–663, 1987.
- [102009]17760.
- [102010]17760.
- [102011]Cartegni, L.; Raffaele di Barletta, M.; Barresi, R.; Squarzoni, S.; Sabatelli, P.; Maraldi, N.; Mora, M.; Di Blasi, C.; Corneo, F.; Merlini, L.; Villa, A.; Cobiauchi, F.; Toniolo, D.: Heart-specific localization of emerin: new insights into Emery–Dreifuss muscular dystrophy. Hum. Molec. Genet. 6: 2257–2264, 1997.
- [102012]17761.
- [102013]Consalez, G. G.; Thomas, N. S. T.; Stayton, C. L.; Knight, S. J. L.; Johnson, M.; Hopkins, L. C.; Harper, P. S.; Elsas, L. J.; Warren, S. T.: Assignment of Emery–Dreifuss muscular dystrophy to the distal region of Xq28: the results of a collaborative study. Am. J. Hum. Genet. 61: 1020–1028, 1997.

[102021]Genet. 48: 468–480, 1991.

[102022]17762.

[102023]Ellis, J. A.; Yates, J. R. W.; Kendrick-Jones, J.; Brown, C. A.

[102024]: Changes at P183 of emerin weaken its protein–protein interactions

[102025]resulting in X-linked Emery–Dreifuss muscular dystrophy. Hum. Genet. 104:

[102026]262–268, 1999.

[102027]17763.

[102028]Hodgson, S. V.; Boswinkel, E.; Walker, A.; Bobrow, M.; Davies,

[102029]K.; Dubowitz, V.; Granata, G.; Merlini, L.: Linkage analysis using

[102030]nine DNA polymorphisms along the length of the X chromosome locates

[102031]the gene for Emery–Dreifuss muscular dystrophy to distal Xq. (Abstract) J.

[102032]Med. Genet. 23: 169–170, 1986.

[102033]17764.

[102034]Klauck, S.; Wilgenbus, P.; Yates, J. R. W.; Muller, C. R.; Poustka,

[102035]A.: Identification of novel mutations in three families with
Emery–Dreifuss

[102036]muscular dystrophy. Hum. Mol. Genet. 4: 1853–1857,
1995.

[102037]17765.

[102038]Manilal, S.; Nguyen thi Man; Sewry, C. A.; Morris, G. E.:
The

[102039]Emery–Dreifuss muscular dystrophy protein, emerin, is a
nuclear membrane

[102040]protein. Hum. Molec. Genet. 5: 801–808, 1996.

[102041]17766.

[102042]Manilal, S.; Recan, D.; Sewry, C. A.; Hoeltzenbein, M.;
Llense,

[102043]S.; Leturcq, F.; Deburgrave, N.; Barbot, J.–C.; Nguyen thi
Man; Muntoni,

[102044]F.; Wehnert, M.; Kaplan, J.–C.; Morris, G. E.: Mutations in
Emery–Dreifuss

[102045]muscular dystrophy and their effects on emerin protein
expression. Hum.

[102046]Molec. Genet. 7: 855–864, 1998.

[102047]17767.

- [102048]Nevo, Y.; Al-Lozi, M.; Parsadanian, A. S.; Elliott, J. L.; Connolly,
muscular
[102049]A. M.; Pestronk, A.: Mutation analysis in Emery–Dreifuss
[102050]dystrophy. *Pediat. Neurol.* 21: 456–459, 1999.
[102051]17768.
- [102052]Romeo, G.; Roncuzzi, L.; Sangiorgi, S.; Giacanelli, M.;
Liguori,
[102053]M.; Tessarolo, D.; Rocchi, M.: Mapping of the Emery–
Dreifuss gene
[102054]through reconstruction of crossover points in two Italian
pedigrees. *Hum.*
[102055]*Genet.* 80: 59–62, 1988.
[102056]17769.
- [102057]Thomas, N. S. T.; Williams, H.; Elsas, L. J.; Hopkins, L. C.;
[102058]Sarfarazi, M.; Harper, P. S.: Localisation of the gene for
Emery–Dreifuss
[102059]muscular dystrophy to the distal long arm of the X chromosome. *J.*
[102060]*Med. Genet.* 23: 596–598, 1986.
[102061]17770.

- [102062]Wulff, K.; Parrish, J. E.; Herrmann, F. H.; Wehnert, M.: Six
[102063]novel mutations in the emerin gene causing X-linked
Emery–Dreifuss
[102064]muscular dystrophy. Hum. Mutat. 9: 526–530, 1997.
[102065]17771.
- [102066]Yamada, T.; Kobayashi, T.: A novel emerin mutation in a
Japanese
[102067]patient with Emery–Dreifuss muscular dystrophy. Hum.
Genet. 97:
[102068]693–694, 1996.
[102069]17772.
- [102070]Yates, J. R. W.; Affara, N. A.; Jamieson, D. M.; Ferguson–
Smith,
[102071]M. A.; Hausmanowa–Petrusewicz, I.; Zaremba, J.;
Borkowska, J.; Johnston,
[102072]A. W.; Kelly, K.: Emery–Dreifuss muscular dystrophy: lo-
calisation
[102073]to Xq27.3–qter confirmed by linkage to the factor VIII
gene. J. Med.
[102074]Genet. 23: 587–590, 1986.
[102075]17773.

- [102076]Yates, J. R. W.; Bagshaw, J.; Aksmanovic, V. M. A.;
Coomber, E.;
- [102077]McMahon, R.; Whittaker, J. L.; Morrison, P. J.; Kendrick-
Jones, J.;
- [102078]Ellis, J. A.: Genotype-phenotype analysis in X-linked
Emery-Dreifuss
[102079]muscular dystrophy and identification of a missense mu-
tation associated
[102080]with a milder phenotype. Neuromusc. Disord. 9: 159-165,
1999.
- [102081]17774.
- [102082]Yates, J. R. W.; Warner, J. P.; Smith, J. A.; Deymeer, F.;
Azulay,
[102083]J.-P.; Hausmanowa-Petrusewicz, I.; Zaremba, J.;
Borkowska, J.; Affara,
[102084]N. A.; Ferguson-Smith, M. A.: Emery-Dreifuss muscular
dystrophy:
[102085]linkage to markers in distal Xq28. J. Med. Genet. 30:
108-111, 1993.
- [102086]17775.
- [102087]Yorifuji, H.; Tadano, Y.; Tsuchiya, Y.; Ogawa, M.; Goto, K.;
Umetani,

[102088]A.; Asaka, Y.; Arahata, K.: Emerin, deficiency of which causes Emery–Dreifuss

[102089]muscular dystrophy, is localized at the inner nuclear membrane. Neurogenetics 1:

[102090]135–140, 1997.

[102091]17776.

[102092]Allen, R. C.; Armitage, R. J.; Conley, M. E.; Rosenblatt, H.; Jenkins,

[102093]N. A.; Copeland, N. G.; Bedell, M. A.; Edelhoff, S.; Disteche, C.

[102094]M.; Simoneaux, D. K.; Fanslow, W. C.; Belmont, J.; Spriggs, M. K.

[102095]: CD40 ligand gene defects responsible for X-linked hyper-IgM syndrome. Science 259:

[102096]990–993, 1993.

[102097]17777.

[102098]Aruffo, A.; Farrington, M.; Hollenbaugh, D.; Li, X.; Milatovich,

[102099]A.; Nonoyama, S.; Bajorath, J.; Grosmaire, L. S.; Stenkamp, R.; Neubauer,

[102100]M.; Roberts, R. L.; Noelle, R. J.; Ledbetter, J. A.; Francke, U.;

- [102101] Ochs, H. D.: The CD40 ligand, gp39, is defective in activated T cells
from patients with X-linked hyper-IgM syndrome. Cell 72: 291-300,
1993.
- [102103] 17778.
- [102104] 17778.
- [102105] DiSanto, J. P.; Bonnefoy, J. Y.; Gauchat, J. F.; Fischer, A.; de
Saint Basile, G.: CD40 ligand mutations in X-linked immunodeficiency
with hyper-IgM. Nature 361: 541-543, 1993.
- [102107] 17779.
- [102108] 17779.
- [102109] Gauchat, J.-F.; Aubry, J.-P.; Mazzei, G.; Life, P.; Jomotte, T.;
- [102110] Elson, G.; Bonnefoy, J.-Y.: Human CD40-ligand: molecular cloning,
cellular distribution and regulation of expression by factors controlling
IgE production. FEBS Lett. 315: 259-266, 1993.
- [102112] 17780.
- [102113] 17780.
- [102114] Graf, D.; Korthauer, U.; Mages, H. W.; Senger, G.; Kroczeck, R.

[102115]A.: Cloning of TRAP, a ligand for CD40 on human T cells.
Europ.

[102116]J. Immun. 22: 3191–3194, 1992.

[102117]17781.

[102118]Hollenbaugh, D.; Grosmaire, L. S.; Kullas, C. D.; Chalupny,
N.

[102119]J.; Braesch-Andersen, S.; Noelle, R. J.; Stamenkovic, I.;
Ledbetter,

[102120]J. A.; Aruffo, A.: The human T cell antigen gp39, a mem-
ber of the

[102121]TNF gene family, is a ligand for the CD40 receptor: ex-
pression of

[102122]a soluble form of gp39 with B cell co-stimulatory activity.
EMBO

[102123]J. 11: 4313–4321, 1992.

[102124]17782.

[102125]Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Benne, F.:
Cattle

[102126]gene mapping by somatic cell hybridization. (Abstract)
Cytogenet.

[102127]Cell Genet. 25: 165–166, 1979.

[102128]17783.

[102129]Biamonti, G.; Ruggiu, M.; Saccone, S.; Della Valle, G.; Riva, S.

[102130]: Two homologous genes, originated by duplication, encode the human

[102131]hnRNP proteins A2 and A1. Nucleic Acids Res. 22: 1996–2002, 1994.

[102132]17784.

[102133]Clarke, A.; Burn, J.: Sweat testing to identify female carriers

[102134]of X linked hypohidrotic ectodermal dysplasia. J. Med. Genet. 28:

[102135]330–333, 1991.

[102136]17785.

[102137]Clarke, A.; Sarfarazi, M.; Thomas, N. S. T.; Roberts, K.; Harper,

[102138]P. S.: X-linked hypohidrotic ectodermal dysplasia: DNA probe linkage

[102139]analysis and gene localization. Hum. Genet. 75: 378–380, 1987.

[102140]17786.

[102141]Clarke, A.; Sarfarazi, M.; Zonana, J.; Thomas, N. S. T.;
Roberts,

[102142]K.; Harper, P. S.: Regional localization of X-linked hypo-
hidrotic

[102143]ectodermal dysplasia (EDA). (Abstract) Cytogenet. Cell
Genet. 46:

[102144]594, 1987.

[102145]17787.

[102146]Cohen, M. M.; Lin, C. C.; Sybert, V.; Orecchio, E. J.: Two
human

[102147]X-autosome translocations identified by autoradiography
and fluorescence. Am.

[102148]J. Hum. Genet. 24: 583-597, 1972.

[102149]17788.

[102150]Crawford, P. J. M.; Aldred, M. J.; Clarke, A.: Clinical and ra-
diographic

[102151]dental findings in X linked hypohidrotic ectodermal dys-
plasia. J.

[102152]Med. Genet. 28: 181-185, 1991.

[102153]17789.

[102154]Darwin, C.: The Variation of Animals and Plants under Do-

mestication.

[102155]London: John Murray (pub.) (2nd ed.): 1875. Pp. 319 only.

[102156]17790.

[102157]de la Chapelle, A.: The map of the human X chromosome. In: Crosignani,

[102158]P. G.; Fraccaro, M.; Rubin, B. L.: Genetic Control of Gamete Production

[102159]and Function. New York: Grune and Stratton (pub.) 1982. Pp. 33–48.

[102160]17791.

[102161]de la Chapelle, A.: Personal Communication. Helsinki, Finland

[102162]9/18/1990.

[102163]17792.

[102164]Durmowicz, M. C.; Cui, C.-Y.; Schlessinger, D.: The EDA gene

[102165]is a target of, but does not regulate Wnt signaling. Gene 285: 203–211,

[102166]2002.

[102167]17793.

[102168]Elomaa, O.; Pulkkinen, K.; Hannelius, U.; Mikkola, M.;

Saarialho-Kere,

[102169]U.; Kere, J.: Ectodysplasin is released by proteolytic shedding and

[102170]binds to the EDAR protein. Hum. Molec. Genet. 10: 953–962, 2001.

[102171]17794.

[102172]Ezer, S.; Bayes, M.; Elomaa, O.; Schlessinger, D.; Kere, J.:

[102173]Ectodysplasin is a collagenous trimeric type II membrane protein with

[102174]a tumor necrosis factor-like domain and co-localizes with cytoskeletal

[102175]structures at lateral and apical surfaces of cells. Hum. Molec. Genet. 8:

[102176]2079–2086, 1999.

[102177]17795.

[102178]Familusi, J. B.; Jaiyesimi, F.; Ojo, C. O.; Attah, E. B.: Hereditary

[102179]anhidrotic ectodermal dysplasia: studies in a Nigerian family. Arch.

[102180]Dis. Child. 50: 642–647, 1975.

[102181]17796.

- [102182]Ferguson, B. M.; Brockdorff, N.; Formstone, E.; Ngyuen, T.;
Kronmiller,
[102183]J. E.; Zonana, J.: Cloning of Tabby, the murine homolog of
the human
[102184]EDA gene: evidence for a membrane-associated protein
with a short
[102185]collagenous domain. Hum. Molec. Genet. 6: 1589–1594,
1997.
[102186]17797.
- [102187]Ferguson, B. M.; Thomas, N. S. T.; Munoz, F.; Morgan, D.;
Clarke,
[102188]A.; Zonana, J.: Scarcity of mutations detected in families
with X
[102189]linked hypohidrotic ectodermal dysplasia: diagnostic im-
plications. J.
[102190]Med. Genet. 35: 112–115, 1998.
[102191]17798.
- [102192]Filippi, G.; Rinaldi, A.; Crisponi, G. L.; Siniscalco, M.: X-
mapping
[102193]in man: evidence against measurable linkage between an-
hidrotic ectodermal
[102194]dysplasia and G6PD deficiency. J. Med. Genet. 16:

223-226, 1979.

[102195]17799.

[102196]Fox, L. A.: X-linked anhidrotic ectodermal dysplasia manifesting

[102197]in a female. Birth Defects Orig. Art. Ser. X(4): 319-320, 1974.

[102198]17800.

[102199]Freire-Maia, N.; Pinheiro, M.: So-called 'anhidrotic ectodermal

[102200]dysplasia.'. Int. J. Derm. 19: 455-456, 1980.

[102201]17801.

[102202]Gerald, P. S.; Brown, J. A.: Report of the committee on the genetic

[102203]constitution of the X chromosome. Cytogenet. Cell Genet. 13: 29-34,

[102204]1974.

[102205]17802.

[102206]Gilgenkrantz, S.; Blanchet-Bardon, C.; Nazzaro, V.; Formiga, L.;

[102207]Mujica, P.; Alembik, Y.: Hypohidrotic ectodermal dysplasia: clinical

[102208]study of a family of 30 over three generations. Hum.

Genet. 81:

[102209]120-122, 1989.

[102210]17803.

[102211]Grant, R.; Falls, H. F.: Anodontia: report of a case associated

[102212]with ectodermal dysplasia of the anhidrotic type. Am. J. Orthodont. 30:

[102213]661-672, 1944.

[102214]17804.

[102215]Graves, R.: The Whitaker Negroes. In: Encounters: An anthropology

[102216]from the first ten years of Encounter Magazine.: Spender, S.; Kristol,

[102217]J.; Lasky, M. J., eds. New York: Basic Books, Inc. (pub.) Pp.

[102218]92-105: 1963.

[102219]17805.

[102220]Halperin, S. L.; Curtis, G. M.: Anhidrotic ectodermal dysplasia

[102221]associated with mental deficiency. Am. J. Ment. Defic. 46: 459-463,

[102222]1942.

[102223]17806.

[102224]Hanauer, A.; Alembik, Y.; Arveiler, B.; Formiga, L.;

Gilgenkrantz,

[102225]S.; Mandel, J. L.: Genetic mapping of hypohidrotic ecto-

dermal dysplasia:

[102226]linkage to DXS159 and DXYS1. (Abstract) Cytogenet. Cell

Genet. 46:

[102227]626 only, 1987.

[102228]17807.

[102229]Hanauer, A.; Alembik, Y.; Arveiler, B.; Formiga, L.;

Gilgenkrantz,

[102230]S.; Mandel, J. L.: Genetic mapping of anhidrotic ectoder-

mal dysplasia:

[102231]DXS159, a closely linked proximal marker. Hum. Genet.

80: 177-180,

[102232]1988.

[102233]17808.

[102234]Happle, R.: Association of pigmentary anomalies with

chromosomal

[102235]and genetic mosaicism and chimerism. (Letter) Am. J.

Hum. Genet. 48:

[102236]1013-1014, 1991.

[102237]17809.

[102238]Happle, R.; Frosch, P. J.: Manifestation of the lines of
Blaschko

[102239]in women heterozygous for X-linked hypohidrotic ecto-
dermal dysplasia. Clin.

[102240]Genet. 27: 468-471, 1985.

[102241]17810.

[102242]Harper, P. S.: Personal Communication. Cardiff, Wales
6/24/1986.

[102243]17811.

[102244]Hutt, F. B.: An earlier record of the toothless men of Sind.
J.

[102245]Hered. 26: 65-66, 1935.

[102246]17812.

[102247]Jespersen, H. G.: Hereditary ectodermal dysplasia of an-
hidrotic

[102248]type. Acta Paediat. 51: 712-720, 1962.

[102249]17813.

[102250]Kapalanga, J.; Blecher, S. R.: Effect of the X-linked gene
Tabby

[102251](Ta) on eyelid opening and incisor eruption in neonatal
mice is opposite

[102252]to that of epidermal growth factor. Development 108:
349–355, 1990.

[102253]17814.

[102254]Kere, J.; Srivastava, A. K.; Montonen, O.; Zonana, J.;
Thomas,

[102255]N.; Ferguson, B; Munoz, F.; Morgan, D.; Clarke, A.; Bay-
bayan, P.;

[102256]Chen, E. Y.; Ezer, S.; Saarialho-Kere, U.; de la Chapelle, A.;
Schlessinger,

[102257]D.: X-linked anhidrotic (hypohidrotic) ectodermal dyspla-
sia is caused

[102258]by mutation in a novel transmembrane protein. Nature
Genet. 13:

[102259]409–416, 1996.

[102260]17815.

[102261]Kerr, C. B.; Wells, R. S.; Cooper, K. E.: Gene effect in carri-
ers

[102262]of anhidrotic ectodermal dysplasia. J. Med. Genet. 3:

169–176, 1966.

[102263]17816.

[102264]Kleinebrecht, J.; Degenhardt, K.-H.; Grubisic, A.; Gunther, E.;

[102265]Svejcar, J.: Sweat pore counts in ectodermal dysplasias. Hum. Genet. 57:

[102266]437–439, 1981.

[102267]17817.

[102268]Kline, A. H.; Sidbury, J. B., Jr.; Richter, C. P.: The occurrence

[102269]of ectodermal dysplasia and corneal dysplasia in one family. J. Pediat. 55:

[102270]355–366, 1959.

[102271]17818.

[102272]Kobiela, K.; Kobiela, A.; Roszkiewicz, J.; Wierzba, J.; Limon,

[102273]J.; Trzeciak, W. H.: Mutations in the EDA gene in three unrelated

[102274]families reveal no apparent correlation between phenotype and genotype

[102275]in the patients with an X-linked anhidrotic ectodermal

dysplasia. Am.

[102276]J. Med. Genet. 100: 191–197, 2001.

[102277]17819.

[102278]Stoddart, K. L.; Jermak, C.; Nagaraja, R.; Schlessinger, D.;
Bech–Hansen,

[102279]N. T.: Physical map covering a 2 Mb region in human
Xp11.3 distal

[102280]to DX6849. Gene 227: 111–116, 1999.

[102281]17820.

[102282]Swanson, D. A.; Freund, C. L.; Ploder, L.; McInnes, R. R.;
Valle,

[102283]D.: A ubiquitin C–terminal hydrolase gene on the proximal
short arm

[102284]of the X chromosome: implications for X–linked retinal
disorders. Hum.

[102285]Molec. Genet. 5: 533–538, 1996.

[102286]17821.

[102287]Olinsky, S.; Loop, B. T.; DeKosky, A.; Ripepi, B.; Weng, W.;
Cummins,

[102288]J.; Wenger, S. L.; Yan, Y.; Lagenaur, C.; Narayanan, V.:
Chromosomal

[102289]mapping of the human M6 genes. Genomics 33: 532–536, 1996.

[102290]17822.

[102291]Yan, Y.; Lagenaur, C.; Narayanan, V.: Molecular cloning of M6:

[102292]identification of a PLP/DM20 gene family. Neuron 11: 423–431, 1993.

[102293]17823.

[102294]Ballabio, A.; Parenti, G.; Napolitano, E.; Di Natale, P.; Andria,

[102295]G.: Genetic complementation of steroid sulphatase after somatic cell

[102296]hybridization of X-linked ichthyosis and multiple sulphatase deficiency. Hum.

[102297]Genet. 70: 315–317, 1985.

[102298]17824.

[102299]Nan, X.; Ng, H.–H.; Johnson, C. A.; Laherty, C. D.; Turner, B.

[102300]M.; Eisenman, R. N.; Bird, A.: Transcriptional repression by the

[102301]methyl–CpG-binding protein MeCP2 involves a histone deacetylase complex. Nature 393:

[102302]386-389, 1998.

[102303]17825.

[102304]Nan, X.; Tate, P.; Li, E.; Bird, A.: DNA methylation specifies

[102305]chromosomal localization of MeCP2. Molec. Cell. Biol. 16:
414-421,

[102306]1996.

[102307]17826.

[102308]Arn, P.; Hauser, B.; Maestri, N.; Herman, G.; Thomas, G.
H.; Brusilow,

[102309]S. W.: Hyperammonemic coma in adult females heterozy-
gous at the ornithine

[102310]transcarbamylase locus (OTCDH). (Abstract) Am. J. Hum.
Genet. 45

[102311](suppl.): A3, 1989.

[102312]17827.

[102313]Andersson, M.; Page, D. C.; Pettay, D.; Subrt, I.; Turleau,
C.;

[102314]de Grouchy, J.; de la Chapelle, A.: Y;autosome transloca-
tions and

[102315]mosaicism in the aetiology of 45,X maleness: assignment
of fertility

[102316]factor to distal Yq11. Hum. Genet. 79: 2–7, 1988.

[102317]17828.

[102318]Chandley, A. C.; Gosden, J. R.; Hargreave, T. B.; Spowart, G.;

[102319]Speed, R. M.; McBeath, S.: Deleted Yq in the sterile son of a man

[102320]with a satellited Y chromosome (Yqs). J. Med. Genet. 26: 145–153,

[102321]1989.

[102322]17829.

[102323]Robledo, R.; Melis, P.; Schillinger, E.; Casciano, I.; Balazs,

[102324]I.; Rinaldi, A.; Siniscalco, M.; Filippi, G.: X-linked ichthyosis

[102325]without STS deficiency: clinical, genetical, and molecular studies. Am.

[102326]J, Med. Genet. 59: 143–148, 1995.

[102327]17830.

[102328]Franco, B.; Meroni, G.; Parenti, G.; Levilliers, J.; Bernard, L.;

[102329]Gebbia, M.; Cox, L.; Maroteaux, P.; Sheffield, L.; Rappold, G. A.;

[102330] Andria, G.; Petit, C.; Ballabio, A.: A cluster of sulfatase genes

[102331] on Xp22.3: mutations in chondrodysplasia punctata (CDPX) and implications

[102332] for warfarin embryopathy. Cell 81: 1–20, 1995.

[102333] 17831.

[102334] Meroni, G.; Franco, B.; Archidiacono, N.; Messali, S.; Andolfi,

[102335] G.; Rocchi, M.; Ballabio, A.: Characterization of a cluster of sulfatase

[102336] genes on Xp22.3 suggests gene duplications in an ancestral pseudoautosomal

[102337] region. Hum. Molec. Genet. 5: 423–431, 1996.

[102338] 17832.

[102339] Franco, B.; Meroni, G.; Parenti, G.; Levilliers, J.; Bernard, L.;

[102340] Gebbia, M.; Cox, L.; Maroteaux, P.; Sheffield, L.; Rappold, G. A.;

[102341] Andria, G.; Petit, C.; Ballabio, A.: A cluster of sulfatase genes

[102342] on Xp22.3: mutations in chondrodysplasia punctata (CDPX) and implications

[102343]for warfarin embryopathy. Cell 81: 15–20, 1995.

[102344]17833.

[102345]Puca, A. A.; Zollo, M.; Repetto, M.; Andolfi, G.; Guffanti, A.;

[102346]Simon, G.; Ballabio, A.; Franco, B.: Identification by shot–
gun sequencing,

[102347]genomic organization, and functional analysis of a fourth
arylsulfatase

[102348]gene (ARSF) from the Xp22.3 region. Genomics 42:
192–199, 1997.

[102349]17834.

[102350]Adler, D. A.; Quaderi, N. A.; Brown, S. D. M.; Chapman, V.
M.;

[102351]Moore, J.; Tate, P.; Disteché, C. M.: The X-linked methy–
lated DNA

[102352]binding protein, Mecp2, is subject to X inactivation in the
mouse. Mammalian

[102353]Genome 6: 491–492, 1995.

[102354]17835.

[102355]Amano, K.; Nomura, Y.; Segawa, M.; Yamakawa, K.: Muta–
tional analysis

[102356]of the MECP2 gene in Japanese patients with Rett syn–

drome. J. Hum.

[102357]Genet. 45: 231–236, 2000.

[102358]17836.

[102359]Amir, R. E.; Van den Veyver, I. B.; Wan, M.; Tran, C. Q.;
Francke,

[102360]U.; Zoghbi, H. Y.: Rett syndrome is caused by mutations in
X-linked

[102361]MECP2, encoding methyl–CpG–binding protein 2. Nature
Genet. 23:

[102362]185–188, 1999.

[102363]17837.

[102364]Ballestar, E.; Yusufzai, T. M.; Wolffe, A. P.: Effects of Rett

[102365]syndrome mutations of the methyl–CpG binding domain
of the transcriptional

[102366]repressor MeCP2 on selectivity for association with
methylated DNA. Biochemistry 39:

[102367]7100–7106, 2000.

[102368]17838.

[102369]Balmer, D.; Arredondo, J.; Samaco, R. C.; LaSalle, J. M.:
MECP2

[102370]mutations in Rett syndrome adversely affect lymphocyte

growth, but

[102371]do not affect imprinted gene expression in blood or brain.

Hum. Genet. 110:

[102372]545–552, 2002.

[102373]17839.

[102374]Bienvenu, T.; Carrie, A.; de Roux, N.; Vinet, M.-C.; Jon-
veaux,

[102375]P.; Couvert, P.; Villard, L.; Arzimanoglou, A.; Beldjord, C.;
Fontes,

[102376]M.; Tardieu, M.; Chelly, J.: MECP2 mutations account for
most cases

[102377]of typical forms of Rett syndrome. Hum. Molec. Genet. 9:
1377–1384,

[102378]2000.

[102379]17840.

[102380]van de Vosse, E.; Walpole, S. M.; Nicolaou, A.; van der
Bent, P.;

[102381]Cahn, A.; Vaudin, M.; Ross, M. T.; Durham, J.; Pavitt, R.;
Wilkinson,

[102382]J.; Grafham, D.; Bergen, A. A. B.; van Ommen, G.-J. B.;
Yates, J.

[102383]R. W.; den Dunnen, J. T.; Trump, D.: Characterization of

SCML1, a

[102384]new gene in Xp22, with homology to developmental poly-comb genes. Genomics 49:

[102385]96–102, 1998.

[102386]17841.

[102387]Ballabio, A.; Bardoni, B.; Carrozzo, R.; Andria, G.; Bick, D.;

[102388]Campbell, L.; Hamel, B.; Ferguson-Smith, M. A.; Gimelli, G.; Fraccaro,

[102389]M.; Maraschio, P.; Zuffardi, O.; Guioli, S.; Camerino, G.: Contiguous

[102390]gene syndromes due to deletions in the distal short arm of the human

[102391]X chromosome. Proc. Nat. Acad. Sci. 86: 10001–10005, 1989.

[102392]17842.

[102393]Fujikawa-Adachi, K.; Nishimori, I.; Taguchi, T.; Onishi, S.: Human

[102394]mitochondrial carbonic anhydrase VB: cDNA cloning, mRNA expression,

[102395]subcellular localization, and mapping to chromosome X. J. Biol. Chem. 274:

[102396]21228–21233, 1999.

[102397]17843.

[102398]Shah, G. N.; Hewett-Emmett, D.; Grubb, J. H.; Migas, M. C.; Fleming,

[102399]R. E.; Waheed, A.; Sly, W. S.: Mitochondrial carbonic anhydrase CA

[102400]VB: differences in tissue distribution and pattern of evolution from

[102401]those of CA VA suggest distinct physiological roles. Proc. Nat. Acad.

[102402]Sci. 97: 1677-1682, 2000.

[102403]17844.

[102404]Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayashi, Y.;

[102405]Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura, N.: Prediction

[102406]of the coding sequences of unidentified human genes. VI. The coding

[102407]sequences of 80 new genes (KIAA0201-KIAA0280) deduced by analysis

[102408]of cDNA clones from cell line KG-1 and brain. DNA Res. 3: 321-329,

[102409]1996. Note: Supplement: DNA Res. 3: 341-354, 1996.

[102410]17845.

[102411]Bowling, F.; McGown, I.; McGill, J.; Cowley, D.; Tuchman, M.:

[102412]Maternal gonadal mosaicism causing ornithine transcarbamylase deficiency. Am.

[102413]J. Med. Genet. 85: 452-454, 1999.

[102414]17846.

[102415]Bruton, C. J.; Corsellia, J. A. N.; Russell, A.: Hereditary hyperammonemia. Brain 93:

[102416]423-434, 1970.

[102417]17847.

[102418]Burdakin, J. H.; Norum, R. A.: Recombination between loci for

[102419]ornithine transcarbamylase (OTC) deficiency and G6PD. (Abstract) Am.

[102420]J. Hum. Genet. 33: 38A, 1981.

[102421]17848.

[102422]Asao, H.; Okuyama, C.; Kumaki, S.; Ishii, N.; Tsuchiya, S.; Foster,

[102423]D.; Sugamura, K.: Cutting edge: the common gamma-chain is an indispensable

[102424]subunit of the IL-21 receptor complex. J. Immun. 167:
1-5, 2001.

[102425]17849.

[102426]Arn, P. H.; Hauser, E. R.; Thomas, G. H.; Herman, G.; Hess,
D.;

[102427]Brusilow, S. W.: Hyperammonemia in women with a muta-
tion at the ornithine

[102428]carbamoyltransferase locus: a cause of postpartum coma.
New Eng.

[102429]J. Med. 322: 1652-1655, 1990.

[102430]17850.

[102431]Batshaw, M. L.; Roan, Y.; Jung, A. L.; Rosenberg, L. A.;
Brusilow,

[102432]S. W.: Cerebral dysfunction in asymptomatic carriers of
ornithine

[102433]transcarbamylase deficiency. New Eng. J. Med. 302:
482-485, 1980.

[102434]17851.

[102435]Bonaiti-Pellie, C.; Pelet, A.; Ogier, H.; Nelson, J.-R.;
Largilliere,

[102436]C.; Berthelot, J.; Saudubray, J.-M.; Munnich, A.: A probable
sex

[102437] difference in mutation rates in ornithine transcarbamylase deficiency. Hum.

[102438] Genet. 84: 163–166, 1990.

[102439] 17852.

[102440] Campbell, A. G. M.; Rosenberg, L. E.; Snodgrass, P. J.;
Nuzum,

[102441] C. T.: Lethal neonatal hyperammonaemia due to complete
ornithine–transcarbamylase

[102442] deficiency. (Letter) Lancet II: 217–218, 1971.

[102443] 17853.

[102444] Campbell, A. G. M.; Rosenberg, L. E.; Snodgrass, P. J.;
Nuzum,

[102445] C. T.: Ornithine transcarbamylase deficiency: a cause of
lethal neonatal

[102446] hyperammonemia in males. New Eng. J. Med. 288: 1–6,
1973.

[102447] 17854.

[102448] Carstens, R. P.; Fenton, W. A.; Rosenberg, L. R.: Identifica-
tion

[102449] of RNA splicing errors resulting in human ornithine tran-
scarbamylase

[102450]deficiency. Am. J. Hum. Genet. 48: 1105–1114, 1991.

[102451]17855.

[102452]Cathelineau, L.; Saudubray, J.–M.; Polonovski, C.: Hetero-
geneous

[102453]mutations of the structural gene of human ornithine car-
bamyltransferase

[102454]as observed in five personal cases. Enzyme 18: 103–113,
1974.

[102455]17856.

[102456]Cavard, C.; Grimber, G.; Dubois, N.; Chasse, J.–F.; Ben-
noun, M.;

[102457]Minet–Thuriaux, M.; Kamoun, P.; Briand, P.: Correction of
mouse ornithine

[102458]transcarbamylase deficiency by gene transfer into the
germ line. Nucleic

[102459]Acids Res. 16: 2099–2110, 1988.

[102460]17857.

[102461]Clayton–Smith, J.; Donnai, D.: Fetal valproate syndrome. J.

[102462]Med. Genet. 32: 724–727, 1995.

[102463]17858.

[102464]Corey, L.; Rubin, R. J.; Hattwick, M. A. W.; Noble, G. R.;
Cassidy,

[102465]E.: A nationwide outbreak of Reye's syndrome: its epidemiologic relationship

[102466]to influenza B. Am. J. Med. 61: 615-625, 1976.

[102467]17859.

[102468]DeMars, R.; LeVan, S. L.; Trend, B. L.; Russell, L. B.: Abnormal

[102469]ornithine carbamyltransferase in mice having the sparse-fur mutation. Proc.

[102470]Nat. Acad. Sci. 73: 1693-1697, 1976.

[102471]17860.

[102472]Drogari, E.; Leonard, J. V.: Late onset ornithine carbamoyl transferase

[102473]deficiency in males. Arch. Dis. Child. 63: 1363-1367, 1988.

[102474]17861.

[102475]Faiella, A.; Wernig, M.; Consalez, G. G.; Hostick, U.; Hofmann,

[102476]C.; Hustert, E.; Boncinelli, E.; Balling, R.; Nadeau, J. H.: A mouse

- [102477]model for valproate teratogenicity: parental effects,
homeotic transformations,
[102478]and altered HOX expression. Hum. Molec. Genet. 9:
227–236, 2000.
[102479]17862.
- [102480]Feldmann, D.; Rozet, J.–M.; Pelet, A.; Hentzen, D.; Briand,
P.;
[102481]Hubert, P.; Largilliere, C.; Rabier, D.; Farriaux, J.–P.;
Munnich,
[102482]A.: Site specific screening for point mutations in ornithine
transcarbamylase
[102483]deficiency. J. Med. Genet. 29: 471–475, 1992.
[102484]17863.
- [102485]Finkelstein, J. E.; Francomano, C. A.; Brusilow, S. W.;
Traystman,
[102486]M. D.: Use of denaturing gradient gel electrophoresis for
detection
[102487]of mutation and prospective diagnosis in late onset or–
nithine transcarbamylase
[102488]deficiency. Genomics 7: 167–172, 1990.
[102489]17864.

[102490]Finkelstein, J. E.; Hauser, E.; Brusilow, S. W.: Late onset ornithine

[102491]transcarbamylase deficiency (OTCD) in males. (Abstract)
Am. J. Hum.

[102492]Genet. 45 (suppl.): A5, 1989.

[102493]17865.

[102494]Finkelstein, J. E.; Hauser, E. R.; Leonard, C. O.; Brusilow, S.

[102495]W.: Late-onset ornithine transcarbamylase deficiency in
male patients. J.

[102496]Pediat. 117: 897-902, 1990.

[102497]17866.

[102498]Fox, J. E.; Hack, A. M.; Fenton, W. A.; Golbus, M. S.; Winter,

[102499]S.; Kalousek, F.; Rozen, R.; Brusilow, S. W.; Rosenberg, L.
E.: Prenatal

[102500]diagnosis of ornithine transcarbamylase deficiency with
use of DNA

[102501]polymorphisms. New Eng. J. Med. 315: 1205-1208, 1986.

[102502]17867.

[102503]Fox, J. E.; Hack, A. M.; Fenton, W. A.; Rosenberg, L. E.:
Identification

[102504]and application of additional restriction fragment length polymorphisms

[102505]at the human ornithine transcarbamylase locus. Am. J. Hum. Genet. 38:

[102506]841–847, 1986.

[102507]17868.

[102508]Garcia-Perez, M. A.; Sanjurjo, P.; Rubio, V.: Demonstration of

[102509]the spf-ash mutation in Spanish patients with ornithine transcarbamylase

[102510]deficiency of moderate severity. Hum. Genet. 95: 183–186, 1995.

[102511]17869.

[102512]Gelehrter, T. D.; Rosenberg, L. E.: Ornithine transcarbamylase

[102513]deficiency: unsuccessful therapy of neonatal hyperammonemia with N-carbamyl-L-glutamate

[102514]and L-arginine. New Eng. J. Med. 292: 351–352, 1975.

[102515]17870.

[102516]Genet, S.; Cranston, T.; Middleton-Price, H. R.: Mutation detection

[102517]in 65 families with a possible diagnosis of ornithine carbamoyltransferase
[102518]deficiency including 14 novel mutations. J. Inherit. Metab. Dis. 23:
[102519]669–676, 2000.
[102520]17871.
[102521]Gilbert–Dussardier, B.; Segues, B.; Rozet, J.–M.; Rabier, D.;
[102522]Calvas, P.; de Lumley, L.; Bonnefond, J.–P.; Munnich, A.:
Partial
[102523]duplication [dup. TCAC (178)] and novel point mutations
(T125M, G188R,
[102524]A209V, and H302L) of the ornithine transcarbamylase
gene in congenital
[102525]hyperammonemia. Hum. Mutat. 8: 74–76, 1996.
[102526]17872.
[102527]Gilchrist, J. M.; Coleman, R. A.: Ornithine transcarbamylase
[102528]deficiency: adult onset of severe symptoms. Ann. Intern. Med. 106:
[102529]556–558, 1987.
[102530]17873.

- [102531]Glick, T. H.; Likosky, W. H.; Levitt, L. P.; Mellin, H.;
Reynolds,
[102532]D. W.: Reye's syndrome: an epidemiologic approach. *Pedi-*
iatrics 46:
[102533]371–377, 1970.
[102534]17874.
- [102535]Grompe, M.; Caskey, C. T.; Fenwick, R. G.: Improved
molecular
[102536]diagnostics for ornithine transcarbamylase deficiency. *Am.*
J. Hum.
[102537]*Genet.* 48: 212–222, 1991.
[102538]17875.
- [102539]Grompe, M.; Muzny, D. M.; Caskey, C. T.: Scanning detec-
tions
[102540]of mutations in human ornithine transcarbamylase defi-
ciency (OTC)
[102541]by chemical mismatch cleavage. (Abstract) *Am. J. Hum.*
Genet. 45
[102542](suppl.): A192, 1989.
[102543]17876.
- [102544]Ogata, T.; Wakui, K.; Muroya, K.; Ohashi, H.; Matsuo, N.;

Brown,

[102545]D. M.; Ishii, T.; Fukushima, Y.: Microphthalmia with linear skin

[102546]defects syndrome in a mosaic female infant with monosomy for the Xp22

[102547]region: molecular analysis of the Xp22 breakpoint and the X-inactivation

[102548]pattern. Hum. Genet. 103: 51-56, 1998.

[102549]17877.

[102550]Lankes, W.; Griesmacher, A.; Grunwald, J.; Schwartz-Albiez, R.;

[102551]Keller, R.: A heparin-binding protein involved in inhibition of smooth-muscle

[102552]cell proliferation. Biochem. J. 251: 831-842, 1988.

[102553]17878.

[102554]Lankes, W. T.; Furthmayr, H.: Moesin: a member of the protein

[102555]4.1-talin-ezrin family of proteins. Proc. Nat. Acad. Sci. 88: 8297-8301,

[102556]1991.

[102557]17879.

- [102558] Shcherbina, A.; Bretscher, A.; Rosen, F. S.; Kenney, D. M.; Remold-O'Donnell, E.: The cytoskeletal linker protein moesin: decreased levels in Wiskott-Aldrich syndrome platelets and identification of a cleavage pathway in normal platelets. *Brit. J. Haemat.* 106: 216-223, 1999.
- [102562] 17880.
- [102563] Wilgenbus, K. K.; Hsieh, C.-L.; Lankes, W. T.; Milatovich, A.; Francke, U.; Furthmayr, H.: Structure and localization on the X chromosome of the gene coding for the human filopodial protein moesin (MSN). *Genomics* 19: 326-333, 1994.
- [102567] 17881.
- [102568] Deng, L.; Wang, C.; Spencer, E.; Yang, L.; Braun, A.; You, J.; Slaughter, C.; Pickart, C.; Chen, Z. J.: Activation of the I-kappa-B complex by TRAF6 requires a dimeric ubiquitin-conjugating enzyme complex

[102571]and a unique polyubiquitin chain. Cell 103: 351–361,
2000.

[102572]17882.

[102573]Nakano, H.; Oshima, H.; Chung, W.; Williams–Abbott, L.;
Ware, C.

[102574]F.; Yagita, H.; Okumura, K.: TRAF5, an activator of NF–
kappaB and

[102575]putative signal transducer for the lymphotoxin–beta re–
ceptor. J.

[102576]Biol. Chem. 271: 14661–14664, 1996.

[102577]17883.

[102578]Nakano, H.; Sakon, S.; Koseki, H.; Takemori, T.; Tada, K.;
Matsumoto,

[102579]M.; Munechika, E.; Sakai, T.; Shirasawa, T.; Akiba, H; Ko–
bata, T.;

[102580]Santee, S. M.; Ware, C. F.; Renner, P. D.; Taniguchi, M.;
Yagita,

[102581]H.; Okumura, K.: Targeted disruption of Traf5 gene causes
defects

[102582]in CD40– and CD27–mediated lymphocyte activation.
Proc. Nat. Acad.

[102583]Sci. 96: 9803–9808, 1999.

[102584]17884.

[102585]Nakano, H.; Shindo, M.; Yamada, K.; Yoshida, M. C.; Santee, S.

[102586]M.; Ware, C. F.; Jenkins, N. A.; Gilbert, D. J.; Yagita, H.; Copeland,

[102587]N. G.; Okumura, K.: Human TNF receptor-associated factor 5 (TRAF5):

[102588]cDNA cloning, expression and assignment of the TRAF5 gene to chromosome

[102589]1q32. Genomics 42: 26-32, 1997.

[102590]17885.

[102591]Augarten, A.; Kerem, B.-S.; Yahav, Y.; Noiman, S.; Rivlin, Y.;

[102592]Tal, A.; Blau, H.; Ben-Tur, L.; Szeinberg, A.; Kerem, E.; Gazit, E.

[102593]: Mild cystic fibrosis and normal or borderline sweat test in patients

[102594]with the 3849 + 10 kb C-to-T mutation. Lancet 342: 25-26, 1993.

[102595]17886.

[102596]Bal, J.; Stuhmann, M.; Schloesser, M.; Schmidtke, J.; Reiss, J.

- [102597]: A cystic fibrosis patient homozygous for the nonsense mutation R553X. J.
- [102598]Med. Genet. 28: 715–717, 1991.
- [102599]17887.
- [102600]Ballabio, A.; Gibbs, R. A.; Caskey, C. T.: PCR test for cystic
- [102601]fibrosis deletion. (Letter) Nature 343: 220 only, 1990.
- [102602]17888.
- [102603]Baylin, S. B.; Rosenstein, B. J.; Marton, L. J.; Lockwood, D.
- [102604]H.: Age-related abnormalities of circulating polyamines and diamine
- [102605]oxidase activity in cystic fibrosis heterozygotes and homozygotes. Pediat.
- [102606]Res. 14: 921–925, 1980.
- [102607]17889.
- [102608]Bienvenu, T.; Beldjord, C.; Adjiman, M.; Kaplan, J. C.: Male
- [102609]infertility as the only presenting sign of cystic fibrosis when homozygous
- [102610]for the mild mutation R117H. (Letter) J. Med. Genet. 30: 797 only,
- [102611]1993.

[102612]17890.

[102613]Bobadilla, J. L.; Macek, M., Jr.; Fine, J. P.; Farrell, P. M.

[102614]: Cystic fibrosis: a worldwide analysis of CFTR mutations–
–correlation

[102615]with incidence data and application to screening. Hum.
Mutat. 19:

[102616]575–606, 2002.

[102617]17891.

[102618]Broackes–Carter, F. C.; Mouchel, N.; Gill, D.; Hyde, S.; Bas–
sett,

[102619]J.; Harris, A.: Temporal regulation of CFTR expression
during ovine

[102620]lung development: implications for CF gene therapy. Hum.
Molec. Genet. 11:

[102621]125–131, 2002.

[102622]17892.

[102623]Brown, C. R.; Hong–Brown, L. Q.; Biwersi, J.; Verkman, A.
S.;

[102624]Welsh, W. J.: Chemical chaperones correct the mutant
phenotype of

[102625]the deltaF508 cystic fibrosis transmembrane conductance

regulator

[102626]protein. Cell Stress Chaperones 1: 117–125, 1996.

[102627]17893.

[102628]Burger, J.; Macek, M., Jr.; Stuhmann, M.; Reis, A.;

Krawczak,

[102629]M.; Schmidtke, J.: Genetic influences in the formation of

nasal polyps.

[102630](Letter) Lancet 337: 974, 1991.

[102631]17894.

[102632]Callen, A.; Diener–West, M.; Zeitlin, P. L.; Rubenstein, R. C.

[102633]: A simplified cyclic adenosine monophosphate–mediated

sweat rate

[102634]test for quantitative measure of cystic fibrosis transmem–

brane regulator

[102635](CFTR) function. J. Pediat. 137: 849–855, 2000.

[102636]17895.

[102637]Casals, T.; Ramos, M. D.; Gimenez, J.; Larriba, S.; Nunes,

V.;

[102638]Estivill, X.: High heterogeneity for cystic fibrosis in Span–

ish families

[102639]: 75 mutations account for 90% of chromosomes. Hum.

Genet. 101:

[102640]365–370, 1997.

[102641]17896.

[102642]Casals, T.; Vazquez, C.; Lazaro, C.; Girbau, E.; Gimenez, F.
J.;

[102643]Estivill, X.: Cystic fibrosis in the Basque country: high frequency

[102644]of mutation $\Delta F508$ in patients of Basque origin. Am. J.
Hum. Genet. 50:

[102645]404–410, 1992.

[102646]17897.

[102647]Castaldo, G.; Rippa, E.; Salvatore, D.; Sibillo, R.; Raia, V.;

[102648]de Ritis, G.; Salvatore, F.: Severe liver impairment in a cystic

[102649]fibrosis-affected child homozygous for the G542X mutation. Am. J.

[102650]Med. Genet. 69: 155–158, 1997.

[102651]17898.

[102652]Chalkley, G.; Harris, A.: A cystic fibrosis patient who is homozygous

[102653]for the G85E mutation has very mild disease. J. Med.

Genet. 28:

[102654]875–877, 1991.

[102655]17899.

[102656]Chalkley, G.; Harris, A.: Lymphocyte mRNA as a resource
for detection

[102657]of mutations and polymorphism in the CF gene. J. Med.

Genet. 28:

[102658]777–780, 1991.

[102659]17900.

[102660]Chang, X.; Cui, L.; Hou, Y.; Jensen, T. J.; Aleksandrov, A.
A.;

[102661]Mengos, A.; Riordan, J. R.: Removal of multiple arginine-
framed trafficking

[102662]signals overcomes misprocessing of delta-F508 CFTR
present in most

[102663]patients with cystic fibrosis. Molec. Cell 4: 137–142,
1999.

[102664]17901.

[102665]Chanson, M.; Scerri, I.; Suter, S.: Defective regulation of
gap

[102666]junctional coupling in cystic fibrosis pancreatic duct cells.

J.

[102667]Clin. Invest. 103: 1677–1684, 1999.

[102668]17902.

[102669]Chehab, F. F.; Johnson, J.; Louie, E.; Goossens, M.;
Kawasaki,

[102670]E.; Erlich, H.: A dimorphic 4-bp repeat in the cystic fibro-
sis gene

[102671]is in absolute disequilibrium with the delta-F508 muta-
tion: implications

[102672]for prenatal diagnosis and mutation origin. Am. J. Hum.
Genet. 48:

[102673]223–226, 1991.

[102674]17903.

[102675]Cheng, S. H.; Gregory, R. J.; Marshall, J.; Paul, S.; Souza, D.

[102676]W.; White, G. A.; O'Riordan, C. R.; Smith, A. E.: Defective
intracellular

[102677]transport and processing of CFTR is the molecular basis of
most cystic

[102678]fibrosis. Cell 63: 827–834, 1990.

[102679]17904.

[102680]Chillon, M.; Casals, T.; Gimenez, J.; Nunes, V.; Estivill, X.

- [102681]: A cystic fibrosis patient homozygous for the new
frameshift mutation
- [102682]936delTA: description and clinical data. J. Med. Genet. 31:
369–370,
[102683]1994.
[102684]17905.
- [102685]Choi, J. Y.; Muallem, D.; Kiselyov, K.; Lee, M. G.; Thomas,
P.
[102686]J.; Muallem, S.: Aberrant CFTR-dependent HCO(–3) trans-
port in mutations
[102687]associated with cystic fibrosis. Nature 410: 94–97, 2001.
[102688]17906.
- [102689]Chu, C.–S.; Trapnell, B. C.; Curristin, S.; Cutting, G. R.;
Crystal,
[102690]R. G.: Genetic basis of variable exon 9 skipping in cystic
fibrosis
[102691]transmembrane conductance regulator mRNA. Nature
Genet. 3: 151–156,
[102692]1993.
[102693]17907.
- [102694]Yost, C.; Farr, G. H., III; Pierce, S. B.; Ferkey, D. M.; Chen,

[102695]M. M.; Kimelman, D.: GBP, an inhibitor of GSK-3, is implicated in

[102696]Xenopus development and oncogenesis. Cell 93:
1031-1041, 1998.

[102697]17908.

[102698]Fernandez-Valle, C.; Tang, Y.; Ricard, J.; Rodenas-Ruano,
A.; Taylor,

[102699]A.; Hackler, E.; Biggerstaff, J.; Iacovelli, J.: Paxillin binds
schwannomin

[102700]and regulates its density-dependent localization and effect on cell

[102701]morphology. Nature Genet. 31: 354-362, 2002.

[102702]17909.

[102703]Glenney, J. R., Jr.; Zokas, L.: Novel tyrosine kinase substrates

[102704]from Rous sarcoma virus-transformed cells are present in
the membrane

[102705]skeleton. J. Cell Biol. 108: 2401-2408, 1989.

[102706]17910.

[102707]Mazaki, Y.; Hashimoto, S.; Sabe, H.: Monocyte cells and
cancer

[102708]cells express novel paxillin isoforms with different binding properties

[102709]to focal adhesion proteins. J. Biol. Chem. 272: 7437–7444, 1997.

[102710]17911.

[102711]Salgia, R.; Li, J.-L.; Lo, S. H.; Brunkhorst, B.; Kansas, G. S.;

[102712]Sobhany, E. S.; Sun, Y.; Pisick, E.; Hallek, M.; Ernst, T.; Tantravahi,

[102713]R.; Chen, L. B.; Griffin, J. D.: Molecular cloning of human paxillin,

[102714]a focal adhesion protein phosphorylated by P210(BCR/ABL). J. Biol.

[102715]Chem. 270: 5039–5047, 1995.

[102716]17912.

[102717]Turner, C. E.; Glenney, J. R., Jr.; Burridge, K.: Paxillin: a

[102718]new vinculin-binding protein present in focal adhesions. J. Cell

[102719]Biol. 111: 1059–1068, 1990.

[102720]17913.

[102721]Nguyen, V. T.; Kiss, T.; Michels, A. A.; Bensaude, O.: 7SK small

[102722]nuclear RNA binds to and inhibits the activity of
CDK9/cyclin T complexes. Nature 414:

[102723]322–325, 2001.

[102724]17914.

[102725]Peng, J.; Zhu, Y.; Milton, J. T.; Price, D. H.: Identification

[102726]of multiple cyclin subunits of human P-TEFb. Genes Dev.
12: 755–762,

[102727]1998.

[102728]17915.

[102729]Yang, Z.; Zhu, Q.; Luo, K.; Zhou, Q.: The 7SK small nuclear
RNA

[102730]inhibits the CDK9/cyclin T1 kinase to control transcrip-
tion. Nature 414:

[102731]317–322, 2001.

[102732]17916.

[102733]Adachi, H.; Tsujimoto, M.; Hattori, M.; Arai, H.; Inoue, K.:
Differential

[102734]tissue distribution of the beta- and gamma-subunits of
human cytosolic

[102735]platelet-activating factor acetylhydrolase (isoform 1).
Biochem.

[102736]Biophys. Res. Commun. 233: 10–13, 1997.

[102737]17917.

[102738]Moro, F.; Arrigo, G.; Fogli, A.; Bernard, L.; Carrozzo, R.:

The

[102739]beta and gamma subunits of the human platelet-activating factor acetyl

[102740]hydrolase isoform Ib (PAFAH1B2 and PAFAH1B3) map to chromosome 11q23

[102741]and 19q13.1, respectively. Genomics 51: 157–159, 1998.

[102742]17918.

[102743]Daigo, Y.; Isomura, M.; Nishiwaki, T.; Tamari, M.;

Ishikawa, S.;

[102744]Kai, M.; Murata, Y.; Takeuchi, K.; Yamane, Y.; Hayashi, R.; Minami,

[102745]M.; Fujino, M. A.; Hojo, Y.; Uchiyama, I.; Takagi, T.; Nakamura, Y.

[102746]: Characterization of a 1200-kb genomic segment of chromosome 3p22–p21.3. DNA

[102747]Res. 6: 37–44, 1999.

[102748]17919.

[102749]Erlich, R.; Gleeson, P. A.; Campbell, P.; Dietzsch, E.; Toh,

B.-H.

[102750] Molecular characterization of trans-Golgi p230: a human peripheral

[102751] membrane protein encoded by a gene on chromosome 6p12-22 contains

[102752] extensive coiled-coil alpha-helical domains and a granin motif. J.

[102753] Biol. Chem. 271: 8328-8337, 1996.

[102754] 17920.

[102755] Barr, F. A.; Nakamura, N.; Warren, G.: Mapping the interaction

[102756] between GRASP65 and GM130, components of a protein complex involved

[102757] in the stacking of Golgi cisternae. EMBO J. 17: 3258-3268, 1998.

[102758] 17921.

[102759] Boguski, M. S.; Lowe, T. M. J.; Tolstocher, C. M.: dbEST:database

[102760] for 'expressed sequence tags.' (Letter) Nature Genet. 4: 332-333,

[102761] 1993.

[102762] 17922.

[102763]Izón, D. J.; Aster, J. C.; He, Y.; Weng, A.; Karnell, F. G.; Patriub,

[102764]V.; Xu, L.; Bakkour, S.; Rodriguez, C.; Allman, D.; Pear, W. S.:

[102765]Deltex1 redirects lymphoid progenitors to the B cell lineage by antagonizing

[102766]Notch1. *Immunity* 16: 231–243, 2002.

[102767]17923.

[102768]Lee, L.; Dowhanick–Morrissette, J.; Katz, A.; Jukofsky, L.; Krantz,

[102769]J. D.: Chromosomal localization, genomic characterization, and mapping

[102770]to the Noonan syndrome critical region of the human Deltex (DTX1)

[102771]gene. *Hum. Genet.* 107: 577–581, 2000.

[102772]17924.

[102773]Matsuno, K.; Eastman, D.; Mitsiades, T.; Quinn, A. M.; Cancian,

[102774]M. L.; Ordentlich, P.; Kadesch, T.; Artavanis–Tsakonas, S.: Human

[102775]deltex is a conserved regulator of Notch signalling. *Nature Genet.* 19:

[102776]74–78, 1998.

[102777]17925.

[102778]Chen, J. J.; Reid, C. E.; Band, V.; Androphy, E. J.: Interaction

[102779]of papillomavirus E6 oncoproteins with a putative calcium-binding

[102780]protein. *Science* 269: 529–531, 1995.

[102781]17926.

[102782]Imai, T.; Matsuda, K.; Shimojima, T.; Hashimoto, T.; Masuhiro,

[102783]Y.; Kitamoto, T.; Sugita, A.; Suzuki, K.; Matsumoto, H.; Masushige,

[102784]S.; Nogi, Y.; Muramatsu, M.; Handa, H.; Kato, S.: ERC-55, a binding

[102785]protein for the papilloma virus E6 oncoprotein, specifically interacts

[102786]with vitamin D receptor among nuclear receptors.

Biochem. Biophys.

[102787]*Res. Commun.* 233: 765–769, 1997.

[102788]17927.

[102789]Wang, J. Y.; Zhen, D. K.; Bianchi, D. W.; Androphy, E. J.; Chen,

[102790]J. J.: Assignment of the gene for ERC-55 (RCN2) to human chromosome

[102791]band 15q22.33-q24.1 by in situ hybridization. Cytogenet. Cell Genet. 79:

[102792]60-61, 1997.

[102793]17928.

[102794]Weis, K.; Griffiths, G.; Lamond, A. I.: The endoplasmic reticulum

[102795]calcium-binding protein of 55 kDa is a novel EF-hand protein retained

[102796]in the endoplasmic reticulum by a carboxyl-terminal His-Asp-Glu-Leu

[102797]motif. J. Biol. Chem. 269: 19142-19150, 1994.

[102798]17929.

[102799]Chan, A. M.-L.; Chedid, M.; McGovern, E. S.; Popescu, N. C.; Miki,

[102800]T.; Aaronson, S. A.: Expression cDNA cloning of a serine kinase transforming

[102801]gene. Oncogene 8: 1329-1333, 1993.

[102802]17930.

[102803]Bahn, R. S.; Heufelder, A. E.; Dutton, C. M.: A point muta-

tion

[102804]of the TSH receptor in retro-ocular fibroblasts from a patient with

[102805]Graves' ophthalmopathy. (Abstract) J. Endocr. Invest. 16 (suppl.

[102806]2-6): 30 only, 1993.

[102807]17931.

[102808]Moreland, R. J.; Dresser, M. E.; Rodgers, J. S.; Roe, B. A.; Conaway,

[102809]J. W.; Conaway, R. C.; Hanas, J. S.: Identification of a transcription

[102810]factor IIIA-interacting protein. Nucleic Acids Res. 28: 1986-1993,

[102811]2000.

[102812]17932.

[102813]Biebermann, H.; Schoneberg, T.; Hess, C.; Germak, J.; Gundermann,

[102814]T.; Gruters, A.: The first activating TSH receptor mutation in transmembrane

[102815]domain 1 identified in a family with nonautoimmune hyperthyroidism. J.

[102816]Clin. Endocr. Metab. 86: 4429-4433, 2001.

[102817]17933.

[102818]de Roux, N.; Polak, M.; Couet, J.; Leger, J.; Czernichow, P.;
Milgrom,

[102819]E.; Misrahi, M.: A neomutation of the thyroid-stimulating
hormone

[102820]receptor in a severe neonatal hyperthyroidism. J. Clin. En-
docr. Metab. 81:

[102821]2023-2026, 1996.

[102822]17934.

[102823]Feliciello, A.; Porcellini, A.; Ciullo, I.; Bonavolonta, G.;
Avvedimento,

[102824]E. V.; Fenzi, G.: Expression of thyrotropin-receptor mRNA
in healthy

[102825]and Graves' disease retro-orbital tissue. Lancet 342:
337-338, 1993.

[102826]17935.

[102827]Fuhrer, D.; Holzapfel, H.-P.; Wonerow, P.; Scherbaum, W.
A.; Paschke,

[102828]R.: Somatic mutations in the thyrotropin receptor gene
and not in

[102829]the Gs-alpha protein gene in 31 toxic thyroid nodules. J.
Clin. Endocr.

[102830]Metab. 82: 3885–3891, 1997.

[102831]17936.

[102832]Fuhrer, D.; Wonerow, P.; Willgerodt, H.; Paschke, R.: Identification

[102833]of a new thyrotropin receptor germline mutation
(leu629phe) in a family

[102834]with neonatal onset of autosomal dominant nonautoimmune hyperthyroidism. J.

[102835]Clin. Endocr. Metab. 82: 4234–4238, 1997.

[102836]17937.

[102837]Gabriel, E. M.; Bergert, E. R.; Grant, C. S.; van Heerden, J.

[102838]A.; Thompson, G. B.; Morris, J. C.: Germline polymorphism
of codon

[102839]727 of human thyroid-stimulating hormone receptor is
associated with

[102840]toxic multinodular goiter. J. Clin. Endocr. Metab. 84:
3328–3335,

[102841]1999.

[102842]17938.

[102843]Graves, P.; Pritsker, A.; Davies, T. F.: Post-translational
processing

[102844] of the natural human thyrotropin receptor: demonstration
of more than

[102845] two cleavage sites. J. Clin. Endocr. Metab. 84: 2177–2181,
1999.

[102846] 17939.

[102847] Gruters, A.; Schoneberg, T.; Biebermann, H.; Krude, H.;
Kron,

[102848] H. P.; Dralle, H.; Gudermann, T.: Severe congenital hyper-
thyroidism

[102849] caused by a germ-line neo mutation in the extracellular
portion of

[102850] the thyrotropin receptor. J. Clin. Endocr. Metab. 83:
1431–1436,

[102851] 1998.

[102852] 17940.

[102853] Heldin, N.-E.; Gustavsson, B.; Westermark, K.; Wester-
mark, B.

[102854]: A somatic point mutation in a putative ligand binding
domain of

[102855] the TSH receptor in a patient with autoimmune hyperthy-
roidism. J.

[102856] Clin. Endocr. Metab. 73: 1374–1376, 1991.

[102857]17941.

[102858]Holzapfel, H.-P.; Wonerow, P.; von Petrykowski, W.; Henschen,

[102859]M.; Scherbaum, W. A.; Paschke, R.: Sporadic congenital hyperthyroidism

[102860]due to a spontaneous germline mutation in the thyrotropin receptor

[102861]gene. J. Clin. Endocr. Metab. 82: 3879-3884, 1997.

[102862]17942.

[102863]Khoo, D. H. C.; Parma, J.; Rajasoorya, C.; Ho, S. C.; Vassart,

[102864]G.: A germline mutation of the thyrotropin receptor gene associated

[102865]with thyrotoxicosis and mitral valve prolapse in a Chinese family. J.

[102866]Clin. Endocr. Metab. 84: 1459-1462, 1999.

[102867]17943.

[102868]Kopp, P.; Muirhead, S.; Jourdain, N.; Gu, W.-X.; Jameson, J. L.;

[102869]Rodd, C.: Congenital hyperthyroidism caused by a solitary toxic adenoma

[102870]harboring a novel somatic mutation
(serine281-isoleucine) in the extracellular
[102871]domain of the thyrotropin receptor. J. Clin. Invest. 100:
1634-1639,
[102872]1997.
[102873]17944.
[102874]Kopp, P.; van Sande, J.; Parma, J.; Duprez, L.; Gerber, H.;
Joss,
[102875]E.; Jameson, J. L.; Dumont, J. E.; Vassart, G.: Congenital
hyperthyroidism
[102876]caused by a mutation in the thyrotropin-receptor gene.
New Eng. J.
[102877]Med. 332: 150-154, 1995.
[102878]17945.
[102879]Libert, F.; Lefort, A.; Gerard, C.; Parmentier, M.; Perret, J.;
[102880]Ludgate, M.; Dumont, J. E.; Vassart, G.: Cloning, sequenc-
ing and
[102881]expression of the human thyrotropin (TSH) receptor: evi-
dence for binding
[102882]of autoantibodies. Biochem. Biophys. Res. Commun. 165:
1250-1255,

[102883]1989.

[102884]17946.

[102885]Libert, F.; Passage, E.; Lefort, A.; Vassart, G.; Mattei, M.-G.

[102886]: Localization of human thyrotropin receptor gene to
chromosome region

[102887]14q31 by in situ hybridization. Cytogenet. Cell Genet. 54:
82-83,

[102888]1990.

[102889]17947.

[102890]Loosfelt, H.; Pichon, C.; Jolivet, A.; Misrahi, M.; Caillou, B.;

[102891]Jamous, M.; Vannier, B.; Milgrom, E.: Two-subunit struc-
ture of the

[102892]human thyrotropin receptor. Proc. Nat. Acad. Sci. 89:
3765-3769,

[102893]1992.

[102894]17948.

[102895]Muhlberg, T.; Herrmann, K.; Joba, W.; Kirchberger, M.;
Heberling,

[102896]H.-J.; Heufelder, A. E.: Lack of association of nonautoim-
mune hyperfunctioning

- [102897] thyroid disorders and a germline polymorphism of codon 727 of the
- [102898] human thyrotropin receptor in a European Caucasian population. J.
- [102899] Clin. Endocr. Metab. 85: 2640–2643, 2000.
- [102900] 17949.
- [102901] Garcia-Barcelo, M.; Law, P. T. W.; Tsui, S. K. W.; Fung, K. P.;
- [102902] Lee, C. Y.; Waye, M. M. Y.: Mapping of the human ribosomal large
- [102903] subunit protein gene RPL29 to human chromosome 3q29–qter. Genomics 46:
- [102904] 148–151, 1997.
- [102905] 17950.
- [102906] Kirn-Safran, C. B.; Dayal, S.; Martin-DeLeon, P. A.; Carson, D.
- [102907] D.: Cloning, expression, and chromosome mapping of the murine Hip/Rpl29
- [102908] gene. Genomics 68: 210–219, 2000.
- [102909] 17951.
- [102910] Law, P. T. W.; Tsui, S. K. W.; Lam, W. Y.; Luk, S. C. W.;

Hwaang,

[102911]D. M.; Liew, C. C.; Lee, C. Y.; Fung, K. P.; Waye, M. M. Y.: A novel

[102912]cDNA encoding a human homologue of ribosomal protein L29. Biochim.

[102913]Biophys. Acta 1305: 105–108, 1996.

[102914]17952.

[102915]Liu, S.; Smith, S. E.; Julian, J.; Rohde, L. H.; Karin, N. J.;

[102916]Carson, D. D.: cDNA cloning and expression of HIP, a novel cell surface

[102917]heparan sulfate/heparin-binding protein of human uterine epithelial

[102918]cells and cell lines. J. Biol. Chem. 271: 11817–11823, 1996.

[102919]17953.

[102920]Rohde, L. H.; Julian, J.; Babaknia, A.; Carson, D. D.: Cell surface

[102921]expression of HIP, a novel heparin/heparan sulfate-binding protein,

[102922]of human uterine epithelial cells and cell lines. J. Biol. Chem. 271:

[102923]11824–11830, 1996.

[102924]17954.

[102925]Giachino, C.; Lantelme, E.; Lanzetti, L.; Saccone, S.; Della Valle,

[102926]G.; Migone, N.: A novel SH3-containing human gene family preferentially

[102927]expressed in the central nervous system. Genomics 41: 427-434, 1997.

[102928]17955.

[102929]So, C. W.; Caldas, C.; Liu, M.-M.; Chen, S.-J.; Huang, Q.-H.; Gu,

[102930]L.-J.; Sham, M. H.; Wiedemann, L. M.; Chan, L. C.: EEN encodes for

[102931]a member of a new family of proteins containing an Src homology 3

[102932]domain and is the third gene located on chromosome 19p13 that fuses

[102933]to MLL in human leukemia. Proc. Nat. Acad. Sci. 94: 2563-2568, 1997.

[102934]17956.

[102935]Carling, T.; Kindmark, A.; Hellman, P.; Holmberg, L.; Akerstrom,

[102936]G.; Rastad, J.: Vitamin D receptor alleles b, a, and T: risk factors

[102937]for sporadic primary hyperparathyroidism (HPT) but not HPT of uremia

[102938]or MEN1. Biochem. Biophys. Res. Commun. 231: 329-332, 1997.

[102939]17957.

[102940]Carling, T.; Rastad, J.; Akerstrom, G.; Westin, G.: Vitamin D

[102941]receptor (VDR) and parathyroid hormone messenger ribonucleic acid

[102942]levels correspond to polymorphic VDR alleles in human parathyroid

[102943]tumors. J. Clin. Endocr. Metab. 83: 2255-2259, 1998.

[102944]17958.

[102945]Carling, T.; Ridefelt, P.; Hellman, P.; Rastad, J.; Akerstrom,

[102946]G.: Vitamin D receptor polymorphisms correlate to parathyroid cell

[102947]function in primary hyperparathyroidism. J. Clin. Endocr. Metab. 82:

[102948]1772–1775, 1997.

[102949]17959.

[102950]Correa, P.; Rastad, J.; Schwarz, P.; Westin, G.; Kindmark, A.;

[102951]Lundgren, E.; Akerstrom, G.; Carling, T.: The vitamin D receptor

[102952](VDR) start codon polymorphism in primary hyperparathyroidism and

[102953]parathyroid VDR messenger ribonucleic acid levels. J. Clin. Endocr.

[102954]Metab. 84: 1690–1694, 1999.

[102955]17960.

[102956]Ensrud, K. E.; Stone, K.; Cauley, J. A.; White, C.; Zmuda, J. M.;

[102957]Nguyen, T. V.; Eisman, J. A.; Cummings, S. R.: Vitamin D receptor

[102958]gene polymorphisms and the risk of fractures in older women. J. Bone

[102959]Miner. Res. 14: 1637–1645, 1999.

[102960]17961.

[102961]Ferrari, S.; Manen, D.; Bonjour, J.-P.; Slosman, D.; Rizzoli,

R.

[102962]; Bone mineral mass and calcium and phosphate metabolism in young

[102963]men: relationships with vitamin D receptor allelic polymorphisms. J.

[102964]Clin. Endocr. Metab. 84: 2043–2048, 1999.

[102965]17962.

[102966]Gennari, L.; Becherini, L.; Masi, L.; Mansani, R.; Gonnelli, S.;

[102967]Cepollaro, C.; Martini, S.; Montagnani, A.; Lentini, G.; Becorpi,

[102968]A. M.; Brandi, M. L.: Vitamin D and estrogen receptor allelic variants

[102969]in Italian postmenopausal women: evidence of multiple gene contribution

[102970]to bone mineral density. J. Clin. Endocr. Metab. 83: 939–944, 1998.

[102971]17963.

[102972]Kristjansson, K.; Rut, A. R.; Hewison, M.; O'Riordan, J. L. H.;

[102973]Hughes, M. R.: Two mutations in the hormone binding domain of the

- [102974] vitamin D receptor cause tissue resistance to 1,25 dihydroxyvitamin D(3). J. Clin. Invest. 92: 12-16, 1993.
- [102975] 17964.
- [102977] Labuda, M.; Ross, M. V.; Fujiwara, T. M.; Morgan, K.; Ledbetter, D.; Hughes, M. R.; Glorieux, F. H.: Two hereditary defects related to vitamin D metabolism map to the same region of human chromosome 12q. II(Abtract) Cytogenet. Cell Genet. 58: 1978, 1991.
- [102980] 17965.
- [102982] Jurutka, P. W.; Remus, L. S.; Whitfield, G. K.; Thompson, P. D.; Hsieh, J.-C.; Zitzer, H.; Tavakkoli, P.; Galligan, M. A.; Dang, H. T. L.; Haussler, C. A.; Haussler, M. R.: The polymorphic N terminus in human vitamin D receptor isoforms influences transcriptional activity by modulating interaction with transcription factor IIB. Molec. Endocr. 14:

[102987]401–420, 2000.

[102988]17966.

[102989]Lin, U.-T.; Malloy, P. J.; Sakati, N.; Al-Ashwal, A.; Feldman,

[102990]D.: A novel mutation in the deoxyribonucleic acid-binding
domain

[102991]of the vitamin D receptor causes hereditary
1,25-dihydroxyvitamin

[102992]D-resistant rickets. J. Clin. Endocr. Metab. 81:
2564–2569, 1996.

[102993]17967.

[102994]Lorentzon, M.; Lorentzon, R.; Nordstrom, P.: Vitamin D re-
ceptor

[102995]gene polymorphism is associated with birth height,
growth to adolescence,

[102996]and adult stature in healthy Caucasian men: a cross-
sectional and

[102997]longitudinal study. J. Clin. Endocr. Metab. 85: 1666–1671,
2000.

[102998]17968.

[102999]Makishima, M.; Lu, T. T.; Xie, W.; Whitfield, G. K.; Domoto,
H.;

[103000]Evans, R. M.; Haussler, M. R.; Mangelsdorf, D. J.: Vitamin D receptor

[103001]as an intestinal bile acid sensor. Science 296: 1313–1316, 2002.

[103002]17969.

[103003]Mechica, J. B.; Leite, M. O. R.; Mendonca, B. B.; Frazzatto, E.

[103004]S. T.; Borelli, A.; Latronico, A. C.: A novel nonsense mutation in

[103005]the first zinc finger of the vitamin D receptor causing hereditary

[103006]1,25-dihydroxyvitamin D(3)-resistant rickets. J. Clin. Endocr. Metab. 82:

[103007]3892–3894, 1997.

[103008]17970.

[103009]Furlan, M.; Robles, R.; Galbusera, M.; Remuzzi, G.; Kyrle, P. A.;

[103010]Brenner, B.; Krause, M.; Scharrer, I.; Aumann, V.; Mittler, U.; Solenthaler,

[103011]M.; Lammle, B.: Von Willebrand factor-cleaving protease in thrombotic

[103012]thrombocytopenic purpura and the hemolytic-uremic syn-

drome. New Eng.

[103013]J. Med. 339: 1578–1584, 1998.

[103014]17971.

[103015]Furlan, M.; Robles, R.; Lammle, B.: Partial purification and characterization

[103016]of a protease from human plasma cleaving von Willebrand factor to

[103017]fragments produced by in vivo proteolysis. Blood 87: 4223–4234,

[103018]1996.

[103019]17972.

[103020]Nichols, A. F.; Ong, P.; Linn, S.: Mutations specific to the xeroderma

[103021]pigmentosum group E Ddb– phenotype. J. Biol. Chem. 271: 24317–24320,

[103022]1996.

[103023]17973.

[103024]Itoh, T.; Linn, S.; Kamide, R.; Tokushige, H.; Katori, N.; Hosaka,

[103025]Y.; Yamaizumi, M.: Xeroderma pigmentosum variant heterozygotes show

[103026]reduced levels of recovery of replicative DNA synthesis in
the presence
[103027]of caffeine after ultraviolet irradiation. J. Invest. Derm.
115:
[103028]981–985, 2000.
[103029]17974.
[103030]Shisheva, A.; Sudhof, T. C.; Czech, M. P.: Cloning, charac-
terization,
[103031]and expression of a novel GDP dissociation inhibitor iso-
form from
[103032]skeletal muscle. Molec. Cell. Biol. 14: 3459–3468, 1994.
[103033]17975.
[103034]Fathi, Z.; Corjay, M. H.; Shapira, H.; Wada, E.; Benya, R.;
Jensen,
[103035]R.; Viallet, J.; Sausville, E. A.; Battey, J. F.: BRS-3: a novel
[103036]bombesin receptor subtype selectively expressed in testis
and lung
[103037]carcinoma cells. J. Biol. Chem. 268: 5979–5984, 1993.
[103038]17976.
[103039]Gorbulev, V.; Akhundova, A.; Grzeschik, K. H.; Fahrenholz,
F.:

- [103040] Organization and chromosomal localization of the gene
for the human
- [103041] bombesin receptor subtype expressed in pregnant uterus.
FEBS Lett. 340:
- [103042] 260–264, 1994.
- [103043] 17977.
- [103044] Ohki–Hamazaki, H.; Watase, K.; Yamamoto, K.; Ogura, H.;
Yamano,
- [103045] M.; Yamada, K.; Maeno, H.; Imaki, J.; Kikuyama, S.; Wada,
E.; Wada,
- [103046] K.: Mice lacking bombesin receptor subtype–3 develop
metabolic defects
- [103047] and obesity. Nature 390: 166–169, 1997.
- [103048] 17978.
- [103049] Banfi, S.; Borsani, G.; Bulfone, A.; Ballabio, A.: Drosophila–re-
lated
- [103050] expressed sequences. Hum. Molec. Genet. 6: 1745–1753,
1997.
- [103051] 17979.
- [103052] Castrillon, D. H.; Wasserman, S. A.: Diaphanous is re-
quired for

[103053]cytokinesis in *Drosophila* and shares domains of similarity with the

[103054]products of the limb deformity gene. *Development* 120: 3367–3377,

[103055]1994.

[103056]17980.

[103057]Philippe, C.; Cremers, F. P. M.; Chery, M.; Bach, I.; Abbadi, N.;

[103058]Ropers, H. H.; Gilgenkrantz, S.: Physical mapping of DNA markers

[103059]in the q13–q22 region of the human X chromosome. *Genomics* 17: 147–152,

[103060]1993.

[103061]17981.

[103062]Bione, S.; Sala, C.; Manzini, C.; Arrigo, G.; Zuffardi, O.; Banfi,

[103063]S.; Borsani, G.; Jonveaux, P.; Philippe, C.; Zuccotti, M.; Bal-labio,

[103064]A.; Toniolo, D.: A human homologue of the *Drosophila melanogaster*

[103065]diaphanous gene is disrupted in a patient with premature ovarian failure:

[103066]evidence for conserved function in oogenesis and implications for

[103067]human sterility. *Am. J. Hum. Genet.* 62: 533–541, 1998.

[103068]17982.

[103069]Sala, C.; Arrigo, G.; Torri, G.; Martinazzi, F.; Riva, P.; Larizza,

[103070]L.; Philippe, C.; Jonveaux, P.; Sloan, F.; Labella, T.; Toniolo, D.

[103071]: Eleven X chromosome breakpoints associated with premature ovarian

[103072]failure (POF) map to a 15-Mb YAC contig spanning Xq21. *Genomics* 40:

[103073]123–131, 1997.

[103074]17983.

[103075]Jagla, W. M.; Jagle, H.; Hayashi, T.; Sharpe, L. T.; Deeb, S.

[103076]S.: The molecular basis of dichromatic color vision in males with

[103077]multiple red and green visual pigment genes. *Hum. Molec. Genet.* 11:

[103078]23–32, 2002.

[103079]17984.

- [103080]Huijing, F.; Fernandez, J.: X-chromosomal inheritance of
liver
- [103081]glycogenosis with phosphorylase kinase deficiency. Am. J.
Hum. Genet. 21:
- [103082]275–284, 1969.
- [103083]17985.
- [103084]Franco, B.; Guioli, S.; Pragliola, A.; Incerti, B.; Bardoni, B.;
- [103085]Tonlorenzi, R.; Carrozzo, R.; Maestrini, E.; Pieretti, M.;
- Taillon–Miller,
- [103086]P.; Brown, C. J.; Willard, H. F.; Lawrence, C.; Persico, M. G.;
- Camerino,
- [103087]G.; Ballabio, A.: A gene deleted in Kallmann's syndrome
shares homology
- [103088]with neural cell adhesion and axonal path–finding
molecules. Nature 353:
- [103089]529–536, 1991.
- [103090]17986.
- [103091]Lachner, M.; O'Carroll, D.; Rea, S.; Mechtler, K.; Jenuwein,
T.
- [103092]: Methylation of histone H3 lysine 9 creates a binding site
for HP1

[103093]proteins. Nature 410: 116–120, 2001.

[103094]17987.

[103095]Melcher, M.; Schmid, M.; Aagaard, L.; Selenko, P.; Laible, G.;

[103096]Jenuwein, T.: Structure–function analysis of SUV39H1 reveals a dominant

[103097]role in heterochromatin organization, chromosome segregation, and

[103098]mitotic progression. Molec. Cell Biol. 20: 3728–3841, 2000.

[103099]17988.

[103100]Nielsen, S. J.; Schneider, R.; Bauer, U.–M.; Bannister, A. J.;

[103101]Morrison, A.; O'Carroll, D.; Firestein, R.; Cleary, M.; Jenuwein,

[103102]T.; Herrera, R. E.; Kouzarides, T.: Rb targets histone H3 methylation

[103103]and HP1 to promoters. Nature 412: 561–565, 2001.

[103104]17989.

[103105]Peters, A. H. F. M.; O'Carroll, D.; Scherthan, H.; Mechtler, K.;

[103106]Sauer, S.; Schofer, C.; Weipoltshammer, K.; Pagani, M.;

Lachner, M.;

[103107]Kohlmaier, A.; Opravil, S.; Doyle, M.; Sibilia, M.; Jenuwein, T.:

[103108]Loss of the Suv39h histone methyltransferases impairs mammalian heterochromatin

[103109]and genome stability. Cell 107: 323–337, 2001.

[103110]17990.

[103111]Rea, S.; Eisenhaber, F.; O'Carroll, D.; Strahl, B. D.; Sun, Z.-W.;

[103112]Schmid, M.; Opravil, S.; Mechtler, K.; Ponting, C. P.; Allis, C. D.;

[103113]Jenuwein, T.: Regulation of chromatin structure by site-specific

[103114]histone H3 methyltransferases. Nature 406: 593–599, 2000.

[103115]17991.

[103116]Scott, A. F.: Personal Communication. Baltimore, Md. 8/7/2000.

[103117]17992.

[103118]Haltiwanger, R. S.; Blomberg, M. A.; Hart, G. W.: Glycosylation

[103119]of nuclear and cytoplasmic proteins: purification and

characterization

[103120] of a uridine diphospho-N-acetylglucosamine:polypeptide
beta-N-acetylglucosaminyltransferase. J.

[103121] Biol. Chem. 267: 9005-9013, 1992.

[103122] 17993.

[103123] Kreppel, L. K.; Blomberg, M. A.; Hart, G. W.: Dynamic gly-
cosylation

[103124] of nuclear and cytosolic proteins: cloning and characteri-
zation of

[103125] a unique O-GlcNAc transferase with multiple tetratri-
copeptide repeats. J.

[103126] Biol. Chem. 272: 9308-9315, 1997.

[103127] 17994.

[103128] Lubas, W. A.; Frank, D. W.; Krause, M.; Hanover, J. A.: O-
linked

[103129] GlcNAc transferase is a conserved nucleocytoplasmic pro-
tein containing

[103130] tetratricopeptide repeats. J. Biol. Chem. 272: 9316-9324,
1997.

[103131] 17995.

[103132] Shafi, R.; Iyer, S. P. N.; Ellies, L. G.; O'Donnell, N.; Marek,

- [103133]K. W.; Chui, D.; Hart, G. W.; Marth, J. D.: The O-GlcNAc transferase
- [103134]gene resides on the X chromosome and is essential for embryonic stem
- [103135]cell viability and mouse ontogeny. Proc. Nat. Acad. Sci. 97: 5735–5739,
- [103136]2000.
- [103137]17996.
- [103138]Yang, X.; Zhang, F.; Kudlow, J. E.: Recruitment of O-GlcNAc transferase
- [103139]to promoters by corepressor mSin3A: coupling protein O-GlcNAcylation
- [103140]to transcriptional repression. Cell 110: 69–80, 2002.
- [103141]17997.
- [103142]Danon, M. J.; Oh, S. J.; DiMauro, S.; Manaligod, J. R.; Eastwood,
- [103143]A.; Naidu, S.; Schliselfeld, L. H.: Lysosomal glycogen storage disease
- [103144]with normal acid maltase. Neurology 31: 51–57, 1981.
- [103145]17998.
- [103146]Dworzak, F.; Casazza, F.; Mora, C. M.; De Maria, R.;

Gronda, E.;

[103147]Baroldi, G.; Rimoldi, M.; Morandi, L.; Cornelio, F.: Lysosomal glycogen

[103148]storage with normal acid maltase: a familial study with successful

[103149]heart transplant. Neuromusc. Disord. 4: 243–247, 1994.

[103150]17999.

[103151]Liu, Y.; Hoffmann, A.; Grinberg, A.; Westphal, H.; McDonald, M.

[103152]P.; Miller, K. M.; Crawley, J. N.; Sandhoff, K.; Suzuki, K.; Proia,

[103153]R. L.: Mouse model of GM2 activator deficiency manifests cerebellar

[103154]pathology and motor impairment. Proc. Nat. Acad. Sci. 94: 8138–8143,

[103155]1997.

[103156]18000.

[103157]Mahuran, D. J.: Personal Communication. Toronto, Ontario, Canada

[103158]12/1/1994.

[103159]18001.

[103160]O'Neill, B.; Butler, A. B.; Young, E.; Falk, P. M.; Bass, N. H.

[103161]: Adult-onset Gm2-gangliosidosis: seizures, dementia,
and normal pressure

[103162]hydrocephalus associated with glycolipid storage in the
brain and

[103163]arachnoid granulation. Neurology 28: 1117-1123, 1978.

[103164]18002.

[103165]Phillips, J. A., III: Personal Communication. Baltimore, Md.

[103166]8/11/1983.

[103167]18003.

[103168]Sakuraba, H.; Itoh, K.; Shimmoto, M.; Utsumi, K.; Kase, R.;
Hashimoto,

[103169]Y.; Ozawa, T.; Ohwada, Y.; Imataka, G.; Eguchi, M.; Fu-
rukawa, T.;

[103170]Schepers, U.; Sandhoff, K.: GM2 gangliosidosis AB variant:
clinical

[103171]and biochemical studies of a Japanese patient. Neurology
52: 372-377,

[103172]1999.

[103173]18004.

[103174]Schepers, U.; Glombitza, G.; Lemm, T.; Hoffmann, A.;
Chabas, A.;

[103175]Ozand, P.; Sandhoff, K.: Molecular analysis of a
GM2-activator deficiency

[103176]in two patients with GM2-gangliosidosis AB variant. Am. J.
Hum. Genet. 59:

[103177]1048-1056, 1996.

[103178]18005.

[103179]Schroder, M.; Schnabel, D.; Hurwitz, R.; Young, E.; Suzuki,
K.;

[103180]Sandhoff, K.: Molecular genetics of GM2-gangliosidosis
AB variant:

[103181]a novel mutation and expression in BHK cells. Hum.
Genet. 92: 437-440,

[103182]1993.

[103183]18006.

[103184]Schroder, M.; Schnabel, D.; Suzuki, K.; Sandhoff, K.: A mu-
tation

[103185]in the gene of a glycolipid-binding protein (GM2 activator)
that causes

[103186]GM2-gangliosidosis variant AB. FEBS Lett. 290: 1-3, 1991.

[103187]18007.

[103188]Swallow, D. M.; Islam, I.; Fox, M. F.; Povey, S.; Klima, H.;
Schepers,

[103189]U.; Sandhoff, K.: Regional localization of the gene coding
for the

[103190]GM2 activator protein (GM2A) to chromosome 5q32-33
and confirmation

[103191]of the assignment of GM2AP to chromosome 3. Ann. Hum.
Genet. 57:

[103192]187-193, 1993.

[103193]18008.

[103194]Valentine, C. R.; Heflich, R. H.: The association of non-
sense

[103195]mutations with exon-skipping in hprt mRNA of Chinese
hamster ovary

[103196]cells results from an artifact of RT-PCR. RNA 3: 660-676,
1997.

[103197]18009.

[103198]Xie, B.; Kennedy, J. L.; McInnes, B.; Auger, D.; Mahuran, D.:

[103199]Identification of a processed pseudogene related to the
functional

[103200] gene encoding the G-M2 activator protein: localization of the pseudogene

[103201] to human chromosome 3 and the functional gene to human chromosome

[103202] 5. Genomics 14: 796-798, 1992.

[103203] 18010.

[103204] Xie, B.; Wang, W.; Mahuran, D. J.: A cys138-to-arg substitution

[103205] in the G-M2 activator protein is associated with the AB variant form

[103206] of G-M2 gangliosidosis. Am. J. Hum. Genet. 50: 1046-1052, 1992.

[103207] 18011.

[103208] Yamanaka, S.; Johnson, O. N.; Lyu, M. S.; Kozak, C. A.; Proia,

[103209] R. L.: The mouse gene encoding the G(M2) activator protein (Gm2a):

[103210] cDNA sequence, expression, and chromosome mapping. Genomics 24:

[103211] 601-604, 1994.

[103212] 18012.

[103213]Ainsworth, P. J.; Coulter-Mackie, M. B.: A double mutation
in
[103214]exon 6 of the beta-hexosaminidase alpha subunit in a pa-
tient with
[103215]the B1 variant of Tay-Sachs disease. Am. J. Hum. Genet.
51: 802-809,
[103216]1992.
[103217]18013.
[103218]Akalin, N.; Shi, H.-P.; Vavougios, G.; Hechtman, P.; Lo, W.;
Scriver,
[103219]C. R.; Mahuran, D.; Kaplan, F.: Novel Tay-Sachs disease
mutations
[103220]from China. Hum. Mutat. 1: 40-46, 1992.
[103221]18014.
[103222]Akerman, B. R.; Zielenski, J.; Triggs-Raine, B. L.; Prencz, E.
[103223]M.; Natowicz, M. R.; Lim-Steele, J. S. T.; Kaback, M. M.;
Miles, E.
[103224]H.; Thomas, G. H.; Clarke, J. T. R.; Gravel, R. A.: A muta-
tion common
[103225]in non-Jewish Tay-Sachs disease: frequency and RNA
studies. Hum.

[103226]Mutat. 1: 303–309, 1992.

[103227]18015.

[103228]Akli, S.; Chelly, J.; Lacorte, J.; Poenaru, L.; Kahn, A.: Seven

[103229]novel Tay–Sachs mutations detected by chemical mismatch cleavage of

[103230]PCR–amplified cDNA fragments. Genomics 11: 124–134, 1991.

[103231]18016.

[103232]Akli, S.; Chomel, J.–C.; Lacorte, J.–M.; Bachner, L.; Poenaru,

[103233]A.; Poenaru, L.: Ten novel mutations in the HEXA gene in non–Jewish

[103234]Tay–Sachs patients. Hum. Molec. Genet. 2: 61–67, 1993.

[103235]18017.

[103236]Lyons, L. B.; Cox, R. P.; Dancis, J.: Complementation analysis

[103237]of maple syrup urine disease in heterokaryons derived from cultured

[103238]human fibroblasts. Nature 243: 533–535, 1973.

[103239]18018.

[103240]Singh, S.; Willers, I.; Goedde, H. W.: Heterogeneity in
maple

[103241]syrup urine disease: aspects of cofactor requirement and
complementation

[103242]in cultured fibroblasts. Clin. Genet. 11: 277–284, 1977.

[103243]18019.

[103244]Ohno, K.; Brengman, J.; Tsujino, A.; Engel, A. G.: Human
endplate

[103245]acetylcholinesterase deficiency caused by mutations in the
collagen-like

[103246]tail subunit (ColQ) of the asymmetric enzyme. Proc. Nat.
Acad. Sci. 95:

[103247]9654–9659, 1998.

[103248]18020.

[103249]Schwartz, J. M.; Reiss, A. L.; Jaffe, E. R.: Hereditary methemoglobinemia

[103250]with deficiency of NADH cytochrome b5 reductase. In:
Stanbury, J. B.;

[103251]Wyngaarden, J. B.; Fredrickson, D. S.; Goldstein, J. L.;
Brown, M.

[103252]S.: The Metabolic Basis of Inherited Disease. New York:
McGraw-Hill

[103253](pub.) (5th ed.): 1983. Pp. 1654–1665.

[103254]18021.

[103255]Scott, E. M.: The relationship of diaphorase of human erythrocytes

[103256]to inheritance of methemoglobinemia. J. Clin. Invest. 39: 1176–1179,

[103257]1960.

[103258]18022.

[103259]Scott, E. M.; Lewis, M.; Kaita, H.; Chown, B.; Giblett, E. R.

[103260]: The absence of close linkage of methemoglobinemia and blood group

[103261]loci. Am. J. Hum. Genet. 15: 493–494, 1963.

[103262]18023.

[103263]Scott, E. M.; Wright, R. C.: The absence of close linkage of

[103264]methemoglobinemia and other loci. Am. J. Hum. Genet. 21: 194–195,

[103265]1969.

[103266]18024.

[103267]Shirabe, K.: Personal Communication. Oita, Japan

7/21/1997.

[103268]18025.

[103269]Shirabe, K.; Fujimoto, Y.; Yubisui, T.; Takeshita, M.: An in-frame

[103270]deletion of codon 298 of the NADH-cytochrome b(5) reductase gene results

[103271]in hereditary methemoglobinemia type II (generalized type): a functional

[103272]implication for the role of the COOH-terminal region of the enzyme. J.

[103273]Biol. Chem. 269: 5952-5957, 1994.

[103274]18026.

[103275]Shirabe, K.; Landi, M. T.; Takeshita, M.; Uziel, G.; Fedrizzi,

[103276]E.; Borgese, N.: A novel point mutation in a 3-prime splice site

[103277]of the NADH-cytochrome b5 reductase gene results in immunologically

[103278]undetectable enzyme and impaired NADH-dependent ascorbate regeneration

[103279]in cultured fibroblasts of a patient with type II hereditary methemoglobinemia. Am.

[103280]J. Hum. Genet. 57: 302-310, 1995.

[103281]18027.

[103282]Shirabe, K.; Yubisui, T.; Borgese, N.; Tang, C.; Hultquist,
D.

[103283]E.; Takeshita, M.: Enzymatic instability of NADH-cy-
tochrome b5 reductase

[103284]as a cause of hereditary methemoglobinemia type I (red
cell type). J.

[103285]Biol. Chem. 267: 20416-20421, 1992.

[103286]18028.

[103287]Tanishima, K.; Matsuki, T.; Fukuda, N.; Takeshita, M.;
Yoneyama,

[103288]Y.: NADH-cytochrome b5 reductase in platelets and
leukocytes with

[103289]special reference to normal levels and to levels in carriers
of hereditary

[103290]methemoglobinemia with or without neurological symp-
toms. Acta Haemat. 63:

[103291]7-12, 1980.

[103292]18029.

[103293]Tanishima, K.; Tanimoto, K.; Tomoda, A.; Mawatari, K.;
Matsukawa,

[103294]S.; Yoneyama, Y.; Ohkuwa, H.; Takazakura, E.: Hereditary methemoglobinemia

[103295]due to cytochrome b(5) reductase deficiency in blood cells without

[103296]associated neurologic and mental disorders. Blood 66: 1288–1291,

[103297]1985.

[103298]18030.

[103299]Tomatsu, S.; Kobayashi, Y.; Fukumaki, Y.; Yubisui, T.; Orii, T.;

[103300]Sakaki, Y.: The organization and the complete nucleotide sequence

[103301]of the human NADH-cytochrome b5 reductase gene. Gene 80: 353–361,

[103302]1989.

[103303]18031.

[103304]Townes, P. L.; Morrison, M.: Investigation of the defect in a

[103305]variant of hereditary methemoglobinemia. Blood 19: 60–74, 1962.

[103306]18032.

[103307]Trost, C.: The Blue People of Troublesome Creek. Science
82

[103308](Nov.): 35–39, 1982.

[103309]18033.

[103310]Vieira, L. M.; Kaplan, J.-C.; Kahn, A.; Leroux, A.: Four new
[103311]mutations in the NADH-cytochrome b5 reductase gene
from patients with

[103312]recessive congenital methemoglobinemia type II. Blood
85: 2254–2262,

[103313]1995.

[103314]18034.

[103315]Vives-Corróns, J. L.; Pujades, A.; Vela, E.; Corretger, J. M.;

[103316]Leroux, A.; Kaplan, J. C.: Congenital methemoglobin-reduc-
tase (cytochrome

[103317]b5 reductase) deficiency associated with mental retarda-
tion in a Spanish

[103318]girl. Acta Haemat. 59: 348–353, 1978.

[103319]18035.

[103320]Waller, H. D.: Inherited methemoglobinemia (enzyme defi-
ciencies). Humangenetik 9:

[103321]217–218, 1970.

[103322]18036.

[103323]Wang, Y.; Wu, Y.-S.; Zheng, P.-Z.; Yang, W.-X.; Fang, G.-A.; Tang,

[103324]Y.-C.; Xie, F.; Lan, F.-H.; Zhu, Z.-Y.: A novel mutation in the NADH-cytochrome

[103325]b5 reductase gene of a Chinese patient with recessive congenital methemoglobinemia. Blood 95:

[103326]3250–3255, 2000.

[103327]18037.

[103328]West, C. A.; Gomperts, B. D.; Huehns, E. R.; Kessel, I.; Ashby,

[103329]J. R.: Demonstration of an enzyme variant in a case of congenital

[103330]methaemoglobinaemia. Brit. Med. J. 4: 212–214, 1967.

[103331]18038.

[103332]Worster-Drought, C.; White, J. C.; Sargent, F.: Familial, idiopathic

[103333]methaemoglobinaemia associated with mental deficiency and neurological

[103334]abnormalities. Brit. Med. J. 2: 114–118, 1953.

[103335]18039.

[103336]Wu, Y.-S.; Huang, C.-H.; Wan, Y.; Huang, Q.-J.; Zhu, Z.-Y.:

Identification

[103337]of a novel point mutation (leu72-to-pro) in the NADH-cytochrome b5

[103338]reductase gene of a patient with hereditary methaemoglobinaemia type

[103339]J. Brit. J. Haemat. 102: 575-577, 1998.

[103340]18040.

[103341]Yubisui, T.; Miyata, T.; Iwanaga, S.; Tamura, M.; Takeshita, M.

[103342]: Complete amino acid sequence of NADH-cytochrome b5 reductase purified

[103343]from human erythrocytes. J. Biochem. 99: 407-422, 1986.

[103344]18041.

[103345]Yubisui, T.; Naitoh, Y.; Zenno, S.; Tamura, M.; Takeshita, M.;

[103346]Sakaki, Y.: Molecular cloning of cDNAs of human liver and placenta

[103347]NADH-cytochrome b(5) reductase. Proc. Nat. Acad. Sci. 84: 3609-3613,

[103348]1987.

[103349]18042.

[103350]Yubisui, T.; Shirabe, K.; Takeshita, M.; Kobayashi, Y.;
Fukumaki,

[103351]Y.; Sakaki, Y.; Takano, T.: Structural role of serine 127 in
the

[103352]NADH-binding site of human NADH-cytochrome b5 re-
ductase. J. Biol.

[103353]Chem. 266: 66-70, 1991.

[103354]18043.

[103355]Alvarez, L.; Corrales, F.; Martin-Duce, A.; Mato, J. M.:
Characterization

[103356]of a full-length cDNA encoding human liver S-
adenosylmethionine synthetase:

[103357]tissue-specific gene expression and mRNA levels in hep-
atopathies. Biochem.

[103358]J. 293: 481-486, 1993.

[103359]18044.

[103360]Blom, H. J.; Davidson, A. J.; Finkelstein, J. D.; Luder, A. S.;

[103361]Bernardini, I.; Martin, J. J.; Tangerman, A.; Trijbels, J. M. F.;

- [103362]Mudd, S. H.; Goodman, S. I.; Gahl, W. A.: Persistent hypermethioninaemia
[103363]with dominant inheritance. J. Inherit. Metab. Dis. 15:
188–197,
[103364]1992.
[103365]18045.
- [103366]Cabrero, C.; Puerta, J.; Alemany, S.: Purification and comparison
[103367]of two forms of S-adenosyl-L-methionine synthetase
from rat liver. Europ.
[103368]J. Biochem. 170: 299–304, 1987.
[103369]18046.
- [103370]Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Levy, H. L.;
Chou,
[103371]J. Y.: Dominant inheritance of isolated hypermethionine-
mia is associated
[103372]with a mutation in the human methionine adenosyltrans-
ferase 1A gene. Am.
[103373]J. Hum. Genet. 60: 540–546, 1997.
[103374]18047.
- [103375]Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Thomas, J.;

Pao, V.

[103376]Y.; Nguyen, T. K.; Levy, H. L.; Greene, C.; Freehauf, C.;
Chou, J.

[103377]Y.: Methionine adenosyltransferase I/III deficiency: novel
mutations

[103378]and clinical variations. Am. J. Hum. Genet. 66: 347–355,
2000.

[103379]18048.

[103380]Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Wilson, W. G.;
Leonard,

[103381]J. V.; Chou, J. Y.: Demyelination of the brain is associated
with

[103382]methionine adenosyltransferase I/III deficiency. J. Clin. In-
vest. 98:

[103383]1021–1027, 1996.

[103384]18049.

[103385]Finkelstein, J. D.; Kyle, W. E.; Martin, J. J.: Abnormal me-
thionine

[103386]adenosyltransferase in hypermethioninemia. Biochem.
Biophys. Res.

[103387]Commun. 66: 1491–1497, 1975.

[103388]18050.

- [103389]Lim, L. E.; Duclos, F.; Broux, O.; Bourg, N.; Sunada, Y.; Al-
lamand,
[103390]V.; Meyer, J.; Richard, I.; Moomaw, C.; Slaughter, C.; Tome,
F. M.
[103391]S.; Fardeau, M.; Jackson, C. E.; Beckmann, J. S.; Campbell,
K. P.
[103392]; Beta-sarcoglycan: characterization and role in limb-
girdle muscular
[103393]dystrophy linked to 4q12. Nature Genet. 11: 257-265,
1995.
[103394]18051.
[103395]Passos-Bueno, M. R.; Vainzof, M.; Moreira, E. S.; Zatz, M.:
Seven
[103396]autosomal recessive limb-girdle muscular dystrophies in
the Brazilian
[103397]population: from LGMD2A to LGMD2G. Am. J. Med. Genet.
82: 392-398,
[103398]1999.
[103399]18052.
[103400]Jarvela, I.: Infantile neuronal ceroid lipofuscinosis (CLN1):
[103401]linkage disequilibrium in the Finnish population and evi-
dence that

[103402]variant late infantile form (variant CLN2) represents a nonallelic

[103403]locus. Genomics 10: 333–337, 1991.

[103404]18053.

[103405]Nishiyama, K.; Funai, T.; Katafuchi, R.; Hattori, F.; Onoyama,

[103406]K.; Ichiyama, A.: Primary hyperoxaluria type I due to a point mutation

[103407]of T to C in the coding region of the serine:pyruvate aminotransferase

[103408]gene. Biochem. Biophys. Res. Commun. 176: 1093–1099, 1991.

[103409]18054.

[103410]Pirulli, D.; Puzzer, D.; Ferri, L.; Crovella, S.; Amoroso, A.;

[103411]Ferrettini, C.; Marangella, M.; Mazzola, G.; Florian, F.: Molecular

[103412]analysis of hyperoxaluria type 1 in Italian patients reveals eight

[103413]new mutations in the alanine:glyoxylate aminotransferase gene. Hum.

[103414]Genet. 104: 523–525, 1999.

[103415]18055.

[103416]Watts, R. W. E.; Calne, R. Y.; Rolles, K.; Danpure, C. J.;
Morgan,

[103417]S. H.; Mansell, M. A.; Williams, R.; Purkiss, P.: Successful
treatment

[103418]of primary hyperoxaluria type I by combined hepatic and
renal transplantation. Lancet II:

[103419]474–475, 1987.

[103420]18056.

[103421]Argoff, C. E.; Kaneski, C. R.; Blanchette–Mackie, E. J.;
Comly,

[103422]M.; Dwyer, N. K.; Brown, A.; Brady, R. O.; Pentchev, P. G.:
Type

[103423]C Niemann–Pick disease: documentation of abnormal LDL
processing in

[103424]lymphocytes. Biochem. Biophys. Res. Commun. 171:
38–45, 1990.

[103425]18057.

[103426]Natt, E.; Kida, K.; Odievre, M.; Di Rocco, M.; Scherer, G.:
Point

[103427]mutations in the tyrosine aminotransferase gene in ty-
rosinemia type

[103428]II. Proc. Nat. Acad. Sci. 89: 9297–9301, 1992.

[103429]18058.

[103430]Brenner, V.; Nyakatura, G.; Rosenthal, A.; Platzer, M.: Ge-
nomic

[103431]organization of two novel genes on human Xq28: compact
head to head

[103432]arrangement of IDH-gamma and TRAP-delta is conserved
in rat and mouse. Genomics 44:

[103433]8–14, 1997.

[103434]18059.

[103435]Achen, M. G.; Jeltsch, M.; Kukk, E.; Makinen, T.; Vitali, A.;
Wilks,

[103436]A. F.; Alitalo, K.; Stacker, S. A.: Vascular endothelial
growth factor

[103437]D (VEGF-D) is a ligand for the tyrosine kinases VEGF re-
ceptor 2 (Flk1)

[103438]and VEGF receptor 3 (Flt4). Proc. Nat. Acad. Sci. 95:
548–553, 1998.

[103439]18060.

[103440]Stacker, S. A.; Caesar, C.; Baldwin, M. E.; Thornton, G. E.;
Williams,

[103441]R. A.; Prevo, R.; Jackson, D. G.; Nishikawa, S.; Kubo, H.;
Achen,

[103442]M. G.: VEGF-D promotes the metastatic spread of tumor
cells via the

[103443]lymphatics. Nature Med. 7: 186–191, 2001.

[103444]18061.

[103445]Yamada, Y.; Nezu, J.; Shimane, M.; Hirata, Y.: Molecular
cloning

[103446]of a novel vascular endothelial growth factor, VEGF-D. Ge-
nomics 42:

[103447]483–488, 1997.

[103448]18062.

[103449]Davies, P. A.; Hanna, M. C.; Hales, T. G.; Kirkness, E. F.:
Insensitivity

[103450]to anaesthetic agents conferred by a class of GABA(A) re-
ceptor subunit. Nature 385:

[103451]820–823, 1997.

[103452]18063.

[103453]Sinkkonen, S. T.; Hanna, M. C.; Kirkness, E. F.; Korpi, E. R.:

[103454]GABA-A receptor epsilon and theta subunits display un-
usual structural

[103455]variation between species and are enriched in the rat locus ceruleus. J.

[103456]Neurosci. 20: 3588–3595, 2000.

[103457]18064.

[103458]Wilke, K.; Gaul, R.; Klauck, S. M.; Poustka, A.: A gene in human

[103459]chromosome band Xq28 (GABRE) defines a putative new subunit class

[103460]of the GABA(A) neurotransmitter receptor. Genomics 45: 1–10, 1997.

[103461]18065.

[103462]Mazzarella, R.; Ciccodicola, A.; Esposito, T.; Arcucci, A.; Migliaccio,

[103463]C.; Jones, C.; Schlessinger, D.; D'Urso, M.; D'Esposito, M.: Human

[103464]protein kinase C iota gene (PRKCI) is closely linked to the BTK gene

[103465]in Xq21.3. Genomics 26: 629–631, 1995.

[103466]18066.

[103467]Lafreniere, R. G.; Carrel, L.; Willard, H. F.: A novel trans-membrane

[103468] transporter encoded by the XPCT gene in Xq13.2. Hum.

Molec. Genet. 3:

[103469] 1133–1139, 1994.

[103470] 18067.

[103471] Baric, I.; Sarnavka, V.; Fumic, K.; Maradin, M.; Begovic, D.;

Ruiter,

[103472] J. P. N.; Wanders, R. J. A.: A new case of succinyl–

CoA:acetoacetate

[103473] transferase deficiency: favourable course despite very low
residual

[103474] activity. J. Inherit. Metab. Dis. 24: 81–82, 2001.

[103475] 18068.

[103476] Berry, G. T.; Fukao, T.; Mitchell, G. A.; Mazur, A.; Ciafre,

M.;

[103477] Gibson, J.; Kondo, N.; Palmieri, M. J.: Neonatal hypogly–

caemia in

[103478] severe succinyl–CoA:3–oxoacid CoA–transferase defi–
ciency. J. Inherit.

[103479] Metab. Dis. 24: 587–595, 2001.

[103480] 18069.

[103481] Fukao, T.; Mitchell, G. A.; Song, X.–Q.; Nakamura, H.;

Kassovska-Bratinova,

[103482]S.; Orii, K. E.; Wraith, J. E.; Besley, G.; Wanders, R. J. A.;

Niezen-Koning,

[103483]K. E.; Berry, G. T.; Palmieri, M.; Kondo, N.: Succinyl-

CoA:3-ketoacid

[103484]CoA transferase (SCOT): cloning of the human SCOT gene,
tertiary structural

[103485]modeling of the human SCOT monomer, and characteri-
zation of three

[103486]pathogenic mutations. Genomics 68: 144-151, 2000.

[103487]18070.

[103488]Fukao, T.; Song, X.-Q.; Watanabe, H.; Hirayama, K.;
Sakazaki, H.;

[103489]Shintaku, H.; Imanaka, M.; Orii, T.; Kondo, N.: Prenatal di-
agnosis

[103490]of succinyl-coenzyme A:3-ketoacid coenzyme A trans-
ferase deficiency. Prenatal

[103491]Diag. 16: 471-474, 1996.

[103492]18071.

[103493]Kassovska-Bratinova, S.; Fukao, T.; Song, X.-Q.; Duncan,
A. M.

[103494]Y.; Chen, H. S.; Robert, M.-F.; Perez-Cerda, C.; Ugarte, M.;

Chartrand,

[103495]C.; Vobecky, S.; Kondo, N.; Mitchell, G. A.: Succinyl

CoA:3-oxoacid

[103496]CoA transferase (SCOT): human cDNA cloning, human
chromosomal mapping

[103497]to 5p13, and mutation detection in a SCOT-deficient pa-
tient. Am.

[103498]J. Hum. Genet. 59: 519-528, 1996.

[103499]18072.

[103500]Middleton, B.; Day, R.; Lombes, A.; Saudubray, J. M.: In-
fantile

[103501]ketoacidosis associated with decreased activity of suc-
cinyl-CoA:3-ketoacid

[103502]CoA-transferase. J. Inherit. Metab. Dis. 10 (suppl. 2):
273-275,

[103503]1987.

[103504]18073.

[103505]Mitchell, G. A.; Kassovska-Bratinova, S.; Boukaftane, Y.;
Robert,

[103506]M.-F.; Wang, S. P.; Ashmarina, L.; Lambert, M.; Lapierre,
P.; Potier,

[103507]E.: Medical aspects of ketone body metabolism. Clin. In-

vest. Med. 18:

[103508]193–216, 1995.

[103509]18074.

[103510]Niezen-Koning, K. E.; Wanders, R. J. A.; Ruiter, J. P. N.;
Ijlst,

[103511]L.; Visser, G.; Reitsma-Bierens, W. C. C.; Heymans, H. S.
A.; Reijngoud,

[103512]D. J.; Smit, G. P. A.: Succinyl-CoA:acetoacetate transferase
deficiency:

[103513]identification of a new patient with a neonatal onset and
review of

[103514]the literature. Europ. J. Pediat. 156: 870–873, 1997.

[103515]18075.

[103516]Perez-Cerda, C.; Merinero, B.; Sanz, P.; Jimenez, A.; Her-
nandez,

[103517]C.; Garcia, M. J.; Ugarte, M.: A new case of succinyl-
CoA:acetoacetate

[103518]transferase deficiency. J. Inherit. Metab. Dis. 15: 371–373,
1992.

[103519]18076.

[103520]Sakazaki, H.; Hirayama, K.; Murakami, S.; Yonezawa, S.;

Shintaku,

[103521]H.; Sawada, Y.; Fukao, T.; Watanabe, H.; Orii, T.; Isshiki, G.: A

[103522]new Japanese case of succinyl-CoA:3-ketoacid CoA-transferase deficiency. J.

[103523]Inherit. Metab. Dis. 18: 323-325, 1995.

[103524]18077.

[103525]Snyderman, S. E.; Sansaricq, C.; Middleton, B.: Succinyl-CoA:3-ketoacid

[103526]CoA-transferase deficiency. Pediatrics 101: 709-711, 1998.

[103527]18078.

[103528]D'Adamo, P.; Fassone, L.; Gedeon, A.; Janssen, E. A. M.; Bione,

[103529]S.; Bolhuis, P. A.; Barth, P. G.; Wilson, M.; Haan, E.; Orstavik,

[103530]K. H.; Patton, M. A.; Green, A. J.; Zammarchi, E.; Donati, M. A.;

[103531]Tonio, D.: The X-linked gene G4.5 is responsible for different

[103532]infantile dilated cardiomyopathies. Am. J. Hum. Genet. 61: 862-867,

[103533]1997.

[103534]18079.

[103535]Gedeon, A. K.; Wilson, M. J.; Colley, A. C.; Sillence, D. O.;
Mulley,

[103536]J. C.: X linked fatal infantile cardiomyopathy maps to Xq28
and is

[103537]possibly allelic to Barth syndrome. J. Med. Genet. 32:
383–388,

[103538]1995.

[103539]18080.

[103540]Gecz, J.; Baker, E.; Donnelly, A.; Ming, J. E.; McDonald–
McGinn,

[103541]D. M.; Spinner, N. B.; Zackai, E. H.; Sutherland, G. R.; Mul–
ley, J.

[103542]C.: Fibroblast growth factor homologous factor 2 (FHF2):
gene structure,

[103543]expression and mapping to the Borjeson–Forss–
man–Lehmann syndrome region

[103544]in Xq26 delineated by a duplication breakpoint in a BFLS–
like patient. Hum.

[103545]Genet. 104: 56–63, 1999.

[103546]18081.

[103547]Lovec, H.; Hartung, H.; Verdier, A.-S.; Mattei, M.-G.; Birnbaum,

[103548]D.; Goldfarb, M.; Coulier, F.: Assignment of FGF13 to human chromosome

[103549]band Xq21 by in situ hybridization. Cytogenet. Cell Genet. 76: 183-184,

[103550]1997.

[103551]18082.

[103552]Smallwood, P. M.; Munoz-Sanjuan, I.; Tong, P.; Macke, J. P.; Hendry,

[103553]S. H. C.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.; Nathans,

[103554]J.: Fibroblast growth factor (FGF) homologous factors: new members

[103555]of the FGF family implicated in nervous system development. Proc.

[103556]Nat. Acad. Sci. 93: 9850-9857, 1996.

[103557]18083.

[103558]Jones, M. H.; Furlong, R. A.; Burkin, H.; Chalmers, I. J.; Brown,

[103559]G. M.; Khwaja, O.; Affara, N. A.: The Drosophila developmental gene

[103560]fat facets has a human homologue in Xp11.4 which escapes X-inactivation

[103561]and has related sequences on Yq11.2. Hum. Molec. Genet. 5: 1695-1701,

[103562]1996.

[103563]18084.

[103564]Caldas, C.; Hahn, S. A.; da Costa, L. T.; Redston, M. S.; Schutte,

[103565]M.; Seymour, A. B.; Weinstein, C. L.; Hruban, R. H.; Yeo, C. J.; Kern,

[103566]S. E.: Frequent somatic mutations and homozygous deletions of the

[103567]p16 (MTS1) gene in pancreatic adenocarcinoma. Nature Genet. 8: 27-32,

[103568]1994.

[103569]18085.

[103570]Liu, Q.; Yan, Y.-X.; McClure, M.; Nakagawa, H.; Fujimura, F.;

[103571]Rustgi, A. K.: MTS-1 (CDKN2) tumor suppressor gene deletions are

[103572]a frequent event in esophagus squamous cancer and pancreatic adenocarcinoma

[103573]cell lines. *Oncogene* 10: 619–622, 1995.

[103574]18086.

[103575]Whelan, A. J.; Bartsch, D.; Goodfellow, P. J.: Brief report:

[103576]a familial syndrome of pancreatic cancer and melanoma with a mutation

[103577]in the CDKN2 tumor-suppressor gene. *New Eng. J. Med.* 333: 975–977,

[103578]1995.

[103579]18087.

[103580]Schutte, M.; da Costa, L. T.; Hahn, S. A.; Moskaluk, C.; Hoque,

[103581]A. T. M. S.; Rozenblum, E.; Weinstein, C. L.; Bittner, M.; Meltzer,

[103582]P. S.; Trent, J. M.; Yeo, C. J.; Hruban, R. H.; Kern, S. E.: Identification

[103583]by representational difference analysis of a homozygous deletion in

[103584]pancreatic carcinoma that lies with the BRCA2 region. *Proc. Nat.*

[103585]Acad. Sci. 92: 5950–5954, 1995.

[103586]18088.

[103587]Stoffers, D. A.; Zinkin, N. T.; Stanojevic, V.; Clarke, W. L.;

[103588]Habener, J. F.: Pancreatic agenesis attributable to a single nucleotide

[103589]deletion in the human IPF1 gene coding sequence. Nature Genet. 15:

[103590]106–110, 1997.

[103591]18089.

[103592]Tacconi, P.; Ferrigno, P.; Cocco, L.; Cannas, A.; Tamburini, G.;

[103593]Bergonzi, P.; Giagheddu, M.: Sclerosteosis: report of a case in a

[103594]black African man. Clin. Genet. 53: 497–501, 1998.

[103595]18090.

[103596]Tanzi, R. E.; Petrukhin, K.; Chernov, I.; Pellequer, J. L.; Wasco,

[103597]W.; Ross, B.; Romano, D. M.; Parano, E.; Pavone, L.; Brzustowicz,

[103598]L. M.; Devoto, M.; Peppercorn, J.; Bush, A. I.; Sternlieb, I.; Pirastu,

[103599]M.; Gusella, J. F.; Evgrafov, O.; Penchaszadeh, G. K.;
Honig, B.;

[103600]Edelman, I. S.; Soares, M. B.; Scheinberg, I. H.; Gilliam, T.
C.:

[103601]The Wilson disease gene is a copper transporting ATPase
with homology

[103602]to the Menkes disease gene. Nature Genet. 5: 344–350,
1993.

[103603]18091.

[103604]Terada, K.; Nakako, T.; Yang, X.-L.; Iida, M.; Aiba, N.; Mi-
namiya,

[103605]Y.; Nakai, M.; Sakaki, T.; Miura, N.; Sugiyama, T.: Restora-
tion of

[103606]holoceruloplasmin synthesis in LEC rat after infusion of
recombinant

[103607]adenovirus bearing WND cDNA. J. Biol. Chem. 273:
1815–1820, 1998.

[103608]18092.

[103609]Thomas, G. R.; Forbes, J. R.; Roberts, E. A.; Walshe, J. M.;
Cox,

[103610]D. W.: The Wilson disease gene: spectrum of mutations
and their consequences. Nature

- [103611]Genet. 9: 210–217, 1995. Note: Correction. Nature Genet.
9: 451 only,
[103612]1995.
[103613]18093.
- [103614]Thomas, G. R.; Jensson, O.; Gudmundsson, G.;
Thorsteinsson, L.;
[103615]Cox, D. W.: Wilson disease in Iceland: a clinical and ge-
netic study. Am.
[103616]J. Hum. Genet. 56: 1140–1146, 1995.
[103617]18094.
- [103618]Thomas, G. R.; Roberts, E. A.; Walshe, J. M.; Cox, D. W.:
Haplotypes
[103619]and mutations in Wilson disease. Am. J. Hum. Genet. 56:
1315–1319,
[103620]1995.
[103621]18095.
- [103622]Beutler, E.; Kuhl, W.; Comings, D.: Hexosaminidase
isozyme in
[103623]type O Gm2 gangliosidosis (Sandhoff–Jatzkewitz disease).
Am. J. Hum.
[103624]Genet. 27: 628–638, 1975.

[103625]18096.

[103626]Boustany, R.-M. N.; Tanaka, A.; Nishimoto, J.; Suzuki, K.:
Genetic

[103627]cause of a juvenile form of Tay-Sachs disease in a
Lebanese child. Ann.

[103628]Neurol. 29: 104-107, 1991.

[103629]18097.

[103630]Brown, C. A.; Mahuran, D. J.: Beta-hexosaminidase
isozymes from

[103631]cells cotransfected with alpha and beta cDNA constructs:
analysis

[103632]of the alpha-subunit missense mutation associated with
the adult form

[103633]of Tay-Sachs disease. Am. J. Hum. Genet. 5: 497-508,
1993.

[103634]18098.

[103635]Chern, C. J.; Beutler, E.; Kuhl, W.; Gilbert, F.; Mellman, W.

[103636]J.; Croce, C. M.: Characterization of heteropolymeric hex-
osaminidase

[103637]A in human x mouse hybrid cells. Proc. Nat. Acad. Sci. 73:
3637-3640,

[103638]1976.

[103639]18099.

[103640]Chern, C. J.; Kennett, R.; Engel, E.; Mellman, W. J.; Croce, C.

[103641]M.: Assignment of the structural genes for the alpha sub-unit of hexosaminidase

[103642]A, mannosephosphate isomerase and pyruvate kinase to the region of

[103643]22-qter of human chromosome 15. Somat. Cell Genet. 3: 553-560, 1977.

[103644]18100.

[103645]Coulter-Mackie, M. B.: Molecular characterization of both alleles

[103646]in an unusual Tay-Sachs disease B1 variant. (Letter) Am. J. Hum.

[103647]Genet. 54: 1126-1127, 1994.

[103648]18101.

[103649]De Braekeleer, M.; Hechtman, P.; Andermann, E.; Kaplan, F.: The

[103650]French Canadian Tay-Sachs disease deletion mutation: identification

[103651]of probable founders. Hum. Genet. 89: 83–87, 1992.

[103652]18102.

[103653]De Gasperi, R.; Gama Sosa, M. A.; Battistini, S.; Yeretsian, J.;

[103654]Raghavan, S.; Zelnik, N.; Leshinsky, E.; Kolodny, E. H.:
Late-onset

[103655]G(M2)–gangliosidosis: Ashkenazi Jewish family with an
exon 5 mutation

[103656](tyr180–to–his) in the Hex A alpha–chain gene. Neurology
47: 547–552,

[103657]1996.

[103658]18103.

[103659]dos Santos, M. R.; Tanaka, A.; sa Miranda, M. C.; Ribeiro, M.

[103660]G.; Maia, M.; Suzuki, K.: G(M2)–gangliosidosis B1 variant:
analysis

[103661]of beta–hexosaminidase alpha gene mutations in 11 pa-
tients from a

[103662]defined region in Portugal. Am. J. Hum. Genet. 49:
886–890, 1991.

[103663]18104.

[103664]Drucker, L.; Navon, R.: Tay-Sachs disease in an Israeli Arab

[103665]family: trp26-to-stop in the alpha-subunit of hexosaminidase A. Hum.

[103666]Mutat. 2: 415-417, 1993.

[103667]18105.

[103668]Drucker, L.; Proia, R. L.; Navon, R.: Identification and rapid
[103669]detection of three Tay-Sachs mutations in the Moroccan Jewish population. Am.

[103670]J. Hum. Genet. 51: 371-377, 1992.

[103671]18106.

[103672]Fernandes, M.; Kaplan, F.; Natowicz, M.; Prence, E.;
Kolodny,

[103673]E.; Kaback, M.; Hechtman, P.: A new Tay-Sachs disease B1
allele in

[103674]exon 7 in two compound heterozygotes each with a second novel mutation. Hum.

[103675]Molec. Genet. 1: 759-761, 1992.

[103676]18107.

[103677]Fernandes, M. J. G.; Hechtman, P.; Boulay, B.; Kaplan, F.: A

[103678]chronic GM(2) gangliosidosis variant with a HEXA splicing defect:

[103679]quantitation of HEXA mRNAs in normal and mutant fibroblasts. *Europ.*

[103680]*J. Hum. Genet.* 5: 129–136, 1997.

[103681]18108.

[103682]Formiga, L. de F.; Poenaru, L.; Couronne, F.; Flori, E.; Eibel,

[103683]J. L.; Deminatti, M. M.; Savary, J. B.; Lai, J. L.; Gilgenkrantz,

[103684]S.; Pierson, M.: Interstitial deletion of chromosome 15: two cases. *Hum.*

[103685]*Genet.* 80: 401–404, 1988.

[103686]18109.

[103687]Goebel, H. H.; Stolte, G.; Kustermann-Kuhn, B.; Harzer, K.: B(1)

[103688]variant of G(M2) gangliosidosis in a 12-year-old patient. *Pediat.*

[103689]*Res.* 25: 89–93, 1989.

[103690]18110.

[103691]Gordon, B. A.; Gordon, K. E.; Hinton, G. G.; Cadera, W.; Feleki,

[103692]V.; Bayleran, J.; Hechtman, P.: Tay Sachs disease: B1 variant. *Pediat.*

[103693]*Neurol.* 4: 54–57, 1988.

[103694]18111.

[103695]Greenberg, D. A.; Kaback, M. M.: Estimation of the frequency

[103696]of hexosaminidase A variant alleles in the American Jewish population. *Am.*

[103697]*J. Hum. Genet.* 34: 444–451, 1982.

[103698]18112.

[103699]Natt, E.; Westphal, E.–M.; Toth–Fejel, S. E.; Magenis, R. E.;

[103700]Buist, N. R. M.; Rettenmeier, R.; Scherer, G.: Inherited and de novo

[103701]deletion of the tyrosine aminotransferase gene locus at 16q22.1–q22.3

[103702]in a patient with tyrosinemia type II. *Hum. Genet.* 77: 352–358,

[103703]1987.

[103704]18113.

[103705]Pelet, B.; Antener, I.; Faggioni, R.; Spahr, A.; Gautier, E.:

- [103706]Tyrosinemia without liver or renal damage with plantar and palmar
- [103707]keratosis and keratitis (hypertyrosinemia type II). *Helv. Paediat.*
- [103708]*Acta* 34: 177–183, 1979.
- [103709]18114.
- [103710]Rehak, A.; Selim, M. M.; Yadav, G.: Richner–Hanhart syndrome
- [103711](tyrosinaemia–II) (report of four cases without ocular involvement). *Brit.*
- [103712]*J. Derm.* 104: 469–475, 1981.
- [103713]18115.
- [103714]Rettenmeier, R.; Natt, E.; Zentgraf, H.; Scherer, G.: Isolation
- [103715]and characterization of the human tyrosine aminotransferase gene. *Nucleic*
- [103716]*Acids Res.* 18: 3853–3861, 1990.
- [103717]18116.
- [103718]Richner, H.: Hornhautaffektion bei Keratoma palmare et plantare
- [103719]hereditarium. *Klin. Mbl. Augenheilk.* 100: 580–588, 1938.

[103720]18117.

[103721]Schmid, W.; Muller, G.; Schutz, G.; Gluecksohn-Waelsch, S.: Deletions

[103722]near the albino locus on chromosome 7 of the mouse affect the level

[103723]of tyrosine aminotransferase mRNA. Proc. Nat. Acad. Sci. 82: 2866-2869,

[103724]1985.

[103725]18118.

[103726]Tallab, T.M.: Richner-Hanhart syndrome: importance of early diagnosis

[103727]and early intervention. J. Am. Acad. Dermatol. 35: 857-859, 1996.

[103728]18119.

[103729]Ventura, G.; Biasini, G.; Petrozzi, M.: Cheratomia palmo-plantare

[103730]dissipatum associato a lesioni corneali in due fratelli. Boll. Oculist. 44:

[103731]497-510, 1965.

[103732]18120.

[103733]Waardenburg, P. J.; Franceschetti, A.; Klein, D.: Genetics

and

[103734]Ophthalmology. Springfield, Ill.: Charles C Thomas (pub.)

1:

[103735]1961. Pp. 515-517.

[103736]18121.

[103737]Westphal, E.-M.; Natt, E.; Grimm, T.; Odievre, M.; Scherer,
G.

[103738]: The human tyrosine aminotransferase gene: characteri-
zation of restriction

[103739]fragment length polymorphisms and haplotype analysis in
a family with

[103740]tyrosinemia type II. Hum. Genet. 79: 260-264, 1988.

[103741]18122.

[103742]Perheentupa, J.: Autoimmune polyendocrinopathy-candido-
sis-ectodermal

[103743]dystrophy (APECED).In: Eriksson, A. W.; Forsius, H. R.;
Nevanlinna,

[103744]H. R.; Workman, P. L.; Norio, R. K.: Population Structure
and Genetic

[103745]Disorders. New York: Academic Press (pub.) 1980. Pp.
583-587.

[103746]18123.

[103747]Prader, A.: Personal Communication. Zurich, Switzerland
1972.

[103748]18124.

[103749]Ramsay, C.; Winqvist, O.; Puhakka, L.; Halonen, M.; Moro,
A.;

[103750]Kampe, O.; Eskelin, P.; Pelto-Huikko, M.; Peltonen, L.: Aire
deficient

[103751]mice develop multiple features of APECED phenotype and
show altered

[103752]immune response. Hum. Molec. Genet. 11: 397-409,
2002.

[103753]18125.

[103754]Rinderle, C.; Christensen, H.-M.; Schweiger, S.; Lehrach,
H.;

[103755]Yaspo, M.-L.: AIRE encodes a nuclear protein co-localizing
with cytoskeletal

[103756]filaments: altered sub-cellular distribution of mutants
lacking PHD

[103757]zinc fingers. Hum. Molec. Genet. 8: 277-290, 1999.

[103758]18126.

[103759]Rosatelli, M. C.; Meloni, A.; Meloni, A.; Devoto, M.; Cao,
A.;

- [103760] Scott, H. S.; Peterson, P.; Heino, M.; Krohn, K. J.;
Nagamine, K.;
- [103761] Kudoh, J.; Shimizu, N.; Antonarakis, S. E.: A common mu-
tation in
- [103762] Sardinian autoimmune polyendocrinopathy-candidia-
sis-ectodermal dystrophy
- [103763] patients. Hum. Genet. 103: 428-434, 1998.
- [103764] 18127.
- [103765] Saenger, P.: Primary hypoaldosteronism due to zona
glomerulosa
- [103766] defect. (Letter) New Eng. J. Med. 310: 1394 only, 1984.
- [103767] 18128.
- [103768] Saenger, P.; Levine, L. S.; Irvine, W. J.; Gottesdiener, K.;
- Rauh,
- [103769] W.; Sonino, N.; Chow, D.; New, M. I.: Progressive adrenal
failure
- [103770] in polyglandular autoimmune disease. J. Clin. Endocr.
Metab. 54:
- [103771] 863-868, 1982.
- [103772] 18129.
- [103773] Scott, H. S.; Heino, M.; Peterson, P.; Mittaz, L.; Lalioti, M.

[103774]D.; Betterle, C.; Cohen, A.; Seri, M.; Lerone, M.; Romeo, G.; Collin,
 [103775]P.; Salo, M.; and 9 others: Common mutations in autoimmune polyendocrinopathy–candidiasis–ectodermal
 [103776]dystrophy patients of different origins. *Molec. Endocr.* 12: 1112–1119,
 [103777]1998.
 [103778]18130.
 [103779]Shapiro, M. S.; Zamir, R.; Weiss, E.; Radnay, J.; Shenkman, L.
 [103780]: The polyglandular deficiency syndrome: a new variant in Persian
 [103781]Jews. *J. Endocr. Invest.* 10: 1–7, 1987.
 [103782]18131.
 [103783]Spinner, M. W.; Blizzard, R. M.; Childs, B.: Clinical and genetic
 [103784]heterogeneity in idiopathic Addison's disease and hypoparathyroidism. *J.*
 [103785]*Clin. Endocr.* 28: 795–804, 1968.
 [103786]18132.
 [103787]Sweetnam, W. P.: Juvenile familial endocrinopathy. *Lancet*

I:

[103788]463–465, 1966.

[103789]18133.

[103790]Wang, C.-Y.; Davoodi-Semiromi, A.; Huang, W.; Connor, E.; Shi,

[103791]J.-D.; She, J.-X.: Characterization of mutations in patients with

[103792]autoimmune polyglandular syndrome type 1 (APS1). Hum. Genet. 103:

[103793]681–685, 1998.

[103794]18134.

[103795]Whitaker, J. A.; Landing, B. H.; Esselborn, V. M.; Williams, R.

[103796]R.: Syndrome of familial juvenile hypoadrenocorticism, hypoparathyroidism

[103797]and superficial moniliasis. J. Clin. Endocr. 16: 1374–1387, 1956.

[103798]18135.

[103799]Wirfalt, A.: Genetic heterogeneity in autoimmune polyglandular

[103800]failure. Acta Med. Scand. 210: 7–13, 1981.

[103801]18136.

[103802]Zlotogora, J.; Shapiro, M. S.: Polyglandular autoimmune syndrome

[103803]type I among Iranian Jews. J. Med. Genet. 29: 824–826, 1992.

[103804]18137.

[103805]Hediger, M. A.: New view at C. Nature Med. 8: 445–446, 2002.

[103806]18138.

[103807]Sotiriou, S.; Gispert, S.; Cheng, J.; Wang, Y.; Chen, A.; Hoogstraten–Miller,

[103808]S.; Miller, G. F.; Kwon, O.; Levine, M.; Guttentag, S. H.; Nussbaum,

[103809]R. L.: Ascorbic–acid transporter Slc23a1 is essential for vitamin

[103810]C transport into the brain and for perinatal survival. Nature Med. 8:

[103811]514–517, 2002.

[103812]18139.

[103813]Chuang, D. T.; Fisher, C. W.; Lau, K. S.; Griffin, T. A.; Wynn,

[103814]R. M.; Cox, R. P.: Maple syrup urine disease: domain structure, mutations

[103815]and exon skipping in the dihydrolipoyl transacylase (E2) component

[103816]of the branched-chain alpha-keto acid dehydrogenase complex. Molec.

[103817]Biol. Med. 8: 49-63, 1991.

[103818]18140.

[103819]Chuang, J. L.; Cox, R. P.; Chuang, D. T.: E2 transacylase-deficient

[103820](type II) maple syrup urine disease: aberrant splicing of E2 mRNA

[103821]caused by internal intronic deletions and association with thiamine-responsive

[103822]phenotype. J. Clin. Invest. 100: 736-744, 1997.

[103823]18141.

[103824]Fisher, C. W.; Fisher, C. R.; Chuang, J. L.; Lau, K. S.; Chuang,

[103825]D. T.; Cox, R. P.: Occurrence of a 2-bp (AT) deletion allele and

[103826]nonsense (G-to-T) mutant allele at the E2 (DBT) locus of six patients

[103827]with maple syrup urine disease: multiple-exon skipping as a secondary

[103828]effect of the mutations. Am. J. Hum. Genet. 52: 414-424, 1993.

[103829]18142.

[103830]Fisher, C. W.; Lau, K. S.; Fisher, C. R.; Wynn, R. M.; Cox, R.

[103831]P.; Chuang, D. T.: A 17-bp insertion and a phe215-to-cys missense

[103832]mutation in the dihydrolipoyl transacylase (E2) mRNA from a thiamine-responsive

[103833]maple syrup urine disease patient WG-34. Biochem. Biophys. Res. Commun. 174:

[103834]804-809, 1991.

[103835]18143.

[103836]Herring, W. J.; Litwer, S.; Weber, J. L.; Danner, D. J.: Molecular

[103837]genetic basis of maple syrup urine disease in a family with two defective

[103838]alleles for branched chain acyltransferase and localization of the

[103839]gene to human chromosome 1. Am. J. Hum. Genet. 48: 342-350, 1991.

[103840]18144.

[103841]Hummel, K. B.; Litwer, S.; Bradford, A. P.; Aitken, A.; Dan-
ner,

[103842]D. J.; Yeaman, S. J.: Nucleotide sequence of a cDNA for
branched

[103843]chain acyltransferase with analysis of the deduced protein
structure. J.

[103844]Biol. Chem. 263: 6165–6168, 1988.

[103845]18145.

[103846]Indo, Y.; Akaboshi, I.; Nobukuni, Y.; Endo, F.; Matsuda, I.:
Maple

[103847]syrup urine disease: a possible biochemical basis for the
clinical

[103848]heterogeneity. Hum. Genet. 80: 6–10, 1988.

[103849]18146.

[103850]Lau, K. S.; Eddy, R. L.; Shows, T. B.; Fisher, C. W.; Chuang,

[103851]D. T.; Cox, R. P.: Localization of the dihydrolipoamide
branched–chain

[103852]transacylase gene (DBT) of the human branched–chain
keto acid dehydrogenase

[103853]complex to chromosome 1. Cytogenet. Cell Genet. 56:

33–35, 1991.

[103854]18147.

[103855]Lau, K. S.; Griffin, T. A.; Hu, C.-W. C.; Chuang, D. T.: Conservation

[103856]of primary structure in the lipoyl-bearing and dihydrolipoyl dehydrogenase

[103857]binding domains of mammalian branched-chain alpha-keto acid dehydrogenase

[103858]complex: molecular cloning of human and bovine transacylase (E2) cDNAs. *Biochemistry* 27:

[103859]1972–1981, 1988.

[103860]18148.

[103861]Lau, K. S.; Herring, W. J.; Chuang, J. L.; McKean, M.; Danner,

[103862]D. J.; Cox, R. P.; Chuang, D. T.: Structure of the gene encoding

[103863]dihydrolipoyl transacylase (E2) component of human branched chain

[103864]alpha-keto acid dehydrogenase and characterization of an E2 pseudogene. *J.*

[103865]*Biol. Chem.* 267: 24090–24096, 1992.

[103866]18149.

[103867]Lebo, R. V.; Shapiro, L. R.; Fenerci, E. Y.; Hoover, J. M.;
Chuang,
[103868]J. L.; Chuang, D. T.; Kronn, D. F.: Rare etiology of autosomal recessive
[103869]disease in a child with noncarrier parents. Am. J. Hum.
Genet. 67:
[103870]750–754, 2000.
[103871]18150.
[103872]Litwer, S.; Danner, D. J.: Identification of a cDNA clone in
[103873]lambda-gt11 for the transacylase component of branched
chain ketoacid
[103874]dehydrogenase. Biochem. Biophys. Res. Commun. 131:
961–967, 1985.
[103875]18151.
[103876]Litwer, S.; Danner, D. J.: Mitochondrial import and processing
[103877]of an in vitro synthesized human prebranched chain acyltransferase
[103878]fragment. Am. J. Hum. Genet. 43: 764–769, 1988.
[103879]18152.
[103880]Mitsubuchi, H.; Nobukuni, Y.; Akaboshi, I.; Indo, Y.; Endo,

F.;

[103881]Matsuda, I.: Maple syrup urine disease caused by a partial deletion

[103882]in the inner E2 core domain of the branched chain alpha-keto acid

[103883]dehydrogenase complex due to aberrant splicing: a single base deletion

[103884]at a 5-prime-splice donor site of an intron of the E2 gene disrupts

[103885]the consensus sequence in this region. J. Clin. Invest. 87: 1207-1211,

[103886]1991.

[103887]18153.

[103888]Tsuruta, M.; Mitsubuchi, H.; Mardy, S.; Miura, Y.; Hayashida,

[103889]Y.; Kinugasa, A.; Ishitsu, T.; Matsuda, I.; Indo, Y.: Molecular basis

[103890]of intermittent maple syrup urine disease: novel mutations in the

[103891]E2 gene of the branched-chain alpha-keto acid dehydrogenase complex. J.

[103892]Hum. Genet. 43: 91-100, 1998.

[103893]18154.

[103894]Zneimer, S. M.; Lau, K. S.; Eddy, R. L.; Shows, T. B.;
Chuang,

[103895]J. L.; Chuang, D. T.; Cox, R. P.: Regional assignment of two
genes

[103896]of the human branched-chain alpha-keto acid dehydro-
genase complex:

[103897]the E1-beta gene (BCKDHB) to chromosome 6p21-22 and
the E2 gene (DBT)

[103898]to chromosome 1p31. Genomics 10: 740-747, 1991.

[103899]18155.

[103900]Gahl, W.; Finkelstein, J.; Martin, J.; Mullen, K.; Hoofnagle,
J.;

[103901]Backlund, P.; Bernar, J.; Bernardini, I.: Hepatic methionine
adenosyltransferase

[103902]deficiency in a 31-year-old man. (Abstract) Am. J. Hum.
Genet. 37:

[103903]A8 only, 1985.

[103904]18156.

[103905]Gahl, W. A.; Bernardini, I.; Finkelstein, J. D.; Tangerman,
A.;

- [103906]Martin, J. J.; Blom, H. J.; Mullen, K. D.; Mudd, S. H.:
Transsulfuration
[103907]in an adult with hepatic methionine adenosyltransferase
deficiency. J.
[103908]Clin. Invest. 81: 390–397, 1988.
[103909]18157.
[103910]Gahl, W. A.; Finkelstein, J. D.; Mullen, K. D.; Bernardini, I.;
[103911]Martin, J. J.; Backlund, P.; Ishak, K. G.; Hoofnagle, J. H.;
Mudd,
[103912]S. H.: Hepatic methionine adenosyltransferase deficiency
in a 31-year-old
[103913]man. Am. J. Hum. Genet. 40: 39–49, 1987.
[103914]18158.
[103915]Gaull, G. E.; Bender, A. N.; Vulovic, D.; Tallan, H. H.;
Schaffner,
[103916]F.: Methioninemia and myopathy: a new disorder. Ann.
Neurol. 9:
[103917]423–432, 1981.
[103918]18159.
[103919]Gaull, G. E.; Tallan, H. H.: Methionine adenosyltransferase
deficiency:

- [103920]new enzymatic defect associated with hypermethionine-
mia. Science 186:
- [103921]59-60, 1974.
- [103922]18160.
- [103923]Gaull, G. E.; Tallan, H. H.; Lonsdale, D.; Przyrembel, H.;
Schaffner,
- [103924]F.; von Bassewitz, D. B.: Hypermethioninemia associated
with methionine
- [103925]adenosyltransferase deficiency: clinical, morphologic, and
biochemical
- [103926]observations in four patients. J. Pediat. 98: 734-741,
1981.
- [103927]18161.
- [103928]Gout, J.-P.; Serre, J.-C.; Dieterlen, M.; Antener, I.; Frappat,
- [103929]P.; Bost, M.; Beaudoin, A.: Une nouvelle cause
d'hypermethioninemie
- [103930]de l'enfant: le deficit en S-adenosyl-
methionine-synthetase. Arch.
- [103931]Franc. Pediat. 34: 416-423, 1977.
- [103932]18162.
- [103933]Hazelwood, S.; Bernardini, I.; Shotelersuk, V.; Tangerman,

A.;

[103934]Guo, J.; Mudd, H.; Gahl, W. A.: Normal brain myelination in a patient

[103935]homozygous for a mutation that encodes a severely truncated methionine

[103936]adenosyltransferase I/III. Am. J. Med. Genet. 75: 395-400, 1998.

[103937]18163.

[103938]Horikawa, S.; Tsukada, K.: Molecular cloning and nucleotide sequence

[103939]of cDNA encoding the human liver S-adenosylmethionine synthetase. Biochem.

[103940]Int. 25: 81-90, 1991.

[103941]18164.

[103942]Hug, G.; Cussen, L. J.; Schubert, W. K.; Chuck, G.: 'Fingerprints'

[103943]in the liver: loss of methionine-activating enzyme. J. Clin. Invest. 47:

[103944]49A-50A, 1968.

[103945]18165.

[103946]Meny, R. G.; Gutberlet, R. L.; Ozand, P.; Morris, C.; Kim, C.

[103947]H.: Hypermethioninemia in an infant. Am. J. Dis. Child.
132: 261–262,

[103948]1978.

[103949]18166.

[103950]Mudd, S. H.; Levy, H. L.; Tangerman, A.; Boujet, C.; Buist,
N.;

[103951]Davidson–Mundt, A.; Hudgins, L.; Oyanagi, K.; Nagao, M.;
Wilson, W.

[103952]G.: Isolated persistent hypermethioninemia. Am. J. Hum.
Genet. 57:

[103953]882–892, 1995.

[103954]18167.

[103955]Natori, Y.: Studies on ethionine. VI. Sex-dependent behavior

[103956]of methionine and ethionine in rats. J. Biol. Chem. 238: 2075–2080,

[103957]1963.

[103958]18168.

[103959]Sakata, S. F.; Shelly, L. L.; Ruppert, S.; Schutz, G.; Chou, J.

[103960]Y.: Cloning and expression of murine S-adenosylmethionine synthetase. J.

[103961]Biol. Chem. 268: 13978–13986, 1993.

[103962]18169.

[103963]Surtees, R.; Leonard, J.; Austin, S.: Association of demyelination

[103964]with deficiency of cerebrospinal-fluid S-adenosylmethionine in inborn

[103965]errors of methyl-transfer pathway. Lancet 338: 1550–1554, 1991.

[103966]18170.

[103967]Ubagai, T.; Lei, K.-J.; Huang, S.; Mudd, S. H.; Levy, H. L.;

Chou,

[103968]J. Y.: Molecular mechanisms of an inborn error of methionine pathway:

[103969]methionine adenosyltransferase deficiency. J. Clin. Invest. 96:

[103970]1943–1947, 1995.

[103971]18171.

[103972]Cahane, M.; Treister, G.; Abraham, F. A.; Melamed, S.:
Glaucoma

[103973]in siblings with Morquio syndrome. Brit. J. Ophthal. 74:
382–383,

[103974]1990.

[103975]18172.

[103976]DiFerrante, N. M.; Ginsburg, L. C.; Donnelly, P. V.; DiFerrante,

[103977]D. T.; Caskey, C. T.: Deficiencies of glucosamine–6–sulfate
or galactosamine–6–sulfate

[103978]sulfatases are responsible for different mucopolysacchari-
doses. Science 199:

[103979]79–81, 1978.

[103980]18173.

- [103981]Fujimoto, A.; Horwitz, A. L.: Biochemical defect of non-keratan-sulfate-excreting
- [103982]Morquio syndrome. Am. J. Med. Genet. 15: 265-273, 1983.
- [103983]18174.
- [103984]Fukuda, S.; Tomatsu, S.; Masue, M.; Sukegawa, K.; Iwata, H.; Ogawa,
- [103985]T.; Nakashima, Y.; Hori, T.; Yamagishi, A.; Hanyu, Y.; Morooka, K.;
- [103986]Kiman, T.; Hashimoto, T.; Orie, T.: Mucopolysaccharidosis type IVA:
- [103987]N-acetyl-galactosamine-6-sulfate sulfatase exonic point mutations in
- [103988]classical Morquio and mild cases. J. Clin. Invest. 90: 1049-1053,
- [103989]1992.
- [103990]18175.
- [103991]Fukuda, S.; Tomatsu, S. Masuno, M.; Ogawa, T.; Yamagishi, A.;
- [103992]Rezvi, G. M. M.; Sukegawa, K.; Shimozawa, N.; Suzuki, Y.; Kondo, N.;
- [103993]Imaizumi, K.; Kuroki, Y.; Okabe, T.; Orie, T.: Mucopolysac-

charidosis

[103994]IVA: submicroscopic deletion of 16q24.3 and a novel
R386C mutation

[103995]of N-acetylgalactosamine-6-sulfate sulfatase gene in a
classical Morquio

[103996]disease. Hum. Mutat. 7: 123-134, 1996.

[103997]18176.

[103998]Gadbois, P.; Moreau, J.; Laberge, C.: La maladie de
Morquio dans

[103999]la province de Quebec. Un. Med. Canada 102: 602-607,
1973.

[104000]18177.

[104001]Gibson, G. J.; Saccone, G. T. P.; Brooks, D. A.; Clements, P.

[104002]R.; Hopwood, J. J.: Human N-acetyl-
galactosamine-4-sulphate sulphatase:

[104003]purification, monoclonal antibody production and native
and subunit

[104004]M(r) values. Biochem. J. 248: 755-764, 1987.

[104005]18178.

[104006]Glossl, J.; Kresse, H.; Mendla, K.; Cantz, M.; Rosenkranz,
W.

[104007]: Partial deficiency of glycoprotein neuraminidase in some patients

[104008]with Morquio disease type A. *Pediat. Res.* 18: 302–305, 1984.

[104009]18179.

[104010]Glossl, J.; Lembeck, K.; Gamse, G.; Kresse, H.: Morquio's disease

[104011]type A: absence of material cross reacting with antibodies against

[104012]N-acetylglactosamine-6-sulfate sulfatase. *Hum. Genet.* 54: 87–91,

[104013]1980.

[104014]18180.

[104015]Greenberg, A. D.: Atlantoaxial dislocations. *Brain* 91: 655–684,

[104016]1968.

[104017]18181.

[104018]Guiney, W. B., Jr.; Stevenson, R. E.: Morquio disease (MPS IV)

[104019]with survival to age 67 years. *Proc. Greenwood Genet. Center* 1:

[104020]84–87, 1982.

[104021]18182.

[104022]Hecht, J. T.; Scott, C. I., Jr.; Smith, T. K.; Williams, J. C.

[104023]: Mild manifestations of the Morquio syndrome. (Letter)
Am. J. Med.

[104024]Genet. 18: 369–371, 1984.

[104025]18183.

[104026]Holzgreve, W.; Grobe, H.; von Figura, K.; Kresse, H.; Beck,
H.;

[104027]Mattei, J. F.: Morquio syndrome: clinical findings in 11 pa-
tients

[104028]with MPS IVA and 2 patients with MPS IVB. Hum. Genet.
57: 360–365,

[104029]1981.

[104030]18184.

[104031]Hori, T.; Tomatsu, S.; Nakashima, Y.; Uchiyama, A.;
Fukuda, S.;

[104032]Sukegawa, K.; Shimozawa, N.; Suzuki, Y.; Kondo, N.; Hori-
uchi, T.;

[104033]Ogura, S.; Orii, T.: Mucopolysaccharidosis type IVA: com-
mon double

[104034]deletion in the N-acetylgalactosamine-6-sulfatase gene
(GALNS). Genomics 26:

[104035]535-542, 1995.

[104036]18185.

[104037]Hussels, I.: Morquio syndrome in a woman with two normal children.In:

[104038]Bergsma, D.: Skeletal Dysplasias. Amsterdam: Excerpta
Medica (pub.)

[104039]1974. Pp. 465-466.

[104040]18186.

[104041]Kato, Z.; Fukuda, S.; Tomatsu, S.; Vega, H.; Yasunaga, T.;
Yamagishi,

[104042]A.; Yamada, N.; Valencia, A.; Barrera, L. A.; Sukegawa, K.;
Orii,

[104043]T.; Kondo, N.: A novel common missense mutation G301C
in the N-acetylgalactosamine-6-sulfate

[104044]sulfatase gene in mucopolysaccharidosis IVA. Hum. Genet.
101: 97-101,

[104045]1997.

[104046]18187.

[104047]Langer, L. O., Jr.; Carey, L. S.: The roentgenographic fea-

tures

[104048]of the KS mucopolysaccharidosis of Morquio

(Morquio-Brailsford's disease). Am.

[104049]J. Roentgen. 97: 1-20, 1966.

[104050]18188.

[104051]Levin, L. S.; Jorgenson, R. J.; Salinas, C. F.: Oral findings

[104052]in the Morquio syndrome (mucopolysaccharidosis IV). Oral

Surg. Oral

[104053]Med. Oral Path. 39: 390-395, 1975.

[104054]18189.

[104055]Linker, A.; Evans, L. R.; Langer, L. O., Jr.: Morquio's dis-

ease

[104056]and mucopolysaccharide excretion. J. Pediat. 77:

1039-1047, 1970.

[104057]18190.

[104058]Maroteaux, P.; Lamy, M.: Opacités corneennes et troubles

metaboliques

[104059]dans la maladie de Morquio. Rev. Franc. Etud. Clin. Biol. 6:

481-483,

[104060]1961.

[104061]18191.

[104062]Masuno, M.; Tomatsu, S.; Nakashima, Y.; Hori, T.; Fukuda, S.;

[104063]Masue, M.; Sukegawa, K.; Orie, T.: Mucopolysaccharidosis
IV A: assignment

[104064]of the human N-acetylgalactosamine-6-sulfate sulfatase
(GALNS) gene

[104065]to chromosome 16q24. Genomics 16: 777-778, 1993.
[104066]18192.

[104067]Matalon, R.; Arbogast, B.; Dorfman, A.: Morquio's syn-
drome: a

[104068]deficiency of chondroitin sulfate N-acetylhexosamine sul-
fate sulfatase.

[104069](Abstract) Pediat. Res. 8: 436, 1974.
[104070]18193.

[104071]McKusick, V. A.: Osler as a medical geneticist. Johns Hop-
kins

[104072]Med. J. 139: 163-174, 1976.
[104073]18194.

[104074]McKusick, V. A.: Heritable Disorders of Connective Tissue.
Non-keratan-sulfate-excreting

[104075]Morquio syndrome. St. Louis: C. V. Mosby (pub.) 1972. Pp.

600-604.

[104076]Note: Fig. 11-30...

[104077]18195.

[104078]Morquio, L.: Sur une forme de dystrophie osseuse familiale. Bull.

[104079]Soc. Pediat. Paris 27: 145-152, 1929.

[104080]18196.

[104081]Morris, C. P.; Guo, X.-H.; Apostolou, S.; Hopwood, J. J.; Scott,

[104082]H. S.: Morquio A syndrome: cloning, sequence, and structure of the

[104083]human N-acetylgalactosamine 6-sulfatase (GALNS) gene. Genomics 22:

[104084]652-654, 1994.

[104085]18197.

[104086]Nakashima, Y.; Tomatsu, S.; Hori, T.; Fukuda, S.; Sukegawa, K.;

[104087]Kondo, N.; Suzuki, Y.; Shimozawa, N.; Orii, T.: Mucopolysaccharidosis

[104088]IV A: molecular cloning of the human N-acetylgalactosamine-6-sulfatase

[104089]gene (GALNS) and analysis of the 5-prime-flanking region. Genomics 20:

[104090]99-104, 1994.

[104091]18198.

[104092]Nelson, J.; Broadhead, D.; Mossman, J.: Clinical findings in

[104093]12 patients with MPS IV A (Morquio's disease): further evidence for

[104094]heterogeneity. Part I: clinical and biochemical findings. Clin. Genet. 33:

[104095]111-120, 1988.

[104096]18199.

[104097]Nelson, J.; Kinirons, M.: Clinical findings in 12 patients with

[104098]MPS IV A (Morquio's disease): further evidence for heterogeneity.

[104099]Part II: dental findings. Clin. Genet. 33: 121-125, 1988.

[104100]18200.

[104101]Nelson, J.; Thomas, P. S.: Clinical findings in 12 patients with

[104102]MPS IV A (Morquio's disease): further evidence for heterogeneity.

- [104103]Part III: odontoid dysplasia. Clin. Genet. 33: 126–130,
1988.
- [104104]18201.
- [104105]Osler, W.: Sporadic cretinism in America. Am. J. Med. Sci.
114:
[104106]377–401, 1897.
- [104107]18202.
- [104108]Nauseef, W. M.; Brigham, S.; Cogley, M.: Hereditary
myeloperoxidase
[104109]deficiency due to a missense mutation of arginine 569 to
tryptophan. J.
[104110]Biol. Chem. 269: 1212–1216, 1994.
- [104111]18203.
- [104112]Bartholomew, L. G.; Baggenstoss, A. H.; Morlock, C. G.;
Comfort,
[104113]M. W.: Primary atrophy and lipomatosis of the pancreas.
Gastroenterology 36:
[104114]563–572, 1959.
- [104115]18204.
- [104116]Bodian, M.; Sheldon, W.; Lightwood, R.: Congenital hy-
poplasia

[104117]of the exocrine pancreas. Acta Paediat. 53: 282-293,
1964.

[104118]18205.

[104119]Cipolli, M.; D'Orazio, C.; Delmarco, A.; Marchesini, C.; Mi-
ano,

[104120]A.; Mastella, G.: Shwachman's syndrome: pathomorphosis
and long-term

[104121]outcome. J. Pediat. Gastroent. Nutr. 29: 265-272, 1999.

[104122]18206.

[104123]Danks, D. M.; Haslam, R. H. A.; Mayne, V.; Kaufmann, H. J.;
Holtzapfel,

[104124]P. G.: Metaphyseal chondrodysplasia, neutropenia, and
pancreatic

[104125]insufficiency presenting with respiratory distress in the
neonatal

[104126]period. Arch. Dis. Child. 51: 697-701, 1976.

[104127]18207.

[104128]Davie, A. M.: The 'singles' method for segregation analysis
under

[104129]incomplete ascertainment. Ann. Hum. Genet. 42:
507-512, 1979.

[104130]18208.

[104131]Dokal, I.; Rule, S.; Chen, F.; Potter, M.; Goldman, J.: Adult

[104132]onset of acute myeloid leukaemia (M6) in patients with
Shwachman–Diamond

[104133]syndrome. Brit. J. Haemat. 99: 171–173, 1997.

[104134]18209.

[104135]Dror, Y.; Freedman, M. H.: Shwachman–Diamond syn-
drome. Brit.

[104136]J. Haemat. 118: 701–713, 2002.

[104137]18210.

[104138]Dror, Y.; Freedman, M. H.: Shwachman–Diamond syn-
drome: an inherited

[104139]preleukemic bone marrow failure disorder with aberrant
hematopoietic

[104140]progenitors and faulty marrow microenvironment. Blood
94: 3048–3054,

[104141]1999.

[104142]18211.

[104143]Dror, Y.; Freedman, M. H.: Shwachman–Diamond syn-
drome marrow

[104144]cells show abnormally increased apoptosis mediated through the Fas

[104145]pathway. Blood 97: 3011–3016, 2001.

[104146]18212.

[104147]Dror, Y.; Ginzberg, H.; Dalal, I.; Cherepanov, V.; Downey, G.;

[104148]Durie, P.; Roifman, C. M.; Freedman, M. H.: Immune function in patients

[104149]with Shwachman–Diamond syndrome. Brit. J. Haemat. 114: 712–717,

[104150]2001.

[104151]18213.

[104152]Fraccaro, M.; Scappaticci, S.; Arico, M.: Shwachman syndrome

[104153]and chromosome breakage. (Letter) Hum. Genet. 79: 194, 1988.

[104154]18214.

[104155]Genieser, N. B.; Halac, E. R.; Greco, M. A.; Selvarani Richards,

[104156]H. M.: Shwachman–Bodian syndrome. J. Comput. Assist. Tomogr. 6:

[104157]1191–1192, 1982.

[104158]18215.

[104159]Ginzberg, H.; Shin, J.; Ellis, L.; Goobie, S.; Morrison, J.;
Corey,

[104160]M.; Durie, P. R.; Rommens, J. M.: Segregation analysis in
Shwachman–Diamond

[104161]syndrome: evidence for recessive inheritance. Am. J. Hum.
Genet. 66:

[104162]1413–1416, 2000.

[104163]18216.

[104164]Ginzberg, H.; Shin, J.; Ellis, L.; Morrison, J.; Ip, W.; Dror,

[104165]Y.; Freedman, M.; Heitlinger, L. A.; Belt, M. A.; Corey, M.;
Rommens,

[104166]J. M.; Durie, P. R.: Shwachman syndrome: phenotypic
manifestations

[104167]of sibling sets and isolated cases in a large patient cohort
are similar. J.

[104168]Pediat. 135: 81–88, 1999.

[104169]18217.

[104170]Goldstein, R.: Congenital lipomatosis of the pancreas.
Malabsorption,

[104171]dwarfism, leukopenia with relative granulocytopenia and thrombocytopenia. Clin.

[104172]Pediat. 7: 419–422, 1968.

[104173]18218.

[104174]Goobie, S.; Morrison, J.; Ginzberg, H.; Ellis, L.; Corey, M.;

[104175]Masuno, M.; Imaizumi, K.; Kuroki, Y.; Fujiwara, T. M.; Morgan, K.;

[104176]Durie, P. R.; Rommens, J. M.: Exclusion of linkage of Shwachman–Diamond

[104177]syndrome to chromosome regions 6q and 12q implicated by a de novo

[104178]translocation. Am. J. Med. Genet. 85: 171–174, 1999.

[104179]18219.

[104180]Li, C. C.; Mantel, N.: A simple method of estimating the segregation

[104181]ratio under complete ascertainment. Am. J. Hum. Genet. 20: 61–81,

[104182]1968.

[104183]18220.

[104184]Mack, D. R.; Forstner, G. G.; Wilschanski, M.; Freedman, M. H.;

[104185]Durie, P. R.: Shwachman syndrome: exocrine pancreatic dysfunction

[104186]and variable phenotypic expression. Gastroenterology 111: 1593–1602,

[104187]1996.

[104188]18221.

[104189]Masuno, M.; Imaizumi, K.; Nishimura, G.; Nakamura, M.; Saito,

[104190]I.; Akagi, K.; Kuroki, Y.: Shwachman syndrome associated with de

[104191]novo reciprocal translocation t(6;12)(q16.2;q21.2). J. Med. Genet. 32:

[104192]894–895, 1995.

[104193]18222.

[104194]McLennan, T. W.; Steinbach, H. L.: Shwachman's syndrome: the

[104195]broad spectrum of bony abnormalities. Radiology 112: 167–173, 1974.

[104196]18223.

[104197]Goobie, S.; Popovic, M.; Morrison, J.; Ellis, L.; Ginzberg, H.;

[104198]Boocock, G. R. B.; Ehtesham, N.; Betard, C.; Brewer, C. G.;

Roslin,

[104199]N. M.; Hudson, T. J.; Morgan, K.; Fujiwara, T. M.; Durie, P.
R.; Rommens,

[104200]J. M.: Shwachman–Diamond syndrome with exocrine pan-
creatic dysfunction

[104201]and bone marrow failure maps to the centromeric region
of chromosome

[104202]7. Am. J. Hum. Genet. 68: 1048–1054, 2001.

[104203]18224.

[104204]Nezelof, C.; Watchi, M.: L'hypoplasie congenitale lipoma-
teuse

[104205]du pancreas exocrine chez l'enfant (Deux observations et
revue de

[104206]la litterature). Arch. Franc. Pediat. 18: 1135–1172, 1961.

[104207]18225.

[104208]Pringle, E. M.; Young, W. F.; Haworth, E. M.: Syndrome of
pancreatic

[104209]insufficiency, blood dyscrasia and metaphyseal dysplasia.
Proc. Roy.

[104210]Soc. Med. 61: 776–777, 1968.

[104211]18226.

[104212]Rothbaum, R. J.; Williams, D. A.; Daugherty, C. C.: Unusual surface

[104213]distribution of concanavalin A reflects a cytoskeletal defect in neutrophils

[104214]in Shwachman's syndrome. Lancet II: 800–801, 1982.

[104215]18227.

[104216]Saint-Martin, J.; Fournet, J. P.; Charlas, J.; Schaison, G.; Nodot,

[104217]A.; Meyer, B.; Vialatte, J.: Insuffisance pancreatique externe avec

[104218]granulopenie chronique. Arch. Franc. Pediat. 26: 861–871, 1969.

[104219]18228.

[104220]Saunders, E. F.; Gall, G.; Freedman, M. H.: Granulopoiesis in

[104221]Shwachman's syndrome (pancreatic insufficiency and bone marrow dysfunction). Pediatrics 64:

[104222]515–519, 1979.

[104223]18229.

[104224]Cazeneuve, C.; Sarkisian, T.; Pecheux, C.; Dervichian, M.; Nedelec,

- [104225]B.; Reinert, P.; Ayvazyan, A.; Kouyoumdjian, J.-C.;
Ajrapietian, H.;
- [104226]Delpech, M.; Goossens, M.; Dode, C.; Grateau, G.; Amse-
lem, S.: MEFV-gene
- [104227]analysis in Armenian patients with familial Mediterranean
fever: diagnostic
- [104228]value and unfavorable renal prognosis of the M694V ho-
mozygous genotype--genetic
- [104229]and therapeutic implications. Am. J. Hum. Genet. 65:
88-97, 1999.
- [104230]18230.
- [104231]Centola, M.; Wood, G.; Frucht, D. M.; Galon, J.; Aringer, M.;
- [104232]Farrell, C.; Kingma, D. W.; Horwitz, M. E.; Mansfield, E.;
- Holland,
- [104233]S. M.; O'Shea, J. J.; Rosenberg, H. F.; Malech, H. L.; Kast-
ner, D.
- [104234]L.: The gene for familial Mediterranean fever, MEFV, is ex-
pressed
- [104235]in early leukocyte development and is regulated in re-
sponse to inflammatory
- [104236]mediators. Blood 95: 3223-3231, 2000.
- [104237]18231.

[104238]Carpenter, S.; Karpati, G.; Andermann, F.; Jakob, J. C.; Andermann,

[104239]E.: The ultrastructural characteristics of the abnormal cytosomes

[104240]in Batten-Kuf's disease. Brain 100: 137-156, 1977.

[104241]18232.

[104242]Klockars, T.; Savukoski, M.; Isosomppi, J.; Laan, M.; Jarvela,

[104243]I.; Petrukhin, K.; Palotie, A.; Peltonen, L.: Efficient construction

[104244]of a physical map by fiber-fish of the CLN5 region: refined assignment

[104245]and long-range contig covering the critical region on 13q22. Genomics 35:

[104246]71-78, 1996.

[104247]18233.

[104248]Santavuori, P.; Rapola, J.; Nuutila, A.; Raininko, R.; Lappi, M.;

[104249]Launes, J.; Herva, R.; Sainio, K.: The spectrum of Jansky-Bielschowsky

[104250]disease. Neuropediatrics 22: 92-96, 1991.

[104251]18234.

[104252]Santavuori, P.; Rapola, J.; Sainio, K.; Raitta, C.: A variant
[104253]of Jansky-Bielschowsky disease. Neuropediatrics 13:
135–141, 1982.

[104254]18235.

[104255]Savukoski, M.; Kestila, M.; Williams, R.; Jarvela, I.; Sharp, J.;
[104256]Harris, J.; Santavuori, P.; Gardiner, M.; Peltonen, L.: De-
fined chromosomal

[104257]assignment of CLN5 demonstrates that at least four loci
are involved

[104258]in the pathogenesis of human ceroid lipofuscinoses. Am.
J. Hum. Genet. 55:

[104259]695–701, 1994.

[104260]18236.

[104261]Savukoski, M.; Klockars, T.; Holmberg, V.; Santavuori, P.;
Lander,

[104262]E. S.; Peltonen, L.: CLN5, a novel gene encoding a putative
transmembrane

[104263]protein mutated in Finnish variant late infantile neuronal
ceroid

[104264]lipofuscinosis. Nature Genet. 19: 286–288, 1998.

[104265]18237.

[104266]Varilo, T.; Savukoski, M.; Norio, R.; Santavuori, P.; Peltonen,

[104267]L.; Jarvela, I.: The age of human mutation: genealogical and linkage

[104268]disequilibrium analysis of the CLN5 mutation in the Finnish population. Am.

[104269]J. Hum. Genet. 58: 506–512, 1996.

[104270]18238.

[104271]Williams, R.; Santavuori, P.; Peltonen, L.; Gardiner, R. M.; Jarvela,

[104272]I.: A variant form of late infantile neuronal ceroid lipofuscinosis

[104273](CLN5) is not an allelic form of Batten (Spielmeyer–Vogt–Sjogren,

[104274]CLN3) disease: exclusion of linkage to the CLN3 region of chromosome

[104275]16. Genomics 20: 289–290, 1994.

[104276]18239.

[104277]Busard, B. L. S. M.; Renier, W. O.; Gabreels, F. J. M.; Jaspar,

[104278]H. H. J.; van Haelst, U. J. G.; Slooff, J. L.: Lafora's disease:

[104279]comparison of inclusion bodies in skin and in brain. Arch.
Neurol. 43:

[104280]296–299, 1986.

[104281]18240.

[104282]Busard, H. L. S. M.; Gabreels–Festen, A. A. W. M.; Renier,
W. O.;

[104283]Gabreels, F. J. M.; Stadhouders, A. M.: Axilla skin biopsy: a
reliable

[104284]test for the diagnosis of Lafora's disease. Ann. Neurol. 21:
599–601,

[104285]1987.

[104286]18241.

[104287]Fluharty, A. L.; Porter, M. T.; Hirsh, G. A.; Pevida, E.; Ki-
hara,

[104288]H.: Metachromasia in fibroblasts from a patient with
Lafora's disease.

[104289](Letter) Lancet II: 109–110, 1970.

[104290]18242.

[104291]Ganesh, S.; Agarwala, K. L.; Ueda, K.; Akagi, T.; Shoda, K.;
Usui,

[104292]T.; Hashikawa, T.; Osada, H.; Delgado–Escueta, A. V.; Ya–

makawa, K.

[104293] Laforin, defective in the progressive myoclonus epilepsy of Lafora

[104294] type, is a dual-specificity phosphatase associated with polyribosomes. Hum.

[104295] Molec. Genet. 9: 2251–2261, 2000.

[104296] 18243.

[104297] Gomez–Garre, P.; Sanz, Y.; Rodriguez de Cordoba, S.; Ser-ratosa,

[104298] J. M.: Mutational spectrum of the EPM2A gene in progressive myoclonus

[104299] epilepsy of Lafora: high degree of allelic heterogeneity and prevalence

[104300] of deletions. Europ. J. Hum. Genet. 8: 946–954, 2000.

[104301] 18244.

[104302] Harriman, D. G. F.; Millar, J. H. D.: Progressive familial myoclonic

[104303] epilepsy in 3 families: its clinical features and pathological basis. Brain 78:

[104304] 325–349, 1955.

[104305] 18245.

- [104306]Janeway, R.; Ravens, J. R.; Pearce, L. A.; Odor, D. L.; Suzuki,
Suzuki,
[104307]K.: Progressive myoclonus epilepsy with Lafora inclusion
bodies.
[104308]I. Clinical, genetic, histopathologic and biochemical as-
pects. Arch.
[104309]Neurol. 16: 565–582, 1967.
[104310]18246.
[104311]Lehesjoki, A.–E.; Koskiniemi, M.; Pandolfo, M.; Antonelli,
A.;
[104312]Kyllerman, M.; Wahlstrom, J.; Nergardh, A.; Burmeister, M.;
Sistonen,
[104313]P.; Norio, R.; de la Chapelle, A.: Linkage studies in pro-
gressive
[104314]myoclonus epilepsy: Unverricht–Lundborg and Lafora’s
diseases. Neurology 42:
[104315]1545–1550, 1992.
[104316]18247.
[104317]Maddox, L. O.; Descartes, M.; Collins, J.; Keating, J.;
Rosenfeld,
[104318]S.; Palmer, C.; Carroll, A. J.; Kuzniecky, R.: Identification of

[104319]a recombination event narrowing the Lafora disease gene region. J.

[104320]Med. Genet. 34: 590–591, 1997.

[104321]18248.

[104322]Harding, B. N.; Leonard, J. V.; Erdohazi, M.: Ornithine carbamoyl

[104323]transferase deficiency: a neuropathological study. Europ. J. Pediat. 141:

[104324]215–220, 1984.

[104325]18249.

[104326]Amadou, C.; Ribouchon, M. T.; Mattei, M. G.; Jenkins, N. A.; Gilbert,

[104327]D. J.; Copeland, N. G.; Avoustin, P.; Pontarotti, P.: Localization

[104328]of new genes and markers to the distal part of the human major histocompatibility

[104329]complex (MHC) region and comparison with the mouse: new insights into

[104330]the evolution of mammalian genomes. Genomics 26: 9–20, 1995.

[104331]18250.

- [104332]Solinas-Toldo, S.; Lengauer, C.; Fries, R.: Comparative genome map of human and cattle. Genomics 27: 489–496, 1995.
- [104333]18251.
- [104334]Olives, B.; Martial, S.; Mattei, M.–G.; Matassi, G.; Rousset, G.; Ripoche, P.; Cartron, J.–P.; Bailly, P.: Molecular characterization of a new urea transporter in the human kidney. FEBS Lett. 386: 156–160, 1996.
- [104335]18252.
- [104336]Ranade, K.; Wu, K.–W.; Hwu, C.–M.; Ting, C.–T.; Pei, D.; Pesich, R.; Hebert, J.; Chen, Y.–D. I.; Pratt, R.; Olshen, R.; Masaki, K.; Risch, N.; Cox, D. R.; Botstein, D.: Genetic variation in the human urea transporter–2 is associated with variation in blood pressure. Hum. Molec. Genet. 10: 2157–2164, 2001.

[104345]18253.

[104346]Galko, M. J.; Tessier-Lavigne, M.: Function of an axonal chemoattractant

[104347]modulated by metalloprotease activity. Science 289: 1365–1367, 2000.

[104348]18254.

[104349]Stein, E.; Tessier-Lavigne, M.: Hierarchical organization of guidance

[104350]receptors: silencing of netrin attraction by Slit through a Robo/DCC

[104351]receptor complex. Science 291: 1928–1938, 2001.

[104352]18255.

[104353]Stein, E.; Zou, Y.; Poo, M.; Tessier-Lavigne, M.: Binding of

[104354]DCC by netrin-1 to mediate axon guidance independent of adenosine

[104355]A2B receptor activation. Science 291: 1976–1982, 2001.

[104356]18256.

[104357]Orr, H. T.; Zoghbi, H. Y.: SCA1 molecular genetics: a history

[104358]of a 13 year collaboration against glutamines. Hum. Molec. Genet. 10:

[104359]2307–2311, 2001.

[104360]18257.

[104361]Lubahn, D. B.; Moyer, J. S.; Golding, T. S.; Couse, J. F.; Ko-
rach,

[104362]K. S.; Smithies, O.: Alteration of reproductive function but
not

[104363]prenatal sexual development after insertional disruption
of the mouse

[104364]estrogen receptor gene. Proc. Nat. Acad. Sci. 90:
11162–11166, 1993.

[104365]18258.

[104366]Hanna, I. H.; Dawling, S.; Roodi, N.; Guengerich, F. P.; Parl,

[104367]F. F.: Cytochrome P450 1B1 (CYP1B1) pharmacogenetics:
association

[104368]of polymorphisms with functional differences in estrogen
hydroxylation

[104369]activity. Cancer Res. 60: 3440–3444, 2000.

[104370]18259.

[104371]Dai, G.; Levy, O.; Carrasco, N.: Cloning and characteriza-
tion

[104372]of the thyroid iodide transporter. Nature 379: 458–460,

1996.

[104373]18260.

[104374]Costanzo, A.; Merlo, P.; Pediconi, N.; Fulco, M.; Sartorelli, V.;

[104375]Cole, P. A.; Fontemaggi, G.; Fanciulli, M.; Schiltz, L.; Blandino,

[104376]G.; Balsano, C.; Levrero, M.: DNA damage-dependent acetylation of

[104377]p73 dictates the selective activation of apoptotic target genes. Molec.

[104378]Cell 9: 175–186, 2002.

[104379]18261.

[104380]Flores, E. R.; Tsai, K. Y.; Crowley, D.; Sengupta, S.; Yang, A.;

[104381]McKeon, F.; Jacks, T.: p63 and p73 are required for p53-dependent

[104382]apoptosis in response to DNA damage. Nature 416: 560–564, 2002.

[104383]18262.

[104384]Jost, C. A.; Marin, M. C.; Kaelin, W. G., Jr.: p73 is a human

[104385]p53-related protein that can induce apoptosis. Nature

389: 191–194,

[104386]1997.

[104387]18263.

[104388]Kaghad, M.; Bonnet, H.; Yang, A.; Creancier, L.; Biscan, J.-C.;

[104389]Valent, A.; Minty, A.; Chalon, P.; Lelias, J.-M.; Dumont, X.; Ferrara,

[104390]P.; McKeon, F.; Caput, D.: Monoallelically expressed gene related

[104391]to p53 at 1p36, a region frequently deleted in neuroblastoma and other

[104392]human cancers. Cell 90: 809–819, 1997.

[104393]18264.

[104394]Marin, M. C.; Jost, C. A.; Brooks, L. A.; Irwin, M. S.; O'Nions,

[104395]J.; Tidy, J. A.; James, N.; McGregor, J. M.; Harwood, C. A.; Yulug,

[104396]I. G.; Vousden, K. H.; Allday, M. J.; Gusterson, B.; Ikawa, S.; Hinds,

[104397]P. W.; Crook, T.; Kaelin, W. G., Jr. A common polymorphism acts

[104398]as an intragenic modifier of mutant p53 behaviour. Nature

Genet. 25:

[104399]47–54, 2000.

[104400]18265.

[104401]Morrison, R. S.; Kinoshita, Y.: p73--guilt by association?

Science 289:

[104402]257–258, 2000.

[104403]18266.

[104404]Pozniak, C. D.; Radinovic, S.; Yang, A.; McKeon, F.; Kaplan,
D.

[104405]R.; Miller, F. D.: An anti-apoptotic role for the p53 family
member,

[104406]p73, during developmental neuron death. Science 289:
304–306, 2000.

[104407]18267.

[104408]Yang, A.; Walker, N.; Bronson, R.; Kaghad, M.; Ooster-
wegel, M.;

[104409]Bonnin, J.; Vagner, C.; Bonnet, H.; Dikkes, P.; Sharpe, A.;
McKeon,

[104410]F.; Caput, D.: p73-deficient mice have neurological,
pheromonal and

[104411]inflammatory defects but lack spontaneous tumours. Na-

ture 404: 99–103,

[104412]2000.

[104413]18268.

[104414]Moore, J. T.; McKee, D. D.; Slentz–Kesler, K.; Moore, L. B.;
Jones,

[104415]S. A.; Horne, E. L.; Su, J.–L.; Kliewer, S. A.; Lehmann, J. M.;
Willson,

[104416]T. M.: Cloning and characterization of human estrogen re-
ceptor beta

[104417]isoforms. *Biochem. Biophys. Res. Commun.* 247: 75–78,
1998.

[104418]18269.

[104419]Mosselman, S.; Polman, J.; Dijkema, R.: ER–beta: identifi-
cation

[104420]and characterization of a novel human estrogen receptor.
FEBS Lett. 392:

[104421]49–53, 1996.

[104422]18270.

[104423]Ogawa, S.; Emi, M.; Shiraki, M.; Hosoi, T.; Ouchi, Y.; Inoue,

[104424]S.: Association of estrogen receptor beta (ESR2) gene
polymorphism

[104425]with blood pressure. J. Hum. Genet. 45: 327–330, 2000.

[104426]18271.

[104427]Ogawa, S.; Inoue, S.; Watanabe, T.; Orimo, A.; Hosoi, T.;
Ouchi,

[104428]Y.; Muramatsu, M.: Molecular cloning and characterization
of human

[104429]estrogen receptor beta-cx: a potential inhibitor of estrogen
action

[104430]in human. Nucleic Acids Res. 26: 3505–3512, 1998.

[104431]18272.

[104432]Osterlund, M. K.; Gustafsson, J. A.; Keller, E.; Hurd, Y. L.:

[104433]Estrogen receptor beta (ER-beta) messenger ribonucleic
acid (mRNA)

[104434]expression within the human forebrain: distinct distribution
pattern

[104435]to ER-alpha mRNA. J. Clin. Endocr. Metab. 85: 3840–3846,
2000.

[104436]18273.

[104437]Pasquali, D.; Rossi, V.; Esposito, D.; Abbondanza, C.; Puca,
G.

[104438]A.; Bellastella, A.; Sinisi, A. A.: Loss of estrogen receptor

beta

[104439]expression in malignant human prostate cells in primary cultures and

[104440]in prostate cancer tissues. J. Clin. Endocr. Metab. 86: 2051–2055,

[104441]2001.

[104442]18274.

[104443]Petersen, D. N.; Tkalcevic, G. T.; Koza–Taylor, P. H.; Turi, T.

[104444]G.; Brown, T. A.: Identification of estrogen receptor beta–2, a functional

[104445]variant of estrogen receptor beta expressed in normal rat tissues. Endocrinology 139:

[104446]1082–1092, 1998.

[104447]18275.

[104448]Rissman, E. F.; Heck, A. L.; Leonard, J. E.; Shupnik, M. A.; Gustafsson,

[104449]J.–A.: Disruption of estrogen receptor beta gene impairs spatial

[104450]learning in female mice. Proc. Nat. Acad. Sci. 99: 3996–4001, 2002.

[104451]18276.

[104452]Rosenkranz, K.; Hinney, A.; Ziegler, A.; Hermann, H.;
Fichter,
[104453]M.; Mayer, H.; Siegfried, W.; Young, J. K.; Remschmidt, H.;
Hebebrand,
[104454]J.: Systematic mutation screening of the estrogen receptor
beta gene
[104455]in probands of different weight extremes: identification of
several
[104456]genetic variants. J. Clin. Endocr. Metab. 83: 4524-4527,
1998.
[104457]18277.
[104458]Speirs, V.; Adams, I. P.; Walton, D. S.; Atkin, S. L.: Identifi-
cation
[104459]of wild-type and exon 5 deletion variants of estrogen re-
ceptor beta
[104460]in normal human mammary gland. J. Clin. Endocr. Metab.
85: 1601-1605,
[104461]2000.
[104462]18278.
[104463]Tsukamoto, K.; Inoue, S.; Hosoi, T.; Orimo, H.; Emi, M.:
Isolation
[104464]and radiation hybrid mapping of dinucleotide repeat poly-

morphism at

[104465]the human estrogen receptor beta locus. J. Hum. Genet.

43: 73–74,

[104466]1998.

[104467]18279.

[104468]Weihua, Z.; Saji, S.; Makinen, S.; Cheng, G.; Jensen, E. V.;

Warner,

[104469]M.; Gustafsson, J.–A.: Estrogen receptor (ER)–beta, a mod-

ulator of

[104470]ER–alpha in the uterus. Proc. Nat. Acad. Sci. 97:

5936–5941, 2000.

[104471]18280.

[104472]Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.;

Landen,

[104473]M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Poly-

morphisms

[104474]of the androgen receptor gene and the estrogen receptor

gene are associated

[104475]with androgen levels in women. J. Clin. Endocr. Metab. 86:

2562–2568,

[104476]2001.

[104477]18281.

[104478]Zhu, Y.; Bian, Z.; Lu, P.; Karas, R. H.; Bao, L.; Cox, D.; Hodgins,

[104479]J.; Shaul, P. W.; Thoren, P.; Smithies, O.; Gustafsson, J.-A.; Mendelsohn,

[104480]M. E.: Abnormal vascular function and hypertension in mice deficient

[104481]in estrogen receptor beta. *Science* 295: 505–508, 2002.

[104482]18282.

[104483]DesGroseillers, L.; Lemieux, N.: Localization of a human double-stranded

[104484]RNA-binding protein gene (STAU) to band 20q13.1 by fluorescence in

[104485]situ hybridization. *Genomics* 36: 527–529, 1996.

[104486]18283.

[104487]Marion, R. M.; Fortes, P.; Beloso, A.; Dotti, C.; Ortin, J.: A

[104488]human sequence homologue of stau68 is an RNA-binding protein that

[104489]is associated with polysomes and localizes to the rough endoplasmic

[104490]reticulum. *Molec. Cell. Biol.* 19: 2212–2219, 1999.

[104491]18284.

[104492]Wickham, L.; Duchaine, T.; Luo, M.; Nabi, I. R.; DesGro-
seillers,

[104493]L.: Mammalian stau-1 is a double-stranded-RNA- and
tubulin-binding

[104494]protein which localizes to the rough endoplasmic reticu-
lum. Molec.

[104495]Cell. Biol. 19: 2220-2230, 1999.

[104496]18285.

[104497]Lee, J. E.; Hollenberg, S. M.; Snider, L.; Turner, D. L.; Lip-
nick,

[104498]N.; Weintraub, H.: Conversion of Xenopus ectoderm into
neurons by

[104499]NeuroD, a basic helix-loop-helix protein. Science 268:
836-844,

[104500]1995.

[104501]18286.

[104502]Plasilova, M.; Stoilov, I.; Sarfarazi, M.; Kadasi, L.; Ferakova,

[104503]E.; Ferak, V.: Identification of a single ancestral CYP1B1
mutation

[104504]in Slovak Gypsies (Roms) affected with primary congenital
glaucoma. J.

[104505]Med. Genet. 36: 290–294, 1999.

[104506]18287.

[104507]Schwartzman, M. L.; Balazy, M.; Masferrer, J.; Abraham, N.
G.;

[104508]McGiff, J. C.; Murphy, R. C.: 12(R)–hydroxyicosatetraenoic
acid:

[104509]a cytochrome P450–dependent arachidonate metabolite
that inhibits

[104510]Na⁺,K⁺–ATPase in the cornea. Proc. Nat. Acad. Sci. 84:
8125–8129,

[104511]1987.

[104512]18288.

[104513]Stoilov, I.; Akarsu, A. N.; Alozie, I.; Child, A.; Barsoum–
Homsy,

[104514]M.; Turacli, M. E.; Or, M.; Lewis, R. A.; Ozdemir, N.; Brice,
G.;

[104515]Aktan, S. G.; Chevrette, L.; Coca–Prados, M.; Sarfarazi, M.:
Sequence

[104516]analysis and homology modeling suggest that primary
congenital glaucoma

[104517]on 2p21 results from mutations disrupting either the
hinge region

- [104518] or the conserved core structures of cytochrome P4501B1.
Am. J. Hum.
- [104519] Genet. 62: 573–584, 1998.
- [104520] 18289.
- [104521] Stoilov, I.; Akarsu, A. N.; Sarfarazi, M.: Identification of
three
[104522] different truncating mutations in cytochrome P4501B1
(CYP1B1) as the
[104523] principal cause of primary congenital glaucoma
(buphthalmos) in families
[104524] linked to the GLC3A locus on chromosome 2p21. Hum.
Molec. Genet. 6:
[104525] 641–647, 1997.
- [104526] 18290.
- [104527] Sutter, T. R.; Tang, Y. M.; Hayes, C. L.; Wo, Y.-Y. P.; Jabs,
E.
[104528] W.; Li, X.; Yin, H.; Cody, C. W.; Greenlee, W. F.: Complete
cDNA
[104529] sequence of a human dioxin-inducible mRNA identifies a
new gene subfamily
[104530] of cytochrome P450 that maps to chromosome 2. J. Biol.
Chem. 269:

[104531]13092–13099, 1994.

[104532]18291.

[104533]Tang, Y. M.; Wo, Y.-Y. P.; Stewart, J.; Hawkins, A. L.; Griffin,
fin,

[104534]C. A.; Sutter, T. R.; Greenlee, W. F.: Isolation and characterization
terization

[104535]of the human cytochrome P450 CYP1B1 gene. J. Biol.
Chem. 271: 28324–28330,

[104536]1996.

[104537]18292.

[104538]Vincent, A.; Billingsley, G.; Priston, M.; Williams–Lyn, D.;
Sutherland,

[104539]J.; Glaser, T.; Oliver, E.; Walter, M. A.; Heathcote, G.; Levin,
A.;

[104540]Heon, E.: Phenotypic heterogeneity of CYP1B1: mutations
in a patient

[104541]with Peters' anomaly. J. Med. Genet. 38: 324–326, 2001.

[104542]18293.

[104543]Chen, H. T.; Bhandoola, A.; Difilippantonio, M. J.; Zhu, J.;
Brown,

[104544]M. J.; Tai, X.; Rogakou, E. P.; Brotz, T. M.; Bonner, W. M.;

Ried,

[104545]T.; Nussenzweig, A.: Response to RAG-mediated V(D)J cleavage by NBS1

[104546]and gamma-H2AX. Science 290: 1962–1964, 2000.

[104547]18294.

[104548]Petersen, S.; Casellas, R.; Reina-San-Martin, B.; Chen, H. T.;

[104549]Difilippantonio, M. J.; Wilson, P. C.; Hanitsch, L.; Celeste, A.;

[104550]Muramatsu, M.; Pilch, D. R.; Redon, C.; Ried, T.; Bonner, W. M.; Honjo,

[104551]T.; Nussenzweig, M. C.; Nussenzweig, A.: AID is required to initiate

[104552]Nbs1/gamma-H2AX focus formation and mutations at sites of class switching. Nature 414:

[104553]660–665, 2001.

[104554]18295.

[104555]Lan, M. S.; Lu, J.; Goto, Y.; Notkins, A. L.: Molecular cloning

[104556]and identification of a receptor-type protein tyrosine phosphatase,

[104557]IA-2, from human insulinoma. DNA and Cell Biol. 13:

505-514, 1994.

[104558]18296.

[104559]Leiter, E. H.; Tsumura, H.; Serreze, D. V.; Chapman, H. D.;
Rabin,

[104560]D. U.; Lan, M. S.; Notkins, A. L.: Mapping to chromosomes
1 and 12

[104561]of mouse homologs of human protein tyrosine phos-
phatase, receptor-type,

[104562]related genes encoding pancreatic beta cell autoantigens.
Mammalian

[104563]Genome 8: 949-950, 1997.

[104564]18297.

[104565]Rabin, D. U.; Pleasic, S. M.; Shapiro, J. A.; Yoo-Warren, H.;
Oles,

[104566]J.; Hicks, J. M.; Goldstein, D. E.; Rae, P. M.: Islet cell anti-
gen

[104567]512 is a diabetes-specific islet autoantigen related to pro-
tein tyrosine

[104568]phosphatases. J. Immun. 152: 3183-3188, 1994.

[104569]18298.

[104570]Xie, J.; Zhang, B.; Lan, M. S.; Notkins, A. L.: Genomic

structure

[104571]and promoter sequence of the insulin-dependent diabetes mellitus autoantigen,

[104572]IA-2 (PTPRN). Genomics 54: 338-343, 1998.

[104573]18299.

[104574]Demetriou, M.; Granovsky, M.; Quaggin, S.; Dennis, J. W.:

Negative

[104575]regulation of T-cell activation and autoimmunity by Mgat5 N-glycosylation. Nature 409:

[104576]733-739, 2001.

[104577]18300.

[104578]van den Maagdenberg, A. M. J. M.; Weghuis, D. O.; Rijss,

J.; van

[104579]de Wetering, R. A. C.; Wieringa, B.; van Kessel, A. G.; Hendriks,

[104580]W. J. A. J.: Assignment of the human gene for receptor-type protein

[104581]tyrosine phosphatase IA-2 (PTPRN) to chromosome region 2q35-q36.1

[104582]and identification of an intragenic genetic marker. Cytogenet. Cell

[104583]Genet. 73: 145-148, 1996.

[104584]18301.

[104585]Granovsky, M.; Fata, J.; Pawling, J.; Muller, W. J.; Khokha, R.;

[104586]Dennis, J. W.: Suppression of tumor growth and metastasis in Mgat5-deficient

[104587]mice. Nature Med. 6: 306-12, 2000.

[104588]18302.

[104589]Saito, H.; Nishikawa, A.; Gu, J.; Ihara, Y.; Soejima, H.; Wada,

[104590]Y.; Sekiya, C.; Niikawa, N.; Taniguchi, N.: cDNA cloning and chromosomal

[104591]mapping of human N-acetylglucosaminyltransferase V+. Biochem. Biophys.

[104592]Res. Commun. 198: 318-327, 1994.

[104593]18303.

[104594]Morasso, M. I.; Yonescu, R.; Griffin, C. A.; Sargent, T. D.: Localization

[104595]of human DLX8 to chromosome 17q21.3-q22 by fluorescence in situ hybridization. Mammalian

[104596]Genome 8: 302-303, 1997.

[104597]18304.

[104598]Quinn, L. M.; Johnson, B. V.; Nicholl, J.; Sutherland, G. R.;
Kalionis,
[104599]B.: Isolation and identification of homeobox genes from
the human
[104600]placenta including a novel member of the Distal-less fam-
ily, DLX4. Gene 187:
[104601]55-61, 1997.
[104602]18305.
[104603]Boddy, M. N.; Howe, K.; Etkin, L. D.; Solomon, E.;
Freemont, P.
[104604]S.: PIC 1, a novel ubiquitin-like protein which interacts
with the
[104605]PML component of a multiprotein complex that is dis-
rupted in acute
[104606]promyelocytic leukaemia. Oncogene 13: 971-982, 1996.
[104607]18306.
[104608]Desterro, J. M. P.; Rodriguez, M. S.; Hay, R. T.: SUMO-1
modification
[104609]of I-kappa-B-alpha inhibits NF-kappa-B activation. Molec.
Cell 2:
[104610]233-239, 1998.

[104611]18307.

[104612]Lapenta, V.; Chiurazzi, P.; van der Spek, P.; Pizzuti, A.;
Hanaoka,

[104613]F.; Brahe, C.: SMT3A, a human homologue of the *S. cere-*
visiae SMT3

[104614]gene, maps to chromosome 21qter and defines a novel
gene family. *Genomics* 40:

[104615]362–366, 1997.

[104616]18308.

[104617]Mao, Y.; Sun, M.; Desai, S. D.; Liu, L. F.: SUMO–1 conjuga-
tion

[104618]to topoisomerase I: a possible repair response to topoiso-
merase-mediated

[104619]DNA damage. *Proc. Nat. Acad. Sci.* 97: 4046–4051, 2000.

[104620]18309.

[104621]Okura, T.; Gong, L.; Kamitani, T.; Wada, T.; Okura, I.; Wei,
C.-F.;

[104622]Chang, H.-M.; Yeh, E. T. H.: Protection against Fas/
APO–1– and tumor

[104623]necrosis factor-mediated cell death by a novel protein,
sentrin. *J.*

[104624]Immun. 157: 4277–4281, 1996.

[104625]18310.

[104626]Shen, Z.; Pardington–Purtymun, P. E.; Comeaux, J. C.;
Moyzis, R.

[104627]K.; Chen, D. J.: UBL1, a human ubiquitin–like protein asso-
ciating

[104628]with human RAD51/RAD52 proteins. Genomics 36:
271–279, 1996.

[104629]18311.

[104630]Bengtsson, E.; Neame, P. J.; Heinegard, D.; Sommarin, Y.:
The

[104631]primary structure of a basic leucine–rich repeat protein,
PRELP, found

[104632]in connective tissues. J. Biol. Chem. 270: 25639–25644,
1995.

[104633]18312.

[104634]Grover, J.; Chen, X.–N.; Korenberg, J. R.; Recklies, A. D.;
Roughley,

[104635]P. J.: The gene organization, chromosome location, and
expression

[104636]of a 55–kDa matrix protein (PRELP) of human articular
cartilage. Genomics 38:

[104637]109–117, 1996.

[104638]18313.

[104639]Bigg, H. F.; Shi, Y. E.; Liu, Y. E.; Steffensen, B.; Overall, C.

[104640]M.: Specific, high affinity binding of tissue inhibitor of metalloproteinases–4

[104641](TIMP–4) to the COOH–terminal hemopexin–like domain of human gelatinase

[104642]A. J. Biol. Chem. 272: 15496–15500, 1997.

[104643]18314.

[104644]Greene, J.; Wang, M.; Liu, Y. E.; Raymond, L. A.; Rosen, C.; Shi,

[104645]Y. E.: Molecular cloning and characterization of human tissue inhibitor

[104646]of metalloproteinase 4. J. Biol. Chem. 271: 30375–30380, 1996.

[104647]18315.

[104648]Leco, K. J.; Apte, S. S.; Taniguchi, G. T.; Hawkes, S. P.; Khokha,

[104649]R.; Schultz, G. A.; Edwards, D. R.: Murine tissue inhibitor of metalloproteinases–4

[104650](Timp–4): cDNA isolation and expression in adult mouse

tissues. FEBS

[104651]Lett. 401: 213–217, 1997.

[104652]18316.

[104653]Olson, T. M; Hirohata, S.; Ye, J.; Leco, K.; Seldin, M. F.;
Apte,

[104654]S. S.: Cloning of the human tissue inhibitor of metallopro-
teinase-4

[104655]gene (TIMP4) and localization of the TIMP4 and Timp4
gene to human

[104656]chromosome 3p25 and mouse chromosome 6, respec-
tively. Genomics 51:

[104657]148–151, 1998.

[104658]18317.

[104659]Wang, M.; Liu, Y. E.; Greene, J.; Sheng, S.; Fuchs, A.;
Rosen,

[104660]E. M.; Shi, Y. E.: Inhibition of tumor growth and metastasis
of human

[104661]breast cancer cells transfected with tissue inhibitor of
metalloproteinase

[104662]4. Oncogene 14: 2767–2774, 1997.

[104663]18318.

- [104664]Hsu, L. C.; Chang, W.-C.: Sequencing and expression of the human
- [104665]ALDH8 encoding a new member of the aldehyde dehydrogenase family. Gene 174:
- [104666]319–322, 1996.
- [104667]18319.
- [104668]Hsu, L. C.; Chang, W.-C.; Lin, S. W.; Yoshida, A.: Cloning and
- [104669]characterization of genes encoding four additional human aldehyde
- [104670]dehydrogenase isozymes. Adv. Exp. Med. Biol. 372: 159–168, 1995.
- [104671]18320.
- [104672]Fujiwara, H.; Tatsumi, K.-I.; Miki, K.; Harada, T.; Okada, S.;
- [104673]Nose, O.; Kodama, S.; Amino, N.: Recurrent T354P mutation of the
- [104674]Na⁺/I⁻ symporter in patients with iodide transport defect. J. Clin.
- [104675]Endocr. Metab. 83: 2940–2943, 1998.
- [104676]18321.
- [104677]Kosugi, S.; Bhayana, S.; Dean, H. J.: A novel mutation in

the

[104678]sodium/iodide symporter gene in the largest family with
iodide transport

[104679]defect. J. Clin. Endocr. Metab. 84: 3248–3253, 1999.

[104680]18322.

[104681]Kosugi, S.; Inoue, S.; Matsuda, A.; Jhiang, S. M.: Novel,
missense

[104682]and loss-of-function mutations in the sodium/iodide
symporter gene

[104683]causing iodide transport defect in three Japanese patients.
J. Clin.

[104684]Endocr. Metab. 83: 3373–3376, 1998.

[104685]18323.

[104686]Kosugi, S.; Sato, Y.; Matsuda, A.; Ohyama, Y.; Fujieda, K.;
Inomata,

[104687]H.; Kameya, T.; Isozaki, O.; Jhiang, S. M.: High prevalence
of T354P

[104688]sodium/iodide symporter gene mutation in Japanese pa-
tients with iodide

[104689]transport defect who have heterogeneous clinical pictures.
J. Clin.

[104690]Endocr. Metab. 83: 4123–4129, 1998.

[104691]18324.

[104692]Levy, O.; Ginter, C. S.; De la Vieja, A.; Levy, D.; Carrasco,

[104693]N.: Identification of a structural requirement for thyroid

Na(+)/I(-)

[104694]symporter (NIS) function from analysis of a mutation that

causes human

[104695]congenital hypothyroidism. FEBS Lett. 429: 36-40, 1998.

[104696]18325.

[104697]Matsuda, A.; Kosugi, S.: A homozygous missense muta-

tion of the

[104698]sodium/iodide symporter gene causing iodide transport

defect. J.

[104699]Clin. Endocr. Metab. 82: 3966-3971, 1997.

[104700]18326.

[104701]Ohmori, M.; Endo, T.; Harii, N.; Onaya, T.: A novel thyroid

transcription

[104702]factor is essential for thyrotropin-induced up-regulation

of Na+ /I-

[104703]symporter gene expression. Molec. Endocr. 12: 727-736,

1998.

[104704]18327.

- [104705]Pohlenz, J.; Medeiros-Neto, G.; Gross, J. L.; Silveiro, S. P.;
- [104706]Knobel, M.; Refetoff, S.: Hypothyroidism in a Brazilian kindred due
- [104707]to iodide trapping defect caused by a homozygous mutation in the sodium/iodide
- [104708]symporter gene. *Biochem. Biophys. Res. Commun.* 240: 488–491, 1997.
- [104709]18328.
- [104710]Pohlenz, J.; Rosenthal, I. M.; Weiss, R. E.; Jhiang, S. M.; Barrant,
- [104711]C.; Refetoff, S.: Congenital hypothyroidism due to mutations in the
- [104712]sodium/iodide symporter: identification of a nonsense mutation producing
- [104713]a downstream cryptic 3-prime splice site. *J. Clin. Invest.* 101: 1028–1035, 1998.
- [104714]18329.
- [104715]18329.
- [104716]Smanik, P. A.; Liu, Q.; Furminger, T. L.; Ryu, K.; Xing, S.; Mazzaferri,
- [104717]E. L.; Jhiang, S. M.: Cloning of the human sodium iodide symporter. *Biochem.*

[104718]Biophys. Res. Commun. 226: 339–345, 1996.

[104719]18330.

[104720]Smanik, P. A.; Ryu, K.-Y.; Theil, K. S.; Mazzaferri, E. L.; Jhi-
ang,

[104721]S. M.: Expression, exon–intron organization, and chromo-
some mapping

[104722]of the human sodium iodide symporter. Endocrinology
138: 3555–3558,

[104723]1997.

[104724]18331.

[104725]Spitzweg, C.; Joba, W.; Schriever, K.; Goellner, J. R.; Morris,

[104726]J. C.; Heufelder, A. E.: Analysis of human sodium iodide
symporter

[104727]immunoreactivity in human exocrine glands. J. Clin. En-
docr. Metab. 84:

[104728]4178–4184, 1999.

[104729]18332.

[104730]Venkataraman, G. M.; Yatin, M.; Marcinek, R.; Ain, K. B.:
Restoration

[104731]of iodide uptake in dedifferentiated thyroid carcinoma: re-
lationship

- [104732]to human Na(+)/I(-) symporter gene methylation status. J. Clin. Endocr.
- [104733]Metab. 84: 2449–2457, 1999.
- [104734]18333.
- [104735]Lyon, M. F.; Peters, J.; Glenister, P. H.; Ball, S.; Wright, E.
- [104736]: The scurfy mouse mutant has previously unrecognized hematological abnormalities and resembles Wiskott–Aldrich syndrome. Proc. Nat.
- [104738]Acad. Sci. 87: 2433–2437, 1990.
- [104739]18334.
- [104740]Rawson, R. B.; Zelenski, N. G.; Nijhawan, D.; Ye, J.; Sakai, J.;
- [104741]Hasan, M. T.; Chang, T. Y.; Brown, M. S.; Goldstein, J. L.: Complementation
- [104742]cloning of S2P, a gene encoding a putative metalloprotease required
- [104743]for intramembrane cleavage of SREBPs. Molec. Cell 1: 47–57, 1997.
- [104744]18335.
- [104745]Ye, J.; Rawson, R. B.; Komuro, R.; Chen, X.; Dave, U. P.;

Prywes,

[104746]R.; Brown, M. S.; Goldstein, J. L.: ER stress induces cleavage of

[104747]membrane-bound ATF6 by the same proteases that process SREBPs. Molec.

[104748]Cell 6: 1355–1364, 2000.

[104749]18336.

[104750]Zelenski, N. G.; Rawson, R. B.; Brown, M. S.; Goldstein, J. L.

[104751]: Membrane topology of S2P, a protein required for intramembraneous

[104752]cleavage of sterol regulatory element-binding proteins. J. Biol.

[104753]Chem. 274: 21973–21980, 1999.

[104754]18337.

[104755]Baytel, D.; Shalom, S.; Madgar, I.; Weissenberg, R.; Don, J.:

[104756]The human Pim-2 proto-oncogene and its testicular expression. Biochim.

[104757]Biophys. Acta 1442: 274–285, 1998.

[104758]18338.

[104759]Cocchia, M.; Huber, R.; Pantano, S.; Chen, E. Y.; Ma, P.;

Forabosco,

[104760]A.; Ko, M. S. H.; Schlessinger, D.: PLAC1, an Xq26 gene
with placenta-specific

[104761]expression. Genomics 68: 305–312, 2000.

[104762]18339.

[104763]Kim, V. N.; Kataoka, N.; Dreyfuss, G.: Role of the non-
sense-mediated

[104764]decay factor hUpf3 in the splicing-dependent exon-exon
junction complex. Science 293:

[104765]1832–1836, 2001.

[104766]18340.

[104767]Lykke-Andersen, J.; Shu, M.-D.; Steitz, J. A.: Human Upf
proteins

[104768]target an mRNA for nonsense-mediated decay when
bound downstream of

[104769]a termination codon. Cell 103: 1121–1131, 2000.

[104770]18341.

[104771]Scott, A. F.: Personal Communication. Baltimore, Md.
2/9/2001.

[104772]18342.

[104773]Serin, G.; Gersappe, A.; Black, J. D.; Aronoff, R.; Maquat, L.

- [104774]E.: Identification and characterization of human orthologues to *Saccharomyces*
- [104775]*cerevisiae* Upf2 protein and Upf3 protein (*Caenorhabditis elegans* SMG-4). *Molec.*
- [104776]*Cell. Biol.* 21: 209–223, 2001.
- [104777]18343.
- [104778]Kelley, W. N.; Rosenbloom, F. M.; Henderson, J. F.; Seeg-
- millar,
- [104779]J. E.: A specific enzyme defect in gout associated with
- overproduction
- [104780]of uric acid. *Proc. Nat. Acad. Sci.* 57: 1735–1739, 1967.
- [104781]18344.
- [104782]McDonald, J. A.; Kelley, W. N.: Lesch–Nyhan syndrome: al-
- tered
- [104783]kinetic properties of mutant enzyme. *Science* 171:
- 689–691, 1971.
- [104784]18345.
- [104785]de Leeuw, B.; Balemans, M.; Geurts van Kessel, A.: A novel
- Kruppel-associated
- [104786]box containing the SSX gene (SSX3) on the human X chro-
- mosome is not

[104787] implicated in t(X;18)-positive synovial sarcomas. Cyto-genet. Cell

[104788] Genet. 73: 179-183, 1996.

[104789] 18346.

[104790] Gure, A. O.; Tureci, O.; Sahin, U.; Tsang, S.; Scanlan, M. J.;

[104791] Jager, E.; Knuth, A.; Pfreundschuh, M.; Old, L. J.; Chen, Y.-T.:

[104792] SSX: a multigene family with several members transcribed in normal

[104793] testis and human cancer. Int. J. Cancer 72: 965-971, 1997.

[104794] 18347.

[104795] Sossey-Alaoui, K.; Lyon, J. A.; Jones, L.; Abidi, F. E.; Hartung,

[104796] A. J.; Hane, B.; Schwartz, C. E.; Stevenson, R. E.; Srivastava, A.

[104797] K.: Molecular cloning and characterization of TRPC5 (HTRP5), the

[104798] human homologue of a mouse brain receptor-activated capacitative Ca(2+)

[104799] entry channel. Genomics 60: 330-340, 1999.

[104800]18348.

[104801]Tipnis, S. R.; Hooper, N. M.; Hyde, R.; Karran, E.; Christie, G.;

[104802]Turner, A. J.: A human homolog of angiotensin-converting enzyme:

[104803]cloning and functional expression as a captopril-insensitive carboxypeptidase. J.

[104804]Biol. Chem. 275: 33238-33243, 2000.

[104805]18349.

[104806]Johnson, R. E.; Kondratick, C. M.; Prakash, S.; Prakash, L.: hRAD30

[104807]mutations in the variant form of xeroderma pigmentosum. Science 285:

[104808]263-265, 1999.

[104809]18350.

[104810]Masutani, C.; Kusumoto, R.; Yamada, A.; Dohmae, N.; Yokoi, M.;

[104811]Yuasa, M.; Araki, M.; Iwai, S.; Takio, K.; Hanaoka, F.: The XPV (xeroderma

[104812]pigmentosum variant) gene encodes human DNA polymerase ϵ . Nature 399:

[104813]700–704, 1999.

[104814]18351.

[104815]Neitz, M.; Neitz, J.: Numbers and ratios of visual pigment genes

[104816]for normal red–green color vision. Science 267: 1013–1016, 1995.

[104817]18352.

[104818]Reyniers, E.; Van Thienen, M.–N.; Meire, F.; De Boulle, K.; Devries,

[104819]K.; Kestelijn, P.; Willems, P. J.: Gene conversion between red and

[104820]defective green opsin gene in blue cone monochromacy. Genomics 29:

[104821]323–328, 1995.

[104822]18353.

[104823]Ruberg, F. L.; Skene, D. J.; Hanifin, J. P.; Rollag, M. D.; English,

[104824]J.; Arendt, J.; Brainard, G. C.: Melatonin regulation in humans with

[104825]color vision deficiencies. J. Clin. Endocr. Metab. 81: 2980–2985,

[104826]1996.

[104827]18354.

[104828]Ueyama, H.; Kuwayama, S.; Imai, H.; Tanabe, S.; Oda, S.;
Nishida,

[104829]Y.; Wada, A.; Shichida, Y.; Yamade, S.: Novel missense
mutations

[104830]in red/green opsin genes in congenital color-vision defi-
ciencies. Biochem.

[104831]Biophys. Res. Commun. 294: 205-209, 2002.

[104832]18355.

[104833]Brunner, W.: Ueber den Vererbungsmodus der verschiede-
nen Typen

[104834]der angeborenen Rotgruenblindheit. Albrecht von Graefes
Arch. Ophthal. 124:

[104835]1-52, 1932.

[104836]18356.

[104837]Crone, R. A.: Spectral sensitivity in color-defective sub-
jects

[104838]and heterozygous carriers. Am. J. Ophthal. 48: 231-238,
1959.

[104839]18357.

[104840]Deeb, S. S.; Lindsey, D. T.; Hibiya, Y.; Sanocki, E.; Winder-
ickx,

[104841]J.; Teller, D. Y.; Motulsky, A. G.: Genotype-phenotype re-
lationships

[104842]in human red/green color-vision defects: molecular and
psychophysical

[104843]studies. Am. J. Hum. Genet. 51: 687-700, 1992.

[104844]18358.

[104845]Drummond-Borg, M.; Deeb, S.; Motulsky, A. G.: Molecular
basis

[104846]of abnormal red-green color vision: a family with three
types of color

[104847]vision defects. Am. J. Hum. Genet. 43: 675-683, 1988.

[104848]18359.

[104849]Emmerson, B. T.; Thompson, L.; Wallace, D. C.; Spence, M.
A.:

[104850]Absence of measurable linkage between the loci for hy-
poxanthine-guanine

[104851]phosphoribosyltransferase and deutan color blindness.
Am. J. Hum.

[104852]Genet. 26: 78-82, 1974.

[104853]18360.

[104854]Keating, J. P.; Brown, B. I.; White, N. H.; DiMauro, S.: X-linked

[104855]glycogen storage disease: a cause of hypotonia, hyperuricemia, and

[104856]growth retardation. Am. J. Dis. Child. 139: 609-613, 1985.

[104857]18361.

[104858]Krebs, E. G.; Love, D. S.; Bratvold, G. E.; Trayser, K. A.; Meyer,

[104859]W. L.; Fischer, E. H.: Purification and properties of rabbit skeletal

[104860]muscle phosphorylase B kinase. Biochemistry 3: 1022-1033, 1964.

[104861]18362.

[104862]Kumatori, A.; Faizunnessa, N. N.; Suzuki, S.; Moriuchi, T.; Kurozumi,

[104863]H.; Nakamura, M.: Nonhomologous recombination between the cytochrome

[104864]b-558 heavy chain gene (CYBB) and LINE-1 causes an X-linked chronic

[104865]granulomatous disease. Genomics 53: 123-128, 1998.

[104866]18363.

[104867]Lyon, J. B., Jr.; Porter, J.; Robertson, M.: Phosphorylase B

[104868]kinase inheritance in mice. Science 155: 1550–1551,
1967.

[104869]18364.

[104870]Malthus, R.; Clark, D. G.; Watts, C.; Sneyd, J. G. T.: Glyco-
gen-storage

[104871]disease in rats, a genetically determined deficiency of liver
phosphorylase

[104872]kinase. Biochem. J. 188: 99–106, 1980.

[104873]18365.

[104874]McAdams, A. J.; Hug, G.; Bove, K. E.: Glycogen storage
disease,

[104875]type I to X: criteria for morphologic diagnosis. Hum. Path.
5: 463–487,

[104876]1974.

[104877]18366.

[104878]Migeon, B. R.; Huijing, F.: Glycogen-storage disease asso-
ciated

[104879]with phosphorylase kinase deficiency: evidence for X inac-
tivation. Am.

[104880]J. Hum. Genet. 26: 360–368, 1974.

[104881]18367.

[104882]Nagai, T.; Matsuo, N.; Tsuchiya, Y.; Cho, H.; Hasegawa, Y.;
Igarashi,

[104883]Y.: Proximal renal tubular acidosis associated with glyco-
gen storage

[104884]disease, type 9. Acta Paediat. Scand. 77: 460–463, 1988.

[104885]18368.

[104886]Ryder–Cook, A. S.; Derry, J. M. J.; Barnard, P. J.: Localiza-
tion

[104887]of the phosphorylase kinase alpha subunit gene on the
mouse X chromosome.

[104888](Abstract) Cytogenet. Cell Genet. 51: 1071–1072, 1989.

[104889]18369.

[104890]Schimke, R. N.; Zakheim, R. M.; Corder, R. C.; Hug, G.:
Glycogen

[104891]storage disease type IX: benign glycogenosis of liver and
hepatic

[104892]phosphorylase kinase deficiency. J. Pediat. 83:
1031–1034, 1973.

[104893]18370.

[104894]Schneider, A.; Davidson, J. J.; Wullrich, A.; Kilimann, M. W.

[104895]Phosphorylase kinase deficiency in I-strain mice is associated with

[104896]a frameshift mutation in the alpha-subunit muscle isoform. Nature

[104897]Genet. 5: 381-385, 1993.

[104898]18371.

[104899]van den Berg, I. E. T.; van Beurden, E. A. C. M.; Malingre, H.

[104900]E. M.; Ploos van Amstel, H. K.; Poll-The, B. T.; Smeitink, J. A. M.;

[104901]Lamers, W. H.; Berger, R.: X-linked liver phosphorylase kinase deficiency

[104902]is associated with mutations in the human liver phosphorylase kinase

[104903]alpha subunit. Am. J. Hum. Genet. 56: 381-387, 1995.

[104904]18372.

[104905]Varsanyi, M.; Vrbica, A.; Heilmeyer, L. M. G., Jr.: X-linked

[104906]dominant inheritance of partial phosphorylase kinase deficiency in

[104907]mice. Biochem. Genet. 18: 247-261, 1980.

[104908]18373.

[104909]Wallis, P. G.; Sidbury, J. B., Jr.; Harris, R. C.: Hepatic phosphorylase

[104910]defect. Studies on peripheral blood. Am. J. Dis. Child. 111: 278–282,

[104911]1966.

[104912]18374.

[104913]Wauters, J. G.; Bossuyt, P. J.; Davidson, J.; Hendrickx, J.; Kilimann,

[104914]M. W.; Willems, P. J.: Regional mapping of a liver alpha-subunit

[104915]gene of phosphorylase kinase (PHKA) to the distal region of human

[104916]chromosome Xp. Cytogenet. Cell Genet. 60: 194–196, 1992.

[104917]18375.

[104918]Willems, P. J.; Gerver, W. J. M.; Berger, R.; Fernandes, J.:

[104919]The natural history of liver glycogenosis due to phosphorylase kinase

[104920]deficiency: a longitudinal study of 41 patients. Europ. J. Pediat. 149:

[104921]268-271, 1990.

[104922]18376.

[104923]Willems, P. J.; Hendrickx, J.; Van der Auwera, B. J.; Vits, L.;

[104924]Raeymaekers, P.; Coucke, P. J.; Van den Bergh, I.; Berger, R.; Smit,

[104925]G. P. A.; Van Broeckhoven, C.; Kilimann, M. W.; Van Elsen, A. F.;

[104926]Fernandes, J. F.: Mapping of the gene for X-linked liver glycogenosis

[104927]due to phosphorylase kinase deficiency to human chromosome region

[104928]Xp22. Genomics 9: 565-569, 1991.

[104929]18377.

[104930]Williams, H. E.; Field, J. B.: Low leukocyte phosphorylase in

[104931]hepatic phosphorylase deficient glycogen storage disease. J. Clin.

[104932]Invest. 40: 1841-1845, 1961.

[104933]18378.

[104934]Zander, N. F.; Meyer, H. E.; Hoffmann-Posorske, E.; Crabb, J.

- [104935]W.; Heilmeyer, L. M. G., Jr.; Kilimann, M. W.: cDNA cloning and complete
[104936]primary structure of skeletal muscle phosphorylase kinase (alpha subunit). Proc.
[104937]Nat. Acad. Sci. 85: 2929–2933, 1988.
[104938]18379.
- [104939]Berta, P.; Hawkins, J. R.; Sinclair, A. H.; Taylor, A.; Griffiths,
[104940]B. L.; Goodfellow, P. N.; Fellous, M.: Genetic evidence equating
[104941]SRY and the testis-determining factor. Nature 348: 448–450, 1990.
[104942]18380.
- [104943]Disteche, C. M.; Casanova, M.; Saal, H.; Friedman, C.; Sybert,
[104944]V.; Graham, J.; Thuline, H.; Page, D.; Fellous, M.: Small deletions
[104945]of the short arm of the Y chromosome in 46,XY females. Proc. Nat.
[104946]Acad. Sci. 83: 7841–7844, 1986.
[104947]18381.
- [104948]Georgopoulos, N. A.; Pralong, F. P.; Seidman, C. E.; Seid-

man,

[104949]J. G.; Crowley, W. F., Jr.; Vallejo, M.: Genetic heterogeneity evidenced

[104950]by low incidence of KAL-1 gene mutations in sporadic cases of gonadotropin-releasing

[104951]hormone deficiency. J. Clin. Endocr. Metab. 82: 213-217, 1997.

[104952]18382.

[104953]Guioli, S.; Incerti, B.; Zanaria, E.; Bardoni, B.; Franco, B.;

[104954]Taylor, K.; Ballabio, A.; Camerino, G.: Kallmann syndrome due to

[104955]a translocation resulting in an X/Y fusion gene. Nature Genet. 1:

[104956]337-340, 1992.

[104957]18383.

[104958]Hardelin, J.-P.; Levilliers, J.; Blanchard, S.; Carel, J.-C.;

[104959]Leutenegger, M.; Pinard-Bertelletto, J.-P.; Bouloux, P.; Petit, C.

[104960]: Heterogeneity in the mutations responsible for X chromosome-linked

[104961]Kallmann syndrome. Hum. Molec. Genet. 2: 373-377,

1993.

[104962]18384.

[104963]Hardelin, J.-P.; Levilliers, J.; del Castillo, I.; Cohen-Salmon,

[104964]M.; Legouis, R.; Blanchard, S.; Compain, S.; Bouloux, P.;

Kirk, J.;

[104965]Moraine, C.; Chaussain, J.-L.; Weissenbach, J.; Petit, C.: X

chromosome-linked

[104966]Kallmann syndrome: stop mutations validate the candi-

date gene. Proc.

[104967]Nat. Acad. Sci. 89: 8190-8194, 1992.

[104968]18385.

[104969]Henkin, R. I.: Abnormalities of taste and olfaction in pa-

tients

[104970]with chromatin negative gonadal dysgenesis. J. Clin. En-

docr. 27:

[104971]1436-1440, 1967.

[104972]18386.

[104973]Hermanussen, M.; Sippell, W. G.: Heterogeneity of Kallmann's

[104974]syndrome. Clin. Genet. 28: 106–111, 1985.

[104975]18387.

[104976]Hipkin, L. J.; Casson, I. F.; Davis, J. C.: Identical twins discordant

[104977]for Kallmann's syndrome. J. Med. Genet. 27: 198–199, 1990.

[104978]18388.

[104979]Incerti, B.; Guioli, S.; Pragliola, A.; Zanaria, E.; Borsani,

[104980]G.; Tonlorenzi, R.; Bardoni, B.; Franco, B.; Wheeler, D.; Balabio,

[104981]A.; Camerino, G.: Kallmann syndrome gene on the X and Y chromosomes:

[104982]implications for evolutionary divergence of human sex chromosomes. Nature

[104983]Genet. 2: 311–314, 1992.

[104984]18389.

[104985]Kirk, J. M. W.; Grant, D. B.; Besser, G. M.; Shalet, S.; Quinton,

[104986]R.; Smith, C. S.; White, M.; Edwards, O.; Bouloux, P.–M. G.:

Unilateral

[104987]renal aplasia in X-linked Kallmann's syndrome. Clin.

Genet. 46:

[104988]260–262, 1994.

[104989]18390.

[104990]Krams, M.; Quinton, R.; Ashburner, J.; Friston, K. J.; Frack-
owiak,

[104991]R. S. J.; Bouloux, P.-M. G.; Passingham, R. E.: Kallmann's
syndrome:

[104992]mirror movements associated with bilateral corticospinal
tract hypertrophy. Neurology 52:

[104993]816–822, 1999.

[104994]18391.

[104995]Legouis, R.; Cohen-Salmon, M.; del Castillo, I.; Levilliers,
J.;

[104996]Capy, L.; Mornon, J.-P.; Petit, C.: Characterization of the
chicken

[104997]and quail homologues of the human gene responsible for
the X-linked

[104998]Kallmann syndrome. Genomics 17: 516–518, 1993.

[104999]18392.

- [105000]Legouis, R.; Hardelin, J.-P.; Levilliers, J.; Claverie, J.-M.;
- [105001]Compain, S.; Wunderle, V.; Millasseau, P.; Le Paslier, D.;
Cohen,
- [105002]D.; Caterina, D.; Bougueleret, L.; Delemarre-Van de Waal,
H.; Lutfalla,
- [105003]G.; Weissenbach, J.; Petit, C.: The candidate gene for the
X-linked
- [105004]Kallmann syndrome encodes a protein related to adhesion
molecules. Cell 67:
- [105005]423-435, 1991.
- [105006]18393.
- [105007]Males, J. L.; Townsend, J. L.; Schneider, R. A.: Hypogo-
nadotropic
- [105008]hypogonadism with anosmia--Kallmann's syndrome: a
disorder of olfactory
- [105009]and hypothalamic function. Arch. Intern. Med. 131:
501-507, 1973.
- [105010]18394.
- [105011]Maya-Nunez, G.; Covarrubias, S. C.; Zenteno, J. C.; Ulloa-
Aguirre,
- [105012]A.; Kofman-Alfaro, S.; Mendez, J. P.: Contiguous gene
syndrome due

[105013] to deletion of the first three exons of the Kallmann gene
and complete

[105014] deletion of the steroid sulphatase gene. Clin. Endocr. 48:
713–718,

[105015] 1998.

[105016] 18395.

[105017] Maya–Nunez, G.; Zenteno, J. C.; Ulloa–Aguirre, A.; Kof–
man–Alfaro,

[105018] S.; Mendez, J. P.: A recurrent missense mutation in the
KAL gene

[105019] in patients with X–linked Kallmann's syndrome. J. Clin. En–
docr. Metab. 83:

[105020] 1650–1653, 1998.

[105021] 18396.

[105022] Meitinger, T.; Heye, B.; Petit, C.; Levilliers, J.; Golla, A.;

[105023] Moraine, C.; Dalla Piccola, B.; Sippell, W. G.; Murken, J.;

Ballabio,

[105024] A.: Definitive localization of X–linked Kallmann syndrome
(hypogonadotropic

[105025] hypogonadism and anosmia) to Xp22.3: close linkage to
the hypervariable

- [105026]repeat sequence CRI-S232. Am. J. Hum. Genet. 47:
664-669, 1990.
- [105027]18397.
- [105028]Nagata, K.; Yamamoto, T.; Chikumi, H.; Ikeda, T.; Yamamoto, H.;
- [105029]Hashimoto, K.; Yoneda, K.; Nanba, E.; Ninomiya, H.; Ishitobi, K.:
- [105030]A novel interstitial deletion of KAL1 in a Japanese family with Kallmann
- [105031]syndrome. J. Hum. Genet. 45: 237-240, 2000.
- [105032]18398.
- [105033]Nass, R.: Mirror movement asymmetries in congenital hemiparesis:
- [105034]the inhibition hypothesis revisited. Neurology 35:
1059-1062, 1985.
- [105035]18399.
- [105036]Nowakowski, H.; Lenz, W.: Genetic aspects in male hypogonadism. Recent
- [105037]Prog. Horm. Res. 17: 53-95, 1961.
- [105038]18400.
- [105039]Oppermann, D.; Happ, J.; Mayr, W. R.: Stimulation of sper-

matogenesis

[105040]and biological paternity by intranasal (low dose) gonadotropin-releasing

[105041]hormone (GnRH) in a male with Kallmann's syndrome: intra-individual

[105042]comparison of GnRH and gonadotropins for stimulation of spermatogenesis. J.

[105043]Clin. Endocr. Metab. 65: 1060-1066, 1987.

[105044]18401.

[105045]Parenti, G.; Rizzolo, M. G.; Ghezzi, M.; Di Maio, S.; Sperandio,

[105046]M. P.; Incerti, B.; Franco, B.; Ballabio, A.; Andria, G.: Variable

[105047]penetrance of hypogonadism in a sibship with Kallmann syndrome due

[105048]to a deletion of the KAL gene. Am. J. Med. Genet. 57: 476-478, 1995.

[105049]18402.

[105050]Pawlowitzki, I. H.; Diekstatt, P.; Schadel, A.; Miny, P.: Estimating

[105051]frequency of Kallmann syndrome among hypogonadic and among anosmic

[105052]patients. Am. J. Med. Genet. 26: 473–479, 1987.

[105053]18403.

[105054]Petit, C.; Levilliers, J.; Weissenbach, J.: Long-range restriction

[105055]map of the terminal part of the short arm of the human X chromosome. Proc.

[105056]Nat. Acad. Sci. 87: 3680–3684, 1990.

[105057]18404.

[105058]Pittman, J.: Personal Communication. Boston, Mass. 1966.

[105059]18405.

[105060]Prager, D.; Braunstein, G. D.: X-chromosome-linked Kallmann's

[105061]syndrome: pathology at the molecular level. J. Clin. Endocr. Metab. 76:

[105062]824–826, 1993.

[105063]18406.

[105064]Imperato-McGinley, J.; Peterson, R. E.; Leshin, M.; Griffin, J.

[105065]E.; Cooper, G.; Draghi, S.; Berenyi, M.; Wilson, J. D.: Steroid

[105066]alpha-reductase deficiency in a 65-year-old male pseudohermaphrodite:

[105067]the natural history, ultrastructure of the testes, and evidence for

[105068]inherited enzyme heterogeneity. J. Clin. Endocr. Metab. 50: 15-22,

[105069]1980.

[105070]18407.

[105071]Jenkins, E. P.; Andersson, S.; Imperato-McGinley, J.; Wilson,

[105072]J. D.; Russell, D. W.: Genetic and pharmacological evidence for more

[105073]than one human steroid 5-alpha-reductase. J. Clin. Invest. 89: 293-300,

[105074]1992.

[105075]18408.

[105076]Francke, U.; Taggart, R. T.: Assignment of the gene for cytoplasmic

[105077]superoxide dismutase (Sod-1) to a region of chromosome 16 and HPRT

[105078]to a region of the X-chromosome in the mouse. Proc. Nat. Acad. Sci. 76:

[105079]5230-5233, 1979.

[105080]18409.

[105081]Francke, U.; Taggart, R. T.: Regional mapping of SOD-1 on
mouse

[105082]chromosome 16, and of HPRT and alpha-GAL (Ags) on the
mouse X, using

[105083]Chinese hamster-mouse T(X;16)16H somatic cell hybrids.
(Abstract) Cytogenet.

[105084]Cell Genet. 25: 155-156, 1979.

[105085]18410.

[105086]Fujimori, S.; Davidson, B. L.; Kelley, W. N.; Palella, T. D.:

[105087]Identification of a single nucleotide change in the hypox-
anthine-guanine

[105088]phosphoribosyltransferase gene (HPRT-Yale) responsible
for Lesch-Nyhan

[105089]syndrome. J. Clin. Invest. 83: 11-13, 1989.

[105090]18411.

[105091]Fujimori, S.; Hidaka, Y.; Davidson, B. L.; Palella, T. D.; Kel-
ley,

[105092]W. N.: Identification of a single nucleotide change in a
mutant gene

- [105093]for hypoxanthine-guanine phosphoribosyltransferase (HPRT-Ann Arbor). Hum. Genet. 79: 39-43, 1988.
- [105094]18412.
- [105095]18412.
- [105096]Fujimori, S.; Kamatani, N.; Nishida, Y.; Ogasawara, N.; Akaoka, I.: Hypoxanthine guanine phosphoribosyltransferase deficiency: nucleotide substitution causing Lesch-Nyhan syndrome identified for the first time among Japanese. Hum. Genet. 84: 483-486, 1990.
- [105097]18413.
- [105098]18413.
- [105099]Fujimori, S.; Tagaya, T.; Kamatani, N.; Akaoka, I.: A germ line mutation within the coding sequence for the putative 5-phosphoribosyl-1-pyrophosphate binding site of hypoxanthine-guanine phosphoribosyltransferase (HPRT) in a Lesch-Nyhan patient: missense mutations within a functionally important region probably cause disease. Hum. Genet. 90: 385-388,

[105106]1992.

[105107]18414.

[105108]Fujimori, S.; Tagaya, T.; Yamaoka, N.; Kamatani, N.;
Akaoka, I.

[105109]: Identification of mutations leading to the Lesch–Nyhan
syndrome

[105110]in two unrelated Japanese patients. (Abstract) Jpn. J. Hum.
Genet. 36:

[105111]58 only, 1991.

[105112]18415.

[105113]Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors–Cayla,
M.

[105114]C.; Gillois, M.: Pig gene mapping: syntenic, attempt to as-
sign the

[105115]histocompatibility complex (SLA). (Abstract) Cytogenet.
Cell Genet. 25:

[105116]159 only, 1979.

[105117]18416.

[105118]Gibbs, R. A.; Caskey, C. T.: Identification and localization

[105119]of mutations at the Lesch–Nyhan locus by ribonuclease A
cleavage. Science 236:

[105120]303–305, 1987.

[105121]18417.

[105122]Gibbs, R. A.; Nguyen, P.–N.; Edwards, A.; Civitello, A. B.;
Caskey,

[105123]C. T.: Multiplex DNA deletion detection and exon se-
quencing of the

[105124]hypoxanthine phosphoribosyltransferase gene in Lesch-
Nyhan families. *Genomics* 7:

[105125]235–244, 1990.

[105126]18418.

[105127]Gibbs, R. A.; Nguyen, P.–N.; McBride, L. J.; Koepf, S. M.;
Caskey,

[105128]C. T.: Identification of mutations leading to the Lesch-
Nyhan syndrome

[105129]by automated direct DNA sequencing of in vitro amplified
cDNA. *Proc.*

[105130]*Nat. Acad. Sci.* 86: 1919–1923, 1989.

[105131]18419.

[105132]Gordon, R. B.; Dawson, P. A.; Sculley, D. G.; Emmerson, B.
T.;

[105133]Caskey, C. T.; Gibbs, R. A.: The molecular characterisation

of HPRT(Chermside)

[105134]and HPRT(Coorparoo): two Lesch–Nyhan patients with reduced amounts

[105135]of mRNA. Gene 108: 299–304, 1991.

[105136]18420.

[105137]Gordon, R. B.; Sculley, D. G.; Dawson, P. A.; Beacham, I. R.;

[105138]Emmerson, B. T.: Identification of a single nucleotide substitution

[105139]in the coding sequence of in vitro amplified cDNA from a patient with

[105140]partial HPRT deficiency (HPRT–Brisbane). J. Inherit. Metab. Dis. 13:

[105141]692–700, 1990.

[105142]18421.

[105143]Greene, M. L.: Clinical features of patients with the 'partial'

[105144]deficiency of the X-linked uricaciduria enzyme. Arch. Intern. Med. 130:

[105145]193–198, 1972.

[105146]18422.

[105147]Gutensohn, W.; Jahn, H.: Partial deficiency of hypoxan–

thine-phosphoribosyltransferase:

[105148]evidence for a structural mutation in a patient with gout.

Europ.

[105149]J. Clin. Invest. 9: 43-47, 1979.

[105150]18423.

[105151]Hashmi, S.; Miller, O. J.: Further evidence of X-linkage of hypoxanthine

[105152]phosphoribosyl-transferase in the mouse. Cytogenet. Cell Genet. 17:

[105153]35-41, 1976.

[105154]18424.

[105155]Anguiano, A.; Oates, R. D.; Amos, J. A.; Dean, M.; Gerrard, B.;

[105156]Stewart, C.; Maher, T. A.; White, M. B.; Milunsky, A.: Congenital

[105157]bilateral absence of the vas deferens: a primarily genital form of

[105158]cystic fibrosis. J.A.M.A. 267: 1794-1797, 1992.

[105159]18425.

[105160]Chillon, M.; Casals, T.; Mercier, B.; Bassas, L.; Lissens, W.;

[105161]Silber, S.; Romey, M.-C.; Ruiz-Romero, J.; Verlingue, C.;

Claustres,

[105162]M.; Nunes, V.; Ferec, C.; Estivill, X.: Mutations in the cystic fibrosis

[105163]gene in patients with congenital absence of the vas deferens. New

[105164]Eng. J. Med. 332: 1475–1480, 1995.

[105165]18426.

[105166]Dumur, V.; Gervais, R.; Rigot, J.-M.; Delomel-Vinner, E.; Decaestecker,

[105167]B.; Lafitte, J.-J.; Roussel, P.: Congenital bilateral absence of

[105168]the vas deferens (CBAVD) and cystic fibrosis transmembrane regulator

[105169](CFTR): correlation between genotype and phenotype. Hum. Genet. 97:

[105170]7–10, 1996.

[105171]18427.

[105172]Andersson, M.; Page, D. C.; de la Chapelle, A.: Chromosome Y-specific

[105173]DNA is transferred to the short arm of X chromosome in human XX males. Science 233:

[105174]786–788, 1986.

[105175]18428.

[105176]Ferguson-Smith, M. A.: X-Y chromosomal interchange in
the aetiology

[105177]of true hermaphroditism and of XX Klinefelter's syndrome.
Lancet II:

[105178]475-476, 1966.

[105179]18429.

[105180]Tiepolo, L.; Zuffardi, O.: Localization of factors controlling

[105181]spermatogenesis in the nonfluorescent portion of the hu-
man Y chromosome

[105182]long arm. Hum. Genet. 34: 119-124, 1976.

[105183]18430.

[105184]Sertie, A. L.; Sossi, V.; Camargo, A. A.; Zatz, M.; Brahe, C.;

[105185]Passos-Bueno, M. R.: Collagen XVIII, containing an en-
dogenous inhibitor

[105186]of angiogenesis and tumor growth, plays a critical role in
the maintenance

[105187]of retinal structure and in neural tube closure (Knobloch
syndrome). Hum.

[105188]Molec. Genet. 9: 2051-2058, 2000.

[105189]18431.

[105190]Fujiwara, H.; Tatsumi, K.; Miki, K.; Harada, T.; Miyai, K.;
Takai,

[105191]S.; Amino, N.: Congenital hypothyroidism caused by a
mutation in

[105192]the Na(+)/I(-) symporter. (Letter) Nature Genet. 16:
124-125, 1997.

[105193]18432.

[105194]Muroi, J.; Yorifuji, T.; Uematsu, A.; Shigematsu, Y.; Oni-
gata,

[105195]K.; Maruyama, H.; Nobutoki, T.; Kitamura, A.; Nakahata,
T.: Molecular

[105196]and clinical analysis of Japanese patients with
3-hydroxy-3-methylglutaryl

[105197]CoA lyase (HL) deficiency. Hum. Genet. 107: 320-326,
2000.

[105198]18433.

[105199]Ozand, P. T.; Al Aqeel, A.; Gascon, G.; Brismar, J.; Thomas,
E.;

[105200]Gleispach, H.: 3-Hydroxy-3-methylglutaryl-coenzyme A
(HMG-CoA) lyase

[105201]deficiency in Saudi Arabia. J. Inherit. Metab. Dis. 14:

174-188,

[105202]1991.

[105203]18434.

[105204]Ozand, P. T.; Devol, E. B.; Gascon, G. G.: Neurometabolic diseases

[105205]at a national referral center: five years experience at the King Faisal

[105206]Specialist Hospital and Research Centre. J. Child Neurol. 7 (suppl.):

[105207]S4-S11, 1992.

[105208]18435.

[105209]Ribes, A.; Briones, P.; Vilaseca, M. A.; Baraibar, R.; Gairi,

[105210]J. M.: Sudden death in an infant with 3-hydroxy-3-methylglutaryl-CoA

[105211]lyase deficiency. J. Inherit. Metab. Dis. 13: 752-753, 1990.

[105212]18436.

[105213]Robinson, B. H.; Oei, J.; Sherwood, W. G.; Slyper, A. H.; Heininger,

[105214]J.; Mamer, O. A.: Hydroxymethylglutaryl CoA lyase deficiency: features

[105215]resembling Reye syndrome. Neurology 30: 714-718,

1980.

[105216]18437.

[105217]Roe, C. R.; Millington, D. S.; Maltby, D. A.: Identification

[105218]of 3-methylglutaryl carnitine: a new diagnostic metabolite
of 3-hydroxy-3-methylglutaryl-coenzyme

[105219]A lyase deficiency. J. Clin. Invest. 77: 1391-1394, 1986.

[105220]18438.

[105221]Shilkin, R.; Wilson, G.; Owles, E.:

3-Hydroxy-3-methylglutaryl

[105222]coenzyme A lyase deficiency: follow-up of first described
case. Acta

[105223]Paediat. Scand. 70: 265-268, 1981.

[105224]18439.

[105225]Shutgens, R. B. H.; Haymans, H.; Ketel, A.: Lethal hypo-
glycemia

[105226]in a child with a deficiency of 3-hydroxy-3-methylglutaryl
coenzyme

[105227]A lyase. J. Pediat. 94: 89-91, 1979.

[105228]18440.

[105229]Sovik, O.; Sweetman, L.; Gibson, K. M.; Nyhan, W. L.: Ge-

netic

[105230] complementation analysis of

3-hydroxy-3-methylglutaryl-coenzyme A

[105231] lyase deficiency in cultured fibroblasts. Am. J. Hum.

Genet. 36:

[105232] 791-801, 1984.

[105233] 18441.

[105234] Wanders, R. J. A.; Schutgens, R. B. H.; Zoeters, P. H. M.:

3-Hydroxy-3-methylglutaryl-CoA

[105235] lyase in human skin fibroblasts: study of its properties
and deficient

[105236] activity in 3-hydroxy-3-methylglutaric aciduria patients
using a simple

[105237] spectrophotometric method. Clin. Chim. Acta 171:
95-102, 1988.

[105238] 18442.

[105239] Wang, S.; Nadeau, J. H.; Duncan, A.; Robert, M.-F.;
Fontaine,

[105240] G.; Schappert, K.; Johnson, K. R.; Zietkiewicz, E.; Hruz, P.;
Miziorko,

[105241] H.; Mitchell, G. A.: 3-Hydroxy-3-methylglutaryl coenzyme
A lyase

[105242](HL): cloning and characterization of a mouse liver HL
cDNA and subchromosomal

[105243]mapping of the human and mouse HL genes. Mammalian
Genome 4: 382-387,

[105244]1993.

[105245]18443.

[105246]Wang, S. P.; Marth, J. D.; Oligny, L. L.; Vachon, M.; Robert,

[105247]M.-F.; Ashmarina, L.; Mitchell, G. A.:

3-Hydroxy-3-methylglutaryl-CoA

[105248]lyase (HL): gene targeting causes prenatal lethality in HL-
deficient

[105249]mice. Hum. Molec. Genet. 7: 2057-2062, 1998.

[105250]18444.

[105251]Wang, S. P.; Robert, M.-F.; Gibson, K. M.; Wanders, R. J. A.;

[105252]Mitchell, G. A.: 3-Hydroxy-3-methylglutaryl CoA lyase
(HL): mouse

[105253]and human HL gene (HMGCL) cloning and detection of
large gene deletions

[105254]in two unrelated HL-deficient patients. Genomics 33:
99-104, 1996.

[105255]18445.

[105256]Wilson, W. G.; Cass, M. B.; Sovik, O.; Gibson, K. M.; Sweetman,

[105257]L.: A child with acute pancreatitis and recurrent hypoglycemia due

[105258]to 3-hydroxy-3-methylglutaryl-CoA lyase deficiency. Europ. J. Pediat. 142:

[105259]289-291, 1984.

[105260]18446.

[105261]Wysocki, S. J.; Hahnel, R.: 3-Hydroxy-3-methylglutaric aciduria:

[105262]3-hydroxy-3-methylglutaryl-coenzyme A lyase levels in leucocytes. Clin.

[105263]Chim. Acta 73: 373-375, 1976.

[105264]18447.

[105265]Wysocki, S. J.; Hahnel, R.:

3-Hydroxy-3-methylglutaryl-coenzyme

[105266]A lyase deficiency: a review. J. Inherit. Metab. Dis. 9: 225-233,

[105267]1986.

[105268]18448.

[105269]Laporte, J.; Kioschis, P.; Hu, L.-J.; Kretz, C.; Carlsson, B.;

[105270]Poustka, A.; Mandel, J. L.; Dahl, N.: Cloning and characterization
[105271]of an alternatively spliced gene in proximal Xq28 deleted
in two patients
[105272]with intersexual genitalia and myotubular myopathy. Genomics 41:
[105273]458-462, 1997.
[105274]18449.

[105275]Caspi, M.; Atlas, R.; Kantor, A.; Sapir, T.; Reiner, O.: Interaction
[105276]between LIS1 and doublecortin, two lissencephaly gene
products. Hum.
[105277]Molec. Genet. 9: 2205-2213, 2000.
[105278]18450.

[105279]des Portes, V.; Francis, F.; Pinard, J.-M.; Desguerre, I.;
Moutard,
[105280]M.-L.; Snoeck, I.; Meiners, L. C.; Capron, F.; Cusmai, R.;
Ricci,
[105281]S.; Motte, J.; Echenne, B.; Ponsot, G.; Dulac, O.; Chelly, J.;
Beldjord,
[105282]C.: Doublecortin is the major gene causing X-linked sub-
cortical laminar

[105283]heterotopia (SCLH). Hum. Molec. Genet. 7: 1063–1070, 1998.

[105284]18451.

[105285]Gleeson, J. G.; Lin, P. T.; Flanagan, L. A.; Walsh, C. A.: Doublecortin

[105286]is a microtubule-associated protein and is expressed widely by migrating

[105287]neurons. Neuron 23: 257–271, 1999.

[105288]18452.

[105289]Gleeson, J. G.; Minnerath, S.; Kuzniecky, R. I.; Dobyns, W. B.;

[105290]Young, I. D.; Ross, M. E.; Walsh, C. A.: Somatic and germline mosaic

[105291]mutations in the doublecortin gene are associated with variable phenotypes. Am.

[105292]J. Hum. Genet. 67: 574–581, 2000.

[105293]18453.

[105294]Gleeson, J. G.; Minnerath, S. R.; Fox, J. W.; Allen, K. M.; Luo,

[105295]R. F.; Hong, S. E.; Berg, M. J.; Kuzniecky, R.; Reitnauer, P. J.;

- [105296]Borgatti, R.; Puche Mira, A.; Guerrini, R.; and 14 others:
Characterization
[105297]of mutations in the gene doublecortin in patients with
double cortex
[105298]syndrome. *Ann. Neurol.* 45: 146–153, 1999.
[105299]18454.
- [105300]Pilz, D. T.; Kuc, J.; Matsumoto, N.; Bodurtha, J.; Bernadi, B.;
[105301]Tassinari, C. A.; Dobyns, W. B.; Ledbetter, D. H.: Subcorti-
cal band
[105302]heterotopia in rare affected males can be caused by mis-
sense mutations
[105303]in DCX (XLIS) or LIS1. *Hum. Molec. Genet.* 8: 1757–1760,
1999.
[105304]18455.
- [105305]Sapir, T.; Horesh, D.; Caspi, M.; Atlas, R.; Burgess, H. A.;
Wolf,
[105306]S. G.; Francis, F.; Chelly, J.; Elbaum, M.; Pietrokovski, S.;
Reiner,
[105307]O.: Doublecortin mutations cluster in evolutionarily con-
served functional
[105308]domains. *Hum. Molec. Genet.* 9: 703–712, 2000.

[105309]18456.

[105310]Shmueli, O.; Gdalyahu, A.; Sorokina, K.; Nevo, E.; Avivi, A.;

[105311]Reiner, O.: DCX in PC12 cells: CREB-mediated transcription and neurite

[105312]outgrowth. Hum. Molec. Genet. 10: 1061–1070, 2001.

[105313]18457.

[105314]Sossey-Alaoui, K.; Hartung, A. J.; Guerrini, R.; Manchester, D.

[105315]K.; Posar, A.; Puche-Mira, A.; Andermann, E.; Dobyns, W. B.; Srivastava,

[105316]A. K.: Human doublecortin (DCX) and the homologous gene in mouse

[105317]encode a putative Ca(2+)-dependent signaling protein which is mutated

[105318]in human X-linked neuronal migration defects. Hum. Molec. Genet. 7:

[105319]1327–1332, 1998.

[105320]18458.

[105321]Yourc'h, P.; Petit, E.; Muh, J. P.; Andres, C.; Bienvenu, T.;

[105322]Beldjord, C.; Chelly, J.; Barthelemy, C.: Exclusion of the coding

- [105323] sequence of the doublecortin gene as a susceptibility locus in autistic disorder. (Letter) Am. J. Med. Genet. 108: 164–167, 2002.
- [105324] 18459.
- [105326] Farnsworth, D.: The Farnsworth–Munsell 100 hue and dichotomy tests for colour vision. J. Ophthal. Soc. Am. 33: 568–578, 1943.
- [105328] 18460.
- [105329] Franceschetti, A.; Klein, D.: Two families with parents of different types of red–green blindness. Acta Genet. Statist. Med. 7: 255–259, 1957.
- [105331] 18461.
- [105332] 18461.
- [105333] Fraser, G. R.: Estimation of the recombination fraction between the protan and deutan loci. Am. J. Hum. Genet. 21: 593–599, 1969.
- [105335] 18462.
- [105336] Kalmus, H.: Diagnosis and Genetics of Defective Colour Vision.

[105337]Oxford: Pergamon Press (pub.) 1965.

[105338]18463.

[105339]Kondo, T.: Untersuchungen bei angeborenen Farbensinn-Anomalien.

[105340]Ueber das Zustandekommen und Wesen der angeborenen Farbensinn-Anomalien. Acta

[105341]Soc. Ophthal. Jpn. 45: 659, 1941.

[105342]18464.

[105343]Nemoto, H.; Murao, M.: A genetic study of colorblindness. Jpn.

[105344]J. Hum. Genet. 6: 165-173, 1961.

[105345]18465.

[105346]Schmidt, I.: A sign of manifest heterozygosity in carriers of

[105347]color deficiency. Am. J. Optom. 32: 404-408, 1955.

[105348]18466.

[105349]Siniscalco, M.; Filippi, G.; Latte, B.: Recombination between

[105350]protan and deutan genes: data on their relative positions in respect

[105351]of the G6PD locus. Nature 204: 1062–1064, 1964.

[105352]18467.

[105353]Sloan, L. L.: Evaluation of the Tokyo Medical College color vision

[105354]test. Am. J. Ophthal. 52: 650–659, 1961.

[105355]18468.

[105356]Thuline, H. C.; Hodgkin, W. E.; Fraser, G. R.; Motulsky, A. G.

[105357]: Genetics of protan and deutan color–vision anomalies: an instructive

[105358]family. Am. J. Hum. Genet. 21: 581–592, 1969.

[105359]18469.

[105360]Vanderdonck, R.; Verriest, G.: Femme protanomale et heterozygote

[105361]mixte (genes de la protanomalie et de la deuteranopie en position

[105362]de repulsion) ayant deux fils deuteranopes, un fils protanomal et

[105363]deux fils normaux. Biotypologie 21: 110–120, 1960.

[105364]18470.

[105365] von Kries, J.; Nagel, W.: Ueber den Einfluss von Licht-
staerke

[105366] und Adaptation auf das Sehen des Dichromaten
(Gruenblinden). Ztschr.

[105367] Psychol. Physiol. Sinnesorg. 12: 1–38, 1896.

[105368] 18471.

[105369] von Planta, P.: Die Haeufigkeit der angeborenen Far-
bensinnstoerungen

[105370] bei Knaben und Maedchen und ihre Feststellung durch die
ueblichen

[105371] klinischen Proben. Albrecht von Graefes Arch. Klin. Exp.
Ophthal. 120:

[105372] 253–281, 1928.

[105373] 18472.

[105374] Waaler, G. H.: Ueber die Erblichkeitsverhaeltnisse der ver-
schieden

[105375] Arten von angeborener Rotgruenblindheit. Ztschr. F. In-
dukt. Abstammungs-u.

[105376] Vererbungsbl. 45: 279–333, 1927.

[105377] 18473.

[105378] Winderickx, J.; Lindsey, D. T.; Sanocki, E.; Teller, D. Y.;

Motulsky,

[105379]A. G.; Deeb, S. S.: Polymorphism in red photopigment underlies variation

[105380]in colour matching. *Nature* 356: 431–433, 1992.

[105381]18474.

[105382]Bordarier, C.; Robain, O.; Rethore, M.–O.; Dulac, O.; Dhellemmes,

[105383]C.: Inverted neurons in agyria: a Golgi study of a case with abnormal

[105384]chromosome 17. *Hum. Genet.* 73: 374–378, 1986.

[105385]18475.

[105386]Chong, S. S.; Lo Nigro, C.; Roschke, A. V.; Tanigami, A.; Pack,

[105387]S. D.; Smith, A. C. M.; Carrozzo, R.; Dobyns, W. B.; Ledbetter, D.

[105388]H.: Point mutations and an intragenic deletion in three ILS patients

[105389]confirm LIS1 as the lissencephaly causative gene in isolated lissencephaly

[105390]sequence and Miller–Dieker syndrome. (Abstract) *Am. J. Hum. Genet.* 59

[105391](suppl.): A23 only, 1996.

[105392]18476.

[105393]Hanahan, D. J. A.: Platelet activating factor: a biologically

[105394]active phosphoglyceride. Annu. Rev. Biochem. 55:

483–509, 1986.

[105395]18477.

[105396]Hattori, M.; Adachi, H.; Tsujimoto, M.; Arai, H.; Inoue, K.:

[105397]Miller–Dieker lissencephaly gene encodes a subunit of

brain platelet–activating

[105398]factor. Nature 370: 216–218, 1994.

[105399]18478.

[105400]Hattori, M.; Arai, H.; Inoue, K.: Purification and characteri-
zation

[105401]of bovine brain platelet–activating factor acetylhydrolase.

J. Biol.

[105402]Chem. 268: 18748–18753, 1993.

[105403]18479.

[105404]Ye, L.; Nakura, J.; Mitsuda, N.; Fujioka, Y.; Kamino, K.;

Ohta,

[105405]T.; Jinno, Y.; Niikawa, N.; Miki, T.; Ogihara, T.: Genetic as-

sociation

[105406]between chromosome 8 microsatellite (MS8-134) and
Werner syndrome

[105407](WRN): chromosome microdissection and homozygosity
mapping. Genomics 28:

[105408]566-569, 1995.

[105409]18480.

[105410]Yu, C.-E.; Oshima, J.; Fu, Y.-H.; Wijsman, E. M.; Hisama,
F.;

[105411]Alisch, R.; Matthews, S.; Nakura, J.; Miki, T.; Quais, S.;
Martin,

[105412]G. M.; Mulligan, J.; Schellenberg, G. D.: Positional cloning
of the

[105413]Werner's syndrome gene. Science 272: 258-262, 1996.

[105414]18481.

[105415]Yu, C.-E.; Oshima, J.; Goddard, K. A. B.; Miki, T.; Nakura,
J.;

[105416]Ogihara, T.; Poot, M.; Hoehn, H.; Fraccaro, M.; Piussan, C.;
Martin,

[105417]G. M.; Schellenberg, G. D.; Wijsman, E. M.: Linkage dise-
quilibrium

[105418]and haplotype studies of chromosome 8p11.1-21.1
markers and Werner

- [105419]syndrome. Am. J. Hum. Genet. 55: 356–364, 1994.
- [105420]18482.
- [105421]Yu, C.-E.; Oshima, J.; Wijsman, E. M.; Nakura, J.; Miki, T.;
Piussan,
[105422]C.; Matthews, S.; Fu, Y.-H.; Mulligan, J.; Martin, G. M.;
Schellenberg,
[105423]G. D.; Werner's Syndrome Collaborative Group: Mutations
in the
[105424]consensus helicase domains of the Werner syndrome
gene. Am. J. Hum.
[105425]Genet. 60: 330–341, 1997.
- [105426]18483.
- [105427]Gibbons, R. J.; Brueton, L.; Buckle, V. J.; Burn, J.; Clayton-
Smith,
[105428]J.; Davison, B. C. C.; Gardner, R. J. M.; Homfray, T.; Kear-
ney, L.;
[105429]Kingston, H. M.; Newbury-Ecob, R.; Porteous, M. E. P.;
Wilkie, A.
[105430]O. M.; Higgs, D. R.: Clinical and hematologic aspects of
the X-linked
[105431]alpha-thalassemia/mental retardation syndrome (ATR-X).
Am. J. Med.

[105432]Genet. 55: 288–299, 1995.

[105433]18484.

[105434]Marchese, A.; Sawzdargo, M.; Nguyen, T.; Cheng, R.;
Heng, H. H.

[105435]Q.; Nowak, T.; Im, D-S.; Lynch, K. R.; George, S. R.;
O'Dowd, B. F.

[105436]: Discovery of three novel orphan G-protein-coupled re-
ceptors. Genomics 56:

[105437]12–21, 1999.

[105438]18485.

[105439]Sanchis, D.; Fleury, C.; Chomiki, N.; Gubern, M.; Huang,
Q.; Neverova,

[105440]M.; Gregoire, F.; Easlick, J.; Raimbault, S.; Levi-Meyrueis,
C.; Miroux,

[105441]B.; Collins, S.; Seldin, M.; Richard, D.; Warden, C.; Bouil-
laud, F.;

[105442]Ricquier, D.: BMCP1, a novel mitochondrial carrier with
high expression

[105443]in the central nervous system of humans and rodents, and
respiration

[105444]uncoupling activity in recombinant yeast. J. Biol. Chem.
273: 34611–34615,

[105445]1998.

[105446]18486.

[105447]Blanco, P.; Sargent, C. A.; Boucher, C. A.; Mitchell, M.; Af-
fara,

[105448]N. A.: Conservation of PCDHX in mammals; expression of
X/Y genes

[105449]predominantly in brain. Mammalian Genome 11: 906–914,
2000.

[105450]18487.

[105451]Ciccodicola, A.; D'Esposito, M.; Esposito, T.; Gian-
francesco, F.;

[105452]Migliaccio, C.; Miano, M. G.; Matarazzo, M. R.; Vacca, M.;
Franze,

[105453]A.; Cuccurese, M.; Cocchia, M.; Curci, A.; and 9 others:
Differentially

[105454]regulated and evolved genes in the fully sequenced Xq/Yq
pseudoautosomal

[105455]region. Hum. Molec. Genet. 9: 395–401, 2000.

[105456]18488.

[105457]Mumm, S.; Molini, B.; Terrell, J.; Srivastava, A.; Sch-
lessinger,

[105458]D.: Evolutionary features of the 4-Mb Xq21.3 XY homology region revealed
[105459]by a map at 60-kb resolution. *Genome Res.* 7: 307–314, 1997.
[105460]18489.
[105461]Schwartz, A.; Chan, D. C.; Brown, L. G.; Alagappan, R.; Pettay,
[105462]D.; Disteche, C.; McGillivray, B.; de la Chapelle, A.; Page, D. C.
[105463]: Reconstructing hominid Y evolution: X-homologous block, created
[105464]by X–Y transposition, was disrupted by Yp inversion through LINE
[105465]recombination. *Hum. Molec. Genet.* 7: 1–11, 1998.
[105466]18490.
[105467]Tilford, C. A.; Kuroda-Kawaguchi, T.; Skaletsky, H.; Rozen, S.;
[105468]Brown, L. G.; Rosenberg, M.; McPherson, J. D.; Wylie, K.; Sekhon,
[105469]M.; Kucaba, T. A.; Waterston, R. H.; Page, D. C.: A physical map
[105470]of the human Y chromosome. *Nature* 409: 943–945,

2001.

[105471]18491.

[105472]des Portes, V.; Pinard, J. M.; Billuart, P.; Vinet, M. C.;

Koulakoff,

[105473]A.; Carrie, A.; Gelot, A.; Dupuis, E.; Motte, J.; Berwald-Net-
ter,

[105474]Y.; Catala, M.; Kahn, A.; Beldjord, C.; Chelly, J.: A novel
CNS gene

[105475]required for neuronal migration and involved in X-linked
subcortical

[105476]laminal heterotopia and lissencephaly syndrome. Cell 92:
51-61,

[105477]1998.

[105478]18492.

[105479]Gleeson, J. G.; Allen, K. M.; Fox, J. W.; Lamperti, E. D.;
Berkovic,

[105480]S.; Scheffer, I.; Cooper, E. C.; Dobyns, W. B.; Minnerath, S.
R.;

[105481]Ross, M. E.; Walsh, C. A.: Doublecortin, a brain-specific
gene mutated

[105482]in human X-linked lissencephaly and double cortex syn-
drome, encodes

[105483]a putative signaling protein. Cell 92: 63–72, 1998.

[105484]18493.

[105485]Matsumoto, N.; Leventer, R. J.; Kuc, J. A.; Mewborn, S. K.;
Dudliceck,

[105486]L. L.; Ramocki, M. B.; Pilz, D. T.; Mills, P. L.; Das, S.; Ross,
M.

[105487]E.; Ledbetter, D. H.; Dobyns, W. B.: Mutation analysis of
the DCX

[105488]gene and genotype/phenotype correlation in subcortical
band heterotopia. Europ.

[105489]J. Hum. Genet. 9: 5–12, 2001.

[105490]18494.

[105491]Batch, J. A.; Evans, B. A. J.; Hughes, I. A.; Patterson, M. N.

[105492]: Mutations of the androgen receptor gene identified in
perineal hypospadias. J.

[105493]Med. Genet. 30: 198–201, 1993.

[105494]18495.

[105495]Gibbons, R. J.; McDowell, T. L.; Raman, S.; O'Rourke, D. M.;
Garrick,

[105496]D.; Ayyub, H.; Higgs, D. R.: Mutations in ATRX, encoding a
SWI/SNF-like

[105497]protein, cause diverse changes in the pattern of DNA methylation. Nature

[105498]Genet. 24: 368–371, 2000.

[105499]18496.

[105500]Gibbons, R. J.; Picketts, D. J.; Villard, L.; Higgs, D. R.: Mutations

[105501]in a putative global transcriptional regulator cause X-linked mental

[105502]retardation with alpha-thalassemia (ATR-X syndrome). Cell 80: 837–845,

[105503]1995.

[105504]18497.

[105505]Kiesewetter, S.; Macek, M., Jr.; Davis, C.; Curristin, S. M.;

[105506]Chu, C.-S.; Graham, C.; Shrimpton, A. E.; Cashman, S. M.; Tsui, L.-C.;

[105507]Mickle, J.; Amos, J.; Highsmith, W. E.; Shuber, A.; Witt, D. R.; Crystal,

[105508]R. G.; Cutting, G. R.: A mutation in CFTR produces different phenotypes

[105509]depending on chromosomal background. Nature Genet. 5: 274–278, 1993.

[105510]18498.

[105511]Lossi, A. M.; Millan, J. M.; Villard, L.; Orellana, C.; Cardoso,

[105512]C.; Prieto, F.; Fontes, M.; Martinez, F.: Mutation of the
XNP/ATR-X

[105513]gene in a family with severe mental retardation, spastic
paraplegia

[105514]and skewed pattern of X inactivation: demonstration that
the mutation

[105515]is involved in the inactivation bias. (Letter) Am. J. Hum.
Genet. 65:

[105516]558-562, 1999.

[105517]18499.

[105518]Martinez, F.; Tomas, M.; Millan, J. M.; Fernandez, A.; Palau,

[105519]F.; Prieto, F.: Genetic localisation of mental retardation
with spastic

[105520]diplegia to the pericentromeric region of the X chromo-
some: X inactivation

[105521]in female carriers. J. Med. Genet. 35: 284-287, 1998.

[105522]18500.

[105523]Mattei, J. F.; Collignon, P.; Ayme, S.; Giraud, F.: X-linked

[105524]mental retardation, growth retardation, deafness and microgenitalism:

[105525]a second familial report. Clin. Genet. 23: 70-74, 1983.

[105526]18501.

[105527]McDowell, T. L.; Gibbons, R. J.; Sutherland, H.; O'Rourke, D.

[105528]M.; Bickmore, W. A.; Pombo, A.; Turley, H.; Gatter, K.; Picketts,

[105529]D. J.; Buckle, V. J.; Chapman, L.; Rhodes, D.; Higgs, D. R.: Localization

[105530]of a putative transcriptional regulator (ATRX) at pericentromeric

[105531]heterochromatin and the short arms of acrocentric chromosomes. Proc.

[105532]Nat. Acad. Sci. 96: 13983-13988, 1999.

[105533]18502.

[105534]Pask, A.; Renfree, M. B.; Graves, J. A. M.: The human sex-reversing

[105535]ATRX gene has a homologue on the marsupial Y chromosome, ATRY: implications

[105536]for the evolution of mammalian sex determination. Proc. Nat. Acad.

[105537]Sci. 97: 13198–13202, 2000.

[105538]18503.

[105539]Picketts, D. J.; Higgs, D. R.; Bachoo, S.; Blake, D. J.; Quarrell,

[105540]O. W. J.; Gibbons, R. J.: ATRX encodes a novel member of the SNF2

[105541]family of proteins: mutations point to a common mechanism underlying

[105542]the ATR-X syndrome. Hum. Molec. Genet. 5: 1899–1907, 1996.

[105543]18504.

[105544]Picketts, D. J.; Tastan, A. O.; Higgs, D. R.; Gibbons, R. J.:

[105545]Comparison of the human and murine ATRX gene identifies highly conserved,

[105546]functionally important domains. Mammalian Genome 9: 400–403, 1998.

[105547]18505.

[105548]Reardon, W.; Gibbons, R. J.; Winter, R. M.; Baraitser, M.: Male

[105549]pseudohermaphroditism in sibs with the alpha-thalassemia/mental retardation

- [105550](ATR-X) syndrome. Am. J. Med. Genet. 55: 285–287, 1995.
- [105551]18506.
- [105552]Stayton, C. L.; Dabovic, B.; Gulisano, M.; Gecz, J.; Broccoli, V.; Giovanazzi, S.; Bossolasco, M.; Monaco, L.; Rastan, S.; Boncinelli, E.; Bianchi, M. E.; Consalez, G. G.: Cloning and characterization of a new human Xq13 gene, encoding a putative helicase. Hum. Molec. Genet. 3: 1957–1964, 1994.
- [105556]18507.
- [105558]Villard, L.; Bonino, M.-C.; Abidi, F.; Ragusa, A.; Belougne, J.; Lossi, A.-M.; Seaver, L.; Bonnefont, J.-P.; Romano, C.; Fichera, M.; Lacombe, D.; Hanauer, A.; Philip, N.; Schwartz, C.; Fontes, M.: Evaluation of a mutation screening strategy for sporadic cases of ATR-X syndrome. J. Med. Genet. 36: 183–186, 1999.

[105563]18508.

[105564]Villard, L.; Fontes, M.; Ades, L. C.; Gecz, J.: Identification
[105565]of a mutation in the XNP/ATR-X gene in a family reported
as Smith-Fineman-Myers
[105566]syndrome. (Letter) Am. J. Med. Genet. 91: 83-85, 2000.

[105567]18509.

[105568]Villard, L.; Gecz, J.; Mattei, J. F.; Fontes, M.; Saugier-Weber,
[105569]P.; Munnich, A.; Lyonnet, S.: XNP mutation in a large fam-
ily with
[105570]Juberg-Marsidi syndrome. (Letter) Nature Genet. 12:
359-360, 1996.

[105571]18510.

[105572]Villard, L.; Lacombe, D.; Fontes, M.: A point mutation in
the
[105573]XNP gene, associated with an ATR-X phenotype without
alpha-thalassemia. Europ.

[105574]J. Hum. Genet. 4: 316-320, 1996.

[105575]18511.

[105576]Villard, L.; Lossi, A.-M.; Cardoso, C.; Proud, V.; Chiaroni,
P.;

[105577]Colleaux, L.; Schwartz, C.; Fontes, M.: Determination of the genomic structure of the XNP/ATRX gene encoding a potential zinc finger helicase. Genomics 43: 149–155, 1997.

[105580]18512.

[105581]Villard, L.; Toutain, A.; Lossi, A.-M.; Gecz, J.; Houdayer, C.; Moraine, C.; Fontes, M.: Splicing mutation in the ATR-X gene can lead to a dysmorphic mental retardation phenotype without alpha-thalassemia. Am. J. Hum. Genet. 58: 499–505, 1996.

[105585]18513.

[105586]Wada, T.; Kubota, T.; Fukushima, Y.; Saitoh, S.: Molecular genetic study of Japanese patients with X-linked alpha-thalassemia/mental retardation syndrome (ATR-X). Am. J. Med. Genet. 94: 242–248, 2000.

[105589]18514.

[105590]Wang, L. H.; Collins, A.; Lawrence, S.; Keats, B. J.; Morton,

[105591]N. E.: Integration of gene maps: chromosome X. Genomics
22: 590-604,

[105592]1994.

[105593]18515.

[105594]Yntema, H. G.; Poppelaars, F. A.; Derksen, E.; Oudakker, A.
R.;

[105595]van Roosmalen, T.; Jacobs, A.; Obbema, H.; Brunner, H. G.;
Hamel,

[105596]B. C. J.; van Bokhoven, H.: Expanding phenotype of XNP
mutations:

[105597]mild to moderate mental retardation. Am. J. Med. Genet.
110: 243-247,

[105598]2002.

[105599]18516.

[105600]Borkhardt, A.; Repp, R.; Haas, O. A.; Leis, T.; Harbott, J.;
Kreuder,

[105601]J.; Hammermann, J.; Henn, T.; Lampert, F.: Cloning and
characterization

[105602]of AFX, the gene that fuses to MLL in acute leukemias with
a t(X;11)(q13;q23). Oncogene 14:

[105603]195-202, 1997.

[105604]18517.

[105605]Katz, F.; Hinshelwood, S.; Rutland, P.; Jones, A.; Kinnon, C.;

[105606]Morgan, G.: Mutation analysis in CD40 ligand deficiency leading to

[105607]X-linked hypogammaglobulinemia with hyper IgM syndrome. Hum. Mutat. 8:

[105608]223–228, 1996.

[105609]18518.

[105610]Kolvraa, S.; Kruse, T. A.; Jensen, P. K. A.; Linde, K. H.; Vestergaard,

[105611]S. R.; Bolund, L.: Close linkage between X-linked ectodermal dysplasia

[105612]and a cloned DNA sequence detecting a two allele restriction fragment

[105613]length polymorphism in the region Xp11–q12. Hum. Genet. 74: 284–287,

[105614]1986.

[105615]18519.

[105616]Korthauer, U.; Graf, D.; Mages, H. W.; Briere, F.; Padayachee,

- [105617]M.; Malcolm, S.; Ugazio, A. G.; Notarangelo, L. D.; Levin-sky, R. J.;
- [105618]Kroczek, R. A.: Defective expression of T-cell CD40 ligand causes
- [105619]X-linked immunodeficiency with hyper-IgM. Nature 361: 539-541, 1993.
- [105620]18520.
- [105621]Kraakman, M. E. M.; de Weers, M.; Espanol, T.; Schuurman, R. K.
- [105622]B.; Hendriks, R. W.: Identification of a CD40L gene mutation and
- [105623]genetic counselling in a family with immunodeficiency with hyperimmunoglobulinemia
- [105624]M. Clin. Genet. 48: 46-48, 1995.
- [105625]18521.
- [105626]Kroczek, R. A.; Graf, D.; Brugnani, D.; Giliani, S.; Korthauer,
- [105627]U.; Ugazio, A.; Senger, G.; Mages, H. W.; Villa, A.; Notarangelo,
- [105628]L. D.: Defective expression of CD40 ligand on T cell causes 'X-linked
- [105629]immunodeficiency with hyper-IgM (HIGM1)'. Immun. Rev.

138: 39–59,

[105630]1994.

[105631]18522.

[105632]Lin, Q.; Rohrer, J.; Allen, R. C.; Larche, M.; Greene, J. M.;

[105633]Shigeoka, A. O.; Gatti, R. A.; Derauf, D. C.; Belmont, J. W.;

Conley,

[105634]M. E.: A single strand conformation polymorphism study
of CD40 ligand:

[105635]efficient mutation analysis and carrier detection of X-
linked hyper

[105636]IgM syndrome. J. Clin. Invest. 97: 196–201, 1996.

[105637]18523.

[105638]Nonoyama, S.; Shimadzu, M.; Toru, H.; Seyama, K.; Nunoi,
H.; Neubauer,

[105639]M.; Yata, J.; Och, H. D.: Mutations of the CD40 ligand gene
in 13

[105640]Japanese patients with X-linked hyper-IgM syndrome.
Hum. Genet. 99:

[105641]624–627, 1997.

[105642]18524.

[105643]Padayachee, M.; Feighery, C.; Finn, A.; McKeown, C.;

Levinsky,

[105644]R. J.; Kinnon, C.; Malcolm, S.: Mapping of the X-linked
form of hyper-IgM

[105645]syndrome (HIGM1) to Xq26 by close linkage to HPRT. Ge-
nomics 14:

[105646]551-553, 1992.

[105647]18525.

[105648]Macchi, P.; Villa, A.; Strina, D.; Sacco, M. G.; Morali, F.;
Brugnoni,

[105649]D.; Giliani, S.; Mantuano, E.; Fasth, A.; Andersson, B.;
Zegers, B.

[105650]J. M.; Cavagni, G.; Reznick, I.; Levy, J.; Zan-Bar, I.; Porat,
Y.;

[105651]Airo, P.; Plebani, A.; Vezzoni, P.; Notarangelo, L. D.: Char-
acterization

[105652]of nine novel mutations in the CD40 ligand gene in pa-
tients with X-linked

[105653]hyper IgM syndrome of various ancestry. Am. J. Hum.
Genet. 56: 898-906,

[105654]1995.

[105655]18526.

[105656]Padayachee, M.; Levinsky, R. J.; Kinnon, C.; Finn, A.; McKe-

own,

[105657]C.; Feighery, C.; Notarangelo, L. D.; Hendriks, R. W.; Read, A. P.;

[105658]Malcolm, S.: Mapping of the X linked form of hyper IgM syndrome (HIGM1). J.

[105659]Med. Genet. 30: 202–205, 1993.

[105660]18527.

[105661]Pilia, G.; Porta, B.; Padayachee, M.; Malcolm, S.; Zucchi, I.;

[105662]Villa, A.; Macchi, P.; Vezzoni, P.; Schlessinger, D.: Human CD40L

[105663]gene maps between DXS144E and DXS300 in Xq26. Genomics 22: 249–251,

[105664]1994.

[105665]18528.

[105666]Ramesh, N.; Fuleihan, R.; Geha, R.: Molecular pathology of X-linked

[105667]immunoglobulin deficiency with normal or elevated IgM (HIGMX-1). Immun.

[105668]Rev. 138: 87–104, 1994.

[105669]18529.

[105670]Seyama, K.; Kira, S.; Ishidoh, K.; Souma, S.; Miyakawa, T.;

Kominami,

[105671]E.: Genomic structure and PCR-SSCP analysis of the human CD40 ligand

[105672]gene: its application to prenatal screening for X-linked hyper-IgM

[105673]syndrome. Human Genet. 97: 180-185, 1996.

[105674]18530.

[105675]Seyama, K.; Nonoyama, S.; Gangsaas, I.; Hollenbaugh, D.; Pabst,

[105676]H. F.; Aruffo, A.; Ochs, H. D.: Mutations of the CD40 ligand and gene

[105677]and its effect on CD40 ligand expression in patients with X-linked

[105678]hyper IgM syndrome. Blood 92: 2421-2434, 1998.

[105679]18531.

[105680]Tan, J.; Town, T.; Crawford, F.; Mori, T.; DelleDonne, A.; Crescentini,

[105681]R.; Obregon, D.; Flavell, R. A.; Mullan, M. J.: Role of CD40 ligand

[105682]in amyloidosis in transgenic Alzheimer's mice. Nature Neurosci. 28Oct,

[105683]2002. Note: Advance Electronic Publication.

[105684]18532.

[105685]Chatterjee, A.; Faust, C. J.; Herman, G. E.: Genetic and physical

[105686]mapping of the biglycan gene on the mouse X chromosome. Mammalian

[105687]Genome 4: 33–36, 1993.

[105688]18533.

[105689]Das, S.; Metzenberg, A.; Pai, G. S.; Gitschier, J.: Mutational

[105690]analysis of the biglycan gene excludes it as a candidate for X-linked

[105691]dominant chondrodysplasia punctata, dyskeratosis congenita, and incontinentia

[105692]pigmenti. (Letter) Am. J. Hum. Genet. 54: 922–925, 1994.

[105693]18534.

[105694]Fisher, L. W.; Heegaard, A.-M.; Vetter, U.; Vogel, W.; Just, W.;

[105695]Termine, J. D.; Young, M. F.: Human biglycan gene: putative promoter,

[105696]intron–exon junctions, and chromosomal localization. J. Biol. Chem. 266:

[105697]14371–14377, 1991.

[105698]18535.

[105699]Fisher, L. W.; Termine, J. D.; Young, M. F.: Deduced-protein sequence

[105700]of bone small proteoglycan I (biglycan) shows homology with proteoglycan

[105701]II (decorin) and several nonconnective tissue proteins in a variety

[105702]of species. J. Biol. Chem. 264: 4571-4576, 1989.

[105703]18536.

[105704]Geerkens, C.; Vetter, U.; Just, W.; Fedarko, N. S.; Fisher, L.

[105705]W.; Young, M. F.; Termine, J. D.; Gehron Robey, P.; Wohrle, D.; Vogel,

[105706]W.: The X-chromosomal human biglycan gene BGN is subject to X inactivation

[105707]but is transcribed like an X-Y homologous gene. Hum. Genet. 96:

[105708]44-52, 1995.

[105709]18537.

[105710]McBride, O. W.; Fisher, L. W.; Young, M. F.: Localization of PGI

[105711](biglycan, BGN) and PGII (decorin, DCN, PG-40) genes on

human chromosomes

[105712]Xq13-qter and 12q, respectively. Genomics 6: 219-225,
1990.

[105713]18538.

[105714]Traupe, H.; van den Ouweland, A. M. W.; van Oost, B. A.;
Vogel,

[105715]W.; Vetter, U.; Warren, S. T.; Rocchi, M.; Darlison, M. G.;
Ropers,

[105716]H.-H.: Fine mapping of the human biglycan (BGN) gene
within the Xq28

[105717]region employing a hybrid cell panel. Genomics 13:
481-483, 1992.

[105718]18539.

[105719]Wegrowski, Y.; Pillarisetti, J.; Danielson, K. G.; Suzuki, S.;

[105720]Iozzo, R. V.: The murine biglycan: complete cDNA cloning,
genomic

[105721]organization, promoter function, and expression. Ge-
nomics 30: 8-17,

[105722]1995.

[105723]18540.

[105724]Xu, T.; Bianco, P.; Fisher, L. W.; Longenecker, G.; Smith, E.;

[105725]Goldstein, S.; Bonadio, J.; Boskey, A.; Heegaard, A.-M.;
Sommer, B.;

[105726]Satomura, K.; Dominguez, P.; Zhao, C.; Kulkarni, A. B.;
Robey, P.

[105727]G.; Young, M. F.: Targeted disruption of the biglycan gene
leads

[105728]to an osteoporosis-like phenotype in mice. Nature Genet.
20: 78-82,

[105729]1998.

[105730]18541.

[105731]Sandhoff, K.; Harzer, K.; Wasse, W.; Jatzkewitz, H.: En-
zyme

[105732]alterations and lipid storage in three variants of Tay-Sachs
disease. J.

[105733]Neurochem. 18: 2469-2489, 1971.

[105734]18542.

[105735]Bowcock, A. M.; Farrer, L. A.; Hebert, J. M.; Agger, M.;
Sternlieb,

[105736]I.; Scheinberg, I. H.; Buys, C. H. C. M.; Scheffer, H.; Fryd-
man, M.;

[105737]Chajek-Saul, T.; Bonne-Tamir, B.; Cavalli-Sforza, L. L.:
Eight closely

[105738]linked loci place the Wilson disease locus within
13q14-q21. Am.

[105739]J. Hum. Genet. 43: 664-674, 1988.

[105740]18543.

[105741]Yoshida, K.; Sugano, S.: Identification of a novel protocad-
herin

[105742]gene (PCDH11) on the human XY homology region in
Xq21.3. Genomics 62:

[105743]540-543, 1999.

[105744]18544.

[105745]de la Chapelle, A.; Sankila, E.-M.; Lindlof, M.; Aula, P.; No-
rio,

[105746]R.: Norrie disease caused by a gene deletion allowing car-
rier detection

[105747]and prenatal diagnosis. Clin. Genet. 28: 317-320, 1985.

[105748]18545.

[105749]Denney, R. M.; Fritz, R. R.; Patel, N. T.; Abell, C. W.: Hu-
man

[105750]liver MAO-A and MAO-B separated by immunoaffinity
chromatography with

[105751]MAO-B-specific monoclonal antibody. Science 215:

1400–1403, 1982.

[105752]18546.

[105753]Kochersperger, L. M.; Parker, E. L.; Siciliano, M.; Darling-
ton,

[105754]G. J.; Denney, R. M.: Assignment of genes for human
monoamine oxidases

[105755]A and B to the X chromosome. J. Neurosci. Res. 16:
601–616, 1986.

[105756]18547.

[105757]Sims, K. B.; de la Chapelle, A.; Norio, R.; Sankila, E.–M.;
Hsu,

[105758]Y.–P. P.; Rinehart, W. B.; Corey, T. J.; Ozelius, L.; Powell, J.
F.;

[105759]Bruns, G.; Gusella, J. F.; Murphy, D. L.; Breakefield, X. O.:
Monoamine

[105760]oxidase deficiency in males with an X chromosome dele-
tion. Neuron 2:

[105761]1069–1076, 1989.

[105762]18548.

[105763]Binda, C.; Newton–Vinson, P.; Hubalek, F.; Edmondson, D.
E.; Mattevi,

[105764]A.: Structure of human monoamine oxidase B, a drug tar-

get for the

[105765]treatment of neurological disorders. Nature Struct. Biol. 9:
22–26,

[105766]2002.

[105767]18549.

[105768]Checkoway, H.; Franklin, G. M.; Costa-Mallen, P.; Smith-
Weller,

[105769]T.; Dilley, J.; Swanson, P. D.; Costa, L. G.: A genetic poly-
morphism

[105770]of MAO-B modifies the association of cigarette smoking
and Parkinson's

[105771]disease. Neurology 50: 1458–1461, 1998.

[105772]18550.

[105773]Fowler, J. S.; Volkow, N. D.; Wang, G.-J.; Pappas, N.; Lo-
gan, J.;

[105774]MacGregor, R.; Alexoff, D.; Shea, C.; Schlyer, D.; Wolf, A.
P.; Warner,

[105775]D.; Zezulkova, I.; Cilento, R.: Inhibition of monoamine oxi-
dase B

[105776]in the brains of smokers. Nature 379: 733–736, 1996.

[105777]18551.

- [105778] Hamano, Y.; Kodama, H.; Fujikawa, Y.; Tanaka, Y.;
Nishimura, K.;
- [105779] Yanagisawa, M.: Use of immunocytochemical analysis of a
duodenal
- [105780] biopsy specimen to identify a carrier of ornithine tran-
scarbamylase
- [105781] deficiency. New Eng. J. Med. 318: 1521-1523, 1988.
- [105782] 18552.
- [105783] Bull, P. C.; Thomas, G. R.; Rommens, J. M.; Forbes, J. R.;
- Cox,
- [105784] D. W.: The Wilson disease gene is a putative copper trans-
porting
- [105785] P-type ATPase similar to the Menkes gene. Nature Genet.
5: 327-337,
- [105786] 1993.
- [105787] 18553.
- [105788] Forbes, J. R.; Cox, D. W.: Copper-dependent trafficking of
Wilson
- [105789] disease mutant ATP7B proteins. Hum. Molec. Genet. 9:
1927-1935,
- [105790] 2000.

[105791]18554.

[105792]Ha-Hao, D.; Hefter, H.; Stremmel, W.; Castaneda-Guillot, C.; Hernandez,

[105793]A. H.; Cox, D. W.; Auburger, G.: His1069-to-gln and six novel Wilson

[105794]disease mutations: analysis of relevance for early diagnosis and phenotype. Europ.

[105795]J. Hum. Genet. 6: 616-623, 1998.

[105796]18555.

[105797]Grino, P. B.; Griffin, J. E.; Cushard, W. G., Jr.; Wilson, J.

[105798]D.: A mutation of the androgen receptor associated with partial androgen

[105799]resistance, familial gynecomastia, and fertility. J. Clin. Endocr.

[105800]Metab. 66: 754-761, 1988.

[105801]18556.

[105802]Holterhus, P.-M.; Sinnecker, G. H. G.; Hiort, O.: Phenotypic

[105803]diversity and testosterone-induced normalization of mutant L712F androgen

[105804]receptor function in a kindred with androgen insensitivity. J. Clin.

[105805]Endocr. Metab. 85: 3245–3250, 2000.

[105806]18557.

[105807]Hughes, I. A.; Evans, B. A. J.: Complete androgen insensitivity

[105808]syndrome characterized by increased concentration of a normal androgen

[105809]receptor in genital skin fibroblasts. J. Clin. Endocr. Metab. 63:

[105810]309–315, 1986.

[105811]18558.

[105812]Imperato-McGinley, J.; Ip, N. Y.; Gautier, T.; Neuweiler, J.;

[105813]Gruenspan, H.; Liao, S.; Chang, C.; Balazs, I.: DNA linkage analysis

[105814]and studies of the androgen receptor gene in a large kindred with

[105815]complete androgen insensitivity. Am. J. Med. Genet. 36: 104–108,

[105816]1990.

[105817]18559.

[105818]Jukier, L.; Kaufman, M.; Pinsky, L.; Peterson, R. E.: Partial

- [105819] androgen resistance associated with secondary
5-alpha-reductase deficiency:
- [105820] identification of a novel qualitative androgen receptor de-
fect and
- [105821] clinical implications. J. Clin. Endocr. Metab. 59: 679-688,
1984.
- [105822] 18560.
- [105823] Kaufman, M.; Pinsky, L.; Bowin, A.; Au, M. W. S.: Familial
external
- [105824] genital ambiguity due to a transformation defect of an-
drogen-receptor
- [105825] complexes that is expressed with
- 5-alpha-dihydrotestosterone and the
- [105826] synthetic androgen methyltrienolone. Am. J. Med. Genet.
18: 493-507,
- [105827] 1984.
- [105828] 18561.
- [105829] Kaufman, M.; Straisfeld, C.; Pinsky, L.: Male pseudo-
hermaphroditism
- [105830] presumably due to target organ unresponsiveness to an-
drogens: deficient
- [105831] 5-alpha-dihydrotestosterone binding in cultured skin fi-

broblasts. J.

[105832]Clin. Invest. 58: 345-350, 1976.

[105833]18562.

[105834]Keenan, B. S.; Meyer, W. J., III; Hadjian, A. J.; Jones, H. W.;

[105835]Migeon, C. J.: Syndrome of androgen insensitivity in man:
absence

[105836]of 5- α -dihydrotestosterone binding protein in skin fi-
broblasts. J.

[105837]Clin. Endocr. 38: 1143-1146, 1974.

[105838]18563.

[105839]Liao, S.; Witte, D.: Autoimmune anti-androgen-receptor
antibodies

[105840]in human serum. Proc. Nat. Acad. Sci. 82: 8345-8348,
1985.

[105841]18564.

[105842]Lin, S.-Y.; Ohno, S.: The binding of androgen receptor to
DNA

[105843]and RNA. Biochim. Biophys. Acta 654: 181-186, 1981.

[105844]18565.

[105845]Aradhya, S.; Bardaro, T.; Galgoczy, P.; Yamagata, T.; Es-

posito,

[105846]T.; Patlan, H.; Ciccodicola, A.; Munnich, A.; Kenwrick, S.;

Platzer,

[105847]M.; D'Urso, M.; Nelson, D. L.: Multiple pathogenic and benign genomic

[105848]rearrangements occur at a 35 kb duplication involving the NEMO and

[105849]LAGE2 genes. Hum. Molec. Genet. 10: 2557-2567, 2001.

[105850]18566.

[105851]Aradhya, S.; Courtois, G.; Rajkovic, A.; Lewis, R. A.; Levy, M.;

[105852]Israel, A.; Nelson, D. L.: Atypical forms of incontinentia pigmenti

[105853]in male individuals result from mutations of a cytosine tract in exon

[105854]10 of NEMO (IKK-gamma). Am. J. Hum. Genet. 68: 765-771, 2001.

[105855]18567.

[105856]Aradhya, S.; Woffendin, H.; Jakins, T.; Bardaro, T.; Esposito,

[105857]T.; Smahi, A.; Shaw, C.; Levy, M.; Munnich, A.; D'Urso, M.; Lewis,

[105858]R. A.; Kenwrick, S.; Nelson, D. L.: A recurrent deletion in
the ubiquitously
[105859]expressed NEMO (IKK- γ) gene accounts for the vast
majority of
[105860]incontinentia pigmenti mutations. Hum. Molec. Genet. 10:
2171–2179,
[105861]2001.
[105862]18568.
[105863]Doffinger, R.; Smahi, A.; Bessia, C.; Geissmann, F.; Fein-
berg,
[105864]J.; Durandy, A.; Bodemer, C.; Kenwrick, S.; Dupuis-Girod,
S.; Blanche,
[105865]S.; Wood, P.; Rabia, S. H.; and 16 others: X-linked an-
hidrotic
[105866]ectodermal dysplasia with immunodeficiency is caused by
impaired NF- κ B
[105867]signaling. Nature Genet. 27: 277–285, 2001.
[105868]18569.
[105869]Jain, A.; Ma, C. A.; Liu, S.; Brown, M.; Cohen, J.; Strober, W.
[105870]: Specific missense mutations in NEMO result in hyper-IgM
syndrome

[105871]with hypohydrotic (sic) ectodermal dysplasia. Nature Immun. 2: 223–228,

[105872]2001.

[105873]18570.

[105874]Jin, D. Y.; Jeang, K. T.: Isolation of full-length cDNA and chromosomal

[105875]localization of human NF-kappaB modulator NEMO to Xq28. J. Biomed.

[105876]Sci. 6: 115–120, 1999.

[105877]18571.

[105878]Kosaki, K.; Shimasaki, N.; Fukushima, H.; Hara, M.; Ogata, T.;

[105879]Matsuo, N.: Female patient showing hypohidrotic ectodermal dysplasia

[105880]and immunodeficiency (HED-ID). (Letter) Am. J. Hum. Genet. 69: 664–665,

[105881]2001.

[105882]18572.

[105883]Li, Q.; Van Antwerp, D.; Mercurio, F.; Lee, K.-F.; Verma, I. M.

[105884]: Severe liver degeneration in mice lacking the I-kappa-B

kinase 2

[105885]gene. Science 284: 321–325, 1999.

[105886]18573.

[105887]Li, Y.; Kang, J.; Friedman, J.; Tarassishin, L.; Ye, J.; Kovalenko,

[105888]A.; Wallach, D.; Horwitz, M. S.: Identification of a cell protein

[105889](FIP-3) as a modulator of NF-kappa-B activity and as a target of an

[105890]adenovirus inhibitor of tumor necrosis factor alpha-induced apoptosis. Proc.

[105891]Nat. Acad. Sci. 96: 1042–1047, 1999.

[105892]18574.

[105893]Makris, C.; Godfrey, V. L.; Krahn-Senftleben, G.; Takahashi, T.;

[105894]Roberts, J. L.; Schwarz, T.; Feng, L.; Johnson, R. S.; Karin, M.:

[105895]Female mice heterozygous for IKK-gamma/NEMO deficiencies develop a

[105896]dermatopathy similar to the human X-linked disorder incontinentia

[105897]pigmenti. Molec. Cell 5: 969–979, 2000.

[105898]18575.

[105899]May, M. J.; D'Acquisto, F.; Madge, L. A.; Glockner, J.;
Poher,

[105900]J. S.; Ghosh, S.: Selective inhibition of NF-kappa-B activa-
tion by

[105901]a peptide that blocks the interaction of NEMO with the I-
kappa-B kinase

[105902]complex. Science 289: 1550-1554, 2000.

[105903]18576.

[105904]Roberts, J. L.; Morrow, B.; Vega-Rich, C.; Salafia, C. M.; Ni-
towsky,

[105905]H. M.: Incontinentia pigmenti in a newborn male infant
with DNA confirmation. Am.

[105906]J. Med. Genet. 75: 159-163, 1998.

[105907]18577.

[105908]Rothwarf, D. M.; Zandi, E.; Natoli, G.; Karin, M.: IKK-
gamma

[105909]is an essential regulatory subunit of the I-kappa-B kinase
complex. Nature 395:

[105910]297-300, 1998.

[105911]18578.

[105912]Rudolph, D.; Yeh, W.-C.; Wakeham, A.; Rudolph, B.; Nal-
lainathan,
[105913]D.; Potter, J.; Elia, A. J.; Mak, T. W.: Severe liver degenera-
tion
[105914]and lack of NF-kappa-B activation in NEMO/
IKK-gamma-deficient mice. Genes
[105915]Dev. 14: 854-862, 2000.
[105916]18579.
[105917]Schmidt-Supprian, M.; Bloch, W.; Courtois, G.; Addicks, K.;
Israel,
[105918]A.; Rajewsky, K.; Pasparakis, M.: NEMO/
IKK-gamma-deficient mice model
[105919]incontinentia pigmenti. Molec. Cell 5: 981-992, 2000.
[105920]18580.
[105921]The International Incontinentia Pigmenti Consortium: Ge-
nomic
[105922]rearrangement in NEMO impairs NF-kappa-B activation
and is a cause
[105923]of incontinentia pigmenti. Nature 405: 466-472, 2000.
[105924]18581.
[105925]The International IP Consortium: Survival of male patients

[105926]with incontinentia pigmenti carrying a lethal mutation can
be explained

[105927]by somatic mosaicism or Klinefelter syndrome. Am. J.
Hum. Genet. 69:

[105928]1210–1217, 2001.

[105929]18582.

[105930]Yamaoka, S.; Courtois, G.; Bessia, C.; Whiteside, S. T.;
Weil,

[105931]R.; Agou, F.; Kirk, H. E.; Kay, R. J.; Israel, A.: Complemen-
tation

[105932]cloning of NEMO, a component of the I-kappa-B kinase
complex essential

[105933]for NF-kappa-B activation. Cell 93: 1231–1240, 1998.

[105934]18583.

[105935]Zonana, J.; Elder, M. E.; Schneider, L. C.; Orlow, S. J.; Moss,

[105936]C.; Golabi, M.; Shapira, S. K.; Farndon, P. A.; Wara, D. W.;
Emmal,

[105937]S. A.; Ferguson, B. M.: A novel X-linked disorder of im-
mune deficiency

[105938]and hypohidrotic ectodermal dysplasia is allelic to inconti-
nencia

[105939] pigmenti and due to mutations in IKK-gamma (NEMO).

Am. J. Hum. Genet. 67:

[105940] 1555–1562, 2000.

[105941] 18584.

[105942] Bauer, M. F.; Gempel, K.; Reichert, A. S.; Rappold, G. A.;

Lichtner,

[105943] P.; Gerbitz, K.-D.; Neupert, W.; Brunner, M.; Hofmann, S.:

Genetic

[105944] and structural characterization of the human mitochondrial inner membrane

[105945] translocase. J. Molec. Biol. 289: 69–82, 1999.

[105946] 18585.

[105947] Kruse, T. A.; Kolvraa, S.; Bolund, L.; Kaitila, I.; Soderholm,

[105948] A. L.; de la Chapelle, A.; MacDermot, K. D.: X-linked anhidrotic

[105949] ectodermal dysplasia (EDA): multipoint linkage analysis.

(Abstract) Cytogenet.

[105950] Cell Genet. 51: 1026, 1989.

[105951] 18586.

[105952] Limon, J.; Filipiuk, J.; Nedoszytko, B.; Mrozek, K.; Castren,

[105953]M.; Larramendy, M.; Roszkiewicz, J.: X-linked anhidrotic
ectodermal
[105954]dysplasia and de novo t(X;1) in a female. Hum. Genet. 87:
338–340,
[105955]1991.
[105956]18587.
[105957]MacDermot, K. D.; Hulten, M.: Female with hypohidrotic
ectodermal
[105958]dysplasia and de novo (X;9) translocation: clinical docu-
mentation
[105959]of the AnLy cell line case. Hum. Genet. 84: 577–579,
1990.
[105960]18588.
[105961]MacDermot, K. D.; Malcolm, S.: X linked hypohidrotic ec-
todermal
[105962]dysplasia (CST syndrome): clinical and molecular study
using X chromosome
[105963]markers. (Abstract) J. Med. Genet. 23: 167, 1986.
[105964]18589.
[105965]MacDermot, K. D.; Morgan, S. H.; Cheshire, J. K.; Wilson, T.
M.;

- [105966] Kaitila, I.; Soderholm, L.; de la Chapelle, A.; Kolvraa, S.; Bolund, L.; Kruse, T. A.: X-linked anhidrotic ectodermal dysplasia (EDA): two point and multipoint linkage analysis using 7 polymorphic DNA markers. (Abstract) Cytogenet. Cell Genet. 46: 653-654, 1987.
- [105970] 18590.
- [105971] MacDermot, K. D.; Winter, R. M.; Malcolm, S.: Gene localization of X-linked hypohidrotic ectodermal dysplasia (C-S-T syndrome). Hum. Genet. 74: 172-173, 1986.
- [105974] 18591.
- [105975] Malagon, V.; Taveras, J. E.: Congenital anhidrotic ectodermal and mesodermal dysplasia. Arch. Derm. 74: 253-258, 1956.
- [105977] 18592.
- [105978] Monreal, A. W.; Zonana, J.; Ferguson, B.: Identification of a

- [105979] new splice form of the EDA1 gene permits detection of nearly all X-linked
- [105980] hypohidrotic ectodermal dysplasia mutations. *Am. J. Hum. Genet.* 63:
- [105981] 380–389, 1998.
- [105982] 18593.
- [105983] Pedrini, V.; Lennzi, L.; Zamtotti, V.: Isolation and identification
- [105984] of keratosulphate in urine of patients affected by Morquio–Ullrich
- [105985] disease. *Proc. Soc. Exp. Biol. Med.* 110: 847–849, 1962.
- [105986] 18594.
- [105987] Riedner, E. D.; Levin, L. S.: Hearing patterns in Morquio's syndrome
- [105988] (mucopolysaccharidosis IV). *Arch. Otolaryng.* 103: 518–520, 1977.
- [105989] 18595.
- [105990] Robins, M. M.; Stevens, H. F.; Linker, A.: Morquio's disease:
- [105991] an abnormality of mucopolysaccharide metabolism. *J. Pediat.* 62:

[105992]881–889, 1963.

[105993]18596.

[105994]Singh, J.; DiFerrante, N. M.; Niebes, P.; Tavella, D.: N-acetylgalactosamine-6-sulfate

[105995]sulfatase in man: absence of the enzyme in Morquio disease. J. Clin.

[105996]Invest. 57: 1036–1040, 1976.

[105997]18597.

[105998]Sukegawa, K.; Nakamura, H.; Kato, Z.; Tomatsu, S.; Montano, A.

[105999]M.; Fukao, T.; Toietta, G.; Tortora, P.; Orii, T.; Kondo, N.: Biochemical

[106000]and structural analysis of missense mutations in N-acetylgalactosamine-6-sulfate

[106001]sulfatase causing mucopolysaccharidosis IVA phenotypes. Hum. Molec.

[106002]Genet. 9: 1283–1290, 2000.

[106003]18598.

[106004]Tomatsu, S.; Fukuda, S.; Cooper, A.; Wraith, J. E.; Ferreira,

[106005]P.; Di Natale, P.; Tortora, P.; Fujimoto, A.; Kato, Z.; Ya-

mada, N.;

[106006]Isogai, K.; Yamagishi, A.; Sukegawa, K.; Suzuki, Y.; Shimozawa, N.;

[106007]Kondo, N.; Sly, W. S.; Orii, T.: Fourteen novel mucopolysaccharidosis

[106008]IVA producing mutations in GALNS gene. Hum. Mutat. 10: 368-375,

[106009]1997.

[106010]18599.

[106011]Tomatsu, S.; Fukuda, S.; Cooper, A.; Wraith, J. E.; Rezvi, G.

[106012]M. M.; Yamagishi, A.; Yamada, N.; Kato, Z.; Isogai, K.; Sukegawa,

[106013]K.; Kondo, N.; Suzuki, Y.; Shimozawa, N.; Orii, T.: Mucopolysaccharidosis

[106014]IVA: identification of a common missense mutation I113F in the N-acetylgalactosamine-6-sulfate

[106015]sulfatase gene. Am. J. Hum. Genet. 57: 556-563, 1995.

[106016]18600.

[106017]Tomatsu, S.; Fukuda, S.; Cooper, A.; Wraith, J. E.; Yamada, N.;

[106018]Isogai, K.; Kato, Z.; Sukegawa, K.; Kondo, N.; Suzuki, Y.;

Shimozawa,

[106019]N.; Orii, T.: Two new mutations, Q473X and N487S, in a
Caucasian

[106020]patient with mucopolysaccharidosis IVA (Morquio disease).
Hum. Mutat. 6:

[106021]195–196, 1995.

[106022]18601.

[106023]Tomatsu, S.; Fukuda, S.; Masue, M.; Sukegawa, K.; Fukao,
T.; Yamagishi,

[106024]A.; Hori, T.; Iwata, H.; Ogawa, T.; Nakashima, Y.; Hanyu,
Y.; Hashimoto,

[106025]T.; Titani, K.; Oyama, R.; Suzuki, M.; Yagi, K.; Hayashi, Y.;
Orii,

[106026]T.: Morquio disease: isolation, characterization and ex-
pression of

[106027]full-length cDNA for human N-acetyl-
galactosamine-6-sulfate sulfatase. Biochem.

[106028]Biophys. Res. Commun. 181: 677–683, 1991.

[106029]18602.

[106030]Tomatsu, S.; Fukuda, S.; Masue, M.; Sukegawa, K.; Ma-
suno, M.;

[106031]Orii, T.: Mucopolysaccharidosis type IVA: characterization

and chromosomal

[106032] localization of N-acetylgalactosamine-6-sulfate sulfatase gene and

[106033] genetic heterogeneity. (Abstract) Am. J. Hum. Genet. 51 (suppl.):

[106034] A178, 1992.

[106035] 18603.

[106036] Tomatsu, S.; Fukuda, S.; Yamagishi, A.; Cooper, A.; Wraith, J.

[106037] E.; Hori, T.; Kato, Z.; Yamada, N.; Isogai, K.; Sukegawa, K.; Kondo,

[106038] N.; Suzuki, Y.; Shimozawa, N.; Orii, T.: Mucopolysacchari-
dosis IVA:

[106039] four new exonic mutations in patients with N-
acetylgalactosamine-6-sulfate

[106040] sulfatase deficiency. Am. J. Hum. Genet. 58: 950-962,
1996.

[106041] 18604.

[106042] Tyłki-Szymanska, A.; Czartoryska, B.; Bunge, S.; van
Diggelen,

[106043] O. P.; Kleijer, W. J.; Poorthuis, B. J. H. M.; Huijmans, J. G.
M.;

[106044]Gorska, D.: Clinical, biochemical and molecular findings in
a two-generation

[106045]Morquio A family. Clin. Genet. 53: 369–374, 1998.

[106046]18605.

[106047]Von Noorden, G. K.; Zellweger, H.; Ponseti, I. V.: Ocular
findings

[106048]in Morquio–Ullrich's disease. Arch. Ophthal. 64: 585–591,
1960.

[106049]18606.

[106050]Wang, L.; Ou, X.; Sebesta, I.; Vondrak, K.; Krijt, J.; Elleder,

[106051]M.; Poupetova, H.; Ledvinova, J.; Zeman, J.; Simmonds, H.
A.; Tischfield,

[106052]J. A.; Sahota, A.: Combined adenine phosphoribosyltrans-
ferase and

[106053]N-acetylgalactosamine-6-sulfate sulfatase deficiency.
Molec. Genet.

[106054]Metab. 68: 78–85, 1999.

[106055]18607.

[106056]Wiedemann, H.–R.: The pioneers of pediatric medicine:
Luis Morquio

[106057](1867–1935). Europ. J. Pediat. 151: 549, 1992.

[106058]18608.

[106059]Yamada, N.; Fukuda, S.; Tomatsu, S.; Muller, V.; Hopwood, J. J.;

[106060]Nelson, J.; Kato, Z.; Yamagishi, A.; Sukegawa, K.; Kondo, N.; Orii,

[106061]T.: Molecular heterogeneity in mucopolysaccharidosis IVA in Australia

[106062]and Northern Ireland: nine novel mutations including T312S, a common

[106063]allele that confers a mild phenotype. Hum. Mutat. 11: 202-208, 1998.

[106064]18609.

[106065]Yuen, M.; Fensom, A. H.: Diagnosis of classical Morquio's disease:

[106066]N-acetylgalactosamine 6-sulphate sulphatase activity in cultured fibroblasts,

[106067]leukocytes, amniotic cells and chorionic villi. J. Inherit. Metab.

[106068]Dis. 8: 80-86, 1985.

[106069]18610.

[106070]Zellweger, H.; Ponseti, I. V.; Pedrini, V.; Stamler, F. S.; Von

[106071]Noorden, G. K.: Morquio-Ullrich's disease: report of 2 cases. J.

[106072]Pediat. 59: 549-561, 1961.

[106073]18611.

[106074]Shmerling, D. H.; Prader, A.; Hitzig, W. H.; Giedion, A.; Hadorn,

[106075]B.; Kuhni, M.: The syndrome of exocrine pancreatic insufficiency,

[106076]neutropenia, metaphyseal dysostosis and dwarfism. Helv. Paediat.

[106077]Acta 24: 547-575, 1969.

[106078]18612.

[106079]Moller, D. E.; Xia, C. H.; Tang, W.; Zhu, A. X.; Jakubowski, M.

[106080]; Human rsk isoforms: cloning and characterization of tissue-specific

[106081]expression. Am. J. Physiol. 266: C351-C359, 1994.

[106082]18613.

[106083]Trivier, E.; De Cesare, D.; Jacquot, S.; Pannetier, S.; Zackai,

[106084]E.; Young, I.; Mandel, J.-L.; Sassone-Corsi, P.; Hanauer, A.: Mutations

[106085]in the kinase Rsk-2 associated with Coffin-Lowry syndrome. Nature 384:

[106086]567-570, 1996.

[106087]18614.

[106088]Zhao, Y.; Bjorbaek, C.; Weremowicz, S.; Morton, C. C.; Moller,

[106089]D. E.: RSK3 encodes a novel pp90rsk isoform with a unique N-terminal

[106090]sequence: growth factor-stimulated kinase function and nuclear translocation. Molec.

[106091]Cell. Biol. 15: 4353-4363, 1995.

[106092]18615.

[106093]Liu, Z.; Sun, C.; Olejniczak, E. T.; Meadows, R. P.; Betz, S. F.;

[106094]Oost, T.; Herrmann, J.; Wu, J. C.; Fesik, S. W.: Structural basis

[106095]for binding of Smac/DIABLO to the XIAP BIR3 domain. Nature 408:

[106096]1004-1008, 2000.

[106097]18616.

[106098]Srinivasula, S. M.; Hegde, R.; Saleh, A.; Datta, P.; Shiozaki,

[106099]E.; Chai, J.; Lee, R.-A.; Robbins, P. D.; Fernandes-Alnemri, T.; Shi, Y.; Alnemri, E. S.: A conserved XIAP-interaction motif in caspase-9 and Smac/DIABLO regulates caspase activity and apoptosis. *Nature* 410: 112-116, 2001.

[106102]18617.

[106103]Shwachman, H.; Diamond, L. K.; Oski, F. A.; Khaw, K. T.: The syndrome of pancreatic insufficiency and bone marrow dysfunction. *J. Pediat.* 65: 645-663, 1964.

[106106]18618.

[106107]Shwachman, H.; Holsclaw, D.: Some clinical observations on the Shwachman syndrome (pancreatic insufficiency and bone marrow hypoplasia). *Birth Defects Orig. Art. Ser.* 8(3): 46-49, 1972.

[106110]18619.

[106111]Smith, O. P.; Chan, M. Y.; Evans, J.; Veys, P.: Shwachman-Di-

among

[106113] syndrome and matched unrelated donor BMT. Bone Marrow Transplant. 16:

[106114] 717–718, 1995.

[106115] 18620.

[106116] Smith, O. P.; Hann, I. M.; Chessells, J. M.; Reeves, B. R.; Milla,

[106117] P.: Haematological abnormalities in Shwachman–Diamond syndrome. Brit.

[106118] J. Haemat. 94: 279–284, 1996.

[106119] 18621.

[106120] Tada, H.; Ri, T.; Yoshida, H.; Ishimoto, K.; Kaneko, M.; Yamashiro,

[106121] Y.; Shinohara, T.: A case of Shwachman syndrome with increased spontaneous

[106122] chromosome breakage. Hum. Genet. 77: 289–291, 1987.

[106123] 18622.

[106124] Taybi, H.; Mitchell, A. D.; Friedman, G. D.: Metaphyseal dysostosis

[106125] and associated syndrome of pancreatic insufficiency and blood disorders. Radiology 93:

[106126]563–571, 1969.

[106127]18623.

[106128]Woods, W. G.; Roloff, J. S.; Lukens, J. N.; Krivit, W.: The occurrence

[106129]of leukemia in patients with the Shwachman syndrome. J. Pediat. 99:

[106130]425–428, 1981.

[106131]18624.

[106132]Wulfeck, B.; Trauner, D.; Marsden, D.; Nyhan, W.; Tallal, P.:

[106133]Neurobehavioral profiles in two sisters with Shwachman–Diamond syndrome. Dysmorph.

[106134]Clin. Genet. 5: 15–22, 1991.

[106135]18625.

[106136]Britton, C. H.; Schultz, R. A.; Zhang, B.; Esser, V.; Foster, D.

[106137]W.; McGarry, J. D.: Human liver mitochondrial carnitine palmitoyltransferase

[106138]: characterization of its cDNA and chromosomal localization and partial

[106139]analysis of the gene. Proc. Nat. Acad. Sci. 92: 1984–1988, 1995.

[106140]18626.

[106141]Song, X.-Q.; Fukao, T.; Watanabe, H.; Shintaku, H.; Hirayama,

[106142]K.; Kassovska-Bratinova, S.; Kondo, N.; Mitchell, G. A.: Succinyl-CoA:3-ketoacid

[106143]CoA transferase (SCOT) deficiency: two pathogenic mutations, V133E

[106144]and C456F, in Japanese siblings. Hum. Mutat. 12: 83-88, 1998.

[106145]18627.

[106146]Spence, M. W.; Murphy, M. G.; Cook, H. W.; Ripley, B. A.; Embil,

[106147]J. A.: Succinyl CoA:3-ketoacid CoA transferase deficiency: a 'new'

[106148]phenotype? (Abstract) Pediat. Res. 7: 394 only, 1973.

[106149]18628.

[106150]Tildon, J. T.; Cornblath, M.: Succinyl-CoA: 3-ketoacid CoA-transferase

[106151]deficiency. A cause for ketoacidosis in infancy. J. Clin. Invest. 51:

[106152]493-498, 1972.

[106153]18629.

[106154]Cannizzaro, L. A.; Chen, Y. Q.; Rafi, M. A.; Wenger, D. A.:
Regional

[106155]mapping of the human galactocerebrosidase gene (GALC)
to 14q31 by

[106156]in situ hybridization. Cytogenet. Cell Genet. 66: 244–245,
1994.

[106157]18630.

[106158]De Gasperi, R.; Sosa, M. A. G.; Sartorato, E. L.; Battistini,
S.;

[106159]MacFarlane, H.; Gusella, J. F.; Krivit, W.; Kolodny, E. H.:
Molecular

[106160]heterogeneity of late-onset forms of globoid-cell
leukodystrophy. Am.

[106161]J. Hum. Genet. 59: 1233–1242, 1996.

[106162]18631.

[106163]Duchen, L. W.; Eicher, E. M.; Jacobs, J. M.; Scaravilli, F.;
Teixeira,

[106164]F.: Hereditary leucodystrophy in the mouse: the new mu-
tant twitcher. Brain 103:

[106165]695–710, 1980.

[106166]18632.

[106167]Trauner, D. A.; Page, T.; Greco, C.; Sweetman, L.; Kulovich, S.;

[106168]Nyhan, W. L.: Progressive neurodegenerative disorder in a patient

[106169]with nonketotic hyperglycinemia. J. Pediat. 98: 272–275, 1981.

[106170]18633.

[106171]Minassian, B. A.; Lee, J. R.; Herbrick, J.–A.; Huizenga, J.; Soder,

[106172]S.; Mungall, A. J.; Dunham, I.; Gardner, R.; Fong, C. G.; Carpenter,

[106173]S.; Jardim, L.; Satishchandra, P.; Andermann, E.; Snead, O. C., III;

[106174]Lopes–Cendes, I.; Tsui, L.–C.; Delgado–Escueta, A. V.; Rouleau, G.

[106175]A.; Scherer, S. W.: Mutations in a gene encoding a novel protein

[106176]tyrosine phosphatase cause progressive myoclonus epilepsy. Nature

[106177]Genet. 20: 171–174, 1998.

[106178]18634.

- [106179]Berger, R.; Smit, G. P. A.; Schierbeek, H.; Bijsterveld, K.; le
[106180]Coutre, R.: Mevalonic aciduria: an inborn error of cholesterol biosynthesis? Clin.
[106181]Chim. Acta 152: 219–222, 1985.
[106182]18635.
- [106183]Cuisset, L.; Drenth, J. P. H.; Simon, A.; Vincent, M. F.; van
der
[106184]Velde Visser, S.; van der Meer, J. W. M.; Grateau, G.;
Delpech, M.;
[106185]International Hyper-IgD Study Group: Molecular analysis
of MVK
[106186]mutations and enzymatic activity in hyper-IgD and periodic fever syndrome. Europ.
[106187]J. Hum. Genet. 9: 260–266, 2001.
[106188]18636.
- [106189]Drenth, J. P. H.; Cuisset, L.; Grateau, G.; Vasseur, C.; van
de
[106190]Velde-Visser, S. D.; de Jong, J. G. N.; Beckmann, J. S.; van
der Meer,
[106191]J. W. M.; Delpech, M.; International Hyper-IgD Study
Group: Mutations

[106192]in the gene encoding mevalonate kinase cause hyper-IgD
and periodic

[106193]fever syndrome. *Nature Genet.* 22: 178–181, 1999.

[106194]18637.

[106195]Drenth, J. P. H.; Haagsma, C. J.; van der Meer, J. W. M.; In-
ternational

[106196]Hyper-IgD Study Group: Hyperimmunoglobulinemia D and
periodic fever

[106197]syndrome: the clinical spectrum in a series of 50 patients.
Medicine 73:

[106198]133–144, 1994.

[106199]18638.

[106200]Gibson, K. M.; Hoffmann, G.; Nyhan, W. L.; Sweetman, L.;
Berger,

[106201]R.; le Coultre, R.; Smit, G. P. A.: Mevalonate kinase defi-
ciency

[106202]in a child with cerebellar ataxia hypotonia and mevalonic
aciduria. *Europ.*

[106203]*J. Pediat.* 148: 250–252, 1988.

[106204]18639.

[106205]Gibson, K. M.; Hoffmann, G.; Nyhan, W. L.; Sweetman, L.;

Brandt,

[106206]J. K.; Wappner, R. S.; Bader, P. I.: Mevalonic aciduria: family studies

[106207]in mevalonate kinase deficiency, an inborn error of cholesterol biosynthesis. J.

[106208]Inherit. Metab. Dis. 10 (suppl. 2): 282–285, 1987.

[106209]18640.

[106210]Gibson, K. M.; Hoffmann, G. F.; Tanaka, R. D.; Bishop, R. W.; Chambliss,

[106211]K. L.: Mevalonate kinase map position 12q24. Chromosome Res. 5:

[106212]150 only, 1997.

[106213]18641.

[106214]Hinson, D. D.; Rogers, Z. R.; Hoffmann, G. F.; Schachtele, M.;

[106215]Fingerhut, R.; Kohlschutter, A.; Kelley, R. I.; Gibson, K. M.: Hematological

[106216]abnormalities and cholestatic liver disease in two patients with mevalonate

[106217]kinase deficiency. Am. J. Med. Genet. 78: 408–412, 1998.

[106218]18642.

- [106219]Hinson, D. D.; Ross, R. M.; Krisans, S.; Shaw, J. L.; Kozich, V.;
- [106220]Rolland, M.-O.; Divry, P.; Mancini, J.; Hoffmann, G. F.; Gibson, K.
- [106221]M.: Identification of a mutation cluster in mevalonate kinase deficiency,
- [106222]including a new mutation in a patient of Mennonite ancestry. Am.
- [106223]J. Hum. Genet. 65: 327-335, 1999.
- [106224]18643.
- [106225]Hoffmann, G.; Gibson, K. M.; Brandt, I. K.; Bader, P. I.; Wappner,
- [106226]R. S.; Sweetman, L.: Mevalonic aciduria--an inborn error of cholesterol
- [106227]and nonsterol isoprene biosynthesis. New Eng. J. Med. 314: 1610-1614,
- [106228]1986.
- [106229]18644.
- [106230]Hoffmann, G. F.; Charpentier, C.; Mayatepek, E.; Mancini, J.;
- [106231]Leichsenring, M.; Gibson, K. M.; Divry, P.; Hrebicek, M.; Lehnert,

[106232]W.; Sartor, K.; Trefz, F. K.; Rating, D.; Bremer, H. J.; Nyhan, W.

[106233]L.: Clinical and biochemical phenotype in 11 patients with mevalonic

[106234]aciduria. *Pediatrics* 91: 915–921, 1993.

[106235]18645.

[106236]Houten, S. M; Koster, J.; Romeijn, G.-J.; Frenkel, J.; Di Rocco,

[106237]M.; Caruso, U.; Landrieu, P.; Kelley, R. I.; Kuis, W.; Poll-The, B.

[106238]T.; Gibson, K. M.; Wanders, R. J. A.; Waterham, H. R.: Organization

[106239]of the mevalonate kinase (MVK) gene and identification of novel mutations

[106240]causing mevalonic aciduria and hyperimmunoglobulinemia D and periodic

[106241]fever syndrome. *Europ. J. Hum. Genet.* 9: 253–259, 2001.
Note: Erratum:

[106242]*Europ. J. Med. Genet.* 9: 651 only, 2001.

[106243]18646.

[106244]Krisans, S. K.; Ericsson, J.; Edwards, P. A.; Keller, G. A.:

[106245]Farnesyl-diphosphate synthase is localized in peroxisomes. J. Biol.

[106246]Chem. 269: 14165-14169, 1994.

[106247]18647.

[106248]Mancini, J.; Philip, N.; Chabrol, B.; Divry, P.; Rolland, M.-O.;

[106249]Pinsard, N.: Mevalonic aciduria in 3 siblings: a new recognizable

[106250]metabolic encephalopathy. Pediat. Neurol. 9: 243-246, 1993.

[106251]18648.

[106252]Schafer, B. L.; Bishop, R. W.; Kratunis, V. J.; Kalinowski, S.

[106253]S.; Mosley, S. T.; Gibson, K. M.; Tanaka, R. D.: Molecular cloning

[106254]of human mevalonate kinase and identification of a missense mutation

[106255]in the genetic disease mevalonic aciduria. J. Biol. Chem. 267: 13229-13238,

[106256]1992.

[106257]18649.

[106258]van der Meer, J. W. M.; Vossen, J. M.; Radl, J.; van

Nieuwkoop,

[106259]J. A.; Meyer, C. J. L. M.; Lobatto, S.; van Furth, R.: Hyper-immunoglobulinaemia

[106260]D and periodic fever: a new syndrome. Lancet I: 1087-1090, 1984.

[106261]18650.

[106262]Houten, S. M.; Kuis, W.; Duran, M.; de Koning, T. J.; van Royen-Kerkhof,

[106263]A.; Romeijn, G. J.; Frenkel, J.; Dorland, L.; de Barse, M. M. J.;

[106264]Huijbers, W. A. R.; Rijkers, G. T.; Waterham, H. R.; Wanders, R. J.

[106265]A.; Poll-The, B. T.: Mutations in MVK, encoding mevalonate kinase,

[106266]cause hyperimmunoglobulinaemia D and periodic fever syndrome. Nature

[106267]Genet. 22: 175-177, 1999.

[106268]18651.

[106269]Houten, S. M.; Romeijn, G. J.; Koster, J.; Gray, R. G. F.; Darbyshire,

[106270]P.; Smit, G. P. A.; de Klerk, J. B. C.; Duran, M.; Gibson, K. M.;

- [106271]Wanders, R. J. A.; Waterham, H. R.: Identification and characterization
[106272]of three novel missense mutations in mevalonate kinase
cDNA causing
[106273]mevalonic aciduria, a disorder of isoprene biosynthesis.
Hum. Molec.
[106274]Genet. 8: 1523–1528, 1999.
[106275]18652.
- [106276]Norio, R.; Koskiniemi, M.: Progressive myoclonus epilepsy:
genetic
[106277]and nosological aspects with special reference to 107
Finnish patients. Clin.
[106278]Genet. 15: 382–398, 1979.
[106279]18653.
- [106280]Ortiz-Hidalgo, C.: The man behind Lafora's bodies. Am. J.
Surg.
[106281]Path. 10: 358–361, 1986.
[106282]18654.
- [106283]Sainz, J.; Minassian, B. A.; Serratosa, J. M.; Gee, M. N.;
Sakamoto,
[106284]L. M.; Iranmanesh, R.; Bohlega, S.; Baumann, R. J.; Ryan,

S.; Sparkes,

[106285]R. S.; Delgado-Escueta, A. V.: Lafora progressive myoclonus epilepsy:

[106286]narrowing the chromosome 6q24 locus by recombinations and homozygosities.

[106287](Letter) Am. J. Hum. Genet. 61: 1205-1209, 1997.

[106288]18655.

[106289]Sarlin, M. B.; Kloepfer, H. W.; Mickle, W. A.; Heath, R. G.:

[106290]The detection of carriers in hereditary myoclonic epilepsy.
Acta

[106291]Genet. Med. Gemellol. 9: 466-471, 1960.

[106292]18656.

[106293]Schwarz, G. A.; Yanoff, M.: Lafora's disease, distinct clinico-pathologic

[106294]form of Unverricht's syndrome. Arch. Neurol. 12:
172-188, 1965.

[106295]18657.

[106296]Serratos, J. M.; Delgado-Escueta, A. V.; Posada, I.; Shih, S.;

[106297]Drury, I.; Berciano, J.; Zabala, J. A.; Antunez, M. C.;
Sparkes, R.

- [106298]S.: The gene for progressive myoclonus epilepsy of the Lafora type maps to chromosome 6q. Hum. Molec. Genet. 4: 1657–1663, 1995.
- [106300]18658.
- [106301]Serratos, J. M.; Gomez–Garre, P.; Gallardo, M. E.; Anta, B.; Beltran–Valero de Bernabe, D.; Lindhout, D.; Augustijn, P. B.; Tassinari, C. A.; Michelucci, R.; Malafosse, A.; Topcu, M.; Grid, D.; Dravet, C.; Berkovic, S. F.; Rodriguez de Cordoba, S.: A novel protein tyrosine phosphatase gene is mutated in progressive myoclonus epilepsy of the Lafora type (EPM2). Hum. Molec. Genet. 8: 345–352, 1999.
- [106307]18659.
- [106308]Yanoff, M.; Schwarz, G. A.: Lafora's disease: a distinct genetically determined form of Unverricht's syndrome. Genet. Hum. 14: 235–244,
- [106310]1965.

[106311]18660.

[106312]Yokoi, S.; Austin, J.; Witmer, F.; Sakai, M.: Studies in myoclonus

[106313]epilepsy (Lafora body forms). I. Isolation and preliminary characterization

[106314]of Lafora bodies in two cases. Arch. Neurol. 19: 15–33, 1968.

[106315]18661.

[106316]Mowat, D. R.; Croaker, G. D. H.; Cass, D. T.; Kerr, B. A.; Chaitow,

[106317]J.; Ades, L. C.; Chia, N. L.; Wilson, M. J.: Hirschsprung disease,

[106318]microcephaly, mental retardation, and characteristic facial features:

[106319]delineation of a new syndrome and identification of a locus at chromosome

[106320]2q22–q23. J. Med. Genet. 35: 617–623, 1998.

[106321]18662.

[106322]Wakamatsu, N.; Yamada, Y.; Yamada, K.; Ono, T.; Nomura, N.; Taniguchi,

[106323]H.; Kitoh, H.; Mutoh, N.; Yamanaka, T.; Mushiaki, K.; Kato, K.; Sonta,

[106324]S.; Nagaya, M.: Mutations in SIP1, encoding Smad interacting protein-1,
[106325]cause a form of Hirschsprung disease. Nature Genet. 27:
369-370,
[106326]2001.
[106327]18663.
[106328]Zweier, C.; Albrecht, B.; Mitulla, B.; Behrens, R.; Beese, M.;
[106329]Gillesen-Kaesbach, G.; Rott, H.-D.; Rauch, A.: 'Mowat-
Wilson' syndrome
[106330]with and without Hirschsprung disease is a distinct, recognizable
[106331]multiple congenital anomalies-mental retardation syndrome caused by
[106332]mutations in the zinc finger homeo box 1B gene. Am. J. Med. Genet. 108:
177-181, 2002.
[106333]18664.
[106334]18664.
[106335]Yamada, K.; Yamada, Y.; Nomura, N.; Miura, K.; Wakako, R.; Hayakawa,
[106336]C.; Matsumoto, A.; Kumagai, T.; Yoshimura, I.; Miyazaki, S.; Kato,

[106337]K.; Sonta, S.; Ono, H.; Yamanaka, T.; Nagaya, M.; Wakamatsu, N.:

[106338]Nonsense and frameshift mutations in ZFX1B, encoding Smad-interacting

[106339]protein 1, cause a complex developmental disorder with a great variety

[106340]of clinical features. Am. J. Hum. Genet. 69: 1178–1185, 2001.

[106341]18665.

[106342]Alm, J.; Holmgren, G.; Larsson, A.; Schimpfessel, L.: Histidinaemia

[106343]in Sweden: report on a neonatal screening programme. Clin. Genet. 20:

[106344]229–233, 1981.

[106345]18666.

[106346]Anakura, M.; Matsuda, I.; Arashima, S.; Fukushima, N.; Oka, Y.

[106347]: Histidinemia--classical and atypical form in siblings. Am. J. Dis.

[106348]Child. 129: 858–861, 1975.

[106349]18667.

- [106350]Auerbach, V. H.; DiGeorge, A. M.; Baldrige, R. C.;
Tourtellotte,
[106351]C. D.; Brigham, M. P.: Histidinemia: a deficiency in histi-
dase resulting
[106352]in the urinary excretion of histidine and of imidazolepyru-
vic acid. J.
[106353]Pediat. 60: 487-497, 1962.
[106354]18668.
- [106355]Bruckman, C.; Berry, H. K.; Dasenbrock, R. J.: Histidinemia
in
[106356]two successive generations. Am. J. Dis. Child. 119:
221-227, 1970.
[106357]18669.
- [106358]Coulombe, J. T.; Kammerer, B. L.; Levy, H. L.; Hirsch, B. Z.;
Sriver,
[106359]C. R.: Histidinaemia. Part III: Impact; a prospective study. J.
[106360]Inherit. Metab. Dis. 6: 58-61, 1983.
[106361]18670.
- [106362]Kacser, H. K.; Bulfield, G.; Wallace, M. E.: Histidinaemia
mutant
[106363]in the mouse. Nature 244: 77-79, 1973.

[106364]18671.

[106365]Kappelman, M.; Thomas, G. H.; Howell, R. R.: Histidinemia
in a

[106366]Negro child. Am. J. Dis. Child. 122: 212–214, 1971.

[106367]18672.

[106368]La Du, B. N.: Histidinemia: current status. Am. J. Dis.
Child. 113:

[106369]88–92, 1967.

[106370]18673.

[106371]La Du, B. N.: Histidinemia.In: Stanbury, J. B.; Wyngaarden,
J.

[106372]B.; Fredrickson, D. S.: Metabolic Basis of Inherited Disease.
New

[106373]York: McGraw–Hill (pub.) (4th ed.): 1978. Pp. 317–327.

[106374]18674.

[106375]La Du, B. N.; Howell, R. R.; Jacoby, G. A.; Seegmiller, J. E.;

[106376]Sober, E. K.; Zannoni, V. G.; Canby, J. P.; Ziegler, L. K.:
Clinical

[106377]and biochemical studies on two cases of histidinemia. Pe-
diatrics 32:

[106378]216–227, 1963.

[106379]18675.

[106380]Lam, W. K.; Cleary, M. A.; Wraith, J. E.; Walter, J. H.: Histid-
inaemia:

[106381]a benign metabolic disorder. Arch. Dis. Child. 74:
343–346, 1996.

[106382]18676.

[106383]Lemieux, B.; Auray-Blais, C.; Giguere, R.; Shapcott, D.;
Scriver,

[106384]C. R.: Newborn urine screening experience with over one
million infants

[106385]in the Quebec network of genetic medicine. J. Inherit.
Metab. Dis. 11:

[106386]45–55, 1988.

[106387]18677.

[106388]Levy, H. L.; Shin, V. E.; Madigan, P. M.: Routine newborn
screening

[106389]for histidinemia: clinical and biochemical results. New
Eng. J. Med. 291:

[106390]1214–1219, 1974.

[106391]18678.

[106392]Neville, B. G. R.; Bentovim, A.; Clayton, B. E.; Sheperd, J.:

[106393]Histidinaemia: study of relation between clinical and biological findings

[106394]in 7 subjects. Arch. Dis. Child. 47: 190-200, 1972.

[106395]18679.

[106396]Neville, B. G. R.; Harris, R. F.; Stern, D. J.; Stern, J.: Maternal

[106397]histidinaemia. Arch. Dis. Child. 46: 119-121, 1971.

[106398]18680.

[106399]Popkin, J. S.; Clow, C. L.; Scriver, C. R.; Grove, J.: Is hereditary

[106400]histidinemia harmful? Lancet I: 721-722, 1974.

[106401]18681.

[106402]Rosenblatt, D. S.; Mohyuddin, F.; Scriver, C. R.: Histidinemia

[106403]discovered by urine screening after renal transplantation. Pediatrics 46:

[106404]47-53, 1970.

[106405]18682.

[106406]Barton, R. W.; Neufeld, E. F.: A distinct biochemical deficit

- [106407]in the Maroteaux–Lamy syndrome (mucopolysaccharidosis VI). J. Pediat. 80: 114–116, 1972.
- [106408]18683.
- [106409]Hechtman, P.; Boulay, B.; De Braekeleer, M.; Andermann, E.; Melancon, S.; Larochelle, J.; Prevost, C.; Kaplan, F.: The intron 7 donor splice site transition: a second Tay–Sachs disease mutation in French Canada. Hum. Genet. 90: 402–406, 1992.
- [106410]18684.
- [106411]Hechtman, P.; Kaplan, F.; Bayleran, J.; Boulay, B.; Andermann, E.; de Braekeleer, M.; Melancon, S.; Lambert, M.; Potier, M.; Gagne, R.; Kolodny, E.; Clow, C.; Capua, A.; Prevost, C.; Sriver, C.: More than one mutant allele causes infantile Tay–Sachs disease in French–Canadians. Am. J. Hum. Genet. 47: 815–822, 1990.

[106420]18685.

[106421]Hellkuhl, B.; Mayr, W. R.; Grzeschik, K.-H.: Localization of

[106422]MPI, PK-M2, IDH-M, and the alpha subunit of hex-
osaminidase (HEX-A)

[106423]to the q21-qter region of human chromosome 15. Cyto-
genet. Cell Genet. 22:

[106424]503-505, 1978.

[106425]18686.

[106426]Hou, Y.; Vavougios, G.; Hinek, A.; Wu, K. K.; Hechtman, P.;
Kaplan,

[106427]F.; Mahuran, D. J.: The val192-to-leu mutation in the al-
pha-subunit

[106428]of beta-hexosaminidase A is not associated with the
B1-variant form

[106429]of Tay-Sachs disease. Am. J. Hum. Genet. 59: 52-58,
1996.

[106430]18687.

[106431]Kappler, J.; Gieselmann, V.; Propping, P.: Hexosaminidase-
-pseudodeficiency?.

[106432](Letter) Am. J. Hum. Genet. 47: 880-881, 1990.

[106433]18688.

[106434]Keats, B. J. B.; Elston, R. C.; Andermann, E.: Pedigree discriminant

[106435]analysis of two French Canadian Tay-Sachs families.
Genet. Epidemiol. 4:

[106436]77-85, 1987.

[106437]18689.

[106438]Kelly, T. E.; Chase, G. A.; Kaback, M. M.; Kumor, K.; McKusick,

[106439]V. A.: Tay-Sachs disease: high gene frequency in a non-Jewish population. Am.

[106440]J. Hum. Genet. 27: 287-291, 1975.

[106441]18690.

[106442]Kelly, T. E.; Reynolds, L. W.; O'Brien, J. S.: Segregation within

[106443]a family to two mutant alleles for hexosaminidase A. Clin.
Genet. 9:

[106444]540-543, 1976.

[106445]18691.

[106446]Lalley, P. A.; Rattazzi, M. C.; Shows, T. B.: Human beta-D-N-acetylhexosaminidases

[106447]A and B: expression and linkage relationships in somatic

cell hybrids. Proc.

[106448]Nat. Acad. Sci. 71: 1569–1573, 1975.

[106449]18692.

[106450]Korneluk, R. G.; Mahuran, D. J.; Neote, K.; Klavins, M. H.;
O'Dowd,

[106451]B. F.; Tropak, M.; Willard, H. F.; Anderson, M.-J.; Lowden,
J. A.;

[106452]Gravel, R. A.: Isolation of cDNA clones coding for the al-
pha-subunit

[106453]of human beta-hexosaminidase: extensive homology be-
tween the alpha-

[106454]and beta-subunits and studies on Tay-Sachs disease. J.
Biol. Chem. 261:

[106455]8407–8413, 1986.

[106456]18693.

[106457]Landels, E. C.; Green, P. M.; Ellis, I. H.; Fensom, A. H.; Bo-
brow,

[106458]M.: Beta-hexosaminidase splice site mutation has a high
frequency

[106459]among non-Jewish Tay-Sachs disease carriers from the
British Isles. J.

[106460]Med. Genet. 29: 563–567, 1992.

[106461]18694.

[106462]Landels, E. C.; Green, P. M.; Ellis, I. H.; Fensom, A. H.;
Kaback,

[106463]M. M.; Lim-Steele, J.; Zeiger, K.; Levy, N.; Bobrow, M.: Fur-
ther

[106464]investigation of the HEXA gene intron 9 donor splice site
mutation

[106465]frequently found in non-Jewish Tay-Sachs disease pa-
tients from the

[106466]British Isles. J. Med. Genet. 30: 479-481, 1993.

[106467]18695.

[106468]Lau, M. M. H.; Neufeld, E. F.: A frameshift mutation in a
patient

[106469]with Tay-Sachs disease causes premature termination and
defective

[106470]intracellular transport of the alpha-subunit of beta-
hexosaminidase. J.

[106471]Biol. Chem. 264: 21376-21380, 1989.

[106472]18696.

[106473]McDowell, G. A.; Mules, E. H.; Fabacher, P.; Shapira, E.;
Blitzer,

- [106474]M. G.: The presence of two different infantile Tay-Sachs disease mutations in a Cajun population. Am. J. Hum. Genet. 51: 1071-1077, 1992.
- [106476]18697.
- [106477]18697.
- [106478]Lyerla, T. A.; Konola, J. T.; Skiba, M. C.; Raghavan, S.: Galactocerebrosidase activity in somatic cell hybrids derived from twitcher mouse/control human fibroblasts is associated with human chromosome 17. Am. J. Hum. Genet. 44: 198-207, 1989.
- [106480]18698.
- [106481]18698.
- [106482]Sakai, N.; Inui, K.; Fujii, N.; Fukushima, H.; Nishimoto, J.; Yanagihara, I.; Isegawa, Y.; Iwamatsu, A.; Okada, S.: Krabbe disease: isolation and characterization of a full-length cDNA for human galactocerebrosidase. Biochem. Biophys. Res. Commun. 198: 485-491, 1994.
- [106486]18699.

[106488]Sweet, H.: Twitcher (twi) is on chromosome 12. Mouse
Newsletter 75:

[106489]30, 1986.

[106490]18700.

[106491]Black, S. H.; Pelias, M. Z.; Miller, J. B.; Blitzer, M. G.;
Shapira,

[106492]E.: Maroteaux-Lamy syndrome in a large consanguineous
kindred: biochemical

[106493]and immunological studies. Am. J. Med. Genet. 25:
273-279, 1986.

[106494]18701.

[106495]Brooks, D. A.; McCourt, P. A. G.; Gibson, G. J.; Ashton, L.
J.;

[106496]Shutter, M.; Hopwood, J. J.: Analysis of N-
acetylgalactosamine-4-sulfatase

[106497]protein and kinetics in mucopolysaccharidosis type VI pa-
tients. Am.

[106498]J. Hum. Genet. 48: 710-719, 1991.

[106499]18702.

[106500]Brooks, D. A.; McCourt, P. A. G.; Gibson, G. J.; Hopwood, J.
J.

[106501]: Immunoquantification of the low abundance lysosomal enzyme N-acetylgalactosamine

[106502]4-sulphatase. J. Inherit. Metab. Dis. 13: 108-120, 1990.

[106503]18703.

[106504]Cantor, L. B.; Disseler, J. A.; Wilson, F. M., II: Glaucoma in

[106505]the Maroteaux-Lamy syndrome. Am. J. Ophthal. 108: 426-430, 1989.

[106506]18704.

[106507]Crawley, A. C.; Brooks, D. A.; Muller, V. J.; Petersen, B. A.;

[106508]Isaac, E. L.; Bielicki, J.; King, B. M.; Boulter, C. D.; Moore, A.

[106509]J.; Fazzalari, N. L.; Anson, D. S.; Byers, S.; Hopwood, J. J.: Enzyme

[106510]replacement therapy in a feline model of Maroteaux-Lamy syndrome. J.

[106511]Clin. Invest. 97: 1864-1873, 1996.

[106512]18705.

[106513]Crawley, A. C.; Niedzielski, K. H.; Isaac, E. L.; Davey, R. C.

[106514]A.; Byers, S.; Hopwood, J. J.: Enzyme replacement therapy from birth

- [106515]in a feline model of mucopolysaccharidosis type VI. J. Clin. Invest. 99: 651-662, 1997.
- [106517]18706.
- [106518]Crawley, A. C.; Yogalingam, G.; Muller, V. J.; Hopwood, J. J.: Two mutations within a feline mucopolysaccharidosis type VI colony cause three different clinical phenotypes. J. Clin. Invest. 101: 109-119, 1998.
- [106522]18707.
- [106523]DeLuca, C.; Brown, J. A.; Shows, T. B.: Lysosomal arylsul-fatase deficiencies in humans: chromosome assignment for aryl-sulfatase A and B. Proc. Nat. Acad. Sci. 76: 1957-1961, 1979.
- [106526]18708.
- [106527]DiFerrante, N. M.; Hyman, B. H.; Kish, W.; Donnelly, P. V.; Nichols, B. L., Jr.; Dutton, R. V.: Mucopolysaccharidosis VI

(Maroteaux-Lamy

[106529]disease) clinical and biochemical study of a mild variant
case. Johns

[106530]Hopkins Med. J. 135: 42-53, 1974.

[106531]18709.

[106532]Dudin, G.; Alexander, D.; Talj, F.; Deeb, M.; Musallam, S.;
Der

[106533]Kaloustian, V. M.: Interstitial deletion of band q12 of
chromosome

[106534]5. Clin. Genet. 25: 455-458, 1984.

[106535]18710.

[106536]Evers, M.; Saftig, P.; Schmidt, P.; Hafner, A.; McLoughlin, D.

[106537]B.; Schmahl, W.; Hess, B.; von Figura, K.; Peters, C.: Tar-
geted disruption

[106538]of the arylsulfatase B gene results in mice resembling the
phenotype

[106539]of mucopolysaccharidosis VI. Proc. Nat. Acad. Sci. 93:
8214-8219,

[106540]1996.

[106541]18711.

[106542]Fidzianska, E.; Abramowicz, T.; Czartoryska, B.; Glo-

gowska, I.;

[106543]Gorska, D.; Mazurozak, T.; Rodo, M.; Wehr, H.; Witkowska,
J.: Localization

[106544]of the human arylsulfatase B gene locus to chromosome 5
by use of

[106545]hamster-man somatic cell hybrids.(Abstract) 7th Int.
Cong. Hum. Genet.,

[106546]Berlin 611-612, 1986.

[106547]18712.

[106548]Fidzianska, E.; Abramowicz, T.; Czartoryska, B.; Glo-
owska, I.;

[106549]Gorska, D.; Rodo, M.: Assignment of the gene for human
arylsulfatase

[106550]B, ARSB, to chromosome region 5p11-5qter. Cytogenet.
Cell Genet. 38:

[106551]150-151, 1984.

[106552]18713.

[106553]Fox, M. F.; DuToit, D. L.; Warnich, L.; Retief, A. E.: Regional

[106554]localization of alpha-galactosidase (GLA) to Xpter-q22,
hexosaminidase

[106555]B (HEXB) to 5q13-qter, and arylsulfatase B (ARSB) to

5pter-q13. Cytogenet.

[106556]Cell Genet. 38: 45-49, 1984.

[106557]18714.

[106558]Gitzelmann, R.: Personal Communication. Zurich, Switzerland

[106559]4/27/1990.

[106560]18715.

[106561]Goldberg, M. F.; Scott, C. I., Jr.; McKusick, V. A.: Hydrocephalus

[106562]and papilledema in the Maroteaux-Lamy syndrome (mucopolysaccharidosis

[106563]type VI). Am. J. Ophthal. 69: 969-975, 1970.

[106564]18716.

[106565]Haskins, M. E.; Jezyk, P. F.; Patterson, D. F.: Mucopolysaccharide

[106566]storage disease in three families of cats with arylsulfatase B deficiency:

[106567]leukocyte studies and carrier identification. Pediat. Res. 13: 1203-1210,

[106568]1979.

[106569]18717.

[106570]Hellkuhl, B.; Grzeschik, K.-H.: Assignment of a gene for
arylsulfatase

[106571]B to human chromosome 5 (EC 3.1.6.1) using human-
mouse somatic cell

[106572]hybrids. Cytogenet. Cell Genet. 22: 203-206, 1978.

[106573]18718.

[106574]Hopwood, J. J.: Personal Communication. North Adelaide,
South

[106575]Australia 5/9/1991.

[106576]18719.

[106577]Isbrandt, D.; Hopwood, J. J.; von Figura, K.; Peters, C.: Two

[106578]novel frameshift mutations causing premature stop
codons in a patient

[106579]with the severe form of Maroteaux-Lamy syndrome. Hum.
Mutat. 7:

[106580]361-363, 1996.

[106581]18720.

[106582]Jezyk, P. F.; Haskins, M. E.; Patterson, D. F.; Mellman, W. J.;

[106583]Greenstein, M.: Mucopolysaccharidosis in a cat with aryl-
sulfatase

[106584]B deficiency: a model of Maroteaux-Lamy syndrome. Science 198: 834-836,
[106585]1977.
[106586]18721.
[106587]Jin, W.-D.; Jackson, C. E.; Desnick, R. J.; Schuchman, E. H.:
[106588]Mucopolysaccharidosis type VI: identification of three mutations in
[106589]the arylsulfatase B gene of patients with the severe and mild phenotypes
[106590]provides molecular evidence for genetic heterogeneity. Am. J. Hum.
[106591]Genet. 50: 795-800, 1992.
[106592]18722.
[106593]Krivit, W.; Pierpont, M. E.; Ayaz, K.; Tsai, M.; Ramsay, N. K.
[106594]C.; Kersey, J. H.; Weisdorf, S.; Sibley, R.; Snover, D.; McGovern,
[106595]M. M.; Schwartz, M. F.; Desnick, R. J.: Bone-marrow transplantation
[106596]in the Maroteaux-Lamy syndrome (mucopolysaccharidosis type VI): biochemical
[106597]and clinical status 24 months after transplantation. New

Eng. J.

[106598]Med. 311: 1606–1611, 1984.

[106599]18723.

[106600]Kunieda, T.; Simonaro, C. M.; Yoshida, M.; Ikadai, H.;
Levan,

[106601]G.; Desnick, R. J.; Schuchman, E. H.: Mucopolysaccharido-
sis type

[106602]VI in rats: isolation of cDNAs encoding arylsulfatase B,
chromosomal

[106603]localization of the gene, and identification of the muta-
tion. Genomics 29:

[106604]582–587, 1995.

[106605]18724.

[106606]Levy, L. A.; Lewis, J. C.; Sumner, T. E.: Ultrastructures of

[106607]Reilly bodies (metachromatic granules) in the Maroteaux–
Lamy syndrome

[106608](mucopolysaccharidosis VI): a histochemical study. Am. J.
Clin. Path. 73:

[106609]416–422, 1980.

[106610]18725.

[106611]Litjens, T.; Baker, E. G.; Beckmann, K. R.; Morris, C. P.;

Hopwood,

[106612]J. J.; Callen, D. F.: Chromosomal localization of ARSB, the
gene

[106613]for human N-acetylgalactosamine-4-sulphatase. Hum.
Genet. 82: 67-68,

[106614]1989.

[106615]18726.

[106616]Litjens, T.; Brooks, D. A.; Peters, C.; Gibson, G. J.; Hop-
wood,

[106617]J. J.: Identification, expression, and biochemical character-
ization

[106618]of N-acetylgalactosamine-4-sulfatase mutations and re-
lationship with

[106619]clinical phenotype in MPS-VI patients. Am. J. Hum. Genet.
58: 1127-1134,

[106620]1996.

[106621]18727.

[106622]Litjens, T.; Hopwood, J. J.: Mucopolysaccharidosis type VI:
structural

[106623]and clinical implications of mutations in N-
acetylgalactosamine-4-sulfatase. Hum.

[106624]Mutat. 18: 282-295, 2001.

[106625]18728.

[106626]Litjens, T.; Morris, C. P.; Robertson, E. F.; Peters, C.; von

[106627]Figura, K.; Hopwood, J. J.: An N-acetyl-

galactosamine-4-sulfatase

[106628]mutation (delta-G-238) results in a severe Maroteaux-

Lamy phenotype. Hum.

[106629]Mutat. 1: 397-402, 1992.

[106630]18729.

[106631]Taroni, F.; Verderio, E.; Dworzak, F.; Willems, P. J.; Cava-
dini,

[106632]P.; DiDonato, S.: Identification of a common mutation in
the carnitine

[106633]palmitoyltransferase II gene in familial recurrent myo-
globinuria patients. Nature

[106634]Genet. 4: 314-320, 1993.

[106635]18730.

[106636]Gellera, C.; Verderio, E.; Floridia, G.; Finocchiaro, G.; Mon-
termini,

[106637]L.; Cavadini, P.; Zuffardi, O.; Taroni, F.: Assignment of the
human

[106638]carnitine palmitoyltransferase II gene (CTP1) to chromo-

some 1p32. Genomics 24:

[106639]195–197, 1994.

[106640]18731.

[106641]Agmon, D.; Green, J.; Platau, E.; Better, O. S.: Isolated
adrenal

[106642]mineralocorticoid deficiency due to amyloidosis associ-
ated with familial

[106643]Mediterranean fever. Am. J. Med. Sci. 288: 40–43, 1984.

[106644]18732.

[106645]Akarsu, A. N.; Saatci, U.; Ozen, S.; Bakkaloglu, A.; Besbas,
N.;

[106646]Sarfarazi, M.: Genetic linkage study of familial Mediter-
ranean fever

[106647](FMF) to 16p13.3 and evidence for genetic heterogeneity
in the Turkish

[106648]population. J. Med. Genet. 34: 573–578, 1997.

[106649]18733.

[106650]Lukong, K. E.; Elsliger, M.–A.; Chang, Y.; Richard, C.;
Thomas,

[106651]G.; Carey, W.; Tyłki-Szymanska, A.; Czartoryska, B.; Buch-
holz, T.;

[106652]Rodriguez Criado, G.; Palmeri, S.; Pshezhetsky, A. V.:
Characterization

[106653]of the sialidase molecular defects in sialidosis patients
suggests

[106654]the structural organization of the lysosomal multienzyme
complex. Hum.

[106655]Molec. Genet. 9: 1075–1085, 2000.

[106656]18734.

[106657]Fernandes, M.; Poirier, C.; Lespinasse, F.; Carle, G. F.: The

[106658]mouse homologs of human GIF, DDB1, and CFL1 genes
are located on chromosome

[106659]19. Mammalian Genome 9: 339–342, 1998.

[106660]18735.

[106661]Aminoff, M.; Carter, J. E.; Chadwick, R. B.; Johnson, C.;
Grasbeck,

[106662]R.; Abdelaal, M. A.; Broch, H.; Jenner, L. B.; Verroust, P. J.;
Moestrup,

[106663]S. K.; de la Chapelle, A.; Krahe, R.: Mutations in CUBN, en-
coding

[106664]the intrinsic factor–vitamin B12 receptor, cubilin, cause
hereditary

[106665]megaloblastic anaemia 1. Nature Genet. 21: 309–313,
1999.

[106666]18736.

[106667]Aminoff, M.; Tahvanainen, E.; Grasbeck, R.; Weissenbach,
J.; Broch,

[106668]H.; de la Chapelle, A.: Selective intestinal malabsorption of
vitamin

[106669]B12 displays recessive mendelian inheritance: assignment
of a locus

[106670]to chromosome 10 by linkage. Am. J. Hum. Genet. 57:
824–831, 1995.

[106671]18737.

[106672]Kozyraki, R.; Kristiansen, M.; Silahtaroglu, A.; Hansen, C.;
Jacobsen,

[106673]C.; Tommerup, N.; Verroust, P. J.; Moestrup, S. K.: The hu-
man intrinsic

[106674]factor–vitamin B12 receptor, cubilin: molecular characteri-
zation and

[106675]chromosomal mapping of the gene to 10p within the au-
tosomal recessive

[106676]megaloblastic anemia (MGA1) region. Blood 91:
3593–3600, 1998.

[106677]18738.

[106678]Carney, J. P.; Maser, R. S.; Olivares, H.; Davis, E. M.; Le
Beau,

[106679]M.; Yates, J. R., III; Hays, L.; Morgan, W. F.; Petrini, J. H. J.

[106680]: The hMre11/hRad50 protein complex and Nijmegen
breakage syndrome:

[106681]linkage of double-strand break repair to the cellular DNA
damage response. Cell 93:

[106682]477-486, 1998.

[106683]18739.

[106684]Matsuo, T.; Egawa, I.; Okada, S.; Suetsugu, M.; Yamamoto,
K.;

[106685]Watanabe, M.: Sialidosis type 2 in Japan: clinical study in
two siblings'

[106686]cases and review of literature. J. Neurol. Sci. 58: 45-55,
1983.

[106687]18740.

[106688]Miyatake, T.; Atsumi, T.; Obayashi, T.; Mizuno, Y.; Ando,
S.;

[106689]Ariga, T.; Matsui-Nakamura, K.; Yamada, T.: Adult type
neuronal storage

[106690]disease with neuraminidase deficiency. Ann. Neurol. 6:
232–243,

[106691]1979.

[106692]18741.

[106693]Mueller, O. T.; Henry, W. M.; Haley, L. L.; Byers, M. G.;
Eddy,

[106694]R. L.; Shows, T. B.: Sialidosis and galactosialidosis: chro-
mosomal

[106695]assignment of two genes associated with neuraminidase
deficiency disorders. Proc.

[106696]Nat. Acad. Sci. 83: 1817–1821, 1986.

[106697]18742.

[106698]Naganawa, Y.; Itoh, K.; Shimmoto, M.; Takiguchi, K.; Doi,
H.;

[106699]Nishizawa, Y.; Kobayashi, T.; Kamei, S.; Lukong, K. E.;
Pshezhetsky,

[106700]A. V.; Sakuraba, H.: Molecular and structural studies of
Japanese

[106701]patients with sialidosis type 1. J. Hum. Genet. 45:
241–249, 2000.

[106702]18743.

[106703] Nakamura, Y.; Takahashi, Y.; Yamaguchi, S.; Omiya, S.;
Orii, T.;

[106704] Yara, A.; Gushiken, M.: Severe infantile sialidosis--the
characteristics

[106705] of oligosaccharides isolated from the urine and the ab-
dominal ascites. Tohoku

[106706] J. Exp. Med. 166: 407-415, 1992.

[106707] 18744.

[106708] O'Brien, J. S.: Neuraminidase deficiency in the cherry red
spot-myoclonus

[106709] syndrome. Biochem. Biophys. Res. Commun. 79:
1136-1141, 1977.

[106710] 18745.

[106711] O'Brien, J. S.: The cherry red spot-myoclonus syndrome: a
newly

[106712] recognized inherited lysosomal storage disease due to
acid neuraminidase

[106713] deficiency. Clin. Genet. 14: 55-60, 1978.

[106714] 18746.

[106715] O'Brien, J. S.; Warner, T. G.: Sialidosis: delineation of sub-
types

[106716]by neuraminidase assay. Clin. Genet. 17: 35–38, 1980.

[106717]18747.

[106718]Oohira, T.; Nagata, N.; Akaboshi, I.; Matsuda, I.; Naito, S.:

[106719]The infantile form of sialidosis type II associated with
congenital

[106720]adrenal hyperplasia: possible linkage between HLA and
the neuraminidase

[106721]deficiency gene. Hum. Genet. 70: 341–343, 1985.

[106722]18748.

[106723]Oohira, T.; Yoshida, M. C.; Matsuda, I.: Additional evi-
dence

[106724]for the location of the alpha-neuraminidase gene on
chromosome 6. Jpn.

[106725]J. Hum. Genet. 31: 309–310, 1986.

[106726]18749.

[106727]Orii, T.; Minami, R.; Sukegawa, K.; Sato, S.; Tsugawa, S.;
Horino,

[106728]K.; Miura, R.; Nakao, T.: A new type of mucopolidosis with
beta-galactosidase

[106729]deficiency and glycopeptiduria. Tohoku J. Exp. Med. 107:
303–315,

[106730]1972.

[106731]18750.

[106732]Paschke, E.; Trinkl, G.; Erwa, W.; Pavelka, M.; Mutz, I.;
Roscher,

[106733]A.: Infantile type of sialic acid storage disease with
sialuria. Clin.

[106734]Genet. 29: 417-424, 1986.

[106735]18751.

[106736]Peters, J.; Swallow, D. M.; Andrews, S. J.; Evans, L.: A gene

[106737](Neu-1) on chromosome 17 of the mouse affects acid al-
pha-glucosidase

[106738]and codes for neuraminidase. Genet. Res. Camb. 38:
47-55, 1981.

[106739]18752.

[106740]Pincus, J. H.; Rossi, J. P.; Daroff, R. B.: Delayed develop-
ment

[106741]of disturbed mucopolysaccharide metabolism in a Hurler
variant. Arch.

[106742]Neurol. 16: 244-253, 1967.

[106743]18753.

[106744]Potier, M.; Beauregard, G.; Gelisle, M.; Mameli, L.; Hong, V.

[106745]N.; Melancon, S. B.; Dallaire, L.: Neuraminidase activity in
the

[106746]mucopolysaccharidoses (types I, II and III) and the cherry-red spot
myoclonus

[106747]syndrome. Clin. Chim. Acta 99: 97-105, 1979.

[106748]18754.

[106749]Pshchetsky, A. V.; Richard, C.; Michaud, L.; Igouda, S.;
Wang,

[106750]S.; Elsliger, M.-A.; Qu, J.; Leclerc, D.; Gravel, R.; Dallaire,
L.;

[106751]Potier, M.: Cloning, expression and chromosomal map-
ping of human

[106752]lysosomal sialidase and characterization of mutations in
sialidosis. Nature

[106753]Genet. 15: 316-320, 1997.

[106754]18755.

[106755]Rottier, R. J.; Bonten, E.; d'Azzo, A.: A point mutation in
the

[106756]neu-1 locus causes the neuraminidase defect in the SM/J
mouse. Hum.

[106757]Molec. Genet. 7: 313–321, 1998.

[106758]18756.

[106759]Sanfilippo, S. J.; Yunis, J. J.; Worthen, H. G.: An unusual storage

[106760]disease resembling the Hurler–Hunter syndrome.
(Abstract) Am. J.

[106761]Dis. Child. 104: 553 only, 1962.

[106762]18757.

[106763]Sasagasako, N.; Miyahara, S.; Saito, N.; Shinnoh, N.;
Kobayashi,

[106764]T.; Goto, I.: Prenatal diagnosis of congenital sialidosis.
Clin.

[106765]Genet. 44: 8–11, 1993.

[106766]18758.

[106767]Sergi, C.; Penzel, R.; Uhl, J.; Zoubaa, S.; Dietrich, H.;
Decker,

[106768]N.; Rieger, P.; Kopitz, J.; Otto, H. F.; Kiessling, M.; Cantz,
M.

[106769]: Prenatal diagnosis and fetal pathology in a Turkish family harboring

[106770]a novel nonsense mutation in the lysosomal alpha–

N-acetyl-neuraminidase

[106771](sialidase) gene. Hum. Genet. 109: 421-428, 2001.

[106772]18759.

[106773]Shows, T. B.: Personal Communication. Buffalo, N. Y.

10/9/1985.

[106774]18760.

[106775]Spranger, J.: Advances in bone dysplasias. (Abstract) Sixth

[106776]Int. Cong. Hum. Genet., Jerusalem , 1981.

[106777]18761.

[106778]Spranger, J.: Mucopolidosis I: phenotype and nosology.

Perspect.

[106779]Inherit. Metab. Dis. 4: 303-315, 1981.

[106780]18762.

[106781]Spranger, J.; Cantz, M.: Mucopolidosis I, the cherry red

spot-myoclonus

[106782]syndrome and neuraminidase deficiency. Birth Defects

Orig. Art. Ser. XIV(6B):

[106783]105-112, 1978.

[106784]18763.

[106785]Spranger, J. W.: Personal Communication. Kiel, Germany
1972.

[106786]18764.

[106787]Spranger, J. W.; Gehler, J.; Cantz, M.: Mucopolidosis I--a
sialidosis. Am.

[106788]J. Med. Genet. 1: 21-29, 1977.

[106789]18765.

[106790]Spranger, J. W.; Wiedemann, H. R.: The genetic mucolipi-
doses.

[106791]Diagnosis and differential diagnosis. Humangenetik 9:
113-139, 1970.

[106792]18766.

[106793]Spranger, J. W.; Wiedemann, H. R.; Tolksdorf, M.; Graucob,
E.;

[106794]Caesar, R.: Lipomucopolysaccharidose: eine neue Spe-
icherkrankheit. Z.

[106795]Kinderheilk. 103: 285-306, 1968.

[106796]18767.

[106797]Steinmann, L.; Tharp, B. R.; Dorfman, L. J.; Forno, L. S.;
Sogg,

[106798]R. L.; Kelts, K. A.; O'Brien, J. S.: Peripheral neuropathy in

the

[106799]cherry red spot–myoclonus syndrome (sialidosis type I).

Ann. Neurol. 7:

[106800]450–456, 1980.

[106801]18768.

[106802]Stevenson, R. E.; Lubinsky, M.; Taylor, H. A.; Wenger, D.

A.;

[106803]Schroer, R. J.; Olmstead, P. M.: Sialic acid storage disease

with

[106804]sialuria: clinical and biochemical features in the severe in-

fantile

[106805]type. Pediatrics 72: 441–449, 1983.

[106806]18769.

[106807]Swallow, D. M.; Evans, L.; Stewart, G.; Thomas, P. K.;

Abrams,

[106808]J. D.: Sialidosis type I: cherry red spot–myoclonus syn-

drome with

[106809]sialidase deficiency and altered electrophoretic mobility of

some

[106810]enzymes known to be glycoproteins. II. Enzyme studies.

Ann. Hum.

[106811]Genet. 43: 27–35, 1979.

[106812]18770.

[106813]Swallow, D. M.; Hoogeveen, A. T.; Verheijen, F. W.; Galjaard,

[106814]H.: Complementation analysis of human sialidase deficiency using

[106815]natural substrates. Ann. Hum. Genet. 45: 105–112, 1981.

[106816]18771.

[106817]Kitagawa, T.: An animal model of human acid sphingomyelinase

[106818]deficiency (Niemann–Pick disease) and the study of its enzyme replacement. Jpn.

[106819]J. Hum. Genet. 32: 55–69, 1987.

[106820]18772.

[106821]Kurimasa, A.; Ohno, K.; Oshimura, M.: Restoration of the cholesterol

[106822]metabolism in 3T3 cell lines derived from the sphingomyelinosis mouse

[106823](spm/spm) by transfer of a human chromosome 18. Hum. Genet. 92:

[106824]157–162, 1993.

[106825]18773.

- [106826]Miyawaki, S.; Yoshida, H.; Mitsuoka, S.; Enomoto, H.; Ikehara,
hara,
[106827]S.: A mouse model for Niemann–Pick disease: influence of
genetic
[106828]background on disease expression in spm/spm mice. J.
Hered. 77:
[106829]379–384, 1986.
[106830]18774.
[106831]Mailly, F.; Palmen, J.; Muller, D. P. R.; Gibbs, T.; Lloyd, J.;
[106832]Brunzell, J.; Durrington, P.; Mitropoulos, K.; Betteridge, J.;
Watts,
[106833]G.; Lithell, H.; Angelico, F.; Humphries, S. E.; Talmud, P. J.:
Familial
[106834]lipoprotein lipase (LPL) deficiency: a catalogue of LPL gene
mutations
[106835]identified in 20 patients from the UK, Sweden, and Italy.
Hum. Mutat. 10:
[106836]465–473, 1997.
[106837]18775.
[106838]Normand, T.; Bergeron, J.; Fernandez–Margallo, T.;
Bharucha, A.;

[106839]Ven Murthy, M. R.; Julien, P.; Gagne, C.; Dionne, C.; De
Braekeleer,
[106840]M.; Ma, R.; Hayden, M. R.; Lupien, P. J.: Geographic distri-
bution
[106841]and genealogy of mutation 207 of the lipoprotein lipase
gene in the
[106842]French Canadian population of Quebec. Hum. Genet. 89:
671-675, 1992.
[106843]18776.

[106844]Monsalve, M. V.; Henderson, H.; Roederer, G.; Julien, P.;
Deeb,
[106845]S.; Kastelein, J. J. P.; Peritz, L.; Devlin, R.; Bruin, T.; Murthy,
[106846]M. R. V.; Gagne, C.; Davignon, J.; Lupien, P. J.; Brunzell, J.
D.;
[106847]Hayden, M. R.: A missense mutation at codon 188 of the
human lipoprotein
[106848]lipase gene is a frequent cause of lipoprotein lipase defi-
ciency in
[106849]persons of different ancestries. J. Clin. Invest. 86:
728-734, 1990.
[106850]18777.

[106851]Campbell, C.; Cucci, R. A.; Prasad, S.; Green, G. E.; Edeal, J.

[106852]B.; Galer, C. E.; Karniski, L. P.; Sheffield, V. C.; Smith, R. J.
 [106853]H.: Pendred syndrome, DFNB4, and PDS/SLC26A4 identification of eight
 [106854]novel mutations and possible genotype-phenotype correlations. Hum.
 [106855]Mutat. 17: 403-411, 2001.
 [106856]18778.
 [106857]Everett, L. A.; Glaser, B.; Beck, J. C.; Idol, J. R.; Buchs, A.;
 [106858]Heyman, M.; Adawi, F.; Hazani, E.; Nassir, E.; Baxevanis, A.
 D.; Sheffield,
 [106859]V. C.; Green, E. D.: Pendred syndrome is caused by mutations in a
 [106860]putative sulphate transporter gene (PDS). Nature Genet.
 17: 411-422,
 [106861]1997.
 [106862]18779.
 [106863]Fugazzola, L.; Mannavola, D.; Cerutti, N.; Maghnie, M.;
 Pagella,
 [106864]F.; Bianchi, P.; Weber, G.; Persani, L.; Beck-Peccoz, P.:
 Molecular
 [106865]analysis of the Pendred's syndrome gene and magnetic

resonance imaging

[106866]studies of the inner ear are essential for the diagnosis of
true Pendred's

[106867]syndrome. J. Clin. Endocr. Metab. 85: 2469–2475, 2000.

[106868]18780.

[106869]Kopp, P.; Karamanoglu Arseven, O.; Sabacan, L.; Kotlar, T.;
Dupuis,

[106870]J.; Cavaliere, H.; Santos, C. L. S.; Jameson, J. L.; Medeiros–
Neto,

[106871]G.: Phenocopies for deafness and goiter development in a
large inbred

[106872]Brazilian kindred with Pendred's syndrome associated with
a novel

[106873]mutation in the PDS gene. J. Clin. Endocr. Metab. 84:
336–341, 1999.

[106874]18781.

[106875]Masmoudi, S.; Charfedine, I.; Hmani, M.; Grati, M.; Ghor–
bel, A.

[106876]M.; Elgaied–Boulila, A.; Drira, M.; Hardelin, J.–P.; Ayadi, M.:
Pendred

[106877]syndrome: phenotypic variability in two families carrying
the same

[106878]PDS missense mutation. Am. J. Med. Genet. 90: 38–44,
2000.

[106879]18782.

[106880]Boughman, J. A.; Conneally, P. M.; Nance, W. E.: Population
genetic

[106881]studies of retinitis pigmentosa. Am. J. Hum. Genet. 32:
223–235,

[106882]1980.

[106883]18783.

[106884]Gomez–Lira, M.; Perusi, C.; Mottes, M.; Pignatti, P. F.;
Manfredi,

[106885]M.; Rizzuto, N.; Salviati, A.: Molecular genetic characteri-
zation

[106886]of two metachromatic leukodystrophy patients who carry
the T799G mutation

[106887]and show different phenotypes; description of a novel
null-type mutation. Hum.

[106888]Genet. 102: 459–463, 1998.

[106889]18784.

[106890]Neer, E. J.; Schmidt, C. J.; Smith, T.: LIS is more. Nature

[106891]Genet. 5: 3–4, 1993.

[106892]18785.

[106893]Reiner, O.; Carrozzo, R.; Shen, Y.; Wehnert, M.;

Faustinella,

[106894]F.; Dobyns, W. B.; Caskey, C. T.; Ledbetter, D. H.: Isolation
of

[106895]a Miller-Dieker lissencephaly gene containing G protein
beta-subunit-like

[106896]repeats. Nature 364: 717-721, 1993.

[106897]18786.

[106898]Greene, H.; Hug, G.; Schubert, W. K.: Arylsulfatase A in the

[106899]urine and metachromatic leukodystrophy. J. Pediat. 71:
709-711,

[106900]1967.

[106901]18787.

[106902]Greenfield, J. G.: Form of progressive cerebral sclerosis in

[106903]infants associated with primary degeneration of interfascicular
glia. Proc.

[106904]Roy. Soc. Med. 26: 690-697, 1933.

[106905]18788.

[106906]Gustavson, K.-H.; Hagberg, B.: The incidence and genetics

of

[106907]metachromatic leukodystrophy in northern Sweden. Acta
Paediat. Scand. 60:

[106908]585–590, 1971.

[106909]18789.

[106910]Hagberg, B.; Sourander, P.; Svennerholm, L.: Sulfatide lipi-
dosis

[106911]in childhood. Am. J. Dis. Child. 104: 644–656, 1962.

[106912]18790.

[106913]Haltia, T.; Palo, J.; Haltia, M.; Icen, A.: Juvenile metachro-
matic

[106914]leukodystrophy: clinical, biochemical, and neuropatho-
logic studies

[106915]in nine new cases. Arch. Neurol. 37: 42–46, 1980.

[106916]18791.

[106917]Harvey, J. S.; Carey, W. F.; Morris, C. P.: Importance of the

[106918]glycosylation and polyadenylation variants in metachro-
matic leukodystrophy

[106919]pseudodeficiency phenotype. Hum. Molec. Genet. 7:
1215–1219, 1998.

[106920]18792.

- [106921]Harvey, J. S.; Nelson, P. V.; Carey, W. F.; Robertson, E. F.;
[106922]Morris, C. P.: An arylsulfatase A (ARSA) missense mutation
(T274M)
[106923]causing late-infantile metachromatic leukodystrophy.
Hum. Mutat. 2:
[106924]261–267, 1993.
[106925]18793.
[106926]Hasegawa, Y.; Kawame, H.; Eto, Y.: Mutations in the aryl-
sulfatase
[106927]A gene of Japanese patients with metachromatic
leukodystrophy. DNA
[106928]Cell Biol. 12: 493–498, 1993.
[106929]18794.
[106930]Hasegawa, Y.; Kawame, H.; Ida, H.; Ohashi, T.; Eto, Y.: Sin-
gle
[106931]exon mutation in arylsulfatase A gene has two effects:
loss of enzyme
[106932]activity and aberrant splicing. Hum. Genet. 93: 415–420,
1994.
[106933]18795.
[106934]Heinisch, U.; Zlotogora, J.; Kafert, S.; Gieselmann, V.: Mul-

tiple

[106935] mutations are responsible for the high frequency of
metachromatic

[106936] leukodystrophy in a small geographic area. Am. J. Hum.
Genet. 56:

[106937] 51–57, 1995.

[106938] 18796.

[106939] Herz, B.; Bach, G.: Arylsulfatase A in pseudodeficiency.
Hum.

[106940] Genet. 66: 147–150, 1984.

[106941] 18797.

[106942] Hess, B.; Saftig, P.; Hartmann, D.; Coenen, R.; Lullmann–
Rauch,

[106943] R.; Goebel, H. H.; Evers, M.; von Figura, K.; D'Hooge, R.;
Nagels,

[106944] G.; De Deyn, P.; Peters, C.; Gieselmann, V.: Phenotype of
arylsulfatase

[106945] A–deficient mice: relationship to human metachromatic
leukodystrophy. Proc.

[106946] Nat. Acad. Sci. 93: 14821–14826, 1996.

[106947] 18798.

[106948]Hohenschutz, C.; Eich, P.; Friedl, W.; Waheed, A.; Conzelmann,

[106949]E.; Propping, P.: Pseudodeficiency of arylsulfatase A: a common genetic

[106950]polymorphism with possible disease implications. Hum. Genet. 82:

[106951]45–48, 1989.

[106952]18799.

[106953]Hohenschutz, C.; Friedl, W.; Schlor, K.–H.; Waheed, A.; Conzelmann,

[106954]E.; Sandhoff, K.; Propping, P.: Probable metachromatic leukodystrophy/pseudodeficiency

[106955]compound heterozygote at the arylsulfatase A locus with neurological

[106956]and psychiatric symptomatology. Am. J. Med. Genet. 31: 169–175,

[106957]1988.

[106958]18800.

[106959]Holve, S.; Hu, D.; McCandless, S. E.: Metachromatic leukodystrophy

[106960]in the Navajo: fallout of the American–Indian Wars of the nineteenth

[106961]century. Am. J. Med. Genet. 101: 203–208, 2001.

[106962]18801.

[106963]Honke, K.; Kobayashi, T.; Fujii, T.; Gasa, S.; Xu, M.; Takamaru,

[106964]Y.; Kondo, R.; Tsuji, S.; Makita, A.: An adult-type metachromatic

[106965]leukodystrophy caused by substitution of serine for glycine-122 in

[106966]arylsulfatase A. Hum. Genet. 92: 451–456, 1993.

[106967]18802.

[106968]Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Weil, D.; Frezal,

[106969]J.: Confirmation of the assignment of the gene for arylsulfatase

[106970]A to chromosome 22 using somatic cell hybrids. Hum. Genet. 49: 33–39,

[106971]1979.

[106972]18803.

[106973]Jervis, G. A.: Infantile metachromatic leukodystrophy (Greenfield's

[106974]disease). J. Neuropath. Exp. Neurol. 19: 323–341, 1960.

[106975]18804.

[106976]Kaback, M. M.; Howell, R. R.: Infantile metachromatic leukodystrophy:

[106977]heterozygote detection in skin fibroblasts and possible applications

[106978]to intrauterine diagnosis. New Eng. J. Med. 282: 1336–1340, 1970.

[106979]18805.

[106980]Kappler, J.; Sommerlade, H. J.; von Figura, K.; Gieselmann, V.

[106981]: Complex arylsulfatase A alleles causing metachromatic leukodystrophy. Hum.

[106982]Mutat. 4: 119–127, 1994.

[106983]18806.

[106984]Kappler, J.; von Figura, K.; Gieselmann, V.: Late-onset metachromatic

[106985]leukodystrophy: molecular pathology in two siblings. Ann. Neurol. 31:

[106986]256–261, 1992.

[106987]18807.

[106988]Kihara, H.: Genetic heterogeneity in metachromatic

leukodystrophy. Am.

[106989]J. Hum. Genet. 34: 171-181, 1982.

[106990]18808.

[106991]Kihara, H.; Fluharty, A. L.; O'Brien, J. S.; Fish, C. H.:

Metachromatic

[106992]leukodystrophy caused by a partial cerebroside sulfatase defect. Clin.

[106993]Genet. 21: 253-261, 1982.

[106994]18809.

[106995]Kihara, H.; Meek, W. E.; Fluharty, A. L.: Attenuated activities

[106996]and structural alterations of arylsulfatase A in tissues from subjects

[106997]with pseudo arylsulfatase A deficiency. Hum. Genet. 74: 59-62, 1986.

[106998]18810.

[106999]Kohn, H.; Manowitz, P.; Miller, M.; Kling, A.: Neuropsychological

[107000]deficits in obligatory heterozygotes for metachromatic leukodystrophy. Hum.

[107001]Genet. 79: 8-12, 1988.

[107002]18811.

[107003]Kondo, R.; Wakamatsu, N.; Yoshino, H.; Fukuhara, N.; Miy-
atake,

[107004]T.; Tsuji, S.: Identification of a mutation in the arylsulfa-
tase

[107005]A gene of a patient with adult-type metachromatic
leukodystrophy. Am.

[107006]J. Hum. Genet. 48: 971-978, 1991.

[107007]18812.

[107008]Kreysing, J.; Bohne, W.; Bosenberg, C.; Marchesini, S.;
Turpin,

[107009]J. C.; Baumann, N.; von Figura, K.; Gieselmann, V.: High
residual

[107010]aryl sulfatase A (ARSA) activity in a patient with late-
infantile metachromatic

[107011]leukodystrophy. Am. J. Hum. Genet. 53: 339–346, 1993.

[107012]18813.

[107013]Krivit, W.; Shapiro, E.; Kennedy, W.; Lipton, M.; Lockman, L.;

[107014]Smith, S.; Summers, C. G.; Wenger, D. A.; Tsai, M. Y.; Ramsay, N.

[107015]K. C.; Kersey, J. H.; Yao, J. K.; Kaye, E.: Treatment of late infantile

[107016]metachromatic leukodystrophy by bone marrow transplantation. New

[107017]Eng. J. Med. 322: 28–32, 1990.

[107018]18814.

[107019]Langenbeck, U.; Dunker, P.; Heipertz, R.; Pilz, H.: Inheritance

[107020]of metachromatic leukodystrophy. (Letter) Am. J. Hum. Genet. 29:

[107021]639–640, 1977.

[107022]18815.

[107023]Li, Z. G.; Waye, J. S.; Chang, P. L.: Diagnosis of arylsulphatase

[107024]A deficiency. Am. J. Med. Genet. 43: 976–982, 1992.

[107025]18816.

[107026]Lugowska, A.; Berger, J.; Tylki-Szymanska, A.; Czartowska, B.;

[107027]Loschl, B.; Molzer, B.: High prevalence of I179S mutation in patients

[107028]with late-onset metachromatic leukodystrophy. (Letter) Clin. Genet. 61:

[107029]389–390, 2002.

[107030]18817.

[107031]Lyon, G.; Arthiu, M.; Thieffry, S.: Leucodystrophie metachromatique

[107032]infantile familiale: etude de deux observations, dont une avec examen

[107033]anatomique et chimique. Rev. Neurol. 104: 508–533, 1961.

[107034]18818.

[107035]Masters, P. L.; MacDonald, W. B.; Ryan, M. M. P.; Cumings, J.

[107036]N.: Familial leucodystrophy. Arch. Dis. Child. 39: 345–355, 1964.

[107037]18819.

- [107038] Moser, H. W.: Sulfatide lipidosis: metachromatic leukodys-
trophy. In:
- [107039] Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The
Metabolic
- [107040] Basis of Inherited Disease. New York: McGraw-Hill (pub.)
(3rd
- [107041] ed.): 1972. Pp. 688-729.
- [107042] 18820.
- [107043] Kestila, M.; Lenkkeri, U.; Mannikko, M.; Lamerdin, J.; Mc-
Cready,
- [107044] P.; Putaala, H.; Ruotsalainen, V.; Morita, T.; Nissinen, M.;
Herva,
- [107045] R.; Kashtan, C. E.; Peltonen, L.; Holmberg, C.; Olsen, A.;
Tryggvason,
- [107046] K.: Positionally cloned gene for a novel glomerular pro-
tein--nephrin--is
- [107047] mutated in congenital nephrotic syndrome. Molec. Cell 1:
575-582,
- [107048] 1998.
- [107049] 18821.
- [107050] Muller, D.; Pilz, H.; Ter Meulen, V.: Studies on adult
metachromatic

[107051]leukodystrophy. I. Clinical, morphological and histochemical observations

[107052]in two cases. J. Neurol. Sci. 9: 567–584, 1969.

[107053]18822.

[107054]Narahara, K.; Takahashi, Y.; Murakami, M.; Tsuji, K.; Yokoyama,

[107055]Y.; Murakami, R.; Ninomiya, S.; Seino, Y.: Terminal 22q deletion

[107056]associated with a partial deficiency of arylsulphatase A. J. Med.

[107057]Genet. 29: 432–433, 1992.

[107058]18823.

[107059]Nelson, P. V.; Carey, W. F.; Morris, C. P.: Population frequency

[107060]of the arylsulphatase A pseudo-deficiency allele. Hum. Genet. 87:

[107061]87–88, 1991.

[107062]18824.

[107063]Nicholls, R. G.; Roy, A. G.: Arylsulfatases. In: Boyer, P. D.

[107064]: The Enzymes. New York: Academic Press (pub.) 5: 1971. Pp. 21–41.

[107065]18825.

[107066]Ott, R.; Waye, J. S.; Chang, P. L.: Evolutionary origins of
two

[107067]tightly linked mutations in arylsulfatase-A pseudodeficiency. Hum.

[107068]Genet. 101: 135–140, 1997.

[107069]18826.

[107070]Pastor-Soler, N. M.; Rafi, M. A.; Hoffman, J. D.; Hu, D.;
Wenger,

[107071]D. A.: Metachromatic leukodystrophy in the Navajo Indian
population:

[107072]a splice site mutation in intron 4 of the arylsulfatase A
gene. Hum.

[107073]Mutat. 4: 199–207, 1994.

[107074]18827.

[107075]Percy, A. K.; Brady, R. O.: Metachromatic leukodystrophy:
diagnosis

[107076]with samples of venous blood. Science 161: 594–595,
1968.

[107077]18828.

[107078]Percy, A. K.; Kaback, M. M.: Infantile and adult-onset

metachromatic

[107079]leukodystrophy: biochemical comparisons and predictive diagnosis. New

[107080]Eng. J. Med. 285: 785-787, 1971.

[107081]18829.

[107082]Percy, A. K.; Kaback, M. M.; Herndon, R. M.: Metachromatic leukodystrophy:

[107083]comparison of early- and late-onset forms. Neurology 27: 933-941,

[107084]1977.

[107085]18830.

[107086]Pilz, H.; Duensing, I.; Heipertz, R.; Seidel, D.; Lowitsch, K.;

[107087]Hopf, H. C.; Goebel, H. H.: Adult metachromatic leukodystrophy. I.

[107088]Clinical manifestation in a female aged 44 years, previously diagnosed

[107089]in the preclinical state. Europ. Neurol. 15: 301-307, 1977.

[107090]18831.

[107091]Pilz, H.; Muller, D.: Studies on adult metachromatic leukodystrophy.

[107092]II. Biochemical aspects of adult cases of metachromatic

leukodystrophy. J.

[107093]Neurol. Sci. 9: 585–595, 1969.

[107094]18832.

[107095]Poenaru, L.; Castelnau, L.; Besancon, A.–M.; Nicolesco, H.;
Akli,

[107096]S.; Theophil, D.: First trimester prenatal diagnosis of
metachromatic

[107097]leukodystrophy on chorionic villi by 'immunoprecipitation–elec-
trophoresis.'. J.

[107098]Inherit. Metab. Dis. 11: 123–130, 1988.

[107099]18833.

[107100]Polten, A.; Fluharty, A. L.; Fluharty, C. B.; Kappler, J.; von

[107101]Figura, K.; Gieselmann, V.: Molecular basis of different
forms of

[107102]metachromatic leukodystrophy. New Eng. J. Med. 324:
18–22, 1991.

[107103]18834.

[107104]Porter, M. T.; Fluharty, A. L.; Kihara, H.: Correction of ab-
normal

[107105]cerebroside sulfate metabolism in cultured metachromatic
leukodystrophy

[107106]fibroblasts. Science 172: 1263–1265, 1971.

[107107]18835.

[107108]Porter, M. T.; Fluharty, A. L.; Trammell, J.; Kihara, H.: A correlation

[107109]of intracellular cerebroside sulfatase activity in fibroblasts with

[107110]latency in metachromatic leukodystrophy. Biochem. Biophys. Res. Commun. 44:

[107111]660–666, 1971.

[107112]18836.

[107113]Propping, P.; Friedl, W.; Huschka, M.; Schlor, K.–H.; Reimer,

[107114]F.; Lee–Vaupel, M.; Conzelmann, E.; Sandhoff, K.: The influence of

[107115]low arylsulfatase A activity on neuropsychiatric morbidity: a large–scale

[107116]screening in patients. Hum. Genet. 74: 244–248, 1986.

[107117]18837.

[107118]Quigley, H. A.; Green, W. R.: Clinical and ultrastructural ocular

[107119]histopathologic studies of adult–onset metachromatic

leukodystrophy. Am.

[107120]J. Ophthalm. 82: 472-479, 1976.

[107121]18838.

[107122]Regis, S.; Corsolini, F.; Stroppiano, M.; Cusano, R.; Filo-
carno,

[107123]M.: Contribution of arylsulfatase A mutations located on
the same

[107124]allele to enzyme activity reduction and metachromatic
leukodystrophy

[107125]severity. Hum. Genet. 110: 351-355, 2002.

[107126]18839.

[107127]Ricketts, M. H.; Poretz, R. D.; Manowitz, P.: The R496H
mutation

[107128]of arylsulfatase A does not cause metachromatic
leukodystrophy. Hum.

[107129]Mutat. 12: 238-239, 1998.

[107130]18840.

[107131]Sanguinetti, N.; Marsh, J.; Jackson, M.; Fensom, A. H.;
Warren,

[107132]R. C.; Rodeck, C. H.: The arylsulphatases of chorionic villi:
potential

- [107133] problems in the first-trimester diagnosis of metachromatic leucodystrophy
- [107134] and Maroteaux-Lamy disease. Clin. Genet. 30: 302-308, 1986.
- [107135] 18841.
- [107136] Schaap, T.; Zlotogora, J.; Elian, E.; Barak, Y.; Bach, G.: The
- [107137] genetics of the aryl sulfatase A locus. Am. J. Hum. Genet. 33: 531-539, 1981.
- [107138] 1981.
- [107139]
- [107140] 100. Schutta, H. S.; Pratt, R. T. C.; Metz, H.; Evans, K. A.; Carter,
- [107141] C. O.: A family study of the late infantile and juvenile forms of
- [107142] metachromatic leukodystrophy. J. Med. Genet. 3: 86-91, 1966.
- [107143]
- [107144] 101. Shen, N.; Li, Z.-G.; Waye, J. S.; Francis, G.; Chang, P. L.:
- [107145] Complications in the genotypic molecular diagnosis of pseudo arylsulfatase

[107146]A deficiency. Am. J. Med. Genet. 45: 631–637, 1993.

[107147]

[107148]102. Sourander, P.; Svennerholm, L.: Sulphatide lipidosis
in the

[107149]adult with the clinical picture of progressive organic de-
mentia with

[107150]epileptic seizures. Acta Neuropath. 1: 384–396, 1962.

[107151]

[107152]103. Stein, C.; Gieselmann, V.; Kreysing, J.; Schmidt, B.;
Pohlmann,

[107153]R.; Waheed, A.; Meyer, H. E.; O'Brien, J. S.; von Figura, K.:
Cloning

[107154]and expression of human arylsulfatase A. J. Biol. Chem.
264: 1252–1259,

[107155]1989.

[107156]

[107157]104. Stumpf, D. A.; Austin, J.: Metachromatic leukodystro-
phy (MLD).

[107158]IX. Qualitative and quantitative differences in urinary aryl-
sulfatase

[107159]A in different forms of MLD. Arch. Neurol. 24: 117–124,

1971.

[107160]

[107161]105. Tonnesen, T.; Bro, P. V.; Nielsen, K. B.; Lykkelund, C.:

Metachromatic

[107162]leukodystrophy and pseudoarylsulfatase A deficiency in a

Danish family. Acta

[107163]Paediat. Scand. 72: 175–178, 1983.

[107164]

[107165]106. Tonnesen, T.; Vrang, C.; Wiesmann, U. N.; Christom-

anou, H.; Lou,

[107166]H. O.: Atypical metachromatic leukodystrophy? Problems
with the biochemical

[107167]diagnosis. Hum. Genet. 67: 170–173, 1984.

[107168]

[107169]107. Van Bogaert, L. V.; Dewulf, A.: Diffuse progressive

leukodystrophy

[107170]in the adult with production of metachromatic degenerative
products

[107171](Alzheimer-Baroncini). Arch. Neurol. Psychiat. 42:

1083–1097, 1939.

[107172]

[107173]108. von Figura, K.; Steckel, F.; Conary, J.; Hasilik, A.; Shaw, E.

[107174]: Heterogeneity in late-onset metachromatic leukodystrophy: effect

[107175]of inhibitors of cysteine proteinases. Am. J. Hum. Genet. 39: 371-382,

[107176]1986.

[107177]

[107178]109. von Figura, K.; Steckel, F.; Hasilik, A.: Juvenile and adult

[107179]metachromatic leukodystrophy: partial restoration of aryl-sulfatase

[107180]A (cerebroside sulfatase) activity by inhibitors of thiol proteinases. Proc.

[107181]Nat. Acad. Sci. 80: 6066-6070, 1983.

[107182]

[107183]110. Waheed, A.; Steckel, F.; Hasilik, A.; von Figura, K.: Two allelic

[107184]forms of human arylsulfatase A with different numbers of asparagine-linked

[107185]oligosaccharides. Am. J. Hum. Genet. 35: 228-233, 1983.

[107186]

[107187]111. Waltz, G.; Harik, S. I.; Kaufman, B.: Adult metachromatic leukodystrophy:

[107188]value of computed tomographic scanning and magnetic resonance imaging

[107189]of the brain. Arch. Neurol. 44: 225–227, 1987.

[107190]

[107191]112. Yatziv, S.; Russell, A.: An unusual form of metachromatic leukodystrophy

[107192]in three siblings. Clin. Genet. 19: 222–227, 1981.

[107193]

[107194]113. Zlotogora, J.; Bach, G.: Deficiency of lysosomal hydrolases

[107195]in apparently healthy individuals. Am. J. Med. Genet. 14: 73–80,

[107196]1983.

[107197]

[107198]114. Zlotogora, J.; Bach, G.; Barak, Y.; Elian, E.: Metachromatic

[107199]leukodystrophy in the Habbanite Jews: high frequency in a genetic

[107200]isolate and screening for heterozygotes. Am. J. Hum.
Genet. 32:

[107201]663–669, 1980.

[107202]

[107203]115. Zlotogora, J.; Furman–Shaharabani, Y.; Harris, A.;
Barth, M.

[107204]L.; von Figura, K.; Gieselmann, V.: A single origin for the
most

[107205]frequent mutation causing late infantile metachromatic
leucodystrophy. J.

[107206]Med. Genet. 31: 672–674, 1994.

[107207]

[107208]116. Zlotogora, J.; Gieselman, V.; von Figura, K.; Zeigler,
M.; Bach,

[107209]G.: Late infantile metachromatic leukodystrophy in Israel.
Biomed.

[107210]Pharmacother. 48: 347–350, 1994.

[107211]18842.

[107212]Rozet, J.–M.; Gerber, S.; Ghazi, I. Perrault, I.; Ducroq, D.;

[107213]Souied, E.; Cabot, A.; Dufier, J.–L.; Munnich, A.; Kaplan, J.:
Mutations

[107214] of the retinal specific ATP binding transporter gene (ABCR)
in a single
[107215] family segregating both autosomal recessive retinitis pig-
mentosa RP19
[107216] and Stargardt disease: evidence of clinical heterogeneity
at this
[107217] locus. J. Med. Genet. 36: 447-451, 1999.
[107218] 18843.
[107219] Rozet, J.-M.; Gerber, S.; Souied, E.; Perrault, I.; Chatelin,
[107220] S.; Ghazi, I.; Leowski, C.; Dufier, J.-L.; Munnich, A.; Ka-
plan, J.
[107221]: Spectrum of ABCR gene mutations in autosomal reces-
sive macular dystrophies. Europ.
[107222] J. Hum. Genet. 6: 291-295, 1998.
[107223] 18844.
[107224] Coppa, G. V.; Giorgi, P. L.; Felici, L.; Gabrielli, O.; Donti,
[107225] E.; Bernasconi, S.; Kresse, H.; Paschke, E.; Mastropaolo, C.:
Clinical
[107226] heterogeneity in Sanfilippo disease
(mucopolysaccharidosis III) type
[107227] D: presentation of two new cases. Europ. J. Pediat. 140:

130–133,

[107228]1983.

[107229]18845.

[107230]Dhoot, G. K.; Gustafsson, M. K.; Ai, X.; Sun, W.; Standiford, D.

[107231]M.; Emerson, C. P., Jr.: Regulation of Wnt signaling and embryo patterning

[107232]by an extracellular sulfatase. Science 293: 1663–1666, 2001.

[107233]18846.

[107234]Freeman, C.; Clements, P. R.; Hopwood, J. J.: Human liver N-acetylglucosamine-6-sulphate

[107235]sulphatase: purification and characterization. Biochem. J. 246:

[107236]347–354, 1987.

[107237]18847.

[107238]Freeman, C.; Hopwood, J. J.: Human liver N-acetylglucosamine-6-sulphate

[107239]sulphatase: catalytic properties. Biochem. J. 246: 355–365, 1987.

[107240]18848.

[107241]Gatti, R.; Borrone, C.; Durand, P.; De Virgiliis, S.; Sanna, G.;

[107242]Cao, A.; von Figura, K.; Kresse, H.; Paschke, E.: Sanfilippo type

[107243]D disease: clinical findings in two patients with a new variant of

[107244]mucopolysaccharidosis III. Europ. J. Pediat. 138: 168–171, 1982.

[107245]18849.

[107246]Kaplan, P.; Wolfe, L. S.: Sanfilippo syndrome type D. J. Pediat. 110:

[107247]267–271, 1987.

[107248]18850.

[107249]Kresse, H.; Paschke, E.; von Figura, K.; Gilberg, W.; Fuchs, W.

[107250]: Sanfilippo disease type D: deficiency of N-acetylglucosamine-6-sulfate

[107251]sulfatase required for heparan sulfate degradation. Proc. Nat. Acad.

[107252]Sci. 77: 6822–6826, 1980.

[107253]18851.

- [107254]Robertson, D. A.; Callen, D. F.; Baker, E. G.; Morris, C. P.;
Hopwood,
[107255]J. J.: Chromosomal localization of the gene for human glu-
cosamine-6-sulphatase
[107256]to 12q14. Hum. Genet. 79: 175-178, 1988.
[107257]18852.
- [107258]Robertson, D. A.; Freeman, C.; Nelson, P. V.; Morris, C. P.;
Hopwood,
[107259]J. J.: Human glucosamine-6-sulfatase cDNA reveals ho-
mology with steroid
[107260]sulfatase. Biochem. Biophys. Res. Commun. 157:
218-224, 1988.
[107261]18853.
- [107262]Siciliano, L.; Fiumara, A.; Pavone, L.; Freeman, C.; Robert-
son,
[107263]D.; Morris, C. P.; Hopwood, J. J.; Di Natale, P.; Musumeci,
S.; Horwitz,
[107264]A. L.: Sanfilippo syndrome type D in two adolescent sis-
ters. J.
[107265]Med. Genet. 28: 402-405, 1991.
[107266]18854.

[107267]Thompson, J. N.; Jones, M. Z.; Dawson, G.; Huffman, P. S.:

N-acetylglucosamine

[107268]6-sulphatase deficiency in a Nubian goat: a model of San-filippo syndrome

[107269]type D (mucopolysaccharidosis IIID). J. Inherit. Metab. Dis.
15:

[107270]760-768, 1992.

[107271]18855.

[107272]Baker, E.; Guo, X.-H.; Orsborn, A. M.; Sutherland, G. R.;
Callen,

[107273]D. F.; Hopwood, J. J.; Morris, C. P.: The Morquio A syndrome (mucopolysaccharidosis

[107274]IVA) gene maps to 16q24.3. Am. J. Hum. Genet. 52:
96-98, 1993.

[107275]18856.

[107276]Beck, M.; Braun, S.; Coerdts, W.; Merz, E.; Young, E.; Sewell,
A.

[107277]C.: Fetal presentation of Morquio disease type A. Prenatal
Diag. 12:

[107278]1019-1029, 1992.

[107279]18857.

[107280]Blaw, M. E.; Langer, L. O., Jr.: Spinal cord compression in
Morquio–Brailsford's

[107281]disease. J. Pediat. 74: 593–600, 1969.

[107282]18858.

[107283]Brailsford, J. F.: Chondro–osteo–dystrophy: roentgeno–
graphic and

[107284]clinical features of child with dislocation of vertebrae. Am.
J.

[107285]Surg. 7: 404–410, 1929.

[107286]18859.

[107287]Bunge, S.; Kleijer, W. J.; Tylki–Szymanska, A.; Steglich, C.;
Beck,

[107288]M.; Tomatsu, S.; Fukuda, S.; Poorthuis, B. J. H. M.; Czarto–
ryska,

[107289]B.; Orii, T.; Gal, A.: Identification of 31 novel mutations in
the

[107290]N–acetylgalactosamine–6–sulfatase gene reveals exces–
sive allelic heterogeneity

[107291]among patients with Morquio A syndrome. Hum. Mutat.
10: 223–232,

[107292]1997.

[107293]18860.

[107294]Engel, A. G.; Lambert, E. H.; Gomez, M. R.: A new myasthenic syndrome

[107295]with end-plate acetylcholinesterase deficiency, small nerve terminals,

[107296]and reduced acetylcholine release. Ann. Neurol. 1: 315-330, 1977.

[107297]18861.

[107298]Faiella, A.; Brunelli, S.; Granata, T.; D'Incerti, L.; Cardini,

[107299]R.; Lenti, C.; Battaglia, G.; Boncinelli, E.: A number of schizencephaly

[107300]patients including 2 brothers are heterozygous for germline mutations

[107301]in the homeobox gene EMX2. Europ. J. Hum. Genet. 5: 186-190, 1997.

[107302]18862.

[107303]Li, Y.-T.; Hirabayashi, Y.; Li, S.-C.: Differentiation of two

[107304]variants of type-AB Gm-2-gangliosidosis using chromogenic substrates. Am.

[107305]J. Hum. Genet. 35: 520-522, 1983.

[107306]18863.

- [107307]Rousseau-Merck, M.-F.; Duro, D.; Berger, R.; Thiesen, H. J.: Chromosomal
[107308]localization of two KOX zinc finger genes on chromosome
bands 7q21-q22. Ann.
[107309]Genet. 38: 81-84, 1995.
[107310]18864.
[107311]Tommerup, N.; Vissing, H.: Isolation and fine mapping of
16 novel
[107312]human zinc finger-encoding cDNAs identify putative can-
didate genes
[107313]for developmental and malignant disorders. Genomics 27:
259-264,
[107314]1995.
[107315]18865.
[107316]Canman, C. E.; Radany, E. H.; Parsels, L. A.; Davis, M. A.;
Lawrence,
[107317]T. S.; Maybaum, J.: Induction of resistance to fluo-
rodeoxyuridine
[107318]cytotoxicity and DNA damage in human tumor cells by
expression of
[107319]Escherichia coli deoxyuridinetriphosphatase. Cancer Res.
54: 2296-2298,

[107320]1994.

[107321]18866.

[107322]Canman, C. E.; Tang, H. Y.; Normolle, D. P.; Lawrence, T. S.; Maybaum,

[107323]J.: Variations in patterns of DNA damage induced in human colorectal

[107324]tumor cells by 5-fluorodeoxyuridine: implications for mechanisms of

[107325]resistance and cytotoxicity. Proc. Nat. Acad. Sci. 89: 10474-10478,

[107326]1992.

[107327]18867.

[107328]Chu, R.; Lin, Y.; Rao, M. S.; Reddy, J. K.: Cloning and identification

[107329]of rat deoxyuridine triphosphatase as an inhibitor of peroxisome proliferator-activated

[107330]receptor alpha. J. Biol. Chem. 271: 27670-27676, 1996.

[107331]18868.

[107332]Cohen, D.; Heng, H. H. Q.; Shi, X.-M.; McIntosh, E. M.; Tsui, L.-C.;

[107333]Pearlman, R. E.: Assignment of the human dUTPase gene

(DUT) to chromosome

[107334]15q15-q21.1 by fluorescence in situ hybridization. Genomics 40:

[107335]213-215, 1997.

[107336]18869.

[107337]el-Hajj, H. H.; Zhang, H.; Weiss, B.: Lethality of a dut (deoxyuridine

[107338]triphosphatase) mutation in Escherichia coli. J. Bacteriol. 170:

[107339]1069-1075, 1988.

[107340]18870.

[107341]Ladner, R. D.; Caradonna, S. J.: The human dUTPase gene encodes

[107342]both nuclear and mitochondrial isoforms: differential expression of

[107343]the isoforms and characterization of a cDNA encoding the mitochondrial

[107344]species. J. Biol. Chem. 272: 19072-19080, 1997.

[107345]18871.

[107346]Ladner, R. D.; McNulty, D. E.; Carr, S. A.; Roberts, G. D.; Caradonna,

[107347]S. J.: Characterization of distinct nuclear and mitochondrial forms
of human deoxyuridine triphosphate nucleotidohydrolase.
J. Biol.
Chem. 271: 7745–7751, 1996.
18872.
Lindahl, T.: DNA repair enzymes. Annu. Rev. Biochem. 51:
61–87,
1982.
18873.
McIntosh, E. M.; Ager, D. D.; Gadsden, M. H.; Haynes, R.
H.: Human
dUTP pyrophosphatase: cDNA sequence and potential biological importance
of the enzyme. Proc. Nat. Acad. Sci. 89: 8020–8024,
1992. Note:
Erratum: Proc. Nat. Acad. Sci.:90: 4328 only, 1993.
18874.
Boring, L.; Gosling, J.; Cleary, M.; Charo, I. F.: Decreased
lesion
formation in CCR2 –/– mice reveals a role for chemokines

in the initiation

[107361]of atherosclerosis. Nature 394: 894–897, 1998.

[107362]18875.

[107363]Charo, I. F.; Myers, S. J.; Herman, A.; Franci, C.; Connolly, A.

[107364]J.; Coughlin, S. R.: Molecular cloning and functional expression

[107365]of two monocyte chemoattractant protein 1 receptors reveals alternative

[107366]splicing of the carboxyl-terminal tails. Proc. Nat. Acad. Sci. 91:

[107367]2752–2756, 1994.

[107368]18876.

[107369]Sun, D.–S.; Chang, A. C.; Jenkins, N. A.; Gilbert, D. J.; Copeland,

[107370]N. G.; Chang, N.–C. A.: Identification, molecular characterization,

[107371]and chromosomal localization of the cDNA encoding a novel leucine

[107372]zipper motif-containing protein. Genomics 36: 54–62, 1996.

[107373]18877.

[107374]Barrett, T. E.; Savva, R.; Panayotou, G.; Barlow, T.; Brown, T.;

[107375]Jiricny, J.; Pearl, L. H.: Crystal structure of a G:T/U mismatch-specific

[107376]DNA glycosylase: mismatch recognition by complementary-strand interactions. Cell 92:

[107377]117-129, 1998.

[107378]18878.

[107379]De Gregorio, L.; Gallinari, P.; Gariboldi, M.; Manenti, G.; Pierotti,

[107380]M. A.; Jiricny, J.; Dragani, T. A.: Genetic mapping of thymine DNA

[107381]glycosylase (Tdg) gene and of one pseudogene in the mouse. Mammalian

[107382]Genome 7: 909-910, 1996.

[107383]18879.

[107384]Lindahl, T.: DNA repair enzymes. Ann. Rev. Biochem. 51: 61-87,

[107385]1982.

[107386]18880.

[107387]Neddermann, P.; Gallinari, P.; Lettieri, T.; Schmid, D.;

Truong,

[107388]O.; Hsuan, J. J.; Wiebauer, K.; Jiricny, J.: Cloning and expression

[107389]of human G/T mismatch-specific thymine-DNA glycosylase. J. Biol.

[107390]Chem. 271: 12767-12774, 1996.

[107391]18881.

[107392]Sard, L.; Tornielli, S.; Gallinari, P.; Minoletti, F.; Jiricny,

[107393]J.; Lettieri, T.; Pierotti, M. A.; Sozzi, G.; Radice, P.: Chromosomal

[107394]localizations and molecular analysis of TDG gene-related sequences. Genomics 44:

[107395]222-226, 1997.

[107396]18882.

[107397]DiMarco, S. P.; Glover, T. W.; Miller, D. E.; Reines, D.; Warren,

[107398]S. T.: Transcription elongation factor SII (TCEA) maps to human chromosome

[107399]3p22-p21.3. Genomics 36: 185-188, 1996.

[107400]18883.

[107401]Park, H.; Baek, K.; Jeon, C.; Agarwal, K.; Yoo, O.: Charac-

terization

[107402]of the gene encoding the human transcriptional elongation factor TFIIIS. Gene 139:

[107403]263–267, 1994.

[107404]18884.

[107405]Reines, D.: Nascent RNA cleavage by transcription elongation complexes.In:

[107406]Conaway, R. C.; Conaway, J. W. (eds.): Transcription: Mechanisms

[107407]and Regulation. New York: Raven Press 1994. Pp. 263–278.

[107408]18885.

[107409]Thomas, M. J.; Platas, A. A.; Hawley, D. K.: Transcriptional fidelity

[107410]and proofreading by RNA polymerase II. Cell 93: 627–637, 1998.

[107411]18886.

[107412]Applequist, S. E.; Selg, M.; Raman, C.; Jack, H.–M.: Cloning and

[107413]characterization of HUPF1, a human homolog of the Saccharomyces cerevisiae

[107414]nonsense mRNA–reducing UPF1 protein. Nucleic Acids

Res. 25: 814–821,

[107415]1997.

[107416]18887.

[107417]Medghalchi, S. M.; Frischmeyer, P. A.; Mendell, J. T.; Kelly, A.

[107418]G.; Lawler, A. M.; Dietz, H. C.: Rent1, a trans-effector of nonsense-mediated

[107419]mRNA decay, is essential for mammalian embryonic viability. Hum.

[107420]Molec. Genet. 10: 99–105, 2001.

[107421]18888.

[107422]Mendell, J. T.; ap Rhys, C. M. J.; Dietz, H. C.: Separable roles

[107423]for rent1/hUpf1 in altered splicing and decay of nonsense transcripts. Science 298:

[107424]419–371, 2002.

[107425]18889.

[107426]Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Dietz, H. C.

[107427]: Cloning and characterization of a human regulator of nonsense transcript

[107428]stability. (Abstract) Am. J. Hum. Genet. 59 (suppl.): A32

only,

[107429]1996.

[107430]18890.

[107431]Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Kendzior,
R.

[107432]J., Jr.; Dietz, H. C.: Mammalian orthologues of a yeast reg-
ulator

[107433]of nonsense transcript stability. Proc. Nat. Acad. Sci. 93:
10928-10932,

[107434]1996.

[107435]18891.

[107436]Sun, X.; Perlick, H. A.; Dietz, H. C.; Maquat, L. E.: A mu-
tated

[107437]human homologue to yeast Upf1 protein has a dominant-
negative effect

[107438]on the decay of nonsense-containing mRNAs in mam-
malian cells. Proc.

[107439]Nat. Acad. Sci. 95: 10009-10014, 1998.

[107440]18892.

[107441]Eng, C.; Myers, S. M.; Kogon, M. D.; Sanicola, M.; Hession,

C.;

- [107442]Cate, R. L.; Mulligan, L. M.: Genomic structure and chromosomal localization
[107443]of the human GDNFR- α gene. *Oncogene* 16: 597-601, 1998.
[107444]18893.
- [107445]GFR- α Nomenclature Committee: Nomenclature of GPI-linked
[107446]receptors for the GDNF ligand family. *Neuron* 19: 485 only, 1997.
[107447]18894.
- [107448]Gorodinsky, A.; Zimonjic, D. B.; Popescu, N. C.; Milbrandt, J.
[107449]: Assignment of the GDNF family receptor α -1 (GFRA1) to human
[107450]chromosome band 10q26 by in situ hybridization. *Cytogenet. Cell Genet.* 78:
[107451]289-290, 1997.
[107452]18895.
- [107453]Jing, S.; Wen, D.; Yu, Y.; Holst, P. L.; Luo, Y.; Fang, M.; Tamir,
[107454]R.; Antonio, L.; Hu, Z.; Cupples, R.; Louis, J.-C.; Hu, S.; Al-

trock,

[107455]B. W.; Fox, G. M.: GDNF-induced activation of the Ret protein tyrosine

[107456]kinase is mediated by GDNFR-alpha, a novel receptor for GDNF. Cell 85:

[107457]1113-1124, 1996.

[107458]18896.

[107459]Paratcha, G.; Ledda, F.; Baars, L.; Couplier, M.; Besset, V.; Anders,

[107460]J.; Scott, R.; Ibanez, C. F.: Released GFR-alpha-1 potentiates downstream

[107461]signaling, neuronal survival, and differentiation via a novel mechanism

[107462]of recruitment of c-Ret to lipid rafts. Neuron 29: 171-184, 2001.

[107463]18897.

[107464]Puliti, A.; Cinti, R.; Seri, M.; Ceccherini, I.; Romeo, G.: Assignment

[107465]of mouse Gfra1, the homologue of a new human HSCR candidate gene,

[107466]to the telomeric region of mouse chromosome 19. Cytogenet. Cell Genet. 78:

[107467]291–294, 1997.

[107468]18898.

[107469]Shefelbine, S. E.; Khorana, S.; Schultz, P. N.; Huang, E.;
Thobe,

[107470]N.; Hu, Z. J.; Fox, G. M.; Jing, S.; Cote, G. J.; Gagel, R. F.:
Mutational

[107471]analysis of the GDNF/RET–GDNFR–alpha signaling com-
plex in a kindred

[107472]with vesicoureteral reflux. Hum. Genet. 102: 474–478,
1998.

[107473]18899.

[107474]Bardelli, A.; Longati, P.; Albero, D.; Goruppi, S.; Schneider,

[107475]C.; Ponzetto, C.; Comoglio, P. M.: HGF receptor associates
with the

[107476]anti-apoptotic protein BAG–1 and prevents cell death.

EMBO J. 15:

[107477]6205–6212, 1996.

[107478]18900.

[107479]Takayama, S.; Kochel, K.; Irie, S.; Inazawa, J.; Abe, T.; Sato,

[107480]T.; Druck, T.; Huebner, K.; Reed, J. C.: Cloning of cDNAs
encoding

[107481]the human BAG1 protein and localization of the human
BAG1 gene to

[107482]chromosome 9p12. Genomics 35: 494–498, 1996.

[107483]18901.

[107484]Takayama, S.; Sato, T.; Krajewski, S.; Kochel, K.; Irie, S.;
Millan,

[107485]J. A.; Reed, J. C.: Cloning and functional analysis of BAG–1:
a novel

[107486]Bcl–2–binding protein with anti–cell death activity. Cell
80: 279–284,

[107487]1995.

[107488]18902.

[107489]Wang, H.–G.; Takayama, S.; Rapp, U. R.; Reed, J. C.: Bcl–2
interacting

[107490]protein, BAG–1, binds to and activates the kinase Raf–1.
Proc. Nat.

[107491]Acad. Sci. 93: 7063–7068, 1996.

[107492]18903.

[107493]Yang, X.; Pater, A.; Tang, S.–C.: Cloning and characteriza-
tion

[107494]of the human BAG–1 gene promoter: upregulation by tu–

mor-derived p53

[107495]mutants. *Oncogene* 18: 4546–4553, 1999.

[107496]18904.

[107497]Chen, H.; Chrast, R.; Rossier, C.; Gos, A.; Antonarakis, S.
E.;

[107498]Kudoh, J.; Yamaki, A.; Shindoh, N.; Maeda, H.; Minoshima,
S.; Shimizu,

[107499]N.: Single-minded and Down syndrome? (Letter) *Nature*
Genet. 10:

[107500]9–10, 1995.

[107501]18905.

[107502]Chrast, R.; Scott, H. S.; Madani, R.; Huber, L.; Wolfer, D. P.;

[107503]Prinz, M.; Aguzzi, A.; Lipp, H.-P.; Antonarakis, S. E.: Mice
trisomic

[107504]for a bacterial artificial chromosome with the single-
minded 2 gene

[107505](Sim2) show phenotypes similar to some of those present
in the partial

[107506]trisomy 16 mouse models of Down syndrome. *Hum.*
Molec. Genet. 9:

[107507]1853–1864, 2000.

[107508]18906.

[107509]Dahmane, N.; Charron, G.; Lopes, C.; Yaspo, M.-L.; Maunoury, C.;

[107510]Decorte, L.; Sinet, P.-M.; Bloch, B.; Delabar, J.-M.: Down syndrome-critical

[107511]region contains a gene homologous to *Drosophila* sim expressed during

[107512]rat and human central nervous system development. Proc. Nat. Acad.

[107513]Sci. 92: 9191-9195, 1995.

[107514]18907.

[107515]Ema, M.; Ikegami, S.; Hosoya, T.; Mimura, J.; Ohtani, H.; Nakao,

[107516]K.; Inokuchi, K.; Katsuki, M.; Fujii-Kuriyama, Y.: Mild impairment

[107517]of learning and memory in mice overexpressing the mSim2 gene located

[107518]on chromosome 16: an animal model of Down's syndrome. Hum. Molec.

[107519]Genet. 8: 1409-1415, 1999.

[107520]18908.

- [107521]Moffett, P.; Dayo, M.; Reece, M.; McCormick, M. K.; Pelletier,
J.: Characterization of msim, a murine homologue of the
Drosophila
sim transcription factor. Genomics 35: 144–155, 1996.
[107524]18909.
- [107525]Muenke, M.; Bone, L. J.; Mitchell, H. F.; Hart, I.; Walton, K.;
Hall-Johnson, K.; Ippel, E. F.; Dietz-Band, J.; Kvaloy, K.;
Fan, C.-M.;
Tessier-Lavigne, M.; Patterson, D.: Physical mapping of
the holoprosencephaly
critical region in 21q22.3, exclusion of SIM2 as a candidate gene
for holoprosencephaly, and mapping of SIM2 to a region
of chromosome
21 important for Down syndrome. Am. J. Hum. Genet. 57:
1074–1079,
[107531]1995.
[107532]18910.
- [107533]Yamaki, A.; Noda, S.; Kudoh, J.; Shindoh, N.; Maeda, H.;
Minoshima,

[107534]S.; Kawasaki, K.; Shimizu, Y.; Shimizu, N.: The mammalian single-minded

[107535](SIM) gene: mouse cDNA structure and diencephalic expression indicate

[107536]a candidate gene for Down syndrome. Genomics 35: 136–143, 1996.

[107537]18911.

[107538]de Winter, J. P.; Leveille, F.; van Berkel, C. G. M.; Rooimans,

[107539]M. A.; van der Weel, L.; Steltenpool, J.; Demuth, I.; Morgan, N. V.;

[107540]Alon, N.; Bosnoyan–Collins, L.; Lightfoot, J.; Leegwater, P. A.; Waisfisz,

[107541]Q.; Komatsu, K.; Arwert, F.; Pronk, J. C.; Mathew, C. G.; Digweed,

[107542]M.; Buchwald, M.; Joenje, H.: Isolation of a cDNA representing the

[107543]Fanconi anemia complementation group E gene. Am. J. Hum. Genet. 67:

[107544]1306–1308, 2000. Note: Erratum: Am. J. Hum. Genet. 67: 1365 only,

[107545]2000.

[107546]18912.

[107547]Hirotsume, S.; Fleck, M. W.; Gambello, M. J.; Bix, G. J.;
Chen,

[107548]A.; Clark, G. D.; Ledbetter, D. H.; McBain, C. J.; Wynshaw-
Boris,

[107549]A.: Graded reduction of Pafah1b1 (Lis1) activity results in
neuronal

[107550]migration defects and early embryonic lethality. *Nature*
Genet. 19:

[107551]333–339, 1998.

[107552]18913.

[107553]Kurahashi, H.; Sakamoto, M.; Ono, J.; Honda, A.; Okada,
S.; Nakamura,

[107554]Y.: Molecular cloning of the chromosomal breakpoint in
the LIS1 gene

[107555]of a patient with isolated lissencephaly and balanced
t(8;17). *Hum.*

[107556]*Genet.* 103: 189–192, 1998.

[107557]18914.

[107558]Liu, Z.; Steward, R.; Luo, L.: *Drosophila* Lis1 is required for

[107559]neuroblast proliferation, dendritic elaboration and axonal

transport. Nature

[107560]Cell Biol. 2: 776–783, 2000.

[107561]18915.

[107562]Lo Nigro, C.; Chong, S. S.; Smith, A. C. M.; Dobyns, W. B.;
Carrozzo,

[107563]R.; Ledbetter, D. H.: Point mutations and an intragenic
deletion

[107564]in LIS1, the lissencephaly causative gene in isolated
lissencephaly

[107565]sequence and Miller–Dieker syndrome. Hum. Molec.
Genet. 6: 157–164,

[107566]1997.

[107567]18916.

[107568]Reiner, O.; Bar–Am, I.; Sapir, T.; Shmueli, O.; Carrozzo, R.;

[107569]Lindsay, E. A.; Baldini, A.; Ledbetter, D. H.; Cahana, A.:
LIS2,

[107570]gene and pseudogene, homologous to LIS1 (lissencephaly
1), located

[107571]on the short and long arms of chromosome 2. Genomics
30: 251–256,

[107572]1995.

[107573]18917.

[107574]Sakamoto, M.; Ono, J.; Okada, S.; Masuno, M.; Nakamura, Y.; Kurahashi,

[107575]H.: Alteration of the LIS1 gene in Japanese patients with isolated

[107576]lissencephaly sequence or Miller–Dieker syndrome. Hum. Genet. 103:

[107577]586–589, 1998.

[107578]18918.

[107579]Smith, D. S.; Niethammer, M.; Ayala, R.; Zhou, Y.; Gambello, M.

[107580]J.; Wynshaw–Boris, A.; Tsai, L.–H.: Regulation of cytoplasmic dynein

[107581]behaviour and microtubule organization by mammalian Lis1. Nature

[107582]Cell Biol. 2: 767–775, 2000.

[107583]18919.

[107584]Sosa–Pineda, B.; Wigle, J. T.; Oliver, G.: Hepatocyte migration

[107585]during liver development requires Prox1. Nature Genet. 25: 254–255,

[107586]2000.

[107587]18920.

[107588]Wigle, J. T.; Chowdhury, K.; Gruss, P.; Oliver, G.: Prox1
function

[107589]is crucial for mouse lens-fibre elongation. Nature Genet.
21: 318-322,

[107590]1999.

[107591]18921.

[107592]Wigle, J. T.; Oliver, G.: Prox1 function is required for the
development

[107593]of the murine lymphatic system. Cell 98: 769-778, 1999.

[107594]18922.

[107595]Zinovieva, R. D.; Duncan, M. K.; Johnson, T. R.; Torres, R.;
Polymeropoulos,

[107596]M. H.; Tomarev, S. I.: Structure and chromosomal localization of
tion of

[107597]the human homeobox gene Prox 1. Genomics 35:
517-522, 1996.

[107598]18923.

[107599]Ladenburger, E. M.; Fackelmayer, F. O.; Hameister, H.;
Knippers,

- [107600]R.: MCM4 and PRKDC, human genes encoding proteins
MCM4 and DNA-PKcs,
[107601]are close neighbours located on chromosome 8q12-q13.
Cytogenet. Cell
[107602]Genet. 77: 268-270, 1997.
[107603]18924.
- [107604]Ma, Y.; Pannicke, U.; Schwarz, K.; Lieber, M. R.: Hairpin
opening
[107605]and overhang processing by an Artemis/DNA-dependent
protein kinase
[107606]complex in nonhomologous end joining and V(D)J recom-
bination. Cell 108:
[107607]781-794, 2002.
[107608]18925.
- [107609]Satoh, T.; Tsuruga, H.; Yabuta, N.; Ishidate, M., Jr.; No-
jima,
[107610]H.: Assignment of the human CDC21 (MCM4) gene to
chromosome 8q11.2. Genomics 46:
[107611]525-526, 1997.
[107612]18926.
- [107613]Barresi, R.; Di Blasi, C.; Negri, T.; Brugnoli, R.; Vitali, A.;

- [107614]Felisari, G.; Salandi, A.; Daniel, S.; Cornelio, F.; Morandi, L.;
- [107615]Mora, M.: Disruption of heart sarcoglycan complex and severe cardiomyopathy
- [107616]caused by beta sarcoglycan mutations. J. Med. Genet. 37: 102–107,
- [107617]2000.
- [107618]18927.
- [107619]Bonnemann, C. G.; Modi, R.; Noguchi, S.; Mizuno, Y.; Yoshida, M.;
- [107620]Gussoni, E.; McNally, E. M.; Duggan, D. J.; Angelini, C.; Hoffman,
- [107621]E. P.; Ozawa, E.; Kunkel, L. M.: Beta-sarcoglycan (A3b) mutations
- [107622]cause autosomal recessive muscular dystrophy with loss of the sarcoglycan
- [107623]complex. Nature Genet. 11: 266–273, 1995. Note: Erratum: Nature
- [107624]Genet. 12: 110 only, 1996.
- [107625]18928.
- [107626]Bonnemann, C. G.; Passos-Bueno, M. R.; McNally, E. M.; Vainzof,

- [107627]M.; Moreira, E. S.; Marie, S. K.; Pavanello, R. C. M.;
Noguchi, S.;
- [107628]Ozawa, E.; Zatz, M.; Kunkel, L. M.: Genomic screening for
beta-sarcoglycan
- [107629]gene mutations: missense mutations may cause severe
limb-girdle muscular
- [107630]dystrophy type 2E (LGMD 2E). Hum. Molec. Genet. 5:
1953-1961, 1996.
- [107631]18929.
- [107632]Bonnemann, C. G.; Wong, J.; Ben Hamida, C.; Ben Hamida,
M.; Hentati,
- [107633]F.; Kunkel, L. M.: LGMD 2E in Tunisia is caused by a ho-
mozygous missense
- [107634]mutation in beta-sarcoglycan exon 3. Neuromusc. Dis. 8:
193-197,
- [107635]1998.
- [107636]18930.
- [107637]Durbeej, M.; Cohn, R. D.; Hrstka, R. F.; Moore, S. A.; Alla-
mand,
- [107638]V.; Davidson, B. L.; Williamson, R. A.; Campbell, K. P.: Dis-
ruption
- [107639]of the beta-sarcoglycan gene reveals pathogenetic com-

plexity of limb-girdle

[107640]muscular dystrophy type 2E. Molec. Cell 5: 141–151,
2000.

[107641]18931.

[107642]Joenje, H.; Lo Ten Foe, J. R.; Oostra, A. B.; van Berkel, C. G.

[107643]M.; Rooimans, M. A.; Schroeder-Kurth, T.; Wegner, R.-D.;
Gille, J.

[107644]J. P.; Buchwald, M.; Arwert, F.: Classification of Fanconi
anemia

[107645]patients by complementation analysis: evidence for a fifth
genetic

[107646]subtype. Blood 86: 2156–2160, 1995.

[107647]18932.

[107648]Fielder, P. J.; Guevara-Aguirre, J.; Rosenbloom, A. L.;
Carlsson,

[107649]L.; Hintz, R. L.; Rosenfeld, R. G.: Expression of serum in-
sulin-like

[107650]growth factors, insulin-like growth factor-binding pro-
teins, and the

[107651]growth hormone-binding protein in heterozygote relatives
of Ecuadorian

[107652]growth hormone receptor deficient patients. J. Clin. En-

docr. Metab. 74:

[107653]743–750, 1992.

[107654]18933.

[107655]Fisker, S.; Kristensen, K.; Rosenfalck, A. M.; Pedersen, S.
B.;

[107656]Ebdrup, L.; Richelsen, B.; Hilsted, J.; Christiansen, J. S.;
Jorgensen,

[107657]J. O. L.: Gene expression of a truncated and the full-
length growth

[107658]hormone (GH) receptor in subcutaneous fat and skeletal
muscle in GH-deficient

[107659]adults: impact of GH treatment. J. Clin. Endocr. Metab. 86:
792–796,

[107660]2001.

[107661]18934.

[107662]Gastier, J. M.; Berg, M. A.; Vesterhus, P.; Reiter, E. O.;
Francke,

[107663]U.: Diverse deletions in the growth hormone receptor
gene cause growth

[107664]hormone insensitivity syndrome. Hum. Mutat. 16:
323–333, 2000.

[107665]18935.

- [107666] Godowski, P. J.; Leung, D. W.; Meacham, L. R.; Galgani, J. P.;
- [107667] Hellmiss, R.; Keret, R.; Rotwein, P. S.; Parks, J. S.; Laron, Z.;
- [107668] Wood, W. I.: Characterization of the human growth hormone receptor gene and demonstration of a partial gene deletion in two patients
- [107669] with Laron-type dwarfism. Proc. Nat. Acad. Sci. 86: 8083–8087, 1989.
- [107671] 18936.
- [107672] Iida, K.; Takahashi, Y.; Kaji, H.; Nose, O.; Okimura, Y.; Abe,
- [107673] H.; Chihara, K.: Growth hormone (GH) insensitivity syndrome with
- [107674] high serum GH-binding protein levels caused by a heterozygous splice
- [107675] site mutation of the GH receptor gene producing a lack of intracellular
- [107676] domain. J. Clin. Endocr. Metab. 83: 531–537, 1998.
- [107677] 18937.
- [107678] Iida, K.; Takahashi, Y.; Kaji, H.; Takahashi, M. O.; Okimura,

[107679]Y.; Nose, O.; Abe, H.; Chihara, K.: Functional characterization of
 truncated growth hormone (GH) receptor-(1-277) causing
 partial GH
 insensitivity syndrome with high GH-binding protein. J.
 Clin. Endocr.
 Metab. 84: 1011-1016, 1999.
 18938.

[107684]Jorge, A. A.; Souza, S. C.; Arnhold, I. J.; Mendonca, B. B.:
 Poor reproducibility of IGF-I and IGF binding protein-3
 generation
 test in children with short stature and normal coding re-
 gion of the
 GH receptor gene. J. Clin. Endocr. Metab. 87: 469-472,
 2002.
 18939.

[107689]Kaji, H.; Nose, O.; Tajiri, H.; Takahashi, Y.; Iida, K.; Takahashi,
 hashi,
 T.; Okimura, Y.; Abe, H.; Chihara, K.: Novel compound
 heterozygous
 mutations of growth hormone (GH) receptor gene in a pa-
 tient with GH

[107692]insensitivity syndrome. J. Clin. Endocr. Metab. 82:
3705–3709, 1997.

[107693]18940.

[107694]Kranzler, J. H.; Rosenbloom, A. L.; Martinez, V.; Guevara-
Aguirre,

[107695]J.: Normal intelligence with severe insulin-like growth fac-
tor I

[107696]deficiency due to growth hormone receptor deficiency: a
controlled

[107697]study in a genetically homogeneous population. J. Clin.
Endocr.

[107698]Metab. 83: 1953–1958, 1998.

[107699]18941.

[107700]Laron, Z.; Klinger, B.; Erster, B.; Silbergeld, A.: Serum GH

[107701]binding protein activities identifies the heterozygous car-
riers for

[107702]Laron type dwarfism. Acta Endocr. 121: 603–608, 1989.

[107703]18942.

[107704]Leung, D. W.; Spencer, S. A.; Cachianes, G.; Hammonds, R.
G.;

[107705]Collins, C.; Henzel, W. J.; Barnard, R.; Waters, M. J.; Wood,

W. I.

[107706]: Growth hormone receptor and serum binding protein:
purification,

[107707]cloning and expression. Nature 330: 537-543, 1987.

[107708]18943.

[107709]Marzullo, P.; Buckway, C.; Pratt, K. L.; Colao, A.; Guevara-
Aguirre,

[107710]J.; Rosenfeld, R. G.: Leptin concentrations in GH defi-
ciency: the

[107711]effect of GH insensitivity. J. Clin. Endocr. Metab. 87:
540-545,

[107712]2002.

[107713]18944.

[107714]Menon, R. K.; Cheng, H.; Singh, M.: Identification and
characterization

[107715]of single strand DNA-binding protein that represses
growth hormone

[107716]receptor gene expression. Molec. Endocr. 11: 1291-1304,
1997.

[107717]18945.

[107718]Metherell, L. A.; Akker, S. A.; Munroe, P. B.; Rose, S. J.;

Caulfield,

[107719]M.; Savage, M. O.; Chew, S. L.; Clark, A. J. L.: Pseudoexon activation

[107720]as a novel mechanism for disease resulting in atypical growth-hormone

[107721]insensitivity. Am. J. Hum. Genet. 69: 641-646, 2001.

[107722]18946.

[107723]Rosenbloom, A. L.; Guevara-Aguirre, J.; Berg, M. A.; Francke,

[107724]U.: Stature in Ecuadorians heterozygous for growth hormone receptor

[107725]gene E180 splice mutation does not differ from that of homozygous

[107726]normal relatives. J. Clin. Endocr. Metab. 83: 2373-2375, 1998.

[107727]18947.

[107728]Sanchez, J. E.; Perera, E.; Baumbach, L.; Cleveland, W. W.: Growth

[107729]hormone receptor mutations in children with idiopathic short stature. J.

[107730]Clin. Endocr. Metab. 83: 4079-4083, 1998.

[107731]18948.

- [107732]Shen, X. Y.; Holt, R. I. G.; Miell, J. P.; Justice, S.; Portmann,
[107733]B.; Postel-Vinay, M.-C.; Ross, R. J. M.: Cirrhotic liver ex-
presses
[107734]low levels of the full-length and truncated growth hor-
mone receptors. J.
[107735]Clin. Endocr. Metab. 83: 2532-2538, 1998.
[107736]18949.
- [107737]Spencer, S. A.; Hammonds, R. G.; Henzel, W. J.; Rodriguez,
H.;
[107738]Waters, M. J.; Wood, W. I.: Rabbit liver growth hormone re-
ceptor
[107739]and serum binding protein: purification, characterization,
and sequence. J.
[107740]Biol. Chem. 263: 7862-7867, 1988.
[107741]18950.
- [107742]Walker, J. L.; Crock, P. A.; Behncken, S. N.; Rowlinson, S.
W.;
[107743]Nicholson, L. M.; Boulton, T. J. C.; Waters, M. J.: A novel
mutation
[107744]affecting the interdomain link region of the growth hor-
mone receptor

[107745]in a Vietnamese girl, and response to long-term treatment with recombinant

[107746]human insulin-like growth factor-I and luteinizing hormone-releasing

[107747]hormone analogue. J. Clin. Endocr. Metab. 83: 2554-2561, 1998.

[107748]18951.

[107749]Denier, C.; Ducros, A.; Durr, A.; Eymard, B.; Chassande, B.; Tournier-Lasserre,

[107750]E.: Missense CACNA1A mutation causing episodic ataxia type 2. Arch.

[107751]Neurol. 58: 292-295, 2001.

[107752]18952.

[107753]Byk, T.; Dobransky, T.; Cifuentes-Diaz, C.; Sobel, A.: Identification

[107754]and molecular characterization of Unc-33-like phosphoprotein (Ulip),

[107755]a putative mammalian homolog of the axonal guidance-associated unc-33

[107756]gene product. J. Neurosci. 16: 688-701, 1996.

[107757]18953.

- [107758]Choe, H.; Farzan, M.; Sun, Y.; Sullivan, N.; Rollins, B.;
Ponath,
[107759]P. D.; Wu, L.; Mackay, C. R.; LaRosa, G.; Newman, W.; Ger-
ard, N.;
[107760]Gerard, C.; Sodroski, J.: The beta-chemokine receptors
CCR3 and CCR5
[107761]facilitate infection by primary HIV-1 isolates. Cell 85:
1135-1148,
[107762]1996.
[107763]18954.
[107764]Diriong, S.; Lory, P.; Williams, M. E.; Ellis, S. B.; Harpold,
[107765]M. M.; Taviaux, S.: Chromosomal localization of the hu-
man genes for
[107766]alpha-1A, alpha-1B, and alpha-1E voltage-dependent
Ca(2+) channel
[107767]subunits. Genomics 30: 605-609, 1995.
[107768]18955.
[107769]Ducros, A.; Denier, C.; Joutel, A.; Cecillon, M.; Lescoat, C.;
[107770]Vahedi, K.; Darcel, F.; Vicaut, E.; Bousser, M.-G.; Tournier-
Lasserre,
[107771]E.: The clinical spectrum of familial hemiplegic migraine

associated

[107772]with mutations in a neuronal calcium channel. New Eng. J.
Med. 345:

[107773]17–24, 2001.

[107774]18956.

[107775]Dunlap, K.; Luebke, J. I.; Turner, T. J.: Exocytotic Ca(2+)
channels

[107776]in mammalian central neurons. Trends Neurosci. 18:
89–98, 1995.

[107777]18957.

[107778]Fletcher, C. F.; Lutz, C. M.; O'Sullivan, T. N.; Shaughnessy,
J.

[107779]D, Jr.; Hawkes, R.; Frankel, W. N.; Copeland, N. G.; Jenkins,
N. A.

[107780]: Absence epilepsy in tottering mutant mice is associated
with calcium

[107781]channel defects. Cell 87: 607–617, 1996.

[107782]18958.

[107783]Friend, K. L.; Crimmins, D.; Phan, T. G.; Sue, C. M.; Colley,
A.;

[107784]Fung, V. S. C.; Morris, J. G. L.; Sutherland, G. R.; Richards,

R.

[107785]J.: Detection of a novel missense mutation and second re-
current mutation

[107786]in the CACNA1A gene in individuals with EA-2 and FHM.

Hum. Genet. 105:

[107787]261–265, 1999.

[107788]18959.

[107789]Hess, E. J.: Migraines in mice? Cell 87: 1149–1151, 1996.

[107790]18960.

[107791]Hoffman, E. P.; Gardner, K.: Ion channels--molecular di-
vining

[107792]rods hit their clinical mark. New Eng. J. Med. (Editorial)

336:

[107793]1599–1600, 1997.

[107794]18961.

[107795]Jodice, C.; Mantuano, E.; Veneziano, L.; Trettel, F.; Sabba-
dini,

[107796]G.; Calandriello, L.; Francia, A.; Spadaro, M.; Pierelli, F.;
Salvi,

[107797]F.; Ophoff, R. A.; Frants, R. R.; Frontali, M.: Episodic ataxia
type

[107798]2 (EA2) and spinocerebellar ataxia type 6 (SCA6) due to
CAG repeat

[107799]expansion in the CACNA1A gene on chromosome 19p.
Hum. Molec. Genet. 6:

[107800]1973–1978, 1997.

[107801]18962.

[107802]Kim, J.-S.; Yue, Q.; Jen, J. C.; Nelson, S. F.; Baloh, R. W.:

[107803]Familial migraine with vertigo: no mutations found in
CACNA1A. Am.

[107804]J. Med. Genet. 79: 148–151, 1998.

[107805]18963.

[107806]Kraus, R. L.; Sinnegger, M. J.; Glossmann, H.; Hering, S.;
Striessnig,

[107807]J.: Familial hemiplegic migraine mutations change al-
pha(1A)Ca(2+)

[107808]channel kinetics. J. Biol. Chem. 273: 5586–5590, 1998.

[107809]18964.

[107810]Guida, S.; Trettel, F.; Pagnutti, S.; Mantuano, E.; Tottene,
A.;

[107811]Veneziano, L.; Fellin, T.; Spadaro, M.; Stauderman, K. A.;
Williams,

[107812]M. E.; Volsen, S.; Ophoff, R. A.; Frants, R. R.; Jodice, C.;
Frontali,
[107813]M.; Pietrobon, D.: Complete loss of P/Q calcium channel
activity
[107814]caused by a CACNA1A missense mutation carried by pa-
tients with episodic
[107815]ataxia type 2. Am. J. Hum. Genet. 68: 759–764, 2001.
[107816]18965.
[107817]Ishikawa, K.; Tanaka, H.; Saito, M.; Ohkoshi, N.; Fujita, T.;
[107818]Yoshizawa, K.; Ikeuchi, T.; Watanabe, M.; Hayashi, A.;
Takiyama, Y.;
[107819]Nishizawa, M.; Nakano, I.; Matsubayashi, K.; Miwa, M.;
Shoji, S.;
[107820]Kanazawa, I.; Tsuji, S.; Mizusawa, H.: Japanese families
with autosomal
[107821]dominant pure cerebellar ataxia map to chromosome
19p13.1–p13.2 and
[107822]are strongly associated with mild CAG expansions in the
spinocerebellar
[107823]ataxia type 6 gene in chromosome 19p13.1. Am J. Hum.
Genet. 61:
[107824]336–346, 1997.

[107825]18966.

[107826]Kors, E. E.; Terwindt, G. M.; Vermeulen, F. L. M. G.; Fitzsimons,

[107827]R. B.; Jardine, P. E.; Heywood, P.; Love, S.; van den Maagdenberg,

[107828]A. M. J. M.; Haan, J.; Frants, R. R.; Ferrari, M. D.: Delayed cerebral

[107829]edema and fatal coma after minor head trauma: role of the CACNA1A

[107830]calcium channel subunit gene and relationship with familial hemiplegic

[107831]migraine. Ann. Neurol. 49: 753–760, 2001.

[107832]18967.

[107833]Kraus, R. L.; Sinnegger, M. J.; Koschak, A.; Glossmann, H.; Stenirri,

[107834]S.; Carrera, P.; Striessnig, J.: Three new familial hemiplegic migraine

[107835]mutants affect P/Q-type Ca^{2+} channel kinetics. J. Biol. Chem. 275:

[107836]9239–9243, 2000.

[107837]18968.

- [107838]Lehmann-Horn, F.; Jurkat-Rott, K.: Voltage-gated ion channels
[107839]and hereditary disease. *Physiol. Rev.* 79: 1317–1372, 1999.
[107840]18969.
- [107841]Matsuyama, Z.; Kawakami, H.; Maruyama, H.; Izumi, Y.; Komure,
[107842]O.; Udaoka, F.; Kameyama, M.; Nishio, T.; Kuroda, Y.; Nishimura, M.;
[107843]Nakamura, S.: Molecular features of the CAG repeats of spinocerebellar
[107844]ataxia 6 (SCA6). *Hum. Molec. Genet.* 6: 1283–1287, 1997.
[107845]18970.
- [107846]Riess, O.; Schols, L.; Bottger, H.; Nolte, D.; Viera-Saecker, A. M. M.; Schimming, C.; Kreuz, F.; Macek, M., Jr.; Krebsova, A.;
[107848]Macek, M., Sr.; Klockgether, T.; Zuhlke, C.; Laccone, F. A.: SCA6
[107849]is caused by moderate CAG expansion in the α 1A-voltage-dependent
[107850]calcium channel gene. *Hum. Molec. Genet.* 6: 1289–1293, 1997.

[107851]18971.

[107852]Sasaki, H.; Kojima, H.; Yabe, I.; Tashiro, K.; Hamada, T.;
Sawa,

[107853]H.; Hiraga, H.; Nagashima, K.: Neuropathological and
molecular studies

[107854]of spinocerebellar ataxia type 6 (SCA6). Acta. Neuropath.
95: 199–204,

[107855]1998.

[107856]18972.

[107857]Scoggan, K. A.; Chandra, T.; Nelson, R.; Hahn, A. F.; Bul-
man,

[107858]D. E.: Identification of two novel mutations in the
CACNA1A gene

[107859]responsible for episodic ataxia type 2. (Letter) J. Med.
Genet. 38:

[107860]249–253, 2001.

[107861]18973.

[107862]Thibault, O.; Landfield, P. W.: Increase in single L-type
calcium

[107863]channels in hippocampal neurons during aging. Science
272: 1017–1019,

[107864]1996.

[107865]18974.

[107866]Toru, S.; Murakoshi, T.; Ishikawa, K.; Saegusa, H.; Fuji-
gasaki,

[107867]H.; Uchihara, T.; Nagayama, S.; Osanai, M.; Mizusawa, H.;
Tanabe,

[107868]T.: Spinocerebellar ataxia type 6 mutation alters P-type
calcium

[107869]channel function. J. Biol. Chem. 275: 10893–10898, 2000.

[107870]18975.

[107871]Trettel, F.; Mantuano, E.; Calabresi, V.; Veneziano, L.;
Olsen,

[107872]A. S.; Georgescu, A.; Gordon, L.; Sabbadini, G.; Frontali,
M.; Jodice,

[107873]C.: A fine physical map of the CACNA1A gene region on
19p13.1–p13.2

[107874]chromosome. Gene 241: 45–50, 2000.

[107875]18976.

[107876]Vahedi, K.; Denier, C.; Ducros, A.; Bousson, V.; Levy, C.;
Chabriat,

[107877]H.; Haguenau, M.; Tournier-Lasserre, E.; Bousser, M. G.:

CACNA1A

[107878] gene de novo mutation causing hemiplegic migraine, coma, and cerebellar

[107879] atrophy. *Neurology* 55: 1040–1042, 2000.

[107880] 18977.

[107881] Ben-Porath, I.; Benvenisty, N.: Characterization of a tumor-associated

[107882] gene, a member of a novel family of genes encoding membrane glycoproteins. *Gene* 183:

[107883] 69–75, 1996.

[107884] 18978.

[107885] Bosch, E. P.; Murphy, M. J.; Cancilla, P. A.: Peripheral neurofibromatosis

[107886] and peroneal muscular atrophy. *Neurology* 31: 1408–1414, 1981.

[107887] 18979.

[107888] Bridges, C. B.: The Bar 'gene' a duplication. *Science* 83: 210–211,

[107889] 1936.

[107890] 18980.

[107891] Gabreels-Festen, A. A. W. M.; Joosten, E. M. G.; Gabreels,

F. J.

[107892]M.; Jennekens, F. G. I.; Janssen-van Kempen, T. W.: Early morphological

[107893]features in dominantly inherited demyelinating motor and sensory neuropathy

[107894](HMSN type I). J. Neurol. Sci. 107: 145–154, 1992.

[107895]18981.

[107896]Green, M. C.: Genetic variants and strains of the laboratory mouse.

[107897]New York: Oxford Univ. Press (pub.) 1989.

[107898]18982.

[107899]Hartl, D. L.: The most unkindest cut of all. Nature Genet. 12:

[107900]227–229, 1996.

[107901]18983.

[107902]Hoogendijk, J. E.; Hensels, G. W.; Zorn, I.; Valentijn, L.; Janssen,

[107903]E. A. M.; de Visser, M.; Barker, D. F.; Ongerboer de Visser, B. W.;

[107904]Baas, F.; Bolhuis, P. A.: The duplication in Charcot-Marie-Tooth

[107905]disease type 1a spans at least 1100 kb on chromosome
17p11.2. Hum.

[107906]Genet. 88: 215–218, 1991.

[107907]18984.

[107908]Hoogendijk, J. E.; Janssen, E. A. M.; Gabreels–Festen, A. A.
W.

[107909]M.; Hensels, G. W.; Joosten, E. M. G.; Gabreels, F. J. M.;
Zorn, I.;

[107910]Valentijn, L. J.; Baas, F.; Ongerboer de Visser, B. W.; de
Visser,

[107911]M.; Bolhuis, P. A.: Allelic heterogeneity in hereditary motor
and

[107912]sensory neuropathy type 1a (Charcot–Marie–Tooth disease
type 1a). Neurology 43:

[107913]1010–1015, 1993.

[107914]18985.

[107915]Ikegami, T.; Ikeda, H.; Aoyama, M.; Matsuki, T.; Imota, T.;
Fukuuchi,

[107916]Y.; Amano, T.; Toyoshima, I.; Ishihara, Y.; Endoh, H.;
Hayasaka, K.

[107917]: Novel mutations of the peripheral myelin protein 22
gene in two

[107918]pedigrees with Dejerine–Sottas disease. Hum. Genet. 102:
294–298,
[107919]1998.
[107920]18986.
[107921]Ikegami, T.; Ikeda, H.; Chance, P. F.; Kiyosawa, H.; Ya-
mamoto,
[107922]M.; Sobue, G.; Ohnishi, A.; Tachi, N.; Hayasaka, K.: Facili-
tated
[107923]diagnosis of CMT1A duplication in chromosome
17p11.2–12: analysis
[107924]with a CMT1A–REP repeat probe and photostimulated lu-
minescence imaging. Hum.
[107925]Mutat. 9: 563–566, 1997.
[107926]18987.
[107927]Ionasescu, V. V.; Searby, C.; Greenberg, S. A.: Dejerine–Sot-
tas
[107928]disease with sensorineural hearing loss, nystagmus, and
peripheral
[107929]facial nerve weakness: de novo dominant point mutation
of the PMP22
[107930]gene. J. Med. Genet. 33: 1048–1049, 1996.

[107931]18988.

[107932]Isaacs, A. M.; Davies, K. E.; Hunter, A. J.; Nolan, P. M.; Vi-
zor,

[107933]L.; Peters, J.; Gale, D. G.; Kelsell, D. P.; Latham, I. D.;
Chase,

[107934]J. M.; Fisher, E. M. C.; Bouzyk, M. M.; and 11 others: Iden-
tification

[107935]of two new Pmp22 mouse mutants using large-scale mu-
tagenesis and a

[107936]novel rapid mapping strategy. Hum. Molec. Genet. 9:
1865-1871, 2000.

[107937]18989.

[107938]King, P. H.; Waldrop, R.; Lupski, J. R.; Shaffer, L. G.: Char-
cot-Marie-Tooth

[107939]phenotype produced by a duplicated PMP22 gene as part
of a 17p trisomy-translocation

[107940]to the X chromosome. Clin. Genet. 54: 413-416, 1998.

[107941]18990.

[107942]Kousseff, B. G.; Hadro, T. A.; Treiber, D. L.; Wollner, T.;
Morris,

[107943]C.: Charcot-Marie-Tooth disease with sensorineural hear-
ing loss--an

- [107944]autosomal dominant trait. Birth Defects Orig. Art. Ser. 18:
223–228,
[107945]1982.
- [107946]18991.
- [107947]Kovach, M. J.; Lin, J.–P.; Boyadjiev, S.; Campbell, K.;
Mazzeo,
[107948]L.; Herman, K.; Rimer, L. A.; Frank, W.; Llewellyn, B.; Jabs,
E. W.;
- [107949]Gelber, D.; Kimonis, V. E.: A unique point mutation in the
PMP22
[107950]gene is associated with Charcot–Marie–Tooth disease and
deafness. Am.
[107951]J. Hum. Genet. 64: 1580–1593, 1999.
- [107952]18992.
- [107953]Liehr, T.; Rautenstrauss, B.; Grehl, H.; Bathke, K. D.; Ekici,
[107954]A.; Rauch, A.; Rott, H.–D.: Mosaicism for the Charcot–
Marie–Tooth
[107955]disease type 1A duplication suggests somatic reversion.
Hum. Genet. 98:
[107956]22–28, 1996.
- [107957]18993.

[107958]Gaetano, C; Matsuo, T.; Thiele, C. J.: Identification and
characterization
[107959]of a retinoic acid-regulated human homologue of the
unc-33-like phosphoprotein
[107960]gene (hUlip) from neuroblastoma cells. J. Biol. Chem. 272:
12195-12201,
[107961]1997.
[107962]18994.
[107963]Matsuo, T.; Stauffer, J. K.; Walker, R. L.; Meltzer, P.; Thiele,
[107964]C. J.: Structure and promoter analysis of the human unc-
33-like phosphoprotein
[107965]gene: E-box required for maximal expression in neurob-
lastoma and myoblasts. J.
[107966]Biol. Chem. 275: 16560-16568, 2000.
[107967]18995.
[107968]Behm, F. G.; Smith, F. O.; Raimondi, S. C.; Pui, C.-H.;
Bernstein,
[107969]J. D.: Human homologue of the rat chondroitin sulfate
proteoglycan,
[107970]NG2, detected by monoclonal antibody 7.1, identifies
childhood acute

- [107971] lymphoblastic leukemias with t(4;11)(q21;q23) or t(11;19)(q23;p13) and MLL gene rearrangements. Blood 87: 1134–1139, 1996.
- [107972] 18996.
- [107973] Pluschke, G.; Vanek, M.; Evans, A.; Dittmar, T.; Schmid, P.; Itin, P.; Filardo, E. J.; Reisfeld, R. A.: Molecular cloning of a human melanoma-associated chondroitin sulfate proteoglycan. Proc. Nat. Acad. Sci. 93: 9710–9715, 1996.
- [107974] 18997.
- [107975] Rettig, W. J.; Dracopoli, N. C.; Goetzger, T. A.; Spengler, B. A.; Biedler, J. L.; Oettgen, H. F.; Old, L. J.: Somatic cell genetic analysis of human cell surface antigens: chromosomal assignments and regulation of expression in rodent–human hybrid cells. Proc. Nat. Acad. Sci. 81: 6437–6441, 1984.

[107984]18998.

[107985]Rettig, W. J.; Real, F. X.; Spengler, B. A.; Biedler, J. L.; Old,

[107986]L. J.: Human melanoma proteoglycan: expression in hybrids controlled

[107987]by intrinsic and extrinsic signals. Science 231:
1281–1284, 1986.

[107988]18999.

[107989]Smith, F. O.; Rauch, C.; Williams, D. E.; March, C. J.;
Arthur,

[107990]D.; Hilden, J.; Lampkin, B. C.; Buckley, J. D.; Buckley, C. V.;
Woods,

[107991]W. G.; Dinndorf, P. A.; Sorensen, P.; Kersey, J.; Hammond,
D.; Bernstein,

[107992]I. D.: The human homologue of rat NG2, a chondroitin
sulfate proteoglycan,

[107993]is not expressed on the cell surface of normal hematopoietic cells

[107994]but is expressed by acute myeloid leukemia blasts from
poor-prognosis

[107995]patients with abnormalities of chromosome band 11q23.
Blood 87:

[107996]1123–1133, 1996.

[107997]19000.

[107998]Clark, J.; Moore, L.; Krasinskas, A.; Way, J.; Battey, J.;
Tamkun,

[107999]J.; Kahn, R. A.: Selective amplification of additional mem-
bers of

[108000]the ADP-ribosylation factor (ARF) family: cloning of addi-
tional human

[108001]and Drosophila ARF-like genes. Proc. Nat. Acad. Sci. 90:
8952-8956,

[108002]1993.

[108003]19001.

[108004]Gipp, J. J.; Bailey, H. H.; Mulcahy, R. T.: Cloning and se-
quencing

[108005]of the cDNA for the light subunit of human liver gamma-
glutamylcysteine

[108006]synthetase and relative mRNA levels for heavy and light
subunits in

[108007]human normal tissues. Biochem. Biophys. Res. Commun.
206: 584-589,

[108008]1995.

[108009]19002.

- [108010]Huang; C.-S.; Anderson; M. E.; Meister, A.: Amino acid sequence
- [108011]and function of the light subunit of rat kidney gamma-glutamylcysteine
- [108012]synthetase. J. Biol. Chem. 268: 20578-20583, 1993.
- [108013]19003.
- [108014]Sierra-Rivera, E.; Dasouki, M.; Summar, M. L.; Krishnamani, M.
- [108015]R. S.; Meredith, M.; Rao, P. N.; Phillips, J. A., III; Freeman, M.
- [108016]L.: Assignment of the human gene (GLCLR) that encodes the regulatory
- [108017]subunit of gamma-glutamylcysteine synthetase to chromosome 1p21. Cytogenet.
- [108018]Cell Genet. 72: 252-254, 1996.
- [108019]19004.
- [108020]Tsuchiya, K.; Mulcahy, R. T.; Reid, L. L.; Disteché, C. M.; Kavanagh,
- [108021]T. J.: Mapping of the glutamate-cysteine ligase catalytic subunit
- [108022]gene (GLCLC) to human chromosome 6p12 and mouse chromosome 9D-E and

[108023]of the regulatory subunit gene (GLCLR) to human chromosome 1p21–p22

[108024]and mouse chromosome 3H1–3. Genomics 30: 630–632, 1995.

[108025]19005.

[108026]Rozet, J.–M.; Gerber, S.; Perrault, I.; Calvas, P.; Souied, E.;

[108027]Chatelin, S.; Viegas–Pequignot, E.; Molina–Gomez, D.; Munnich, A.;

[108028]Kaplan, J.: Structure and refinement of the physical mapping of the

[108029]gamma–glutamylcysteine ligase regulatory subunit (GLCLR) gene to chromosome

[108030]1pp22.1 within the critically deleted region of human malignant mesothelioma. Cytogenet.

[108031]Cell Genet. 82: 91–94, 1998.

[108032]19006.

[108033]Ellis, J. A.; Luzio, J. P.: Identification and characterization

[108034]of a novel protein (p137) which transcytoses bidirectionally in Caco–2

[108035]cells. J. Biol. Chem. 270: 20717–20723, 1995.

[108036]19007.

[108037]Gessler, M.; Klamt, B.; Tsaoussidou, S.; Ellis, J. A.; Luzio, J.

[108038]P.: The gene encoding the GPI-anchored membrane protein p137(GPI)

[108039](M11S1) maps to human chromosome 11p13 and is highly conserved in

[108040]the mouse. Genomics 32: 169–170, 1996.

[108041]19008.

[108042]Rouault, J.–P; Falette, N.; Guehenneux, F.; Guillot, C.; Rimokh,

[108043]R.; Wang, Q.; Berthet, C.; Moyret–Lalle, C.; Savatier, P.; Pain, B.;

[108044] Shaw, P.; Berger, R.; Samarut, J.; Magaud, J.-P.; Ozturk, M.; Samarut, C.; Puisieux, A.: Identification of BTG2, an antiproliferative p53-dependent component of the DNA damage cellular response pathway. Nature Genet. 14: 482-486, 1996.

[108047] 19009.

[108048] Ring, H. Z.; Vameghi-Meyers, V.; Wang, W.; Crabtree, G. R.; Francke, U.: Five SWI/SNF-related, matrix-associated, actin-dependent regulator of chromatin (SMARC) genes are dispersed in the human genome. Genomics 51: 140-143, 1998.

[108052] 19010.

[108053] Wang, W.; Xue, Y.; Zhou, S.; Kuo, A.; Cairns, B. R.; Crabtree, G. R.: Diversity and specialization of mammalian SWI/SNF complexes. Genes Dev. 10: 2117-2130, 1996.

[108057]19011.

[108058]Kim, B.-T.; Kitagawa, H.; Tamura, J.; Saito, T.; Kusche-Gullberg,

[108059]M.; Lindahl, U.; Sugahara, K.: Human tumor suppressor
EXT gene family

[108060]members EXTL1 and EXTL3 encode alpha-
1,4-N-acetylglucosaminyltransferases

[108061]that likely are involved in heparan sulfate/heparin biosynthesis. Proc.

[108062]Nat. Acad. Sci. 98: 7176-7181, 2001.

[108063]19012.

[108064]Wise, C. A.; Clines, G. A.; Massa, H.; Trask, B. J.; Lovett, M.

[108065]: Identification and localization of the gene for EXTL, a
third member

[108066]of the multiple exostoses gene family. Genome Res. 7:
10-16, 1997.

[108067]19013.

[108068]Moskow, J. J.; Bullrich, F.; Huebner, K.; Daar, I. O.; Buch-
berg,

[108069]A. M.: Meis1, a PBX1-related homeobox gene involved in
myeloid leukemia

[108070]in BXH-2 mice. *Molec. Cell. Biol.* 15: 5434–5443, 1995.

[108071]19014.

[108072]Steelman, S.; Moskow, J. J.; Muzynski, K.; North, C.; Druck, T.;

[108073]Montgomery, J. C.; Huebner, K.; Daar, I. O.; Buchberg, A. M.: Identification

[108074]of a conserved family of Meis1-related homeobox genes. *Genome Res.* 7:

[108075]142–156, 1997.

[108076]19015.

[108077]Thorsteinsdottir, U.; Kroon, E.; Jerome, L.; Blasi, F.; Sauvageau,

[108078]G.: Defining roles for HOX and MEIS1 genes in induction of acute

[108079]myeloid leukemia. *Molec. Cell. Biol.* 21: 224–234, 2001.

[108080]19016.

[108081]Capdevila, J.; Tsukui, T.; Esteban, C. R.; Zappavigna, V.; Belmonte,

[108082]J. C. I.: Control of vertebrate limb outgrowth by the proximal factor

[108083]Meis2 and distal antagonism of BMPs by Gremlin. *Molec.*

- Cell 4: 839–849,
[108084]1999.
- [108085]19017.
- [108086]Nakamura, T.; Jenkins, N. A.; Copeland, N. G.: Identification
of a new family of Pbx-related homeobox genes. *Oncogene* 13: 2235–2242,
[108088]1996.
- [108089]19018.
- [108090]Smith, J. E.; Afonja, O.; Yee, H. T.; Inghirami, G.;
Takeshita,
[108091]K.: Chromosomal mapping to 15q14 and expression analysis of the human
[108092]MEIS2 homeobox gene. *Mammalian Genome* 8: 951–952,
1997.
- [108093]19019.
- [108094]Kobayashi, S.; Uemura, H.; Kohda, T.; Nagai, T.; Chinen,
Y.; Naritomi,
[108095]K.; Kinoshita, E.; Ohashi, H.; Imaizumi, K.; Tsukahara, M.;
Sugio,
[108096]Y.; Tonoki, H.; Kishino, T.; Tanaka, T.; Yamada, M.; Tsut-

sumi, O.;

[108097]Niikawa, N.; Kaneko-Ishino, T.; Ishino, F.: No evidence of
PEG1/MEST

[108098]gene mutations in Silver-Russell syndrome patients. Am. J.
Med. Genet. 104:

[108099]225-231, 2001.

[108100]19020.

[108101]Mizuno, K.; Hasegawa, K.; Katagiri, T.; Ogimoto, M.;
Ichikawa,

[108102]T.; Yakura, H.: MPTP-delta, a putative murine homolog of
HPTP-delta,

[108103]is expressed in specialized regions of the brain and in the
B-cell

[108104]lineage. Molec. Cell. Biol. 13: 5513-5523, 1993.

[108105]19021.

[108106]Uetani, N.; Kato, K.; Ogura, H.; Mizuno, K.; Kawano, K.;
Mikoshiba,

[108107]K.; Yakura, H.; Asano, M.; Iwakura, Y.: Impaired learning
with enhanced

[108108]hippocampal long-term potentiation in PTP-
delta-deficient mice. EMBO

[108109]J. 19: 2775-2785, 2000.

[108110]19022.

[108111]Heighway, J.; Betticher, D. C.; Hoban, P. R.; Altermatt, H. J.;

[108112]Cowen, R.: Coamplification in tumors of KRAS2, type 2 in-
ositol 1,4,5

[108113]triphosphate receptor gene, and a novel human gene,
KRAg. Genomics 35:

[108114]207-214, 1996.

[108115]19023.

[108116]Scott, A. F.; Elizaga, A.; Morrell, J.; Bergen, A.; Penno, M. B.

[108117]: Characterization of a gene coamplified with Ki-ras in Y1
murine

[108118]adrenal carcinoma cells that codes for a putative mem-
brane protein. Genomics 20:

[108119]227-230, 1994.

[108120]19024.

[108121]Monte, D.; Baert, J. L.; Defossez, P. A.; de Launoit, Y.; Ste-
helin,

[108122]D.: Molecular cloning and characterization of human ERM,
a new member

[108123]of the Ets family closely related to mouse PEA3 and ER81
transcription

[108124]factors. *Oncogene* 9: 1397–1406, 1994.

[108125]19025.

[108126]Monte, D.; Coutte, L.; Dewitte, F.; Defossez, P.-A.; Le Coniat,

[108127]M.; Stehelin, D.; Berger, R.; de Launoit, Y.: Genomic organization

[108128]of the human ERM (ETV5) gene, a PEA3 group member of ETS transcription

[108129]factors. *Genomics* 35: 236–240, 1996.

[108130]19026.

[108131]Protopopova, M. V.; Vorobieva, N. V.; Protopopov, A. I.; Gizatullin,

[108132]R. Z.; Kashuba, V. I.; Klein, G.; Zabarovsky, E. R.; Graphodatsky,

[108133]A. S.: Assignment of the ERM gene (ETV5) coding for the ets-related

[108134]protein to human chromosome band 3q28 by in situ hybridization. *Cytogenet.*

[108135]*Cell Genet.* 74: 220 only, 1996.

[108136]19027.

[108137]Bosher, J. M.; Williams, T.; Hurst, H. C.: The developmen-

tally

[108138]regulated transcription factor AP-2 is involved in c-erbB-2 overexpression

[108139]in human mammary carcinoma. Proc. Nat. Acad. Sci. 92: 744-747, 1995.

[108140]19028.

[108141]Brown, M. E.; Pollak, M.; Seidman, C. E.; Seidman, J. G.; Chou,

[108142]Y.-H. W.; Riccardi, D.; Herbert, S. C.: Calcium-ion-sensing cell-surface

[108143]receptors. New Eng. J. Med. 333 -234-240, 1995.

[108144]19029.

[108145]Carling, T.; Szabo, E.; Bai, M.; Ridefelt, P.; Westin, G.; Gustavsson,

[108146]P.; Trivedi, S.; Hellman, P.; Brown, E. M.; Dahl, N.; Rastad, J.:

[108147]Familial hypercalcemia and hypercalciuria caused by a novel mutation

[108148]in the cytoplasmic tail of the calcium receptor. J. Clin. Endocr.

[108149]Metab. 85: 2042-2047, 2000.

[108150]19030.

[108151]Cetani, F.; Pinchera, A.; Pardi, E.; Cianferotti, L.; Vignali,
[108152]E.; Picone, A.; Miccoli, P.; Viacava, P.; Marcocci, C.: No evi-
dence
[108153]for mutations in the calcium-sensing receptor gene in
sporadic parathyroid
[108154]adenomas. J. Bone Miner. Res. 14: 878-882, 1999.
[108155]19031.
[108156]D'Souza-Li, L.; Canaff, L.; Janicic, N.; Cole, D. E. C.; Hendy,
[108157]G. N.: An acceptor splice site mutation in the calcium-
sensing receptor
[108158](CASR) gene in familial hypocalciuric hypercalcemia and
neonatal severe
[108159]hyperparathyroidism. Hum. Mutat. 18: 411-421, 2001.
[108160]19032.
[108161]De Luca, F.; Ray, K.; Mancilla, E. E.; Fan, G.-F.; Winer, K. K.;
[108162]Gore, P.; Spiegel, A. M.; Baron, J.: Sporadic hypoparathy-
roidism
[108163]caused by de novo gain-of-function mutations of the
Ca(2+)-sensing
[108164]receptor. J. Clin. Endocr. Metab. 82: 2710-2715, 1997.

[108165]19033.

[108166]Hendy, G. N.; D'Souza-Li, L.; Yang, B.; Canaff, L.; Cole, D. E.

[108167]C.: Mutations of the calcium-sensing receptor (CASR) in familial

[108168]hypocalciuric hypercalcemia, neonatal severe hyperparathyroidism,

[108169]and autosomal dominant hypocalcemia. Hum. Mutat. 16: 281-296, 2000.

[108170]19034.

[108171]Janicic, N.; Pausova, Z.; Cole, D. E. C.; Hendy, G. N.: Insertion

[108172]of an Alu sequence in the Ca(2+)-sensing receptor gene in familial

[108173]hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. Am.

[108174]J. Hum. Genet. 56: 880-886, 1995.

[108175]19035.

[108176]Jap, T.-S.; Wu, Y.-C.; Jenq, S.-F.; Won, G.-S.: A novel mutation

[108177]in the calcium-sensing receptor gene in a Chinese subject with persistent

[108178]hypercalcemia and hypocalciuria. J. Clin. Endocr. Metab.
86: 13–15,

[108179]2001.

[108180]19036.

[108181]Kobayashi, M.; Tanaka, H.; Tsuzuki, K.; Tsuyuki, M.; Igaki,
H.;

[108182]Ichinose, Y.; Aya, K.; Nishioka, N.; Seino, Y.: Two novel
missense

[108183]mutations in calcium–sensing receptor gene associated
with neonatal

[108184]severe hyperparathyroidism. J. Clin. Endocr. Metab. 82:
2716–2719,

[108185]1997.

[108186]19037.

[108187]Lienhardt, A.; Bai, M.; Lagarde, J.–P.; Rigaud, M.; Zhang,
Z.;

[108188]Jiang, Y.; Kottler, M.–L.; Brown, E. M.; Garabedian, M.: Ac–
tivating

[108189]mutations of the calcium–sensing receptor: management
of hypocalcemia. J.

[108190]Clin. Endocr. Metab. 86: 5313–5323, 2001.

[108191]19038.

[108192]Lienhardt, A.; Garabedian, M.; Bai, M.; Sinding, C.; Zhang, Z.;

[108193]Lagarde, J.-P.; Boulesteix, J.; Rigaud, M.; Brown, E. M.; Kottler,

[108194]M.-L.: A large homozygous or heterozygous in-frame deletion within

[108195]the calcium-sensing receptor's carboxylterminal cytoplasmic tail that

[108196]causes autosomal dominant hypocalcemia. J. Clin. Endocr. Metab. 85:

[108197]1695-1702, 2000.

[108198]19039.

[108199]Lovlie, R.; Eiken, H. G.; Sorheim, J. I.; Boman, H.: The Ca(2+)-sensing

[108200]receptor gene (PCAR1) mutation T151M in isolated autosomal dominant

[108201]hypoparathyroidism. Hum. Genet. 98: 129-133, 1996.

[108202]19040.

[108203]Okazaki, R.; Chikatsu, N.; Nakatsu, M.; Takeuchi, Y.; Ajima, M.;

- [108204]Miki, J.; Fujita, T.; Arai, M.; Totsuka, Y.; Tanaka, K.; Fukumoto,
moto,
- [108205]S.: A novel activating mutation in calcium-sensing receptor gene
tor gene
- [108206]associated with a family of autosomal dominant hypocalcemia. J. Clin.
cemia. J. Clin.
- [108207]Endocr. Metab. 84: 363–366, 1999.
- [108208]19041.
- [108209]Combadiere, C.; Ahuja, S. K.; Murphy, P. M.: Cloning,
chromosomal
- [108210]localization, and RNA expression of a human beta
chemokine receptor-like
- [108211]gene. DNA Cell Biol. 14: 673–680, 1995.
- [108212]19042.
- [108213]Combadiere, C.; Ahuja, S. K.; Van Damme, J.; Tiffany, H.
L.; Gao,
- [108214]J.-L.; Murphy, P. M.: Monocyte chemoattractant protein-3
is a functional
- [108215]ligand for CC chemokine receptors 1 and 2B. J. Biol.
Chem. 270:
- [108216]29671–29675, 1995.

[108217]19043.

[108218]Doranz, B. J.; Rucker, J.; Yi, Y.; Smyth, R. J.; Samson, M.;
Peiper,

[108219]S. C.; Parmentier, M.; Collman, R. G.; Doms, R. W.: A dual-
tropic

[108220]primary HIV-1 isolate that uses fusin and the beta-
chemokine receptors

[108221]CKR-5, CKR-3, and CKR-2b as fusion cofactors. Cell 85:
1149-1158,

[108222]1996.

[108223]19044.

[108224]Mummidi, S.; Ahuja, S. S.; Gonzalez, E.; Anderson, S. A.;
Santiago,

[108225]E. N.; Stephan, K. T.; Craig, F. E.; O'Connell, P.; Tryon, V.;
Clark,

[108226]R. A.; Dolan, M. J.; Ahuja, S. K.: Genealogy of the CCR5 lo-
cus and

[108227]chemokine system gene variants associated with altered
rates of HIV-1

[108228]disease progression. Nature Med. 4: 786-793, 1998.

[108229]19045.

[108230]Peters, W.; Dupuis, M.; Charo, I. F.: A mechanism for the impaired

[108231]IFN- γ production in C-C chemokine receptor 2 (CCR2) knockout mice:

[108232]role of CCR2 in linking the innate and adaptive immune responses. J.

[108233]Immun. 165: 7072–7077, 2000.

[108234]19046.

[108235]Peters, W.; Scott, H. M.; Chambers, H. F.; Flynn J. L.; Charo,

[108236]I. F.; Ernst, J. D.: Chemokine receptor 2 serves an early and essential

[108237]role in resistance to Mycobacterium tuberculosis. Proc. Nat. Acad.

[108238]Sci. 98: 7958–7963, 2001.

[108239]19047.

[108240]Samson, M.; Labbe, O.; Mollereau, C.; Vassart, G.; Parmentier,

[108241]M.: Molecular cloning and functional expression of a new human CC-chemokine

[108242]receptor gene. Biochemistry 35: 3362–3367, 1996.

[108243]19048.

[108244]Samson, M.; Soularue, P.; Vassart, G.; Parmentier, M.: The
genes
[108245]encoding the human CC-chemokine receptors CC-CCR1
to CC-CCR5 (CMKBR1-CMKBR5)
[108246]are clustered in the p21.3-p24 region of chromosome 3.
Genomics 36:
[108247]522-526, 1996.
[108248]19049.
[108249]Sanders, S. K.; Crean, S. M.; Boxer, P. A.; Kellner, D.;
LaRosa,
[108250]G. J.; Hunt, S. W., III.: Functional differences between
monocyte
[108251]chemotactic protein-1 receptor A and monocyte chemo-
tactic protein-1
[108252]receptor B expressed in a Jurkat T cell. J. Immun. 165:
4877-4883,
[108253]2000.
[108254]19050.
[108255]Smith, M. W.; Dean, M.; Carrington, M.; Winkler, C.; Hutt-
ley,
[108256]G. A.; Lomb, D. A.; Goedert, J. J.; O'Brien, T. R.; Jacobson,
L. P.;

- [108257]Kaslow, R.; Buchbinder, S.; Vittinghoff, E.; Vlahov, D.;
Hoots, K.;
- [108258]Hilgartner, M. W.; Hemophilia Growth and Development
Study (HGDS);
- [108259]Multicenter AIDS Cohort Study (MACS); Multicenter
Hemophilia Cohort
- [108260]Study (MHCS); San Francisco City Cohort (SFCC); ALIVE
Study;
- [108261]O'Brien, S. J.: Contrasting genetic influence of CCR2 and
CCR5 variants
- [108262]on HIV-1 infection and disease progression. *Science* 277:
959-965,
- [108263]1997.
- [108264]19051.
- [108265]Wong, L.-M.; Myers, S. J.; Tsou, C.-L.; Gosling, J.; Arai, H.;
- [108266]Charo, I. F.: Organization and differential expression of
the human
- [108267]monocyte chemoattractant protein 1 receptor gene: evi-
dence for the
- [108268]role of the carboxyl-terminal tail in receptor trafficking. *J.*
Biol.
- [108269]*Chem.* 272: 1038-1045, 1997.

[108270]19052.

[108271]Robertson, N. G.; Khetarpal, U.; Gutierrez-Espeleta, G. A.;
Bieber,

[108272]F. R.; Morton, C. C.: Isolation of novel and known genes
from a human

[108273]fetal cochlear cDNA library using subtractive hybridization
and differential

[108274]screening. Genomics 23: 42-50, 1994.

[108275]19053.

[108276]Rothschild, C. B.; Freedman, B. I.; Hodge, R.; Rao, P. N.;
Pettenati,

[108277]M. J.; Anderson, R. A.; Akots, G.; Qadri, A.; Roh, B.; Fajans,
S.

[108278]S.; Reis, A.; Morris, D. J.; Usala, A.; Hayward, C.; Brock, D.;
Colle,

[108279]E.; Spray, B. J.; Rich, S. S.; Bowden, D. W.: Fructose-
1,6-bisphosphatase:

[108280]genetic and physical mapping to human chromosome
9q22.3 and evaluation

[108281]in non-insulin-dependent diabetes mellitus. Genomics 29:
187-194,

[108282]1995.

[108283]19054.

[108284]Simunovic, M. P.; Regan, B. C.; Mollon, J. D.: Is color vision

[108285]deficiency an advantage under scotopic conditions? Invest.
Ophthalm.

[108286]Vis. Sci. 42: 3357–3364, 2001.

[108287]19055.

[108288]Austen, K. F.: Inborn errors of the complement system of
man. New

[108289]Eng. J. Med. 276: 1363–1367, 1967.

[108290]19056.

[108291]Austen, K. F.; Becker, E. L.; Bero, C. E.; Borsos, T.; Dal-
masso,

[108292]A. P.; Dias Da Silva, D.: Nomenclature of complement. Bull.
WHO 39:

[108293]935–938, 1968.

[108294]19057.

[108295]Awdeh, Z. L.; Raum, D. D.; Glass, D.; Agnello, V.; Schur, P.
H.;

[108296]Johnston, R. B., Jr.; Gelfand, E. W.; Ballow, M.; Yunis, E.;
Alper,

- [108297]C. A.: Complement–human histocompatibility antigen
haplotypes in
- [108298]C2 deficiency. J. Clin. Invest. 67: 581–583, 1981.
- [108299]19058.
- [108300]Belin, D. C.; Bordwell, B. J.; Einarson, M. E.; McLean, R. H.;
- [108301]Weinstein, A.; Yunis, E. J.; Rothfield, N. F.: Familial discoid
lupus
- [108302]erythematosus associated with heterozygote C2 defi-
ciency. Arthritis
- [108303]Rheum. 23: 898–903, 1980.
- [108304]19059.
- [108305]Bentley, D. R.: Primary structure of human complement
component
- [108306]C2: homology to two unrelated protein families. Biochem.
J. 239:
- [108307]339–345, 1986.
- [108308]19060.
- [108309]Bentley, D. R.; Porter, R. R.: Isolation of cDNA clones for
human
- [108310]complement component C2. Proc. Nat. Acad. Sci. 81:
1212–1215, 1984.

[108311]19061.

[108312]Bottger, E. C.; Hoffmann, T.; Hadding, U.; Bitter-Suer-
mann, D.

[108313]: Guinea pigs with inherited deficiencies of complement
components

[108314]C2 or C4 have characteristics of immune complex disease.
J. Clin.

[108315]Invest. 78: 689–695, 1986.

[108316]19062.

[108317]Callen, J. P.; Hodge, S. J.; Kulick, K. B.; Stelzer, G.; Buchino,

[108318]J. J.: Subacute cutaneous lupus erythematosus in multiple
members

[108319]of a family with C2 deficiency. Arch. Derm. 123: 66–70,
1987.

[108320]19063.

[108321]Cole, F. S.; Whitehead, A. S.; Auerbach, H. S.; Lint, T.;
Zeitz,

[108322]H. J.; Kilbridge, P.; Colten, H. R.: The molecular basis for
genetic

[108323]deficiency of the second component of human comple-
ment. New Eng.

[108324]J. Med. 313: 11-16, 1985.

[108325]19064.

[108326]Donald, L. J.; Wang, H. S.; Hamerton, J. L.: Assignment of the

[108327]gene for cystathionase (CYS) to human chromosome 16.
(Abstract) Cytogenet.

[108328]Cell Genet. 32: 268 only, 1982.

[108329]19065.

[108330]Frimpter, G. W.: Cystathioninuria: nature of the defect.
Science 149:

[108331]1095-1096, 1965.

[108332]19066.

[108333]Frimpter, G. W.: Cystathioninuria, sulfite oxidase deficiency,

[108334]and 'beta-mercaptolactate-cysteine disulfiduria.In: Stanbury, J. B.;

[108335]Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic Basis
of Inherited

[108336]Disease. New York: McGraw-Hill (pub.) (3rd ed.): 1972. Pp.
413-425.

[108337]19067.

- [108338]Frimpter, G. W.; Haymovitz, A.; Horwith, M.: Cystathioninuria. New
- [108339]Eng. J. Med. 268: 333–339, 1963.
- [108340]19068.
- [108341]Harris, H.; Penrose, L. S.; Thomas, D. H. H.: Cystathioninuria. Ann.
- [108342]Hum. Genet. 23: 442–453, 1959.
- [108343]19069.
- [108344]Lu, Y.; O'Dowd, B. F.; Orrego, H.; Israel, Y.: Cloning and nucleotide
- [108345]sequence of human liver cDNA encoding for cystathionine gamma-lyase. Biochem.
- [108346]Biophys. Res. Commun. 189: 749–758, 1992.
- [108347]19070.
- [108348]Lyon, I. C. T.; Procopis, P. G.; Turner, B.: Cystathioninuria
- [108349]in a well baby population. Acta Paediat. Scand. 60:
- 324–328, 1971.
- [108350]19071.
- [108351]Mongeau, J.-G.; Hilgartner, M.; Worthen, H. G.; Frimpter, G. W.

[108352]: Cystathioninuria: study of an infant with normal mental-
ity, thrombocytopenia,

[108353]and renal calculi. J. Pediat. 69: 1113-1120, 1967.

[108354]19072.

[108355]Pascal, T. A.; Gaull, G. E.; Beratis, N. G.; Gillam, B. M.; Tal-
lan,

[108356]H. H.: Cystathionase deficiency: evidence for genetic het-
erogeneity

[108357]in primary cystathioninuria. Pediat. Res. 12: 125-133,
1978.

[108358]19073.

[108359]Perry, T. L.; Hardwick, D. F.; Hansen, S.; Love, D. L.; Is-
raels,

[108360]S.: Cystathioninuria in two healthy siblings. New Eng. J.
Med. 278:

[108361]590-592, 1968.

[108362]19074.

[108363]Schneiderman, L. J.: Latent cystathioninuria. J. Med. Genet.
4:

[108364]260-263, 1967.

[108365]19075.

- [108366] Scott, C. R.; Dassell, S. W.; Clark, S. H.; Chiang-Teng, C.;
Swedberg,
[108367] K. R.: Cystathioninemia: a benign genetic condition. J. Pe-
diat. 76:
[108368] 571-577, 1970.
[108369] 19076.
- [108370] Shaw, K. N. F.; Lieberman, E.; Koch, R.; Donnell, G. N.:
Cystathioninuria. Am.
[108371] J. Dis. Child. 113: 119-128, 1967.
[108372] 19077.
- [108373] Tada, K.; Yoshida, T.; Yokoyama, Y.; Sato, T.; Nakagawa,
H.; Arakawa,
[108374] T.: Cystathioninuria not associated with vitamin B6 de-
pendency: a
[108375] probably new type of cystathioninuria. Tohoku J. Exp.
Med. 95: 235-242,
[108376] 1968.
[108377] 19078.
- [108378] Whelan, D. T.; Scriver, C. R.: Cystathioninuria and renal
iminoglycinuria
[108379] in a pedigree: a perspective on counseling. New Eng. J.

Med. 278:

[108380]924-927, 1968.

[108381]19079.

[108382]Abeliovich, D.; Lavon, I. P.; Lerer, I.; Cohen, T.; Springer, C.;

[108383]Avital, A.; Cutting, G. R.: Screening for five mutations de-
fects

[108384]97% of cystic fibrosis (CF) chromosomes and predicts a
carrier frequency

[108385]of 1:29 in the Jewish Ashkenazi population. Am. J. Hum.
Genet. 51:

[108386]951-956, 1992.

[108387]19080.

[108388]Hobart, M. J.; Joysey, V.; Lachmann, P. J.: Inherited struc-
tural

[108389]variation and linkage relationships of C7. J. Immunogenet.
5: 157-163,

[108390]1978.

[108391]19081.

[108392]Jeremiah, S. J.; Abbott, C. M.; Murad, Z.; Povey, S.;
Thomas,

- [108393]H. J.; Solomon, E.; DiScipio, R. G.; Fey, G. H.: The assignment of
the genes coding for human complement components C6
and C7 to chromosome
5. *Ann. Hum. Genet.* 54: 141–147, 1990.
19082.
- [108397]Schulpis, K. H.; Papaconstantinou, E. D.; Koidou, A.;
Michelakakis,
H.; Tzamouranis, J.; Patsouras, A.; Shin, Y.: UDP galac-
tose-4-epimerase
deficiency in a 5.5-year-old girl with a unilateral cataract.
J.
Inherit. Metab. Dis. 16: 903–904, 1993.
19083.
- [108402]Thoden, J. B.; Wohlers, T. M.; Fridovich-Keil, J. L.; Holden,
H. M.: Molecular basis for severe epimerase deficiency
galactosemia:
x-ray structure of the human V94M-substituted UDP-
galactose 4-epimerase. J.
Biol. Chem. 276: 20617–20623, 2001.
19084.

[108407] Wohlers, T. M.; Christacos, N. C.; Harreman, M. T.;
Fridovich-Keil,
[108408] J. L.: Identification and characterization of a mutation, in
the
[108409] human UDP-galactose-4-epimerase gene, associated with
generalized
[108410] epimerase-deficiency galactosemia. Am. J. Hum. Genet.
64: 462-470,
[108411] 1999.
[108412] 19085.
[108413] Cheadle, J.; Al-Jader, L.; Goodchild, M.; Meredith, A. L.:
Mild
[108414] pulmonary disease in a cystic fibrosis child homozygous
for R553X. J.
[108415] Med. Genet. 29: 597, 1992.
[108416] 19086.
[108417] Cheadle, J. P.; Meredith, A. L.; Al-Jader, L. N.: A new mis-
sense
[108418] mutation (R1283M) in exon 20 of the cystic fibrosis trans-
membrane
[108419] conductance regulator gene. Hum. Molec. Genet. 1:
123-125, 1992.

[108420]19087.

[108421]Colledge, W. H.; Abella, B. S.; Southern, K. W.; Ratcliff, R.;

[108422]Jiang, C.; Cheng, S. H.; MacVinish, L. J.; Anderson, J. R.;
Cuthbert,

[108423]A. W.; Evans, M. J.: Generation and characterisation of a
delta-F508

[108424]cystic fibrosis mouse model. Nature Genet. 10: 445-452,
1995.

[108425]19088.

[108426]Haworth, J. C.; Gourley, B.; Hadorn, B.; Sumida, C.: Malab-
sorption

[108427]and growth failure due to intestinal enterokinase defi-
ciency. J.

[108428]Pediat. 78: 481-490, 1971.

[108429]19089.

[108430]Haworth, J. C.; Hadorn, B.; Gourley, B.; Prasad, A.;
Troesch, V.

[108431]: Intestinal enterokinase deficiency: occurrence in two sibs
and age

[108432]dependency of clinical expression. Arch. Dis. Child. 50:
277-282,

[108433]1975.

[108434]19090.

[108435]Holzinger, A.; Maier, E. M.; Buck, C.; Mayerhofer, P. U.;
Kappler,

[108436]M.; Haworth, J. C.; Moroz, S. P.; Hadorn, H.-B.; Sadler, J.
E.; Roscher,

[108437]A. A.: Mutations in the proenteropeptidase gene are the
molecular

[108438]cause of congenital enteropeptidase deficiency. Am. J.
Hum. Genet. 70:

[108439]20–25, 2002.

[108440]19091.

[108441]DiGeorge, A. M.; Harley, R. D. L.: The association of
aniridia,

[108442]Wilms tumor and genital abnormalities. Arch. Ophthalm. 75:
796–798,

[108443]1966.

[108444]19092.

[108445]Dressler, G. R.; Douglass, E. C.: Pax-2 is a DNA-binding
protein

[108446]expressed in embryonic kidney and Wilms tumor. Proc.

Nat. Acad. Sci. 89:

[108447]1179–1183, 1992.

[108448]19093.

[108449]Fearon, E. R.; Vogelstein, B.; Feinberg, A. P.: Somatic deletion

[108450]and duplication of genes on chromosome 11 in Wilms' tumours. Nature 309:

[108451]176–178, 1984.

[108452]19094.

[108453]Fitzgerald, W. L.; Hardin, H. C., Jr.: Bilateral Wilms' tumor

[108454]in a Wilms tumor family: case report. J. Urol. 73: 468–474, 1955.

[108455]19095.

[108456]Francke, U.: A gene for Wilms tumour? Nature 343: 692–694,

[108457]1990.

[108458]19096.

[108459]Francke, U.; Holmes, L. B.; Atkins, L.; Riccardi, V. M.: Aniridia–Wilms'

[108460]tumor association: evidence for specific deletion of

11p13. Cytogenet.

[108461]Cell Genet. 24: 185–192, 1979.

[108462]19097.

[108463]Francke, U.; Riccardi, V. M.; Hittner, H. M.; Borges, W.: In-
terstitial

[108464]del(11p) as a cause of the aniridia–Wilms tumor associa-
tion: band

[108465]localization and a heritable basis. (Abstract) Am. J. Hum.
Genet. 30:

[108466]81A, 1978.

[108467]19098.

[108468]Fraumeni, J. F., Jr.; Glass, A. G.: Wilms' tumor and congen-
ital

[108469]aniridia. J.A.M.A. 206: 825–828, 1968.

[108470]19099.

[108471]Haber, D. A.; Buckler, A. J.; Glaser, T.; Call, K. M.; Pelletier,

[108472]J.; Sohn, R. L.; Douglass, E. C.; Housman, D. E.: An internal
deletion

[108473]within an 11p13 zinc finger gene contributes to the de-
velopment of

[108474]Wilms' tumor. Cell 61: 1257–1269, 1990.

[108475]19100.

[108476]Haber, D. A.; Park, S.; Maheswaran, S.; Englert, C.; Re, G.
G.;

[108477]Hazen-Martin, D. J.; Sens, D. A.; Garvin, A. J.:
WT1-mediated growth

[108478]suppression of Wilms tumor cells expressing a WT1 splic-
ing variant. Science 262:

[108479]2057-2059, 1993.

[108480]19101.

[108481]Haber, D. A.; Timmers, H. T. M.; Pelletier, J.; Sharp, P. A.;

[108482]Housman, D. E.: A dominant mutation in the Wilms tumor
gene WT1 cooperates

[108483]with the viral oncogene E1A in transformation of primary
kidney cells. Proc.

[108484]Nat. Acad. Sci. 89: 6010-6014, 1992.

[108485]19102.

[108486]Haning, R. V., Jr.; Chesney, R. W.; Moorthy, A. V.; Gilbert,
E.

[108487]F.: A syndrome of chronic renal failure and XY gonadal
dysgenesis

[108488]in young phenotypic females without genital ambiguity.

Am. J. Kidney

[108489]Dis. 6: 40–48, 1985.

[108490]19103.

[108491]Hastie, N. D.: Dominant negative mutations in the Wilms
tumour

[108492](WT1) gene cause Denys–Drash syndrome--proof that a
tumour-suppressor

[108493]gene plays a crucial role in normal genitourinary develop-
ment. Hum.

[108494]Molec. Genet. 1: 293–295, 1992.

[108495]19104.

[108496]Huff, V.; Meadows, A.; Riccardi, V. M.; Strong, L. C.; Saun-
ders,

[108497]G. F.: Parental origin of de novo constitutional deletions of
chromosomal

[108498]band 11p13. Am. J. Hum. Genet. 47: 155–160, 1990.

[108499]19105.

[108500]Jeanpierre, C.; Antignac, C.; Beroud, C.; Lavedan, C.;
Henry,

[108501]J.; Saunders, G.; Williams, B.; Glaser, T.; Junien, C.: Consti-
tutional

[108502]and somatic deletions of two different regions of maternal chromosome

[108503]11 in Wilms tumor. Genomics 7: 434-438, 1990.

[108504]19106.

[108505]Jolles, B.: Wilms' tumor in father and son. Lancet I: 207,

[108506]1973.

[108507]19107.

[108508]Juberg, R. C.; St. Martin, E. C.; Hundley, J. R.: Familial occurrence

[108509]of Wilms's tumor: nephroblastoma in one of monozygous twins and in

[108510]another sibling. Am. J. Hum. Genet. 27: 155-169, 1975.

[108511]19108.

[108512]Junien, C.; Huerre, C.; Despoisse, S.; Gilgenkrantz, S.; Lenoir,

[108513]G. M.: c-Ha-ras1 is not deleted in del (11p13) Wilms' tumor (WAGR)

[108514]and maps to 11p15.1-11p15.5. (Abstract) Cytogenet. Cell Genet. 37:

[108515]503, 1984.

[108516]19109.

[108517]Junien, C.; Turleau, C.; de Grouchy, J.; Said, R.; Rethore, M.-O.;

[108518]Tenconi, R.; Dufier, J. L.: Regional assignment of catalase (CAT)

[108519]gene to band 11p13: association with the aniridia-Wilms' tumor-gonadoblastoma

[108520](WAGR) complex. Ann. Genet. 23: 165-168, 1980.

[108521]19110.

[108522]Kakati, S.; Xiao, H.; Siddiqui, S. Y.; Sreekantaiah, C.; Weier,

[108523]H.-U. G.; Green, D. M.; Fisher, J. E.; Allen, J. E.: Constitutional

[108524]extra chromosomal element in a family with Wilms' tumor. Hum. Genet. 87:

[108525]183-188, 1991.

[108526]19111.

[108527]Kaneko, Y.; Celida Egues, M.; Rowley, J. D.: Interstitial deletion

[108528]of short arm of chromosome 11 limited to Wilms' tumor cells in a patient

[108529]without aniridia. Cancer Res. 41: 4577-4578, 1981.

[108530]19112.

[108531]Kaufman, R. L.; Vietti, T. J.; Wabner, C. I.: Wilms' tumor in
[108532]father and son. Lancet I: 43, 1973.

[108533]19113.

[108534]Kinberg, J. A.; Angle, C. R.; Wilson, R. B.: Nephropathy-go-
nadal
[108535]dysgenesis, type 2: renal failure in three siblings with XY
dysgenesis

[108536]in one. Am. J. Kidney Dis. 9: 507-510, 1987.

[108537]19114.

[108538]Knudson, A. G., Jr.; Strong, L. C.: Mutation and cancer: a
model

[108539]for Wilms' tumor of the kidney. J. Nat. Cancer Inst. 48:
313-324,

[108540]1972.

[108541]19115.

[108542]Knudson, A. G., Jr.; Strong, L. C.: Familial Wilms's tumor.
(Letter) Am.

[108543]J. Hum. Genet. 27: 809-810, 1975.

[108544]19116.

- [108545]Kolata, G. B.: Genes and cancer: the story of Wilms tumor.
Science 207:
[108546]970–971, 1980.
[108547]19117.
- [108548]Kontras, S. B.; Newton, W. A., Jr.: Familial Wilms' tumor.
Birth
[108549]Defects Orig. Art. Ser. X(4): 187–188, 1974.
[108550]19118.
- [108551]Koufos, A.; Hansen, M. F.; Lampkin, B. C.; Workman, M. L.;
Copeland,
[108552]N. G.; Jenkins, N. A.; Cavenee, W. K.: Loss of alleles at loci
on
[108553]human chromosome 11 during genesis of Wilms' tumour.
Nature 309:
[108554]170–172, 1984.
[108555]19119.
- [108556]Kozman, H. M.; Clarke, J. M.; Little, M. H.; Smith, P. J.:
Molecular
[108557]genetic evidence for common pathogenesis of childhood
and adult Wilms'
[108558]tumor. Cancer Genet. Cytogenet. 38: 121–125, 1989.

[108559]19120.

[108560]Kumar, S.; Harrison, C. J.; Heighway, J.; Marsden, H. B.;
West,

[108561]D. C.; Jones, P. M.: A cell line from Wilms' tumour with
deletion

[108562]in short arm of chromosome 11. Int. J. Cancer 40:
499–504, 1987.

[108563]19121.

[108564]Ladda, R. L.; Atkins, L.; Littlefield, J.: Computer-assisted

[108565]analysis of chromosomal abnormalities: detection of a
deletion in

[108566]aniridia Wilms' tumor syndrome. Science 185: 784–787,
1974.

[108567]19122.

[108568]Berx, G.; Staes, K.; van Hengel, J.; Molemans, F.; Busse-
makers,

[108569]M. J. G.; van Bokhoven, A.; van Roy, F.: Cloning and char-
acterization

[108570]of the human invasion suppressor gene E-cadherin
(CDH1). Genomics 26:

[108571]281–289, 1995.

[108572]19123.

[108573]Boggon, T. J.; Murray, J.; Chappuis-Flament, S.; Wong, E.;
Gumbiner,

[108574]B. M.; Shapiro, L.: C-cadherin ectodomain structure and
implications

[108575]for cell adhesion mechanisms. Science 296: 1308–1313,
2002.

[108576]19124.

[108577]Cano, A.; Perez-Moreno, M. A.; Rodrigo, I.; Locascio, A.;
Blanco,

[108578]M. J.; del Barrio, M. G.; Portillo, F.; Nieto, M. A.: The tran-
scription

[108579]factor Snail controls epithelial–mesenchymal transitions
by repressing

[108580]E-cadherin expression. Nature Cell Biol. 2: 76–83, 2000.

[108581]19125.

[108582]Ceccherini, I.; Romeo, G.; Lawrence, S.; Breuning, M. H.;
Harris,

[108583]P. C.; Himmelbauer, H.; Frischauf, A. M.; Sutherland, G. R.;
Germino,

[108584]G. G.; Reeders, S. T.; Morton, N. E.: Construction of a map
of chromosome

- [108585]16 by using radiation hybrids. Proc. Nat. Acad. Sci. 89:
104–108,
[108586]1992.
[108587]19126.
[108588]Chun, Y. S.; Lindor, N. M.; Smyrk, T. C.; Petersen, B. T.;
Burgart,
[108589]L. J.; Guilford, P. J.; Donohue, J. H.: Germline E-cadherin
gene
[108590]mutations: is prophylactic total gastrectomy indicated?
Cancer 92:
[108591]181–187, 2001.
[108592]19127.
[108593]Eistetter, H. R.; Adolph, S.; Ringwald, M.; Simon–Cha-
zottes, D.;
[108594]Schuh, R.; Guenet, J. L.; Kemler, R.: Chromosomal map-
ping of the
[108595]structural gene coding for the mouse cell adhesion
molecule uvomorulin. Proc.
[108596]Nat. Acad. Sci. 85: 3489–3493, 1988.
[108597]19128.
[108598]Gayther, S. A.; Goringe, K. L.; Ramus, S. J.; Huntsman, D.;

Roviello,

[108599]F.; Grehan, N.; Machado, J. C.; Pinto, E.; Seruca, R.; Halling, K.;

[108600]MacLeod, P.; Powell, S. M.; Jackson, C. E.; Ponder, B. A. J.; Caldas,

[108601]C.: Identification of germ-line E-cadherin mutations in gastric cancer

[108602]families of European origin. Cancer Res. 58: 4086–4089, 1998.

[108603]19129.

[108604]Grady, W. M.; Willis, J.; Guilford, P. J.; Dunbier, A. K.; Toro,

[108605]T. T.; Lynch, H.; Wiesner, G.; Ferguson, K.; Eng, C.; Park, J.-G.;

[108606]Kim, S.-J.; Markowitz, S.: Methylation of the CDH1 promoter as the

[108607]second genetic hit in hereditary diffuse gastric cancer. Nature Genet. 26:

[108608]16–17, 2000.

[108609]19130.

[108610]Humar, B.; Toro, T.; Graziano, F.; Muller, H.; Dobbie, Z.; Kwang-Yang,

[108611]H.; Eng, C.; Hampel, H.; Gilbert, D.; Winship, I.; Parry, S.;

Ward,

[108612]R.; Findlay, M.; Christian, A.; Tucker, M.; Tucker, K.; Merri-
man,

[108613]T.; Guilford, P.: Novel germline CDH1 mutations in heredi-
tary diffuse

[108614]gastric cancer families. Hum. Mutat. 19: 518–525, 2002.

[108615]19131.

[108616]Ilyas, M.; Tomlinson, I. P. M.; Hanby, A.; Talbot, I. C.; Bod-
mer,

[108617]W. F.: Allele loss, replication errors and loss of expression
of

[108618]E-cadherin in colorectal cancers. Gut 40: 654–659, 1997.

[108619]19132.

[108620]Jones, E. G.: Familial gastric cancer. New Zealand Med. J.
63:

[108621]287–296, 1964.

[108622]19133.

[108623]Lecuit, M.; Dramsi, S.; Gottardi, C.; Fedor–Chaiken, M.;
Gumbiner,

[108624]B.; Cossart, P.: A single amino acid in E-cadherin respon-
sible for

[108625] host specificity towards the human pathogen *Listeria monocytogenes*. *EMBO*

[108626] *J.* 18: 3956–3963, 1999.

[108627] 19134.

[108628] Guilford, P. J.; Hopkins, J. B. W.; Grady, W. M.; Markowitz, S.

[108629] D.; Willis, J.; Lynch, H.; Rajput, A.; Wiesner, G. L.; Lindor, N.

[108630] M.; Burgart, L. J.; Toro, T. T.; Lee, D.; Limacher, J.-M.; Shaw, D.

[108631] W.; Findlay, M. P. N.; Reeve, A. E.: E-cadherin germline mutations

[108632] define an inherited cancer syndrome dominated by diffuse gastric cancer. *Hum.*

[108633] *Mutat.* 14: 249–255, 1999.

[108634] 19135.

[108635] Lecuit, M.; Vandormael-Pournin, S.; Lefort, J.; Huerre, M.; Gounon,

[108636] P.; Dupuy, C.; Babinet, C.; Cossart, P.: A transgenic model for listeriosis:

[108637] role of internalin in crossing the intestinal barrier. *Science* 292:

[108638]1722–1725, 2001.

[108639]19136.

[108640]Lorber, B.: Listeriosis. Clin. Infect. Dis. 24: 1–11, 1997.

[108641]19137.

[108642]Lynch, H. T.; Grady, W.; Lynch, J. F.; Tsuchiya, K. D.; Wiesner,

[108643]G.; Markowitz, S. D.: E-cadherin mutation-based genetic counseling

[108644]and hereditary diffuse gastric carcinoma. Cancer Genet. Cytogenet. 122:

[108645]1–6, 2000.

[108646]19138.

[108647]Mansouri, A.; Goodfellow, P. N.; Kemler, R.: Molecular cloning

[108648]and chromosomal localization of the human cell adhesion molecule uvomorulin

[108649](UVO). (Abstract) Cytogenet. Cell Genet. 46: 655, 1987.

[108650]19139.

[108651]Mansouri, A.; Spurr, N.; Goodfellow, P. N.; Kemler, R.: Characterization

[108652]and chromosomal localization of the gene encoding the human cell adhesion

[108653]molecule uvomorulin. Differentiation 38: 67–71, 1988.

[108654]19140.

[108655]Natt, E.; Magenis, R. E.; Zimmer, J.; Mansouri, A.; Scherer, G.

[108656]; Regional assignment of the human loci for uvomorulin (UVO) and chymotrypsinogen

[108657]B (CTRB) with the help of two overlapping deletions on the long arm

[108658]of chromosome 16. Cytogenet. Cell Genet. 50: 145–148, 1989.

[108659]19141.

[108660]Oda, T.; Kanai, Y.; Oyama, T.; Yoshiura, K.; Shimoyama, Y.; Birchmeier,

[108661]W.; Sugimura, T.; Hirohashi, S.: E-cadherin gene mutations in human

[108662]gastric carcinoma cell lines. Proc. Nat. Acad. Sci. 91: 1858–1862,

[108663]1994.

[108664]19142.

- [108665]Oliveira, C.; Bordin, M. C.; Grehan, N.; Huntsman, D.; Suriano,
ano,
- [108666]G.; Machado, J. C.; Kiviluoto, T.; Aaltonen, L.; Jackson, C.
E.; Seruca,
- [108667]R.; Caldas, C.: Screening E-cadherin in gastric cancer families reveals
germline mutations only in hereditary diffuse gastric cancer kindred. Hum.
Mutat. 19: 510–517, 2002.
- [108670]19143.
- [108671]Overduin, M.; Harvey, T. S.; Bagby, S.; Tong, K. I.; Yau, P.;
- [108672]Takeichi, M.; Ikura, M.: Solution structure of the epithelial
cadherin
domain responsible for selective cell adhesion. Science
267: 386–389,
1995.
- [108675]19144.
- [108676]Perl, A.-K.; Wilgenbus, P.; Dahl, U.; Semb, H.; Christofori,
G.
- [108677]: A causal role for E-cadherin in the transition from adenoma to carcinoma. Nature 392:

[108678]190–193, 1998.

[108679]19145.

[108680]Joenje, H.: Fanconi anaemia complementation groups in
Germany

[108681]and The Netherlands. Hum. Genet. 97: 280–282, 1996.

[108682]19146.

[108683]Joenje, H.; Oostra, A. B.; Wijker, M.; di Summa, F. M.; van
Berkel,

[108684]C. G. M.; Rooimans, M. A.; Ebell, W.; van Weel, M.; Pronk, J.
C.;

[108685]Buchwald, M.; Arwert, F.: Evidence for at least eight Fan-
coni anemia

[108686]genes. Am. J. Hum. Genet. 61: 940–944, 1997.

[108687]19147.

[108688]Rassool, F. V.; Le Beau, M. M.; Shen, M.–L.; Neilly, M. E.;
Espinosa,

[108689]R., III; Ong, S. T.; Boldog, F.; Drabkin, H.; McCarroll, R.;
McKeithan,

[108690]T. W.: Direct cloning of DNA sequences from the common
fragile site

[108691]region at chromosome band 3p14.2. Genomics 35:

109–117, 1996.

[108692]19148.

[108693]Rudduck, C.; Franzen, G.: A new heritable fragile site on human

[108694]chromosome 3. *Hereditas* 98: 297–299, 1983.

[108695]19149.

[108696]Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association

[108697]of FHIT (fragile histidine triad), a candidate tumour suppressor gene,

[108698]with the ubiquitin-conjugating enzyme hUBC9. *Biochem. J.* 352: 443–448,

[108699]2000.

[108700]19150.

[108701]Shiraishi, T.; Druck, T.; Mimori, K.; Flomenberg, J.; Berk, L.;

[108702]Alder, H.; Miller, W.; Huebner, K.; Croce, C. M.: Sequence conservation

[108703]at human and mouse orthologous common fragile regions, FRA3B/FHIT

[108704]and Fra14A2/Fhit. *Proc. Nat. Acad. Sci.* 98: 5722–5727, 2001.

[108705]19151.

[108706]Sozzi, G.; Veronese, M. L.; Negrini, M.; Baffa, R.; Cotticelli,

[108707]M. G.; Inoue, H.; Tornielli, S.; Pilotti, S.; De Gregorio, L.;
Pastorino,

[108708]U.; Pierotti, M. A.; Ohta, M.; Huebner, K.; Croce, C. M.: The
FHIT

[108709]gene at 3p14.2 is abnormal in lung cancer. Cell 85:
17-26, 1996.

[108710]19152.

[108711]Stein, C. K.; Glover, T. W.; Palmer, J. L.; Glisson, B. S.: Di-
rect

[108712]correlation between FRA3B expression and cigarette
smoking. Genes

[108713]Chromosomes Cancer 34: 333-340, 2002.

[108714]19153.

[108715]Virgilio, L.; Shuster, M.; Gollin, S. M.; Veronese, M. L.;
Ohta,

[108716]M.; Huebner, K.; Croce, C. M.: FHIT gene alterations in
head and

[108717]neck squamous cell carcinomas. Proc. Nat. Acad. Sci. 93:
9770-9775,

[108718]1996.

[108719]19154.

[108720]Wegner, R.-D.: A new inducible fragile site on chromosome 3 (p14.2)

[108721]in human lymphocytes. Hum. Genet. 63: 297-298, 1983.

[108722]19155.

[108723]Wegner, R.-D.: Reply to the letter of A. Markkanen, S. Knuutila,

[108724]and A. de la Chapelle. (Letter) Hum. Genet. 65: 218 only, 1983.

[108725]19156.

[108726]Zanesi, N.; Fidanza, V.; Fong, L. Y.; Mancini, R.; Druck, T.;

[108727]Valtieri, M.; Rudiger, T.; McCue, P. A.; Croce, C. M.; Huebner, K.

[108728]: The tumor spectrum in FHIT-deficient mice. Proc. Nat. Acad. Sci. 98:

[108729]10250-10255, 2001.

[108730]19157.

[108731]Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayasi, Y.;

[108732]Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura, N.: Prediction of the coding sequences of unidentified human genes. VI. The coding sequences of 80 new genes (KIAA0201–KIAA0280) deduced by analysis of cDNA clones from cell line KG–1 and brain. DNA Res. 3: 321–329, 1996.

[108736]19158.

[108737]19158.

[108738]Lee, E.; Marcucci, M.; Daniell, L.; Pypaert, M.; Weisz, O. A.; Ochoa, G.–C.; Farsad, K.; Wenk, M. R.; De Camilli, P.: Amphiphysin 2 (Bin1) and T-tubule biogenesis in muscle. Science 297: 1193–1196, 2002.

[108741]2002.

[108742]19159.

[108743]Negorev, D.; Riethman, H.; Wechsler–Reya, R.; Sakamuro, D.; Prendergast, G. C.; Simon, D.: The Bin1 gene localizes to human chromosome 2q14

[108745]by PCR analysis of somatic cell hybrids and fluorescence
in situ hybridization. Genomics 33:

[108746]329–331, 1996.

[108747]19160.

[108748]Sakamuro, D.; Elliott, K. J.; Wechsler-Reya, R.; Prender-
gast, G.

[108749]C.: BIN1 is a novel MYC–interacting protein with features
of a tumour

[108750]suppressor. Nature Genet. 14: 69–77, 1996.

[108751]19161.

[108752]Endo, Y.; Sato, Y.; Matsushita, M.; Fujita, T.: Cloning and
characterization

[108753]of the human lectin P35 gene and its related gene. Ge-
nomics 36:

[108754]515–521, 1996.

[108755]19162.

[108756]Betz, R. C.; Schoser, B. G. H.; Kasper, D.; Ricker, K.;
Ramirez,

[108757]A.; Stein, V.; Torbergesen, T.; Lee, Y.–A.; Nothen, M. M.;
Wienker,

[108758]T. F.; Malin, J.–P.; Propping, P.; Reis, A.; Mortier, W.;

Jentsch,

[108759]T. J.; Vorgerd, M.; Kubisch, C.: Mutations in CAV3 cause mechanical

[108760]hyperirritability of skeletal muscle in rippling muscle disease. Nature

[108761]Genet. 28: 218–219, 2001.

[108762]19163.

[108763]Drumheller, T.; McGillivray, B. C.; Behrner, D.; MacLeod, P.; McFadden,

[108764]D. E.; Roberson, J.; Venditti, C.; Chorney, K.; Chorney, M.; Smith,

[108765]D. I.: Precise localisation of 3p25 breakpoints in four patients

[108766]with the 3p– syndrome. J. Med. Genet. 33: 842–847, 1996.

[108767]19164.

[108768]Galbiati, F.; Volonte, D.; Chu, J. B.; Li, M.; Fine, S. W.; Fu,

[108769]M.; Bermudez, J.; Pedemonte, M.; Weidenheim, K. M.; Pestell, R. G.;

[108770]Minetti, C.; Lisanti, M. P.: Transgenic overexpression of caveolin-3

[108771]in skeletal muscle fibers induces a Duchenne-like muscu-

lar dystrophy

[108772]phenotype. Proc. Nat. Acad. Sci. 97: 9689–9694, 2000.

[108773]19165.

[108774]Wayne, S.; Robertson, N. G.; DeClau, F.; Chen, N.; Verhoeven, K.;

[108775]Prasad, S.; Tranebjarg, L.; Morton, C. C.; Ryan, A. F.; Van Camp,

[108776]G.; Smith, R. J. H.: Mutations in the transcriptional activator EYA4

[108777]cause late-onset deafness at the DFNA10 locus. Hum. Molec. Genet. 10:

[108778]195–200, 2001.

[108779]19166.

[108780]Kanai, Y.; Miura, K.; Uehara, T.; Amagai, M.; Takeda, O.; Tanuma,

[108781]S.; Kurosawa, Y.: Natural occurrence of Nuc in the sera of autoimmune-prone

[108782]MRL/lpr mice. Biochem. Biophys. Res. Commun. 196: 729–736, 1993.

[108783]19167.

[108784]Miura, K.; Hirai, M.; Kanai, Y.; Kurosawa, Y.: Organization

of

[108785]the human gene for nucleobindin (NUC) and its chromosomal assignment

[108786]to 19q13.2–q13.4. Genomics 34: 181–186, 1996.

[108787]19168.

[108788]Miura, K.; Titani, K.; Kurosawa, Y.; Kanai, Y.: Molecular cloning

[108789]of nucleobindin, a novel DNA-binding protein that contains both a

[108790]signal peptide and a leucine zipper structure. Biochem. Biophys.

[108791]Res. Commun. 187: 375–380, 1992.

[108792]19169.

[108793]Brewer, G.: Personal Communication. Winston–Salem, N.C. 3/4/1999.

[108794]19170.

[108795]Brewer, G.: An A+U-rich element RNA-binding factor regulates c-myc

[108796]mRNA stability in vitro. Molec. Cell. Biol. 11: 2460–2466, 1991.

[108797]19171.

[108798]Dempsey, L. A.; Li, M.; DePace, A.; Bray-Ward, P.; Maizels, N.

[108799]: The human HNRPD locus maps to 4q21 and encodes a highly conserved

[108800]protein. Genomics 49: 378–384, 1998.

[108801]19172.

[108802]Wagner, B. J.; DeMaria, C. T.; Sun, Y.; Wilson, G. M.; Brewer,

[108803]G.: Structure and genomic organization of the human AUF1 gene: alternative

[108804]pre-mRNA splicing generates four protein isoforms. Genomics 48:

[108805]195–202, 1998.

[108806]19173.

[108807]Wagner, B. J.; Long, L.; Rao, P. N.; Pettenati, M. J.; Brewer,

[108808]G.: Localization and physical mapping of genes encoding the A+U-rich

[108809]element RNA-binding protein AUF1 to human chromosomes 4 and X. Genomics 34:

[108810]219–222, 1996.

[108811]19174.

[108812]Zhang, W.; Wagner, B. J.; Ehrenman, K.; Schaefer, A. W.;
DeMaria,
[108813]C. T.; Crater, D.; DeHaven, K.; Long, L.; Brewer, G.: Purifi-
cation,
[108814]characterization and cDNA cloning of an AU-rich element
RNA-binding
[108815]protein, AUF1. Molec. Cell. Biol. 13: 7652-7665, 1993.
[108816]19175.
[108817]Connelly, M. A.; Grady, R. C.; Mushinski, J. F.; Marcu, K. B.:
[108818]PANG, a gene encoding a neuronal glycoprotein, is ectopi-
cally activated
[108819]by intracisternal A-type particle long terminal repeats in
murine
[108820]plasmacytomas. Proc. Nat. Acad. Sci. 91: 1337-1341,
1994.
[108821]19176.
[108822]Mock, B. A.; Connelly, M. A.; McBride, O. W.; Kozak, C. A.;
Marcu,
[108823]K. B.: Plasmacytoma-associated neuronal glycoprotein,
Pang, maps
[108824]to mouse chromosome 6 and human chromosome 3. Ge-
nomics 34: 226-228,

[108825]1996.

[108826]19177.

[108827]Bolino, A.; Seri, M.; Caroli, F.; Eubanks, J.; Srinivasan, J.;

[108828]Mandich, P.; Schenone, A.; Quattrone, A.; Romeo, G.; Catterall, W.

[108829]A.; Devoto, M.: Exclusion of the SCN2B gene as candidate for CMT4B. Europ.

[108830]J. Hum. Genet. 6: 629-634, 1998.

[108831]19178.

[108832]Eubanks, J.; Srinivasan, J.; Dinulos, M. B.; Distèche, C. M.; Catterall,

[108833]W. A.: Structure and chromosomal localization of the beta2 subunit

[108834]of the human brain sodium channel. Neuroreport 8: 2775-2779, 1997.

[108835]19179.

[108836]Imai, K.; Kingsley, D. M.: Mouse chromosome 9. Mammalian Genome 5:

[108837]S139-S153, 1994.

[108838]19180.

[108839]Noben–Trauth, K.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[108840]A.; Sonoda, G.; Testa, J. R.; Klempnauer, K.–H.: Mybl2 (Bmyb) maps

[108841]to mouse chromosome 2 and human chromosome 20q13.1. Genomics 35:

[108842]610–612, 1996.

[108843]19181.

[108844]Ando, A.; Kikuti, Y. Y.; Shigenari, A.; Kawata, H.; Okamoto, N.;

[108845]Shiina, T.; Chen, L.; Ikemura, T.; Abe, K.; Kimura, M.; Inoko, H.

[108846]: cDNA cloning of the human homologues of the mouse Ke4 and Ke6 genes

[108847]at the centromeric end of the human MHC region. Genomics 35: 600–602,

[108848]1996.

[108849]19182.

[108850]Aziz, N.; Maxwell, M. M.; Brenner, B. M.: Coordinate regulation

[108851]of 11–beta–HSD and Ke6 gene in cpk mouse: implications for steroid

[108852]metabolic defect in PKD. Am. J. Physiol. 267: F791–F797,
1994.

[108853]19183.

[108854]Aziz, N.; Maxwell, M. M.; St.–Jacques, B.; Brenner, B. M.:
Downregulation

[108855]of Ke 6, a novel gene encoded within the major histocom-
patibility

[108856]complex, in murine polycystic kidney disease. Molec. Cell.
Biol. 13:

[108857]1847–1853, 1993.

[108858]19184.

[108859]Kikuti, Y. Y.; Tamiya, G.; Ando, A.; Chen, L.; Kimura, M.;
Ferreira,

[108860]E.; Tsuji, K.; Trowsdale, J.; Inoko, H.: Physical mapping
220 kb

[108861]centromeric of the human MHC and DNA sequence analy-
sis of the 43–kb

[108862]segment including the RING1, HKE6, and HKE4 genes. Ge-
nomics 42:

[108863]422–435, 1997.

[108864]19185.

[108865]Macalma, T.; Otte, J.; Hensler, M. E.; Bockholt, S. M.; Louis,
[108866]H. A.; Kalff-Suske, M.; Grzeschik, K.-H.; von der Ahe, D.;
Beckerle,
[108867]M. C.: Molecular characterization of human zyxin. J. Biol.
Chem. 271:
[108868]31470-31478, 1996.
[108869]19186.
[108870]Zumbrunn, J.; Trueb, B.: Assignment of the ZYX gene for
the LIM
[108871]protein zyxin to human chromosome bands 7q34-q35 by
in situ hybridization. Cytogenet.
[108872]Cell Genet. 81: 283-284, 1998.
[108873]19187.
[108874]Jacobsen, L.; Madsen, P.; Moestrup, S. K.; Lund, A. H.;
Tommerup,
[108875]N.; Nykjaer, A.; Sottrup-Jensen, L.; Gliemann, J.; Petersen,
C. M.
[108876]: Molecular characterization of a novel human hybrid-type
receptor
[108877]that binds the alpha(2)-macroglobulin receptor-associ-
ated protein. J.

[108878]Biol. Chem. 271: 31379–31383, 1996.

[108879]19188.

[108880]Senechal, K.; Halpern, J.; Sawyers, C. L.: The CRKL adaptor protein

[108881]transforms fibroblasts and functions in transformation by the BCR-ABL

[108882]oncogene. J. Biol. Chem. 38: 23255–23261, 1996.

[108883]19189.

[108884]ten Hoeve, J.; Morris, C.; Heisterkamp, N.; Groffen, J.: Iso-lation

[108885]and chromosomal localization of CRKL, a human crk-like gene. Oncogene 8:

[108886]2469–2474, 1993.

[108887]19190.

[108888]Cheung, A. H.; Stewart, R. J.; Marsden, P. A.: Endothelial Tie2/Tek

[108889]ligands angiopoietin-1 (ANGPT1) and angiopoietin-2 (ANGPT2): regional

[108890]localization of the human genes to 8q22.3–q23 and 8p23. Genomics 48:

[108891]389–391, 1998.

[108892]19191.

[108893]Davis, S.; Aldrich, T. H.; Jones, P. F.; Acheson, A.; Compton,

[108894]D. L.; Jain, V.; Ryan, T. E.; Bruno, J.; Radziejewski, C.;
Maisonpierre,

[108895]P. C.; Yancopoulos, G. D.: Isolation of angiopoietin-1, a
ligand

[108896]for the TIE2 receptor, by secretion-trap expression
cloning. Cell 87:

[108897]1161-1169, 1996.

[108898]19192.

[108899]Folkman, J.; D'Amore, P. A.: Blood vessel formation: what
is its

[108900]molecular basis? Cell 87: 1153-1155, 1996.

[108901]19193.

[108902]Marziliano, N.; Crovella, S.; Audero, E.; Pecile, V.; Bus-
solino,

[108903]F.; Amoroso, A.; Garagna, S.: Genetic mapping of the
mouse homologue

[108904]of the human angiopoietin-1 gene (Agpt) to mouse chro-
mosome 9E2 by

[108905]in situ hybridization. Cytogenet. Cell Genet. 87: 199–200, 1999.

[108906]19194.

[108907]Sato, T. N.; Tozawa, Y.; Deutsch, U.; Wolburg–Buchholz, K.; Fujiwara,

[108908]Y.; Gendron–Maguire, M.; Gridley, T.; Wolburg, H.; Risau, W.; Qin,

[108909]Y.: Distinct roles of the receptor tyrosine kinases Tie–1 and Tie–2

[108910]in blood vessel formation. Nature 376: 70–73, 1995.

[108911]19195.

[108912]Suri, C.; Jones, P. F.; Patan, S.; Bartunkova, S.; Maison–pierre,

[108913]P. C.; Davis, S.; Sato, T. N.; Yancopoulos, G. D.: Requisite role

[108914]of angiopoietin–1, a ligand for the TIE2 receptor, during embryonic

[108915]angiogenesis. Cell 87: 1171–1180, 1996.

[108916]19196.

[108917]Suri, C.; McClain, J.; Thurston, G.; McDonald, D. M.; Zhou, H.;

[108918]Oldmixon, E. H.; Sato, T. N.; Yancopoulos, G. D.: Increased vascularization

[108919]in mice overexpressing angiopoietin-1. Science 282: 468-471, 1998.

[108920]19197.

[108921]Valenzuela, D. M.; Griffiths, J. A.; Rojas, J.; Aldrich, T. H.;

[108922]Jones, P. F.; Zhou, H.; McClain, J.; Copeland, N. G.; Gilbert, D.

[108923]J.; Jenkins, N. A.; Huang, T.; Papadopoulos, N.; Maison-pierre, P.

[108924]C.; Davis, S.; Yancopoulos, G. D.: Angiopoietins 3 and 4: diverging

[108925]gene counterparts in mice and humans. Proc. Nat. Acad. Sci. 96:

[108926]1904-1909, 1999.

[108927]19198.

[108928]Chestkov, A. V.; Baka, I. D.; Kost, M. V.; Georgiev, G. P.; Buchman,

[108929]V. L.: The d4 gene family in the human genome. Genomics 36: 174-177,

[108930]1996.

[108931]19199.

[108932]Gabig, T. G.; Crean, C. D.; Klenk, A.; Long, H.; Copeland, N. G.;

[108933]Gilbert, D. J.; Jenkins, N. A.; Quincey, D.; Parente, F.; Lespinasse,

[108934]F.; Carle, G. F.; Gaudray, P.; and 13 others: Expression and chromosomal

[108935]localization of the Requiem gene. Mammalian Genome 9: 660–665, 1998.

[108936]19200.

[108937]Gabig, T. G.; Mantel, P. L.; Rosli, R.; Crean, C. D.: Requiem:

[108938]a novel zinc finger gene essential for apoptosis in myeloid cells. J.

[108939]Biol. Chem. 269: 29515–29519, 1994.

[108940]19201.

[108941]Smith, J. E., Jr.; Bollekens, J. A.; Inghirami, G.; Takeshita,

[108942]K.: Cloning and mapping of the MEIS1 gene, the human homolog of a

[108943]murine leukemogenic gene. Genomics 43: 99–103, 1997.

[108944]19202.

[108945]Nakanishi-Matsui, M.; Zheng, Y.-W.; Sulciner, D. J.; Weiss, E.

[108946]J.; Ludeman, M. J.; Coughlin, S. R.: PAR3 is a cofactor for PAR4

[108947]activation by thrombin. Nature 404: 609-613, 2000.

[108948]19203.

[108949]Sambrano, G. R.; Weiss, E. J.; Zheng, Y.-W.; Huang, W.; Coughlin,

[108950]S. R.: Role of thrombin signalling in platelets in haemostasis and

[108951]thrombosis. Nature 413: 74-78, 2001.

[108952]19204.

[108953]Crosnier, C.; Driancourt, C.; Raynaud, N.; Dhorne-Pollet, S.; Pollet,

[108954]N.; Bernard, O.; Hadchouel, M.; Meunier-Rotival, M.: Mutations in

[108955]JAGGED1 gene are predominantly sporadic in Alagille syndrome. Gastroenterology 116:

[108956]1141-1148, 1999.

[108957]19205.

[108958]Liese, J.; Kloos, S.; Jendrossek, V.; Petropoulou, T.; Win-

tergerst,

[108959]U.; Notheis, G.; Gahr, M.; Belohradsky, B. H.: Long-term follow-up

[108960]and outcome of 39 patients with chronic granulomatous disease. J.

[108961]Pediat. 137: 687-693, 2000.

[108962]19206.

[108963]Parkos, C. A.; Allen, R. A.; Cochrane, C. G.; Jesaitis, A. J.

[108964]: Purified cytochrome b from human granulocyte plasma membrane is

[108965]comprised of two polypeptides with relative molecular weights of 91,000

[108966]and 22,000. J. Clin. Invest. 80: 732-742, 1987.

[108967]19207.

[108968]Parkos, C. A.; Dinanuer, M. C.; Walker, L. E.; Allen, R. A.; Jesaitis,

[108969]A. J.; Orkin, S. H.: Primary structure and unique expression of the

[108970]22-kilodalton light chain of human neutrophil cytochrome b. Proc.

[108971]Nat. Acad. Sci. 85: 3319-3323, 1988.

[108972]19208.

[108973]Schapiro, B. L.; Newburger, P. E.; Klempner, M. S.; Dinauer, M.

[108974]C.: Chronic granulomatous disease presenting in a 69-year-old man. New

[108975]Eng. J. Med. 325: 1786-1790, 1991.

[108976]19209.

[108977]Stasia, M. J.; Bordigoni, P.; Martel, C.; Morel, F.: A novel

[108978]and unusual case of chronic granulomatous disease in a child with

[108979]a homozygous 36-bp deletion in the CYBA gene (A22-0) leading to the

[108980]activation of a cryptic splice site in intron 4. Hum. Genet. 110:

[108981]444-450, 2002.

[108982]19210.

[108983]Ushio-Fukai, M.; Zafari, A. M.; Fukui, T.; Ishizaka, N.; Griendling,

[108984]K.: p22-phox is a critical component of the superoxide-generating

[108985]NADH/NADPH oxidase system and regulates angiotensin

II-induced hypertrophy

[108986]in vascular smooth muscle cells. J. Biol. Chem. 271:
23317-23321,

[108987]1996.

[108988]19211.

[108989]Weening, R. S.; Corbeel, L.; de Boer, M.; Lutter, R.; van
Zwieten,

[108990]R.; Hamers, M. N.; Roos, D.: Cytochrome b deficiency in an
autosomal

[108991]form of chronic granulomatous disease: a third form of
chronic granulomatous

[108992]disease recognized by monocyte hybridization. J. Clin. In-
vest. 75:

[108993]915-920, 1985.

[108994]19212.

[108995]Yamada, M.; Ariga, T.; Kawamura, N.; Ohtsu, M.; Imajoh-
Ohmi, S.;

[108996]Ohshika, E.; Tatsuzawa, O.; Kobayashi, K.; Sakiyama, Y.:
Genetic

[108997]studies of three Japanese patients with
p22-phox-deficient chronic

[108998]granulomatous disease: detection of a possible common

mutant CYBA

[108999]allele in Japan and a genotype–phenotype correlation in these patients. Brit.

[109000]J. Haemat. 108: 511–517, 2000.

[109001]19213.

[109002]Francke, U.; Hsieh, C.–L.; Foellmer, B. E.; Lomax, K. J.; Malech,

[109003]H. L.; Leto, T. L.: Genes for two autosomal recessive forms of chronic

[109004]granulomatous disease assigned to 1q25 (NCF2) and 7q11.23 (NCF1). Am.

[109005]J. Hum. Genet. 47: 483–492, 1990.

[109006]19214.

[109007]Chasse, J. F.; Paul, V.; Escanez, R.; Kamoun, P.; London, J.:

[109008]Human cystathionine beta–synthase: gene organization and expression

[109009]of different 5–prime alternative splicing. Mammalian Genome 8: 917–921,

[109010]1997.

[109011]19215.

[109012]Paulweber, B.; Wiebusch, H.; Miesenboeck, G.; Funke, H.;

Assmann,

[109013]G.; Hoelzl, B.; Sippl, M. J.; Friedl, W.; Patsch, J. R.; Sand-
hofer,

[109014]F.: Molecular basis of lipoprotein lipase deficiency in two
Austrian

[109015]families with type I hyperlipoproteinemia. Atherosclerosis
86: 239-250,

[109016]1991.

[109017]19216.

[109018]Andersson, S.; Berman, D. M.; Jenkins, E. P.; Russell, D. W.:

[109019]Deletion of steroid 5-alpha-reductase 2 gene in male
pseudohermaphroditism. Nature 354:

[109020]159-161, 1991.

[109021]19217.

[109022]Boudon, C.; Lobaccaro, J. M.; Lumbroso, S.; Ogur, G.; Ocal,
G.;

[109023]Belon, C.; Sultan, C.: A new deletion of the
5-alpha-reductase type

[109024]2 gene in a Turkish family with 5-alpha-reductase defi-
ciency. Clin.

[109025]Endocr. 43: 183-188, 1995.

[109026]19218.

[109027]Cai, L.-Q.; Zhu, Y.-S.; Katz, M. D.; Herrera, C.; Baez, J.;
DeFillo-Ricart,

[109028]M.; Shackleton, C. H. L.; Imperato-McGinley, J.:
5-Alpha-reductase-2

[109029]gene mutations in the Dominican Republic. J. Clin. Endocr.
Metab. 81:

[109030]1730-1735, 1996.

[109031]19219.

[109032]Can, S.; Zhu, Y.-S.; Cai, L.-Q.; Ling, Q.; Katz, M. D.; Ak-
gun,

[109033]S.; Shackleton, C. H. L.; Imperato-McGinley, J.: The identi-
fication

[109034]of 5-alpha-reductase-2 and 17-beta-hydroxysteroid de-
hydrogenase-3

[109035]gene defects in male pseudohermaphrodites from a Turk-
ish kindred. J.

[109036]Clin. Endocr. Metab. 83: 560-569, 1998.

[109037]19220.

[109038]Chavez, B.; Valdez, E.; Vilchis, F.: Uniparental disomy in
steroid

[109039]5-alpha-reductase 2 deficiency. J. Clin. Endocr. Metab. 85:
3147-3150,

[109040]2000.

[109041]19221.

[109042]De Vaal, O. M.: Genital intersexuality in three brothers,
connected

[109043]with consanguineous marriages in the three previous gen-
erations. Acta

[109044]Paediat. 44: 35-39, 1955.

[109045]19222.

[109046]Forti, G.; Falchetti, A.; Santoro, S.; Davis, D. L.; Wilson, J.

[109047]D.; Russell, D. W.: Steroid 5 alpha-reductase 2 deficiency:
virilization

[109048]in early infancy may be due to partial function of mutant
enzyme. Clin.

[109049]Endocr. 44: 477–482, 1996.

[109050]19223.

[109051]Greene, S. A.; Symes, E.; Brook, C. G. D.:

5-Alpha-reductase deficiency

[109052]causing male pseudohermaphroditism. Arch. Dis. Child.

53: 751–753,

[109053]1978.

[109054]19224.

[109055]Hochberg, Z.; Chayen, R.; Reiss, N.; Falik, Z.; Makler, A.;

Munichor,

[109056]M.; Farkas, A.; Goldfarb, H.; Ohana, N.; Hiort, O.: Clinical,

biochemical,

[109057]and genetic findings in a large pedigree of male and fe-

male patients

[109058]with 5-alpha-reductase 2 deficiency. J. Clin. Endocr.

Metab. 81:

[109059]2821–2827, 1996.

[109060]19225.

[109061]Imperato-McGinley, J.; Gautier, T.: Inherited

5-alpha-reductase

[109062]deficiency in man. Trends Genet. 2: 130–133, 1986.

[109063]19226.

[109064]Imperato-McGinley, J.; Guerrero, L.; Gautier, T.; German, J.
L.;

[109065]Peterson, R. E.: Steroid-5-alpha-reductase deficiency in
man. An

[109066]inherited form of male pseudohermaphroditism.In:
Bergsma, D.: Genetic

[109067]Forms of Hypogonadism. New York: National Foundation-
March of Dimes

[109068](pub.) 1975. Pp. 91-103.

[109069]19227.

[109070]Imperato-McGinley, J.; Guerrero, L.; Gautier, T.; Peterson,
R.

[109071]E.: Steroid 5-alpha-reductase deficiency in man: an inher-
ited form

[109072]of male pseudohermaphroditism. Science 186:
1213-1215, 1974.

[109073]19228.

[109074]Imperato-McGinley, J.; Miller, M.; Wilson, J. D.; Peterson,
R.

[109075]E.; Shackleton, C.; Gajdusek, D. C.: A cluster of male
pseudohermaphrodites

[109076]with 5- α -reductase deficiency in Papua New Guinea.

Clin. Endocr. 34:

[109077]293-298, 1991.

[109078]19229.

[109079]Imperato-McGinley, J.; Peterson, R. E.; Gautier, T.; Sturla, E.

[109080]; Androgens and the evolution of male-gender identity among male pseudohermaphrodites

[109081]with 5- α -reductase deficiency. New Eng. J. Med. 300: 1233-1237,

[109082]1979.

[109083]19230.

[109084]Chrzanowska, B. L.; Nitzan, M.; Phillips, L. S.; Schulman, J.

[109085]D.: Homocysteic acid: an examination of its possible growth hormone-like

[109086]activity. Metabolism 28: 80-84, 1979.

[109087]19231.

[109088]Cochran, F. B.; Sweetman, L.; Schmidt, K.; Barsh, G.; Kraus, J.;

[109089]Packman, S.: Pyridoxine-unresponsive homocystinuria with an unusual

[109090]clinical course. Am. J. Med. Genet. 35: 519–522, 1990.

[109091]19232.

[109092]Collins, J. E.; Brenton, D. P.: Pancreatitis and homocystinuria. J.

[109093]Inherit. Metab. Dis. 13: 232–233, 1990.

[109094]19233.

[109095]Di Minno, G.; Davi, G.; Margaglione, M.; Cirillo, F.; Grandone,

[109096]E.; Ciabattini, G.; Catalano, I.; Strisciuglio, P.; Andria, G.; Patrono,

[109097]C.; Mancini, M.: Abnormally high thromboxane biosynthesis in homozygous

[109098]homocystinuria: evidence for platelet involvement and probucol-sensitive

[109099]mechanism. J. Clin. Invest. 92: 1400–1406, 1993.

[109100]19234.

[109101]Falcon, C. R.; Cattaneo, M.; Panzeri, D.; Martinelli, I.; Mannucci,

[109102]P. M.: High prevalence of hyperhomocyst(e)inemia in patients with

[109103]juvenile venous thrombosis. Arteriosclerosis Thromb. 14:

- 1080–1083,
[109104]1994.
- [109105]19235.
- [109106]Field, C. M. B.; Carson, N. A. J.; Cusworth, D. C.; Dent, C.
E.;
- [109107]Neill, D. W.: Homocystinuria, a new disorder of
metabolism. (Abstract) Proc.
- [109108]10th Int. Cong. Pediat., Lisbon 274–275, 1962.
- [109109]19236.
- [109110]Fowler, B.; Kraus, J.; Packman, S.; Rosenberg, L. E.: Homo-
cystinuria:
- [109111]evidence for three distinct classes of cystathionine beta-
synthetase
- [109112]mutants in cultured fibroblasts. J. Clin. Invest. 61:
645–653, 1978.
- [109113]19237.
- [109114]Franco, R. F.; Elion, J.; Lavinha, J.; Krishnamoorthy, R.;
Tavella,
- [109115]M. H.; Zago, M. A.: Heterogeneous ethnic distribution of
the 844ins68
- [109116]in the cystathionine beta-synthase gene. Hum. Hered. 48:

338–342,

[109117]1998.

[109118]19238.

[109119]Franken, D. G.; Boers, G. H. J.; Blom, H. J.; Cruysberg, J. R.

[109120]M.; Trijbels, F. J. M.; Hamel, B. C. J.: Prevalence of familial
mild

[109121]hyperhomocysteinemia. Atherosclerosis 125: 71–80,
1996.

[109122]19239.

[109123]Frimpter, G. W.: Homocystinuria: vitamin B6 dependent or
not?

[109124](Editorial) Ann. Intern. Med. 71: 209–211, 1969.

[109125]19240.

[109126]Fryer, R. H.; Wilson, B. D.; Gubler, D. B.; Fitzgerald, L. A.;

[109127]Rodgers, G. M.: Homocysteine, a risk factor for premature
vascular

[109128]disease and thrombosis, induces tissue factor activity in
endothelial

[109129]cells. Arteriosclerosis Thromb. 13: 1327–1333, 1993.

[109130]19241.

- [109131]Gallagher, P. M.; Naughten, E.; Hanson, N. Q.; Schwichtenberg,
berg,
- [109132]K.; Bignell, M.; Yuan, M.; Ward, P.; Yap, S.; Whitehead, A.
S.; Tsai,
M. Y.: Characterization of mutations in the cystathionine
beta-synthase
gene in Irish patients with homocystinuria. Molec. Genet.
Metab. 65:
298–302, 1998.
- [109135]19242.
- [109136]19242.
- [109137]Gallagher, P. M.; Ward, P.; Tan, S.; Naughten, E.; Kraus, J.
P.;
- [109138]Sellar, G. C.; McConnell, D. J.; Graham, I.; Whitehead, A. S.:
High
frequency (71%) of cystathionine beta-synthase mutation
G307S in Irish
homocystinuria patients. Hum. Mutat. 6: 177–180, 1995.
- [109140]19243.
- [109141]19243.
- [109142]Gaustadnes, M.; Ingerslev, J.; Rutiger, N.: Prevalence of
congenital
homocystinuria in Denmark. (Letter) New Eng. J. Med. 340:
1513 only,

[109144]1999.

[109145]19244.

[109146]Gaustadnes, M.; Rudiger, N.; Rasmussen, K.; Ingerslev, J.:
Intermediate

[109147]and severe hyperhomocysteinemia with thrombosis: a
study of genetic

[109148]determinants. Thromb. Haemost. 83: 554-558, 2000.

[109149]19245.

[109150]Gaustadnes, M.; Wilcken, B.; Oliveriusova, J.; McGill, J.;
Fletcher,

[109151]J.; Kraus, J. P.; Wilcken, D. E.: The molecular basis of cys-
tathionine

[109152]beta-synthase deficiency in Australian patients: genotype-
phenotype

[109153]correlations and response to treatment. Hum. Mutat. 20:
117-126,

[109154]2002.

[109155]19246.

[109156]Gerding, H.: Ocular complications and a new surgical ap-
proach

[109157]to lens dislocation in homocystinuria due to cystathion-

ine-beta-synthetase

[109158]deficiency. Europ. J. Pediat. 157 (suppl. 2): S94-S101,
1998.

[109159]19247.

[109160]Gerritsen, T.; Vaughn, J. G.; Waisman, H. A.: The identifi-
cation

[109161]of homocystine in the urine. Biochem. Biophys. Res. Com-
mun. 9: 493-496,

[109162]1962.

[109163]19248.

[109164]Goldstein, J. L.; Campbell, B. K.; Gartler, S. M.: Homo-
cystinuria:

[109165]heterozygote detection using phytohemagglutinin-stimu-
lated lymphocytes. J.

[109166]Clin. Invest. 52: 218, 1973.

[109167]19249.

[109168]Goldstein, J. L.; Campbell, B. K.; Gartler, S. M.: Cystathion-
ine

[109169]synthetase activity in human lymphocytes: induction by
phytohemagglutinin. J.

[109170]Clin. Invest. 51: 1034-1037, 1972.

[109171]19250.

[109172]Gu, Z.; Ramesh, V.; Kozich, V.; Korson, M. S.; Kraus, J. P.;
Shih,

[109173]V. E.: Identification of a molecular genetic defect in homo-
cystinuria

[109174]due to cystathionine beta-synthase deficiency. (Abstract)
Am. J.

[109175]Hum. Genet. 49: 406 only, 1991.

[109176]19251.

[109177]Guttormsen, A. B.; Ueland, P. M.; Kruger, W. D.; Kim, C. E.;
Ose,

[109178]L.; Folling, I.; Refsum, H.: Disposition of homocysteine in
subjects

[109179]heterozygous for homocystinuria due to cystathionine
beta-synthase

[109180]deficiency: relationship between genotype and phenotype.
Am. J. Med.

[109181]Genet. 100: 204-213, 2001.

[109182]19252.

[109183]Harker, L. A.; Ross, R.; Slichter, S. J.; Scott, C. R.: Homo-
cystine-induced

- [109184]arteriosclerosis: the role of endothelial cell injury and platelet
- [109185]response in its genesis. J. Clin. Invest. 58: 731–741, 1976.
- [109186]19253.
- [109187]Harker, L. A.; Slichter, S. J.; Scott, C. R.; Ross, R.: Homocystinuria:
- [109188]vascular injury and arterial thrombosis. New Eng. J. Med. 291: 537–543,
- [109189]1974.
- [109190]19254.
- [109191]Harrison, D. A.; Mullaney, P. B.; Mesfer, S. A.; Awad, A. H.;
- [109192]Dhindsa, H.: Management of ophthalmic complications of homocystinuria. Ophthalmology 105:
- [109193]1886–1890, 1998.
- [109194]19255.
- [109195]Hooft, C.; Carton, D.; Samyn, W.: Pyridoxine treatment in homocystinuria.
- [109196](Letter) Lancet I: 1384 only, 1967.
- [109197]19256.
- [109198]Hu, F. L.; Gu, Z.; Kozich, V.; Kraus, J. P.; Ramesh, V.; Shih,

[109199]V. E.: Molecular basis of cystathionine beta-synthase deficiency

[109200]in pyridoxine responsive and nonresponsive homocystinuria. Hum. Molec.

[109201]Genet. 2: 1857-1860, 1993.

[109202]19257.

[109203]Janosik, M.; Oliveriusova, J.; Janosikova, B.; Sokolova, J.; Kraus,

[109204]E.; Kraus, J. P.; Kozich, V.: Impaired heme binding and aggregation

[109205]of mutant cystathionine beta-synthase subunits in homocystinuria. Am.

[109206]J. Hum. Genet. 68: 1506-1513, 2001.

[109207]19258.

[109208]Kaeser, A. C.; Rodnight, R.; Ellis, B. A.: Psychiatric and biochemical

[109209]aspects of a case of homocystinuria. J. Neurol. Neurosurg. Psychiat. 32:

[109210]88-93, 1969.

[109211]19259.

[109212]Kang, S.-S.; Wong, P. W. K.; Cook, H. Y.; Norusis, M.;

Messer,

[109213]J. V.: Protein-bound homocyst(e)ine: a possible risk factor
for coronary

[109214]artery disease. J. Clin. Invest. 77: 1482–1486, 1986.

[109215]19260.

[109216]Kery, V.; Poneleit, L.; Meyer, J. D.; Manning, M. C.; Kraus, J.

[109217]P.: Binding of pyridoxal 5-prime-phosphate to the heme
protein human

[109218]cystathionine beta-synthase. Biochemistry 38:
2716–2724, 1999.

[109219]19261.

[109220]Kim, C. E.; Gallagher, P. M.; Guttormsen, A. B.; Refsum, H.;
Ueland,

[109221]P. M.; Ose, L.; Folling, I.; Whitehead, A. S.; Tsai, M. Y.;
Kruger,

[109222]W. D.: Functional modeling of vitamin responsiveness in
yeast: a

[109223]common pyridoxine-responsive cystathionine beta-
synthase mutation

[109224]in homocystinuria. Hum. Molec. Genet. 6: 2213–2221,
1997.

[109225]19262.

- [109226]Giralt, M.; Park, E. A.; Gurney, A. L.; Liu, J.; Hakimi, P.;
Hanson,
[109227]R. W.: Identification of a thyroid hormone response ele-
ment in the
[109228]phosphoenolpyruvate carboxykinase (GTP) gene: evidence
for synergistic
[109229]interaction between thyroid hormone and cAMP cis-
regulatory elements. J.
[109230]Biol. Chem. 266: 21991-21996, 1991.
[109231]19263.
[109232]Pilz, A. J.; Willer, E.; Povey, S.; Abbott, C. M.: The genes
coding
[109233]for phosphoenolpyruvate carboxykinase-1 (PCK1) and
neuronal nicotinic
[109234]acetylcholine receptor alpha-4 subunit (CHRNA4) map to
human chromosome
[109235]20, extending the known region of homology with mouse
chromosome 2. Ann.
[109236]Hum. Genet. 56: 289-293, 1992.
[109237]19264.
[109238]Olswang, Y.; Cohen, H.; Papo, O.; Cassuto, H.; Croniger, C.
M.;

[109239]Hakimi, P.; Tilghman, S. M.; Hanson, R. W.; Reshef, L.: A mutation
[109240]in the peroxisome proliferator-activated receptor gamma-binding site
[109241]in the gene for the cytosolic form of phosphoenolpyruvate carboxykinase
[109242]reduces adipose tissue size and fat content in mice. Proc. Nat. Acad.
[109243]Sci. 99: 625-630, 2002.
[109244]19265.
[109245]Stoffel, M.; Xiang, K.; Espinosa, R., III; Cox, N. J.; Le Beau,
[109246]M. M.; Bell, G. I.: cDNA sequence and localization of polymorphic
[109247]human cytosolic phosphoenolpyruvate carboxykinase gene (PCK1) to chromosome
[109248]20, band q13.31: PCK1 is not tightly linked to maturity-onset diabetes
[109249]of the young. Hum. Molec. Genet. 2: 1-4, 1993.
[109250]19266.
[109251]Ting, C.-N.; Burgess, D. L.; Chamberlain, J. S.; Keith, T. P.;
[109252]Falls, K.; Meisler, M. H.: Phosphoenolpyruvate carboxyki-

nase (GTP):

[109253] characterization of the human PCK1 gene and localization
distal to

[109254] MODY on chromosome 20. Genomics 16: 698–706, 1993.

[109255] 19267.

[109256] Vidnes, J.; Sovik, O.: Gluconeogenesis in infancy and
childhood.

[109257] III. Deficiency of the extramitochondrial form of hepatic
phosphoenolpyruvate

[109258] carboxykinase in a case of persistent neonatal hypogly-
caemia. Acta

[109259] Paediat. Scand. 65: 301–312, 1976.

[109260] 19268.

[109261] Yu, H.; Thun, R.; Chandrasekharappa, S.; Trent, J. M.;
Zhang, J.;

[109262] Meisler, M. H.: Human PCK1 encoding phosphoenolpyru-
vate carboxykinase

[109263] is located on chromosome 20q13.2. Genomics 15:
219–221, 1993.

[109264] 19269.

[109265] Abarbanel, J. M.; Bashan, N.; Potashnik, R.; Osimani, A.;

Moses,

[109266]S. W.; Herishanu, Y.: Adult muscle phosphorylase 'b' kinase deficiency. *Neurology* 36:

[109267]560–562, 1986.

[109268]19270.

[109269]Pepe, G.; Chimienti, G.; Resta, F.; Di Perna, V.; Tarricone, C.;

[109270]Lovecchio, M.; Colacicco, A. M.; Capurso, A.: A new Italian case

[109271]of lipoprotein lipase deficiency: a leu365-to-val change resulting

[109272]in loss of enzyme activity. *Biochem. Biophys. Res. Commun.* 199:

[109273]570–576, 1994.

[109274]19271.

[109275]Reymer, P. W. A.; Groenemeyer, B. E.; Gagne, E.; Miao, L.; Appelman,

[109276]E. E. G.; Seidel, J. C.; Kromhout, D.; Bijvoet, S. M.; van de Oever,

[109277]K.; Bruin, T.; Hayden, M. R.; Kastelein, J. J. P.: A frequently occurring

[109278]mutation in the lipoprotein lipase gene (Asn291Ser) con-

tributes to

[109279]the expression of familial combined hyperlipidemia. Hum.

Molec. Genet. 4:

[109280]1543–1549, 1995.

[109281]19272.

[109282]Samuels, M. E.; Forbey, K. C.; Reid, J. E.; Abkevich, V.;

Bulka,

[109283]K.; Wardell, B. R.; Bowen, B. R.; Hopkins, P. N.; Hunt, S. C.;

Ballinger,

[109284]D. G.; Skolnick, M. H.; Wagner, S.: Identification of a com-

mon variant

[109285]in the lipoprotein lipase gene in a large Utah kindred as-

certained

[109286]for coronary heart disease: the -93G/D9N variant predis-

poses to low

[109287]HDL-C/high triglycerides. Clin. Genet. 59: 88–98, 2001.

[109288]19273.

[109289]Schreibman, P. H.; Arons, D. L.; Saudek, C. D.; Arky, R. A.:

[109290]Abnormal lipoprotein lipase in familial exogenous hyper-

triglyceridemia. J.

[109291]Clin. Invest. 52: 2074–2082, 1973.

[109292]19274.

[109293]Sprecher, D. L.; Kobayashi, J.; Rymaszewski, M.; Goldberg,
I.

[109294]J.; Harris, B. V.; Bellet, P. S.; Ameis, D.; Yunker, R. L.; Black,
[109295]D. M.; Stein, E. A.; Schotz, M. C.; Wiginton, D. A.:

Trp64-to-nonsense

[109296]mutation in the lipoprotein lipase gene. J. Lipid Res. 33:
859-866,

[109297]1992.

[109298]19275.

[109299]Sternowsky, H. J.; Gaertner, U.; Stahnkel, N.; Kaukel, E.:
Juvenile

[109300]familial hypertriglyceridemia and growth retardation: clin-
ical and

[109301]biochemical observations in three siblings. Europ. J. Pe-
diat. 125:

[109302]59-70, 1977.

[109303]19276.

[109304]Takagi, A.; Ikeda, Y.; Tsutsumi, Z.; Shoji, T.; Yamamoto,
A.:

[109305]Molecular studies on primary lipoprotein lipase (LPL) defi-

ciency:

[109306]one base deletion (G-916) in exon 5 of LPL gene causes
no detectable

[109307]LPL protein due to the absence of LPL mRNA transcript. J.
Clin. Invest. 89:

[109308]581-591, 1992.

[109309]19277.

[109310]Weinstock, P. H.; Bisgaier, C. L.; Aalto-Setälä, K.; Radner,
H.;

[109311]Ramakrishnan, R.; Levak-Frank, S.; Essenburg, A. D.;
Zechner, R.;

[109312]Breslow, J. L.: Severe hypertriglyceridemia, reduced high
density

[109313]lipoprotein, and neonatal death in lipoprotein lipase
knockout mice:

[109314]mild hypertriglyceridemia with impaired very low density
lipoprotein

[109315]clearance in heterozygotes. J. Clin. Invest. 96: 2555-2568,
1995.

[109316]19278.

[109317]Wessler, S.; Avioli, L. A.: Familial hyperlipoproteinemia.
J.A.M.A. 207:

[109318]929-937, 1969.

[109319]19279.

[109320]Wilson, D. E.; Edwards, C. Q.; Chan, I.-F.: Phenotypic heterogeneity

[109321]in the extended pedigree of a proband with lipoprotein lipase deficiency. Metabolism 32:

[109322]1107-1114, 1983.

[109323]19280.

[109324]Wilson, D. E.; Emi, M.; Iverius, P.-H.; Hata, A.; Wu, L. L.; Hillas,

[109325]E.; Williams, R. R.; Lalouel, J.-M.: Phenotypic expression of heterozygous

[109326]lipoprotein lipase deficiency in the extended pedigree of a proband

[109327]homozygous for a missense mutation. J. Clin. Invest. 86: 735-750,

[109328]1990.

[109329]19281.

[109330]Wilson, D. E.; Hata, A.; Kwong, L. K.; Lingam, A.; Shuhua, J.;

[109331]Ridinger, D. N.; Yeager, C.; Kaltenborn, K. C.; Iverius,

P.-H.; Lalouel,
[109332]J.-M.: Mutations in exon 3 of the lipoprotein lipase gene
segregating
[109333]in a family with hypertriglyceridemia, pancreatitis, and
non-insulin-dependent
[109334]diabetes. J. Clin. Invest. 92: 203-211, 1993.
[109335]19282.
[109336]Wion, K. L.; Kirchgessner, T. G.; Lusi, A. J.; Schotz, M. C.;
[109337]Lawn, R. M.: Human lipoprotein lipase complementary
DNA sequence. Science 235:
[109338]1638-1641, 1987.
[109339]19283.
[109340]Wittrup, H. H.; Tybjaerg-Hansen, A.; Abildgaard, S.; Steffensen,
[109341]R.; Schnohr, P.; Nordestgaard, B. G.: A common substitution (asn291ser)
[109342]in lipoprotein lipase is associated with increased risk of
ischemic
[109343]heart disease. J. Clin. Invest. 99: 1606-1613, 1997.
[109344]19284.
[109345]Wittrup, H. H.; Tybjaerg-Hansen, A.; Steffensen, R.; Deeb,

S.

- [109346]S.; Brunzell, J. D.; Jensen, G.; Nordestgaard, B. G.: Mutations in
the lipoprotein lipase gene associated with ischemic heart
disease
in men: the Copenhagen City Heart Study. *Arterioscler. Thromb. Vasc. Biol.* 19: 1535–1540, 1999.
- [109349]19285.
- [109351]Wood, S.; Schertzer, M.; Hayden, M.; Ma, Y.: Support for
founder
effect for two lipoprotein lipase (LPL) gene mutations in
French Canadians
by analysis of GT microsatellites flanking the LPL gene. *Hum. Genet.* 91: 312–316, 1993.
- [109355]19286.
- [109356]Burke, V.; Colebatch, J. H.; Anderson, C. M.; Simons, M. J.:
Association
of pancreatic insufficiency and chronic neutropenia in
childhood. *Arch. Dis. Child.* 42: 147–157, 1967.

[109359]19287.

[109360]Ando, T.; Klingberg, W. G.; Ward, A. N.; Rasmussen, K.;
Nyhan,

[109361]W. L.: Isovaleric acidemia presenting with altered
metabolism of

[109362]glycine. *Pediat. Res.* 5: 478-486, 1971.

[109363]19288.

[109364]Ando, T.; Nyhan, W. L.; Bachmann, C.; Rasmussen, K.;
Scott, R.;

[109365]Smith, E. K.: Isovaleric acidemia: identification of iso-
valerylglycine,

[109366]and 3-hydroxyisovalerate in urine of a patient previously
reported

[109367]as having butyric and hexanoic acidemia. *J. Pediat.* 82:
243-248,

[109368]1973.

[109369]19289.

[109370]Budd, M. A.; Tanaka, K. R.; Holmes, L. B.; Efron, M. L.;
Crawford,

[109371]J. D.; Isselbacher, K. J.: Isovaleric acidemia: clinical feature

[109372]of a new genetic defect of leucine metabolism. *New Eng. J.*

Med. 277:

[109373]321-327, 1967.

[109374]19290.

[109375]Cohn, R. M.; Yudkoff, R.; Rothman, R.; Segal, S.: Isovaleric
acidemia:

[109376]Use of glycine therapy in neonates. New Eng. J. Med. 299:
996-999,

[109377]1978.

[109378]19291.

[109379]de Sousa, C.; Chalmers, R. A.; Stacey, T. E.; Tracey, B. M.;
Weaver,

[109380]C. M.; Bradley, D.: The response to L-carnitine and glycine
therapy

[109381]in isovaleric acidaemia. Europ. J. Pediat. 144: 451-456,
1986.

[109382]19292.

[109383]Duran, M.; van Sprang, F. J.; Drewes, J. G.; Bruinvis, L.;
Ketling,

[109384]D.; Wadman, S. K.: Two sisters with isovaleric acidemia,
multiple

[109385]attacks of ketoacidosis and normal development. Europ. J.

Pediat. 131:

[109386]205–211, 1979.

[109387]19293.

[109388]Efron, M. L.: Isovaleric acidemia. Am. J. Dis. Child. 113:
74–76,

[109389]1967.

[109390]19294.

[109391]Kupfer, G. M.; Naf, D.; Suliman, A.; Pulsipher, M.;
D'Andrea, A.

[109392]D.: The Fanconi anaemia proteins, FAA and FAC, interact
to form a

[109393]nuclear complex. Nature Genet. 17: 487–490, 1997.

[109394]19295.

[109395]Kupfer, G. M.; Yamashita, T.; Naf, D.; Suliman, A.; Asano,
S.;

[109396]D'Andrea, A. D.: The Fanconi anemia polypeptide, FAC,
binds to the

[109397]cyclin-dependent kinase, cdc2. Blood 90: 1047–1054,
1997.

[109398]19296.

[109399]Liebetrau, W.; Budde, A.; Savoia, A.; Grummt, F.; Hoehn,

H.:

[109400]p53 activates Fanconi anemia group C gene expression.

Hum. Molec.

[109401]Genet. 6: 277–283, 1997.

[109402]19297.

[109403]Lo Ten Foe, J. R.; Kwee, M. L.; Rooimans, M. A.; Oostra, A.

B.;

[109404]Veermans, A. J. P.; van Weel, M.; Pauli, R. M.; Shahidi, N.

T.; Dokal,

[109405]I.; Roberts, I.; Altay, C.; Gluckman, E.; Gibson, R. A.;

Mathew, C.

[109406]G.; Arwert, F.; Joenje, H.: Somatic mosaicism in Fanconi anemia:

[109407]molecular basis and clinical significance. Europ. J. Hum.

Genet. 5:

[109408]137–148, 1997.

[109409]19298.

[109410]Lo Ten Foe, J. R.; Rooimans, M. A.; Joenje, H.; Arwert, F.:

Novel

[109411]frameshift mutation (1806insA) in exon 14 of the Fanconi anemia C

[109412]gene, FAC. Hum. Mutat. 7: 264–265, 1996.

[109413]19299.

[109414]Murer-Orlando, M.; Llerena, J. C., Jr.; Birjandi, F.; Gibson,

[109415]R. A.; Mathew, C. G.: FACC gene mutations and early pre-natal diagnosis

[109416]of Fanconi's anaemia. (Letter) Lancet 342: 686 only, 1993.

[109417]19300.

[109418]Pace, P.; Johnson, M.; Tan, W. M.; Mosedale, G.; Sng, C.; Hoatlin,

[109419]M.; de Winter, J.; Joenje, H.; Gergely, F.; Patel, K. J.: FANCE:

[109420]the link between Fanconi anaemia complex assembly and activity. EMBO

[109421]J. 21: 3414-3423, 2002.

[109422]19301.

[109423]Pang, Q.; Christianson, T. A.; Keeble, W.; Diaz, J.; Faulkner,

[109424]G. R.; Reifsteck, C.; Olson, S.; Bagby, G. C.: The Fanconi anemia

[109425]complementation group C gene product: structural evidence of multifunctionality. Blood 98:

[109426]1392-1401, 2001.

[109427]19302.

[109428]Pang, Q.; Fagerlie, S.; Christianson, T. A.; Keeble, W.;
Faulkner,

[109429]G.; Diaz, J.; Rathbun, R. K.; Bagby, G. C.: The Fanconi ane-
mia protein

[109430]FANCC binds to and facilitates the activation of STAT1 by
gamma interferon

[109431]and hematopoietic growth factors. Molec. Cell. Biol. 20:
4724-4735,

[109432]2000.

[109433]19303.

[109434]Riley, J.; Butler, R.; Ogilvie, D.; Finniear, R.; Jenner, D.;

[109435]Powell, S.; Anand, R.; Smith, J. C.; Markham, A. F.: A novel,
rapid

[109436]method for the isolation of terminal sequences from yeast
artificial

[109437]chromosome (YAC) clones. Nucleic Acids Res. 18:
2887-2890, 1990.

[109438]19304.

[109439]Roberts, R. G.; Coffey, A. J.; Bobrow, M.; Bentley, D. R.:
Determination

[109440]of the exon structure of the distal portion of the dys-
trophin gene

[109441]by vectorette PCR. Genomics 13: 942–950, 1992.

[109442]19305.

[109443]Savoia, A.; Centra, M.; Ianzano, L.; de Cillis, G. P.; Zelante,

[109444]L.; Buchwald, M.: Characterization of the 5-prime region
of the Fanconi

[109445]anaemia group C (FACC) gene. Hum. Molec. Genet. 4:
1321–1326, 1995.

[109446]19306.

[109447]Segal, G. M.; Magenis, R. E.; Brown, M.; Keeble, W.; Smith,
T.

[109448]D.; Heinrich, M. C.; Bagby, G. C., Jr.: Repression of Fanconi
anemia

[109449]gene (FACC) expression inhibits growth of hematopoietic
progenitor

[109450]cells. J. Clin. Invest. 94: 846–852, 1994.

[109451]19307.

[109452]Strathdee, C. A.; Duncan, A. M. V.; Buchwald, M.: Evidence
for

[109453]at least four Fanconi anaemia genes including FACC on

chromosome 9. Nature

[109454]Genet. 1: 196–198, 1992.

[109455]19308.

[109456]Verlander, P. C.; Kaporis, A.; Liu, Q.; Zhang, Q.; Seligsohn,

[109457]U.; Auerbach, A. D.: Carrier frequency of the IVS4 + 4 A-
to-T mutation

[109458]of the Fanconi anemia gene FAC in the Ashkenazi Jewish
population. Blood 86:

[109459]4034–4038, 1995.

[109460]19309.

[109461]Verlinsky, Y.; Rechitsky, S.; Schoolcraft, W.; Strom, C.;
Kuliev,

[109462]A.: Preimplantation diagnosis for Fanconi anemia com-
bined with HLA

[109463]matching. J.A.M.A. 285: 3130–3133, 2001.

[109464]19310.

[109465]Waisfisz, Q.; Morgan, N. V.; Savino, M.; de Winter, J. P.; van

[109466]Berkel, C. G. M.; Hoatlin, M. E.; Ianzano, L.; Gibson, R. A.;
Arwert,

[109467]F.; Savoia, A.; Mathew, C. G.; Pronk, J. C.; Joenje, H.: Spon-

taneous

[109468]functional correction of homozygous Fanconi anaemia alleles reveals

[109469]novel mechanistic basis for reverse mosaicism. Nature Genet. 22:

[109470]379–383, 1999.

[109471]19311.

[109472]Wevrick, R.; Barker, J. E.; Eppig, J. T.; Nadeau, J. H.; Buchwald,

[109473]M.: Expression of the Fanconi anemia group C gene in the mouse. (Abstract) Am.

[109474]J. Hum. Genet. 51 (suppl.): A137 only, 1992.

[109475]19312.

[109476]Wevrick, R.; Barker, J. E.; Nadeau, J. H.; Szpirer, C.; Buchwald,

[109477]M.: Mapping of the murine and rat Facc genes and assessment of flexed-tail

[109478]as a candidate mouse homolog of Fanconi anemia group C. Mammalian

[109479]Genome 4: 440–444, 1993.

[109480]19313.

[109481]Whitney, M. A.; Jakobs, P.; Kaback, M.; Moses, R. E.;

Grompe,

[109482]M.: The Ashkenazi Jewish Fanconi anemia mutation: incidence among

[109483]patients and carrier frequency in the at-risk population.

Hum. Mutat. 3:

[109484]339–341, 1994.

[109485]19314.

[109486]Whitney, M. A.; Royle, G.; Low, M. J.; Kelly, M. A.; Axthelm,

[109487]M. K.; Reifsteck, C.; Olson, S.; Braun, R. E.; Heinrich, M. C.;

Rathbun,

[109488]R. K.; Bagby, G. C.; Grompe, M.: Germ cell defects and hematopoietic

[109489]hypersensitivity to gamma-interferon in mice with a targeted disruption

[109490]of the Fanconi anemia C gene. Blood 88: 49–58, 1996.

[109491]19315.

[109492]Whitney, M. A.; Saito, H.; Jakobs, P. M.; Gibson, R. A.;

Moses,

[109493]R. E.; Grompe, M.: A common mutation in the FACC gene causes Fanconi

[109494] anaemia in Ashkenazi Jews. Nature Genet. 4: 202–205,
1993.

[109495] 19316.

[109496] Yamashita, T.; Barber, D. L.; Zhu, Y.; Wu, N.; D'Andrea, A.
D.

[109497]: The Fanconi anemia polypeptide FACC is localized to the
cytoplasm. Proc.

[109498] Nat. Acad. Sci. 91: 6712–6716, 1994.

[109499] 19317.

[109500] Yamashita, T.; Wu, N.; Kupfer, G.; Corless, C.; Joenje, H.;
Grompe,

[109501] M.; D'Andrea, A. D.: Clinical variability of Fanconi anemia
(type

[109502] C) results from expression of an amino terminal truncated
Fanconi

[109503] anemia complementation group C polypeptide with partial
activity. Blood 87:

[109504] 4424–4432, 1996.

[109505] 19318.

[109506] Youssoufian, H.: Cytoplasmic localization of FAC is essen-
tial

[109507]for the correction of a prerepair defect in Fanconi anemia
group C

[109508]cells. J. Clin. Invest. 97: 2003–2010, 1996.

[109509]19319.

[109510]Yousoufian, H.: Localization of Fanconi anemia C protein
to

[109511]the cytoplasm of mammalian cells. Proc. Nat. Acad. Sci.
91: 7975–7979,

[109512]1994.

[109513]19320.

[109514]Yousoufian, H.; Li, Y.; Martin, M. E.; Buchwald, M.: Induc-
tion

[109515]of Fanconi anemia cellular phenotype in human 293 cells
by overexpression

[109516]of a mutant FAC allele. J. Clin. Invest. 97: 957–962, 1996.

[109517]19321.

[109518]Hodge, S. E.; Anderson, C. E.; Neiswanger, K.; Field, L. L.;
Spence,

[109519]M. A.; Sparkes, R. S.; Sparkes, M. C.; Crist, M.; Terasaki, P.
I.;

[109520]Rimoin, D. L.; Rotter, J. I.: Close genetic linkage between

diabetes

[109521]mellitus and Kidd blood group. Lancet II: 893–895, 1981.

[109522]19322.

[109523]Marcolongo, P.; Banhegyi, G.; Benedetti, A.; Hinds, C. J.;
Burchell,

[109524]A.: Liver microsomal transport of glucose–6–phosphate,
glucose, and

[109525]phosphate in type 1 glycogen storage diseases. J. Clin.
Endocr. Metab. 83:

[109526]224–229, 1998.

[109527]19323.

[109528]Michels, V. V.; Beaudet, A. L.: Hemorrhagic pancreatitis in
a

[109529]patient with glycogen storage disease type I. Clin. Genet.
17: 220–222,

[109530]1980.

[109531]19324.

[109532]Nakai, K.; Sakamoto, H.: Construction of a novel database
containing

[109533]aberrant splicing mutations of mammalian genes. Gene
141: 171–177,

[109534]1994.

[109535]19325.

[109536]Napole, V. M.; Campbell, W. G., Jr.: Hepatoblastoma in infant

[109537]sister and brother. Cancer 39: 2647–2650, 1977.

[109538]19326.

[109539]Nuoffer, J. M.; Mullis, P. E.; Wiesmann, U. N.: Treatment with

[109540]low-dose diazoxide in two growth-retarded prepubertal girls with glycogen

[109541]storage disease type Ia resulted in catch-up growth. J. Inherit.

[109542]Metab. Dis. 20: 790–798, 1997.

[109543]19327.

[109544]Obara, K.; Saito, T.; Sato, H.; Ogawa, M.; Igarashi, Y.; Yoshinaga,

[109545]K.: Renal histology in two adult patients with type I glycogen storage

[109546]disease. Clin. Nephrol. 39: 59–64, 1993.

[109547]19328.

[109548]Parvari, R.; Lei, K.-J.; Bashan, N.; HersHKovitz, E.; Korman,
[109549]S. H.; Barash, V.; Lerman-Sagie, T.; Mandel, H.; Chou, J. Y.;
Moses,
[109550]S. W.: Glycogen storage disease type 1a in Israel: bio-
chemical, clinical,
[109551]and mutational studies. Am. J. Med. Genet. 72: 286-290,
1997.
[109552]19329.
[109553]Pizzo, C. J.: Type I glycogen storage disease with focal
nodular
[109554]hyperplasia of the liver and vasoconstrictive pulmonary
hypertension. Pediatrics 65:
[109555]341-343, 1980.
[109556]19330.
[109557]Qu, Y.; Abdenur, J. E.; Eng, C. M.; Desnick, R. J.: Molecular
[109558]prenatal diagnosis of glycogen storage disease type Ia.
Prenatal
[109559]Diag. 15: 333-336, 1996.
[109560]19331.
[109561]Reitsma-Bierens, W. C. C.: Renal complications in glyco-
gen storage

[109562]disease type I. *Europ. J. Pediat.* 152 (suppl. 1): S60–S62, 1993.

[109563]19332.

[109564]Rennert, O. M.; Mukhopadhyay, D.: Diazoxide in Von Gierke's disease. *Arch.*

[109565]*Dis. Child.* 43: 358–361, 1968.

[109566]19333.

[109567]Restaino, I.; Kaplan, B. S.; Stanley, C.; Baker, L.: Nephrolithiasis,

[109568]hypocitraturia, and a distal renal tubular acidification defect in

[109569]type 1 glycogen storage disease. *J. Pediat.* 122: 392–396, 1993.

[109570]19334.

[109571]Ryan, I. P.; Havel, R. J.; Laros, R. K., Jr.: Three consecutive

[109572]pregnancies in a patient with glycogen storage disease type IA (von

[109573]Gierke's disease). *Am. J. Obstet. Gynec.* 170: 1687–1691, 1994.

[109574]19335.

[109575]Selby, R.; Starzl, T. E.; Yunis, E.; Todo, S.; Tzakis, A. G.;

[109576]Brown, B. I.; Kendall, R. S.: Liver transplantation for type I
and

[109577]type IV glycogen storage disease. Europ. J. Pediat. 152
(suppl.

[109578]1): S71–S76, 1993.

[109579]19336.

[109580]Senior, B.; Loridan, L.: Liver glycogenoses: metabolism of
intravenously

[109581]administered glycerol. New Eng. J. Med. 279: 958–965,
1968.

[109582]19337.

[109583]Senior, B.; Loridan, L.: Functional differentiation of
glycogenoses

[109584]of the liver with respect to the use of glycerol. New Eng. J.
Med. 279:

[109585]965–970, 1968.

[109586]19338.

[109587]Seydewitz, H. H.; Matern, D.: Molecular genetic analysis of
40

[109588]patients with glycogen storage disease type Ia: 100% mu-
tation detection

[109589]rate and 5 novel mutations. (Abstract) Hum. Mutat. 15:
115–116,

[109590]2000.

[109591]19339.

[109592]Shelly, L. L.; Lei, K.-J.; Pan, C.-J.; Sakata, S. F.; Ruppert,

[109593]S.; Schutz, G.; Chou, J. Y.: Isolation of the gene for murine
glucose-6-phosphatase,

[109594]the enzyme deficient in glycogen storage disease type 1A.
J. Biol.

[109595]Chem. 268: 21482–21485, 1993.

[109596]19340.

[109597]Sidbury, J. B., Jr.: The genetics of the glycogen storage
disease.In:

[109598]Steinberg, A. G.; Bearn, A. G.: Progress in Medical Genet-
ics. New

[109599]York: Grune and Stratton (pub.) 4: 1965. Pp. 32–58.

[109600]19341.

[109601]Smit, G. P. A.: The long-term outcome of patients with
glycogen

[109602]storage disease type Ia. Europ. J. Pediat. 152 (suppl. 1):
S52–S55,

[109603]1993.

[109604]19342.

[109605]Spencer-Peet, J.; Norman, M. E.; Lake, B. D.; McNamara, J.;
Patrick,

[109606]A. D.: Hepatic glycogen storage disease. Clinical and laboratory
findings in 23 cases. Quart. J. Med. 40: 95-114, 1971.

[109607]

[109608]19343.

[109609]Stamm, W. E.; Webb, D. I.: Partial deficiency of hepatic
glucose-6-phosphatase

[109610]in an adult patient. Arch. Intern. Med. 135: 1107-1109,
1975.

[109611]19344.

[109612]Stevenson, R. E.; Ben-Menachem, Y.; Dudrick, S.; Howell,
R. R.

[109613]: Hepatocellular carcinoma in type 1 glycogen storage disease. Proc.

[109614]Greenwood Genet. Center 3: 39-46, 1984.

[109615]19345.

[109616]Talente, G. M.; Coleman, R. A.; Alter, C.; Baker, L.; Brown,
B.

[109617]I.; Cannon, R. A.; Chen, Y.-T.; Crigler, J. F., Jr.; Ferreira, P.;
 [109618]Haworth, J. C.; Herman, G. E.; Issenman, R. M.; Keating, J.
 P.; Linde,
 [109619]R.; Roe, T. F.; Senior, B.; Wolfsdorf, J. I.: Glycogen storage
 disease
 [109620]in adults. Ann. Intern. Med. 120: 218-226, 1994.
 [109621]19346.
 [109622]Trioche, P.; Francoual, J.; Capel, L.; Odievre, M.; Linden-
 baum,
 [109623]A.; Labrune, P.: Apolipoprotein E polymorphism and
 serum concentrations
 [109624]in patients with glycogen storage disease type Ia. J. In-
 herit. Metab.
 [109625]Dis. 107-112, 2000.
 [109626]19347.
 [109627]Ullrich, K.; Smit, G. P. A.: Clinical aspects of glycogen
 storage
 [109628]disease type I: summary of the discussions. Europ. J. Pe-
 diat. 152
 [109629](suppl. 1): S87-S88, 1993.
 [109630]19348.

[109631]Weinstein, D. A.; Somers, M. J. G.; Wolfsdorf, J. I.: De-
creased

[109632]urinary citrate excretion in type 1a glycogen storage dis-
ease. J.

[109633]Pediat. 138: 378–382, 2001.

[109634]19349.

[109635]Wierzbicki, A. S.; Watts, G. F.; Lynas, J.; Winder, A. F.;
Wray,

[109636]R.: Very low-density lipoprotein apolipoprotein B-100
turnover in

[109637]glycogen storage disease type Ia (von Gierke disease). J.
Inherit.

[109638]Metab. Dis. 24: 527–534, 2001.

[109639]19350.

[109640]Zangeneh, F.; Limbeck, G. A.; Brown, B. I.; Emch, J. R.; Ar-
casoy,

[109641]M. M.; Goldenberg, V. E.; Kelley, V. C.: Hepatorenal
glycogenosis

[109642](type I glycogenosis) and carcinoma of the liver. J. Pediat.
74:

[109643]73–83, 1969.

[109644]19351.

[109645]Maroteaux, P.; Lamy, M.: Hurler's disease, Morquio's disease,

[109646]and related mucopolysaccharidoses. J. Pediat. 67: 312-323, 1965.

[109647]19352.

[109648]Adams, P. C.: Intrafamilial variation in hereditary hemochromatosis. Digest.

[109649]Dis. Sci. 37: 361-363, 1992.

[109650]19353.

[109651]Aguilar-Martinez, P.; Bismuth, M.; Picot, M. C.; Thelcide, C.;

[109652]Pageaux, G.-P.; Blanc, F.; Blanc, P.; Schved, J.-F.; Larrey, D.:

[109653]Variable phenotypic presentation of iron overload in H63D homozygotes:

[109654]are genetic modifiers the cause? Gut 48: 836-842, 2001.

[109655]19354.

[109656]Anand, S.; Schade, R. R.; Bendetti, C.; Kelly, R.; Rabin, B. S.;

[109657]Krause, J.; Starzl, T. E.; Iwatsuki, S. I.; Van Thiel, D. H.: Id-

iopathic

[109658]alpha-hemochromatosis (sic) and 1-antitrypsin deficiency: coexistence

[109659]in a family with progressive liver disease in the proband.

Hepatology 3:

[109660]714-718, 1983.

[109661]19355.

[109662]Bahram, S.; Gilfillan, S.; Kuhn, L. C.; Moret, R.; Schulze, J.

[109663]B.; Lebeau, A.; Schumann, K.: Experimental hemochromatosis due to

[109664]MHC class I HFE deficiency: immune status and iron metabolism. Proc.

[109665]Nat. Acad. Sci. 96: 13312-13317, 1999.

[109666]19356.

[109667]Balcerzak, S. P.; Westerman, M. P.; Lee, R. E.; Doyle, A. P.:

[109668]Idiopathic hemochromatosis: a study of three families. Am. J. Med. 40:

[109669]857-873, 1966.

[109670]19357.

[109671]Barton, J. C.; Acton, R. T.: Inheritance of two HFE muta-

tions

[109672]in African Americans: cases with hemochromatosis phenotypes and estimates

[109673]of hemochromatosis phenotype frequency. Genet. Med. 3: 294–300,

[109674]2001.

[109675]19358.

[109676]Barton, J. C.; McDonnell, S. M.; Adams, P. C.; Brissot, P.; Powell,

[109677]L. W.; Edwards, C. Q.; Cook, J. D.; Kowdley, K. V.; Hemochromatosis

[109678]Management Working Group: Management of hemochromatosis. Ann. Intern.

[109679]Med. 129: 932–939, 1998.

[109680]19359.

[109681]Barton, J. C.; Patton, M. A.; Edwards, C. Q.; Griffen, L. M.; Kushner,

[109682]J. P.; Meeks, R. G.; Leggett, R. W.: Blood lead concentrations in

[109683]hereditary hemochromatosis. J. Lab. Clin. Med. 124: 193–198, 1994.

[109684]19360.

[109685]Barton, J. C.; Rothenberg, B. E.; Bertoli, L. F.; Acton, R. T.

[109686]: Diagnosis of hemochromatosis in family members of probands: a comparison

[109687]of phenotyping and HFE genotyping. Genet. Med. 1: 89–93, 1999.

[109688]19361.

[109689]Barton, J. C.; Sawada–Hirai, R.; Rothenberg, B. E.; Acton, R.

[109690]T.: Two novel missense mutations of the HFE gene (I105T and G93R)

[109691]and identification of the S65C mutation in Alabama hemochromatosis

[109692]probands. Blood Cells Molec. Dis. 25: 146–154, 1999.

[109693]19362.

[109694]Bassett, M. L.; Doran, T. J.; Halliday, J. W.; Bashir, H. V.;

[109695]Powell, L. W.: Idiopathic hemochromatosis: demonstration of homozygous–heterozygous

[109696]mating by HLA typing of families. Hum. Genet. 60: 352–356, 1982.

[109697]19363.

[109698]Beaumont, C.; Simon, M.; Fauchet, R.; Hespel, J.–P.; Bris–sot,

[109699]P.; Genetet, B.; Bourel, M.: Serum ferritin as a possible marker

[109700]of the hemochromatosis allele. New Eng. J. Med. 301: 169–174, 1979.

[109701]19364.

[109702]Beaumont, C.; Simon, M.; Smith, P. M.; Worwood, M.: Hepatic and

[109703]serum ferritin concentrations in patients with idiopathic hemochromatosis. Gastroenterology 79:

[109704]877–883, 1980.

[109705]19365.

[109706]Beckman, L. E.; Saha, N.; Spitsyn, V.; Van Landeghem, G.; Beckman,

[109707]L.: Ethnic differences in the HFE codon 282 (Cys/Tyr) polymorphism. Hum.

[109708]Hered. 47: 263–267, 1997.

[109709]19366.

[109710]Beutler, E.: Targeted disruption of the HFE gene. Proc. Nat.

[109711]Acad. Sci. 95: 2033–2034, 1998.

[109712]19367.

[109713]Beutler, E.: The significance of the 187G (H63D) mutation
in

[109714]hemochromatosis. (Letter) Am. J. Hum. Genet. 61:
762–764, 1997.

[109715]19368.

[109716]Beutler, E.; Felitti, V. J.; Koziol, J. A.; Ho, N. J.; Gelbart,

[109717]T.: Penetrance of 845G–A (C282Y) HFE hereditary
haemochromatosis

[109718]mutation in the USA. Lancet 359: 211–218, 2002.

[109719]19369.

[109720]Beutler, E.; Gelbart, T.: HLA–H mutations in the Ashkenazi
Jewish

[109721]population. Blood Cells Molec. Dis. 23: 95–98, 1997.

[109722]19370.

[109723]Beutler, E.; Gelbart, T.; West, C.; Lee, P.; Adams, M.; Black–
stone,

[109724]R.; Pockros, P.; Kosty, M.; Venditti, C. P.; Phatak, P. D.;
Seese,

[109725]N. K.; Chorney, K. A.; Ten Elshof, A. E.; Gerhard, G. S.;
Chorney,

[109726]M.: Mutation analysis in hereditary hemochromatosis.

Blood Cells

[109727]Molec. Dis. 22: 187-194, 1996.

[109728]19371.

[109729]Beutler, E.; West, C.; Gelbart, T.: HLA-H and associated proteins

[109730]in patients with hemochromatosis. Molec. Med. 3: 397-402, 1997.

[109731]19372.

[109732]Bodmer, J. G.; Parham, P.; Albert, E. D.; Marsh, S. G. E.: Putting

[109733]a hold on 'HLA-H.' (Letter) Nature Genet. 15: 234-235, 1997.

[109734]19373.

[109735]Borecki, I. B.; Lathrop, G. M.; Bonney, G. E.; Yaouanq, J.; Rao,

[109736]D. C.: Combined segregation and linkage analysis of genetic hemochromatosis

[109737]using affection status, serum iron, and HLA. Am. J. Hum. Genet. 47:

[109738]542-550, 1990.

[109739]19374.

[109740]Borecki, I. B.; Rao, D. C.; Yaouanq, J.; Lalouel, J. M.: Per-
cent

[109741]transferrin saturation in segregating hemochromatosis.
Am. J. Med.

[109742]Genet. 36: 301-305, 1990.

[109743]19375.

[109744]Borecki, I. B.; Rao, D. C.; Yaouanq, J.; Lalouel, J. M.: Segre-
gation

[109745]of genetic hemochromatosis indexed by latent capacity of
transferrin. Am.

[109746]J. Hum. Genet. 45: 465-470, 1989.

[109747]19376.

[109748]Boretto, J.; Jouanolle, A.-M.; Yaouanq, J.; El Kahloun, A.;
Mauvieux,

[109749]V.; Blayau, M.; Perichon, M.; Le Treut, A.; Clayton, J.; Borot,
N.;

[109750]Le Gall, J.-Y.; Pontarotti, P.; David, V.: Anonymous mark-
ers located

[109751]on chromosome 6 in the HLA-A class I region: allelic dis-
tribution

[109752]in genetic haemochromatosis. Hum. Genet. 89: 33-36,
1992.

[109753]19377.

[109754]Maroteaux, P.; Leveque, B.; Marie, J.; Lamy, M.: Une nouvelle

[109755]dysostose avec elimination urinaire de chondroïtine-sulfate B. Presse

[109756]Med. 71: 1849-1852, 1963.

[109757]19378.

[109758]McGovern, M. M.; Mandell, N.; Haskins, M.; Desnick, R. J.: Animal

[109759]model studies of allelism: characterization of arylsulfatase B mutations

[109760]in homoallelic and heteroallelic (genetic compound) homozygotes with

[109761]feline mucopolysaccharidosis VI. Genetics 110: 733-749, 1985.

[109762]19379.

[109763]Migeon, B. R.; Sprenkle, J. A.; Liebaers, I.; Scott, J. F.; Neufeld,

[109764]E. F.: X-linked Hunter syndrome: the heterozygous phenotype in cell

[109765]culture. Am. J. Hum. Genet. 29: 448-454, 1977.

[109766]19380.

[109767]Neufeld, E. F.; Muenzer, J.: The mucopolysaccharidoses. In:
Scriver,

[109768]C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds.): The
Metabolic

[109769]and Molecular Bases of Inherited Disease New York: Mc-
Graw-Hill

[109770](7th ed.): 1995. Pp. 2465-2494.

[109771]19381.

[109772]Paterson, D. E.; Rad, M.; Harper, G.; Weston, H. J.; Mattin-
gley,

[109773]J.: Maroteaux-Lamy syndrome, mild form--MPS VIb. Brit.
J. Radiol. 55:

[109774]805-812, 1982.

[109775]19382.

[109776]Peterson, D. I.; Bacchus, H.; Seaich, L.; Kelly, T. E.:
Myelopathy

[109777]associated with Maroteaux-Lamy syndrome. Arch. Neurol.
32: 127-129,

[109778]1975.

[109779]19383.

- [109780]Poser, C. M.; Ojemann, R. G.; Sobel, R. A.: MGH CPC case
44–1983. New
- [109781]Eng. J. Med. 309: 1109–1117, 1983.
- [109782]19384.
- [109783]Quigley, H. A.; Kenyon, K. R.: Ultrastructural and histo-
chemical
- [109784]studies of a newly recognized form of systemic mu-
copolysaccharidosis
- [109785](Maroteaux–Lamy syndrome, mild phenotype). Am. J.
Ophthal. 77: 809–818,
- [109786]1974.
- [109787]19385.
- [109788]Saul, R. A.; Stevenson, R. E.; Taylor, H. A.: Atypical presen-
tation
- [109789]with normal stature in Maroteaux–Lamy syndrome (MPS
VI). Proc. Greenwood
- [109790]Genet. Center 3: 49–52, 1984.
- [109791]19386.
- [109792]Schuchman, E. H.; Jackson, C. E.; Desnick, R. J.: Human
arylsulfatase
- [109793]B: MOPAC cloning, nucleotide sequence of a full-length

cDNA, and regions

[109794] of amino acid identity with arylsulfatases A and C. Genomics 6:

[109795] 149–158, 1990.

[109796] 19387.

[109797] Schwartz, G. P.; Cohen, E. J.: Hydrocephalus in Maroteaux–Lamy

[109798] syndrome. Arch. Ophthalmol. 116: 400 only, 1998.

[109799] 19388.

[109800] Spranger, J. W.; Koch, F.; McKusick, V. A.; Natzschka, J.; Wiedemann,

[109801] H. R.; Zellweger, H.: Mucopolysaccharidosis VI (Maroteaux–Lamy's

[109802] disease). Helv. Paediat. Acta 25: 337–362, 1970.

[109803] 19389.

[109804] Stumpf, D. A.; Austin, J. H.; Crocker, A. C.; LaFrance, M.: Mucopolysaccharidosis

[109805] type VI (Maroteaux–Lamy syndrome). I. Sulfatase B deficiency in tissues. Am.

[109806] J. Dis. Child. 126: 747–755, 1973.

[109807] 19390.

- [109808]Tan, C. T. T.; Schaff, H. V.; Miller, F. A., Jr.; Edwards, W.
[109809]D.; Karnes, P. S.: Valvular heart disease in four patients
with Maroteaux-Lamy
[109810]syndrome. *Circulation* 85: 188-195, 1992.
[109811]19391.
- [109812]Tonnesen, T.; Gregersen, H. N.; Guttler, F.: Normal MPS
excretion,
[109813]but dermatan sulphaturia, combined with a mild
Maroteaux-Lamy phenotype. *J.*
[109814]*Med. Genet.* 28: 499-501, 1991.
[109815]19392.
- [109816]van Dorpe, J.; Moerman, P.; Pecceu, A.; van den Steen, P.;
Fryns,
[109817]J. P.: Non-immune hydrops fetalis caused by beta-
glucuronidase deficiency
[109818](mucopolysaccharidosis VII): study of a family with 3 af-
fected siblings. *Genet.*
[109819]*Counsel.* 7: 105-112, 1996.
[109820]19393.
- [109821]Van Dyke, D. L.; Fluharty, A. L.; Schafer, I. A.; Shapiro, L.

[109822]J.; Kihara, H.; Weiss, L.: Prenatal diagnosis of Maroteaux-Lamy syndrome. Am.

[109823]J. Med. Genet. 8: 235-242, 1981.

[109824]19394.

[109825]Vestermarck, S.; Tonnesen, T.; Schultz Andersen, M.; Guttler, F.

[109826]: Mental retardation in a patient with Maroteaux-Lamy. Clin. Genet. 31:

[109827]114-117, 1987.

[109828]19395.

[109829]Villani, G. R. D.; Balzano, N.; Vitale, D.; Saviano, M.; Pavone,

[109830]V. Di Natale, P.: Maroteaux-Lamy syndrome: five novel mutations and

[109831]their structural localization. Biochim. Biophys. Acta 1453: 185-192,

[109832]1999.

[109833]19396.

[109834]Vine, D. T.; McGovern, M. M.; Haskins, M. E.; Desnick, R. J.:

[109835]Feline mucopolysaccharidosis. VI. Purification and charac-

terization

[109836] of the residual arylsulfatase B activity. Am. J. Hum. Genet.

33:

[109837] 916–927, 1981.

[109838] 19397.

[109839] Weller, P. F.; Austen, K. F.: Human eosinophil arylsulfatase

[109840] B: structure and activity of the purified tetrameric lysosomal hydrolase. J.

[109841] Clin. Invest. 71: 114–123, 1983.

[109842] 19398.

[109843] Wicker, G.; Prill, V.; Brooks, D.; Gibson, G.; Hopwood, J.;
von

[109844] Figura, K.; Peters, C.: Mucopolysaccharidosis VI
(Maroteaux–Lamy

[109845] syndrome): an intermediate clinical phenotype caused by
substitution

[109846] of valine for glycine at position 137 of arylsulfatase B. J.
Biol.

[109847] Chem. 266: 21386–21391, 1991.

[109848] 19399.

[109849] Wilson, C. S.; Mankin, H. T.; Pluth, J. R.: Aortic stenosis

and

[109850] mucopolysaccharidosis. Ann. Intern. Med. 92: 496–498,
1980.

[109851] 19400.

[109852] Yogalingam, G.; Hopwood, J. J.; Crawley, A.; Anson, D. S.:
Mild

[109853] feline mucopolysaccharidosis type VI: identification of an
N-acetylgalactosamine-4-sulfatase

[109854] mutation causing instability and increased specific activ-
ity. J.

[109855] Biol. Chem. 273: 13421–13429, 1998.

[109856] 19401.

[109857] Yogalingam, G.; Litjens, T.; Bielicki, J.; Crawley, A. C.;
Muller,

[109858] V.; Anson, D. S.; Hopwood, J. J.: Feline mucopolysacchari-
dosis type

[109859] VI. J. Biol. Chem. 271: 27259–27265, 1996.

[109860] 19402.

[109861] Yoshida, M.; Noguchi, J.; Ikadaï, H.; Takahashi, M.; Na-
gase, S.

[109862] Arylsulfatase B-deficient mucopolysaccharidosis in rats.

J. Clin.

[109863]Invest. 91: 1099–1104, 1993.

[109864]19403.

[109865]Yoshida, M.; Tachibana, M.; Kobayashi, E.; Ikadai, H.; Kuni-
nieda,

[109866]T.: The locus responsible for mucopolysaccharidosis VI
(Maroteaux–Lamy

[109867]syndrome) is located on rat chromosome 2. Genomics 20:
145–146,

[109868]1994.

[109869]19404.

[109870]Young, R.; Kleinman, G.; Ojemann, R. G.; Kolodny, E.;
Davis, K.;

[109871]Halperin, J.; Zalneraitis, E.; DeLong, G. R.: Compressive
myelopathy

[109872]in Maroteaux–Lamy syndrome: clinical and pathological
findings. Ann.

[109873]Neurol. 8: 336–340, 1980.

[109874]19405.

[109875]Bradley, W. G.; Price, D. L.; Watanabe, C. K.: Familial cen-
tronuclear

[109876]myopathy. J. Neurol. Neurosurg. Psychiat. 33: 687–693,
1970.

[109877]19406.

[109878]Touchman, J. W.; Anikster, Y.; Dietrich, N. L.; Braden
Maduro,

[109879]V. V.; McDowell, G.; Shotelersuk, V.; Bouffard, G. G.; Beck-
strom–Sternberg,

[109880]S. M.; Gahl, W. A.; Green, E. D.: The genomic region en-
compassing

[109881]the nephropathic cystinosis gene (CTNS): complete se-
quencing of a

[109882]200-kb segment and discovery of a novel gene within the
common cystinosis–causing

[109883]deletion. Genome Res. 10: 165–173, 2000.

[109884]19407.

[109885]Shapiro, M. B.; Senapathy, P.: RNA splice junctions of dif-
ferent

[109886]classes of eukaryotes: sequence statistics and functional
implications

[109887]in gene expression. Nucleic Acids Res. 15: 7155–7174,
1987.

[109888]19408.

[109889]Applegarth, D. A.; Toone, J. R.: Nonketotic hyperglycemia (glycine
encephalopathy): laboratory diagnosis. *Molec. Genet. Metab.* 74:
139–146, 2001.
19409.

[109893]Kure, S.; Kojima, K.; Kudo, T.; Kanno, K.; Aoki, Y.; Suzuki, Y.;

[109894]Shinka, T.; Sakata, Y.; Narisawa, K.; Matsubara, Y.: Chromosomal
localization, structure, single-nucleotide polymorphisms, and expression
of the human H-protein gene of the glycine cleavage system (GCSH),
a candidate gene for nonketotic hyperglycinemia. *J. Hum. Genet.* 46:
378–384, 2001.
19410.

[109900]Sakata, Y.; Owada, Y.; Sato, K.; Kojima, K.; Hisanaga, K.; Shinka,

[109901]T.; Suzuki, Y.; Aoki, Y.; Satoh, J.; Kondo, H.; Matsubara, Y.; Kure,

[109902]S.: Structure and expression of the glycine cleavage system in rat

[109903]central nervous system. Molec. Brain Res. 94: 119–130, 2001.

[109904]19411.

[109905]Toone, J. R.; Applegarth, D. A.; Coulter–Mackie, M. B.; James,

[109906]E. R.: Recurrent mutations in P– and T–proteins of the glycine cleavage

[109907]complex and a novel T–protein mutation (N145I): a strategy for the

[109908]molecular investigation of patients with nonketotic hyperglycinemia

[109909](NKH). Molec. Genet. Metab. 72: 322–325, 2001.

[109910]19412.

[109911]Bamforth, J. S.; Hughes, I. A.; Lazarus, J. H.; Weaver, C. M.;

[109912]Harper, P. S.: Congenital hypothyroidism, spiky hair, and cleft palate. J.

[109913]Med. Genet. 26: 49–60, 1989.

[109914]19413.

[109915]Jobard, F.; Lefevre, C.; Karaduman, A.; Blanchet–Bardon,

C.; Emre,

[109916]S.; Weissenbach, J.; Ozguc, M.; Lathrop, M.; Prud'homme,
J.-F.; Fischer,

[109917]J.: Lipoxxygenase-3 (ALOXE3) and 12(R)-lipoxxygenase
(ALOX12B) are

[109918]mutated in non-bullous congenital ichthyosiform erythro-
derma (NCIE)

[109919]linked to chromosome 17p13.1. Hum. Molec. Genet. 11:
107-113, 2002.

[109920]19414.

[109921]Dinarello, C. A.; Wolff, S. M.; Goldfinger, S. E.; Dale, D. C.;

[109922]Alling, D. W.: Colchicine therapy for familial Mediterranean
fever:

[109923]a double-blind trial. New Eng. J. Med. 291: 934-937,
1974.

[109924]19415.

[109925]Akaboshi, S.; Yano, T.; Miyawaki, S.; Ohno, K.; Takeshita,
K.:

[109926]A C57BL/KsJ mouse model of Niemann-Pick disease (spm)
belongs to the

[109927]same complementation group as the major childhood type
of Niemann-Pick

[109928]disease type C. Hum. Genet. 99: 350–353, 1997.

[109929]19416.

[109930]Levy, E. N.; Shen, Y.; Kupelian, A.; Kruglyak, L.; Aksentijevich,

[109931]I.; Pras, E.; Balow, J. E., Jr.; Linzer, B.; Chen, X.; Shelton, D.

[109932]A.; Gumucio, D.; Pras, M.; Shohat, M.; Rotter, J. I.; Fischel-Ghodsian,

[109933]N.; Richards, R. I.; Kastner, D. L.: Linkage disequilibrium mapping

[109934]places the gene causing familial Mediterranean fever close to D16S246. Am.

[109935]J. Hum. Genet. 58: 523–534, 1996.

[109936]19417.

[109937]Majeed, H. A.; Barakat, M.: Familial Mediterranean fever (recurrent

[109938]hereditary polyserositis) in children: analysis of 88 cases. Europ.

[109939]J. Pediat. 148: 636–641, 1989.

[109940]19418.

[109941]Varon, R.; Vissinga, C.; Platzer, M.; Cerosaletti, K. M.; Chrzanowska,

[109942]K. H.; Saar, K.; Beckmann, G.; Seemanova, E.; Cooper, P.
R.; Nowak,

[109943]N. J.; Stumm, M.; Weemaes, C. M. R.; Gatti, R. A.; Wilson,
R. K.;

[109944]Digweed, M.; Rosenthal, A.; Sperling, K.; Concannon, P.;
Reis, A.

[109945]; Nibrin, a novel DNA double-strand break repair protein,
is mutated

[109946]in Nijmegen breakage syndrome. Cell 93: 467-476, 1998.

[109947]19419.

[109948]Becker, J. A.; Vlach, J.; Raben, N.; Nagaraju, K.; Adams, E.
M.;

[109949]Hermans, M. M.; Reuser, A. J. J.; Brooks, S. S.; Tifft, C. J.;
Hirschhorn,

[109950]R.; Huie, M. L.; Nicolino, M.; Plotz, P. H.: The African origin
of

[109951]the common mutation in African American patients with
glycogen-storage

[109952]disease type II. (Letter) Am. J. Hum. Genet. 62: 991-994,
1998.

[109953]19420.

[109954]Boerkoel, C. F.; Exelbert, R.; Nicastrì, C.; Nichols, R. C.;

Miller,

[109955]F. W.; Plotz, P. H.; Raben, N.: Leaky splicing mutation in the acid

[109956]maltase gene is associated with delayed onset of glycogenosis type

[109957]II. Am. J. Hum. Genet. 56: 887–897, 1995.

[109958]19421.

[109959]Byrne, E.; Dennett, X.; Crotty, B.; Trounce, I.; Sands, J. M.;

[109960]Hawkins, R.; Hammond, J.; Anderson, S.; Haan, E. A.; Pol-lard, A.:

[109961]Dominantly inherited cardioskeletal myopathy with lyso-somal glycogen

[109962]storage and normal acid maltase levels. Brain 109: 523–536, 1986.

[109963]19422.

[109964]Friduss, S. R.; Sadoff, W. I.; Hern, A. E.; Fivenson, D. P.:

[109965]Fatal pyoderma gangrenosum in association with C7 defi-ciency. J.

[109966]Am. Acad. Derm. 27: 356–359, 1992.

[109967]19423.

[109968]Lee, T. J.; Utsinger, P. D.; Snyderman, R.; Yount, W. J.;

Sparling,

[109969]P. F.: Familial deficiency of the seventh component of complement

[109970]associated with recurrent bacteremic infections due to *Neisseria*. J.

[109971]*Infect. Dis.* 138: 359–368, 1978.

[109972]19424.

[109973]Nakamura, S.; Ooue, O.; Abe, K.: Genetic polymorphism of the

[109974]seventh component of complement in a Japanese population. *Hum. Genet.* 66:

[109975]279–281, 1984.

[109976]19425.

[109977]Nishimukai, H.; Tamaki, Y.: Genetic polymorphism of the seventh

[109978]component of complement: a new variant. *Vox Sang.* 51: 60–62, 1986.

[109979]19426.

[109980]Nishizaka, H.; Horiuchi, T.; Zhu, Z. B.; Fukumori, Y.; Volanakis,

[109981]J. E.: Genetic bases of human complement C7 deficiency. J.

Immun. 157:

[109982]4239–4243, 1996.

[109983]19427.

[109984]Nurnberger, W.; Pietsch, H.; Seger, R.; Bufon, T.; Wahn, V.:

[109985]Familial deficiency of the seventh component of complement associated

[109986]with recurrent meningococcal infections. Europ. J. Pediat. 148:

[109987]758–760, 1989.

[109988]19428.

[109989]Petersen, B. H.; Lee, T. J.; Snyderman, R.; Brooks, G. F.:
Neisseria

[109990]meningitidis and Neisseria gonorrhoeae bacteremia associated with

[109991]C6, C7, or C8 deficiency. Ann. Intern. Med. 90: 917–920, 1979.

[109992]19429.

[109993]Rittner, C.; Opferkuch, W.; Wellek, B.; Grosse-Wilde, H.;
Wernet,

[109994]P.: Lack of linkage between gene(s) controlling the synthesis of

[109995]the seventh component of complement and the HLA re-
gion on chromosome

[109996]no. 6 in man. Hum. Genet. 34: 137-142, 1976.

[109997]19430.

[109998]Schlesinger, M.; Nave, Z.; Levy, Y.; Slater, P. E.; Fishelson,

[109999]Z.: Prevalence of hereditary properdin, C7 and C8 defi-
ciencies in

[110000]patients with meningococcal infections. Clin. Exp. Immun.
81: 423-427,

[110001]1990.

[110002]19431.

[110003]Washio, K.; Tokunaga, K.; Omoto, K.; Misawa, S.: Human
C7 polymorphism:

[110004]classification and association analysis with C6. Jpn. J.
Hum. Genet. 31:

[110005]345-352, 1986.

[110006]19432.

[110007]Wurzner, R.; Hobart, M. J.; Orren, A.; Tokunaga, K.; Nitze,
R.;

[110008]Gotze, O.; Lachmann, P. J.: A novel protein polymorphism
of human

[110009]complement C7 detected by a monoclonal antibody. Immunogenetics 35:

[110010]398–402, 1992.

[110011]19433.

[110012]Wurzner, R.; Nitze, R.; Gotze, O.: C7*9, a new frequent C7 allele

[110013]detected by an allotype-specific monoclonal antibody. Complement

[110014]Inflamm. 7: 290–297, 1990.

[110015]19434.

[110016]Zimran, A.; Rudensky, B.; Kramer, M. R.; Tedesco, F.; Ehrenfeld,

[110017]M.; Raz, R.; Greif, Z.; Gelber, M.; Lishner, M.; Golan, E.; Hershko,

[110018]C.: Hereditary complement deficiency in survivors of meningococcal

[110019]disease: high prevalence of C7/C8 deficiency in Sephardic (Moroccan)

[110020]Jews. Quart. J. Med. 63: 349–358, 1987.

[110021]19435.

[110022]Hadano, S.; Hand, C. K.; Osuga, H.; Yanagisawa, Y.;

Otomo, A.;

[110023]Devon, R. S.; Miyamoto, N.; Showguchi-Miyata, J.; Okada, Y.; Singaraja,

[110024]R.; Figlewicz, D. A.; Kwiatkowski, T.; and 9 others: A gene encoding

[110025]a putative GTPase regulator is mutated in familial amyotrophic lateral

[110026]sclerosis 2. Nature Genet. 29: 166-173, 2001. Note: Erratum: Nature

[110027]Genet. 29: 352 only, 2001.

[110028]19436.

[110029]Colten, H. R.; Alper, C. A.; Rosen, F. S.: Current concepts in

[110030]immunology: genetics and biosynthesis of complement proteins. New

[110031]Eng. J. Med. 304: 653-656, 1981.

[110032]19437.

[110033]Howlett, N. G.; Taniguchi, T.; Olson, S.; Cox, B.; Waisfisz, Q.;

[110034]de Die-Smulders, C.; Persky, N.; Grompe, M.; Joenje, H.; Pals, G.;

[110035]Ikeda, H.; Fox, E. A.; D'Andrea, A. D.: Biallelic inactivation

of

[110036]BRCA2 in Fanconi anemia. Science 297: 606–609, 2002.

[110037]19438.

[110038]Mazoyer, S.; Dunning, A. M.; Serova, O.; Dearden, J.;
Puget, N.;

[110039]Healey, C. S.; Gayther, S. A.; Mangion, J.; Stratton, M. R.;
Lynch,

[110040]H. T.; Goldgar, D. E.; Ponder, B. A.; Lenoir, G. M.: A poly-
morphic

[110041]stop codon in BRCA2. Nature Genet. 14: 253–254, 1996.

[110042]19439.

[110043]Lowry, R. B.; Applegarth, D. A.; Toone, J. R.; MacDonald,
E.;

[110044]Thunem, N. Y.: An update on the frequency of mu-
copolysaccharide syndromes

[110045]in British Columbia. Hum. Genet. 85: 389–390, 1990.

[110046]19440.

[110047]ten Hove, T.; Corbaz, A.; Amitai, H.; Aloni, S.; Belzer, I.;
Graber,

[110048]P.; Drillenburger, P.; van Deventer, S. J.; Chvatchko, Y.; Te
Velde,

[110049]A. A.: Blockade of endogenous IL-18 ameliorates TNBS-induced colitis
[110050]by decreasing local TNF-alpha production in mice. Gastroenterology 121:
[110051]1372-1379, 2001.
[110052]19441.
[110053]Roux, A.-F.; Rommens, J.; McDowell, C.; Anson-
Cartwright, L.; Bell,
[110054]S.; Schappert, K.; Fishman, G. A.; Musarella, M.: Identification
[110055]of a gene from Xp21 with similarity to the tctex-1 gene of
the murine
[110056]t complex. Hum. Molec. Genet. 3: 257-263, 1994.
[110057]19442.
[110058]Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.;
Tanaka, A.;
[110059]Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding
sequences
[110060]of unidentified human genes. X. The complete sequences
of 100 new
[110061]cDNA clones from brain which can code for large proteins
in vitro. DNA

[110062]Res. 5: 169–176, 1998.

[110063]19443.

[110064]Kipreos, E. T.; Lander, L. E.; Wing, J. P.; He, W. W.; Hedge-
cock,

[110065]E. M.: *cul-1* is required for cell cycle exit in *C. elegans* and
identifies

[110066]a novel gene family. *Cell* 85: 829–839, 1996.

[110067]19444.

[110068]Rasooly, R. S.: Personal Communication. Baltimore, Md.
9/29/1998.

[110069]19445.

[110070]Wang, P. J.; McCarrey, J. R.; Yang, F.; Page, D. C.: An abundance

[110071]of X-linked genes expressed in spermatogonia. Nature
Genet. 27:

[110072]422-426, 2001.

[110073]19446.

[110074]Herold, A.; Suyama, M.; Rodrigues, J. P.; Braun, I. C.; Kuntay,

[110075]U.; Carmo-Fonseca, M.; Bork, P.; Izaurralde, E.: TAP (NXF1)
belongs

[110076]to a multigene family of putative RNA export factors with
a conserved

[110077]modular architecture. Molec. Cell. Biol. 20: 8996-9008,
2000.

[110078]19447.

[110079]Yang, J.; Bogerd, H. P.; Wang, P. J.; Page, D. C.; Cullen, B.
R.

[110080]: Two closely related human nuclear export factors utilize
entirely

[110081]distinct export pathways. Molec. Cell 8: 397–406, 2001.

[110082]19448.

[110083]Ikeda, M.; Ishida, O.; Hinoi, T.; Kishida, S.; Kikuchi, A.:

Identification

[110084]and characterization of a novel protein interacting with

Ral-binding

[110085]protein 1, a putative effector protein of Ral. J. Biol. Chem.

273:

[110086]814–821, 1998.

[110087]19449.

[110088]Koshiba, S.; Kigawa, T.; Iwahara, J.; Kikuchi, A.; Yokoyama,

S.

[110089]: Solution structure of the Eps15 homology domain of a

human POB1

[110090](partner of RalBP1). FEBS Lett. 442: 138–142, 1999.

[110091]19450.

[110092]Crackower, M. A.; Sarao, R.; Oudit, G. Y.; Yagil, C.; Kozier-

adzki,

[110093]J.; Scanga, S. E.; Oliveira-dos-Santos, A. J.; da Costa, J.;

Zhang,

[110094]L.; Pei, Y.; Scholey, J.; Ferrario, C. M.; Manoukian, A. S.;

Chappell,

[110095]M. C.; Backx, P. H.; Yagil, Y.; Penninger, J. M.: An-
giotensin-converting

[110096]enzyme 2 is an essential regulator of heart function. Na-
ture 417:

[110097]822-828, 2002.

[110098]19451.

[110099]Scanlan, M. J.; Gordan, J. D.; Williamson, B.; Stockert, E.;
Bander,

[110100]N. H.; Jongeneel, V.; Gure, A. O.; Jager, D.; Jager, E.;
Knuth, A.;

[110101]Chen, Y.-T.; Old, L. J.: Antigens recognized by autologous
antibody

[110102]in patients with renal-cell carcinoma. Int. J. Cancer 83:
456-464,

[110103]1999.

[110104]19452.

[110105]Yan, Z.; Fedorov, S. A.; Mumby, M. C.; Williams, R. S.:
PR48,

[110106]a novel regulatory subunit of protein phosphatase 2A, in-
teracts with

[110107]Cdc6 and modulates DNA replication in human cells.

Molec. Cell. Biol. 20:

[110108]1021–1029, 2000.

[110109]19453.

[110110]Gitzelmann, R.: Hereditary galactokinase deficiency, a
newly recognized

[110111]cause of juvenile cataracts. *Pediat. Res.* 1: 14–23, 1967.

[110112]19454.

[110113]Kalaydjieva, L.; Perez–Lezaun, A.; Angelicheva, D.;
Onengut, S.;

[110114]Dye, D.; Bosshard, N. U.; Jordanova, A.; Savov, A.;
Yanakiev, P.;

[110115]Kremensky, I.; Radeva, B.; Hallmayer, J.; Markov, A.; Ned-
kova, V.;

[110116]Tournèv, I.; Aneva, L.; Gitzelmann, R.: A founder mutation
in the

[110117]GK1 gene is responsible for galactokinase deficiency in
Roma (Gypsies). *Am.*

[110118]*J. Hum. Genet.* 65: 1299–1307, 1999.

[110119]19455.

[110120]Kim, Y. J.; Rosenberg, L. E.: On the mechanism of pyridox-
ine

- [110121] responsive homocystinuria. II. Properties of normal and mutant cystathionine
- [110122] beta-synthase from cultured fibroblasts. Proc. Nat. Acad. Sci. 71:
- [110123] 4821–4825, 1974.
- [110124] 19456.
- [110125] Kluijtmans, L. A. J.; Boers, G. H. J.; Kraus, J. P.; van den Heuvel,
- [110126] L. P. W. J.; Cruysberg, J. R. M.; Trijbels, F. J. M.; Blom, H. J.
- [110127]: The molecular basis of cystathionine beta-synthase deficiency in
- [110128] Dutch patients with homocystinuria: effect of CBS genotype on biochemical
- [110129] and clinical phenotype and on response to treatment. Am. J. Hum.
- [110130] Genet. 65: 59–67, 1999.
- [110131] 19457.
- [110132] Kluijtmans, L. A. J.; Boers, G. H. J.; Stevens, E. M. B.; Re-
- [110133] nier, W. O.; Kraus, J. P.; Trijbels, F. J. M.; van den Heuvel, L. P. W.

[110134]J.; Blom, H. J.: Defective cystathionine beta-synthase regulation
[110135]by S-adenosylmethionine in a partially pyridoxine responsive homocystinuria
[110136]patient. J. Clin. Invest. 98: 285-289, 1996.
[110137]19458.
[110138]Komrower, G. M.: Dietary treatment of homocystinuria.
Am. J.
[110139]Dis. Child. 113: 98-100, 1967.
[110140]19459.
[110141]Kozich, V.; de Franchis, R.; Kraus, J. P.: Molecular defect in
[110142]a patient with pyridoxine-responsive homocystinuria.
Hum. Molec.
[110143]Genet. 2: 815-816, 1993.
[110144]19460.
[110145]Kozich, V.; Kraus, E.; de Franchis, R.; Fowler, B.; Boers, G.
[110146]H. J.; Graham, I.; Kraus, J. P.: Hyperhomocysteinemia in
premature
[110147]arterial disease: examination of cystathionine beta-synthase alleles

[110148]at the molecular level. Hum. Molec. Genet. 4: 623–629,
1995.

[110149]19461.

[110150]Kozich, V.; Kraus, J. P.: Screening for mutations by ex-
pressing

[110151]patient cDNA segments in E. coli: homocystinuria due to
cystathionine

[110152]beta-synthase deficiency. Hum. Mutat. 1: 113–123, 1992.

[110153]19462.

[110154]Kraus, J.; Packman, S.; Fowler, B.; Rosenberg, L. E.: Purifi-
cation

[110155]and properties of cystathionine beta-synthase from hu-
man liver: evidence

[110156]for identical subunits. J. Biol. Chem. 253: 6523–6528,
1978.

[110157]19463.

[110158]Kluijtmans, L. A. J.; van den Heuvel, L. P. W. J.; Boers, G. H.

[110159]J.; Frosst, P.; Stevens, E. M. B.; van Oost, B. A.; den Heijer,
M.;

[110160]Trijbels, F. J. M.; Rozen, R.; Blom, H. J.: Molecular genetic
analysis

[110161]in mild hyperhomocysteinemia: a common mutation in the methylenetetrahydrofolate

[110162]reductase gene is a genetic risk factor for cardiovascular disease. Am.

[110163]J. Hum. Genet. 58: 35–41, 1996.

[110164]19464.

[110165]Kraus, J. P.: Molecular basis of phenotype expression in homocystinuria. J.

[110166]Inherit. Metab. Dis. 17: 383–390, 1994.

[110167]19465.

[110168]Kraus, J. P.; Janosik, M.; Kozich, V.; Mandell, R.; Shih, V.;

[110169]Sperandeo, M. P.; Sebastio, G.; de Franchis, R.; Andria, G.; Kluijtmans,

[110170]L. A. J.; Blom, H.; Boers, G. H. J.; Gordon, R. B.; Kamoun, P.; Tsai,

[110171]M. Y.; Kruger, W. D.; Koch, H. G.; Ohura, T.; Gaustadnes, M.: Cystathionine

[110172]beta-synthase mutations in homocystinuria. Hum. Mutat. 13: 362–375,

[110173]1999.

[110174]19466.

[110175]Kraus, J. P.; Oliveriusova, J.; Sokolova, J.; Kraus, E.; Vlcek,

[110176]C.; de Franchis, R.; Maclean, K. N.; Bao, L.; Bukovska, G.;
Patterson,

[110177]D.; Paces, V.; Ansorge, W.; Kozich, V.: The human cystathionine beta-synthase

[110178](CBS) gene: complete sequence, alternative splicing, and
polymorphisms. Genomics 52:

[110179]312–324, 1998.

[110180]19467.

[110181]Kruger, W. D.; Cox, D. R.: A yeast assay for functional detection

[110182]of mutations in the human cystathionine beta-synthase
gene. Hum.

[110183]Molec. Genet. 4: 1155–1161, 1995.

[110184]19468.

[110185]Kruger, W. D.; Cox, D. R.: A yeast system for expression of
human

[110186]cystathionine beta-synthase: structural and functional
conservation

[110187]of the human and yeast genes. Proc. Nat. Acad. Sci. 91:
6614–6618,

[110188]1994.

[110189]19469.

[110190]Kurczynski, T. W.; Muir, W. A.; Fleisher, L. D.; Palomaki, J.

[110191]F.; Gaull, G. E.; Rassin, D. K.; Abramowsky, C.: Maternal
homocystinuria:

[110192]studies of an untreated mother and fetus. Arch. Dis. Child.
55:

[110193]721–723, 1980.

[110194]19470.

[110195]Loughridge, L. W.: Renal abnormalities in the Marfan syn-
drome. Quart.

[110196]J. Med. 112: 531–544, 1959.

[110197]19471.

[110198]Lubec, B.; Fang–Kircher, S.; Lubec, T.; Blom, H. J.; Boers,
G.

[110199]H. J.: Evidence for McKusick's hypothesis of deficient col-
lagen cross–linking

[110200]in patients with homocystinuria. Biochim. Biophys. Acta
1315: 159–162,

[110201]1996.

[110202]19472.

[110203]Maclean, K. N.; Gaustadnes, M.; Oliveriusova, J.; Janosik, M.;

[110204]Kraus, E.; Kozich, V.; Kery, V.; Skovby, F.; Rudiger, N.; Ingerslev,

[110205]J.; Stabler, S. P.; Allen, R. H.; Kraus, J. P.: High homocysteine

[110206]and thrombosis without connective tissue disorders are associated

[110207]with a novel class of cystathionine beta-synthase (CBS) mutations. Hum.

[110208]Mutat. 19: 641-655, 2002.

[110209]19473.

[110210]Malinow, M. R.; Stampfer, M. J.: Role of plasma homocyst(e)ine

[110211]in arterial occlusive disease. (Editorial) Clin. Chem. 40: 857-858,

[110212]1994.

[110213]19474.

[110214]Mandel, H.; Brenner, B.; Berant, M.; Rosenberg, N.; Lanir, N.;

- [110215]Jakobs, C.; Fowler, B.; Seligsohn, U.: Coexistence of hereditary
- [110216]homocystinuria and factor V Leiden: effect on thrombosis. New Eng.
- [110217]J. Med. 334: 763–768, 1996.
- [110218]19475.
- [110219]McCully, K. S.; Ragsdale, B. D.: Production of arteriosclerosis
- [110220]by homocystinuria. Am. J. Path. 61: 1–12, 1970.
- [110221]19476.
- [110222]McKusick, V. A.: Heritable Disorders of Connective Tissue St.
- [110223]Louis: C. V. Mosby (pub.) (3rd ed.): 1966. Pp. 155.
- [110224]19477.
- [110225]Mudd, S. H.: Vascular disease and homocysteine metabolism. (Editorial) New
- [110226]Eng. J. Med. 313: 751–753, 1985.
- [110227]19478.
- [110228]Mudd, S. H.; Edwards, W. A.; Loeb, P. M.; Brown, M. S.; Laster,

[110229]L.: Homocystinuria due to cystathionine synthase deficiency: the

[110230]effect of pyridoxine. J. Clin. Invest. 49: 1762–1773, 1970.

[110231]19479.

[110232]Mudd, S. H.; Finkelstein, J. D.; Irreverre, F.; Laster, L.: Homocystinuria:

[110233]an enzymatic defect. Science 143: 1443–1445, 1964.

[110234]19480.

[110235]Mudd, S. H.; Havlik, R.; Levy, H. L.; McKusick, V. A.; Feinleib,

[110236]M.: A study of cardiovascular risk in heterozygotes for homocystinuria. Am.

[110237]J. Hum. Genet. 33: 883–893, 1981.

[110238]19481.

[110239]Mudd, S. H.; Levy, H. L.: Disorders of transulfuration. In: Stanbury,

[110240]J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: Metabolic Basis of

[110241]Inherited Disease. New York: McGraw–Hill (pub.) (4th ed.): 1978.

[110242]pp. 458–503.

[110243]19482.

[110244]Hsieh, C. L.; Leto, T. L.; Lomax, K. J.; Malech, H. L.;
Francke,

[110245]U.: The genes for two neutrophil cytosol factors that are
deficient

[110246]in autosomal forms of chronic granulomatous disease are
on human chromosome

[110247]10 (47 kD), and chromosome 1 cen-q32 (65 kD).
(Abstract) Cytogenet.

[110248]Cell Genet. 51: 1015-1016, 1989.

[110249]19483.

[110250]Karnovsky, M. L.: Steps toward an understanding of
chronic granulomatous

[110251]disease. (Editorial) New Eng. J. Med. 308: 274-275, 1983.

[110252]19484.

[110253]Segal, A. W.: Variations on the theme of chronic granulo-
matous

[110254]disease. Lancet I: 1378-1383, 1985.

[110255]19485.

[110256]Segal, A. W.: The molecular and cellular pathology of
chronic

[110257]granulomatous disease. Europ. J. Clin. Invest. 18:
433–443, 1988.

[110258]19486.

[110259]Aoshima, M.; Nunoi, H.; Shimazu, M.; Shimizu, S.; Tatsuzawa, O.;

[110260]Kenney, R. T.; Kanegasaki, S.: Two-exon skipping due to a
point mutation

[110261]in p67-phox-deficient chronic granulomatous disease.
Blood 88: 1841–1845,

[110262]1996.

[110263]19487.

[110264]Bonizzato, A.; Russo, M. P.; Donini, M.; Dusi, S.: Identification

[110265]of a double mutation (D160V–K161E) (sic) in the p67phox
gene of a

[110266]chronic granulomatous disease patient. Biochem. Biophys.
Res. Commun. 231:

[110267]861–863, 1997.

[110268]19488.

[110269]de Boer, M.; Hilarius–Stokman, P. M.; Hossle, J.–P.; Verhoeven,

- [110270]A. J.; Graf, N.; Kenney, R. T.; Seger, R.; Roos, D.: Autosomal recessive
- [110271]chronic granulomatous disease with absence of the 67-kD cytosolic
- [110272]NADPH oxidase component: identification of mutation and detection
- [110273]of carriers. Blood 83: 531–536, 1994.
- [110274]19489.
- [110275]Dancis, J.; Hutzler, J.; Woody, N. C.; Cox, R. P.: Multiple enzyme
- [110276]defects in familial hyperlysinemia. Pediat. Res. 10: 686–691, 1976.
- [110277]19490.
- [110278]Fischer, A. Q.; Challa, V. R.; Burton, B. K.; McLean, W. T.: Cerebellar
- [110279]hemorrhage complicating isovaleric acidemia: a case report. Neurology 31:
- [110280]746–748, 1981.
- [110281]19491.
- [110282]Sacksteder, K. A.; Biery, B. J.; Morrell, J. C.; Goodman, B. K.;

[110283]Geisbrecht, B. V.; Cox, R. P.; Gould, S. J.; Geraghty, M. T.:

Identification

[110284]of the alpha-aminoadipic semialdehyde synthase gene,
which is defective

[110285]in familial hyperlysinemia. Am. J. Hum. Genet. 66:
1736-1743, 2000.

[110286]19492.

[110287]Gerdes, A.-M.; Gregersen, N.; Ludvigsson, P.; Guttler, F.: A

[110288]Scandinavian case of isovaleric acidemia. J. Inherit.
Metab. Dis. 11:

[110289]219-220, 1988.

[110290]19493.

[110291]Ikeda, Y.; Keese, S. M.; Tanaka, K.: Molecular heterogeneity

[110292]of variant isovaleryl-CoA dehydrogenase from cultured
isovaleric acidemia

[110293]fibroblasts. Proc. Nat. Acad. Sci. 82: 7081-7085, 1985.

[110294]19494.

[110295]Kelleher, J. F., Jr.; Yudkoff, M.; Hutchinson, R.; August, C.

[110296]S.; Cohn, R. M.: The pancytopenia of isovaleric acidemia.
Pediatrics 65:

[110297]1023-1027, 1980.

[110298]19495.

[110299]Naglak, M.; Salvo, R.; Madsen, K.; Dembure, P.; Elsas, L.:

The

[110300]treatment of isovaleric acidemia with glycine supplement.

Pediat.

[110301]Res. 24: 9-13, 1988.

[110302]19496.

[110303]Newman, C. G. H.; Wilson, B. D. R.; Callaghan, P.; Young,

L.:

[110304]Neonatal death associated with isovalericacidaemia.

Lancet II: 439-441,

[110305]1967.

[110306]19497.

[110307]Rhead, W. J.; Tanaka, K.: Demonstration of a specific mi-

tochondrial

[110308]isovaleryl-CoA dehydrogenase deficiency in fibroblasts

from patients

[110309]with isovaleric acidemia. Proc. Nat. Acad. Sci. 77:

580-583, 1980.

[110310]19498.

[110311]Sidbury, J. B., Jr.; Smith, E. K.; Harlan, W.: An inborn error
[110312]of short-chain fatty acid metabolism: the odor-
of-sweaty-feet syndrome. J.
[110313]Pediat. 70: 8-15, 1967.
[110314]19499.
[110315]Tanaka, K.; Budd, M. A.; Efron, M. L.; Isselbacher, K. J.:
Isovaleric
[110316]acidemia: a new genetic defect of leucine metabolism.
Proc. Nat.
[110317]Acad. Sci. 56: 236-242, 1966.
[110318]19500.
[110319]Tanaka, K.; Orr, J.; Isselbacher, K. J.: Identification of b-
hydroxyisovaleric
[110320]acid in the urine of a patient with isovaleric acidemia.
Biochim.
[110321]Biophys. Acta 152: 638-641, 1968.
[110322]19501.
[110323]Vockley, J.; Parimoo, B.; Tanaka, K.: Molecular characteri-
zation
[110324]of four different classes of mutations in the isovaleryl-
CoA dehydrogenase

[110325] gene responsible for isovaleric acidemia. Am. J. Hum. Genet. 49:

[110326] 147–157, 1991.

[110327] 19502.

[110328] Mohandas, T. K.; Chen, X.-N.; Rowe, L. B.; Birkenmeier, E. H.;

[110329] Fanning, A. S.; Anderson, J. M.; Korenberg, J. R.: Localization of

[110330] the tight junction protein gene TJP1 to human chromosome 15q13, distal

[110331] to the Prader-Willi/Angelman region, and to mouse chromosome 7. Genomics 30:

[110332] 594–597, 1995.

[110333] 19503.

[110334] Willott, E.; Balda, M. S.; Fanning, A. S.; Jameson, B.; Van Itallie,

[110335] C.; Anderson, J. M.: The tight junction protein ZO-1 is homologous

[110336] to the Drosophila discs-large tumor suppressor protein of septate

[110337] junctions. Proc. Nat. Acad. Sci. 90: 7834–7838, 1993.

[110338]19504.

[110339]Blancar, M. A.; Crossley, P. H.; Peters, K. G.; Steingrimsson, E.;

[110340]Copeland, N. G.; Jenkins, N. A.; Martin, G. R.; Rutter, W. J.:
Meso1,

[110341]a basic helix-loop-helix protein involved in mammalian
presomitic

[110342]mesoderm development. Proc. Nat. Acad. Sci. 92:
5870-5874, 1995.

[110343]19505.

[110344]Burgess, R.; Rawls, A.; Brown, D.; Bradley, A.; Olson, E. N.:

[110345]Requirement of the paraxis gene for somite formation and
musculoskeletal

[110346]patterning. Nature 384: 570-573, 1996.

[110347]19506.

[110348]Hidai, H.; Quertermous, E. E.; Espinosa, R., III; Le Beau, M.
M.;

[110349]Quertermous, T.: Genomic organization and chromosomal
localization

[110350]of the gene TCF15 encoding the early mesodermal basic
helix-loop-helix

[110351]factor bHLH-EC2. Genomics 30: 598–601, 1995.

[110352]19507.

[110353]Quertermous, E. E.; Hidai, H.; Blonar, M. A.; Quertermous, T.:

[110354]Cloning and characterization of a basic helix–loop–helix protein expressed

[110355]in early mesoderm and the developing somites. Proc. Nat. Acad. Sci. 91:

[110356]7066–7070, 1994.

[110357]19508.

[110358]Quertermous, T.: Personal Communication. Nashville, Tenn. 1/22/1996.

[110359]19509.

[110360]Clevidence, D. E.; Overdier, D. G.; Peterson, R. S.; Porcella,

[110361]A.; Ye, H.; Paulson, K. E.; Costa, R. H.: Members of the HNF–3/forkhead

[110362]family of transcription factors exhibit distinct cellular expression

[110363]patterns in lung and regulate the surfactant protein B promoter. Dev.

[110364]Biol. 166: 195–209, 1994.

[110365]19510.

[110366]Hellqvist, M.; Mahlapuu, M.; Blixt, A.; Enerback, S.; Carlsson,

[110367]P.: The human forkhead protein FREAC-2 contains two functionally

[110368]redundant activation domains and interacts with TBP and TFIIIB. J.

[110369]Biol. Chem. 273: 23335-23343, 1998.

[110370]19511.

[110371]Hellqvist, M.; Mahlapuu, M.; Samuelsson, L.; Enerback, S.; Carlsson,

[110372]P.: Differential activation of lung-specific genes by two forkhead

[110373]proteins, FREAC-1 and FREAC-2. J. Biol. Chem. 271: 4482-4490, 1996.

[110374]19512.

[110375]Larsson, C.; Hellqvist, M.; Pierrou, S.; White, I.; Enerback, S.;

[110376]Carlsson, P.: Chromosomal localization of six human forkhead genes,

[110377]freac-1 (FKHL5), -3 (FKHL7), -4 (FKHL8), -5 (FKHL9), -6 (FKHL10),

[110378]and -8 (FKHL12). Genomics 30: 464-469, 1995.

[110379]19513.

[110380]Pierrou, S.; Hellqvist, M.; Samuelsson, L.; Enerback, S.;
Carlsson,

[110381]P.: Cloning and characterization of seven human forkhead
proteins:

[110382]binding site specificity and DNA bending. EMBO J. 13:
5002-5012,

[110383]1994.

[110384]19514.

[110385]Ernstsson, S.; Pierrou, S.; Hulander, M.; Cederberg, A.;
Hellqvist,

[110386]M.; Carlsson, P.; Enerback, S.: Characterization of the hu-
man forkhead

[110387]gene FREAC-4. J. Biol. Chem. 271: 21094-21099, 1996.

[110388]19515.

[110389]Lorenzetti, D.; Pareyson, D.; Sghirlanzoni, A.; Roa, B. B.;
Abbas,

[110390]N. E.; Pandolfo, M.; Di Donato, S.; Lupski, J. R.: A 1.5-Mb
deletion

[110391]in 17p11.2-p12 is frequently observed in Italian families

with hereditary

[110392]neuropathy with liability to pressure palsies. Am. J. Hum.

Genet. 56:

[110393]91–98, 1995.

[110394]19516.

[110395]Lupski, J. R.; Garcia, C. A.; Parry, G. J.; Patel, P. I.: Charcot–

Marie–Tooth

[110396]polyneuropathy syndrome: clinical, electrophysiological,

and genetic

[110397]aspects.In: Appel, S.: Current Neurology. Chicago: Mosby–

Yearbook

[110398](pub.) 1991. Pp. 1–25.

[110399]19517.

[110400]Lupski, J. R.; Pentao, L.; Williams, L. L.; Patel, P. I.: Stable

[110401]inheritance of the CMT1A DNA duplication in two patients

with CMT1

[110402]and NF1. Am. J. Med. Genet. 45: 92–96, 1993.

[110403]19518.

[110404]Lupski, J. R.; Wise, C. A.; Kuwano, A.; Pentao, L.; Parke, J.

[110405]T.; Glaze, D. G.; Ledbetter, D. H.; Greenberg, F.; Patel, P.

I.:

[110406]Gene dosage is a mechanism for Charcot-Marie-Tooth disease type 1A. Nature

[110407]Genet. 1: 29-33, 1992.

[110408]19519.

[110409]MacMillan, J. C.; Upadhyaya, M.; Harper, P. S.: Charcot-Marie-Tooth

[110410]disease type 1a (CMT1a): evidence for trisomy of the region p11.2

[110411]of chromosome 17 in south Wales families. J. Med. Genet. 29: 12-13,

[110412]1992.

[110413]19520.

[110414]Manfioletti, G.; Ruaro, M. E.; Del Sal, G.; Philipson, L.; Schneider,

[110415]C.: A growth arrest-specific (gas) gene codes for a membrane protein. Molec.

[110416]Cell. Biol. 10: 2924-2930, 1990.

[110417]19521.

[110418]Marques, W., Jr.; Thomas, P. K.; Sweeney, M. G.; Carr, L.; Wood,

- [110419]N. W.: Dejerine-Sottas neuropathy and PMP22 point mutations: a new
base pair substitution and a possible "hot spot" on ser72.
Ann. Neurol. 43:
680-683, 1998.
- [110422]19522.
- [110423]Martinotti, A.; Cariani, C. T.; Melani, C.; Sozzi, G.; Spurr,
N. K.; Pierotti, M. A.; Colombo, M. P.: Isolation and mapping to
17p12-13 of the human homologous of the murine
growth arrest specific
Gas-3 gene. Hum. Molec. Genet. 1: 331-334, 1992.
- [110427]19523.
- [110428]Matise, T. C.; Chakravarti, A.; Patel, P. I.; Lupski, J. R.;
Nelis,
E.; Timmerman, V.; Van Broeckhoven, C.; Weeks, D. E.:
Detection of
tandem duplications and implications for linkage analysis.
Am. J.
Hum. Genet. 54: 1110-1121, 1994.
- [110432]19524.

[110433]Matsunami, N.; Smith, B.; Ballard, L.; Lensch, M. W.;
Robertson,
[110434]M.; Albertsen, H.; Hanemann, C. O.; Muller, H. W.; Bird, T.
D.; White,
[110435]R.; Chance, P. F.: Peripheral myelin protein-22 gene maps
in the
[110436]duplication in chromosome 17p11.2 associated with
Charcot-Marie-Tooth
[110437]1A. Nature Genet. 1: 176-179, 1992.
[110438]19525.
[110439]Nelis, E.; De Jonghe, P.; De Vriendt, E.; Patel, P. I.; Martin,
[110440]J.-J.; Van Broeckhoven, C.: Mutation analysis of the nerve
specific
[110441]promoter of the peripheral myelin protein 22 gene in
CMT1 disease
[110442]and HNPP. J. Med. Genet. 35: 590-593, 1998.
[110443]19526.
[110444]Nicholson, G. A.; Valentijn, L. J.; Cherryson, A. K.; Kenner-
son,
[110445]M. L.; Bragg, T. L.; DeKroon, R. M.; Ross, D. A.; Pollard, J.
D.;

[110446]McLeod, J. G.; Bolhuis, P. A.; Baas, F.: A frame shift mutation in

[110447]the PMP22 gene in hereditary neuropathy with liability to pressure

[110448]palsies. Nature Genet. 6: 263–266, 1994.

[110449]19527.

[110450]Niemann, S.; Sereda, M. W.; Suter, U.; Griffiths, I. R.; Nave,

[110451]K.-A.: Uncoupling of myelin assembly and Schwann cell differentiation

[110452]by transgenic overexpression of peripheral myelin protein 22. J.

[110453]Neurosci. 20: 4120–4128, 2000.

[110454]19528.

[110455]Palau, F.; Lofgren, A.; De Jonghe, P.; Bort, S.; Nelis, E.; Sevilla,

[110456]T.; Martin, J.-J.; Vilchez, J.; Prieto, F.; Van Broeckhoven, C.:

[110457]Origin of the de novo duplication in Charcot–Marie–Tooth disease type

[110458]1A: unequal nonsister chromatid exchange during spermatogenesis. Hum.

[110459]Molec. Genet. 2: 2031–2035, 1993.

[110460]19529.

[110461]Patel, P. I.; Roa, B. B.; Welcher, A. A.; Schoener-Scott, R.;

[110462]Trask, B. J.; Pentao, L.; Snipes, G. J.; Garcia, C. A.; Francke,
U.;

[110463]Shooter, E. M.; Lupski, J. R.; Suter, U.: The gene for the
peripheral

[110464]myelin protein PMP-22 is a candidate for Charcot-
Marie-Tooth disease

[110465]type 1A. Nature Genet. 1: 159-165, 1992.

[110466]19530.

[110467]Pentao, L.; Wise, C. A.; Chinault, A. C.; Patel, P. I.; Lupski,

[110468]J. R.: Charcot-Marie-Tooth type 1A duplication appears to
arise from

[110469]recombination at repeat sequences flanking the 1.5 Mb
monomer unit. Nature

[110470]Genet. 2: 292-300, 1992.

[110471]19531.

[110472]Raeymaekers, P.; Timmerman, V.; Nelis, E.; De Jonghe, P.;
Hoogendijk,

[110473]J. E.; Baas, F.; Barker, D. F.; Martin, J. J.; De Visser, M.; Bol-
huis,

[110474]P. A.; Van Broeckhoven, C.; HMSN Collaborative Research Group:

[110475]Duplication in chromosome 17p11.2 in Charcot-Marie-Tooth neuropathy

[110476]type 1a (CMT 1a). Neuromusc. Disord. 1: 93-97, 1991.
[110477]19532.

[110478]Raeymaekers, P.; Timmerman, V.; Nelis, E.; Van Hul, W.; De Jonghe,

[110479]P.; Martin, J.-J.; Van Broeckhoven, C.; HMSN Collaborative Research

[110480]Group: Estimation of the size of the chromosome 17p11.2 duplication

[110481]in Charcot-Marie-Tooth neuropathy type 1a (CMT1a). J. Med. Genet. 29:

[110482]5-11, 1992.

[110483]19533.

[110484]Reiter, L. T.; Hastings, P. J.; Nelis, E.; De Jonghe, P.; Van

[110485]Broeckhoven, C.; Lupski, J. R.: Human meiotic recombination products

[110486]revealed by sequencing a hotspot for homologous strand exchange in

[110487]multiple HNPP deletion patients. Am. J. Hum. Genet. 62:
1023–1033,

[110488]1998.

[110489]19534.

[110490]Reiter, L. T.; Murakami, T.; Koeuth, T.; Pentao, L.; Muzny,
D.

[110491]M.; Gibbs, R. A.; Lupski, J. R.: A recombination hotspot re-
sponsible

[110492]for two inherited peripheral neuropathies is located near a
mariner

[110493]transposon-like element. Nature Genet. 12: 288–297,
1996. Note:

[110494]Erratum: Nature Genet. 19: 303 only, 1998.

[110495]19535.

[110496]Roa, B. B.; Garcia, C. A.; Pentao, L.; Killian, J. M.; Trask,

[110497]B. J.; Suter, U.; Snipes, G. J.; Ortiz-Lopez, R.; Shooter, E.
M.;

[110498]Patel, P. I.; Lupski, J. R.: Evidence for a recessive PMP22
point

[110499]mutation in Charcot-Marie-Tooth disease type 1A. Nature
Genet. 5:

[110500]189–194, 1993.

[110501]19536.

[110502]Roa, B. B.; Garcia, C. A.; Suter, U.; Kulpa, D. A.; Wise, C. A.;

[110503]Mueller, J.; Welcher, A. A.; Snipes, G. J.; Shooter, E. M.; Patel,
tel,

[110504]P. I.; Lupski, J. R.: Charcot–Marie–Tooth disease type 1A:
association

[110505]with a spontaneous point mutation in the PMP22 gene.
New Eng. J.

[110506]Med. 329: 96–101, 1993.

[110507]19537.

[110508]LeGuern, E.; Gouider, R.; Lopes, J.; Abbas, N.; Gugenheim,
M.;

[110509]Tardieu, S.; Ravise, N.; Leger, J.–M.; Vallat, J.–M.; Bouche,
P.;

[110510]Agid, Y.; Brice, A.; French CMT Collaborative Research
Group: Constant

[110511]rearrangement of the CMT1A–REP sequences in HNPP pa-
tients with a deletion

[110512]in chromosome 17p11.2: a study of 30 unrelated cases.
Hum. Molec.

[110513]Genet. 4: 1673–1674, 1995.

[110514]19538.

[110515]De Smaele, E.; Zazzeroni, F.; Papa, S.; Nguyen, D. U.; Jin, R.;

[110516]Jones, J.; Cong, R.; Franzoso, G.: Induction of gadd45–beta by NF–kappa–B

[110517]downregulates pro–apoptotic JNK signaling. Nature 414: 308–313,

[110518]2001.

[110519]19539.

[110520]Garcia–Cardena, G.; Martasek, P.; Masters, B. S. S.; Skidd, P.

[110521]M.; Couet, J. C.; Li, S.; Lisanti, M. P.; Sessa, W. C.: Dissect– ing

[110522]the interaction between nitric oxide synthase (NOS) and caveolin:

[110523]functional significance of the NOS caveolin binding do– main in vivo. J.

[110524]Biol. Chem. 272: 25437–25440, 1997.

[110525]19540.

[110526]Isom, L. L.; Ragsdale, D. S.; De Jongh, K. S.; Westenbroek,

R.

[110527]E.; Reber, B. F. X.; Scheuer, T.; Catterall, W. A.: Structure
and
[110528]function of the beta-2 subunit of brain sodium channels,
a transmembrane
[110529]glycoprotein with a CAM motif. Cell 83: 433-442, 1995.
[110530]19541.

[110531]Mattei, M. G.; Stoetzel, C.; Perrin-Schmitt, F.: The B-HLH
protein
[110532]encoding the M-twist gene is located by in situ hybridiza-
tion on murine
[110533]chromosome 12. Mammalian Genome 4: 127-128, 1993.
[110534]19542.

[110535]Rose, C. S. P.; Patel, P.; Reardon, W.; Malcolm, S.; Winter,
R.
[110536]M.: The TWIST gene, although not disrupted in Saethre-
Chotzen patients
[110537]with apparently balanced translocations of 7p21, is mu-
tated in familial
[110538]and sporadic cases. Hum. Molec. Genet. 6: 1369-1373,
1997.
[110539]19543.

[110540]Shishido, E.; Higashijima, S.; Emori, Y.; Saigo, K.: Two
FGF-receptor
[110541]homologues of Drosophila: one is expressed in mesoder-
mal primordium
[110542]in early embryos. Development 117: 751-761, 1993.
[110543]19544.
[110544]Struhl, K.: Histone acetylation and transcriptional regula-
tory
[110545]mechanisms. Genes Dev. 12: 599-606, 1998.
[110546]19545.
[110547]Tsuji, K.; Narahara, K.; Yokoyama, Y.; Grzeschik, K.-H.;
Kunz,
[110548]J.: The breakpoint on 7p in a patient with t(6;7) and cran-
iosynostosis
[110549]is spanned by a YAC clone containing the D7S503 locus.
Hum. Genet. 95:
[110550]303-307, 1995.
[110551]19546.
[110552]Yousfi, M.; Lasmoles, F.; El Ghouzzi, V.; Marie, P. J.: Twist
[110553]haploinsufficiency in Saethre-Chotzen syndrome induces
calvarial osteoblast

- [110554]apoptosis due to increased TNF expression and caspase-2 activation. Hum. Molec. Genet. 11: 359-369, 2002.
- [110555]19547.
- [110556]Albrecht, U.; Sutcliffe, J. S.; Cattanach, B. M.; Beechey, C. V.;
- [110557]Armstrong, D.; Eichele, G.; Beaudet, A. L.: Imprinted expression of the murine Angelman syndrome gene, Ube3a, in hippocampal and Purkinje neurons. Nature Genet. 17: 75-78, 1997.
- [110558]19548.
- [110559]Fang, P.; Lev-Lehman, E.; Tsai, T.-F.; Matsuura, T.; Benton, C. S.; Sutcliffe, J. S.; Christian, S. L.; Kubota, T.; Halley, D. J.;
- [110560]Meijers-Heijboer, H.; Langlois, S.; Graham, J. M., Jr.; Beuten, J.;
- [110561]Willems, P. J.; Ledbetter, D. H.; Beaudet, A. L.: The spectrum of mutations in UBE3A causing Angelman syndrome. Hum. Molec. Genet. 8:

[110567]129–135, 1999.

[110568]19549.

[110569]Fung, D. C. Y.; Yu, B.; Cheong, K. F.; Smith, A.; Trent, R. J.

[110570]: UBE3A 'mutations' in two unrelated and phenotypically different

[110571]Angelman syndrome patients. Hum. Genet. 102: 487–492, 1998.

[110572]19550.

[110573]Huang, L.; Kinnucan, E.; Wang, G.; Beaudenon, S.; Howley, P. M.;

[110574]Huibregtse, J. M.; Pavletich, N. P.: Structure of an E6AP-Ubch7 complex:

[110575]insights into ubiquitination by the E2–E3 enzyme cascade. Science 286:

[110576]1321–1326, 1999.

[110577]19551.

[110578]Huibregtse, J. M.; Scheffner, M.; Howley, P. M.: A cellular protein

[110579]mediates association of p53 with the E6 oncoprotein of human papillomavirus

[110580]types 16 or 18. EMBO J. 10: 4129–4135, 1991.

[110581]19552.

[110582]Huibregtse, J. M.; Scheffner, M.; Howley, P. M.: Cloning
and expression

[110583]of the cDNA for E6-AP, a protein that mediates the inter-
action of

[110584]the human papillomavirus E6 oncoprotein with p53.

Molec. Cell. Biol. 13:

[110585]775-784, 1993.

[110586]19553.

[110587]Kishino, T.; Wagstaff, J.: Genomic organization of the
UBE3A/E6-AP

[110588]gene and related pseudogenes. Genomics 47: 101-107,
1998.

[110589]19554.

[110590]Malzac, P.; Webber, H.; Moncla, A.; Graham, J. M., Jr.;
Kukolich,

[110591]M.; Williams, C.; Pagon, R. A.; Ramsdell, L. A.; Kishino, T.;
Wagstaff,

[110592]J.: Mutation analysis of UBE3A in Angelman syndrome pa-
tients. Am.

[110593]J. Hum. Genet. 62: 1353-1360, 1998.

[110594]19555.

[110595]Rougeulle, C.; Glatt, H.; Lalande, M.: The Angelman syndrome

[110596]candidate gene, UBE3A/E6-AP, is imprinted in brain.
(Letter) Nature

[110597]Genet. 17: 14-15, 1997.

[110598]19556.

[110599]Dong, C.; Yang, D. D.; Tournier, C.; Whitmarsh, A. J.; Xu,
J.;

[110600]Davis, R. J.; Flavell, R. A.: JNK is required for effector T-cell
[110601]function but not for T-cell activation. Nature 405: 91-94,
2000.

[110602]19557.

[110603]Gupta, S.; Barrett, T.; Whitmarsh, A. J.; Cavanagh, J.; Sluss,
[110604]H. K.; Derijard, B.; Davis, R. J.: Selective interaction of JNK
protein

[110605]kinase isoforms with transcription factors. EMBO J. 15:
2760-2770,

[110606]1996.

[110607]19558.

[110608]Tournier, C.; Hess, P.; Yang, D. D.; Xu, J.; Turner, T. K.;
 Nimnual,
 [110609]A.; Bar-Sagi, D.; Jones, S. N.; Flavell, R. A.; Davis, R. J.: Re-
 quirement
 [110610]of JNK for stress-induced activation of the cytochrome c-
 mediated
 [110611]death pathway. *Science* 288: 870–874, 2000.
 [110612]19559.

[110613]Daugherty, B. L.; Springer, M. S.: The beta-chemokine re-
 ceptor
 [110614]genes CCR1 (CMKBR1), CCR2 (CMKBR2), and CCR3
 (CMKBR3) cluster within
 [110615]285 kb on human chromosome 3p21. *Genomics* 41:
 294–295, 1997.
 [110616]19560.

[110617]Ayyanathan, K.; Naylor, S. L.; Kunapuli, S. P.: Structural
 characterization
 [110618]and fine chromosomal mapping of the human P2Y(1)
 purinergic receptor
 [110619]gene (P2RY1). *Somat. Cell Molec. Genet.* 22: 419–424,
 1996.
 [110620]19561.

[110621]Ayyanathan, K.; Webbs, T. E.; Sandhu, A. K.; Athwal, R. S.;
Barnard,

[110622]E. A.; Kunapuli, S. P.: Cloning and chromosomal localization of the

[110623]human P2Y₁ purinoceptor. Biochem. Biophys. Res. Commun. 218: 783–788,

[110624]1996.

[110625]19562.

[110626]Leon, C.; Hechler, B.; Freund, M.; Eckly, A.; Vial, C.;
Ohlmann,

[110627]P.; Dierich, A.; LeMeur, M.; Cazenave, J.-P.; Gachet, C.:
Defective

[110628]platelet aggregation and increased resistance to thrombosis in purinergic

[110629]P2Y₁ receptor-null mice. J. Clin. Invest. 104: 1731–1737,
1999.

[110630]19563.

[110631]Leon, C.; Hechler, B.; Vial, C.; Leray, C.; Cazenave, J.-P.;
Gachet,

[110632]C.: The P2Y₁(1) receptor is an ADP receptor antagonized by
ATP and

[110633]expressed in platelets and megakaryoblastic cells. FEBS

Letts. 403:

[110634]26–30, 1997.

[110635]19564.

[110636]Leon, C.; Vial, C.; Cazenave, J.-P.; Gachet, C.: Cloning and sequencing

[110637]of a human cDNA encoding endothelial P2Y1 purinoceptor. Gene 171:

[110638]295–297, 1996.

[110639]19565.

[110640]Hagiwara, Y.; Sasaoka, T.; Araishi, K.; Imamura, M.; Yori-fuji,

[110641]H.; Nonaka, I.; Ozawa, E.; Kikuchi, T.: Caveolin-3 deficiency causes

[110642]muscle degeneration in mice. Hum. Molec. Genet. 9: 3047–3054, 2000.

[110643]19566.

[110644]Herrmann, R.; Straub, V.; Blank, M.; Kutzick, C.; Franke, N.; Jacob,

[110645]E. N.; Lenard, H.-G.; Kroger, S.; Voit, T.: Dissociation of the dystroglycan

[110646]complex in caveolin-3-deficient limb girdle muscular dys-

trophy. Hum.

[110647]Molec. Genet. 9: 2335–2340, 2000.

[110648]19567.

[110649]Matsuda, C.; Hayashi, Y. K.; Ogawa, M.; Aoki, M.; Murayama, K.;

[110650]Nishino, I.; Nonaka, I.; Arahata, K.; Brown, R. H., Jr.: The sarcolemmal

[110651]proteins dysferlin and caveolin-3 interact in skeletal muscle. Hum.

[110652]Molec. Genet. 10: 1761–1766, 2001.

[110653]19568.

[110654]McNally, E. M.: Personal Communication. Chicago, Ill. 6/8/1998.

[110655]19569.

[110656]McNally, E. M.; de Sa Moreira, E.; Duggan, D. J.; Bonnemann, C.

[110657]G.; Lisanti, M. P.; Lidov, H. G. W.; Vainzof, M.; Passos-Bueno, M.

[110658]R.; Hoffman, E. P.; Zatz, M.; Kunkel, L. M.: Caveolin-3 in muscular

[110659]dystrophy. Hum. Molec. Genet. 7: 871–877, 1998.

[110660]19570.

[110661]Minetti, C.; Sotgia, F.; Bruno, C.; Scartezzini, P.; Broda, P.;

[110662]Bado, M.; Masetti, E.; Mazzocco, M.; Egeo, A.; Donati, M.

A.; Volonte,

[110663]D.; Galbiati, F.; Cordone, G.; Bricarelli, F. D.; Lisanti, M. P.;

[110664]Zara, F.: Mutations in the caveolin-3 gene cause autosomal dominant
mal dominant

[110665]limb-girdle muscular dystrophy. Nature Genet. 18:
365-368, 1998.

[110666]19571.

[110667]Sotgia, F.; Minetti, C.; Lisanti, M. P.: Localization of the

[110668]human caveolin-3 gene to the D3S18/D3S4163/D3S4539
locus (3p25), in

[110669]close proximity to the human oxytocin receptor gene:
identification

[110670]of the caveolin-3 gene as a candidate for deletion in
3p-syndrome. FEBS

[110671]Lett. 452: 177-180, 1999.

[110672]19572.

[110673]Sunada, Y.; Ohi, H.; Hase, A.; Ohi, H.; Hosono, T.; Arata,
S.;

[110674]Higuchi, S.; Matsumura, K.; Shimizu, T.: Transgenic mice expressing
[110675]mutant caveolin-3 show severe myopathy associated with
increased nNOS
[110676]activity. Hum. Molec. Genet. 10: 173-178, 2001.
[110677]19573.

[110678]Tang, Z.; Scherer, P. E.; Okamoto, T.; Song, K.; Chu, C.;
Kohtz,
[110679]D. S.; Nishimoto, I.; Lodish, H. F.; Lisanti, M. P.: Molecular
cloning
[110680]of caveolin-3, a novel member of the caveolin gene family
expressed
[110681]predominantly in muscle. J. Biol. Chem. 271: 2255-2261,
1996.
[110682]19574.

[110683]Vorgerd, M.; Bolz, H.; Patzold, T.; Kubisch, C.; Malin, J.-P.;
[110684]Mortier, W.: Phenotypic variability in rippling muscle dis-
ease. Neurology 52:
[110685]1453-1459, 1999.
[110686]19575.

[110687]Furlong, R. A.; Zhou, C. Y.; Ferguson-Smith, M. A.; Affara,

N.

[110688]A.: Characterization of a kinesin-related gene ATSV,
within the tuberous

[110689]sclerosis locus (TSC1) candidate region on chromosome
9q34. Genomics 33:

[110690]421-429, 1996.

[110691]19576.

[110692]Keller, M. P.; Seifried, B. A.; Rabin, B. A.; Chance, P. F.:
Mapping

[110693]of the kinesin-related gene ATSV to chromosome 2q37.
Hum. Genet. 104:

[110694]254-256, 1999.

[110695]19577.

[110696]Kikkawa, M.; Okada, Y.; Hirokawa, N.: 15-angstrom reso-
lution model

[110697]of the monomeric kinesin motor, KIF1A. Cell 100:
241-252, 2000.

[110698]19578.

[110699]Kikkawa, M.; Sablin, E. P.; Okada, Y.; Yajima, H.; Fletterick,

[110700]R. J.; Hirokawa, N.: Switch-based mechanism of kinesin
motors. Nature 411:

[110701]439–445, 2001.

[110702]19579.

[110703]Klopfenstein, D. R.; Tomishige, M.; Stuurman, N.; Vale, R.
D.:

[110704]Role of phosphatidylinositol(4,5)bispophosphate organiza-
tion in membrane

[110705]transport by the Unc104 kinesin motor. Cell 109:
347–358, 2002.

[110706]19580.

[110707]Okada, Y.; Yamazaki, H.; Sekine–Aizawa, Y.; Hirokawa, N.:
The

[110708]neuron–specific kinesin superfamily protein KIF1A is a
unique monomeric

[110709]motor for anterograde axonal transport of synaptic vesicle
precursors. Cell 81:

[110710]769–780, 1995.

[110711]19581.

[110712]Yonekawa, Y.; Harada, A.; Okada, Y.; Funakoshi, T.; Kanai,
Y.;

[110713]Takei, Y.; Terada, S.; Noda, T.; Hirokawa, N.: Defect in
synaptic

[110714]vesicle precursor transport and neuronal cell death in
KIF1A motor
[110715]protein-deficient mice. J. Cell Biol. 141: 431–441, 1998.
[110716]19582.

[110717]Godbout, R.; Squire, J.: Amplification of a DEAD box pro-
tein gene
[110718]in retinoblastoma cell lines. Proc. Nat. Acad. Sci. 90:
7578–7582,
[110719]1993.
[110720]19583.

[110721]Jones, J. M.; Meisler, M. H.; Isom, L. L.: Scn2b, a voltage-
gated
[110722]sodium channel beta-2 gene on mouse chromosome 9.
Genomics 34: 258–259,
[110723]1996.
[110724]19584.

[110725]Bernard, O.; Burkitt, V.; Webb, G. C.; Bottema, C. D. K.;
Nicholl,
[110726]J.; Sutherland, G. R.; Matthew, P.: Structure and chromoso-
mal localization
[110727]of the genomic locus encoding the Kiz1 LIM-kinase gene.

Genomics 35:

[110728]593–596, 1996.

[110729]19585.

[110730]Mao, X.; Jones, T. A.; Williamson, J.; Gutowski, N. J.;
Proschel,

[110731]C.; Noble, M.; Sheer, D.: Assignment of the human and
mouse LIM-kinase

[110732]genes (LIMK1; Limk1) to chromosome bands 7q11.23 and
5G1, respectively,

[110733]by in situ hybridization. Cytogenet. Cell Genet. 74:
190–191, 1996.

[110734]19586.

[110735]Mizuno, K.; Okano, I.; Ohashi, K.; Nunoue, K.; Kuma, K.;
Miyata,

[110736]T.; Nakamura, T.: Identification of a human cDNA encod-
ing a novel

[110737]protein kinase with two repeats of the LIM/double zinc
finger motif. Oncogene 9:

[110738]1605–1612, 1994.

[110739]19587.

[110740]Okano, I.; Hiraoka, J.; Otera, H.; Nunoue, K.; Ohashi, K.;

Iwashita,

[110741]S.; Hirai, M.; Mizuno, K.: Identification and characterization of

[110742]a novel family of serine/threonine kinases containing two N-terminal

[110743]LIM motifs. J. Biol. Chem. 270: 31321–31330, 1995.

[110744]19588.

[110745]Proschel, C.; Blouin, M. J.; Gutowski, N. J.; Ludwig, R.; Noble,

[110746]M.: Limk1 is predominantly expressed in neural tissues and phosphorylates

[110747]serine, threonine and tyrosine residues in vitro. Oncogene 11: 1271–1281,

[110748]1995.

[110749]19589.

[110750]Sotiropoulos, A.; Gineitis, D.; Copeland, J.; Treisman, R.: Signal-regulated

[110751]activation of serum response factor is mediated by changes in actin

[110752]dynamics. Cell 98: 159–169, 1999.

[110753]19590.

[110754]Tassabehji, M.; Metcalfe, K.; Fergusson, W. D.; Carette, M.
J.

[110755]A.; Dore, J. K.; Donnai, D.; Read, A. P.; Proschel, C.;
Gutowski,

[110756]N. J.; Mao, X.; Sheer, D.: LIM-kinase deleted in Williams
syndrome.

[110757](Letter) Nature Genet. 13: 272–273, 1996.

[110758]19591.

[110759]de Sauvage, F. J.; Camerato, T. R.; Goeddel, D. V.: Primary
structure

[110760]and functional expression of the human receptor for Es-
cherichia coli

[110761]heat-stable enterotoxin. J. Biol. Chem. 266:
17912–17918, 1991.

[110762]19592.

[110763]Mann, E. A.; Swenson, E. S.; Copeland, N. G.; Gilbert, D. J.;
Jenkins,

[110764]N. A.; Taguchi, T.; Testa, J. R.; Giannella, R. A.: Localiza-
tion

[110765]of the guanylyl cyclase C gene to mouse chromosome 6
and human chromosome

[110766]12p12. Genomics 34: 265–267, 1996.

[110767]19593.

[110768]Singh, S.; Singh, G.; Heim, J. M.; Gerzer, R.: Isolation and expression

[110769]of a guanylate cyclase-coupled heat stable enterotoxin receptor cDNA

[110770]from a human colonic cell line. Biochem. Biophys. Res. Commun. 179:

[110771]1455-1463, 1991.

[110772]19594.

[110773]Yamamoto, Y.; Huibregtse, J. M.; Howley, P. M.: The human E6-AP

[110774]gene (UBE3A) encodes three potential protein isoforms generated by

[110775]differential splicing. Genomics 41: 263-266, 1997.

[110776]19595.

[110777]Han, J.; Knops, J. F.; Longshore, J. W.; King, P. H.: Localization

[110778]of human elav-like neuronal protein 1 (Hel-N1) on chromosome 9p21

[110779]by chromosome microdissection polymerase chain reaction and fluorescence

[110780]in situ hybridization. Genomics 36: 189–191, 1996.

[110781]19596.

[110782]King, P. H.: Hel–N2: a novel isoform of Hel–N1 which is conserved

[110783]in rat neural tissue and produced in early embryogenesis. Gene 151:

[110784]261–265, 1994.

[110785]19597.

[110786]King, P. H.; Levine, T. D.; Fremeau, R. T., Jr.; Keene, J. D.:

[110787]Mammalian homologs of Drosophila ELAV localized to a neuronal subset

[110788]can bind in vitro to the 3' UTR of mRNA encoding the Id transcriptional

[110789]repressor. J. Neurosci. 14: 1943–52, 1994.

[110790]19598.

[110791]Abel, K. J.; Brody, L. C.; Valdes, J. M.; Erdos, M. R.; McKinley,

[110792]D. R.; Castilla, L. H.; Merajver, S. D.; Couch, F. J.; Friedman, L.

[110793]S.; Ostermeyer, E. A.; Lynch, E. D.; King, M.–C.; Welcsh, P. L.; Osborne–Lawrence,

[110794]S.; Spillman, M.; Bowcock, A. M.; Collins, F. S.; Weber, B.
L.: Characterization

[110795]of EZH1, a human homolog of Drosophila enhancer of
zeste near BRCA1. Genomics 37:

[110796]161–171, 1996.

[110797]19599.

[110798]Quirk, J.; van den Heuvel, M.; Henrique, D.; Marigo, V.;
Jones,

[110799]T. A.; Tabin, C.; Ingham, P. W.: The Smoothed gene and
Hedgehog

[110800]signal transduction in Drosophila and vertebrate develop-
ment. Cold

[110801]Spring Harbor Symp. Quant. Biol. 62: 217–226, 1997.

[110802]19600.

[110803]Sublett, J. E.; Entrekin, R. E.; Look, A. T.; Reardon, D. A.:

[110804]Chromosomal localization of the human Smoothed
gene (SMOH) to 7q32.3

[110805]by fluorescence in situ hybridization and radiation hybrid
mapping. Genomics 50:

[110806]112–114, 1998.

[110807]19601.

- [110808]Xie, J.; Murone, M.; Luoh, S.-M.; Ryan, A.; Gu, Q.; Zhang, C.;
- [110809]Bonifas, J. M.; Lam, C.-W.; Hynes, M.; Goddard, A.; Rosenthal, A.;
- [110810]Epstein, E. H., Jr.; de Sauvage, F. J.: Activating Smoothed mutations
- [110811]in sporadic basal-cell carcinoma. *Nature* 391: 90-92, 1998.
- [110812]19602.
- [110813]Ferbis, D.; Le Chalony, C.; Prosperi, M.-T.; Muleris, M.; Vincent-Salomon,
- [110814]A.; Goubin, G.: Identification, nuclear localization, and binding
- [110815]activities of OZF, a human protein solely composed of zinc finger
- [110816]motifs. *Europ. J. Biochem.* 236: 991-995, 1996.
- [110817]19603.
- [110818]Le Chalony, C.; Apiou, F.; Pibouin, L.; Dutrillaux, B.; Goubin,
- [110819]G.: Constitutive amplification of a zinc finger protein gene in cattle. *DNA*
- [110820]*Cell Biol.* 15: 83-88, 1996.

[110821]19604.

[110822]Le Chalony, C.; Prosperi, M.-T.; Haluza, R.; Apiou, F.;
Dutrillaux,

[110823]B.; Goubin, G.: The OZF gene encodes a protein consisting
essentially

[110824]of zinc-finger motifs. J. Molec. Biol. 236: 399-404, 1994.

[110825]19605.

[110826]Sereda, M.; Griffiths, I. Puhlhofer, A.; Stewart, H.; Rossner,

[110827]M. J.; Zimmermann, F.; Magyar, J. P.; Schneider, A.; Hund,
E.; Meinck,

[110828]H.-M.; Suter, U.; Nave, K.-A.: A transgenic rat model of
Charcot-Marie-Tooth

[110829]disease. Neuron 16: 1049-1060, 1996.

[110830]19606.

[110831]Silander, K.; Meretoja, P.; Nelis, E.; Timmerman, V.; Van
Broeckhoven,

[110832]C.; Aula, P.; Savontaus, M.-L.: A de novo duplication in
17p11.2

[110833]and a novel mutation in the P(0) gene in two Dejerine-Sot-
tas syndrome

[110834]patients. Hum. Mutat. 8: 304-310, 1996.

[110835]19607.

[110836]Stogbauer, F.; Young, P.; Kerschensteiner, M.; Ringelstein, E.

[110837]B.; Assmann, G.; Funke, H.: Recurrent brachial plexus palsies as

[110838]the only clinical expression of hereditary neuropathy with liability

[110839]to pressure palsies associated with a de novo deletion of the peripheral

[110840]myelin protein-22 gene. Muscle Nerve 21: 1199-1201, 1998.

[110841]19608.

[110842]Suter, U.; Moskow, J. J.; Welcher, A. A.; Snipes, G. J.; Kosaras,

[110843]B.; Sidman, R. L.; Buchberg, A. M.; Shooter, E. M.: A leucine-to-proline

[110844]mutation in the putative first transmembrane domain of the 22-kDa

[110845]peripheral myelin protein in the trembler-J mouse. Proc. Nat. Acad.

[110846]Sci. 89: 4382-4386, 1992.

[110847]19609.

- [110848]Suter, U.; Welcher, A. A.; Ozcelik, T.; Snipes, G. J.; Kosaras,
[110849]B.; Francke, U.; Billings-Gagliardi, S.; Sidman, R. L.;
Shooter, E.
[110850]M.: Trembler mouse carries a point mutation in a myelin
gene. Nature 356:
[110851]241-244, 1992.
[110852]19610.
[110853]Takahashi, E.; Takeda, O.; Himoro, M.; Nanao, K.; Takada,
G.;
[110854]Hayasaka, K.: Localization of PMP-22 gene (candidate
gene for the
[110855]Charcot-Marie-Tooth disease 1A) to band 17p11.2 by di-
rect R-banding
[110856]fluorescence in situ hybridization. Jpn. J. Hum. Genet. 37:
303-306,
[110857]1992.
[110858]19611.
[110859]Timmerman, V.; Nelis, E.; Van Hul, W.; Nieuwenhuijsen, B.
W.;
[110860]Chen, K. L.; Wang, S.; Othman, K. B.; Cullen, B.; Leach, R.
J.; Hanemann,

[110861]C. O.; De Jonghe, P.; Raeymaekers, P.; van Ommen, G.-J. B.; Martin,

[110862]J.-J.; Muller, H. W.; Vance, J. M.; Fischbeck, K. H.; Van Broeckhoven,

[110863]C.: The peripheral myelin protein gene PMP-22 is contained within

[110864]the Charcot-Marie-Tooth disease type 1A duplication. Nature Genet. 1:

[110865]171-175, 1992.

[110866]19612.

[110867]Tobler, A. R.; Liu, N.; Mueller, L.; Shooter, E. M.: Differential

[110868]aggregation of the Trembler and Trembler J mutants of peripheral myelin

[110869]protein 22. Proc. Nat. Acad. Sci. 99: 483-488, 2002.

[110870]19613.

[110871]Upadhyaya, M.; Roberts, S. H.; Farnham, J.; MacMillan, J. C.;

[110872]Clarke, A.; Heath, J. P.; Hodges, I. C. G.; Harper, P. S.: Charcot-Marie-Tooth

[110873]disease 1A (CMT1A) associated with a maternal duplication of chromosome

[110874]17p11.2–12. Hum. Genet. 91: 392–394, 1993.

[110875]19614.

[110876]Valentijn, L. J.; Baas, F.; Wolterman, R. A.; Hoogendijk, J. E.;

[110877]van den Bosch, N. H. A.; Zorn, I.; Gabreels–Festen, A. A. W. M.; de

[110878]Visser, M.; Bolhuis, P. A.: Identical point mutations of PMP–22 in

[110879]Trembler–J mouse and Charcot–Marie–Tooth disease type 1A. Nature

[110880]Genet. 2: 288–291, 1992.

[110881]19615.

[110882]Valentijn, L. J.; Bolhuis, P. A.; Zorn, I.; Hoogendijk, J. E.;

[110883]van den Bosch, N.; Hensels, G. W.; Stanton, V. P., Jr.; Housman, D.

[110884]E.; Fischbeck, K. H.; Ross, D. A.; Nicholson, G. A.; Meer–shoek, E.

[110885]J.; Dauwerse, H. G.; van Ommen, G.–J. B.; Baas, F.: The peripheral

[110886]myelin gene PMP–22/GAS–3 is duplicated in Charcot–Marie–Tooth disease

[110887]type 1A. Nature Genet. 1: 166–170, 1992.

[110888]19616.

[110889]Valentijn, L. J.; Ouvrier, R. A.; van den Bosch, N. H. A.; Bolhuis,

[110890]P. A.; Baas, F.; Nicholson, G. A.: Dejerine–Sottas neuropathy is

[110891]associated with a de novo PMP22 mutation. Hum. Mutat. 5: 76–80,

[110892]1995.

[110893]19617.

[110894]Vance, J. M.: Hereditary motor and sensory neuropathies. J.

[110895]Med. Genet. 28: 1–5, 1991.

[110896]19618.

[110897]Wise, C. A.; Garcia, C. A.; Davis, S. N.; Heju, Z.; Pentao, L.;

[110898]Patel, P. I.; Lupski, J. R.: Molecular analyses of unrelated Charcot–Marie–Tooth

[110899](CMT) disease patients suggest a high frequency of the CMT1A duplication. Am.

[110900]J. Hum. Genet. 53: 853–863, 1993.

[110901]19619.

[110902]Wulfsberg, E. A.; Weaver, R. P.; Cunniff, C. M.; Jones, M. C.;

[110903]Jones, K. L.: Chromosome 10qter deletion syndrome: a review and report

[110904]of three new cases. Am. J. Med. Genet. 32: 364–367, 1989.

[110905]19620.

[110906]Young, P.; Wiebusch, H.; Stogbauer, F.; Ringelstein, B.; Assmann,

[110907]G.; Funke, H.: A novel frameshift mutation in PMP22 accounts for

[110908]hereditary neuropathy with liability to pressure palsies. Neurology 48:

[110909]450–452, 1997.

[110910]19621.

[110911]Angrist, M.; Wells, D. E.; Chakravarti, A.; Pandey, A.: Chromosomal

[110912]localization of the mouse Src-like adapter protein (Slap) gene and

[110913]its putative human homolog SLA. Genomics 30: 623–625, 1995.

[110914]19622.

[110915]Holland, S. J.; Liao, X. C.; Mendenhall, M. K.; Zhou, X.;
Pardo,

[110916]J.; Chu, P.; Spencer, C.; Fu, A.; Sheng, N.; Yu, P.; Pali, E.;
Nagin,

[110917]A.; and 14 others: Functional cloning of Src-like adapter
protein-2

[110918](SLAP-2), a novel inhibitor of antigen receptor signaling. J.
Exp.

[110919]Med. 194: 1263-1276, 2001.

[110920]19623.

[110921]Kratchmarova, I.; Sosinowski, T.; Weiss, A.; Witter, K.; Vin-
cenz,

[110922]C.; Pandey, A.: Characterization of promoter region and
genomic structure

[110923]of the murine and human genes encoding Src like adapter
protein. Gene 262:

[110924]267-273, 2001.

[110925]19624.

[110926]Meijerink, P. H. S.; Yanakiev, P.; Zorn, I.; Grierson, A. J.;
Bikker,

- [110927]H.; Dye, D.; Kalaydjieva, L.; Baas, F.: The gene for the human Src-like
[110928]adaptor protein (hSLAP) is located within the 64-kb intron
of the
[110929]thyroglobulin gene. *Europ. J. Biochem.* 254: 297–303,
1998.
[110930]19625.
- [110931]Sosinowski, T.; Pandey, A.; Dixit, V. M.; Weiss, A.: Src-like
[110932]adaptor protein (SLAP) is a negative regulator of T cell re-
ceptor
[110933]signaling. *J. Exp. Med.* 191: 463–474, 2000.
[110934]19626.
- [110935]Bailey, S. M.; Cornforth, M. N.; Kurimasa, A.; Chen, D. J.;
Goodwin,
[110936]E. H.: Strand-specific postreplicative processing of mam-
malian telomeres. *Science* 293:
[110937]2462–2465, 2001.
[110938]19627.
- [110939]Junghans, R. P.; Anderson, C. L.: The protection receptor
for
[110940]IgG catabolism is the beta-2-microglobulin-containing

neonatal intestinal

[110941]transport receptor. Proc. Nat. Acad. Sci. 93: 5512–5516,
1996.

[110942]19628.

[110943]Junghans, R. P.; Ebralidze, A.; Tiwari, B.: Does (CUG)_n repeat

[110944]in DMPK mRNA 'paint' chromosome 19 to suppress distant
genes to create

[110945]the diverse phenotype of myotonic dystrophy? A new hypothesis of long-range

[110946]cis autosomal inactivation. Neurogenetics 3: 59–67, 2001.

[110947]19629.

[110948]Waldmann, T. A.; Strober, W.: Metabolism of immunoglobulins. Prog.

[110949]Allergy 13: 1–110, 1969.

[110950]19630.

[110951]Beddow, A. L.; Richards, S. A.; Orem, N. R.; Macara, I. G.:
The

[110952]Ran/TC4 GTPase-binding domain: identification by expression cloning

[110953]and characterization of a conserved sequence motif. Proc.

Nat. Acad.

[110954]Sci. 92: 3328–3332, 1995.

[110955]19631.

[110956]Fauser, S.; Aslanukov, A.; Roepman, R.; Ferreira, P. A.: Ge-
nomic

[110957]organization, expression, and localization of murine Ran-
binding protein

[110958]2 (RanBP2) gene. Mammalian Genome 12: 406–415, 2001.

[110959]19632.

[110960]Krebber, H.; Bastians, H.; Hoheisel, J.; Lichter, P.; Ponst-
ingl,

[110961]H.; Joos, S.: Localization of the gene encoding the Ran-
binding protein

[110962]RanBP2 to human chromosome 2q11–q13 by fluorescence
in situ hybridization. Genomics 43:

[110963]247–248, 1997.

[110964]19633.

[110965]Pichler, A.; Gast, A.; Seeler, J. S.; Dejean, A.; Melchior, F.

[110966]: The nucleoporin RanBP2 has SUMO1 E3 ligase activity.
Cell 108:

[110967]109–120, 2002.

[110968]19634.

[110969]Wu, J.; Matunis, M. J.; Kraemer, D.; Blobel, G.; Coutavas, E.:

[110970]Nup358, a cytoplasmically exposed nucleoporin with peptide repeats,

[110971]Ran-GTP binding sites, zinc fingers, a cyclophilin A homologous domain,

[110972]and a leucine-rich region. J. Biol. Chem. 270: 14209-14213, 1995.

[110973]19635.

[110974]Yokoyama, N.; Hayashi, N.; Seki, T.; Pante, N.; Ohba, T.; Nishii,

[110975]K.; Kuma, K.; Hayashida, T.; Miyata, T.; Abei, U.; Fukui, M.; Nishimoto,

[110976]T.: A giant nucleopore protein that binds Ran/TC4. Nature 376:

[110977]184-188, 1995.

[110978]19636.

[110979]Chang, A. C.-M.; Janosi, J.; Hulsbeek, M.; de Jong, D.; Jeffrey,

[110980]K. J.; Noble, J. R.; Reddel, R. R.: A novel human cDNA highly homologous

[110981]to the fish hormone stanniocalcin. *Molec. Cell. Endocr.*
112: 241–247,
[110982]1995.
[110983]19637.
[110984]Chang, A. C.–M.; Jeffrey, K. J.; Tokutake, Y.; Shimamoto,
A.; Neumann,
[110985]A. A.; Dunham, M. A.; Cha, J.; Sugawara, M.; Furuichi, Y.;
Reddel,
[110986]R. R.: Human stanniocalcin (STC): genomic structure,
chromosomal
[110987]localization, and the presence of CAG trinucleotide re-
peats. *Genomics* 47:
[110988]393–398, 1998.
[110989]19638.
[110990]Jellinek, D. A.; Chang, A. C.; Larsen, M. R.; Wang, X.;
Robinson,
[110991]P. J.; Reddel, R. R.: Stanniocalcin 1 and 2 are secreted as
phosphoproteins
[110992]from human fibrosarcoma cells. *Biochem. J.* 350:
453–461, 2000.
[110993]19639.

- [110994]Olsen, H. S.; Cepeda, M. A.; Zhang, Q.-Q.; Rosen, C. A.;
Vozzolo,
[110995]B. L.; Wagner, G. F.: Human stanniocalcin: a possible hor-
monal regulator
[110996]of mineral metabolism. Proc. Nat. Acad. Sci. 93:
1792-1796, 1996.
[110997]19640.
- [110998]Varghese, R.; Wong, C. K. C.; Deol, H.; Wagner, G. F.; Di-
Mattia,
[110999]G. E.: Comparative analysis of mammalian stanniocalcin
genes. Endocrinology 139:
[111000]4714-4725, 1998.
[111001]19641.
- [111002]Wagner, G. F.; Guiraudon, C. C.; Milliken, C.; Copp, D. H.:
Immunological
[111003]and biological evidence for a stanniocalcin-like hormone
in human
[111004]kidney. Proc. Nat. Acad. Sci. 92: 1871-1875, 1995.
[111005]19642.
- [111006]Bartsch, O.; Wagner, A.; Hinkel, G. K.; Lichtner, P.; Murken,
J.;

- [111007]Schuffenhauer, S.: No evidence for chromosomal microdeletions at
the second DiGeorge syndrome locus on 10p near
D10S585. (Letter) Am.
J. Med. Genet. 83: 425–426, 1999.
- [111010]19643.
- [111011]Daw, S. C. M.; Taylor, C.; Kraman, M.; Call, K.; Mao, J.;
Schuffenhauer,
S.; Meitinger, T.; Lipson, T.; Goodship, J.; Scambler, P.: A
common
region of 10p deleted in DiGeorge and velocardiofacial
syndromes. Nature
Genet. 13: 458–461, 1996.
- [111014]19644.
- [111015]19644.
- [111016]Lichtner, P.; Konig, R.; Hasegawa, T.; Van Esch, H.;
Meitinger,
T.; Schuffenhauer, S.: An HDR (hypoparathyroidism, deaf-
ness, renal
dysplasia) syndrome locus maps distal to the DiGeorge
syndrome region
on 10p13/14. J. Med. Genet. 37: 33–37, 2000.

[111020]19645.

[111021]Artzt, K.; Shin, H. S.; Bennett, D.: Gene mapping within the
T/t-complex

[111022]of the mouse: II: anomalous position of the H-2 complex
in t-haplotypes. Cell 28:

[111023]471-476, 1982.

[111024]19646.

[111025]Watanabe, T. K.; Fujiwara, T.; Shimizu, F.; Okuno, S.;
Suzuki,

[111026]M.; Takahashi, E.; Nakamura, Y.; Hirai, Y.: Cloning, ex-
pression,

[111027]and mapping of TCTEL1, a putative human homologue of
murine Tcte1,

[111028]to 6q. Cytogenet. Cell Genet. 73: 153-156, 1996.

[111029]19647.

[111030]Davidson, J. D.; Riley, B.; Burright, E.N.; Duvick, L.A.;
Zoghbi,

[111031]H.Y.; Orr, H. T.: Identification and characterization of an
ataxin-1-interactin

[111032]g protein: A1Up, a ubiquitin-like nuclear protein. Hum.
Molec. Genet. 9:

[111033]2305–2312, 2000.

[111034]19648.

[111035]Genis, D.; Matilla, T.; Volpini, V.; Rosell, J.; Davalos, A.; Ferrer,

[111036]I.; Molins, A.; Estivill, X.: Clinical, neuropathologic, and genetic

[111037]studies of a large spinocerebellar ataxia type 1 (SCA1) kindred: (CAG)_n

[111038]expansion and early premonitory signs and symptoms. Neurology 45:

[111039]24–30, 1995.

[111040]19649.

[111041]Klanke, C. A.; Su, Y. R.; Callen, D. F.; Wang, Z.; Meneton, P.;

[111042]Baird, N.; Kandasamy, R. A.; Orlowski, J.; Otterud, B. E.; Leppert,

[111043]M.; Shull, G. E.; Menton, A. G.: Molecular cloning and physical and

[111044]genetic mapping of a novel human Na⁺/H⁺ exchanger (NHE5/SLC9A5) to

[111045]chromosome 16q22.1. Genomics 25: 615–622, 1995.

[111046]19650.

[111047]Chen, C.-Y.; Gherzi, R.; Ong, S.-E.; Chan, E. L.; Rajmakers, R.;

[111048]Pruijn, G. J. M.; Stoecklin, G.; Moroni, C.; Mann, M.; Karin, M.:

[111049]AU binding proteins recruit the exosome to degrade ARE-containing

[111050]mRNAs. Cell 107: 451-464, 2001.

[111051]19651.

[111052]Doyle, K.; Zhang, Y.; Baer, R.; Bina, M.: Distinguishable patterns

[111053]of protein-DNA interactions involving complexes of basic helix-loop-helix

[111054]proteins. J. Biol. Chem. 269: 12099-12105, 1994.

[111055]19652.

[111056]Hu, J.-S.; Olson, E. N.; Kingston, R. E.: HEB, a helix-loop-helix

[111057]protein related to E2A and ITF2 that can modulate the DNA-binding

[111058]ability of myogenic regulatory factors. Molec. Cell. Biol. 12: 1031-1042,

[111059]1992.

[111060]19653.

[111061]Sawada, S.; Littman, D. R.: A heterodimer of HEB and an
E12-related

[111062]protein interacts with the CD4 enhancer and regulates its
activity

[111063]in T-cell lines. Molec. Cell. Biol. 13: 5620-5628, 1993.

[111064]19654.

[111065]Zhang, Y.; Babin, J.; Feldhaus, A. L.; Singh, H.; Sharp, P. A.;

[111066]Bina, M.: HTF4: a new human helix-loop-helix protein.
Nucleic Acids

[111067]Res. 19: 4555 only, 1991.

[111068]19655.

[111069]Zhang, Y.; Bina, M.: The nucleotide sequence of the hu-
man transcription

[111070]factor HTF4a cDNA. DNA Sequence 2: 397-403, 1992.

[111071]19656.

[111072]Zhang, Y.; Flejter, W. L.; Barcroft, C. L.; Riviere, M.; Szpirer,

[111073]J.; Szpirer, C.; Bina, M.: Localization of the human HTF4

transcription

[111074] factors 4 gene (TCF12) to chromosome 15q21. Cytogenet.

Cell Genet. 68:

[111075] 235–238, 1995.

[111076] 19657.

[111077] Hua, X.; Yokoyama, C.; Wu, J.; Briggs, M. R.; Brown, M. S.;

Goldstein,

[111078] J. L.; Wang, X.: SREBP-2, a second basic-he-

lix-loop-helix-leucine

[111079] zipper protein that stimulates transcription by binding to

a sterol

[111080] regulatory element. Proc. Nat. Acad. Sci. 90:

11603–11607, 1993.

[111081] 19658.

[111082] Miserez, A. R.; Cao, G.; Probst, L. C.; Hobbs, H. H.: Struc-

ture

[111083] of the human gene encoding sterol regulatory element

binding protein

[111084] 2 (SREBF2). Genomics 40: 31–40, 1997.

[111085] 19659.

[111086] Yang, J.; Sato, R.; Goldstein, J. L.; Brown, M. S.: Sterol-

resistant

[111087]transcription in CHO cells caused by gene rearrangement
that truncates

[111088]SREBP-2. Genes Dev. 8: 1910–1919, 1994.

[111089]19660.

[111090]Rowen, L.; Young, J.; Birditt, B.; Kaur, A.; Madan, A.;
Philipps,

[111091]D. L.; Qin, S.; Minx, P.; Wilson, R. K.; Hood, L.; Graveley, B.
R.

[111092]: Analysis of the human neurexin genes: alternative splic-
ing and the

[111093]generation of protein diversity. Genomics 79: 587–597,
2002.

[111094]19661.

[111095]Tabuchi, K.; Sudhof, T. C.: Structure and evolution of
neurexin

[111096]genes: insight into the mechanism of alternative splicing.
Genomics 79:

[111097]849–859, 2002.

[111098]19662.

[111099]Ullrich, B.; Ushkaryov, Y. A.; Sudhof, T. C.: Cartography of
neurexins:

[111100]more than 1000 isoforms generated by alternative splicing and expressed

[111101]in distinct subsets of neurons. Neuron 14: 497–507, 1995.

[111102]19663.

[111103]Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T. C.

[111104]; Neurexins: synaptic cell surface proteins related to the alpha-latrotoxin

[111105]receptor and laminin. Science 257: 50–56, 1992.

[111106]19664.

[111107]Bergman, L.; Silins, G.; Grimmond, S.; Hummerich, H.; Stewart,

[111108]C.; Little, P.; Hayward, N.: A 500-kb sequence-ready cosmid contig

[111109]and transcript map of the MEN1 region on 11q13. Genomics 55: 49–56,

[111110]1999.

[111111]19665.

[111112]Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T. C.:

[111113]Neurexins: synaptic cell surface proteins related to the alpha-latrotoxin

[111114]receptor and laminin. Science 257: 50–56, 1992.

[111115]19666.

[111116]Scheiffele, P.; Fan, J.; Choih, J.; Fetter, R.; Serafini, T.:

[111117]Neurologin expressed in nonneuronal cells triggers presynaptic development

[111118]in contacting axons. Cell 101: 657–669, 2000.

[111119]19667.

[111120]Abderrahmani, A.; Steinmann, M.; Plaisance, V.; Niederhauser, G.;

[111121]Haeffliger, J.-A.; Mooser, V.; Bonny, C.; Nicod, P.; Waeber, G.: The

[111122]transcriptional repressor REST determines the cell-specific expression

[111123]of the human MAPK8IP1 gene encoding IB1 (JIP-1). Molec. Cell. Biol. 21:

[111124]7256–7627, 2001.

[111125]19668.

[111126]Cavaloc, Y.; Popielarz, M.; Fuchs, J.-P.; Gattoni, R.; Stevenin,

[111127]J.: Characterization and cloning of the human splicing factor 9G8:

[111128]a novel 35 kDa factor of the serine/arginine protein family. EMBO

[111129]J. 13: 2639–2649, 1994.

[111130]19669.

[111131]Popielarz, M.; Cavaloc, Y.; Mattei, M.–G.; Gattoni, R.; Stevenin,

[111132]J.: The gene encoding human splicing factor 9G8: structure, chromosomal

[111133]localization, and expression of alternatively processed transcripts. J.

[111134]Biol. Chem. 270: 17830–17835, 1995.

[111135]19670.

[111136]Chiang, C.–M.; Roeder, R. G.: Cloning of an intrinsic human TFIID

[111137]subunit that interacts with multiple transcriptional activators. Science 267:

[111138]531–536, 1995.

[111139]19671.

[111140]Wu, Q.; Zhang, T.; Cheng, J.–F.; Kim, Y.; Grimwood, J.;

Schmutz,

[111141]J.; Dickson, M.; Noonan, J. P.; Zhang, M. Q.; Myers, R. M.;

Maniatis,

[111142]T.: Comparative DNA sequence analysis of mouse and human protocadherin

[111143]gene clusters. *Genome Res.* 11: 389–404, 2001.

[111144]19672.

[111145]Kurahashi, H.; Akagi, K.; Inazawa, J.; Ohta, T.; Niikawa, N.; Kayatani,

[111146]F.; Sano, T.; Okada, S.; Nishisho, I.: Isolation and characterization

[111147]of a novel gene deleted in DiGeorge syndrome. *Hum.*

Molec. Genet. 4:

[111148]541–549, 1995.

[111149]19673.

[111150]Aoki, K.; Inazawa, J.; Takahashi, T.; Nakahara, K.; Kasai, M.:

[111151]Genomic structure and chromosomal localization of the gene encoding

[111152]translin, a recombination hotspot binding protein. *Genomics* 43:

[111153]237–241, 1997.

[111154]19674.

[111155]Aoki, K.; Nakahara, K.; Ikegawa, C.; Seto, M.; Takahashi, T.; Minowada,

[111156]J.; Strominger, J. L.; Maziarz, R. T.; Kasai, M.: Nuclear proteins

[111157]binding to a novel target sequence within the recombination hotspot

[111158]regions of Bcl-2 and the immunoglobulin D(H) gene family. Oncogene 9:

[111159]1109-1115, 1994.

[111160]19675.

[111161]Sprecher, E.; Bergman, R.; Szargel, R.; Friedman-Birnbaum, R.;

[111162]Cohen, N.: Identification of a genetic defect in the hairless gene

[111163]in atrichia with papular lesions: evidence for phenotypic heterogeneity

[111164]among inherited atrichias. Am. J. Hum. Genet. 64: 1323-1329, 1999.

[111165]19676.

[111166]Sprecher, E.; Bergman, R.; Szargel, R.; Raz, T.; Labay, V.; Ramon,

[111167]M.; Baruch-Gershoni, R.; Friedman-Birnbaum, R.; Cohen, N.: Atrichia

[111168]with papular lesions maps to 8p in the region containing the human

[111169]hairless gene. Am. J. Med. Genet. 80: 546-550, 1998.

[111170]19677.

[111171]Shannon, R. S.; Mann, J. R.; Harper, E.; Harnden, D. G.; Morten,

[111172]J. E. N.; Herbert, A.: Wilms's tumour and aniridia: clinical and

[111173]cytogenetic features. Arch. Dis. Child. 57: 685-690, 1982.

[111174]19678.

[111175]Strom, T.: A Wilms' tumor family. Acta Paediat. 46: 601-604,

[111176]1957.

[111177]19679.

[111178]Blair, I. P.; Gibson, R. R.; Bennett, C. L.; Chance, P. F.: Search

[111179]for genes involved in Joubert syndrome: evidence that one or more

[111180]major loci are yet to be identified and exclusion of candi-

date genes

[111181]EN1, EN2, FGF8, and BARHL1. Am. J. Med. Genet. 107:
190–196, 2002.

[111182]19680.

[111183]Sweetman, L.; Nyhan, W. L.: Inheritable biotin-treatable
disorders

[111184]and associated phenomena. Ann. Rev. Nutr. 6: 317–343,
1986.

[111185]19681.

[111186]D'Cruz, D.; Taylor, J.; Ahmed, T.; Asherson, R.;
Khamashta, M.;

[111187]Hughes, G. R. V.: Complement factor 2 deficiency: a clinical and

[111188]serological family study. Ann. Rheum. Dis. 51:
1254–1256, 1992.

[111189]19682.

[111190]Day, N. K.; L'Esperance, R.; Good, R. A.; Michael, A. F.;
Hansen,

[111191]J. A.; Dupont, B.; Jersild, C.: Hereditary C2 deficiency: genetic

[111192]studies and association with the HL-A system. J. Exp. Med.
141:

[111193]1464–1469, 1975.

[111194]19683.

[111195]Day, N. K.; Rubinstein, P.; Case, D. J. A.; Good, R. A.; Walker,

[111196]M. E.; Tulchin, N.; Dupont, B.; Jersild, C.: Linkage of gene for

[111197]C2 deficiency and the major histocompatibility complex MHC in man:

[111198]family study of a further case. Vox Sang. 31: 96–102, 1976.

[111199]19684.

[111200]Dewald, G.; Rittner, C.: Polymorphism of the second component

[111201]of human complement (C2): observation of the rare phenotype C2 (=C2

[111202]B) and data on the localization of the C2 locus in the HLA region. Vox

[111203]Sang. 37: 47–54, 1979.

[111204]19685.

[111205]Efthimiou, J.; D'Cruz, D.; Kaplan, P.; Isenberg, D.: Heterozygous

- [111206]C2 deficiency associated with angioedema, myasthenia gravis, and systemic
- [111207]lupus erythematosus. Ann. Rheum. Dis. 45: 428–430, 1986.
- [111208]19686.
- [111209]Einstein, L. P.; Alper, C. A.; Bloch, K. J.; Herrin, J. T.; Rosen, F. S.; David, J. R.; Colten, H. R.: Biosynthetic defect in monocytes from human beings with genetic deficiency of the second component of complement. New Eng. J. Med. 292: 1169–1171, 1975.
- [111212]of complement. New Eng. J. Med. 292: 1169–1171, 1975.
- [111213]19687.
- [111214]Friend, P. S.; Handwerger, B. S.; Kim, Y.; Michael, A. F.; Yunis, E. J.: C2 deficiency in man. Genetic relationship to a mixed lymphocyte reaction determinant (7a*). Immunogenetics 2: 569–576, 1975.
- [111216]reaction determinant (7a*). Immunogenetics 2: 569–576, 1975.
- [111217]19688.
- [111218]Fu, S. M.; Kunkel, H. G.; Brusman, H. P.; Allen, F. H., Jr.; Fotino,

- [111219]M.: Evidence for linkage between HL-A histocompatibility genes and
- [111220]those involved in the synthesis of the second component of complement. J.
- [111221]Exp. Med. 140: 1108-1111, 1974.
- [111222]19689.
- [111223]Gewurz, A.; Lint, T. F.; Roberts, J. L.; Zeitz, H.; Gewurz, H.
- [111224]: Homozygous C2 deficiency with fulminant lupus erythematosus: severe
- [111225]nephritis via the alternative complement pathway. Arthritis Rheum. 21:
- [111226]28-36, 1978.
- [111227]19690.
- [111228]Gibson, D. J.; Glass, D.; Carpenter, C. B.; Schur, P. H.: Hereditary
- [111229]C2 deficiency: diagnosis and HLA gene complex associations. J. Immunogenet. 116:
- [111230]1065-1070, 1976.
- [111231]19691.
- [111232]Johnson, C. A.; Densen, P.; Hurford, R.; Colten, H. R.; Wetzel,

- [111233]R. A.: Deficiency of human complement C2, type I: a 28bp gene deletion
- [111234]leads to a splicing defect. (Abstract) Clin. Res. 39: 304A, 1991.
- [111235]19692.
- [111236]Johnson, C. A.; Densen, P.; Hurford, R. K., Jr.; Colten, H. R.;
- [111237]Wetsel, R. A.: Type I human complement C2 deficiency: a 28-base pair
- [111238]gene deletion causes skipping of exon 6 during RNA splicing. J. Biol.
- [111239]Chem. 267: 9347-9353, 1992.
- [111240]19693.
- [111241]Johnson, C. A.; Densen, P.; Wetsel, R. A.; Cole, F. S.; Goeken,
- [111242]N. E.; Colten, H. R.: Molecular heterogeneity of C2 deficiency. New
- [111243]Eng. J. Med. 326: 871-874, 1992.
- [111244]19694.
- [111245]Kim, Y.; Friend, P. S.; Dresner, I. G.; Yunis, E. J.; Michael,
- [111246]A. F.: Inherited deficiency of the second component of

complement

[111247](C2) with membranoproliferative glomerulonephritis. Am. J. Med. 62:

[111248]765-771, 1977.

[111249]19695.

[111250]Klemperer, M. R.: Hereditary deficiency of the second component

[111251]of complement in man: an immunochemical study. J. Immunogenet. 102:

[111252]168-171, 1969.

[111253]19696.

[111254]Klemperer, M. R.; Austen, K. F.; Rosen, F. S.: Hereditary deficiency

[111255]of second component of complement (C-prime-2) in man: further observations

[111256]on a second kindred. J. Immunogenet. 98: 72-78, 1967.

[111257]19697.

[111258]Klemperer, M. R.; Woodworth, H. C.; Rosen, F. S.; Austen, K. F.

[111259]: Hereditary deficiency of second component of complement (C-prime-2)

[111260]in man. J. Clin. Invest. 45: 880–890, 1966.

[111261]19698.

[111262]Leddy, J. P.; Griggs, R. C.; Klemperer, M. R.; Frank, M. M.:

[111263]Hereditary complement (C2) deficiency with dermato-
myositis. Am. J.

[111264]Med. 58: 83–91, 1975.

[111265]19699.

[111266]Loirat, C.; Levy, M.; Peltier, A. P.; Broyer, M.; Checoury, A.;

[111267]Mathieu, H.: Deficiency of the second component of com-
plement: its

[111268]occurrence with membranoproliferative glomerulonephri-
tis. Arch. Path.

[111269]Lab. Med. 100: 467–472, 1980.

[111270]19700.

[111271]Mahowald, M. L.; Dalmaso, A. P.; Petzel, R. A.; Yunis, E. J.

[111272]: Linkage relationship of C2 deficiency, HLA and glyox-
alase I loci. Vox

[111273]Sang. 37: 321–328, 1979.

[111274]19701.

- [111275]McCarty, D. J.; Tan, E. M.; Zvaifler, N. J.; Koethe, E.;
Duquesnoy,
[111276]R. J.: Serologic studies in a family with heterozygous C2
deficiency. Am.
[111277]J. Med. 71: 945-948, 1981.
[111278]19702.
[111279]Meo, T.; Atkinson, J.; Bernoco, M.; Bernoco, D.; Ceppellini,
R.
[111280]: Mapping of the HLA locus controlling C2 structural vari-
ants and
[111281]linkage disequilibrium between alleles C2-2 and Bw15.
Europ. J. Immun. 6:
[111282]916-919, 1976.
[111283]19703.
[111284]Mortensen, J. P.; Buskjaer, L.; Lamm, L. U.: Studies on the
C2-deficiency
[111285]gene in man. Immunology 39: 541-549, 1980.
[111286]19704.
[111287]Polley, M. J.: Inherited C-prime-2 deficiency in man: lack
of
[111288]immunochemically detectable C-prime-2 protein in

serums from deficient

[111289]individuals. Science 161: 1149–1151, 1968.

[111290]19705.

[111291]Provost, T. T.; Arnett, F. C.; Reichlin, M.: Homozygous C2 deficiency,

[111292]lupus erythematosus, and anti-Ro (SSA) antibodies.

Arthritis Rheum. 26:

[111293]1279–1282, 1983.

[111294]19706.

[111295]Riggs, J. E.; Griggs, R. C.; Rosenfeld, S. I.; May, A. G.; Penn,

[111296]A. S.: Heterozygous C2–deficiency and myasthenia gravis.

Neurology 30:

[111297]871–873, 1980.

[111298]19707.

[111299]Ruddy, S.; Austen, K. F.: Inherited abnormalities of the complement

[111300]system in man. Prog. Med. Genet. 7: 69–95, 1970.

[111301]19708.

[111302]Seligmann, M.; Brouet, J.–C.; Sasportes, M.: Hereditary C2 deficiency

[111303]associated with common variable immunodeficiency. Ann.
Intern. Med. 91:

[111304]216–217, 1979.

[111305]19709.

[111306]Curtis, A.; Nelson, R.; Porteous, M.; Burn, J.; Bhattacharya,

[111307]S. S.: Association of less common cystic fibrosis mutations
with

[111308]a mild phenotype. J. Med. Genet. 28: 34–37, 1991.

[111309]19710.

[111310]Cutting, G. R.; Curristin, S. M.; Nash, E.; Rosenstein, B. J.;

[111311]Lerer, I.; Abeliovich, D.; Hill, A.; Graham, C.: Analysis of
four

[111312]diverse population groups indicates that a subset of cystic
fibrosis

[111313]mutations occur in common among Caucasians. Am. J.
Hum. Genet. 50:

[111314]1185–1194, 1992.

[111315]19711.

[111316]Devoto, M.; Ronchetto, P.; Fanen, P.; Orriols, J. J. T.;
Romeo,

[111317]G.; Goossens, M.; Ferrari, M.; Magnani, C.; Seia, M.; Cremonesi, L.

[111318] Screening for non-delta-F508 mutations in five exons of the cystic

[111319] fibrosis transmembrane conductance regulator (CFTR) gene in Italy. Am.

[111320] J. Hum. Genet. 48: 1127-1132, 1991.

[111321] 19712.

[111322] de Vries, H. G.; Collee, J. M.; de Walle, H. E. K.; van Veldhuizen,

[111323] M. H. R.; Smit Sibinga, C. T.; Scheffer, H.; ten Kate, L. P.: Prevalence

[111324] of delta-F508 cystic fibrosis carriers in The Netherlands: logistic

[111325] regression on sex, age, region of residence and number of offspring. Hum.

[111326] Genet. 99: 74-79, 1997.

[111327] 19713.

[111328] Dorin, J. R.; Dickinson, P.; Alton, E. W. F. W.; Smith, S. N.;

[111329] Geddes, D. M.; Stevenson, B. J.; Kimber, W. L.; Fleming, S.; Clarke,

- [111330]A. R.; Hooper, M. L.; Anderson, L.; Beddington, R. S. P.;
Porteous,
- [111331]D. J.: Cystic fibrosis in the mouse by targeted insertional
mutagenesis. *Nature* 359:
- [111332]211–215, 1992.
- [111333]19714.
- [111334]Dork, T.; El-Harith, E.-H. A.; Stuhmann, M.; Macek, M.,
Jr.;
- [111335]Egan, M.; Cutting, G. R.; Tzetzis, M.; Kanavakis, E.; Carles,
S.; Claustres,
- [111336]M.; Padoa, C.; Ramsay, M.; Schmidtke, J.: Evidence for a
common ethnic
- [111337]origin of cystic fibrosis mutation 3120+1G-to-A in di-
verse populations.
- [111338](Letter) *Am. J. Hum. Genet.* 63: 656–662, 1998.
- [111339]19715.
- [111340]Dork, T.; Wulbrand, U.; Richter, T.; Neumann, T.; Wolfes,
H.;
- [111341]Wulf, B.; Maass, G.; Tummeler, B.: Cystic fibrosis with three
mutations
- [111342]in the cystic fibrosis transmembrane conductance regula-
tor gene. *Hum.*

[111343]Genet. 87: 441–446, 1991.

[111344]19716.

[111345]Dumur, V.; Lafitte, J. J.; Gervais, R.; Debaecker, D.;
Kesteloot,

[111346]M.; Lalau, G.; Roussel, P.: Abnormal distribution of cystic
fibrosis

[111347]delta-F508 allele in adults with chronic bronchial hyper-
secretion. Lancet 335:

[111348]1340, 1990.

[111349]19717.

[111350]Ferec, C.; Audrezet, M. P.; Mercier, B.; Guillermit, H.;
Moullier,

[111351]P.; Quere, I.; Verlingue, C.: Detection of over 98% cystic fi-
brosis

[111352]mutations in a Celtic population. Nature Genet. 1:
188–191, 1992.

[111353]19718.

[111354]Kelly, T. E.: Personal Communication. Charlottesville, Va.

[111355]1977.

[111356]19719.

- [111357]Lindor, N. M.; Furuichi, Y.; Kitao, S.; Shimamoto, A.; Arndt,
[111358]C.; Jalal, S.: Rothmund–Thomson syndrome due to RECQ4
helicase mutations:
[111359]report and clinical and molecular comparisons with Bloom
syndrome
[111360]and Werner syndrome. Am. J. Med. Genet. 90: 223–228,
2000.
[111361]19720.
- [111362]Bodzioch, M.; Orso, E.; Klucken, J.; Langmann, T.;
Bottcher, A.;
[111363]Diederich, W.; Drobnik, W.; Barlage, S.; Buchler, C.;
Porsch–Ozcurumez,
[111364]M.; Kaminski, W. E.; Hahmann, H. W.; Oette, K.; Rothe, G.;
Aslanidis,
[111365]C.; Lackner, K. J.; Schmitz, G.: The gene encoding ATP–
binding cassette
[111366]transporter 1 is mutated in Tangier disease. Nature Genet.
22: 347–351,
[111367]1999.
[111368]19721.
- [111369]Brooks–Wilson, A.; Marcil, M.; Clee, S. M.; Zhang, L.–H.;
Roomp,

- [111370]K.; van Dam, M.; Yu, L.; Brewer, C.; Collins, J. A.; Molhuizen, H.
- [111371]O. F.; Loubser, O.; Ouelette, B. F. F.; and 14 others: Mutations
- [111372]in ABC1 in Tangier disease and familial high-density lipoprotein deficiency. *Nature*
- [111373]*Genet.* 22: 336–345, 1999.
- [111374]19722.
- [111375]Hoogendijk, J. E.; Hensels, G. W.; Gabreels–Festen, A. A. W. M.;
- [111376]Gabreels, F. J. M.; Janssen, E. A. M.; De Jonghe, P.; Martin, J.–J.;
- [111377]Van Broeckhoven, C.; Valentijn, L. J.; Baas, F.; de Visser, M.; Bolhuis,
- [111378]P. A.: De–novo mutation in hereditary motor and sensory neuropathy
- [111379]type I. *Lancet* 339: 1081–1082, 1992.
- [111380]19723.
- [111381]Menasche, G.; Pastural, E.; Feldmann, J.; Certain, S.; Ersoy, F.;
- [111382]Dupuis, S.; Wulffraat, N.; Bianchi, D.; Fischer, A.; Le Deist, F.;

[111383] de Saint Basile, G.: Mutations in RAB27A cause Griscelli syndrome
[111384] associated with haemophagocytic syndrome. Nature
Genet. 25: 173–176,
[111385] 2000.
[111386] 19724.
[111387] Pastural, E.; Barrat, F. J.; Dufourcq-Lagelouse, R.; Certain, S.;
[111388] Sanal, O.; Jabado, N.; Seger, R.; Griscelli, C.; Fischer, A.; de Saint
[111389] Basile, G.: Griscelli disease maps to chromosome 15q21 and is associated
[111390] with mutations in the myosin-Va gene. Nature Genet. 16:
289–292,
[111391] 1997.
[111392] 19725.
[111393] Asamoah, A.; Wilson, A. F.; Elston, R. C.; Dalferes, E., Jr.; Berenson,
[111394] G. S.: Segregation and linkage analyses of dopamine-beta-hydroxylase
[111395] activity in a six-generation pedigree. Am. J. Med. Genet. 27: 613–621,

[111396]1987.

[111397]19726.

[111398]Biaggioni, I.; Goldstein, D. S.; Atkinson, T.; Robertson, D.:

[111399]Dopamine-beta-hydroxylase deficiency in humans. Neu-
rology 40: 370-373,

[111400]1990.

[111401]19727.

[111402]Biaggioni, I.; Robertson, D.: Endogenous restoration of
noradrenaline

[111403]by precursor therapy in dopamine-beta-hydroxylase defi-
ciency. Lancet II:

[111404]1170-1172, 1987.

[111405]19728.

[111406]Craig, S. P.; Buckle, V. J.; Lamouroux, A.; Mallet, J.; Craig,

[111407]I. W.: Localization of the human dopamine beta hydroxy-
lase (DBH)

[111408]gene to chromosome 9q34. Cytogenet. Cell Genet. 48:
48-50, 1988.

[111409]19729.

[111410]Dunnette, J.; Weinshilboum, R.: Human serum dopamine

beta-hydroxylase:

[111411]correlation of enzymatic activity with immunoreactive protein in genetically

[111412]defined samples. Am. J. Hum. Genet. 28: 155-166, 1976.

[111413]19730.

[111414]Dunnette, J.; Weinshilboum, R.: Inheritance of low immunoreactive

[111415]human plasma dopamine-beta-hydroxylase: radioimmunoassay studies. J.

[111416]Clin. Invest. 60: 1080-1087, 1977.

[111417]19731.

[111418]Elston, R. C.; Namboodiri, K. K.; Hames, C. G.: Segregation and

[111419]linkage analysis of dopamine-beta-hydroxylase activity. Hum. Hered. 29:

[111420]284-292, 1979.

[111421]19732.

[111422]Gershon, E. S.; Goldin, L. R.: Segregation and linkage studies

[111423]of plasma dopamine-beta-hydroxylase (DBH), erythrocyte catechol-O-methyltransferase

[111424](COMT) and platelet monoamine oxidase (MAO): possible linkage between

[111425]the ABO locus and a gene controlling DBH activity.

(Abstract) Am.

[111426]J. Hum. Genet. 33: 136A only, 1981.

[111427]19733.

[111428]Goldin, L. R.; Gershon, E. S.; Lake, C. R.; Murphy, D. L.; McGinniss,

[111429]M.; Sparkes, R. S.: Segregation and linkage studies of plasma dopamine-beta-hydroxylase

[111430](DBH), erythrocyte catechol-O-methyltransferase (COMT), and platelet

[111431]monoamine oxidase (MAO): possible linkage between the ABO locus and

[111432]a gene controlling DBH activity. Am. J. Hum. Genet. 34: 250-262,

[111433]1982.

[111434]19734.

[111435]Joh, T. H.; Baetge, E. E.; Reis, D. J.: Evidence for the existence

[111436]of a single gene or linked genes coding for catecholamine biosynthetic

[111437]enzymes. Trans. Assoc. Am. Phys. 96: 38–43, 1983.

[111438]19735.

[111439]Joh, T. H.; Baetge, E. E.; Ross, M. E.; Albert, V. R.; Moon, H.

[111440]M.; Reis, D. J.: Existence of catecholamine biosynthetic
enzyme gene

[111441]family. (Abstract) Clin. Res. 31: 528 only, 1983.

[111442]19736.

[111443]Joh, T. H.; Baetge, E. E.; Ross, M. E.; Reis, D. J.: Biochem-
istry

[111444]and molecular biology of catecholamine neurons: a single
gene or gene

[111445]family hypothesis. Clin. Exp. Hypertension 6A: 11–21,
1984.

[111446]19737.

[111447]Kobayashi, K.; Kurosawa, Y.; Fujita, K.; Nagatsu, T.: Hu-
man dopamine

[111448]beta-hydroxylase gene: two mRNA types having different
3-prime-terminal

[111449]regions are produced through alternative polyadenylation.
Nucleic

[111450]Acids Res. 17: 1089–1102, 1989.

[111451]19738.

[111452]Lamouroux, A.; Vigny, A.; Faucon Biguet, N.; Darmon, M.
C.; Franck,

[111453]R.; Henry, J.-P.; Mallet, J.: The primary structure of human
dopamine-beta-hydroxylase:

[111454]insights into the relationship between the soluble and the
membrane-bound

[111455]forms of the enzyme. EMBO J. 6: 3931-3937, 1987.

[111456]19739.

[111457]Lea, R. A.; Dohy, A.; Jordan, K.; Quinlan, S.; Brimage, P. J.;

[111458]Griffiths, L. R.: Evidence for allelic association of the
dopamine

[111459]beta-hydroxylase gene (DBH) with susceptibility to typical
migraine. Neurogenetics 3:

[111460]35-40, 2000.

[111461]19740.

[111462]Mathias, C. J.; Bannister, R. B.; Cortelli, P.; Heslop, K.; Po-
lak,

[111463]J. M.; Raimbach, S.; Springall, D. R.; Watson, L.: Clinical,
autonomic

[111464]and therapeutic observations in two siblings with postural

hypotension

[111465]and sympathetic failure due to an inability to synthesize
noradrenaline

[111466]from dopamine because of a deficiency of dopamine beta
hydroxylase. Quart.

[111467]J. Med. 75: 617-633, 1990.

[111468]19741.

[111469]McKinney, E. F.; Walton, R. T.; Yudkin, P.; Fuller, A.; Hal-
dar,

[111470]N. A.; Mant, D.; Murphy, M.; Welsh, K. I.; Marshall, S. E.:
Association

[111471]between polymorphisms in dopamine metabolic enzymes
and tobacco consumption

[111472]in smokers. Pharmacogenetics 10: 483-491, 2000.

[111473]19742.

[111474]O'Malley, K. L.; Mauron, A.; Raese, J.; Barchas, J. D.; Kedes,

[111475]L.: Genes for catecholamine biosynthesis: cloning by ex-
pression and

[111476]identification of the cDNA for rat dopamine beta-
hydroxylase. Proc.

[111477]Nat. Acad. Sci. 80: 2161-2165, 1983.

[111478]19743.

[111479]Ogihara, T.; Nugent, C. A., Jr.; Shen, S.-W.; Goldfein, S.:
Serum

[111480]dopamine-beta-hydroxylase activity in parents and chil-
dren. J. Lab.

[111481]Clin. Med. 85: 566-573, 1975.

[111482]19744.

[111483]Perry, S. E.; Summar, M. L.; Phillips, J. A., III; Robertson,

[111484]D.: Linkage analysis of the human dopamine beta-
hydroxylase gene. Genomics 10:

[111485]493-495, 1991.

[111486]19745.

[111487]Robertson, D.; Haile, V.; Perry, S. E.; Robertson, R. M.;
Phillips,

[111488]J. A., III; Biaggioni, I.: Dopamine beta-hydroxylase defi-
ciency:

[111489]a genetic disorder of cardiovascular regulation. Hyperten-
sion 18:

[111490]1-8, 1991.

[111491]19746.

- [111492] Ross, S. B.; Wetterberg, L.; Myrhed, M.: Genetic control of plasma dopamine-beta-hydroxylase. Life Sci. 12: 529-532, 1973.
- [111493] 19747.
- [111494] Schanberg, S. M.; Stone, R. A.; Kirshner, N.; Gunnells, J. C.; Robinson, R. R.: Plasma dopamine beta-hydroxylase: a possible aid in the study and evaluation of hypertension. Science 183: 523-525, 1974.
- [111495] 19748.
- [111496] Hoefsloot, L. H.; van der Ploeg, A. T.; Kroos, M. A.; Hoogeveen-Westerveld, M.; Oostra, B. A.; Reuser, A. J. J.: Adult and infantile glycogenosis type II in one family, explained by allelic diversity. Am. J. Hum. Genet. 46: 45-52, 1990.
- [111497] 19749.
- [111498] de Villiers, J. N. P.; Hillermann, R.; Loubser, L.; Kotze, M.

[111506]J.: Spectrum of mutations in the HFE gene implicated in haemochromatosis

[111507]and porphyria. Hum. Molec. Genet. 8: 1517–1522, 1999.

[111508]19750.

[111509]Marlar, R. A.; Griffin, J. H.: Deficiency of protein C inhibitor

[111510]in combined factor V/VIII deficiency disease. J. Clin. Invest. 66:

[111511]1186–1189, 1980.

[111512]19751.

[111513]Meijers, J. C.; Chung, D. W.: Organization of the gene coding

[111514]for human protein C inhibitor (plasminogen activator inhibitor-3):

[111515]assignment of the gene to chromosome 14. J. Biol. Chem. 266: 15028–15034,

[111516]1991.

[111517]19752.

[111518]Nichols, W. C.; Seligsohn, U.; Zivelin, A.; Terry, V. H.; Arnold,

[111519]N. D.; Siemieniak, D. R.; Kaufman, R. J.; Ginsburg, D.: Linkage of

[111520]combined factors V and VIII deficiency to chromosome
18q by homozygosity

[111521]mapping. J. Clin. Invest. 99: 596–601, 1997.

[111522]19753.

[111523]Suzuki, K.; Deyashiki, Y.; Nishioka, J.; Kurachi, K.; Akira,
M.;

[111524]Yamamoto, S.; Hashimoto, S.: Characterization of a cDNA
for human

[111525]protein C inhibitor: a new member of the plasma serine
protease inhibitor

[111526]superfamily. J. Biol. Chem. 262: 611–616, 1987.

[111527]19754.

[111528]Abramson, R. D.; Barbosa, P.; Kalumuck, K.; O'Brien, W. E.:
Characterization

[111529]of the human argininosuccinate lyase gene and analysis of
exon skipping. Genomics 10:

[111530]126–132, 1991.

[111531]19755.

[111532]Allan, J. D.; Cusworth, D. C.; Dent, C. E.; Wilson, V. K.: A
disease,

[111533]probably hereditary, characterized by severe mental defi-

ciency and

[111534]a constant gross abnormality of amino acid metabolism.

Lancet I:

[111535]182–187, 1958.

[111536]19756.

[111537]Asai, K.; Ishii, S.; Ohta, S.; Furusho, K.: Fatal hyperam-
monaemia

[111538]in argininosuccinic aciduria following enflurane anaesthe-
sia. (Letter) Europ.

[111539]J. Paediat. 157: 169–170, 1997.

[111540]19757.

[111541]Bohles, H.; Heid, H.; Harms, D.; Schmid, D.; Fekl, W.:
Argininosuccinic

[111542]aciduria: metabolic studies and effects of treatment with
keto-analogues

[111543]of essential amino acids. Europ. J. Pediat. 128: 225–233,
1978.

[111544]19758.

[111545]Brusilow, S. W.; Batshaw, M. L.: Arginine therapy of argini-
nosuccinase

[111546]deficiency. Lancet I: 124–127, 1979.

[111547]19759.

[111548]Cathelineau, L.; Dinh, D. P.; Briand, P.; Kamoun, P.: Studies

[111549]on complementation in argininosuccinate synthetase and
argininosuccinate

[111550]lyase deficiencies in human fibroblasts. Hum. Genet. 57:
282-284,

[111551]1981.

[111552]19760.

[111553]Collins, F. S.; Summer, G. K.; Schwartz, R. P.; Parke, J. C.,
Jr.

[111554]: Neonatal argininosuccinic aciduria--survival after early
diagnosis

[111555]and dietary management. J. Pediat. 96: 429-431, 1980.

[111556]19761.

[111557]Coryell, M. E.; Hall, W. K.; Thevaos, T. G.; Welter, D. A.;
Gatz,

[111558]A. J.; Horton, B. F.; Sisson, B. D.; Looper, J. W., Jr.; Farrow,
R.

[111559]T.: Familial study of human enzyme defect, argininosuc-
cinic aciduria. Biochem.

[111560]Biophys. Res. Commun. 14: 307-312, 1964.

[111561]19762.

[111562]Fleisher, L. D.; Rassin, D. K.; Desnick, R. J.; Salwen, H. R.;

[111563]Rogers, P.; Bean, M.; Gaull, G. E.: Argininosuccinic
aciduria: prenatal

[111564]studies in a family at risk. Am. J. Hum. Genet. 31:
439-445, 1979.

[111565]19763.

[111566]Glick, N. R.; Snodgrass, P. J.; Schafer, I. A.: Neonatal
argininosuccinic

[111567]aciduria with normal brain and kidney but absent liver
argininosuccinate

[111568]lyase activity. Am. J. Hum. Genet. 28: 22-30, 1976.

[111569]19764.

[111570]Goodman, S. I.; Mace, J. W.; Turner, B.; Garrett, W. J.: An-
tenatal

[111571]diagnosis of argininosuccinic aciduria. Clin. Genet. 4:
236-240,

[111572]1973.

[111573]19765.

[111574]Kint, J. A.; Carton, D.: Deficient argininosuccinase activity

[111575]in brain in argininosuccinicaciduria. (Letter) Lancet II: 635
only,
[111576]1968.
[111577]19766.
[111578]Kvedar, J. C.; Baden, H. P.; Baden, L. A.; Shih, V. E.;
Kolodny,
[111579]E. H.: Dietary management reverses grooving and abnormal polarization
[111580]of hair shafts in argininosuccinase deficiency. Am. J. Med.
Genet. 40:
[111581]211-213, 1991.
[111582]19767.
[111583]Levin, B.: Argininosuccinic aciduria. Am. J. Dis. Child. 113:
[111584]162-165, 1967.
[111585]19768.
[111586]Levin, B.; MacKay, H. M.; Oberholzer, V. G.: Argininosuccinic
[111587]aciduria: an inborn error of amino acid metabolism. Arch.
Dis. Child. 36:
[111588]622-632, 1961.

[111589]19769.

[111590]Lewis, P. D.; Miller, A. L.: Argininosuccinic aciduria: case

[111591]report with neuropathological findings. Brain 93:

413-422, 1970.

[111592]19770.

[111593]Matuo, S.; Tatsuno, M.; Kobayashi, K.; Saheki, T.; Miyata,

T.;

[111594]Iwanaga, S.; Amaya, Y.; Mori, M.: Isolation of cDNA clones

of human

[111595]argininosuccinate lyase and corrected amino acid se-

quence. FEBS Lett. 234:

[111596]395-399, 1988.

[111597]19771.

[111598]McInnes, R. R.; Shih, V.; Chilton, S.: Interallelic comple-

mentation

[111599]in an inborn error of metabolism: genetic heterogeneity in

argininosuccinate

[111600]lyase deficiency. Proc. Nat. Acad. Sci. 81: 4480-4484,

1984.

[111601]19772.

[111602]Moser, H. W.; Efron, M. L.; Brown, H.; Diamond, R.; Neu-

mann, C.

[111603]G.: Argininosuccinic aciduria: report of two cases and demonstration

[111604]of intermittent elevation of blood ammonia. Am. J. Med. 42: 9–26,

[111605]1967.

[111606]19773.

[111607]Sinclair, D. A.; Mills, K.; Guarente, L.: Accelerated aging and

[111608]nucleolar fragmentation in yeast sgs1 mutants. Science. 277: 1313–1316,

[111609]1997.

[111610]19774.

[111611]Koster, J. F.; Busch, H. F. M.; Slee, R. G.; van Weerden, T. W.

[111612]; Glycogenosis type II: the infantile– and late–onset acid maltase

[111613]deficiency observed in one family. Clin. Chim. Acta 87: 451–453,

[111614]1978.

[111615]19775.

- [111616] Kroos, M. A.; Van der Kraan, M.; Van Diggelen, O. P.; Kleijer,
[111617] W. J.; Reuser, A. J. J.: Two extremes of the clinical spectrum of
[111618] glycogen storage disease type II in one family: a matter of
genotype. Hum.
[111619] Mutat. 9: 17–22, 1997.
[111620] 19776.
[111621] Kroos, M. A.; Van der Kraan, M.; Van Diggelen, O. P.; Kleijer,
[111622] W. J.; Reuser, A. J. J.; Van den Boogaard, M. J.; Ausems, M.
G. E.
[111623] M.; Ploos van Amstel, H. K.; Poenaru, L.; Nicolino, M.;
Wevers, R.
[111624]: Glycogen storage disease type II: frequency of three
common mutant
[111625] alleles and their associated clinical phenotypes studied in
121 patients. J.
[111626] Med. Genet. 32: 836–837, 1995.
[111627] 19777.
[111628] Loonen, M. C. B.; Schram, A. W.; Koster, J. F.; Niermeijer,
M.

[111629]F.; Busch, H. F. M.; Martin, J. J.; Brouwer-Kelder, B.; Mekes, W.;

[111630]Slee, R. G.; Tager, J. M.: Identification of heterozygotes for glycogenosis

[111631]2 (acid maltase deficiency). Clin. Genet. 19: 55-63, 1981.

[111632]19778.

[111633]Martiniuk, F.; Bodkin, M.; Tzall, S.; Hirschhorn, R.: Identification

[111634]of the base-pair substitution responsible for a human acid alpha glucosidase

[111635]allele with lower 'affinity' for glycogen (GAA 2) and transient gene

[111636]expression in deficient cells. Am. J. Hum. Genet. 47: 440-445, 1990.

[111637]19779.

[111638]Martiniuk, F.; Chen, A.; Mack, A.; Arvanitopoulos, E.; Chen, Y.;

[111639]Rom, W. N.; Codd, W. J.; Hanna, B.; Alcabes, P.; Raben, N.; Plotz,

[111640]P.: Carrier frequency for glycogen storage disease type II in New

[111641]York and estimates of affected individuals born with the

disease.

[111642](Letter) Am. J. Med. Genet. 79: 69–72, 1998.

[111643]19780.

[111644]Diamond, T.; Stiel, D.; Posen, S.: Osteoporosis in
hemochromatosis:

[111645]iron excess, gonadal deficiency, or other factors? Ann. In-
tern. Med. 110:

[111646]430–436, 1989.

[111647]19781.

[111648]Dokal, I.; Lord, D.; Rhodes, D.; Bydder, G.; Cox, T.: Detec-
tion

[111649]of hereditary haemochromatosis in an HLA-identical pedi-
gree showing

[111650]discordance between HLA class I genes and the disease
locus. Hum.

[111651]Genet. 88: 209–214, 1991.

[111652]19782.

[111653]Dugast, I. J.; Papadopoulos, P.; Zappone, E.; Jones, C.;
Theriault,

[111654]K.; Handelman, G. J.; Benarous, R.; Drysdale, J. W.: Identi-
fication

[111655] of two human ferritin H genes on the short arm of chromosome 6. Genomics 6:

[111656] 204–211, 1990.

[111657] 19783.

[111658] Edwards, C. Q.; Carroll, M.; Bray, P. F.; Cartwright, G. E.:

[111659] Hereditary hemochromatosis: diagnosis in siblings and children. New

[111660] Eng. J. Med. 297: 7–13, 1977.

[111661] 19784.

[111662] Edwards, C. Q.; Cartwright, G. E.; Skolnick, M. H.; Amos, D. B.

[111663]: Homozygosity for hemochromatosis: clinical manifestations. Ann.

[111664] Intern. Med. 93: 519–525, 1980.

[111665] 19785.

[111666] Edwards, C. Q.; Griffen, L. M.; Dadone, M. M.; Skolnick, M. H.;

[111667] Kushner, J. P.: Mapping the locus for hereditary hemochromatosis:

[111668] localization between HLA-B and HLA-A. Am. J. Hum. Genet. 38: 805–811,

[111669]1986.

[111670]19786.

[111671]Edwards, C. Q.; Griffen, L. M.; Dadone, M. M.; Skolnick, M.
H.;

[111672]Kushner, J. P.: The locus for hereditary hemochromatosis
maps between

[111673]HLA-A and HLA-B. (Abstract) Cytogenet. Cell Genet. 40:
620, 1985.

[111674]19787.

[111675]Edwards, C. Q.; Griffen, L. M.; Goldgar, D.; Drummond, C.;
Skolnick,

[111676]M. H.; Kushner, J. P.: Prevalence of hemochromatosis
among 11,065

[111677]presumably healthy blood donors. New Eng. J. Med. 318:
1355-1362,

[111678]1988.

[111679]19788.

[111680]Edwards, C. Q.; Skolnick, M. H.; Kushner, J. P.: Hereditary
hemochromatosis:

[111681]contributions of genetic analyses. Prog. Hemat. 12:
43-71, 1981.

[111682]19789.

[111683]Edwards, J. H.: Allelic association in man.In: Eriksson, A.
W.;

[111684]Forsius, H. R.; Nevanlinna, H. R.; Workman, P. L.; Norio, R.
K. (eds.)

[111685]: Population Structure and Genetic Disorders: Seventh
Sigrid Juselius

[111686]Foundation Symposium: Mariehamn, Aland Islands, Fin-
land, August 1978.

[111687]London: Academic Press 1980. Pp. 239–255.

[111688]19790.

[111689]El Kahloun, A.; Chauvel, B.; Mauvieux, V.; Dorval, I.;
Jouanolle,

[111690]A.–M.; Gicquel, I.; Le Gall, J.–Y.; David, V.: Localization of
seven

[111691]new genes around the HLA–A locus. Hum. Molec. Genet.
2: 55–60, 1993.

[111692]19791.

[111693]Elzouki, A.–N. Y.; Hultcrantz, R.; Stal, P.; Befrits, R.; Eriks-
son,

[111694]S.: Increased PiZ gene frequency for alpha(1) antitrypsin in
patients

[111695]with genetic haemochromatosis. Gut 36: 922–926, 1995.

[111696]19792.

[111697]Eriksson, S.; Lindmark, B.; Hanik, L.: A Swedish family with

[111698]alpha-1-antitrypsin deficiency, haemochromatosis,
haemoglobinopathy

[111699]D and early death in liver cirrhosis. J. Hepatol. 2: 65–72,
1986.

[111700]19793.

[111701]Escobar, G. J.; Heyman, M. B.; Smith, W. B.; Thaler, M. M.:
Primary

[111702]hemochromatosis in childhood. Pediatrics 80: 549–554,
1987.

[111703]19794.

[111704]European Haemochromatosis Consortium: Polymorphism
in intron

[111705]4 of HFE does not compromise haemochromatosis muta-
tion results. (Letter) Nature

[111706]Genet. 23: 271 only, 1999.

[111707]19795.

[111708]Feder, J. N.; Penny, D. M.; Irrinki, A.; Lee, V. K.; Lebron, J.

- [111709]A.; Watson, N.; Tsuchihashi, Z.; Sigal, E.; Bjorkman, P. J.; Schatzman,
- [111710]R. C.: The hemochromatosis gene product complexes with the transferrin
- [111711]receptor and lowers its affinity for ligand binding. Proc. Nat. Acad.
- [111712]Sci. 95: 1472–1477, 1998.
- [111713]19796.
- [111714]Feder, J. N.; Tsuchihashi, Z.; Irrinki, A.; Lee, V. K.; Mapa,
- [111715]F. A.; Morikang, E.; Prass, C. E.; Starnes, S. M.; Wolff, R. K.; Parkkila,
- [111716]S.; Sly, W. S.; Schatzman, R. C. :J. Biol. Chem. 272: 14025–14028,
- [111717]1997.
- [111718]19797.
- [111719]Feller, E. R.; Pont, A.; Wands, J. R.; Carter, E. A.; Foster,
- [111720]G.; Kourides, I. A.; Isselbacher, K. J.: Familial hemochromatosis:
- [111721]physiologic studies in the precirrhotic stage of the disease. New
- [111722]Eng. J. Med. 296: 1422–1426, 1977.

[111723]19798.

[111724]Felts, J. H.; Nelson, J. R.; Herndon, C. N.; Spurr, C. L.:
Hemochromatosis

[111725]in two young sisters: case studies and a family survey.
Ann. Intern.

[111726]Med. 67: 117–123, 1967.

[111727]19799.

[111728]Feder, J. N.; Gnirke, A.; Thomas, W.; Tsuchihashi, Z.;
Ruddy,

[111729]D. A.; Basava, A.; Dormishian, F.; Domingo, R., Jr.; Ellis, M.
C.;

[111730]Fullan, A.; Hinton, L. M.; Jones, N. L.; Kimmel, B. E.; Kron-
mal, G.

[111731]S.; Lauer, P.; Lee, V. K.; Loeb, D. B.; Mapa, F. A.; McClel-
land, E.;

[111732]Meyer, N. C.; Mintier, G. A.; Moeller, N.; Moore, T.;
Morikang, E.;

[111733]Prass, C. E.; Quintana, L.; Starnes, S. M.; Schatzman, R. C.;
Brunke,

[111734]K. J.; Drayna, D. T.; Risch, N. J.; Bacon, B. R.; Wolff, R. K.: A

[111735]novel MHC class I-like gene is mutated in patients with
hereditary

[111736]haemochromatosis. Nature Genet. 13: 399–408, 1996.

[111737]19800.

[111738]Fleming, R. E.; Holden, C. C.; Tomatsu, S.; Waheed, A.;
Brunt,

[111739]E. M.; Britton, R. S.; Bacon, B. R.; Roopenian, D. C.; Sly, W.
S.

[111740] Mouse strain differences determine severity of iron accu-
mulation

[111741]in Hfe knockout model of hereditary hemochromatosis.
Proc. Nat. Acad.

[111742]Sci. 98: 2707–2711, 2001.

[111743]19801.

[111744]Fleming, R. E.; Migas, M. C.; Zhou, X. Y.; Jiang, J.; Britton,

[111745]R. S.; Brunt, E. M.; Tomatsu, S.; Waheed, A.; Bacon, B. R.;
Sly, W.

[111746]S.: Mechanism of increased iron absorption in murine
model of hereditary

[111747]hemochromatosis: increased duodenal expression of the
iron transporter

[111748]DMT1. Proc. Nat. Acad. Sci. 96: 3143–3148, 1999.

[111749]19802.

[111750]Goei, V. L.; Parimoo, S.; Capossela, A.; Chu, T. W.; Gruen, J.

[111751]R.: Isolation of novel non-HLA gene fragments from the hemochromatosis

[111752]region (6p21.3) by cDNA hybridization selection. Am. J. Hum. Genet. 54:

[111753]244-251, 1994.

[111754]19803.

[111755]Gomez, P. S.; Parks, S.; Ries, R.; Tran, T. C.; Gomez, P. F.;

[111756]Press, R. D.: Polymorphism in intron 4 of HFE does not compromise

[111757]haemochromatosis mutation results (Letter) Nature Genet. 23: 272

[111758]only, 1999.

[111759]19804.

[111760]Gordeuk, V.; Mukiibi, J.; Hasstedt, S.; Samowitz, W.; Edwards,

[111761]C. Q.; West, G.; Ndambire, S.; Emmamial, J.; Nkanza, N.; Chapanduka,

[111762]Z.; Randall, M.; Boone, P.; Romano, P.; Martell, R. W.; Yamashita,

[111763]T.; Effler, P.; Brittenham, G. :New Eng. J. Med. 326:

95–100,

[111764]1992.

[111765]19805.

[111766]Griffiths, W.; Cox, T.: Haemochromatosis: novel gene discovery

[111767]and the molecular pathophysiology of iron metabolism.
Hum. Molec.

[111768]Genet. 9: 2377–2382, 2000.

[111769]19806.

[111770]Grove, J.; Daly, A. K.; Burt, A. D.; Guzail, M.; James, O. F.

[111771]W.; Bassendine, M. F.; Day, C. P.: Heterozygotes for HFE mutations

[111772]have no increased risk of advanced alcoholic liver disease.
Gut 43:

[111773]262–266, 1998.

[111774]19807.

[111775]Halliday, J. W.; Cowlshaw, J. L.; Russo, A. M.; Powell, L. W.

[111776]: Serum-ferritin in diagnosis of haemochromatosis: a study of 43 families. Lancet II:

[111777]621-624, 1977.

[111778]19808.

[111779]Hardy, L.; Hansen, J. L.; Kushner, J. P.; Knisely, A. S.:

Neonatal

[111780]hemochromatosis: genetic analysis of transferrin-receptor, H-apoferritin,

[111781]and L-apoferritin loci and of the human leukocyte antigen class I

[111782]region. Am. J. Path. 137: 149-153, 1990.

[111783]19809.

[111784]Beattie, E. C.; Stellwagen, D.; Morishita, W.; Bresnahan, J. C.;

[111785]Ha, B. K.; Von Zastrow, M.; Beattie, M. S.; Malenka, R. C.: Control

[111786]of synaptic strength by glial TNF-alpha. Science 295: 2282-2285,

[111787]2002.

[111788]19810.

[111789]Beutler, B.; Krochin, N.; Milsark, I. W.; Luedke, C.; Cerami, A.

[111790]: Control of cachectin (tumor necrosis factor) synthesis:

mechanisms

[111791] of endotoxin resistance. Science 232: 977–980, 1986.

[111792] 19811.

[111793] Brenner, D. A.; O'Hara, M.; Angel, P.; Chojkier, M.; Karin, M.

[111794]; Prolonged activation of JUN and collagenase genes by tumour necrosis

[111795] factor- α . Nature 337: 661–663, 1989.

[111796] 19812.

[111797] Broudy, V. C.; Kaushansky, K.; Segal, G. M.; Harlan, J. M.; Adamson,

[111798] J. W.: Tumor necrosis factor type α stimulates human endothelial

[111799] cells to produce granulocyte/macrophage colony-stimulating factor. Proc.

[111800] Nat. Acad. Sci. 83: 7467–7471, 1986.

[111801] 19813.

[111802] Bruce, A. J.; Boling, W.; Kindy, M. S.; Peschon, J.; Kraemer, P.

[111803] J.; Carpenter, M. K.; Holtsberg, F. W.; Mattson, M. P.: Altered neuronal

- [111804]and microglial responses to excitotoxic and ischemic
brain injury
- [111805]in mice lacking TNF receptors. Nature Med. 2: 788–794,
1996.
- [111806]19814.
- [111807]Cabrera, M.; Shaw, M. A.; Sharples, C.; Williams, H.;
Castes, M.;
- [111808]Convit, J.; Blackwell, J. M.: Polymorphism in tumor necro-
sis factor
- [111809]genes associated with mucocutaneous leishmaniasis. J.
Exp. Med. 182:
- [111810]1259–1264, 1995.
- [111811]19815.
- [111812]Conway, D. J.; Holland, M. J.; Bailey, R. L.; Campbell, A. E.;
- [111813]Mahdi, O. S.; Jennings, R.; Mbena, E.; Mabey, D. C.: Scar-
ring trachoma
- [111814]is associated with polymorphism in the tumor necrosis
factor alpha
- [111815](TNF- α) gene promoter and with elevated TNF- α levels in tear
- [111816]fluid. Infect. Immun. 65: 1003–1006, 1997.

[111817]19816.

[111818]Davis, J. M.; Narachi, M. A.; Alton, N. K.; Arakawa, T.:
Structure

[111819]of human tumor necrosis factor alpha derived from re-
combinant DNA. Biochemistry 26:

[111820]1322-1326, 1987.

[111821]19817.

[111822]Escobar-Morreale, H. F.; Calvo, R. M.; Sancho, J.; San Mil-
lan,

[111823]J. L.: TNF-alpha and hyperandrogenism: a clinical, bio-
chemical, and

[111824]molecular genetic study. J. Clin. Endocr. Metab. 86:
3761-3767,

[111825]2001.

[111826]19818.

[111827]Gorman, J. D.; Sack, K. E.; Davis, J. C., Jr.: Treatment of
ankylosing

[111828]spondylitis by inhibition of tumor necrosis factor-alpha.
New Eng.

[111829]J. Med. 346: 1349-1356, 2002.

[111830]19819.

- [111831]Herrmann, S.-M.; Ricard, S.; Nicaud, V.; Mallet, C.;
Arveiler,
[111832]D.; Evans, A.; Ruidavets, J.-B.; Luc, G.; Bara, L.; Parra,
H.-J.;
- [111833]Poirier, O.; Cambien, F.: Polymorphisms of the tumour
necrosis factor- α
[111834]gene, coronary heart disease and obesity. *Europ. J. Clin.*
Invest. 28:
[111835]59-66, 1998.
[111836]19820.
- [111837]Inoko, H.; Trowsdale, J.: Linkage of TNF genes to the HLA-
B locus. *Nucleic*
[111838]*Acids Res.* 15: 8957-8962, 1987.
[111839]19821.
- [111840]Knight, J. C.; Udalova, I.; Hill, A. V. S.; Greenwood, B. M.;
- [111841]Peshu, N.; Marsh, K.; Kwiatkowski, D.: A polymorphism
that affects
[111842]OCT-1 binding to the TNF promoter region is associated
with severe
[111843]malaria. *Nature Genet.* 22: 145-150, 1999.
[111844]19822.

[111845]Koss, K.; Satsangi, J.; Fanning, G. C.; Welsh, K. I.; Jewell,

[111846]D. P.: Cytokine (TNF- α , LT- α , and IL-10) polymorphisms in

[111847]inflammatory bowel diseases and normal controls: differential effects

[111848]on production and allele frequencies. Genes Immun. 1: 185-190, 2000.

[111849]19823.

[111850]Marino, M. W.; Dunn, A.; Grail, D.; Inglese, M.; Noguchi, Y.;

[111851]Richards, E.; Jungbluth, A.; Wada, H.; Moore, M.; Williamson, B.;

[111852]Basu, S.; Old, L. J.: Characterization of tumor necrosis factor-deficient

[111853]mice. Proc. Nat. Acad. Sci. 94: 8093-8098, 1997.

[111854]19824.

[111855]McCusker, S. M.; Curran, M. D.; Dynan, K. B.; McCullagh, C. D.;

[111856]Urquhart, D. D.; Middleton, D.; Patterson, C. C.; McIlroy, S. P.;

[111857]Passmore, A. P.: Association between polymorphism in regulatory region

[111858] of gene encoding tumour necrosis factor- α and risk
of Alzheimer's

[111859] disease and vascular dementia: a case-control study.

Lancet 357:

[111860] 436-439, 2001.

[111861] 19825.

[111862] Remaley, A. T.; Rust, S.; Rosier, M.; Knapper, C.; Naudin,
L.;

[111863] Broccardo, C.; Peterson, K. M.; Koch, C.; Arnould, I.;
Prades, C.;

[111864] Duverger, N.; Funke, H.; Assman, G.; Dinger, M.; Dean, M.;
Chimini,

[111865] G.; Santamarina-Fojo, S.; Fredrickson, D. S.; Deneffe, P.
Brewer,

[111866] H. B., Jr.: Human ATP-binding cassette transporter 1
(ABC1): genomic

[111867] organization and identification of the genetic defect in the
original

[111868] Tangier disease kindred. Proc. Nat. Acad. Sci. 96:
12685-12690,

[111869] 1999.

[111870] 19826.

- [111871]Rust, S.; Rosier, M.; Funke, H.; Real, J.; Amoura, Z.; Piette,
[111872]J.-C.; Deleuze, J.-F.; Brewer, H. B.; Duverger, N.; Deneffe,
P.; Assmann,
[111873]G.: Tangier disease is caused by mutations in the gene
encoding ATP-binding
[111874]cassette transporter 1. *Nature Genet.* 22: 352–355, 1999.
[111875]19827.
[111876]Young, S. G.; Fielding, C. J.: The ABCs of cholesterol ef-
flux. *Nature*
[111877]*Genet.* 22: 316–318, 1999.
[111878]19828.
[111879]Garcia-Higuera, I.; Taniguchi, T.; Ganesan, S.; Meyn, M. S.;
Timmers,
[111880]C.; Hejna, J.; Grompe, M.; D'Andrea, A. D.: Interaction of
the Fanconi
[111881]anemia proteins and BRCA1 in a common pathway. *Molec.*
Cell 7: 249–262,
[111882]2001.
[111883]19829.
[111884]Mudd, S. H.; Levy, H. L.; Abeles, R. H.: A derangement in
B12

[111885]metabolism leading to homocystinemia, cystathioninemia and methylmalonic

[111886]aciduria. *Biochem. Biophys. Res. Commun.* 35: 121–126, 1969.

[111887]19830.

[111888]Mudd, S. H.; Levy, H. L.; Skovby, F.: Disorders of transsulfuration. In:

[111889]Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds.): *The Metabolic and Molecular Bases of Inherited Disease*. Vol. 1. New

[111891]York: McGraw–Hill (7th ed.): 1995. Pp. 1279–1327.

[111892]19831.

[111893]Mudd, S. H.; Skovby, F.; Levy, H. L.; Pettigrew, K. D.; Wilcken,

[111894]B.; Pyeritz, R. E.; Andria, G.; Boers, G. H. J.; Bromberg, I. L.;

[111895]Cerone, R.; Fowler, B.; Grobe, H.; Schmidt, H.; Schweitzer, L.: *The*

[111896]*natural history of homocystinuria due to cystathionine beta-synthase*

[111897]*deficiency*. *Am. J. Hum. Genet.* 37: 1–31, 1985.

[111898]19832.

- [111899] Munke, M.; Kraus, J.; Watkins, P.; Tanzi, R.; Gusella, J.;
Millington
- [111900] Ward, A.; Watson, M.; Francke, U.: Homocystinuria gene
on human chromosome
- [111901] 21 mapped with cloned cystathionine beta-synthase
probe and in situ
- [111902] hybridization of other chromosome 21 probes. (Abstract)
Cytogenet.
- [111903] Cell Genet. 40: 706-707, 1985.
- [111904] 19833.
- [111905] Munke, M.; Kraus, J. P.; Ohura, T.; Francke, U.: The gene
for
- [111906] cystathionine beta-synthase (CBS) maps to the subtelom-
eric region
- [111907] on human chromosome 21q and to proximal mouse chro-
mosome 17. Am.
- [111908] J. Hum. Genet. 42: 550-559, 1988.
- [111909] 19834.
- [111910] Munnich, A.; Saudubray, J.-M.; Dautzenberg, M.-D.; Parvy,
P.;
- [111911] Ogier, H.; Girot, R.; Manigne, P.; Frezal, J.: Diet-responsive
proconvertin

[111912](factor VII) deficiency in homocystinuria. J. Pediat. 102:
730-734,

[111913]1983.

[111914]19835.

[111915]Nishinaga, M.; Ozawa, T.; Shimada, K.: Homocysteine, a
thrombogenic

[111916]agent, suppresses anticoagulant heparan sulfate expres-
sion in cultured

[111917]porcine aortic endothelial cells. J. Clin. Invest. 92:
1381-1386,

[111918]1993.

[111919]19836.

[111920]Nugent, A.; Hadden, D. R.; Carson, N. A. J.: Long-term
survival

[111921]of homocystinuria: the first case. Lancet 352: 624-625,
1998.

[111922]19837.

[111923]Perry, T. L.; Hansen, S.; Love, D. L.; Crawford, L. E.; Tis-
chler,

[111924]B.: Treatment of homocystinuria with a low-methionine
diet, supplemental

[111925]cystine and a methyl donor. Lancet II: 474–478, 1968.

[111926]19838.

[111927]Peterschmitt, M. J.; Simmons, J. R.; Levy, H. L.: Reduction
of

[111928]false negative results in screening of newborns for homo-
cystinuria. New

[111929]Eng. J. Med. 341: 1572–1576, 1999.

[111930]19839.

[111931]Pogribna, M.; Melnyk, S.; Pogribny, I.; Chango, A.; Yi, P.;
James,

[111932]S. J.: Homocysteine metabolism in children with Down
syndrome: in

[111933]vitro modulation. Am. J. Hum. Genet. 69: 88–95, 2001.

[111934]19840.

[111935]Ratnoff, O. D.: Activation of Hageman factor by L-
homocystine. Science 162:

[111936]1007–1009, 1968.

[111937]19841.

[111938]Reish, O.; Townsend, D.; Berry, S. A.; Tsai, M. Y.; King, R.
A.

- [111939]: Tyrosinase inhibition due to interaction of homo-
cyst(e)ine with
- [111940]copper: the mechanism for reversible hypopigmentation in
homocystinuria
- [111941]due to cystathionine beta-synthase deficiency. Am. J.
Hum. Genet. 57:
- [111942]127-132, 1995.
- [111943]19842.
- [111944]Rosenquist, T. H.; Ratashak, S. A.; Selhub, J.: Homocys-
teine
- [111945]induces congenital defects of the heart and neural tube:
effect of
- [111946]folic acid. Proc. Nat. Acad. Sci. 93: 15227-15232, 1996.
- [111947]19843.
- [111948]Saudubray, J. M.: Personal Communication. Paris, France
5/20/1997.
- [111949]19844.
- [111950]Schimke, R. N.; McKusick, V. A.; Huang, T.; Pollack, A. D.:
Homocystinuria:
- [111951]studies of 20 families with 38 affected members. J.A.M.A.
193: 711-719,

[111952]1965.

[111953]19845.

[111954]Schnyder, G.; Roffi, M.; Pin, R.; Flammer, Y.; Lange, H.;
Eberli,

[111955]F. R.; Meier, B.; Turi, Z. G.; Hess, O. M.: Decreased rate of
coronary

[111956]restenosis after lowering of plasma homocysteine levels.
New Eng.

[111957]J. Med. 345: 1593–1600, 2001.

[111958]19846.

[111959]Sebastio, G.: Personal Communication. Padua, Italy
3/20/1997.

[111960]19847.

[111961]Sebastio, G.; Sperandeo, M. P.; Panico, M.; de Franchis, R.;
Kraus,

[111962]J. P.; Andria, G.: The molecular basis of homocystinuria
due to cystathionine

[111963]beta-synthase deficiency in Italian families, and report of
four novel

[111964]mutations. Am. J. Hum. Genet. 56: 1324–1333, 1995.

[111965]19848.

[111966]Shan, X.; Dunbrack, R. L., Jr.; Christopher, S. A.; Kruger, W.
[111967]D.: Mutations in the regulatory domain of cystathionine
beta-synthase
[111968]can functionally suppress patient-derived mutations in
cis. Hum.
[111969]Molec. Genet. 10: 635–643, 2001.
[111970]19849.
[111971]Shan, X.; Kruger, W. D.: Correction of disease-causing CBS
mutations
[111972]in yeast. Nature Genet. 19: 91–93, 1998.
[111973]19850.
[111974]Taniguchi, T.; Garcia-Higuera, I.; Xu, B.; Andreassen, P. R.;
Gregory,
[111975]R. C.; Kim, S.-T.; Lane, W. S.; Kastan, M. B.; D'Andrea, A.
D.: Convergence
[111976]of the Fanconi anemia and ataxia telangiectasia signaling
pathways. Cell 109:
[111977]459–472, 2002.
[111978]19851.
[111979]Timmers, C.; Taniguchi, T.; Hejna, J.; Reifsteck, C.; Lucas,
L.;

- [111980] Bruun, D.; Thayer, M.; Cox, B.; Olson, S.; D'Andrea, A. D.;
Moses,
- [111981] R.; Grompe, M.: Positional cloning of a novel Fanconi ane-
mia gene,
- [111982] FANCD2. Molec. Cell 7: 241–248, 2001.
- [111983] 19852.
- [111984] Whitney, M.; Thayer, M.; Reifsteck, C.; Olson, S.; Smith, L.;
Jakobs,
- [111985] P. M.; Leach, R.; Naylor, S.; Joenje, H.; Grompe, M.: Micro-
cell mediated
- [111986] chromosome transfer maps the Fanconi anaemia group D
gene to chromosome
- [111987] 3p. Nature Genet. 11: 341–343, 1995.
- [111988] 19853.
- [111989] Hejna, J. A.; Timmers, C. D.; Reifsteck, C.; Bruun, D. A.;
Lucas,
- [111990] L. W.; Jakobs, P. M.; Toth–Fejel, S.; Unsworth, N.; Clemens,
S. L.;
- [111991] Garcia, D. K.; Naylor, S. L.; Thayer, M. J.; Olson, S. B.;
Grompe,
- [111992] M.; Moses, R. E.: Localization of the Fanconi anemia com-
plementation

- [111993]group D gene to a 200-kb region on chromosome 3p25.3.
Am. J. Hum.
- [111994]Genet. 66: 1540–1551, 2000.
- [111995]19854.
- [111996]Buchwald, M.: Complementation groups: one or more per
gene? Nature
- [111997]Genet. 11: 228–230, 1995.
- [111998]19855.
- [111999]Stambolian, D.; Ai, Y.; Sidjanin, D.; Nesburn, K.; Sathe, G.;
- [112000]Rosenberg, M.; Bergsma, D. J.: Cloning of the galactoki-
nase cDNA
- [112001]and identification of mutations in two families with
cataracts. Nature
- [112002]Genet. 10: 307–312, 1995.
- [112003]19856.
- [112004]Alano, A.; Almashanu, S.; Chinsky, J. M.; Costeas, P.;
- Blitzer,
- [112005]M. G.; Wulfsberg, E. A.; Cowan, T. M.: Molecular character-
ization
- [112006]of a unique patient with epimerase-deficiency galac-
tosaemia. J. Inherit.

[112007]Metab. Dis. 21: 341–350, 1998.

[112008]19857.

[112009]Daude, N.; Gallaher, T. K.; Zeschnigk, M.; Starzinski-Powitz, A.;

[112010]Petry, K. G.; Haworth, I. S.; Reichardt, J. K. V.: Molecular cloning,

[112011]characterization, and mapping of a full-length cDNA encoding human

[112012]UDP-galactose 4-prime-epimerase. Biochem. Molec. Med. 56: 1–7, 1995.

[112013]19858.

[112014]Henderson, M. J.; Holton, J. B.; MacFaul, R.: Further observations

[112015]in a case of uridine diphosphate galactose-4-epimerase deficiency

[112016]with a severe clinical presentation. J. Inherit. Metab. Dis. 6:

[112017]17–20, 1983.

[112018]19859.

[112019]Holton, J. B.; Gillett, M. G.; MacFaul, R.; Young, R.: Galactosaemia:

- [112020] a new severe variant due to uridine diphosphate galactose-4-epimerase
- [112021] deficiency. Arch. Dis. Child. 56: 885-887, 1981.
- [112022] 19860.
- [112023] Kingsley, D. M.; Kozarsky, K. F.; Hobbie, L.; Krieger, M.: Reversible
- [112024] defects in O-linked glycosylation and LDL receptor expression in a
- [112025] UDP-Gal/UDP-GalNAc 4-epimerase deficient mutant. Cell 44: 749-759,
- [112026] 1986.
- [112027] 19861.
- [112028] Maceratesi, P.; Daude, N.; Dallapiccola, B.; Novelli, G.; Allen,
- [112029] R.; Okano, Y.; Reichardt, J.: Human UDP-galactose 4-prime epimerase
- [112030] (GALE) gene and identification of five missense mutations in patients
- [112031] with epimerase-deficiency galactosemia. Molec. Genet. Metab. 63:
- [112032] 26-30, 1998.

[112033]19862.

[112034]Piller, F.; Hanlon, M. H.; Hill, R. L.: Co-purification and
characterization

[112035]of UDP-glucose 4-epimerase and UDP-
N-acetylglucosamine 4-epimerase

[112036]from porcine submaxillary glands. J. Biol. Chem. 258:
10774-10778,

[112037]1983.

[112038]19863.

[112039]Quimby, B. B.; Alano, A.; Almashanu, S.; DeSandro, A. M.;
Cowan,

[112040]T. M.; Fridovich-Keil, J. L.: Characterization of two muta-
tions associated

[112041]with epimerase-deficiency galactosemia, by use of a yeast
expression

[112042]system for human UDP-galactose-4-epimerase. Am. J.
Hum. Genet. 61:

[112043]590-598, 1997.

[112044]19864.

[112045]Dubiel, B.; Dabrowski, C.; Wetts, R.; Tanaka, K.: Comple-
mentation

[112046]studies of isovaleric acidemia and glutaric aciduria type II
using

[112047]cultured skin fibroblasts. J. Clin. Invest. 72: 1543–1552,
1983.

[112048]19865.

[112049]McKusick, V. A.: Ellis–van Creveld syndrome and the
Amish. Nature

[112050]Genet. 24: 203–204, 2000.

[112051]19866.

[112052]McKusick, V. A.; Egeland, J. A.; Eldridge, R.; Krusen, D. E.:

[112053]Dwarfism in the Amish. I. The Ellis–van Creveld syndrome.
Bull. Johns

[112054]Hopkins Hosp. 115: 306–336, 1964.

[112055]19867.

[112056]Coppola, G.; De Michele, G.; Cavalcanti, F.; Pianese, L.;
Perretti,

[112057]A.; Santoro, L.; Vita, G.; Toscano, A.; Amboni, M.;
Grimaldi, G.;

[112058]Salvatore, E.; Caruso, G.; Filla, A.: Why do some Friedre-
ich's ataxia

[112059]patients retain tendon reflexes? A clinical, neurophysio-

logical and

[112060]molecular study. J. Neurol. 246: 353–357, 1999.

[112061]19868.

[112062]Delatycki, M. B.; Knight, M.; Koenig, M.; Cossee, M.;
Williamson,

[112063]R.; Forrest, S. M.: G130V, a common FRDA point mutation,
appears

[112064]to have arisen from a common founder. Hum. Genet. 105:
343–346,

[112065]1999.

[112066]19869.

[112067]Delatycki, M. B.; Paris, D. B. B. P.; Gardner, R. J. M.; Nichol-
son,

[112068]G. A.; Nassif, N.; Storey, E.; MacMillan, J. C.; Collins, V.;
Williamson,

[112069]R.; Forrest, S. M.: Clinical and genetic study of Friedreich
ataxia

[112070]in an Australian population. Am. J. Med. Genet. 87:
168–174, 1999.

[112071]19870.

[112072]Durr, A.; Cossee, M.; Agid, Y.; Campuzano, V.; Mignard,

C.; Penet,

[112073]C.; Mandel, J.-L.; Brice, A.; Koenig, M.: Clinical and genetic abnormalities

[112074]in patients with Friedreich's ataxia. New Eng. J. Med. 335: 1169-1175,

[112075]1996.

[112076]19871.

[112077]Filla, A.; De Michele, G.; Cavalcanti, F.; Pianese, L.; Monticelli,

[112078]A.; Campanella, G.; Coccozza, S.: The relationship between trinucleotide

[112079](GAA) repeat length and clinical features in Friedreich ataxia. Am.

[112080]J. Hum. Genet. 59: 554-560, 1996.

[112081]19872.

[112082]Fujita, R.; Hanauer, A.; Vincent, A.; Mandel, J.-L.; Koenig, M.

[112083]: Physical mapping of two loci (D9S5 and D9S15) tightly linked to

[112084]Friedreich ataxia locus (FRDA) and identification of nearby CpG islands

[112085]by pulse-field gel electrophoresis. Genomics 10:

915–920, 1991.

[112086]19873.

[112087]Gray, J. V.; Johnson, K. J.: Waiting for frataxin. *Nature Genet.* 16:

[112088]323–325, 1997.

[112089]19874.

[112090]Kenney, R. T.; Malech, H. L.; Epstein, N. D.; Roberts, R. L.;
Leto,

[112091]T. L.: Characterization of the p67–phox gene: genomic
organization

[112092]and restriction fragment length polymorphism analysis for
prenatal

[112093]diagnosis in chronic granulomatous disease. *Blood* 82:
3739–3744,

[112094]1993.

[112095]19875.

[112096]Noack, D.; Rae, J.; Cross, A. R.; Munoz, J.; Salmen, S.;
Mendoza,

[112097]J. A.; Rossi, N.; Curnutte, J. T.; Heyworth, P. G.: Autosomal
recessive

[112098]chronic granulomatous disease caused by novel mutations

in NCF-2,

[112099]the gene encoding the p67-phox component of phagocyte NADPH oxidase. Hum.

[112100]Genet. 105: 460-467, 1999.

[112101]19876.

[112102]Nunoi, H.; Iwata, M.; Tatsuzawa, S.; Onoe, Y.; Shimizu, S.; Kanegasaki,

[112103]S.; Matsuda, I.: AG dinucleotide insertion in a patient with chronic

[112104]granulomatous disease lacking cytosolic 67-kD protein. Blood 86:

[112105]329-333, 1995.

[112106]19877.

[112107]Okamura, N.; Babior, B. M.; Mayo, L. A.; Peveri, P.; Smith, R.

[112108]M.; Curnutte, J. T.: The p67-phox cytosolic peptide of the respiratory

[112109]burst oxidase from human neutrophils: functional aspects. J. Clin.

[112110]Invest. 85: 1583-1587, 1990.

[112111]19878.

[112112]Patino, P. J.; Rae, J.; Noack, D.; Erickson, R.; Ding, J.; Garcia

[112113]de Olarte, D.; Curnutte, J. T.: Molecular characterization of autosomal

[112114]recessive chronic granulomatous disease caused by a defect of the

[112115]nicotinamide adenine dinucleotide phosphate (reduced form) oxidase

[112116]component p67-phox. Blood 94: 2505-2514, 1999.

[112117]19879.

[112118]Tanugi-Cholley, L. C.; Issartel, J.-P.; Lunardi, J.; Freycon,

[112119]F.; Morel, F.; Vignais, P. V.: A mutation located at the

5-prime

[112120]splice junction sequence of intron 3 in the p67-phox gene causes the

[112121]lack of p67-phox mRNA in a patient with chronic granulomatous disease. Blood 85:

[112122]242-249, 1995.

[112123]19880.

[112124]Forsius, H.; Eriksson, A.; Nuutila, A.; Vainio-Mattila, B.; Krause,

[112125]U.: A genetic study of three rare retinal disorders: dystrophia retinae

[112126]dysacusis syndrome, X-chromosomal retinoschisis and grouped pigments

[112127]of the retina. Birth Defects Orig. Art. Ser. VII(3): 83-98, 1971.

[112128]19881.

[112129]Hassin, G. B.: Amaurotic family idiocy: late infantile type (Bielschowsky)

[112130]with the clinical picture of decerebrate rigidity. Arch. Neurol.

[112131]Psychiat. 16: 708-727, 1926.

[112132]19882.

[112133]Jarvela, I.: Infantile neuronal ceroid lipofuscinosis (CLN1):

[112134]linkage disequilibrium in the Finnish population and evidence that

[112135]variant late infantile form (variant CLN2) represents a non-allelic

[112136]locus. Genomics 10: 333–337, 1991.

[112137]19883.

[112138]Liu, C.-G.; Sleat, D. E.; Donnelly, R. J.; Lobel, P.: Structural

[112139]organization and sequence of CLN2, the defective gene in classical

[112140]late infantile neuronal ceroid lipofuscinosis. Genomics 50: 206–212,

[112141]1998.

[112142]19884.

[112143]MacLeod, P. M.; Dolman, C. L.; Nickel, R. E.; Chang, E.; Nag,

[112144]S.; Zonana, J.; Silvey, K.: Prenatal diagnosis of neuronal ceroid-lipofuscinoses. Am.

[112145]J. Med. Genet. 22: 781–789, 1985.

[112146]19885.

- [112147] MacLeod, P. M.; Dolman, C. L.; Nickel, R. E.; Chang, E.; Zonana,
nana,
- [112148] J.; Silvey, K.: Prenatal diagnosis of neuronal ceroid lipofus-
cinosis
- [112149] (Letter) New Eng. J. Med. 310: 595 only, 1984.
- [112150] 19886.
- [112151] Seitelberger, F.; Vogel, G.; Stepan, H.: Späta infantile
amaurotische
- [112152] Idiotie. Arch. Psychiat. Nervenkr. 196: 154–190, 1957.
- [112153] 19887.
- [112154] Sharp, J.; Savukoski, M.; Wheeler, R. B.; Harris, J.; Jarvela,
Jarvela,
- [112155] J.; Peltonen, L.; Gardiner, M.; Williams, R.: Linkage analysis
of
- [112156] late-infantile neuronal ceroid-lipofuscinosis. Am. J. Med.
Genet. 57:
- [112157] 348–349, 1995.
- [112158] 19888.
- [112159] Sharp, J. D.; Wheeler, R. B.; Lake, B. D.; Savukoski, M.;
Jarvela,
- [112160] J. E.; Peltonen, L.; Gardiner, R. M.; Williams, R. E.: Loci for
classical

- [112161]and a variant late infantile neuronal ceroid lipofuscinosis map to
- [112162]chromosomes 11p15 and 15q21–23. Hum. Molec. Genet. 6: 591–595, 1997.
- [112163]19889.
- [112164]Sleat, D. E.; Donnelly, R. J.; Lackland, H.; Liu, C.–G.; Sohar,
- [112165]J.; Pullarkat, R. K.; Lobel, P.: Association of mutations in a lysosomal
- [112166]protein with classical late–infantile neuronal ceroid lipofuscinosis. Science 277:
- [112167]1802–1805, 1997.
- [112168]19890.
- [112169]Sleat, D. E.; Gin, R. M.; Sohar, I.; Wisniewski, K.; Sklower–Brooks,
- [112170]S.; Pullarkat, R. K.; Palmer, D. N.; Lerner, T. J.; Boustany, R.–M.;
- [112171]Uldall, P.; Siakotos, A. N.; Donnelly, R. J.; Lobel, P.: Mutational
- [112172]analysis of the defective protease in classic late–infantile neuronal
- [112173]ceroid lipofuscinosis, a neurodegenerative lysosomal storage disorder. Am.

[112174]J. Hum. Genet. 64: 1511–1523, 1999.

[112175]19891.

[112176]Sohar, I.; Sleat, D. E.; Jadot, M.; Lobel, P.: Biochemical
characterization

[112177]of a lysosomal protease deficient in classical late infantile
neuronal

[112178]ceroid lipofuscinosis (LINCL) and development of an en-
zyme-based assay

[112179]for diagnosis and exclusion of LINCL in human specimens
and animal

[112180]models. J. Neurochem. 73: 700–711, 1999.

[112181]19892.

[112182]Volk, B. W.; Wallace, B. J.; Schneck, L.; Saifer, A.: Late in-
fantile

[112183]amaurotic idiocy. Ultramicroscopic and histochemical
studies on a

[112184]case. Arch. Path. 78: 483–500, 1964.

[112185]19893.

[112186]Williams, R.; Vesa, J.; Jarvela, I.; McKay, T.; Mitchison, H.;

[112187]Hellsten, E.; Thompson, A.; Callen, D.; Sutherland, G.;
Luna-Battadano,

[112188]D.; Stallings, R.; Peltonen, L.; Gardiner, M.: Genetic heterogeneity
[112189]in neuronal ceroid lipofuscinosis (NCL): evidence that the
late-infantile
[112190]subtype (Jansky-Bielschowsky disease; CLN2) is not an allelic form
[112191]of the juvenile or infantile subtypes. Am. J. Hum. Genet.
53: 931-935,
[112192]1993.
[112193]19894.
[112194]Yan, W.; Boustany, R.-M. N.; Konradi, C.; Ozelius, L.;
Lerner,
[112195]T.; Trofatter, J. A.; Julier, C.; Breakefield, X. O.; Gusella, J.
[112196]F.; Haines, J. L.: Localization of juvenile, but not late-infantile,
[112197]neuronal ceroid lipofuscinosis on chromosome 16. Am. J.
Hum. Genet. 52:
[112198]89-95, 1993.
[112199]19895.
[112200]Zhong, N.; Wisniewski, K. E.; Hartikainen, J.; Ju, W.; Moroziewicz,

- [112201]D. N.; McLendon, L.; Sklower Brooks, S.; Brown, W. T.: Two common
- [112202]mutations in the CLN2 gene underlie late infantile neuronal ceroid
- [112203]lipofuscinosis. Clin. Genet. 54: 234–238, 1998.
- [112204]19896.
- [112205]Harnden, D. G.: Ataxia–telangiectasia syndrome: cytogenetic and
- [112206]cancer aspects.In: German, J.: Chromosomes and Cancer. New York:
- [112207]Wiley (pub.) 1974. Pp. 619–636.
- [112208]19897.
- [112209]Hartwell, L.: Defects in a cell cycle checkpoint may be responsible
- [112210]for the genomic instability of cancer cells. Cell 71: 543–546, 1992.
- [112211]19898.
- [112212]Hawley, R. S.; Friend, S. H.: Strange bedfellows in even stranger
- [112213]places: the role of ATM in meiotic cells, lymphocytes, tumors, and

[112214]its functional links to p53. Genes Dev. 10: 2383–2388, 1996.

[112215]19899.

[112216]Hecht, F.; Koler, R. D.; Rigas, D. A.; Dahnke, G. S.; Case, M.

[112217]P.; Tisdale, V.; Miller, R. W.: Leukemia and lymphocytes in ataxia–telangiectasia.

[112218](Letter) Lancet II: 1193, 1966.

[112219]19900.

[112220]Henderson, L.; Cole, H.; Arlett, C.; James, S. E.; Cole, J.; Lehmann,

[112221]A.; Rosenbloom, L.; Redmond, T.; Meller, S.: Diagnosis of ataxia–telangiectasia

[112222]by T–lymphocyte cloning assay. (Letter) Lancet II: 1242, 1985.

[112223]19901.

[112224]Hernandez, D.; McConville, C. M.; Stacey, M.; Woods, C. G.; Brown,

[112225]M. M.; Shutt, P.; Rysiecki, G.; Taylor, A. M. R.: A family showing

[112226]no evidence of linkage between the ataxia telangiectasia gene and

[112227]chromosome 11q22-23. J. Med. Genet. 30: 135-140,
1993.

[112228]19902.

[112229]Hoar, D. I.; Sargent, P.: Chemical mutagen hypersensitivity
in

[112230]ataxia-telangiectasia. Nature 261: 590-592, 1976.

[112231]19903.

[112232]Hodge, S. E.; Berkel, A. I.; Gatti, R. A.; Boder, E.; Spence,

[112233]M. A.: Ataxia-telangiectasia and xeroderma pigmentosum:
no evidence

[112234]of linkage to HLA. Tissue Antigens 15: 313-317, 1980.

[112235]19904.

[112236]Huang, P. C.; Sheridan, R. B., III: Genetic and biochemical
studies

[112237]with ataxia telangiectasia. Hum. Genet. 59: 1-9, 1981.

[112238]19905.

[112239]Humphreys, M. W.; Nevin, N. C.; Wooldridge, M. A. W.: Cy-
togenetic

[112240]investigations in a family with ataxia telangiectasia. Hum.
Genet. 83:

[112241]79–82, 1989.

[112242]19906.

[112243]Imai, T.: Personal Communication. Chiba, Japan
9/12/1996.

[112244]

[112245]100. Ishiguro, T.; Taketa, K.; Gatti, R. A.: Tissue of origin
of

[112246]elevated alpha-fetoprotein in ataxia-telangiectasia. Dis.
Markers 4:

[112247]293–297, 1986.

[112248]

[112249]101. Jaspers, N. G. J.; Bootsma, D.: Genetic heterogeneity
in ataxia-telangiectasia

[112250]studied by cell fusion. Proc. Nat. Acad. Sci. 79:
2641–2644, 1982.

[112251]

[112252]102. Jaspers, N. G. J.; Gatti, R. A.; Baan, C.; Linssen, P. C.
M.

[112253]L.; Bootsma, D.: Genetic complementation analysis of
ataxia telangiectasia

[112254]and Nijmegen breakage syndrome: a survey of 50 pa-

tients. Cytogenet.

[112255]Cell Genet. 49: 259–263, 1988.

[112256]

[112257]103. Jaspers, N. G. J.; Painter, R. B.; Paterson, M. C.; Kidson, C.;

[112258]Inoue, T.: Complementation analysis of ataxia–telangiectasia.In:

[112259]Gatti, R. A.; Swift, M.: Ataxia–telangiectasia: Genetics, Neuropathology

[112260]and Immunology of a Degenerative Disease of Childhood. New York:

[112261]Alan R. Liss (pub.) 1985. Pp. 147–162.

[112262]

[112263]104. Johnson, J. P.; Gatti, R. A.; Sears, T. S.; White, R. L.: Inverted

[112264]duplication of J(H) associated with chromosome 14 translocation and

[112265]T–cell leukemia in ataxia–telangiectasia. Am. J. Hum. Genet. 39:

[112266]787–796, 1986.

[112267]

- [112268]105. Johnson, J. P.; White, R. L.; Gatti, R. A.: Rearrange-
ment of
[112269]J(H) genes in a patient with ataxia telangiectasia, chromo-
some 14
[112270]translocation, and T-cell leukemia. (Abstract) Am. J. Hum.
Genet. 37:
[112271]A100, 1985.
[112272]
[112273]106. Jung, M.; Zhang, Y.; Lee, S.; Dritschilo, A.: Correction
of
[112274]radiation sensitivity in ataxia telangiectasia cells by a
truncated
[112275]I-kappa-B-alpha. Science 268: 1619-1621, 1995.
[112276]
[112277]107. Kapp, L. N.; Painter, R. B.; Yu, L.-C.; van Loon, N.;
Richard,
[112278]C. W., III; James, M. R.; Cox, D. R.; Murnane, J. P.: Cloning
of
[112279]a candidate gene for ataxia-telangiectasia group D. Am. J.
Hum. Genet. 51:
[112280]45-54, 1992.

[112281]

[112282]108. Kastan, K.: Clinical implications of basic research:
ataxia-telangiectasia--broad

[112283]implications for a rare disorder. New Eng. J. Med. 333:
662-663,

[112284]1995.

[112285]

[112286]109. Kastan, M. B.; Onyekwere, O.; Sidransky, D.; Vogel-
stein, B.;

[112287]Craig, R. W.: Participation of p53 protein in the cellular re-
sponse

[112288]to DNA damage. Cancer Res. 51: 6304-6311, 1991.

[112289]

[112290]110. Kastan, M. B.; Zhan, Q.; El-Deiry, W. S.; Carrier, F.;

[112291]Jacks,
T.; Walsh, W. V.; Plunkett, B. S.; Vogelstein, B.; Fornace, A.
J.,

[112292]Jr.: A mammalian cell cycle checkpoint pathway utilizing
p53 and

[112293]GADD45 is defective in ataxia-telangiectasia. Cell 71:
587-597,

[112294]1992.

[112295]

[112296]111. Kennaugh, A. A.; Butterworth, S. V.; Hollis, R.; Baer, R.; Rabbitts,

[112297]T. H.; Taylor, A. M. R.: The chromosome breakpoint at 14q32 in an

[112298]ataxia telangiectasia t(14;14) T cell clone is different from the

[112299]14q32 breakpoint in Burkitts and an inv(14) T cell lymphoma. Hum.

[112300]Genet. 73: 254-259, 1986.

[112301]

[112302]112. Khanna, K. K.; Keating, K. E.; Kozlov, S.; Scott, S.; Gatei,

[112303]M.; Hobson, K.; Taya, Y.; Gabrielli, B.; Chan, D.; Lees-Miller, S.

[112304]P.; Lavin, M. F.: ATM associates with and phosphorylates p53: mapping

[112305]the region of interaction. Nature Genet. 20: 398-400, 1998.

[112306]

[112307]113. Klein, C.; Wenning, G. K.; Quinn, N. P.; Marsden, C.
D.: Ataxia

[112308]without telangiectasia masquerading as benign hereditary
chorea. Mov.

[112309]Disord. 11: 217–220, 1996.

[112310]

[112311]114. Kojis, T. L.; Schreck, R. R.; Gatti, R. A.; Sparkes, R. S.:

[112312]Tissue specificity of chromosomal rearrangements in
ataxia–telangiectasia. Hum.

[112313]Genet. 83: 347–352, 1989.

[112314]

[112315]115. Komatsu, K.; Kodama, S.; Okumura, Y.; Koi, M.; Os-
himura, M.:

[112316]Restoration of radiation resistance in ataxia–telangiectasia
cells

[112317]by the introduction of normal human chromosome 11.
Mutat. Res. 235:

[112318]59–63, 1990.

[112319]

[112320]116. Komatsu, K.; Matsuura, S.; Tauchi, H.; Endo, S.; Ko-
dama, S.;

[112321]Smeets, D.; Weemaes, C.; Oshimura, M.: The gene for Ni-jmegen breakage

[112322]syndrome (V2) is not located on chromosome 11. (Letter)
Am. J. Hum.

[112323]Genet. 58: 885–888, 1996.

[112324]

[112325]117. Korein, J.; Steinman, P. A.; Senz, E. H.: Ataxia–telangiectasia:

[112326]report of a case and review of the literature. Arch. Neurol.
4:

[112327]272–280, 1961.

[112328]

[112329]118. Krishna Kumar, G.; Al Saadi, A.; Yang, S. S.; Mc-Caughey, R. S.

[112330]: Ataxia–telangiectasia and hepatocellular carcinoma. Am.
J. Med.

[112331]Sci. 278: 157–160, 1979.

[112332]

[112333]119. Laake, K.; Jansen, L.; Hahnemann, J. M.; Brondum–Nielsen, K.;

[112334]Lonnqvist, T.; Kaariainen, H.; Sankila, R.; Lahdesmaki, A.;

Hammarstrom,

[112335]L.; Yuen, J.; Tretli, S.; Heiberg, A.; Olsen, J. H.; Tucker, M.;
Kleinerman,

[112336]R.; Borresen-Dale, A.-L.: Characterization of ATM muta-
tions in 41

[112337]Nordic families with ataxia telangiectasia. Hum. Mutat. 16:
232-246,

[112338]2000.

[112339]

[112340]120. Laake, K.; Telatar, M.; Geitvik, G. A.; Hansen, R. O.;
Heiberg,

[112341]A.; Andresen, A. M.; Gatti, R.; Borresen-Dale, A.-L.: Identical
mutation

[112342]in 55% of the ATM alleles in 11 Norwegian AT families: evidence for

[112343]a founder effect. Europ. J. Hum. Genet. 6: 235-244, 1998.

[112344]

[112345]121. Lambert, C.; Schultz, R. A.; Smith, M.; Wagner-
McPherson, C.;

[112346]McDaniel, L. D.; Donlon, T.; Stanbridge, E. J.; Friedberg, E.
C.:

[112347]Functional complementation of ataxia-telangiectasia

group D (AT-D)

[112348]cells by microcell-mediated chromosome transfer and mapping of the

[112349]AT-D locus to the region 11q22-23. Proc. Nat. Acad. Sci. 88: 5907-5911,

[112350]1991.

[112351]

[112352]122. Levin, S.; Perlov, S.: Ataxia-telangiectasia in Israel, with

[112353]observations on its relationship to malignant disease. Israel J. rael J.

[112354]Med. Sci. 7: 1535-1541, 1971.

[112355]

[112356]123. Li, A.; Swift, M.: Mutations at the ataxia-telangiectasia locus

[112357]and clinical phenotypes of A-T patients. Am. J. Med. Genet. 92:

[112358]170-177, 2000.

[112359]

[112360]124. Li, S.; Ting, N. S. Y.; Zheng, L.; Chen, P.-L.; Ziv, Y.; Shiloh,

[112361]Y.; Lee, E. Y.–H. P.; Lee, W.–H.: Functional link of BRCA1 and ataxia

[112362]telangiectasia gene product in DNA damage response.
Nature 406:

[112363]210–215, 2000.

[112364]

[112365]125. Lim, D.–S.; Kim, S.–T.; Xu, B.; Maser, R. S.; Lin, J.;
Petrini,

[112366]J. H. J.; Kastan, M. B.: ATM phosphorylates p95/nbs1 in an
S-phase

[112367]checkpoint pathway. Nature 404: 613–617, 2000.

[112368]

[112369]126. Lim, D.–S.; Kirsch, D. G.; Canman, C. E.; Ahn, J.–H.;
Ziv, Y.;

[112370]Newman, L. S.; Darnell, R. B.; Shiloh, Y.; Kastan, M. B.:
ATM binds

[112371]to beta-adaptin in cytoplasmic vesicles. Proc. Nat. Acad.
Sci. 95:

[112372]10146–10151, 1998.

[112373]

[112374]127. Lipkowitz, S.; Garry, V. F.; Kirsch, I. R.: Interlocus V–J

recombination

[112375]measures genomic instability in agriculture workers at risk
for lymphoid

[112376]malignancies. Proc. Nat. Acad. Sci. 89: 5301–5305, 1992.

[112377]

[112378]128. Lipkowitz, S.; Stern, M.–H.; Kirsch, I. R.: Hybrid T cell
receptor

[112379]genes formed by interlocus recombination in normal and
ataxia–telangiectasia

[112380]lymphocytes. J. Exp. Med. 172: 409–418, 1990.

[112381]

[112382]129. Lisker, R.; Cobo, A.: Chromosome breakage in
ataxia–telangiectasia.

[112383](Letter) Lancet I: 618, 1970.

[112384]

[112385]130. Littlefield, L. G.; Colyer, S. P.; Joiner, E. E.; DuFrain, R.

[112386]J.; Frome, E.; Cohen, M. M.: Chromosomal radiation sensi-
tivity in

[112387]ataxia telangiectasia long–term lymphoblastoid cell lines.
Cytogenet.

[112388]Cell Genet. 31: 203–213, 1981.

[112389]

[112390]131. Llerena, J., Jr.; Murer-Orlando, M.; McGuire, M.; Zahed, L.;

[112391]Sheridan, R. J.; Berry, A. C.; Bobrow, M.: Spontaneous and induced

[112392]chromosome breakage in chorionic villus samples: a cytogenetic approach

[112393]to first trimester prenatal diagnosis of ataxia telangiectasia syndrome. J.

[112394]Med. Genet. 26: 174-178, 1989.

[112395]

[112396]132. Maserati, E.; Ottolini, A.; Veggiotti, P.; Lanzi, G.; Pasquali,

[112397]F.: Ataxia-without-telangiectasia in two sisters with rearrangements

[112398]of chromosomes 7 and 14. Clin. Genet. 34: 283-287, 1988.

[112399]

[112400]133. Matsuda, Y.; Imai, T.; Shiomi, T.; Saito, T.; Yamauchi, M.; Fukao,

[112401]T.; Akao, Y.; Seki, N.; Ito, H.; Hori, T.: Comparative genome mapping

[112402]of the ataxia-telangiectasia region in mouse, rat, and Syrian hamster. Genomics 34:

[112403]347-352, 1996.

[112404]

[112405]134. McCaw, B. K.; Hecht, F.; Harden, D. G.; Teplitz, R. L.:
Somatic

[112406]rearrangement of chromosome 14 in human lymphocytes.
Proc. Nat. Acad.

[112407]Sci. 72: 2071-2075, 1975.

[112408]

[112409]135. McConville, C. M.; Formstone, C. J.; Hernandez, D.;
Thick, J.;

[112410]Taylor, A. M.: Fine mapping of the chromosome 11q22-23
region using

[112411]PFGE, linkage and haplotype analysis; localization of the
gene for

[112412]ataxia telangiectasia to a 5cM region flanked by NCAM/
DRD2 and STMY/CJ52.75,

[112413]phi 2.22. Nucleic Acids Res. 18: 4335-4343, 1990.

[112414]

[112415]136. McConville, C. M.; Stankovic, T.; Byrd, P. J.; McGuire,

G. M.;

[112416] Yao, Q.-Y.; Lennox, G. G.; Taylor, A. M. R.: Mutations associated

[112417] with variant phenotypes in ataxia-telangiectasia. Am. J. Hum. Genet. 59:

[112418] 320-330, 1996.

[112419]

[112420] 137. McConville, C. M.; Woods, C. G.; Farrall, M.; Metcalfe, J. A.;

[112421] Taylor, A. M. R.: Analysis of 7 polymorphic markers at chromosome

[112422] 11q22-23 in 35 ataxia telangiectasia families; further evidence of

[112423] linkage. Hum. Genet. 85: 215-220, 1990.

[112424]

[112425] 138. McFarlin, D. E.; Strober, W.; Waldmann, T. A.: Ataxia-telangiectasia. Medicine 51:

[112426] 281-314, 1972.

[112427]

[112428] 139. McKusick, V. A.; Cross, H. E.: Ataxia-telangiectasia and Swiss-type

[112429] agammaglobulinemia. Two genetic disorders of the immune mechanism

[112430] in related Amish sibships. J.A.M.A. 195: 739–745, 1966.

[112431]

[112432] 140. Meyn, M. S.: High spontaneous intrachromosomal recombination

[112433] rates in ataxia–telangiectasia. Science 260: 1327–1330, 1993.

[112434]

[112435] 141. Miller, M. E.; Chatten, J.: Ovarian changes in ataxia–telangiectasia. Acta

[112436] Paediat. Scand. 56: 559–561, 1967.

[112437]

[112438] 142. Mohamed, R.; Singh, S. P.; Kumar, S.; Lavin, M. F.: A defect

[112439] in DNA topoisomerase II activity in ataxia–telangiectasia cells. Biochem.

[112440] Biophys. Res. Commun. 149: 233–238, 1987.

[112441]

[112442] 143. Morrell, D.; Chase, C. L.; Swift, M.: Cancers in 44 families

[112443]with ataxia-telangiectasia. Cancer Genet. Cytogenet. 50:
119-123,

[112444]1990.

[112445]

[112446]144. Morrow, D. M.; Tagle, D. A.; Shiloh, Y.; Collins, F. S.;
Hieter,

[112447]P.: TEL1, an *S. cerevisiae* homolog of the human gene mu-
tated in ataxia

[112448]telangiectasia, is functionally related to the yeast check-
point gene

[112449]MEC1. Cell 82: 831-840, 1995.

[112450]

[112451]145. Murnane, J. P.; Painter, R. B.: Complementation of the
effects

[112452]in DNA synthesis in irradiated and unirradiated ataxia-
telangiectasia

[112453]cells. Proc. Nat. Acad. Sci. 79: 1960-1963, 1982.

[112454]

[112455]146. Naito, T.; Matsuura, A.; Ishikawa, F.: Circular chro-
mosome formation

[112456]in a fission yeast mutant defective in two ATM homo-

logues. Nature

[112457]Genet. 20: 203–206, 1998.

[112458]

[112459]147. Nguyen, T. D.; Boyd, J. B.; Green, M. M.: Sensitivity of
Drosophila

[112460]mutants to chemical carcinogens. Mutat. Res. 63: 67–77,
1979.

[112461]

[112462]148. Oxelius, V.–A.; Berkel, A. I.; Hanson, L. A.: IgG2 defi-
ciency

[112463]in ataxia–telangiectasia. New Eng. J. Med. 306: 515–517,
1982.

[112464]

[112465]149. Oxford, J. M.; Harnden, D. G.; Parrington, J. M.; Del-
hanty, J.

[112466]D. A.: Specific chromosome aberrations in ataxia–
telangiectasia. J.

[112467]Med. Genet. 12: 251–262, 1975.

[112468]

[112469]150. Pagani, F.; Buratti, E.; Stuani, C.; Bendix, R.; Dork, T.;
Baralle,

[112470]F. E.: A new type of mutation causes a splicing defect in
ATM. Nature

[112471]Genet. 30: 426–429, 2002.

[112472]

[112473]151. Painter, R. B.; Cramer, P.; Howard, R.; Young, B. R.:
Two forms

[112474]of inhibition of DNA replicon initiation in human cells.In:
Harris,

[112475]C. C.; Cerutti, P. C.: Mechanisms of Chemical Carcinogen-
esis. New

[112476]York: Alan R. Liss (pub.) 1982. Pp. 383–386.

[112477]

[112478]152. Painter, R. B.; Young, B. R.: Radiosensitivity in ataxia-
telangiectasia:

[112479]a new explanation. Proc. Nat. Acad. Sci. 77: 7315–7317,
1980.

[112480]

[112481]153. Paterson, M. C.; Smith, B. P.; Knight, P. A.; Anderson,
A. K.

[112482]: Ataxia telangiectasia: an inherited human disease involv-
ing radiosensitivity,

[112483]malignancy and defective DNA repair.In: Castellani, A.
(ed.): Research

[112484]in Photobiology. New York: Plenum (pub.) 1977. Pp.
207–218.

[112485]

[112486]154. Paterson, M. C.; Smith, B. P.; Lohman, P. H. M.; An-
derson, A.

[112487]K.; Fishman, L.: Defective excision repair of gamma-
ray-damaged DNA

[112488]in human (ataxia-telangiectasia) fibroblasts. Nature 260:
444–447,

[112489]1976.

[112490]

[112491]155. Paterson, M. C.; Smith, P. J.: Ataxia telangiectasia: an
inherited

[112492]human disorder involving hypersensitivity to ionizing ra-
diation and

[112493]related DNA-damaging chemicals. Ann. Rev. Genet. 13:
291–318, 1979.

[112494]

[112495]156. Paulovich, A. G.; Hartwell, L. H.: A checkpoint regu-
lates the

[112496]rate of progression through S phase in *S. cerevisiae* in response to

[112497]DNA damage. *Cell* 82: 841–847, 1995.

[112498]

[112499]157. Pecker, I.; Avraham, K. B.; Gilbert, D. J.; Savitsky, K.; Rotman,

[112500]G.; Harnik, R.; Fukao, T.; Schrock, E.; Hirotsume, S.; Tagle, D. A.;

[112501]Collins, F. S.; Wynshaw-Boris, A.; Ried, T.; Copeland, N. G.; Jenkins,

[112502]N. A.; Shiloh, Y.; Ziv, Y.: Identification and chromosomal localization

[112503]of Atm, the mouse homolog of the ataxia-telangiectasia gene. *Genomics* 35:

[112504]39–45, 1996.

[112505]

[112506]158. Peterson, R. D. A.; Funkhouser, J. D.: Speculations on ataxia-telangiectasia:

[112507]defective regulation of the immunoglobulin gene superfamily. *Immun.*

[112508]*Today* 10: 313–315, 1989.

[112509]

[112510]159. Peterson, R. D. A.; Funkhouser, J. D.: Ataxia-
telangiectasia:

[112511]an important clue. (Editorial) New Eng. J. Med. 322:
124-125, 1990.

[112512]

[112513]160. Peterson, R. D. A.; Kelly, W. D.; Good, R. A.: Ataxia-
telangiectasia:

[112514]its association with a defective thymus, immunological-defi-
ciency

[112515]disease and malignancy. Lancet I: 1189-1193, 1964.

[112516]

[112517]161. Pippard, E. C.; Hall, A. J.; Barker, D. J. P.; Bridges, B.
A.

[112518]; Cancer in homozygotes and heterozygotes of ataxia-
telangiectasia

[112519]and xeroderma pigmentosum in Britain. Cancer Res. 48:
2929-2932,

[112520]1988.

[112521]

[112522]162. Rary, J. M.; Bender, M. A.; Kelly, T. E.: A 14/14

marker chromosome

[112523] lymphocyte clone in ataxia telangiectasis. J. Hered. 66:
33–35,

[112524] 1975.

[112525]

[112526] 163. Rary, J. M.; Bender, M. A.; Kelly, T. E.: Cytogenetic
studies

[112527] of ataxia–telangiectasia. (Abstract) Am. J. Hum. Genet. 26:
70,

[112528] 1974.

[112529]

[112530] 164. Reye, C.; Mosman, N. S. W.: Ataxia–telangiectasia.
Am. J. Dis.

[112531] Child. 99: 238–247, 1960.

[112532]

[112533] 165. Richkind, K. E.; Boder, E.; Teplitz, R. L.: Fetal proteins
in

[112534] ataxia–telangiectasia. J.A.M.A. 248: 1346–1347, 1982.

[112535]

[112536] 166. Rosen, F. S.; Harris, N. L.: Case records of the Mas–

sachusetts

[112537]General Hospital: a 30-year-old man with ataxia-telangiectasia and

[112538]dysphagia. New Eng. J. Med. 316: 91-100, 1987.

[112539]

[112540]167. Rosin, M. P.; Ochs, H. D.: In vivo chromosomal instability in

[112541]ataxia-telangiectasia homozygotes and heterozygotes. Hum. Genet. 74:

[112542]335-340, 1986.

[112543]

[112544]168. Rosin, M. P.; Ochs, H. D.; Gatti, R. A.; Boder, E.: Heterogeneity

[112545]of chromosomal breakage levels in epithelial tissue of ataxia-telangiectasia

[112546]homozygotes and heterozygotes. Hum. Genet. 83: 133-138, 1989.

[112547]

[112548]169. Russo, G.; Isobe, M.; Gatti, R.; Finan, J.; Batuman, O.; Huebner,

[112549]K.; Nowell, P. C.; Croce, C. M.: Molecular analysis of a

t(14;14)

[112550]translocation in leukemic T-cells of an ataxia telangiectasia patient. Proc.

[112551]Nat. Acad. Sci. 86: 602-606, 1989.

[112552]

[112553]170. Sanal, O.; Lange, E.; Telatar, M.; Sobel, E.; Salazar-Novak,

[112554]J.; Ersoy, F.; Morrison, A.; Concannon, P.; Tolun, A.; Gatti, R. A.

[112555]: Ataxia-telangiectasia: linkage analysis of chromosome 11q22-23 markers

[112556]in Turkish families. FASEB J. 6: 2848-2852, 1992.

[112557]

[112558]171. Sanal, O.; Wei, S.; Foroud, T.; Malhotra, U.; Concannon, P.;

[112559]Charmley, P.; Salser, W.; Lange, K.; Gatti, R. A.: Further mapping

[112560]of an ataxia-telangiectasia locus to the chromosome 11q23 region. Am.

[112561]J. Hum. Genet. 47: 860-866, 1990.

[112562]

[112563]172. Sandoval, N.; Platzer, M.; Rosenthal, A.; Dork, T.;
Bendix, R.;

[112564]Skawran, B.; Stuhmann, M.; Wegner, R.-D.; Sperling, K.;
Banin, S.;

[112565]Shiloh, Y.; Baumer, A.; Bernthaler, U.; Sennefelder, H.;
Brohm, M.;

[112566]Weber, B. H. F.; Schindler, D.: Characterization of ATM
gene mutations

[112567]in 66 ataxia telangiectasia families. Hum. Molec. Genet. 8:
69–79,

[112568]1999.

[112569]

[112570]173. Sasaki, T.; Tian, H.; Kukita, Y.; Inazuka, M.; Tahira,
T.; Imai,

[112571]T.; Yamauchi, M.; Saito, T.; Hori, T.; Hashimoto–Tamaoki,
T.; Komatsu,

[112572]K.; Nikaido, O.; Hayashi, K.: ATM mutations in patients
with ataxia

[112573]telangiectasia screened by a hierarchical strategy. Hum.
Mutat. 12:

[112574]186–195, 1998.

[112575]

[112576]174. Savitsky, K.; Bar-Shira, A.; Gilad, S.; Rotman, G.; Ziv, Y.;

[112577]Vanagaite, L.; Tagle, D. A.; Smith, S.; Uziel, T.; Sfez, S.; Ashkenazi,

[112578]M.; Pecker, I.; and 18 others: A single ataxia telangiectasia gene

[112579]with a product similar to PI-3 kinase. Science 268: 1749-1753, 1995.

[112580]

[112581]175. Savitsky, K.; Sfez, S.; Tagle, D. A.; Ziv, Y.; Sartiell, A.; Collins,

[112582]F. S.; Shiloh, Y.; Rotman, G.: The complete sequence of the coding

[112583]region of the ATM gene reveals similarity to cell cycle regulators

[112584]in different species. Hum. Molec. Genet. 4: 2025-2032, 1995.

[112585]

[112586]176. Saxon, A.; Stevens, R. H.; Golde, D. W.: Helper and suppressor

[112587]T-lymphocyte leukemia in ataxia-telangiectasia. New Eng. J. Med. 300:

[112588]700–704, 1979.

[112589]

[112590]177. Schaffner, C.; Idler, I.; Stilgenbauer, S.; Dohner, H.;
Lichter,

[112591]P.: Mantle cell lymphoma is characterized by inactivation
of the

[112592]ATM gene. Proc. Nat. Acad. Sci. 97: 2773–2778, 2000.

[112593]

[112594]178. Schaffner, C.; Stilgenbauer, S.; Rappold, G. A.;
Dohner, H.;

[112595]Lichter, P.: Somatic ATM mutations indicate a pathogenic
role of

[112596]ATM in B-cell chronic lymphocytic leukemia. Blood 94:
748–753, 1999.

[112597]

[112598]179. Schalch, D. S.; McFarlin, D. E.; Barlow, M. H.: An un-
usual form

[112599]of diabetes mellitus in ataxia–telangiectasia. New Eng. J.
Med. 282:

[112600]1396–1402, 1970.

[112601]

- [112602]180. Scheres, J. M. J. C.; Hustinx, T. W. J.; Weemaes, C. M. R.:
[112603]Chromosome 7 in ataxia-telangiectasia. J. Pediat. 97:
440-441, 1980.
[112604]
- [112605]181. Scott, S. P.; Bendix, R.; Chen, P.; Clark, R.; Dork, T.;
Lavin,
[112606]M. F.: Missense mutations but not allelic variants alter the
function
[112607]of ATM by dominant interference in patients with breast
cancer. Proc.
[112608]Nat. Acad. Sci. 99: 925-930, 2002.
[112609]
- [112610]182. Sedgwick, R. P.; Boder, E.: Ataxia-telangiectasia.In:
Vinken,
[112611]P. J.; Bruyn, G. W. (eds.): Handbook of Clinical Neurology.
Amsterdam:
[112612]North-Holland Publishing Co. (pub.) 14: 1972. Pp.
267-339.
[112613]
- [112614]183. Shaham, M.; Becker, Y.: The ataxia telangiectasia
clastogenic

[112615]factor is a low molecular weight peptide. Hum. Genet. 58:
422–424,
[112616]1981.
[112617]
[112618]184. Shiloh, Y.; Parshad, R.; Frydman, M.; Sanford, K. K.;
Portnoi,
[112619]S.; Ziv, Y.; Jones, G. M.: G(2) chromosomal radiosensitivity
in families
[112620]with ataxia–telangiectasia. Hum. Genet. 84: 15–18, 1989.
[112621]
[112622]185. Shultz, L. D.; Sweet, H. O.; Davisson, M. T.; Coman,
D. R.:
[112623]'Wasted,' a new mutant of the mouse with abnormalities
characteristic
[112624]of ataxia telangiectasia. Nature 297: 402–404, 1982.
[112625]
[112626]186. Shuster, J.; Hart, Z.; Stimson, C. W.; Brough, A. J.;
Poulik,
[112627]M. D.: Ataxia–telangiectasia with cerebellar tumor. Pedi-
atrics 37:
[112628]776–786, 1966.

[112629]

[112630]187. Sobel, E.; Lange, E.; Jaspers, N. G. J.; Chessa, L.;
Sanal, O.;

[112631]Shiloh, Y.; Taylor, A. M. R.; Weemaes, C. M. A.; Lange, K.;
Gatti,

[112632]R. A.: Ataxia-telangiectasia: linkage evidence for genetic
heterogeneity.

[112633](Letter) Am. J. Hum. Genet. 50: 1343-1348, 1992.

[112634]

[112635]188. Sourander, P.; Bonnevier, J. O.; Olsson, Y.: A case of
ataxia-telangiectasia

[112636]with lesions in the spinal cord. Acta Neurol. Scand. 42:
354-366,

[112637]1966.

[112638]

[112639]189. Spring, K.; Ahangari, F.; Scott, S. P.; Waring, P.; Pur-
die, D.

[112640]M.; Chen, P. C.; Hourigan, K.; Ramsay, J.; McKinnon, P. J.;
Swift,

[112641]M.; Lavin, M. F.: Mice heterozygous for mutation in Atm,
the gene

[112642]involved in ataxia-telangiectasia, have heightened susceptibility

[112643]to cancer. *Nature Genet.* 32: 185–190, 2002.

[112644]

[112645]190. Stankovic, T.; Kidd, A. M. J.; Sutcliffe, A.; McGuire, G. M.;

[112646]Robinson, P.; Weber, P.; Bedenham, T.; Bradwell, A. R.; Easton, D.

[112647]F.; Lennox, G. G.; Haites, N.; Byrd, P. J.; Taylor, A. M. R.: ATM

[112648]mutations and phenotypes in ataxia-telangiectasia families in the

[112649]British Isles: expression of mutant ATM and the risk of leukemia,

[112650]lymphoma, and breast cancer. *Am. J. Hum. Genet.* 62: 334–345, 1998.

[112651]

[112652]191. Stern, M.-H.; Zhang, F.; Griscelli, C.; Thomas, G.; Aurias, A.

[112653]: Molecular characterization of different ataxia telangiectasia T-cell

[112654]clones. I. A common breakpoint at the 14q11.2 band

splits the T-cell

[112655]receptor alpha-chain gene. Hum. Genet. 78: 33-36, 1988.

[112656]

[112657]192. Stern, M.-H.; Zhang, F.; Thomas, G.; Griscelli, C.; Aurias, A.

[112658]; Molecular characterization of ataxia telangiectasia T cell clones.

[112659]III. Mapping the 14q32.1 distal breakpoint. Hum. Genet. 81: 18-22,

[112660]1988.

[112661]

[112662]193. Stewart, G. S.; Maser, R. S.; Stankovic, T.; Bressan, D. A.;

[112663]Kaplan, M. I.; Jaspers, N. G. J.; Raams, A.; Byrd, P. J.; Petrini,

[112664]J. H. J.; Taylor, A. M. R.: The DNA double-strand break repair gene

[112665]hMRE11 is mutated in individuals with an ataxia-telangiectasia-like

[112666]disorder. Cell 99: 577-587, 1999.

[112667]

[112668]194. Stilgenbauer, S.; Schaffner, C.; Litterst, A.; Liebisch, P.;

[112669]Gilad, S.; Bar-Shira, A.; James, M. R.; Lichter, P.; Dohner, H.:

[112670]Biallelic mutations in the ATM gene in T-prolymphocytic leukemia. Nature

[112671]Med. 3: 1155–1159, 1997.

[112672]

[112673]195. Stilgenbauer, S.; Winkler, D.; Ott, G.; Schaffner, C.; Leupolt,

[112674]E.; Bentz, M.; Moller, P.; Muller-Hermelink, H.-K.; James, M. R.;

[112675]Lichter, P.; Dohner, H.: Molecular characterization of 11q deletions

[112676]points to a pathogenic role of the ATM gene in mantle cell lymphoma. Blood 94:

[112677]3262–3264, 1999.

[112678]

[112679]196. Sugimoto, T.; Kidowaki, T.; Sawada, T.; Ohtsuka-Urano, T.; Kusunoki,

[112680]T.: Ataxia-telangiectasia associated with non-T, non-B cell acute

[112681]lymphocytic leukemia. Acta Paediat. Scand. 71: 509–510,
1982.

[112682]

[112683]197. Swift, M.: Cancer risk counseling. (Letter) Science
210: 1074,

[112684]1980.

[112685]

[112686]198. Swift, M.; Morrell, D.; Cromartie, E.; Chamberlin, A.
R.; Skolnick,

[112687]M. H.; Bishop, D. T.: The incidence and gene frequency of
ataxia–telangiectasia

[112688]in the United States. Am. J. Hum. Genet. 39: 573–583,
1986.

[112689]

[112690]199. Swift, M.; Morrell, D.; Massey, R. B.; Chase, C. L.: In-
cidence

[112691]of cancer in 161 families affected by ataxia–telangiectasia.
New

[112692]Eng. J. Med. 325: 1831–1836, 1991.

[112693]

[112694]200. Swift, M.; Reitnauer, P. J.; Morrell, D.; Chase, C. L.:

Breast

[112695]and other cancers in families with ataxia–telangiectasia.

New Eng.

[112696]J. Med. 316: 1289–1294, 1987.

[112697]

[112698]201. Swift, M. R.; Sholman, L.; Perry, M.; Chase, C.: Malignant neoplasms

[112699]in the families of patients with ataxia–telangiectasia. Cancer Res. 36:

[112700]209–215, 1976.

[112701]

[112702]202. Tadjoeidin, M. K.; Fraser, F. C.: Heredity of ataxia–telangiectasia

[112703](Louis–Bar syndrome). Am. J. Dis. Child. 110: 64–68, 1965.

[112704]

[112705]203. Taniguchi, T.; Garcia–Higuera, I.; Xu, B.; Andreassen, P. R.;

[112706]Gregory, R. C.; Kim, S.–T.; Lane, W. S.; Kastan, M. B.; D'Andrea,

[112707]A. D.: Convergence of the Fanconi anemia and ataxia

telangiectasia

[112708]signaling pathways. Cell 109: 459–472, 2002.

[112709]

[112710]204. Taylor, A. M. R.; Flude, E.; Laher, B.; Stacey, M.; McKay, E.;

[112711]Watt, J.; Green, S. H.; Harding, A. E.: Variant forms of ataxia telangiectasia. J.

[112712]Med. Genet. 24: 669–677, 1987.

[112713]

[112714]205. Taylor, A. M. R.; Harnden, D. G.; Arlett, C. F.; Harcourt, S.

[112715]A.; Lehmann, A. R.; Stevens, S.; Bridges, B. A.: Ataxia–telangiectasia:

[112716]a human mutation with abnormal radiation sensitivity. Nature 258:

[112717]427–429, 1975.

[112718]

[112719]206. Taylor, A. M. R.; Metcalfe, J. A.; Oxford, J. M.; Harnden, D.

[112720]G.: Is chromatid–type damage in ataxia–telangiectasia after irradiation

[112721]at G(0) a consequence of defective repair? Nature 260:
441–443,

[112722]1976.

[112723]

[112724]207. Tchirkov, A.; Bay, J.–O.; Pernin, D.; Bignon, Y.–J. Rio,
P.;

[112725]Granchó, M.; Kwiatkowski, F.; Giollant, M.; Malet, P.; Ver-
relle, P.

[112726]: Detection of heterozygous carriers of the ataxia–
telangiectasia

[112727](ATM) gene by G(2) phase chromosomal radiosensitivity of
peripheral

[112728]blood lymphocytes. Hum. Genet. 101: 312–316, 1997.

[112729]

[112730]208. Telatar, M.; Teraoka, S.; Wang, Z.; Chun, H. H.; Liang,
T.; Castellvi–Bel,

[112731]S.; Udar, N.; Borresen–Dale, A.–L.; Chessa, L.; Berna-
towska–Matuskiewicz,

[112732]E.; Porras, O.; Watanabe, M.; Junker, A.; Concannon, P.;
Gatti, R.

[112733]A.: Ataxia–telangiectasia: identification and detection of
founder–effect

[112734] mutations in the ATM gene in ethnic populations. Am. J. Hum. Genet. 62:

[112735] 86–97, 1998.

[112736]

[112737] 209. Telatar, M.; Wang, Z.; Castellvi–Bel, S.; Tai, L.–Q.; Sheikhavandi,

[112738] S.; Regueiro, J. R.; Porras, O.; Gatti, R. A.: A model for ATM heterozygote

[112739] identification in a large population: four founder–effect ATM mutations

[112740] identify most of Costa Rican patients with ataxia telangiectasia. Molec.

[112741] Genet. Metab. 64: 36–43, 1998.

[112742]

[112743] 210. Teplitz, R. L.: Ataxia–telangiectasia. Arch. Neurol. 35: 553–554,

[112744] 1978.

[112745]

[112746] 211. Teraoka, S. N.; Telatar, M.; Becker–Catania, S.; Liang, T.; Onengut,

[112747] S.; Tolun, A.; Chessa, L.; Sanal, O.; Bernatowska, E.; Gatti,

R. A.;

[112748]Concannon, P.: Splicing defects in the ataxia-telangiectasia gene,

[112749]ATM: underlying mutations and consequences. Am. J. Hum. Genet. 64:

[112750]1617-1631, 1999.

[112751]

[112752]212. Thibaut, S.; Sass, U.; Khoury, A.; Simonart, J.-M.: Ataxia-telangiectasia

[112753]and necrobiosis lipoidica: an explainable association. Europ. J.

[112754]Derm. 4: 509-513, 1994.

[112755]

[112756]213. Toledano, S. R.; Lang, B. J.: Ataxia-telangiectasia and acute

[112757]lymphoblastic leukemia. Cancer 45: 1675-1678, 1980.

[112758]

[112759]214. Toyoshima, M.; Hara, T.; Zhang, H.; Yamamoto, T.; Akaboshi, S.;

[112760]Nanba, E.; Ohno, K.; Hori, N.; Sato, K.; Takeshita, K.: Ataxia-telangiectasia

[112761] without immunodeficiency: novel point mutations within
and adjacent

[112762] to the phosphatidylinositol 3-kinase-like domain. Am. J.
Med. Genet. 75:

[112763] 141-144, 1998.

[112764]

[112765] 215. Tse, A. G. D.; Barclay, A. N.; Watts, A.; Williams, A. F.:
A

[112766] glycopospholipid tail at the carboxyl terminus of the
Thy-1 glycoprotein

[112767] of neurons and thymocytes. Science 230: 1003-1008,
1985.

[112768]

[112769] 216. Uziel, T.; Savitsky, K.; Platzer, M.; Ziv, Y.; Helbitz, T.;
Nehls,

[112770] M.; Boehm, T.; Rosenthal, A.; Shiloh, Y.; Rotman, G.: Ge-
nomic organization

[112771] of the ATM gene. Genomics 33: 317-320, 1996.

[112772]

[112773] 217. van Belzen, M. J.; Hiel, J. A. P.; Weemaes, C. M. R.;
Gabreels,

[112774]F. J. M.; van Engelen, B. G. M.; Smeets, D. F. C. M.; van den Heuvel,

[112775]L. P. W. J.: A double missense mutation in the ATM gene of a Dutch

[112776]family with ataxia telangiectasia. Hum. Genet. 102: 187–191, 1998.

[112777]

[112778]218. Vincent, R. A., Jr.; Sheridan, R. B., III; Huang, P. C.: DNA

[112779]strand breakage repair in ataxia–telangiectasia fibroblast–like cells. Mutat.

[112780]Res. 33: 357–366, 1975.

[112781]

[112782]219. Vorechovsky, I.; Luo, L.; Dyer, M. J. S.; Catovsky, D.; Amlot,

[112783]P. L.; Yaxley, J. C.; Foroni, L.; Hammarstrom, L.; Webster, A. D.

[112784]B.; Yuille, M. A. R.: Clustering of missense mutations in the ataxia–telangiectasia

[112785]gene in a sporadic T–cell leukaemia. Nature Genet. 17: 96–99, 1997.

[112786]

[112787]220. Vorechovsky, I.; Luo, L.; Lindblom, A.; Negrini, M.; Webster,

[112788]A. D.; Croce, C. M.; Hammarstrom, L.: ATM mutations in cancer families. Cancer

[112789]Res. 56: 4130–4133, 1996.

[112790]

[112791]221. Waldmann, T. A.; McIntire, K. R.: Serum- α -pha-fetoprotein levels

[112792]in patients with ataxia-telangiectasia. Lancet II: 1112–1115, 1972.

[112793]

[112794]222. Waldmann, T. A.; Misiti, J.; Nelson, D. L.; Kraemer, K. H.:

[112795]Ataxia-telangiectasia: a multisystem hereditary disease with immunodeficiency,

[112796]impaired organ maturation, x-ray hypersensitivity, and a high incidence

[112797]of neoplasia. Ann. Intern. Med. 99: 367–379, 1983.

[112798]

[112799]223. Wang, Y.; Cortez, D.; Yazdi, P.; Neff, N.; Elledge, S. J.; Qin,

[112800]J.: BASC, a super complex of BRCA1-associated proteins involved in

[112801]the recognition and repair of aberrant DNA structures.

Genes Dev. 14:

[112802]927–939, 2000.

[112803]

[112804]224. Wang, Y. A.; Elson, A.; Leder, P.: Loss of p21 in-
creases sensitivity

[112805]to ionizing radiation and delays the onset of lymphoma in
atm-deficient

[112806]mice. Proc. Nat. Acad. Sci. 94: 14590–14595, 1997.

[112807]

[112808]225. Watanabe, A.; Hanazono, H.; Sogawa, H.; Takaya, H.:
Stomach

[112809]cancer in a 14-year-old-boy with ataxia-telangiectasia.
Tohoku J.

[112810]Exp. Med. 121: 127–131, 1977.

[112811]

[112812]226. Waterman, M. J. F.; Stavridi, E. S.; Waterman, J. L. F.;
Halazonetis,

[112813]T. D.: ATM-dependent activation of p53 involves dephos-

phorylation

[112814]and association with 14-3-3 proteins. Nature Genet. 19:
175-178,

[112815]1998.

[112816]

[112817]227. Watts, J. A.; Morley, M.; Burdick, J. T.; Fiori, J. L.;
Ewens,

[112818]W. J.; Spielman, R. S.; Cheung, V. G.: Gene expression
phenotype

[112819]in heterozygous carriers of ataxia telangiectasia. Am. J.
Hum. Genet. 71:

[112820]791-800, 2002.

[112821]

[112822]228. Weinstein, S.; Scottolini, A. G.; Loo, S. Y. T.; Caldwell,
P.

[112823]C.; Bhagavan, N. V.: Ataxia telangiectasia with hepatocel-
lular carcinoma

[112824]in a 15-year-old girl and studies of her kindred. Arch.
Path. Lab.

[112825]Med. 109: 1000-1004, 1985.

[112826]

[112827]229. Welshimer, K.; Swift, M.: Congenital malformations and developmental disabilities in ataxia-telangiectasia, Fanconi anemia, and xeroderma pigmentosum families. Am. J. Hum. Genet. 34: 781-793, 1982.

[112830]

[112831]230. Westphal, C. H.; Rowan, S.; Schmaltz, C.; Elson, A.; Fisher, D. E.; Leder, P.: atm and p53 cooperate in apoptosis and suppression

[112833]of tumorigenesis, but not in resistance to acute radiation toxicity. Nature

[112834]Genet. 16: 397-401, 1997.

[112835]

[112836]231. Woods, C. G.; Bunday, S. E.; Taylor, A. M. R.: Unusual features

[112837]in the inheritance of ataxia telangiectasia. Hum. Genet. 84: 555-562,

[112838]1990.

[112839]

[112840]232. Woods, C. G.; Taylor, A. M. R.: Ataxia telangiectasia
in the

[112841]British Isles: the clinical and laboratory features of 70 af-
fected

[112842]individuals. Quart. J. Med. 82: 169–179, 1992.

[112843]

[112844]233. Wooster, R.; Ford, D.; Mangion, J.; Ponder, B. A. J.;
Peto, J.;

[112845]Easton, D. F.; Stratton, M. R.: Absence of linkage to the
ataxia

[112846]telangiectasia locus in familial breast cancer. Hum. Genet.
92:

[112847]91–94, 1993.

[112848]

[112849]234. Worgul, B. V.; Smilenov, L.; Brenner, D. J.; Junk, A.;
Zhou,

[112850]W.; Hall, E. J.: Atm heterozygous mice are more sensitive
to radiation-induced

[112851]cataracts than are their wild-type counterparts. Proc. Nat.
Acad.

[112852]Sci. 99: 9836–9839, 2002.

[112853]

[112854]235. Wright, J.; Teraoka, S.; Onengut, S.; Tolun, A.; Gatti, R. A.;

[112855]Ochs, H. D.; Concannon, P.: A high frequency of distinct ATM gene

[112856]mutations in ataxia-telangiectasia. Am. J. Hum. Genet. 59: 839-846,

[112857]1996.

[112858]

[112859]236. Wu, X.; Ranganathan, V.; Weisman, D. S.; Heine, W. F.; Ciccone,

[112860]D. N.; O'Neill, T. B.; Crick, K. E.; Pierce, K. A.; Lane, W. S.; Rathbun,

[112861]G.; Livingston, D. M.; Weaver, D. T.: ATM phosphorylation of Nijmegen

[112862]breakage syndrome protein is required in a DNA damage response. Nature 405:

[112863]477-482, 2000.

[112864]

[112865]237. Xia, Y.-R.; Welch, C. L.; Warden, C. H.; Lange, E.; Fukao, T.;

[112866]Lusis, A. J.; Gatti, R. A.: Assignment of the mouse ataxia-telangiectasia

[112867]gene (Atm) to mouse chromosome 9. Mammalian Genome 7: 554-555, 1996.

[112868]

[112869]238. Xu, Y.; Ashley, T; Brainerd, E. E.; Bronson, R. T.; Meyn, M.

[112870]S.; Baltimore, D.: Targeted disruption of ATM leads to growth retardation,

[112871]chromosomal fragmentation during meiosis, immune defects, and thymic

[112872]lymphoma. Genes Dev. 10: 2411-2422, 1996.

[112873]

[112874]239. Xu, Y.; Baltimore, D.: Dual roles of ATM in the cellular response

[112875]to radiation and in cell growth control. Genes Dev. 10: 2401-2410,

[112876]1996.

[112877]

[112878]240. Ying, K. L.; Decoteau, W. E.: Cytogenetic anomalies in a patient

[112879]with ataxia, immune deficiency, and high alpha-fetoprotein in the

[112880]absence of telangiectasia. Cancer Genet. Cytogenet. 4: 311–317,

[112881]1981.

[112882]

[112883]241. Yount, W. J.: IgG2 deficiency and ataxia-telangiectasia. (Editorial) New

[112884]Eng. J. Med. 306: 541–543, 1982.

[112885]

[112886]242. Zadik, Z.; Levin, S.; Prager-Lewin, R.; Laron, Z.: Gonadal dysfunction

[112887]in patients with ataxia telangiectasia. Acta Paediat. Scand. 67:

[112888]477–479, 1978.

[112889]

[112890]243. Zakian, V. A.: ATM-related genes: what do they tell us about

[112891]functions of the human gene? Cell 82: 685–687, 1995.

[112892]

[112893]244. Zhang, F.; Stern, M.-H.; Thomas, G.; Aurias, A.:

Molecular characterization

[112894]of ataxia telangiectasia T cell clones. II. The clonal inv(14)

in

[112895]ataxia telangiectasia differs from the inv(14) in T cell lym-

phoma. Hum.

[112896]Genet. 78: 316-319, 1988.

[112897]

[112898]245. Zhang, N.; Chen, P.; Khanna, K. K.; Scott, S.; Gatei,

M.; Kozlov,

[112899]S.; Watters, D.; Spring, K.; Yen, T.; Lavin, M. F.: Isolation of

[112900]full-length ATM cDNA and correction of the ataxia-

telangiectasia cellular

[112901]phenotype. Proc. Nat. Acad. Sci. 94: 8021-8026, 1997.

[112902]

[112903]246. Zhao, S.; Weng, Y.-C.; Yuan, S.-S. F.; Lin, Y.-T.; Hsu,

H.-C.;

[112904]Lin, S.-C. J.; Gerbino, E.; Song, M.; Zdzienicka, M. Z.; Gatti,

R.

[112905]A.; Shay, J. W.; Ziv, Y.; Shiloh, Y.; Lee, E. Y.-H. P.: Func-

tional

[112906]link between ataxia–telangiectasia and Nijmegen breakage syndrome

[112907]gene products. Nature 405: 473–477, 2000.

[112908]

[112909]247. Ziv, Y.; Amiel, A.; Jaspers, N. G. J.; Berkel, A. I.; Shiloh,

[112910]Y.: Ataxia–telangiectasia: a variant with altered in vitro phenotype

[112911]of fibroblast cells. Mutat. Res. 210: 211–219, 1989.

[112912]

[112913]248. Ziv, Y.; Frydman, M.; Lange, E.; Zelnik, N.; Rotman, G.; Julier,

[112914]C.; Jaspers, N. G. J.; Dagan, Y.; Abeliovicz, D.; Dar, H.; Borochowitz,

[112915]Z.; Lathrop, M.; Gatti, R. A.; Shiloh, Y.: Ataxia–telangiectasia:

[112916]linkage analysis in highly inbred Arab and Druze families and differentiation

[112917]from an ataxia–microcephaly–cataract syndrome. Hum. Genet. 88: 619–626,

[112918]1992.

[112919]

[112920]249. Ziv, Y.; Rotman, G.; Frydman, M.; Dagan, J.; Cohen, T.; Foroud,

[112921]T.; Gatti, R. A.; Shiloh, Y.: The ATC (ataxia-telangiectasia complementation

[112922]group C) locus localizes to 11q22-q23. Genomics 9: 373-375, 1991.

[112923]19907.

[112924]Date, H.; Onodera, O.; Tanaka, H.; Iwabuchi, K.; Uekawa, K.; Igarashi,

[112925]S.; Koike, R.; Hiroi, T.; Yuasa, T.; Awaya, Y.; Sakai, T.; and 9

[112926]others: Early-onset ataxia with ocular motor apraxia and hypoalbuminemia

[112927]is caused by mutations in a new HIT superfamily gene. Nature Genet. 29:

[112928]184-188, 2001.

[112929]19908.

[112930]Moreira, M.-C.; Barbot, C.; Tachi, N.; Kozuka, N.; Uchida, E.;

[112931]Gibson, T.; Mendonca, P.; Costa, M.; Barros, J.; Yanagisawa, T.; Watanabe,

[112932]M.; Ikeda, Y.; Aoki, M.; Nagata, T.; Coutinho, P.; Sequeiros, J.;

[112933]Koenig, M.: The gene mutated in ataxia–oculomotor apraxia 1 encodes

[112934]the new HIT/Zn–finger protein aprataxin. *Nature Genet.* 29: 189–193,

[112935]2001.

[112936]19909.

[112937]Shelley, W. B.; Rawnsley, H. M.; Morrow, G., III: Pyridoxine–dependent

[112938]hair pigmentation in association with homocystinuria. *Arch. Derm.* 106:

[112939]228–230, 1972.

[112940]

[112941]100. Shih, V. E.; Efron, M. L.: Pyridoxine–unresponsive homocystinuria. *New*

[112942]*Eng. J. Med.* 283: 1206–1208, 1970.

[112943]

[112944]101. Shih, V. E.; Fringer, J. M.; Mandell, R.; Kraus, J. P.; Berry,

[112945]G. T.; Heidenreich, R. A.; Korson, M. S.; Levy, H. L.;

Ramesh, V.

[112946] A missense mutation (I278T) in the cystathionine beta-synthase gene

[112947] prevalent in pyridoxine-responsive homocystinuria and associated with

[112948] mild clinical phenotype. *Am. J. Hum. Genet.* 57: 34–39, 1995.

[112949]

[112950] 102. Shipman, R. T.; Townley, R. R. W.; Danks, D. M.: Homocystinuria,

[112951] Addisonian pernicious anaemia, and partial deletion of a G chromosome. *Lancet* II:

[112952] 693–694, 1969.

[112953]

[112954] 103. Skovby, F.: Homocystinuria: clinical, biochemical and genetic

[112955] aspects of cystathionine beta-synthase and its deficiency in man. *Acta*

[112956] *Paediat. Scand.* 321 (suppl.): 1–21, 1985.

[112957]

[112958] 104. Skovby, F.; Krassikoff, N.; Francke, U.: Assignment of

the gene

[112959]for cystathionine beta-synthase to human chromosome
21 in somatic

[112960]cell hybrids. Hum. Genet. 65: 291-294, 1984.

[112961]

[112962]105. Skovby, F.; Krassikoff, N.; Francke, U.: Assignment of
the gene

[112963]for cystathionine beta-synthase (CBS) to human chromo-
some 21 in somatic

[112964]cell hybrids. (Abstract) Cytogenet. Cell Genet. 37: 585
only, 1984.

[112965]

[112966]106. Skovby, F.; Kraus, J.; Redlich, C.; Rosenberg, L. E.:
Immunochemical

[112967]studies on cultured fibroblasts from patients with homo-
cystinuria

[112968]due to cystathionine beta-synthase deficiency. Am. J.
Hum. Genet. 34:

[112969]73-83, 1982.

[112970]

[112971]107. Skovby, F.; Kraus, J. P.; Rosenberg, L. E.: Homo-

cystinuria:

[112972]biogenesis of cystathionine beta-synthase subunits in
cultured fibroblasts

[112973]and in an in vitro translation system programmed with fi-
broblast messenger

[112974]RNA. Am. J. Hum. Genet. 36: 452-459, 1984.

[112975]

[112976]108. Spaeth, G. L.; Barber, G. W.: Prevalence of homo-
cystinuria among

[112977]the mentally retarded: evaluation of a specific screening
test. Pediatrics 40:

[112978]586-589, 1967.

[112979]

[112980]109. Sperandio, M. P.; de Franchis, R.; Andria, G.; Sebas-
tio, G.:

[112981]A 68-bp insertion found in a homocystinuric patient is a
common variant

[112982]and is skipped by alternative splicing of the cystathionine
beta-synthase

[112983]mRNA. (Letter) Am. J. Hum. Genet. 59: 1391-1393, 1996.

[112984]

- [112985]110. Sperandio, M. P.; Panico, M.; Pepe, A.; Candito, M.;
de Franchis,
[112986]R.; Kraus, J. P.; Andria, G.; Sebastio, G.: Molecular analysis
of
[112987]patients affected by homocystinuria due to cystathionine
beta-synthase
[112988]deficiency: report of a new mutation in exon 8 and a dele-
tion in intron
[112989]11. J. Inherit. Metab. Dis. 18: 211-214, 1995.
[112990]
[112991]111. Streeten, B. W.: The nature of the ocular zonule.
Trans. Am.
[112992]Ophthal. Soc. 80: 823-854, 1982.
[112993]
[112994]112. Stubbs, L.; Kraus, J.; Lehrach, H.: The alpha-
A-crystallin and
[112995]cystathionine beta-synthase genes are physically very
closely linked
[112996]in proximal mouse chromosome 17. Genomics 7:
284-288, 1990.
[112997]

[112998]113. Tada, K.; Yoshida, T.; Hirono, H.; Arakawa, T.: Homocystinuria:

[112999]amino acid pattern of the liver. Tohoku J. Exp. Med. 92: 325-332,

[113000]1967.

[113001]

[113002]114. Tsai, M. Y.; Bignell, M.; Schwichtenberg, K.; Hanson, N. Q.:

[113003]High prevalence of a mutation in the cystathionine beta-synthase gene. Am.

[113004]J. Hum. Genet. 59: 1262-1267, 1996.

[113005]

[113006]115. Uhlemann, E. R.; TenPas, J. H.; Lucky, A. W.; Schulman, J. D.;

[113007]Mudd, S. H.; Shulman, N. R.: Platelet survival and morphology in

[113008]homocystinuria due to cystathionine synthase deficiency. New Eng.

[113009]J. Med. 295: 1283-1286, 1976.

[113010]

[113011]116. Uhlendorf, B. W.; Mudd, S. H.: Cystathionine synthase

in tissue

[113012]culture derived from human skin: enzyme defect in homocystinuria. Science 160:

[113013]1007–1009, 1968.

[113014]

[113015]117. Visy, J. M.; Le Coz, P.; Chadeaux, B.; Fressinaud, C.; Woimant,

[113016]F.; Marquet, J.; Zittoun, J.; Visy, J.; Vallat, J. M.; Haguenau, M.

[113017]: Homocystinuria due to 5,10-methylenetetrahydrofolate reductase deficiency

[113018]revealed by stroke in adult siblings. Neurology 41: 1313–1315, 1991.

[113019]

[113020]118. Wadman, S. K.; Cats, B. P.; de Bree, P. K.: Sulfite oxidase

[113021]deficiency and the detection of urinary sulfite. (Letter) Europ.

[113022]J. Pediat. 141: 62–63, 1983.

[113023]

[113024]119. Watanabe, M.; Osada, J.; Aratani, Y.; Kluckman, K.;

Reddick,

[113025]R.; Malinow, M. R.; Maeda, N.: Mice deficient in cystathionine beta-synthase:

[113026]animal models for mild and severe homocyst(e)inemia.

Proc. Nat. Acad.

[113027]Sci. 92: 1585-1589, 1995.

[113028]

[113029]120. Wilcken, D. E.; Wilcken, B.: The pathogenesis of coronary artery

[113030]disease: a possible role for methionine metabolism. J.

Clin. Invest. 57:

[113031]1079-1082, 1976.

[113032]

[113033]121. Wilcken, D. E. L.; Dudman, N. P. B.; Tyrrell, P. A.: Homocystinuria

[113034]due to cystathionine beta-synthase deficiency--the effects of betaine

[113035]treatment in pyridoxine-responsive patients. Metabolism 34: 1115-1121,

[113036]1985.

[113037]

[113038]122. Wilcken, D. E. L.; Wilcken, B.; Dudman, N. P. B.; Tyrrell, P.

[113039]A.: Homocystinuria--the effects of betaine in the treatment of patients

[113040]not responsive to pyridoxine. New Eng. J. Med. 309: 448-453, 1983.

[113041]

[113042]123. Wong, P. W. K.; Schwarz, V.; Komrower, G. M.: The biosynthesis

[113043]of cystathionine in patients with homocystinuria. Pediat. Res. 2:

[113044]149-160, 1968.

[113045]

[113046]124. Yaghmai, R.; Kashani, A. H.; Geraghty, M. T.; Okoh, J.; Pomper,

[113047]M.; Tangerman, A.; Wagner, C.; Stabler, S. P.; Allen, R. H.; Mudd,

[113048]S. H.; Braverman, N.: Progressive cerebral edema associated with

[113049]high methionine levels and betaine therapy in a patient with cystathionine

[113050]beta-synthase (CBS) deficiency. Am. J. Med. Genet. 108:

57-63, 2002.

[113051]

[113052]125. Yap, S.; Naughten, E.: Homocystinuria due to cystathionine beta-synthase

[113053]deficiency in Ireland: 25 years' experience of a newborn screened

[113054]and treated population with reference to clinical outcome and biochemical

[113055]control. J. Inherit. Metab. Dis. 21: 738-747, 1998.

[113056]

[113057]126. Yap, S.; Rushe, H.; Howard, P. M.; Naughten, E. R.: The intellectual

[113058]abilities of early-treated individuals with pyridoxine-nonresponsive

[113059]homocystinuria due to cystathionine beta-synthase deficiency. J.

[113060]Inherit. Metab. Dis. 24: 437-447, 2001.

[113061]19910.

[113062]Burchell, A.; Jung, R. T.; Lang, C. C.; Bennet, W.; Shepherd, A.

[113063]N.: Diagnosis of type 1a and type 1c glycogen storage

diseases in

[113064]adults. Lancet I: 1059-1062, 1987.

[113065]19911.

[113066]Burchell, A.; Waddell, I. D.: Diagnosis of a novel glycogen storage

[113067]disease: type 1aSP. J. Inherit. Metab. Dis. 13: 247-249, 1990.

[113068]19912.

[113069]Chen, Y.-T.; Coleman, R. A.; Scheinman, J. I.; Kolbeck, P. C.;

[113070]Sidbury, J. B.: Renal disease in type I glycogen storage disease. New

[113071]Eng. J. Med. 318: 7-11, 1988.

[113072]19913.

[113073]Chen, Y.-T.; Cornblath, M.; Sidbury, J. B.: Cornstarch therapy

[113074]in type I glycogen-storage disease. New Eng. J. Med. 310: 171-175,

[113075]1984.

[113076]19914.

[113077]Chen, Y.-T.; Scheinman, J. I.; Park, H. K.; Coleman, R. A.;

Roe,

[113078]C. R.: Amelioration of proximal renal tubular dysfunction
in type

[113079]glycogen storage disease with dietary therapy. New Eng.
J. Med. 323:

[113080]590–593, 1990.

[113081]19915.

[113082]Chevalier–Porst, F.; Bozon, D.; Bonardot, A.–M.; Bruni, N.;
Mithieux,

[113083]G.; Mathieu, M.; Maire, I.: Mutation analysis in 24 French
patients

[113084]with glycogen storage disease type 1a. J. Med. Genet. 33:
358–360,

[113085]1996.

[113086]19916.

[113087]Chou, J. Y.: Personal Communication. Bethesda, Md.
11/4/1993.

[113088]19917.

[113089]Chou, J.Y.; Mansfield, B.C.: Molecular genetics of type 1
glycogen

[113090]storage diseases. Trends Endocr. Metab. 10: 104–113,

1999.

[113091]19918.

[113092]Cohen, J. L.; Vinik, A.; Faller, J.; Fox, I. H.: Hyperuricemia

[113093]in glycogen storage disease type I: contributions by hypoglycemia

[113094]and hyperglucagonemia to increased urate production. J.

Clin. Invest. 75:

[113095]251–257, 1985.

[113096]19919.

[113097]Cori, G. T.; Cori, C. F.: Glucose-6-phosphatase of the liver

[113098]in glycogen storage disease. J. Biol. Chem. 199: 661–667, 1952.

[113099]19920.

[113100]Emmett, M.; Narins, R. G.: Renal transplantation in type I glycogenosis:

[113101]failure to improve glucose metabolism. J.A.M.A. 239: 1642–1644,

[113102]1978.

[113103]19921.

[113104]Faivre, L.; Houssin, D.; Valayer, J.; Brouard, J.; Hadchouel,

[113105]M.; Bernard, O.: Long-term outcome of liver transplantation in patients

[113106]with glycogen storage disease type Ia. J. Inherit. Metab. Dis. 22:

[113107]723–732, 1999.

[113108]19922.

[113109]Fine, R. N.; Wilson, W. A.; Donnell, G. N.: Retinal changes in

[113110]glycogen storage disease type I. Am. J. Dis. Child. 115: 328–331,

[113111]1968.

[113112]19923.

[113113]Fraumeni, J. F.; Rosen, P. J.; Hull, E. W.; Barth, R. F.; Shapiro,

[113114]S. R.; O'Connor, J. F.: Hepatoblastoma in infant sisters. Cancer 24:

[113115]1086–1090, 1969.

[113116]19924.

[113117]Furukawa, N.; Kinugasa, A.; Inoue, F.; Imashuku, S.; Takamatsu,

[113118]T.; Sawada, T.: Type I glycogen storage disease with vaso-

constrictive

[113119]pulmonary hypertension. J. Inherit. Metab. Dis. 13:
102–107, 1990.

[113120]19925.

[113121]Gibson, Q. H.: The reduction of methaemoglobin in red
blood cells

[113122]and studies on the cause of idiopathic methaemoglobi-
naemia. Biochem.

[113123]J. 42: 13–23, 1948.

[113124]19926.

- [113125]Greene, H. L.; Slonim, A. E.; O'Neill, J. A., Jr.; Burr, I. M.
- [113126]: Continuous nocturnal intragastric feeding for management of type
- [113127]1 glycogen-storage disease. New Eng. J. Med. 294: 423-425, 1976.
- [113128]19927.
- [113129]Howell, R. R.: The glycogen storage diseases.In: Stanbury, J.
- [113130]B.; Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic Basis of
- [113131]Inherited Disease. New York: McGraw-Hill (pub.) (4th ed.): 1978.
- [113132]Pp. 137-159.
- [113133]19928.
- [113134]Howell, R. R.: The interrelationship of glycogen storage disease
- [113135]and gout. Arthritis Rheum. 8: 780-785, 1965.
- [113136]19929.
- [113137]Howell, R. R.; Stevenson, R. E.; Ben-Menachem, Y.; Phyliky, R.
- [113138]L.; Berry, D. H.: Hepatic adenomata with type I glycogen

storage

[113139]disease. J. Nucl. Med. 19: 354–358, 1978.

[113140]19930.

[113141]Ito, E.; Sato, Y.; Kawauchi, K.; Munakata, H.; Kamata, Y.;
Yodono,

[113142]H.; Yokoyama, M.: Type 1a glycogen storage disease with
hepatoblastoma

[113143]in siblings. Cancer 59: 1776–1780, 1987.

[113144]19931.

[113145]Kajihara, S.; Matsushashi, S.; Yamamoto, K.; Kido, K.; Tsuji,
K.;

[113146]Tanae, A.; Fujiyama, S.; Itoh, T.; Tanigawa, K.; Uchida, M.;
Setoguchi,

[113147]Y.; Motomura, M.; Mizuta, T.; Sakai, T.: Exon redefinition
by a point

[113148]mutation within exon 5 of the glucose–6–phosphatase
gene is the major

[113149]cause of glycogen storage disease type 1a in Japan. Am. J.
Hum. Genet. 57:

[113150]549–555, 1995.

[113151]19932.

- [113152] Kikuchi, M.; Hasegawa, K.; Handa, I.; Watabe, M.; Nari-
sawa, K.;
- [113153] Tada, K.: Chronic pancreatitis in a child with glycogen
storage disease
- [113154] type 1. *Europ. J. Pediat.* 150: 852–853, 1991.
- [113155] 19933.
- [113156] Lee, P. J.; Dixon, M. A.; Leonard, J. V.: Uncooked corn-
starch--efficacy
- [113157] in type I glycogenosis. *Arch. Dis. Child.* 74: 546–547,
1996.
- [113158] 19934.
- [113159] Lee, W.-J.; Lee, H.-M.; Chi, C.-S.; Shu, S.-G.; Lin, L.-Y.;
- Lin,
- [113160] W.-H.: Genetic analysis of the glucose-6-phosphatase
mutation of
- [113161] type 1a glycogen storage disease in a Chinese family. *Clin.*
Genet. 50:
- [113162] 206–211, 1996.
- [113163] 19935.
- [113164] Lei, K.-J.; Chen, H.; Pan, C.-J.; Ward, J. M.; Mosinger, B.,
Jr.;

[113165]Lee, E. J.; Westphal, H.; Mansfield, B. C.; Chou, J. Y.: Glucose-6-phosphatase

[113166]dependent substrate transport in the glycogen storage disease type-1a

[113167]mouse. *Nature Genet.* 13: 203-209, 1996.

[113168]19936.

[113169]Lei, K.-J.; Chen, Y.-T.; Chen, H.; Wong, L.-J. C.; Liu, J.-L.;

[113170]McConkie-Rosell, A.; Van Hove, J. L. K.; Ou, H. C.-Y.; Yeh, N. J.;

[113171]Pan, L. Y.; Chou, J. Y.: Genetic basis of glycogen storage disease

[113172]type 1a: prevalent mutations at the glucose-6-phosphatase locus. *Am.*

[113173]*J. Hum. Genet.* 57: 766-771, 1995.

[113174]19937.

[113175]Lei, K.-J.; Pan, C.-J.; Shelly, L. L.; Liu, J.-L.; Chou, J. Y.

[113176]: Identification of mutations in the gene for glucose-6-phosphatase,

[113177]the enzyme deficient in glycogen storage disease type 1A. *J. Clin.*

[113178]*Invest.* 93: 1994-1999, 1994.

[113179]19938.

[113180]Lei, K.-J.; Shelly, L. L.; Lin, B.; Sidbury, J. B.; Chen, Y.-T.;

[113181]Nordlie, R. C.; Chou, J. Y.: Mutations in the glucose-
6-phosphatase

[113182]gene are associated with glycogen storage disease types
1a and 1aSP

[113183]but not 1b and 1c. J. Clin. Invest. 95: 234-240, 1995.

[113184]19939.

[113185]Lei, K.-J.; Shelly, L. L.; Pan, C.-J.; Sidbury, J. B.; Chou, J.

[113186]Y.: Mutations in the glucose-6-phosphatase gene that
cause glycogen

[113187]storage disease type 1a. Science 262: 580-583, 1993.

[113188]19940.

[113189]Limmer, J.; Fleig, W. E.; Leupold, D.; Bittner, R.;
Ditschuneit,

[113190]H.; Beger, H.-G.: Hepatocellular carcinoma in type I glyco-
gen storage

[113191]disease. Hepatology 8: 531-537, 1988.

[113192]19941.

[113193]Malatack, J. J.; Finegold, D. N.; Iwatsuki, S.; Shaw, B. W.,

Jr.;

[113194]Gartner, J. C.; Zitelli, B. J.; Roe, T.; Starzl, T. E.: Liver trans-
plantation

[113195]for type I glycogen storage disease. Lancet I: 1073–1075,
1983.

[113196]19942.

[113197]Aksentijevich, I.; Gruberg, L.; Pras, E.; Balow, J. E., Jr.;
Kovo,

[113198]M.; Gazit, E.; Dean, M.; Pras, M.; Kastner, D. L.: Evidence
for linkage

[113199]of the gene causing familial Mediterranean fever to chro-
mosome 17q

[113200]in non-Ashkenazi Jewish families: second locus or type I
error? Hum.

[113201]Genet. 91: 527–534, 1993.

[113202]19943.

[113203]Aksentijevich, I.; Pras, E.; Gruberg, L.; Shen, Y.; Holman,
K.;

[113204]Helling, S.; Prosen, L.; Sutherland, G. R.; Richards, R. I.;
Ramsburg,

[113205]M.; Dean, M.; Pras, M.; Amos, C. I.; Kastner, D. L.: Refined
mapping

[113206]of the gene causing familial Mediterranean fever, by linkage and homozygosity

[113207]studies. Am. J. Hum. Genet. 53: 451-461, 1993.

[113208]19944.

[113209]Aksentijevich, I.; Pras, E.; Gruberg, L.; Shen, Y.; Richards, R.

[113210]I.; Prosen, L.; Helling, S.; Pras, M.; Kastner, D. L.: Linkage disequilibrium

[113211]in the familial Mediterranean fever candidate region.
(Abstract) Am.

[113212]J. Hum. Genet. 51 (suppl.): A181, 1992.

[113213]19945.

[113214]Aksentijevich, I.; Torosyan, Y.; Samuels, J.; Centola, M.; Pras,

[113215]E.; Chae, J. J.; Oddoux, C.; Wood, G.; Azzaro, M. P.; Palumbo, G.;

[113216]Giustolisi, R.; Pras, M.; Ostrer, H.; Kastner, D. L.: Mutation and

[113217]haplotype studies of familial Mediterranean fever reveal new ancestral

[113218]relationships and evidence for a high carrier frequency with reduced

[113219]penetrance in the Ashkenazi Jewish population. Am. J. Hum. Genet. 64:

[113220]949–962, 1999.

[113221]19946.

[113222]Armenian, H. K.: Enrollment bias and variation in clinical manifestations:

[113223]a review of consecutive cases of familial paroxysmal polyserositis. J.

[113224]Chronic Dis. 36: 209–212, 1983.

[113225]19947.

[113226]Armenian, H. K.; Khachadurian, A. K.: Familial paroxysmal polyserositis:

[113227]clinical and laboratory findings in 120 cases. Leban. Med. J. 26:

[113228]605–614, 1973.

[113229]19948.

[113230]Ayesh, S. K.; Ferne, M.; Flechner, I.; Babior, B. M.; Matzner,

[113231]Y.: Partial characterization of a C5a-inhibitor in peritoneal fluid. J.

[113232]Immun. 144: 3066–3070, 1990.

[113233]19949.

[113234]Babior, B. M.; Matzner, Y.: The familial Mediterranean fever

[113235]gene--cloned at last. New Eng. J. Med. 337: 1548-1549, 1997.

[113236]19950.

[113237]Barakat, M. H.; El-Khawad, A. O.; Gumaa, K. A.; El-Sobki, N. I.;

[113238]Fenech, F. F.: Metaraminol provocative test: a specific diagnostic

[113239]test for familial Mediterranean fever. Lancet I: 656-657, 1984.

[113240]19951.

[113241]Barakat, M. H.; El-Sobki, N. I.; El-Khawad, A. O.; Gumma, K. A.;

[113242]Fenech, F. F.: Diagnosing familial Mediterranean fever. (Letter) Lancet II:

[113243]41-42, 1984.

[113244]19952.

[113245]Barakat, M. H.; Karnik, A. M.; Majeed, H. W. A.; El-Sobki, N.

[113246]I.; Fenech, F. F.: Familial Mediterranean fever (recurrent hereditary

[113247]polyserositis) in Arabs--a study of 175 patients and review of the

[113248]literature. Quart. J. Med. 60: 837-847, 1986.

[113249]19953.

[113250]Barakat, M. H.; Mustafa, H. T.; Shakir, R. A.: Mollaret's meningitis:

[113251]a variant of recurrent hereditary polyserositis, both provoked by

[113252]metaraminol. Arch. Neurol. 45: 926-927, 1988.

[113253]19954.

[113254]Ben-Chetrit, E.; Scherrmann, J.-M.; Levy, M.: Colchicine in breast

[113255]milk of patients with familial Mediterranean fever. Arthritis Rheum. 39:

[113256]1213-1217, 1996.

[113257]19955.

[113258]Benson, M. D.; Skinner, M.; Cohen, A. S.: Amyloid deposition

[113259]in a renal transplant in familial Mediterranean fever. Ann.

Intern.

[113260]Med. 87: 31–34, 1977.

[113261]19956.

[113262]Bernot, A.; da Silva, C.; Petit, J.–L.; Cruaud, C.; Caloustian,

[113263]C.; Castet, V.; Ahmed–Arab, M.; Dross, C.; Dupont, M.;

Cattan, D.;

[113264]Smaoui, N.; Dode, C.; and 10 others: Non–founder mutations in the

[113265]MEFV gene establish this gene as the cause of familial Mediterranean

[113266]fever (FMF). Hum. Molec. Genet. 7: 1317–1325, 1998.

[113267]19957.

[113268]Booth, D. R.; Booth, S. E.; Gillmore, J. D.; Hawkins, P. N.; Pepys,

[113269]M. B.: SAA1 alleles as risk factors in reactive systemic AA amyloidosis. Amyloid 5:

[113270]262–265, 1998.

[113271]19958.

[113272]Brik, R.; Litmanovitz, D.; Berkowitz, D.; Shamir, R.; Rosenthal,

[113273]E.; Shinawi, M.; Gershoni–Baruch, R.: Incidence of familial

Mediterranean

[113274]fever (FMF) mutations among children of Mediterranean
extraction with

[113275]functional abdominal pain. J. Pediat. 138: 759–762, 2001.

[113276]19959.

[113277]Cattan, D.; Dervichian, M.; Courillon, A.; Nurit, Y.:

Metaraminol

[113278]provocation test for familial Mediterranean fever. (Letter)
Lancet I:

[113279]1130–1131, 1984.

[113280]19960.

[113281]Cattan, D.; Notarnicola, C.; Molinari, N.; Touitou, I.: In-
flammatory

[113282]bowel disease in non-Ashkenazi Jews with familial
Mediterranean fever. Lancet 355:

[113283]378–379, 2000.

[113284]19961.

[113285]Chae, J. J.; Centola, M.; Aksentijevich, I.; Dutra, A.; Tran,

[113286]M.; Wood, G.; Nagaraju, K.; Kingma, D. W.; Liu, P. P.; Kast-
ner, D.

[113287]L.: Isolation, genomic organization, and expression analy-

sis of the

[113288]mouse and rat homologs of MEFV, the gene for familial
Mediterranean

[113289]fever. Mammalian Genome 11: 428–435, 2000.

[113290]19962.

[113291]Chen, X.; Fischel–Ghodsian, N.; Cercek, A.; Hamon, M.;
Ogur, G.;

[113292]Lotan, R.; Danon, Y.; Shohat, M.: Assessment of pyrin
gene mutations

[113293]in Turks with familial Mediterranean fever (FMF). Hum.
Mutat. 11:

[113294]456–460, 1998.

[113295]19963.

[113296]Daniels, M.; Shohat, T.; Brenner–Ullman, A.; Shohat, M.:
Familial

[113297]Mediterranean fever: high gene frequency among the
non–Ashkenazic

[113298]and Ashkenazic Jewish populations in Israel. Am. J. Med.
Genet. 55:

[113299]311–314, 1995.

[113300]19964.

- [113301] Dewalle, M.; Domingo, C.; Rozenbaum, M.; Ben-Chetrit, E.; Cattan,
- [113302] D.; Bernot, A.; Dross, C.; Dupont, M.; Notarnicola, C.; Levy, M.;
- [113303] Rosner, I.; Demaille, J.; Touitou, I.: Phenotype-genotype correlation
- [113304] in Jewish patients suffering from familial Mediterranean fever (FMF). Europ.
- [113305] J. Hum. Genet. 6: 95-97, 1998.
- [113306] 19965.
- [113307] Ito, K.; Suzuki, H.; Hirohashi, T.; Kume, K.; Shimizu, T.; Sugiyama,
- [113308] Y.: Molecular cloning of canalicular multispecific organic anion
- [113309] transporter defective in EHBR. Am. J. Physiol. 272: G16-G22, 1997.
- [113310] 19966.
- [113311] van Kuijk, M. A.; Kool, M.; Merks, G. F. M.; Geurts van Kessel,
- [113312] A.; Bindels, R. J. M.; Deen, P. M. T.; van Os, C. H.: Assignment
- [113313] of the canalicular multispecific organic anion transporter

gene (CMOAT)

[113314] to human chromosome 10q24 and mouse chromosome
19D2 by fluorescent

[113315] in situ hybridization. Cytogenet. Cell Genet. 77: 285–287,
1997.

[113316] 19967.

[113317] Wada, M.; Toh, S.; Taniguchi, K.; Nakamura, T.; Uchiumi,
T.; Kohno,

[113318] K.; Yoshida, I.; Kimura, A.; Sakisaka, S.; Adachi, Y.;
Kuвано, M.

[113319]: Mutations in the canalicular multispecific organic anion
transporter

[113320] (cMOAT) gene, a novel ABC transporter, in patients with
hyperbilirubinemia

[113321] II/Dubin–Johnson syndrome. Hum. Molec. Genet. 7:
203–207, 1998.

[113322] 19968.

[113323] Spranger, J. W.: Personal Communication. Kiel, Germany
1974.

[113324] 19969.

[113325] Victoria, T.; Rafi, M. A.; Wenger, D. A.: Cloning of the ca-
nine

[113326]GALC cDNA and identification of the mutation causing globoid cell

[113327]leukodystrophy in West Highland White and Cairn terriers.

Genomics 33:

[113328]457–462, 1996.

[113329]19970.

[113330]Wenger, D. A.; Rafi, M. A.; Luzi, P.: Molecular genetics of Krabbe

[113331]disease (globoid cell leukodystrophy): diagnostic and clinical implications. Hum.

[113332]Mutat. 10: 268–279, 1997.

[113333]19971.

[113334]Zlotogora, J.; Chakraborty, S.; Knowlton, R. G.; Wenger, D. A.

[113335]; Krabbe disease locus mapped to chromosome 14 by genetic linkage. Am.

[113336]J. Hum. Genet. 47: 37–44, 1990.

[113337]19972.

[113338]Zlotogora, J.; Regev, R.; Zeigler, M.; Iancu, T. C.; Bach, G.

[113339]; Krabbe disease: increased incidence in a highly inbred community. Am.

[113340]J. Med. Genet. 21: 765–770, 1985.

[113341]19973.

[113342]Rushton, A. R.; Dawson, G.: Genetic linkage studies of the
human

[113343]glycosphingolipid beta-galactosidases. Biochem. Genet.
15: 1071–1082,

[113344]1977.

[113345]19974.

[113346]Balciunaite, G.; Keller, M. P.; Balciunaite, E.; Piali, L.;
Zuklys,

[113347]S.; Mathieu, Y. D.; Gill, J.; Boyd, R.; Sussman, D. J.; Hollan-
der,

[113348]G. A.: Wnt glycoproteins regulate the expression of
FoxN1, the gene

[113349]defective in nude mice. Nature Immun. 15Oct, 2002. Note:
Advance

[113350]Electronic Publication.

[113351]19975.

[113352]Frank, J.; Pignata, C.; Panteleyev, A. A.; Prowse, D. M.;
Baden,

[113353]H.; Weiner, L.; Gaetaniello, L.; Ahmad, W.; Pozzi, N.; Caer-

halmi-Friedman,

[113354]P. B.; Aita, V. M.; Uyttendaele, H.; Gordon, D.; Ott, J.; Brissette,

[113355]J. L.; Christiano, A. M.: Exposing the human nude phenotype. *Nature* 398:

[113356]473–474, 1999.

[113357]19976.

[113358]Nehls, M.; Pfeifer, D.; Schorpp, M.; Hedrich, H.; Boehm, T.:
New

[113359]member of the winged-helix protein family disrupted in
mouse and rat

[113360]nude mutations. *Nature* 372: 103–107, 1994.

[113361]19977.

[113362]Pignata, C.; Fiore, M.; Guzzetta, V.; Castaldo, A.; Sebastio,
G.;

[113363]Porta, F.; Guarino, A.: Congenital alopecia and nail dystrophy associated

[113364]with severe functional T-cell immunodeficiency in two
sibs. *Am. J.*

[113365]*Med. Genet.* 65: 167–170, 1996.

[113366]19978.

[113367]Schorpp, M.; Hofmann, M.; Dear, T. N.; Boehm, T.: Characterization
[113368]of mouse and human nude genes. Immunogenetics 46:
509–515, 1997.
[113369]19979.

[113370]Segre, J. A.; Nemhauser, J. L.; Taylor, B. A.; Nadeau, J. H.;
Lander,
[113371]E. S.: Positional cloning of the nude locus: genetic, physiological, and
[113372]transcription maps of the region and mutations in the
mouse and rat. Genomics 28:
[113373]549–559, 1995.
[113374]19980.

[113375]Delpire, E.; Lu, J.; England, R.; Dull, C.; Thorne, T.: Deafness
[113376]and imbalance associated with inactivation of the secretory Na–K–2Cl
[113377]co-transporter. Nature Genet. 22: 192–195, 1999.
[113378]19981.

[113379]Evans, R. L.; Park, K.; Turner, R. J.; Watson, G. E.; Nguyen, H.–V.;

- [113380]Dennett, M. R.; Hand, A. R.; Flagella, M.; Shull, G. E.;
Melvin, J.
- [113381]E.: Severe impairment of salivation in Na⁺/K⁺/2Cl⁻ co-
transporter
- [113382](NKCC1)-deficient mice. J. Biol. Chem. 275:
26720-26726, 2000.
- [113383]19982.
- [113384]Payne, J. A.; Xu, J.-C.; Haas, M.; Lytle, C. Y.; Ward, D.; For-
bush,
- [113385]B., III: Primary structure, functional expression, and chro-
mosomal
- [113386]localization of the bumetanide-sensitive Na-K-Cl cotrans-
porter in
- [113387]human colon. J. Biol. Chem. 270: 17977-17985, 1995.
- [113388]19983.
- [113389]Quaggin, S. E.; Payne, J. A.; Forbush, B., III; Igarashi, P.:
- [113390]Localization of the renal Na-K-Cl cotransporter gene
(Slc12a1) on
- [113391]mouse chromosome 2. Mammalian Genome 6: 557-561,
1995.
- [113392]19984.

[113393] Xu, J.-C.; Lytle, C.; Zhu, T. T.; Payne, J. A.; Benz, E., Jr.;
 [113394] Forbush, B., III: Molecular cloning and functional expres-
 sion of
 [113395] the bumetanide-sensitive Na-K-Cl cotransporter. Proc.
 Nat. Acad.
 [113396] Sci. 91: 2201-2205, 1994.
 [113397] 19985.
 [113398] Sedlacek, Z.; Konecki, D. S.; Korn, B.; Klauck, S. M.;
 Poustka,
 [113399] A.: Evolutionary conservation and genomic organization of
 XAP-4,
 [113400] an Xq28 located gene coding for a human rab GDP-
 dissociation inhibitor
 [113401] (GDI). Mammalian Genome 5: 633-639, 1995.
 [113402] 19986.
 [113403] Sedlacek, Z.; Munstermann, E.; Mincheva, A.; Lichter, P.;
 Poustka,
 [113404] A.: The human rab GDI beta gene with long retroposon-
 rich introns
 [113405] maps to 10p15 and its pseudogene to 7p11-p13. Mam-
 malian Genome 9:

[113406]78–80, 1998.

[113407]19987.

[113408]Desseyn, J.–L.; Aubert, J.–P.; van Seuning, I.; Porchet, N.;

[113409]Laine, A.: Genomic organization of the 3–prime region of
the human

[113410]mucin gene MUC5B. J. Biol. Chem. 272: 16873–16883,
1997.

[113411]19988.

[113412]Desseyn, J.–L.; Guyonnet–Duperat, V.; Porchet, N.; Aubert,
J.–P.;

[113413]Laine, A.: Human mucin gene MUC5B, the 10.7–kb large
central exon

[113414]encodes various alternate subdomains resulting in a su-
per-repeat:

[113415]structural evidence for a 11p15.5 gene family. J. Biol.
Chem. 272:

[113416]3168–3178, 1997.

[113417]19989.

[113418]Gipson, I. K.; Moccia, R.; Spurr–Michaud, S.; Argueso, P.;
Gargiulo,

[113419]A. R.; Hill, J. A., III; Offner, G. D.; Keutmann, H. T.: The

amount

[113420]of MUC5B in cervical mucus peaks at midcycle. J. Clin. Endocr. Metab. 86:

[113421]594–600, 2001.

[113422]19990.

[113423]Keates, A. C.; Nunes, D. P.; Afdhal, N. H.; Troxler, R. F.; Offner,

[113424]G. D.: Molecular cloning of a major human gall bladder mucin: complete

[113425]C-terminal sequence and genomic organization of MUC5B. Biochem. J. 324:

[113426]295–303, 1997.

[113427]19991.

[113428]Troxler, R. F.; Offner, G. D.; Zhang, F.; Iontcheva, I.; Oppenheim,

[113429]F. G.: Molecular cloning of a novel high molecular weight mucin (MG1)

[113430]from human sublingual gland. Biochem. Biophys. Res. Commun. 217:

[113431]1112–1119, 1995.

[113432]19992.

[113433] Birck, C.; Poch, O.; Romier, C.; Ruff, M.; Mengus, G.; Lavigne,
 [113434] A.-C.; Davidson, I.; Moras, D.: Human TAFII28 and TAFII18
 interact
 [113435] through a histone fold encoded by atypical evolutionary
 conserved
 [113436] motifs also found in the SPT3 family. Cell 94: 239–249,
 1998.
 [113437] 19993.
 [113438] Mengus, G.; May, M.; Jacq, X.; Staub, A.; Tora, L.; Chambon, P.;
 [113439] Davidson, I.: Cloning and characterization of hTAFII18,
 hTAFII20
 [113440] and hTAFII28: three subunits of the human transcription
 factor TFIID. EMBO
 [113441] J. 14: 1520–1531, 1995.
 [113442] 19994.
 [113443] Waisfisz, Q.; Saar, K.; Morgan, N. V.; Altay, C.; Leegwater,
 P.
 [113444] A.; de Winter, J. P.; Komatsu, K.; Evans, G. R.; Wegner,
 R.-D.; Reis,
 [113445] A.; Joenje, H.; Arwert, F.; Mathew, C. G.; Pronk, J. C.; Dig-

weed,

[113446]M.: The Fanconi anemia group E gene, FANCE, maps to chromosome 6p. Am.

[113447]J. Hum. Genet. 64: 1400–1405, 1999.

[113448]19995.

[113449]Wegner, R.-D.; Henrichs, I.; Joenje, H.; Schroeder-Kurth, T.:

[113450]Fanconi anemia complementation group E: clinical and cytogenetic data

[113451]of the first patient. Clin. Genet. 50: 479–482, 1996.

[113452]19996.

[113453]Fink, J. M.; Hirsch, B.; Zheng, C.; Deitz, G.; Hatten, M. E.; Ross,

[113454]M. E.: The CNS neuronal migration gene, astrotactin, is mapped to

[113455]human chromosome band 1q25 by fluorescence in situ hybridization (FISH).

[113456](Abstract) Am. J. Hum. Genet. 57: A133 only, 1995.

[113457]19997.

[113458]Baker, C. H.; Matsuda, S. P. T.; Liu, D. R.; Corey, E. J.: Molecular

[113459]cloning of the human gene encoding lanosterol synthase
from a liver

[113460]cDNA library. Biochem. Biophys. Res. Commun. 213:
154–160, 1995.

[113461]19998.

[113462]Roessler, E.; Mittaz, L.; Du, Y.; Scott, H. S.; Chang, J.;
Rossier,

[113463]C.; Guipponi, M.; Matsuda, S. P. T.; Muenke, M.; An-
tonarakis, S. E.

[113464]: Structure of the human lanosterol synthase gene and its
analysis

[113465]as a candidate for holoprosencephaly (HPE1). Hum. Genet.
105: 489–495,

[113466]1999.

[113467]19999.

[113468]Young, M.; Chen, H.; Lalioti, M. D.; Antonarakis, S. E.: The
human

[113469]lanosterol synthase gene maps to chromosome 21q22.3.
Hum. Genet. 97:

[113470]620–624, 1996.

[113471]20000.

[113472]Green, P.; Lipman, D.; Hillier, L.; Waterston, R.; States, D.;

[113473]Claverie, J.-M.: Ancient conserved regions in new gene sequences

[113474]and the protein databases. Science 259: 1711–1716, 1993.

[113475]20001.

[113476]Schwartz, F.; Eisenman, R.; Knoll, J.; Gessler, M.; Bruns, G.:

[113477]cDNA sequence, genomic organization, and evolutionary conservation

[113478]of a novel gene from the WAGR region. Genomics 29: 526–532, 1995.

[113479]20002.

[113480]Schwartz, F.; Neve, R.; Eisenman, R.; Gessler, M.; Bruns, G.:

[113481]A WAGR region gene between PAX-6 and FSHB expressed in fetal brain. Hum.

[113482]Genet. 94: 658–664, 1994.

[113483]20003.

[113484]Joseph, R. E.; Walker, J.; Norris, F. A.: Assignment of the inositol

[113485]polyphosphate 4-phosphatase type I gene (INPP4A) to hu-

man chromosome

[113486]band 2q11.2 by in situ hybridization. Cytogenet. Cell
Genet. 87:

[113487]276–277, 1999.

[113488]20004.

[113489]Norris, F. A.; Auethavekiat, V.; Majerus, P. W.: The isolation

[113490]and characterization of cDNA encoding human and rat
brain inositol

[113491]polyphosphate 4–phosphatase. J. Biol. Chem. 270:
16128–16133, 1995.

[113492]20005.

[113493]van den Maagdenberg, A. M. J. M.; van den Hurk, H. H.;
Olde Weghuis,

[113494]D.; Wieringa, B.; Geurts van Kessel, A.; Hendriks, W. J. A.
J.: Assignment

[113495]of the human protein tyrosine phosphatase epsilon
(PTPRE) gene to

[113496]chromosome 10q26 by fluorescence in situ hybridization.
Genomics 30:

[113497]128–129, 1995.

[113498]20006.

[113499]Yue, Q.; Jen, J. C.; Nelson, S. F.; Baloh, R. W.: Progressive
[113500]ataxia due to a missense mutation in a calcium-channel
gene. Am.
[113501]J. Hum. Genet. 61: 1078–1087, 1997.
[113502]20007.
[113503]Wojcik, J.; Berg, M. A.; Esposito, N.; Geffner, M. E.; Sakati,
[113504]N.; Reiter, E. O.; Dower, S.; Francke, U.; Postel-Vinay,
M.–C.; Finidori,
[113505]J.: Four contiguous amino acid substitutions, identified in
patients
[113506]with Laron syndrome, differently affect the binding affinity
and intracellular
[113507]trafficking of the growth hormone receptor. J. Clin. En-
docr. Metab. 83:
[113508]4481–4489, 1998.
[113509]20008.
[113510]Azim, A. C.; Knoll, J. H. M.; Marfatia, S. M.; Peel, D. J.;
Bryant,
[113511]P. J.; Chishti, A. H.: DLG1: chromosome location of the
closest human
[113512]homologue of the Drosophila discs large tumor suppres-

sor gene. Genomics 30:

[113513]613–616, 1995.

[113514]20009.

[113515]Burgess, D. L.; Rafael, J. A.; Meisler, M. H.; Chamberlain, J.

[113516]S.: Dlgh1, a mouse homolog of the Drosophila discs-large gene, is

[113517]located on chromosome 16. Mammalian Genome 7:
623–624, 1996.

[113518]20010.

[113519]Hanada, T.; Lin, L.; Chandy, K. G.; Oh, S. S.; Chishti, A. H.:

[113520]Human homologue of the Drosophila discs large tumor suppressor binds

[113521]to p56lck tyrosine kinase and Shaker type Kv1.3 potassium channel

[113522]in T lymphocytes. J. Biol. Chem. 272: 26899–26904,
1997.

[113523]20011.

[113524]Lue, R. A.; Marfatia, S. M.; Branton, D.; Chishti, A. H.:
Cloning

[113525]and characterization of hdlg: the human homologue of the
Drosophila

[113526]discs large tumor suppressor binds to protein 4.1. Proc.
Nat. Acad.

[113527]Sci. 91: 9818–9822, 1994.

[113528]20012.

[113529]Mori, K.; Iwao, K.; Miyoshi, Y.; Nakagawara, A.; Kofu, K.;
Akiyama,

[113530]T.; Arita, N.; Hayakawa, T.; Nakamura, Y.: Identification of
brain-specific

[113531]splicing variants of the hDLG1 gene and altered splicing in
neuroblastoma

[113532]cell lines. J. Hum. Genet. 43: 123–127, 1998.

[113533]20013.

[113534]Peters, L. L.; Ciciotte, S. L.; Lin, L.; Chishti, A. H.: The
mouse

[113535]homolog of the Drosophila discs large tumor suppressor
gene maps to

[113536]chromosome 16. Mammalian Genome 7: 619–620, 1996.

[113537]20014.

[113538]Woods, K. A.; Dastot, F.; Preece, M. A.; Clark, A. J. L.; Pos-
tel-Vinay,

[113539]M.-C.; Chatelain, P. G.; Ranke, M. B.; Rosenfeld, R. G.;

Amselem,

[113540]S.; Savage, M. O.: Phenotype:genotype relationships in growth hormone

[113541]insensitivity syndrome. J. Clin. Endocr. Metab. 82: 3529–3535, 1997.

[113542]20015.

[113543]Bertaux, F.; Sharp, A. H.; Ross, C. A.; Lehrach, H.; Bates, G.

[113544]P.; Wanker, E.: HAP1–huntingtin interactions do not contribute to

[113545]the molecular pathology in Huntington's disease transgenic mice. FEBS

[113546]Lett. 426: 229–232, 1998.

[113547]20016.

[113548]Engelender, S.; Sharp, A. H.; Colomer, V.; Tokito, M. K.; Lanahan,

[113549]A.; Worley, P.; Holzbaur, E. L. F.; Ross, C. A.: Huntingtin-associated

[113550]protein 1 (HAP1) interacts with the p150(Glued) subunit of dynactin. Hum.

[113551]Molec. Genet. 6: 2205–2212, 1997.

[113552]20017.

[113553]Li, S.-H.; Hosseini, S. H.; Gutekunst, C.-A.; Hersch, S. M.;
Ferrante,
[113554]R. J.; Li, X.-J.: A human HAP1 homologue: cloning, expres-
sion, and
[113555]interaction with huntingtin. J. Biol. Chem. 273:
19220-19227, 1998.
[113556]20018.
[113557]Li, X.-J.; Li, S.-H.; Sharp, A. H.; Nucifora, F. C., Jr.;
Schilling,
[113558]G.; Lanahan, A.; Worley, P.; Snyder, S. H.; Ross, C. A.: A
huntingtin-associated
[113559]protein enriched in brain with implications for pathology.
Nature 378:
[113560]398-402, 1995.
[113561]20019.
[113562]Nasir, J.; Duan, K.; Nichol, K.; Engelender, S.; Ashworth, R.;
[113563]Colomer, V.; Thomas, S.; Distèche, C. M.; Hayden, M. R.;
Ross, C.
[113564]A.: Gene structure and map location of the murine ho-
molog of the
[113565]Huntington-associated protein, Hap1. Mammalian
Genome 9: 565-570,

[113566]1998.

[113567]20020.

[113568]Nasir, J.; Lafuente, M.-J.; Duan, K.; Colomer, V.; Engelen-
der,

[113569]S.; Ingersoll, R.; Margolis, R. L.; Ross, C. A.; Hayden, M. R.:
Human

[113570]huntingtin-associated protein (HAP-1) gene: genomic or-
ganisation and

[113571]an intragenic polymorphism. Gene 254: 181-187, 2000.

[113572]20021.

[113573]Coon, S. L.; Mazuruk, K.; Bernard, M.; Roseboom, P. H.;
Klein,

[113574]D. C.; Rodriguez, I. R.: The human serotonin N-
acetyltransferase

[113575](EC 2.3.1.87) gene (AANAT): structure, chromosomal lo-
calization, and

[113576]tissue expression. Genomics 34: 76-84, 1996.

[113577]20022.

[113578]Coon, S. L.; Roseboom, P. H.; Baler, R.; Weller, J. L.; Nam-
boodiri,

[113579]M. A. A.; Koonin, E. V.; Klein, D. C.: Pineal serotonin N-

acetyltransferase:

[113580]expression cloning and molecular analysis. Science 270:

1681–1683,

[113581]1995.

[113582]20023.

[113583]Joy, J. E.; Johnson, G. S.; Lazar, T.; Ralph, M. R.;

Hochstrasser,

[113584]A. C.; Menaker, M.; Merrill, C. R.: Protein differences in tau

mutant

[113585]hamsters: candidate clock proteins. Brain Res. Mol. Brain

Res. 15:

[113586]8–14, 1992.

[113587]20024.

[113588]Ralph, M. R.; Menaker, M.: A mutation of the circadian

system

[113589]in golden hamsters. Science 241: 1225–1227, 1988.

[113590]20025.

[113591]Tosini, G.; Menaker, M.: Circadian rhythms in cultured

mammalian

[113592]retina. Science 272: 419–421, 1996.

[113593]20026.

- [113594]Yoshimura, T.; Nagabukuro, A.; Matsuda, Y.; Suzuki, T.; Kuroiwa, A.; Iigo, M.; Namikawa, T.; Ebihara, S.: Chromosomal mapping of the gene encoding serotonin N-acetyltransferase to rat chromosome 10q32.3 and mouse chromosome 11E2. Cytogenet. Cell Genet. 79: 172–175, 1997.
- [113598]20027.
- [113599]Broccoli, D.; Chong, L.; Oelmann, S.; Fernald, A. A.; Marziliano, N.; van Steensel, B.; Kipling, D.; Le Beau, M. M.; de Lange, T.: Comparison of the human and mouse genes encoding the telomeric protein, TRF1: chromosomal localization, expression and conserved protein domains. Hum. Molec. Genet. 6: 69–76, 1997.
- [113604]20028.
- [113605]Chong, L.; van Steensel, B.; Broccoli, D.; Erdjument-Bromage, H.; Hanish, J.; Tempst, P.; de Lange, T.: A human telomeric

protein. Science 270:

[113607]1663–1667, 1995.

[113608]20029.

[113609]Fairall, L.; Chapman, L.; Moss, H.; de Lange, T.; Rhodes,
D.:

[113610]Structure of the TRFH dimerization domain of the human
telomeric proteins

[113611]TRF1 and TRF2. Molec. Cell 8: 351–361, 2001.

[113612]20030.

[113613]Griffith, J. D.; Comeau, L.; Rosenfield, S.; Stansel, R. M.;
Bianchi,

[113614]A.; Moss, H.; de Lange, T.: Mammalian telomeres end in a
large duplex

[113615]loop. Cell 97: 503–514, 1999.

[113616]20031.

[113617]Kim, S.; Kaminker, P.; Campisi, J.: TIN2, a new regulator of
telomere

[113618]length in human cells. Nature Genet. 23: 405–412, 1999.

[113619]20032.

[113620]Marcand, S.; Gilson, E.; Shore, D.: A protein–counting

mechanism

[113621] for telomere length regulation in yeast. *Science* 275:
986–990, 1997.

[113622] 20033.

[113623] Okabe, J.; Eguchi, A.; Masago, A.; Hayakawa, T.; Nakan-
ishi, M.

[113624] TRF1 is a critical trans-acting factor required for de novo
telomere

[113625] formation in human cells. *Hum. Molec. Genet.* 9:
2639–2650, 2000.

[113626] 20034.

[113627] Young, A. C.; Chavez, M.; Giambernardi, T. A.; Mattern, V.;
McGill,

[113628] J. R.; Harris, J. M.; Sarosdy, M. F.; Patel, P.; Sakaguchi, A. Y.

[113629] Organization and expression of human telomere repeat
binding factor

[113630] genes. *Somat. Cell Molec. Genet.* 23: 275–286, 1997.

[113631] 20035.

[113632] Zakian, V. A.: Telomeres: beginning to understand the
end. *Science* 270:

[113633] 1601–1607, 1995.

[113634]20036.

[113635]Brodsky, G.; Otterson, G. A.; Parry, B. B.; Hart, I.; Patterson,
son,

[113636]D.; Kaye, F. J.: Localization of STCH to human chromosome 21q11.1. Genomics 30:

[113637]627-628, 1995.

[113638]20037.

[113639]Otterson, G. A.; Flynn, G. C.; Kratzke, R. A.; Coxon, A.; Johnston,

[113640]P. G.; Kaye, F. J.: Stch encodes the 'ATPase core' of a mitochondrial
crosomal

[113641]stress70 protein. EMBO J. 13: 1216-1225, 1994.

[113642]20038.

[113643]Reeves, R. H.; Rue, E.; Yu, J.; Kao, F.-T.: Stch maps to mouse
mouse

[113644]chromosome 16, extending the conserved synteny with human chromosome
human chromosome

[113645]21. Genomics 49: 156-157, 1998.

[113646]20039.

[113647]Nielsen, P. J.; Rochelle, J. M.; Seldin, M. F.: The functional

[113648] genes for protein synthesis initiation factor 4A1 and 4A2
map to
[113649] mouse chromosomes 11 and 16. Mammalian Genome 4:
185–186, 1993.
[113650] 20040.

[113651] Nielsen, P. J.; Trachsel, H.: The mouse protein synthesis
initiation
[113652] factor 4A gene family includes two related functional
genes which
[113653] are differentially expressed. EMBO J. 7: 2097–2105, 1988.
[113654] 20041.

[113655] Sudo, K.; Takahashi, E.; Nakamura, Y.: Isolation and map-
ping of
[113656] the human EIF4A2 gene homologous to the murine pro-
tein synthesis initiation
[113657] factor 4A-II gene Eif4a2. Cytogenet. Cell Genet. 71:
385–388, 1995.
[113658] 20042.

[113659] Gibson, M. A.; Hatzinikolas, G.; Kumaratilake, J. S.; Sand-
berg,
[113660] L. B.; Nicholl, J. K.; Sutherland, G. R.; Cleary, E. G.: Further

[113661] characterization of proteins associated with elastic fiber
microfibrils

[113662] including the molecular cloning of MAGP-2 (MP25). J. Biol.
Chem. 271:

[113663] 1096-1103, 1996.

[113664] 20043.

[113665] Gelb, B. D.; Willner, J. P.; Verloes, A.; Herens, C.; Desnick,
[113666] R. J.: Mutation analysis of pycnodysostosis reveals uni-
parental disomy

[113667] of chromosome 1. (Abstract) Am. J. Hum. Genet. 61
(suppl.): A28

[113668] only, 1997.

[113669] 20044.

[113670] Gladyshev, V. N.; Jeang, K.-T.; Stadtman, T. C.: Selenocys-
teine,

[113671] identified as the penultimate C-terminal residue in human
T-cell thioredoxin

[113672] reductase, corresponds to TGA in the human placental
gene. Proc.

[113673] Nat. Acad. Sci. 93: 6146-6151, 1996.

[113674] 20045.

- [113675]Zhang, K.; Lindsberg, P. J.; Tatlisumak, T.; Kaste, M.; Olsen,
[113676]H. S.; Andersson, L. C.: Stanniocalcin: a molecular guard
of neurons
[113677]during cerebral ischemia. Proc. Nat. Acad. Sci. 97:
3637–3642, 2000.
[113678]20046.
- [113679]Gillett, G. T.; Fox, M. F.; Rowe, P. S. N.; Casimir, C. M.;
Povey,
[113680]S.: Mapping of human non-muscle type cofilin (CFL1) to
chromosome
[113681]11q13 and muscle-type cofilin (CFL2) to chromosome 14.
Ann. Hum.
[113682]Genet. 60: 201–211, 1996.
[113683]20047.
- [113684]Ogawa, K.; Tashima, M.; Yumoto, Y.; Okuda, T.; Sawada,
H.; Okuma,
[113685]M.; Maruyama, Y.: Coding sequence of human placenta
cofilin cDNA. Nuc.
[113686]Acids Res. 18: 7169 only, 1990.
[113687]20048.

[113688]Ono, S.; Minami, N.; Abe, H.; Obinata, T.: Characterization
of
[113689]a novel cofilin isoform that is predominantly expressed in
mammalian
[113690]skeletal muscle. J. Biol. Chem. 269: 15280–15286, 1994.
[113691]20049.

[113692]Georgakopoulos, T.; Thireos, G.: Two distinct yeast tran-
scriptional
[113693]activators require the function of the GCN5 protein to
promote normal
[113694]levels of transcription. EMBO J. 11: 4145–4152, 1992.
[113695]20050.

[113696]Inoue, M.; Isomura, M.; Ikegawa, S.; Fujiwara, T.; Shin, S.;
Moriya,
[113697]H.; Nakamura, Y.: Isolation and characterization of a hu-
man cDNA
[113698]clone (GCN5L1) homologous to GCN5, a yeast transcrip-
tion factor. Cytogenet.
[113699]Cell Genet. 73: 134–136, 1996.
[113700]20051.

[113701]Emahazion, T.; Beskow, A.; Gyllenstein, U.; Brookes, A. J.:

Intron

[113702]based radiation hybrid mapping of 15 complex I genes of the human

[113703]electron transport chain. Cytogenet. Cell Genet. 82: 115–119, 1998.

[113704]20052.

[113705]Byrd, P. J.; Cooper, P. R.; Stankovic, T.; Kullar, H. S.; Watts,

[113706]G. D. J.; Robinson, P. J.; Taylor, M. R.: A gene transcribed from

[113707]the bidirectional ATM promoter coding for a serine rich protein: amino

[113708]acid sequence, structure and expression studies. Hum. Molec. Genet. 5:

[113709]1785–1791, 1996.

[113710]20053.

[113711]Chen, X.; Yang, L.; Udar, N.; Liang, T.; Uhrhammer, N.; Xu, S.;

[113712]Bay, J.–O.; Wang, Z.; Dandakar, S.; Chiplunkar, S.; Klisak, I.; Telatar,

[113713]M.; Yang, H.; Concannon, P.; Gatti, R. A.: CAND3: a ubiquitously

[113714]expressed gene immediately adjacent and in opposite

transcriptional

[113715]orientation to the ATM gene at 11q23.1. Mammalian
Genome 8: 129–133,

[113716]1997.

[113717]20054.

[113718]Imai, T.: Personal Communication. Chiba, Japan
9/12/1996.

[113719]20055.

[113720]Imai, T.; Sugawara, T.; Nishiyama, A.; Shimada, R.; Ohki,
R.; Seki,

[113721]N.; Sagara, M.; Ito, H.; Yamauchi, M.; Hori, T.: The struc-
ture and

[113722]organization of the human NPAT gene. Genomics 42:
388–392, 1997.

[113723]20056.

[113724]Imai, T.; Yamauchi, M.; Seki, N.; Sugawara, T.; Saito, T.;
Matsuda,

[113725]Y.; Ito, H.; Nagase, T.; Nomura, N.; Hori, T.: Identification
and

[113726]characterization of a new gene physically linked to the
ATM gene. Genome

[113727]Res. 6: 439–447, 1996.

[113728]20057.

[113729]Borowsky, B.; Hoffman, B. J.: Analysis of a gene encoding two

[113730]glycine transporter variants reveals alternative promoter usage and

[113731]a novel gene structure. J. Biol. Chem. 273: 29077–29085, 1998.

[113732]20058.

[113733]Borowsky, B.; Mezey, E.; Hoffman, B. J.: Two glycine transporter

[113734]variants with distinct localization in the CNS and peripheral tissues

[113735]are encoded by a common gene. Neuron 10: 851–863, 1993.

[113736]20059.

[113737]Jones, E. M. C.; Fernald, A.; Bell, G. I.; Le Beau, M. M.: Assignment

[113738]of SLC6A9 to human chromosome band 1p33 by in situ hybridization. Cytogenet.

[113739]Cell Genet. 71: 211, 1995.

[113740]20060.

[113741]Kim, K.-M.; Kingsmore, S. F.; Han, H.; Yang-Feng, T. L.;
Godinot,

[113742]N.; Seldin, M. F.; Caron, M. G.; Giros, B.: Cloning of the
human

[113743]glycine transporter type 1: molecular and pharmacological
characterization

[113744]of novel isoform variants and chromosomal localization of
the gene

[113745]in the human and mouse genomes. Molec. Pharmacol. 45:
608-617, 1994.

[113746]20061.

[113747]Arai, Y.; Hosoda, F.; Kobayashi, H.; Arai, K.; Hayashi, Y.;
Kamada,

[113748]N.; Kaneko, Y.; Ohki, M.: The inv(11)(p15q22) chromo-
some translocation

[113749]of de novo and therapy-related myeloid malignancies re-
sults in fusion

[113750]of the nucleoporin gene, NUP98, with the putative RNA
helicase gene,

[113751]DDX10. Blood 89: 3936-3944, 1997.

[113752]20062.

[113753]Dash, A. B.; Williams, I. R.; Kutok, J. L.; Tomasson, M. H.;

Anastasiadou,

[113754]E.; Lindahl, K.; Li, S.; Van Etten, R. A.; Borrow, J.; Housman, D.;

[113755]Druker, B.; Gilliland, D. G.: A murine model of CML blast crisis

[113756]induced by cooperation between BCR/ABL and NUP98/HOXA9. Proc. Nat.

[113757]Acad. Sci. 99: 7622–7627, 2002.

[113758]20063.

[113759]Enninga, J.; Levy, D. E.; Blobel, G.; Fontoura, B. M. A.: Role

[113760]of nucleoporin induction in releasing an mRNA nuclear export block. Science 295:

[113761]1523–1525, 2002.

[113762]20064.

[113763]Fontoura, B. M.; Blobel, G.; Matunis, M. J.: A conserved biogenesis

[113764]pathway for nucleoporins: proteolytic processing of a 186–kilodalton

[113765]precursor generates Nup98 and the novel nucleoporin, Nup96. J. Cell

[113766]Biol. 144: 1097–1112, 1999.

[113767]20065.

[113768]Hodel, A. E.; Hodel, M. R.; Griffis, E. R.; Hennig, K. A.; Ratner,

[113769]G. A.; Xu, S.; Powers, M. A.: The three-dimensional structure of

[113770]the autoproteolytic, nuclear pore-targeting domain of the human nucleoporin

[113771]Nup98. *Molec. Cell* 10: 347–358, 2002.

[113772]20066.

[113773]Jaju, R. J.; Fidler, C.; Haas, O. A.; Strickson, A. J.; Watkins,

[113774]F.; Clark, K.; Cross, N. C. P.; Cheng, J.-F.; Aplan, P. D.; Kearney,

[113775]L.; Boultonwood, J.; Wainscoat, J. S.: A novel gene, NSD1, is fused

[113776]to NUP98 in the t(5;11)(q35;p15.5) in de novo childhood acute myeloid

[113777]leukemia. *Blood* 98: 1264–1267, 2001.

[113778]20067.

[113779]Jaju, R. J.; Haas, C. A.; Neat, M.; et al: A new recurrent translocation,

[113780]t(5;11)(q35p15.5), associated with del(5q) in childhood

acute myeloid

[113781]leukemia. Blood 94: 773–780, 1999.

[113782]20068.

[113783]Mizuno, T.; Kaibuchi, K.; Yamamoto, T.; Kawamura, M.;
Sakoda,

[113784]T.; Fujioka, H.; Matsuura, Y.; Takai, Y.: A stimulatory GDP/
GTP exchange

[113785]protein for smg p21 is active on the post-translationally
processed

[113786]form of c-Ki-ras p21 and rhoA p21. Proc. Nat. Acad. Sci.
88: 6442–6446,

[113787]1991.

[113788]20069.

[113789]Nakamura, T.; Largaespada, D. A.; Lee, M. P.; Johnson, L.
A.;

[113790]Ohyashiki, K.; Toyama, K.; Chen, S. J.; Willman, C. L.;
Chen, I.-M.;

[113791]Feinberg, A. P.; Jenkins, N. A.; Copeland, N. G.; Shaugh-
nessy, J.

[113792]D., Jr.: Fusion of the nucleoporin gene NUP98 to HOXA9
by the chromosome

[113793]translocation t(7;11)(p15;p15) in human myeloid

leukaemia. Nature

[113794]Genet. 12: 154–158, 1996.

[113795]20070.

[113796]Radu, A.; Moore, M. S.; Blobel, G.: The peptide repeat domain

[113797]of nucleoporin Nup98 functions as a docking site in transport across

[113798]the nuclear pore complex. Cell 81: 215–222, 1995.

[113799]20071.

[113800]Rosati, R.; La Starza, R.; Veronese, A.; Aventin, A.; Schwienbacher,

[113801]C.; Vallespi, T.; Negrini, M.; Martelli, M. F.; Mecucci, C.: NUP98

[113802]is fused to the NSD3 gene in acute myeloid leukemia associated with

[113803]t(8;11)(p11.2;p15). Blood 99: 3857–3860, 2002.

[113804]20072.

[113805]von Kobbe, C.; van Deursen, J. M. A.; Rodrigues, J. P.; Sitterlin,

[113806]D.; Bachi, A.; Wu, X.; Wilm, M.; Carmo-Fonseca, M.; Izaurralde, E.

[113807]: Vesicular stomatitis virus matrix protein inhibits host cell gene

[113808]expression by targeting the nucleoporin Nup98. Molec. Cell 6: 1243–1252,

[113809]2000.

[113810]20073.

[113811]Cheng, Y.; Austin, S. C.; Rocca, B.; Koller, B. H.; Coffman, T.

[113812]M.; Grosser, T.; Lawson, J. A.; FitzGerald, G. A.: Role of prostacyclin

[113813]in the cardiovascular response to thromboxane A2. Science 296: 539–541,

[113814]2002.

[113815]20074.

[113816]Fuse, I.; Mito, M.; Hattori, A.; Higuchi, W.; Shibata, A.; Ushikubi,

[113817]F.; Okuma, M.; Yahata, K.: Defective signal transduction induced

[113818]by thromboxane A2 in a patient with a mild bleeding disorder: impaired

[113819]phospholipase C activation despite normal phospholipase A2 activation. Blood 81:

[113820]994–1000, 1993.

[113821]20075.

[113822]Hirata, M.; Hayashi, Y.; Ushikubi, F.; Yokota, Y.; Kageyama, R.;

[113823]Nakanishi, S.; Narumiya, S.: Cloning and expression of cDNA for a

[113824]human thromboxane A2 receptor. Nature 349: 617–620, 1991.

[113825]20076.

[113826]Hirata, T.; Kakizuka, A.; Ushikubi, F.; Fuse, I.; Okuma, M.; Narumiya,

[113827]S.: Arg60-to-leu mutation of the human thromboxane A2 receptor in

[113828]a dominantly inherited bleeding disorder. J. Clin. Invest. 94: 1662–1667,

[113829]1994.

[113830]20077.

[113831]Watanabe, O.; Maruyama, I.; Arimura, K.; Kitajima, I.; Arimura,

[113832]H.; Hanatani, M.; Matsuo, K.; Arisato, T.; Osame, M.: Over-production

[113833] of vascular endothelial growth factor/vascular permeability factor

[113834] is causative in Crow-Fukase (POEMS) syndrome. Muscle Nerve 21: 1390-1397,

[113835] 1998.

[113836] 20078.

[113837] Wei, M.-H.; Popescu, N. C.; Lerman, M. I.; Merrill, M. J.; Zimonjic,

[113838] D. B.: Localization of the human vascular endothelial growth factor

[113839] gene, VEGF, at chromosome 6p12. Hum. Genet. 97: 794-797, 1996.

[113840] 20079.

[113841] Wong, A. K.; Alfert, M.; Castrillon, D. H.; Shen, Q.; Holash,

[113842] J.; Yancopoulos, G. D.; Chin, L.: Excessive tumor-elaborated VEGF

[113843] and its neutralization define a lethal paraneoplastic syndrome. Proc.

[113844] Nat. Acad. Sci. 98: 7481-7486, 2001.

[113845] 20080.

[113846] Wulff, C.; Wilson, H.; Lague, P.; Duncan, W. C.; Arm-

strong, D.

[113847]G.; Fraser, H. M.: Angiogenesis in the human corpus luteum: localization

[113848]and changes in angiopoietins, Tie-2, and vascular endothelial growth

[113849]factor messenger ribonucleic acid. J. Clin. Endocr. Metab. 85: 4302-4309,

[113850]2000.

[113851]20081.

[113852]Ylikorkala, A.; Rossi, D. J.; Korsisaari, N.; Luukko, K.; Alitalo,

[113853]K.; Henkemeyer, M.; Makela, T. P.: Vascular abnormalities and deregulation

[113854]of VEGF in Lkb1-deficient mice. Science 293: 1323-1326, 2001.

[113855]20082.

[113856]Bodner, M.; Fridkin, M.; Gozes, I.: Coding sequences for vasoactive

[113857]intestinal peptide and PHM-27 peptide are located on two adjacent

[113858]exons in the human genome. Proc. Nat. Acad. Sci. 82: 3548-3551,

[113859]1985.

[113860]20083.

[113861]Delgado, M.; Abad, C.; Martinez, C.; Leceta, J.; Gomariz, R.
P.

[113862]; Vasoactive intestinal peptide prevents experimental
arthritis by

[113863]downregulating both autoimmune and inflammatory com-
ponents of the

[113864]disease. Nature Med. 7: 563–568, 2001.

[113865]20084.

[113866]Gotoh, E.; Yamagami, T.; Yamamoto, H.; Okamoto, H.:
Chromosomal

[113867]assignment of human VIP/PHM-27 gene to 6q26–q27 re-
gion by spot blot

[113868]hybridization and in situ hybridization. Biochem. Int. 17:
555–562,

[113869]1988.

[113870]20085.

[113871]Gozes, I.; Avidor, R.; Yahav, Y.; Katznelson, D.; Croce, C.
M.;

[113872]Huebner, K.: The gene encoding vasoactive intestinal pep-

tide is located

[113873]on human chromosome 6p21–6qter. Hum. Genet. 75:
41–44, 1987.

[113874]20086.

[113875]Gozes, I.; Nakai, H.; Byers, M.; Avidor, R.; Weinstein, Y.;
Shani,

[113876]Y.; Shows, T. B.: Sequential expression in the nervous sys-
tem of

[113877]C-MYB and VIP genes, located in human chromosomal re-
gion 6q24. Somat.

[113878]Cell Molec. Genet. 13: 305–313, 1987.

[113879]20087.

[113880]Heinz-Erian, P.; Dey, R. D.; Flux, M.; Said, S. I.: Deficient

[113881]vasoactive intestinal peptide innervation in sweat glands
of cystic

[113882]fibrosis patients. Science 229: 1407–1408, 1985.

[113883]20088.

[113884]Itoh, N.; Obata, K.; Yanaihara, N.; Okamoto, H.: Human
preprovasoactive

[113885]intestinal polypeptide contains a novel PHI-27-like pep-
tide, PHM-27. Nature 304:

[113886]547-549, 1983.

[113887]20089.

[113888]Linder, S.; Barkhem, T.; Norberg, A.; Persson, H.;
Schalling, M.;

[113889]Hokfelt, T.; Magnusson, G.: Structure and expression of
the gene

[113890]encoding the vasoactive intestinal peptide precursor. Proc.
Nat.

[113891]Acad. Sci. 84: 605-609, 1987.

[113892]20090.

[113893]Omary, M. B.; Kagnoff, M. F.: Identification of nuclear re-
ceptors

[113894]for VIP on a human colonic adenocarcinoma cell line. Sci-
ence 238:

[113895]1578-1581, 1987.

[113896]20091.

[113897]Couvineau, A.; Rouyer-Fessard, C.; Darmoul, D.; Maoret,
J.-J.;

[113898]Carrero, I.; Ogier-Denis, E.; Laburthe, M.: Human intesti-
nal VIP

[113899]receptor: cloning and functional expression of two cDNA

encoding proteins

[113900]with different N-terminal domains. Biochem. Biophys. Res. Commun. 200:

[113901]769–776, 1994.

[113902]20092.

[113903]Hashimoto, H.; Nishino, A.; Shintani, N.; Hagihara, N.; Copeland,

[113904]N. G.; Jenkins, N. A.; Yamamoto, K.; Matsuda, T.; Ishihara, T.; Nagata,

[113905]S.; Baba, A.: Genomic organization and chromosomal location of the

[113906]mouse vasoactive intestinal polypeptide 1 (VPAC-1) receptor. Genomics 58:

[113907]90–93, 1999.

[113908]20093.

[113909]Sreedharan, S. P.; Huang, J.-X.; Cheung, M.-C.; Goetzl, E. J.:

[113910]Structure, expression, and chromosomal localization of the type I

[113911]human vasoactive intestinal peptide receptor gene. Proc. Nat. Acad.

[113912]Sci. 92: 2939–2943, 1995.

[113913]20094.

[113914]Sreedharan, S. P.; Patel, D. R.; Huang, J.-X.; Goetzl, E. J.:

[113915]Cloning and functional expression of a human neuroen-
docrine vasoactive

[113916]intestinal peptide receptor. Biochem. Biophys. Res. Com-
mun. 193:

[113917]546-553, 1993.

[113918]20095.

[113919]Sreedharan, S. P.; Robichon, A.; Peterson, K. E.; Goetzl, E.
J.

[113920]: Cloning and expression of the human vasoactive intesti-
nal peptide

[113921]receptor. Proc. Nat. Acad. Sci. 88: 4986-4990, 1991.

[113922]20096.

[113923]Anderson, S. R.; Geertinger, P.; Larsen, H.-W.; Mikkelsen,
M.;

[113924]Parving, A.; Vestermark, S.; Warburg, M.: Aniridia, cataract
and

[113925]gonadoblastoma in a mentally retarded girl with deletion
of chromosome

[113926]11: a clinicopathological case report. Ophthalmologica

176: 171-177,

[113927]1978.

[113928]20097.

[113929]Babaian, R. J.; Skinner, D. G.; Waisman, J.: Wilms' tumor in
the

[113930]adult patient: diagnosis, management, and review of the
world medical

[113931]literature. Cancer 45: 1713-1719, 1980.

[113932]20098.

[113933]Beckwith, J. B.: Nephrogenic rests and the pathogenesis of
Wilms

[113934]tumor: developmental and clinical considerations. Am. J.
Med. Genet. 79:

[113935]268-273, 1998.

[113936]20099.

[113937]Blanchet, P.; Daloze, P.; Lesage, R.; Papas, S.; van Camp-
enhout,

[113938]J.: XY gonadal dysgenesis with gonadoblastoma discov-
ered after kidney

[113939]transplantation. Am. J. Obstet. Gynec. 129: 221-222,
1977.

[113940]20100.

[113941]Bond, J. V.: Bilateral Wilms' tumor: age at diagnosis, associated

[113942]congenital anomalies, and possible pattern of inheritance.
Lancet II:

[113943]482–484, 1975.

[113944]20101.

[113945]Bonetta, L.; Huang, A.; Gregoris, M.; Yeger, H.; Williams, B.
R.

[113946]G.: Characterization of a homozygous deletion mapping to
the Wilms

[113947]tumor region on 11p13. (Abstract) Cytogenet. Cell Genet.
51: 965,

[113948]1989.

[113949]20102.

[113950]Bove, K. E.: Wilms' tumor and related abnormalities in the
fetus

[113951]and newborn. Semin. Perinatol. 23: 310–318, 1999.

[113952]20103.

[113953]Breslow, N. E.; Beckwith, J. R.: Epidemiological features of
Wilms'

[113954]tumor: results of the national Wilms' tumor study. J. Nat.
Cancer

[113955]Inst. 68: 429-436, 1982.

[113956]20104.

[113957]Brown, W. T.; Puranik, S. R.; Altman, D. H.; Hardin, H. C.,
Jr.

[113958]: Wilms' tumor in three successive generations. Surgery
72: 756-761,

[113959]1972.

[113960]20105.

[113961]Cordero, J. F.; Li, F. P.; Holmes, L. B.; Gerald, P. S.: Wilms

[113962]tumor in five cousins. Pediatrics 66: 716-719, 1980.

[113963]20106.

[113964]Dao, D. D.; Schroeder, W. T.; Chao, L.-Y.; Kikuchi, H.;
Strong,

[113965]L. C.; Riccardi, V. M.; Pathak, S.; Nichols, W. W.; Lewis, W.
H.;

[113966]Saunders, G. F.: Genetic mechanisms of tumor-specific
loss of 11p

[113967]DNA sequences in Wilms tumor. Am. J. Hum. Genet. 41:
202-217, 1987.

[113968]20107.

[113969]de Martinville, B.; Francke, U.: HRAS1, insulin and beta-globin

[113970]map outside of 11p11.2-p14.1. (Abstract) Cytogenet. Cell Genet. 37:

[113971]530, 1984.

[113972]20108.

[113973]Denys, P.; Malvaux, P.; van den Berghe, H.; Tanghe, W.; Proesmans,

[113974]W.: Association d'un syndrome anatomo-pathologique de pseudohermaphrodisme

[113975]masculin, d'une tumeur de Wilms, d'une nephropathie parenchymateuse

[113976]et d'un mosaicisme XX/XY. Arch. Franc. Pediat. 24: 729-739, 1967.

[113977]20109.

[113978]Ueno, S.; Maruki, Y.; Nakamura, M.; Tomemori, Y.; Kamae, K.; Tanabe,

[113979]H.; Yamashita, Y.; Matsuda, S.; Kaneko, S.; Sano, A.: The gene encoding

[113980]a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. Nature

[113981]Genet. 28: 121–122, 2001.

[113982]20110.

[113983]Wolf, B.; Feldman, G. L.: The biotin-dependent carboxylase deficiencies. Am.

[113984]J. Hum. Genet. 34: 699–716, 1982.

[113985]20111.

[113986]Hirata, T.; Ushikubi, F.; Kakizuka, A.; Okuma, M.; Narumiya, S.

[113987]: Two thromboxane A(2) receptor isoforms in human platelets: opposite

[113988]coupling to adenylyl cyclase with different sensitivity to arg60-to-leu

[113989]mutation. J. Clin. Invest. 97: 949–956, 1996.

[113990]20112.

[113991]Nusing, R. M.; Hirata, M.; Kakizuka, A.; Eki, T.; Ozawa, K.; Narumiya,

[113992]S.: Characterization and chromosomal mapping of the human thromboxane

[113993]A2 receptor gene. J. Biol. Chem. 268: 25253–25259, 1993.

[113994]20113.

[113995]Schwengel, D. A.; Nouri, N.; Meyers, D. A.; Levitt, R. C.:
Linkage

[113996]mapping of the human thromboxane A2 receptor
(TBXA2R) to chromosome

[113997]19p13.3 using transcribed 3-prime untranslated DNA se-
quence polymorphisms. Genomics 18:

[113998]212-215, 1993.

[113999]20114.

[114000]Thomas, D. W.; Mannon, R. B.; Mannon, P. J.; Latour, A.;
Oliver,

[114001]J. A.; Hoffman, M.; Smithies, O.; Koller, B. H.; Coffman, T.
M.:

[114002]Coagulation defects and altered hemodynamic responses
in mice lacking

[114003]receptors for thromboxane A(2). J. Clin. Invest. 102:
1994-2001,

[114004]1998.

[114005]20115.

[114006]Unoki, M.; Furuta, S.; Onouchi, Y.; Watanabe, O.; Doi, S.;
Fujiwara,

[114007]H.; Miyatake, A.; Fujita, K.; Tamari, M.; Nakamura, Y.: As-
sociation

- [114008] studies of 33 single nucleotide polymorphisms (SNPs) in
29 candidate
- [114009] genes for bronchial asthma: positive association a T924C
polymorphism
- [114010] in the thromboxane A2 receptor gene. Hum. Genet. 106:
440–446, 2000.
- [114011] 20116.
- [114012] Ushikubi, F.; Nakajima, M.; Hirata, M.; Okuma, M.; Fuji-
wara, M.;
- [114013] Narumiya, S.: Purification of the thromboxane
A2/prostaglandin H2
- [114014] receptor from human blood platelets.. J. Biol. Chem. 264:
16496–16501,
- [114015] 1989.
- [114016] 20117.
- [114017] Ushikubi, F.; Okuma, M.; Kanaji, K.; Sugiyama, T.;
- Ogorochi, T.;
- [114018] Narumiya, S.; Uchino, H.: Hemorrhagic thrombocytopathy
with platelet
- [114019] thromboxane A2 abnormality: defective signal transduc-
tion with normal
- [114020] binding activity. Thromb. Haemost. 57: 158–164, 1987.

[114021]20118.

[114022]Ades, E. W.; Zwerner, R. K.; Acton, R. T.; Balch, C. M.: Iso-
lation

[114023]and partial characterisation of the human homologue of
Thy-1. J.

[114024]Exp. Med. 151: 400-406, 1980.

[114025]20119.

[114026]Bonewald, L.; Ades, E. W.; Tung, E.; Marchalonis, J. J.;
Wang,

[114027]A. C.: Biochemical characterization of human Thy-1. J. Im-
munogenet. 11:

[114028]283-296, 1984.

[114029]20120.

[114030]Gatti, R. A.; Lathrop, G. M.; Salser, W.; Silver, J.; Lalouel,

[114031]J. M.; White, R.: Location of Thy-1 with respect to a pri-
mary linkage

[114032]map of chromosome 11q. (Abstract) Cytogenet. Cell
Genet. 46: 618

[114033]only, 1987.

[114034]20121.

[114035]Gatti, R. A.; Shaked, R.; Wei, S.; Koyama, M.; Salser, W.;
Silver,

[114036]J.: DNA polymorphism in the human Thy-1 gene. Hum.
Immun. 22:

[114037]145-150, 1988.

[114038]20122.

[114039]Greenspan, R. J.; O'Brien, M. C.: Genetic evidence for the
role

[114040]of Thy-1 in neurite outgrowth in the mouse. J. Neuro-
genet. 5: 25-36,

[114041]1989.

[114042]20123.

[114043]Grzeschik, K.-H.; Kazazian, H. H.: Report of the committee
on

[114044]the genetic constitution of chromosomes 10, 11, and 12.
Cytogenet.

[114045]Cell Genet. 40: 179-205, 1985.

[114046]20124.

[114047]Letarte-Muirhead, M.; Barclay, A. N.; Williams, A. F.: Purifi-
cation

[114048]of the Thy-1 molecule, a major cell surface glycoprotein

of rat thymocytes.

[114049]Biochem. J. 151: 685–697, 1975.

[114050]20125.

[114051]Mansour, M. H.; Negm, H. I.; Cooper, E. L.: Thy-1 evolution. Dev.

[114052]Comp. Immun. 11: 3–15, 1987.

[114053]20126.

[114054]McKenzie, J. L.; Fabre, J. W.: Human Thy-1: unusual localization

[114055]and possible functional significance in lymphoid tissues. J. Immun. 126:

[114056]843–850, 1981.

[114057]20127.

[114058]Morris, R.: Thy-1 in developing nervous tissue. Dev. Neurosci. 7:

[114059]133–160, 1985.

[114060]20128.

[114061]Raff, M. C.: Surface antigenic markers for distinguishing T and

[114062]B lymphocytes in mice. Transplant. Rev. 6: 52–80, 1971.

[114063]20129.

[114064]Rettig, W. J.; Dracopoli, N. C.; Chesa, P. G.; Spengler, B. A.;

[114065]Beresford, H. R.; Davies, P.; Biedler, J. L.; Old, L. J.: Role of

[114066]human chromosome 11 in determining surface antigenic
phenotype of

[114067]normal and malignant cells: somatic cell genetic analysis
of eight

[114068]antigens, including putative human Thy-1. J. Exp. Med.
162: 1603-1619,

[114069]1985.

[114070]20130.

[114071]Rettig, W. J.; Dracopoli, N. C.; Silver, J.; Old, L. J.: Human

[114072]THY-1: regional mapping on chromosome 11 and com-
parison with other

[114073]chromosome 11-encoded cell surface glycoproteins.
(Abstract) Cytogenet.

[114074]Cell Genet. 40: 731 only, 1985.

[114075]20131.

[114076]Rettig, W. J.; Dracopoli, N. C.; Spengler, B. A.; Biedler, J.

[114077]L.; Old, L. J.: Somatic cell genetic analysis of human cell

surface

[114078]antigens, including putative human Thy-1: eight distinct antigenic

[114079]systems controlled by chromosome 11. (Abstract) Cytogenet. Cell

[114080]Genet. 40: 732 only, 1985.

[114081]20132.

[114082]Seki, T.; Spurr, N.; Obata, F.; Goyert, S.; Goodfellow, P.; Silver,

[114083]J.: The human Thy-1 gene: structure and chromosomal location. Proc.

[114084]Nat. Acad. Sci. 82: 6657-6661, 1985.

[114085]20133.

[114086]Tse, A. G. D.; Barclay, A. N.; Watts, A.; Williams, A. F.: A

[114087]glycophospholipid tail at the carboxyl terminus of the Thy-1 glycoprotein

[114088]of neurons and thymocytes. Science 230: 1003-1008, 1985.

[114089]20134.

[114090]Chen, Z.; Friedrich, G. A.; Soriano, P.: Transcriptional enhancer

[114091]factor 1 disruption by a retroviral gene trap leads to heart defects

[114092]and embryonic lethality in mice. Genes Dev. 8: 2293–2301, 1994.

[114093]20135.

[114094]Aso, T.; Tsai, P.; Kawaguchi, T.; Menninger, J. C.; Kitajima, S.;

[114095]Yasukochi, Y.; Ward, D. C.; Weissman, S. M.: Assignment of the human

[114096]GTF2F1 gene to chromosome 19p13.3. Genomics 16: 252–253, 1993.

[114097]20136.

[114098]Aso, T.; Vasavada, H. A.; Kawaguchi, T.; Germino, F. J.; Ganguly,

[114099]S.; Kitajima, S.; Weissman, S. M.; Yasukochi, Y.: Characterization

[114100]of cDNA for the large subunit of the transcription initiation factor

[114101]TFIIF. Nature 355: 461–464, 1992.

[114102]20137.

[114103]Finkelstein, A.; Kostrub, C. F.; Li, J.; Chavez, D. P.; Wang, B.

- [114104]Q.; Fang, S. M.; Greenblatt, J.; Burton, Z. F.: A cDNA encoding RAP74,
[114105]a general initiation factor for transcription by RNA polymerase II. Nature 355:
[114106]464–467, 1992.
[114107]20138.
- [114108]Joliot, V.; Demma, M.; Prywes, R.: Interaction with RAP74 subunit
[114109]of TFIIF is required for transcriptional activation by serum response
[114110]factor. Nature 373: 632–635, 1995.
[114111]20139.
- [114112]Sopta, M.; Burton, Z. F.; Greenblatt, J.: Structure and associated
[114113]DNA–helicase activity of a general transcription initiation factor
[114114]that binds to RNA polymerase II. Nature 341: 410–414, 1989.
[114115]20140.
- [114116]Field, S. J.; Tsai, F.–Y.; Kuo, F.; Zubiaga, A. M.; Kaelin, W.
[114117]G., Jr.; Livingston, D. M.; Orkin, S. H.; Greenberg, M. E.:

E2F-1

[114118]functions in mice to promote apoptosis and suppress proliferation. Cell 85:

[114119]549–561, 1996.

[114120]20141.

[114121]Helin, K.; Lees, J. A.; Vidal, M.; Dyson, N.; Harlow, E.; Fataey,

[114122]A.: A cDNA encoding a pRB-binding protein with properties of the

[114123]transcription factor E2F. Cell 70: 337–350, 1992.

[114124]20142.

[114125]Irwin, M.; Marin, M. C.; Phillips, A. C.; Seelan, R. S.; Smith,

[114126]D. I.; Liu, W.; Flores, E. R.; Tsai, K. Y.; Jacks, T.; Vousden, K.

[114127]H.; Kaelin, W. G., Jr.: Role for the p53 homologue p73 in E2F-1-induced

[114128]apoptosis. Nature 407: 645–648, 2000.

[114129]20143.

[114130]Jacks, T.; Fazeli, A.; Schmitt, E. M.; Bronson, R. T.; Goodell,

[114131]M. A.; Weinberg, R. A.: Effects of an Rb mutation in the

mouse. Nature 359:

[114132]295–300, 1992.

[114133]20144.

[114134]Lees, J. A.; Saito, M.; Valentine, M.; Look, T.; Harlow, E.;
Dyson,

[114135]N.; Helin, K.: The retinoblastoma protein binds to a family
of E2F

[114136]transcription factors. Molec. Cell. Biol. 13: 7813–7825,
1993.

[114137]20145.

[114138]Leone, G.; Sears, R.; Huang, E.; Rempel, R.; Nuckolls, F.;
Park,

[114139]C.–H.; Giangrande, P.; Wu, L.; Saavedra, H. I.; Field, S. J.;
Thompson,

[114140]M. A.; Yang, H.; Fujiwara, Y.; Greenberg, M. E.; Orkin, S.;
Smith,

[114141]C.; Nevins, J. R.: Myc requires distinct E2F activities to in-
duce

[114142]S phase and apoptosis. Molec. Cell 8: 105–113, 2001.

[114143]20146.

[114144]Lissy, N. A.; Davis, P. K.; Irwin, M.; Kaelin, W. G.; Dowdy, S.

[114145]F.: A common E2F-1 and p73 pathway mediates cell death
induced by

[114146]TCR activation. *Nature* 407: 642-645, 2000.

[114147]20147.

[114148]Neuman, E.; Sellers, W. R.; McNeil, J. A.; Lawrence, J. B.;
Kaelin,

[114149]W. G., Jr.: Structure and partial genomic sequence of the
human E2F1

[114150]gene. *Gene* 173: 163-169, 1996.

[114151]20148.

[114152]Nevins, J. R.: The Rb/E2F pathway and cancer. *Hum.*
Molec. Genet. 10:

[114153]699-703, 2001.

[114154]20149.

[114155]Nevins, J. R.: E2F: a link between the Rb tumor suppressor
protein

[114156]and viral oncoproteins. *Science* 258: 424-429, 1992.

[114157]20150.

[114158]Ohtani, K.; DeGregori, J.; Nevins, J. R.: Regulation of the

cyclin

[114159]E gene by transcription factor E2F1. Proc. Nat. Acad. Sci.
92: 12146–12150,

[114160]1995.

[114161]20151.

[114162]Arden, K. C.; Boutin, J.–M.; Djiane, J.; Kelly, P. A.; Cavenee,

[114163]W. K.: The receptors for prolactin and growth hormone are
localized

[114164]in the same region of human chromosome 5. Cytogenet.
Cell Genet. 53:

[114165]161–165, 1990.

[114166]20152.

[114167]Arden, K. C.; Cavenee, W. K.; Boutin, J.–M.; Kelly, P. A.:
The

[114168]genes encoding the receptors for prolactin and growth
hormone map

[114169]to human chromosome 5. (Abstract) Am. J. Hum. Genet.
45 (suppl.):

[114170]A129 only, 1989.

[114171]20153.

[114172]Kelly, A.; Powis, S. H.; Glynne, R.; Radley, E.; Beck, S.;

Trowsdale,

[114173]J.: Second proteasome-related gene in the human MHC class II region. *Nature* 353:

[114174]667–668, 1991.

[114175]20154.

[114176]Martinez, C. K.; Monaco, J. J.: Homology of proteasome subunits

[114177]to a major histocompatibility complex-linked LMP gene. *Nature* 353:

[114178]664–667, 1991.

[114179]20155.

[114180]Petes, T. D.: Meiotic recombination hot spots and cold spots. *Nature*

[114181]*Rev. Genet.* 2: 360–369, 2001.

[114182]20156.

[114183]Van Kaer, L.; Ashton-Rickardt, P. G.; Eichelberger, M.; Gaczynska,

[114184]M.; Nagashima, K.; Rock, K. L.; Goldberg, A. L.; Doherty, P. C.; Tonegawa,

[114185]S.: Altered peptidase and viral-specific T cell response in LMP2

[114186]mutant mice. Immunity 1: 533–541, 1994.

[114187]20157.

[114188]Zhou, P.; Zanelli, E.; Smart, M.; David, C.: Genomic organization

[114189]and tissue expression of mouse proteasome gene Lmp-2. Genomics 16:

[114190]664–668, 1993.

[114191]20158.

[114192]Deng, G. Y.; Muir, A.; Maclaren, N. K.; She, J.-X.: Association

[114193]of LMP2 and LMP7 genes within the major histocompatibility complex

[114194]with insulin-dependent diabetes mellitus: population and family studies. Am.

[114195]J. Hum. Genet. 56: 528–534, 1995.

[114196]20159.

[114197]Bouhassira, E. E.; Schwartz, R. S.; Yawata, Y.; Ata, K.; Kan-

[114198]A.; Qui, J. J.-H.; Nagel, R. L.; Rybicki, A. C.: An alanine-to-threonine

[114199]substitution in protein 4.2 cDNA is associated with a

Japanese form

[114200]of hereditary hemolytic anemia (protein 4.2-Nippon).

Blood 79: 1846-1854,

[114201]1992.

[114202]20160.

[114203]Doolittle, R. F.; Hunkapiller, M. W.; Hood, L. E.; Devare, S.

[114204]G.; Robbins, K. C.; Aaronson, S. A.; Antoniades, H. N.:

Simian sarcoma

[114205]virus onc gene, v-sis, is derived from the gene (or genes)
encoding

[114206]a platelet-derived growth factor. Science 221: 275-277,
1983.

[114207]20161.

[114208]Frolova, L. Y.; Sudomoina, M. A.; Grigorieva, A. Y.; Zi-
novieva,

[114209]O. L.; Kisselev, L. L.: Cloning and nucleotide sequence of
the structural

[114210]gene encoding for human tryptophanyl-tRNA synthetase.
Gene 109:

[114211]291-296, 1991.

[114212]20162.

[114213]Vassart, G.: Personal Communication. Brussels, Belgium
1/15/1992.

[114214]20163.

[114215]Wenger, G. D.: Personal Communication. Columbus, Ohio
8/3/1993.

[114216]20164.

[114217]Abbes, A. P.; Bruggeman, B.; van den Akker, E. L. T.; de
Groot,

[114218]M. R.; Franken, A. A. M.; Drexhage, V. R.; Engel, H.: Identi-
fication

[114219]of two distinct mutations at the same nucleotide position,
concomitantly

[114220]with a novel polymorphism in the vasopressin-neuro-
physin II gene (AVP-NP

[114221]II) in two Dutch families with familial neurohypophyseal
diabetes

[114222]insipidus. Clin. Chem. 46: 1699-1702, 2000.

[114223]20165.

[114224]Bahnsen, U.; Oosting, P.; Swaab, D. F.; Nahke, P.; Richter,
D.;

[114225]Schmale, H.: A missense mutation in the vasopressin-
neurophysin precursor

[114226]gene cosegregates with human autosomal dominant neurohypophyseal diabetes

[114227]insipidus. EMBO J. 11: 19–23, 1992.

[114228]20166.

[114229]Calvo, B.; Bilbao, J. R.; Rodriguez, A.; Rodriguez–Arnao, M. D.;

[114230]Castano, L.: Molecular analysis in familial neurohypophyseal diabetes

[114231]insipidus: early diagnosis of an asymptomatic carrier. J. Clin. Endocr.

[114232]Metab. 84: 3351–3354, 1999.

[114233]20167.

[114234]Calvo, B.; Bilbao, J. R.; Urrutia, I.; Eizaguirre, J.; Gaztam-bide,

[114235]S.; Castano, L.: Identification of a novel nonsense mutation and

[114236]a missense substitution in the vasopressin–neurophysin II gene in

[114237]two Spanish kindreds with familial neurohypophyseal diabetes insipidus. J.

[114238]Clin. Endocr. Metab. 83: 995–997, 1998.

[114239]20168.

[114240]Doherty-Fuller, E.; Copeland, K. C.: Sweat tests in patients
with

[114241]diabetes insipidus. Clin. Pediat. 27: 330-332, 1988.

[114242]20169.

[114243]Evans, D. A. P.; van der Kleij, A. A. M.; Sonnemans, M. A.
F.;

[114244]Burbach, J. P. H.; van Leeuwen, F. W.: Frameshift muta-
tions at two

[114245]hotspots in vasopressin transcripts in post-mitotic neu-
rons. Proc.

[114246]Nat. Acad. Sci. 91: 6059-6063, 1994.

[114247]20170.

[114248]Gagliardi, P. C.; Bernasconi, S.; Repaske, D. R.: Autosomal
dominant

[114249]neurohypophyseal diabetes insipidus associated with a
missense mutation

[114250]encoding gly(23) to val in neurophysin II. J. Clin. Endocr.
Metab. 82:

[114251]3643-3646, 1997.

[114252]20171.

[114253] Ganz, M. B.; Boyarsky, G.; Sterzel, R. B.; Boron, W. F.: Arginine

[114254] Vasopressin enhances pH(i) regulation in the presence of HCO₃⁻ by

[114255] stimulating three acid-base transport systems. Nature 337: 648-651,

[114256] 1989.

[114257] 20172.

[114258] Grant, F. D.; Ahmadi, A.; Hosley, C. M.; Majzoub, J. A.: Two

[114259] novel mutations of the vasopressin gene associated with familial diabetes

[114260] insipidus and identification of an asymptomatic carrier infant. J.

[114261] Clin. Endocr. Metab. 83: 3958-3964, 1998.

[114262] 20173.

[114263] Heppner, C.; Kotzka, J.; Bullmann, C.; Krone, W.; Muller-Wieland,

[114264] D.: Identification of mutations of the arginine vasopressin-neurophysin

[114265] II gene in two kindreds with familial central diabetes insipidus. J.

[114266]Clin. Endocr. Metab. 83: 693–696, 1998.

[114267]20174.

[114268]Ito, M.; Jameson, J. L.; Ito, M.: Molecular basis of autosomal

[114269]dominant neurohypophyseal diabetes insipidus: cellular toxicity caused

[114270]by the accumulation of mutant vasopressin precursors within the endoplasmic

[114271]reticulum. J. Clin. Invest. 99: 1897–1905, 1997.

[114272]20175.

[114273]Ito, M.; Mori, Y.; Oiso, Y.; Saito, H.: A single base substitution

[114274]in the coding region for neurophysin II associated with familial central

[114275]diabetes insipidus. J. Clin. Invest. 87: 725–728, 1991.

[114276]20176.

[114277]Ito, M.; Oiso, Y.; Murase, T.; Kondo, K.; Saito, H.; Chinzei,

[114278]T.; Racchi, M.; Lively, M. O.: Possible involvement of inefficient

[114279]cleavage of preprovasopressin by signal peptidase as a cause for familial

- [114280]central diabetes insipidus. J. Clin. Invest. 91: 2565–2571, 1993.
- [114281]20177.
- [114282]Korbonits, M.; Kaltsas, G.; Perry, L. A.; Putignano, P.; Grossman, A. B.; Besser, G. M.; Trainer, P. J.: The growth hormone secretagogue hexarelin stimulates the hypothalamo–pituitary–adrenal axis via arginine vasopressin. J. Clin. Endocr. Metab. 84: 2489–2495, 1999.
- [114283]20178.
- [114284]Krishnamani, M. R. S.; Phillips, J. A., III; Copeland, K. C.: Detection of a novel arginine vasopressin defect by dideoxy fingerprinting. J. Clin. Endocr. Metab. 77: 596–598, 1993.
- [114285]20179.
- [114286]Land, H.; Schutz, G.; Schmale, H.; Richter, D.: Nucleotide sequence of cloned cDNA encoding bovine arginine vasopressin–neurophysin II precursor. Nature 295: 299–303, 1982.

[114294]20180.

[114295]Mahoney, C. P.; Weinberger, E.; Bryant, C.; Ito, M.; Jameson,

[114296]J. L.; Ito, M.: Effects of aging on vasopressin production in a kindred

[114297]with autosomal dominant neurohypophyseal diabetes insipidus due to

[114298]the delta-E47 neurophysin mutation. J. Clin. Endocr.

Metab. 87:

[114299]870–876, 2002.

[114300]20181.

[114301]McLeod, J. F.; Kovacs, L.; Gaskill, M. B.; Rittig, S.; Bradley,

[114302]G. S.; Robertson, G. L.: Familial neurohypophyseal diabetes insipidus

[114303]associated with a signal peptide mutation. J. Clin. Endocr.

Metab. 77:

[114304]599A–599G, 1993.

[114305]20182.

[114306]Gardner, T. L.; Elston, D. M.; Wotowic, P. J.: A familial dermatofibrosarcoma

[114307]protuberans. J. Am. Acad. Derm. 39: 504–505, 1998.

[114308]20183.

[114309]Groffen, J.; Heisterkamp, N.; Stephenson, J. R.; Geurts van Kessel,

[114310]A.; de Klein, A.; Grosveld, G.; Bootsma, D.: c-sis is translocated

[114311]from chromosome 22 to chromosome 9 in chronic myelocytic leukemia. J.

[114312]Exp. Med. 158: 9-15, 1983.

[114313]20184.

[114314]Hermansson, M.; Nister, M.; Betsholtz, C.; Heldin, C.-H.; Westermark,

[114315]B.; Funa, K.: Endothelial cell hyperplasia in human glioblastoma:

[114316]coexpression of mRNA for platelet-derived growth factor (PDGF) B chain

[114317]and PDGF receptor suggests autocrine growth stimulation. Proc. Nat.

[114318]Acad. Sci. 85: 7748-7752, 1988.

[114319]20185.

[114320]Josephs, S. F.; Dalla-Favera, R.; Gelmann, E. P.; Gallo, R. C.;

[114321]Wong-Staal, F.: 5-prime viral and human cellular sequences corresponding
[114322]to the transforming gene of simian sarcoma virus. Science
219: 503-505,
[114323]1983.
[114324]20186.
[114325]Josephs, S. F.; Guo, C.; Ratner, L.; Wong-Staal, F.: Human
proto-oncogene
[114326]nucleotide sequences corresponding to the transforming
region of simian
[114327]sarcoma virus. Science 223: 487-491, 1984.
[114328]20187.
[114329]Josephs, S. F.; Ratner, L.; Clarke, M. F.; Westin, E. H.; Reitz,
[114330]M. S.; Wong-Staal, F.: Transforming potential of human c-
sis nucleotide
[114331]sequences encoding platelet-derived growth factor. Science 225:
[114332]636-639, 1984.
[114333]20188.
[114334]Kelly, J. D.; Raines, E. W.; Ross, R.; Murray, M. J.: The B
chain

[114335]of PDGF alone is sufficient for mitogenesis. EMBO J. 4:
3399–3405,

[114336]1985.

[114337]20189.

[114338]Kiuru-Kuhlefelt, S.; El-Rifai, W.; Fanburg-Smith, J.; Kere,
J.;

[114339]Miettinen, M.; Knuutila, S.: Concomitant DNA copy number
amplification

[114340]at 17q and 22q in dermatofibrosarcoma protuberans. Cy-
togenet. Cell

[114341]Genet. 92: 192–195, 2001.

[114342]20190.

[114343]Kozak, C. A.; Sears, J. F.; Hoggan, M. D.: Genetic mapping
of

[114344]the mouse proto-oncogene c-sis to chromosome 15. Sci-
ence 221: 867–869,

[114345]1983.

[114346]20191.

[114347]Lindahl, P.; Johansson, B. R.; Leveen, P.; Betsholtz, C.: Per-
icyte

[114348]loss and microaneurysm formation in PDGF-B-deficient

mice. Science 277:

[114349]242–245, 1997.

[114350]20192.

[114351]Owen, A. J.; Pantazis, P.; Antoniades, H. N.: Simian sarcoma

[114352]virus-transformed cells secrete a mitogen identical to platelet-derived

[114353]growth factor. Science 225: 54–56, 1984.

[114354]20193.

[114355]Rao, C. D.; Igarashi, H.; Chiu, I.-M.; Robbins, K. C.; Aaronson,

[114356]S. A.: Structure and sequence of the human c-sis/platelet-derived

[114357]growth factor 2 (SIS/PDGF2) transcriptional unit. Proc. Nat. Acad.

[114358]Sci. 83: 2392–2396, 1986.

[114359]20194.

[114360]Robbins, K. C.; Antoniades, H. N.; Devare, S. G.; Hunkapiller,

[114361]M. W.; Aaronson, S. A.: Structural and immunological similarities

[114362]between simian sarcoma virus gene product(s) and human platelet-derived

[114363]growth factor. *Nature* 305: 605–608, 1983.

[114364]20195.

[114365]Robbins, K. C.; Devare, S. G.; Reddy, E. P.; Aaronson, S. A.:

[114366]In vivo identification of the transforming gene product of simian

[114367]sarcoma virus. *Science* 218: 1131–1133, 1982.

[114368]20196.

[114369]Simon, M.–P.; Navarro, M.; Roux, D.; Pouyssegur, J.: Structural

[114370]and functional analysis of a chimeric protein COL1A1–PDGFB generated

[114371]by the translocation t(17;22)(q22;q13.1) in dermatofibrosarcoma protuberans

[114372](DP). *Oncogene* 20: 2965–2975, 2001.

[114373]20197.

[114374]Smidt, M.; Kirsch, I.; Ratner, L.: Deletion of Alu sequences

[114375]in the fifth c-sis intron in individuals with meningiomas. *J. Clin.*

[114376]Invest. 86: 1151-1157, 1990.

[114377]20198.

[114378]Turc-Carel, C.; Philip, I.; Berger, M. P.; Philip, T.; Lenoir,

[114379]G. M.: Chromosomal translocations in Ewing's sarcoma.

(Letter) New

[114380]Eng. J. Med. 309: 497-498, 1983.

[114381]20199.

[114382]Shimizu, A.; O'Brien, K. P.; Sjoblom, T.; Pietras, K.; Buch-
dunger,

[114383]E.; Collins, V. P.; Heldin, C.-H.; Dumanski, J. P.; Ostman,
A.: The

[114384]dermatofibrosarcoma protuberans-associated collagen
type I-alpha-1/platelet-derived

[114385]growth factor (PDGF) beta-chain fusion gene generates a
transforming

[114386]protein that is processed to functional PDGF-BB. Cancer
Res. 59:

[114387]3719-3723, 1999.

[114388]20200.

[114389]Simon, M.-P.; Pedeutour, F.; Sirvent, N.; Grosgeorge, J.;
Minoletti,

[114390]F.; Coindre, J.-M.; Terrier-Lacombe, M.-J.; Mandahl, N.;
Craver, R.

[114391]D.; Blin, N.; Sozzi, G.; Turc-Carel, C.; O'Brien, K. P.; Kedra,
D.;

[114392]Fransson, I.; Guilbaud, C.; Dumanski, J. P.: Deregulation of
the

[114393]platelet-derived growth factor B-chain gene via fusion
with collagen

[114394]gene COL1A1 in dermatofibrosarcoma protuberans and
giant-cell fibroblastoma. Nature

[114395]Genet. 15: 95-98, 1997.

[114396]20201.

[114397]Waterfield, M. D.; Scrace, G. T.; Whittle, N.; Stroobant, P.;

[114398]Johnsson, A.; Wasteson, A.; Westermarck, B.; Heldin, C.-H.;
Huang,

[114399]J. S.; Deuel, T. F.: Platelet-derived growth factor is struc-
turally

[114400]related to the putative transforming protein p28(sis) of
simian sarcoma

[114401]virus. Nature 304: 35-39, 1983.

[114402]20202.

[114403]Graphodatsky, A.; Frolova, L.; Biltueva, L.; Eremina, V.;
Lushnikova,
[114404]T.; Sudomoina, M.; Zinovieva, O.; Kisselev, L.: Localization
of the
[114405]tryptophanyl tRNA synthetase gene (WARS) on human and
bovine chromosomes
[114406]by in situ hybridization. Mammalian Genome 4: 183–184,
1993.
[114407]20203.
[114408]Jensen, L. L.; Nielsen, M. M.; Justesen, J.; Hansen, L. L.:
Assignment
[114409]of human NADH dehydrogenase (ubiquinone) 1 beta sub-
complex 3 (NDUFB3)
[114410]and of its four pseudogenes to human chromosomes
2q31.3, 1p13.3-p13.1,
[114411]9q32-q34.1, 14q22.3-q23.1 and 14q32.2 by radiation
hybrid mapping. Cytogenet.
[114412]Cell Genet. 93: 147–150, 2001.
[114413]20204.
[114414]Otani, A.; Slike, B. M.; Dorrell, M. I.; Hood, J.; Kinder, K.;
[114415]Ewalt, K. L.; Cheresch, D.; Schimmel, P.; Friedlander, M.: A
fragment

- [114416] of human TrpRS as a potent antagonist of ocular angiogenesis. Proc.
- [114417] Nat. Acad. Sci. 99: 178–183, 2002.
- [114418] 20205.
- [114419] Shimizu, N.; Kucherlapati, R. S.; Ruddle, F. H.: Assignment of
- [114420] a human gene for tryptophanyl-tRNA synthetase to chromosome 14 using
- [114421] human-mouse somatic cell hybrids. Somat. Cell Genet. 2: 345–357,
- [114422] 1976.
- [114423] 20206.
- [114424] Tolstrup, A. B.; Bejder, A.; Fleckner, J.; Justesen, J.: Transcriptional
- [114425] regulation of the interferon-gamma-inducible tryptophanyl-tRNA synthetase
- [114426] includes alternative splicing. J. Biol. Chem. 270: 397–403, 1995.
- [114427] 20207.
- [114428] Turpaev, K. T.; Zakhariev, V. M.; Sokolova, I. V.; Narovlyansky,

- [114429]A. N.; Amchenkova, A. M.; Justesen, J.; Frolova, L. Y.: Alternative
[114430]processing of the tryptophanyl-tRNA synthetase mRNA
from interferon-treated
[114431]human cells. *Europ. J. Biochem.* 240: 732–737, 1996.
[114432]20208.
- [114433]Wakasugi, K.; Slike, B. M.; Hood, J.; Otani, A.; Ewalt, K. L.;
[114434]Friedlander, M.; Cheresch, D. A.; Schimmel, P.: A human
aminoacyl-tRNA
[114435]synthetase as a regulator of angiogenesis. *Proc. Nat. Acad.
Sci.* 99:
[114436]173–177, 2002.
[114437]20209.
- [114438]Comings, D. E.; Comings, B. G.; Dietz, G.; Muhleman, D.:
Localization
[114439]of human tryptophan oxygenase to 4q31: possible relevance to alcoholism,
[114440]depression and Tourette syndrome. (Abstract) *Am. J. Hum.
Genet.* 45
[114441](suppl.): A135 only, 1989.
[114442]20210.

[114443]Comings, D. E.; Dietz, G.; Muhlman, D.: Localization of human
tryptophan oxygenase to 4q25-q31. (Abstract) Cytogenet.
Cell Genet. 51:
[114445]979 only, 1989.
[114446]20211.
[114447]Comings, D. E.; Muhleman, D.; Dietz, G.; Sherman, M.;
Forest, G.
[114448]L.: Sequence of human tryptophan 2,3-dioxygenase
(TDO2): presence
[114449]of a glucocorticoid response-like element composed of a
GTT repeat
[114450]and an intronic CCCCT repeat. Genomics 29: 390-396,
1995.
[114451]20212.
[114452]Comings, D. E.; Muhleman, D.; Dietz, G. W., Jr.; Donlon,
T.: Human
[114453]tryptophan oxygenase localized to 4q31: possible impli-
cations for
[114454]alcoholism and other behavioral disorders. Genomics 9:
301-308,
[114455]1991.

[114456]20213.

[114457]Donlon, T. A.; Muhleman, D.; Dietz, G.; Comings, D. E.;
Spak, D.

[114458]K.: Localization of human tryptophan oxygenase to
4q31-q32 by in

[114459]situ hybridization. (Abstract) Cytogenet. Cell Genet. 51:
992 only,

[114460]1989.

[114461]20214.

[114462]Miller, J. S.; Moxley, G.; Schwartz, L. B.: Cloning and char-
acterization

[114463]of a second complementary DNA for human tryptase. J.
Clin. Invest. 86:

[114464]864-870, 1990.

[114465]20215.

[114466]Miller, J. S.; Westin, E. H.; Schwartz, L. B.: Cloning and
characterization

[114467]of complementary DNA for human tryptase. J. Clin. Invest.
84: 1188-1195,

[114468]1989.

[114469]20216.

- [114470]Vanderslice, P.; Ballinger, S. M.; Tam, E. K.; Goldstein, S. M.;
- [114471]Craig, C. S.; Caughey, G. H.: Human mast cell tryptase: multiple
- [114472]cDNAs and genes reveal a multigene serine protease family. Proc.
- [114473]Nat. Acad. Sci. 87: 3811–3815, 1990.
- [114474]20217.
- [114475]Lewis, W. H.; Yeger, H.: Characterization of the aniridia–Wilms
- [114476]tumor association region of chromosome 11. (Abstract) Cytogenet.
- [114477]Cell Genet. 46: 650, 1987.
- [114478]20218.
- [114479]Mach, B.; Steimle, V.; Reith, W.: MHC class II–deficient combined
- [114480]immunodeficiency: a disease of gene regulation. Immun. Rev. 138:
- [114481]207–221, 1994.
- [114482]20219.
- [114483]Little, M.; Holmes, G.; Bickmore, W.; van Heyningen, V.;

Hastie,

[114484]N.; Wainwright, B.: DNA binding capacity of the WT1 protein is abolished

[114485]by Denys-Drash syndrome WT1 point mutations. Hum. Molec. Genet. 4:

[114486]351-358, 1995.

[114487]20220.

[114488]Little, M. H.; Williamson, K. A.; Mannens, M.; Kelsey, A.; Gosden,

[114489]C.; Hastie, N. D.; van Heyningen, V.: Evidence that WT1 mutations

[114490]in Denys-Drash syndrome patients may act in a dominant-negative fashion. Hum.

[114491]Molec. Genet. 2: 259-264, 1993.

[114492]20221.

[114493]Mannens, M.; Slater, R. M.; Heyting, C.; Blik, J.; de Kraker,

[114494]J.; Coad, N.; de Pagter-Holthuisen, P.; Pearson, P. L.: Molecular

[114495]nature of genetic changes resulting in loss of heterozygosity of chromosome

[114496]11 in Wilms' tumours. Hum. Genet. 81: 41-48, 1988.

[114497]20222.

[114498]Marshall, L. S.; Qureshi, A. R.; DiGeorge, A. M.; Kistenschalmacher,

[114499]M. L.; Punnett, H. H.: Aniridia and the 11p13 deletion.
(Abstract) Am.

[114500]J. Hum. Genet. 34: 74A, 1982.

[114501]20223.

[114502]Matsunaga, E.: Genetics of Wilms' tumor. Hum. Genet. 57:
231-246,

[114503]1981.

[114504]20224.

[114505]McCoy, F. E., Jr.; Franklin, W. A.; Aronson, A. J.; Spargo, B.

[114506]H.: Glomerulonephritis associated with male pseudo-hermaphroditism

[114507]and nephroblastoma. Am. J. Surg. Path. 7: 387-395, 1983.

[114508]20225.

[114509]Meadows, A. T.; Lichtenfeld, J. L.; Koop, C. E.: Wilms' tumor

[114510]in three children of a woman with congenital hemihypertrophy. New

[114511]Eng. J. Med. 291: 23-24, 1974.

[114512]20226.

[114513]Michalopoulos, E. E.; Bevilacqua, P. J.; Stokoe, N.; Powers, V.

[114514]E.; Willard, H. F.; Lewis, W. H.: Molecular analysis of gene deletion

[114515]in aniridia--Wilms tumor association. Hum. Genet. 70: 157-162, 1985.

[114516]20227.

[114517]Miller, R. W.; Fraumeni, J. F., Jr.; Manning, M. D.: Association

[114518]of Wilms' tumor with aniridia, hemihypertrophy and other congenital

[114519]malformations. New Eng. J. Med. 270: 922-927, 1964.

[114520]20228.

[114521]Miyagawa, K.; Kent, J.; Moore, A.; Charlieu, J.-P.; Little, M.

[114522]H.; Williamson, K. A.; Kelsey, A.; Brown, K. W.; Hassam, S.; Briner,

[114523]J.; Hayashi, Y.; Hirai, H.; Yazaki, Y.; van Heyningen, V.; Hastie,

[114524]N. D.: Loss of WT1 function leads to ectopic myogenesis

in Wilms'

[114525]tumour. (Letter) Nature Genet. 18: 15–17, 1998.

[114526]20229.

[114527]Nakagome, Y.; Ise, T.; Sakurai, M.; Nakajo, T.; Okamoto,
E.; Takano,

[114528]T.; Nakahori, Y.; Tsuchida, Y.; Nagahara, N.; Takada, Y.;
Ohsawa,

[114529]Y.; Sawaguchi, S.; Toyosaka, A.; Kobayashi, N.; Matsunaga,
E.; Saito,

[114530]S.: High-resolution studies in patients with aniridia-Wilms
tumor

[114531]association, Wilms tumor or related congenital abnormali-
ties. Hum.

[114532]Genet. 67: 245–248, 1984.

[114533]20230.

[114534]Neidhardt, M.: Wilms-tumor und Aniridie--ein genetisch
fixiertes

[114535]Syndrome? Klin. Paediat. 184: 312–316, 1972.

[114536]20231.

[114537]Nordenskjold, A.; Fricke, G.; Anvret, M.: Absence of muta-
tions

[114538]in the WT1 gene in patients with XY gonadal dysgenesis.

Hum. Genet. 96:

[114539]102–104, 1995.

[114540]20232.

[114541]Orkin, S. H.; Goldman, D. S.; Sallan, S. E.: Development of
homozygosity

[114542]for chromosome 11p markers in Wilms' tumour. Nature
309: 172–174,

[114543]1984.

[114544]20233.

[114545]Park, S.; Bernard, A.; Bove, K. E.; Sens, D. A.; Hazen–Mar-
tin,

[114546]D. J.; Garvin, A. J.; Haber, D. A.: Inactivation of WT1 in
nephrogenic

[114547]rests, genetic precursors to Wilms' tumour. Nature Genet.
5: 363–367,

[114548]1993.

[114549]20234.

[114550]Park, S.; Tomlinson, G.; Nisen, P.; Haber, D. A.: Altered
trans–activational

[114551]properties of a mutated WT1 gene product in a WAGR–

associated Wilms'

[114552]tumor. Cancer Res. 53: 4757-4760, 1993.

[114553]20235.

[114554]Puissant, H.; Azoulay, M.; Serre, J.-L.; Piet, L.; Junien, C.

[114555]: Molecular analysis of a reciprocal translocation

t(5;11)(q11;p13)

[114556]in a WAGR patient. Hum. Genet. 79: 280-282, 1988.

[114557]20236.

[114558]Rahman, N.; Arbour, L.; Tonin, P.; Renshaw, J.; Pelletier, J.;

[114559]Baruchel, S.; Pritchard-Jones, K.; Stratton, M. R.; Narod, S.

A.:

[114560]Evidence for a familial Wilms' tumour gene (FWT1) on

chromosome 17q12-q21. Nature

[114561]Genet. 13: 461-463, 1996.

[114562]20237.

[114563]Reeve, A. E.; Housiaux, P. J.; Gardner, R. J. M.; Chewings,

W.

[114564]E.; Grindley, R. M.; Millow, L. J.: Loss of a Harvey ras allele

in

[114565]sporadic Wilms' tumour. Nature 309: 174-176, 1984.

[114566]20238.

[114567]Riccardi, V. M.; Hittner, H. M.; Strong, L. C.; Fernbach, D.
J.;

[114568]Lebo, R.; Ferrell, R. E.: Wilms tumor with aniridia/iris dys-
plasia

[114569]and apparently normal chromosomes. J. Pediat. 100:
574–577, 1982.

[114570]20239.

[114571]Patek, C. E.; Little, M. H.; Fleming, S.; Miles, C.; Charlieu,

[114572]J.-P.; Clarke, A. R.; Miyagawa, K.; Christie, S.; Doig, J.;
Harrison,

[114573]D. J.; Porteous, D. J.; Brookes, A. J.; Hooper, M. L.; Hastie,
N.

[114574]D.: A zinc finger truncation of murine WT1 results in the
characteristic

[114575]urogenital abnormalities of Denys–Drash syndrome. Proc.
Nat. Acad.

[114576]Sci. 96: 2931–2936, 1999.

[114577]20240.

[114578]Riccardi, V. M.; Sujansky, E.; Smith, A. C.; Francke, U.:
Chromosomal

[114579]imbalance in the aniridia--Wilms' tumor association: 11p
interstitial

[114580]deletion. Pediatrics 61: 604-610, 1978.

[114581]20241.

[114582]Salazar, H.; Kanbour, A.; Burgess, F.: Ultrastructure and
observations

[114583]on the histogenesis of mesotheliomas 'adenomatoid tu-
mors' of the female

[114584]genital tract. Cancer 29: 141-152, 1972.

[114585]20242.

[114586]Schmickel, R. D.: Chromosomal deletions and enzyme de-
ficiencies. J.

[114587]Pediat. 108: 244-246, 1986.

[114588]20243.

[114589]Schroeder, W. T.; Chao, L.-Y.; Dao, D. D.; Strong, L. C.;
Pathak,

[114590]S.; Riccardi, V.; Lewis, W. H.; Saunders, G. F.: Nonrandom
loss of

[114591]maternal chromosome 11 alleles in Wilms tumors. Am. J.
Hum. Genet. 40:

[114592]413-420, 1987.

[114593]20244.

[114594]Schumacher, V.; Schneider, S.; Figge, A.; Wildhardt, G.;
Harms,

[114595]D.; Schmidt, D.; Weirich, A.; Ludwig, R.; Royer-Pokora, B.:
Correlation

[114596]of germ-line mutations and two-hit inactivation of the
WT1 gene with

[114597]Wilms tumors of stromal-predominant histology. Proc.
Nat. Acad. Sci. 94:

[114598]3972-3977, 1997.

[114599]20245.

[114600]Babcock, M.; de Silva, D.; Oaks, R.; Davis-Kaplan, S.; Ji-
ralerspong,

[114601]S.; Montermini, L.; Pandolfo, M.; Kaplan, J.: Regulation of
mitochondrial

[114602]iron accumulation by Yfh1p, a putative homolog of
frataxin. Science 276:

[114603]1709-1712, 1997.

[114604]20246.

[114605]Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: The GAA
triplet-repeat

[114606] expansion in Friedreich ataxia interferes with transcription and may

[114607] be associated with an unusual DNA structure. Am. J. Hum. Genet. 62:

[114608] 111–121, 1998.

[114609] 20247.

[114610] Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: Atypical Friedreich

[114611] ataxia caused by compound heterozygosity for a novel missense mutation

[114612] and the GAA triplet-repeat expansion. (Letter) Am. J. Hum. Genet. 60:

[114613] 1251–1256, 1997.

[114614] 20248.

[114615] Cavadini, P.; Gellera, C.; Patel, P. I.; Isaya, G.: Human frataxin

[114616] maintains mitochondrial iron homeostasis in *Saccharomyces cerevisiae*. Hum.

[114617] Molec. Genet. 9: 2523–2530, 2000.

[114618] 20249.

[114619] MacPherson, A. I. S.; Richmond, J.; Donaldson, G. W. K.;

Muir,

[114620]A. R.: The role of the spleen in congenital spherocytosis.

Am. J.

[114621]Med. 50: 35–41, 1971.

[114622]20250.

[114623]Masera, G.; Mieli, G.; Petrone, M.; Porcelli, P.: Transient
aplastic

[114624]crisis in hereditary spherocytosis. Acta Haemat. 63:
28–31, 1980.

[114625]20251.

[114626]Miraglia del Giudice, E.; Perrotta, S.; Nobili, B.; Specchia,

[114627]C.; d'Urzo, G.; Iolascon, A.: Coinheritance of Gilbert syn-
drome increases

[114628]risk for developing gallstones in patients with hereditary
spherocytosis. Blood 94:

[114629]2259–2262, 1999.

[114630]20252.

[114631]Miwa, S.; Tanaka, K. R.; Valentine, W. N.: Enolase activity
of

[114632]erythrocytes in hereditary spherocytosis. Nature 195:
613–614, 1962.

[114633]20253.

[114634]Mohler, D. N.; Wheby, M. S.: Patients with hereditary spherocytosis

[114635]may have clinically significant iron overload when they are also heterozygous

[114636]for hemochromatosis. Trans. Am. Clin. Climatol. Assoc. 96: 34-40,

[114637]1984.

[114638]20254.

[114639]Mohler, D. N.; Wheby, M. S.: Hemochromatosis heterozygotes may

[114640]have significant iron overload when they also have hereditary spherocytosis. Am.

[114641]J. Med. Sci. 292: 320-324, 1986.

[114642]20255.

[114643]Moiseyev, V. S.; Korovina, E. A.; Polotskaya, E. L.; Poliyan-skaya,

[114644]I. S.; Yazdovsky, V. V.: Hypertrophic cardiomyopathy associated with

[114645]hereditary spherocytosis in three generations of one family.(Letter) Lancet II:

[114646]853–854, 1987.

[114647]20256.

[114648]Morton, N. E.; MacKinney, A. A.; Kosower, N. S.; Schilling, R.

[114649]F.; Gray, M. P.: Genetics of spherocytosis. Am. J. Hum. Genet. 14:

[114650]170–184, 1962.

[114651]20257.

[114652]Motulsky, A. G.; Anderson, R.; Sparkes, R. S.; Huestis, R. H.

[114653]Marrow transplantation in newborn mice with hereditary spherocytosis.

[114654]A model system. Trans. Assoc. Am. Phys. 75: 64–72, 1962.

[114655]20258.

[114656]Nakashima, K.; Yamauchi, K.; Miwa, S.; Fujimura, K.; Mizutani,

[114657]A.; Kuramoto, A.: Glutathione reductase deficiency in a kindred with

[114658]hereditary spherocytosis. Am. J. Hemat. 4: 141–150, 1978.

[114659]20259.

[114660]Ng, J.-P.; Cumming, R. L. C.; Horn, E. H.; Hogg, R. B.:
Hereditary

[114661]spherocytosis revealed by human parvovirus infection.(Letter) Brit.

[114662]J. Haemat. 65: 379-380, 1987.

[114663]20260.

[114664]Nozawa, Y.; Noguchi, T.; Iida, H.; Fukushima, H.; Sekiya,
T.;

[114665]Ito, Y.: Erythrocyte membrane of hereditary spherocytosis:
alteration

[114666]in surface ultrastructure and membrane proteins, as inferred by scanning

[114667]electron microscopy and SDS-disc gel electrophoresis.
Clin. Chim.

[114668]Acta 55: 81-86, 1974.

[114669]20261.

[114670]Okamoto, N.; Wada, Y.; Nakamura, Y.; Nakayama, M.;
Chiyo, H.;

[114671]Murayama, K.; Inoue, T.; Kanzaki, A.; Yawata, Y.; Hirono,
A.; Miwa,

[114672]S.: Hereditary spherocytic anemia with deletion of the
short arm
[114673]of chromosome 8. Am. J. Med. Genet. 58: 225–229, 1995.
[114674]20262.

[114675]Peretz, E.; Hallel–Halevy, D.; Grunwald, M. H.; Halevy, S.:
Hereditary
[114676]spherocytosis with leg ulcers which healed after splenec-
tomy. Europ.
[114677]J. Derm. 7: 527–528, 1997.
[114678]20263.

[114679]Rao, K. R. P.; Patel, A. R.; Anderson, M. J.; Hodgson, J.;
Jones,
[114680]S. E.; Pattison, J. R.: Infection with parvovirus–like agent
and
[114681]aplastic crisis in adults with chronic hemolytic anemia.
Ann. Intern.
[114682]Med. 98: 930–932, 1983.
[114683]20264.

[114684]Reznikoff–Etievant, M. F.; Bonaiti, C.; Maigret, P.;
Malvoisin,
[114685]A.; Maynier, M.; Mesnard, G.; Haupman, G.: Hereditary

spherocytosis

[114686]linkage. Brit. J. Haemat. 46: 153–155, 1980.

[114687]20265.

[114688]Sengar, D. P. S.; McLeish, W. A.; Smiley, R. K.; Luke, B.:

HLA

[114689]and hereditary spherocytosis. Vox Sang. 33: 278–279,
1977.

[114690]20266.

[114691]Shohet, S. B.: Reconstitution of spectrin–deficient spherocytic

[114692]mouse erythrocyte membranes. J. Clin. Invest. 64:
483–494, 1979.

[114693]20267.

[114694]Stratton, R. F.; Crudo, D. F.; Varela, M.; Shapira, E.: Deletion

[114695]of the proximal short arm of chromosome 8. Am. J. Med. Genet. 42:

[114696]15–18, 1992.

[114697]20268.

[114698]Tse, W. T.; Meninger, J.; Ward, D.; John, K.; Lux, S. E.; Forget,

[114699]B. G.: Genomic cloning and chromosomal sublocalization
of the human

[114700]ankyrin gene.(Abstract) Clin. Res. 38: 266A, 1990.

[114701]20269.

[114702]Tsukada, T.; Koike, T.; Koike, R.; Sanada, M.; Takahashi,
M.;

[114703]Shibata, A.; Nunoue, T.: Epidemic of aplastic crisis in pa-
tients

[114704]with hereditary spherocytosis in Japan.(Letter) Lancet 1:
1401,

[114705]1985.

[114706]20270.

[114707]White, R.; Barker, J.: Normoblastosis, a mutant mouse with
severe

[114708]hemolytic anemia.(Abstract) Blood 70S: 57a, 1987.

[114709]20271.

[114710]White, R. A.; Birkenmeier, C. S.; Lux, S. E.; Barker, J. E.:

[114711]Ankyrin and the hemolytic anemia mutation, nb, map to
mouse chromosome

[114712]8: presence of the nb allele is associated with a truncated
erythrocyte

[114713]ankyrin. Proc. Nat. Acad. Sci. 87: 3117–3121, 1990.

[114714]20272.

[114715]Wiley, J. S.: Co-ordinated increase of sodium leak and sodium

[114716]pump in hereditary spherocytosis. Brit. J. Haemat. 22: 529–542,

[114717]1972.

[114718]20273.

[114719]Wiley, J. S.; Firkin, B. G.: An unusual variant of hereditary

[114720]spherocytosis. Am. J. Med. 48: 63–71, 1970.

[114721]20274.

[114722]Zail, S. S.; Krawitz, E.; Viljoen, E.; Kramer, S.; Metz, J.:

[114723]Atypical hereditary spherocytosis: biochemical studies and sites of

[114724]erythrocyte destruction. Brit. J. Haemat. 13: 323–334, 1967.

[114725]20275.

[114726]Sukhatme, V. P.; Vollmer, A. C.; Erikson, J.; Isobe, M.; Croce,

[114727]C.; Parnes, J. R.: Gene for the human T cell differentiation

antigen

[114728]Leu-2/T8 is closely linked to the kappa light chain locus
on chromosome

[114729]2. J. Exp. Med. 161: 429-434, 1985.

[114730]20276.

[114731]Traver, D.; Akashi, K.; Manz, M.; Merad, M.; Miyamoto, T.;
Engleman,

[114732]E. G.; Weissman, I. L.: Development of CD8-alpha-positive
dendritic

[114733]cells from a common myeloid progenitor. Science 290:
2152-2154,

[114734]2000.

[114735]20277.

[114736]Weichhold, G. M.; Huber, C.; Parnes, J. R.; Zachau, H. G.:
The

[114737]CD8-alpha locus is located on the telomere side of the
immunoglobulin-kappa

[114738]locus at a distance of 2 Mb. Genomics 16: 512-514, 1993.

[114739]20278.

[114740]Angel, J. M.; Moore, J. L.; Pelphrey, A.; Richie, E. R.: The
mouse

[114741]homolog of the rhombotin (Ttg-1) gene maps on chromosome 7 distal

[114742]to the beta-globin (Hbb) locus. Mammalian Genome 4: 281-282, 1993.

[114743]20279.

[114744]Boehm, T.; Foroni, L.; Kennedy, M.; Rabbitts, T. H.: The rhombotin

[114745]gene belongs to a class of transcriptional regulators with a potential

[114746]novel protein dimerisation motif. Oncogene 5: 1103-1105, 1990.

[114747]20280.

[114748]Chen, Q.; Cheng, J. T.; Tasi, L. H.; Schneider, N.; Buchanan, G.;

[114749]Carroll, A.; Crist, W.; Ozanne, B.; Siciliano, M. J.; Baer, R.: The

[114750]TAL gene undergoes chromosome translocation in T cell leukemia and

[114751]potentially encodes a helix-loop-helix protein. EMBO J. 9: 415-424,

[114752]1990.

[114753]20281.

[114754]Greenberg, J.; Boehm, T.; Surane, A.; Keynes, R.; Sofronew, M.;

[114755]Rabbitts, T. H.: A human T cell translocation involving chromosome

[114756]11p15 identifies a gene expressed in developing rhombomeres of the

[114757]mouse hind brain. (Abstract) Cytogenet. Cell Genet. 51: 1008 only,

[114758]1989.

[114759]20282.

[114760]Honey, K.; Benlagha, K.; Beers, C.; Forbush, K.; Teyton, L.; Kleijmeer,

[114761]M. J.; Rudensky, A. Y.; Bendelac, A.: Thymocyte expression of cathepsin

[114762]L is essential for NKT cell development. Nature Immun. 7Oct, 2002.

[114763]Note: Advance Electronic Publication.

[114764]20283.

[114765]Adkison, L. R.; Taylor, S.; Beamer, W. G.: Mutant gene-induced

[114766]disorders of structure, function and thyroglobulin synthesis in congenital

[114767]goitre (cog/cog) in mice. J. Endocr. 126: 51–58, 1990.

[114768]20284.

[114769]Avvedimento, V. E.; Di Lauro, R.; Monticelli, A.; Bernardi, F.;

[114770]Patracchini, P.; Calzolari, E.; Martini, G.; Varrone, S.: Mapping

[114771]of human thyroglobulin gene on the long arm of chromosome 8 by in

[114772]situ hybridization. Hum. Genet. 71: 163–166, 1985.

[114773]20285.

[114774]Baas, F.; Bikker, H.; Geurts van Kessel, A.; Melsert, R.; Pearson,

[114775]P. L.; de Vijlder, J. J. M.; van Ommen, G.–J. B.: The human thyroglobulin

[114776]gene: a polymorphic marker localized distal to C–MYC on chromosome

[114777]8 band q24. Hum. Genet. 69: 138–143, 1985.

[114778]20286.

[114779]Baas, F.; van Ommen, G.–J. B.; Bikker, H.; Arnberg, A. C.; de Vijlder,

[114780]J. J. M.: The human thyroglobulin gene is over 300 kb long

and contains

[114781]introns of up to 64 kb. Nucleic Acids Res. 14: 5171–5186,
1986.

[114782]20287.

[114783]Beamer, W. G.; Maltais, L. J.; DeBaets, M. H.; Eicher, E. M.:

[114784]Inherited congenital goiter in mice. Endocrinology 120:
838–840,

[114785]1987.

[114786]20288.

[114787]Berge-Lefranc, J.-L.; Cartouzou, G.; Mattei, M.-G.; Pas-
sage, E.;

[114788]Malezet-Desmoulins, C.; Lissitzky, S.: Localization of the
thyroglobulin

[114789]gene by in situ hybridization to human chromosomes.
Hum. Genet. 69:

[114790]28–31, 1985.

[114791]20289.

[114792]Bernardi, F.; Patracchini, P.; Monticelli, A.; Varrone, S.;

Aiello,

[114793]V.; Calzolari, E.; Marchetti, G.; Avvedimento, V. E.: Human
thyroglobulin

[114794] gene is located on the terminal part of the long arm of chromosome

[114795] 8. (Abstract) Cytogenet. Cell Genet. 40: 582–583, 1985.

[114796] 20290.

[114797] Brocas, H.; Szpirer, J.; Lebo, R. V.; Levan, G.; Szpirer, C.; Cheung,

[114798] M. C.; Vassart, G.: The thyroglobulin gene resides on chromosome

[114799] 8 in man and on chromosome 7 in the rat. Cytogenet. Cell Genet. 39:

[114800] 150–153, 1985.

[114801] 20291.

[114802] Cochaux, P.; Ieiri, T.; Targovnik, H.; Suzuki, M.; Shimoda, S.-I.;

[114803] Perret, J.; Vassart, G.: Identification of a splicing mutation responsible

[114804] for a human hereditary goiter with hypothyroidism. (Abstract) Am.

[114805] J. Hum. Genet. 49 (suppl.): 131 only, 1991.

[114806] 20292.

[114807] Hashimoto, K.; Hirai, M.; Kurosawa, Y.: A gene outside the

human

[114808]MHC related to classical HLA class I genes. Science 269:
693–695,

[114809]1995.

[114810]20293.

[114811]Hashimoto, K.; Hirai, M.; Kurosawa, Y.: Identification of a
mouse

[114812]homolog for the human hereditary haemochromatosis
candidate gene. Biochem.

[114813]Biophys. Res. Commun. 230: 35–39, 1997.

[114814]20294.

[114815]Hickman, P. E.; Hourigan, L. F.; Powell, L. W.; Cordingley,
F.;

[114816]Dimeski, G.; Ormiston, B.; Shaw, J.; Ferguson, W.; Johnson,
M.; Ascough,

[114817]J.; McDonell, K.; Pink, A.; Crawford, D. H. G.: Automated
measurement

[114818]of unsaturated iron binding capacity is an effective
screening strategy

[114819]for C282Y homozygous haemochromatosis. Gut 46:
405–409, 2000.

[114820]20295.

[114821]Hill, W. G.; Robertson, A.: Linkage disequilibrium in finite
[114822]populations. Theor. Appl. Genet. 38: 226–231, 1968.

[114823]20296.

[114824]Jazwinska, E. C.; Cullen, L. M.; Busfield, F.; Pyper, W. R.;
Webb,

[114825]S. I.; Powell, L. W.; Morris, C. P.; Walsh T. P.: Haemochro-
matosis

[114826]and HLA-H. (Letter) Nature Genet. 14: 249–251, 1996.

[114827]20297.

[114828]Jazwinska, E. C.; Lee, S. C.; Webb, S. I.; Halliday, J. W.;
Powell,

[114829]L. W.: Localization of the hemochromatosis gene close to
D6S105. Am.

[114830]J. Hum. Genet. 53: 347–352, 1993.

[114831]20298.

[114832]Jazwinska, E. C.; Pyper, W. R.; Burt, M. J.; Francis, J. L.;
Goldwurm,

[114833]S.; Webb, S. I.; Lee, S. C.; Halliday, J. W.; Powell, L. W.:
Haplotype

[114834]analysis in Australian hemochromatosis patients: evidence
for a predominant

- [114835]ancestral haplotype exclusively associated with
hemochromatosis. Am.
- [114836]J. Hum. Genet. 56: 428–433, 1995.
- [114837]20299.
- [114838]Jeffrey, G. P.; Chakrabarti, S.; Hegele, R. A.; Adams, P. C.:
- [114839]Polymorphism in intron 4 of HFE may cause overestima-
tion of C282Y
- [114840]homozygote prevalence in haemochromatosis. (Letter)
Nature Genet. 22:
- [114841]325–326, 1999.
- [114842]20300.
- [114843]Johnson, G. B., Jr.; Frey, W. G., III: Familial aspects of idio-
pathic
- [114844]hemochromatosis. J.A.M.A. 179: 747–751, 1962.
- [114845]20301.
- [114846]Jouanolle, A.-M.; Yaouanq, J.; Blayau, M.; Perichon, M.;
Fauchet,
- [114847]R.; Font, M.-P.; Le Gall, J.-Y.; David, V.: HLA class I gene
polymorphism
- [114848]in genetic hemochromatosis. Hum. Genet. 85: 279–282,
1990.

[114849]20302.

[114850]Jouanolle, A. M.; Fergelot, P.; Gandon, G.; Yaouanq, J.; Le Gall,

[114851]J. Y.; David, V.: A candidate gene for hemochromatosis: frequency

[114852]of the C282Y and H63D mutations. Hum. Genet. 100: 544–547, 1997.

[114853]

[114854]100. Jouanolle, A. M.; Gandon, G.; Jezequel, P.; Blayau, M.; Champion,

[114855]M. L.; Yaouanq, J.; Mosser, J.; Fergelot, P.; Chauvel, B.; Bouric,

[114856]P.; Carn, G.; Andrieux, N.; Gicquel, I.; Le Gall, J.-Y.; David, V.

[114857]: Haemochromatosis and HLA-H. (Letter) Nature Genet. 14: 251–252,

[114858]1996.

[114859]

[114860]101. Kaikov, Y.; Wadsworth, L. D.; Hassall, E.; Dimmick, J. E.; Rogers,

[114861]P. C. J.: Primary hemochromatosis in children: report of three newly

[114862]diagnosed cases and review of the pediatric literature. Pediatrics 90:

[114863]37–42, 1992.

[114864]

[114865]102. Karlsson, M.; Ikkala, E.; Reunanen, A.; Takkunen, H.; Vuori,

[114866]E.; Makinen, J.: Prevalence of hemochromatosis in Finland. Acta

[114867]Med. Scand. 224: 385–390, 1988.

[114868]

[114869]103. Kidd, K. K.: Genetic linkage and hemochromatosis. (Editorial) New

[114870]Eng. J. Med. 301: 209–210, 1979.

[114871]

[114872]104. Kravitz, K.; Skolnick, M.; Cannings, C.; Carmelli, D.; Baty,

[114873]B.; Amos, B.; Johnson, A.; Mendell, N.; Edwards, C.; Cartwright, G.

[114874]: Genetic linkage between hereditary hemochromatosis and HLA. Am.

[114875]J. Hum. Genet. 31: 601–619, 1979.

[114876]

[114877]105. Krikker, M. A.: A foundation for hemochromatosis.
(Letter) Ann.

[114878]Intern. Med. 97: 782-783, 1982.

[114879]

[114880]106. Kuhl, P.; Kaltwasser, J. P.; Seidl, S.: HLA antigens in
patients

[114881]with idiopathic hemochromatosis (IH). Tissue Antigens 12:
398-401,

[114882]1978.

[114883]

[114884]107. Kushner, J. P.; Skolnick, M. H.; Edwards, C. Q.;
Goldgar, D.;

[114885]Griffen, L. M.; Drummond, C.: Advantageous hemochro-
matosis. (Letter) New

[114886]Eng. J. Med. 319: 1156, 1988.

[114887]

[114888]108. Lalouel, J. M.; Le Mignon, L.; Simon, M.; Fauchet, R.;
Bourel,

[114889]M.; Rao, D. C.; Morton, N. E.: Genetic analysis of idiopathic
hemochromatosis

[114890] using both qualitative (disease status) and quantitative
(serum iron)

[114891] information. Am. J. Hum. Genet. 37: 700–718, 1985.

[114892]

[114893] 109. Lamon, J. M.; Marynick, S. P.; Roseblatt, R.; Donnelly,
S.:

[114894] Idiopathic hemochromatosis in a young female: a case
study and review

[114895] of the syndrome in young people. Gastroenterology 76:
178–183, 1979.

[114896]

[114897] 110. Lebron, J. A.; Bennett, M. J.; Vaughn, D. E.; Chirino, A.
J.;

[114898] Snow, P. M.; Mintier, G. A.; Feder, J. N.; Bjorkman, P. J.:
Crystal

[114899] structure of the hemochromatosis protein HFE and char-
acterization

[114900] of its interaction with transferrin receptor. Cell 93:
111–123,

[114901] 1998.

[114902]

[114903]111. Leggett, B. A.; Halliday, J. W.; Brown, N. N.; Bryant, S.;
Powell,

[114904]L. W.: Prevalence of haemochromatosis amongst asymptomatic
Australians. Brit.

[114905]J. Haemat. 74: 525–530, 1990.

[114906]

[114907]112. Le Mignon, L.; Simon, M.; Fauchet, R.; Edan, G.; Le
Reun, M.;

[114908]Brissot, P.; Genetet, B.; Bourel, M.: An HLA–A1 association
with

[114909]the hemochromatosis allele? Clin. Genet. 24: 171–176,
1983.

[114910]

[114911]113. Levy, J. E.; Montross, L. K.; Cohen, D. E.; Fleming, M.
D.; Andrews,

[114912]N. C.: The C282Y mutation causing hereditary hemochro-
matosis does

[114913]not produce a null allele. Blood 94: 9–11, 1999.

[114914]

[114915]114. Lipinski, M.; Hors, J.; Saleun, J.–P.; Saddi, R.; Passa,
P.;

[114916] Lafaurie, S.; Feingold, N.; Dausset, J.: Idiopathic hemochromatosis:

[114917] linkage with HLA. Tissue Antigens 11: 471–474, 1978.

[114918]

[114919] 115. Lonjou, C.; Collins, A.; Ajioka, R. S.; Jorde, L. B.; Kushner,

[114920] J. P.; Morton, N. E.: Allelic association under map error and recombinational

[114921] heterogeneity: a tale of two sites. Proc. Nat. Acad. Sci. 95: 11366–11370,

[114922] 1998.

[114923]

[114924] 116. Looker, A. C.; Johnson, C. J.: Prevalence of elevated serum

[114925] transferrin saturation in adults in the United States. Ann. Intern.

[114926] Med. 129: 940–945, 1998.

[114927]

[114928] 117. Lord, D. K.; Dunham, I.; Campbell, R. D.; Bomford, A.; Strachan,

[114929] T.; Cox, T. M.: Molecular analysis of the human MHC class

I region

[114930]in hereditary haemochromatosis: a study by pulsed-field gel electrophoresis. Hum.

[114931]Genet. 85: 531–536, 1990.

[114932]

[114933]118. Lucotte, G.: Celtic origin of the C282Y mutation of hemochromatosis. Blood

[114934]Cells Molecules Dis. 24: 433–438, 1998.

[114935]

[114936]119. Lucotte, G.; Coulondre, C.: Association between a 10 kb PvuII

[114937]restriction fragment of genomic DNA with the hemochromatosis gene. Exp.

[114938]Clin. Immunogenet. 3: 219–223, 1986.

[114939]

[114940]120. MacSween, R. N. M.; Scott, A. R.: Hepatic cirrhosis: a clinico-pathological

[114941]review of 520 cases. J. Clin. Path. 26: 936–942, 1973.

[114942]

[114943]121. Maddrey, W. C.; Hamilton, S. R.; Belitsos, N. J.: Famil–

ial hemochromatosis. Johns

[114944]Hopkins Med. J. 144: 66–69, 1979.

[114945]

[114946]122. McGill, J. R.; Naylor, S. L.; Sakaguchi, A. Y.; Moore, C.
M.;

[114947]Boyd, D.; Barrett, K. J.; Shows, T. B.; Drysdale, J. W.: Hu-
man ferritin

[114948]H and L sequences lie on ten different chromosomes.
Hum. Genet. 76:

[114949]66–72, 1987.

[114950]

[114951]123. Mercier, B.; Mura, C.; Ferec, C.: Putting a hold on
'HLA-H.'

[114952](Letter) Nature Genet. 15: 234 only, 1997.

[114953]

[114954]124. Merryweather-Clarke, A. T.; Pointon, J. J.; Shearman,
J. D.;

[114955]Robson, K. J. H.: Global prevalence of putative
haemochromatosis

[114956]mutations. J. Med. Genet. 34: 275–278, 1997.

[114957]

- [114958]125. Merryweather-Clarke, A. T.; Simonsen, H.; Shearman, J. D.; Pointon, J. J.; Norgaard-Pedersen, B.; Robson, K. J. H.: A retrospective anonymous pilot study in screening newborns for HFE mutations in Scandinavian populations. Hum. Mutat. 13: 154-159, 1999.
- [114962]
- [114963]126. Meyer, T. E.; Ballot, D.; Bothwell, T. H.; Green, A.; Derman, D. P.; Baynes, R. D.; Jenkins, T.; Jooste, P. L.; du Toit, E. D.; Jacobs, P. J.: The HLA linked iron loading gene in an Afrikaner population. J. Med. Genet. 24: 348-356, 1987.
- [114967]
- [114968]127. Milman, N.: Hereditary haemochromatosis in Denmark 1950-1985: clinical, biochemical and histological features in 179 patients and 13 preclinical cases. Danish Med. Bull. 38: 385-393, 1991.

[114971]

[114972]128. Milman, N.; Eiberg, H.; Thymann, M.; Fenger, K.:
Transferrin

[114973]subtypes in 51 Danish patients with hereditary
haemochromatosis and

[114974]in 847 normal subjects. Hum. Genet. 88: 475–476, 1992.

[114975]

[114976]129. Milman, N.; Graudal, N.; Nielsen, L. S.; Fenger, K.:
HLA determinants

[114977]in 70 Danish patients with idiopathic haemochromatosis.
Clin. Genet. 33:

[114978]286–292, 1988.

[114979]

[114980]130. Milman, N.; Graudal, N.; Nielsen, L. S.; Fenger, K.: An
HLA

[114981]study in 74 Danish haemochromatosis patients and in 21
of their families. Clin.

[114982]Genet. 41: 6–11, 1992.

[114983]

[114984]131. Milman, N.; Graudal, N.; Nielsen, L. S.; Mathiassen,
B.; Tauris,

- [114985]P.; Lund, B.; Kristensen, J. S.; Fenger, K.: Family studies of hereditary hemochromatosis in Denmark and the Faroe Islands. Hum. Genet. 85: 228–232, 1990.
- [114988]
- [114989]132. Muir, W. A.; McLaren, G. D.; Braun, W.; Askari, A.: Evidence for heterogeneity in hereditary hemochromatosis: evaluation of 174 persons in nine families. Am. J. Med. 76: 806–814, 1984.
- [114992]
- [114993]133. Muir, W. A.; McLaren, G. D.; Braun, W. E.; Askari, A. K.: Hereditary hemochromatosis: dominant inheritance and probable linkage to HLA. (Abstract) Am. J. Hum. Genet. 30: 61A, 1978.
- [114996]
- [114997]134. Mura, C.; Le Gac, G.; Scotet, V.; Ragueneas, O.; Mercier, A.-Y.; Ferec, C.: Variation of iron loading expression in C282Y

homozygous

[114999]haemochromatosis probands and sib pairs. (Letter) J. Med. Genet. 38:

[115000]632–636, 2001.

[115001]

[115002]135. Mura, C.; Nousbaum, J.-B.; Verger, P.; Moalic, M.-T.; Raguenees,

[115003]O.; Mercier, A.-Y.; Ferec, C.: Phenotype–genotype correlation in

[115004]haemochromatosis subjects. Hum. Genet. 101: 271–276, 1997.

[115005]

[115006]136. Mura, C.; Raguenees, O.; Ferec, C.: HFE mutations analysis in

[115007]711 hemochromatosis probands: evidence for S65C implication in mild

[115008]form of hemochromatosis. Blood 93: 2502–2505, 1999.

[115009]

[115010]137. Murphy, P. A.: Personal Communication. Baltimore, Md. 12/12/1987.

[115011]

[115012]138. Niederau, C.; Fischer, R.; Sonnenberg, A.; Stremmel, W.; Trampisch,

[115013]H. J.; Strohmeyer, G.: Survival and causes of death in cirrhotic

[115014]and in noncirrhotic patients with primary hemochromatosis. New Eng.

[115015]J. Med. 313: 1256–1262, 1985.

[115016]

[115017]139. Noll, W. W.; Belloni, D. R.; Stenzel, T. T.; Grody, W. W.: Polymorphism

[115018]in intron 4 of HFE does not compromise haemochromatosis mutation results

[115019](Letter) Nature Genet. 23: 271–272, 1999.

[115020]

[115021]140. Nussbaumer, T.; Plattner, H. C.; Rywlin, A. M.: Hemochromatose

[115022]juvenile chez trois soeurs et un frere avec consanguinite des parents:

[115023]etude anatomoclinique et genetique du syndrome endocrinohepato-myocardique. J.

[115024]Genet. Hum. 1: 53–59, 1952.

[115025]

[115026]141. Olsson, K. S.; Eriksson, K.; Ritter, B.; Heedman, P. A.:
Screening
[115027]for iron overload using transferrin saturation. Acta Med.
Scand. 215:
[115028]105–112, 1984.

[115029]

[115030]142. Olsson, K. S.; Ritter, B.; Rosen, U.; Heedman, P. A.;
Staugard,
[115031]F.: Prevalence of iron overload in central Sweden. Acta
Med. Scand. 213:
[115032]145–150, 1983.

[115033]

[115034]143. Olynyk, J. K.; Cullen, D. J.; Aquilia, S.; Rossi, E.; Sum-
merville,
[115035]L.; Powell, L. W.: A population-based study of the clinical
expression
[115036]of the hemochromatosis gene. New Eng. J. Med. 341:
718–724, 1999.

[115037]

[115038]144. Parkkila, S.; Waheed, A.; Britton, R. S.; Bacon, B. R.;

Zhou,

[115039]X. Y.; Tomatsu, S.; Fleming, R. E.; Sly, W. S.: Association of
the

[115040]transferrin receptor in human placenta with HFE, the pro-
tein defective

[115041]in hereditary hemochromatosis. Proc. Nat. Acad. Sci. 94:
13198-13202,

[115042]1997.

[115043]

[115044]145. Parkkila, S.; Waheed, A.; Britton, R. S.; Feder, J. N.;
Tsuchihashi,

[115045]Z.; Schatzman, R. C.; Bacon, B. R.; Sly, W. S.: Immunohis-
tochemistry

[115046]of HLA-H, the protein defective in patients with hereditary
hemochromatosis,

[115047]reveals unique pattern of expression in gastrointestinal
tract. Proc.

[115048]Nat. Acad. Sci. 94: 2534-2539, 1997.

[115049]

[115050]146. Perkins, K. W.; McInnes, I. W. S.; Blackburn, C. R. B.;
Beal,

[115051]R. W.: Idiopathic hemochromatosis in children: report of a

family. Am.

[115052]J. Med. 39: 118–126, 1965.

[115053]

[115054]147. Phatak, P. D.; Sham, R. L.; Raubertas, R. F.; Dunnigan, K.; O'Leary,

[115055]M. T.; Braggins, C.; Cappuccio, J. D.: Prevalence of hereditary hemochromatosis

[115056]in 16,031 primary care patients. Ann. Intern. Med. 129: 954–961,

[115057]1998.

[115058]

[115059]148. Pollycove, M.: Hemochromatosis. In: Stanbury, J. B.; Wyngaarden,

[115060]J. B.; Fredrickson, D. S.: The Metabolic Basis of Inherited Disease.

[115061]New York: McGraw–Hill (pub.) (3rd ed.): 1972. Pp. 1051–1084.

[115062]

[115063]149. Poullis, A.; Moodie, S. J.; Maxwell, J. D.: Clinical haemochromatosis

[115064]in HFE mutation carriers. Lancet 360: 411–412, 2002.

[115065]

[115066]150. Powell, L. W.; Ferluga, J.; Halliday, J. W.; Bassett, M. L.;

[115067]Kohonen-Corish, M.; Serjeantson, S.: Genetic hemochromatosis and

[115068]HLA linkage. Hum. Genet. 77: 55-56, 1987.

[115069]

[115070]151. Powell, L. W.; George, D. K.; McDonnell, S. M.; Kowdley, K. V.

[115071]: Diagnosis of hemochromatosis. Ann. Intern. Med. 129: 925-931,

[115072]1998.

[115073]

[115074]152. Pozzato, G.; Zorat, F.; Nascimben, F.; Gregorutti, M.; Comar,

[115075]C.; Baracetti, S.; Vatta, S.; Bevilacqua, E.; Begnano, A.; Crovella,

[115076]S.; Amoroso, A.: Haemochromatosis gene mutations in a clustered Italian

[115077]population: evidence of high prevalence in people of Celtic ancestry. Europ.

[115078]J. Hum. Genet. 9: 445–451, 2001.

[115079]

[115080]153. Rabinovitz, M.; Gavalier, J. S.; Kelly, R. H.; Van Thiel, D. H.

[115081]: Association between heterozygous alpha-1-antitrypsin deficiency

[115082]and genetic hemochromatosis. Hepatology 16: 145–148, 1992.

[115083]

[115084]154. Raha-Chowdhury, R.; Bowen, D. J.; Worwood, M.: A new highly

[115085]polymorphic marker in the 5-prime untranslated region of HLA-F shows

[115086]strong allelic association with haemochromatosis. Hum. Genet. 97:

[115087]228–231, 1996.

[115088]

[115089]155. Rhodes, D. A.; Raha-Chowdhury, R.; Cox, T. M.; Trowsdale, J.

[115090]: Homozygosity for the predominant Cys282Tyr mutation and absence

[115091] of disease expression in hereditary haemochromatosis. J.

Med. Genet. 34:

[115092] 761–764, 1997.

[115093]

[115094] 156. Risch, N.: Haemochromatosis, HFE and genetic complexity. (Letter) Nature

[115095] Genet. 17: 375–376, 1997.

[115096]

[115097] 157. Ritter, B.; Safwenberg, J.; Olsson, K. S.: HLA as a marker of

[115098] the hemochromatosis gene in Sweden. Hum. Genet. 68: 62–66, 1984.

[115099]

[115100] 158. Robson, K. J. H.; Shearman, J. D.; Merryweather–Clarke, A. T.;

[115101] Pointon, J. J.; Rosenberg, W. M.; Walker, A. P.; Dooley, J. S.; Bomford,

[115102] A.; Raha–Chowdhury, R.; Worwood, M.: Haemochromatosis: a gene at

[115103] last? J. Med. Genet. 34: 148–151, 1997.

[115104]

- [115105]159. Rochette, J.; Pointon, J. J.; Fisher, C. A.; Perera, G.; Arambepola, M.; Kodikara Arichchi, D. S.; De Silva, S.; Vandwalle, J. L.; Monti, J. P.; Old, J. M.; Merryweather-Clarke, A. T.; Weatherall, D. J.; Robson, K. J. H.: Multicentric origin of hemochromatosis gene (HFE) mutations. *Am. J. Hum. Genet.* 64: 1056–1062, 1999.
- [115110]
- [115111]160. Roldan, F. P.; Echenagusia, A. A.; Carro, P. G.: Hemochromatosis presenting as acute liver failure after iron supplementation. (Letter) *New Eng. J. Med.* 339: 269–270, 1998.
- [115114]
- [115115]161. Roth, M.-P.; Giraldo, P.; Hariti, G.; Poloni, E. S.; Sanchez-Mazas, A.; De Stefano, G. F.; Dugoujon, J.-M.; Coppin, H.: Absence of the hemochromatosis gene Cys282Tyr mutation in three ethnic groups from

[115118]Algeria (Mzab), Ethiopia, and Senegal. Immunogenetics
46: 222–225,

[115119]1997.

[115120]

[115121]162. Rothenberg, B. E.; Volland, J. R.: Beta-2 knockout
mice develop

[115122]parenchymal iron overload: a putative role for class I
genes of the

[115123]major histocompatibility complex in iron metabolism.
Proc. Nat. Acad.

[115124]Sci. 93: 1529–1534, 1996.

[115125]

[115126]163. Rowe, J. W.; Wands, J. R.; Mezey, S. E.; Waterbury, L.
A.; Wright,

[115127]J. R.; Tobin, J.; Andres, R.: Familial hemochromatosis:
characteristics

[115128]of the precirrhotic stage in a large kindred. Medicine 56:
197–211,

[115129]1977.

[115130]

[115131]164. Roy, C. N.; Andrews, N. C.: Recent advances in disor–

ders of

[115132]iron metabolism: mutations, mechanisms and modifiers.

Hum. Molec.

[115133]Genet. 10: 2181–2186, 2001.

[115134]

[115135]165. Roy, C. N.; Penny, D. M.; Feder, J. N.; Enns, C. A.: The
hereditary

[115136]hemochromatosis protein, HFE, specifically regulates
transferrin-mediated

[115137]iron uptake in HeLa cells. J. Biol. Chem. 274: 9022–9028,
1999.

[115138]

[115139]166. Ryan, E.; O'Keane, C.; Crowe, J.: Hemochromatosis in
Ireland

[115140]and HFE. Blood Cells Molecules Dis. 24: 428–432, 1998.

[115141]

[115142]167. Saddi, R.; Feingold, J.: Idiopathic haemochromatosis:
an autosomal

[115143]recessive disease. Clin. Genet. 5: 234–241, 1974.

[115144]

[115145]168. Sargent, T., III; Lim, T. H.; Jenson, R. L.: Reduced

chromium

[115146]retention in patients with hemochromatosis, a possible
basis of hemochromatotic

[115147]diabetes. Metabolism 28: 70–79, 1979.

[115148]

[115149]169. Sheldon, J. H.: Haemochromatosis. London: Oxford
Univ. Press

[115150](pub.) 1935. Pp. 19 only.

[115151]

[115152]170. Siemons, L. J.; Mahler, C.: Hypogonadotropic hypog-
onadism in

[115153]hemochromatosis: recovery of reproductive function after
iron depletion. J.

[115154]Clin. Endocr. Metab. 65: 585–587, 1987.

[115155]

[115156]171. Simon, M.; Alexandre, J. L.; Bourel, M.; Le Marec, B.;
Scordia,

[115157]C.: Heredity of idiopathic haemochromatosis: a study of
106 families. Clin.

[115158]Genet. 11: 327–341, 1977.

[115159]

[115160]172. Simon, M.; Bourel, M.; Fauchet, R.; Genetet, B.: Association

[115161]of HLA-A3 and HLA-B14 antigens with idiopathic
haemochromatosis. Gut 17:

[115162]332-334, 1976.

[115163]

[115164]173. Simon, M.; Bourel, M.; Genetet, B.; Fauchet, R.:

Heredity of

[115165]idiopathic haemochromatosis. (Letter) Lancet I: 706, 1977.

[115166]

[115167]174. Simon, M.; Bourel, M.; Genetet, B.; Fauchet, R.: Idio-
pathic

[115168]hemochromatosis: demonstration of recessive transmis-
sion and early

[115169]detection by family HLA typing. New Eng. J. Med. 297:
1017-1021,

[115170]1977.

[115171]

[115172]175. Simon, M.; Fauchet, R.; Hespel, J. P.; Beaumont, C.;
Brissot,

[115173]P.; Hary, B.; De Nercy, H. Y. H.; Genetet, B.; Bourel, M.: Id-
iopathic

[115174]hemochromatosis: a study of biochemical expression in
247 heterozygous

[115175]members of 63 families: evidence for a single major HLA-
linked gene. Gastroenterology 78:

[115176]703–708, 1980.

[115177]

[115178]176. Simon, M.; Fauchet, R.; Le Gall, J. Y.; Brissot, P.;
Bourel,

[115179]M.: Immunogenetics of idiopathic hemochromatosis and
secondary iron

[115180]overload.In: Farid, N. R.: Immunogenetics of Endocrine
Disorders.

[115181]New York: Alan R. Liss (pub.) 1988. Pp. 345–371.

[115182]

[115183]177. Simon, M.; Le Mignon, L.; Fauchet, R.; Yaouanq, J.;
David, V.;

[115184]Edan, G.; Bourel, M.: A study of 609 HLA haplotypes
marking for the

[115185]hemochromatosis gene: (1) mapping of the gene near the
HLA-A locus

[115186]and characters required to define a heterozygous popula-
tion and (2)

[115187]hypothesis concerning the underlying cause of hemochro-
matosis-HLA

[115188]association. Am. J. Hum. Genet. 41: 89–105, 1987.

[115189]

[115190]178. Skolnick, M.: Personal Communication. Salt Lake City,
Utah

[115191]9/27/1983.

[115192]

[115193]179. Sorsby, A.: Clinical Genetics. St. Louis: C. V. Mosby
(pub.)

[115194]1953. Pp. 206 only.

[115195]

[115196]180. Sproule, T. J.; Jazwinska, E. C.; Britton, R. S.; Bacon,
B. R.;

[115197]Fleming, R. E.; Sly, W. S.; Roopenian, D. C.: Naturally vari-
ant autosomal

[115198]and sex-linked loci determine the severity of iron over-
load in beta-2-microglobulin-deficient

[115199]mice. Proc. Nat. Acad. Sci. 98: 5170-5174, 2001.

[115200]

[115201]181. Stevens, F. M.; Walters, J. M.; Watt, D. W.; McCarthy,
C. F.

[115202]: Inheritance of idiopathic haemochromatosis. (Letter)
Lancet I:

[115203]1107, 1977.

[115204]

[115205]182. Summers, K. M.; Tam, K. S.; Bartley, P. B.; Drysdale, J.; Zoghbi,

[115206]H. Y.; Halliday, J. W.; Powell, L. W.: Fine mapping of a human chromosome

[115207]6 ferritin heavy chain pseudogene: relevance to haemochromatosis. Hum.

[115208]Genet. 88: 175–178, 1991.

[115209]

[115210]183. Summers, K. M.; Tam, K. S.; Halliday, J. W.; Powell, L. W.:

[115211]HLA determinants in an Australian population of hemochromatosis patients

[115212]and their families. Am. J. Hum. Genet. 45: 41–48, 1989.

[115213]

[115214]184. Thenie, A. C.; Gicquel, I. M.; Hardy, S.; Ferran, H.; Fergelot,

[115215]P. Le Gall, J.-Y.; Mosser, J.: Identification of an endogenous RNA

[115216]transcribed from the antisense strand of the HFE gene.

Hum. Molec.

[115217]Genet. 10: 1859–1866, 2001.

[115218]

[115219]185. The UK Haemochromatosis Consortium: A simple genetic test

[115220]identifies 90% of UK patients with haemochromatosis. Gut 41: 841–844,

[115221]1997.

[115222]

[115223]186. Totaro, A.; Rommens, J. M.; Grifa, A.; Lunardi, C.; Carella,

[115224]M.; Huizenga, J. J.; Roetto, A.; Camaschella, C.; De Sandre, G.; Gasparini,

[115225]P.: Hereditary hemochromatosis: generation of a transcription map

[115226]within a refined and extended map of the HLA class I region. Genomics 31:

[115227]319–326, 1996.

[115228]

[115229]187. Townsend, A.; Drakesmith, H.: Role of HFE in iron metabolism,

[115230]hereditary haemochromatosis, anaemia of chronic disease, and secondary

[115231]iron overload. Lancet 359: 786–790, 2002.

[115232]

[115233]188. Trousseau, A.: Clinical Medicine Lectures Delivered at the Hotel-Dieu

[115234]Paris by A. Trousseau. Philadelphia: Blakiston (pub.) II: 1882.

[115235]pp. 313–315. Note: Translation: 3rd revised and enlarged edition by

[115236]Sir John Rose Cormack and P. Victor Bazire.....

[115237]

[115238]189. Trousseau, A.: Clinique Med. de l'Hotel de Paris. (pub.)

[115239]II: 1865. Pp. 663–698.

[115240]

[115241]190. Valberg, L. S.; Lloyd, D. A.; Ghent, C. N.; Flanagan, P. R.;

[115242]Sinclair, N. R.; Stiller, C. R.; Chamberlain, M. J.: Clinical and

[115243]biochemical expression of the genetic abnormality in idio-

pathic hemochromatosis. Gastroenterology 79:

[115244]884–892, 1980.

[115245]

[115246]191. Waheed, A.; Parkkila, S.; Saarnio, J.; Fleming, R. E.;
Zhou,

[115247]X. Y.; Tomatsu, S.; Britton, R. S.; Bacon, B. R.; Sly, W. S.:
Association

[115248]of HFE protein with transferrin receptor in crypt entero-
cytes of human

[115249]duodenum. Proc. Nat. Acad. Sci. 96: 1579–1584, 1999.

[115250]

[115251]192. Waheed, A.; Parkkila, S.; Zhou, X. Y.; Tomatsu, S.;
Tsuchihashi,

[115252]Z.; Feder, J. N.; Schatzman, R. C.; Britton, R. S.; Bacon, B.
R.;

[115253]Sly, W. S.: Hereditary hemochromatosis: effects of C282Y
and H63D

[115254]mutations on association with beta-2-microglobulin, in-
tracellular

[115255]processing, and cell surface expression of the HFE protein
in COS-7

[115256]cells. Proc. Nat. Acad. Sci. 94: 12384–12389, 1997.

[115257]

[115258]193. Wands, J. R.; Rowe, J. A.; Mezey, S. E.; Waterbury, L. A.; Wright,

[115259]J. R.; Halliday, J. W.; Isselbacher, K. J.; Powell, L. W.: Normal

[115260]serum ferritin concentrations in precirrhotic hemochromatosis. New

[115261]Eng. J. Med. 294: 302-305, 1976.

[115262]

[115263]194. Willis, G.; Wimperis, J. Z.; Lonsdale, R.; Fellows, I. W.; Watson,

[115264]M. A.; Skipper, L. M.; Jennings, B. A.: Incidence of liver disease

[115265]in people with HFE mutations. Gut 46: 401-404, 2000.

[115266]

[115267]195. Worwood, M.; Darke, C.; Trenchard, P.: Hereditary haemochromatosis

[115268]and blood donation. (Letter) Brit. Med. J. 302: 593, 1991.

[115269]

[115270]196. Yaouang, J.; Perichon, M.; Chorney, M.; Pontarotti, P.; Le Treut,

- [115271]A.; El Kahloun, A.; Mauvieux, V.; Blayau, M.; Jouanolle, A.
M.; Chauvel,
- [115272]B.; Moirand, R.; Nouel, O.; Le Gall, J. Y.; Feingold, J.; David,
V.
- [115273]: Anonymous marker loci within 400 kb of HLA-A gener-
ate haplotypes
- [115274]in linkage disequilibrium with the hemochromatosis gene
(HFE). Am.
- [115275]J. Hum. Genet. 54: 252-263, 1994.
- [115276]
- [115277]197. Yaouanq, J.; El Kahloun, A.; Chorney, M.; Jouanolle,
A. M.; Mauvieux,
- [115278]V.; Perichon, M.; Blayau, M.; Pontarotti, P.; Le Gall, J. Y.;
David,
- [115279]V.: Familial screening for genetic haemochromatosis by
means of DNA
- [115280]markers. J. Med. Genet. 29: 320-322, 1992.
- [115281]
- [115282]198. Zappone, E.; Dugast, I.; Papadopoulos, P.; Theriault,
K.; David,
- [115283]V.; LeGall, J.-Y.; Summers, K.; Powell, L.; Drysdale, J.:
Polymorphism

[115284]in a ferritin H gene from chromosome 6p. Hum. Genet.

86: 557-561,

[115285]1991.

[115286]

[115287]199. Zhou, X. Y.; Tomatsu, S.; Fleming, R. E.; Parkkila, S.;

Waheed,

[115288]A.; Jiang, J.; Fei, Y.; Brunt, E. M.; Ruddy, D. A.; Prass, C. E.;

[115289]Schatzman, R. C.; O'Neill, R.; Britton, R. S.; Bacon, B. R.;

Sly,

[115290]W. S.: HFE gene knockout produces mouse model of

hereditary hemochromatosis. Proc.

[115291]Nat. Acad. Sci. 95: 2492-2497, 1998.

[115292]

[115293]200. Zoller, H.; Pietrangelo, A.; Vogel, W.; Weiss, G.: Duo-

denal

[115294]metal-transporter (DMT-1, NRAMP-2) expression in pa-

tients with hereditary

[115295]haemochromatosis. Lancet 353: 2120-2123, 1999.

[115296]20303.

[115297]Azarnia, R.; Reddy, S.; Kmiecik, T. E.; Shalloway, D.;

Loewenstein,

- [115298]W. R.: The cellular src gene product regulates junctional cell-to-cell communication. Science 239: 398–401, 1988.
- [115300]20304.
- [115301]Czernilofsky, A. P.; Levinson, A. D.; Varmus, H. E.; Bishop, J.
- [115302]M.; Tischer, E.; Goodman, H.: Correction to the nucleotide sequence of the src gene of Rous sarcoma virus. Nature 301: 736–738, 1983.
- [115304]20305.
- [115305]Gibbs, C. P.; Tanaka, A.; Anderson, S. K.; Radul, J.; Baar, J.;
- [115306]Ridgway, A.; Kung, H.-J.; Fujita, D. J.: Isolation and structural mapping of a human c-src gene homologous to the transforming gene (v-src) of Rous sarcoma virus. J. Virol. 53: 19–24, 1985.
- [115309]20306.
- [115310]Irby, R. B.; Mao, W.; Coppola, D.; Kang, J.; Loubeau, J. M.; Trudeau, W.; Karl, R.; Fujita, D. J.; Jove, R.; Yeatman, T. J.: Activating

[115312]SRC mutation in a subset of advanced human colon cancers. Nature

[115313]Genet. 21: 187-190, 1999.

[115314]20307.

[115315]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J. D.:

[115316]c-src is consistently conserved in the chromosomal deletion (20q)

[115317]observed in myeloid disorders. Proc. Nat. Acad. Sci. 82: 6692-6696,

[115318]1985.

[115319]20308.

[115320]Lowe, C.; Yoneda, T.; Boyce, B. F.; Chen, H.; Mundy, G. R.; Soriano,

[115321]P.: Osteopetrosis in Src-deficient mice is due to an autonomous defect

[115322]of osteoclasts. Proc. Nat. Acad. Sci. 90: 4485-4489, 1993.

[115323]20309.

[115324]Morris, C. M.; Honeybone, L. M.; Hollings, P. E.; Fitzgerald,

[115325]P. H.: Localization of the SRC oncogene to chromosome band 20q11.2

[115326]and loss of this gene with deletion (20q) in two leukemic patients. Blood 74:

[115327]1768–1773, 1989.

[115328]20310.

[115329]Sakaguchi, A. Y.; Mohandas, T.; Naylor, S. L.: A human c–src

[115330]gene resides on the proximal long arm of chromosome 20 (cen–q13.1). Cancer

[115331]Genet. Cytogenet. 18: 123–129, 1985.

[115332]20311.

[115333]Sakaguchi, A. Y.; Naylor, S. L.; Weinberg, R. A.; Shows, T. B.

[115334]: Organization of human proto–oncogenes. (Abstract) Am. J. Hum. Genet. 34:

[115335]175A, 1982.

[115336]20312.

[115337]Sakaguchi, A. Y.; Zabel, B. U.; Grzeschik, K. H.; Law, M. L.;

[115338]Naylor, S. L.: Human proto–oncogene assignments. (Abstract) Cytogenet.

[115339]Cell Genet. 37: 572–573, 1984.

[115340]20313.

[115341]Wong, B. R.; Besser, D.; Kim, N.; Arron, J. R.; Vologodskaia,

[115342]M.; Hanafusa, H.; Choi, Y.: TRANCE, a TNF family member,
activates

[115343]Akt/PKB through a signaling complex involving TRAF6 and
c-Src. Molec.

[115344]Cell 4: 1041–1049, 1999.

[115345]20314.

[115346]Starzl, T. E.; Demetris, A. J.; Trucco, M.; Ricordi, C.; Ild-
stad,

[115347]S.; Terasaki, P. I.; Murase, N.; Kendall, R. S.; Kocova, M.;
Rudert,

[115348]W. A.; Zeevi, A.; Van Thiel, D.: Chimerism after liver trans-
plantation

[115349]for type IV glycogen storage disease and type 1 Gaucher's
disease. New

[115350]Eng. J. Med. 328: 745–749, 1993.

[115351]20315.

[115352]Trend, P. St. J.; Wiles, C. M.; Spencer, G. T.; Morgan-
Hughes,

[115353]J. A.; Lake, B. D.; Patrick, A. D.: Acid maltase deficiency in

adults:

[115354]diagnosis and management in five cases. Brain 108:
845–860, 1985.

[115355]20316.

[115356]Wokke, J. H. J.; Ausems, M. G. E. M.; van den Boogaard,
M.–J.

[115357]H.; Ippel, E. F.; van Diggelen, O.; Kroos, M. A.; Boer, M.;
Jennekens,

[115358]F. G. I.; Reuser, A. J. J.; Ploos van Amstel, H. K.: Genotype–
phenotype

[115359]correlation in adult–onset acid maltase deficiency. Ann.
Neurol. 38:

[115360]450–454, 1995.

[115361]20317.

[115362]Bao, Y.; Dawson, T. L., Jr.; Chen, Y.–T.: Human glycogen
debranching

[115363]enzyme gene (AGL): complete structural organization and
characterization

[115364]of the 5–prime flanking region. Genomics 38: 155–165,
1996.

[115365]20318.

[115366]Brunberg, J. A.; McCormick, W. F.; Schochet, S. S., Jr.: Type

[115367]III glycogenosis. An adult with diffuse weakness and muscle wasting. Arch.

[115368]Neurol. 25: 171-178, 1971.

[115369]20319.

[115370]Chen, Y.-T.; He, J.-K.; Ding, J.-H.; Brown, B. I.: Glycogen debranching

[115371]enzyme: purification, antibody characterization, and immunoblot analyses

[115372]of type III glycogen storage disease. Am. J. Hum. Genet. 41: 1002-1015,

[115373]1987.

[115374]20320.

[115375]Chen, Y. T.; Burchell, A.: Glycogen storage diseases. In: Scriver,

[115376]C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds.): The Metabolic

[115377]and Molecular Bases of Inherited Disease. New York: McGraw-Hill

[115378](7th ed.): 1995. Pp. 935-965.

[115379]20321.

[115380]Cohen, J.; Friedman, M.: Renal tubular acidosis associated

with

[115381]type III glycogenosis. *Acta Paediat. Scand.* 68: 779–782,
1979.

[115382]20322.

[115383]Cohn, J.; Wang, P.; Hauge, M.; Henningsen, K.; Jensen, B.;
Svejgaard,

[115384]A.: Amylo-1,6-glucosidase deficiency (glycogenosis type
III) in the

[115385]Faroe Island. *Hum. Hered.* 25: 115–126, 1975.

[115386]20323.

[115387]Coleman, R. A.; Winter, H. S.; Wolf, B.; Chen, Y.-T.: Glyco-
gen

[115388]debranching enzyme deficiency: long-term study of
serum enzyme activities

[115389]and clinical features. *J. Inherit. Metab. Dis.* 15: 869–881,
1992.

[115390]20324.

[115391]Confino, E.; Pauzner, D.; Lidor, A.; Yedwab, G.; David, M.:
Pregnancy

[115392]associated with amylo-1,6-glucosidase deficiency (Forbes'
disease):

[115393]case report. *Brit. J. Obstet. Gynaec.* 91: 494–497, 1984.

[115394]20325.

[115395]DiMauro, S.; Hartwig, G. B.; Hays, A.; Eastwood, A. B.;
Franco,

[115396]R.; Olarte, M.; Chang, M.; Roses, A. D.; Fetell, M.; Schoen-
feldt,

[115397]R. S.; Stern, L. Z.: Debrancher deficiency: neuromuscular
disorder

[115398]in 5 adults. Ann. Neurol. 5: 422-436, 1979.

[115399]20326.

[115400]Fellows, I. W.; Lowe, J. S.; Ogilvie, A.; Stevens, A.; Toghill,

[115401]P. J.; Atkinson, M.: Type III glycogenosis presenting as
liver disease

[115402]in adults with atypical histological features. J. Clin. Path.
36:

[115403]431-434, 1983.

[115404]20327.

[115405]Fernandes, J.: The history of the glycogen storage disease.
(Letter) Europ.

[115406]J. Pediat. 154: 423-424, 1995.

[115407]20328.

- [115408]Garancis, J. C.; Panares, R. R.; Good, T. A.; Kuzma, J. F.:
Type
- [115409]3 glycogenosis. A biochemical and electron microscopic
study. Lab.
- [115410]Invest. 22: 468–477, 1970.
- [115411]20329.
- [115412]Hadjigeorgiou, G. M.; Comi, G. P.; Bordoni, A.; Shen, J.;
Chen,
- [115413]Y.–T.; Salani, S.; Toscano, A.; Fortunato, F.; Lucchiari, S.;
Bresolin,
- [115414]N.; Rodolico, C.; Piscaglia, M. G.; Franceschina, L.; Pa-
padimitriou,
- [115415]A.; Scarlato, G.: Novel donor splice site mutations of AGL
gene in
- [115416]glycogen storage disease type IIIa. J. Inherit. Metab. Dis.
22:
- [115417]762–763, 1999.
- [115418]20330.
- [115419]Levin, S.; Moses, S. W.; Chayoth, R.; Jadoga, N.; Steinitz, K.
- [115420]: Glycogen storage disease in Israel. A clinical, biochemi-
cal and

- [115421]genetic study. Israel J. Med. Sci. 3: 397–410, 1967.
- [115422]20331.
- [115423]Lucchiari, S.; Fogh, I.; Prella, A.; Parini, R.; Bresolin, N.;
- [115424]Melis, D.; Fiori, L.; Scarlato, G.; Comi, G. P.: Clinical and
genetic
- [115425]variability of glycogen storage disease type IIIa: seven
novel AGL
- [115426]gene mutations in the Mediterranean area. Am. J. Med.
Genet. 109:
- [115427]183–190, 2002.
- [115428]20332.
- [115429]Markowitz, A. J.; Chen, Y.-T.; Muenzer, J.; Delbuono, E. A.;
- Lucey,
- [115430]M. R.: A man with type III glycogenosis associated with
cirrhosis
- [115431]and portal hypertension. Gastroenterology 105:
1882–1885, 1993.
- [115432]20333.
- [115433]Miranda, A. F.; DiMauro, S.; Antler, A.; Stern, L. Z.; Row-
- land,
- [115434]L. P.: Glycogen debrancher deficiency is reproduced in

muscle culture. Ann.

[115435]Neurol. 9: 283–288, 1981.

[115436]20334.

[115437]Momoi, T.; Sano, H.; Yamanaka, C.; Sasaki, H.; Mikawa, H.:

Glycogen

[115438]storage disease type III with muscle involvement: reap-
praisal of phenotypic

[115439]variability and prognosis. Am. J. Med. Genet. 42: 696–699,
1992.

[115440]20335.

[115441]Moses, S. W.; Wanderman, K. L.; Myroz, A.; Frydman, M.:
Cardiac

[115442]involvement in glycogen storage disease type III. Europ. J.
Pediat. 148:

[115443]764–766, 1989.

[115444]20336.

[115445]Okubo, M.; Aoyama, Y.; Murase, T.: A novel donor splice
site

[115446]mutation in the glycogen debranching enzyme gene is as-
sociated with

[115447]glycogen storage disease type III. Biochem. Biophys. Res.

Commun. 224:

[115448]493–499, 1996.

[115449]20337.

[115450]Okubo, M.; Horinishi, A.; Makamura, N.; Aoyama, Y.;
Hashimoto,

[115451]M.; Endo, Y.; Murase, T.: A novel point mutation in an ac-
ceptor splice

[115452]site of intron 32 (IVS32 A–12–to–G) but no exon 3 muta-
tions in the

[115453]glycogen debranching enzyme gene in a homozygous pa-
tient with glycogen

[115454]storage disease type IIb. Hum. Genet. 102: 1–5, 1998.

[115455]20338.

[115456]Okubo, M.; Horinishi, A.; Suzuki, Y.; Murase, T.; Hayasaka,
K.

[115457]: Compound heterozygous patient with glycogen storage
disease type

[115458]III: identification of two novel AGL mutations, a donor
splice site

[115459]mutation of Chinese origin and a 1–bp deletion of
Japanese origin. Am.

[115460]J. Med. Genet. 93: 211–214, 2000.

[115461]20339.

[115462]Okubo, M.; Horinishi, A.; Takeuchi, M.; Suzuki, Y.; Sakura, N.;

[115463]Hasegawa, Y.; Igarashi, T.; Goto, K.; Tahara, H.; Uchimoto, S.; Omichi,

[115464]K.; Kanno, H.; Hayasaka, K.; Murase, T.: Heterogeneous mutations

[115465]in the glycogen-debranching enzyme gene are responsible for glycogen

[115466]storage disease type IIIa in Japan. Hum. Genet. 106: 108–115, 2000.

[115467]20340.

[115468]Attard-Montalto, S.; Evans, N.; Sherwood, R. A.: Carotenaemia

[115469]with low vitamin A levels and low retinol-binding protein. J. Inherit.

[115470]Metab. Dis. 15: 929–930, 1992.

[115471]20341.

[115472]Chainani, M.; Sampsell, B.; Elliott, R. W.: Localization of the

[115473]gene for plasma retinol binding protein to the distal half of mouse

[115474]chromosome 19. Genomics 9: 376–379, 1991.

[115475]20342.

[115476]Matsuo, T.: Familial retinol-binding-protein deficiency?
(Letter) Lancet II:

[115477]910 only, 1987.

[115478]20343.

[115479]Matsuo, T.; Matsuo, N.; Shiraga, F.; Koide, N.: Familial
retinol-binding-protein

[115480]deficiency. (Letter) Lancet II: 402–403, 1987.

[115481]20344.

[115482]Morgan, F. F.; Canfield, R. E.; Goodman, D. S.: The partial
structure

[115483]of human plasma prealbumin and retinol-binding protein.
Biochim.

[115484]Biophys. Acta 236: 798–801, 1971.

[115485]20345.

[115486]Rask, L.; Anundi, H.; Fohlman, J.; Peterson, P. A.: The
complete

[115487]amino acid sequence of human serum retinol-binding
protein. Upsala

[115488]J. Med. Sci. 92: 115–146, 1987.

[115489]20346.

[115490]Rask, L.; Vahlquist, A.; Peterson, P. A.: Studies on two physiological

[115491]forms of the human retinol-binding protein differing in vitamin A

[115492]and arginine content. J. Biol. Chem. 246: 6638–6646, 1971.

[115493]20347.

[115494]Rocchi, M.; Covone, A.; Romeo, G.; Faraonio, R.; Colantuoni, V.

[115495]: Regional mapping of RBP4 to 10q23–q24 and RBP1 to 3q21–q22 in man. Somat.

[115496]Cell Molec. Genet. 15: 185–190, 1989.

[115497]20348.

[115498]Seeliger, M. W.; Biesalski, H. K.; Wissinger, B.; Gollnick, H.;

[115499]Gielen, S.; Frank, J.; Beck, S.; Zrenner, E.: Phenotype in retinol

[115500]deficiency due to a hereditary defect in retinol binding protein synthesis. Invest.

[115501]Ophthal. Vis. Sci. 40: 3–11, 1999.

[115502]20349.

[115503]Thuluvath, P. J.: Familial retinol-binding-protein deficiency?

[115504](Letter) Lancet II: 910 only, 1987.

[115505]20350.

[115506]Carson, N. L.; Simpson, N. E.: A physical map of 13 markers on

[115507]chromosome 10 from dosage studies on abnormal cell lines. (Abstract) Cytogenet.

[115508]Cell Genet. 51: 974-975, 1989.

[115509]20351.

[115510]Chin, K. S.; Mathew, C. G. P.; Fong, S. L.; Bridges, C. D. B.;

[115511]Ponder, B. A. J.: Styl RFLP recognised by a human IRBP cDNA localised

[115512]to chromosome 10. Nucleic Acids Res. 16: 1645 only, 1988.

[115513]20352.

[115514]Danciger, M.; Kozak, C. A.; Nickerson, J.; Redmond, T. M.; Farber,

[115515]D. B.: Localization of the gene for interphotoreceptor retinoid-binding

[115516]protein to mouse chromosome 14 near Np-1. Genomics
8: 727-731, 1990.

[115517]20353.

[115518]Farrer, L. A.; Castiglione, C. M.; Kidd, J. R.; Myers, S.; Car-
son,

[115519]N.; Simpson, N. E.; Kidd, K. K.: A linkage group of five
DNA markers

[115520]on human chromosome 10. Genomics 3: 72-77, 1988.

[115521]20354.

[115522]Fong, S.-L.; Liou, G. I.; Landers, R. A.; Alvarez, R. A.; Gon-
zalez-Fernandez,

[115523]F.; Glazebrook, P. A.; Lam, D. M. K.; Bridges, C. D. B.:
Characterization,

[115524]localization, and biosynthesis of an interstitial retinol-
binding

[115525]glycoprotein in the human eye. J. Neurochem. 42:
1667-1676, 1984.

[115526]20355.

[115527]Liou, G. I.; Fong, S.-L.; Beattie, W. G.; Cook, R. G.; Leone,
J.;

[115528]Landers, R. A.; Alvarez, R. A.; Wang, C.; Li, Y.; Bridges, C.
D. B.

[115529] Bovine interstitial retinol-binding protein (IRBP)--isolation and
 [115530] sequence analysis of cDNA clones, characterization and in vitro translation
 [115531] of mRNA. Vision Res. 26: 1645-1653, 1986.
 [115532] 20356.

[115533] Liou, G. I.; Fong, S.-L.; Gosden, J.; vanTuinen, P.; Ledbetter,
 [115534] D. H.; Christie, S.; Rout, D.; Bhattacharya, S.; Cook, R. G.; Li,
 [115535] Y.; Wang, C.; Bridges, C. D. B.: Human interstitial retinol-binding
 [115536] protein (IRBP): cloning, partial sequence, and chromosomal localization. Somat.
 [115537] Cell Molec. Genet. 13: 315-323, 1987.
 [115538] 20357.

[115539] Liou, G. I.; Li, Y.; Wang, C.; Fong, S.-L.; Bhattacharya, F. S.;
 [115540] Bridges, C. D. B.: Bgl II RFLP recognized by a human IRBP cDNA localized
 [115541] to chromosome 10. Nucleic Acids Res. 15: 3196 only, 1987.

[115542]20358.

[115543]McGuire, W.; Hill, A. V. S.; Allsopp, C. E. M.; Greenwood, B.

[115544]M.; Kwiatkowski, D.: Variation in the TNF- α promoter
region associated

[115545]with susceptibility to cerebral malaria. Nature 371:
508-511, 1994.

[115546]20359.

[115547]Bruford, E. A.; Riise, R.; Teague, P. W.; Porter, K.; Thom-
son,

[115548]K. L.; Moore, A. T.; Jay, M.; Warburg, M.; Schinzel, A.;
Tommerup,

[115549]N.; Tornqvist, K.; Rosenberg, T.; Patton, M.; Mansfield, D.
C.; Wright,

[115550]A. F.: Linkage mapping in 29 Bardet-Biedl syndrome fami-
lies confirms

[115551]loci in chromosomal regions 11q13, 15q22.3-q23, and
16q21. Genomics 41:

[115552]93-99, 1997.

[115553]20360.

[115554]Benbrook, D.; Pfahl, M.: A novel thyroid hormone receptor
encoded

[115555]by a cDNA clone from a human testis library. Science 238:
788-791,

[115556]1987.

[115557]20361.

[115558]Bernal, J.; Refetoff, S.; DeGroot, L. J.: Abnormalities of tri-
iodothyronine

[115559]binding to lymphocyte and fibroblast nuclei from a patient
with peripheral

[115560]tissue resistance to thyroid hormone action. J. Clin. En-
docr. Metab. 47:

[115561]1266-1272, 1978.

[115562]20362.

[115563]Dayton, A. I.; Selden, J. R.; Laws, G.; Dorney, D. J.; Finan,
J.;

[115564]Tripputi, P.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.;
Croce, C.

[115565]M.: A human c-erbA oncogene homologue is closely
proximal to the

[115566]chromosome 17 breakpoint in acute promyelocytic
leukemia. Proc. Nat.

[115567]Acad. Sci. 81: 4495-4499, 1984.

[115568]20363.

[115569]Debuire, B.; Henry, C.; Benaissa, M.; Biserte, G.; Claverie, J.

[115570]M.; Saule, S.; Martin, P.; Stehelin, D.: Sequencing the erbA
gene

[115571]of avian erythroblastosis virus reveals a new type of onco-
gene. Science 224:

[115572]1456–1459, 1984.

[115573]20364.

[115574]Ferro, M. T.; San Roman, C.: Constitutional t(15;17). Can-
cer

[115575]Genet. Cytogenet. 4: 89–91, 1981.

[115576]20365.

[115577]Gullberg, H.; Rudling, M.; Forrest, D.; Angelin, B.;
Vennstrom,

[115578]B.: Thyroid hormone receptor beta-deficient mice show
complete loss

[115579]of the normal cholesterol 7- α -hydroxylase (CYP7A)
response to

[115580]thyroid hormone but display enhanced resistance to di-
etary cholesterol. Molec.

[115581]Endocr. 14: 1739–1749, 2000.

[115582]20366.

[115583]Ichikawa, K.; Hughes, I. A.; Horwitz, A. L.; DeGroot, L. J.:

Characterization

[115584]of nuclear thyroid hormone receptors of cultured skin fibroblasts

[115585]from patients with resistance to thyroid hormone.

Metabolism 36:

[115586]392–399, 1987.

[115587]20367.

[115588]Iskaros, J.; Pickard, M.; Evans, I.; Sinha, A.; Hardiman, P.;

Ekins,

[115589]R.: Thyroid hormone receptor gene expression in first trimester human

[115590]fetal brain. J. Clin. Endocr. Metab. 85: 2620–2623, 2000.

[115591]20368.

[115592]Jansson, M.; Philipson, L.; Vennstrom, B.: Isolation and characterization

[115593]of multiple human genes homologous to the oncogenes of avian erythroblastosis

[115594]virus. EMBO J. 2: 561–565, 1983.

[115595]20369.

[115596]Jhanwar, S. C.; Chaganti, R. S. K.; Croce, C. M.: Germ-line chromosomal

[115597]localization of human c-erb-A oncogene. Somat. Cell Molec. Genet. 11:

[115598]99–102, 1985.

[115599]20370.

[115600]Kaneshige, M.; Suzuki, H.; Kaneshige, K.; Cheng, J.; Wimbrow,

[115601]H.; Barlow, C.; Willingham, M. C.; Cheng, S.: A targeted dominant

[115602]negative mutation of the thyroid hormone alpha-1 receptor causes increased

[115603]mortality, infertility, and dwarfism in mice. Proc. Nat. Acad. Sci. 98:

[115604]15095–15100, 2001.

[115605]20371.

[115606]Laudet, V.; Begue, A.; Henry-Duthoit, C.; Joubel, A.; Martin,

[115607]P.; Stehelin, D.; Saule, S.: Genomic organization of the human thyroid

[115608]hormone receptor alpha (c-erbA-1) gene. Nucleic Acids Res. 19: 1105–1112,

[115609]1991.

[115610]20372.

[115611]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J. D.;

[115612]Oren, M.: Translocation of the p53 gene in t(15;17) in acute promyelocytic

[115613]leukaemia. Nature 316: 826–828, 1985.

[115614]20373.

[115615]Mathieu–Mahul, D.; Xu, D. Q.; Saule, S.; Lidereau, R.; Galibert,

[115616]F.; Berger, R.; Mauchauffe, M.; Larsen, C. J.: An EcoRI restriction

[115617]fragment length polymorphism (RFLP) in the human c-erbA locus. Hum.

[115618]Genet. 71: 41–44, 1985.

[115619]20374.

[115620]McCabe, C. J.; Gittoes, N. J.; Sheppard, M. C.; Franklyn, J. A.

[115621]: Thyroid receptor alpha–1 and alpha–2 mutations in non-functioning

[115622]pituitary tumors. J. Clin. Endocr. Metab. 84: 649–653,

1999.

[115623]20375.

[115624]Menezes-Ferreira, M. M.; Eil, C.; Wortsman, J.; Weintraub, B.

[115625]D.: Decreased nuclear uptake of

[125-I]triiodo-L-thyronine in fibroblasts

[115626]from patients with peripheral thyroid hormone resistance.
J. Clin.

[115627]Endocr. Metab. 59: 1081-1087, 1984.

[115628]20376.

[115629]Mitelman, F.; Manolov, G.; Manolova, Y.; Billstrom, R.;
Heim,

[115630]S.; Kristoffersson, U.; Mandahl, N.; Ferro, M. T.; San Roman, C.:

[115631]High resolution chromosome analysis of constitutional
and acquired

[115632]t(15;17) maps c-erbA to subband 17q11.2. Cancer Genet.
Cytogenet. 22:

[115633]95-98, 1986.

[115634]20377.

[115635]Miyajima, N.; Horiuchi, R.; Shibuya, Y.; Fukushige, S.; Mat-

subara,

[115636]K.; Toyoshima, K.; Yamamoto, T.: Two erbA homologs encoding proteins

[115637]with different T(3) binding capacities are transcribed from opposite

[115638]DNA strands of the same genetic locus. Cell 57: 31-39, 1989.

[115639]20378.

[115640]Nagaya, T.; Nomura, Y.; Fujieda, M.; Seo, H.: Heterodimerization

[115641]preferences of thyroid hormone receptor alpha isoforms. Biochem.

[115642]Biophys. Res. Commun. 226: 426-430, 1996.

[115643]20379.

[115644]Nakai, A.; Seino, S.; Sakurai, A.; Szilak, I.; Bell, G. I.; DeGroot,

[115645]L. J.: Characterization of a thyroid hormone receptor expressed in

[115646]human kidney and other tissues. Proc. Nat. Acad. Sci. 85: 2781-2785,

[115647]1988.

[115648]20380.

- [115649]Ng, L.; Rusch, A.; Amma, L. L.; Nordstrom, K.; Erway, L. C.; Vennstrom, B.; Forrest, D.: Suppression of the deafness and thyroid dysfunction in Thr β -null mice by an independent mutation in the Thra thyroid hormone receptor gene. Hum. Molec. Genet. 10: 2701–2708, 2001.
- [115652]20381.
- [115654]Puzianowska-Kuznicka, M.; Krystyniak, A.; Madej, A.; Cheng, S.-Y.; Nauman, J.: Functionally impaired TR mutants are present in thyroid papillary cancer. J. Clin. Endocr. Metab. 87: 1120–1128, 2002.
- [115657]20382.
- [115658]Refetoff, S.; DeGroot, L. J.; Benard, B.; DeWind, L. T.: Studies of a sibship with apparent hereditary resistance to the intracellular action of thyroid hormone. Metabolism 21: 723–756, 1972.
- [115661]20383.

[115662]Rider, S. H.; Bailey, C. J.; Voss, R.; Sheer, D.; Hiorns, L. R.;

[115663]Solomon, E.: RFLP for the human erb-A1 gene. Nucleic
Acids Res. 15:

[115664]863 only, 1987.

[115665]20384.

[115666]Sakurai, A.; Nakai, A.; DeGroot, L. J.: Expression of three
forms

[115667]of thyroid hormone receptor in human tissues. Molec. En-
docr. 3:

[115668]392-399, 1989.

[115669]20385.

[115670]Wolf, F. W.; Marks, R. M.; Sarma, V.; Byers, M. G.; Katz, R.
W.;

[115671]Shows, T. B.; Dixit, V. M.: Characterization of a novel tu-
mor necrosis

[115672]factor-alpha-induced endothelial primary response gene.
J. Biol.

[115673]Chem. 267: 1317-1326, 1992.

[115674]20386.

[115675]Pandey, A.; Lindberg, R. A.; Dixit, V. M.: Receptor orphans
find

[115676]a family. Curr. Biol. 5: 986–989, 1995.

[115677]20387.

[115678]Aaltonen, L.–M.; Chen, R. W.; Roth, S.; Makitie, A. A.;
Rihkanen,

[115679]H.; Vaheri, A.; Aaltonen, L. A.: Role of TP53 P72R poly-
morphism in

[115680]human papillomavirus associated premalignant laryngeal
neoplasm. J.

[115681]Med. Genet. 38: 327 only, 2001.

[115682]20388.

[115683]Aas, T.; Borresen, A.–L.; Geisler, S.; Smith–Sorenson, B.;
Johnsen,

[115684]H.; Varhaug, J. E.; Akslen, L. A.; Lonning, P. E.: Specific P53
mutations

[115685]are associated with de novo resistance to doxorubicin in
breast cancer

[115686]patients. Nature Med. 2: 811–814, 1996.

[115687]20389.

[115688]Aguilar, F.; Harris, C. C.; Sun, T.; Hollstein, M.; Cerutti, P.

[115689]: Geographic variation of p53 mutational profile in non-
malignant human

[115690]liver. Science 264: 1317–1319, 1994.

[115691]20390.

[115692]Aguilar, F.; Hussain, S. P.; Cerutti, P.: Aflatoxin B1 induces

[115693]the transversion of G-to-T in codon 249 of the p53 tumor
suppressor

[115694]gene in human hepatocytes. Proc. Nat. Acad. Sci. 90:
8586–8590,

[115695]1993.

[115696]20391.

[115697]Ara, S.; Lee, P. S. Y.; Hansen, M. F.; Saya, H.: Codon 72
polymorphism

[115698]of the TP53 gene. Nucleic Acids Res. 18: 4961, 1990.

[115699]20392.

[115700]Arai, M.; Shimizu, S.; Imai, Y.; Nakatsuru, Y.; Oda, H.;
Oohara,

[115701]T.; Ishikawa, T.: Mutations of the Ki-ras, p53 and APC
genes in adenocarcinomas

[115702]of the human small intestine. Int. J. Cancer 70: 390–395,
1997.

[115703]20393.

[115704] Baker, S. J.; Fearon, E. R.; Nigro, J. M.; Hamilton, S. R.;
Preisinger,
[115705] A. C.; Jessup, J. M.; vanTuinen, P.; Ledbetter, D. H.; Barker,
D.
[115706] F.; Nakamura, Y.; White, R.; Vogelstein, B.: Chromosome
17 deletions
[115707] and p53 gene mutations in colorectal carcinomas. Science
244: 217–221,
[115708] 1989.
[115709] 20394.
[115710] Benchimol, S.; Lamb, P.; Crawford, L. V.; Sheer, D.; Shows,
T.
[115711] B.; Bruns, G. A. P.; Peacock, J.: Transformation associated
p53 protein
[115712] is encoded by a gene on human chromosome 17. Somat.
Cell Molec. Genet. 11:
[115713] 505–509, 1985.
[115714] 20395.
[115715] Bernal, J. A.; Luna, R.; Espina, A.; Lazaro, I.; Ramos–
Morales,
[115716] F.; Romero, F.; Arias, C.; Silva, A.; Tortolero, M.; Pintor–
Toro,

[115717]J. A.: Human securin interacts with p53 and modulates
p53-mediated
[115718]transcriptional activity and apoptosis. *Nature Genet.* 32:
306–311,
[115719]2002.
[115720]20396.
[115721]Beroud, C.; Verdier, F.; Soussi, T.: p53 gene mutation:
software
[115722]and database. *Nucleic Acids Res.* 24: 147–150, 1996.
[115723]20397.
[115724]Borresen, A.-L.; Hovig, E.; Smith-Sorensen, B.; Malkin, D.;
Lystad,
[115725]S.; Andersen, T. I.; Nesland, J. M.; Isselbacher, K. J.; Friend,
S.
[115726]H.: Constant denaturant gel electrophoresis as a rapid
screening
[115727]technique for p53 mutations. *Proc. Nat. Acad. Sci.* 88:
8405–8409,
[115728]1991.
[115729]20398.
[115730]Borresen, A. L.; Andersen, T. I.; Garber, J.; Barbier-Piroux,

[115731]N.; Thorlacius, S.; Eyfjord, J.; Ottestad, L.; Smith-Sorensen, B.;

[115732]Hovig, E.; Malkin, D.; Friend, S. H.: Screening for germ line TP53

[115733]mutations in breast cancer patients. Cancer Res. 52: 3234-3236,

[115734]1992.

[115735]20399.

[115736]Brash, D. E.; Rudolph, J. A.; Simon, J. A.; Lin, A.; McKenna,

[115737]G. J.; Baden, H. P.; Halperin, A. J.; Ponten, J.: A role for sunlight

[115738]in skin cancer: UV-induced p53 mutations in squamous cell carcinoma. Proc.

[115739]Nat. Acad. Sci. 88: 10124-10128, 1991.

[115740]20400.

[115741]Brennan, J. A.; Boyle, J. O.; Koch, W. M.; Goodman, S. N.; Hruban,

[115742]R. H.; Eby, Y. J.; Couch, M. J.; Forastiere, A. A.; Sidransky, D.

[115743]: Association between cigarette smoking and mutation of the p53 gene

[115744]in squamous-cell carcinoma of the head and neck. New
Eng. J. Med. 332:

[115745]712–717, 1995.

[115746]20401.

[115747]Bressac, B.; Kew, M.; Wands, J.; Ozturk, M.: Selective G to T

[115748]mutations of p53 gene in hepatocellular carcinoma from
southern Africa. Nature 350:

[115749]429–431, 1991.

[115750]20402.

[115751]Brodsky, M. H.; Nordstrom, W.; Tsang, G.; Kwan, E.; Rubin,
G.

[115752]M.; Abrams, J. M.: Drosophila p53 binds a damage re-
sponse element

[115753]at the reaper locus. Cell 101: 103–113, 2000.

[115754]20403.

[115755]Buetow, K. H.; Sheffield, V. C.; Zhu, M.; Zhou, T.; Shen, F.;

[115756]Hino, O.; Smith, M.; McMahon, B. J.; Lanier, A. P.; London,
W. T.;

[115757]Redeker, A. G.; Govindarajan, S.: Low frequency of p53
mutations

[115758]observed in a diverse collection of primary hepatocellular carcinomas. Proc.

[115759]Nat. Acad. Sci. 89: 9622–9626, 1992.

[115760]20404.

[115761]Bunz, F.; Dutriaux, A.; Lengauer, C.; Waldman, T.; Zhou, S.; Brown,

[115762]J. P.; Sedivy, J. M.; Kinzler, K. W.; Vogelstein, B.: Requirement

[115763]for p53 and p21 to sustain G2 arrest after DNA damage. Science 282:

[115764]1497–1501, 1998.

[115765]20405.

[115766]Caelles, C.; Helmberg, A.; Karin, M.: p53–dependent apoptosis

[115767]in the absence of transcriptional activation of p53–target genes. Nature 370:

[115768]220–223, 1994.

[115769]20406.

[115770]Colucci–Guyon, E.; Portier, M.–M.; Dunia, I.; Paulin, D.; Pournin,

[115771]S.; Babinet, C.: Mice lacking vimentin develop and repro–

duce without

[115772]an obvious phenotype. Cell 79: 679–694, 1994.

[115773]20407.

[115774]Ferrari, S.; Battini, R.; Kaczmarek, L.; Rittling, S.; Cal-
abretta,

[115775]B.; de Riel, J. K.; Philiponis, V.; Weil, J.-F.; Baserga, R.:
Coding

[115776]sequence and growth regulation of the human vimentin
gene. Molec.

[115777]Cell. Biol. 6: 3614–3620, 1986.

[115778]20408.

[115779]Gieser, L.; Swaroop, A.: Expressed sequence tags and
chromosomal

[115780]localization of cDNA clones from a subtracted retinal pig-
ment epithelium

[115781]library. Genomics 13: 873–876, 1992.

[115782]20409.

[115783]Lilienbaum, A.; Legagneux, V.; Portier, M.-M.; Dellagi, K.;
Paulin,

[115784]D.: Vimentin gene: expression in human lymphocytes and
in Burkitt's

[115785]lymphoma cells. EMBO J. 5: 2809–2814, 1986.

[115786]20410.

[115787]Marcus, E. M.; Smith, B. A.; Telenius, H.; Landsvater, R. M.;
Buys,

[115788]C. H. C. M.; Ferrari, S.; Ponder, B. A. J.; Mathew, C. G. P.:
Bcll

[115789]RFLP for the human vimentin gene. Nucleic Acids Res. 16:
9068 only,

[115790]1988.

[115791]20411.

[115792]Mathew, C. G.; Wakeling, W.; Jones, E.; Easton, D.; Fisher,
R.;

[115793]Strong, C.; Smith, B.; Chin, K.; Little, P.; Nakamura, Y.;
Shows,

[115794]T. B.; Jones, C.; Goodfellow, P. J.; Povey, S.; Ponder, B. A.
J.:

[115795]Regional localization of polymorphic markers on chromo-
some 10 by physical

[115796]and genetic mapping. Ann. Hum. Genet. 54: 121–129,
1990.

[115797]20412.

- [115798]Perreau, J.; Lilienbaum, A.; Vasseur, M.; Paulin, D.: Nu-
cleotide
[115799]sequence of the human vimentin gene and regulation of
its transcription
[115800]in tissues and cultured cells. Gene 62: 7-16, 1988.
[115801]20413.
- [115802]Gimona, M.; Small, J. V.; Moeremans, M.; Van Damme, J.;
Puype,
[115803]M.; Vandekerckhove, J.: Porcine vinculin and metavinculin
differ
[115804]by a 68-residue insert located close to the carboxy-ter-
minal part
[115805]of the molecule. EMBO J. 7: 2329-2334, 1988.
[115806]20414.
- [115807]Mulligan, L. M.; Gardner, E.; Telenius, H.; Ponder, B. A. J.:
[115808]Complementary physical and genetic techniques map the
vinculin (VCL)
[115809]gene on chromosome 10q. Genomics 13: 1347-1349,
1992.
[115810]20415.
- [115811]Strasser, P.; Gimona, M.; Herzog, M.; Geiger, B.; Small, J.

V.

[115812] Variable and constant regions in the C-terminus of vinculin and

[115813]metavinculin: cloning and expression of fragments in E. coli. FEBS

[115814]Lett. 317: 189-194, 1993.

[115815]20416.

[115816]Koteliansky, V. E.; Ogryzko, E. P.; Zhidkova, N. I.; Weller, P.

[115817]A.; Critchley, D. R.; Vancompernelle, K.; Vandekerckhove, J.; Strasser,

[115818]P.; Way, M.; Gimona, M.; Small, J. V.: An additional exon in the

[115819]human vinculin gene specifically encodes meta-vinculin-specific difference

[115820]peptide: cross-species comparison reveals variable and conserved motifs

[115821]in the meta-vinculin insert. Europ. J. Biochem. 204: 767-772, 1992.

[115822]20417.

[115823]Turner, C. E.; Burridge, K.: Detection of metavinculin in human

[115824]platelets using a modified talin overlay assay. Europ. J.

Cell Biol. 49:

[115825]202–206, 1989.

[115826]20418.

[115827]Weller, P. A.; Ogryzko, E. P.; Corben, E. B.; Zhidkova, N. I.;

[115828]Patel, B.; Price, G. J.; Spurr, N. K.; Koteliensky, V. E.;

Critchley,

[115829]D. R.: Complete sequence of human vinculin and assignment of the

[115830]gene to chromosome 10. Proc. Nat. Acad. Sci. 87: 5667–5671, 1990.

[115831]20419.

[115832]Katsanis, N.; Ansley, S. J.; Badano, J. L.; Eichers, E. R.;

Lewis,

[115833]R. A.; Hoskins, B. E.; Scambler, P. J.; Davidson, W. S.; Beales, P.

[115834]L.; Lupski, J. R.: Triallelic inheritance in Bardet–Biedl syndrome,

[115835]a mendelian recessive disorder. Science 293: 2256–2259, 2001.

[115836]20420.

[115837]Katsanis, N.; Beales, P. L.; Woods, M. O.; Lewis, R. A.;

Green,

[115838]J. S.; Parfrey, P. S.; Ansley, S. J.; Davidson, W. S.; Lupski, J.

[115839]R.: Mutations in MKKS cause obesity, retinal dystrophy and renal

[115840]malformations associated with Bardet-Biedl syndrome.

Nature Genet. 26:

[115841]67-70, 2000.

[115842]20421.

[115843]Kwitek-Black, A. E.; Carmi, R.; Duyk, G. M.; Buetow, K. H.; Elbedour,

[115844]K.; Parvari, R.; Yandava, C. N.; Stone, E. M.; Sheffield, V. C.:

[115845]Linkage of Bardet-Biedl syndrome to chromosome 16q and evidence for

[115846]non-allelic genetic heterogeneity. Nature Genet. 5: 392-396, 1993.

[115847]20422.

[115848]Ishii, H.; Majerus, P. W.: Thrombomodulin is present in human

[115849]plasma and urine. J. Clin. Invest. 76: 2178-2181, 1985.

[115850]20423.

- [115851] Gabreels, B. A. T. F.; Swaab, D. F.; de Kleijn, D. P. V.; Seidah,
dah,
- [115852] N. G.; Van de Loo, J.-W.; Van de Ven, W. J. M.; Martens, G.
J. M.;
- [115853] van Leeuwen, F. W.: Attenuation of the polypeptide 7B2,
prohormone
- [115854] convertase PC2, and vasopressin in the hypothalamus of
some Prader-Willi
- [115855] patients: indications for a processing defect. J. Clin. En-
docr. Metab. 83:
- [115856] 591-599, 1998.
- [115857] 20424.
- [115858] Bell, K.; Hopper, K. E.; McKenzie, H. A.; Murphy, W. H.;
Shaw,
- [115859] D. C.: A comparison of bovine alpha-lactalbumin A and B
of Droughtmaster. Biochim.
- [115860] Biophys. Acta 214: 437-444, 1970.
- [115861] 20425.
- [115862] Bell, K.; McKenzie, H. A.; Murphy, W. H.; Shaw, D. C.: Beta-
lactoglobulin
- [115863] (Droughtmaster): a unique protein variant. Biochim. Bio-
phys. Acta 214:

[115864]427-436, 1970.

[115865]20426.

[115866]Julkunen, M.; Seppala, M.; Janne, O. A.: Complete amino acid sequence

[115867]of human placental protein 14: a progesterone-regulated uterine protein

[115868]homologous to beta-lactoglobulins. Proc. Nat. Acad. Sci. 85: 8845-8849,

[115869]1988.

[115870]20427.

[115871]Morris, H. R.; Dell, A.; Easton, R. L.; Panico, M.; Koistinen,

[115872]H.; Koistinen, R.; Oehninger, S.; Patankar, M. S.; Seppala, M.; Clark,

[115873]G. F.: Gender-specific glycosylation of human glycodefin affects

[115874]its contraceptive activity. J. Biol. Chem. 271: 32159-32167, 1996.

[115875]20428.

[115876]Song, M.; Ramaswamy, S.; Ramachandran, S.; Flowers, L. C.; Horowitz,

[115877]J. R.; Rock, J. A.; Parthasarathy, S.: Angiogenic role for gly-

codelin

[115878]in tumorigenesis. Proc. Nat. Acad. Sci. 98: 9265–9270,
2001.

[115879]20429.

[115880]Van Cong, N.; Vaisse, C.; Gross, M.–S.; Slim, R.; Milgrom,
E.;

[115881]Bernheim, A.: The human placental protein 14 (PP14) gene
is localized

[115882]on chromosome 9q34. Hum. Genet. 86: 515–518, 1991.

[115883]20430.

[115884]Lee, F. S.; Fox, E. A.; Zhou, H.–M.; Strydom, D. J.; Vallee, B.

[115885]L.: Primary structure of human placental ribonuclease in-
hibitor. Biochemistry 27:

[115886]8545–8553, 1988.

[115887]20431.

[115888]Weremowicz, S.; Fox, E. A.; Morton, C. C.; Vallee, B. L.: The

[115889]placental ribonuclease inhibitor (RNH) gene is located on
chromosome

[115890]subband 11p15.5. Genomics 8: 717–721, 1990.

[115891]20432.

[115892]Zneimer, S. M.; Crawford, D.; Schneider, N. R.; Beutler, B.:
Mapping

[115893]of the human ribonuclease inhibitor gene (RNH) to chromosome 11p15

[115894]by in situ hybridization. Genomics 8: 175–178, 1990.

[115895]20433.

[115896]Coughlin, P.; Nicholl, J.; Sun, J.; Salem, H.; Bird, P.; Sutherland,

[115897]G. R.: Chromosomal mapping of the human proteinase inhibitor 6 (PI6)

[115898]gene to 6p25 by fluorescence in situ hybridization. Genomics 26:

[115899]431–433, 1995.

[115900]20434.

[115901]Coughlin, P.; Sun, J.; Cerruti, L.; Salem, H. H.; Bird, P.:
Cloning

[115902]and molecular characterization of a human intracellular serine proteinase

[115903]inhibitor. Proc. Nat. Acad. Sci. 90: 9417–9421, 1993.

[115904]20435.

[115905]Zhang, K.; Westberg, J. A.; Paetau, A.; von Boguslawsky,

K.; Lindsberg,

[115906]P.; Erlander, M.; Guo, H.; Su, J.; Olsen, H. S.; Andersson, L.
C.

[115907]: High expression of stanniocalcin in differentiated brain
neurons. Am.

[115908]J. Path. 153: 439–445, 1998.

[115909]20436.

[115910]Hamilton, S. E.; Hurley, J. B.: A phosphodiesterase in-
hibitor

[115911]specific to a subset of bovine retinal cones. J. Biol. Chem.
265:

[115912]11259–11264, 1990.

[115913]20437.

[115914]Shimizu–Matsumoto, A.; Itoh, K.; Inazawa, J.; Nishida, K.;
Matsumoto,

[115915]Y.; Kinoshita, S.; Matsubara, K.; Okubo, K.: Isolation and
chromosomal

[115916]localization of the human cone cGMP phosphodiesterase
gamma cDNA (PDE6H). Genomics 32:

[115917]121–124, 1996.

[115918]20438.

[115919]Isomura, M.; Okui, K.; Fujiwara, T.; Shin, S.; Nakamura, Y.:

Cloning

[115920]and mapping of a novel human cDNA homologous to

DROER, the enhancer

[115921]of the *Drosophila melanogaster* rudimentary gene. *Ge-*

nomics 32: 125-127,

[115922]1996.

[115923]20439.

[115924]Chen, J.; Engle, S. J.; Seilhamer, J. J.; Tischfield, J. A.:

Cloning

[115925]and recombinant expression of a novel human low molec-

ular weight Ca^{2+} -dependent

[115926]phospholipase A2. *J. Biol. Chem.* 269: 2365-2368, 1994.

[115927]20440.

[115928]Tischfield, J. A.; Xia, Y.-R.; Shih, D. M.; Klisak, I.; Chen, J.;

[115929]Engle, S. J.; Siakotos, A. N.; Winstead, M. V.; Seilhamer, J.

J.;

[115930]Allamand, V.; Gyapay, G.; Lusi, A. J.: Low molecular

weight, calcium-dependent

[115931]phospholipase A(2) genes are linked and map to homolo-

gous chromosome

[115932]regions in mouse and human. Genomics 32: 328–333, 1996.

[115933]20441.

[115934]Hayashi, M.; Fujimoto, S.; Takano, H.; Ushiki, T.; Abe, K.; Ishikura,

[115935]H.; Yoshida, M. C.; Kirchhoff, C.; Ishibashi, T.; Kasahara, M.: Characterization

[115936]of a human glycoprotein with a potential role in sperm-egg fusion:

[115937]cDNA cloning, immunohistochemical localization, and chromosomal assignment

[115938]of the gene (AEGL1). Genomics 32: 367–374, 1996.

[115939]20442.

[115940]Muzio, M.; Bosisio, D.; Polentarutti, N.; D'amico, G.; Stoppacciaro,

[115941]A.; Mancinelli, R.; van't Veer, C.; Penton-Rol, G.; Ruco, L. P.; Allavena,

[115942]P.; Mantovani, A.: Differential expression and regulation of Toll-like

[115943]receptors (TLR) in human leukocytes: selective expression of TLR3

[115944]in dendritic cells. J. Immun. 164: 5998–6004, 2000.

[115945]20443.

[115946]Rock, F. L.; Hardiman, G.; Timans, J. C.; Kastelein, R. A.;
Bazan,

[115947]J. F.: A family of human receptors structurally related to
Drosophila

[115948]Toll. Proc. Nat. Acad. Sci. 95: 588–593, 1998.

[115949]20444.

[115950]Perrimon, N.; Mahowald, A. P.: Multiple functions of seg-
ment polarity

[115951]genes in Drosophila. Dev. Biol. 119: 587–600, 1987.

[115952]20445.

[115953]Pizzuti, A.; Novelli, G.; Mari, A.; Ratti, A.; Colosimo, A.;
Amati,

[115954]F.; Penso, D.; Sangiuolo, F.; Calabrese, G.; Palka, G.; Silani,
V.;

[115955]Gennarelli, M.; Mingarelli, R.; Scarlato, G.; Scambler, P.;
Dallapiccola,

[115956]B.: Human homologue sequences to the Drosophila di-
shevelled segment-polarity

[115957]gene are deleted in the DiGeorge syndrome. Am. J. Hum.
Genet. 58:

[115958]722–729, 1996.

[115959]20446.

[115960]Onyango, P.; Lubyova, B.; Gardellin, P.; Kurzbauer, R.;
Weith,

[115961]A.: Molecular cloning and expression analysis of five novel
genes

[115962]in chromosome 1p36. Genomics 50: 187–198, 1998.

[115963]20447.

[115964]Barbier, M.; Attoub, S.; Calvez, R.; Laffargue, M.; Jarry, A.;

[115965]Mareel, M.; Altruda, F.; Gespach, C.; Wu, D.; Lu, B.; Hirsch,
E.;

[115966]Wymann, M. P.: Weakening link to colorectal cancer? Na-
ture 413:

[115967]796 only, 2001.

[115968]20448.

[115969]Crackower, M. A.; Oudit, G. Y.; Kozieradzki, I.; Sarao, R.;
Sun,

[115970]H.; Sasaki, T.; Hirsch, E.; Suzuki, A.; Shioi, T.; Irie-Sasaki,
J.;

[115971]Sah, R.; Cheng, H.–Y. M.; and 13 others: Regulation of
myocardial

[115972]contractility and cell size by distinct PI3K–PTEN signaling pathways. Cell 110:

[115973]737–749, 2002.

[115974]20449.

[115975]Hirsch, E.; Katanaev, V. L.; Garlanda, C.; Azzolino, O.;
Pirola,

[115976]L.; Silengo, L.; Sozzani, S.; Mantovani, A.; Altruda, F.;
Wymann,

[115977]M. P.: Central role for G protein–coupled phosphoinositide
3–kinase

[115978]gamma in inflammation. Science 287: 1049–1053, 2000.

[115979]20450.

[115980]Jiang, K.; Zhong, B.; Gilvary, D. L.; Corliss, B. C. Hong–
Geller,

[115981]E.; Wei, S; Djeu, J. Y.: Pivotal role of phosphoinositide–3
kinase

[115982]in regulation of cytotoxicity in natural killer cells. Nature
Immun. 1:

[115983]419–425, 2000.

[115984]20451.

[115985]Kikuta, Y.; Kato, M.; Yamashita, Y.; Miyauchi, Y.; Tanaka,

K.;

[115986]Kamada, N.; Kusunose, M.: Human leukotriene B4 omega-hydroxylase

[115987](CYP4F3) gene: molecular cloning and chromosomal localization. DNA

[115988]Cell Biol. 17: 221-230, 1998.

[115989]20452.

[115990]Kikuta, Y.; Kusunose, E.; Endo, K.; Yamamoto, S.; Sogawa, K.; Fujii-Kuriyama,

[115991]Y.; Kusunose, M.: A novel form of cytochrome P-450 family 4 in human

[115992]polymorphonuclear leukocytes: cDNA cloning and expression of leukotriene

[115993]B4 omega-hydroxylase. J. Biol. Chem. 268: 9376-9380, 1993.

[115994]20453.

[115995]Hamra, F. K.; Forte, L. R.; Eber, S. L.; Pidhorodeckyj, N. V.;

[115996]Krause, W. J.; Freeman, R. H.; Chin, D. T.; Tompkins, J. A.; Fok,

[115997]K. F.; Smith, C. E.; Duffin, K. L.; Siegel, N. R.; Currie, M. G.:

[115998]Uroguanylin: structure and activity of a second endoge-

nous peptide

[115999] that stimulates intestinal guanylate cyclase. Proc. Nat.

Acad. Sci. 90:

[116000] 10464–10468, 1993.

[116001] 20454.

[116002] Kita, T.; Smith, C. E.; Fok, K. F.; Duffin, K. L.; Moore, W. M.;

[116003] Karabatsos, P. J.; Kachur, J. F.; Hamra, F. K.; Pid-
horodeckyj, N.

[116004] V.; Forte, L. R.; Currie, M. G.: Characterization of human
uroguanylin:

[116005] a member of the guanylin peptide family. Am. J. Physiol.
266: F342–F348,

[116006] 1994.

[116007] 20455.

[116008] Miyazato, M.; Nakazato, M.; Matsukura, S.; Kangawa, K.;

Matsuo,

[116009] H.: Genomic structure and chromosomal localization of
human uroguanylin. Genomics 43:

[116010] 359–365, 1997.

[116011] 20456.

[116012] Whitaker, T. L.; Steinbrecher, K. A.; Copeland, N. G.;

Gilbert,

[116013]D. J.; Jenkins, N. A.; Cohen, M. B.: The uroguanylin gene
(Guca1b)

[116014]is linked to guanylin (Guca2) on mouse chromosome 4.

Genomics 45:

[116015]348–354, 1997.

[116016]20457.

[116017]Haft, C. R.; de la Luz Sierra, M.; Barr, V. A.; Haft, D. H.;
Taylor,

[116018]S. I.: Identification of a family of sorting nexin molecules
and

[116019]characterization of their association with receptors. Molec.
Cell.

[116020]Biol. 18: 7278–7287, 1998.

[116021]20458.

[116022]Kurten, R. C.; Cadena, D. L.; Gill, G. N.: Enhanced degra-
dation

[116023]of EGF receptors by a sorting nexin, SNX1. Science 272:
1008–1010,

[116024]1996.

[116025]20459.

[116026]Zhong, Q.; Lazar, C. S.; Tronchere, H.; Sato, T.; Meerloo, T.;

[116027]Yeo, M.; Songyang, Z.; Emr, S. D.; Gill, G. N.: Endosomal localization

[116028]and function of sorting nexin 1. *Proc. Nat. Acad. Sci.* 99: 6767–6772,

[116029]2002.

[116030]20460.

[116031]Klingensmith, J.; Nusse, R.; Perrimon, N.: The *Drosophila* segment

[116032]polarity gene *dishevelled* encodes a novel protein required for response

[116033]to wingless signal. *Genes Dev.* 8: 118–130, 1994.

[116034]20461.

[116035]Holmes, S. E.; Riazzi, M. A.; Gong, W.; McDermid, H. E.; Sellinger,

[116036]B. T.; Hua, A.; Chen, F.; Wang, Z.; Zhang, G.; Roe, B.; Gonzalez,

[116037]J.; McDonald–McGinn, D. M.; Zackai, E.; Emanuel, B. S.; Budarf, M.

[116038]L.: Disruption of the clathrin heavy chain–like gene (CLTCL) associated

[116039]with features of DGS/VCFS: a balanced (21;22)(p12;q11) translocation. Hum.

[116040]Molec. Genet. 6: 357–367, 1997.

[116041]20462.

[116042]Kedra, D.; Peyrard, M.; Fransson, I.; Collins, J. E.; Dunham, I.;

[116043]Roe, B. A.; Dumanski, J. P.: Characterization of a second human clathrin

[116044]heavy chain polypeptide gene (CLH-22) from chromosome 22q11. Hum.

[116045]Molec. Genet. 5: 625–631, 1996.

[116046]20463.

[116047]Long, K. R.; Trofatter, J. A.; Ramesh, V.; McCormick, M. K.; Buckler,

[116048]A. J.: Cloning and characterization of a novel human clathrin heavy

[116049]chain gene (CLTCL). Genomics 35: 466–472, 1996.

[116050]20464.

[116051]Lagenaur, C.; Kunemund, V.; Fischer, G.; Fushiki, S.; Schachner,

[116052]M.: Monoclonal M6 antibody interferes with neurite exten-

sion of cultured

[116053]neurons. J. Neurobiol. 23: 71–88, 1992.

[116054]20465.

[116055]Shimizu, F.; Watanabe, T. K.; Fujiwara, T.; Takahashi, E.;
Nakamura,

[116056]Y.; Maekawa, H.: Isolation and mapping of the human gly-
coprotein

[116057]M6 gene (GPM6A) to 4q33–to–q34. Cytogenet. Cell Genet.
74: 138–139,

[116058]1996.

[116059]20466.

[116060]Bui, T. D.; Beier, D. R.; Jonssen, M.; Smith, K.; Dorrington,
S.

[116061]M.; Kaklamanis, L.; Kearney, L.; Regan, R.; Sussman, D. J.;
Harris,

[116062]A. L.: cDNA cloning of a human dishevelled DVL–3 gene,
mapping to

[116063]3q27, and expression in human breast and colon carcino-
mas. Biochem.

[116064]Biophys. Res. Commun. 239: 510–516, 1997.

[116065]20467.

[116066]Scheffner, M.; Huibregtse, J. M.; Vierstra, R. D.; Howley, P.

[116067]M.: The HPV-16 E6 and E6-AP complex functions as a ubiquitin-protein

[116068]ligase in the ubiquitination of p53. Cell 75: 495-505, 1993.

[116069]20468.

[116070]Cheng, Y.-S. E.; Patterson, C. E.; Staeheli, P.: Interferon-induced

[116071]guanylate-binding proteins lack an N(T)KXD consensus motif and bind

[116072]GMP in addition to GDP and GTP. Molec. Cell. Biol. 11: 4717-4725,

[116073]1991.

[116074]20469.

[116075]Kumar, S.; Li, Q.; Dua, A.; Ying, Y.-K.; Bagchi, M. K.; Bagchi,

[116076]I. C.: Messenger ribonucleic acid encoding interferon-inducible guanylate

[116077]binding protein 1 is induced in human endometrium within the putative

[116078]window of implantation. J. Clin. Endocr. Metab. 86: 2420-2427, 2001.

[116079]20470.

[116080]Prochazka, M.; Staeheli, P.; Holmes, R. S.; Haller, O.: Interferon-induced

[116081]guanylate-binding proteins: mapping of the murine Gbp-1 locus to chromosome

[116082]3. Virology 145: 273-279, 1985.

[116083]20471.

[116084]Strehlow, I.; Lohmann-Matthes, M. L.; Decker, T.: The interferon-inducible

[116085]GBP1 gene: structure and mapping to human chromosome 1. Gene 144:

[116086]295-299, 1994.

[116087]20472.

[116088]Scheffner, M.; Werness, B. A.; Huibregtse, J. M.; Levine, A. J.;

[116089]Howley, P. M.: The E6 oncoprotein encoded by human papillomavirus

[116090]types 16 and 18 promotes the degradation of p53. Cell 63: 1129-1136,

[116091]1990.

[116092]20473.

[116093]Tsai, T.-F.; Raas-Rothschild, A.; Ben-Neriah, Z.; Beaudet, A.

[116094]L.: Prenatal diagnosis and carrier detection for a point mutation

[116095]in UBE3A causing Angelman syndrome. (Letter) Am. J. Hum. Genet. 63:

[116096]1561–1563, 1998.

[116097]20474.

[116098]Yu, T. H.; Hoffman, A. R.: Imprinting of the Angelman syndrome

[116099]gene, UBE3A, is restricted to brain. (Letter) Nature Genet. 17:

[116100]12–13, 1997.

[116101]20475.

[116102]Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek,

[116103]F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.; Copeland, N.

[116104]G.: Murine chromosomal location of four hepatocyte-enriched transcription

[116105]factors: HNF-3-alpha, HNF3-beta, HNF-3-gamma, and HFN-4. Genomics 13:

[116106]264–268, 1992.

[116107]20476.

[116108]Chartier, F. L.; Bossu, J.-P.; Laudet, V.; Fruchart, J.-C.:

Cloning

[116109]and sequencing of cDNAs encoding the human hepatocyte
nuclear factor

[116110]4 indicate the presence of two isoforms in human liver.

Gene 147:

[116111]269–272, 1994.

[116112]20477.

[116113]Eeckhoutte, J.; Formstecher, P.; Laine, B.: Maturity-onset
diabetes

[116114]of the young type 1 (MODY1)–associated mutations R154X
and E276Q in

[116115]hepatocyte nuclear factor 4–alpha (HNF4–alpha) gene im-
pair recruitment

[116116]of p300, a key transcriptional coactivator. Molec. Endocr.
15: 1200–1210,

[116117]2001.

[116118]20478.

[116119]Furuta, H.; Iwasaki, N.; Oda, N.; Hinokio, Y.; Horikawa, Y.;

Yamagata,

[116120]K.; Yano, N.; Sugahiro, J.; Ogata, M.; Ohgawara, H.; Omori,
Y.; Iwamoto,

[116121]Y.; Bell, G. I.: Organization and partial sequence of the
hepatocyte

[116122]nuclear factor-4-alpha/MODY1 gene and identification of
a missense

[116123]mutation, R127W, in a Japanese family with MODY. Dia-
betes 46: 1652-1657,

[116124]1997.

[116125]20479.

[116126]Hani, E. H.; Suaud, L.; Boutin, P.; Chevre, J.-C.; Durand, E.;

[116127]Philippi, A.; Demenais, F.; Vionnet, N.; Furuta, H.; Velho,
G.; Bell,

[116128]G. I.; Laine, B.; Froguel, P.: A missense mutation in hepa-
tocyte

[116129]nuclear factor-4-alpha, resulting in a reduced transacti-
vation activity,

[116130]in human late-onset non-insulin-dependent diabetes
mellitus. J. Clin.

[116131]Invest. 101: 521-526, 1998.

[116132]20480.

- [116133]Lausen, J.; Thomas, H.; Lemm, I.; Bulman, M.;
Borgschulze, M.;
- [116134]Lingott, A.; Hattersley, A. T.; Ryffel, G. U.: Naturally occurring
ring
- [116135]mutations in the human HNF4- α gene impair the
function of the
- [116136]transcription factor to a varying degree. Nucleic Acids Res.
28:
- [116137]430-437, 2000.
- [116138]20481.
- [116139]Li, J.; Ning, G.; Duncan, S. A.: Mammalian hepatocyte differentiation
ferentiation
- [116140]requires the transcription factor HNF-4- α . Genes Dev.
14: 464-474,
- [116141]2000.
- [116142]20482.
- [116143]Stoffel, M.; Duncan, S. A.: The maturity-onset diabetes of
the
- [116144]young (MODY1) transcription factor HNF4- α regulates
expression
- [116145]of genes required for glucose transport and metabolism.
Proc. Nat.

- [116146]Acad. Sci. 94: 13209–13214, 1997.
- [116147]20483.
- [116148]Yamagata, K.; Furuta, H.; Oda, N.; Kaisaki, P. J.; Menzel, S.;
- [116149]Cox, N. J.; Fajans, S. S.; Signorini, S.; Stoffel, M.; Bell, G. I.
- [116150]; Mutations in the hepatocyte nuclear factor-4- α gene
in maturity-onset
- [116151]diabetes of the young (MODY1). Nature 384: 458–460,
1996.
- [116152]20484.
- [116153]Zouali, H.; Hani, E. H.; Philippi, A.; Vionnet, N.; Beckmann,
- [116154]J. S.; Demenais, F.; Froguel, P.: A susceptibility locus for
early-onset
- [116155]non-insulin dependent (type 2) diabetes mellitus maps to
chromosome
- [116156]20q, proximal to the phosphoenolpyruvate carboxykinase
gene. Hum.
- [116157]Molec. Genet. 6: 1401–1408, 1997.
- [116158]20485.
- [116159]Frolova, L.; Le Goff, X.; Rasmussen, H. H.; Cheperegine, S.;
- Drugeon,

[116160]G.; Kress, M.; Arman, I.; Haenni, A.-L.; Celis, J. E.; Philippe, M.;

[116161]Justesen, J.; Kisselev, L.: A highly conserved eukaryotic protein

[116162]family possessing properties of polypeptide chain release factor. Nature 372:

[116163]701–703, 1994.

[116164]20486.

[116165]Guenet, L.; Henry, C.; Toutain, B.; Dubourg, C.; Le Gall, J. Y.;

[116166]David, V.; Le Treut, A.: Eukaryotic translation termination factor

[116167]gene (ETF1/eRF1) maps at D5S500 in a commonly deleted region of chromosome

[116168]5q31 in malignant myeloid diseases. Cytogenet. Cell Genet. 88: 82–86,

[116169]2000.

[116170]20487.

[116171]Hansen, L. L.; Jakobsen, C. G.; Justesen, J.: Assignment of the

[116172]human translation termination factor 1 (ETF1) to 5q31.1 and of the

[116173]proximal marker D5S1995 by radiation hybrid mapping.

Cytogenet. Cell

[116174]Genet. 87: 256–257, 1999.

[116175]20488.

[116176]Le Goff, X.; Philippe, M.; Jean-Jean, O.: Overexpression of
human

[116177]release factor 1 alone has an antisuppressor effect in hu-
man cells. Molec.

[116178]Cell Biol. 17: 3164–3172, 1997.

[116179]20489.

[116180]Song, H.; Mugnier, P.; Das, A. K.; Webb, H. M.; Evans, D.
R.; Tuite,

[116181]M. F.; Hemmings, B. A.; Barford, D.: The crystal structure
of human

[116182]eukaryotic release factor eRF1--mechanism of stop codon recognition

[116183]and peptidyl-tRNA hydrolysis. Cell 100: 311-321, 2000.

[116184]20490.

[116185]Liu, Q.; Fischer, U.; Wang, F.; Dreyfuss, G.: The spinal muscular

[116186]atrophy disease gene product, SMN, and its associated protein SIP1

[116187]are in a complex with spliceosomal snRNP proteins. Cell 90: 1013-1021,

[116188]1997.

[116189]20491.

[116190]Meister, G.; Buhler, D.; Laggerbauer, B.; Zobawa, M.; Lottspeich,

[116191]F.; Fischer, U.: Characterization of a nuclear 20S complex containing

[116192]the survival of motor neurons (SMN) protein and a specific subset

[116193]of spliceosomal Sm proteins. Hum. Molec. Genet. 9: 1977-1986, 2000.

[116194]20492.

- [116195] Aramburu, J.; Garcia-Cozar, F.; Raghavan, A.; Okamura, H.; Rao, A.; Hogan, P. G.: Selective inhibition of NFAT activation by a peptide spanning the calcineurin targeting site of NFAT. Molec. Cell 1: 627–637, 1998.
- [116198] 20493.
- [116199] Castigli, E.; Pahwa, R.; Good, R. A.; Geha, R. S.; Chatila, T. A.: Molecular basis of a multiple lymphokine deficiency in a patient with severe combined immunodeficiency. Proc. Nat. Acad. Sci. 90: 4728–4732, 1993.
- [116203] 20494.
- [116205] Chatila, T.; Castigli, E.; Pahwa, R.; Pahwa, S.; Chirmule, N.; Oyaizu, N.; Good, R. A.; Geha, R. S.: Primary combined immunodeficiency resulting from defective transcription of multiple T-cell lymphokine genes. Proc. Nat. Acad. Sci. 87: 10033–10037, 1990.

[116209]20495.

[116210]Horsley, V.; Pavlath, G. K.: NFAT: ubiquitous regulator of cell

[116211]differentiation and adaptation. J. Cell Biol. 156: 771–774, 2002.

[116212]20496.

[116213]Jauliac, S.; Lopez-Rodriguez, C.; Shaw, L. M.; Brown, L. F.; Rao,

[116214]A.; Toker, A.: The role of NFAT transcription factors in integrin-mediated

[116215]carcinoma invasion. Nature Cell Biol. 4: 540–544, 2002.

[116216]20497.

[116217]Li, X.; Ho, S. N.; Luna, J.; Giacalone, J.; Thomas, D. J.; Timmerman,

[116218]L. A.; Crabtree, G. R.; Francke, U.: Cloning and chromosomal localization

[116219]of the human and murine genes for the T-cell transcription factors

[116220]NFATc and NFATp. Cytogenet. Cell Genet. 68: 185–191, 1995.

[116221]20498.

[116222]Northrop, J. P.; Ho, S. N.; Chen, L.; Thomas, D. J.; Timmer-
man,
[116223]L. A.; Nolan, G. P.; Admon, A.; Crabtree, G. R.: NF-AT
components
[116224]define a family of transcription factors targeted in T-cell
activation. Nature 369:
[116225]497-502, 1994.
[116226]20499.
[116227]Okamura, H.; Aramburu, J.; Garcia-Rodriguez, C.; Viola, J.
P. B.;
[116228]Raghavan, A.; Tashirani, M.; Zhang, X.; Qin, J.; Hogan, P.
G.; Rao,
[116229]A.: Concerted dephosphorylation of the transcription fac-
tor NFAT1
[116230]induces a conformational switch that regulates transcrip-
tional activity. Molec.
[116231]Cell 6: 539-550, 2000.
[116232]20500.
[116233]Pahwa, R.; Chatila, T.; Pahwa, S.; Paradise, C.; Day, N. K.;
Geha,
[116234]R.; Schwartz, S. A.; Slade, H.; Oyaizu, N.; Good, R. A.: Re-
combinant

[116235]interleukin 2 therapy in severe combined immunodeficiency disease. Proc.

[116236]Nat. Acad. Sci. 86: 5069–5073, 1989.

[116237]20501.

[116238]Park, J.; Takeuchi, A.; Sharma, S.: Characterization of a new

[116239]isoform of the NFAT (nuclear factor of activated T cells) gene family

[116240]member NFATc. J. Biol. Chem. 271: 20914–20921, 1996.

[116241]20502.

[116242]Peng, S. L.; Gerth, A. J.; Ranger, A. M.; Glimcher, L. H.: NFATc1

[116243]and NFATc2 together control both T and B cell activation and differentiation. Immunity 14:

[116244]13–20, 2001.

[116245]20503.

[116246]Aoki, K.; Suzuki, K.; Sugano, T.; Tasaka, T.; Nakahara, K.; Kuge,

[116247]O.; Omori, A.; Kasai, M.: A novel gene, 'Translin,' encodes a recombination

[116248]hotspot binding protein associated with chromosomal

translocations. Nature

[116249]Genet. 10: 167–174, 1995.

[116250]20504.

[116251]Badge, R. M.; Yardley, J.; Jeffreys, A. J.; Armour, J. A. L.:

[116252]Crossover breakpoint mapping identifies a subtelomeric
hotspot for

[116253]male meiotic recombination. Hum. Molec. Genet. 9:
1239–1244, 2000.

[116254]20505.

[116255]Hosaka, T.; Kanoe, H.; Nakayama, T.; Murakami, H.; Ya-
mamoto, H.;

[116256]Nakamata, T.; Tsuboyama, T.; Oka, M.; Kasai, M.; Sasaki,
M. S.; Nakamura,

[116257]T.; Toguchida, J.: Translin binds to the sequences adjacent
to the

[116258]breakpoints of the TLS and CHOP genes in liposarcomas
with translocation

[116259]t(12;16). Oncogene 19: 5821–5825, 2000.

[116260]20506.

[116261]Kasai, M.; Aoki, K.; Matsuo, Y.; Minowada, J.; Maziarz, R.
T.;

[116262]Strominger, J. L.: Recombination hotspot associated factors specifically

[116263]recognize novel target sequences at the site of interchromosomal rearrangements

[116264]in T-ALL patients with t(8;14)(q24;q11) and (t(1;14)(p32;q11). *Int.*

[116265]*Immun.* 6: 1017–1025, 1994.

[116266]20507.

[116267]Cirillo, L. A.; Lin, F. R.; Cuesta, I.; Friedman, D.; Jarnik, M.;

[116268]Zaret, K. S.: Opening of compacted chromatin by early developmental

[116269]transcription factors HNF3(FoxA) and GATA-4. *Molec. Cell* 9: 279–289,

[116270]2002.

[116271]20508.

[116272]Laitinen, M. P. E.; Anttonen, M.; Ketola, I.; Wilson, D. B.; Ritvos,

[116273]O.; Butzow, R.; Heikinheimo, M.: Transcription factors GATA-4 and

[116274]GATA-6 and a GATA family cofactor, FOG-2, are expressed in human ovary

[116275]and sex cord-derived ovarian tumors. J. Clin. Endocr.

Metab. 85:

[116276]3476–3483, 2000.

[116277]20509.

[116278]Freyd, G.; Kim, S. K.; Horvitz, H. R.: Novel cysteine-rich
motif

[116279]and homeodomain in the product of the *Caenorhabditis*
elegans cell

[116280]lineage gene *lin-11*. Nature 344: 876–879, 1990.

[116281]20510.

[116282]Mbikay, M.; Tadros, H.; Seidah, N. G.; Simpson, E. M.:
Linkage

[116283]mapping of the gene for the LIM-homeoprotein LIM3
(locus *Lhx3*) to

[116284]mouse chromosome 2. Mammalian Genome 6: 818–819,
1995.

[116285]20511.

[116286]Sharma, K.; Leonard, A. E.; Lettieri, K.; Pfaff, S. L.: Genetic

[116287]and epigenetic mechanisms contribute to motor neuron
pathfinding. Nature 406:

[116288]515–519, 2000.

[116289]20512.

[116290]Sharma, K.; Sheng, H. Z.; Lettieri, K.; Li, H.; Karavanov, A.;

[116291]Potter, S.; Westphal, H.; Pfaff, S. L.: LIM homeodomain factors Lhx3

[116292]and Lhx4 assign subtype identities for motor neurons.

Cell 95: 817–828,

[116293]1998.

[116294]20513.

[116295]Sheng, H. Z.; Zhadanov, A. B.; Mosinger, B., Jr.; Fujii, T.; Bertuzzi,

[116296]S.; Grinberg, A.; Lee, E. J.; Huang, S.–P.; Mahon, K. A.; Westphal,

[116297]H.: Specification of pituitary cell lineages by the LIM homeobox

[116298]gene Lhx3. Science 272: 1004–1007, 1996.

[116299]20514.

[116300]Sloop, K. W.; Meier, B. C.; Bridwell, J. L.; Parker, G. E.; Schiller,

[116301]A. M.; Rhodes, S. J.: Differential activation of pituitary hormone

[116302]genes by human Lhx3 isoforms with distinct DNA binding

properties. Molec.

[116303]Endocr. 13: 2212–2225, 1999.

[116304]20515.

[116305]Sloop, K. W.; Showalter, A. D.; Von Kap–Herr, C.; Pettenati,
M.

[116306]J.; Rhodes, S. J.: Analysis of the human LHX3 neuroen-
docrine transcription

[116307]factor gene and mapping to the subtelomeric region of
chromosome 9. Gene 245:

[116308]237–243, 2000.

[116309]20516.

[116310]Sloop, K. W.; Walvoord, E. C.; Showalter, A. D.; Pescovitz,
O.

[116311]H.; Rhodes, S. J.: Molecular analysis of LHX3 and PROP–1
in pituitary

[116312]hormone deficiency patients with posterior pituitary ec-
topia. J.

[116313]Clin. Endocr. Metab. 85: 2701–2708, 2000.

[116314]20517.

[116315]Brenneman, M. A.; Wagener, B. M.; Miller, C. A.; Allen, C.;
Nickoloff,

[116316]J. A.: XRCC3 controls the fidelity of homologous recombination: roles
for XRCC3 in late stages of recombination. Molec. Cell 10:
387–395,
2002.
20518.
20518.
Liu, N.; Lamerdin, J. E.; Tebbs, R. S.; Schild, D.; Tucker, J.
D.; Shen, M. R.; Brookman, K. W.; Siciliano, M. J.; Walter, C.
A.;
Fan, W.; Narayana, L. S.; Zhou, Z.-Q.; Adamson, A. W.;
Sorensen, K.
J.; Chen, D. J.; Jones, N. J.; Thompson, L. H.: XRCC2 and
XRCC3,
new human Rad51-family members, promote chromosome stability and protect
against DNA cross-links and other damages. Molec. Cell
1: 783–793,
1998.
20519.
Tebbs, R. S.; Zhao, Y.; Tucker, J. D.; Scheerer, J. B.; Siciliano,

[116329]M. J.; Hwang, M.; Liu, N.; Legerski, R. J.; Thompson, L. H.:

Correction

[116330]of chromosomal instability and sensitivity to diverse mutagens by

[116331]a cloned cDNA of the XRCC3 DNA repair gene. Proc. Nat. Acad. Sci. 92:

[116332]6354–6358, 1995.

[116333]20520.

[116334]Winsey, S. L.; Haldar, N. A.; Marsh, H. P.; Bunce, M.; Marshall,

[116335]S. E.; Harris, A. L.; Wojnarowska, F.; Welsh, K. I.: A variant within

[116336]the DNA repair gene XRCC3 is associated with the development of melanoma

[116337]skin cancer. Cancer Res. 60: 5612–5616, 2000.

[116338]20521.

[116339]Hayashida, S.; Yamasaki, K.; Asada, Y.; Soeda, E.; Niikawa, N.;

[116340]Kishino, T.: Construction of a physical and transcript map flanking

[116341]the imprinted MEST/PEG1 region at 7q32. Genomics 66: 221–225, 2000.

[116342]20522.

[116343]de Wind, N.; Dekker, M.; Claij, N.; Jansen, L.; van Klink, Y.;

[116344]Radman, M.; Riggins, G.; van der Valk, M.; van't Wout, K.;
te Riele,

[116345]H.: HNPCC-like cancer predisposition in mice through si-
multaneous

[116346]loss of Msh3 and Msh6 mismatch-repair protein func-
tions. Nature Genet. 23:

[116347]359-362, 1999.

[116348]20523.

[116349]Chen, Y. H.; Hansen, L.; Chen, M. X.; Bjorbaek, C.; Vester-
gaard,

[116350]H.; Hansen, T.; Cohen, P. T. W.; Pedersen, O.: Sequence of
the human

[116351]glycogen-associated regulatory subunit of type 1 protein
phosphatase

[116352]and analysis of its coding region and mRNA level in mus-
cle from patients

[116353]with NIDDM. Diabetes 43: 1234-1241, 1994.

[116354]20524.

[116355]Braun-Dullaes, R. C.; Mann, M. J.; Ziegler, A.; von der

Leyen,

[116356]H. E.; Dzaou, V. J.: A novel role for the cyclin-dependent kinase

[116357]inhibitor p27(Kip1) in angiotensin II-stimulated vascular smooth muscle

[116358]cell hypertrophy. J. Clin. Invest. 104: 815–823, 1999.

[116359]20525.

[116360]Carrano, A. C.; Eytan, E.; Hershko, A.; Pagano, M.: SKP2 is required

[116361]for ubiquitin-mediated degradation of the CDK inhibitor p27. Nature

[116362]Cell Biol. 1: 193–199, 1999.

[116363]20526.

[116364]Di Cristofano, A.; De Acetis, M.; Koff, A.; Cordon-Cardo, C.; Pandolfi,

[116365]P. P.: Pten and p27(KIP1) cooperate in prostate cancer tumor suppression

[116366]in the mouse. Nature Genet. 27: 222–224, 2001.

[116367]20527.

[116368]Fero, M. L.; Rivkin, M.; Tasch, M.; Porter, P.; Carow, C. E.; Firpo,

[116369]E.; Polyak, K.; Tsai, L.-H.; Broudy, V.; Perlmutter, R. M.;
Kaushansky,
[116370]K.; Roberts, J. M.: A syndrome of multiorgan hyperplasia
with features
[116371]of gigantism, tumorigenesis, and female sterility in
p27(Kip1)-deficient
[116372]mice. Cell 85: 733-744, 1996.
[116373]20528.

[116374]Gowen, M.; Lazner, F.; Dodds, R.; Kapadia, R.; Feild, J.;
Tavaria,
[116375]M.; Bertoncello, I.; Drake, F.; Zavarselk, S.; Tellis, I.; Hert-
zog,
[116376]P.; Debouck, C.; Kola, I.: Cathepsin K knockout mice de-
velop osteopetrosis
[116377]due to a deficit in matrix degradation but not demineral-
ization. J.
[116378]Bone Miner. Res. 14: 1654-1663, 1999.
[116379]20529.

[116380]Buyse, I. M.; Shao, G.; Huang, S.: The retinoblastoma pro-
tein
[116381]binds to RIZ, a zinc-finger protein that shares an epitope
with the

[116382]adenovirus E1A protein. Proc. Nat. Acad. Sci. 92:
4467-4471, 1995.

[116383]20530.

[116384]Buyse, I. M.; Takahashi, E.; Huang, S.: Physical mapping of
the

[116385]retinoblastoma interacting zinc finger gene RIZ to D1S228
on chromosome

[116386]1p36. Genomics 34: 119-121, 1996.

[116387]20531.

[116388]Chadwick, R. B.; Jiang, G.-L.; Bennington, G. A.; Yuan, B.;
Johnson,

[116389]C. K.; Stevens, M. W.; Niemann, T. H.; Peltomaki, P.;
Huang, S.; de

[116390]la Chapelle, A.: Candidate tumor suppressor RIZ is fre-
quently involved

[116391]in colorectal carcinogenesis. Proc. Nat. Acad. Sci. 97:
2662-2667,

[116392]2000.

[116393]20532.

[116394]Mock, B. A.; Coleman, M. P.; Huang, S.: Riz maps to distal
chromosome

[116395]4 near genes involved in tumorigenesis and nerve degeneration. Mammalian

[116396]Genome 7: 637 only, 1996.

[116397]20533.

[116398]Poetsch, M.; Dittberner, T.; Woenckhaus, C.: Frameshift mutations

[116399]of RIZ, but no point mutations in RIZ1 exons in malignant melanomas

[116400]with deletions in 1p36. Oncogene 21: 3038–3042, 2002.

[116401]20534.

[116402]Malek, N. P.; Sundberg, H.; McGrew, S.; Nakayama, K.; Kyriakidis,

[116403]T. R.; Roberts, J. M.: A mouse knock-in model exposes sequential

[116404]proteolytic pathways that regulate p27(Kip1) in G1 and S phase. Nature 413:

[116405]323–327, 2001.

[116406]20535.

[116407]Martin, E.; Cacheux, V.; Cave, H.; Lapierre, J. M.; Le\Paslier,

[116408]D.; Grandchamp, B.: Localization of the CDKN4/p27(Kip1)

gene to human

[116409]chromosome 12p12.3. Hum. Genet. 96: 668–670, 1995.

[116410]20536.

[116411]Mitsuhashi, T.; Aoki, Y.; Eksioglu, Y. Z.; Takahashi, T.;

Bhide,

[116412]P. G.; Reeves, S. A.; Caviness, V. S., Jr.: Overexpression of

p27(Kip1)

[116413]lengthens the G1 phase in a mouse model that targets in-

ducible gene

[116414]expression to central nervous system progenitor cells.

Proc. Nat.

[116415]Acad. Sci. 98: 6435–6440, 2001.

[116416]20537.

[116417]Polyak, K.; Lee, M.-H.; Erdjument-Bromage, H.; Koff, A.;

Roberts,

[116418]J. M.; Tempst, P.; Massague, J.: Cloning of p27(Kip1), a cy-

clin-dependent

[116419]kinase inhibitor and a potential mediator of extracellular

antimitogenic

[116420]signals. Cell 78: 59–66, 1994.

[116421]20538.

[116422]Saito, T.; Seki, N.; Hattori, A.; Hayashi, A.; Abe, M.; Araki,

[116423]R.; Fujimori, A.; Fukumura, R.; Kozuma, S.; Matsuda, Y.:

Structure,

[116424]expression profile, and chromosomal location of a mouse
gene homologous

[116425]to human DNA-PK(cs) interacting protein (KIP) gene.

Mammalian Genome 10:

[116426]315–317, 1999.

[116427]20539.

[116428]Sherr, C. J.; Roberts, J. M.: Inhibitors of mammalian G1 cy-
clin-dependent

[116429]kinases. Genes Dev. 9: 1149–1163, 1995.

[116430]20540.

[116431]Toyoshima, H.; Hunter, T.: p27, a novel inhibitor of G1 cy-
clin-Cdk

[116432]protein kinase activity, is related to p21. Cell 78: 67–74,
1994.

[116433]20541.

[116434]Khakh, B. S.; Zhou, X.; Sydes, J.; Galligan, J. J.; Lester, H.

[116435]A.: State-dependent cross-inhibition between transmit-
ter-gated cation

[116436]channels. Nature 406: 405–410, 2000.

[116437]20542.

[116438]Chen, J. D.; Evans, R. M.: A transcriptional co-repressor
that

[116439]interacts with nuclear hormone receptors. Nature 377:
454–457, 1995.

[116440]20543.

[116441]Fischle, W.; Dequiedt, F.; Hendzel, M. J.; Guenther, M. G.;
Lazar,

[116442]M. A.; Voelter, W.; Verdin, E.: Enzymatic activity associated
with

[116443]class II HDACs is dependent on a multiprotein complex
containing HDAC3

[116444]and SMRT/N-CoR. Molec. Cell 9: 45–57, 2002.

[116445]20544.

[116446]Horlein, A. J.; Naar, A. M.; Heinzel, T.; Torchia, J.; Gloss,
B.;

[116447]Kurokawa, R.; Ryan, A.; Kamel, Y.; Soderstrom, M.; Glass,
C. K.; Rosenfeld,

[116448]M. G.: Ligand-independent repression by the thyroid hor-
mone receptor

[116449]mediated by a nuclear receptor co-repressor. Nature 377:
397-403,

[116450]1995.

[116451]20545.

[116452]Hu, X.; Lazar, M. A.: The CoRNR motif controls the recruit-
ment

[116453]of corepressors by nuclear hormone receptors. Nature
402: 93-96,

[116454]1999.

[116455]20546.

[116456]Jiang, Q.; Galiegue-Zouitina, S.; Roumier, C.; Hildebrand,
M. P.;

[116457]Thomas, S.; Coignet, L. J.: Genomic organization and re-
fined mapping

[116458]of the human nuclear corepressor 2 (NCOR2)/ silencing
mediator of

[116459]retinoid and thyroid hormone receptor (SMRT) gene on
chromosome 12q24.3. Cytogenet.

[116460]Cell Genet. 92: 217-220, 2001.

[116461]20547.

[116462]Ordentlich, P.; Downes, M.; Xie, W.; Genin, A.; Spinner, N.

B.;

[116463]Evans, R. M.: Unique forms of human and mouse nuclear
receptor corepressor

[116464]SMRT. Proc. Nat. Acad. Sci. 96: 2639–2644, 1999.

[116465]20548.

[116466]Hansen, L.; Hansen, T.; Vestergaard, H.; BJORBAEK, C.; ECH-
wald,

[116467]S. M.; Clausen, J. O.; Chen, Y. H.; Chen, M. X.; Cohen, P. T.
W.;

[116468]Pedersen, O.: A widespread amino acid polymorphism at
codon 905 of

[116469]the glycogen-associated regulatory subunit of protein
phosphatase-1

[116470]is associated with insulin resistance and hypersecretion of
insulin. Hum.

[116471]Molec. Genet. 4: 1313–1320, 1995.

[116472]20549.

[116473]Hegele, R. A.; Harris, S. B.; Zinman, B.; Wang, J.; Cao, H.;
Hanley,

[116474]A. J. G.; Tsui, L.-C.; Scherer, S. W.: Variation in the
AU(AT)-rich

[116475]element within the 3-prime-untranslated region of

PPP1R3 is associated

[116476]with variation in plasma glucose in aboriginal Canadians.

J. Clin.

[116477]Endocr. Metab. 83: 3980–3983, 1998.

[116478]20550.

[116479]Savage, D. B.; Agostini, M.; Barroso, I.; Gurnell, M.; Luan,

J.;

[116480]Meirhaeghe, A.; Harding, A.-H.; Ihrke, G.; Rajanayagam,

O.; Soos,

[116481]M. A.; George, S.; Berger, D.; and 9 others: Digenic inheri-

tance

[116482]of severe insulin resistance in a human pedigree. Nature

Genet. 31:

[116483]379–384, 2002.

[116484]20551.

[116485]Tang, P. M.; Bondor, J. A.; Swiderek, K. M.; DePaoli-Roach,

A.

[116486]A.: Molecular cloning and expression of the regulatory

(RG1) subunit

[116487]of the glycogen-associated protein phosphatase. J. Biol.

Chem. 266:

[116488]15782–15789, 1991.

[116489]20552.

[116490]International Cystinuria Consortium: Functional analysis of

[116491]mutations in SLC7A9, and genotype-phenotype correlation in non-Type

[116492]I cystinuria. Hum. Molec. Genet. 10: 305-316, 2001.

[116493]20553.

[116494]International Cystinuria Consortium: Non-type I cystinuria caused

[116495]by mutations in SLC7A9, encoding a subunit (b(0,+)-AT) of rBAT. Nature

[116496]Genet. 23: 52-57, 1999.

[116497]20554.

[116498]Garcia, J. G.; Lazar, V.; Gilbert-McClain, L. I.; Gallagher, P.

[116499]J.; Verin, A. D.: Myosin light chain kinase in endothelium: molecular

[116500]cloning and regulation. Am. J. Resp. Cell Molec. Biol. 16: 489-494,

[116501]1997.

[116502]20555.

[116503]Giorgi, D.; Brand-Arpon, V.; Rouquier, S.: The functional

myosin

[116504]light chain kinase (MYLK) gene localizes with marker
D3S3552 on human

[116505]chromosome 3q21 in a greater than 5-Mb yeast artificial
chromosome

[116506]region and is not linked to olfactory receptor genes. Cyto-
genet.

[116507]Cell Genet. 92: 85-88, 2001.

[116508]20556.

[116509]Lazar, V.; Garcia, J. G. N.: A single human myosin light
chain

[116510]kinase gene (MLCK; MYLK) transcribes multiple nonmuscle
isoforms. Genomics 57:

[116511]256-267, 1999.

[116512]20557.

[116513]Potier, M.-C.; Chelot, E.; Pekarsky, Y.; Gardiner, K.;
Rossier,

[116514]J.; Turnell, W. G.: The human myosin light chain kinase
(MLCK) from

[116515]hippocampus: cloning, sequencing, expression, and local-
ization to

[116516]3cen-q21. Genomics 29: 562-570, 1995.

[116517]20558.

[116518]Walker, L. A.; MacDonald, J. A.; Liu, X.; Nakamoto, R. K.;
Haystead,

[116519]T. A. J.; Somlyo, A. V.; Somlyo, A. P.: Site-specific phos-
phorylation

[116520]and point mutations of telokin modulate its
Ca(2+)-desensitizing effect

[116521]in smooth muscle. J. Biol. Chem. 276: 24519-24524,
2001.

[116522]20559.

[116523]Watterson, D. M.; Schavocky, J. P.; Guo, L.; Weiss, C.;
Chlenski,

[116524]A.; Shirinsky, V. P.; Van Eldik, L. J.; Haiech, J.: Analysis of
the

[116525]kinase-related protein gene found at human chromosome
3q21 in a multi-gene

[116526]cluster: organization, expression, alternative splicing, and
polymorphic

[116527]marker. J. Cell. Biochem. 75: 481-491, 1999.

[116528]20560.

[116529]Druck, T.; Gu, Y.; Prabhala. G.; Cannizzaro, L. A.; Park,
S.-H.;

[116530] Huebner, K.; Keen, J. H.: Chromosome localization of human genes
[116531] for clathrin adaptor polypeptides AP2-beta and AP50 and
the clathrin-binding
[116532] protein, VCP. Genomics 30: 94-97, 1995.
[116533] 20561.
[116534] Liu, Q.; Feng, Y.; Forgac, M.: Activity and in vitro reassembly
of the coated vesicle (H⁺)-ATPase requires the 50-kDa
subunit of the
[116535] clathrin assembly complex AP-2. J. Biol. Chem. 269:
31592-31597,
[116537] 1994.
[116538] 20562.
[116539] Thuriereau, C.; Brosius, J.; Burne, C.; Jolles, P.; Keen, J. H.;
[116540] Mattaliano, R. J.; Chow, E. P.; Ramachandran, K. L.; Kirchhausen,
[116541] T.: Molecular cloning and complete amino acid sequence
of AP50, an
[116542] assembly protein associated with clathrin-coated vesicles.
DNA 7:

[116543]663–669, 1988.

[116544]20563.

[116545]Ponnambalam, S.; Robinson, M. S.; Jackson, A. P.; Peiperl, L.;

[116546]Parham, P.: Conservation and diversity in families of coated vesicle

[116547]adaptins. *J. Biol. Chem.* 265: 4814–4820, 1990.

[116548]20564.

[116549]Waelter, S.; Scherzinger, E.; Hasenbank, R.; Nordhoff, E.; Lurz,

[116550]R.; Goehler, H.; Gauss, C.; Sathasivam, K.; Bates, G. P.; Lehrach,

[116551]H.; Wanker, E. E.: The huntingtin interacting protein HIP1 is a clathrin

[116552]and alpha-adaptin-binding protein involved in receptor-mediated endocytosis. *Hum.*

[116553]*Molec. Genet.* 10: 1807–1817, 2001.

[116554]20565.

[116555]Kaneko-Ishino, T.; Kuroiwa, Y.; Miyoshi, N.; Kohda, T.; Suzuki,

[116556]R.; Yokoyama, M.; Viville, S.; Barton, S. C.; Ishino, F.;

Surani,

[116557]M. A.: Peg1/Mest imprinted gene on chromosome 6 identified by cDNA

[116558]subtraction hybridization. Nature Genet. 11: 52–59, 1995.

[116559]20566.

[116560]Kobayashi, S.; Kohda, T.; Miyoshi, N.; Kuroiwa, Y.; Aisaka, K.;

[116561]Tsutsumi, O.; Kaneko-Ishino, T.; Ishino, F.: Human PEG1/MEST, an

[116562]imprinted gene on chromosome 7. Hum. Molec. Genet. 6: 781–786, 1997.

[116563]20567.

[116564]Kosaki, K.; Kosaki, R.; Craigen, W. J.; Matsuo, N.: Isoform-specific

[116565]imprinting of the human PEG1/MEST gene. (Letter) Am. J. Hum. Genet. 66:

[116566]309–312, 2000.

[116567]20568.

[116568]Kotzot, D.; Schmitt, S.; Bernasconi, F.; Robinson, W. P.; Lurie,

[116569]J. W.; Ilyina, H.; Mehes, K.; Hamel, B. C. J.; Otten, B. J.;

Hergersberg,

[116570]M.; Werder, E.; Shoenle, E.; Schinzel, A.: Uniparental disomy 7 in

[116571]Silver–Russell syndrome and primordial growth retardation. Hum. Molec.

[116572]Genet. 4: 583–587, 1995.

[116573]20569.

[116574]Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Keverne,

[116575]E. B.; Surani, M. A.: Abnormal maternal behaviour and growth retardation

[116576]associated with loss of the imprinted gene Mest. Nature Genet. 20:

[116577]163–169, 1998.

[116578]20570.

[116579]Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Surani, M.

[116580]A.: Genomic structure and parent–of–origin–specific methylation of

[116581]Peg1. Hum. Molec. Genet. 6: 1907–1915, 1997.

[116582]20571.

[116583]Ayala–Madrigal, M. L.; Doerr, S.; Ramirez–Duenas, M. L.;

Hansmann,

[116584]J.: Assignment of karyopherin alpha 1 (KPNA1) to human chromosome

[116585]band 3q21 by in situ hybridization. Cytogenet. Cell Genet. 90: 58-59,

[116586]2000.

[116587]20572.

[116588]Conti, E.; Uy, M.; Leighton, L.; Blobel, G.; Kuriyan, J.: Crystallographic

[116589]analysis of the recognition of a nuclear localization signal by the

[116590]nuclear import factor karyopherin alpha. Cell 94: 193-204, 1998.

[116591]20573.

[116592]Cortes, P.; Ye, Z.-S.; Baltimore, D.: RAG-1 interacts with the

[116593]repeated amino acid motif of the human homologue of the yeast protein

[116594]SRP1. Proc. Nat. Acad. Sci. 91: 7633-7637, 1994.

[116595]20574.

[116596]Aplan, P. D.; Johnson, B. E.; Russell, E.; Chervinsky, D. S.;

Kirsch,

[116597]J. R.: Cloning and characterization of TCTA, a gene located
at the

[116598]site of a t(1;3) translocation. Cancer Res. 55: 1917–1921,
1995.

[116599]20575.

[116600]Ghetti, A.; Pinol-Roma, S.; Michael, W. M.; Morandi, C.;
Dreyfuss,

[116601]G.: HNRNP I, the polypyrimidine tract-binding protein:
distinct nuclear

[116602]localization and association with hnRNAs. Nucleic Acids
Res. 20:

[116603]3671–3678, 1992.

[116604]20576.

[116605]Raimondi, E.; Romanelli, M. G.; Moralli, D.; Gamberi, C.;
Russo,

[116606]M. P.; Morandi, C.: Assignment of the human gene encod-
ing heterogeneous

[116607]nuclear RNA ribonucleoprotein I (PTB) to chromosome
14q23–q24.1. Genomics 27:

[116608]553–555, 1995.

[116609]20577.

[116610]Romanelli, M. G.; Lorenzi, P.; Morandi, C.: Organization of the

[116611]human gene encoding heterogeneous nuclear ribonucleo-
protein type I

[116612](hnRNP I) and characterization of hnRNP I related pseudo-
gene. Gene 255:

[116613]267-272, 2000.

[116614]20578.

[116615]Hirota, H.; Chen, J.; Betz, U. A. K.; Rajewsky, K.; Gu, Y.;
Ross,

[116616]J., Jr.; Muller, W.; Chien, K. R.: Loss of a gp130 cardiac
muscle

[116617]cell survival pathway is a critical event in the onset of
heart failure

[116618]during biochemical stress. Cell 97: 189-198, 1999.

[116619]20579.

[116620]Kidd, V. J.; Nesbitt, J. E.; Fuller, G. M.: Chromosomal local-
ization

[116621]of the IL-6 receptor signal transducing subunit, gp130
(IL6ST). Somat.

[116622]Cell Molec. Genet. 18: 477-483, 1992.

[116623]20580.

[116624]Kishimoto, T.; Taga, T.; Akira, S.: Cytokine signal trans-
duction. Cell 76:

[116625]253–262, 1994.

[116626]20581.

[116627]Rodriguez, C.; Grosgeorge, J.; Nguyen, V. C.; Gaudray, P.;
Theillet,

[116628]C.: Human gp130 transducer chain gene (IL6ST) is local-
ized to chromosome

[116629]band 5q11 and possesses a pseudogene on chromosome
band 17p11. Cytogenet.

[116630]Cell Genet. 70: 64–67, 1995.

[116631]20582.

[116632]Haagerup, A.; Hertz, J. M.; Christensen, M. F.; Binderup,
H.; Kruse,

[116633]T. A.: Cathepsin K gene mutations and 1q21 haplotypes in
patients

[116634]with pycnodysostosis in an outbred population. Europ. J.
Hum. Genet. 8:

[116635]431–436, 2000.

[116636]20583.

- [116637]Ho, N.; Punturieri, A.; Wilkin, D.; Szabo, J.; Johnson, M.;
Whaley,
[116638]J.; Davis, J.; Clark, A.; Weiss, S.; Francomano, C.: Mutations
of
[116639]CTSK result in pycnodysostosis via a reduction in cathep-
sin K protein. J.
[116640]Bone Miner. Res. 14: 1649–1653, 1999.
[116641]20584.
[116642]Inaoka, T.; Bilbe, G.; Ishibashi, O.; Tezuka, K.; Kumegawa,
M.;
[116643]Kokubo, T.: Molecular cloning of human cDNA for cathep-
sin K: novel
[116644]cysteine proteinase predominantly expressed in bone.
Biochem. Biophys.
[116645]Res. Commun. 206: 89–96, 1995.
[116646]20585.
[116647]Johnson, M. R.; Polymeropoulos, M. H.; Vos, H. L.; Ortiz de
Luna,
[116648]R. I.; Francomano, C. A.: A nonsense mutation in the
cathepsin K
[116649]gene observed in a family with pycnodysostosis. Genome
Res. 6: 1050–1055,

[116650]1996.

[116651]20586.

[116652]Hoogenraad, C. C.; Koekkoek, B.; Akhmanova, A.; Krugers, H.; Dortland,

[116653]B.; Miedema, M.; van Alphen, A.; Kistler, W. M.; Jaegle, M.; Koutsourakis,

[116654]M.; Van Camp, N.; Verhoye, M.; van der Linden, A.; Kaverina, I.; Grosveld,

[116655]F.; De Zeeuw, C. I.; Galjart, N.: Targeted mutation of Cyln2 in the

[116656]Williams syndrome critical region links CLIP-115 haploinsufficiency

[116657]to neurodevelopmental abnormalities in mice. Nature Genet. 32: 116-127,

[116658]2002. Note: Erratum: Nature Genet. 32: 331 only, 2002.

[116659]20587.

[116660]Bannerman, R. M.; Edwards, J. A.; Kreimer-Birnbaum, M.; McFarland,

[116661]E.; Russell, E. S.: Hereditary microcytic anaemia in the mouse; studies

[116662]in iron distribution and metabolism. Brit. J. Haemat. 23: 235-245,

[116663]1972.

[116664]20588.

[116665]Silverstein, A. M.: Essential hypocomplementemia: report
of a

[116666]case. Blood 16: 1338–1341, 1960.

[116667]20589.

[116668]Hintner, H.; Wolff, K.: Generalized atrophic benign epider-
molysis

[116669]bullosa. Arch. Derm. 118: 375–384, 1982.

[116670]20590.

[116671]McGrath, J. A.; Gatalica, B.; Christiano, A. M.; Li, K.;
Owaribe,

[116672]K.; McMillan, J. R.; Eady, R. A. J.; Uitto, J.: Mutations in the

[116673]180-kD bullous pemphigoid antigen (BPAG2), a
hemidesmosomal transmembrane

[116674]collagen (COL17A1), in generalized atrophic benign epi-
dermolysis bullosa. Nature

[116675]Genet. 11: 83–86, 1995.

[116676]20591.

[116677]Vincent, J. B.; Herbrick, J.-A.; Gurling, H. M. D.; Bolton, P.

[116678]F.; Roberts, W.; Scherer, S. W.: Identification of a novel
gene on

[116679]chromosome 7q31 that is interrupted by a translocation
breakpoint

[116680]in an autistic individual. Am. J. Hum. Genet. 67: 510–514,
2000.

[116681]20592.

[116682]Vourc'h, P.; Bienvenu, T.; Beldjord, C.; Chelly, J.;
Barthelemy,

[116683]C.; Muh, J. P.; Andres, C.: No mutations in the coding re-
gion of

[116684]the Rett syndrome gene MECP2 in 59 autistic patients. Eu-
rop. J. Hum.

[116685]Genet. 9: 556–558, 2001.

[116686]20593.

[116687]Jaspers, M.; Zhang, Z.; Marynen, P.; Vekemans, S.; Aly, M.
S.;

[116688]Cuppens, H.; Hillicker, C.; Cassiman, J.-J.: Localization of
the

[116689]genes encoding the alpha-2 and alpha-4 subunits of the
human VLA-receptors

[116690]to chromosome 5q23–31 and 2q31–32 respectively.

(Abstract) Cytogenet.

[116691]Cell Genet. 58: 1870 only, 1991.

[116692]20594.

[116693]Kaplan, C.; Morel-Kopp, M. C.; Kroll, H.; Kiefel, V.;
Schlegel,

[116694]N.; Chesnel, N.; Mueller-Eckhardt, C.: HPA-5b (Br-a)
neonatal alloimmune

[116695]thrombocytopenia: clinical and immunological analysis of
39 cases. Brit.

[116696]J. Haemat. 78: 425-429, 1991.

[116697]20595.

[116698]Kiefel, V.; Santoso, S.; Katzmann, B.; Mueller-Eckhardt, C.:
A

[116699]new platelet-specific alloantigen Br(a): report of 4 cases
with neonatal

[116700]alloimmune thrombocytopenia. Vox Sang. 54: 101-106,
1988.

[116701]20596.

[116702]Kiefel, V.; Shechter, Y.; Atias, D.; Kroll, H.; Santoso, S.;
Mueller-Eckhardt,

[116703]C.: Neonatal alloimmune thrombocytopenia due to anti-

Br(b) (HPA-5a):

[116704]report of three cases in two families. Vox Sang. 60:
244-245, 1991.

[116705]20597.

[116706]Kritzik, M.; Savage, B.; Nugent, D. J.; Santoso, S.; Ruggeri,
Z.

[116707]M.; Kunicki, T. J.: Nucleotide polymorphisms in the alpha-
2 gene

[116708]define multiple alleles that are associated with differences
in platelet

[116709]alpha-2/beta-1 density. Blood 92: 2382-2388, 1998.

[116710]20598.

[116711]Kunicki, T. J.; Kritzik, M.; Annis, D. S.; Nugent, D. J.:
Hereditary

[116712]variation in platelet integrin alpha-2-beta-1 density is as-
sociated

[116713]with two silent polymorphisms in the alpha-2 gene coding
sequence. Blood 89:

[116714]1939-1943, 1997.

[116715]20599.

[116716]Mueller-Eckhardt, C.; Kiefel, V.; Grubert, A.; Kroll, H.;

Weisheit,

[116717]M.; Schmidt, S.; Mueller-Eckhardt, G.; Santoso, S.: 348

cases of

[116718]suspected neonatal alloimmune thrombocytopenia. Lancet

I: 363-366,

[116719]1989.

[116720]20600.

[116721]Nieuwenhuis, H. K.; Akkerman, J. W. N.; Houdijk, W. P.;

Sixma,

[116722]J. J.: Human blood platelets showing no response to colla-

gen fail

[116723]to express surface glycoprotein Ia. Nature 318: 470-472,

1985.

[116724]20601.

[116725]Santoso, S.; Amrhein, J.; Hofmann, H. A.; Sachs, U. J. H.;

Walka,

[116726]M. M.; Kroll, H.; Kiefel, V.: A point mutation thr799met on

the alpha-2

[116727]integrin leads to the formation of new human platelet al-

loantigen

[116728]Sit(a) and affects collagen-induced aggregation. Blood 94:

4103-4111,

[116729]1999.

[116730]20602.

[116731]Santoso, S.; Kalb, R.; Walka, M.; Kiefel, V.; Mueller-Eckhardt,

[116732]C.; Newman, P. J.: The human platelet alloantigens Br(a) and Br(b)

[116733]are associated with a single amino acid polymorphism on glycoprotein

[116734]Ia (integrin subunit alpha-2). J. Clin. Invest. 92: 2427-2432, 1993.

[116735]20603.

[116736]Santoso, S.; Kunicki, T. J.; Kroll, H.; Haberbosch, W.; Gardemann,

[116737]A.: Association of the platelet glycoprotein Ia C807T gene polymorphism

[116738]with nonfatal myocardial infarction in younger patients. Blood 93:

[116739]2449-2453, 1999.

[116740]20604.

[116741]Takada, Y.; Hemler, M. E.: The primary structure of the VLA-2/collagen

[116742]receptor alpha-2 subunit (platelet GPIa): homology to
other integrins

[116743]and the presence of a possible collagen-binding domain.

J. Cell Biol. 109:

[116744]397-407, 1989.

[116745]20605.

[116746]von Beckerath, N.; Koch, W.; Mehilli, J.; Bottiger, C.;
Schomig,

[116747]A.; Kastrati, A.: Glycoprotein Ia gene C807T polymorphism
and risk

[116748]for major adverse cardiac events within the first 30 days
after coronary

[116749]artery stenting. Blood 95: 3297-3301, 2000.

[116750]20606.

[116751]Fernandez-Ruiz, E.; Pardo-Manuel de Villena, F.; Rubio, M.
A.;

[116752]Corbi, A. L.; Rodriguez de Cordoba, S.; Sanchez-Madrid,
F.: Mapping

[116753]of the human VLA-alpha-4 gene to chromosome
2q31-q32. Europ. J. Immun. 22:

[116754]587-590, 1992.

[116755]20607.

[116756]Roseblatt, M.; Vuillet-Gaugler, M. H.; Leroy, C.;
Coulombel, L.

[116757]: Coexpression of two fibronectin receptors, VLA-4 and
VLA-5, by immature

[116758]human erythroblastic precursor cells. J. Clin. Invest. 87:
6-11,

[116759]1991.

[116760]20608.

[116761]Zhang, Z.; Vekemans, S.; Aly, M. S.; Jaspers, M.; Marynen,
P.;

[116762]Cassiman, J.-J.: The gene for the alpha-4 subunit of the
VLA-4 integrin

[116763]maps to chromosome 2q31-32. Blood 78: 2396-2399,
1991.

[116764]20609.

[116765]Hiesberger, T.; Trommsdorff, M.; Howell, B. W.; Goffinet,
A.; Mumby,

[116766]M. C.; Cooper, J. A.; Herz, J.: Direct binding of reelin to
VLDL

[116767]receptor and apoE receptor 2 induces tyrosine phosphory-
lation of disabled-1

[116768]and modulates tau phosphorylation. Neuron 24: 481–489, 1999.

[116769]20610.

[116770]Trommsdorff, M.; Gotthardt, M.; Hiesberger, T.; Shelton, J.; Stockinger,

[116771]W.; Nimpf, J.; Hammer, R. E.; Richardson, J. A.; Herz, J.: Reeler/Disabled-like

[116772]disruption of neuronal migration in knockout mice lacking the VLDL

[116773]receptor and ApoE receptor 2. Cell 97: 689–701, 1999.

[116774]20611.

[116775]Archidiacono, N.; Grimaldi, G.; Rocchi, M.; Romeo, G.: Mapping

[116776]of three sequenced clones coding for Zn(++) finger proteins. Cytogenet.

[116777]Cell Genet. 51: 952 only, 1989.

[116778]20612.

[116779]Miller, J.; McLachlan, A. D.; Klug, A.: Repetitive zinc-binding

[116780]domains in the protein transcription factor IIIA from *Xenopus* oocytes. EMBO

[116781]J. 4: 1609–1614, 1985.

[116782]20613.

[116783]Rocchi, M.; Gentile, E.; Rosati, M.; Grimaldi, G.: The human
KRAB/FPB

[116784]containing zinc finger gene ZNF2 maps to chromosome
2q11.2. Cytogenet.

[116785]Cell Genet. 86: 305–306, 1999.

[116786]20614.

[116787]Lloyd, S. L.; Sargent, C. A.; Chalmers, J.; Lim, E.; Habeebu,
S.

[116788]S. M.; Affara, N. A.: An X-linked zinc finger gene mapping
to Xq21.1–q21.3

[116789]closely related to ZFX and ZFY: possible origins from a
common ancestral

[116790]gene. Nucleic Acids Res. 19: 4835–4841, 1991.

[116791]20615.

[116792]Rousseau-Merck, M. F.; Tunnacliffe, A.; Berger, R.; Ponder,
B.

[116793]A. J.; Thiesen, H. J.: A cluster of expressed zinc finger pro-
tein

[116794]genes in the pericentromeric region of human chromo-

some 10. Genomics 13:

[116795]845–848, 1992.

[116796]20616.

[116797]Huebner, K.; Druck, T.; Croce, C. M.; Thiesen, H. J.:

Twenty-seven

[116798]nonoverlapping zinc finger cDNAs from human T cells

map to nine different

[116799]chromosomes with apparent clustering. Am. J. Hum.

Genet. 48: 726–740,

[116800]1991.

[116801]20617.

[116802]Rousseau-Merck, M.–F.; Hillion, J.; Jonveaux, P.; Berger,

R.; Thiesen,

[116803]H. J.: Chromosomal localization of two human zinc finger
protein

[116804]genes (Kox), to chromosome 17p13p12. (Abstract) Cyto-
genet. Cell Genet. 58:

[116805]2009, 1991.

[116806]20618.

[116807]Rousseau-Merck, M.–F.; Hillion, J.; Jonveaux, P.; Couillin,
P.;

[116808]Seite, P.; Thiesen, H.-J.; Berger, R.: Chromosomal localization of

[116809]9 KOX zinc finger genes: physical linkages suggest clustering of KOX

[116810]genes on chromosomes 12, 16, and 19. Hum. Genet. 92: 583-587, 1993.

[116811]20619.

[116812]Bray, P.; Lichter, P.; Thiesen, H.-J.; Ward, D. C.; Dawid, I. B.

[116813]: Characterization and mapping of human genes encoding zinc finger

[116814]proteins. Proc. Nat. Acad. Sci. 88: 9563-9567, 1991.

[116815]20620.

[116816]Naylor, S. L.; Klebe, R. J.; Shows, T. B.: Argininosuccinic aciduria:

[116817]assignment of the argininosuccinate lyase gene to the pter-q22 region

[116818]of human chromosome 7 by bioautography. Proc. Nat. Acad. Sci. 75:

[116819]6159-6162, 1978.

[116820]20621.

[116821]Pastural, E.; Ersoy, F.; Yalman, N.; Wulffraat, N.; Grillo, E.;

[116822]Ozkinay, F.; Tezcan, I.; Gedikoglu, G.; Philippe, N.; Fischer, A.;

[116823]de Saint Basile, G.: Two genes are responsible for Griscelli syndrome

[116824]at the same 15q21 locus. Genomics 63: 299–306, 2000.

[116825]20622.

[116826]Lehmann, O. J.; El-ashry, M. F.; Ebenezer, N. D.; Ocaka, L.; Francis,

[116827]P. J.; Wilkie, S. E.; Patel, R. J.; Ficker, L.; Jordan, T.; Khaw,

[116828]P. T.; Bhattacharya, S. S.: A novel keratocan mutation causing autosomal

[116829]recessive cornea plana. Invest. Ophthalm. Vis. Sci. 42: 3118–3122,

[116830]2001.

[116831]20623.

[116832]Hayasaka, K.; Tada, K.; Kikuchi, G.; Winter, S.; Nyhan, W. L.:

[116833]Nonketotic hyperglycinemia: two patients with primary defects of P-protein

[116834]and T-protein, respectively, in the glycine cleavage system. Pediat.

[116835]Res. 17: 967-970, 1983.

[116836]20624.

[116837]Kure, S.; Mandel, H.; Rolland, M.-O.; Sakata, Y.; Shinka, T.;
Drugan,

[116838]A.; Boneh, A.; Tada, K.; Matsubara, Y.; Narisawa, K.: A
missense

[116839]mutation (his42arg) in the T-protein gene from a large Is-
raeli-Arab

[116840]kindred with nonketotic hyperglycinemia. Hum. Genet.
102: 430-434,

[116841]1998.

[116842]20625.

[116843]Kure, S.; Shinka, T.; Sakata, Y.; Osamu, N.; Takayanagi, M.;
Tada,

[116844]K.; Matsubara, Y.; Narisawa, K.: A one-base deletion
(183delC) and

[116845]a missense mutation (D276H) in the T-protein gene from
a Japanese

[116846]family with nonketotic hyperglycinemia. J. Hum. Genet.
43: 135-137,

[116847]1998.

[116848]20626.

[116849]Nanao, K.; Okamura-Ikeda, K.; Motokawa, Y.; Danks, D.
M.; Baumgartner,

[116850]E. R.; Takada, G.; Hayasaka, K.: Identification of the muta-
tions

[116851]in the T-protein gene causing typical and atypical nonke-
totic hyperglycinemia. Hum.

[116852]Genet. 93: 655-658, 1994.

[116853]20627.

[116854]Nanao, K.; Takada, G.; Takahashi, E.; Seki, N.; Komatsu,
Y.; Okamura-Ikeda,

[116855]K.; Motokawa, Y.; Hayasaka, K.: Structure and chromoso-
mal localization

[116856]of the aminomethyltransferase gene (AMT). Genomics 19:
27-30, 1994.

[116857]20628.

[116858]Toone, J. R.; Applegarth, D. A.; Coulter-Mackie, M. B.;
James,

[116859]E. R.: Biochemical and molecular investigations of patients
with

[116860]nonketotic hyperglycemia. Molec. Genet. Metab. 70:
116-121, 2000.

[116861]20629.

[116862]Toone, J. R.; Applegarth, D. A.; Coulter-Mackie, M. B.;
James,

[116863]E. R.: Identification of the first reported splice site mutation

[116864](IVS7-1G-A) in the aminomethyltransferase (T-protein) gene (AMT) of

[116865]the glycine cleavage complex in 3 unrelated families with nonketotic

[116866]hyperglycinemia. (Abstract) Hum. Mutat. 17: 76 only, 2000.

[116867]20630.

[116868]Fujiwara, K.; Okamura-Ikeda, K.; Hayasaka, K.; Motokawa, Y.: The

[116869]primary structure of human H-protein of the glycine cleavage system

[116870]deduced by cDNA cloning. Biochem. Biophys. Res. Commun. 176: 711-716,

[116871]1991.

[116872]20631.

[116873]Hiraga, K.; Kochi, H.; Hayasaka, K.; Kikuchi, G.; Nyhan, W. L.

[116874]: Defective glycine cleavage system in non-ketotic hyperglycinemia:

[116875]occurrence of a less active glycine decarboxylase and an abnormal

[116876]aminomethyl carrier protein. J. Clin. Invest. 68: 525-534, 1981.

[116877]20632.

[116878]Hiraga, K.; Kure, S.; Yamamoto, M.; Ishiguro, Y.; Suzuki, T.:

[116879]Cloning of cDNA encoding human H-protein, a constituent of the glycine

[116880]cleavage system. Biochem. Biophys. Res. Commun. 151: 758-762, 1988.

[116881]20633.

[116882]Koyata, H.; Hiraga, K.: The glycine cleavage system: structure

[116883]of a cDNA encoding human H-protein, and partial characterization of

[116884]its gene in patients with hyperglycinemias. Am. J. Hum. Genet. 48:

[116885]351-361, 1991.

[116886]20634.

[116887]O'Brien, W. E.; McInnes, R.; Kalumuck, K.; Adcock, M.:

Cloning

[116888]and sequence analysis of cDNA for human argininosuccinate lyase. Proc.

[116889]Nat. Acad. Sci. 83: 7211–7215, 1986.

[116890]20635.

[116891]Piatigorsky, J.; O'Brien, W. E.; Norman, B. L.; Kalumuck, K.;

[116892]Wistow, G. J.; Borras, T.; Nickerson, J. M.; Wawrousek, E.
F.: Gene

[116893]sharing by delta-crystallin and argininosuccinate lyase.
Proc. Nat.

[116894]Acad. Sci. 85: 3479–3483, 1988.

[116895]20636.

[116896]Piatigorsky, J.; Wistow, G.: The recruitment of crystallins:

[116897]new functions precede gene duplication. Science 252:
1078–1079,

[116898]1991.

[116899]20637.

[116900]Pijpers, L.; Kleijer, W. J.; Reuss, A.; Jahoda, M. G. J.; Los,

[116901]F. J.; Sachs, E. S.; Wladimiroff, J. W.: Transabdominal

chorionic

[116902] villus sampling in a multiple pregnancy at risk of argininosuccinic

[116903] aciduria: a case report. Am. J. Med. Genet. 36: 449–450, 1990.

[116904] 20638.

[116905] Qureshi, I. A.; Letarte, J.; Ouellet, R.; Lemieux, B.: Enzymologic

[116906] and metabolic studies in two families affected by argininosuccinic

[116907] aciduria. Pediat. Res. 12: 256–262, 1978.

[116908] 20639.

[116909] Shih, V. E.; Efron, M. L.: Urea cycle disorders. In: Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The Metabolic Basis

[116911] of Inherited Disease. New York: McGraw–Hill (pub.) (3rd ed.)

[116912]: 1972. Pp. 370–392.

[116913] 20640.

[116914] Shih, V. E.; Littlefield, J. W.; Moser, H. W.: Argininosuccinase

- [116915]deficiency in fibroblasts cultured from patients with
argininosuccinic
aciduria. *Biochem. Genet.* 3: 81–83, 1969.
- [116917]20641.
- [116918]Simard, L.; O'Brien, W. E.; McInnes, R. R.: Argininosuccinate
lyase deficiency: evidence for heterogeneous structural
gene mutations
by immunoblotting. *Am. J. Hum. Genet.* 39: 38–51, 1986.
- [116921]20642.
- [116922]Todd, S.; McGill, J. R.; McCombs, J. L.; Moore, C. M.; Weider,
J.; Naylor, S. L.: cDNA sequence, interspecies comparison
and gene
mapping analysis of argininosuccinate lyase. *Genomics* 4:
53–59,
1989.
- [116926]20643.
- [116927]Van der Heiden, C.; Gerards, L. J.; van Biervliet, J. P. G. M.;
Desplanque, J.; DeBree, P. K.; Van Sprang, F. J.; Wadman,
S. K.:

[116929] Lethal neonatal argininosuccinate lyase deficiency in four children

[116930] from one sibship. *Helv. Paediat. Acta* 31: 407-417, 1976.

[116931] 20644.

[116932] Walker, D. C.; McCloskey, D. A.; Simard, L. R.; McInnes, R. R.

[116933]: Molecular analysis of human argininosuccinate lyase (ASAL): mutant

[116934] characterization and alternate splicing of the active site. (Abstract) *Am.*

[116935] *J. Hum. Genet.* 45 (suppl.): A227 only, 1989.

[116936] 20645.

[116937] Walker, D. C.; McCloskey, D. A.; Simard, L. R.; McInnes, R. R.

[116938]: Identification of a mutation frequently involved in inter-allelic

[116939] complementation at the human argininosuccinic acid lyase locus. (Abstract) *Am.*

[116940] *J. Hum. Genet.* 47 (suppl.): A169 only, 1990.

[116941] 20646.

[116942] Walker, D. C.; McCloskey, D. A.; Simard, L. R.; McInnes, R.

R.

[116943] Molecular analysis of human argininosuccinate lyase:
mutant characterization

[116944] and alternative splicing of the coding region. Proc. Nat.
Acad. Sci. 87:

[116945] 9625–9629, 1990.

[116946] 20647.

[116947] Widhalm, K.; Koch, S.; Scheibenreiter, S.; Knoll, E.;
Colombo,

[116948] J. P.; Bachmann, C.; Thalhammer, O.: Long-term follow-up
of 12 patients

[116949] with the late-onset variant of argininosuccinic acid lyase
deficiency:

[116950] no impairment of intellectual and psychomotor develop-
ment during therapy. Pediatrics 89:

[116951] 1182–1184, 1992.

[116952] 20648.

[116953] Aloysia, M.; Gelb, A. G.; Fudenberg, H.; Hamper, J.; Tip-
pett, P.;

[116954] Race, R. R.: The expected 'Bombay' group O(H-A1) and
O(H-A2). Transfusion 1:

[116955] 212–217, 1961.

[116956]20649.

[116957]Bhende, Y. M.; Deshpande, C. K.; Bhatia, H. M.; Sanger, R.;
Race,

[116958]R. R.; Morgan, W. T. J.; Watkins, W. M.: A 'new' blood
group character

[116959]related to the ABO system. Lancet I: 903–904, 1952.

[116960]20650.

[116961]Gerard, G.; Vitrac, D.; Le Pendu, J.; Muller, A.; Oriol, R.: H-
deficient

[116962]blood groups (Bombay) of Reunion Island. Am. J. Hum.
Genet. 34:

[116963]937–947, 1982.

[116964]20651.

[116965]Hrubisko, M.; Laluha, J.; Mergancova, O.; Zakovicova, S.:
New

[116966]variants in the ABOH blood group system due to interac-
tion of recessive

[116967]genes controlling the formation of H antigen in erythro-
cytes: the

[116968]'Bombay-like' phenotypes O HM, OB HM, OAB HM. Vox
Sang. 19: 113–122,

[116969]1970.

[116970]20652.

[116971]Kelly, R. J.; Ernst, L. K.; Larsen, R. D.; Bryant, J. G.; Robinson,

[116972]J. S.; Lowe, J. B.: Molecular basis for H blood group deficiency

[116973]in Bombay (Oh) and para-Bombay individuals. Proc. Nat. Acad. Sci. 91:

[116974]5843–5847, 1994.

[116975]20653.

[116976]Koda, Y.; Soejima, M.; Johnson, P. H.; Smart, E.; Kimura, H.:

[116977]Missense mutation of FUT1 and deletion of FUT2 are responsible for

[116978]Indian Bombay phenotype at ABO blood group system. Biochem. Biophys.

[116979]Res. Commun. 238: 21–25, 1997.

[116980]20654.

[116981]Larsen, R. D.; Ernst, L. K.; Nair, R. P.; Lowe, J. B.: Molecular

[116982]cloning, sequence, and expression of a human GDP–L–fucose:beta–D–galactoside

[116983]2- α -L-fucosyltransferase cDNA that can form the H blood group

[116984]antigen. Proc. Nat. Acad. Sci. 87: 6674-6678, 1990.

[116985]20655.

[116986]Le Pendu, J.; Cartron, J. P.; Lemieux, R. U.; Oriol, R.: The presence

[116987]of at least two different H-blood-group-related beta-D-Gal α -2-L-fucosyltransferases

[116988]in human serum and the genetics of blood group H substances. Am.

[116989]J. Hum. Genet. 37: 749-760, 1985.

[116990]20656.

[116991]Le Pendu, J.; Clamagirand-Mulet, C.; Cartron, J.-P.; Gerard, G.;

[116992]Vitrac, D.; Oriol, R.: H-deficient blood groups of Reunion Island.

[116993]III. α -2-L-fucosyltransferase activity in sera of homozygous and

[116994]heterozygous individuals. Am. J. Hum. Genet. 35: 497-507, 1983.

[116995]20657.

[116996]Le Pendu, J.; Gerard, G.; Vitrac, D.; Juszczak, G.; Liberge, G.;

[116997]Rouger, P.; Salmon, C.; Lambert, F.; Dalix, A.-M.; Oriol, R.:
H-deficient

[116998]blood groups of Reunion Island. II. Differences between
Indians (Bombay

[116999]phenotype) and whites (Reunion phenotype). Am. J. Hum.
Genet. 35:

[117000]484-496, 1983.

[117001]20658.

[117002]Levine, P.; Robinson, E.; Celano, M.; Briggs, O.; Falkinburg,

[117003]L.: Gene interaction resulting in suppression of blood
group substance

[117004]B. Blood 10: 1100-1108, 1955.

[117005]20659.

[117006]Oriol, R.: Personal Communication. Paris, France
2/7/1983.

[117007]20660.

[117008]Sathe, M.; Bhatia, H. M.: Bombay (OH) phenotype in two
generations

[117009]in an Indian family. Vox Sang. 30: 312-314, 1976.

[117010]20661.

[117011]Yunis, E. J.; Svoldal, J. M.; Bridges, R. A.: Genetics of the

[117012]Bombay phenotype. Blood 33: 124–132, 1969.

[117013]20662.

[117014]Lees–Miller, J. P.; Helfman, D. M.: The molecular basis for
tropomyosin

[117015]isoform diversity. BioEssays 13: 429–437, 1991.

[117016]20663.

[117017]MacLeod, A. R.; Gooding, C.: Human hTM–alpha gene: ex-
pression

[117018]in muscle and nonmuscle tissue. Molec. Cell. Biol. 8:
433–440, 1988.

[117019]20664.

[117020]Muthuchamy, M.; Pieples, K.; Rethinasamy, P.; Hoit, B.;
Grupp,

[117021]J. L.; Boivin, G. P.; Wolska, B.; Evans, C.; Solaro, R. J.; Wiece-
zorek,

[117022]D. F.: Mouse model of a familial hypertrophic cardiomy-
opathy mutation

[117023]in alpha–tropomyosin manifests cardiac dysfunction. Circ.
Res. 85:

[117024]47–56, 1999.

[117025]20665.

[117026]Schleef, M.; Werner, K.; Satzger, U.; Kaupmann, K.;
Jockusch, H.

[117027]: Chromosomal localization and genomic cloning of the
mouse alpha-tropomyosin

[117028]gene Tpm-1. Genomics 17: 519–521, 1993.

[117029]20666.

[117030]Thierfelder, L.; Watkins, H.; MacRae, C.; Lamas, R.;
McKenna, W.;

[117031]Vosberg, H.-P.; Seidman, J. G.; Seidman, C. E.: Alpha-
tropomyosin

[117032]and cardiac troponin T mutations cause familial hyper-
trophic cardiomyopathy:

[117033]a disease of the sarcomere. Cell 77: 701–712, 1994.

[117034]20667.

[117035]Watkins, H.; Anan, R.; Coviello, D. A.; Spirito, P.; Seidman,

[117036]J. G.; Seidman, C. E.: A de novo mutation in alpha-
tropomyosin that

[117037]causes hypertrophic cardiomyopathy. Circulation 91:
2302–2305, 1995.

[117038]20668.

[117039]Watkins, H.; McKenna, W. J.; Thierfelder, L.; Suk, H. J.;
Anan,

[117040]R.; O'Donoghue, A.; Spirito, P.; Matsumori, A.; Moravec, C.
S.; Seidman,

[117041]J. G.; Seidman, C. E.: Mutations in the genes for cardiac
troponin

[117042]T and alpha-tropomyosin in hypertrophic cardiomyopa-
thy. New Eng.

[117043]J. Med. 332: 1058-1064, 1995.

[117044]20669.

[117045]Gahlmann, R.; Kedes, L.: Cloning, structural analysis, and
expression

[117046]of the human fast twitch skeletal muscle troponin C gene.
J. Biol.

[117047]Chem. 265: 12520-12528, 1990.

[117048]20670.

[117049]Gahlmann, R.; Wade, R.; Gunning, P.; Kedes, L.: Differen-
tial expression

[117050]of slow and fast skeletal muscle troponin C: slow skeletal
muscle

[117051]troponin C is expressed in human fibroblasts. J. Molec.
Biol. 201:

[117052]379–391, 1988.

[117053]20671.

[117054]Townsend, P. J.; Yacoub, M. H.; Barton, P. J. R.: Assign-
ment of

[117055]the human fast skeletal muscle troponin C gene (TNNC2)
between D20S721

[117056]and GCT10F11 on chromosome 20 by somatic cell hybrid
analysis. Ann.

[117057]Hum. Genet. 61: 457–459, 1997.

[117058]20672.

[117059]Bermingham, N.; Hernandez, D.; Balfour, A.; Gilmour, F.;
Martin,

[117060]J. E.; Fisher, E. M. C.: Mapping TNNC1, the gene that en-
codes cardiac

[117061]troponin I in the human and mouse. Genomics 30:
620–622, 1995.

[117062]20673.

[117063]Kwon, B. S.; Halaban, R.; Chintamaneni, C.: Molecular ba-
sis of

[117064]mouse Himalayan mutation. Biochem. Biophys. Res. Commun. 161: 252-260,

[117065]1989.

[117066]20674.

[117067]DiLella, A. G.: Chromosomal assignment of the human immunophilin

[117068]FKBP-12 gene. Biochem. Biophys. Res. Commun. 179: 1427-1433, 1991.

[117069]20675.

[117070]DiLella, A. G.; Hawkins, A.; Craig, R. J.; Schreiber, S. L.; Griffin,

[117071]C. A.: Chromosomal band assignments of the genes encoding human FKBP12

[117072]and FKBP13. Biochem. Biophys. Res. Commun. 189: 819-823, 1992.

[117073]20676.

[117074]Goebl, M. G.: The peptidyl-prolyl isomerase, FK506-binding protein,

[117075]is most likely the 12 kd endogenous inhibitor 2 of protein kinase

[117076]C. (Letter) Cell 64: 1051-1052, 1991.

[117077]20677.

[117078]Jin, Y.-J.; Albers, M. W.; Lane, W. S.; Bierer, B. E.; Schreiber,

[117079]S. L.; Burakoff, S. J.: Molecular cloning of a membrane-associated

[117080]human FK506- and rapamycin-binding protein, FKBP-13.
Proc. Nat. Acad.

[117081]Sci. 88: 6677-6681, 1991.

[117082]20678.

[117083]Maki, N.; Sekiguchi, F.; Nishimaki, J.; Miwa, K.; Hayano, T.;
Takahashi,

[117084]N.; Suzuki, M.: Complementary DNA encoding the human
T-cell FK506-binding

[117085]protein, a peptidylprolyl cis-trans isomerase distinct from
cyclophilin. Proc.

[117086]Nat. Acad. Sci. 87: 5440-5443, 1990.

[117087]20679.

[117088]Peattie, D. A.; Hsaio, K.; Benasutti, M.; Lippke, J. A.: Three

[117089]distinct messenger RNAs can encode the human immunosuppressant-binding

[117090]protein FKBP12. Gene 150: 251-257, 1994.

[117091]20680.

[117092]Shou, W.; Aghdasi, B.; Armstrong, D. L.; Guo, Q.; Bao, S.;
Charng,

[117093]M.-J.; Mathews, L. M.; Schneider, M. D.; Hamilton, S. L.;
Matzuk,

[117094]M. M.: Cardiac defects and altered ryanodine receptor
function in

[117095]mice lacking FKBP12. Nature 391: 489–492, 1998.

[117096]20681.

[117097]Standaert, R. F.; Galat, A.; Verdine, G. L.; Schreiber, S. L.:

[117098]Molecular cloning and overexpression of the human
FK506-binding protein

[117099]FKBP. Nature 346: 671–674, 1990.

[117100]20682.

[117101]Wang, T.; Donahoe, P. K.; Zervos, A. S.: Specific interaction

[117102]of type I receptors of the TGF-beta family with the im-
munophilin FKBP-12. Science 265:

[117103]674–676, 1994.

[117104]20683.

[117105]Bichi, R.; Shinton, S. A.; Martin, E. S.; Koval, A.; Calin, G.

[117106]A.; Cesari, R.; Russo, G.; Hardy, R. R.; Croce, C. M.: Human chronic
[117107]lymphocytic leukemia modeled in mouse by targeted TCL1
expression. Proc.
[117108]Nat. Acad. Sci. 99: 6955–6960, 2002.
[117109]20684.
[117110]Davey, M. P.; Bertness, V.; Nakahara, K.; Johnson, J. P.;
McBride,
[117111]O. W.; Waldmann, T. A.; Kirsch, I. R.: Juxtaposition of the
T-cell
[117112]receptor alpha-chain locus (14q11) and a region (14q32)
of potential
[117113]importance in leukemogenesis by a 14;14 translocation in
a patient
[117114]with T-cell chronic lymphocytic leukemia and ataxia-
telangiectasia. Proc.
[117115]Nat. Acad. Sci. 85: 9287–9291, 1988.
[117116]20685.
[117117]Hecht, F.; Morgan, R.; Hecht, B. K.–M.; Smith, S. D.: Com-
mon region
[117118]on chromosome 14 in T-cell leukemia and lymphoma.
Science 226: 1445–1447,

[117119]1984.

[117120]20686.

[117121]Laine, J.; Kunstle, G.; Obata, T.; Sha, M.; Noguchi, M.: The protooncogene

[117122]TCL1 is an Akt kinase coactivator. *Molec. Cell* 6: 395–407, 2000.

[117123]20687.

[117124]Mathieu–Mahul, D.; Caubet, J. F.; Bernheim, A.; Mauchauffe, M.;

[117125]Palmer, E.; Berger, R.; Larsen, C.–J.: Molecular cloning of a DNA

[117126]fragment from human chromosome 14(14q11) involved in T–cell malignancies. *EMBO*

[117127]*J.* 4: 3427–3433, 1985.

[117128]20688.

[117129]Mengle–Gaw, L.; Albertson, D. G.; Sherrington, P. D.; Rabbits,

[117130]T. H.: Analysis of a T–cell tumor–specific breakpoint cluster at

[117131]human chromosome 14q32. *Proc. Nat. Acad. Sci.* 85: 9171–9175, 1988.

[117132]20689.

[117133]Howard, H. C.; Mount, D. B.; Rochefort, D.; Byun, N.;
Dupre, N.;

[117134]Lu, J.; Fan, X.; Song, L.; Riviere, J.-B.; Prevost, C.; Horst, J.;

[117135]Simonati, A.; and 12 others: The K-Cl cotransporter KCC3
is mutant

[117136]in a severe peripheral neuropathy associated with agene-
sis of the

[117137]corpus callosum. Nature Genet. 7Oct, 2002. Note: Ad-
vance Electronic

[117138]Publication.

[117139]20690.

[117140]Haire, R. N.; Litman, G. W.: The murine form of TXK, a
novel TEC

[117141]kinase expressed in thymus maps to chromosome 5.
Mammalian Genome 6:

[117142]476-480, 1995.

[117143]20691.

[117144]Haire, R. N.; Ohta, Y.; Lewis, J. E.; Fu, S. M.; Kroisel, P.; Lit-
man,

[117145]G. W.: TXK, a novel human tyrosine kinase expressed in T

cells shares

[117146]sequence identity with Tec family kinases and maps to
4p12. Hum.

[117147]Molec. Genet. 3: 897–901, 1994.

[117148]20692.

[117149]Schneider, H.; Schwartzberg, P. L.; Rudd, C. E.: Resting
lymphocyte

[117150]kinase (Rlk/Txk) phosphorylates the YVKM motif and reg-
ulates PI 3-kinase

[117151]binding to T-cell antigen CTLA-4. Biochem. Biophys. Res.
Commun. 252:

[117152]14–19, 1998.

[117153]20693.

[117154]Takeba, Y.; Nagafuchi, H.; Takeno, M.; Kashiwakura, J.;
Suzuki,

[117155]N.: Txk, a member of nonreceptor tyrosine kinase of Tec
family, acts

[117156]as a Th1 cell-specific transcription factor and regulates
IFN-gamma

[117157]gene transcription. J. Immun. 168: 2365–2370, 2002.

[117158]20694.

[117159]Gasdaska, J. R.; Gasdaska, P. Y.; Gallegos, A.; Powis, G.:
Human

[117160]thioredoxin reductase gene localization to chromosomal
position 12q23-q24.1

[117161]and mRNA distribution in human tissue. Genomics 37:
257-259, 1996.

[117162]20695.

[117163]Gasdaska, P. Y.; Gasdaska, J. R.; Cochran, S.; Powis, G.:
Cloning

[117164]and sequencing of a human thioredoxin reductase. FEBS
Lett. 373:

[117165]5-9, 1995.

[117166]20696.

[117167]Russo, M. W.; Sevetson, B. R.; Milbrandt, J.: Identification
of

[117168]NAB1, a repressor of NGFI-A- and Krox20-mediated tran-
scription. Proc.

[117169]Nat. Acad. Sci. 92: 6873-6877, 1995.

[117170]20697.

[117171]Svaren, J.; Apel, E. D.; Simburger, K. S.; Jenkins, N. A.;
Gilbert,

[117172]D. J.; Copeland, N. A.; Milbrandt, J.: The Nab2 and Stat6 genes share

[117173]a common transcription termination region. Genomics 41: 33–39, 1997.

[117174]20698.

[117175]Swirnoff, A. H.; Apel, E. D.; Svaren, J.; Sevetson, B. R.; Zimonjic,

[117176]D. B.; Popescu, N. C.; Milbrandt, J.: Nab1, a corepressor of NGFI-A

[117177](Egr-1), contains an active transcriptional repression domain. Molec.

[117178]Cell. Biol. 18: 512–524, 1998.

[117179]20699.

[117180]Venken, K.; Di Maria, E.; Bellone, E.; Balestra, P.; Cassandrini,

[117181]D.; Mandich, P.; De Jonghe, P.; Timmerman, V.; Svaren, J.: Search

[117182]for mutation in the EGR2 corepressor proteins, NAB1 and NAB2, in human

[117183]peripheral neuropathies. Neurogenetics 4: 37–41, 2002.

[117184]20700.

- [117185]Nichols, R. C.; Blinder, J.; Pai, S. I.; Ge, Q.; Targoff, I. N.;
- [117186]Plotz, P. H.; Liu, P.: Assignment of two human autoantigen genes:
- [117187]isoleucyl-tRNA synthetase locates to 9q21 and lysyl-tRNA synthetase
- [117188]locates to 16q23-q24. Genomics 36: 210-213, 1996.
- [117189]20701.
- [117190]Nichols, R. C.; Raben, N.; Boerkoel, C. F.; Plotz, P. H.: Human
- [117191]isoleucyl-tRNA synthetase: sequence of the cDNA, alternative mRNA
- [117192]splicing, and the characteristics of an unusually long C-terminal
- [117193]extension. Gene 155: 299-304, 1995.
- [117194]20702.
- [117195]Tommerup, N.; Leffers, H.: Assignment of human KH-box-containing
- [117196]genes by in situ hybridization: HNRNPK maps to 9q21.32-q21.33, PCBP1
- [117197]to 2p12-p13, and PCBP2 to 12q13.12-q13.13, distal to FRA12A. Genomics 32:

[117198]297–298, 1996.

[117199]20703.

[117200]Martell, K. J.; Kwak, S.; Hakes, D. J.; Dixon, J. E.; Trent, J.

- [117201]M.: Chromosomal localization of four human VH1-like protein-tyrosine phosphatases. Genomics 22: 462-464, 1994.
- [117202]20704.
- [117203]Chrast, R.; Chen, H.; Morris, M. A.; Antonarakis, S. E.: Mapping of the human transcription factor GABPA (E4TF1-60) gene to chromosome 21. Genomics 28: 119-122, 1995.
- [117204]20705.
- [117205]Gugneja, S.; Virbasius, J. V.; Scarpulla, R. C.: Four structurally distinct, non-DNA-binding subunits of human nuclear respiratory factor 2 share a conserved transcriptional activation domain. Molec. Cell. Biol. 15: 102-111, 1995.
- [117206]20706.
- [117207]Guo, A.; Nie, F.; Wong-Riley, M.: Human nuclear respiratory factor 2-alpha subunit cDNA: isolation, subcloning, sequencing,

and in situ

[117215]hybridization of transcripts in normal and monocularly
deprived macaque

[117216]visual system. J. Comp. Neurol. 417: 221–232, 2000.

[117217]20707.

[117218]Sawada, J.; Goto, M.; Watanabe, H.; Handa, H.; Yoshida, M.
C.:

[117219]Regional mapping of two subunits of transcription factor
E4TF1 to

[117220]human chromosome. Jpn. J. Cancer Res. 86: 10–12, 1995.

[117221]20708.

[117222]Virbasius, J. V.; Scarpulla, R. C.: Activation of the human
mitochondrial

[117223]transcription factor A gene by nuclear respiratory factors:
a potential

[117224]regulatory link between nuclear and mitochondrial gene
expression

[117225]in organelle biogenesis. Proc. Nat. Acad. Sci. 91:
1309–1313, 1994.

[117226]20709.

[117227]Watanabe, H.; Sawada, J.; Yano, K.-I.; Yamaguchi, K.; Goto,

M.;

[117228]Handa, H.: cDNA cloning of transcription factor E4TF1 subunits with

[117229]Ets and notch motifs. *Molec. Cell. Biol.* 13: 1385–1391, 1993.

[117230]20710.

[117231]Chambrud, B.; Radanyi, C.; Camonis, J. H.; Rajkowski, K.; Schumacher,

[117232]M.; Baulieu, E.–E.: Immunophilins, Refsum disease, and lupus nephritis:

[117233]the peroxisomal enzyme phytanoyl–CoA alpha–hydroxylase is a new FKBP–associated

[117234]protein. *Proc. Nat. Acad. Sci.* 96: 2104–2109, 1999.

[117235]20711.

[117236]Dixon, J.; Loftus, S. K.; Gladwin, A. J.; Scambler, P. J.; Wasmuth,

[117237]J. J.; Dixon, M. J.: Cloning of the human heparan sulfate–N–deacetylase/N–sulfotransferase

[117238]gene from the Treacher Collins syndrome candidate region at 5q32–q33.1. *Genomics* 26:

[117239]239–244, 1995.

[117240]20712.

[117241]Gladwin, A. J.; Dixon, J.; Loftus, S. K.; Wasmuth, J. J.;
Dixon,

[117242]M. J.: Genomic organization of the human heparan sul-
fate-N-deacetylase/N-sulfotransferase

[117243]gene: exclusion from a causative role in the pathogenesis
of Treacher

[117244]Collins syndrome. Genomics 32: 471-473, 1996.

[117245]20713.

[117246]Hashimoto, Y.; Orellana, A.; Gil, G.; Hirschberg, C. B.:
Molecular

[117247]cloning and expression of rat liver N-heparan sulfate sul-
fotransferase. J.

[117248]Biol. Chem. 267: 15744-15750, 1995.

[117249]20714.

[117250]Chen, H.; Antonarakis, S. E.: Localisation of a human ho-
mologue

[117251]of the Drosophila *mnb* and rat *Dyrk* genes to chromosome
21q22.2. Hum.

[117252]Genet. 99: 262-265, 1997.

[117253]20715.

[117254]Patil, N.; Cox, D. R.; Bhat, D.; Faham, M.; Myers, R. M.; Pe-

terson,

[117255]A. S.: A potassium channel mutation in weaver mice implicates membrane

[117256]excitability in granule cell differentiation. *Nature Genet.*

11:

[117257]126–129, 1995.

[117258]20716.

[117259]Shindoh, N.; Kudoh, J.; Maeda, H.; Yamaki, A.; Minoshima, S.; Shimizu,

[117260]Y.; Shimizu, N.: Cloning of a human homolog of the *Drosophila* minibrain/rat

[117261]Dyrk gene from 'the Down syndrome critical region' of chromosome 21. *Biochem.*

[117262]*Biophys. Res. Commun.* 225: 92–99, 1996.

[117263]20717.

[117264]Smith, D. J.; Stevens, M. E.; Sudanagunta, S. P.; Bronson, R. T.;

[117265]Makhinson, M.; Watabe, A. M.; O'Dell, T. J.; Fung, J.; Weier, H.–U.

[117266]G.; Cheng, J.–F.; Rubin, E. M.: Functional screening of 2 Mb of human

[117267]chromosome 21q22.2 in transgenic mice implicates mini–

brain in learning

[117268]defects associated with Down syndrome. Nature Genet.
16: 28–36,

[117269]1997.

[117270]20718.

[117271]Song, W.-J.; Chung, S.-H.; Kurnit, D. M.: The murine Dyrk
protein

[117272]maps to chromosome 16, localizes to the nucleus, and
can form multimers. Biochem.

[117273]Biophys. Res. Commun. 231: 640–644, 1997.

[117274]20719.

[117275]Song, W.-J.; Sternberg, L. R.; Kasten-Sportes, C.; Van
Keuren,

[117276]M. L.; Chung, S.-H.; Slack, A. C.; Miller, D. E.; Glover, T.
W.; Chiang,

[117277]P.-W.; Lou, L.; Kurnit, D. M.: Isolation of human and
murine homologues

[117278]of the Drosophila minibrain gene: human homologue
maps to 21q22.2

[117279]in the Down syndrome 'critical region.' Genomics 38:
331–339, 1996.

[117280]20720.

[117281]Tejedor, F.; Zhu, X. R.; Kaltenbach, E.; Ackermann, A.;
Baumann,
[117282]A.; Canal, I.; Heisenberg, M.; Fischbach, K. F.; Pongs, O.:
Minibrain:
[117283]a new protein kinase family involved in postembryonic
neurogenesis
[117284]in *Drosophila*. *Neuron* 14: 287–301, 1995.
[117285]20721.
[117286]Algar, E.; Brickell, S.; Deeble, G.; Amor, D.; Smith, P.:
Analysis
[117287]of CDKN1C in Beckwith Wiedemann syndrome. *Hum. Mu-
tat.* 15: 497–508,
[117288]2000.
[117289]20722.
[117290]Bhuiyan, Z. A.; Yatsuki, H.; Sasaguri, T.; Joh, K.; Soejima,
H.;
[117291]Zhu, X.; Hatada, I. Morisaki, H.; Morisaki, T.; Mukai, T.:
Functional
[117292]analysis of the p57(KIP2) gene mutation in Beckwith-
Wiedemann syndrome. *Hum.*
[117293]*Genet.* 104: 205–210, 1999.

[117294]20723.

[117295]Fitzpatrick, G. V.; Soloway, P. D.; Higgins, M. J.: Regional
loss

[117296]of imprinting and growth deficiency in mice with a tar-
geted deletion

[117297]of KvDMR1. Nature Genet. 9Sept, 2002. Note: Advance
Electronic Publication.

[117298]20724.

[117299]Hatada, I.; Mukai, T.: Genomic imprinting of p57(KIP2), a
cyclin-dependent

[117300]kinase inhibitor, in mouse. Nature Genet. 11: 204-206,
1995.

[117301]20725.

[117302]Hatada, I.; Nabetani, A.; Morisaki, H.; Xin, Z.; Ohishi, S.;
Tonoki,

[117303]H.; Niikawa, N.; Inoue, M.; Komoto, Y.; Okada, A.; Ste-
ichen, E.; Ohashi,

[117304]H.; Fukushima, Y.; Nakayama, M.; Mukai, T.: New
p57(KIP2) mutations

[117305]in Beckwith-Wiedemann syndrome. Hum. Genet. 100:
681-683, 1997.

[117306]20726.

- [117307] Hatada, I.; Ohashi, H.; Fukushima, Y.; Kaneko, Y.; Inoue, M.; Komoto, Y.; Okada, A.; Ohishi, S.; Nabetani, A.; Morisaki, H.; Nakayama, M.; Niikawa, M.; Mukai, T.: An imprinted gene p57(KIP2) is mutated in Beckwith–Wiedemann syndrome. *Nature Genet.* 14: 171–173, 1996.
- [117311] 20727.
- [117312] John, R. M.; Ainscough, J. F.–X.; Barton, S. C.; Surani, M. A. [117313]: Distant cis–elements regulate imprinted expression of the mouse p57(Kip2) (Cdkn1c) gene: implications for the human disorder, Beckwith–Wiedemann [117314] syndrome. *Hum. Molec. Genet.* 10: 1601–1609, 2001.
- [117316] 20728.
- [117317] Lee, M.–H.; Reynisdottir, I.; Massague, J.: Cloning of p57(KIP2), [117318] a cyclin–dependent kinase inhibitor with unique domain structure and [117319] tissue distribution. *Genes Dev.* 9: 639–649, 1995.

[117320]20729.

[117321]Zhang, X.; Zhou, J.; Reeders, S. T.; Tryggvason, K.: Structure

[117322]of the human type IV collagen COL4A6 gene, which is mutated in Alport

[117323]syndrome-associated leiomyomatosis. Genomics 33: 473-479, 1996.

[117324]20730.

[117325]Zhou, J.; Ding, M.; Zhao, Z.; Reeders, S. T.: Complete primary

[117326]structure of the sixth chain of human basement membrane collagen,

[117327]alpha-6(IV): isolation of the cDNAs for alpha-6(IV) and comparison

[117328]with five other type IV collagen chains. J. Biol. Chem. 269: 13193-13199,

[117329]1994.

[117330]20731.

[117331]Nathans, J.; Maumenee, I. H.; Zrenner, E.; Sadowski, B.; Sharpe,

[117332]L. T.; Lewis, R. A.; Hansen, E.; Rosenberg, T.; Schwartz, M.; Heckenlively,

[117333]J. R.; Traboulsi, E.; Klingaman, R.; Bech-Hansen, N. T.;
LaRoche,

[117334]G. R.; Pagon, R. A.; Murphey, W. H.; Weleber, R. G.: Genetic
heterogeneity

[117335]among blue-cone monochromats. Am. J. Hum. Genet. 53:
987-1000, 1993.

[117336]20732.

[117337]Bell, J.; Haldane, J. B. S.: The linkage between the genes
for

[117338]colour-blindness and haemophilia in man. Proc. Roy. Soc.
London 123B:

[117339]119-150, 1937.

[117340]20733.

[117341]Filippi, G.; Rinaldi, A.; Palmarino, R.; Seravalli, E.; Sinis-
calco,

[117342]M.: Linkage disequilibrium for two X-linked genes in Sar-
dinia and

[117343]its bearing on the statistical mapping of the human X
chromosome. Genetics 86:

[117344]199-222, 1977.

[117345]20734.

- [117346]Thompson, D. A.; McHenry, C. L.; Li, Y.; Richards, J. E.;
Othman,
[117347]M. I.; Schwinger, E.; Vollrath, D.; Jacobson, S. G.; Gal, A.:
Retinal
[117348]dystrophy due to paternal isodisomy for chromosome 1 or
chromosome
[117349]2, with homoallelism for mutations in RPE65 or MERTK,
respectively. Am.
[117350]J. Hum. Genet. 70: 224–229, 2002.
[117351]20735.
[117352]Aguirre, G. D.; Baldwin, V.; Weeks, K. M.; Acland, G. M.;
Ray,
[117353]K.: Frequency of the codon 807 mutation in the cGMP
phosphodiesterase
[117354]beta-subunit gene in Irish setters and other dog breeds
with hereditary
[117355]retinal degeneration. J. Hered. 90: 143–147, 1999.
[117356]20736.
[117357]Bateman, J. B.; Klisak, I.; Kojis, T.; Mohandas, T.; Sparkes,
R.
[117358]S.; Li, T.; Applebury, M. L.; Bowes, C.; Farber, D. B.: As-
signment

[117359] of the beta-subunit of rod photoreceptor cGMP phosphodiesterase gene

[117360] PDEB (homolog of the mouse rd gene) to human chromosome 4p16. Genomics 12:

[117361] 601-603, 1992.

[117362] 20737.

[117363] Bayes, M.; Giordano, M.; Balcells, S.; Grinberg, D.; Vigeliliu,

[117364] L.; Martinez, I.; Ayuso, C.; Benitez, J.; Ramos-Arroyo, M. A.; Chivelet,

[117365] P.; Solans, T.; Valverde, D.; Amselem, S.; Goossens, M.; Baiget, M.;

[117366] Gonzalez-Duarte, R.; Besmond, C.: Homozygous tandem duplication within

[117367] the gene encoding the beta-subunit of rod phosphodiesterase as a cause

[117368] for autosomal recessive retinitis pigmentosa. Hum. Mutat. 5: 228-234,

[117369] 1995.

[117370] 20738.

[117371] Bennett, J.; Tanabe, T.; Sun, D.; Zeng, Y.; Kjeldbye, H.; Gouras,

[117372]P.; Maguire, A. M.: Photoreceptor cell rescue in retinal degeneration

[117373](rd) mice by in vivo gene therapy. Nature Med. 2: 649, 1996.

[117374]20739.

[117375]Bowes, C.; Danciger, M.; Kozak, C. A.; Farber, D. B.: Isolation

[117376]of a candidate cDNA for the gene causing retinal degeneration in the

[117377]rd mouse. Proc. Nat. Acad. Sci. 86: 9722-9726, 1989.

Note: Erratum:

[117378]Proc. Nat. Acad. Sci 87:1625, 1990...

[117379]20740.

[117380]Bowes, C.; Li, T.; Danciger, M.; Baxter, L. C.; Applebury, M. L.;

[117381]Farber, D. B.: Retinal degeneration in the rd mouse is caused by

[117382]a defect in the beta subunit of rod cGMP-phosphodiesterase. Nature 347:

[117383]677-680, 1990.

[117384]20741.

- [117385]Bowes, C.; Li, T.; Frankel, W. N.; Danciger, M.; Coffin, J. M.;
- [117386]Applebury, M. L.; Farber, D. B.: Localization of a retroviral element within the rd gene coding for the beta subunit of cGMP phosphodiesterase. Proc.
- [117387]Nat. Acad. Sci. 90: 2955–2959, 1993.
- [117388]20742.
- [117389]Cachon–Gonzalez, M. B.; Fenner, S.; Coffin, J. M.; Moran, C.; Best, S.; Stoye, J. P.: Structure and expression of the hairless gene of mice. Proc. Nat. Acad. Sci. 91: 7717–7721, 1994.
- [117390]20743.
- [117391]Clements, P. J. M.; Gregory, C. Y.; Peterson–Jones, S. M.; Sargan, D. R.; Bhattacharya, S. S.: Confirmation of the rod cGMP phosphodiesterase beta subunit (PDE–beta) nonsense mutation in affected rcd–1 Irish setters in the UK and the development of a diagnostic test. Curr.

[117398]Eye Res. 12: 861–866, 1993.

[117399]20744.

[117400]Collins, C.; Hutchinson, G.; Kowbel, D.; Riess, O.; Weber, B.;

[117401]Hayden, M. R.: The human beta-subunit of rod photoreceptor cGMP phosphodiesterase:

[117402]complete retinal cDNA sequence and evidence for expression in brain. Genomics 13:

[117403]698–704, 1992.

[117404]20745.

[117405]Danciger, M.; Bowes, C.; Kozak, C. A.; LaVail, M. M.; Farber,

[117406]D. B.: Fine mapping of a putative rd cDNA and its cosegregation with

[117407]rd expression. Invest. Ophthalmol. Vis. Sci. 31: 1427–1432, 1990.

[117408]20746.

[117409]Farber, D. B.; Danciger, J. S.; Aguirre, G.: The beta subunit
[117410]of cyclic GMP phosphodiesterase mRNA is deficient in canine rod-cone

[117411]dysplasia 1. Neuron 9: 349–356, 1992.

[117412]20747.

[117413]Farber, D. B.; Lolley, R. N.: Enzyme basis for cyclic GMP
accumulation

[117414]in degenerative photoreceptor cells of mouse retina. J.
Cyclic Nucleotide

[117415]Res. 2: 139–148, 1976.

[117416]20748.

[117417]Farber, D. B.; Lolley, R. N.: Cyclic guanosine monophos-
phate:

[117418]elevation in degenerating photoreceptor cells of the C3H
mouse retina. Science 186:

[117419]449–451, 1974.

[117420]20749.

[117421]Gal, A.; Orth, U.; Baehr, W.; Schwinger, E.; Rosenberg, T.:
Heterozygous

[117422]missense mutation in the rod cGMP phosphodiesterase
beta-subunit gene

[117423]in autosomal dominant stationary night blindness. Nature
Genet. 7:

[117424]64–68, 1994. Note: Correction: Nature Genet. 7: 551 only,
1994...

[117425]20750.

[117426]Bogenmann, E.; Lochrie, M. A.; Simon, M. I.: Cone cell-specific

[117427]genes expressed in retinoblastoma. Science 240: 76-78, 1988.

[117428]20751.

[117429]Bonaiti-Pellie, C.; Briard-Guillemot, M. L.: Segregation analysis

[117430]in hereditary retinoblastoma. Hum. Genet. 57: 411-419, 1981.

[117431]20752.

[117432]Nagasaki, H.; Ito, M.; Yuasa, H.; Saito, H.; Fukase, M.; Hamada,

[117433]K.; Ishikawa, E.; Katakami, H.; Oiso, Y.: Two novel mutations in

[117434]the coding region for neurophysin-II associated with familial central

[117435]diabetes insipidus. J. Clin. Endocr. Metab. 80: 1352-1356, 1995.

[117436]20753.

[117437]Bartram, C. R.; de Klein, A.; Hagemeijer, A.; van Agthoven, T.;

[117438]Geurts van Kessel, A.; Bootsma, D.; Grosveld, G.; Ferguson-Smith,
son-Smith,
[117439]M. A.; Davies, T.; Stone, M.; Heisterkamp, N.; Stephenson,
J. R.;
[117440]Groffen, J.: Translocation of c-abl oncogene correlates
with the
[117441]presence of a Philadelphia chromosome in chronic myelo-
cytic leukaemia. Nature 306:
[117442]277-280, 1983.
[117443]20754.
[117444]Rittig, S.; Robertson, G. L.; Siggaard, C.; Kovacs, L.;
Gregersen,
[117445]N.; Nyborg, J.; Pedersen, E. B.: Identification of 13 new
mutations
[117446]in the vasopressin-neurophysin II gene in 17 kindreds
with familial
[117447]autosomal dominant neurohypophyseal diabetes in-
sipidus. Am. J. Hum.
[117448]Genet. 58: 107-117, 1996.
[117449]20755.
[117450]Russell, J. T.; Brownstein, M. J.; Gainer, H.: Trypsin liber-
ates

[117451]an arginine vasopressin-like peptide and neurophysin
from a M(r)20,000

[117452]putative common precursor. Proc. Nat. Acad. Sci. 76:
6086-6090,

[117453]1979.

[117454]20756.

[117455]Sachs, H.; Fawcett, P.; Takabatake, Y.; Portanova, R.:
Biosynthesis

[117456]and release of vasopressin and neurophysin. Recent Prog.
Horm. Res. 25:

[117457]447-491, 1969.

[117458]20757.

[117459]O'Brien, R. O.; Taske, N. L.; Hansbro, P. M.; Matthaei, K. I.;

[117460]Hogan, S. P.; Denborough, M. A.; Foster, P. S.: Exclusion of
defects

[117461]in the skeletal muscle specific regions of the DHPR alpha-
1 subunit

[117462]as frequent causes of malignant hyperthermia. J. Med.
Genet. 32:

[117463]913-914, 1995.

[117464]20758.

[117465]Wichterle, H.; Hanspal, M.; Palek, J.; Jarolim, P.: Combination
[117466]of two mutant alpha spectrin alleles underlies a severe
[117467]spherocytic
[117467]hemolytic anemia. J. Clin. Invest. 98: 2300–2307, 1996.
[117468]20759.
[117469]Nijenhuis, M.; van den Akker, E. L. T.; Zalm, R.; Franken,
A.
[117470]A. M.; Abbes, A. P.; Engel, H.; de Wied, D.; Burbach, J. P.
H.: Familial
[117471]neurohypophysial diabetes insipidus in a large Dutch kin-
dred: effect
[117472]of the onset of diabetes on growth in children and cell bi-
ological
[117473]defects of the mutant vasopressin prohormone. J. Clin.
Endocr. Metab. 86:
[117474]3410–3420, 2001.
[117475]20760.
[117476]Nijenhuis, M.; Zalm, R.; Burbach, J. P. H.: Mutations in the
[117477]vasopressin prohormone involved in diabetes insipidus
impair endoplasmic

[117478]reticulum export but not sorting. J. Biol. Chem. 274:
21200–21208,

[117479]1999.

[117480]20761.

[117481]Olias, G.; Richter, D.; Schmale, H.: Heterologous expres-
sion

[117482]of human vasopressin–neurophysin precursors in a pitu-
itary cell line:

[117483]defective transport of a mutant protein from patients with
familial

[117484]diabetes insipidus. DNA Cell Biol. 15: 929–935, 1996.

[117485]20762.

[117486]Repaske, D. R.; Summar, M. L.; Krishnamani, M. R. S.; Gul-
tekin,

[117487]E. K.; Arriazu, M. C.; Roubicek, M. E.; Blanco, M.; Isaac, G.
B.;

[117488]Phillips, J. A., III: Recurrent mutations in the vasopressin–
neurophysin

[117489]II gene cause autosomal dominant neurohypophyseal dia-
betes insipidus. J.

[117490]Clin. Endocr. Metab. 81: 2328–2334, 1996.

[117491]20763.

[117492]Riddell, D. C.; Mallonee, R.; Phillips, J. A.; Parks, J. S.; Sex-
ton,

[117493]L. A.; Hamerton, J. L.: Chromosomal assignment of human
sequences

[117494]encoding arginine vasopressin-neurophysin II and growth
hormone releasing

[117495]factor. Somat. Cell Molec. Genet. 11: 189-195, 1985.

[117496]20764.

[117497]Repaske, D. R.; Medlej, R.; Gultekin, E. K.; Krishnamani, M.
R.

[117498]S.; Halaby, G.; Findling, J. W.; Phillips, J. A., III: Hetero-
geneity

[117499]in clinical manifestation of autosomal dominant neurohy-
popyseal diabetes

[117500]insipidus caused by a mutation encoding Ala-1 Val in the
signal peptide

[117501]of the arginine vasopressin/neurophysin II/copeptin pre-
cursor. J.

[117502]Clin. Endocr. Metab. 82: 51-56, 1997.

[117503]20765.

[117504]Rutishauser, J.; Boni-Schnetzler, M.; Boni, J.; Wichmann, W.;

[117505]Huisman, T.; Vallotton, M. B.; Froesch, E. R.: A novel point mutation

[117506]in the translation initiation codon of the pre-pro-vasopressin-neurophysin

[117507]II gene: Cosegregation with morphological abnormalities and clinical

[117508]symptoms in autosomal dominant neurohypophyseal diabetes. J. Clin.

[117509]Endocr. Metab. 81: 192-198, 1996.

[117510]20766.

[117511]Sausville, E.; Carney, D.; Battey, J.: The human vasopressin

[117512]gene is linked to the oxytocin gene and is selectively expressed in

[117513]a cultured lung cancer cell line. J. Biol. Chem. 260: 10236-10241,

[117514]1985.

[117515]20767.

[117516]Scherbaum, W. A.; Bottazzo, G. F.: Autoantibodies to vasopressin

[117517]cells in idiopathic diabetes insipidus: evidence for an autoimmune

[117518]variant. *Lancet* I: 897–901, 1983.

[117519]20768.

[117520]Siggaard, C.; Rittig, S.; Corydon, T. J.; Andreasen, P. H.; Jensen,

[117521]T. G.; Andresen, B. S.; Robertson, G. L.; Gregersen, N.; Bolund, L.;

[117522]Pedersen, E. B.: Clinical and molecular evidence of abnormal processing

[117523]and trafficking of the vasopressin preprohormone in a large kindred

[117524]with familial neurohypophyseal diabetes insipidus due to a signal

[117525]peptide mutation. *J. Clin. Endocr. Metab.* 84: 2933–2941, 1999.

[117526]20769.

[117527]Summar, M. L.; Phillips, J. A., III; Battey, J.; Castiglione,

[117528]C. M.; Kidd, K. K.; Maness, K. J.; Weiffenbach, B.; Gravius, T. C.

[117529]: Linkage relationships of human arginine vasopressin-neurophysin-II

[117530]and oxytocin-neurophysin-I to prodynorphin and other
loci on chromosome

[117531]20. Molec. Endocr. 4: 947-950, 1990.

[117532]20770.

[117533]Sunde, D. A.; Sokol, H. W.: Quantification of rat neuro-
physins

[117534]by polyacrylamide gel electrophoresis (page): application
to the rat

[117535]with hereditary hypothalamic diabetes insipidus. Ann. N.Y.
Acad.

[117536]Sci. 248: 345-364, 1975.

[117537]20771.

[117538]Ueta, Y.; Taniguchi, S.; Yoshida, A.; Murakami, I.; Mitani,
Y.;

[117539]Hisatome, I.; Manabe, I.; Sato, R.; Tsuboi, M.; Ohtahara, A.;
Nanba,

[117540]E.; Shigemasa, C.: A new type of familial central diabetes
insipidus

[117541]caused by a single base substitution in the neurophysin II
coding

[117542]region of the vasopressin gene. J. Clin. Endocr. Metab. 81:
1787-1790,

[117543]1996.

[117544]20772.

[117545]Valtin, H.; Sawyer, W. H.; Sokol, H. W.: Neurohypophyseal principles

[117546]in rats homozygous and heterozygous for hypothalamic diabetes insipidus

[117547](Brattleboro strain). Endocrinology 77: 701-706, 1965.

[117548]20773.

[117549]van Leeuwen, F. W.; de Kleijn, D. P. V.; van den Hurk, H. H.;

[117550]Neubauer, A.; Sonnemans, M. A. F.; Sluijs, J. A.; Koycu, S.; Ramdjielal,

[117551]R. D. J.; Salehi, A.; Martens, G. J. M.; Grosveld, F. G.; Burbach,

[117552]J. P. H.; Hol, E. M.: Frameshift mutants of beta-amyloid precursor

[117553]protein and ubiquitin-B in Alzheimer's and Down patients. Science 279:

[117554]242-247, 1998.

[117555]20774.

[117556]Yuasa, H.; Ito, M.; Nagasaki, H.; Oiso, Y.; Miyamoto, S.;

Sasaki,

[117557]N.; Saito, H.: Glu-47, which forms a salt bridge between neurophysin-II

[117558]and arginine vasopressin, is deleted in patients with familial central

[117559]diabetes insipidus. J. Clin. Endocr. Metab. 77: 600-604, 1993.

[117560]20775.

[117561]Oesterling, J. E.; Chan, D. W.; Epstein, J. I.; Kimball, A. W.,

[117562]Jr.; Bruzek, D. J.; Rock, R. C.; Brendler, C. B.; Walsh, P. C.: Prostate

[117563]specific antigen in the preoperative and postoperative evaluation

[117564]of localized prostatic cancer treated with radical prostatectomy. J.

[117565]Urol. 139: 766-772, 1988.

[117566]20776.

[117567]Riegman, P. H. J.; Vlietstra, R. J.; van der Korput, J. A. G. M.;

[117568]Brinkmann, A. O.; Trapman, J.: The promoter of the prostate-specific

[117569]antigen gene contains a functional androgen responsive

element. Molec.

[117570]Endocr. 5: 1921–1930, 1991.

[117571]20777.

[117572]van den Elsen, P.; Bruns, G.; Gerhard, D. S.; Pravtcheva, D.;

[117573]Jones, C.; Housman, D.; Ruddle, F. A.; Orkin, S.; Terhorst,

C.: Assignment

[117574]of the gene coding for the T3–delta subunit of the

T3–T-cell receptor

[117575]complex to the long arm of human chromosome 11 and

to mouse chromosome

[117576]9. Proc. Nat. Acad. Sci. 82: 2920–2924, 1985.

[117577]20778.

[117578]van Rijs, J.; Giguere, V.; Hurst, J.; van Agthoven, T.; Geurts

[117579]van Kessel, A.; Goyert, S.; Grosveld, F.: Chromosomal lo-

calization

[117580]of the human Thy–1 gene. Proc. Nat. Acad. Sci. 82:

5832–5835, 1985.

[117581]20779.

[117582]Williams, A. F.; Gagnon, J.: Neuronal cell Thy–1 glycopro-

tein:

[117583]homology with immunoglobulin. Science 216: 696–703,

1982.

[117584]20780.

[117585]Johansson, M.; Karlsson, A.: Cloning of the cDNA and
chromosome

[117586]localization of the gene for human thymidine kinase 2. J.
Biol. Chem. 272:

[117587]8454–8458, 1997.

[117588]20781.

[117589]Mandel, H.; Szargel, R.; Labay, V.; Elpeleg, O.; Saada, A.;
Shalata,

[117590]A.; Anbinder, Y.; Berkowitz, D.; Hartman, C.; Barak, M.;
Eriksson,

[117591]S.; Cohen, N.: The deoxyguanosine kinase gene is mu-
tated in individuals

[117592]with depleted hepatocerebral mitochondrial DNA. Nature
Genet. 29:

[117593]337–341, 2001. Note: Erratum: Nature Genet. 29: 491
only, 2001.

[117594]20782.

[117595]Saada, A.; Shaag, A.; Mandel, H.; Nevo, Y.; Eriksson, S.;
Elpeleg,

[117596]O.: Mutant mitochondrial thymidine kinase in mitochon-

drial DNA depletion

[117597]myopathy. Nature Genet. 29: 342–344, 2001.

[117598]20783.

[117599]Wang, L.; Munch–Petersen, B.; Herrstrom Sjoberg, A.; Hell–
man, U.;

[117600]Bergman, T.; Jornvall, H.; Eriksson, S.: Human thymidine
kinase 2:

[117601]molecular cloning and characterisation of the enzyme ac–
tivity with

[117602]antiviral and cytostatic nucleoside substrates. FEBS Lett.
443:

[117603]170–174, 1999.

[117604]20784.

[117605]Willecke, K.; Reuber, T.; Kucherlapati, R. S.; Ruddle, F. H.:

[117606]Human mitochondrial thymidine kinase is coded for by a
gene on chromosome

[117607]16 of the nucleus. Somat. Cell Genet. 3: 237–245, 1977.

[117608]20785.

[117609]Klobutcher, L. A.; Ruddle, F. H.: Phenotype stabilisation
and

[117610]integration of transferred material in chromosome–medi–

ated gene transfer. Nature 280:

[117611]657-660, 1979.

[117612]20786.

[117613]Kuo, W.-L.; Hirschhorn, R.; Huie, M. L.; Hirschhorn, K.: Localization

[117614]and ordering of acid alpha-glucosidase (GAA) and thymidine kinase

[117615](TK1) by fluorescence in situ hybridization. Hum. Genet. 97: 404-406,

[117616]1996.

[117617]20787.

[117618]Chang, J. G.; Scarpa, A.; Eddy, R. L.; Byers, M. G.; Harris, A.

[117619]S.; Morrow, J. S.; Watkins, P.; Shows, T. B.; Forget, B. G.: Cloning

[117620]of a portion of the chromosomal gene and cDNA for human beta-fodrin,

[117621]the nonerythroid form of beta-spectrin. Genomics 17: 287-293, 1993.

[117622]20788.

[117623]Watkins, P. C.; Eddy, R.; Forget, B. G.; Chang, J. G.; Rochelle,

[117624]R.; Shows, T. B.: Assignment of a non-erythroid spectrin gene to

[117625]human chromosome 2. (Abstract) Am. J. Hum. Genet. 43: A161, 1988.

[117626]20789.

[117627]Birkenmeier, C. S.; McFarland-Starr, E. C.; Barker, J. E.: Chromosomal

[117628]location of three spectrin genes: relationship to the inherited hemolytic

[117629]anemias of mouse and man. Proc. Nat. Acad. Sci. 85: 8121-8125, 1988.

[117630]20790.

[117631]Agre, P.; Asimos, A.; Casella, J. F.; McMillan, C.: Inheritance

[117632]pattern and clinical response to splenectomy as a reflection of erythrocyte

[117633]spectrin deficiency in hereditary spherocytosis. New Eng. J. Med. 315:

[117634]1579-1583, 1986.

[117635]20791.

[117636]Alloisio, N.; Guetarni, D.; Morle, L.; Pothier, B.; Ducluzeau,

[117637]M. T.; Soun, A.; Colonna, P.; Clerc, M.; Philippe, N.; Delaunay, J.

[117638]: SP-alpha(I/65) hereditary elliptocytosis in North Africa.
Am. J.

[117639]Hemat. 23: 113-122, 1986.

[117640]20792.

[117641]Alloisio, N.; Morle, L.; Pothier, B.; Roux, A.-F.; Marechal, J.;

[117642]Ducluzeau, M.-T.; Benhadji-Zouaoui, Z.; Delaunay, J.:
Spectrin Oran

[117643](alpha II/21), a new spectrin variant concerning the alpha
II domain

[117644]and causing severe elliptocytosis in the homozygous
state. Blood 71:

[117645]1039-1047, 1988.

[117646]20793.

[117647]Alloisio, N.; Wilmotte, R.; Marechal, J.; Texier, P.; Denoroy,

[117648]L.; Feo, C.; Benhadji-Zouaoui, Z.; Delaunay, J.: A splice site
mutation

[117649]of alpha-spectrin gene causing skipping of exon 18 in
hereditary elliptocytosis. Blood 81:

[117650]2791-2798, 1993.

[117651]20794.

[117652]Alloisio, N.; Wilmotte, R.; Morle, L.; Baklouti, F.; Marechal,

[117653]J.; Ducluzeau, M.-T.; Denoroy, L.; Feo, C.; Forget, B. G.;

Kastally,

[117654]R.; Delaunay, J.: Spectrin Jendouba: an alpha-II/31 spectrin variant

[117655]that is associated with elliptocytosis and carries a mutation distant

[117656]from the dimer self-association site. Blood 80: 809-815, 1992.

[117657]20795.

[117658]Baklouti, F.; Marechal, J.; Morle, L.; Alloisio, N.; Wilmotte,

[117659]R.; Pothier, B.; Ducluzeau, M.-T.; Kastally, R.; Delaunay, J.:

Occurrence

[117660]of the alpha-I 22 arg-to-his (CGT-to-CAT) spectrin mutation in Tunisia:

[117661]potential association with severe elliptopoikilocytosis. Brit. J.

[117662]Haemat. 78: 108-113, 1991.

[117663]20796.

[117664]Boulanger, L.; LeCompte, M.-C.; Dhermy, D.; Garbarz, M.:

Fast

[117665]screening methods to detect mutations of spectrin in subjects with

[117666]hereditary elliptocytosis. (Letter) Am. J. Hum. Genet. 51: 440–442,

[117667]1992.

[117668]20797.

[117669]Burke, J. P. W. G.; Van Zyl, D.; Zail, S. S.; Coetzer, T. L.:

[117670]Reduced spectrin–ankyrin binding in a South African hereditary elliptocytosis

[117671]kindred homozygous for spectrin St Claude. (Letter) Blood 92: 2591–2592,

[117672]1998.

[117673]20798.

[117674]Coetzer, T.; Zail, S.: Spectrin tetramer–dimer equilibrium in

[117675]hereditary elliptocytosis. Blood 59: 900–905, 1982.

[117676]20799.

[117677]Coetzer, T. L.; Sahr, K.; Prchal, J.; Blacklock, H.; Peterson,

[117678]L.; Koler, R.; Doyle, J.; Manaster, J.; Palek, J.: Four different

[117679] mutations in codon 28 of alpha spectrin are associated with structurally

[117680] and functionally abnormal spectrin alpha(I/74) in hereditary elliptocytosis. J.

[117681] Clin. Invest. 88: 743-749, 1991.

[117682] 20800.

[117683] Dalla Venezia, N.; Wilmotte, R.; Morle, L.; Forissier, A.; Parquet,

[117684] N.; Garbarz, M.; Rousset, T.; Dhermy, D.; Alloisio, N.; Delaunay,

[117685] J.: An alpha-spectrin mutation responsible for hereditary elliptocytosis

[117686] associated in cis with the alpha-V/41 polymorphism. Hum. Genet. 90:

[117687] 641-644, 1993.

[117688] 20801.

[117689] Delaunay, J.; Dhermy, D.: Mutations involving the spectrin heterodimer

[117690] contact site: clinical expression and alterations in specific function. Seminars

[117691] Hemat. 30: 21-33, 1993.

[117692]20802.

[117693]del Giudice, E. M.; Ducluzeau, M. T.; Alloisio, N.; Wilmotte,

[117694]R.; Delaunay, J.; Perrotta, S.; Cutillo, S.; Iolascon, A.: Al-
pha(I/65)

[117695]hereditary elliptocytosis in southern Italy: evidence for an
African

[117696]origin. Hum. Genet. 89: 553-556, 1992.

[117697]20803.

[117698]Dhermy, D.; Lecomte, M. C.; Garbarz, M.; Bournier, O.;
Galand,

[117699]C.; Gautero, H.; Feo, C.; Alloisio, N.; Delaunay, J.; Boivin,
P.:

[117700]Spectrin beta-chain variant associated with hereditary el-
liptocytosis. J.

[117701]Clin. Invest. 70: 707-715, 1982.

[117702]20804.

[117703]Elgsaeter, A.; Stokke, B. T.; Mikkelsen, A.; Branton, D.: The

[117704]molecular basis of erythrocyte shape. Science 234:
1217-1223, 1986.

[117705]20805.

[117706]Miceli-Richard, C.; Lesage, S.; Rybojad, M.; Prieur, A.-M.;
Manouvrier-Hanu,

[117707]S.; Hafner, R.; Chamailard, M.; Zouali, H.; Thomas, G.;
Hugot, J.-P.

[117708]: CARD15 mutations in Blau syndrome. *Nature Genet.* 29:
19-20, 2001.

[117709]20806.

[117710]Bernards, A.; Rubin, C. M.; Westbrook, C. A.; Paskind, M.;
Baltimore,

[117711]D.: The first intron in the human c-abl gene is at least 200
kilobases

[117712]long and is a target for translocations in chronic myeloge-
nous leukemia. *Molec.*

[117713]*Cell. Biol.* 7: 3231-3236, 1987.

[117714]20807.

[117715]Clark, S. S.; McLaughlin, J.; Crist, W. M.; Champlin, R.;
Witte,

[117716]O. N.: Unique forms of the abl tyrosine kinase distinguish
Ph-1-positive

[117717]CML from Ph-1-positive ALL. *Science* 235: 85-88, 1986.

[117718]20808.

[117719]Collins, S. J.; Groudine, M. T.: Rearrangement and amplification
[117720]of c-abl sequences in the human chronic myelogenous
leukemia cell
[117721]line K-562. Proc. Nat. Acad. Sci. 80: 4813-4817, 1983.
[117722]20809.

[117723]Cong, F.; Spencer, S.; Cote, J.-F.; Wu, Y.; Tremblay, M. L.;
Lasky,
[117724]L. A.; Goff, S. P.: Cytoskeletal protein PSTPIP1 directs the
PEST-type
[117725]protein tyrosine phosphatase to the c-Abl kinase to mediate Abl dephosphorylation. Molec.
[117726]Cell 6: 1413-1423, 2000.
[117727]20810.

[117728]Fainstein, E.; Marcelle, C.; Rosner, A.; Canaani, E.; Gale, R.
[117729]P.; Dreazen, O.; Smith, S. D.; Croce, C. M.: A new fused
transcript
[117730]in Philadelphia chromosome positive acute lymphocytic
leukaemia. Nature 330:
[117731]386-388, 1987.
[117732]20811.

[117733]Goff, S. P.; D'Eustachio, P.; Ruddle, F. H.; Baltimore, D.:
Chromosomal

[117734]assignment of endogenous protooncogene c-abl. Science
218: 1317-1319,

[117735]1982.

[117736]20812.

[117737]Gong, J.; Costanzo, A.; Yang, H.-Q.; Melino, G.; Kaelin, W.
G.,

[117738]Jr.; Levvero, M.; Wang, J. Y. J.: The tyrosine kinase c-Abl
regulates

[117739]p73 in apoptotic response to cisplatin-induced DNA dam-
age. Nature 399:

[117740]806-809, 1999.

[117741]20813.

[117742]Han, L. Wong, D.; Dhaka, A.; Afar, D.; White, M.; Xie, W.;
Herschman,

[117743]H.; Witte, O.; Colicelli, J.: Protein binding and signaling
properties

[117744]of RIN1 suggest a unique effector function. Proc. Nat.
Acad. Sci. 94:

[117745]4954-4959, 1997.

[117746]20814.

[117747]Heisterkamp, N.; Groffen, J.; Stephenson, J. R.; Spurr, N.
K.;

[117748]Goodfellow, P. N.; Solomon, E.; Carritt, B.; Bodmer, W. F.:
Chromosomal

[117749]localization of human cellular homologues of two viral
oncogenes. Nature 299:

[117750]747-749, 1982.

[117751]20815.

[117752]Heisterkamp, N.; Stephenson, J. R.; Groffen, J.; Hansen, P.
F.;

[117753]de Klein, A.; Bartram, C. R.; Grosveld, G.: Localization of
the c-abl

[117754]oncogene adjacent to a translocation breakpoint in
chronic myelocytic

[117755]leukaemia. Nature 306: 239-242, 1983.

[117756]20816.

[117757]Hoffman-Falk, H.; Einat, P.; Shilo, B.-Z.; Hoffmann, F. M.:
Drosophila

[117758]melanogaster DNA clones homologous to vertebrate
oncogenes: evidence

[117759] for a common ancestor to the src and abl cellular genes.

Cell 32:

[117760] 589–598, 1983.

[117761] 20817.

[117762] Huntly, B. J. P.; Reid, A. G.; Bench, A. J.; Campbell, L. J.;

[117763] Telford, N.; Shepherd, P.; Szer, J.; Prince, H. M.; Turner, P.;

Grace,

[117764] C.; Nacheva, E. P.; Green, A. R.: Deletions of the derivative
chromosome

[117765] 9 occur at the time of the Philadelphia translocation and
provide

[117766] a powerful and independent prognostic indicator in
chronic myeloid

[117767] leukemia. Blood 98: 1732–1738, 2001.

[117768] 20818.

[117769] Jhanwar, S. C.; Neel, B. G.; Hayward, W. S.; Chaganti, R. S.
K.

[117770]: Localization of the cellular oncogenes ABL, SIS, and FES
on human

[117771] germ-line chromosomes. Cytogenet. Cell Genet. 38:
73–75, 1984.

[117772]20819.

[117773]Konopka, J. B.; Watanabe, S. M.; Singer, J. W.; Collins, S. J.;

[117774]Witte, O. N.: Cell lines and clinical isolates derived from
Ph1-positive

[117775]chronic myelogenous leukemia patients express c-abl
proteins with

[117776]a common structural alteration. Proc. Nat. Acad. Sci. 82:
1810-1814,

[117777]1985.

[117778]20820.

[117779]Kurzrock, R.; Shtalrid, M.; Romero, P.; Kloetzer, W. S.; Tal-
pas,

[117780]M.; Trujillo, J. M.; Blick, M.; Beran, M.; Gutterman, J. U.: A
novel

[117781]c-abl protein product in Philadelphia-positive acute lym-
phoblastic

[117782]leukaemia. Nature 325: 631-635, 1987.

[117783]20821.

[117784]Lane, M.-A.; Neary, D.; Cooper, G. M.: Activation of a cel-
lular

[117785]transforming gene in tumours induced by Abelson murine

leukaemia virus. Nature 300:

[117786]659–661, 1982.

[117787]20822.

[117788]Li, B.; Boast, S.; de los Santos, K.; Schieren, I.; Quiroz, M.;

[117789]Teitelbaum, S. L.; Tondravi, M. M.; Goff, S. P.: Mice deficient in

[117790]Abl are osteoporotic and have defects in osteoblast maturation. Nature

[117791]Genet. 24: 304–308, 2000.

[117792]20823.

[117793]Melo, J. V.; Gordon, D. E.; Cross, N. C. P.; Goldman, J. M.:

[117794]The ABL–BCR fusion gene is expressed in chronic myeloid leukemia. Blood 81:

[117795]158–165, 1993.

[117796]20824.

[117797]Pluk, H.; Dorey, K.; Superti-Furga, G.: Autoinhibition of c–Abl. Cell 108:

[117798]247–259, 2002.

[117799]20825.

- [117800]Schindler, T.; Bornmann, W.; Pellicena, P.; Miller, W. T.; Clarkson, B.; Kuriyan, J.: Structural mechanism for STI-571 inhibition of Abelson tyrosine kinase. Science 289: 1938–1941, 2000.
- [117803]20826.
- [117804]Schwartzberg, P. L.; Stall, A. M.; Hardin, J. D.; Bowdish, K. S.; Humaran, T.; Boast, S.; Harbison, M. L.; Robertson, E. J.; Goff, S. P.: Mice homozygous for the abl(m1) mutation show poor viability and depletion of selected B and T cell populations. Cell 65: 1165–1175, 1991.
- [117808]1991.
- [117809]20827.
- [117810]Scott, M. L.; Van Etten, R. A.; Daley, G. Q.; Baltimore, D.: bcr-abl causes hematopoietic disease distinct from that caused by bcr-abl. Proc. Nat. Acad. Sci. 88: 6506–6510, 1991.
- [117812]20828.

- [117814]Bouhassira, E. E.; Schwartz, R. S.; Yawata, Y.; Ata, K.;
Nagel,
[117815]R. L.; Rybicki, A. C.: An alanine to threonine substitution
in protein
[117816]4.2 cDNA is associated with a Japanese form of hereditary
hemolytic
[117817]anemia. (Abstract) Clin. Res. 39: 313A, 1991.
[117818]20829.
[117819]Davies, K. A.; Lux, S. E.: Hereditary disorders of the red
cell
[117820]membrane skeleton. Trends Genet. 5: 222–227, 1989.
[117821]20830.
[117822]Ghanem, A.; Pothier, B.; Marechal, J.; Ducluzeau, M. T.;
Morle,
[117823]L.; Alloisio, N.; Feo, C.; Ben Abdeladhim, A.; Fattoum, S.;
Delaunay,
[117824]J.: A haemolytic syndrome associated with the complete
absence of
[117825]red cell membrane protein 4.2 in two Tunisian siblings.
Brit. J.
[117826]Haemat. 75: 414–420, 1990.

[117827]20831.

[117828]Grenard, P.; Bates, M. K.; Aeschlimann, D.: Evolution of
transglutaminase

[117829]genes: identification of a transglutaminase gene cluster
on human

[117830]chromosome 15q15: structure of the gene encoding
transglutaminase

[117831]X and a novel gene family member, transglutaminase Z. J.
Biol. Chem. 276:

[117832]33066–33078, 2001.

[117833]20832.

[117834]Gwynn, B.; Korsgren, C.; Cohen, C. M.; Ciciotte, S. L.; Pe-
ters,

[117835]L. L.: The gene encoding protein 4.2 is distinct from the
mouse platelet

[117836]storage pool deficiency mutation pallid. Genomics 42:
532–535, 1997.

[117837]20833.

[117838]Hayashi, S.; Koomoto, R.; Yano, A.; Ishigami, S.; Tsujino,
G.;

[117839]Saeki, S.; Tanaka, T.: Abnormality in a specific protein of
the erythrocyte

[117840]membrane in hereditary spherocytosis. Biochem. Biophys.
Res. Commun. 57:

[117841]1038–1044, 1974.

[117842]20834.

[117843]Hayette, S.; Dhermy, D.; dos Santos, M.–E.; Bozon, M.;
Drenckhahn,

[117844]D.; Alloisio, N.; Texier, P.; Delaunay, J.; Morle, L.: A dele-
tional

[117845]frameshift mutation in protein 4.2 gene (allele 4.2 Lisboa)
associated

[117846]with hereditary hemolytic anemia. Blood 85: 250–256,
1995.

[117847]20835.

[117848]Hayette, S.; Morle, L.; Bozon, M.; Ghanem, A.; Risinger, M.;
Korsgren,

[117849]C.; Tanner, M. J. A.; Fattoum, S.; Cohen, C. M.; Delaunay,
J.: A

[117850]point mutation in the protein 4.2 gene (allele 4.2 Tozeur)
associated

[117851]with hereditary haemolytic anaemia. Brit. J. Haemat. 89:
762–770,

[117852]1995.

[117853]20836.

[117854]Ideguchi, H.; Nishimura, J.; Nawata, H.; Hamasaki, N.: A
genetic

[117855]defect of erythrocyte band 4.2 protein associated with
hereditary

[117856]spherocytosis. Brit. J. Haemat. 74: 347–353, 1990.

[117857]20837.

[117858]Iwamoto, S.; Kajii, E.; Omi, T.; Kamesaki, T.; Akifuji, Y.;
Ikemoto,

[117859]S.: Point mutation in the band 4.2 gene associated with
autosomal

[117860]recessively inherited erythrocyte band 4.2 deficiency. Eu-
rop. J.

[117861]Haemat. 50: 286–291, 1993.

[117862]20838.

[117863]Korsgren, C.; Cohen, C. M.: Organization of the gene for
human

[117864]erythrocyte membrane protein 4.2: structural similarities
with the

[117865]gene for the A subunit of factor XIII. Proc. Nat. Acad. Sci.
88:

[117866]4840-4844, 1991.

[117867]20839.

[117868]Korsgren, C.; Cohen, C. M.: cDNA sequence, gene sequence, and

[117869]properties of murine pallidin (band 4.2), the protein implicated in

[117870]the murine pallid mutation. Genomics 21: 478-485, 1994.

[117871]20840.

[117872]Korsgren, C.; Lawler, J.; Lambert, S.; Speicher, D.; Cohen, C.

[117873]M.: Complete amino acid sequence and homologies of human erythrocyte

[117874]membrane protein band 4.2. Proc. Nat. Acad. Sci. 87: 613-617, 1990.

[117875]20841.

[117876]Matsuda, M.; Hatano, N.; Ideguchi, H.; Takahira, H.; Fukumaki,

[117877]Y.: A novel mutation causing an aberrant splicing in the protein

[117878]4.2 gene associated with hereditary spherocytosis (protein 4.2-Notame). Hum.

[117879]Molec. Genet. 4: 1187–1191, 1995.

[117880]20842.

[117881]Najfeld, V.; Ballard, S. G.; Menninger, J.; Ward, D. C.;
Bouhassira,

[117882]E. E.; Schwartz, R. S.; Nagel, R. L.; Rybicki, A. C.: The gene
for

[117883]human erythrocyte protein 4.2 maps to chromosome
15q15. Am. J. Hum.

[117884]Genet. 50: 71–75, 1992.

[117885]20843.

[117886]Nozawa, Y.; Hogushi, T.; Iida, H.; Fukushima, H.; Sekiya,
T.;

[117887]Ito, Y.: Erythrocyte membranes in hereditary spherocyto-
sis: alterations

[117888]in surface ultrastructure and membrane proteins as in-
ferred by scanning

[117889]electron microscopy and SDS gel electrophoresis. Clin.
Chim. Acta 55:

[117890]81–85, 1974.

[117891]20844.

[117892]Perrotta, S.; Iolascon, A.; Polito, R.; d'Urzo, G.; Conte, M.

[117893]L.; Miraglia del Giudice, E.: 4.2 Nippon mutation in a non-Japanese

[117894]patient with hereditary spherocytosis. (Letter) Haematologica 84:

[117895]660-662, 1999.

[117896]20845.

[117897]Rybicki, A. C.; Heath, R.; Wolf, J. L.; Lubin, B.; Schwartz, R.

[117898]S.: Deficiency of protein 4.2 in erythrocytes from a patient with

[117899]a Coombs negative hemolytic anemia: evidence for a role of protein

[117900]4.2 in stabilizing ankyrin on the membrane. J. Clin. Invest. 81:

[117901]893-901, 1988.

[117902]20846.

[117903]Sung, L. A.; Chien, S.; Chang, L.-S.; Lambert, K.; Bliss, S. A.;

[117904]Bouhassira, E. E.; Nagel, R. L.; Schwartz, R. S.; Rybicki, A. C.:

[117905]Molecular cloning of human protein 4.2: a major component of the erythrocyte

[117906]membrane. Proc. Nat. Acad. Sci. 87: 955–959, 1990.

[117907]20847.

[117908]Sung, L. A.; Fan, Y. S.; Lambert, K.; Chien, S.; Lin, C.: Map-
ping

[117909]of gene for human erythrocyte protein 4.2 to chromoso-
mal region 15q15–q21.

[117910](Abstract) Cytogenet. Cell Genet. 58: 1996, 1991.

[117911]20848.

[117912]White, R. A.; Dowler, L. L.; Hummel, G. S.; Adkison, L. R.:
Exclusion

[117913]of Ebp4.2 as a candidate for the mouse mutant pallid.

Mouse News

[117914]Letter 95: 492–494, 1997.

[117915]20849.

[117916]White, R. A.; Peters, L. L.; Adkison, L. R.; Korsgren, C.; Co-
hen,

[117917]C. M.; Lux, S. E.: The murine pallid mutation is a platelet
storage

[117918]pool disease associated with the protein 4.2 (pallidin)
gene. Nature

[117919]Genet. 2: 80–83, 1992.

[117920]20850.

[117921]Blank, V.; Andrews, N. C.: The Maf transcription factors:
regulators

[117922]of differentiation. Trends Biochem. Sci. 22: 437–441,
1997.

[117923]20851.

[117924]Jamieson, R. V.; Perveen, R.; Kerr, B.; Carette, M.; Yardley,
J.;

[117925]Heon, E.; Wirth, M. G.; van Heyningen, V.; Donnai, D.; Mu-
nier, F.;

[117926]Black, G. C. M.: Domain disruption and mutation of the
bZIP transcription

[117927]factor, MAF, associated with cataract, ocular anterior seg-
ment dysgenesis

[117928]and coloboma. Hum. Molec. Genet. 11: 33–42, 2002.

[117929]20852.

[117930]Kim, J. I.; Li, T.; Ho, I. C.; Grusby, M. J.; Glimcher, L. H.:

[117931]Requirement for the c-Maf transcription factor in crystallin
gene

[117932]regulation and lens development. Proc. Nat. Acad. Sci. 96:
3781–3785,

[117933]1999.

[117934]20853.

[117935]Nishizawa, M.; Kataoka, K.; Goto, N.; Fujiwara, K. T.;
Kawai, S.

[117936]: v-maf, a viral oncogene that encodes a 'leucine zipper'
motif. Proc.

[117937]Nat. Acad. Sci. 86: 7711-7715, 1989.

[117938]20854.

[117939]Yoshida, M. C.; Nishizawa, M.; Kataoka, K.; Goto, N.; Fuji-
wara,

[117940]K. T.; Kawai, S.: Localization of the human MAF pro-
tooncogene on

[117941]chromosome 16 to bands q22-q23. (Abstract) Cytogenet.
Cell Genet. 58:

[117942]2003 only, 1991.

[117943]20855.

[117944]Harbers, K.; Kuehn, M.; Delius, H.; Jaenisch, R.: Insertion
of

[117945]retrovirus into the first intron of alpha 1 (I) collagen gene
to embryonic

[117946]lethal mutation in mice. Proc. Nat. Acad. Sci. 81:

1504–1508, 1984.

[117947]20856.

[117948]Howard, I. K.: Clyde Edgar Keeler (1900–1994): geneticist,
artist,

[117949]cultural historian. J. Hered. 86: 489–491, 1995.

[117950]20857.

[117951]Jenkins, N. A.; Copeland, N. G.; Taylor, B. A.; Lee, B. K.:
Dilute

[117952](d) coat colour mutation of DBA/2J mice is associated with
the site

[117953]of integration of an ecotropic MLV genome. Nature 293:
370–374,

[117954]1981.

[117955]20858.

[117956]Keeler, C. E.: The inheritance of a retinal abnormality in
white

[117957]mice. Proc. Nat. Acad. Sci. 10: 329–333, 1924.

[117958]20859.

[117959]Khramtsov, N. V.; Feshchenko, E. A.; Suslova, V. A.;
Shmukler,

[117960]B. E.; Terpufov, B. E.; Rakitina, T. V.; Atabekova, N. V.;

Lipkin,

[117961]V. M.: The human rod photoreceptor cGMP phosphodiesterase beta-subunit:

[117962]structural studies of its cDNA and gene. FEBS Lett. 327: 275-278,

[117963]1993.

[117964]20860.

[117965]Lem, J.; Flannery, J. G.; Li, T.; Applebury, M. L.; Farber, D.

[117966]B.; Simon, M. I.: Retinal degeneration is rescued in transgenic rd

[117967]mice by expression of the cGMP phosphodiesterase beta subunit. Proc.

[117968]Nat. Acad. Sci. 89: 4422-4426, 1992.

[117969]20861.

[117970]McKusick, V. A.: The effect of lithium on the electrocardiogram

[117971]of animals and relation of this effect to the ratio of the intracellular

[117972]and extracellular concentrations of potassium. J. Clin. Invest. 33:

[117973]598-610, 1954.

[117974]20862.

[117975]McLaughlin, M. E.; Ehrhart, T. L.; Berson, E. L.; Dryja, T. P.

[117976]: Mutation spectrum of the gene encoding the beta subunit of rod phosphodiesterase

[117977]among patients with autosomal recessive retinitis pigmentosa. Proc.

[117978]Nat. Acad. Sci. 92: 3249-3253, 1995.

[117979]20863.

[117980]McLaughlin, M. E.; Sandberg, M. A.; Berson, E. L.; Dryja, T. P.

[117981]: Recessive mutations in the gene encoding the beta-subunit of rod

[117982]phosphodiesterase in patients with retinitis pigmentosa. Nature Genet. 4:

[117983]130-134, 1993.

[117984]20864.

[117985]Pittler, S. J.; Baehr, W.: Identification of a nonsense mutation

[117986]in the rod photoreceptor cGMP phosphodiesterase beta-subunit gene

[117987]of the rd mouse. Proc. Nat. Acad. Sci. 88: 8322-8326,

1991.

[117988]20865.

[117989]Pittler, S. J.; Keeler, C. E.; Sidman, R. L.; Baehr, W.: PCR

[117990]analysis of DNA from 70-year-old sections of rodless
retina demonstrates

[117991]identity with the mouse rd defect. Proc. Nat. Acad. Sci. 90:
9616-9619,

[117992]1993.

[117993]20866.

[117994]Ray, K.; Baldwin, V. J.; Acland, G. M.; Blanton, S. H.;
Aguirre,

[117995]G. D.: Cosegregation of codon 807 mutation of the canine
rod cGMP

[117996]phosphodiesterase beta gene and rcd1. Invest. Ophthalm.
Vis. Sci. 35:

[117997]4291-4299, 1994.

[117998]20867.

[117999]Riess, O.; Kozak, C.; Van Ommen, G.-J.: Report of the
fourth

[118000]international workshop on human chromosome 4 map-
ping 1996. Cytogenet.

[118001]Cell Genet. 74: 57–69, 1996.

[118002]20868.

[118003]Riess, O.; Noerremoelle, A.; Collins, C.; Mah, D.; Weber, B.;

[118004]Hayden, M. R.: Exclusion of DNA changes in the beta-subunit of the

[118005]c-GMP phosphodiesterase gene as the cause for Huntington's disease. Nature

[118006]Genet. 1: 104–108, 1992.

[118007]20869.

[118008]Riess, O.; Noerremoelle, A.; Weber, B.; Musarella, M. A.; Hayden,

[118009]M. R.: The search for mutations in the gene for the beta subunit

[118010]of the cGMP phosphodiesterase (PDEB) in patients with autosomal recessive

[118011]retinitis pigmentosa. Am. J. Hum. Genet. 51: 755–762, 1992.

[118012]20870.

[118013]Sidman, R. L.; Green, M. C.: Retinal degeneration in the mouse. J.

[118014]Hered. 56: 23–29, 1965.

[118015]20871.

[118016]Suber, M. L.; Pittler, S. J.; Qin, N.; Wright, G. C.; Holcombe,

[118017]Y.; Lee, R. H.; Craft, C. M.; Lolley, R. N.; Baehr, W.; Hur-
witz, R.

[118018]L.: Irish setter dogs affected with rod/cone dysplasia con-
tain a

[118019]nonsense mutation in the rod cGMP phosphodiesterase
beta-subunit gene. Proc.

[118020]Nat. Acad. Sci. 90: 3968-3972, 1993.

[118021]20872.

[118022]Weber, B.; Collins, C.; Kowbel, D.; Riess, O.; Hayden, M. R.:

[118023]Identification of multiple CpG islands and associated con-
served sequences

[118024]in a candidate region for the Huntington disease gene.

Genomics 11:

[118025]1113-1124, 1991.

[118026]20873.

[118027]Weber, B.; Riess, O.; Hutchinson, G.; Collins, C.; Lin, B.;
Kowbel,

[118028]D.; Andrew, S.; Schappert, K.; Hayden, M. R.: Genomic or-
ganization

[118029]and complete sequence of the human gene encoding the
beta-subunit

[118030]of the cGMP phosphodiesterase and its localisation to
4p16.3. Nucleic

[118031]Acids Res. 19: 6263-6268, 1991.

[118032]20874.

[118033]Bear, J. C.; Nemec, T. F.; Kennedy, J. C.; Marshall, W. H.;
Power,

[118034]A. A.; Kolonel, V. M.; Burke, G. B.: Inbreeding in outport
Newfoundland. Am.

[118035]J. Med. Genet. 29: 649-660, 1988.

[118036]20875.

[118037]Fontaine, B.; Vale-Santos, J.; Jurkat-Rott, K.; Reboul, J.;
Plassart,

[118038]E.; Rime, C.-S.; Elbaz, A.; Heine, R.; Guimaraes, J.; Weis-
senbach,

[118039]J.; Baumann, N.; Fardeau, M.; Lehmann-Horn, F.: Mapping
of the hypokalaemic

[118040]periodic paralysis (HypoPP) locus to chromosome
1q31-32 in three European

[118041]families. Nature Genet. 6: 267-272, 1994.

[118042]20876.

[118043]Jurkat-Rott, K.; Mitrovic, N.; Hang, C.; Kouzmekine, A.;
laizzo,

[118044]P.; Herzog, J.; Lerche, H.; Nicole, S.; Vale-Santos, J.; Chau-
veau,

[118045]D.; Fontaine, B.; Lehmann-Horn, F.: Voltage-sensor
sodium channel

[118046]mutations cause hypokalemic periodic paralysis type 2 by
enhanced

[118047]inactivation and reduced current. Proc. Nat. Acad. Sci. 97:
9549-9554,

[118048]2000.

[118049]20877.

[118050]Ptacek, L. J.; Tawil, R.; Griggs, R. C.; Engel, A. G.; Layzer,

[118051]R. B.; Kwiecinski, H.; McManis, P. G.; Santiago, L.; Moore,
M.; Fouad,

[118052]G.; Bradley, P.; Leppert, M. F.: Dihydropyridine receptor
mutations

[118053]cause hypokalemic periodic paralysis. Cell 77: 863-868,
1994.

[118054]20878.

[118055]Armstrong, F. S.: Hyperkalemic familial periodic paralysis
(adynamia

[118056]episodica hereditaria). Ann. Intern. Med. 57: 455–461,
1962.

[118057]20879.

[118058]Bennani-Baiti, I. M.; Jones, B. K.; Liebhaber, S. A.; Cooke,
N.

[118059]E.: Physical linkage of the human growth hormone gene
cluster and

[118060]the skeletal muscle sodium channel alpha-subunit gene
(SCN4A) on chromosome

[118061]17. Genomics 29: 647–652, 1995.

[118062]20880.

[118063]Borg, K.; Hovmoller, M.; Larsson, L.; Edstrom, L.: Paramy-
otonia

[118064]congenita (Eulenberg): clinical, neurophysiological and
muscle biopsy

[118065]observations in a Swedish family. Acta Neurol. Scand. 87:
37–42,

[118066]1993.

[118067]20881.

- [118068]Burama, O. J. S.; Schipperheyn, J. J.: Periodic paralysis.In:
[118069]Vinken, P. J.; Bruyn, G. W.: Handbook of Clinical Neurology. Part
[118070]II, Diseases of Muscle. Amsterdam: Elsevier/North Holland
(pub.)
[118071]41: 1979. Pp. 147–174.
[118072]20882.
[118073]Eulenberg, A.: Ueber eine familiäre, durch 6 generationen
verfolgbare
[118074]Form congenitaler Paramyotonie. Neurologisches Central-
blatt 12:
[118075]265–272, 1886.
[118076]20883.
[118077]Feero, W. G.; Wang, J.; Barany, F.; Zhou, J.; Todorovic, S.
M.;
[118078]Conwit, R.; Galloway, G.; Hausmanowa-Petrusewicz, I.;
Fidzianska,
[118079]A.; Arahata, K.; Wessel, H. B.; Wadelius, C.; Marks, H. G.;
Hartlage,
[118080]P.; Hayakawa, H.; Hoffman, E. P.: Hyperkalemic periodic
paralysis:

[118081]rapid molecular diagnosis and relationship of genotype to phenotype

[118082]in 12 families. Neurology 43: 668–673, 1993.

[118083]20884.

[118084]Fontaine, B.; Khurana, T. S.; Hoffman, E. P.; Bruns, G. A. P.;

[118085]Haines, J. L.; Trofatter, J. A.; Hanson, M. P.; Rich, J.; McFarlane,

[118086]H.; Yasek, D. M.; Romano, D.; Gusella, J. F.; Brown, R. H., Jr.:

[118087]Hyperkalemic periodic paralysis and the adult muscle sodium channel

[118088]alpha-subunit gene. Science 250: 1000–1002, 1990.

[118089]20885.

[118090]George, A. L., Jr.; Ledbetter, D. H.; Kallen, R. G.; Barchi, R.

[118091]L.: Assignment of a human skeletal muscle sodium channel alpha-subunit

[118092]gene (SCN4A) to 17q23.1–25.3. Genomics 9: 555–556, 1991.

[118093]20886.

[118094]Ammann, A. J.; Cain, W. A.; Ishizaka, K.; Hong, R.; Good, R. A.

[118095]: Immunoglobulin E deficiency in ataxia–telangiectasia.

New Eng.

[118096]J. Med. 281: 469–472, 1969.

[118097]20887.

[118098]Amromin, G. D.; Boder, E.; Teplitz, R.: Ataxia–
telangiectasia

[118099]with a 32–year survival: a clinicopathological report. J.
Neuropath.

[118100]Exp. Neurol. 38: 621–643, 1979.

[118101]20888.

[118102]Athma, P.; Rappaport, R.; Swift, M.: Molecular genotyping
shows

[118103]that ataxia–telangiectasia heterozygotes are predisposed
to breast

[118104]cancer. Cancer Genet. Cytogenet. 92: 130–134, 1996.

[118105]20889.

[118106]Aurias, A.; Croquette, M. F.; Nuyts, J. P.; Griscelli, C.;
Dutrillaux,

[118107]B.: New data on clonal anomalies of chromosome 14 in
ataxia telangiectasia:

[118108]tct(14;14) and inv(14). Hum. Genet. 72: 22–24, 1986.

[118109]20890.

[118110]Aurias, A.; Dutrillaux, B.: A possible new type of chromosome
some

[118111]rearrangement: telomere-centromere translocation (tct)
followed by

[118112]double duplication. Hum. Genet. 72: 25-26, 1986.

[118113]20891.

[118114]Aurias, A.; Dutrillaux, B.: Probable involvement of im-
munoglobulin

[118115]superfamily genes in most recurrent chromosomal rear-
rangements from

[118116]ataxia telangiectasia. Hum. Genet. 72: 210-214, 1986.

[118117]20892.

[118118]Aurias, A.; Dutrillaux, B.; Buriot, D.; Lejeune, J.: High fre-
quencies

[118119]of inversions and translocations of chromosomes 7 and
14 in ataxia-telangiectasia. Mutat.

[118120]Res. 69: 369-374, 1980.

[118121]20893.

[118122]Aurias, A.; Dutrillaux, B.; Griscelli, C.: Tandem transloca-
tion

[118123]t(14;14) in isolated and clonal cells in ataxia telangiectasia
are

[118124]different. Hum. Genet. 63: 320–322, 1983.

[118125]20894.

[118126]Baker, B. S.; Boyd, J. B.; Carpenter, A. T. C.; Green, M. M.;
Nguyen,

[118127]T. D.; Ripoll, P.; Smith, P. D.: Genetic controls of meiotic
recombination

[118128]and somatic DNA metabolism in *Drosophila melanogaster*.
Proc. Nat.

[118129]Acad. Sci. 73: 4140–4144, 1976.

[118130]20895.

[118131]Banga, S. S.; Shenkar, R.; Boyd, J. B.: Hypersensitivity of
Drosophila

[118132]mei-41 mutants to hydroxyurea is associated with re-
duced mitotic chromosome

[118133]stability. Mutat. Res. 163: 157–165, 1986.

[118134]20896.

[118135]Banin, S.; Moyal, L.; Shieh, S.-Y.; Taya, Y.; Anderson, C.
W.;

[118136]Chessa, L.; Smorodinsky, N. I.; Prives, C.; Reiss, Y.; Shiloh,

Y.;

[118137]Ziv, Y.: Enhanced phosphorylation of p53 by ATM in response to DNA

[118138]damage. Science 281: 1674–1677, 1998.

[118139]20897.

[118140]Bao, S.; Tibbetts, R. S.; Brumbaugh, K. M.; Fang, Y.; Richardson,

[118141]D. A.; Ali, A.; Chen, S. M.; Abraham, R. T.; Wang, X.-F.: ATR/ATM-mediated

[118142]phosphorylation of human Rad17 is required for genotoxic stress responses. Nature 411:

[118143]969–974, 2001.

[118144]20898.

[118145]Bar, R. S.; Levis, W. R.; Rechler, M. M.; Harrison, L. C.; Siebert,

[118146]C.; Podskalny, J.; Roth, J.; Muggeo, M.: Extreme insulin resistance

[118147]in ataxia telangiectasia: defect in affinity of insulin receptors. New

[118148]Eng. J. Med. 298: 1164–1171, 1978.

[118149]20899.

[118150]Barlow, C.; Dennerly, P. A.; Shigenaga, M. K.; Smith, M. A.;
Morrow,
[118151]J. D.; Roberts, L. J., II; Wynshaw-Boris, A.; Levine, R. L.:
Loss
[118152]of the ataxia-telangiectasia gene product causes oxidative
damage
[118153]in target organs. Proc. Nat. Acad. Sci. 96: 9915-9919,
1999.
[118154]20900.
[118155]Barlow, C.; Eckhaus, M. A.; Schaffer, A. A.; Wynshaw-Boris,
A.
[118156]: Atm haploinsufficiency results in increased sensitivity to
sublethal
[118157]doses of ionizing radiation in mice. Nature Genet. 21:
359-360,
[118158]1999.
[118159]20901.
[118160]Barlow, C.; Hirotsume, S.; Paylor, R.; Liyanage, M.; Eckhaus,
[118161]M.; Collins, F.; Shiloh, Y.; Crawley, J. N.; Ried, T.; Tagle, D.;
[118162]Wynshaw-Boris, A.: Atm-deficient mice: a paradigm of
ataxia telangiectasia. Cell 86:

[118163]159–171, 1996.

[118164]20902.

[118165]Bay, J.–O.; Uhrhammer, N.; Pernin, D.; Presneau, N.;
Tchirkov,

[118166]A.; Vuillaume, M.; Laplace, V.; Grancho, M.; Verrelle, P.;
Hall, J.;

[118167]Bignon, Y.–J.: High incidence of cancer in a family segregating a

[118168]mutation of the ATM gene: possible role of ATM heterozygosity in cancer. Hum.

[118169]Mutat. 14: 485–492, 1999.

[118170]20903.

[118171]Becker, Y.; Tabor, E.; Asher, Y.: Ataxia–telangiectasia fibroblasts

[118172]have less fibronectin mRNA than control cells but have the same levels

[118173]of integrin and beta–actin mRNA. Hum. Genet. 81:
165–170, 1989.

[118174]20904.

[118175]Bender, M. A.; Rary, J. M.; Kale, R. P.: G(2) chromosomal radiosensitivity

[118176]in ataxia telangiectasia lymphocytes. Mutat.Res. 152:
39-47, 1985.

[118177]20905.

[118178]Bender, M. A.; Rary, J. M.; Kale, R. P.: G(0) chromosomal
radiosensitivity

[118179]in ataxia telangiectasia lymphocytes. Mutat. Res. 150:
277-282,

[118180]1985.

[118181]20906.

[118182]Bernstein, R.; Pinto, M.; Jenkins, T.: Ataxia telangiectasia

[118183]with evolution of monosomy 14 and emergence of
Hodgkin's disease. Cancer

[118184]Genet. Cytogenet. 4: 31-37, 1981.

[118185]20907.

[118186]Bigbee, W. L.; Langlois, R. G.; Swift, M.; Jensen, R. H.: Evi-
dence

[118187]for an elevated frequency of in vivo somatic cell mutations
in ataxia

[118188]telangiectasia. Am. J. Hum. Genet. 44: 402-408, 1989.

[118189]20908.

[118190] Bishop, D. T.; Hopper, J.: AT-tributable risks? Nature
Genet. 15:

[118191] 226 only, 1997.

[118192] 20909.

[118193] Bochkov, N. P.; Lopukhin, Y. M.; Kuleshov, N. P.; Ko-
valchuk, L.

[118194] V.: Cytogenetic study of patients with ataxia-
telangiectasia. Humangenetik 24:

[118195] 115-128, 1974.

[118196] 20910.

[118197] Boder, E.: Ataxia-telangiectasia: an overview. In: Gatti, R.
A.;

[118198] Swift, M.: Ataxia-telangiectasia: Genetics, Neuropathology
and Immunology

[118199] of a Degenerative Disease of Childhood. New York: Alan R.
Liss (pub.)

[118200] 1985. Pp. 1-63.

[118201] 20911.

[118202] Boder, E.: Ataxia-telangiectasia: some historic, clinical and

[118203] pathologic observations. In: Bergsma, D. (ed.): Immunode-
ficiency in

[118204]Man and Animals. New York: National Foundation–March
of Dimes (pub.)

[118205]1975. Pp. 255–300.

[118206]20912.

[118207]Boder, E.; Sedgwick, R. P.: Ataxia–telangiectasia: a familial

[118208]syndrome of progressive cerebellar ataxia, oculocuta-
neous telangiectasia

[118209]and frequent pulmonary infection. Pediatrics 21: 526–554,
1958.

[118210]20913.

[118211]Borghesani, P. R.; Alt, F. W.; Bottaro, A.; Davidson, L.; Ak-
soy,

[118212]S.; Rathbun, G. A.; Roberts, T. M.; Swat, W.; Segal, R. A.;
Gu, Y.

[118213]: Abnormal development of Purkinje cells and lymphocytes
in Atm mutant

[118214]mice. Proc. Nat. Acad. Sci. 97: 3336–3341, 2000.

[118215]20914.

[118216]Boyd, J. B.; Golino, M. D.; Nguyen, T. D.; Green, M. M.: Iso-
lation

[118217]and characterization of X-linked mutants of Drosophila

melanogaster

[118218] which are sensitive to mutagens. Genetics 84: 485–506,
1976.

[118219] 20915.

- [118220]Bridges, B. A.; Arlett, C. F.: Risk of breast cancer in ataxia-telangiectasia.
- [118221](Letter) New Eng. J. Med. 326: 1357-1361, 1992.
- [118222]20916.
- [118223]Bridges, B. A.; Harnden, D. G.: Ataxia-telangiectasia: A Cellular
- [118224]and Molecular Link between Cancer, Neuropathology, and Immune Deficiency.
- [118225]New York: John Wiley (pub.) 1982.
- [118226]20917.
- [118227]Barbosa, M. D. F. S.; Barrat, F. J.; Tchernev, V. T.; Nguyen, Q.
- [118228]A.; Mishra, V. S.; Colman, S. D.; Pastural, E.; Dufourcq-Lagelouse,
- [118229]R.; Fischer, A.; Holcombe, R. F.; Wallace, M. R.; Brandt, S. J.; de
- [118230]Saint Basile, G.; Kingsmore S. F.: Identification of mutations in
- [118231]two major mRNA isoforms of the Chediak-Higashi syndrome gene in human
- [118232]and mouse. Hum. Molec. Genet. 6: 1091-1098, 1997.

[118233]20918.

[118234]Barrat, F. J.; Auloge, L.; Pastural, E.; Dufourcq Lagelouse, R.;

[118235]Vilmer, E.; Cant, A. J.; Weissenbach, J.; Le Paslier, D.; Fischer,

[118236]A.; de Saint Basile, G.: Genetic and physical mapping of the Chediak-Higashi

[118237]syndrome on chromosome 1q42-43. Am. J. Hum. Genet. 59: 625-632,

[118238]1996.

[118239]20919.

[118240]Dufourcq-Lagelouse, R.; Lambert, N.; Duval, M.; Viot, G.; Vilmer,

[118241]E.; Fischer, A.; Prieur, M.; de Saint Basile, G.: Chediak-Higashi

[118242]syndrome associated with maternal uniparental isodisomy of chromosome

[118243]1. Europ. J. Hum. Genet. 7: 633-637, 1999.

[118244]20920.

[118245]Faigle, W.; Raposo, G.; Tenza, D.; Pinet, V.; Vogt, A. B.; Kropshofer,

- [118246]H.; Fischer, A.; de Saint-Basile, G.; Amigorena, S.: Deficient peptide loading and MHC class II endosomal sorting in a human genetic immunodeficiency disease: the Chediak-Higashi syndrome. J. Cell Biol. 141: 1121-1134, 1998.
- [118249]1998.
- [118250]20921.
- [118251]Karim, M. A.; Suzuki, K.; Fukai, K.; Oh, J.; Nagle, D. L.; Moore, K. J.; Barbosa, E.; Falik-Borenstein, T.; Filipovich, A.; Ishida, Y. Kivrikko, S.; Klein, C.; and 8 others: Apparent genotype-phenotype correlation in childhood, adolescent, and adult Chediak-Higashi syndrome. Am. J. Med. Genet. 108: 16-22, 2002.
- [118255]J. Med. Genet. 108: 16-22, 2002.
- [118256]20922.
- [118257]Kunieda, T.; Ide, H.; Nakagiri, M.; Yoneda, K.; Konfortov, B.; Ogawa, H.: Localization of the locus responsible for Chediak-Higashi

[118259]syndrome in cattle to bovine chromosome 28. Anim.

Genet. 31: 87–90,

[118260]2000.

[118261]20923.

[118262]El-Ashry, M. F.; El-Aziz, M. M. A.; Wilkins, S.; Cheetham,
M. E.;

[118263]Wilkie, S. E.; Hardcastle A. J.; Halford, S.; Bayoumi, A. Y.;

Ficker,

[118264]L. A.; Tuft, S.; Bhattacharya, S. S.; Ebenezer, N. D.: Identi-
fication

[118265]of novel mutations in the carbohydrate sulfotransferase
gene (CHST6)

[118266]causing macular corneal dystrophy. Invest. Ophthal. Vis.
Sci. 43:

[118267]377–382, 2002.

[118268]20924.

[118269]Akama, T. O.; Nishida, K.; Nakayama, J.; Watanabe, H.;

Ozaki, K.;

[118270]Nakamura, T.; Dota, A.; Kawasaki, S.; Inoue, Y.; Maeda, N.;

Yamamoto,

[118271]S.; Fujiwara, T.; Thonar, E. J.–M. A.; Shimomura, Y.; Ki-
noshita, S.;

[118272]Tanigami, A.; Fukuda, M. N.: Macular corneal dystrophy
type I and
[118273]type II are caused by distinct mutations in a new sulpho-
transferase
[118274]gene. Nature Genet. 26: 237–241, 2000.
[118275]20925.
[118276]Hasegawa, N.; Torii, T.; Kato, T.; Miyajima, H.; Furuhashi,
A.;
[118277]Nakayasu, K.; Kanai, A.; Habuchi, O.: Decreased GlcNAc
6-O-sulfotransferase
[118278]activity in the cornea with macular corneal dystrophy. In-
vest. Ophthalmol.
[118279]Vis. Sci. 41: 3670–3677, 2000.
[118280]20926.
[118281]Enomoto, A.; Kimura, H.; Chairoungdua, A.; Shigeta, Y.;
Jutabha,
[118282]P.; Cha, S. H.; Hosoyamada, M.; Takeda, M.; Sekine, T.;
Igarashi,
[118283]T.; Matsuo, H.; Kikuchi, Y.; Oda, T.; Ichida, K.; Hosoya, T.;
Shimokata,
[118284]K.; Niwa, T.; Kanai, Y.; Endou, H.: Molecular identification
of a

[118285]renal urate-anion exchanger that regulates blood urate levels. Nature 417:

[118286]447-452, 2002.

[118287]20927.

[118288]Anderson, S. L.; Coli, R.; Daly, I. W.; Kichula, E. A.; Rork, M.

[118289]J.; Volpi, S. A.; Ekstein, J.; Rubin, B. Y.: Familial dysautonomia

[118290]is caused by mutations of the IKAP gene. Am. J. Hum. Genet. 68:

[118291]753-758, 2001.

[118292]20928.

[118293]Blumenfeld, A.; Slaugenhaupt, S. A.; Axelrod, F. B.; Lucette, D.

[118294]E.; Maayan, C.; Liebert, C. B.; Ozelius, L. J.; Trofatter, J. A.;

[118295]Haines, J. L.; Breakefield, X. O.; Gusella, J. F.: Localization of

[118296]the gene for familial dysautonomia on chromosome 9 and definition

[118297]of DNA markers for genetic diagnosis. Nature Genet. 4: 160-164,

[118298]1993.

[118299]20929.

[118300]Blumenfeld, A.; Slaughter, S. A.; Liebert, C. B.; Temper, V.;

[118301]Maayan, C.; Gill, S.; Lucente, D. E.; Idelson, M.; McCormack, K.;

[118302]Monahan, M. A.; Mull, J.; Leyne, M.; Mendillo, M.; Schiripo, T.; Mishori,

[118303]E.; Breakefield, X.; Axelrod, F. B.; Gusella, J. F.: Precise genetic

[118304]mapping and haplotype analysis of the familial dysautonomia gene on

[118305]human chromosome 9q31. Am. J. Hum. Genet. 64: 1110–1118, 1999.

[118306]20930.

[118307]Xia, F.; Taghian, D. G.; DeFrank, J. S.; Zeng, Z.-C.; Willers,

[118308]H.; Iliakis, G.; Powell, S. N.: Deficiency of human BRCA2 leads to

[118309]impaired homologous recombination but maintains normal nonhomologous

[118310]end joining. Proc. Nat. Acad. Sci. 98: 8644–8649, 2001.

[118311]20931.

[118312]Goedert, M.; Hasegawa, J.; Craxton, M.; Leversha, M. A.;
Clegg,
[118313]S.: Assignment of the human stress-activated protein ki-
nase-3 gene
[118314](SAPK3) to chromosome 22q13.3 by fluorescence in situ
hybridization. Genomics 41:
[118315]501-502, 1997.
[118316]20932.
[118317]Wooster, R.; Bignell, G.; Lancaster, J.; Swift, S.; Seal, S.;
[118318]Mangion, J.; Collins, N.; Gregory, S.; Gumbs, C.; Micklem,
G.; Barfoot,
[118319]R.; Hamoudi, R.; Patel, S.; Rice, C.; Biggs, P.; Hashim, Y.;
Smith,
[118320]A.; Connor, F.; Arason, A.; Gudmundsson, J.; Ficenec, D.;
Keisell,
[118321]D.; Ford, D.; Tonin, P.; Bishop D. T.; Spurr, N. K.; Ponder, B.
A.
[118322]J.; Eeles, R.; Peto, J.; Devilee, P.; Cornelisse, C.; Lynch, H.;
Narod,
[118323]S.; Lenoir, G.; Egilsson, V.; Barkadottir, R. B.; Easton, D. F.;
Bentley,
[118324]D. R.; Futreal, P. A.; Ashworth, A.; Stratton, M. R.: Identifi-

cation

[118325]of the breast cancer susceptibility gene BRCA2. Nature
378: 789–792,

[118326]1995.

[118327]20933.

[118328]Wooster, R.; Neuhausen, S. L.; Mangion, J.; Quirk, Y.; Ford,
D.;

[118329]Collins, N.; Nguyen, K.; Seal, S.; Tran, T.; Averill, D.; Fields,

[118330]P.; Marshall, G.; Narod, S.; Lenoir, G. M.; Lynch, H.; Feun-
teun, J.;

[118331]Devilee, P.; Cornelisse, C. J.; Menko, F. H.; Daly, P. A.;
Ormiston,

[118332]W.; McManus, R.; Pye, C.; Lewis, C. M.; Cannon–Albright,
L. A.; Peto,

[118333]J.; Ponder, B. A. J.; Skolnick, M. H.; Easton, D. F.; Goldgar,
D.

[118334]E.; Stratton, M. R.: Localization of a breast cancer suscep-
tibility

[118335]gene, BRCA2, to chromosome 13q12–13. Science 265:
2088–2090, 1994.

[118336]20934.

[118337]Yang, H.; Jeffrey, P. D.; Miller, J.; Kinnucan, E.; Sun, Y.;

Thoma,

[118338]N. H.; Zheng, N.; Chen, P.-L.; Lee, W.-H.; Pavletich, N. P.:

BRCA2

[118339]function in DNA binding and recombination from a

BRCA2-DSS1-ssDNA

[118340]structure. Science 297: 1837-1848, 2002.

[118341]20935.

[118342]Hartley, D. A.; Preiss, A.; Artavanis-Tsakonas, S.: A deduced

[118343]gene product from the Drosophila neurogenic locus, enhancer of split,

[118344]shows homology to mammalian G-protein beta subunit.

Cell 55: 785-795,

[118345]1988.

[118346]20936.

[118347]Hou, E. W.; Li, S.-L.: Genomic organization and chromosome localization

[118348]to band 19p13.3 of the human AES gene: gene product exhibits strong

[118349]similarity to the N-terminal domain of Drosophila enhancer of split

[118350]Groucho protein. DNA Cell Biol. 17: 911-913, 1998.

[118351]20937.

[118352]Miyasaka, H.; Choudhury, B. K.; Hou, E. W.; Li, S. S.-L.:
Molecular

[118353]cloning and expression of mouse and human cDNA en-
coding AES and ESG

[118354]proteins with strong similarity to Drosophila enhancer of
split groucho

[118355]protein. Europ. J. Biochem. 216: 343-352, 1993.

[118356]20938.

[118357]Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek,

[118358]F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.;
Copeland, N.

[118359]G.: Murine chromosomal location of four hepatocyte-en-
riched transcription

[118360]factors: HNF-3-alpha, HNF-3-beta, HNF-3-gamma, and
HFN-4. Genomics 13:

[118361]264-268, 1992.

[118362]20939.

[118363]Deleuze, J. F.; Dhorne, S.; Hazan, J.; Borghi, E.; Raynaud,
N.;

[118364]Pollet, N.; Meunier-Rotival, M.; Deschatrette, J.; Alagille,

D.; Hadchouel,

[118365]M.: Deleted chromosome 20 from a patient with Alagille syndrome isolated

[118366]in a cell hybrid through leucine transport selection: study of three

[118367]candidate genes. Mammalian Genome 5: 663–669, 1994.

[118368]20940.

[118369]Kaestner, K. H.; Hiemisch, H.; Luckow, B.; Schutz, G.: The HNF-3

[118370]gene family of transcription factors in mice: gene structure, cDNA

[118371]sequence, and mRNA distribution. Genomics 20: 377–385, 1994.

[118372]20941.

[118373]Mincheva, A.; Lichter, P.; Schutz, G.; Kaestner, K. H.: Assignment

[118374]of the human genes for hepatocyte nuclear factor 3-alpha, -beta, and

[118375]-gamma (HNF3A, HNF3B, HNF3G) to 14q12–q13, 20p11, and 19q13.2–q13.4. Genomics 39:

[118376]417–419, 1997.

[118377]20942.

[118378]Adams, R. H.; Porras, A.; Alonso, G.; Jones, M.; Vintersten, K.;

[118379]Panelli, S.; Valladares, A.; Perez, L.; Klein, R.; Nebreda, A. R.

[118380]: Essential role of p38- α MAP kinase in placental but not embryonic

[118381]cardiovascular development. *Molec. Cell* 6: 109–116, 2000.

[118382]20943.

[118383]Ge, B.; Gram, H.; Di Padova, F.; Huang, B.; New, L.; Ulevitch,

[118384]R. J.; Luo, Y.; Han, J.: MAPKK-independent activation of p38- α

[118385]mediated by TAB1-dependent autophosphorylation of p38- α . *Science* 295:

[118386]1291–1294, 2002.

[118387]20944.

[118388]Han, J.; Lee, J.-D.; Bibbs, L.; Ulevitch, R. J.: A MAP kinase

[118389]targeted by endotoxin and hyperosmolarity in mammalian cells. *Science* 265:

[118390]808–811, 1994.

[118391]20945.

[118392]Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.;
Garsin,

[118393]D. A.; Inoue, H.; Tanaka-Hino, M.; Hisamoto, N.; Mat-
sumoto, K.; Tan,

[118394]M.-W.; Ausubel, F. M.: A conserved p38 MAP kinase path-
way in *Caenorhabditis*

[118395]*elegans* innate immunity. *Science* 297: 623-626, 2002.

[118396]20946.

[118397]Kumar, S.; McLaughlin, M. M.; McDonnell, P. C.; Lee, J. C.;
Livi,

[118398]G. P.; Young, P. R.: Human mitogen-activated protein ki-
nase CSBP1,

[118399]but not CSBP2, complements a *hog1* deletion in yeast. *J.*
Biol. Chem. 270:

[118400]29043-29046, 1995.

[118401]20947.

[118402]Lee, J. C.; Laydon, J. T.; McDonnell, P. C.; Gallagher, T. F.;

[118403]Kumar, S.; Green, D.; McNulty, D.; Blumenthal, M. J.; Heys,
J. R.;

[118404]Landvatter, S. W.; Stickler, J. E.; McLaughlin, M. M.;

Siemens, I.

[118405]R.; Fisher, S. M.; Livi, G. P.; White, J. R.; Adams, J. L.;

Young,

[118406]P. R.: A protein kinase involved in the regulation of in-

flammatory

[118407]cytokine biosynthesis. *Nature* 372: 739–746, 1994.

[118408]20948.

[118409]Liao, P.; Georgakopoulos, D.; Kovacs, A.; Zheng, M.;

Lerner, D.;

[118410]Pu, H.; Saffitz, J.; Chien, K.; Xiao, R.-P.; Kass, D. A.; Wang,

Y.

[118411]: The in vivo role of p38 MAP kinases in cardiac remodel-

ing and restrictive

[118412]cardiomyopathy. *Proc. Nat. Acad. Sci.* 98: 12283–12288,

2001.

[118413]20949.

[118414]Maizels, E. T.; Mukherjee, A.; Sithanandam, G.; Peters, C.

A.;

[118415]Cottom, J.; Mayo, K. E.; Hunzicker-Dunn, M.: Develop-

mental regulation

[118416]of mitogen-activated protein kinase-activated kinases-2

and -3 (MAPKAPK-2/-3)

[118417]in vivo during corpus luteum formation in the rat. Molec.

Endocr. 15:

[118418]716–733, 2001.

[118419]20950.

[118420]McDonnell, P. C.; DiLella, A. G.; Lee, J. C.; Young, P. R.:

Localization

[118421]of the human stress responsive MAP kinase–like CSAIDs
binding protein

[118422](CSBP) gene to chromosome 6p21.3/21.2. Genomics 29:
301–302, 1995.

[118423]20951.

[118424]New, L.; Jiang, Y.; Zhao, M.; Liu, K.; Zhu, W.; Flood, L. J.;

[118425]Kato, Y.; Parry, G. C. N.; Han, J.: PRAK, a novel protein ki-
nase

[118426]regulated by the p38 MAP kinase. EMBO J. 17: 3372–3384,
1998.

[118427]20952.

[118428]Ni, H.; Wang, X. S.; Diener, K.; Yao, Z.: MAPKAPK5, a novel
mitogen–activate

[118429]d protein kinase (MAPK)–activated protein kinase, is a
substrate of

[118430]the extracellular-regulated kinase (ERK) and p38 kinase.

Biochem.

[118431]Biophys. Res. Commun. 243: 492–496, 1998.

[118432]20953.

[118433]Takekawa, M.; Maeda, T.; Saito, H.: Protein phosphatase
2C- α

[118434]inhibits the human stress-responsive p38 and JNK MAPK
pathways. EMBO

[118435]J. 17: 4744–4752, 1998.

[118436]20954.

[118437]Tamura, K.; Sudo, T.; Senftleben, U.; Dadak, A. M.; John-
son, R.;

[118438]Karin, M.: Requirement for p38- α in erythropoietin
expression:

[118439]a role for stress kinases in erythropoiesis. Cell 102:
221–231,

[118440]2000.

[118441]20955.

[118442]Haber, N.; Stengel, D.; Defer, N.; Roeckel, N.; Mattei,
M.–G.;

[118443]Hanoune, J.: Chromosomal mapping of human adenylyl

cyclase genes

[118444]type III, type V and type VI. Hum. Genet. 94: 69–73, 1994.

[118445]20956.

[118446]Sinnarajah, S.; Dessauer, C. W.; Srikumar, D.; Chen, J.;

Yuen,

[118447]J.; Yilma, S.; Dennis, J. C.; Morrison, E. E.; Vodyanoy, V.;

Kehrl,

[118448]J. H.: RGS2 regulates signal transduction in olfactory neu-

rons by

[118449]attenuating activation of adenylyl cyclase III. Nature 409:

1051–1055,

[118450]2001.

[118451]20957.

[118452]Thaler, J. P.; Lee, S.-K.; Jurata, L. W.; Gill, G. N.; Pfaff,

[118453]S. L.: LIM factor Lhx3 contributes to the specification of

motor

[118454]neuron and interneuron identity through cell-

type-specific protein-protein

[118455]interactions. Cell 110: 237–249, 2002.

[118456]20958.

[118457]Tebbutt, N. C.; Giraud, A. S.; Inglese, M.; Jenkins, B.; War-

ing,

[118458]P.; Clay, F. J.; Malki, S.; Alderman, B. M.; Grail, D.; Hol-
lande,

[118459]F.; Heath, J. K.; Ernst, M.: Reciprocal regulation of gas-
trointestinal

[118460]homeostasis by SHP2 and STAT-mediated trefoil gene ac-
tivation in gp130

[118461]mutant mice. Nature Med. 8: 1089-1097, 2002.

[118462]20959.

[118463]Brown, K. A.; Leek, J. P.; Lench, N. J.; Moynihan, L. M.;
Markham,

[118464]A. F.; Mueller, R. F.: Human sequences homologous to the
gene for

[118465]the cochlear protein Ocp-II do not map to currently known
non-syndromic

[118466]hearing loss loci. Ann. Hum. Genet. 60: 385-389, 1996.

[118467]20960.

[118468]Chen, H.; Thalmann, I.; Adams, J. C.; Avraham, K. B.;
Copeland,

[118469]N. G.; Jenkins, N. A.; Beier, D. R.; Corey, D. P.; Thalmann,
R.; Duyk,

[118470]G. M.: cDNA cloning, tissue distribution, and chromoso-

mal localization

[118471]of Ocp2, a gene encoding a putative transcription-associated factor

[118472]predominantly expressed in the auditory organs. Genomics 27: 389–398,

[118473]1995.

[118474]20961.

[118475]Demetrick, D. J.; Zhang, H.; Beach, D. H.: Chromosomal mapping

[118476]of the genes for the human CDK2/cyclin A-associated proteins p19 (SKP1A

[118477]and SKP1B) and p45 (SKP2). Cytogenet. Cell Genet. 73: 104–107, 1996.

[118478]20962.

[118479]Liang, Y.; Chen, H.; Asher, J. H., Jr.; Chang, C.-C.; Friedman,

[118480]T. B.: Human inner ear OCP2 cDNA maps to 5q22–5q35.2 with related

[118481]sequences on chromosomes 4p16.2–4p14, 5p13–5q22, 7pter–q22, 10 and

[118482]12p13–12qter. Gene 184: 163–167, 1997.

[118483]20963.

[118484] Sowden, J.; Morrison, K.; Schofield, J.; Putt, W.; Edwards, Y.

[118485] A novel cDNA with homology to an RNA polymerase II
elongation factors

[118486] maps to human chromosome 5q31 (TCEB1L) and to
mouse chromosome 11

[118487] (Tceb1l). Genomics 29: 145–151, 1995.

[118488] 20964.

[118489] Thalmann, I.; Rosenthal, H. L.; Moore, B. W.; Thalmann, R.:
Organ

[118490] of Corti-specific polypeptides: OCP-I and OCP-II. Arch.
Oto-Rhino-Laryngol. 226:

[118491] 123–128, 1980.

[118492] 20965.

[118493] Thalmann, I.; Takahashi, K.; Varghese, J.; Comegys, T. H.;
Thalmann,

[118494] R.: Biochemical features of major organ of Corti proteins
(OCP-I

[118495] and OCP-II) including partial amino acid sequence. Laryn-
goscope 100:

[118496] 99–105, 1990.

[118497] 20966.

[118498]Zhang, H.; Kobayashi, R.; Galaktionov, K.; Beach, D.:

p19Skl

[118499]and p45Skl are essential elements of the cyclin A-CDK2
S phase kinase. Cell 82:

[118500]915-925, 1995.

[118501]20967.

[118502]FitzPatrick, D. R.; Germain-Lee, E.; Valle, D.: Isolation and

[118503]characterization of rat and human cDNAs encoding a
novel putative

[118504]peroxisomal enoyl-CoA hydratase. Genomics 27:
457-466, 1995.

[118505]20968.

[118506]Saijo, M.; Sakai, Y.; Kishino, T.; Niikawa, N.; Matsuura, Y.;
Morino,

[118507]K.; Tamai, K.; Taya, Y.: Molecular cloning of a human pro-
tein that

[118508]binds to the retinoblastoma protein and chromosomal
mapping. Genomics 27:

[118509]511-519, 1995.

[118510]20969.

[118511]Anand, A.; Chada, K.: In vivo modulation of Hmgic reduces

obesity. Nature

[118512]Genet. 24: 377–380, 2000.

[118513]20970.

[118514]Arlotta, P.; Tai, A. K.-F.; Manfioletti, G.; Clifford, C.; Jay,

[118515]G.; Ono, S. J.: Transgenic mice expressing a truncated
form of the

[118516]high mobility group I-C protein develop adiposity and an
abnormally

[118517]high prevalence of lipomas. J. Biol. Chem. 275:
14394–14400, 2000.

[118518]20971.

[118519]Ashar, H. R.; Cherath, L.; Przybysz, K. M.; Chada, K.: Ge-
nomic

[118520]characterization of human HMGIC, a member of the ac-
cessory transcription

[118521]factor family found at translocation breakpoints in lipo-
mas. Genomics 31:

[118522]207–214, 1996.

[118523]20972.

[118524]Ashar, H. R.; Schoenberg Fejzo, M.; Tkachenko, A.; Zhou,
X.; Fletcher,

[118525]J. A.; Weremowicz, S.; Morton, C. C.; Chada, K.: Disruption
of the
[118526]architectural factor HMGI-C: DNA-binding AT hook motifs
fused in lipomas
[118527]to distinct transcriptional regulatory domains. Cell 82:
57-65,
[118528]1995.
[118529]20973.
[118530]Chau, K.-Y.; Patel, U. A.; Lee, K.-L. D.; Lam, H.-Y. P.;
Crane-Robinson,
[118531]C.: The gene for the human architectural transcription
factor HMGI-C
[118532]consists of five exons each coding for a distinct functional
element. Nucleic
[118533]Acids Res. 23: 4262-4266, 1995.
[118534]20974.
[118535]Danforth, E., Jr.: Failure of adipocyte differentiation causes
[118536]type II diabetes mellitus. Nature Genet. 26: 13 only, 2000.
[118537]20975.
[118538]Friedmann, M.; Holth, L. T.; Zoghbi, H. Y.; Reeves, R.: Or-
ganization,

[118539]inducible-expression and chromosome localization of the human HMG-I(Y)

[118540]nonhistone protein gene. Nucleic Acids Res. 21: 4259-4267, 1993.

[118541]20976.

[118542]Ishwad, C. S.; Shriver, M. D.; Lassige, D. M.; Ferrell, R. E.:

[118543]The high mobility group I-C gene (HMGI-C): polymorphism and genetic

[118544]localization. Hum. Genet. 99: 103-105, 1997.

[118545]20977.

[118546]Kazmierczak, B.; Dal Cin, P.; Wanschura, S.; Bartnitzke, S.; Van

[118547]den Berghe, H.; Bullerdiek, J.: Cloning and molecular characterization

[118548]of part of a new gene fused to HMGIC in mesenchymal tumors. Am. J.

[118549]Path. 152: 431-435, 1998.

[118550]20978.

[118551]Kazmierczak, B.; Pohnke, Y.; Bullerdiek, J.: Fusion transcripts

[118552]between the HMGIC gene and RTVL-H-related sequences

in mesenchymal

[118553] tumors without cytogenetic aberrations. Genomics 38:
223-226, 1996.

[118554] 20979.

[118555] Manfioletti, G.; Rustighi, A.; Mantovani, F.; Goodwin, G. H.;

[118556] Giancotti, V.: Isolation and characterization of the gene
coding

[118557] for murine high-mobility-group protein HMGI-C. Gene
167: 249-253,

[118558] 1995.

[118559] 20980.

[118560] Mine, N.; Kurose, K.; Nagai, H.; Doi, D.; Ota, Y.; Yoneyama,
K.;

[118561] Konishi, H.; Araki, T.; Emi, M.: Gene fusion involving
HMGIC is a

[118562] frequent aberration in uterine leiomyomas. J. Hum. Genet.
46: 408-412,

[118563] 2001.

[118564] 20981.

[118565] Nucci, M. R.; Weremowicz, S.; Neskey, D. M.; Sornberger,
K.; Tallini,

[118566]G.; Morton, C. C.; Quade, B. J.: Chromosomal translocation
t(8;12)

[118567]induces aberrant HMGIC expression in aggressive an-
giomyxoma of the

[118568]vulva. Genes Chromosomes Cancer 32: 172-176, 2001.

[118569]20982.

[118570]Oka, J.; Matsumoto, A.; Hosokawa, Y.; Inoue, S.: Molecular
cloning

[118571]of human cytosolic purine 5-prime-nucleotidase.
Biochem. Biophys.

[118572]Res. Commun. 205: 917-922, 1994.

[118573]20983.

[118574]Kingsmore, S. F.; Suh, D.; Seldin, M. F.: Genetic mapping
of the

[118575]glycine receptor alpha-3 subunit on mouse chromosome
8. Mammalian

[118576]Genome 5: 831-832, 1994.

[118577]20984.

[118578]Kuhse, J.; Schmieden, V.; Betz, H.: Identification and func-
tional

[118579]expression of a novel ligand binding subunit of the in-

hibitory glycine

[118580]receptor. J. Biol. Chem. 265: 22317–22320, 1990.

[118581]20985.

[118582]Nikolic, Z.; Laube, B.; Weber, R. G.; Lichter, P.; Kioschis, P.;

[118583]Poustka, A.; Mulhardt, C.; Becker, C.–M.: The human
glycine receptor

[118584]subunit alpha-3: GLRA3 gene structure, chromosomal localization, and

[118585]functional characterization of alternative transcripts. J.
Biol.

[118586]Chem. 273: 19708–19714, 1998.

[118587]20986.

[118588]Way, J. C.; Chalfie, M.: mec-3, a homeobox-containing
gene that

[118589]specifies differentiation of the touch receptor neurons in
C. elegans. Cell 54:

[118590]5–16, 1988.

[118591]20987.

[118592]Zhadanov, A. B.; Bertuzzi, S.; Taira, M.; Dawid, I. B.; West-
phal,

[118593]H.: Expression pattern of the murine LIM class homeobox

gene Lhx3

[118594]in subsets of neural and neuroendocrine tissues. Dev.

Dyn. 202:

[118595]354–364, 1995.

[118596]20988.

[118597]Zhadanov, A. B.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.

[118598]A.; Westphal, H.: Genomic structure and chromosomal localization

[118599]of the mouse LIM/homeobox gene Lhx3. Genomics 27: 27–32, 1995.

[118600]20989.

[118601]Seimiya, H.; Sawabe, T.; Inazawa, J.; Tsuruo, T.: Cloning, expression

[118602]and chromosomal localization of a novel gene for protein tyrosine

[118603]phosphatase (PTP-U2) induced by various differentiation-inducing agents. Oncogene 10:

[118604]1731–1738, 1995.

[118605]20990.

[118606]Wiggins, R. C.; Wiggins, J. E.; Goyal, M.; Wharram, B. L.; Thomas,

[118607]P. E.: Molecular cloning of cDNAs encoding human
GLEPP1, a membrane
[118608]protein tyrosine phosphatase: characterization of the
GLEPP1 protein
[118609]distribution in human kidney and assignment of the
GLEPP1 gene to
[118610]human chromosome 12p12-p13. Genomics 27: 174-181,
1995.
[118611]20991.

[118612]Jia, S.; VanDusen, W. J.; Diehl, R. E.; Kohl, N. E.; Dixon, R.
[118613]A. F.; Elliston, K. O.; Stern, A. M.; Friedman, P. A.: cDNA
cloning
[118614]and expression of bovine aspartyl (asparaginy) beta-
hydroxylase. J.
[118615]Biol. Chem. 267: 14322-14327, 1992.
[118616]20992.

[118617]Korioth, F.; Gieffers, C.; Frey, J.: Cloning and characteriza-
tion
[118618]of the human gene encoding aspartyl beta-hydroxylase.
Gene 150:
[118619]395-399, 1994.

[118620]20993.

[118621]Lavaissiere, L.; Jia, S.; Nishiyama, M.; de la Monte, S.;
Stern,

[118622]A. M.; Wands, J. R.; Friedman, P. A.: Overexpression of human aspartyl(asparaginy)-beta-hydroxylase

[118623]in hepatocellular carcinoma and cholangiocarcinoma. J.
Clin. Invest. 98:

[118624]1313-1323, 1996.

[118625]20994.

[118626]Lim, K. Y.; Hong, C.-S.; Kim, D. H.: cDNA cloning and
characterization

[118627]of human cardiac junctin. Gene 255: 35-42, 2000.

[118628]20995.

[118629]Scott, A. F.: Personal Communication. Baltimore, Md.
2/19/2001.

[118630]20996.

[118631]Treves, S.; Feriotto, G.; Moccagatta, L.; Gambari, R.;
Zorzato,

[118632]F.: Molecular cloning, expression, functional characterization,
chromosomal

[118633]localization, and gene structure of junctate, a novel inte-

gral calcium

[118634]binding protein of sarco(endo)plasmic reticulum membrane. J. Biol.

[118635]Chem. 275: 39555–39568, 2000.

[118636]20997.

[118637]Wetzel, G. T.; Ding, S.; Chen, F.: Molecular cloning of junctin

[118638]from human and developing rabbit heart. Molec. Genet. Metab. 69:

[118639]252–258, 2000.

[118640]20998.

[118641]Sato, K.; Mano, H.; Ariyama, T.; Inazawa, J.; Yazaki, Y.; Hirai,

[118642]H.: Molecular cloning and analysis of the human Tec protein-tyrosine

[118643]kinase. Leukemia 8: 1663–1672, 1994.

[118644]20999.

[118645]Zhou, P.; Sun, L. J.; Dotsch, V.; Wagner, G.; Verdine, G. L.:

[118646]Solution structure of the core NFATC1/DNA complex. Cell 92: 687–696,

[118647]1998.

[118648]21000.

[118649]McCaffrey, P. G.; Luo, C.; Kerppola, T. K.; Jain, J.; Badalian,

[118650]T. M.; Ho, A. M.; Burgeon, E.; Lane, W. S.; Lambert, J. N.;

Curran,

[118651]T.; Verdine, G. L.; Rao, A.; Hogan, P. G.: Isolation of the

cyclosporin-sensitive

[118652]T cell transcription factor NFATp. Science 262: 750-754,

1993.

[118653]21001.

[118654]Rengarajan, J.; Tang, B.; Glimcher, L. H.: NFATc2 and

NFATc3 regulate

[118655]TH2 differentiation and modulate TCR-responsiveness of

naive TH cells. Nature

[118656]Immun. 3: 48-54, 2002.

[118657]21002.

[118658]Abrams, W. R.; Ma, R.-I.; Kucich, U.; Bashir, M. M.; Decker,

S.;

[118659]Tsipouras, P.; McPherson, J. D.; Wasmuth, J. J.; Rosen-

bloom, J.:

[118660]Molecular cloning of the microfibrillar protein MFAP3 and

assignment

[118661]of the gene to human chromosome 5q32-q33.2. Ge-
nomics 26: 47-54,

[118662]1995.

[118663]21003.

[118664]Atanasoski, S.; Toldo, S. S.; Malipiero, U.; Schreiber, E.;
Fries,

[118665]R.; Fontana, A.: Isolation of the human genomic brain-
2/N-Oct 3 gene

[118666](POUF3) and assignment to chromosome 6q16. Genomics
26: 272-280,

[118667]1995.

[118668]21004.

[118669]Schreiber, E.; Tobler, A.; Malipiero, U.; Schaffner, W.;
Fontana,

[118670]A.: cDNA cloning of human N-Oct 3, a nervous-system
specific POU

[118671]domain transcription factor binding to the octamer DNA
motif. Nucleic

[118672]Acids Res. 21: 253-258, 1993.

[118673]21005.

[118674]Xia, Y.-R.; Andersen, B.; Mehrabian, M.; Diep, A. T.; War-

den, C.

[118675]H.; Mohandas, T.; McEvilly, R. J.; Rosenfeld, M. G.; Lusic, A.
J.

[118676]; Chromosomal organization of mammalian POU domain
factors. Genomics 18:

[118677]126–130, 1993.

[118678]21006.

[118679]Brass, N.; Heckel, D.; Sahin, U.; Pfreundschuh, M.;
Sybrecht, G.

[118680]W.; Meese, E.: Translation initiation factor eIF-4gamma is
encoded

[118681]by an amplified gene and induces an immune response in
squamous cell

[118682]lung carcinoma. Hum. Molec. Genet. 6: 33–39, 1997.

[118683]21007.

[118684]Gradi, A.; Imataka, H.; Svitkin, Y. V.; Rom, E.; Raught, B.;
Morino,

[118685]S.; Sonenberg, N.: A novel functional human eukaryotic
translation

[118686]initiation factor 4G. Molec. Cell. Biol. 18: 334–342, 1998.

[118687]21008.

[118688]Imataka, H.; Gradi, A.; Sonenberg, N.: A newly identified
N-terminal

[118689]amino acid sequence of human eIF4G binds
poly(A)-binding protein and

[118690]functions in poly(A)-dependent translation. EMBO J. 17:
7480-7489,

[118691]1998.

[118692]21009.

[118693]Imataka, H.; Sonenberg, N.: Human eukaryotic translation
initiation

[118694]factor 4G (eIF4G) possesses two separate and independent
binding sites

[118695]for eIF4A. Molec. Cell. Biol. 17: 6940-6947, 1997.

[118696]21010.

[118697]Yan, R.; Rhoads, R. E.: Human protein synthesis initiation
factor

[118698]eIF-4-gamma is encoded by a single gene (EIF4G) that
maps to chromosome

[118699]3q27-qter. Genomics 26: 394-398, 1995.

[118700]21011.

[118701]Yan, R.; Rychlik, W.; Etchison, D.; Rhoads, R. E.: Amino

acid

[118702]sequence of the human protein synthesis initiation factor
eIF-4-gamma. J.

[118703]Biol. Chem. 267: 23226-23231, 1992.

[118704]21012.

[118705]Corrigall, A. V.; Hift, R. J.; Hancock, V.; Meissner, D.;
Davids,

[118706]L.; Kirsch, R. E.; Meissner, P. N.: Identification and charac-
terisation

[118707]of a deletion (537delAT) in the protoporphyrinogen oxi-
dase gene in

[118708]a South African variegate porphyria family. Hum. Mutat.
12: 403-407,

[118709]1998.

[118710]21013.

[118711]Aso, T.; Lane, W. S.; Conaway, J. W.; Conaway, R. C.: Elong-
gin

[118712](SIII): a multisubunit regulator of elongation by RNA poly-
merase II. Science 269:

[118713]1439-1443, 1995.

[118714]21014.

[118715] Baldwin, C. T.; Weiss, S.; Farrer, L.; De Stefano, A.; Adair, R.;

[118716] Franklyn, B.; Kidd, K. K.; Korostishevsky, M.; Bonne-Tamir, B.: Linkage

[118717] of congenital, recessive deafness (DFNB4) to chromosome 7q31 and evidence

[118718] for genetic heterogeneity in the Middle Eastern Druze population. Hum.

[118719] Molec. Genet. 4: 1637-1642, 1995.

[118720] 21015.

[118721] Li, X. C.; Everett, L. A.; Lalwani, A. K.; Desmukh, D.; Friedman,

[118722] T. B.; Green, E. D.; Wilcox, E. R.: A mutation in PDS causes non-syndromic

[118723] recessive deafness. (Letter) Nature Genet. 18: 215-217, 1998.

[118724] 21016.

[118725] Boon, L. M.; Mulliken, J. B.; Vikkula, M.; Watkins, H.; Seidman,

[118726] J.; Olsen, B. R.; Warman, M. L.: Assignment of a locus for dominantly

[118727] inherited venous malformations to chromosome 9p. Hum.

Molec. Genet. 3:

[118728]1583–1587, 1994.

[118729]21017.

[118730]Calvert, J. T.; Riney, T. J.; Kontos, C. D.; Cha, E. H.; Prieto,

[118731]V. G.; Shea, C. R.; Berg, J. N.; Nevin, N. C.; Simpson, S. A.;

Pasyk,

[118732]K. A.; Speer, M. C.; Peters, K. G.; Marchuk, D. A.: Allelic

and locus

[118733]heterogeneity in inherited venous malformations. Hum.

Molec. Genet. 8:

[118734]1279–1289, 1999.

[118735]21018.

[118736]Vikkula, M.; Boon, L. M.; Carraway, K. L., III; Calvert, J. T.;

[118737]Diamonti, A. J.; Goumnerov, B.; Pasyk, K. A.; Marchuk, D.

A.; Warman,

[118738]M. L.; Cantley, L. C.; Mulliken, J. B.; Olsen, B. R.: Vascular

dysmorphogenesis

[118739]caused by an activating mutation in the receptor tyrosine

kinase TIE2. Cell 87:

[118740]1181–1190, 1996.

[118741]21019.

[118742]Iwata, T.; Kogame, K.; Toki, T.; Yokoyama, M.; Yamamoto, M.; Ito,

[118743]E.: Structure and chromosome mapping of the human small maf genes

[118744]MAFG and MAFK. Cytogenet. Cell Genet. 82: 88–90, 1998.
[118745]21020.

[118746]Motohashi, H.; Katsuoka, F.; Shavit, J. A.; Engel, J. D.; Yamamoto,

[118747]M.: Positive or negative MARE-dependent transcriptional regulation

[118748]is determined by the abundance of small Maf proteins. Cell 103:

[118749]865–875, 2000.
[118750]21021.

[118751]Peters, L. L.; Eicher, E. M.: The ubiquitous subunit of the globin

[118752]enhancer-binding protein NF-E2 (Nfe2u) maps to mouse chromosome 5. Genomics 22:

[118753]490–491, 1994.
[118754]21022.

[118755]Shavit, J. A.; Motohashi, H.; Onodera, K.; Akasaka, J.; Ya-

mamoto,

[118756]M.; Engel, J. D.: Impaired megakaryopoiesis and behavioral defects

[118757]in mafG-null mutant mice. Genes Dev. 12: 2164–2174, 1998.

[118758]21023.

[118759]Toki, T.; Itoh, J.; Kitazawa, J.; Arai, K.; Hatakeyama, K.; Akasaka,

[118760]J.; Igarashi, K.; Nomura, N.; Yokoyama, M.; Yamamoto, M.; Ito, E.

[118761]: Human small Maf proteins form heterodimers with CNC family transcription

[118762]factors and recognize the NF-E2 motif. Oncogene 14: 1901–1910, 1997.

[118763]21024.

[118764]Yang, B.; He, B.; Abdel-Halim, S. M.; Tibell, A.; Brendel, M. D.;

[118765]Bretzel, R. G.; Efendic, S.; Hillert, J.: Molecular cloning of a

[118766]full-length cDNA for human type 3 adenylyl cyclase and its expression

[118767]in human islets. Biochem. Biophys. Res. Commun. 254: 548–551, 1999.

[118768]21025.

[118769]Chawengsaksophak, K.; James, R.; Hammond, V. E.; Kontgen, F.; Beck,

[118770]F.: Homeosis and intestinal tumours in Cdx2 mutant mice. Nature 386:

[118771]84–87, 1997.

[118772]21026.

[118773]Drummond, F.; Putt, W.; Fox, M.; Edwards, Y. H.: Cloning and chromosome

[118774]assignment of the human CDX2 gene. Ann. Hum. Genet. 61: 393–400,

[118775]1997.

[118776]21027.

[118777]German, M. S.; Wang, J.; Fernald, A. A.; Espinosa, R., III; Le

[118778]Beau, M. M.; Bell, G. I.: Localization of the genes encoding two

[118779]transcription factors, LMX1 and CDX3, regulating insulin gene expression

[118780]to human chromosomes 1 and 13. Genomics 24: 403–404, 1994.

[118781]21028.

- [118782]Woodford–Richens, K. L.; Halford, S.; Rowan, A.; Bevan, S.;
Aaltonen,
[118783]L. A.; Wasan, H.; Bicknell, D.; Bodmer, W. F.; Houlston, R.
S.; Tomlinson,
[118784]J. P. M.: CDX2 mutations do not account for juvenile poly-
posis or
[118785]Peutz–Jeghers syndrome and occur infrequently in spo-
radic colorectal
[118786]cancers. *Brit. J. Cancer* 84: 1314–1316, 2001.
[118787]21029.
- [118788]Hiroi, Y.; Kudoh, S.; Monzen, K.; Ikeda, Y.; Yazaki, Y.; Na-
gai,
[118789]R.; Komuro, I.: Tbx5 associates with Nkx2–5 and syner-
gistically promotes
[118790]cardiomyocyte differentiation. *Nature Genet.* 28: 276–280,
2001.
[118791]21030.
- [118792]Dubbink, H. J.; Verkaik, N. S.; Faber, P. W.; Trapman, J.;
Schroder,
[118793]F. H.; Romijn, J. C.: Tissue-specific and androgen-regu-
lated expression
[118794]of human prostate-specific transglutaminase. *Biochem. J.*

315: 901-908,

[118795]1996.

[118796]21031.

[118797]Gentile, V.; Grant, F. J.; Porta, R.; Baldini, A.: Localization

[118798]of the human prostate transglutaminase (type IV) gene
(TGM4) to chromosome

[118799]3p21.33-p22 by fluorescence in situ hybridization. Ge-
nomics 27:

[118800]219-220, 1995.

[118801]21032.

[118802]Grant, F. J.; Taylor, D. A.; Sheppard, P. O.; Mathewes, S. L.;

[118803]Lint, W.; Vanaja, E.; Bishop, P. D.; O'Hara, P. J.: Molecular
cloning

[118804]and characterization of a novel transglutaminase cDNA
from a human

[118805]prostate cDNA library. Biochem. Biophys. Res. Commun.
203: 1117-1123,

[118806]1994.

[118807]21033.

[118808]Takai, S.; Long, J. E.; Yamada, K.; Miki, T.: Chromosomal

localization

[118809]of the human ECT2 proto-oncogene to 3q26.1-q26.2 by
somatic cell analysis

[118810]and fluorescence in situ hybridization. Genomics 27:
220-222, 1995.

[118811]21034.

[118812]Takai, S.; Lorenzi, M. V.; Long, J. E.; Yamada, K.; Miki, T.:

[118813]Assignment of the Ect2 protooncogene to mouse chromo-
some band 3B by

[118814]in situ hybridization. Cytogenet. Cell Genet. 81: 83-84,
1998.

[118815]21035.

[118816]Barker, H. M.; Brewis, N. D.; Street, A. J.; Spurr, N. K.; Co-
hen,

[118817]P. T. W.: Three genes for protein phosphatase 1 map to
different

[118818]human chromosomes: sequence, expression and gene lo-
calisation of protein

[118819]serine/threonine phosphatase 1 beta (PPP1CB). Biochim.
Biophys. Acta 1220:

[118820]212-218, 1994.

[118821]21036.

[118822]Saadat, M.; Kakinoki, Y.; Mizuno, Y.; Kikuchi, K.; Yoshida, M.

[118823]C.: Chromosomal localization of human, rat, and mouse protein phosphatase

[118824]type 1 beta catalytic subunit genes (PPP1CB) by fluorescence in situ

[118825]hybridization. Jpn. J. Genet. 69: 697–700, 1994.

[118826]21037.

[118827]Padilla, C. A.; Bajalica, S.; Lagercrantz, J.; Holmgren, A.: The

[118828]gene for human glutaredoxin (GLRX) is localized to human chromosome

[118829]5q14. Genomics 32: 455–457, 1996.

[118830]21038.

[118831]Padilla, C. A.; Martinez-Galisteo, E.; Barcena, J. A.; Spyrou,

[118832]G.; Holmgren, A.: Purification from placenta, amino acid sequence,

[118833]structure comparisons and cDNA cloning of human glutaredoxin. Europ.

[118834]J. Biochem. 227: 27–34, 1995.

[118835]21039.

[118836]Raghavachari, N.; Krysan, K.; Xing K.; Lou, M. F.: Regulation
of thioltransferase expression in human lens epithelial
cells. Invest.
Ophthalm. Vis. Sci. 42: 1002–1008, 2001.
21040.

[118840]Liang, B. T.; Jacobson, K. A.: A physiological role of the
adenosine
A3 receptor: sustained cardioprotection. Proc. Nat. Acad.
Sci. 95:
6995–6999, 1998.
21041.

[118844]Monitto, C. L.; Levitt, R. C.; DiSilvestre, D.; Holroyd, K. J.
Localization of the A(3) adenosine receptor gene
(ADORA3) to human
chromosome 1p. Genomics 26: 637–638, 1995.
21042.

[118848]Sajjadi, F. G.; Firestein, G. S.: cDNA cloning and sequence
analysis
of the human A3 adenosine receptor. Biochim. Biophys.
Acta 1179:

[118850]105–107, 1993.

[118851]21043.

[118852]Zhao, Z.; Ravid, S.; Ravid, K.: Chromosomal mapping of
the mouse

[118853]A3 adenosine receptor gene, Adora3. Genomics 30:
118–119, 1995.

[118854]21044.

[118855]Zhou, Q.-Y.; Li, C.; Olah, M. E.; Johnson, R. A.; Stiles, G. L.;

[118856]Civelli, O.: Molecular cloning and characterization of an
adenosine

[118857]receptor: the A3 adenosine receptor. Proc. Nat. Acad. Sci.
89: 7432–7436,

[118858]1992.

[118859]21045.

[118860]Corset, V.; Nguyen-Ba-Charvet, K. T.; Forcet, C.; Moyse,
E.; Chedotal,

[118861]A.; Mehlen, P.: Netrin-1-mediated axon outgrowth and
cAMP production

[118862]requires interaction with adenosine A2b receptor. Nature
407: 747–750,

[118863]2000.

[118864]21046.

[118865]Mino, R. P.; Spoerri, P. E.; Caballero, S.; Player, D.; Belardinelli,

[118866]L.; Biaggioni, I.; Grant, M. B.: Adenosine receptor antagonists and

[118867]retinal neovascularization in vivo. Invest. Ophthalm. Vis. Sci. 42:

[118868]3320–3324, 2001.

[118869]21047.

[118870]Stein, E.; Zou, Y.; Poo, M.; Tessier-Lavigne, M.: Binding of DCC

[118871]by netrin-1 to mediate axon guidance independent of adenosine A2B

[118872]receptor activation. Science 291: 1976–1982, 2001.

[118873]21048.

[118874]Townsend-Nicholson, A.; Baker, E.; Sutherland, G. R.; Schofield,

[118875]P. R.: Localization of the adenosine A2b receptor subtype gene (ADORA2B)

[118876]to chromosome 17p11.2–p12 by FISH and PCR screening of somatic cell

[118877]hybrids. Genomics 25: 605–607, 1995.

[118878]21049.

[118879]von Gall, C.; Garabette, M. L.; Kell, C. A.; Frenzel, S.; De-
hghani,

[118880]F.; Schumm–Draeger, P.–M.; Weaver, D. R.; Korf, H.–W.;
Hastings, M.

[118881]H.; Stehle, J. H.: Rhythmic gene expression in pituitary de-
pends

[118882]on heterologous sensitization by the neurohormone mela-
tonin. Nature

[118883]Neurosci. 5: 234–238, 2002.

[118884]21050.

[118885]Hirai, S.; Izawa, M.; Osada, S.; Spyrou, G.; Ohno, S.: Acti-
vation

[118886]of the JNK pathway by distantly related protein kinases,
MEKK and

[118887]MUK. Oncogene 12: 641–650, 1996.

[118888]21051.

[118889]Holzman, L. B.; Merritt, S. E.; Fan, G.: Identification,
molecular

[118890]cloning, and characterization of dual leucine zipper bear-

ing kinase.

[118891]A novel serine/threonine protein kinase that defines a second subfamily

[118892]of mixed lineage kinases. J. Biol. Chem. 269: 30808–30817, 1994.

[118893]21052.

[118894]Mata, M.; Merritt, S. E.; Fan, G.; Yu, G. G.; Holzman, L. B.:

[118895]Characterization of dual leucine zipper-bearing kinase, a mixed lineage

[118896]kinase present in synaptic terminals whose phosphorylation state is

[118897]regulated by membrane depolarization via calcineurin. J. Biol. Chem. 271:

[118898]16888–16896, 1996.

[118899]21053.

[118900]Reddy, U. R.; Nycum, L.; Slavc, I.; Biegel, J. A.: Localization

[118901]of the human zipper protein kinase gene (ZPK) to chromosome 12q13

[118902]by fluorescence in situ hybridization and somatic cell hybrid analysis. Genomics 25:

[118903]597–598, 1995.

[118904]21054.

[118905]Reddy, U. R.; Pleasure, D.: Cloning of a novel putative protein

[118906]kinase having a leucine zipper domain from human brain. Biochem.

[118907]Biophys. Res. Commun. 202: 613–620, 1994.

[118908]21055.

[118909]Becq, F.; Hamon, Y.; Bajetto, A.; Gola, M.; Verrier, B.; Chimini,

[118910]G.: ABC1, an ATP binding cassette transporter required for phagocytosis

[118911]of apoptotic cells, generates a regulated anion flux after expression

[118912]in *Xenopus laevis* oocytes. J. Biol. Chem. 272: 2695–2699, 1997.

[118913]21056.

[118914]Decottignies, A.; Goffeau, A.: Complete inventory of the yeast

[118915]ABC proteins. Nature Genet. 15: 137–145, 1997.

[118916]21057.

[118917]Guo, Z.; Inazu, A.; Yu, W.; Suzumura, T.; Okamoto, M.;

Nohara,

[118918]A.; Higashikata, T.; Sano, R.; Wakasugi, K.; Hayakawa, T.;
Yoshida,

[118919]K.; Suehiro, T.; Schmitz, G.; Mabuchi, H.: Double deletions
and missense

[118920]mutations in the first nucleotide-binding fold of the ATP-
binding

[118921]cassette transporter A1 (ABCA1) gene in Japanese patients
with Tangier

[118922]disease. J. Hum. Genet. 47: 325-329, 2002.

[118923]21058.

[118924]Hong, S. H.; Rhyne, J.; Zeller, K.; Miller, M.: Novel ABCA1
compound

[118925]variant associated with HDL cholesterol deficiency.
Biochim. Biophys.

[118926]Acta 1587: 60-64, 2002.

[118927]21059.

[118928]Huang, W.; Moriyama, K.; Koga, T.; Hua, H.; Ageta, M.;
Kawabata,

[118929]S.; Mawatari, K.; Imamura, T.; Eto, T.; Kawamura, M.; Ter-
amoto, T.;

[118930]Sasaki, J.: Novel mutations in ABCA1 gene in Japanese pa-

tients with

[118931]Tangier disease and familial high density lipoprotein deficiency with

[118932]coronary heart disease. *Biochim. Biophys. Acta* 1537: 71–78, 2001.

[118933]21060.

[118934]Ishii, J.; Nagano, M.; Kujiraoka, T.; Ishihara, M.; Egashira,

[118935]T.; Takada, D.; Tsuji, M.; Hattori, H.; Emi, M.: Clinical variant

[118936]of Tangier disease in Japan: mutation of the ABCA1 gene in hypoalphalipoproteinemia

[118937]with corneal lipidosis. *J. Hum. Genet.* 47: 366–369, 2002.

[118938]21061.

[118939]Jennings, M. W.; Jones, R. W.; Wood, W. G.; Weatherall, D. J.

[118940]: Analysis of an inversion within the human beta globin gene cluster. *Nucleic*

[118941]*Acids Res.* 13: 2897–2906, 1985.

[118942]21062.

[118943]Kulozik, A. E.; Bellan-Koch, A.; Kohne, E.; Kleihauer, E.: A

[118944]deletion/inversion rearrangement of the beta-globin gene

cluster in

[118945]a Turkish family with delta-beta(0)-thalassemia intermedia. Blood 79:

[118946]2455-2459, 1992.

[118947]21063.

[118948]Langmann, T.; Klucken, J.; Reil, M.; Liebisch, G.; Luciani, M.-F.;

[118949]Chimini, G.; Kaminski, W. E.; Schmitz, G.: Molecular cloning of the

[118950]human ATP-binding cassette transporter 1 (hABC1): evidence for sterol-dependent

[118951]regulation in macrophages. Biochem. Biophys. Res. Commun. 257: 29-33,

[118952]1999.

[118953]21064.

[118954]Lawn, R. M.; Wade, D. P.; Garvin, M. R.; Wang, X.; Schwartz, K.;

[118955]Porter, J. G.; Seilhamer, J. J.; Vaughan, A. M.; Oram, J. F.: The

[118956]Tangier disease gene product ABC1 controls the cellular apolipoprotein-mediated

[118957]lipid removal pathway. J. Clin. Invest. 104: R25-R31,

1999.

[118958]21065.

[118959]Luciani, M. F.; Denizot, F.; Savary, S.; Mattei, M. G.; Chi-
mini,

[118960]G.: Cloning of two novel ABC transporters mapping on
human chromosome

[118961]9. Genomics 21: 150–159, 1994.

[118962]21066.

[118963]Lapicka–Bodzioch, K.; Bodzioch, M.; Krull, M.; Kielar, D.;
Probst,

[118964]M.; Kiec, B.; Andrikovics, H.; Bottcher, A.; Hubacek, J.;
Aslanidis,

[118965]C.; Suttorp, N.; Schmitz, G.: Homogeneous assay based on
52 primer

[118966]sets to scan for mutations of the ABCA1 gene and its ap-
plication in

[118967]genetic analysis of a new patient with familial high-
density lipoprotein

[118968]deficiency syndrome. Biochim. Biophys. Acta 1537: 42–48,
2001.

[118969]21067.

[118970]Marcil, M.; Boucher, B.; Krimbou, L.; Solymoss, B. C.; Davi-

gnon,
 [118971]J.; Frohlich, J.; Genest, J., Jr.: Severe familial HDL deficiency
 [118972]in French-Canadian kindreds: clinical, biochemical, and
 molecular
 [118973]characterization. *Arterioscler. Thromb. Vasc. Biol.* 15:
 1015-1024,
 [118974]1995.
 [118975]21068.
 [118976]Marcil, M.; Yu, L.; Krimbou, L.; Boucher, B.; Oram, J. F.;
 Cohn,
 [118977]J. S.; Genest, J., Jr.: Cellular cholesterol transport and ef-
 flux
 [118978]in fibroblasts are abnormal in subjects with familial HDL
 deficiency. *Arterioscler.*
 [118979]*Thromb. Vasc. Biol.* 19: 159-169, 1999.
 [118980]21069.
 [118981]McNeish, J.; Aiello, R. J.; Guyot, D.; Turi, T.; Gabel, C.;
 Aldinger,
 [118982]C.; Hoppe, K. L.; Roach, M. L.; Royer, L. J.; de Wet, J.; Broc-
 cardo,
 [118983]C.; Chimini, G.; Francone, O. L.: High density lipoprotein

deficiency

[118984]and foam cell accumulation in mice with targeted disruption of ATP-binding

[118985]cassette transporter-1. Proc. Nat. Acad. Sci. 97: 4245-4250, 2000.

[118986]21070.

[118987]Pullinger, C. R.; Hakamata, H.; Duchateau, P. N.; Eng, C.; Aouizerat,

[118988]B. E.; Cho, M. H.; Fielding, C. J.; Kane, J. P.: Analysis of hABC1

[118989]gene 5-prime end: additional peptide sequence, promoter region, and

[118990]four polymorphisms. Biochem. Biophys. Res. Commun. 271: 451-455,

[118991]2000.

[118992]21071.

[118993]Petit, M. M. R.; Schoenmakers, E. F. P. M.; Huysmans, C.; Geurts,

[118994]J. M. W.; Mandahl, N.; Van de Ven, W. J. M.: LHFP, a novel translocation

[118995]partner gene of HMGIC in a lipoma, is a member of a new family of

[118996]LHFP-like genes. Genomics 57: 438–441, 1999.

[118997]21072.

[118998]Nishita, Y.; Yoshida, I.; Sado, T.; Takagi, N.: Genomic imprinting

[118999]and chromosomal localization of the human MEST gene. Genomics 36:

[119000]539–542, 1996.

[119001]21073.

[119002]Kohen, R.; Metcalf, M. A.; Khan, N.; Druck, T.; Huebner, K.; Lachowicz,

[119003]J. E.; Meltzer, H. Y.; Sibley, D. R.; Roth, B. L.; Hamblin, M. W.

[119004]: Cloning, characterization, and chromosomal localization of a human

[119005]5-HT-6 serotonin receptor. J. Neurochem. 66: 47–56, 1996.

[119006]21074.

[119007]Pearce, S. H. S.; Williamson, C.; Kifor, O.; Bai, M.; Coulthard,

[119008]M. G.; Davies, M.; Lewis-Barned, N.; McCredie, D.; Powell, H.; Kendall-Taylor,

- [119009]P.; Brown, E. M.; Thakker, R. V.: A familial syndrome of hypocalcemia with hypercalciuria due to mutations in the calcium-sensing receptor. New Eng. J. Med. 335: 1115–1122, 1996.
- [119012]21075.
- [119013]Frank, J.; Aita, V. M.; Ahmad, W.; Lam, H.; Wolff, C.; Christiano, A. M.: Identification of a founder mutation in the protoporphyrinogen oxidase gene in variegate porphyria patients from Chile. Hum. Hered. 51: 160–168, 2001.
- [119017]21076.
- [119018]Frank, J.; Jugert, F. K.; Breitkopf, C.; Goerz, G.; Merk, H. F.; Christiano, A. M.: Recurrent missense mutation in the protoporphyrinogen oxidase gene underlies variegate porphyria. Am. J. Med. Genet. 79: 22–26, 1998.
- [119022]21077.

- [119023] Orso, E.; Broccardo, C.; Kaminski, W. E.; Bottcher, A.;
Liebisch,
[119024] G.; Drobnik, W.; Gotz, A.; Chambenoit, O.; Diederich, W.;
Langmann,
[119025] T.; Spruss, T.; Luciani, M.-F.; Rothe, G.; Lackner, K. J.; Chi-
mini,
[119026] G.; Schmitz, G.: Transport of lipids from Golgi to plasma
membrane
[119027] is defective in Tangier disease patients and Abc1-deficient
mice. Nature
[119028] Genet. 24: 192-196, 2000.
[119029] 21078.
[119030] Hift, R. J.; Meissner, P. N.; Corrigall, A. V.; Ziman, M. R.;
Petersen,
[119031] L. A.; Meissner, D. M.; Davidson, B. P.; Sutherland, J.; Dai-
ley, H.
[119032] A.; Kirsch, R. E.: Variegate porphyria in South Africa,
1688-1996:
[119033] new developments in an old disease. S. Afr. Med. J. 87:
722-731,
[119034] 1997.
[119035] 21079.

- [119036]Hift, R. J.; Meissner, P. N.; Todd, G.; Kirby, P.; Bilsland, D.;
- [119037]Collins, P.; Ferguson, J.; Moore, M. R.: Homozygous variegate porphyria:
- [119038]an evolving clinical syndrome. *Postgrad. Med. J.* 69: 781–786, 1993.
- [119039]21080.
- [119040]Lam, H.; Dragan, L.; Tsou, H. C.; Merk, H.; Peacocke, M.; Goerz,
- [119041]G.; Sassa, S.; Poh-Fitzpatrick, M.; Bickers, D. R.; Christiano, A.
- [119042]M.: Molecular basis of variegate porphyria: a de novo insertion mutation
- [119043]in the protoporphyrinogen oxidase gene. *Hum. Genet.* 99: 126–129,
- [119044]1997.
- [119045]21081.
- [119046]Puy, H.; Robreau, A.-M.; Rosipal, R.; Nordmann, Y.; Deybach, J.-C.
- [119047]: Protoporphyrinogen oxidase: complete genomic sequence and polymorphisms
- [119048]in the human gene. *Biochem. Biophys. Res. Comm.* 226: 226–230, 1996.

[119049]21082.

[119050]Roberts, A. G.; Puy, H.; Dailey, T. A.; Morgan, R. R.; What-
ley,

[119051]S. D.; Dailey, H. A.; Martasek, P.; Nordmann, Y.; Deybach,
J.-C.;

[119052]Elder, G. H.: Molecular characterization of homozygous
variegate

[119053]porphyria. Hum. Molec. Genet. 7: 1921-1925, 1998.

[119054]21083.

[119055]Francavilla, A.; Hagiya, M.; Porter, K. A.; Polimeno, L.;
Ihara,

[119056]J.; Starzl, T. E.: Augmenter of liver regeneration: its place
in

[119057]the universe of hepatic growth factors. Hepatology 20:
747-757,

[119058]1994.

[119059]21084.

[119060]Giorda, R.; Hagiya, M.; Seki, T.; Shimonishi, M.; Sakai, H.;
Michaelson,

[119061]J.; Francavilla, A.; Starzl, T. E.; Trucco, M.: Analysis of the
structure

- [119062]and expression of the augmenter of liver regeneration
(ALR) gene. Molec.
[119063]Med. 2: 97–108, 1996.
- [119064]21085.
- [119065]Hagiya, M.; Francavilla, A.; Polimeno, L.; Ihara, I.; Sakai, H.;
- [119066]Seki, T.; Shimonishi, M.; Porter, K. A.; Starzl, T. E.: Cloning
and
[119067]sequence analysis of the rat augmenter of liver regenera-
tion (ALR)
[119068]gene: expression of biologically active recombinant ALR
and demonstration
[119069]of tissue distribution. Proc. Nat. Acad. Sci. 91:
8142–8146, 1994.
- [119070]Note: Erratum Proc. Nat. Acad. Sci. 92, p. 3076 only,
1995.
- [119071]21086.
- [119072]Lisowsky, T.; Weinstat–Saslow, D. L.; Barton, N.; Reeders,
S. T.;
- [119073]Schneider, M. C.: A new human gene located in the PKD1
region of
[119074]chromosome 16 is a functional homologue to ERV1 of
yeast. Genomics 29:

[119075]690-697, 1995.

[119076]21087.

[119077]Honda, H.; Inazawa, J.; Nishida, J.; Yazaki, Y.; Hirai, H.:

Molecular

[119078]cloning, characterization, and chromosomal localization
of a novel

[119079]protein-tyrosine phosphatase, HPTP eta. Blood 84:
4186-4194, 1994.

[119080]21088.

[119081]Ostman, A.; Yang, Q.; Tonks, N. K.: Expression of DEP-1, a
receptor-like

[119082]protein-tyrosine-phosphatase, is enhanced with increas-
ing cell density. Proc.

[119083]Nat. Acad. Sci. 91: 9680-9684, 1994.

[119084]21089.

[119085]Jacoby, A. S.; Webb, G. C.; Liu, M. L.; Kofler, B.; Hort, Y. J.;

[119086]Fathi, Z.; Bottema, C. D. K.; Shine, J.; Iismaa, T. P.: Struc-
tural

[119087]organization of the mouse and human GALR1 galanin re-
ceptor genes (Galnr

[119088]and GALNR) and chromosomal localization of the mouse

gene. Genomics 45:

[119089]496–508, 1997.

[119090]21090.

[119091]Tomoda, T.; Kurashige, T.; Moriki, T.; Yamamoto, H.; Fujimoto,

[119092]S.; Taniguchi, T.: Enhanced expression of
poly(ADP-ribose) synthetase

[119093]gene in malignant lymphoma. Am. J. Hemat. 37: 223–227,
1991.

[119094]21091.

[119095]Vasquez, K. M.; Marburger, K.; Intody, Z.; Wilson, J. H.:
Manipulating

[119096]the mammalian genome by homologous recombination.
Proc. Nat. Acad.

[119097]Sci. 98: 8403–8410, 2001.

[119098]21092.

[119099]Yu, S.-W.; Wang, H.; Poitras, M. F.; Coombs, C.; Bowers, W.
J.;

[119100]Federoff, H. J.; Poirier, G. G.; Dawson, T. M.; Dawson, V. L.:
Mediation

[119101]of poly(ADP-ribose) polymerase-1-dependent cell death

by apoptosis-inducing

[119102]factor. Science 297: 259–263, 2002.

[119103]21093.

[119104]Zabel, B. U.; Herzog, H.; Schneider, R.; Auer, B.; Hirsch-Kauffmann,

[119105]M.; Schweiger, M.: Chromosomal sublocalization of the gene for human

[119106]poly (ADP-ribose) polymerase (NAD⁺ ADP-ribosyltransferase) at 1q41–42.

[119107](Abstract) Cytogenet. Cell Genet. 51: 1115, 1989.

[119108]21094.

[119109]Davidson, M. K.; Le Beau, M. M.; Eddy, R. L.; Shows, T. B.; DiPietro,

[119110]L. A.; Kingzette, M.; Hanly, W. C.: Genetic mapping of the human

[119111]polymeric immunoglobulin receptor gene to chromosome region 1q31–q41. Cytogenet.

[119112]Cell Genet. 48: 107–111, 1988.

[119113]21095.

[119114]Hood, L.; Kronenberg, M.; Hunkapiller, T.: T cell antigen receptors

[119115]and the immunoglobulin supergene family. Cell 40:
225–229, 1985.

[119116]21096.

[119117]Krajci, P.; Gedde–Dahl, T., Jr.; Hoyheim, B.; Rogde, S.;
Olaisen,

[119118]B.; Brandtzaeg, P.: The gene encoding human transmem-
brane secretory

[119119]component (locus PIGR) is linked to D1S58 on chromo-
some 1. Hum. Genet. 90:

[119120]215–219, 1992.

[119121]21097.

[119122]Krajci, P.; Grzeschik, K. H.; Geurts van Kessel, A. H. M.;
Olaisen,

[119123]B.; Brandtzaeg, P.: The human transmembrane secretory
component (poly-Ig

[119124]receptor): molecular cloning, restriction fragment length
polymorphism

[119125]and chromosomal sublocalization. Hum. Genet. 87:
642–648, 1991.

[119126]21098.

[119127]Krajci, P.; Kvale, D.; Tasken, K.; Brandtzaeg, P.: Molecular
cloning

[119128]and exon–intron mapping of the gene encoding human transmembrane secretory

[119129]component (the poly-Ig receptor). *Europ. J. Immun.* 22: 2309–2315,

[119130]1992.

[119131]21099.

[119132]Zhang, J.–R.; Mostov, K. E.; Lamm, M. E.; Nanno, M.; Shimida, S.;

[119133]Ohwaki, M.; Tuomanen, E.: The polymeric immunoglobulin receptor translocates

[119134]pneumococci across human nasopharyngeal epithelial cells. *Cell* 102:

[119135]827–837, 2000.

[119136]21100.

[119137]Bear, J. C.; McManamon, P.; Morgan, J.; Payne, R. H.; Lewis, H.;

[119138]Gault, M. H.; Churchill, D. N.: Age at clinical onset and at ultrasonographic

[119139]detection of adult polycystic kidney disease: data for genetic counselling. *Am.*

[119140]*J. Med. Genet.* 18: 45–53, 1984.

[119141]21101.

[119142]Bear, J. C.; Parfrey, P. S.; Morgan, J. M.; Martin, C. J.;
Cramer,

[119143]B. C.: Autosomal dominant polycystic kidney disease: new
information

[119144]for genetic counselling. Am. J. Med. Genet. 43: 548-553,
1992.

[119145]21102.

[119146]Boichis, H.; Passwell, J.; David, R.; Miller, H.: Congenital
hepatic

[119147]fibrosis and nephronophthisis: a family study. Quart. J.
Med. 42:

[119148]221-233, 1973.

[119149]21103.

[119150]Breuning, M. H.; Snijdwint, F. G. M.; Dauwerse, J. G.;
Saris,

[119151]J. J.; Bakker, E.; Pearson, P. L.; van Ommen, G. J. B.: Two
step

[119152]procedure for early diagnosis of polycystic kidney disease
with polymorphic

[119153]DNA markers on both sides of the gene. J. Med. Genet. 27:
614-617,

[119154]1990.

[119155]21104.

[119156]Breuning, M. H.; Snijdwint, F. G. M.; Brunner, H.; Verwest, A.;

[119157]Ijdo, J. W.; Saris, J. J.; Dauwerse, J. G.; Blondin, L.; Keith, T.;

[119158]Callen, D. F.; Hyland, V. J.; Xiao, G. H.; Scherer, G.; Higgs, D.

[119159]R.; Harris, P.; Bachner, L.; Reeders, S. T.; Germino, G.; Pearson,

[119160]P. L.; van Ommen, G. J. B.: Map of 16 polymorphic loci on the short

[119161]arm of chromosome 16 close to the polycystic kidney disease gene (PKD1). J.

[119162]Med. Genet. 27: 603-613, 1990.

[119163]21105.

[119164]Burn, T. C.; Connors, T. D.; Dackowski, W. R.; Petry L. R.; Van

[119165]Raay, T. J.; Millholland, J. M.; Venet, M.; Miller, G.; Hakim, R.

[119166]M.; Landes, G. M.; Klinger, K. W.; Qian, F.; Onuchic, L. F.; Watnick,

- [119167]T.; Germino, G. G.; Doggett, N. A.: Analysis of the genomic sequence
for the autosomal dominant polycystic kidney disease (PKD1) gene predicts
the presence of a leucine-rich repeat. Hum. Molec. Genet. 4: 575-582,
1995.
- [119170]21106.
- [119171]21106.
- [119172]Chanmugam, D.; Rasaretnam, R.; Karunaratne, K. E. S.: Hereditary
spherocytosis and polycystic disease of the kidneys in four members
of a family. Am. J. Hum. Genet. 23: 66, 1971.
- [119174]21107.
- [119175]21107.
- [119176]Coto, E.; Aguado, S.; Alvarez, J.; Menendez Diaz, M. J.; Lopez-Larrea,
C.: Genetic and clinical studies in autosomal dominant polycystic
kidney disease type 1 (ADPKD1). J. Med. Genet. 29: 243-246, 1992.
- [119177]21108.
- [119179]21108.

[119180] de Villiers, J. N. P.; Hillermann, R.; Loubser, L.; Kotze, M. J.

[119181]: Spectrum of mutations in the HFE gene implicated in
haemochromatosis

[119182] and porphyria. Hum. Molec. Genet. 8: 1517–1522, 1999.

[119183] 21109.

[119184] Deybach, J.-C.; Puy, H.; Robreau, A.-M.; Lamoril, J.; Da
Silva,

[119185] V.; Grandchamp, B.; Nordmann, Y.: Mutations in the pro-
toporphyrinogen

[119186] oxidase gene in patients with variegate porphyria. Hum.
Molec. Genet. 5:

[119187] 407–410, 1996.

[119188] 21110.

[119189] Meissner, P. N.; Dailey, T. A.; Hift, R. J.; Ziman, M.; Corri-
gall,

[119190] A. V.; Roberts, A. G.; Meissner, D. M.; Kirsch, R. E.; Dailey,
H.

[119191] A.: A R59W mutation in human protoporphyrinogen oxi-
dase results in

[119192] decreased enzyme activity and is prevalent in South
Africans with

[119193]variegate porphyria. *Nature Genet.* 13: 95–97, 1996.

[119194]21111.

[119195]Kalman, K.; Nguyen, A.; Tseng–Crank, J.; Dukes, I. D.;
Chandy,

[119196]G.; Hustad, C. M.; Copeland, N. G.; Jenkins, N. A.; Mohren-
weiser,

[119197]H.; Brandriff, B.; Cahalan, M.; Gutman, G. A.; Chandy, K.
G.: Genomic

[119198]organization, chromosomal localization, tissue distribu-
tion, and biophysical

[119199]characterization of a novel mammalian Shaker–related
voltage–gated

[119200]potassium channel, Kv1.7. *J. Biol. Chem.* 273: 5851–5857,
1998.

[119201]21112.

[119202]Orkiszewski, K. G.; Tedesco, T. A.; Mellman, W. J.; Croce,
C.

[119203]M.: Linkage relationship between the genes for thymidine
kinase and

[119204]galactokinase in different primates. *Somat. Cell Genet.* 2:
21–26,

[119205]1976.

[119206]21113.

[119207]Dunham, I.; Collins, J.; Wade, R.; Scambler, P.: Possible
role

[119208]for COMT in psychosis associated with velo-cardio-facial
syndrome.

[119209](Letter) Lancet 340: 1361-1362, 1992.

[119210]21114.

[119211]Pulkka, A.; Ihalainen, R.; Suorsa, A.; Riviere, M.; Szpirer, J.;

[119212]Pajunen, A.: Structures and chromosomal localizations of
two rat

[119213]genes encoding S-adenosylmethionine decarboxylase.
Genomics 16:

[119214]342-349, 1993.

[119215]21115.

[119216]Radford, D. M.; Eddy, R.; Haley, L.; Henry, W. M.; Pegg, A.
E.;

[119217]Pajunen, A.; Shows, T. B.: Gene sequences coding for S-
adenosylmethionine

[119218]decarboxylase are present on human chromosome 6 and
the X and are

[119219]not amplified in colon neoplasia. Cytogenet. Cell Genet.

49: 285–288,

[119220]1989.

[119221]21116.

[119222]Kleymenova, E.; Ibraghimov-Beskrovnaya, O.; Kugoh, H.;
Everitt,

[119223]J.; Xu, H.; Kiguchi, K.; Landes, G.; Harris, P.; Walker, C.:

Tuberin-dependent

[119224]membrane localization of polycystin-1: a functional link
between polycystic

[119225]kidney disease and the TSC2 tumor suppressor gene.

Molec. Cell 7:

[119226]823–832, 2001.

[119227]21117.

[119228]Sandford, R.; Sgotto, B.; Burn, T.; Brenner, S.: The tuberin

[119229](TSC2), autosomal dominant polycystic kidney disease
(PKD1), and somatostatin

[119230]type V receptor (SSTR5) genes form a synteny group in the
Fugu genome. Genomics 38:

[119231]84–86, 1996.

[119232]21118.

[119233]Pedeutour, F.; Szpirer, C.; Nahon, J.–L.: Assignment of the

human

[119234]pro-melanin-concentrating hormone gene (PMCH) to
chromosome 12q23-q24

[119235]and two variant genes (PMCHL1 and PMCHL2) to chromo-
some 5p14 and 5q12-q13.

[119236]Genomics 19: 31-37, 1994.

[119237]21119.

[119238]Chen, S.-J.; Zelent, A.; Tong, J.-H.; Yu, H.-Q.; Wang, Z.-Y.;
Derre,

[119239]J.; Berger, R.; Waxman, S.; Chen, Z.: Rearrangements of the retinoic acid receptor alpha and promyelocytic leukemia zinc finger genes resulting from t(11;17)(q23;q21) in a patient with acute promyelocytic leukemia. J. Clin. Invest. 91: 2260–2267, 1993.

[119243]21120.

[119244]Chen, Z.; Brand, N. J.; Chen, A.; Chen, S.-J.; Tong, J.-H.; Wang, Z.-Y.; Waxman, S.; Zelent, A.: Fusion between a novel Kruppel-like zinc finger gene and the retinoic acid receptor-alpha locus due to a variant t(11;17) translocation associated with acute promyelocytic leukaemia. EMBO J. 12: 1161–1167, 1993.

[119249]21121.

[119250]Chen, Z.; Guidez, F.; Rousselot, P.; Agadir, A.; Chen, S.-J.; Wang, Z.-Y.; Degos, L.; Zelent, A.; Waxman, S.; Chomienne, C.: PLZF-RAR-alpha

- [119252] fusion proteins generated from the variant
t(11;17)(q23;q21) translocation
- [119253] in acute promyelocytic leukemia inhibit ligand-dependent
transactivation
- [119254] of wild-type retinoic acid receptors. Proc. Nat. Acad. Sci.
91:
1178-1182, 1994.
- [119256] 21122.
- [119257] Grignani, F.; De Matteis, S.; Nervi, C.; Tomassoni, L.; Gel-
metti,
[119258] V.; Cioce, M.; Fanelli, M.; Ruthardt, M.; Ferrara, F. F.; Za-
mir, I.;
[119259] Seiser, C.; Grignani, F.; Lazar, M. A.; Minucci, S.; Pelicci, P.
G.
- [119260]: Fusion proteins of the retinoic acid receptor-alpha re-
cruit histone
- [119261] deacetylase in promyelocytic leukaemia. Nature 391:
815-818, 1998.
- [119262] 21123.
- [119263] Lin, R. J.; Nagy, L.; Inoue, S.; Shao, W.; Miller, W. H., Jr.;
- [119264] Evans, R. M.: Role of the histone deacetylase complex in
acute promyelocytic

[119265]leukaemia. Nature 391: 811–814, 1998.

[119266]21124.

[119267]Burstedt, M. S.; Sandgren, O.; Holmgren, G.; Forsman-Semb, K.:

[119268]Bothnia dystrophy caused by mutations in the cellular retinaldehyde-binding

[119269]protein gene (RLBP1) on chromosome 15q26. Invest. Ophthalmol. Vis. Sci. 40:

[119270]995–1000, 1999.

[119271]21125.

[119272]Fournier, C. M.; Nicolas, G.; Gallagher, P. G.; Dhermy, D.; Grandchamp,

[119273]B.; Lecomte, M.-C.: Spectrin St Claude, a splicing mutation of the

[119274]human alpha-spectrin gene associated with severe poikilocytic anemia. Blood 89:

[119275]4584–4590, 1997.

[119276]21126.

[119277]Unger, S.; Korkko, J.; Krakow, D.; Lachman, R. S.; Rimoin, D.

[119278]L.; Cohn, D. H.: Double heterozygosity for pseudoachon-

droplasia and

[119279]spondyloepiphyseal dysplasia congenita. Am. J. Med. Genet. 104:

[119280]140–146, 2001.

[119281]21127.

[119282]Burstedt, M. S. I.; Forsman-Semb, K.; Golovleva, I.; Janunger,

[119283]T.; Wachtmeister, L.; Sandgren, O.: Ocular phenotype of Bothnia dystrophy,

[119284]an autosomal recessive retinitis pigmentosa associated with an R234W

[119285]mutation in the RLBP1 gene. Arch Ophthal. 119: 260–267, 2001.

[119286]21128.

[119287]Crabb, J. W.; Goldflam, S.; Harris, S. E.; Saari, J. C.: Cloning

[119288]of the cDNAs encoding the cellular retinaldehyde-binding protein from

[119289]bovine and human retina and comparison of the protein structures. J.

[119290]Biol. Chem. 263: 18688–18692, 1988.

[119291]21129.

[119292]Eichers, E. R.; Green, J. S.; Stockton, D. W.; Jackman, C. S.;

[119293]Whelan, J.; McNamara, J. A.; Johnson, G. J.; Lupski, J. R.;
Katsanis,

[119294]N.: Newfoundland rod-cone dystrophy, an early-onset
retinal dystrophy,

[119295]is caused by splice-junction mutations in RLBP1. Am. J.
Hum. Genet. 70:

[119296]955-964, 2002.

[119297]21130.

[119298]Granse, L.; Abrahamson, M.; Ponjavic, V.; Andreasson, S.:
Electrophysiological

[119299]findings in two young patients with Bothnia dystrophy and
a mutation

[119300]in the RLBP1 gene. Ophthal. Genet. 22: 97-105, 2001.

[119301]21131.

[119302]Intres, R.; Goldflam, S.; Cook, J. R.; Crabb, J. W.: Molecular

[119303]cloning and structural analysis of the human gene encod-
ing cellular

[119304]retinaldehyde-binding protein. J. Biol. Chem. 269:
25411-25418,

[119305]1994.

[119306]21132.

[119307]Katsanis, N.; Shroyer, N. F.; Lewis, R. A.; Cavender, J. C.;
Al-Rajhi,

[119308]A. A.; Jabak, M.; Lupski, J. R.: Fundus albipunctatus and
retinitis

[119309]punctata albescens in a pedigree with an R150Q mutation
in RLBP1. Clin.

[119310]Genet. 59: 424–429, 2001.

[119311]21133.

[119312]Maw, M. A.; Kennedy, B.; Knight, A.; Bridges, R.; Roth, K.
E.;

[119313]Mani, E. J.; Mukkadan, J. K.; Nancarrow, D.; Crabb, J. W.;
Denton,

[119314]M. J.: Mutation of the gene encoding cellular retinalde-
hyde-binding

[119315]protein in autosomal recessive retinitis pigmentosa. Na-
ture Genet. 17:

[119316]198–200, 1997.

[119317]21134.

[119318]Morimura, H.; Berson, E. L.; Dryja, T. P.: Recessive muta-
tions

[119319]in the RLBP1 gene encoding cellular retinaldehyde-binding protein

[119320]in a form of retinitis punctata albescens. Invest. Ophthalm. Vis.

[119321]Sci. 40: 1000–1004, 1999.

[119322]21135.

[119323]Sparkes, R. S.; Heinzmann, C.; Goldflam, S.; Kojis, T.; Saari,

[119324]J. C.; Mohandas, T.; Klisak, I.; Bateman, J. B.; Crabb, J. W.: Assignment

[119325]of the gene (RLBP1) for cellular retinaldehyde-binding protein (CRALBP)

[119326]to human chromosome 15q26 and mouse chromosome 7. Genomics 12: 58–62,

[119327]1992.

[119328]21136.

[119329]Blanton, S. H.; Heckenlively, J. R.; Cottingham, A. W.; Friedman,

[119330]J.; Sadler, L. A.; Wagner, M.; Friedman, L. H.; Daiger, S. P.: Linkage

[119331]mapping of autosomal dominant retinitis pigmentosa (RP1) to the pericentric

[119332]region of human chromosome 8. Genomics 11: 857–869, 1991.

[119333]21137.

[119334]Farrar, G. J.; McWilliam, P.; Bradley, D. G.; Kenna, P.; Lawler,

[119335]M.; Sharp, E. M.; Humphries, M. M.; Eiberg, H.; Conneally, P. M.;

[119336]Trofatter, J. A.; Humphries, P.: Autosomal dominant retinitis pigmentosa:

[119337]linkage to rhodopsin and evidence for genetic heterogeneity. Genomics 8:

[119338]35–40, 1990.

[119339]21138.

[119340]Wang, S.-Q.; Song, L.-S.; Lakatta, E. G.; Cheng, H.: Ca(2+) signalling

[119341]between single L-type Ca(2+) channels and ryanodine receptors in heart

[119342]cells. Nature 410: 592–596, 2001.

[119343]21139.

[119344]Laitinen, P. J.; Brown, K. M.; Piippo, K.; Swan, H.; Devaney, J.

- [119345]M.; Brahmabhatt, B.; Donarum, E. A.; Marino, M.; Tiso, N.; Viitasalo,
[119346]M.; Toivonen, L.; Stephan, D. A.; Kontula, K.: Mutations of
the cardiac
[119347]ryanodine receptor (RyR2) gene in familial polymorphic
ventricular
[119348]tachycardia. Circulation 103: 485–490, 2001.
[119349]21140.
- [119350]Marx, S. O.; Reiken, S.; Hisamatsu, Y.; Jayaraman, T.;
Burkhoff,
[119351]D.; Rosemblyt, N.; Marks, A. R.: PKA phosphorylation dis-
sociates
[119352]FKBP12.6 from the calcium release channel (ryanodine re-
ceptor): defective
[119353]regulation in failing hearts. Cell 101: 365–376, 2000.
[119354]21141.
- [119355]Mattei, M. G.; Giannini, G.; Moscatelli, F.; Sorrentino, V.:
Chromosomal
[119356]localization of murine ryanodine receptor genes RYR1,
RYR2, and RYR3
[119357]by in situ hybridization. Genomics 22: 202–204, 1994.

[119358]21142.

[119359]Otsu, K.; Willard, H. F.; Khanna, V. K.; Zorzato, F.; Green, N.

[119360]M.; MacLennan, D. H.: Molecular cloning of cDNA encoding the Ca^{2+}

[119361]release channel (ryanodine receptor) of rabbit cardiac muscle sarcoplasmic

[119362]reticulum. J. Biol. Chem. 265: 13472-13483, 1990.

[119363]21143.

[119364]Priori, S. G.; Napolitano, C.; Tiso, N.; Memmi, M.; Vignati, G.;

[119365]Bloise, R.; Sorrentino, V.; Danieli, G. A.: Mutations in the cardiac

[119366]ryanodine receptor gene (hRyR2) underlie catecholaminergic polymorphic

[119367]ventricular tachycardia. Circulation 102: r49-r53, 2000.

[119368]21144.

[119369]Zorzato, F.; Fujii, J.; Otsu, K.; Phillips, M.; Green, N. M.; Lai,

[119370]F. A.; Meissner, G.; MacLennan, D. H.: Molecular cloning of cDNA

- [119371] encoding human and rabbit forms of the Ca^{2+} release channel (ryanodine
- [119372] receptor) of skeletal muscle sarcoplasmic reticulum. J. Biol. Chem. 265:
- [119373] 2244–2256, 1990.
- [119374] 21145.
- [119375] Bertocchini, F.; Ovitt, C. E.; Conti, A.; Barone, V.; Scholer, H. R.; Bottinelli, R.; Reggiani, C.; Sorrentino, V.: Requirement
- [119376] for the ryanodine receptor type 3 for efficient contraction in neonatal
- [119377] skeletal muscles. EMBO J. 16: 6956–6963, 1997.
- [119379] 21146.
- [119380] Futatsugi, A.; Kato, K.; Ogura, H.; Li, S.-T.; Nagata, E.; Kuwajima,
- [119381] G.; Tanaka, K.; Itohara, S.; Mikoshiba, K.: Facilitation of NMDAR-independent
- [119382] LTP and spatial learning in mutant mice lacking ryanodine receptor
- [119383] type 3. Neuron 24: 701–713, 1999.
- [119384] 21147.

- [119385]Hakamata, Y.; Nakai, J.; Takeshima, H.; Imoto, K.: Primary structure
and distribution of a novel ryanodine receptor/calcium release channel
from rabbit brain. FEBS Lett. 312: 229–235, 1992.
21148.
- [119389]Leeb, T.; Brenig, B.: cDNA cloning and sequencing of the human
ryanodine receptor type 3 (RYR3) reveals a novel alternative splice
site in the RYR3 gene. FEBS Lett. 423: 367–370, 1998.
21149.
- [119393]Sorrentino, V.; Giannini, G.; Malzac, P.; Mattei, M. G.: Localization
of a novel ryanodine receptor gene (RYR3) to human chromosome 15q14–q15
by in situ hybridization. Genomics 18: 163–165, 1993.
21150.
- [119397]Takeshima, H.; Ikemoto, T.; Nishi, M.; Nishiyama, N.; Shimuta,
M.; Sugitani, Y.; Kuno, J.; Saito, I.; Saito, H.; Endo, M.; Iino,

- [119399]M.; Noda, T.: Generation and characterization of mutant mice lacking
- [119400]ryanodine receptor type 3. J. Biol. Chem. 271: 19649–19652, 1996.
- [119401]21151.
- [119402]Tiso, N.; Stephan, D. A.; Nava, A.; Bagattin, A.; Devaney, J. M.;
- [119403]Stanchi, F.; Larderet, G.; Brahmbhatt, B.; Brown, K.; Bauce, B.; Muriago,
- [119404]M.; Basso, C.; Thiene, G.; Danieli, G. A.; Rampazzo, A.: Identification
- [119405]of mutations in the cardiac ryanodine receptor gene in families affected
- [119406]with arrhythmogenic right ventricular cardiomyopathy type 2 (ARVD2). Hum.
- [119407]Molec. Genet. 10: 189–194, 2001.
- [119408]21152.
- [119409]Gallagher, P. G.; Forget, B. G.: Hematologically important mutations:
- [119410]spectrin and ankyrin variants in hereditary spherocytosis. Blood
- [119411]Cells Molec. Dis. 24: 539–543, 1998.

[119412]21153.

[119413]Gallagher, P. G.; Forget, B. G.: Hematologically important mutations:

[119414]spectrin variants in hereditary elliptocytosis and hereditary pyropoikilocytosis. Blood

[119415]Cells Molec. Dis. 22: 254–258, 1996.

[119416]21154.

[119417]Gallagher, P. G.; Forget, B. G.: Spectrin genes in health and disease. Seminars Hemat. 30: 4–20, 1993.

[119419]21155.

[119420]Gallagher, P. G.; Tse, W. T.; Coetzer, T.; Lecomte, M.–C.; Garbarz,

[119421]M.; Zarkowsky, H. S.; Baruchel, A.; Ballas, S. K.; Dhermy, D.; Palek,

[119422]J.; Forget, B. G.: A common type of the spectrin alpha-I 46–50a–kD

[119423]peptide abnormality in hereditary elliptocytosis and pyropoikilocytosis

[119424]is associated with a mutation distant from the proteolytic cleavage

[119425]site: evidence for the functional importance of the triple

helical

[119426]model of spectrin. J. Clin. Invest. 89: 892–898, 1992.

[119427]21156.

[119428]Gallagher, P. G.; Tse, W. T.; Marchesi, S. L.; Zarkowsky, H. S.;

[119429]Forget, B. G.: A defect in alpha-spectrin mRNA accumulation in hereditary

[119430]pyropoikilocytosis. Trans. Assoc. Am. Phys. 104: 32–39, 1991.

[119431]21157.

[119432]Garbarz, M.; Lecomte, M.-C.; Feo, C.; Devaux, I.; Picat, C.; Lefebvre,

[119433]C.; Galibert, F.; Gautero, H.; Bournier, O.; Galand, C.; Forget, B.

[119434]G.; Boivin, P.; Dhermy, D.: Hereditary pyropoikilocytosis and elliptocytosis

[119435]in a white French family with the spectrin alpha(I/74) variant related

[119436]to a CGT to CAT codon change (arg to his) at position 22 of the spectrin

[119437]alpha-I domain. Blood 75: 1691–1698, 1990.

[119438]21158.

[119439]Garbarz, M.; Lecomte, M. C.; Dhermy, D.; Feo, C.; Chave-
roche,

[119440]I.; Gautero, H.; Bournier, O.; Picat, C.; Goepp, A.; Boivin,
P.:

[119441]Double inheritance of an alpha I/65 spectrin variant in a
child with

[119442]homozygous elliptocytosis. Blood 67: 1661–1667, 1986.

[119443]21159.

[119444]Garbarz, M.; Lecomte, M. C.; Lefebvre, C.; Devaux, I.; Gal-
ibert,

[119445]F.; Boivin, P.; Forget, B. G.; Dhermy, D.: The DNAs from HE
and HPP

[119446]patients in a Caucasian French family with the spectrin al-
pha-I/74-kD

[119447]variant contain a CGT-to-CAT codon change (arg-to-his)
at position

[119448]22 of the alpha-I/80-kD domain. (Abstract) Blood 74:
181A, 1989.

[119449]21160.

[119450]Guetarni, D.; Roux, A.-F.; Alloisio, N.; Morle, F.;
Ducluzeau,

[119451]M. T.; Forget, B. G.; Colonna, P.; Delaunay, J.; Godet, J.:

Evidence

[119452] that expression of Sp-alpha-(I/65) hereditary elliptocytosis is compounded

[119453] by a genetic factor that is linked to the homologous alpha-spectrin

[119454] allele. Hum. Genet. 85: 627-630, 1990.

[119455] 21161.

[119456] Hassoun, H.; Coetzer, T. L.; Vassiliadis, J. N.; Sahr, K. E.;

[119457] Maalouf, G. J.; Saad, S. T. O.; Catanzariti, L.; Palek, J.: A novel

[119458] mobile element inserted in the alpha-spectrin gene: spectrin Dayton--a

[119459] truncated alpha-spectrin associated with hereditary elliptocytosis. J.

[119460] Clin. Invest. 94: 643-648, 1994.

[119461] 21162.

[119462] Huebner, K.; Palumbo, A. P.; Isobe, M.; Kozak, C. A.; Monaco,

[119463] S.; Rovera, G.; Croce, C. M.; Curtis, P. J.: The alpha-spectrin gene

[119464] is on chromosome 1 in mouse and man. Proc. Nat. Acad.

Sci. 82: 3790–3793,

[119465]1985.

[119466]21163.

[119467]Iarocci, T. A.; Wagner, G. M.; Mohandas, N.; Lane, P. A.;
Mentzer,

[119468]W. C.: Hereditary poikilocytic anemia associated with the
co-inheritance

[119469]of two alpha spectrin abnormalities. Blood 71:
1390–1396, 1988.

[119470]21164.

[119471]Knowles, W. J.; Bologna, M. L.; Chasis, J. A.; Marchesi, S.
L.;

[119472]Marchesi, V. T.: Common structural polymorphisms in hu-
man erythrocyte

[119473]spectrin. J. Clin. Invest. 73: 973–979, 1984.

[119474]21165.

[119475]Kotula, L.; Laury-Kleintop, L. D.; Showe, L.; Sahr, K.; Lin-
nenbach,

[119476]A. J.; Forget, B.; Curtis, P. J.: The exon–intron organization
of

[119477]the human erythrocyte alpha–spectrin gene. Genomics 9:

131-140,

[119478]1991.

[119479]21166.

[119480]Lambert, S.; Zail, S.: A new variant of the alpha-subunit of

[119481]spectrin in hereditary elliptocytosis. Blood 69: 473-478,

1987.

[119482]21167.

[119483]Lane, P. A.; Shew, R. L.; Iarocci, T. A.; Mohandas, N.; Hays,

[119484]T.; Mentzer, W. C.: Unique alpha-spectrin mutant in a kin-

dred with

[119485]common hereditary elliptocytosis. J. Clin. Invest. 79:

989-996,

[119486]1987.

[119487]21168.

[119488]Lawler, J.; Coetzer, T. L.; Mankad, V. N.; Moore, R. B.; Pr-

chal,

[119489]J. T.; Palek, J.: Spectrin-alpha(I/61): a new structural vari-

ant

[119490]of alpha-spectrin in a double-heterozygous form of

hereditary pyropoikilocytosis. Blood 72:

[119491]1412-1415, 1988.

[119492]21169.

[119493]Lawler, J.; Coetzer, T. L.; Palek, J.; Jacob, H. S.; Luban, N.

[119494]: Sp alpha(I/65): a new variant of the alpha subunit of spectrin in

[119495]hereditary elliptocytosis. Blood 66: 706–709, 1985.

[119496]21170.

[119497]Lawler, J.; Liu, S.-C.; Palek, J.; Prchal, J.: A molecular defect

[119498]in spectrin with a subset of patients with hereditary elliptocytosis:

[119499]alterations in the alpha-subunit domain involved in spectrin self-association. J.

[119500]Clin. Invest. 73: 1688–1695, 1984.

[119501]21171.

[119502]Lecomte, M.-C.; Dhermy, D.; Garbarz, M.; Feo, C.; Gautero, H.;

[119503]Bournier, O.; Picat, C.; Chaveroche, I.; Ester, A.; Galand, C.; Boivin,

[119504]P.: Pathologic and nonpathologic variants of the spectrin molecule

[119505]in two black families with hereditary elliptocytosis. Hum. Genet. 71:

[119506]351–357, 1985.

[119507]21172.

[119508]Lecomte, M.–C.; Dhermy, D.; Solis, C.; Ester, A.; Feo, C.;
Gautero,

[119509]H.; Bournier, O.; Boivin, P.: A new abnormal variant of
spectrin

[119510]in black patients with hereditary elliptocytosis. Blood 65:
1208–1217,

[119511]1985.

[119512]21173.

[119513]Lecomte, M.–C.; Garbarz, M.; Grandchamp, B.; Feo, C.;
Gautero,

[119514]H.; Devaux, I.; Bournier, O.; Galand, C.; d'Auriol, L.; Galib-
ert,

[119515]F.; Sahr, K. E.; Forget, B. G.; Boivin, P.; Dhermy, D.: Sp-
alpha(I/78):

[119516]a mutation of the alpha-I spectrin domain in a white kin-
dred with

[119517]HE and HPP phenotypes. Blood 74: 1126–1133, 1989.

[119518]21174.

[119519]Lecomte, M. C.; Dhermy, D.; Garbarz, M.; Feo, C.; Gautero,

H.;

[119520]Bournier, O.; Picat, C.; Chaveroche, I.; Galand, C.; Boivin, P.:

[119521]Hereditary pyropoikilocytosis and elliptocytosis in a Caucasian family:

[119522]transmission of the same molecular defect in spectrin through three

[119523]generations with different clinical expression. Hum. Genet. 77:

[119524]329-334, 1987.

[119525]21175.

[119526]Lecomte, M. C.; Dhermy, D.; Gautero, H.; Bournier, O.; Galand,

[119527]C.; Boivin, P.: L'elliptocytose hereditaire en Afrique de l'Ouest:

[119528]frequence et repartition des variants de la spectrine. C. R. Acad.

[119529]Sci. 306: 43-46, 1988.

[119530]21176.

[119531]Lecomte, M. C.; Feo, C.; Gautero, H.; Bournier, O.; Galand, C.;

[119532]Garbarz, M.; Boivin, P.; Dhermy, D.: Severe recessive poik-

ilocytic

[119533] anaemia with a new spectrin alpha chain variant. Brit. J.
Haemat. 74:

[119534] 497-507, 1990.

[119535] 21177.

[119536] Linnenbach, A. J.; Speicher, D. W.; Marchesi, V. T.; Forget,
B.

[119537] G.: Cloning of a portion of the chromosomal gene for hu-
man erythrocyte

[119538] alpha-spectrin by using a synthetic gene fragment. Proc.
Nat. Acad.

[119539] Sci. 83: 2397-2401, 1986.

[119540] 21178.

[119541] Bigler, R. D.; Bushkin, Y.; Chiorazzi, N.: S152 (CD27): a
modulating

[119542] disulfide-linked T cell activation antigen. J. Immun. 141:
21-28,

[119543] 1988.

[119544] 21179.

[119545] Hendriks, J.; Gravestien, L. A.; Tesselaar, K.; van Lier, R. A.

[119546] W.; Schumacher, T. N. M.; Borst, J.: CD27 is required for

generation

[119547]and long-term maintenance of T cell immunity. Nature
Immun. 1: 433–440,

[119548]2000.

[119549]21180.

[119550]Prasad, K. V. S.; Ao, Z.; Yoon, Y.; Wu, M. X.; Rizk, M.;
Jacquot,

[119551]S.; Schlossman, S. F.: CD27, a member of the tumor
necrosis factor

[119552]receptor family, induces apoptosis and binds to Siva, a
proapoptotic

[119553]protein. Proc. Nat. Acad. Sci. 94: 6346–6351, 1997.

[119554]21181.

[119555]Moody, D. B.; Ulrichs, T.; Muhlecker, W.; Young, D. C.;
Gurcha,

[119556]S. S.; Grant, E.; Rosat, J.-P.; Brenner, M. B.; Costello, C. E.;
Besra,

[119557]G. S.; Porcelli, S. A.: CD1c-mediated T-cell recognition of
isoprenoid

[119558]glycolipids in Mycobacterium tuberculosis infection. Na-
ture 404:

[119559]884–888, 2000.

[119560]21182.

[119561]Park, S.-H.; Bendelac, A.: CD1-restricted T-cell responses
and

[119562]microbial infection. Nature 406: 788-792, 2000.

[119563]21183.

[119564]Albertson, D. G.; Fishpool, R.; Sherrington, P.; Nacheva, E.;
Milstein,

[119565]C.: Sensitive and high resolution in situ hybridization to
human

[119566]chromosomes using biotin labelled probes: assignment of
the human

[119567]thymocyte CD1 antigen genes to chromosome 1. EMBO J.
7: 2801-2805,

[119568]1988.

[119569]21184.

[119570]Balk, S. P.; Bleicher, P. A.; Terhorst, C.: Isolation and char-
acterization

[119571]of a cDNA and gene coding for a fourth CD1 molecule.
Proc. Nat. Acad.

[119572]Sci. 86: 252-256, 1989.

[119573]21185.

[119574]Calabi, F.; Milstein, C.: A novel family of human major histocompatibility

[119575]complex-related genes not mapping to chromosome 6.
Nature 323: 540-543,

[119576]1986.

[119577]21186.

[119578]Nishimura, K.; Taketani, S.; Inokuchi, H.: Cloning of a human

[119579]cDNA for protoporphyrinogen oxidase by complementation in vivo of

[119580]a hemG mutant of Escherichia coli. J. Biol. Chem. 270:
8076-8080,

[119581]1995.

[119582]21187.

[119583]Roberts, A. G.; Whatley, S. D.; Daniels, J.; Holmans, P.;
Fenton,

[119584]J.; Owen, M. J.; Thompson, P.; Long, C.; Elder, G. H.: Partial
characterization

[119585]and assignment of the gene for protoporphyrinogen oxidase and variegate

[119586]porphyria to human chromosome 1q23. Hum. Molec.
Genet. 4: 2387-2390,

[119587]1995.

[119588]21188.

[119589]Taketani, S.; Inazawa, J.; Abe, T.; Furukawa, T.; Kohno, H.;
Tokunaga,

[119590]R.; Nishimura, K.; Inokuchi, H.: The human protopor-
phyrinogen oxidase

[119591]gene (PPOX): organization and location to chromosome 1.
Genomics 29:

[119592]698–703, 1995.

[119593]21189.

[119594]Warnich, L.; Kotze, M. J.; Groenewald, I. M.; Groenewald, J.
Z.;

[119595]van Brakel, M. G.; van Heerden, C. J.; de Villiers, J. N. P.;
van

[119596]de Ven, W. J. M.; Schoenmakers, E. F. P. M.; Taketani, S.;
Retief,

[119597]A. E.: Identification of three mutations and associated
haplotypes

[119598]in the protoporphyrinogen oxidase gene in South African
families with

[119599]variegate porphyria. Hum. Molec. Genet. 5: 981–984,
1996.

[119600]21190.

[119601]Albrecht, B.; Weber, K.; Pongs, O.: Characterization of a voltage-activated

[119602]K-channel gene cluster on human chromosome 12p13. Receptors Channels 3:

[119603]213-220, 1995.

[119604]21191.

[119605]Klocke, R.; Roberds, S. L.; Tamkun, M. M.; Gronemeier, M.; Augustin,

[119606]A.; Albrecht, B.; Pongs, O.; Jockusch, H.: Chromosomal mapping in

[119607]the mouse of eight K(+)-channel genes representing the four Shaker-like

[119608]subfamilies Shaker, Shab, Shaw, and Shal. Genomics 18: 568-574,

[119609]1993.

[119610]21192.

[119611]Curran, M. E.; Landes, G. M.; Keating, M. T.: Molecular cloning,

[119612]characterization, and genomic localization of a human potassium channel

[119613]gene. Genomics 12: 729–737, 1992.

[119614]21193.

[119615]Al-Chalabi, A.; Andersen, P. M.; Nilsson, P.; Chioza, B.;
Andersson,

[119616]J. L.; Russ, C.; Shaw, C. E.; Powell, J. F.; Leigh, P. N.: Dele-
tions

[119617]of the heavy neurofilament subunit tail in amyotrophic
lateral sclerosis. Hum.

[119618]Molec. Genet. 8: 157–164, 1999.

[119619]21194.

[119620]Collard, J.-F.; Cote, F.; Julien, J.-P.: Defective axonal
transport

[119621]in a transgenic mouse model of amyotrophic lateral scler-
osis. Nature 375:

[119622]61–64, 1995.

[119623]21195.

[119624]Figlewicz, D. A.; Krizus, A.; Martinoli, M. G.; Meiningner, V.;

[119625]Dib, M.; Rouleau, G. A.; Julien, J.-P.: Variants of the heavy
neurofilament

[119626]subunit are associated with the development of amy-
otrophic lateral

- [119627]sclerosis. Hum. Molec. Genet. 3: 1757–1761, 1994.
- [119628]21196.
- [119629]Lees, J. F.; Shneidman, P. S.; Skuntz, S. F.; Carden, M. J.;
Lazzarini,
[119630]R. A.: The structure and organization of the human heavy
neurofilament
[119631]subunit (NF-H) and the gene encoding it. EMBO J. 7:
1947–1955, 1988.
- [119632]21197.
- [119633]Mattei, M.-G.; Dautigny, A.; Pham-Dinh, D.; Passage, E.;
Mattei,
[119634]J.-F.; Jolles, P.: The gene encoding the large human neu-
rofilament
[119635]subunit (NF-H) maps to the q121–q131 region on human
chromosome 22. Hum.
[119636]Genet. 80: 293–295, 1988.
- [119637]21198.
- [119638]Rooke, K.; Figlewicz, D. A.; Han, F.; Rouleau, G. A.: Analy-
sis
[119639]of the KSP repeat of the neurofilament heavy subunit in
familial amyotrophic

[119640]lateral sclerosis. Neurology 46: 789–790, 1996.

[119641]21199.

[119642]Rouleau, G. A.; Merel, P.; Lutchman, M.; Sanson, M.; Zucman, J.;

[119643]Marineau, C.; Hoang-Xuan, K.; Demczuk, S.; Desmaze, C.; Plougastel,

[119644]B.; Pulst, S. M.; Lenoir, G.; Bijlsma, E.; Fashold, R.; Duman-ski,

[119645]J.; de Jong, P.; Parry, D.; Eldrige, R.; Aurias, A.; Delattre, O.;

[119646]Thomas, G.: Alteration in a new gene encoding a putative membrane-organizing

[119647]protein causes neuro-fibromatosis type 2. Nature 363: 515–521, 1993.

[119648]21200.

[119649]Tomkins, J.; Usher, P.; Slade, J. Y.; Ince, P. G.; Curtis, A.;

[119650]Bushby, K.; Shaw, P. J.: Novel insertion in the KSP region of the

[119651]neurofilament heavy gene in amyotrophic lateral sclerosis. Neuroreport 9:

[119652]3967–3970, 1998.

[119653]21201.

[119654]Vechio, J. D.; Bruijn, L. I.; Xu, Z.; Brown, R. H., Jr.; Cleveland,

[119655]D. W.: Sequence variants in human neurofilament proteins: absence

[119656]of linkage to familial amyotrophic lateral sclerosis. Ann. Neurol. 40:

[119657]603–610, 1996.

[119658]21202.

[119659]Watson, C. J.; Gaunt, L.; Evans, G.; Patel, K.; Harris, R.; Strachan,

[119660]T.: A disease-associated germline deletion maps the type 2 neurofibromatosis

[119661](NF2) gene between the Ewing sarcoma region and the leukaemia inhibitory

[119662]factor locus. Hum. Molec. Genet. 2: 701–704, 1993.

[119663]21203.

[119664]Umehara, F.; Kiwaki, T.; Yoshikawa, H.; Nishimura, T.; Nakagawa,

[119665]M.; Matsumoto, W.; Hashimoto, K.; Izumo, S.; Arimura, Y.; Arimura,

- [119666]K.; Kuriyama, M.; Osame, M.: Deletion in chromosome 17p11.2 including the peripheral myelin protein-22 (PMP-22) gene in hereditary neuropathy with liability to pressure palsies. J. Neurol. Sci. 133: 173-176, 1995.
- [119669]21204.
- [119670]21204.
- [119671]Barker, P. E.; Besmer, P.; Ruddle, F. H.: Human c-kit oncogene on human chromosome 4. (Abstract) Am. J. Hum. Genet. 37: A143, 1985.
- [119672]21205.
- [119673]21205.
- [119674]Beghini, A.; Larizza, L.; Cairoli, R.; Morra, E.: c-kit activating mutations and mast cell proliferation in human leukemia. (Letter) Blood 92: 701-702, 1998.
- [119675]701-702, 1998.
- [119676]21206.
- [119677]21206.
- [119678]Beghini, A.; Tibiletti, M.; Roversi, G.; Chiaravalli, A.; Serio, G.; Capella, C.; Larizza, L.: Germline mutation in the jux-

tamembrane

[119680]domain of the KIT gene in a family with gastrointestinal
stromal tumors

[119681]and urticaria pigmentosa. Cancer 92: 657–662, 2001.

[119682]21207.

[119683]Blume-Jensen, P.; Jiang, G.; Hyman, R.; Lee, K.-F.;
O’Gorman, S.;

[119684]Hunter, T.: Kit/stem cell factor receptor–induced activa-
tion of phosphatidylinositol

[119685]3–prime–kinase is essential for male fertility. Nature
Genet. 24:

[119686]157–162, 2000.

[119687]21208.

[119688]Bolognia, J. L.; Pawelek, J. M.: Biology of hypopigmenta-
tion. J.

[119689]Am. Acad. Derm. 19: 217–255, 1988.

[119690]21209.

[119691]Brannan, C. I.; Lyman, S. D.; Williams, D. E.; Eisenman, J.;;
Anderson,

[119692]D. M.; Cosman, D.; Bedell, M. A.; Jenkins, N. A.; Copeland,
N. G.

[119693]: Steel–Dickie mutation encodes a c–kit ligand lacking
transmembrane

[119694]and cytoplasmic domains. Proc. Nat. Acad. Sci. 88:
4671–4674, 1991.

[119695]21210.

[119696]Chabot, B.; Stephenson, D. A.; Chapman, V. M.; Besmer,
P.; Bernstein,

[119697]A.: The proto–oncogene c–kit encoding a transmembrane
tyrosine kinase

[119698]receptor maps to the mouse W locus. Nature 335: 88–89,
1988.

[119699]21211.

[119700]d'Auriol, L.; Mattei, M.–G.; Andre, C.; Galibert, F.: Local–
ization

[119701]of the human c–kit protooncogene on the q11–q12 region
of chromosome

[119702]4. Hum. Genet. 78: 374–376, 1988.

[119703]21212.

[119704]De Miguel, M. P.; Cheng, L.; Holland, E. C.; Federspiel, M.
J.;

[119705]Donovan, P. J.: Dissection of the c–Kit signaling pathway
in mouse

[119706]primordial germ cells by retroviral-mediated gene transfer. Proc.

[119707]Nat. Acad. Sci. 99: 10458–10463, 2002.

[119708]21213.

[119709]Dubreuil, P.; Forrester, L.; Rottapel, R.; Reedijk, M.; Fujita,

[119710]J.; Bernstein, A.: The c-fms gene complements the mitogenic defect

[119711]in mast cells derived from mutant W mice but not mi (microphthalmia)

[119712]mice. Proc. Nat. Acad. Sci. 88: 2341–2345, 1991.

[119713]21214.

[119714]el-Omar, M.; Davies, J.; Gupta, S.; Ross, H.; Thompson, R.: Leiomyosarcoma

[119715]in leiomyomatosis of the small intestine. Postgrad. Med. J. 70:

[119716]661–664, 1994.

[119717]21215.

[119718]Fleischman, R. A.: Human piebald trait resulting from a dominant

[119719]negative mutant allele of the c-kit membrane receptor gene. J. Clin.

[119720]Invest. 89: 1713–1717, 1992.

[119721]21216.

[119722]Fleischman, R. A.; Saltman, D. L.; Stastny, V.; Zneimer, S.:

[119723]Deletion of the c-kit protooncogene in the human developmental defect

[119724]piebald trait. Proc. Nat. Acad. Sci. 88: 10885–10889, 1991.

[119725]21217.

[119726]Fritsche-Polanz, R.; Jordan, J.-H.; Feix, A.; Sperr, W. R.; Sunder-Plassmann,

[119727]G.; Valent, P.; Fodinger, M.: Mutation analysis of C-KIT in patients

[119728]with myelodysplastic syndromes without mastocytosis and cases of systemic

[119729]mastocytosis. Brit. J. Haemat. 113: 357–364, 2001.

[119730]21218.

[119731]Geissler, E. N.; Ryan, M. A.; Housman, D. E.: The dominant-white

[119732]spotting (W) locus of the mouse encodes the c-kit protooncogene. Cell 55:

[119733]185–192, 1988.

[119734]21219.

[119735]Giebel, L. B.; Spritz, R. A.: Mutation of the KIT (mast/stem
[119736]cell growth factor receptor) protooncogene in human
piebaldism. Proc.

[119737]Nat. Acad. Sci. 88: 8696–8699, 1991.

[119738]21220.

[119739]Hirota, S.; Isozaki, K.; Moriyama, Y.; Hashimoto, K.;
Nishida,

[119740]T.; Ishiguro, S.; Kawano, K.; Hanada, M.; Kurata, A.;
Takeda, M.;

[119741]Tunio, G. M.; Matsuzawa, Y.; Kanakura, Y.; Shinomura, Y.;
Kitamura,

[119742]Y.: Gain-of-function mutations of c-kit in human gas-
trointestinal

[119743]stromal tumors. Science 279: 577–580, 1998.

[119744]21221.

[119745]Huizinga, J. D.; Thuneberg, L.; Kluppel, M.; Malysz, J.;
Mikkelsen,

[119746]H. B.; Bernstein, A.: W/kit gene required for interstitial
cells

[119747]of Cajal and for intestinal pacemaker activity. Nature 373:

347–349,

[119748]1995.

[119749]21222.

[119750]Furitsu, T.; Tsujimura, T.; Tono, T.; Ikeda, H.; Kitayama,
H.;

[119751]Koshimizu, U.; Sugahara, H.; Butterfield, J. H.; Ashman, L.
K.; Kanayama,

[119752]Y.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Identifica-
tion of

[119753]mutations in the coding sequence of the proto-oncogene
c-kit in a

[119754]human mast cell leukemia cell line causing ligand-
independent activation

[119755]of c-kit product. J. Clin. Invest. 92: 1736–1744, 1993.

[119756]21223.

[119757]Ingram, D. A.; Yang, F.-C.; Travers, J. B.; Wenning, M. J.;
Hiatt,

[119758]K.; New, S.; Hood, A.; Shannon, K.; Williams, D. A.; Clapp,
D. W.

[119759]: Genetic and biochemical evidence that haploinsufficiency
of the

[119760]Nf1 tumor suppressor gene modulates melanocyte and

mast cell fates

[119761]in vivo. J. Exp. Med. 191: 181–187, 2000.

[119762]21224.

[119763]Isozaki, K.; Terris, B.; Belghiti, J.; Schiffmann, S.; Hirota,

[119764]S.; Vanderwinden, J.–M.: Germline–activating mutation in
the kinase

[119765]domain of KIT gene in familial gastrointestinal stromal tu-
mors. Am.

[119766]J. Path. 157: 1581–1585, 2000.

[119767]21225.

[119768]Joensuu, H.; Roberts, P. J.; Sarlomo–Rikala, M.; Andersson,
L.

[119769]C.; Tervahartiala, P.; Tuveson, D.; Silberman, S. L.;
Capdeville,

[119770]R.; Dimitrijevic, S.; Druker, B.; Demetri, G. D.: Effect of the
tyrosine

[119771]kinase inhibitor STI571 in a patient with a metastatic gas-
trointestinal

[119772]stromal tumor. New Eng. J. Med. 344: 1052–1056, 2001.

[119773]21226.

[119774]Johansson Moller, M.; Chaudhary, R.; Hellmen, E.; Hoy–

heim, B.;

[119775]Chowdhary, B.; Andersson, L.: Pigs with the dominant white coat color

[119776]phenotype carry a duplication of the KIT gene encoding the mast/stem

[119777]cell growth factor receptor. Mammalian Genome 7: 822–830, 1996.

[119778]21227.

[119779]Lasota, J.; Jasinski, M.; Sarlomo-Rikala, M.; Miettinen, M.:

[119780]Mutations in exon 11 of c-kit occur preferentially in malignant versus

[119781]benign gastrointestinal stromal tumors and do not occur in leiomyomas

[119782]or leiomyosarcomas. Am. J. Path. 154: 53–60, 1999.

[119783]21228.

[119784]Longley, B. J.; Tyrrell, L.; Lu, S.-Z.; Ma, Y.-S.; Langley, K.;

[119785]Ding, T.; Duffy, T.; Jacobs, P.; Tang, L. H.; Modlin, I.: Somatic

[119786]c-KIT activating mutation in urticaria pigmentosa and aggressive mastocytosis:

[119787]establishment of clonality in a human mast cell neoplasm.

Nature

[119788]Genet. 12: 312–314, 1996.

[119789]21229.

[119790]Longley, B. J., Jr.; Metcalfe, D. D.; Tharp, M.; Wang, X.;
Tyrrell,

[119791]L.; Lu, S.-Z.; Heitjan, D.; Ma, Y.: Activating and dominant
inactivating

[119792]c-KIT catalytic domain mutations in distinct clinical forms
of human

[119793]mastocytosis. Proc. Nat. Acad. Sci. 96: 1609–1614, 1999.

[119794]21230.

[119795]Marklund, S.; Kijas, J.; Rodriguez-Martinez, H.;
Ronnstrand, L.;

[119796]Funa, K.; Moller, M.; Lange, D.; Edfors-Lilja, I.; Andersson,
L.:

[119797]Molecular basis for the dominant white phenotype in the
domestic pig. Genome

[119798]Res. 8: 826–833, 1998.

[119799]21231.

[119800]Ingalls, A. M.; Dickie, M. M.; Snell, G. D.: Obese, a new
mutation

- [119801]in the house mouse. J. Hered. 41: 317–318, 1950.
- [119802]21232.
- [119803]Ioffe, E.; Moon, B.; Connolly, E.; Friedman, J. M.: Abnormal
- [119804]regulation of the leptin gene in the pathogenesis of obesity. Proc.
- [119805]Nat. Acad. Sci. 95: 11852–11857, 1998.
- [119806]21233.
- [119807]Isse, N.; Ogawa, Y.; Tamura, N.; Masuzaki, H.; Mori, K.;
- Okazaki,
- [119808]T.; Satoh, N.; Shigemoto, M.; Yoshimasa, Y.; Nishi, S.;
- Hosoda, K.;
- [119809]Inazawa, J.; Nakao, K.: Structural organization and chromosomal assignment
- [119810]of the human obese gene. J. Biol. Chem. 270:
- 27728–27733, 1995.
- [119811]21234.
- [119812]Jaquet, D.; Leger, J.; Levy-Marchal, C.; Oury, J. F.; Czerni-
- chow,
- [119813]P.: Ontogeny of leptin in human fetuses and newborns: effect of intrauterine
- [119814]growth retardation on serum leptin concentrations. J. Clin.

Endocr.

[119815]Metab. 83: 1243–1246, 1998.

[119816]21235.

[119817]Karvonen, M. K.; Pesonen, U.; Heinonen, P.; Laakso, M.;
Rissanen,

[119818]A.; Naukkarinen, H.; Valve, R.; Uusitupa, M. I. J.; Koulu, M.:
Identification

[119819]of new sequence variants in the leptin gene. J. Clin. En-
docr. Metab. 83:

[119820]3239–3242, 1998.

[119821]21236.

[119822]Kawai, K.; Sugimoto, K.; Nakashima, K.; Miura, H.; Ni-
nomiya, Y.

[119823]: Leptin as a modulator of sweet taste sensitivities in mice.
Proc.

[119824]Nat. Acad. Sci. 97: 11044–11049, 2000.

[119825]21237.

[119826]Kennedy, A.; Gettys, T. W.; Watson, P.; Wallace, P.; Gan-
away,

[119827]E.; Pan, Q.; Garvey, W. T.: The metabolic significance of
leptin

- [119828]in humans: Gender-based differences in relationship to adiposity,
- [119829]insulin sensitivity, and energy expenditure. J. Clin. Endocr. Metab. 82: 1293–1300, 1997.
- [119831]21238.
- [119832]Koistinen, H. A.; Koivisto, V. A.; Andersson, S.; Karonen, S.-L.;
- [119833]Kontula, K.; Oksanen, L.; Teramo, K. A.: Leptin concentration in
- [119834]cord blood correlates with intrauterine growth. J. Clin. Endocr. Metab. 82: 3328–3330, 1997.
- [119835]21239.
- [119837]Li, W.-D.; Reed, D. R.; Lee, J. H.; Xu, W.; Kilker, R. L.; So-
- [119838]dam, B. R.; Price, R. A.: Sequence variants in the 5-prime flanking region
- [119839]of the leptin gene are associated with obesity in women. Ann. Hum. Genet. 63: 227–234, 1999.

[119841]21240.

[119842]Lonnqvist, F.; Arner, P.; Nordfors, L.; Schalling, M.: Over-expression

[119843]of the obese (ob) gene in adipose tissue of human obese subjects. Nature

[119844]Med. 1: 950-953, 1995.

[119845]21241.

[119846]Lord, G. M.; Matarese, G.; Howard, J. K.; Baker, R. J.; Bloom,

[119847]S. R.; Lechler, R. I.: Leptin modulates the T-cell immune response

[119848]and reverses starvation-induced immunosuppression. Nature 394: 897-901,

[119849]1998.

[119850]21242.

[119851]Lucantoni, R.; Ponti, E.; Berselli, M. E.; Savia, G.; Minocci,

[119852]A.; Calo, G.; de Medici, C.; Liuzzi, A.; Di Blasio, A. M.: The A19G

[119853]polymorphism in the 5-prime untranslated region of the human obese

[119854]gene does not affect leptin levels in severely obese pa-

tients. J.

[119855]Clin. Endocr. Metab. 85: 3589–3591, 2000.

[119856]21243.

[119857]Maffei, M.; Fei, H.; Lee, G.–H.; Dani, C.; Leroy, P.; Zhang, Y.;

[119858]Proenca, R.; Negrel, R.; Ailhaud, G.; Friedman, J. M.: Increased

[119859]expression in adipocytes of ob RNA in mice with lesions of the hypothalamus

[119860]and with mutations at the db locus. Proc. Nat. Acad. Sci. 92: 6957–6960,

[119861]1995.

[119862]21244.

[119863]Mammes, O.; Betoulle, D.; Aubert, R.; Giraud, V.; Tuzet, S.; Petiet,

[119864]A.; Colas–Linhardt, N.; Fumeron, F.: Novel polymorphisms in the 5–prime

[119865]region of the LEP gene: association with leptin levels and response

[119866]to low–calorie diet in human obesity. Diabetes 47: 487–489, 1998.

[119867]21245.

- [119868]Mammes, O.; Betoulle, D.; Aubert, R.; Herbeth, B.; Siest, G.;
- [119869]Fumeron, F.: Association of the G-2548A polymorphism in the 5-prime
- [119870]region of the LEP gene with overweight. Ann. Hum. Genet. 64: 391-394,
- [119871]2000.
- [119872]21246.
- [119873]Mancuso, P.; Gottschalk, A.; Phare, S. M.; Peters-Golden, M.;
- [119874]Lukacs, N. W.; Huffnagle, G. B.: Leptin-deficient mice exhibit impaired
- [119875]host defense in Gram-negative pneumonia. J. Immun. 168: 4018-4024,
- [119876]2002.
- [119877]21247.
- [119878]Mantzoros, C. S.; Flier, J. S.; Rogol, A. D.: A longitudinal
- [119879]assessment of hormonal and physical alterations during normal puberty
- [119880]in boys. v. rising leptin levels may signal the onset of puberty. J.

- [119881]Clin. Endocr. Metab. 82: 1066–1070, 1997.
- [119882]21248.
- [119883]Masuzaki, H.; Ogawa, Y.; Isse, N.; Satoh, N.; Okazaki, T.;
Shigemoto,
- [119884]M.; Mori, K.; Tamura, N.; Hosoda, K.; Yoshimasa, Y.;
Jingami, H.;
- [119885]Kawada, T.; Nakao, K.: Human obese gene expression:
adipocyte-specific
- [119886]expression and regional differences in the adipose tissue.
Diabetes 44:
- [119887]855–858, 1995.
- [119888]21249.
- [119889]Masuzaki, H.; Ogawa, Y.; Sagawa, N.; Hosoda, K.; Mat-
sumoto, T.;
- [119890]Mise, H.; Nishimura, H.; Yoshimasa, Y.; Tanaka, I.; Mori,
T.; Nakao,
- [119891]K.: Nonadipose tissue production of leptin: leptin as a
novel placenta-derived
- [119892]hormone in humans. Nature Med. 3: 1029–1033, 1997.
- [119893]21250.
- [119894]Matkovic, V.; Ilich, J. Z.; Badenhop, N. E.; Skugor, M.;

Clairmont,

[119895]A.; Klisovic, D.; Landoll, J. D.: Gain in body fat is inversely related

[119896]to the nocturnal rise in serum leptin level in young females. J.

[119897]Clin. Endocr. Metab. 82: 1368–1372, 1997.

[119898]21251.

[119899]Matkovic, V.; Ilich, J. Z.; Skugor, M.; Badenhop, N. E.; Goel,

[119900]P.; Clairmont, A.; Klisovic, D.; Nahhas, R. W.; Landoll, J. D.:
Leptin

[119901]is inversely related to age at menarche in human females.
J. Clin.

[119902]Endocr. Metab. 82: 3239–3245, 1997.

[119903]21252.

[119904]Matsuda, J.; Yokota, I.; Iida, M.; Murakami, T.; Naito, E.;
Ito,

[119905]M.; Shima, K.; Kuroda, Y.: Serum leptin concentration in
cord blood:

[119906]relationship to birth weight and gender. J. Clin. Endocr.
Metab. 82:

[119907]1642–1644, 1997.

[119908]21253.

[119909]Miller, S. G.; De Vos, P.; Guerre-Millo, M.; Wong, K.; Hermann,

[119910]T.; Staels, B.; Briggs, M. R.; Auwerx, J.: The adipocyte specific

[119911]transcription factor C/EBP- α modulates human ob gene expression. Proc.

[119912]Nat. Acad. Sci. 93: 5507-5511, 1996.

[119913]21254.

[119914]Minokoshi, Y.; Kim, Y.-B.; Peroni, O. D.; Fryer, L. G. D.; Muller,

[119915]C.; Carling, D.; Kahn, B. B.: Leptin stimulates fatty-acid oxidation

[119916]by activating AMP-activated protein kinase. Nature 415: 339-343,

[119917]2002.

[119918]21255.

[119919]Moffett, S.; Martinson, J.; Shriver, M. D.; Deka, R.; McGarvey,

[119920]S. T.; Barrantes, R.; Ferrell, R. E.: Genetic diversity and evolution

[119921]of the human leptin locus tetranucleotide repeat. Hum.
Genet. 110:

[119922]412–417, 2002.

[119923]21256.

[119924]Montague, C. T.; Farooqi, I. S.; Whitehead, J. P.; Soos, M.
A.;

[119925]Rau, H.; Wareham, N. J.; Sewter, C. P.; Digby, J. E.; Mo-
hammed, S.

[119926]N.; Hurst, J. A.; Cheetham, C. H.; Earley, A. R.; Barnett, A.
H.;

[119927]Prins, J. B.; O'Rahilly, S.: Congenital leptin deficiency is as-
sociated

[119928]with severe early-onset obesity in humans. Nature 387:
903–908,

[119929]1997.

[119930]21257.

[119931]Muzzin, P.; Eisensmith, R. C.; Copeland, K. C.; Woo, S. L.
C.

[119932]: Correction of obesity and diabetes in genetically obese
mice by

[119933]leptin gene therapy. Proc. Nat. Acad. Sci. 93:
14804–14808, 1996.

[119934]21258.

[119935]Kiefer, M. C.; Tucker, J. E.; Joh, R.; Landsberg, K. E.; Saltman,

[119936]D.; Barr, P. J.: Identification of a second human subtilisin-like

[119937]protease gene in the fes/fps region of chromosome 15. DNA Cell Biol. 10:

[119938]757-769, 1991.

[119939]21259.

[119940]Amiel, J.; Audollent, S.; Joly, D.; Dureau, P.; Salomon, R.; Tellier,

[119941]A.-L.; Auge, J.; Bouissou, F.; Antignac, C.; Gubler, M.-C.; Eccles,

[119942]M. R.; Munnich, A.; Vekemans, M.; Lyonnet, S.; Attie-Bitach, T.:

[119943]PAX2 mutations in renal-coloboma syndrome: mutational hotspot and

[119944]germline mosaicism. Europ. J. Hum. Genet. 8: 820-826, 2000.

[119945]21260.

[119946]Dehbi, M.; Ghahremani, M.; Lechner, M.; Dressler, G.; Pelletier,

[119947]J.: The paired-box transcription factor, PAX2, positively modulates

[119948]expression of the Wilms' tumor suppressor gene. *Oncogene* 13: 447-453,

[119949]1996.

[119950]21261.

[119951]Devriendt, K.; Matthijs, G.; Van Damme, B.; Van Caesbroeck, D.;

[119952]Eccles, M.; Vanrenterghem, Y.; Fryns, J.-P.; Leys, A.: Mis-sense mutation

[119953]and hexanucleotide duplication in the PAX2 gene in two unrelated families

[119954]with renal-coloboma syndrome (MIM 120330). *Hum. Genet.* 103: 149-153,

[119955]1998.

[119956]21262.

[119957]Dressler, G. R.; Deutsch, U.; Chowdhury, K.; Nornes, H. O.; Gruss,

[119958]P.: Pax2, a new murine paired-box-containing gene and its expression

[119959]in the developing excretory system. *Development* 109: 787-795, 1990.

[119960]21263.

[119961]Ford, B.; Rupps, R.; Lirenman, D.; Van Allen, M. I.; Far-
quharson,

[119962]D.; Lyons, C.; Friedman, J. M.: Renal-coloboma syndrome:
prenatal

[119963]detection and clinical spectrum in a large family. Am. J.
Med. Genet. 99:

[119964]137-141, 2001.

[119965]21264.

[119966]Keller, S. A.; Jones, J. M.; Boyle, A.; Barrow, L. L.; Killen,

[119967]P. D.; Green, D. G.; Kapousta, N. V.; Hitchcock, P. F.;
Swank, R.

[119968]T.; Meisler, M. H.: Kidney and retinal defects (Krd), a
transgene-induced

[119969]mutation with a deletion of mouse chromosome 19 that
includes the

[119970]Pax2 locus. Genomics 23: 309-320, 1994.

[119971]21265.

[119972]Narahara, K.; Baker, E.; Ito, S.; Yokoyama, Y.; Yu, S.; He-
witt,

[119973]D.; Sutherland, G. R.; Eccles, M. R.; Richards, R. I.: Locali-

sation

[119974] of a 10q breakpoint within the PAX2 gene in a patient with
a de novo

[119975] t(10;13) translocation and optic nerve coloboma–renal
disease. J.

[119976] Med. Genet. 34: 213–216, 1997.

[119977] 21266.

[119978] Pilz, A. J.; Povey, S.; Gruss, P.; Abbott, C. M.: Mapping of
the

[119979] human homologs of the murine paired–box–containing
genes. Mammalian

[119980] Genome 4: 78–82, 1993.

[119981] 21267.

[119982] Favor, J.; Sandulache, R.; Neuhauser–Klaus, A.; Pretsch, W.;
Chatterjee,

[119983] B.; Senft, E.; Wurst, W.; Blanquet, V.; Grimes, P.; Sporle, R.;
Schughart,

[119984] K.: The mouse Pax2(1Neu) mutation is identical to a hu–
man PAX2 mutation

[119985] in a family with renal–coloboma syndrome and results in
developmental

[119986] defects of the brain, ear, eye, and kidney. Proc. Nat. Acad.

Sci. 93:

[119987]13870-13875, 1996.

[119988]21268.

[119989]Seawright, A.; Fletcher, J. M.; Fantes, J. A.; Morrison, H.;
Porteous,

[119990]D. J.; Li, S. S.-L.; Hastie, N. D.; Van Heyningen, V.: Analysis
of

[119991]WAGR deletions and related translocations with gene-
specific DNA probes,

[119992]using FACS-selected cell hybrids. Somat. Cell Molec.
Genet. 14:

[119993]21-30, 1988.

[119994]21269.

[119995]Huerre, C.; Despoisse, S.; Gilgenkrantz, S.; Lenoir, G. M.;
Junien,

[119996]C.: c-Ha-ras1 is not deleted in aniridia-Wilms' tumour as-
sociation. Nature 305:

[119997]638-641, 1983.

[119998]21270.

[119999]Liou, G. I.; Ma, D.-P.; Yang, Y.-W.; Geng, L.; Zhu, C.;
Baehr,

[120000]W.: Human interstitial retinoid-binding protein: gene structure and

[120001]primary sequence. J. Biol. Chem. 264: 8200–8206, 1989.

[120002]21271.

[120003]Nakamura, Y.; Lathrop, M.; Bragg, T.; Leppert, M.; O'Connell,

[120004]P.; Jones, C.; Lalouel, J. M.; White, R.: An extended genetic linkage

[120005]map of markers for human chromosome 10. Genomics 3: 389–392, 1988.

[120006]21272.

[120007]Heitman, J.; Agre, P.: A new face of the rhesus antigen. Nature

[120008]Genet. 26: 258–259, 2000.

[120009]21273.

[120010]Marini, A.–M.; Matassi, G.; Raynal, V.; Andre, B.; Cartron, J.–P.;

[120011]Cherif–Zahar, B.: The human Rhesus-associated RhAG protein and a

[120012]kidney homologue promote ammonium transport in yeast. Nature Genet. 26:

[120013]341–344, 2000.

[120014]21274.

[120015]Sap, J.; Munoz, A.; Damm, K.; Goldberg, Y.; Ghysdael, J.;
Leutz,

[120016]A.; Beug, H.; Vennstrom, B.: The c-erb-A protein is a
high-affinity

[120017]receptor for thyroid hormone. Nature 324: 635–640,
1986.

[120018]21275.

[120019]Carrere, N.; Leblanc, R. M.; Begueret, J.; Blouin, P.; Chey-
rou,

[120020]A.: A new mutation of exon 5 of the P53 gene in breast
cancer. Hum.

[120021]Molec. Genet. 2: 1075, 1993.

[120022]21276.

[120023]Walker, R. L.; Anziano, P.; Meltzer, P. S.: A PAC containing
the

[120024]human mitochondrial DNA polymerase gamma gene
(POLG) maps to chromosome

[120025]15q25. Genomics 40: 376–378, 1997.

[120026]21277.

- [120027]Zullo, S. J.; Butler, L.; Zahorchak, R. J.; Macville, M.; Wilkes,
[120028]C.; Merril, C. R.: Localization by fluorescence in situ hybridization
(FISH) of human mitochondrial polymerase gamma (POLG)
[120029]to human chromosome
[120030]band 15q24-q26, and of mouse mitochondrial polymerase gamma (Polg)
[120031]to mouse chromosome band 7E, with confirmation by direct sequence
[120032]analysis of bacterial artificial chromosomes (BACs). Cytogenet. Cell
[120033]Genet. 78: 281-284, 1997.
[120034]21278.
[120035]Bayliss, W.; Starling, E. H.: The mechanism of pancreatic secretion. J.
[120036]Physiol. (London) 28: 325-353, 1902.
[120037]21279.
[120038]Kopin, A. S.; Wheeler, M. B.; Leiter, A. B.: Secretin: structure
[120039]of the precursor and tissue distribution of the mRNA. Proc. Nat.

[120040]Acad. Sci. 87: 2299–2303, 1990.

[120041]21280.

[120042]Marinelli, R. A.; Pham, L.; Agre, P.; LaRusso, N. F.: Secretin

[120043]promotes osmotic water transport in rat cholangiocytes by
increasing

[120044]aquaporin–1 water channels in plasma membrane. J. Biol.
Chem. 272:

[120045]12984–12988, 1997.

[120046]21281.

[120047]Peel, A. L.; Rao, R. V.; Cottrell, B. A.; Hayden, M. R.;
Ellerby,

[120048]L. M.; Bredesen, D. E.: Double–stranded RNA–dependent
protein kinase,

[120049]PKR, binds preferentially to Huntington's disease (HD)
transcripts

[120050]and is activated in HD tissue. Hum. Molec. Genet. 10:
1531–1538,

[120051]2001.

[120052]21282.

[120053]Squire, J.; Meurs, E. F.; Chong, K. L.; McMillan, N. A. J.;
Hovanessian,

- [120054]A. G.; Williams, B. R. G.: Localization of the human interferon-induced,
[120055]ds-RNA activated p68 kinase gene (PRKR) to chromosome
2p21-p22. Genomics 16:
[120056]768-770, 1993.
[120057]21283.
- [120058]Taylor, D. R.; Shi, S. T.; Romano, P. R.; Barber, G. N.; Lal,
M.
[120059]M. C.: Inhibition of the interferon-inducible protein kinase
PKR
[120060]by HCV E2 protein. Science 285: 107-110, 1999.
[120061]21284.
- [120062]Brott, B. K.; Alessandrini, A.; Largaespada, D. A.;
Copeland, N.
[120063]G.; Jenkins, N. A.; Crews, C. M.; Erikson, R. L.: MEK2 is a
kinase
[120064]related to MEK1 and is differentially expressed in murine
tissues. Cell
[120065]Growth Differ. 4: 921-929, 1993.
[120066]21285.
- [120067]Crews, C. M.; Alessandrini, A.; Erikson, R. L.: The primary

structure

[120068]of MEK, a protein kinase that phosphorylates the ERK gene product. Science 258:

[120069]478–480, 1992.

[120070]21286.

[120071]Favata, M. F.; Horiuchi, K. Y.; Manos, E. J.; Daulerio, A. J.;

[120072]Stradley, D. A.; Feeser, W. S.; Van Dyk, D. E.; Pitts, W. J.; Earl,

[120073]R. A.; Hobbs, F.; Copeland, R. A.; Magolda, R. L.; Scherle, P. A.;

[120074]Trzaskos, J. M.: Identification of a novel inhibitor of mitogen-activated

[120075]protein kinase kinase. J. Biol. Chem. 273: 18623–18632, 1998.

[120076]21287.

[120077]Meloche, S.; Gopalbhai, K.; Beatty, B. G.; Scherer, S. W.; Pellerin,

[120078]J.: Chromosome mapping of the human genes encoding the MAP kinase

[120079]kinase MEK1 (MAP2K1) to 15q21 and MEK2 (MAP2K2) to 7q32. Cytogenet.

[120080]Cell Genet. 88: 249–252, 2000.

[120081]21288.

[120082]Orth, K.; Palmer, L. E.; Bao, Z. Q.; Stewart, S.; Rudolph, A. E.;

[120083]Bliska, J. B.; Dixon, J. E.: Inhibition of the mitogen-activated

[120084]protein kinase kinase superfamily by a Yersinia effector. Science 285:

[120085]1920–1923, 1999.

[120086]21289.

[120087]Perry, R. L. S.; Parker, M. H.; Rudnicki, M. A.: Activated MEK1

[120088]binds the nuclear MyoD transcriptional complex to repress transactivation. Molec.

[120089]Cell 8: 291–301, 2001.

[120090]21290.

[120091]Pleschka, S.; Wolff, T.; Ehrhardt, C.; Hobom, G.; Planz, O.; Rapp,

[120092]U. R.; Ludwig, S.: Influenza virus propagation is impaired by inhibition

[120093]of the Raf/MEK/ERK signalling cascade. Nature Cell Biol. 3: 301–305,

[120094]2001.

[120095]21291.

[120096]Rampoldi, L.; Zimbello, R.; Bortoluzzi, S.; Tiso, N.; Valle, G.;

[120097]Lanfranchi, G.; Danieli, G. A.: Chromosomal localization of four

[120098]MAPK signaling cascade genes: MEK1, MEK3, MEK4 and MEKK5. Cytogenet.

[120099]Cell Genet. 78: 301–303, 1997.

[120100]21292.

[120101]Ryan, K. M.; Ernst, M. K.; Rice, N. R.; Vousden, K. H.: Role of

[120102]NF- κ B in p53-mediated programmed cell death. Nature 404: 892–897,

[120103]2000.

[120104]21293.

[120105]Sebolt-Leopold, J. S.; Dudley, D. T.; Herrera, R.; Van Becelaere,

[120106]K.; Wiland, A.; Gowan, R. C.; Tecle, H.; Barrett, S. D.; Bridges,

[120107]A.; Przybranowski, S.; Leopold, W. R.; Saltiel, A. R.: Block-

ade of

[120108] the MAP kinase pathway suppresses growth of colon tumors in vivo. *Nature*

[120109] *Med.* 5: 810–816, 1999.

[120110] 21294.

[120111] Seger, R.; Krebs, E. G.: The MAPK signaling cascade. *FASEB J.* 9:

[120112] 726–735, 1995.

[120113] 21295.

[120114] Seger, R.; Seger, D.; Lozeman, F. J.; Ahn, N. G.; Graves, L. M.;

[120115] Campbell, J. S.; Ericsson, L.; Harrylock, M.; Jensen, A. M.; Krebs,

[120116] E. G.: Human T-cell mitogen-activated protein kinase kinases are

[120117] related to yeast signal transduction kinases. *J. Biol. Chem.* 267:

[120118] 25628–25631, 1992.

[120119] 21296.

[120120] Zheng, C. F.; Guan, K. L.: Cloning and characterization of two

[120121]distinct human extracellular signal-regulated kinase activator kinases,

[120122]MEK1 and MEK2. J. Biol. Chem. 268: 11435-11439, 1993.

[120123]21297.

[120124]White, P. S.; Maris, J. M.; Beltinger, C.; Sulman, E.; Marshall,

[120125]H. N.; Fujimori, M.; Kaufman, B. A.; Biegel, J. A.; Allen, C.; Hilliard,

[120126]C.; Valentine, M. B.; Look, A. T.; Enomoto, H.; Sakiyama, S.; Brodeur,

[120127]G. M.: A region of consistent deletion in neuroblastoma maps within

[120128]human chromosome 1p36.2-36.3. Proc. Nat. Acad. Sci. 92: 5520-5524,

[120129]1995.

[120130]21298.

[120131]Deng, Z.; Morse, J. H.; Slager, S. L.; Cuervo, N.; Moore, K. J.;

[120132]Venetos, G.; Kalachikov, S.; Cayanis, E.; Fischer, S. G.; Barst, R.

[120133]J.; Hodge, S. E.; Knowles, J. A.: Familial primary pulmonary hypertension

[120134](gene PPH1) is caused by mutations in the bone morpho-
genetic protein

[120135]receptor-II gene. Am. J. Hum. Genet. 67: 737-744, 2000.

[120136]21299.

[120137]International PPH Consortium; Lane, K. B.; Machado, R. D.;
Pauciulo,

[120138]M. W.; Thomson, J. R.; Phillips, J. A., III; Loyd, J. E.; Nichols,

[120139]W. C.; Trembath, R. C.: Heterozygous germline mutations
in BMPR2,

[120140]encoding a TGF-beta receptor, cause familial primary pul-
monary hypertension. Nature

[120141]Genet. 26: 81-84, 2000.

[120142]21300.

[120143]Amin, R. S.; Wert, S. E.; Baughman, R. P.; Tomashevski, J.
F.,

[120144]Jr.; Nogee, L. M.; Brody, A. S.; Hull, W. M.; Whitsett, J. A.:
Surfactant

[120145]protein deficiency in familial interstitial lung disease. J.
Pediat. 139:

[120146]85-92, 2001.

[120147]21301.

- [120148]Glasser, S. W.; Burhans, M. S.; Korfhagen, T. R.; Na, C.-L.;
Sly,
[120149]P. D.; Ross, G. F.; Ikegami, M.; Whitsett, J. A.: Altered sta-
bility
[120150]of pulmonary surfactant in SP-C-deficient mice. Proc. Nat.
Acad.
[120151]Sci. 98: 6366-6371, 2001.
[120152]21302.
[120153]Hussain, S. P.; Raja, K.; Amstad, P. A.; Sawyer, M.; Trudel,
L.
[120154]J.; Wogan, G. N.; Hofseth, L. J.; Shields, P. G.; Billiar, T. R.;
[120155]Trautwein, C.; Hohler, T.; Galle, P. R.; Phillips, D. H.;
Markin,
[120156]R.; Marrogi, A. J.; Harris, C. C.: Increased p53 mutation
load in
[120157]nontumorous human liver of Wilson disease and
hemochromatosis: oxyradical
[120158]overload diseases. Proc. Nat. Acad. Sci. 97: 12770-12775,
2000.
[120159]21303.
[120160]Iavarone, A.; Matthay, K. K.; Steinkirchner, T. M.; Israel, M.

[120161]A.: Germ-line and somatic p53 gene mutations in multifocal osteogenic

[120162]sarcoma. Proc. Nat. Acad. Sci. 89: 4207–4209, 1992.

[120163]21304.

[120164]Iggo, R.; Gatter, K.; Bartek, J.; Lane, D.; Harris, A. L.: Increased

[120165]expression of mutant forms of p53 oncogene in primary lung cancer. Lancet 335:

[120166]675–679, 1990.

[120167]21305.

[120168]Isobe, M.; Emanuel, B. S.; Givol, D.; Oren, M.; Croce, C. M.:

[120169]Localization of gene for human p53 tumour antigen to band 17p13. Nature 320:

[120170]84–85, 1986.

[120171]21306.

[120172]Jeffrey, P. D.; Gorina, S.; Pavletich, N. P.: Crystal structure

[120173]of the tetramerization domain of the p53 tumor suppressor at 1.7 angstroms. Science 267:

[120174]1498–1502, 1995.

[120175]21307.

[120176]Jenkins, J. R.; Rudge, K.; Chumakov, P.; Currie, G. A.: The cellular

[120177]oncogene p53 can be activated by mutagenesis. Nature 317: 816–818,

[120178]1985.

[120179]21308.

[120180]Jimenez, G. S.; Nister, M.; Stommel, J. M.; Beeche, M.; Bar-carse,

[120181]E. A.; Zhang, X.-Q.; O'Gorman, S.; Wahl, G. M.: A transac-tivation-deficient

[120182]mouse model provides insights into Trp53 regulation and function. Nature

[120183]Genet. 26: 37–43, 2000.

[120184]21309.

[120185]Jonkers, J.; Meuwissen, R.; van der Gulden, H.; Peterse, H.; van

[120186]der Valk, M.; Berns, A.: Synergistic tumor suppressor ac-tivity of

[120187]BRCA2 and p53 in a conditional mouse model for breast cancer. Nature

[120188]Genet. 29: 418–425, 2001.

[120189]21310.

[120190]Kaelbling, M.; Burk, R. D.; Atkin, N. B.; Johnson, A. B.;
Klinger,

[120191]H. P.: Loss of heterozygosity on chromosome 17p and
mutant p53 in

[120192]HPV-negative cervical carcinomas. Lancet 340: 140–142,
1992.

[120193]21311.

[120194]Keller, D. M.; Zeng, X.; Wang, Y.; Zhang, Q. H.; Kapoor, M.;
Shu,

[120195]H.; Goodman, R.; Lozano, G.; Zhao, Y.; Lu, H.: A DNA
damage-induced

[120196]p53 serine 392 kinase complex contains CK2, hSpt16, and
SSRP1. Molec.

[120197]Cell 283–292, 2001.

[120198]21312.

[120199]Kemp, C. J.; Wheldon, T.; Balmain, A.: p53-deficient mice
are

[120200]extremely susceptible to radiation-induced tumorigenesis.
Nature

[120201]Genet. 8: 66–69, 1994.

[120202]21313.

[120203]Krawczak, M.; Smith-Sorensen, B.; Schmidtke, J.; Kakkar, V. V.;

[120204]Cooper, D. N.; Hovig, E.: Somatic spectrum of cancer-associated single

[120205]basepair substitutions in the TP53 gene is determined mainly by endogenous

[120206]mechanisms of mutation and by selection. Hum. Mutat. 5: 48-57, 1995.

[120207]21314.

[120208]Lamb, P.; Crawford, L.: Characterization of the human p53 gene. Molec.

[120209]Cell. Biol. 6: 1379-1385, 1986.

[120210]21315.

[120211]Latronico, A. C.; Pinto, E. M.; Domenice, S.; Fragoso, M. C. B.

[120212]V.; Martin, R. M.; Zerbini, M. C.; Lucon, A. M.; Mendonca, B. B.:

[120213]An inherited mutation outside the highly conserved DNA-binding domain

[120214]of the p53 tumor suppressor protein in children and adults with sporadic

[120215]adrenocortical tumors. J. Clin. Endocr. Metab. 86:
4970–4973, 2001.

[120216]21316.

[120217]Lavigueur, A.; Maltby, V.; Mock, D.; Rossant, J.; Pawson,
T.;

[120218]Bernstein, A.: High incidence of lung, bone, and lymphoid
tumors

[120219]in transgenic mice overexpressing mutant alleles of the
p53 oncogene. Molec.

[120220]Cell. Biol. 9: 3982–3991, 1989.

[120221]21317.

[120222]Law, J. C.; Strong, L. C.; Chidambaram, A.; Ferrell, R. E.: A

[120223]germ line mutation in exon 5 of the p53 gene in an ex-
tended cancer

[120224]family. Cancer Res. 51: 6385–6387, 1991.

[120225]21318.

[120226]Lee, J. M.; Bernstein, A.: p53 mutations increase resistance

[120227]to ionizing radiation. Proc. Nat. Acad. Sci. 90: 5742–5746,
1993.

[120228]21319.

[120229]Levine, A. J.: p53, the cellular gatekeeper for growth and division. Cell 88:

[120230]323–331, 1997.

[120231]21320.

[120232]Levine, A. J.; Momand, J.; Finlay, C. A.: The p53 tumour suppressor

[120233]gene. Nature 351: 453–456, 1991.

[120234]21321.

[120235]Li, M.; Chen, D.; Shiloh, A.; Luo, J.; Nikolaev, A. Y.; Qin, J.;

[120236]Gu, W.: Deubiquitination of p53 by HAUSP is an important pathway

[120237]for p53 stabilization. Nature 416: 648–653, 2002.

[120238]

[120239]100. Lu, H.; Levine, A. J.: Human TAFII31 protein is a transcriptional

[120240]coactivator of the p53 protein. Proc. Nat. Acad. Sci. 92: 5154–5158,

[120241]1995.

[120242]

[120243]101. Luo, J.; Nikolaev, A. Y.; Imai, S.; Chen, D.; Su, F.;

Shiloh,

[120244]A.; Guarente, L.; Gu, W.: Negative control of p53 by
Sir2- α promotes

[120245]cell survival under stress. Cell 107: 137–148, 2001.

[120246]

[120247]102. Luo, J.; Su, F.; Chen, D.; Shiloh, A.; Gu, W.: Deacety-
lation

[120248]of p53 modulates its effect on cell growth and apoptosis.
Nature 408:

[120249]377–381, 2000.

[120250]

[120251]103. Malkin, D.; Jolly, K. W.; Barbier, N.; Look, A. T.;
Friend, S.

[120252]H.; Gebhardt, M. C.; Andersen, T. I.; Borresen, A.-L.; Li, F.
P.;

[120253]Garber, J.; Strong, L. C.: Germline mutations of the p53
tumor-suppressor

[120254]gene in children and young adults with second malignant
neoplasms. New

[120255]Eng. J. Med. 326: 1309–1315, 1992.

[120256]

[120257]104. Malkin, D.; Li, F. P.; Strong, L. C.; Fraumeni, J. F., Jr.;
Nelson,

- [120258]C. E.; Kim, D. H.; Kassel, J.; Gryka, M. A.; Bischoff, F. Z.;
Tainsky,
[120259]M. A.; Friend, S. H.: Germ line p53 mutations in a familial
syndrome
[120260]of breast cancer, sarcomas, and other neoplasms. Science
250: 1233–1238,
[120261]1990.
[120262]
[120263]105. Marin, M. C.; Jost, C. A.; Brooks, L. A.; Irwin, M. S.;
O'Nions,
[120264]J.; Tidy, J. A.; James, N.; McGregor, J. M.; Harwood, C. A.;
Yulug,
[120265]J. G.; Vousden, K. H.; Allday, M. J.; Gusterson, B.; Ikawa, S.;
Hinds,
[120266]P. W.; Crook, T.; Kaelin, W. G., Jr. A common polymor-
phism acts
[120267]as an intragenic modifier of mutant p53 behaviour. Nature
Genet. 25:
[120268]47–54, 2000.
[120269]
[120270]106. Marino, S.; Vooijs, M.; van der Gulden, H.; Jonker, J.;
Berns,

[120271]A.: Induction of medulloblastomas in p53-null mutant mice by somatic

[120272]inactivation of Rb in the external granular layer cells of the cerebellum. Genes

[120273]Dev. 14: 994–1004, 2000.

[120274]

[120275]107. Masuda, H.; Miller, C.; Koeffler, H. P.; Battifora, H.; Cline,

[120276]M. J.: Rearrangement of the p53 gene in human osteogenic sarcomas. Proc.

[120277]Nat. Acad. Sci. 84: 7716–7719, 1987.

[120278]

[120279]108. Mathupala, S. P.; Heese, C.; Pedersen, P. L.: Glucose catabolism

[120280]in cancer cells: the type II hexokinase promoter contains functionally

[120281]active response elements for the tumor suppressor p53. J. Biol. Chem. 272:

[120282]22776–22780, 1997.

[120283]

[120284]109. Mazoyer, S.; Lalle, P.; Moyret-Lalle, C.; Marcais, C.;

Schraub,

[120285]S.; Frappaz, D.; Sobol, H.; Ozturk, M.: Two germ-line mutations affecting

[120286]the same nucleotide at codon 257 of p53 gene, a rare site for mutations. Oncogene 9:

[120287]1237-1239, 1994.

[120288]

[120289]110. McBride, O. W.; Merry, D.; Givol, D.: The gene for human p53

[120290]cellular tumor antigen is located on chromosome 17 short arm (17p13). Proc.

[120291]Nat. Acad. Sci. 83: 130-134, 1986.

[120292]

[120293]111. McBride, O. W.; Merry, D. E.; Oren, M.; Givol, D.: Human p53

[120294]cellular tumor antigen is on chromosome 17p13. (Abstract) Cytogenet.

[120295]Cell Genet. 40: 694-695, 1985.

[120296]

[120297]112. Metzger, A. K.; Sheffield, V. C.; Duyk, G.; Daneshvar, L.; Edwards,

[120298]M. S. B.; Cogen, P. H.: Identification of a germ-line mutation in

[120299]the p53 gene in a patient with an intracranial ependy-
moma. Proc.

[120300]Nat. Acad. Sci. 88: 7825-7829, 1991.

[120301]

[120302]113. Miller, C.; Mohandas, T.; Wolf, D.; Prokocimer, M.;
Rotter, V.;

[120303]Koeffler, H. P.: Human p53 gene localized to short arm of
chromosome

[120304]17. Nature 319: 783-784, 1986.

[120305]

[120306]114. Moll, U. M.; Riou, G.; Levine, A. J.: Two distinct
mechanisms

[120307]alter p53 in breast cancer: mutation and nuclear exclu-
sion. Proc.

[120308]Nat. Acad. Sci. 89: 7262-7266, 1992.

[120309]

[120310]115. Monpezat, J. P.; Delattre, O.; Bernard, A.; Grunwald,
D.; Remvikos,

[120311]Y.; Muleris, M.; Salmon, R. J.; Frelat, G.; Dutrillaux, B.;

Thomas,

[120312]G.: Loss of alleles on chromosome 18 and on the short arm of chromosome

[120313]17 in polyploid colorectal carcinomas. *Int. J. Cancer* 41: 404–408,

[120314]1988.

[120315]

[120316]116. Mulligan, L. M.; Matlashewski, G. J.; Scrable, H. J.; Cavenee,

[120317]W. K.: Mechanisms of p53 loss in human sarcomas. *Proc. Nat. Acad.*

[120318]*Sci.* 87: 5863–5867, 1990.

[120319]

[120320]117. Nakamura, T.; Pichel, J. G.; Williams–Simons, L.; Westphal, H.

[120321]: An apoptotic defect in lens differentiation caused by human p53

[120322]is rescued by a mutant allele. *Proc. Nat. Acad. Sci.* 92: 6142–6146,

[120323]1995.

[120324]

[120325]118. Neugut, A. I.; Santos, J.: The association between
cancers of

[120326]the small and large bowel. Cancer Epidemiol. Biomarkers
Prev. 2:

[120327]551–553, 1993.

[120328]

[120329]119. Nicol, C. J.; Harrison, M. L.; Laposa, R. R.;
Gimelshtein, I.

[120330]L.; Wells, P. G.: A teratologic suppressor role for p53 in
benzo[a]pyrene-treated

[120331]transgenic p53-deficient mice. Nature Genet. 10:
181–187, 1995.

[120332]

[120333]120. Nigro, J. M.; Baker, S. J.; Preisinger, A. C.; Jessup, J.
M.;

[120334]Hostetter, R.; Cleary, K.; Bigner, S. H.; Davidson, N.; Baylin,
S.;

[120335]Devilee, P.; Glover, T.; Collins, F. S.; Weston, A.; Modali, R.;

[120336]Harris,
C. C.; Vogelstein, B.: Mutations in the p53 gene occur in
diverse

[120337]human tumour types. Nature 342: 705–708, 1989.

[120338]

[120339]121. Oda, T.; Tsuda, H.; Scarpa, A.; Sakamoto, M.; Hirohashi, S.:

[120340]p53 gene mutation spectrum in hepatocellular carcinoma. Cancer Res. 52:

[120341]6358-6364, 1992.

[120342]

[120343]122. Oliner, J. D.; Kinzler, K. W.; Meltzer, P. S.; George, D. L.;

[120344]Vogelstein, B.: Amplification of a gene encoding a p53-associated

[120345]protein in human sarcomas. Nature 358: 80-83, 1992.

[120346]

[120347]123. Olivier, M.; Eeles, R.; Hollstein, M.; Khan, M. A.; Harris, C.

[120348]C.; Hainaut, P.: The IARC TP53 database: new online mutation analysis

[120349]and recommendations to users. Hum. Mutat. 19: 607-614, 2002.

[120350]

[120351]124. Ollmann, M.; Young, L. M.; Di Como, C. J.; Karim, F.;

Belvin,

[120352]M.; Robertson, S.; Whittaker, K.; Demsky, M.; Fisher, W.
W.; Buchman,

[120353]A.; Duyk, G.; Friedman, L.; Prives, C.; Kopczynski, C.:
Drosophila

[120354]p53 is a structural and functional homolog of the tumor
suppressor

[120355]p53. Cell 101: 91–101, 2000.

[120356]

[120357]125. Olschwang, S.; Laurent–Puig, P.; Vassal, A.; Salmon,
R.–J.; Thomas,

[120358]G.: Characterization of a frequent polymorphism in the
coding sequence

[120359]of the Tp53 gene in colonic cancer patients and a control
population. Hum.

[120360]Genet. 86: 369–370, 1991.

[120361]

[120362]126. Parsa, C. F.; Hoyt, C. S.; Lesser, R. L.; Weinstein, J. M.;
Strother,

[120363]C. M.; Muci–Mendoza, R.; Ramella, M.; Manor, R. S.;
Fletcher, W. A.;

[120364]Repka, M. X.; Garrity, J. A.; Ebner, R. N.; Monteiro, M. L. R.;

McFadzean,

[120365]**R. M.; Rubtsova, I. V.; Hoyt, W. F.: Spontaneous regression
of optic**

[120366]**gliomas: thirteen cases documented by serial neuroimag-
ing. Arch.**

[120367]**Ophthalmol. 119: 516–529, 2001.**

[120368]

[120369]**127. Patel, P.; Stephenson, J.; Scheuer, P. J.; Francis, G. E.:
p53**

[120370]**codon 249–ser mutations in hepatocellular carcinoma pa-
tients with**

[120371]**low aflatoxin exposure. (Letter) Lancet 339: 881, 1992.**

[120372]

[120373]**128. Pavletich, N. P.; Chambers, K. A.; Pabo, C. O.: The
DNA-binding**

[120374]**domain of p53 contains the four conserved regions and
the major mutation**

[120375]**hot spots. Genes Dev. 7: 2556–2564, 1993.**

[120376]

[120377]**129. Pearson, M.; Carbone, R.; Sebastiani, C.; Cioce, M.;
Fagioli,**

[120378]M.; Saito, S.; Higashimoto, Y.; Appella, E.; Minucci, S.;
Pandolfi,

[120379]P. P.; Pelicci, P. G.: PML regulates p53 acetylation and pre-
mature

[120380]senescence induced by oncogenic Ras. Nature 406:
207–210, 2000.

[120381]

[120382]130. Polyak, K.; Xia, Y.; Zweler, J. L.; Kinzler, K. W.; Vogel-
stein,

[120383]B.: A model for p53-induced apoptosis. Nature 389:
300–305, 1997.

[120384]

[120385]131. Potzsch, C.; Schaefer, H.–E.; Lubbert, M.: Familial and
metachronous

[120386]malignant lymphoma: absence of constitutional p53 mu-
tations. Am.

[120387]J. Hemat. 62: 144–149, 1999.

[120388]

[120389]132. Raj, K.; Ogston, P.; Beard, P.: Virus-mediated killing
of cells

[120390]that lack p53 activity. Nature 412: 914–917, 2001. Note:
Addendum:

[120391]Nature 416: 202 only, 2002.

[120392]

[120393]133. Raman, V.; Martensen, S. A.; Reisman, D.; Evron, E.;
Odenwald,

[120394]W. F.; Jaffee, E.; Marks, J.; Sukumar, S.: Compromised
HOXA5 function

[120395]can limit p53 expression in human breast tumours. Nature
405: 974–978,

[120396]2000.

[120397]

[120398]134. Reilly, K. M.; Loisel, D. A.; Bronson, R. T.; McLaugh-
lin, M.

[120399]E.; Jacks, T.: Nf1;Trp53 mutant mice develop glioblastoma
with evidence

[120400]of strain-specific effects. Nature Genet. 26: 109–113,
2000.

[120401]

[120402]135. Reisman, D.; Balint, E.; Loging, W. T.; Rotter, V.; Al-
mon, E.

[120403]: A novel transcript encoded within the 10-kb first intron
of the

[120404]human p53 tumor suppressor gene (D17S2179E) is induced during differentiation

[120405]of myeloid leukemia cells. Genomics 38: 364–370, 1996.

[120406]

[120407]136. Reisman, D.; Greenberg, M.; Rotter, V.: Human p53 oncogene contains

[120408]one promoter upstream of exon 1 and a second, stronger promoter within

[120409]intron 1. Proc. Nat. Acad. Sci. 85: 5146–5150, 1988.

[120410]

[120411]137. Ribeiro, R. C.; Sandrini, F.; Figueiredo, B.; Zambetti, G. P.;

[120412]Michalkiewicz, E.; Lafferty, A. R.; DeLacerda, L.; Rabin, M.; Cadwell,

[120413]C.; Sampaio, G.; Cat, I.; Stratakis, C. A.; Sandrini, R.: An inherited

[120414]p53 mutation that contributes in a tissue-specific manner to pediatric

[120415]adrenal cortical carcinoma. Proc. Nat. Acad. Sci. 98: 9330–9335,

[120416]2001.

[120417]

[120418]138. Rideout, W. M., III; Coetzee, G. A.; Olumi, A. F.; Jones, P.

[120419]A.: 5-Methylcytosine as an endogenous mutagen in the human LDL receptor

[120420]and p53 genes. Science 249: 1288–1290, 1990.

[120421]

[120422]139. Rodrigues, N. R.; Rowan, A.; Smith, M. E. F.; Kerr, I. B.; Bodmer,

[120423]W. F.; Gannon, J. V.; Lane, D. P.: p53 mutations in colorectal cancer. Proc.

[120424]Nat. Acad. Sci. 87: 7555–7559, 1990.

[120425]

[120426]140. Romano, J. W.; Ehrhart, J. C.; Duthu, A.; Kim, C. M.; Appella,

[120427]E.; May, P.: A mutation in the p53 gene of a human osteosarcoma cell

[120428]line. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A32, 1989.

[120429]

[120430]141. Roth, J. A.; Nguyen, D.; Lawrence, D. D.; Kemp, B. L.; Carrasco,

[120431]C. H.; Ferson, D. Z.; Hong, W. K.; Komaki, R.; Lee, J. J.;
Nesbitt,

[120432]J. C.; Pisters, K. M. W.; Putnam, J. B.; Schea, R.; Shin, D. M.;
Walsh,

[120433]G. L.; Dolormente, M. M.; Han, C.-I.; Martin, F. D.; Yen, N.;
Xu,

[120434]K.; Stephens, L. C.; McDonnell, T. J.; Mukhopadhyay, T.;
Cai, D.:

[120435]Retrovirus-mediated wild-type p53 gene transfer to tu-
mors of patients

[120436]with lung cancer. Nature Med. 2: 985-990, 1996.

[120437]

[120438]142. Rowley, J. D.: Personal Communication. Chicago, Ill.
1/3/1986.

[120439]

[120440]143. Rutherford, J.; Chu, C. E.; Duddy, P. M.; Charlton, R.
S.; Chumas,

[120441]P.; Taylor, G. R.; Lu, X.; Barnes, D. M.; Camplejohn, R. S.:
Investigations

[120442]on a clinically and functionally unusual and novel germline
p53 mutation. Brit.

[120443]J. Cancer 86: 1592-1596, 2002.

[120444]

[120445]144. Ryan, K. M.; Ernst, M. K.; Rice, N. R.; Vousden, K. H.:

Role

[120446]of NF- κ B in p53-mediated programmed cell death.

Nature 404:

[120447]892–897, 2000.

[120448]

[120449]145. Sah, V. P.; Arradi, L. D.; Mulligan, G. J.; Williams, B.

O.;

[120450]Bronson, R. T.; Jacks, T.: A subset of p53-deficient em-

bryos exhibit

[120451]exencephaly. Nature Genet. 10: 175–180, 1995.

[120452]

[120453]146. Santibanez-Koref, M. F.; Birch, J. M.; Hartley, A. L.;

Morris

[120454]Jones, P. H.; Craft, A. W.; Eden, T.; Crowther, D.; Kelsey, A.

M.;

[120455]Harris, M.: p53 germline mutations in Li-Fraumeni syn-

drome. Lancet 338:

[120456]1490–1491, 1991.

[120457]

- [120458]147. Schiffer, D.; Cavalla, P.; Di Sapio, A.; Giordana, M. T.;
Mauro,
[120459]A.: Mutations and immunohistochemistry of p53 and pro-
liferation markers
[120460]in astrocytic tumors of childhood. Child's Nerv. Syst. 11:
517–522,
[120461]1995.
[120462]
[120463]148. Schmitt, C. A.; Fridman, J. S.; Yang, M.; Lee, S.; Bara-
nov, E.;
[120464]Hoffman, R. M.; Lowe, S. W.: A senescence program con-
trolled by p53
[120465]and p16–INK4a contributes to the outcome of cancer
therapy. Cell 109:
[120466]335–346, 2002.
[120467]
[120468]149. Sedlacek, Z.; Kodet, R.; Kriz, V.; Seemanova, E.; Vod-
varka, P.;
[120469]Wilgenbus, P.; Mares, J.; Poustka, A.; Goetz, P.: Two Li-
Fraumeni
[120470]syndrome families with novel germline p53 mutations:
loss of the wild-type

[120471]p53 allele in only 50% of tumours. Brit. J. Cancer 77:
1034–1039,

[120472]1998.

[120473]

[120474]150. Shieh, S.–Y.; Ikeda, M.; Taya, Y.; Prives, C.: DNA
damage–induced

[120475]phosphorylation of p53 alleviates inhibition by MDM2. Cell
91: 325–334,

[120476]1997.

[120477]

[120478]151. Sidransky, D.; Von Eschenbach, A.; Tsai, Y. C.; Jones,
P.; Summerhayes,

[120479]I.; Marshall, F.; Paul, M.; Green, P.; Hamilton, S. R.; Frost,
P.;

[120480]Vogelstein, B.: Identification of p53 gene mutations in
bladder cancers

[120481]and urine samples. Science 252: 706–709, 1991.

[120482]

[120483]152. Smela, M. E.; Hamm, M. L.; Henderson, P. T.; Harris,
C. M.; Harris,

[120484]T. M.; Essigmann, J. M.: The aflatoxin B1 formamidopy–

rimidine adduct

[120485] plays a major role in causing the types of mutations observed in human

[120486] hepatocellular carcinoma. Proc. Nat. Acad. Sci. 99: 6655–6660, 2002.

[120487]

[120488] 153. Smith, H. S.: Personal Communication. San Francisco, Calif.

[120489] 11/16/1993.

[120490]

[120491] 154. Smith–Sorensen, B.; Gebhardt, M. C.; Kloen, P.; McIntyre, J.;

[120492] Aguilar, F.; Cerutti, P.; Borresen, A.–L.: Screening for TP53 mutations

[120493] in osteosarcomas using constant denaturant gel electrophoresis (CDGE). Hum.

[120494] Mutat. 2: 274–285, 1993.

[120495]

[120496] 155. Srivastava, S.; Zou, Z.; Pirollo, K.; Blattner, W.; Chang, E.

[120497] H.: Germ–line transmission of a mutated p53 gene in a

cancer-prone

[120498]family with Li-Fraumeni syndrome. Nature 348: 747-749, 1990.

[120499]

[120500]156. Storey, A.; Thomas, M.; Kalita, A.; Harwood, C.; Gardiol, D.;

[120501]Mantovani, F.; Breuer, J.; Leigh, I. M.; Matlashewski, G.; Banks,

[120502]L.: Role of a p53 polymorphism in the development of human papilloma-virus-associated

[120503]cancer. Nature 393: 229-234, 1998.

[120504]

[120505]157. Sun, Y.; Hegamyer, G.; Cheng, Y.-J.; Hildesheim, A.; Chen, J.-Y.;

[120506]Chen, I.-H.; Cao, Y.; Yao, K.-T.; Colburn, N. H.: An infrequent point

[120507]mutation of the p53 gene in human nasopharyngeal carcinoma. Proc.

[120508]Nat. Acad. Sci. 89: 6516-6520, 1992.

[120509]

[120510]158. Takahashi, T.; D'Amico, D.; Chiba, I.; Buchhagen, D.

L.; Minna,

[120511]J. D.: Identification of intronic point mutations as an alternative

[120512]mechanism for p53 inactivation in lung cancer. J. Clin. Invest. 86:

[120513]363–369, 1990.

[120514]

[120515]159. Takahashi, T.; Nau, M. M.; Chiba, I.; Birrer, M. J.; Rosenberg,

[120516]R. K.; Vinocour, M.; Levitt, M.; Pass, H.; Gazdar, A. F.; Minna, J.

[120517]D.: p53: a frequent target for genetic abnormalities in lung cancer. Science 246:

[120518]491–494, 1989.

[120519]

[120520]160. Toguchida, J.; Yamaguchi, T.; Dayton, S. H.; Beauchamp, R. L.;

[120521]Herrera, G. E.; Ishizaki, K.; Yamamuro, T.; Meyers, P. A.; Little,

[120522]J. B.; Sasaki, M. S.; Weichselbaum, R. R.; Yandell, D. W.: Prevalence

[120523]and spectrum of germline mutations of the p53 gene

among patients

[120524]with sarcoma. New Eng. J. Med. 326: 1301–1308, 1992.

[120525]

[120526]161. Tonisson, N.; Zernant, J.; Kurg, A.; Pavel, H.; Slavin, G.; Roomere,

[120527]H.; Meiel, A.; Hainaut, P.; Metspalu, A.: Evaluating the arrayed
rayed

[120528]primer extension resequencing assay of TP53 tumor suppressor gene. Proc.

[120529]Nat. Acad. Sci. 99: 5503–5508, 2002.

[120530]

[120531]162. Tyner, S. D.; Venkatachalam, S.; Choi, J.; Jones, S.; Ghebranious,

[120532]N.; Igelmann, H.; Lu, X.; Soron, G.; Cooper, B.; Brayton, C.; Park,

[120533]S. H.; Thompson, T.; Karsenty, G.; Bradley, A.; Donehower, L. A.:

[120534]p53 mutant mice that display early ageing-associated phenotypes. Nature 415:

[120535]45–53, 2002.

[120536]

[120537]163. Ueda, H.; Ullrich, S. J.; Gangemi, J. D.; Kappel, C. A.;
Ngo,

[120538]L.; Feitelson, M. A.; Jay, G.: Functional inactivation but not
structural

[120539]mutation of p53 causes liver cancer. Nature Genet. 9:
41–47, 1995.

[120540]

[120541]164. Unger, T.; Nau, M. M.; Segal, S.; Minna, J. D.: p53: a
transdominant

[120542]regulator of transcription whose function is ablated by
mutations

[120543]occurring in human cancer. EMBO J. 11: 1383–1390,
1992.

[120544]

[120545]165. vanTuinen, P.; Ledbetter, D. H.: Construction and
utilization

[120546]of a detailed somatic cell hybrid mapping panel for human
chromosome

[120547]17: localization of an anonymous clone to the critical re-
gion of Miller–Dieker

[120548]syndrome, deletion 17p13. (Abstract) Cytogenet. Cell
Genet. 46:

[120549]708–709, 1987.

[120550]

[120551]166. Varley, J. M.; McGowan, G.; Thorncroft, M.; Tricker, K.
J.; Teare,,

[120552]M. D.; Santibanez–Koref, M. F.; Houlston, R. S.; Martin, J.;
Birch,

[120553]J. M.; Evans, D. G. R.: An extended Li–Fraumeni kindred
with gastric

[120554]carcinoma and a codon 175 mutation in TP53. J. Clin.
Genet. 32:

[120555]942–945, 1995.

[120556]

[120557]167. Varley, J. M.; McGown, G.; Thorncroft, M.; Cochrane,
S.; Morrison,

[120558]P.; Woll, P.; Kelsey, A. M.; Mitchell, E. L. D.; Boyle, J.; Birch,

[120559]J. M.; Evans, D. G. R.: A previously undescribed mutation
within

[120560]the tetramerisation domain of TP53 in a family with Li–
Fraumeni syndrome

[120561](sic). Oncogene 12: 2437–2442, 1996.

[120562]

[120563]168. Vaziri, H.; Dessain, S. K.; Eaton, E. N.; Imai, S.-I.;
Frye,

[120564]R. A.; Pandita, T. K.; Guarente, L.; Weinberg, R. A.:
hSIR2-SIRT1

[120565]functions as an NAD-dependent p53 deacetylase. Cell
107: 149–159,

[120566]2001.

[120567]

[120568]169. Vogelstein, B.; Kinzler, K. W.: X-rays strike p53
again. Nature 370:

[120569]174–175, 1994.

[120570]

[120571]170. Vogelstein, B.; Kinzler, K. W.: p53 function and dys-
function. Cell 70:

[120572]523–526, 1992.

[120573]

[120574]171. Wheeler, J. M. D.; Warren, B. F.; Mortensen, N. J. M.;
Kim, H.

[120575]C.; Biddolph, S. C.; Elia, G.; Beck, N. E.; Williams, G. T.;
Shepherd,

[120576]N. A.; Bateman, A. C.; Bodmer, W. F.: An insight into the

genetic

[120577] pathway of adenocarcinoma of the small intestine. Gut 50:
218–223,

[120578] 2002.

[120579]

[120580] 172. Yamanishi, Y.; Boyle, D. L.; Rosengren, S.; Green, D.
R.; Zvaifler,

[120581] N. J.; Firestein, G. S.: Regional analysis of p53 mutations in
rheumatoid

[120582] arthritis synovium. Proc. Nat. Acad. Sci. 99: 10025–10030,
2002.

[120583]

[120584] 173. Yu, A.; Fan, H.-Y.; Liao, D.; Bailey, A. D.; Weiner, A.
M.:

[120585] Activation of p53 or loss of the Cockayne syndrome group
B repair

[120586] protein causes metaphase fragility of human U1, U2, and
5S genes. Molec.

[120587] Cell 5: 801–810, 2000.

[120588]

[120589] 174. Yu, J. L.; Rak, J. W.; Coomber B. L.; Hicklin, D. J.; Ker-

bel,

[120590]R. S.: Effect of p53 status on tumor response to antiangiogenic therapy. Science 295:

[120591]1526–1528, 2002.

[120592]

[120593]175. Zhang, W.; Funk, W. D.; Wright, W. E.; Shay, J. W.; Deisseroth,

[120594]A. B.: Novel DNA binding of p53 mutants and their role in transcriptional

[120595]activation. Oncogene 8: 2555–2559, 1993.

[120596]

[120597]176. Zhang, Y.; Xiong, Y.: A p53 amino-terminal nuclear export signal

[120598]inhibited by DNA damage-induced phosphorylation. Science 292: 1910–1915,

[120599]2001.

[120600]

[120601]177. Zhu, C.; Mills, K. D.; Ferguson, D. O.; Lee, C.; Manis, J.; Fleming,

[120602]J.; Gao, Y.; Morton, C. C.; Alt, F. W.: Unrepaired DNA breaks in

[120603]p53-deficient cells lead to oncogenic gene amplification
subsequent

[120604]to translocations. Cell 109: 811–821, 2002.

[120605]

[120606]178. Ziegler, A.; Jonason, A. S.; Leffell, D. J.; Simon, J. A.;
Sharma,

[120607]H. W.; Kimmelman, J.; Remington, L.; Jacks, T.; Brash, D.
E.: Sunburn

[120608]and p53 in the onset of skin cancer. Nature 372:
773–776, 1994.

[120609]21322.

[120610]Juppner, H.: Molecular cloning and characterization of a
parathyroid

[120611]hormone/parathyroid hormone-related peptide receptor:
a member of

[120612]an ancient family of G protein-coupled receptors. Curr.
Opin. Nephrol.

[120613]Hypertension 3: 371–378, 1994.

[120614]21323.

[120615]Juppner, H.; Abou-Samra, A.-B.; Freeman, M.; Kong, X. F.;
Schipani,

- [120616]E.; Richards, J.; Kolakowski, L. F., Jr.; Hock, J.; Potts, J. T.,
[120617]Jr.; Kronenberg, H. M.; Segre, G. V.: A G protein-linked re-
ceptor
[120618]for parathyroid hormone and parathyroid hormone-re-
lated peptide. Science 254:
[120619]1024–1026, 1991.
[120620]21324.
[120621]Karperien, M.; van der Harten, H. J.; van Schooten, R.;
Farih-Sips,
[120622]H.; den Hollander, N. S.; Kneppers, S. L. J.; Nijweide, P.;
Papapoulos,
[120623]S. E.; Lowik, C. W. G. M.: A frame-shift mutation in the
type I parathyroid
[120624]hormone (PTH)/PTH-related peptide receptor causing
Blomstrand lethal
[120625]osteochondrodysplasia. J. Clin. Endocr. Metab. 84:
3713–3720, 1999.
[120626]21325.
[120627]Lanske, B.; Karaplis, A. C.; Lee, K.; Luz, A.; Vortkamp, A.;
Pirro,
[120628]A.; Karperien, M.; Defize, L. H. K.; Ho, C.; Mulligan, R. C.;
Abou-Samra,

[120629]A.-B.; Juppner, H.; Segre, G. V.; Kronenberg, H. M.: PTH/
PTHrP receptor

[120630]in early development and Indian hedgehog-regulated
bone growth. Science 273:

[120631]663–666, 1996.

[120632]21326.

[120633]Loshkajian, A.; Roume, J; Stanescu, V.; Delezoide, A.;
Stampf,

[120634]F.; Maroteaux, P.: Familial Blomstrand chondrodysplasia
with advanced

[120635]skeletal maturation: further delineation. Am. J. Med.
Genet. 71:

[120636]283–288, 1997.

[120637]21327.

[120638]Mahon, M. J.; Donowitz, M.; Yun, C. C.; Segre, G. V.:
Na⁺/H⁺

[120639]exchanger regulatory factor 2 directs parathyroid hor-
mone 1 receptor

[120640]signalling. Nature 417: 858–861, 2002.

[120641]21328.

[120642]Manen, D.; Palmer, G.; Bonjour, J.-P.; Rizzoli, R.: Positive

[120643]and negative control of the expression of parathyroid hormone (PTH)/PTH-related

[120644]protein receptor via proximal promoter P3 in human osteoblast-like

[120645]cells. J. Clin. Endocr. Metab. 85: 3376–3382, 2000.

[120646]21329.

[120647]McCuaig, K. A.; Clarke, J. C.; White, J. H.: Molecular cloning

[120648]of the gene encoding the mouse parathyroid hormone/parathyroid hormone-related

[120649]peptide receptor. Proc. Nat. Acad. Sci. 91: 5051–5055, 1994.

[120650]21330.

[120651]Minagawa, M.; Watanabe, T.; Kohno, Y.; Mochizuki, H.; Hendy, G.

[120652]N.; Goltzman, D.; White, J. H.; Yasuda, T.: Analysis of the P3 promoter

[120653]of the human parathyroid hormone (PTH)/PTH-related peptide receptor

[120654]gene in pseudohypoparathyroidism type 1b. J. Clin. Endocr. Metab. 86:

[120655]1394–1397, 2001.

[120656]21331.

[120657]Minagawa, M.; Yasuda, T.; Watanabe, T.; Minamitani, K.;
Takahashi,

[120658]Y.; Goltzman, D.; White, J. H.; Hendy, G. N.; Kohno, Y.: As-
sociation

[120659]between AAAG repeat polymorphism in the P3 promoter
of the human parathyroid

[120660]hormone (PTH)/PTH-related peptide receptor gene and
adult height,

[120661]urinary pyridinoline excretion, and promoter activity. J.
Clin. Endocr.

[120662]Metab. 87: 1791-1796, 2002.

[120663]21332.

[120664]Parfitt, A. M.; Schipani, E.; Rao, D. S.; Kupin, W.; Han,
Z.-H.;

[120665]Juppner, H.: Hypercalcemia due to constitutive activity of
the parathyroid

[120666]hormone (PTH)/PTH-related peptide receptor: comparison
with primary

[120667]hyperparathyroidism. J. Clin. Endocr. Metab. 81:
3584-3588, 1996.

[120668]21333.

[120669] Schipani, E.; Karga, H.; Karaplis, A. C.; Potts, J. T, Jr.; Kronenberg,

[120670] H. M.; Segre, G. V.; Abou-Samra, A.-B.; Juppner, H.: Identical complementary

[120671] deoxyribonucleic acids encode a human renal and bone parathyroid hormone

[120672] (PTH)/PTH-related peptide receptor. *Endocrinology* 132: 2157-2165,

[120673] 1993.

[120674] 21334.

[120675] Schipani, E.; Langman, C.; Hunzelman, J.; Le Merrer, M.; Loke,

[120676] K. Y.; Dillon, M. J.; Silve, C.; Juppner, H.: A novel parathyroid

[120677] hormone (PTH)/PTH-related peptide receptor mutation in Jansen's metaphyseal

[120678] chondrodysplasia. *J. Clin. Endocr. Metab.* 84: 3052-3057, 1999.

[120679] 21335.

[120680] Schipani, E.; Langman, C. B.; Parfitt, A. M.; Jensen, G. S.; Kikuchi,

[120681] S.; Kooh, S. W.; Cole, W. G.; Juppner, H.: Constitutively ac-

tivated

[120682]receptors for parathyroid hormone and parathyroid hormone-related

[120683]peptide in Jansen's metaphyseal chondrodysplasia. New Eng. J. Med. 335:

[120684]708-714, 1996.

[120685]21336.

[120686]Pausova, Z.; Bourdon, J.; Clayton, D.; Mattei, M.-G.; Seldin,

[120687]M. F.; Janicic, N.; Riviere, M.; Szpirer, J.; Levan, G.; Szpirer,

[120688]C.; Goltzman, D.; Hendy, G. N.: Cloning of a parathyroid hormone/parathyroid

[120689]hormone-related peptide receptor (PTHrP) cDNA from a rat osteosarcoma

[120690](UMR 106) cell line: chromosomal assignment of the gene in the human,

[120691]mouse, and rat genomes. Genomics 20: 20-26, 1994.

[120692]21337.

[120693]Schipani, E.; Weinstein, L. S.; Bergwitz, C.; Iida-Klein, A.;

[120694]Kong, X. F.; Stuhmann, M.; Kruse, K.; Whyte, M. P.; Murray, T.; Schmidtke,

[120695]J.; van Dop, C.; Brickman, A. S.; Crawford, J. D.; Potts, J. T.,

Jr.;

[120696]Kronenberg, H. M.; Abou-Samra, A. B.; Segre, G. V.; Juppner, H.:

[120697]Pseudohypoparathyroidism type Ib is not caused by mutations in the

[120698]coding exons of the human parathyroid hormone (PTH)/PTH-related peptide

[120699]receptor gene. J. Clin. Endocr. Metab. 80: 1611-1621, 1995.

[120700]21338.

[120701]Suarez, F.; Lebrun, J. J.; Lecossier, D.; Escoubet, B.; Coureau,

[120702]C.; Silve, C.: Expression and modulation of the parathyroid hormone

[120703](PTH)/PTH-related peptide receptor messenger ribonucleic acid in skin

[120704]fibroblasts from patients with type Ib pseudohypoparathyroidism. J.

[120705]Clin. Endocr. Metab. 80: 965-970, 1995.

[120706]21339.

[120707]Zhang, P.; Jobert, A.-S.; Couvineau, A.; Silve, C.: A homozygous

- [120708]inactivating mutation in the parathyroid hormone/
parathyroid hormone-related
- [120709]peptide receptor causing Blomstrand chondrodysplasia. J.
Clin. Endocr.
- [120710]Metab. 83: 3365–3368, 1998.
- [120711]21340.
- [120712]Albright, F.: Case records of the Massachusetts General
Hospital
- [120713](case 27461). New Eng. J. Med. 225: 789–791, 1941.
- [120714]21341.
- [120715]Bakre, M. M.; Zhu, Y.; Yin, H.; Burton, D. W.; Terkeltaub,
R.;
- [120716]Deftos, L. J.; Varner, J. A.: Parathyroid hormone-related
peptide
- [120717]is a naturally occurring, protein kinase A-dependent an-
giogenesis
- [120718]inhibitor. Nature Med. 8: 995–1003, 2002.
- [120719]21342.
- [120720]Broadus, A. E.; Mangin, M.; Ikeda, K.; Insogna, K. L.; Weir,
E.
- [120721]C.; Burtis, W. J.; Stewart, A. F.: Humoral hypercalcemia of

cancer:

[120722]identification of a novel parathyroid hormone-like peptide. New Eng.

[120723]J. Med. 319: 556-563, 1988.

[120724]21343.

[120725]Hammonds, R. G., Jr.; McKay, P.; Winslow, G. A.; Diefenbach-Jagger,

[120726]H.; Grill, V.; Glatz, J.; Rodda, C. P.; Moseley, J. M.; Wood, W. I.;

[120727]Martin, T. J.: Purification and characterization of recombinant human

[120728]parathyroid hormone-related protein. J. Biol. Chem. 264: 14806-14811,

[120729]1989.

[120730]21344.

[120731]Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.; Yasuda,

[120732]T.; Banville, D.; Goltzman, D.: Gene for parathyroid hormone-like

[120733]peptide is on mouse chromosome 6. Cytogenet. Cell Genet. 53: 80-82,

[120734]1990.

[120735]21345.

[120736]Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.;
Yasuda,

[120737]T.; Banville, D.; Goltzman, D.: Gene for parathyroid hor-
mone-like

[120738]peptide is on mouse chromosome 6. (Abstract) Cytogenet.
Cell Genet. 51:

[120739]1003 only, 1989.

[120740]21346.

[120741]Kim, S. K.; Selleri, L.; Lee, J. S.; Zhang, A. Y.; Gu, X.; Ja-
cobs,

[120742]Y.; Cleary, M. L.: Pbx1 inactivation disrupts pancreas de-
velopment

[120743]and in Ipf1-deficient mice promotes diabetes mellitus.
Nature Genet. 30:

[120744]430-435, 2002.

[120745]21347.

[120746]Mercader, N.; Leonardo, E.; Azpiazu, N.; Serrano, A.;
Morata, G.;

[120747]Martinez-A, C.; Torres, M.: Conserved regulation of proxi-
modistal

[120748]limb axis development by Meis1/Hth. Nature 402:
425–429, 1999.

[120749]21348.

[120750]Monica, K.; Galili, N.; Nourse, J.; Saltman, D.; Cleary, M. L.

[120751]: PBX2 and PBX3, new homeobox genes with extensive homology to the

[120752]human proto-oncogene PBX1. Molec. Cell. Biol. 11:
6149–6157, 1991.

[120753]21349.

[120754]Aguado, B.; Campbell, R. D.: The novel gene G17, located
in the

[120755]human major histocompatibility complex, encodes PBX2, a
homeodomain-containing

[120756]protein. Genomics 25: 650–659, 1995.

[120757]21350.

[120758]Katsanis, N.; Fitzgibbon, J.; Fisher, E. M. C.: Paralogy mapping:

[120759]identification of a region in the human MHC triplicated
onto human

[120760]chromosomes 1 and 9 allows the prediction and isolation
of novel PBX

[120761]and NOTCH loci. Genomics 35: 101–108, 1996.

[120762]21351.

[120763]Eisenhofer, G.; Walther, M. M.; Huynh, T.–T.; Li, S.–T.;
Bornstein,

[120764]S. R.; Vortmeyer, A.; Mannelli, M.; Goldstein, D. S.; Line-
han, W.

[120765]M.; Lenders, J. W. M.; Pacak, K.: Pheochromocytomas in
von Hippel–Lindau

[120766]syndrome and multiple endocrine neoplasia type 2 display
distinct

[120767]biochemical and clinical phenotypes. J. Clin. Endocr.
Metab. 86:

[120768]1999–2008, 2001.

[120769]21352.

[120770]Glasser, S. W.; Korfhagen, T. R.; Weaver, T. E.; Clark, J. C.;

[120771]Pilot–Matias, T.; Meuth, J.; Fox, J. L.; Whitsett, J. A.: cDNA,
deduced

[120772]polypeptide structure and chromosomal assignment of
human pulmonary

[120773]surfactant proteolipid, SPL(pVal). J. Biol. Chem. 263: 9–12,
1988.

[120774]21353.

[120775]Ahmad, S.; Banville, D.; Zhao, Z.; Fischer, E. H.; Shen, S.-H.

[120776]: A widely expressed human protein-tyrosine phosphatase
containing

[120777]src homology 2 domains. Proc. Nat. Acad. Sci. 90:
2197-2201, 1993.

[120778]21354.

[120779]Bastien, L.; Ramachandran, C.; Liu, S.; Adam, M.: Cloning,
expression,

[120780]and mutational analysis of SH-PTP2, human protein-ty-
rosine phosphatase

[120781]2-domains. Biochem. Biophys. Res. Commun. 196:
124-133, 1993.

[120782]21355.

[120783]Dechert, U.; Duncan, A. M. V.; Bastien, L.; Duff, C.; Adam,
M.;

[120784]Jirik, F. R.: Protein-tyrosine phosphatase SH-PTP2
(PTPN11) is localized

[120785]to 12q24.1-24.3. Hum. Genet. 96: 609-615, 1995.

[120786]21356.

[120787]Higashi, H.; Tsutsumi, R.; Muto, S.; Sugiyama, T.; Azuma,

T; Asaka,

[120788]M.; Hatakeyama, M.: SHP-2 tyrosine phosphatase as an intracellular

[120789]target of *Helicobacter pylori* CagA protein. *Science* 295: 683–686,

[120790]2002.

[120791]21357.

[120792]Hof, P.; Pluskey, S.; Dhe-Paganon, S.; Eck, M. J.; Shoelson, S.

[120793]E.: Crystal structure of the tyrosine phosphatase SHP-2. *Cell* 92:

[120794]441–450, 1998.

[120795]21358.

[120796]Isobe, M.; Hinoda, Y.; Imai, K.; Adachi, M.: Chromosomal localization

[120797]of an SH2 containing tyrosine phosphatase (SH-PTP3) gene to chromosome

[120798]12q24.1. *Oncogene* 9: 1751–1753, 1994.

[120799]21359.

[120800]Qu, C.-K.; Yu, W.-M.; Azzarelli, B.; Cooper, S.; Broxmeyer, H.

- [120801]E.; Feng, G.-S.: Biased suppression of hematopoiesis and multiple
developmental defects in chimeric mice containing Shp-2
mutant cells. Molec.
Cell. Biol. 18: 6075-6082, 1998.
[120804]21360.
- [120805]Saxton, T. M.; Ciruna, B. G.; Holmyard, D.; Kulkarni, S.;
Harpal,
[120806]K.; Rossant, J.; Pawson, T.: The SH2 tyrosine phosphatase
Shp2 is
[120807]required for mammalian limb development. Nature Genet.
24: 420-423,
[120808]2000.
[120809]21361.
- [120810]Saxton, T. M.; Henkemeyer, M.; Gasca, S.; Shen, R.; Rossi,
D.
[120811]J.; Shalaby, F.; Feng, G.-S.; Pawson, T.: Abnormal meso-
derm patterning
[120812]in mouse embryos mutant for the SH2 tyrosine phos-
phatase Shp-2. EMBO
[120813]J. 16: 2352-2364, 1997.

[120814]21362.

[120815]Tartaglia, M.; Kalidas, K.; Shaw, A.; Song, X.; Musat, D. L.;

[120816]van der Burgt, I.; Brunner, H. G.; Bertola, D. R.; Crosby, A.;
Ion,

[120817]A.; Kucherlapati, R. S.; Jeffery, S.; Patton, M. A.; Gelb, B.
D.:

[120818]PTPN11 mutations in Noonan syndrome: molecular spec-
trum, genotype-phenotype

[120819]correlation, and phenotypic heterogeneity. Am. J. Hum.
Genet. 70:

[120820]1555-1563, 2002.

[120821]21363.

[120822]Zhao, Z. J.; Zhao, R.: Purification and cloning of PZR, a
binding

[120823]protein and putative physiological substrate of tyrosine
phosphatase

[120824]SHP-2. J. Biol. Chem. 273: 29367-29372, 1998.

[120825]21364.

[120826]Allaart, C. F.; Aronson, D. C.; Ruys, T.; Rosendaal, F. R.;
van

[120827]Bockel, J. H.; Bertina, R. M.; Briet, E.: Hereditary protein S

deficiency

[120828]in young adults with arterial occlusive disease. Thromb.

Haemost. 64:

[120829]206–210, 1990.

[120830]21365.

[120831]Beauchamp, N. J.; Daly, M. E.; Makris, M.; Preston, F. E.;

Peake,

[120832]J. R.: A novel mutation in intron K of the PROS1 gene

causes aberrant

[120833]RNA splicing and is a common cause of protein S defi-

ciency in a UK

[120834]thrombophilia cohort. Thromb. Haemost. 79: 1086–1091,

1998.

[120835]21366.

[120836]Bertina, R. M.: Nomenclature proposal for protein S defi-

ciency XXXVI

[120837]Annual meeting of the Scientific and Standardization

Committee of

[120838]the ISTH Barcelona, and Spain, 1990.

[120839]21367.

[120840]Bertina, R. M.; Ploos van Amstel, H. K.; van Wijngaarden,

- A.; Coenen,
[120841]J.; Leemhuis, M. P.; Deutz-Terlouw, P. P.; van der Linden,
I. K.;
[120842]Reitsma, P. H.: Heerlen polymorphism of protein S, an im-
munologic
[120843]polymorphism due to dimorphism of residue 460. Blood
76: 538-548,
[120844]1990.
[120845]21368.
[120846]Bosson, J. L.; Francois, P.; Pernod, G.; Carpentier, P.: Mal-
adie
[120847]thrombo-embolique veineuse chez l'enfant liee a un
deficit transitoire
[120848]en proteine S au decours d'une varicelle. (Letter) Presse
Med. 24:
[120849]415 only, 1995.
[120850]21369.
[120851]Chafa, O.; Fischer, A. M.; Meriane, F.; Chellali, F.; Rahal, S.;
[120852]Sternberg, C.; Benabadji, M.: A new case of 'type II' inher-
ited protein
[120853]S deficiency. Brit. J. Haemat. 73: 501-505, 1989.

[120854]21370.

[120855]Clark, D. A.; Williams, W. L.; Marlar, R. A.: Mesenteric vein
[120856]thrombosis associated with a familial deficiency of free
protein S. Arch.

[120857]Path. Lab. Med. 115: 617–619, 1991.

[120858]21371.

[120859]Comp, P. C.; Esmon, C. T.: Recurrent venous thromboem-
bolism in

[120860]patients with a partial deficiency of protein S. New Eng. J.
Med. 311:

[120861]1525–1528, 1984.

[120862]21372.

[120863]Lorimer, D. D.; Benya, R. V.: Cloning and quantification of
galanin-1

[120864]receptor expression by mucosal cells lining the human
gastrointestinal

[120865]tract. Biochem. Biophys. Res. Commun. 222: 379–385,
1996.

[120866]21373.

[120867]Nicholl, J.; Kofler, B.; Sutherland, G. R.; Shine, J.; Iismaa,

[120868]T. P.: Assignment of the gene encoding human galanin receptor (GALNR)

[120869]to 18q23 by in situ hybridization. Genomics 30: 629–630, 1995.

[120870]21374.

[120871]Parker, E. M.; Izzarelli, D. G.; Nowak, H. P.; Mahle, C. D.; Iben,

[120872]L. G.; Wang, J.; Goldstein, M. E.: Cloning and characterization of

[120873]the rat GALR1 galanin receptor from Rin14B insulinoma cells. Molec.

[120874]Brain Res. 34: 179–189, 1995.

[120875]21375.

[120876]Simoneaux, D. K.; Leach, R. J.; O'Connell, P.: Galanin receptor

[120877]1 gene (Galnr1) is tightly linked to the myelin basic protein gene

[120878]on chromosome 18 in mouse. Mammalian Genome 8: 875 only, 1997.

[120879]21376.

[120880]Walli, R.; Schafer, H.; Morys-Wortmann, C.; Paetzold, G.; Nustede,

[120881]R.; Schmidt, W. E.: Identification and biochemical characterization
of the human brain galanin receptor. J. Molec. Endocr. 13:
347–356,
1994.
21377.

[120885]Futterer, A.; Kruppa, G.; Kramer, B.; Lemke, H.; Kronke,
M.: Molecular
cloning and characterization of human kinectin. Molec.
Biol. Cell 6:
161–170, 1995.
21378.

[120889]Plitz, T.; Pfeffer, K.: Intact lysosome transport and phago-
some
function despite kinectin deficiency. Molec. Cell. Biol. 21:
6044–6055,
2001.
21379.

[120893]Print, C. G.; Leung, E.; Harrison, J. E. B.; Watson, J. D.;
Krissansen,
G. W.: Cloning of a gene encoding a human leukocyte

protein characterised

[120895]by extensive heptad repeats. Gene 144: 221–228, 1994.

[120896]21380.

[120897]Print, C. G.; Morris, C. M.; Spurr, N. K.; Rooke, L.; Kris-
sansen,

[120898]G. W.: The CG-1 gene, a member of the kinectin and ES/
130 family,

[120899]maps to human chromosome band 14q22. Immunogenet-
ics 43: 227–229,

[120900]1996.

[120901]21381.

[120902]Rao, P. N.; Yu, H.; Hodge, R.; Pettenati, M. J.; Sheetz, M. P.

[120903]: Assignment of the human kinectin gene (KTN1), encod-
ing a kinesin-binding

[120904]protein, to chromosome 14 band q22.1 by in situ hy-
bridization. Cytogenet.

[120905]Cell Genet. 79: 196–197, 1997.

[120906]21382.

[120907]Yu, H.; Nicchitta, C. V.; Kumar, J.; Becker, M.; Toyoshima,
I.;

[120908]Sheetz, M. P.: Characterization of kinectin, a kinesin-

binding protein:

[120909]primary sequence and N-terminal topogenic signal analysis. *Molec.*

[120910]*Biol. Cell* 6: 171-183, 1995.

[120911]21383.

[120912]Gordenin, D. A.; Kunkel, T. A.; Resnick, M. A.: Repeat expansion--all

[120913]in a flap? *Nature Genet.* 16: 116-118, 1997.

[120914]21384.

[120915]Greene, A. L.; Snipe, J. R.; Gordenin, D. A.; Resnick, M. A.:

[120916]Functional analysis of human FEN1 in *Saccharomyces cerevisiae* and

[120917]its role in genome stability. *Hum. Molec. Genet.* 8: 2263-2273, 1999.

[120918]21385.

[120919]Nakatsuru, S.; Sudo, K.; Nakamura, Y.: Isolation and mapping of

[120920]a human gene (MCM2) encoding a product homologous to yeast proteins

[120921]involved in DNA replication. *Cytogenet. Cell Genet.* 68: 226-230,

[120922]1995.

[120923]21386.

[120924]Augusseau, S.; Jouk, S.; Jalbert, P.; Prieur, M.: DiGeorge syndrome

[120925]and 22q11 rearrangements. (Letter) Hum. Genet. 74: 206 only, 1986.

[120926]21387.

[120927]Van Esch, H.; Groenen, P.; Daw, S.; Poffyn, A.; Holvoet, M.; Scambler,

[120928]P.; Fryns, J.-P.; Van de Ven, W.; Devriendt, K.: Partial DiGeorge

[120929]syndrome in two patients with a 10p rearrangement. Clin. Genet. 55:

[120930]269-276, 1999.

[120931]21388.

[120932]Wadey, R.; Daw, S.; Taylor, C.; Atif, U.; Kamath, S.; Halford,

[120933]S.; O'Donnell, H.; Wilson, D.; Goodship, J.; Burn, J.; Scambler, P.

[120934]: Isolation of a gene encoding an integral membrane protein from the

[120935]vicinity of a balanced translocation breakpoint associated

with DiGeorge

[120936]syndrome. Hum. Molec. Genet. 4: 1027–1033, 1995.

[120937]21389.

[120938]Zhao, Z.; Lee, C.-C.; Jiralerspong, S.; Juyal, R. C.; Lu, F.;
Baldini,

[120939]A.; Greenberg, F.; Caskey, C. T.; Patel, P. I.: The gene for a
human

[120940]microfibril-associated glycoprotein is commonly deleted
in Smith–Magenis

[120941]syndrome patients. Hum. Molec. Genet. 4: 589–597,
1995.

[120942]21390.

[120943]Contractor, A.; Rogers, C.; Maron, C.; Henkemeyer, M.;
Swanson,

[120944]G. T.; Heinemann, S. F.: Trans-synaptic Eph receptor-
ephrin signaling

[120945]in hippocampal mossy fiber LTP. Science 296: 1864–1869,
2002.

[120946]21391.

[120947]Tang, X. X.; Biegel, J. A.; Nycum, L. M.; Yoshioka, A.;
Brodeur,

[120948]G. M.; Pleasure, D. E.; Ikegaki, N.: cDNA cloning, molecular

characterization,

[120949]and chromosomal localization of NET (EPHT2), a human
EPH-related receptor

[120950]protein-tyrosine kinase gene preferentially expressed in
brain. :Genomics 29:

[120951]426-437, 1995.

[120952]21392.

[120953]Kamp, C.; Hirschmann, P.; Voss, H.; Huellen, K.; Vogt, P.
H.:

[120954]Two long homologous retroviral sequence blocks in prox-
imal Yq11 cause

[120955]AZFa microdeletions as a result of intrachromosomal re-
combination

[120956]events. Hum. Molec. Genet. 9: 2563-2572, 2000.

[120957]21393.

[120958]Krausz, C.; Meyts, E. R.-D.; Frydelund-Larson, L.; Quin-
tana-Murci,

[120959]L.; McElreavey, K.; Skakkebaek, N. E.: Double-blind Y
chromosome

[120960]microdeletion analysis in men with known sperm parame-
ters and reproductive

[120961]hormone profiles: microdeletions are specific for sper-

matogenic failure. J.

[120962]Clin. Endocr. Metab. 86: 2638–2642, 2001.

[120963]21394.

[120964]Simpson, E.; Chandler, P.; Goulmy, E.; Ma, K.; Hargreave, T. B.;

[120965]Chandley, A. C.: Loss of the 'azoospermia factor' (AZF) on Yq in

[120966]man is not associated with loss of HYA. Hum. Molec. Genet. 2: 469–471,

[120967]1993.

[120968]21395.

[120969]Sun, C.; Skaletsky, H.; Rozen, S.; Gromoll, J.; Nieschlag, E.;

[120970]Oates, R.; Page, D. C.: Deletion of azoospermia factor a (AZFa) region

[120971]of human Y chromosome caused by recombination between HERV15 proviruses. Hum.

[120972]Molec. Genet. 9: 2291–2296, 2000.

[120973]21396.

[120974]Krausz, C.; Quintana–Murci, L.; Barboux, S.; Siffroi, J.–P.; Rouba,

[120975]H.; Delafontaine, D.; Souleyreau–Therville, N.; Arvis, G.;

Antoine,

[120976]J. M.; Erdei, E.; Taar, J. P.; Tar, A.; Jeandidier, E.; Plessis, G.;

[120977]Bourgeron, T.; Dadoune, J.-P.; Fellous, M.; McElreavey, K.:
A high

[120978]frequency of Y chromosome deletions in males with noni-
diopathic infertility. J.

[120979]Clin. Endocr. Metab. 84: 3606–3612, 1999.

[120980]21397.

[120981]Ma, K.; Inglis, J. D.; Sharkey, A.; Bickmore, W. A.; Hill, R.

[120982]E.; Prosser, E. J.; Speed, R. M.; Thomson, E. J.; Jobling, M.;
Taylor,

[120983]K.; Wolfe, J.; Cooke, H. J.; Hargreave, T. B.; Chandley, A.
C.: A

[120984]Y chromosome gene family with RNA-binding protein ho-
mology: candidates

[120985]for the azoospermia factor AZF controlling human sper-
matogenesis. Cell 75:

[120986]1287–1295, 1993.

[120987]21398.

[120988]Vergnaud, G.; Page, D. C.; Simmler, M. C.; Brown, L.;

Rouyer,

[120989]F.; Noel, B.; Botstein, D.; de la Chapelle, A.; Weissenbach,
J.:

[120990]A deletion map of the human Y chromosome based on
DNA hybridization. Am.

[120991]J. Hum. Genet. 38: 109–124, 1986.

[120992]21399.

[120993]Vogt, P.; Chandley, A. C.; Hargreave, T. B.; Keil, R.; Ma, K.;

[120994]Sharkey, A.: Microdeletions in interval 6 of the Y chromo-
some of

[120995]males with idiopathic sterility point to disruption of AZF, a
human

[120996]spermatogenesis gene. Hum. Genet. 89: 491–496, 1992.

[120997]21400.

[120998]Vogt, P. H.; Edelmann, A.; Kirsch, S.; Henegariu, O.;
Hirschmann,

[120999]P.; Kiesewetter, F.; Kohn, F. M.; Schill, W. B.; Farah, S.;
Ramos,

[121000]C.; Hartmann, M.; Hartschuh, W.; Meschede, D.; Behre, H.
M.; Castel,

[121001]A.; Nieschlag, E.; Weidner, W.; Grone, H.-J.; Jung, A.; En-

gel, W.;

[121002]Haidl, G.: Human Y chromosome azoospermia factors
(AZF) mapped to

[121003]different subregions in Yq11. Hum. Molec. Genet. 5:
933-943, 1996.

[121004]21401.

[121005]Hawkins, J. R.: Mutational analysis of SRY in XY females.
Hum.

[121006]Mutat. 2: 347-350, 1993.

[121007]21402.

[121008]Hawkins, J. R.; Taylor, A.; Berta, P.; Levilliers, J.; Van der

[121009]Auwera, B.; Goodfellow, P. N.: Mutational analysis of SRY:
nonsense

[121010]and missense mutations in XY sex reversal. Hum. Genet.
88: 471-474,

[121011]1992.

[121012]21403.

[121013]Rohen, C.; Caselitz, J.; Stern, C.; Wanschura, S.; Schoen-
makers,

[121014]E. F.; Van de Ven, W. J.; Barnitzke, S.; Bullerdiek, J.: A
hamartoma

- [121015] of the breast with an aberration of 12q mapped to the
MAR region by
[121016] fluorescence in situ hybridization. Genes Chromosomes
Cancer 84:
[121017] 82–84, 1995.
[121018] 21404.
- [121019] Schoenmakers, E. F. P. M.; Huysmans, C.; Van de Ven, W. J.
M.
[121020]: Allelic knockout of novel splice variants of human re-
combination
[121021] repair gene RAD51B in t(12;14) uterine leiomyomas. Can-
cer Res. 59:
[121022] 19–23, 1999.
[121023] 21405.
- [121024] Zhou, X.; Benson, K. F.; Ashar, H. R.; Chada, K.: Mutation
responsible
[121025] for the mouse pygmy phenotype in the developmentally
regulated factor
[121026] HMGI-C. Nature 377: 771–774, 1995.
[121027] 21406.
- [121028] Petit, M. M. R.; Mols, R.; Schoenmakers, E. F. P. M.; Man-

dahl,

[121029]N.; Van de Ven, W. J. M.: LPP, the preferred fusion partner
gene

[121030]of HMGIC in lipomas, is a novel member of the LIM protein
gene family. Genomics 36:

[121031]118–129, 1996.

[121032]21407.

[121033]Burgess, D. L.; Kohrman, D. C.; Galt, J.; Plummer, N. W.;
Jones,

[121034]J. M.; Spear, B.; Meisler, M. H.: Mutation of a new sodium
channel

[121035]gene, Scn8a, in the mouse mutant 'motor endplate dis-
ease'. Nature

[121036]Genet. 10: 461–465, 1995.

[121037]21408.

[121038]DeRepentigny, Y.; Cote, P. D.; Pool, M.; Bernier, G.; Girard,
S.;

[121039]Vidal, S. M.; Kothary, R.: Pathological and genetic analysis
of the

[121040]degenerating muscle (dmu) mouse: a new allele of Scn8a.
Hum. Molec.

[121041]Genet. 10: 1819–1827, 2001.

[121042]21409.

[121043]Kohrman, D. C.; Harris, J. B.; Meisler, M. H.: Mutation detection

[121044]in the med and med(j) alleles of the sodium channel
Scn8a: unusual

[121045]splicing due to a minor class AT-AC intron. J. Biol. Chem.
271:

[121046]17576-17581, 1996.

[121047]21410.

[121048]Kohrman, D. C.; Plummer, N. W.; Schuster, T.; Jones, J. M.;
Jang,

[121049]W.; Burgess, D. L.; Galt, J.; Spear, B. T.; Meisler, M. H.: In-
sertional

[121050]mutation of the motor endplate disease (med) locus on
mouse chromosome

[121051]15. Genomics 26: 171-177, 1995.

[121052]21411.

[121053]Kohrman, D. C.; Smith, M. R.; Goldin, A. L.; Harris, J.;
Meisler,

[121054]M. H.: A missense mutation in the sodium channel Scn8a
is responsible

[121055]for cerebellar ataxia in the mouse mutant jolting. J. Neurosci. 16:

[121056]5993–5999, 1996.

[121057]21412.

[121058]Meisler, M. H.; Sprunger, L. K.; Plummer, N. W.; Escayg, A.; Jones,

[121059]J. M.: Ion channel mutations in mouse models of inherited neurological

[121060]disease. Ann. Med. 29: 569–574, 1997.

[121061]21413.

[121062]Plummer, N. W.; Galt, J.; Jones, J. M.; Burgess, D. L.; Sprunger,

[121063]L. K.; Kohrman, D. C.; Meisler, M. H.: Exon organization, coding

[121064]sequence, physical mapping, and polymorphic intragenic markers for

[121065]the human neuronal sodium channel gene SCN8A. Genomics 54: 287–296,

[121066]1998.

[121067]21414.

[121068]Plummer, N. W.; McBurney, M. W.; Meisler, M. H.: Alterna-

tive splicing

[121069]of the sodium channel SCN8A predicts a truncated two-domain protein

[121070]in fetal brain and non-neuronal cells. J. Biol. Chem. 272: 24008–24015,

[121071]1997.

[121072]21415.

[121073]Reid, E.; Escayg, A.; Dearlove, A. M.; Lee, D. D.; Meisler, M.

[121074]H.; Rubinsztein, D. C.: The spastic paraplegia SPG10 locus: narrowing

[121075]of critical region and exclusion of sodium channel gene SCN8A as a

[121076]candidate. J. Med. Genet. 38: 65–67, 2001.

[121077]21416.

[121078]Sprunger, L. K.; Escayg, A.; Tallaksen–Greene, S.; Albin, R. L.;

[121079]Meisler, M. H.: Dystonia associated with mutation of the neuronal

[121080]sodium channel Scn8a and identification of the modifier locus Scnm1

[121081]on mouse chromosome 3. Hum. Molec. Genet. 8:

471-479, 1999.

[121082]21417.

[121083]Casey, J. L.; Di Jeso, B.; Rao, K.; Klausner, R. D.; Harford, J.

[121084]B.: Two genetic loci participate in the regulation by iron of
the

[121085]gene for the human transferrin receptor. Proc. Nat. Acad.
Sci. 85:

[121086]1787-1791, 1988.

[121087]21418.

[121088]Enns, C. A.; Suomalainen, H. A.; Gebhardt, J. E.; Schroder,
J.;

[121089]Sussman, H. H.: Human transferrin receptor: expression of
the receptor

[121090]is assigned to chromosome 3. Proc. Nat. Acad. Sci. 79:
3241-3245,

[121091]1982.

[121092]21419.

[121093]Gareau, R.; Gagnon, M. G.; Thellend, C.; Chenard, C.; Au-
dran, M.;

[121094]Chanal, J.-L.; Ayotte, C.; Brisson, G. R.: Transferrin soluble
receptor:

[121095]a possible probe for detection of erythropoietin abuse by athletes. Horm.

[121096]Metab. Res. 26: 311–312, 1994.

[121097]21420.

[121098]Goodfellow, P. N.; Banting, G.; Sutherland, R.; Greaves, M.; Solomon,

[121099]E.; Povey, S.: Expression of human transferrin receptor is controlled

[121100]by a gene on chromosome 3: assignment using species specificity of

[121101]a monoclonal antibody. Somat. Cell Genet. 8: 197–206, 1982.

[121102]21421.

[121103]Larrick, J. W.; Hyman, E. S.: Acquired iron-deficiency anemia

[121104]caused by an antibody against the transferrin receptor. New Eng.

[121105]J. Med. 311: 214–218, 1984.

[121106]21422.

[121107]Levy, J. E.; Jin, O.; Fujiwara, Y.; Kuo, F.; Andrews, N. C.: Transferrin

[121108]receptor is necessary for development of erythrocytes and the nervous

[121109]system. Nature Genet. 21: 396–399, 1999.

[121110]21423.

[121111]Miller, Y. E.; Jones, C.; Scoggin, C.; Morse, H.; Seligman, P.

[121112]; Chromosome 3q (22–ter) encodes the human transferrin receptor. Am.

[121113]J. Hum. Genet. 35: 573–583, 1983.

[121114]21424.

[121115]Newman, R.; Schneider, C.; Sutherland, R.; Vodinelich, L.; Greaves,

[121116]M.: The transferrin receptor. Trends Biochem. Sci. 7: 397–400,

[121117]1982.

[121118]21425.

[121119]Nikinmaa, B.; Schroder, J.: Two antigens, the transferrin receptor

[121120]and p90 assigned to human chromosome 3, are probably the same protein. Hereditas 107:

[121121]55–58, 1987.

[121122]21426.

[121123]Omary, M. B.; Trowbridge, I. S.: Biosynthesis of the human transferrin

[121124]receptor in cultured cells. J. Biol. Chem. 256: 12888–12892, 1981.

[121125]21427.

[121126]Kashuba, V. I.; Gizatullin, R. Z.; Protopopov, A. I.; Allik-mets,

[121127]R.; Korolev, S.; Li, J.; Boldog, F.; Tory, K.; Zabarovska, V.; Marcsek,

[121128]Z.; Sumegi, J.; Klein, G.; Zabarovsky, E. R.; Kisselev, L.: NotI

[121129]linking/jumping clones of human chromosome 3: mapping of the TFRC,

[121130]RAB7 and HAUSP genes to regions rearranged in leukemia and deleted

[121131]in solid tumors. FEBS Lett. 419: 181–185, 1997.

[121132]21428.

[121133]Rabin, M.; McClelland, A.; Kuhn, L.; Ruddle, F. H.: Regional

[121134]localization of the human transferrin receptor gene to 3q26.2–qter. Am.

[121135]J. Hum. Genet. 37: 1112–1116, 1985.

[121136]21429.

[121137]Schneider, C.; Kurkinen, M.; Greaves, M.: Isolation of cDNA clones

[121138]for the human transferrin receptor. EMBO J. 2:
2259–2263, 1983.

[121139]21430.

[121140]Schneider, C.; Owen, M. J.; Banville, D.; Williams, J. G.: Primary

[121141]structure of human transferrin receptor deduced from the mRNA sequence. Nature 311:

[121142]675–678, 1984.

[121143]21431.

[121144]Valenzuela, C. Y.; Avendano, A.; Harb, Z.: Association between

[121145]Rh and plasma iron binding (transferrin). Hum. Genet. 87:
438–440,

[121146]1991.

[121147]21432.

[121148]Vodinelich, L.; Sutherland, R.; Schneider, C.; Newman, R.; Greaves,

[121149]M.: Receptor for transferrin may be a 'target' structure for natural

[121150]killer cells. Proc. Nat. Acad. Sci. 80: 835–839, 1983.

[121151]21433.

[121152]de Martinville, B.; Francke, U.: The c-Ha-ras1, insulin and beta-globin

[121153]loci map outside the deletion associated with aniridia–Wilms' tumour. Nature 305:

[121154]641–643, 1983.

[121155]21434.

[121156]Hamm, C. W.; Ravkilde, J.; Gerhardt, W.; Jorgensen, P.; Peheim,

[121157]E.; Ljungdahl, L.; Goldmann, B.; Katus, H. A.: The prognostic value

[121158]of serum troponin T in unstable angina. New Eng. J. Med. 327: 146–150,

[121159]1992.

[121160]21435.

[121161]Carmeliet, P.; Ferreira, V.; Breler, G.; Pollefeyt, S.; Kleckens,

[121162]L.; Gertsenstein, M.; Fahrig, M.; Vandenhoeck, A.; Harpal,

K.; Eberhardt,

[121163]C.; Declercq, C.; Pawling, J.; Moons, L.; Collen, D.; Risau,
W.; Nagy,

[121164]A.: Abnormal blood vessel development and lethality in
embryos lacking

[121165]a single VEGF allele. Nature 380: 435–439, 1996.

[121166]21436.

[121167]Martin, L. H.; Calabi, F.; Lefebvre, F.–A.; Bilsland, C. A. G.;

[121168]Milstein, C.: Structure and expression of the human thymocyte antigens

[121169]CD1a, CD1b, and CD1c. Proc. Nat. Acad. Sci. 84:
9189–9193, 1987.

[121170]21437.

[121171]Martin, L. H.; Calabi, F.; Milstein, C.: Isolation of CD1
genes:

[121172]a family of major histocompatibility complex–related differentiation

[121173]antigens. Proc. Nat. Acad. Sci. 83: 9154–9158, 1986.

[121174]21438.

[121175]Moseley, W. S.; Watson, M. L.; Kingsmore, S. F.; Seldin, M.
F.

- [121176]: CD1 defines conserved linkage group border between human chromosomes
- [121177]1 and mouse chromosomes 1 and 3. Immunogenetics 30: 378–382, 1989.
- [121178]21439.
- [121179]Samson, F.; de Jong, P. J.; Trask, B. J.; Koza–Taylor, P.; Speer,
- [121180]M. C.; Potter, T.; Roses, A. D.; Gilbert, J. R.: Assignment of the
- [121181]human slow skeletal troponin T gene to 19q13.4 using somatic cell
- [121182]hybrids and fluorescence in situ hybridization analysis. Genomics 13:
- [121183]1374–1375, 1992.
- [121184]21440.
- [121185]Samson, F.; Gilbert, J. R.; Lee, J. E.; Potter, T. G.; Koza–Taylor,
- [121186]P.; Speer, M. C.; Bachinski, L. L.; Siciliano, M. J.; Roses, A. D.
- [121187]: Isolation and localization of a human slow troponin T gene on chromosome
- [121188]19q. (Abstract) Cytogenet. Cell Genet. 58: 2025, 1991.

[121189]21441.

[121190]Samson, F.; Lee, J. E.; Hung, W. Y.; Potter, T. G.; Herbstreith,

[121191]M.; Roses, A. D.; Gilbert, J. R.: Isolation and localisation of
a

[121192]slow troponin (TNT) gene on chromosome 19 by subtraction of a cDNA

[121193]muscle library using myotonic muscle cDNA. J. Neurosci.
Res. 27:

[121194]441–451, 1990.

[121195]21442.

[121196]Samson, F.; Mesnard, L.; Mihovilovic, M.; Potter, T. G.;
Mercadier,

[121197]J.-J.; Roses, A. D.; Gilbert, J. R.: A new human slow skeletal
troponin

[121198]T (TnTs) mRNA isoform derived from alternative splicing
of a single

[121199]gene. Biochem. Biophys. Res. Commun. 199: 841–847,
1994.

[121200]21443.

[121201]Johnston, J. J.; Kelley, R. I.; Crawford, T. O.; Morton, D. H.;

[121202]Agarwala, R.; Koch, T.; Schaffer, A. A.; Francomano, C. A.; Biesecker,

[121203]L. G.: A novel nemaline myopathy in the Amish caused by a mutation

[121204]in troponin T1. Am. J. Hum. Genet. 67: 814–821, 2000.

[121205]21444.

[121206]Novelli, G.; Gennarelli, M.; Rocchi, M.; Dallapiccola, B.: Assignment

[121207]of the slow troponin T (TNNT1) gene to chromosome 19 using polymerase

[121208]chain reaction. Hum. Genet. 88: 697–698, 1992.

[121209]21445.

[121210]Novelli, G.; Gennarelli, M.; Zelano, G.; Sangiuolo, F.; Rocchi,

[121211]M.; Dallapiccola, B.: Isolation and mapping of the slow skeletal

[121212]troponin T using the polymerase chain reaction. (Abstract) Cytogenet.

[121213]Cell Genet. 58: 2023, 1991.

[121214]21446.

[121215]Guenet, J.-L.; Simon-Chazottes, D.; Gravel, M.; Hastings,

K. E.

[121216]M.; Schiaffino, S.: Cardiac and skeletal muscle troponin I isoforms

[121217]are encoded by a dispersed gene family on mouse chromosomes 1 and

[121218]7. Mammalian Genome 7: 13–15, 1996.

[121219]21447.

[121220]Wade, R.; Eddy, R.; Shows, T. B.; Kedes, L.: cDNA sequence, tissue-specific

[121221]expression, and chromosomal mapping of the human slow-twitch skeletal

[121222]muscle isoform of troponin I. Genomics 7: 346–357, 1990.

[121223]21448.

[121224]Bhavsar, P. K.; Brand, N. J.; Yacoub, M. H.; Barton, P. J. R.:

[121225]Isolation and characterization of the human cardiac troponin I gene

[121226](TNNI3). Genomics 35: 11–23, 1996.

[121227]21449.

[121228]Kimura, A.; Harada, H.; Park, J.-E.; Nishi, H.; Satoh, M.; Takahashi,

[121229]M.; Hiroi, S.; Sasaoka, T.; Ohbuchi, N.; Nakamura, T.; Koyanagi, T.;

[121230]Hwang, T.-H.; Choo, J.; Chung, K.-S.; Hasegawa, A.; Nagai, R.; Okazaki,

[121231]O.; Nakamura, H.; Matsuzaki, M.; Sakamoto, T.; Toshima, H.; Koga,

[121232]Y.; Imaizumi, T.; Sasazuki, T.: Mutations in the cardiac troponin

[121233]I gene associated with hypertrophic cardiomyopathy. Nature Genet. 16:

[121234]379–382, 1997.

[121235]21450.

[121236]Labarrere, C. A.; Nelson, D. R.; Cox, C. J.; Pitts, D.; Kirlin,

[121237]P.; Halbrook, H.: Cardiac-specific troponin I levels and risk of

[121238]coronary artery disease and graft failure following heart transplantation. J.A.M.A. 284:

[121239]457–464, 2000.

[121240]21451.

[121241]Lang, R.; Gomes, A. V.; Zhao, J.; Housmans, P. R.; Miller, T.;

- [121242]Potter, J. D.: Functional analysis of a troponin I (R145G) mutation
- [121243]associated with familial hypertrophic cardiomyopathy. J. Biol. Chem. 277: 11670–11678, 2002.
- [121244]21452.
- [121245]MacGeoch, C.; Barton, P. J. R.; Vallins, W. J.; Bhavsar, P.; Spurr, N. K.: The human cardiac troponin I locus: assignment to 19p13.2–19q13.2. Hum. Genet. 88: 101–104, 1991.
- [121246]21453.
- [121247]Mogensen, J.; Kruse, T. A.; Borglum, A. D.: Assignment of the human cardiac troponin I gene (TNNI3) to chromosome 19q13.4 by radiation hybrid mapping. Cytogenet. Cell Genet. 79: 272–273, 1997.
- [121248]21454.
- [121249]Crow, R. S.: Peripheral neuritis in myelomatosis. Brit. Med. J. 2: 802–804, 1956.

[121256]21455.

[121257]Dantz, D.; Bewersdorf, J.; Fruehwald-Schultes, B.; Kern, W.; Jelkmann,

[121258]W.; Born, J.; Fehm, H. L.; Peters, A.: Vascular endothelial growth

[121259]factor: a novel endocrine defensive response to hypoglycemia. J.

[121260]Clin. Endocr. Metab. 87: 835–840, 2002.

[121261]21456.

[121262]Ferrara, N.; Carver-Moore, K.; Chen, H.; Dowd, M.; Lu, L.; O'Shea,

[121263]K. S.; Powell-Braxton, L.; Hillan, K. J.; Moore, M. W.: Heterozygous

[121264]embryonic lethality induced by targeted inactivation of the VEGF gene. Nature 380:

[121265]439–442, 1996.

[121266]21457.

[121267]Folkman, J.: Angiogenesis in cancer, vascular, rheumatoid and

[121268]other disease. Nature Med. 1: 27–31, 1995.

[121269]21458.

[121270]Fukumura, D.; Xavier, R.; Sugiura, T.; Chen, Y.; Park, E.-C.;
Lu,

[121271]N.; Selig, M.; Nielsen, G.; Taksir, T.; Jain, R. K.; Seed, B.:
Tumor

[121272]induction of VEGF promoter activity in stromal cells. Cell
94: 715–725,

[121273]1998.

[121274]21459.

[121275]Gerber, H.-P.; Malik, A. K.; Solar, G. P.; Sherman, D.;
Liang,

[121276]X. H.; Meng, G.; Hong, K.; Marsters, J. C.; Ferrara, N.: VEGF
regulates

[121277]haematopoietic stem cell survival by an internal autocrine loop mechanism. Nature 417: 954–958, 2002.

[121279]21460.

[121280]Gerber, H.-P.; Vu, T. H.; Ryan, A. M.; Kowalski, J.; Werb, Z.; Ferrara, N.: VEGF couples hypertrophic cartilage remodeling, ossification and angiogenesis during endochondral bone formation. Nature Med. 5: 623–628, 1999.

[121284]21461.

[121285]Giordano, F. J.; Gerber, H.-P.; Williams, S.-P.; VanBruggen, N.; Bunting, S.; Ruiz-Lozano, P.; Gu, Y.; Nath, A. K.; Huang, Y.; Hickey, R.; Dalton, N.; Peterson, K. L.; Ross, J., Jr.; Chien, K. R.; Ferrara, N.: A cardiac myocyte vascular endothelial growth factor paracrine pathway is required to maintain cardiac function. Proc. Nat. Acad.

[121290]Sci. 98: 5780–5785, 2001.

[121291]21462.

[121292]Helmlinger, G.; Endo, M.; Ferrara, N.; Hlatky, L.; Jain, R. K.

[121293]: Formation of endothelial cell networks. Nature 405:
139–141, 2000.

[121294]21463.

[121295]Hofman, P.; van Blijswijk, B. C.; Gaillard, P. J.; Vrensen, G.

[121296]F. J. M.; Schlingemann, R. O.: Endothelial cell hypertrophy
induced

[121297]by vascular endothelial growth factor in the retina: new
insights

[121298]into the pathogenesis of capillary nonperfusion. Arch.
Ophthalm. 119:

[121299]861–866, 2001.

[121300]21464.

[121301]Holash, J.; Maisonpierre, P. C.; Compton, D.; Boland, P.;
Alexander,

[121302]C. R.; Zagzag, D.; Yancopoulos, G. D.; Wiegand, S. J.: Ves-
sel cooption,

[121303]regression, and growth in tumors mediated by angiopo-
ietins and VEGF. Science 284:

[121304]1994–1998, 1999.

[121305]21465.

[121306]Jin, K.; Zhu, Y.; Sun, Y.; Mao, X. O.; Xie, L.; Greenberg, D.

[121307]A.: Vascular endothelial growth factor (VEGF) stimulates
neurogenesis

[121308]in vitro and in vivo. Proc. Nat. Acad. Sci. 99:
11946–11950, 2002.

[121309]21466.

[121310]Mattei, M.–G.; Borg, J.–P.; Rosnet, O.; Marme, D.; Birn-
baum, D.

[121311]: Assignment of vascular endothelial growth factor (VEGF)
and placenta

[121312]growth factor (PIGF) genes to human chromosome
6p12–p21 and 14q24–q31

[121313]regions, respectively. Genomics 32: 168–169, 1996.

[121314]21467.

[121315]Miralles, G. D.; O'Fallon, J. R.; Talley, N. J.: Plasma-cell

[121316]dyscrasia with polyneuropathy: the spectrum of POEMS
syndrome. New

[121317]Eng. J. Med. 327: 1919–1923, 1992.

[121318]21468.

[121319]Mueller, M. D.; Vigne, J.-L.; Minchenko, A.; Lebovic, D. I.;
Leitman,

[121320]D. C.; Taylor, R. N.: Regulation of vascular endothelial
growth factor

[121321](VEGF) gene transcription by estrogen receptors alpha and
beta. Proc.

[121322]Nat. Acad. Sci. 97: 10972-10977, 2000.

[121323]21469.

[121324]Nakanishi, T.; Sobue, I.; Toyokura, Y.; Nishitani, H.;
Kuroiwa,

[121325]Y.; Satoyoshi, E.; Tsubaki, T.; Igata, A.; Ozaki, Y.: The
Crow-Fukase

[121326]syndrome: a study of 102 cases in Japan. Neurology 34:
712-720,

[121327]1984.

[121328]21470.

[121329]Niimi, H.; Arimura, K.; Jonosono, M.; Hashiguchi, T.;
Kawabata,

[121330]M.; Osame, M.; Kitajima, I.: VEGF is causative for pul-
monary hypertension

[121331]in a patient with Crow–Fukase (POEMS) syndrome. Intern.

Med. 39:

[121332]1101–1104, 2000.

[121333]21471.

[121334]Oosthuysen, B.; Moons, L.; Storkebaum, E.; Beck, H.;

Nuyens, D.;

[121335]Brusselmans, K.; Van Dorpe, J.; Hellings, P.; Gorselink, M.;

Heymans,

[121336]S.; Theilmeier, G.; Dewerchin, M.; and 20 others: Deletion

of the

[121337]hypoxia–response element in the vascular endothelial

growth factor

[121338]promoter causes motor neuron degeneration. Nature

Genet. 28: 131–138,

[121339]2001.

[121340]21472.

[121341]Poltorak, Z.; Cohen, T.; Sivan, R.; Kandelis, Y.; Spira, G.;

Vlodavsky,

[121342]I.; Keshet, E.; Neufeld, G.: VEGF145, a secreted vascular

endothelial

[121343]growth factor isoform that binds to extracellular matrix. J.

Biol.

[121344]Chem. 272: 7151–7158, 1997.

[121345]21473.

[121346]Shimpo, S.: Solitary myeloma causing polyneuritis and endocrine

[121347]disorders. Nippon Rinsho 26: 2444–2456, 1968.

[121348]21474.

[121349]Soker, S.; Takashima, S.; Miao, H. Q.; Neufeld, G.; Klagsbrun,

[121350]M.: Neuropilin-1 is expressed by endothelial and tumor cells as an

[121351]isoform-specific receptor for vascular endothelial growth factor. Cell 92:

[121352]735–745, 1998.

[121353]21475.

[121354]Sone, H.; Kawakami, Y.; Sakauchi, M.; Nakamura, Y.; Takahashi,

[121355]A.; Shimano, H.; Okuda, Y.; Segawa, T.; Suzuki, H.; Yamada, N.: Neutralization

[121356]of vascular endothelial growth factor prevents collagen-induced arthritis

[121357]and ameliorates established disease in mice. Biochem.

Biophys. Res.

[121358]Commun. 281: 562–568, 2001.

[121359]21476.

[121360]Springer, M. L.; Chen, A. S.; Kraft, P. E.; Bednarski, M.;
Blau,

[121361]H. M.: VEGF gene delivery to muscle: potential role for
vasculogenesis

[121362]in adults. Molec. Cell 2: 549–558, 1998.

[121363]21477.

[121364]Thurston, G.; Suri, C.; Smith, K.; McClain, J.; Sato, T. N.;
Yancopoulos,

[121365]G. D.; McDonald, D. M.: Leakage-resistant blood vessels in
mice transgenically

[121366]overexpressing angiopoietin-1. Science 286: 2511–2514,
1999.

[121367]21478.

[121368]Tischer, E.; Mitchell, R.; Hartman, T.; Silva, M.; Gospo-
darowicz,

[121369]D.; Fiddes, J. C.; Abraham, J. A.: The human gene for vas-
cular endothelial

[121370]growth factor: multiple protein forms are encoded

through alternative

[121371]exon splicing. J. Biol. Chem. 266: 11947–11954, 1991.

[121372]21479.

[121373]Ahlgren, A.; Yanase, T.; Simpson, E. R.; Winter, J. S. D.;
Waterman,

[121374]M. R.: Compound heterozygous mutations (arg239-to-ter,
pro342-to-thr)

[121375]in the CYP17 (P45017-alpha) gene lead to ambiguous ex-
ternal genitalia

[121376]in a male patient with partial combined
17-alpha-hydroxylase/17,20-lyase

[121377]deficiency. J. Clin. Endocr. Metab. 74: 667–672, 1992.

[121378]21480.

[121379]Auchus, R. J.; Miller, W. L.: Molecular modeling of human
P450c17

[121380](17-alpha-hydroxylase/17,20-lyase): insights into reac-
tion mechanisms

[121381]and effects of mutations. Molec. Endocr. 13: 1169–1182,
1999.

[121382]21481.

[121383]Biason, A.; Mantero, F.; Scaroni, C.; Simpson, E. R.; Water-

man,

[121384]M. R.: Deletion within the CYP17 gene together with insertion of

[121385]foreign DNA is the cause of combined complete 17-alpha-hydroxylase/17,20-lyase

[121386]deficiency in an Italian patient. Molec. Endocr. 5: 2037-2045, 1991.

[121387]21482.

[121388]Biaison-Lauber, A.; Kempken, B.; Werder, E.; Forest, M. G.; Einaudi,

[121389]S.; Ranke, M. B.; Matsuo, N.; Brunelli, V.; Schonle, E. J.; Zachmann,

[121390]M.: 17-alpha-hydroxylase/17,20-lyase deficiency as a model to study

[121391]enzymatic activity regulation: role of phosphorylation. J. Clin.

[121392]Endocr. Metab. 85: 1226-1231, 2000.

[121393]21483.

[121394]Biaison-Lauber, A.; Leiberman, E.; Zachmann, M.: A single amino

[121395]acid substitution in the putative redox partner-binding site of P450c17

[121396] as cause of isolated 17,20-lyase deficiency. J. Clin. Endocr. Metab. 82:

[121397] 3807–3812, 1997.

[121398] 21484.

[121399] Biglieri, E.G.: A prismatic case: 17- α -hydroxylase deficiency:

[121400] 1963–1966. J. Clin. Endocr. Metab. 82: 48–50, 1997.

[121401] 21485.

[121402] Osborne, L. R.; Martindale, D.; Scherer, S. W.; Shi, X.-M.; Huizenga,

[121403] J.; Heng, H. H. Q.; Costa, T.; Pober, B.; Lew, L.; Brinkman, J.; Rommens,

[121404] J.; Koop, B.; Tsui, L.-C.: Identification of genes from a 500-kb

[121405] region at 7q11.23 that is commonly deleted in Williams syndrome patients. Genomics 36:

[121406] 328–336, 1996.

[121407] 21486.

[121408] Rampoldi, L.; Dobson-Stone, C.; Rubio, J. P.; Danek, A.; Chalmers,

[121409] R. M.; Wood, N. W.; Verellen, C.; Ferrer, X.; Malandrini, A.;

Fabrizi,

[121410]G. M.; Brown, R.; Vance, J.; Pericak-Vance, M.; Rudolf, G.;
Carre,

[121411]S.; Alonso, E.; Manfredi, M.; Nemeth, A. H.; Monaco, A. P.:
A conserved

[121412]sorting-associated protein is mutant in chorea-
acanthocytosis. Nature

[121413]Genet. 28: 119-120, 2001.

[121414]21487.

[121415]Belden, W. J.; Barlowe, C.: Role of Erv29p in collecting sol-
uble

[121416]secretory proteins into ER-derived transport vesicles. Sci-
ence 294:

[121417]1528-1531, 2001.

[121418]21488.

[121419]Duhig, T.; Ruhrberg, C.; Mor, O.; Fried, M.: The human
surfeit

[121420]locus. Genomics 52: 72-78, 1998.

[121421]21489.

[121422]Reeves, J. E.; Fried, M.: The surf-4 gene encodes a novel
30 kDa

- [121423]integral membrane protein. Molec. Membr. Biol. 12:
201–208, 1995.
- [121424]21490.
- [121425]Archer, B. T., III; Ozcelik, T.; Jahn, R.; Francke, U.; Sudhof,
[121426]T. C.: Structures and chromosomal localizations of two
human genes
[121427]encoding synaptobrevins 1 and 2. J. Biol. Chem. 265:
17267–17273,
[121428]1990.
- [121429]21491.
- [121430]Hu, K.; Carroll, J.; Fedorovich, S.; Rickman, C.; Sukhodub,
A.;
- [121431]Davietov, B.: Vesicular restriction of synaptobrevin sug-
gests a role
[121432]for calcium in membrane fusion. Nature 415: 646–650,
2002.
- [121433]21492.
- [121434]Doggen, C. J. M.; Kunz, G.; Rosendaal, F. R.; Lane, D. A.;
Vos,
- [121435]H. L.; Stubbs, P. J.; Cats, V. M.; Ireland, H.: A mutation in
the

[121436]thrombomodulin gene, 127G to A coding for
ala25-to-thr, and the risk
[121437]of myocardial infarction in men. Thromb. Haemost. 80:
743-748, 1998.
[121438]21493.

[121439]Esmon, N. L.: Thrombomodulin. Semin. Thromb. Hemost.
13: 454-463,
[121440]1987.
[121441]21494.

[121442]Espinosa, R., III; Sadler, J. E.; Le Beau, M. M.: Regional lo-
calization
[121443]of the human thrombomodulin gene to 20p12-cen. Ge-
nomics 5: 649-650,
[121444]1989.
[121445]21495.

[121446]Franchi, F.; Biguzzi, E.; Cetin, I.; Facchetti, F.; Radaelli, T.;
[121447]Bozzo, M.; Pardi, G.; Faioni, E. M.: Mutations in the throm-
bomodulin
[121448]and endothelial protein C receptor genes in women with
late fetal
[121449]loss. Brit. J. Haemat. 114: 641-646, 2001.

[121450]21496.

[121451]Jackman, R. W.; Beeler, D. L.; Fritze, L.; Soff, G.; Rosenberg,

[121452]R. D.: Human thrombomodulin gene is intron depleted:
nucleic acid

[121453]sequences of the cDNA and gene predict protein structure
and suggest

[121454]sites of regulatory control. Proc. Nat. Acad. Sci. 84:
6425–6429,

[121455]1987.

[121456]21497.

[121457]Kunz, G.; Ireland, H. A.; Stubbs, P. J.; Kahan, M.; Coulton,
G.

[121458]C.; Lane, D. A.: Identification and characterization of a
thrombomodulin

[121459]gene mutation coding for an elongated protein with re-
duced expression

[121460]in a kindred with myocardial infarction. Blood 95:
569–576, 2000.

[121461]21498.

[121462]Ohlin, A.-K.; Marlar, R. A.: The first mutation identified in

[121463]the thrombomodulin gene in a 45-year-old man presenting with thromboembolic

[121464]disease. *Blood* 85: 330–336, 1995.

[121465]21499.

[121466]Okura, Y.; Kato, K.; Hanawa, H.; Izumi, T.; Kamishima, T.; Yamato,

[121467]Y.; Emura, I.; Shibata, A.: Pericardial mesothelioma secreting thrombomodulin. *Am.*

[121468]*Heart J.* 132: 1309–1311, 1996.

[121469]21500.

[121470]Wen, D.; Dittman, W. A.; Ye, R. D.; Deaven, L. L.; Majerus, P.

[121471]W.; Sadler, J. E.: Human thrombomodulin: complete cDNA sequence and

[121472]chromosome localization of the gene. *Biochemistry* 26: 4350–4357,

[121473]1987.

[121474]21501.

[121475]Grimm, C.; Wenzel, A.; Hafezi, F.; Yu, S.; Redmond, T. M.; Reme,

[121476]C. E.: Protection of Rpe65-deficient mice identifies

rhodopsin as

[121477]a mediator of light-induced retinal degeneration. Nature
Genet. 25:

[121478]63–66, 2000.

[121479]21502.

[121480]Barone, V.; Bertocchini, F.; Bottinelli, R.; Protasi, F.; Allen,

[121481]P. D.; Armstrong, C. F.; Reggiani, C.; Sorrentino, V.: Con-
tractile

[121482]impairment and structural alterations of skeletal muscles
from knockout

[121483]mice lacking type 1 and type 3 ryanodine receptors. FEBS
Lett. 422:

[121484]160–164, 1998.

[121485]21503.

[121486]Cavanna, J. S.; Greenfield, A. J.; Johnson, K. J.; Marks, A.
R.;

[121487]Nadal-Ginard, B.; Brown, S. D. M.: Establishment of the
mouse chromosome

[121488]7 region with homology to the myotonic dystrophy region
of human chromosome

[121489]19q. Genomics 7: 12–18, 1990.

[121490]21504.

[121491]Cheng, H.; Lederer, W. J.; Cannell, M. B.: Calcium sparks:
elementary

[121492]events underlying excitation-contraction coupling in heart
muscle. Science 262:

[121493]740-744, 1993.

[121494]21505.

[121495]Vallins, W. J.; Brand, N. J.; Dabhade, N.; Butler-Browne, G.;

[121496]Yacoub, M. H.; Barton, P. J. R.: Molecular cloning of human
cardiac

[121497]troponin I using polymerase chain reaction. FEBS Lett.
270: 57-61,

[121498]1990.

[121499]21506.

[121500]Vincenti, V.; Cassano, C.; Rocchi, M.; Persico, M. G.: As-
signment

[121501]of the vascular endothelial growth factor gene to human
chromosome

[121502]6p21.3. Circulation 93: 1493-1495, 1996.

[121503]21507.

- [121504]Biglieri, E. G.; Herron, M. A.; Brust, N.: 17-Hydroxylation deficiency
- [121505]in man. J. Clin. Invest. 45: 1946-1954, 1966.
- [121506]21508.
- [121507]Chung, B.-C.; Picado-Leonard, J.; Haniu, M.; Bienkowski, M.; Hall, M.; Hall, P. F.; Shively, J. E.; Miller, W. L.: Cytochrome P450c17 (steroid
- [121508]17 alpha-hydroxylase/17,20 lyase): cloning of human adrenal and testis
- [121509]cDNAs indicates the same gene is expressed in both tissues. Proc.
- [121510]Nat. Acad. Sci. 84: 407-411, 1987.
- [121511]21509.
- [121512]D'Armiento, M.; Reda, G.; Bisignani, G.; Tabolli, S.; Capel-laci,
- [121513]S.; Lulli, P.; Carbonara, A.; Biglieri, E. G.: No linkage between
- [121514]HLA and congenital adrenal hyperplasia due to 17-alpha-hydroxylase
- [121515]deficiency. (Letter) New Eng. J. Med. 308: 970-971, 1983.
- [121516]

[121517]21510.

[121518]Di Cerbo, A.; Biason-Laubert, A.; Savino, M.; Piemontese, M. R.;

[121519]Di Giorgio, A.; Perona, M.; Savoia, A.: Combined
17- α -hydroxylase/17,20-lyase

[121520]deficiency caused by phe93cys mutation in the CYP17
gene. J. Clin.

[121521]Endocr. Metab. 87: 898-905, 2002.

[121522]21511.

[121523]Fan, Y.-S.; Sasi, R.; Lee, C.; Winter, J. S. D.; Waterman, M.

[121524]R.; Lin, C. C.: Localization of the human CYP17 gene
(cytochrome

[121525]p450-17- α) to 10q24.3 by fluorescence in situ hybridization and

[121526]simultaneous chromosome banding. Genomics 14:
1110-1111, 1992.

[121527]21512.

[121528]Geller, D. H.; Auchus, R. J.; Mendonca, B. B.; Miller, W. L.:

[121529]The genetic and functional basis of isolated 17,20-lyase
deficiency. Nature

[121530]Genet. 17: 201-205, 1997.

[121531]21513.

[121532]Goldsmith, O.; Solomon, D. H.; Horton, R.: Hypogonadism
and mineralocorticoid

[121533]excess: 17-hydroxylase deficiency. New Eng. J. Med. 277:
673-677,

[121534]1967.

[121535]21514.

[121536]Gupta, M. K.; Geller, D. H.; Auchus, R. J.: Pitfalls in charac-
terizing

[121537]P450c17 mutations associated with isolated 17,20-lyase
deficiency. J.

[121538]Clin. Endocr. Metab. 86: 4416-4423, 2001.

[121539]21515.

[121540]Hanley, N. A.; Rainey, W. E.; Wilson, D. I.; Ball, S. G.;
Parker,

[121541]K. L.: Expression profiles of SF-1, DAX1, and CYP17 in the
human

[121542]fetal adrenal gland: potential interactions in gene regula-
tion. Molec.

[121543]Endocr. 15: 57-68, 2001.

[121544]21516.

[121545]Heremans, G. F. P.; Moolenaar, A. J.; Van Gelderen, H. H.:

Female

[121546]phenotype in a male child due to 17-alpha-hydroxylase
deficiency. Arch.

[121547]Dis. Child. 51: 721-723, 1976.

[121548]21517.

[121549]Imai, T.; Yanase, T.; Waterman, M. R.; Simpson, E. R.;
Pratt,

[121550]J. J.: Canadian Mennonites and individuals residing in the
Friesland

[121551]region of the Netherlands share the same molecular basis
of 17-alpha-hydroxylase

[121552]deficiency. Hum. Genet. 89: 95-96, 1992.

[121553]21518.

[121554]Jones, K. L.; Freidenberg, G. R.; Buchta, R.; Derenoncourt,
A.

[121555]: Male pseudohermaphroditism resulting from
17-alpha-monooxygenase

[121556](P-450C-17) deficiency in two unrelated Guamanians. Am.
J. Dis. Child. 146:

[121557]592-595, 1992.

[121558]21519.

[121559]Kagimoto, K.; Waterman, M. R.; Kagimoto, M.; Ferreira, P.;
Simpson,

[121560]E. R.; Winter, J. S. D.: Identification of a common molecular
basis

[121561]for combined 17-alpha-hydroxylase/17,20-lyase defi-
ciency in two Mennonite

[121562]families. Hum. Genet. 82: 285-286, 1989.

[121563]21520.

[121564]Kagimoto, M.; Winter, J. S. D.; Kagimoto, K.; Simpson, E.
R.;

[121565]Waterman, M. R.: Structural characterization of normal
and mutant

[121566]human steroid 17-alpha-hydroxylase genes: molecular
basis of one example

[121567]of combined 17-alpha-hydroxylase/17,20 lyase defi-
ciency. Molec. Endocr. 2:

[121568]564-570, 1988.

[121569]21521.

[121570]Krohn, K.; Uiho, R.; Aavik, E.; Peterson, P.; Savilahti, K.:

[121571]Identification of molecular cloning of an autoantigen as-

sociated with

[121572]Addison's disease as steroid 17- α -hydroxylase.

Lancet 339: 770-773,

[121573]1992.

[121574]21522.

[121575]Laflamme, N.; Leblanc, J.-F.; Mailloux, J.; Faure, N.; Labrie,

[121576]F.; Simard, J.: Mutation R96W in cytochrome P450c17 gene

causes combined

[121577]17- α -hydroxylase/17-20-lyase deficiency in two

French Canadian

[121578]patients. J. Clin. Endocr. Metab. 81: 264-268, 1996.

[121579]21523.

[121580]Lin, C. J.; Martens, J. W. M.; Miller, W. L.: NF-1C, Sp1, and

[121581]Sp3 are essential for transcription of the human gene for

P450c17

[121582](steroid 17- α -hydroxylase/17,20 lyase) in human

adrenal NCI-H295A

[121583]cells. Molec. Endocr. 15: 1277-1293, 2001.

[121584]21524.

[121585]Lin, D.; Harikrishna, J. A.; Moore, C. C. D.; Jones, K. L.;

Miller,

[121586]W. L.: Missense mutation serine106-to-proline causes
17-alpha-hydroxylase

[121587]deficiency. J. Biol. Chem. 266: 15992-15998, 1991.

[121588]21525.

[121589]Mallin, S. R.: Congenital adrenal hyperplasia secondary to
17-hydroxylase

[121590]deficiency. Ann. Intern. Med. 70: 69-76, 1969.

[121591]21526.

[121592]Mantero, F.; Scaroni, C.; Pasini, C. V.; Fagiolo, U.: No link-
age

[121593]between HLA and congenital adrenal hyperplasia due to
17-alpha-hydroxylase

[121594]deficiency. (Letter) New Eng. J. Med. 303: 530, 1980.

[121595]21527.

[121596]Matteson, K. J.; Picado-Leonard, J.; Chung, B.-C.; Mohan-
das, T.

[121597]K.; Miller, W. L.: Assignment of the gene for adrenal
P450C17 (steroid

[121598]17-alpha-hydroxylase/17,20 lyase) to human chromo-
some 10. J. Clin.

[121599]Endocr. Metab. 63: 789–791, 1986.

[121600]21528.

[121601]Miller, W. L.; Chung, B.-C.; Matteson, K. J.; Voutilainen, R.;

[121602]Picado–Leonard, J.: Molecular biology of steroid hormone synthesis.

[121603](Abstract) DNA 5: 61, 1986.

[121604]21529.

[121605]Donlon, T. A.; Krensky, A. M.; Clayberger, C.: Localization of

[121606]the human T lymphocyte activation gene 519 (D2S69E) to chromosome

[121607]2p12–q11. Cytogenet. Cell Genet. 53: 230–231, 1990.

[121608]21530.

[121609]Jongstra, J.; Schall, T. J.; Dyer, B. J.; Clayberger, C.; Jorgensen,

[121610]J.; Davis, M. M.; Krensky, A. M.: The isolation and sequence of a

[121611]novel gene from a human functional T cell line. J. Exp. Med. 165:

[121612]601–614, 1987.

[121613]21531.

[121614]Kaufmann, S. H. E.: CD8+ T lymphocytes in intracellular
microbial

[121615]infections. Immun. Today 9: 168–174, 1988.

[121616]21532.

[121617]Pena, S. V.; Hanson, D. A.; Carr, B. A.; Goralski, T. J.; Kren-
sky,

[121618]A. M.: Processing, subcellular localization, and function of
519

[121619](granulysin), a human late T cell activation molecule with
homology

[121620]to small, lytic granule proteins. J. Immun. 158:
2680–2688, 1997.

[121621]21533.

[121622]Stenger, S.; Hanson, D. A.; Teitelbaum, R.; Dewan, P.; Ni-
azi, K.

[121623]R.; Froelich, C. J.; Ganz, T.; Thoma–Uszynski, S.; Melian,
A.; Bogdan,

[121624]C.; Porcelli, S. A.; Bloom, B. R.; Krensky, A. M.; Modlin, R.
L.:

[121625]An antimicrobial activity of cytolytic T cells mediated by
granulysin. Science 282:

[121626]121–125, 1998.

[121627]21534.

[121628]Stenger, S.; Mazzaccaro, R. J.; Uyemura, K.; Cho, S.;
Barnes, P.

[121629]F.; Rosat, J. P.; Sette, A.; Brenner, M. B.; Porcelli, S. A.;
Bloom,

[121630]B. R.; Modlin, R. L.: Differential effects of cytolytic T cell
subsets

[121631]on intracellular infection. Science 276: 1684–1687, 1997.

[121632]21535.

[121633]Wong–Staal, F.; Dalla–Favera, R.; Franchini, G.; Gelmann,
E. P.;

[121634]Gallo, R. C.: Three distinct genes in human DNA related to
the transforming

[121635]genes of mammalian sarcoma retroviruses. Science 213:
226–228, 1981.

[121636]21536.

[121637]Aurias, A.; Rimbaut, C.; Buffe, D.; Dubousset, J.; Maz–
abraud, A.

[121638]: Chromosomal translocations in Ewing's sarcoma. (Letter)
New Eng.

[121639]J. Med. 309: 496–497, 1983.

[121640]21537.

[121641]Bartram, C. R.; de Klein, A.; Hagemeijer, A.; Grosveld, G.;
Heisterkamp,

[121642]N.; Groffen, J.: Localization of the human c-sis oncogene
in Ph-1-positive

[121643]and Ph-1-negative chronic myelocytic leukemia by in situ
hybridization. Blood 63:

[121644]223–225, 1984.

[121645]21538.

[121646]Bechet, J.-M.; Bornkamm, G.; Freese, U.-K.; Lenoir, G. M.:
The

[121647]c-sis oncogene is not activated in Ewing's sarcoma.
(Letter) New

[121648]Eng. J. Med. 310: 393 only, 1984.

[121649]21539.

[121650]Bishop, J. M.: Enemies within: the genesis of retrovirus
oncogenes. Cell 23:

[121651]5–6, 1981.

[121652]21540.

- [121653]Bolger, G. B.; Stamberg, J.; Kirsch, I. R.; Hollis, G. F.;
Schwarz,
[121654]D. F.; Thomas, G. H.: Chromosomal translocation t(14;22)
and oncogene
[121655](c-sis) variant in a pedigree with familial meningioma.
New Eng.
[121656]J. Med. 312: 564-567, 1985.
[121657]21541.
[121658]Cohen, J. B.; Levinson, A. D.: A point mutation in the last
intron
[121659]responsible for increased expression and transforming
activity of
[121660]the c-Ha-ras oncogene. Nature 334: 119-124, 1988.
[121661]21542.
[121662]Collins, T.; Ginsburg, D.; Boss, J. M.; Orkin, S. H.; Pober, J.
[121663]S.: Cultured human endothelial cells express platelet-de-
rived growth
[121664]factor B chain: cDNA cloning and structural analysis. Na-
ture 316:
[121665]748-750, 1985.
[121666]21543.

[121667]Dalla-Favera, R.; Gallo, R. C.; Giallongo, A.; Croce, C.:
Chromosomal

[121668]localization of the human homolog (c-sis) of the simian
sarcoma virus

[121669]onc gene. Science 218: 686-688, 1982.

[121670]21544.

[121671]Dalla-Favera, R.; Gelmann, E. P.; Gallo, R. C.; Wong-Staal,
F.

[121672]: A human onc gene homologous to the transforming
gene (v-sis) of

[121673]simian sarcoma virus. Nature 292: 31-35, 1981.

[121674]21545.

[121675]Deuel, T. F.; Huang, J. S.; Huang, S. S.; Stroobant, P.; Wa-
terfield,

[121676]M. D.: Expression of a platelet-derived growth factor-like
protein

[121677]in simian sarcoma virus transformed cells. Science 221:
1348-1350,

[121678]1983.

[121679]21546.

[121680]Devare, S. G.; Reddy, E. P.; Law, J. D.; Robbins, K. C.;

Aaronson,

[121681]S. A.: Nucleotide sequence of the simian sarcoma virus genome: demonstration

[121682]that its acquired cellular sequences encode the transforming gene

[121683]product p28-sis. Proc. Nat. Acad. Sci. 80: 731-735, 1983.

[121684]21547.

[121685]Charlet-B., N.; Logan, P.; Singh, G.; Cooper, T. A.: Dynamic antagonism

[121686]between ETR-3 and PTB regulates cell type-specific alternative splicing. Molec.

[121687]Cell 9: 649-658, 2002.

[121688]21548.

[121689]Borglum, A. D.; Flint, T.; Tommerup, N.; Fleckner, J.; Justesen,

[121690]J.; Kruse, T. A.: Assignment of the human tryptophanyl-tRNA synthetase

[121691]gene (WARS) to chromosome 14q32.2-q32.32. Cytogenet. Cell Genet. 73:

[121692]99-103, 1996.

[121693]21549.

[121694]Denney, R. M.; Borgaonkar, D.; Ruddle, F. H.: Order of
genes for

[121695]NP and TRPRS on chromosome 14. Cytogenet. Cell Genet.
22: 493-497,

[121696]1978.

[121697]21550.

[121698]Francke, U.; Denney, R. M.; Ruddle, F. H.: Intrachromoso-
mal gene

[121699]mapping in man: the gene for tryptophanyl-tRNA syn-
thetase maps in

[121700]region q21-qter of chromosome 14. Somat. Cell Genet. 3:
381-389,

[121701]1977.

[121702]21551.

[121703]Sack, G. H.; Huether, C. A.; Garg, N.: Familial spastic para-
plegia--clinical

[121704]and pathologic studies in a large kindred. Johns Hopkins
Med. J. 143:

[121705]117-121, 1978.

[121706]21552.

[121707]Schady, W.; Dick, J. P. R.; Sheard, A.; Crampton, S.: Central

[121708]motor conduction studies in hereditary spastic paraplegia.

J. Neurol.

[121709]Neurosurg. Psychiat. 54: 775–779, 1991.

[121710]21553.

[121711]Schady, W.; Sheard, A.: A quantitative study of sensory
function

[121712]in hereditary spastic paraplegia. Brain 113: 709–720,
1990.

[121713]21554.

[121714]Schady, W.; Smith, C. M. L.: Sensory neuropathy in heredi-
tary

[121715]spastic paraplegia. J. Neurol. Neurosurg. Psychiat. 57:
693–698,

[121716]1994.

[121717]21555.

[121718]Scheltens, P.; Bruyn, R. P. M.; Hazenberg, G. J.: A Dutch
family

[121719]with autosomal dominant pure spastic paraparesis
(Strumpell's disease). Acta

[121720]Neurol. Scand. 82: 169–173, 1990.

[121721]21556.

- [121722]Schwarz, G. A.; Liu, C. N.: Hereditary (familial) spastic paraplegia:
[121723]further clinical and pathologic observations. Arch. Neurol. Psychiat. 75:
[121724]144–162, 1956.
[121725]21557.
- [121726]Skre, H.: Hereditary spastic paraplegia in Western Norway. Clin. Genet. 6: 165–183, 1974.
[121728]21558.
- [121729]Strumpell, A.: Die primaere Seitenstrangsklerose (spastische Spinalparalyse). Dtsch. Z. Nervenheilk. 27: 291–339, 1904.
[121731]21559.
- [121732]Thurmon, T. F.; He, C.; Haskell, C.; Thorpe, P.; Thurmon, S. G.;
[121733]Rosen, D. R.: Genetic anticipation in a large family with pure autosomal
[121734]dominant hereditary spastic paraplegia. Am. J. Med. Genet. 83: 392–396,

[121735]1999.

[121736]21560.

[121737]Thurmon, T. F.; Walker, B. A.: Two distinct types of autosomal

[121738]dominant spastic paraplegia. Birth Defects Orig. Art. Ser. VII(1):

[121739]216–218, 1971.

[121740]21561.

[121741]Van Bogaert, L.: Etude genetique sur les paraplegies spasmotiques

[121742]familiales. J. Genet. Hum. 1: 6–23, 1952.

[121743]21562.

[121744]Zhao, X.; Alvarado, D.; Rainier, S.; Lemons, R.; Hedera, P.; Weber,

[121745]C. H.; Tükel, T.; Apak, M.; Heiman-Patterson, T.; Ming, L.; Bui, M.;

[121746]Fink, J. K.: Mutations in a newly identified GTPase gene cause autosomal

[121747]dominant hereditary spastic paraplegia. Nature Genet. 29: 326–331,

[121748]2001.

[121749]21563.

[121750]Benson, K. F.; Horwitz, M.; Wolff, J.; Friend, K.; Thompson, E.;

[121751]White, S.; Richards, R. I.; Raskind, W. H.; Bird, T. D.: CAG repeat

[121752]expansion in autosomal dominant familial spastic paraparesis: novel

[121753]expansion in a subset of patients. Hum. Molec. Genet. 7: 1779–1786,

[121754]1998.

[121755]21564.

[121756]Fonknechten, N.; Mavel, D.; Byrne, P.; Davoine, C.–S.; Cruaud,

[121757]C.; Boentsch, D.; Samson, D.; Coutinho, P.; Hutchinson, M.; McMonagle,

[121758]P.; Burgunder, J.–M.; Tartaglione, A.; and 10 others: Spectrum

[121759]of SPG4 mutations in autosomal dominant spastic paraplegia. Hum.

[121760]Molec. Genet. 9: 637–644, 2000.

[121761]21565.

- [121762]Nielsen, J. E.; Koefoed, P.; Abell, K.; Hasholt, L.; Eiberg, H.;
- [121763]Fenger, K.; Niebuhr, E.; Sorensen, S. A.: CAG repeat expansion in
- [121764]autosomal dominant pure spastic paraplegia linked to chromosome 2p21-p24. Hum.
- [121765]Molec. Genet. 6: 1811-1816, 1997.
- [121766]21566.
- [121767]Svenson, I. K.; Ashley-Koch, A. E.; Gaskell, P. C.; Riney, T.
- [121768]J.; Cumming, W. J. K.; Kingston, H. M.; Hogan, E. L.; Boustany, R.-M.
- [121769]N.; Vance, J. M.; Nance, M. A.; Pericak-Vance, M. A.; Marchuk, D.
- [121770]A.: Identification and expression analysis of spastin gene mutations
- [121771]in hereditary spastic paraplegia. Am. J. Hum. Genet. 68: 1077-1085,
- [121772]2001.
- [121773]21567.
- [121774]Ala-Kapee, M.; Nevanlinna, H.; Mali, M.; Jalkanen, M.; Schroder,
- [121775]J.: Localization of gene for human syndecan, an integral

membrane

[121776]proteoglycan and a matrix receptor, to chromosome 2.

Somat. Cell

[121777]Molec. Genet. 16: 501–505, 1990.

[121778]21568.

[121779]Alexander, C. M.; Reichsman, F.; Hinkes, M. T.; Lincecum, J.; Becker,

[121780]K. A.; Cumberledge, S.; Bernfield, M.: Syndecan–1 is required for

[121781]Wnt–1–induced mammary tumorigenesis in mice. Nature Genet. 25: 329–332,

[121782]2000.

[121783]21569.

[121784]Mali, M.; Jaakkola, P.; Arvilommi, A.–M.; Jalkanen, M.: Sequence

[121785]of human syndecan indicates a novel gene family of integral membrane

[121786]proteoglycans. J. Biol. Chem. 265: 6884–6889, 1990.

[121787]21570.

[121788]Oettinger, H. F.; Streeter, H.; Lose, E.; Copeland, N. G.; Gilbert,

[121789]D. J.; Justice, M. J.; Jenkins, N. A.; Mohandas, T.; Bernfield, M.

[121790]: Chromosome mapping of the murine syndecan gene. Genomics 11: 334–338,

[121791]1991.

[121792]21571.

[121793]Reizes, O.; Lincecum, J.; Wang, Z.; Goldberger, O.; Huang, L.;

[121794]Kaksonen, M.; Ahima, R.; Hinkes, M. T.; Barsh, G. S.; Rauvala, H.;

[121795]Bernfield, M.: Transgenic expression of syndecan-1 uncovers a physiological

[121796]control of feeding behavior by syndecan-3. Cell 106: 105–116, 2001.

[121797]21572.

[121798]Sanderson, R. D.; Lalor, P.; Bernfield, M.: B lymphocytes express

[121799]and lose syndecan at specific stages of differentiation. Cell Regulation 1:

[121800]27–35, 1989.

[121801]21573.

- [121802] Westman, P.; Hsieh, C.-L.; Mali, M.; Jalkanen, M.; Francke, U.;
- [121803] Schroder, J.: Assignment of the human syndecan (SDC) gene to short arm of chromosome 2. (Abstract) Cytogenet. Cell Genet. 58: 1873-1874, 1991.
- [121805] 1991.
- [121806] 21574.
- [121807] Asch, A. S.; Barnwell, J.; Silverstein, R. L.; Nachman, R. L.: Isolation of the thrombospondin membrane receptor. J. Clin. Invest. 79: 1054-1061, 1987.
- [121809] 1054-1061, 1987.
- [121810] 21575.
- [121811] de Fraipont, F.; El Atifi, M.; Gicquel, C.; Bertagna, X.; Chambaz, E. M.; Feige, J. J.: Expression of the angiogenesis markers vascular endothelial growth factor-A, thrombospondin-1, and platelet-derived endothelial cell growth factor in human sporadic adrenocortical tumors:

[121815]correlation with genotypic alterations. J. Clin. Endocr.
Metab. 85:

[121816]4734–4741, 2000.

[121817]21576.

[121818]Dixit, V. M.; Hennessy, S. W.; Grant, G. A.; Rotwein, P.;
Frazier,

[121819]W. A.: Characterization of a cDNA encoding the heparin
and collagen

[121820]binding domains of human thrombospondin. Proc. Nat.
Acad. Sci. 83:

[121821]5449–5453, 1986.

[121822]21577.

[121823]Frazier, W. A.: Thrombospondin: a modular adhesive gly-
coprotein

[121824]of platelets and nucleated cells. J. Cell Biol. 105: 625–632,
1987.

[121825]21578.

[121826]Jaffe, E.; Bornstein, P.; Disteché, C. M.: Mapping of the
thrombospondin

[121827]gene to human chromosome 15 and mouse chromosome
2 by in situ hybridization. Genomics 7:

[121828]123–126, 1990.

[121829]21579.

[121830]Lawler, J.; Sunday, M.; Thibert, V.; Duquette, M.; George, E. L.;

[121831]Rayburn, H.; Hynes, R. O.: Thrombospondin–1 is required for normal

[121832]murine pulmonary homeostasis and its absence causes pneumonia. J.

[121833]Clin. Invest. 101: 982–992, 1998.

[121834]21580.

[121835]Rodriguez–Manzaneque, J. C.; Lane, T. F.; Ortega, M. A.; Hynes,

[121836]R. O.; Lawler, J.; Iruela–Arispe, M. L.: Thrombospondin–1 suppresses

[121837]spontaneous tumor growth and inhibits activation of matrix metalloproteinase–9

[121838]and mobilization of vascular endothelial growth factor. Proc. Nat.

[121839]Acad. Sci. 98: 12485–12490, 2001.

[121840]21581.

[121841]Wolf, F. W.; Eddy, R. L.; Shows, T. B.; Dixit, V. M.: Structure

[121842]and chromosomal localization of the human thrombospondin gene. Genomics 6:

[121843]685–691, 1990.

[121844]21582.

[121845]Imai, K.; Sarker, A. H.; Akiyama, K.; Ikeda, S.; Yao, M.; Tsutsui,

[121846]K.; Shohmori, T.; Seki, S.: Genomic structure and sequence of a human

[121847]homologue (NTHL1/NTH1) of Escherichia coli endonuclease III with those

[121848]of the adjacent parts of TSC2 and SLC9A3R2 genes. Gene 222: 287–295,

[121849]1998.

[121850]21583.

[121851]Mira, J.-P.; Cariou, A.; Grall, F.; Delclaux, C.; Losser, M.-R.;

[121852]Heshmati, F.; Cheval, C.; Monchi, M.; Teboul, J.-L.; Riche, F.; Leleu,

[121853]G.; Arbibe, L.; Mignon, A.; Delpech, M.; Dhainaut, J.-F.: Association

[121854]of TNF2, a TNF- α promoter polymorphism, with septic shock susceptibility

[121855]and mortality: a multicenter study. J.A.M.A. 282: 561–568,
1999.

[121856]21584.

[121857]Moffatt, M. F.; Cookson, W. O. C. M.: Tumour necrosis fac-
tor

[121858]haplotypes and asthma. Hum. Molec. Genet. 6: 551–554,
1997.

[121859]21585.

[121860]Moraes, M. O.; Duppre, N. C.; Suffys, P. N.; Santos, A. R.;
Almeida,

[121861]A. S.; Nery, J. A. C.; Sampaio, E. P.; Sarno, E. N.: Tumor
necrosis

[121862]factor–alpha promoter polymorphism TNF2 is associated
with a stronger

[121863]delayed–type hypersensitivity reaction in the skin of bor-
derline tuberculoid

[121864]leprosy patients. Immunogenetics 53: 45–47, 2001.

[121865]21586.

[121866]Mulcahy, B.; Waldron–Lynch, F.; McDermott, M. F.; Adams,
C.; Amos,

[121867]C. I.; Zhu, D. K.; Ward, R. H.; Clegg, D. O.; Shanahan, F.;
Molloy,

[121868]M. G.; O'Gara, F.: Genetic variability in the tumor necrosis factor-lymphotoxin

[121869]region influences susceptibility to rheumatoid arthritis.
Am. J.

[121870]Hum. Genet. 59: 676-683, 1996.

[121871]21587.

[121872]Muller, U.; Jongeneel, C. V.; Nedospasov, S. A.; Lindahl, K.
F.;

[121873]Steinmetz, M.: Tumour necrosis factor and lymphotoxin
genes map close

[121874]to H-2D in the mouse major histocompatibility complex.
Nature 325:

[121875]265-267, 1987.

[121876]21588.

[121877]Nadel, S.; Newport, M. J.; Booy, R.; Levin, M.: Variation in

[121878]the tumor necrosis factor-alpha gene promoter region
may be associated

[121879]with death from meningococcal disease. J. Infect. Dis.
174: 878-880,

[121880]1996.

[121881]21589.

[121882]Nedospasov, S. A.; Hirt, B.; Shakhov, A. N.; Dobrynin, V. N.;

[121883]Kawashima, E.; Accolla, R. S.; Jongeneel, C. V.: The genes for tumor

[121884]necrosis factor (TNR-alpha) and lymphotoxin (TNR-beta) are tandemly

[121885]arranged on chromosome 17 of the mouse. Nucleic Acids Res. 14: 7713-7725,

[121886]1986.

[121887]21590.

[121888]Nedwin, G. E.; Naylor, S. L.; Sakaguchi, A. Y.; Smith, D.; Jarrett-Nedwin,

[121889]J.; Pennica, D.; Goeddel, D. V.; Gray, P. W.: Human lymphotoxin and

[121890]tumor necrosis factor genes: structure, homology and chromosomal localization. Nucleic

[121891]Acids Res. 13: 6361-6373, 1985.

[121892]21591.

[121893]Norman, R. A.; Bogardus, C.; Ravussin, E.: Linkage between obesity

[121894]and a marker near the tumor necrosis factor-alpha locus in Pima Indians. J.

[121895]Clin. Invest. 96: 158–162, 1995.

[121896]21592.

[121897]Obayashi, H.; Hasegawa, G.; Fukui, M.; Kamiuchi, K.; Kitamura,

[121898]A.; Ogata, M.; Kanaitzuka, T.; Shigeta, H.; Kitagawa, Y.; Nakano,

[121899]K.; Nishimura, M.; Ohta, M.; Nakamura, N.: Tumor necrosis factor

[121900]microsatellite polymorphism influences the development of insulin

[121901]dependency in adult-onset diabetes patients with the DRB1*1502–DQB1*0601

[121902]allele and anti-glutamic acid decarboxylase antibodies. J. Clin.

[121903]Endocr. Metab. 85: 3348–3351, 2000.

[121904]21593.

[121905]Old, L. J.: Tumor necrosis factor (TNF). Science 230: 630–632,

[121906]1985.

[121907]21594.

[121908]Pennica, D.; Nedwin, G. E.; Hayflick, J. S.; Seeburg, P. H.;

Derynck,

[121909]R.; Palladino, M. A.; Kohr, W. J.; Aggarwal, B. B.; Goeddel, D. V.

[121910]; Human tumour necrosis factor: precursor structure, expression and

[121911]homology to lymphotoxin. Nature 312: 724-729, 1984.

[121912]21595.

[121913]Ragoussis, J.; Bloemer, K.; Weiss, E. H.; Ziegler, A.: Localization

[121914]of the genes for tumor necrosis factor and lymphotoxin between the

[121915]HLA class I and III regions by field inversion gel electrophoresis. Immunogenetics 27:

[121916]66-69, 1988.

[121917]21596.

[121918]Rasmussen, S. K.; Urhammer, S. A.; Jensen, J. N.; Hansen, T.;

[121919]Borch-Johnsen, K.; Pedersen, O.: The -238 and -308 G6A polymorphisms

[121920]of the tumor necrosis factor alpha gene promoter are not associated

[121921]with features of the insulin resistance syndrome or altered

birth

[121922]weight in Danish Caucasians. J. Clin. Endocr. Metab. 85:
1731–1734,

[121923]2000.

[121924]21597.

[121925]Rosmond, R.; Chagnon, M.; Bouchard, C.; Bjorntorp, P.: G–
308A

[121926]polymorphism of the tumor necrosis factor alpha gene
promoter and

[121927]salivary cortisol secretion. J. Clin. Endocr. Metab. 86:
2178–2180,

[121928]2001.

[121929]21598.

[121930]Roy, S.; McGuire, W.; Mascie–Taylor, C. G.; Saha, B.; Hazra,
S.

[121931]K.; Hill, A. V.; Kwiatkowski, D.: Tumor necrosis factor pro-
moter

[121932]polymorphism and susceptibility to lepromatous leprosy.
J. Infect.

[121933]Dis. 176: 530–532, 1997.

[121934]21599.

[121935]Ruuls, S. R.; Sedgwick, J. D.: Unlinking tumor necrosis factor

[121936]biology from the major histocompatibility complex: lessons from human

[121937]genetics and animal models. Am. J. Hum. Genet. 65: 294-301, 1999.

[121938]21600.

[121939]Spies, T.; Morton, C. C.; Nedospasov, S. A.; Fiers, W.; Pious,

[121940]D.; Strominger, J. L.: Genes for the tumor necrosis factors alpha

[121941]and beta are linked to the human major histocompatibility complex. Proc.

[121942]Nat. Acad. Sci. 83: 8699-8702, 1986.

[121943]21601.

[121944]van Hensbroek, M. B.; Palmer, A.; Onyiorah, E.; Schneider, G.;

[121945]Jaffar, S.; Dolan, G.; Memming, H.; Frenkel, J.; Enwere, G.; Bennett,

[121946]S.; Kwiatkowski, D.; Greenwood, B.: The effect of a monoclonal antibody

[121947]to tumor necrosis factor on survival from childhood cere-

bral malaria. J.

[121948] Infect. Dis. 174: 1091–1097, 1996.

[121949] 21602.

[121950] Sashio, H.; Tamura, K.; Ito, R.; Yamamoto, Y.; Bamba, H.;
Kosaka,

[121951] T.; Fukui, S.; Sawada, K.; Fukuda, Y.; Tamura, K.; Satomi,
M.; Shimoyama,

[121952] T.; Furuyama, J.: Polymorphisms of the TNF gene and the
TNF receptor

[121953] superfamily member 1B gene are associated with suscep-
tibility to ulcerative

[121954] colitis and Crohn's disease, respectively. Immunogenetics
53: 1020–1027,

[121955] 2002.

[121956] 21603.

[121957] Van Ostade, X.; Vandenabeele, P.; Everaerd, B.; Loetscher,
H.;

[121958] Gentz, R.; Brockhaus, M.; Lesslauer, W.; Tavernier, J.;
Brouckaert,

[121959] P.; Fiers, W.: Human TNF mutants with selective activity on
the p55

[121960] receptor. Nature 361: 266–269, 1993.

[121961]21604.

[121962]Wang, A. M.; Creasey, A. A.; Ladner, M. B.; Lin, L. S.;
Strickler,

[121963]J.; Van Arsdell, J. N.; Yamamoto, R.; Mark, D. F.: Molecular
cloning

[121964]of the complementary DNA for human tumor necrosis fac-
tor. Science 228:

[121965]149–154, 1985.

[121966]21605.

[121967]Wilson, A. G.; Symons, J. A.; McDowell, T. L.; et al: Effects

[121968]of a polymorphism in the human tumor necrosis factor al-
pha promoter

[121969]on transcriptional activation. Proc. Nat. Acad. Sci. 94:
3195–3199,

[121970]1997.

[121971]21606.

[121972]Winchester, E. C.; Millwood, I. Y.; Rand, L.; Penny, M. A.;
Kessling,

[121973]A. M.: Association of the TNF- α -308 (G-A) polymor-
phism with self-reported

[121974]history of childhood asthma. Hum. Genet. 107: 591–596,

2000.

[121975]21607.

[121976]Witte, J. S.; Palmer, L. J.; O'Connor, R. D.; Hopkins, P. J.;

[121977]Hall, J. M.: Relation between tumour necrosis factor polymorphism

[121978]TNF-alpha-308 and risk of asthma. Europ. J. Hum. Genet. 10: 82-85,

[121979]2002.

[121980]21608.

[121981]Peter, D.; Finn, J. P.; Klisak, I.; Liu, Y.; Kojis, T.; Heinzmann,

[121982]C.; Roghani, A.; Sparkes, R. S.; Edwards, R. H.: Chromosomal localization

[121983]of the human vesicular amine transporter genes. Genomics 18: 720-723,

[121984]1993.

[121985]21609.

[121986]Roghani, A.; Welch, C.; Xia, Y.-R.; Liu, Y.; Peter, D.; Finn, J.

[121987]P.; Edwards, R. H.; Lusis, A. J.: Assignment of the mouse vesicular

[121988]monoamine transporter genes, Slc18a1 and Slc18a2, to

chromosomes 8

[121989]and 19 by linkage analysis. Mammalian Genome 7:
393–394, 1996.

[121990]21610.

[121991]Lania, L.; Donti, E.; Pannuti, A.; Pascucci, A.; Pengue, G.;
Felicciello,

[121992]I.; La Mantia, G.; Lanfranccone, L.; Pelicci, P.-G.: cDNA iso-
lation,

[121993]expression analysis, and chromosomal localization of two
human zinc

[121994]finger genes. Genomics 6: 333–340, 1990.

[121995]21611.

[121996]Han, Z.-G.; Zhang, Q.-H.; Ye, M.; Kan, L.-X.; Gu, B.-W.;
He, K.-L.;

[121997]Shi, S.-L.; Zhou, J.; Fu, G.; Mao, M.; Chen, S.-J.; Yu, L.;
Chen,

[121998]Z.: Molecular cloning of six novel Kruppel-like zinc finger
genes

[121999]from hematopoietic cells and identification of a novel
transregulatory

[122000]domain KRN. J. Biol. Chem. 274: 35741–35748, 1999.

[122001]21612.

[122002]Rousseau-Merck, M. F.; Huebner, K.; Berger, R.; Thiesen, H. J.

[122003]: Chromosomal localization of two human zinc finger protein genes,

[122004]ZNF24 (KOX17) and ZNF29 (KOX26), to 18q12 and 17p13-p12, respectively. Genomics 9:

[122005]154-161, 1991.

[122006]21613.

[122007]Seite, P.; Huebner, K.; Rousseau-Merck, M. F.; Berger, R.; Thiesen,

[122008]H. J.: Two human genes encoding zinc finger proteins, ZNF12 (KOX3)

[122009]and ZNF26 (KOX20), map to chromosomes 7p22-p21 and 12q24.33, respectively. Hum.

[122010]Genet. 86: 585-590, 1991.

[122011]21614.

[122012]King, R. A.: Personal Communication. Minneapolis, Minn. 12/31/1992.

[122013]21615.

[122014]Corral, J.; Martin, C.; Perez, R.; Sanchez, I.; Mories, M. T.;

[122015]San Millan, J. L.; Miralles, J. M.; Gonzalez-Sarmiento, R.:

Thyroglobulin

[122016]gene point mutation associated with non-endemic simple goitre. Lancet 341:

[122017]462-464, 1993.

[122018]21616.

[122019]Zinman, B.; Hanley, A. J. G.; Harris, S. B.; Kwan, J.; Fantus,

[122020]J. G.: Circulating tumor necrosis factor-alpha concentrations in

[122021]a Native Canadian population with high rates of type 2 diabetes mellitus. J.

[122022]Clin. Endocr. Metab. 84: 272-278, 1999.

[122023]21617.

[122024]de Vijlder, J. J. M.; Baas, F.; Koch, C. A. M.; Kok, K.; Gons,

[122025]M. H.: Autosomal dominant inheritance of a thyroglobulin abnormality

[122026]suggests cooperation of subunits in hormone formation. Ann. Endocr. 44:

[122027]36 only, 1983.

[122028]21618.

[122029]Hishinuma, A.; Takamatsu, J.; Ohyama, Y.; Yokozawa, T.;

Kanno,

[122030]Y.; Kuma, K.; Yoshida, S.; Matsuura, N.; Ieiri, T.: Two novel
cysteine

[122031]substitutions (C1263R and C1995S) of thyroglobulin
cause a defect

[122032]in intracellular transport of thyroglobulin in patients with
congenital

[122033]goiter and the variant type of adenomatous goiter. J. Clin.
Endocr.

[122034]Metab. 84: 1438–1444, 1999.

[122035]21619.

[122036]Ieiri, T.; Cochaux, P.; Targovnik, H. M.; Suzuki, M.; Shi-
moda,

[122037]S.-I.; Perret, J.; Vassart, G.: A 3-prime splice site mutation
in

[122038]the thyroglobulin gene responsible for congenital goiter
with hypothyroidism. J.

[122039]Clin. Invest. 88: 1901–1905, 1991.

[122040]21620.

[122041]Kim, P. S.; Ding, M.; Menon, S.; Jung, C.-G.; Cheng, J.-M.;
Miyamoto,

[122042]T.; Li, B.; Furudate, S.; Agui, T.: A missense mutation

G2320R in

[122043]the thyroglobulin gene causes non-goitrous congenital
primary hypothyroidism

[122044]in the WIC-rdw rat. Molec. Endocr. 14: 1944-1953, 2000.

[122045]21621.

[122046]Kim, P. S.; Hossain, S. A.; Park, Y.-N.; Lee, I.; Yoo, S.-E.;

[122047]Arvan, P.: A single amino acid change in the acetyl-
cholinesterase-like

[122048]domain of thyroglobulin causes congenital goiter with hy-
pothyroidism

[122049]in the cog/cog mouse: a model of human endoplasmic
reticulum storage

[122050]disease. Proc. Nat. Acad. Sci. 95: 9909-9913, 1998.

[122051]21622.

[122052]Kok, K.; van Dijk, J. E.; Sterk, A.; Baas, F.; van Ommen,
G.-J.

[122053]B.; de Vijlder, J. J. M.: Autosomal recessive inheritance of
goiter

[122054]in Dutch goats. J. Hered. 78: 298-300, 1987.

[122055]21623.

[122056]Landegent, J. E.; Jansen in de Wal, N.; van Ommen, G.-J.

B.; Baas,

[122057]F.; de Vijlder, J. J. M.; van Duijn, P.; van der Ploeg, M.:

Chromosomal

[122058]localization of a unique gene by non-autoradiographic in situ hybridization. Nature 317:

[122059]175-177, 1985.

[122060]21624.

[122061]Lazar, V.; Bidart, J.-M.; Caillou, B.; Mahe, C.; Lacroix, L.;

[122062]Filetti, S.; Schlumberger, M.: Expression of the Na(+)/I(-) symporter

[122063]gene in human thyroid tumors: a comparison study with other thyroid-specific

[122064]genes. J. Clin. Endocr. Metab. 84: 3228-3234, 1999.

[122065]21625.

[122066]Malthiery, Y.; Lissitzky, S.: Primary structure of human thyroglobulin

[122067]deduced from the sequence of its 8448-base complementary DNA. Europ.

[122068]J. Biochem. 165: 491-498, 1987.

[122069]21626.

[122070]Medeiros-Neto, G.; Bunduki, V.; Tomimori, E.; Gomes, S.;

Knobel,

[122071]M.; Martin, R. T.; Zugaib, M.: Prenatal diagnosis and treatment of

[122072]dyshormonogenetic fetal goiter due to defective thyroglobulin synthesis. J.

[122073]Clin. Endocr. Metab. 82: 4239–4242, 1997.

[122074]21627.

[122075]Ricketts, M. H.; Pohl, V.; de Martynoff, G.; Boyd, C. D.; Bester,

[122076]A. J.; Van Jaarsveld, P. P.; Vassart, G.: Defective splicing of thyroglobulin

[122077]gene transcripts in the congenital goitre of the Afrikaner cattle. EMBO

[122078]J. 4: 731–737, 1985.

[122079]21628.

[122080]Ricketts, M. H.; Simons, M. J.; Parma, J.; Mercken, L.; Dong,

[122081]Q.; Vassart, G.: A nonsense mutation causes hereditary goitre in

[122082]the Afrikaner cattle and unmasks alternative splicing of thyroglobulin

[122083]transcripts. Proc. Nat. Acad. Sci. 84: 3181–3184, 1987.

[122084]21629.

[122085]Savagner, F.; Rodien, P.; Reynier, P.; Rohmer, V.; Bigorgne, J.-C.;

[122086]Malthiery, Y.: Analysis of Tg transcripts by real-time RT-PCR in

[122087]the blood of thyroid cancer patients. J. Clin. Endocr. Metab. 87:

[122088]635–639, 2002.

[122089]21630.

[122090]Targovnik, H.; Propato, F.; Varela, V.; Wajchenberg, B.; Knobel,

[122091]M.; D'Abronzio, H. F.; Medeiros-Neto, G.: Low levels of thyroglobulin

[122092]messenger ribonucleic acid in congenital goitrous hypothyroidism with

[122093]defective thyroglobulin synthesis. J. Clin. Endocr. Metab. 69: 1137–1147,

[122094]1989.

[122095]21631.

[122096]Targovnik, H. M.; Frechtel, G. D.; Mendive, F. M.; Vono, J.; Cochaux,

- [122097]P.; Vassart, G.; Medeiros-Neto, G.: Evidence for the segregation
of three different mutated alleles of the thyroglobulin
gene in a
Brazilian family with congenital goiter and hypothyroidism. *Thyroid* 8:
291-297, 1998.
- [122100]21632.
- [122101]21632.
- [122102]Targovnik, H. M.; Medeiros-Neto, G.; Varela, V.; Cochaux, P.;
- [122103]Wajchenberg, B. L.; Vassart, G.: A nonsense mutation causes human
hereditary congenital goiter with preferential production
of a 171-nucleotide-deleted
thyroglobulin ribonucleic acid messenger. *J. Clin. Endocr. Metab.* 77:
210-215, 1993.
- [122106]21633.
- [122107]21633.
- [122108]Targovnik, H. M.; Vono, J.; Billerbeck, A. E. C.; Cerrone, G.
- [122109]E.; Varela, V.; Mendive, F.; Wajchenberg, B. L.; Medeiros-Neto, G.

[122110]: A 138-nucleotide deletion in the thyroglobulin ribonucleic acid

[122111]messenger in a congenital goiter with defective thyroglobulin synthesis. J.

[122112]Clin. Endocr. Metab. 80: 3356-3360, 1995.

[122113]21634.

[122114]Taylor, B. A.; Rowe, L.: The congenital goiter mutation is linked

[122115]to the thyroglobulin gene in the mouse. Proc. Nat. Acad. Sci. 84:

[122116]1986-1990, 1987.

[122117]21635.

[122118]van de Graaf, S. A. R.; Ris-Stalpers, C.; Veenboer, G. J. M.;

[122119]Cammenga, M.; Santos, C.; Targovnik, H. M.; de Vijlder, J. J. M.;

[122120]Medeiros-Neto, G.: A premature stopcodon in thyroglobulin messenger

[122121]RNA results in familial goiter moderate hypothyroidism. J. Clin.

[122122]Endocr. Metab. 84: 2537-2542, 1999.

[122123]21636.

[122124]Van Ommen, G.-J. B.; Baas, F.; Arnberg, A. C.; Pearson, P. L.;

[122125]De Vijlder, J. J. M.: Chromosome mapping and polymorphism study of

[122126]the human thyroglobulin gene. (Abstract) Cytogenet. Cell Genet. 37:

[122127]562 only, 1984.

[122128]21637.

[122129]Van Ommen, G. B.: Merging autosomal dominance and recessivity.

[122130](Letter) Am. J. Hum. Genet. 41: 689-691, 1987.

[122131]21638.

[122132]Yoshida, S.; Takamatsu, J.; Kuma, K.; Murakami, Y.; Sakane, S.;

[122133]Katayama, S.; Tarutani, O.; Ohsawa, N.: A variant of adenomatous

[122134]goiter with characteristic histology and possible hereditary thyroglobulin

[122135]abnormality. J. Clin. Endocr. Metab. 81: 1961-1966, 1996.

[122136]21639.

[122137]Spurr, N. K.; Goodfellow, P. N.; Sheer, D.; Bodmer, W. F.;

Vennstrom,

[122138]B.: Mapping of cellular oncogenes: ERBA1 is on chromosome 17. (Abstract) Cytogenet.

[122139]Cell Genet. 37: 591 only, 1984.

[122140]21640.

[122141]Thompson, C. C.; Weinberger, C.; Lebo, R.; Evans, R. M.: Identification

[122142]of a novel thyroid hormone receptor expressed in the mammalian central

[122143]nervous system. Science 237: 1610-1614, 1987.

[122144]21641.

[122145]Weinberger, C.; Thompson, C. C.; Ong, E. S.; Lebo, R.; Gruol,

[122146]D. J.; Evans, R. M.: The c-erb-A gene encodes a thyroid hormone receptor. Nature 324:

[122147]641-646, 1986.

[122148]21642.

[122149]Zabel, B. U.; Fournier, R. E. K.; Lalley, P. A.; Naylor, S. L.;

[122150]Sakaguchi, A. Y.: Cellular homologs of the avian erythroblastosis

[122151]virus erb-A and erb-B genes are syntenic in mouse but

asynthetic in

[122152]man. Proc. Nat. Acad. Sci. 81: 4874–4878, 1984.

[122153]21643.

[122154]Elchebly, M.; Payette, P.; Michaliszyn, E.; Cromlish, W.;
Collins,

[122155]S.; Loy, A. L.; Normandin, D.; Cheng, A.; Himms-Hagen, J.;
Chan, C.-C.;

[122156]Ramachandran, C.; Gresser, M. J.; Tremblay, M. L.;
Kennedy, B. P.

[122157]: Increased insulin sensitivity and obesity resistance in
mice lacking

[122158]the protein tyrosine phosphatase-1B gene. Science 283:
1544–1548,

[122159]1999.

[122160]21644.

[122161]Forsell, P. K. A. L.; Boie, Y.; Montalibet, J.; Collins, S.;
Kennedy,

[122162]B. P.: Genomic characterization of the human and mouse
protein tyrosine

[122163]phosphatase-1B genes. Gene 260: 145–153, 2000.

[122164]21645.

[122165]Gu, H. F.; Almgren, P.; Lindholm, E.; Frittitta, L.; Pizzuti, A.;

[122166]Trischitta, V.; Groop, L. C.: Association between the human glycoprotein

[122167]PC-1 gene and elevated glucose and insulin levels in paired-sibling

[122168]analysis. Diabetes 49: 1601-1603, 2000.

[122169]21646.

[122170]Haj, F. G.; Verveer, P. J.; Squire, A.; Neel, B. G.; Bastiaens,

[122171]P. I. H.: Imaging sites of receptor dephosphorylation by PTP1B on

[122172]the surface of the endoplasmic reticulum. Science 295: 1708-1711,

[122173]2002.

[122174]21647.

[122175]Jia, Z.; Barford, D.; Flint, A. J.; Tonks, N. K.: Structural

[122176]basis for phosphotyrosine peptide recognition by protein tyrosine

[122177]phosphatase 1B. Science 268: 1754-1758, 1995.

[122178]21648.

- [122179] Kennedy, B. P.; Ramachandran, C.: Protein tyrosine phosphatase-1B
[122180] in diabetes. *Biochem. Pharm.* 60: 877-883, 2000.
[122181] 21649.
- [122182] Mok, A.; Cao, H.; Zinman, B.; Hanley, A. J. G.; Harris, S. B.;
[122183] Kennedy, B. P.; Hegele, R. A.: A single nucleotide polymorphism in
[122184] protein tyrosine phosphatase PTP-1B is associated with
protection
[122185] from diabetes or impaired glucose tolerance in Oji-Cree. *J.*
Clin.
[122186] *Endocr. Metab.* 87: 724-727, 2002.
[122187] 21650.
- [122188] Tonks, N. K.; Diltz, C. D.; Fischer, E. H.: Purification of the
[122189] major protein-tyrosine-phosphatases of human placenta.
J. Biol. Chem. 263:
[122190] 6722-6730, 1988.
[122191] 21651.
- [122192] Barnea, G.; Silvennoinen, O.; Shaanan, B.; Honegger, A. M.;
Canoll,

[122193]P. D.; D'Eustachio, P.; Morse, B.; Levy, J. B.; Laforgia, S.; Huebner,

[122194]K.; Musacchio, J. M.; Sap, J.; Schlessinger, J.: Identification of

[122195]a carbonic anhydrase-like domain in the extracellular region of RPTP-gamma

[122196]defines a new subfamily of receptor tyrosine phosphatases. Molec.

[122197]Cell. Biol. 13: 1497-1506, 1993.

[122198]21652.

[122199]Kastury, K.; Ohta, M.; Lasota, J.; Moir, D.; Dorman, T.; LaForgia,

[122200]S.; Druck, T.; Huebner, K.: Structure of the human receptor tyrosine

[122201]phosphatase gamma gene (PTPRG) and relation to the familial RCC t(3;8)

[122202]chromosome translocation. Genomics 32: 225-235, 1996.

[122203]21653.

[122204]LaForgia, S.; Morse, B.; Levy, J.; Barnea, G.; Cannizzaro, L. A.;

[122205]Li, F.; Nowell, P. C.; Boghosian-Sell, L.; Glick, J.; Weston, A.;

[122206]Harris, C. C.; Drabkin, H.; Patterson, D.; Croce, C. M.; Schlessinger,
[122207]J.; Huebner, K.: Receptor protein-tyrosine phosphatase
gamma is a
[122208]candidate tumor suppressor gene at human chromosome
region 3p21. Proc.
[122209]Nat. Acad. Sci. 88: 5036-5040, 1991.
[122210]21654.
[122211]Latif, F.; Tory, K.; Modi, W.; Geil, L.; LaForgia, S.; Huebner,
[122212]K.; Zbar, B.; Lerman, M. I.: A MspI polymorphism and linkage
mapping
[122213]of the human protein-tyrosine phosphatase G (PTPRG)
gene. Hum. Molec.
[122214]Genet. 2: 91, 1993.
[122215]21655.
[122216]Cool, D. E.; Tonks, N. K.; Charbonneau, H.; Walsh, K. A.;
Fischer,
[122217]E. H.; Krebs, E. G.: cDNA isolated from a human T-cell library
encodes
[122218]a member of the protein-tyrosine-phosphatase family.
Proc. Nat. Acad.

[122219]Sci. 86: 5257–5261, 1989.

[122220]21656.

[122221]Johnson, C. V.; Cool, D. E.; Glaccum, M. B.; Green, N.; Fischer,

[122222]E. H.; Bruskin, A.; Hill, D. E.; Lawrence, J. B.: Isolation and mapping

[122223]of human T-cell protein tyrosine phosphatase sequences: localization

[122224]of genes and pseudogenes discriminated using fluorescence hybridization

[122225]with genomic versus cDNA probes. Genomics 16: 619–629, 1993.

[122226]21657.

[122227]Sakaguchi, A. Y.; Sylvia, V. L.; Martinez, L.; Lalley, P. A.; Shows,

[122228]T. B.; Han, E. S.; Smith, E. A.; Ghosh Choudhury, G.: Assignment

[122229]of tyrosine-specific T-cell phosphatase to conserved syntenic groups

[122230]on human chromosome 18 and mouse chromosome 18. (Abstract) Cytogenet.

[122231]Cell Genet. 58: 2014–2015, 1991.

[122232]21658.

[122233]Sakaguchi, A. Y.; Sylvia, V. L.; Martinez, L.; Smith, E. A.;
Han,

[122234]E. S.; Lalley, P. A.; Shows, T. B.; Ghosh Choudhury, G.: As-
signment

[122235]of tyrosine-specific T-cell phosphatase to conserved syn-
tenic groups

[122236]on human chromosome 18 and mouse chromosome 18.
Genomics 12: 151-154,

[122237]1992.

[122238]21659.

[122239]Adachi, M.; Sekiya, M.; Arimura, Y.; Takekawa, M.; Itoh, F.;
Hinoda,

[122240]Y.; Imai, K.; Yachi, A.: Protein-tyrosine phosphatase ex-
pression

[122241]in pre-B cell NALM-6. Cancer Res. 52: 737-740, 1992.

[122242]21660.

[122243]Adachi, M.; Sekiya, M.; Isobe, M.; Kumura, Y.; Ogita, Z.;
Hinoda,

[122244]Y.; Imai, K.; Yachi, A.: Molecular cloning and chromosomal
mapping

[122245]of a human protein-tyrosine phosphatase LC-PTP.

Biochem. Biophys.

[122246]Res. Commun. 186: 1607-1615, 1992.

[122247]21661.

[122248]Zanke, B.; Suzuki, H.; Kishihara, K.; Mizzen, L.; Minden, M.; Pawson,

[122249]A.; Mak, T. W.: Cloning and expression of an inducible lymphoid-specific,

[122250]protein tyrosine phosphatase (HePTPase). Europ. J. Immun. 22: 235-239,

[122251]1992.

[122252]21662.

[122253]Schols, L.; Szymanski, S.; Peters, S.; Przuntek, H.; Epplen, J.

[122254]T.; Hardt, C.; Riess, O.: Genetic background of apparently idiopathic

[122255]sporadic cerebellar ataxia. Hum. Genet. 107: 132-137, 2000.

[122256]21663.

[122257]Virgilio, L.; Isobe, M.; Narducci, M. G.; Carotenuto, P.; Camerini,

[122258]B.; Kurosawa, N.; ar-Rushdi, A.; Croce, C. M.; Russo, G.:
Chromosome

[122259]walking on the TCL1 locus involved in T-cell neoplasia.
Proc. Nat.

[122260]Acad. Sci. 90: 9275–9279, 1993.

[122261]21664.

[122262]Virgilio, L.; Lazzeri, C.; Bichi, R.; Nibu, K.; Narducci, M. G.;

[122263]Russo, G.; Rothstein, J. L.; Croce, C. M.: Deregulated ex-
pression

[122264]of TCL1 causes T cell leukemia in mice. Proc. Nat. Acad.
Sci. 95:

[122265]3885–3889, 1998.

[122266]21665.

[122267]Virgilio, L.; Narducci, M. G.; Isobe, M.; Billips, L. G.;
Cooper,

[122268]M. D.; Croce, C. M.; Russo, G.: Identification of the TCL1
gene involved

[122269]in T-cell malignancies. Proc. Nat. Acad. Sci. 91:
12530–12534, 1994.

[122270]21666.

[122271]Comp, P. C.; Nixon, R. R.; Cooper, M. R.; Esmon, C. T.: Fa-

milial

[122272]protein S deficiency is associated with recurrent thrombo-
sis. J.

[122273]Clin. Invest. 74: 2082-2088, 1984.

[122274]21667.

[122275]Cooper, D. N.: Personal Communication. London, England
9/17/1993.

[122276]21668.

[122277]D'Angelo, A.; Della Valle, P.; Crippa, L.; Pattarini, E.;
Grimaldi,

[122278]L. M. E.; D'Angelo, S. V.: Autoimmune protein S deficiency
in a boy

[122279]with severe thromboembolic disease. New Eng. J. Med.
328: 1753-1757,

[122280]1993.

[122281]21669.

[122282]Dahlback, B.: Protein S and C4b-binding protein: compo-
nents involved

[122283]in the regulation of the protein C anticoagulant system.
Thromb.

[122284]Haemost. 66: 49-61, 1991.

[122285]21670.

[122286]Dahlback, B.; Stenflo, J.: High molecular weight complex in human

[122287]plasma between vitamin K dependent protein S and complement component

[122288]C4b-binding protein. Proc. Nat. Acad. Sci. 78: 2512-2516, 1981.

[122289]21671.

[122290]Engesser, L.; Broekmans, A. W.; Briet, E.; Brommer, E. J. P.;

[122291]Bertina, R. M.: Hereditary protein S deficiency: clinical manifestations. Ann.

[122292]Intern. Med. 106: 677-682, 1987.

[122293]21672.

[122294]Espinosa-Parrilla, Y.; Morell, M.; Souto, J. C.; Tirado, I.; Fontcuberta,

[122295]J.; Estivill, X.; Sala, N.: Protein S gene analysis reveals the presence

[122296]of a cosegregating mutation in most pedigrees with type I
but not

[122297]type III PS deficiency. Hum. Mutat. 14: 30–39, 1999.

[122298]21673.

[122299]Fulton, A. B.; Hansen, R. M.; Petersen, R. A.; Vanderveen,
D.

[122300]K.: The rod photoreceptors in retinopathy of prematurity:
an electroretinographic

[122301]study. Arch. Ophthalm. 119: 499–505, 2001.

[122302]21674.

[122303]Girolami, A.; Simioni, P.; Lazzaro, A. R.; Pontara, E.; Ruzza,

[122304]G.: Heterozygous protein-S deficiency: a study of a large
kindred. Acta

[122305]Haemat. 84: 162–168, 1990.

[122306]21675.

[122307]Golub, B. M.; Sibony, P. A.; Coller, B. S.: Protein S defi-
ciency

[122308]associated with central retinal artery occlusion. Arch.
Ophthalm. 108:

[122309]918 only, 1990.

[122310]21676.

[122311]Griffin, J. H.; Gruber, A.; Fernandez, J. A.: Reevaluation of

[122312]total, free, and bound protein S and C4b-binding protein
levels in

[122313]plasma anticoagulated with citrate or hirudin. Blood 79:
3203-3211,

[122314]1992.

[122315]21677.

[122316]Hayashi, T.; Nishioka, J.; Shigekiyo, T.; Saito, S.; Suzuki, K.

[122317]: Protein S Tokushima: abnormal molecule with a substi-
tution of glu

[122318]for lys-155 in the second epidermal growth factor-like
domain of protein

[122319]S. Blood 83: 683-690, 1994.

[122320]21678.

[122321]Heeb, M. J.; Rosing, J.; Bakker, H. M.; Fernandez, J. A.;
Tans,

[122322]G.; Griffin, J. H.: Protein S binds to and inhibits factor Xa.
Proc.

[122323]Nat. Acad. Sci. 91: 2728-2732, 1994.

[122324]21679.

[122325]Hoskins, J.; Norman, D. K.; Beckmann, R. J.; Long, G. L.:
Cloning

[122326]and characterization of human liver cDNA encoding a pro-
tein S precursor. Proc.

[122327]Nat. Acad. Sci. 84: 349–353, 1987.

[122328]21680.

[122329]Kamiya, T.; Sugihara, T.; Ogata, K.; Saito, H.; Suzuki, K.;
Nishioka,

[122330]J.; Hashimoto, S.; Yamagata, K.: Inherited deficiency of
protein

[122331]S in a Japanese family with recurrent venous thrombosis: a
study of

[122332]three generations. Blood 67: 406–410, 1986.

[122333]21681.

[122334]Koller, H.; Stoll, G.; Sitzler, M.; Burk, M.; Schottler, B.; Fre-
und,

[122335]H-J.: Deficiency of both protein C and protein S in a family
with

[122336]ischemic strokes in young adults. Neurology 44:
1238–1240, 1994.

[122337]21682.

[122338]Lundwall, A.; Dackowski, W.; Cohen, E.; Shaffer, M.; Mahr, A.;

[122339]Dahlback, B.; Stenflo, J.; Wydro, R.: Isolation and sequence of the

[122340]cDNA for human protein S, a regulator of blood coagulation. Proc.

[122341]Nat. Acad. Sci. 83: 6716-6720, 1986.

[122342]21683.

[122343]Mahasandana, C.; Suvatte, V.; Chuansumrit, A.; Marlar, R. A.;

[122344]Manco-Johnson, M. J.; Jacobson, L. J.; Hathaway, W. E.: Homozygous

[122345]protein S deficiency in an infant with purpura fulminans. J. Pediat. 117:

[122346]750-753, 1990.

[122347]21684.

[122348]Maillard, C.; Berruyer, M.; Serre, C. M.; Dechavanne, M.; Delmas,

[122349]P. D.: Protein-S, a vitamin K-dependent protein, is a bone matrix

[122350]component synthesized and secreted by osteoblasts. Endocrinology 130:

[122351]1599–1604, 1992.

[122352]21685.

[122353]Makris, M.; Leach, M.; Beauchamp, N. J.; Daly, M. E.; Cooper,

[122354]P. C.; Hampton, K. K.; Bayliss, P.; Peake, I. R.; Miller, G. J.; Preston,

[122355]F. E.: Genetic analysis, phenotypic diagnosis, and risk of venous

[122356]thrombosis in families with inherited deficiencies of protein S. Blood 95:

[122357]1935–1941, 2000.

[122358]21686.

[122359]Malnick, S. D. H.; Sthoeger, Z. M.: Autoimmune protein S deficiency. New

[122360]Eng. J. Med. 329: 1898 only, 1993.

[122361]21687.

[122362]Mannucci, P. M.; Valsecchi, C.; Krachmalnicoff, A.; Faioni, E.

[122363]M.; Tripodi, A.: Familial dysfunction of protein S. Thromb.

Haemost. 62:

[122364]763-766, 1989.

[122365]21688.

[122366]Mintz-Hittner, H. A.; Miyashiro, M. J.; Knight-Nanan, D.
M.; O'Malley,

[122367]R. E.; Marlar, R. A.: Vitreoretinal findings similar to
retinopathy

[122368]of prematurity in infants with compound heterozygous
protein S deficiency. Ophthalmology 106:

[122369]1525-1530, 1999.

[122370]21689.

[122371]Naylor, S. L.; Marshall, A.; Solomon, A.; Long, G.: Mapping
human

[122372]protein C and protein S to chromosomes 2 and 3, respec-
tively. (Abstract) Cytogenet.

[122373]Cell Genet. 46: 669 only, 1987.

[122374]21690.

[122375]Pan, E. Y.; Gomperts, E. D.; Millen, R.; Gilsanz, V.: Bone
mineral

[122376]density and its association with inherited protein S defi-
ciency. Thromb.

[122377]Res. 58: 221-231, 1990.

[122378]21691.

[122379]Phillips, W. G.; Marsden, J. R.; Hill, F. G.: Purpura fulminans

[122380]due to protein S deficiency following chickenpox. Brit. J. Derm. 127:

[122381]30-32, 1992.

[122382]21692.

[122383]Ploos van Amstel, H. K.; Huisman, M. V.; Reitsma, P. H.; ten Cate,

[122384]J.; Bertina, R. M.: Partial protein S gene deletion in a family with

[122385]hereditary thrombophilia. Blood 73: 479-483, 1989.

[122386]21693.

[122387]Ploos van Amstel, H. K.; Reitsma, P. H.; Bertina, R. M.: The

[122388]human protein S locus: identification of the PS-alpha gene as a site

[122389]of liver protein S messenger RNA synthesis. Biochem. Biophys. Res.

[122390]Commun. 157: 1033-1038, 1988.

[122391]21694.

[122392]Ploos van Amstel, J. K.; van der Zanden, A. L.; Bakker, E.;
Reitsma,

[122393]P. H.; Bertina, R. M.: Two genes homologous with human
protein S

[122394]cDNA are located on chromosome 3. Thromb. Haemost.
58: 982-987,

[122395]1987.

[122396]21695.

[122397]Reitsma, P. H.; Ploos van Amstel, H. K.; Bertina, R. M.:
Three

[122398]novel mutations in five unrelated subjects with hereditary
protein

[122399]S deficiency type I. J. Clin. Invest. 93: 486-492, 1994.

[122400]21696.

[122401]Sacco, R. L.; Owen, J.; Mohr, J. P.; Tatemichi, T. K.; Gross-
man,

[122402]B. A.: Free protein S deficiency: a possible association with
cerebrovascular

[122403]occlusion. Stroke 20: 1657-1661, 1989.

[122404]21697.

[122405] Sas, G.; Blasko, G.; Petro, I.; Griffin, J. H.: A protein S deficient

[122406] family with portal vein thrombosis. (Letter) *Thromb. Haemost.* 54:

[122407] 724 only, 1985.

[122408] 21698.

[122409] Chan, J. Y.; Cheung, M.-C.; Moi, P.; Chan, K.; Kan, Y. W.: Chromosomal

[122410] localization of the human NF-E2 family of bZIP transcription factors

[122411] by fluorescence in situ hybridization. *Hum. Genet.* 95: 265-269,

[122412] 1995.

[122413] 21699.

[122414] Chan, J. Y.; Han, X.-L.; Kan, Y. W.: Cloning of Nrf1, an NF-E2-related

[122415] transcription factor, by genetic selection in yeast. *Proc. Nat. Acad.*

[122416] *Sci.* 90: 11371-11375, 1993.

[122417] 21700.

[122418] Chan, J. Y.; Kwong, M.; Lu, R.; Chang, J.; Wang, B.; Yen, T.

S.

[122419]B.; Kan, Y. W.: Targeted disruption of the ubiquitous CNC-
bZIP transcription

[122420]factor, Nrf-1, results in anemia and embryonic lethality in
mice. EMBO

[122421]J. 17: 1779-1787, 1998.

[122422]21701.

[122423]Luna, L.; Johnsen, O.; Skartlien, A.; Pedeutour, F.; Turc-
Carel,

[122424]C.; Prydz, H.; Kolsto, A.-B.: Molecular cloning of a putative
novel

[122425]human bZIP transcription factor on chromosome 17q22.
Genomics 22:

[122426]553-562, 1994.

[122427]21702.

[122428]Luna, L.; Johnsen, O.; Skartlien, A. H.; Pedeutour, F.; Turc-
Carel,

[122429]C.; Prydz, H.; Kolsto, A.-B.: Molecular cloning of a putative
novel

[122430]human bZIP transcription factor on chromosome 17q22.
Genomics 22:

[122431]553-562, 1994.

[122432]21703.

[122433]McKie, J.; Johnstone, K.; Mattei, M.-G.; Scambler, P.:

Cloning

[122434]and mapping of murine Nfe2l1. Genomics 25: 716–719,
1995.

[122435]21704.

[122436]McKie, J.; Scambler, P. J.: The Nfe2l1 gene maps to distal
mouse

[122437]chromosome 11. Mammalian Genome 7: 89–90, 1996.

[122438]21705.

[122439]Dryja, T. P.; Berson, E. L.; Rao, V. R.; Oprian, D. D.: Het-
erozygous

[122440]missense mutation in the rhodopsin gene as a cause of
congenital stationary

[122441]night blindness. Nature Genet. 4: 280–283, 1993.

[122442]21706.

[122443]Gal, A.; Xu, S.; Piczenik, Y.; Eiberg, H.; Duvigneau, C.;
Schwinger,

[122444]E.; Rosenberg, T.: Gene for autosomal dominant congeni-
tal stationary

[122445]night blindness maps to the same region as the gene for

the beta-subunit
[122446]of the rod photoreceptor cGMP phosphodiesterase (PDEB)
in chromosome
[122447]4p16.3. Hum. Molec. Genet. 3: 323-325, 1994.
[122448]21707.
[122449]Alfano, J. E.; Berger, J. P.: Retinitis pigmentosa, ophthalmoplegia,
[122450]and spastic quadriplegia. Am. J. Ophthalm. 43: 231-240,
1957.
[122451]21708.
[122452]Anttinen, A.; Nikoskelainen, E.; Marttila, R. J.; Grenman, R.;
[122453]Falck, B.; Aarnisalo, E.; Kalimo, H.: Familial olivopontocerebellar
[122454]atrophy with macular degeneration: a separate entity
among the olivopontocerebellar
[122455]atrophies. Acta Neurol. Scand. 73: 180-190, 1986.
[122456]21709.
[122457]Benomar, A.; Krols, L.; Stevanin, G.; Cancel, G.; LeGuern,
E.;
[122458]David, G.; Ouhabi, H.; Martin, J.-J.; Durr, A.; Zaim, A.;
Ravise,

- [122459]N.; Busque, C.; Penet, C.; Van Regemorter, N.; Weissenbach, J.; Yahyaoui, M.; Chkili, T.; Agid, Y.; Van Broeckhoven, C.; Brice, A.: The gene for autosomal dominant cerebellar ataxia with pigmentary macular dystrophy maps to chromosome 3p12-p21.1. Nature Genet. 10: 84-88, 1995.
- [122463]21710.
- [122464]Benomar, A.; Le Guern, E.; Durr, A.; Ouhabi, H.; Stevanin, G.; Yahyaoui, M.; Chkili, T.; Agid, Y.; Brice, A.: Autosomal-dominant cerebellar ataxia with retinal degeneration (ADCA type II) is genetically different from ADCA type I. Ann. Neurol. 35: 439-444, 1994.
- [122468]21711.
- [122469]Bjork, A.; Lindblom, U.; Wadensten, L.: Retinal degeneration in hereditary ataxia. J. Neurol. Neurosurg. Psychiat. 19: 186-193,

[122471]1956.

[122472]21712.

[122473]Carpenter, S.; Schumacher, G. A.: Familial infantile cerebellar

[122474]atrophy associated with retinal degeneration. Arch. Neurol. 14:

[122475]82–94, 1966.

[122476]21713.

[122477]Cooles, P.; Michaud, R.; Best, P. V.: A dominantly inherited progressive

[122478]disease in a black family characterized by cerebellar and retinal

[122479]degeneration, external ophthalmoplegia and abnormal mitochondria. J.

[122480]Neurol. Sci. 87: 275–288, 1988.

[122481]21714.

[122482]Cruysberg, J. R. M.; Eerola, K. U.; Vrijland, H. R.; Aandeker,

[122483]A. L.; Kremer, H. P. H.; Deutman, A. F.: Autosomal recessive cerebellar

[122484]ataxia with bull's-eye macular dystrophy. Am. J. Ophthal.

133: 410–413,

[122485]2002.

[122486]21715.

[122487]David, G.; Abbas, N.; Stevanin, G.; Durr, A.; Yvert, G.; Cancel,

[122488]G.; Weber, C.; Imbert, G.; Saudou, F.; Antoniou, E.;
Drabkin, H.;

[122489]Gemmill, R.; Giunti, P.; Benomar, A.; Wood, N.; Ruberg, M.;
Agid,

[122490]Y.; Mandel, J.–L.; Brice, A.: Cloning of the SCA7 gene reveals a

[122491]highly unstable CAG repeat expansion. *Nature Genet.* 17:
65–70, 1997.

[122492]21716.

[122493]David, G.; Giunti, P.; Abbas, N.; Coullin, P.; Stevanin, G.;
Horta,

[122494]W.; Gemmell, R.; Weissenbach, J.; Wood, N.; Cunha, S.;
Drabkin, H.;

[122495]Harding, A. E.; Agid, Y.; Brice, A.: The gene for autosomal
dominant

[122496]cerebellar ataxia type II is located in a 5–cM region in
3p12–p13:

[122497]genetic and physical mapping of the SCA7 locus. Am. J. Hum. Genet. 59:

[122498]1328–1336, 1996.

[122499]21717.

[122500]Duinkerke–Eerola, K. U.; Cruysberg, J. R. M.; Deutman, A. F.:

[122501]Atrophic maculopathy associated with hereditary ataxia. Am. J. Ophthal. 90:

[122502]597–603, 1980.

[122503]21718.

[122504]Enevoldson, T. P.; Sanders, M. D.; Harding, A. E.: Autosomal

[122505]dominant cerebellar ataxia with pigmentary macular dystrophy: a clinical

[122506]and genetic study of eight families. Brain 117: 445–460, 1994.

[122507]21719.

[122508]Foster, J. B.; Ingram, T. T. S.: Familial cerebro–macular degeneration

[122509]and ataxia. J. Neurol. Neurosurg. Psychiat. 25: 63–68, 1962.

[122510]21720.

[122511]Froment, J.; Bonnet, P.; Colrat, A.: Heredo-degenerations
retinienne

[122512]et spino-cerebelleuse: variantes ophtalmoscopiques et
neurologiques

[122513]presentees par trois generations successives. J. Med. Lyon
153-163,

[122514]1937.

[122515]21721.

[122516]Giunti, P.; Stevanin, G.; Worth, P. F.; David, G.; Brice, A.;

[122517]Wood, N. W.: Molecular and clinical study of 18 families
with ADCA

[122518]type II: evidence for genetic heterogeneity and de novo
mutation. Am.

[122519]J. Hum. Genet. 64: 1594-1603, 1999.

[122520]21722.

[122521]Gouw, L. G.; Digre, K. B.; Harris, C. P.; Haines, J. H.;

Ptacek,

[122522]L. J.: Autosomal dominant cerebellar ataxia with retinal
degeneration:

[122523]clinical, neuropathologic, and genetic analysis of a large

kindred. Neurology 44:

[122524]1441-1447, 1994.

[122525]21723.

[122526]Gouw, L. G.; Kaplan, C. D.; Haines, J. H.; Digre, K. B.; Rutledge,

[122527]S. L.; Matilla, A.; Leppert, M.; Zoghbi, H. Y.; Ptacek, L. J.:
Retinal

[122528]degeneration characterizes a spinocerebellar ataxia mapping to chromosome

[122529]3p. Nature Genet. 10: 89-93, 1995.

[122530]21724.

[122531]Gu, W.; Wang, Y.; Liu, X.; Zhou, B.; Zhou, Y.; Wang, G.:
Molecular

[122532]and clinical study of spinocerebellar ataxia type 7 in Chinese kindreds. Arch.

[122533]Neurol. 57: 1513-1518, 2000.

[122534]21725.

[122535]Halsey, J. H., Jr.; Scott, T. R.; Farmer, T. W.: Adult hereditary

[122536]cerebelloretinal degeneration. Neurology 17: 87-90,
1967.

[122537]21726.

[122538]Sobetzko, D.; Eich, G.; Kalff-Suske, M.; Grzeschik, K.-H.;
Superti-Furga,

[122539]A.: Boy with syndactylies, macrocephaly, and severe skeletal dysplasia:

[122540]not a new syndrome, but two dominant mutations (GLI3
E543X and COL2A1

[122541]G973R) in the same individual. Am. J. Med. Genet. 90:
239-242, 2000.

[122542]21727.

[122543]Stroschein, S. L.; Wang, W.; Zhou, S.; Zhou, Q.; Luo, K.:
Negative

[122544]feedback regulation of TGF-beta signaling by the SnoN
oncoprotein. Science 286:

[122545]771-774, 1999.

[122546]21728.

[122547]Goldblum, S. E.; Ding, X.; Funk, S. E.; Sage, E. H.: SPARC
(secreted

[122548]protein acidic and rich in cysteine) regulates endothelial
cell shape

[122549]and barrier function. Proc. Nat. Acad. Sci. 91: 3448-3452,
1994.

[122550]21729.

[122551]Lossie, A. C.; Gordon, D. F.; Camper, S. A.: Localization of thyrotropin-releasing

[122552]hormone receptor and thyrotroph embryonic factor on mouse chromosome

[122553]15. Mammalian Genome 4: 621-623, 1993.

[122554]21730.

[122555]Bongarzone, I.; Butti, M. G.; Coronelli, S.; Borrello, M. G.; Santoro,

[122556]M.; Mondellini, P.; Pilotti, S.; Fusco, A.; Della Porta, G.; Pierotti,

[122557]M. A.: Frequent activation of ret protooncogene by fusion with a

[122558]new activating gene in papillary thyroid carcinomas. Cancer Res. 54:

[122559]2979-2985, 1994.

[122560]21731.

[122561]Bongarzone, I.; Butti, M. G.; Fugazzola, L.; Pacini, F.; Pinchera,

[122562]A.; Vorontsova, T. V.; Demidchik, E. P.; Pierotti, M. A.: Comparison

[122563]of the breakpoint regions of ELE1 and RET genes involved
in the generation

[122564]of RET/PTC3 oncogene in sporadic and in radiation-as-
sociated papillary

[122565]thyroid carcinomas. Genomics 42: 252–259, 1997.

[122566]21732.

[122567]Klein, M.; Vignaud, J.-M.; Hennequin, V.; Toussaint, B.;
Bresler,

[122568]L.; Plenat, F.; Leclere, J.; Duprez, A.; Weryha, G.: Increased
expression

[122569]of the vascular endothelial growth factor is a pejorative
prognosis

[122570]marker in papillary thyroid carcinoma. J. Clin. Endocr.
Metab. 86:

[122571]656–658, 2001.

[122572]21733.

[122573]Katyal, S. L.; Singh, G.; Locker, J.: Characterization of a
second

[122574]human pulmonary surfactant-associated protein SP-A
gene. Am. J. Resp.

[122575]Cell Molec. Biol. 6: 446–452, 1992.

[122576]21734.

[122577]Ramet, M.; Lofgren, J.; Albo, O.-P.; Hallman, M.: Surfactant protein-A

[122578]gene locus associated with recurrent otitis media. J. Pediat. 138:

[122579]266-268, 2001.

[122580]21735.

[122581]Mignon, C.; Okada, A.; Mattei, M. G.; Basset, P.: Assignment of

[122582]the human membrane-type matrix metalloproteinase (MMP14) gene to 14q11-q12

[122583]by in situ hybridization. Genomics 28: 360-361, 1995.

[122584]21736.

[122585]Wilson, C. L.; Ouellette, A. J.; Satchell, D. P.; Ayabe, T.; Lopez-Boado,

[122586]Y. S.; Stratman, J. L.; Hultgren, S. J.; Matrisian, L. M.; Parks,

[122587]W. C.: Regulation of intestinal alpha-defensin activation by the

[122588]metalloproteinase matrilysin in innate host defense. Science 286:

[122589]113-117, 1999.

[122590]21737.

[122591]Inglehearn, C. F.; Bashir, R.; Lester, D. H.; Jay, M.; Bird, A.

[122592]C.; Bhattacharya, S. S.: A 3-bp deletion in the rhodopsin
gene in

[122593]a family with autosomal dominant retinitis pigmentosa.
Am. J. Hum.

[122594]Genet. 48: 26-30, 1991.

[122595]21738.

[122596]Inglehearn, C. F.; Keen, T. J.; Bashir, R.; Jay, M.; Fitzke, F.;

[122597]Bird, A. C.; Crombie, A.; Bhattacharya, S.: A completed
screen for

[122598]mutations of the rhodopsin gene in a panel of patients
with autosomal

[122599]dominant retinitis pigmentosa. Hum. Molec. Genet. 1:
41-45, 1992.

[122600]21739.

[122601]Inglehearn, C. F.; Lester, D. H.; Bashir, R.; Atif, U.; Keen,

[122602]T. J.; Sertedaki, A.; Lindsey, J.; Jay, M.; Bird, A. C.; Farrar,
G.

[122603]J.; Humphries, P.; Bhattacharya, S. S.: Recombination be-
tween rhodopsin

[122604]and locus D3S47 (C17) in rhodopsin retinitis pigmentosa families. Am.

[122605]J. Hum. Genet. 50: 590–597, 1992.

[122606]21740.

[122607]Keen, T. J.; Inglehearn, C. F.; Lester, D. H.; Bashir, R.; Jay,

[122608]M.; Bird, A. C.; Jay, B.; Bhattacharya, S. S.: Autosomal dominant

[122609]retinitis pigmentosa: four new mutations in rhodopsin, one of them

[122610]in the retinal attachment site. Genomics 11: 199–205, 1991.

[122611]21741.

[122612]Khorana, H. G.: Rhodopsin, photoreceptor of the rod cell: an

[122613]emerging pattern for structure and function. J. Biol. Chem. 267:

[122614]1–4, 1992.

[122615]21742.

[122616]Kijas, J. W.; Cideciyan, A. V.; Aleman, T. S.; Pianta, M. J.;

[122617]Pearce–Kelling, S. E.; Miller, B. J.; Jacobson, S. G.; Aguirre, G.

[122618]D.; Acland, G. M.: Naturally occurring rhodopsin mutation in the

[122619]dog causes retinal dysfunction and degeneration mimicking human dominant

[122620]retinitis pigmentosa. Proc. Nat. Acad. Sci. 99: 6328–6333, 2002.

[122621]21743.

[122622]Kiselev, A.; Socolich, M.; Vinos, J.; Hardy, R. W.; Zuker, C.

[122623]S.; Ranganathan, R.: A molecular pathway for light-dependent photoreceptor

[122624]apoptosis in Drosophila. Neuron 28: 139–152, 2000.

[122625]21744.

[122626]Kranich, H.; Bartkowski, S.; Denton, M. J.; Krey, S.; Dickinson,

[122627]P.; Duvigneau, C.; Gal, A.: Autosomal dominant 'sector' retinitis

[122628]pigmentosa due to a point mutation predicting an asn15-to-ser substitution

[122629]of rhodopsin. Hum. Molec. Genet. 2: 813–814, 1993.

[122630]21745.

[122631]Kumar-Singh, R.; Wang, H.; Humphries, P.; Farrar, G. J.:

Autosomal

[122632]dominant retinitis pigmentosa: no evidence for nonallelic genetic

[122633]heterogeneity on 3q. Am. J. Hum. Genet. 52: 319–326, 1993.

[122634]21746.

[122635]Kumaramanickavel, G.; Maw, M.; Denton, M. J.; John, S.; Srisailapathy

[122636]Srikumari, C. R.; Orth, U.; Oehlmann, R.; Gal, A.: Missense rhodopsin

[122637]mutation in a family with recessive RP. (Letter) Nature Genet. 8:

[122638]10–11, 1994.

[122639]21747.

[122640]Lem, J.; Krasnoperova, N. V.; Calvert, P. D.; Kosaras, B.; Cameron,

[122641]D. A.; Nicolo, M.; Makino, C. L.; Sidman, R. L.: Morphological, physiological,

[122642]and biochemical changes in rhodopsin knockout mice. Proc. Nat. Acad.

[122643]Sci. 96: 736–741, 1999.

[122644]21748.

[122645]Li, T.; Sandberg, M. A.; Pawlyk, B. S.; Rosner, B.; Hayes, K.

[122646]C.; Dryja, T. P.; Berson, E. L.: Effect of vitamin A supplementation

[122647]on rhodopsin mutants threonine-17-to-methionine and proline-347-to-serine

[122648]in transgenic mice and in cell cultures. Proc. Nat. Acad. Sci. 95:

[122649]11933-11938, 1998.

[122650]21749.

[122651]Li, T.; Snyder, W. K.; Olsson, J. E.; Dryja, T. P.: Transgenic

[122652]mice carrying the dominant rhodopsin mutation P347S: evidence for

[122653]defective vectorial transport of rhodopsin to the outer segments. Proc.

[122654]Nat. Acad. Sci. 93: 14176-14181, 1996.

[122655]21750.

[122656]Macke, J. P.; Davenport, C. M.; Jacobson, S. G.; Hennessey, J.

[122657]C.; Gonzalez-Fernandez, F.; Conway, B. P.; Heckenlively, J.; Palmer,

[122658]R.; Maumenee, I. H.; Sieving, P.; Gouras, P.; Good, W.;

Nathans, J.

[122659] Identification of novel rhodopsin mutations responsible for retinitis

[122660] pigmentosa: implications for the structure and function of rhodopsin. Am.

[122661] J. Hum. Genet. 53: 80–89, 1993.

[122662] 21751.

[122663] Mendez, A.; Burns, M. E.; Roca, A.; Lem, J.; Wu, L.-W.; Simon,

[122664] M. I.; Baylor, D. A.; Chen, J.: Rapid and reproducible deactivation

[122665] of rhodopsin requires multiple phosphorylation sites.

Neuron 28:

[122666] 153–164, 2000.

[122667] 21752.

[122668] Naash, M. I.; Hollyfield, J. G.; Al-Ubaidi, M. R.; Baehr, W.:

[122669] Simulation of human autosomal dominant retinitis pigmentosa in transgenic

[122670] mice expressing a mutated murine opsin gene. Proc. Nat. Acad. Sci. 90:

[122671] 5499–5503, 1993.

[122672]21753.

[122673]Nathans, J.; Hogness, D. S.: Isolation and nucleotide sequence

[122674]of the gene encoding human rhodopsin. Proc. Nat. Acad. Sci. 81:

[122675]4851–4855, 1984.

[122676]21754.

[122677]Nathans, J.; Piantanida, T. P.; Eddy, R. L.; Shows, T. B.; Hogness,

[122678]D. S.: Molecular genetics of inherited variation in human color vision. Science 232:

[122679]203–210, 1986.

[122680]21755.

[122681]Nathans, J.; Thomas, D.; Hogness, D. S.: Molecular genetics of

[122682]human color vision: the genes encoding blue, green, and red pigments. Science 232:

[122683]193–202, 1986.

[122684]21756.

[122685]Nie, Z.; Chen, S.; Kumar, R.; Zack, D. J.: RER, an evolutionarily

[122686] conserved sequence upstream of the rhodopsin gene, has enhancer activity. J.

[122687] Biol. Chem. 271: 2667–2675, 1996.

[122688] 21757.

[122689] Oh, K. T.; Weleber, R. G.; Lotery, A.; Oh, D. M.; Billingslea,

[122690] A. M.; Stone, E. M.: Description of a new mutation in rhodopsin,

[122691] pro23 to ala, and comparison with electroretinographic and clinical

[122692] characteristics of the pro23-to-his mutation. Arch. Ophthalmol. 118:

[122693] 1269–1276, 2000.

[122694] 21758.

[122695] Olsson, J. E.; Gordon, J. W.; Pawlyk, B. S.; Roof, D.; Hayes,

[122696] A.; Molday, R. S.; Mukai, S.; Cowley, G. S.; Berson, E. L.; Dryja,

[122697] T. P.: Transgenic mice with a rhodopsin mutation (Pro23His): a mouse

[122698] model of autosomal dominant retinitis pigmentosa. Neuron 9: 815–830,

[122699] 1992.

[122700]21759.

[122701]Palczewski, K.; Kumasaka, T.; Hori, T.; Behnke, C. A.; Mo-
toshima,

[122702]H.; Fox, B. A.; Le Trong, I.; Teller, D. C.; Okada, T.;
Stenkamp,

[122703]R. E.; Yamamoto, M.; Miyano, M.: Crystal structure of
rhodopsin:

[122704]a G protein-coupled receptor. Science 289: 739-745,
2000.

[122705]21760.

[122706]Rao, V. R.; Cohen, G. B.; Oprian, D. D.: Rhodopsin muta-
tion G90D

[122707]and a molecular mechanism for congenital night blind-
ness. Nature 367:

[122708]639-642, 1994.

[122709]21761.

[122710]Robinson, P. R.; Cohen, G. B.; Zhukovsky, E. A.; Oprian, D.
D.

[122711]: Constitutively active mutants of rhodopsin. Neuron 9:
719-725,

[122712]1992.

[122713]21762.

[122714]Rosenfeld, P. J.; Cowley, G. S.; Hahn, L. B.; Sandberg, M. A.;

[122715]Berson, E. L.; Dryja, T. P.: A null mutation within the rhodopsin

[122716]gene in a family with autosomal recessive retinitis pigmentosa. (Abstract) Invest.

[122717]Ophthalmol. Vis. Sci. 33: 1397, 1992.

[122718]21763.

[122719]Mason, I. J.; Murphy, D.; Munke, M.; Francke, U.; Elliott, R. W.;

[122720]Hogan, B. L. M.: Developmental and transformation-sensitive expression

[122721]of the SPARC gene on mouse chromosome 11. EMBO J. 5: 1831-1837,

[122722]1986.

[122723]21764.

[122724]Mason, I. J.; Taylor, A.; Williams, J. G.; Sage, H.; Hogan, B.

[122725]L. M.: Evidence from molecular cloning that SPARC, a major product

[122726]of mouse embryo parietal endoderm, is related to an en-

endothelial cell

[122727] 'culture shock' glycoprotein of Mr 43,000. EMBO J. 5:

1465–1472,

[122728] 1986.

[122729] 21765.

[122730] Naylor, S. L.; Helen-Davis, D.; Villarreal, X. C.; Long, G. L.

[122731]: The human osteonectin gene on chromosome 5 is polymorphic. (Abstract) Cytogenet.

[122732] Cell Genet. 51: 1051 only, 1989.

[122733] 21766.

[122734] Schwartz, R. C.; Young, M. F.; Tsipouras, P.: Two RFLPs in the

[122735] 5-prime end of the human osteonectin (ON) gene. Nucleic Acids Res. 16:

[122736] 9076 only, 1988.

[122737] 21767.

[122738] Stenner, D. D.; Tracy, R. P.; Riggs, B. L.; Mann, K. G.: Human

[122739] platelets contain and secrete osteonectin, a major protein of mineralized

[122740] bone. Proc. Nat. Acad. Sci. 83: 6892–6896, 1986.

[122741]21768.

[122742]Swaroop, A.; Francke, U.: Molecular cloning, cDNA sequence, and

[122743]expression of human SPARC (osteonectin). (Abstract) Am. J. Hum. Genet. 41:

[122744]A240 only, 1987.

[122745]21769.

[122746]Swaroop, A.; Hogan, B. L. M.; Francke, U.: Molecular analysis

[122747]of the cDNA for human SPARC/osteonectin/BM-40: sequence, expression,

[122748]and localization of the gene to chromosome 5q31-q33. Genomics 2:

[122749]37-47, 1988.

[122750]21770.

[122751]Termine, J. D.; Kleinman, H. K.; Whitson, S. W.; Conn, K. M.;

[122752]McGarvey, M. L.; Martin, G. R.: Osteonectin, a bone-specific protein

[122753]linking mineral to collagen. Cell 26: 99-105, 1981.

[122754]21771.

[122755]Thony, B.; Heizmann, C. W.; Mattei, M.-G.: Human GTP-cyclohydrolase

[122756] gene and sepiapterin reductase gene map to region 14q21-q22 and

[122757]2p14-p12, respectively, by in situ hybridization. Genomics 26: 168-170,

[122758]1995.

[122759]21772.

[122760]Jin, H.; Oksenberg, D.; Ashkenazi, A.; Peroutka, S. J.; Duncan,

[122761]A. M. V.; Rozmahel, R.; Yang, Y.; Mengod, G.; Palacios, J. M.; O'Dowd,

[122762]B. F.: Characterization of the human 5-hydroxytryptamine(1B) receptor. J.

[122763]Biol. Chem. 267: 5735-5738, 1992.

[122764]21773.

[122765]Levy, F. O.; Gudermann, T.; Birnbaumer, M.; Kaumann, A. J.; Birnbaumer,

[122766]L.: Molecular cloning of a human gene (S31) encoding a novel serotonin

[122767]receptor mediating inhibition of adenylyl cyclase. FEBS Lett. 296:

[122768]201–206, 1992.

[122769]21774.

[122770]Levy, F. O.; Holtgreve-Grez, H.; Tasken, K.; Solberg, R.;
Ried,

[122771]T.; Gudermann, T.: Assignment of the gene encoding the
5-HT-1E serotonin

[122772]receptor (S31) (locus HTR1E) to human chromosome
6q14–q15. Genomics 22:

[122773]637–640, 1994.

[122774]21775.

[122775]McAllister, G.; Charlesworth, A.; Snodin, C.; Beer, M. S.;
Noble,

[122776]A. J.; Middlemiss, D. N.; Iversen, L. L.; Whiting, P.: Molecu-
lar

[122777]cloning of a serotonin receptor from human brain
(5HT1E): a fifth

[122778]5HT1-like subtype. Proc. Nat. Acad. Sci. 89: 5517–5521,
1992.

[122779]21776.

[122780]Weinshank, R. L.; Zgombick, J. M.; Macchi, M. J.; Branchek,
T.

[122781]A.; Hartig, P. R.: Human serotonin 1D receptor is encoded by a subfamily

[122782]of two distinct genes: 5-HT(1D-alpha) and 5-HT(1D-beta). Proc. Nat.

[122783]Acad. Sci. 89: 3630-3634, 1992.

[122784]21777.

[122785]Pabst, T.; Mueller, B. U.; Zhang, P.; Radomska, H. S.; Nar-ravula,

[122786]S.; Schnittger, S.; Behre, G.; Hiddemann, W.; Tenen, D. G.: Dominant-negative

[122787]mutations of CEBPA, encoding CCAAT/enhancer binding protein-alpha

[122788](C/EBP-alpha), in acute myeloid leukemia. Nature Genet. 27: 263-270,

[122789]2001.

[122790]21778.

[122791]Chen, Z.-Y.; Battinelli, E. M.; Fielder, A.; Bunday, S.; Sims,

[122792]K.; Breakefield, X. O.; Craig, I. W.: A mutation in the Norrie disease

[122793]gene (NDP) associated with X-linked familial exudative vitreoretinopathy. Nature

[122794]Genet. 5: 180–183, 1993.

[122795]21779.

[122796]Dudgeon, J.: Familial exudative vitreo–retinopathy. Trans.
Ophthalm.

[122797]Soc. U.K. 99: 45–49, 1979.

[122798]21780.

[122799]Matsuda, I.; Arashima, S.; Nambu, H.; Takekoshi, Y.;
Anakura,

[122800]M.: Hyperammonemia due to a mutant enzyme of or-
nithine transcarbamylase. Pediatrics 48:

[122801]595–600, 1971.

[122802]21781.

[122803]Yang–Feng, T. L.; DeGennaro, L. J.; Francke, U.: Genes for
synapsin

[122804]I, a neuronal phosphoprotein, map to conserved regions
of human and

[122805]murine X chromosomes. Proc. Nat. Acad. Sci. 83:
8679–8683, 1986.

[122806]21782.

[122807]Matsuda, I.; Nagata, N.; Matsuura, T.; Oyanagi, K.; Tada,
K.;

[122808]Narisawa, K.; Kitagawa, T.; Sakiyama, T.; Yamashita, F.;
Yoshino,
[122809]M.: Retrospective survey of urea cycle disorders: Part 1.
Clinical
[122810]and laboratory observations of thirty-two Japanese male
patients with
[122811]ornithine transcarbamylase deficiency. Am. J. Med. Genet.
38: 85-89,
[122812]1991.
[122813]21783.
[122814]Mavinakere, M.; Morizono, H.; Shi, D.; Allewell, N. M.;
Tuchman,
[122815]M.: The clinically variable R40H mutant ornithine car-
bamoyltransferase
[122816]shows cytosolic degradation of the precursor protein in
CHO cells. J.
[122817]Inherit. Metab. Dis. 24: 614-622, 2001.
[122818]21784.
[122819]Michels, V. V.; Potts, E.; Walser, M.; Beaudet, A. L.: Or-
nithine
[122820]transcarbamylase deficiency: long-term survival. Clin.
Genet. 22:

[122821]211-214, 1982.

[122822]21785.

[122823]Matsuda, I.; Matsuura, T.; Nishiyori, A.; Komaki, S.;
Hoshide,

[122824]R.; Matsumoto, T.; Funakoshi, M.; Kiwaki, K.; Endo, F.;
Hata, A.;

[122825]Shimadzu, M.; Yoshino, M.: Phenotypic variability in male
patients

[122826]carrying the mutant ornithine transcarbamylase (OTC) al-
lele, arg40his,

[122827]ranging from a child with an unfavourable prognosis to an
asymptomatic

[122828]older adult. J. Med. Genet. 33: 645-648, 1996.

[122829]21786.

[122830]Morsy, M. A.; Zhao, J. Z.; Ngo, T. T.; Warman, A. W.;
O'Brien,

[122831]W. E.; Graham, F. L.; Caskey, C. T.: Patient selection may
affect

[122832]gene therapy success: dominant negative effects observed
for ornithine

[122833]transcarbamylase in mouse and human hepatocytes. J.
Clin. Invest. 97:

[122834]826-832, 1996.

[122835]21787.

[122836]Nagata, N.; Matsuda, I.; Oyanagi, K.: Estimated frequency
of
[122837]urea cycle enzymopathies in Japan. (Letter) Am. J. Med.
Genet. 39:

[122838]228-229, 1991.

[122839]21788.

[122840]Nussbaum, R. L.; Boggs, B. A.; Beaudet, A. L.; Doyle, S.;
Potter,

[122841]J. L.; O'Brien, W. E.: New mutation and prenatal diagnosis
in ornithine

[122842]transcarbamylase deficiency. Am. J. Hum. Genet. 38:
149-158, 1986.

[122843]21789.

[122844]Ohtake, A.; Takayanagi, M.; Yamamoto, S.; Kakinuma, H.;
Nakajima,

[122845]H.; Tatibana, M.; Mori, M.: Molecular basis of ornithine
transcarbamylase

[122846]deficiency in spf and spf-ash mutant mice. J. Inherit.
Metab. Dis. 9:

[122847]289–291, 1986.

[122848]21790.

[122849]Oizumi, J.; Ng, W. G.; Koch, R.; Shaw, K. N. F.; Sweetman, L.;

[122850]Velazquez, A.; Donnell, G. N.: Partial ornithine transcarbamylase

[122851]deficiency associated with recurrent hyperammonemia, lethargy and

[122852]depressed sensorium. Clin. Genet. 25: 538–542, 1984.

[122853]21791.

[122854]Old, J. M.; Briand, P. L.; Purvis-Smith, S.; Howard, N. J.; Wilcken,

[122855]B.; Hammond, J.; Pearson, P.; Cathelineau, L.; Williamson, R.; Davies,

[122856]K. E.: Prenatal exclusion of ornithine transcarbamylase deficiency

[122857]by direct gene analysis. Lancet I: 73–75, 1985.

[122858]21792.

[122859]Oppliger Leibundgut, E.; Liechti-Gallati, S.; Colombo, J.-P.;

[122860]Wermuth, B.: Ornithine transcarbamylase deficiency: ten new mutations

[122861]and high proportion of de novo mutations in heterozygous females. Hum.

[122862]Mutat. 9: 409–411, 1997.

[122863]21793.

[122864]Oppliger Leibundgut, E.; Liechti-Gallati, S.; Colombo, J. P.;

[122865]Wermuth, B.: Ornithine transcarbamylase deficiency: new sites with

[122866]increased probability of mutation. Hum. Genet. 95: 191–196, 1995.

[122867]21794.

[122868]Oppliger Leibundgut, E.; Wermuth, B.; Colombo, J.-P.; Liechti-Gallati,

[122869]S.: Ornithine transcarbamylase deficiency: characterization of gene

[122870]mutations and polymorphisms. Hum. Mutat. 8: 333–339, 1996.

[122871]21795.

[122872]Pelet, A.; Rotig, A.; Bonaiti-Pellie, C.; Rabier, D.; Cormier,

[122873]V.; Toumas, E.; Hentzen, D.; Saudubray, J.-M.; Munnich, A.: Carrier

[122874]detection in a partially dominant X-linked disease: or-

ornithine transcarbamylase

[122875] deficiency. Hum. Genet. 84: 167–171, 1990.

[122876] 21796.

[122877] Pembrey, M. E.; Old, J. M.; Leonard, J. V.; Rodeck, C. H.;
Warren,

[122878] R.; Davies, K. E.: Prenatal diagnosis of ornithine carbamoyl
transferase

[122879] deficiency using a gene specific probe. J. Med. Genet. 22:
462–465,

[122880] 1985.

[122881] 21797.

[122882] Ploechl, E.; Ploechl, W.; Stoeckler-Ipsiroglu, S.; Pokorny,
H.;

[122883] Wermuth, B.: Late-onset ornithine transcarbamylase defi-
ciency in

[122884] two families with different mutations in the same codon.
Clin. Genet. 59:

[122885] 111–114, 2001.

[122886] 21798.

[122887] Qureshi, I. A.; Letarte, J.; Ouellet, R.: Ornithine transcar-
bamylase

[122888]deficiency in mutant mice. I. Studies on the characterization of enzyme

[122889]defect and suitability as animal model of human disease. *Pediat.*

[122890]*Res.* 13: 807-811, 1979.

[122891]21799.

[122892]Ricciuti, F. C.; Gelehrter, T. D.; Rosenberg, L. E.: X-chromosome

[122893]inactivation in human liver: confirmation of X-linkage of ornithine

[122894]transcarbamylase. *Am. J. Hum. Genet.* 28: 332-338, 1976.

[122895]21800.

[122896]Rodeck, C. H.; Patrick, A. D.; Pembrey, M. E.; Tzannatos, C.;

[122897]Whitfield, A. E.: Fetal liver biopsy for prenatal diagnosis of ornithine

[122898]carbamyl transferase deficiency. *Lancet* II: 297-300, 1982.

[122899]21801.

[122900]Rosenberg, L. E.; Kalousek, F.; Orsulak, M. D.: Biogenesis of

[122901]ornithine transcarbamylase in *spf(ash)* mutant mice: two

cytoplasmic

[122902]precursors, one mitochondrial enzyme. Science 222:
426-428, 1983.

[122903]21802.

[122904]Rowe, P. C.; Newman, S. L.; Brusilow, S. W.: Natural history
[122905]of symptomatic partial ornithine transcarbamylase defi-
ciency. New

[122906]Eng. J. Med. 314: 541-547, 1986.

[122907]21803.

[122908]Rozen, R.; Fox, J.; Fenton, W. A.; Horwich, A. L.; Rosen-
berg,

[122909]L. E.: Gene deletion and restriction fragment length poly-
morphisms

[122910]at the human ornithine transcarbamylase locus. Nature
313: 815-817,

[122911]1985.

[122912]21804.

[122913]Russell, A.; Levin, B.; Oberholzer, V. G.; Sinclair, L.: Hyper-
ammonaemia.

[122914]A new instance of an inborn enzymatic defect of the
biosynthesis of

[122915]urea. Lancet II: 699–700, 1962.

[122916]21805.

[122917]Schwartz, M.; Christensen, E.; Christensen, N. C.; Skovby, F.;

[122918]Davies, K. E.; Old, J. M.: Detection and exclusion of carriers of

[122919]ornithine transcarbamylase deficiency by RFLP analysis. Clin. Genet. 29:

[122920]449–452, 1986.

[122921]21806.

[122922]Scott, C. R.; Chiang–Teng, C.; Goodman, S. I.; Greensher, A.;

[122923]Mace, J. W.: X-linked transmission of ornithine–transcarbamylase

[122924]deficiency. (Letter) Lancet II: 1148, 1972.

[122925]21807.

[122926]Shapiro, J. M.; Schaffner, F.; Tallan, H. H.; Gaull, G. E.: Mitochondrial

[122927]abnormalities of liver in primary ornithine transcarbamylase deficiency. Pediat.

[122928]Res. 14: 735–759, 1980.

[122929]21808.

[122930]Sinke, R. J.; de Leeuw, B.; Janssen, H. A. P.; Olde Weghuis, D.;

[122931]Suijkerbuijk, R. F.; Meloni, A. M.; Gilgenkrantz, S.; Berger, W.;

[122932]Ropers, H. H.; Sandberg, A. A.; Geurts van Kessel, A.: Localization

[122933]of X chromosome short arm markers relative to synovial sarcoma- and

[122934]renal adenocarcinoma-associated translocation break-points. Hum. Genet. 92:

[122935]305-308, 1993.

[122936]21809.

[122937]Brennan, T. J.; Seeley, W. W.; Kilgard, M.; Schreiner, C. E.; Tecott,

[122938]L. H.: Sound-induced seizures in serotonin 5-HT-2C receptor mutant

[122939]mice. Nature Genet. 16: 387-390, 1997.

[122940]21810.

[122941]Gurevich, I.; Tamir, H.; Arango, V.; Dwork, A. J.; Mann, J. J.;

[122942]Schmauss, C.: Altered editing of serotonin 2C receptor

pre-mRNA in

[122943]the prefrontal cortex of depressed suicide victims. *Neuron*
34: 349–356,

[122944]2002.

[122945]21811.

[122946]Hall, C. S.: Genetic differences in fatal audiogenic
seizures:

[122947]between two inbred strains of house mice. *J. Hered.* 38:
3–6, 1947.

[122948]21812.

[122949]Lappalainen, J.; Zhang, L.; Dean, M.; Oz, M.; Ozaki, N.; Yu,
D.;

[122950]Virkkunen, M.; Weight, F.; Linnoila, M.; Goldman, D.: Iden-
tification,

[122951]expression, and pharmacology of a cys(23)–ser(23) sub-
stitution in

[122952]the human 5-HT(2C) receptor gene (HTR2C). *Genomics*
27: 274–279,

[122953]1995.

[122954]21813.

[122955]Milatovich, A.; Hsieh, C.–L.; Bonaminio, G.; Tecott, L.;

Julius,

[122956]D.; Francke, U.: Serotonin receptor 1c gene assigned to X chromosome

[122957]in human (band q24) and mouse (bands D-F4). Hum.

Molec. Genet. 1:

[122958]681-684, 1992.

[122959]21814.

[122960]Tecott, L. H.; Sun, L. M.; Akana, S. F.; Strack, A. M.;

Lowenstein,

[122961]D. H.; Dallman, M. F.; Julius, D.: Eating disorder and epilepsy in

[122962]mice lacking 5-HT_{2C} serotonin receptors. Nature 374: 542-546, 1995.

[122963]21815.

[122964]Blaschke, R. J.; Monaghan, A. P.; Schiller, S.; Schechinger, B.;

[122965]Rao, E.; Padilla-Nash, H.; Ried, T.; Rappold, G. A.: SHOT, a SHOX-related

[122966]homeobox gene, is implicated in craniofacial, brain, heart, and limb

[122967]development. Proc. Nat. Acad. Sci. 95: 2406-2411, 1998.

[122968]21816.

[122969]Castillo, S.; Youlton, R.; Be, C.: Dyschondrosteosis is controlled

[122970]by X and Y linked loci. (Abstract) Cytogenet. Cell Genet.
40: 601-602,

[122971]1985.

[122972]21817.

[122973]Clement-Jones, M.; Schiller, S.; Rao, E.; Blaschke, R. J.; Zungiga,

[122974]A.; Zeller, R.; Robson, S. C.; Binder, G.; Glass, I.; Strachan,
T.;

[122975]Lindsay, S.; Rappold, G. A.: The short stature homeobox
gene SHOX

[122976]is involved in skeletal abnormalities in Turner syndrome.
Hum. Molec.

[122977]Genet. 9: 695-702, 2000.

[122978]21818.

[122979]Ellison, J. W.; Wardak, Z.; Webster, M.; Chiong, W.: PHOG, a
candidate

[122980]gene for involvement in the short stature of Turner syndrome. (Abstract) Am.

[122981]J. Hum. Genet. 59 (suppl.): A32, 1996.

[122982]21819.

[122983]Ellison, J. W.; Wardak, Z.; Young, M. F.; Robey, P. G.; Webster,
ster,

[122984]M.; Chiong, W.: PHOG, a candidate gene for involvement in
the short

[122985]stature of Turner syndrome. Hum. Molec. Genet. 6:
1341–1347, 1997.

[122986]21820.

[122987]Guichet, A.; Briault, S.; Le Merrer, M.; Moraine, C.: Are
t(X;Y)

[122988](p22;q11) translocations in females frequently associated
with Madelung

[122989]deformity? Clin. Dysmorphol. 6: 341–345, 1997.

[122990]21821.

[122991]Henke, A.; Wapenaar, M.; van Ommen, G.-J.; Maraschio,
P.; Camerino,

[122992]G.; Rappold, G.: Deletions within the pseudoautosomal re-
gion help

[122993]map three new markers and indicate a possible role of
this region

[122994]in linear growth. Am. J. Hum. Genet. 49: 811–819, 1991.

[122995]21822.

[122996]Affara, N. A.; Chalmers, I. J.; Ferguson-Smith, M. A.: Analysis

[122997]of the SRY gene in 22 sex-reversed XY females identifies four new

[122998]point mutations in the conserved DNA binding domain. Hum. Molec.

[122999]Genet. 2: 785-789, 1993.

[123000]21823.

[123001]Behlke, M. A.; Bogan, J. S.; Beer-Romero, P.; Page, D. C.: Evidence

[123002]that the SRY protein is encoded by a single exon on the human Y chromosome. Genomics 17:

[123003]736-739, 1993.

[123004]21824.

[123005]Berkovitz, G. D.; Fechner, P. Y.; Zacur, H. W.; Rock, J. A.; Snyder,

[123006]H. M., III; Migeon, C. J.; Perlman, E. J.: Clinical and pathologic

[123007]spectrum of 46,XY gonadal dysgenesis: its relevance to the understanding

[123008]of sex differentiation. *Medicine* 70: 375–383, 1991.

[123009]21825.

[123010]Bowles, J.; Cooper, L.; Berkman, J.; Koopman, P.: Sry re-quires

[123011]a CAG repeat domain for male sex determination in *Mus musculus*. *Nature*

[123012]*Genet.* 22: 405–408, 1999.

[123013]21826.

[123014]Braun, A.; Kammerer, S.; Cleve, H.; Lohrs, U.; Schwarz, H.-P.;

[123015]Kuhnle, U.: True hermaphroditism in a 46,XY individual, caused by

[123016]a postzygotic somatic point mutation in the male gonadal sex-determining

[123017]locus (SRY): molecular genetics and histological findings in a sporadic

[123018]case. *Am. J. Hum. Genet.* 52: 578–585, 1993.

[123019]21827.

[123020]Brown, S.; Yu, C. C.; Lanzano, P.; Heller, D.; Thomas, L.; Warburton,

[123021]D.; Kitajewski, J.; Stadtmauer, L.: A de novo mutation

(gln2stop)

[123022]at the 5-prime end of the SRY gene leads to sex reversal
with partial

[123023]ovarian function. (Letter) Am. J. Hum. Genet. 62: 189-192,
1998.

[123024]21828.

[123025]Canto, P.; de la Chesnaye, E.; Lopez, M.; Cervantes, A.;
Chavez,

[123026]B.; Vilchis, F.; Reyes, E.; Ulloa-Aguirre, A.; Kofman-Alfaro,
S.;

[123027]Mendez, J. P.: A mutation in the 5-prime non-high mobil-
ity group

[123028]box region of the SRY gene in patients with Turner syn-
drome and Y

[123029]mosaicism. J. Clin. Endocr. Metab. 85: 1908-1911, 2000.

[123030]21829.

[123031]Cherfas, J.: Sex and the single gene. Science 252: 782,
1991.

[123032]21830.

[123033]Clepet, C.; Schafer, A. J.; Sinclair, A. H.; Palmer, M. S.;
Lovell-Badge,

[123034]R.; Goodfellow, P. N.: The human SRY transcript. Hum.

Molec. Genet. 2:

[123035]2007–2012, 1993.

[123036]21831.

[123037]Davis, R. M.: Localisation of male determining factors in man:

[123038]a thorough review of structural anomalies of the Y chromosome. J.

[123039]Med. Genet. 18: 161–195, 1981.

[123040]21832.

[123041]Dork, T.; Stuhrmann, M.; Miller, K.; Schmidtke, J.: Independent

[123042]observation of SRY mutation I90M in a patient with complete gonadal

[123043]dysgenesis. (Letter) Hum. Mutat. 11: 90–91, 1998.

[123044]21833.

[123045]Eicher, E. M.; Washburn, L. L.: Genetic control of primary sex

[123046]determination in mice. Annu. Rev. Genet. 20: 327–360, 1986.

[123047]21834.

[123048]Foster, J. W.; Brennan, F. E.; Hampikian, G. K.; Goodfellow,

P.

[123049]N.; Sinclair, A. H.; Lovell-Badge, R.; Selwood, L.; Renfree, M. B.;

[123050]Cooper, D. W.; Graves, J. A. M.: Evolution of sex determination and

[123051]the Y chromosome: SRY-related sequences in marsupials. Nature 359:

[123052]531-533, 1992.

[123053]21835.

[123054]Fuqua, J. S.; McLaughlin, J.; Perlman, E. J.; Berkowitz, G. D.

[123055]: Analysis of the SRY gene in gonadal tissue of subjects with 46,XY

[123056]gonadal dysgenesis (Letter) J. Clin. Endocr. Metab. 82: 701-702,

[123057]1997.

[123058]21836.

[123059]Domenice, S.; Nishi, M. Y.; Billerbeck, A. E. C.; Latronico, A.

[123060]C.; Medeiros, M. A.; Russell, A. J.; Vass, K.; Carvalho, F. M.; Frade,

[123061]E. M. C.; Arnhold, I. J. P.; Mendonca, B. B.: A novel mis-

sense mutation

[123062](S18N) in the 5-prime non-HMG box region of the SRY gene in a patient

[123063]with partial gonadal dysgenesis and his normal male relatives. Hum.

[123064]Genet. 102: 213-215, 1998.

[123065]21837.

[123066]Giese, K.; Pagel, J.; Grosschedl, R.: Distinct DNA-binding properties

[123067]of the high mobility group domain of murine and human SRY sex-determining

[123068]factors. Proc. Nat. Acad. Sci. 91: 3368-3372, 1994.

[123069]21838.

[123070]Goodfellow, P. N.; Lovell-Badge, R.: SRY and sex determination

[123071]in mammals. Annu. Rev. Genet. 27: 71-92, 1993.

[123072]21839.

[123073]Gubbay, J.; Collignon, J.; Koopman, P.; Capel, B.; Economou, A.;

[123074]Munsterberg, A.; Vivian, N.; Goodfellow, P. N.; Lovell-Badge, R.:

[123075]A gene mapping to the sex-determining region of the mouse Y chromosome

[123076]is a member of a novel family of embryonically expressed genes. Nature 346:

[123077]245-250, 1990.

[123078]21840.

[123079]Haqq, C. M.; King, C.-Y.; Ukiyama, E.; Falsafi, S.; Haqq, T. N.;

[123080]Donahoe, P. K.; Weiss, M. A.: Molecular basis of mammalian sexual

[123081]determination: activation of Mullerian inhibiting substance gene expression

[123082]by SRY. Science 266: 1494-1500, 1994.

[123083]21841.

[123084]Ramulu, P.; Kennedy, M.; Xiong, W.-H.; Williams, J.; Cowan, M.;

[123085]Blesh, D.; Yau, K.-W.; Hurley, J. B.; Nathans, J.: Normal light response,

[123086]photoreceptor integrity, and rhodopsin dephosphorylation in mice lacking

[123087]both protein phosphatases with EF hands (PPEF-1 and PPEF-2). Molec.

[123088]Cell. Biol. 21: 8605–8614, 2001.

[123089]21842.

[123090]van de Vosse, E.; Franco, B.; van der Bent, P.; Montini, E.;
Orth,

[123091]U.; Hanauer, A.; Tijmes, N.; van Ommen, G.–J. B.; Ballabio,
A.; den

[123092]Dunnen, J. T.; Bergen, A. A. B.: Exclusion of PPEF as the
gene causing

[123093]X-linked juvenile retinoschisis. Hum. Genet. 101:
235–237, 1997.

[123094]21843.

[123095]Spranger, S.; Schiller, S.; Jauch, A.; Wolff, K.; Rauterberg–Ru-
land,

[123096]I.; Hager, D.; Tariverdian, G.; Troger, J.; Rappold, G.: Leri-
Weill

[123097]syndrome as part of a contiguous gene syndrome at
Xp22.3. Am. J.

[123098]Med. Genet. 83: 367–371, 1999.

[123099]21844.

[123100]Stern, M.–H.; Soulier, J.; Rosenzweig, M.; Nakahara, K.;
Canki–Klain,

[123101]N.; Aurias, A.; Sigaux, F.; Kirsch, I. R.: MTCP-1: a novel gene on

[123102]the human chromosome Xq28 translocated to the T cell receptor alpha/delta

[123103]locus in mature T cell proliferations. *Oncogene* 8: 2475-2483, 1993.

[123104]21845.

[123105]Watanabe, T. K.; Fujiwara, T.; Nakamura, Y.; Hirai, Y.; Maekawa,

[123106]H.; Takahashi, E.: Cloning, expression pattern and mapping to Xq

[123107]of NAP1L3, a gene encoding a peptide homologous to human and yeast

[123108]nucleosome assembly proteins. *Cytogenet. Cell Genet.* 74: 281-285,

[123109]1996.

[123110]21846.

[123111]Prakash, S. K.; Paylor, R.; Jenna, S.; Lamarche-Vane, N.; Armstrong,

[123112]D. L.; Xu, B.; Mancini, M. A.; Zoghbi, H. Y.: Functional analysis

[123113]of ARHGAP6, a novel GTPase-activating protein for RhoA.

Hum. Molec.

[123114]Genet. 9: 477–488, 2000.

[123115]21847.

[123116]Schaefer, L.; Prakash, S.; Zoghbi, H. Y.: Cloning and characterization

[123117]of a novel rho-type GTPase-activating protein gene (ARHGAP6) from

[123118]the critical region for microphthalmia with linear skin defects. Genomics 46:

[123119]268–277, 1997.

[123120]21848.

[123121]Aman, M. J.; Tayebi, N.; Obiri, N. I.; Puri, R. K.; Modi, W. S.;

[123122]Leonard, W. J.: cDNA cloning and characterization of the human interleukin

[123123]13 receptor alpha chain. J. Biol. Chem. 271: 29265–29270, 1996.

[123124]21849.

[123125]Guo, J.; Apiou, F.; Mellerin, M.-P.; Lebeau, B.; Jacques, Y.;

[123126]Minvielle, S.: Chromosome mapping and expression of the human interleukin-13

[123127]receptor. Genomics 42: 141–5, 1997.

[123128]21850.

[123129]Hilton, D. J.; Zhang, J.-G.; Metcalf, D.; Alexander, W. S.; Nicola,

[123130]N. A.; Willson, T. A.: Cloning and characterization of a binding

[123131]subunit of the interleukin 13 receptor that is also a component of

[123132]the interleukin 4 receptor. Proc. Nat. Acad. Sci. 93: 497-501, 1996.

[123133]21851.

[123134]Riggs, J. E.; Schochet, S. S., Jr.; Gutmann, L.; Shanske, S.;

[123135]Neal, W. A.; DiMauro, S.: Lysosomal glycogen storage disease without

[123136]acid maltase deficiency. Neurology 33: 873-877, 1983.

[123137]21852.

[123138]Verloes, A.; Massin, M.; Lombet, J.; Grattagliano, B.; Soyeur,

[123139]D.; Rigo, J.; Koulischer, L.; Van Hoof, F.: Nosology of lysosomal

[123140]glycogen storage diseases without in vitro acid maltase deficiency:

[123141]delineation of a neonatal form. Am. J. Med. Genet. 72:
135–142,

[123142]1997.

[123143]21853.

[123144]Kaye, F. J.; Modi, S.; Ivanovska, I.; Koonin, E. V.; Thress, K.;

[123145]Kubo, A.; Kornbluth, S.; Rose, M. D.: A family of ubiquitin-
like

[123146]proteins binds the ATPase domain of Hsp70-like Stch.
FEBS Lett. 467:

[123147]348–352, 2000.

[123148]21854.

[123149]Kaye, F. J.; Shows, T. B.: Assignment of ubiquilin 2
(UBQLN2)

[123150]to human chromosome xp11.23–p11.1 by GeneBridge ra-
diation hybrids. Cytogenet.

[123151]Cell Genet. 89: 116–117, 2000.

[123152]21855.

[123153]Kleijnen, M. F.; Shih, A. H.; Zhou, P.; Kumar, S.; Soccio, R.
E.;

[123154]Kedersha, N. L.; Gill, G.; Howley, P. M.: The hPLIC proteins
may

[123155]provide a link between the ubiquitination machinery and the proteasome. Molec.

[123156]Cell 6: 409–419, 2000.

[123157]21856.

[123158]Lee, D. K.; Nguyen, T.; Lynch, K. R.; Cheng, R.; Vanti, W. B.;

[123159]Arkhitko, O.; Lewis, T.; Evans, J. F.; George, S. R.; O'Dowd, B. F.

[123160]: Discovery and mapping of ten novel G protein–coupled receptor genes. Gene 275:

[123161]83–91, 2001.

[123162]21857.

[123163]Ades, L. C.; Gedeon, A. K.; Wilson, M. J.; Latham, M.; Partington,

[123164]M. W.; Mulley, J. C.; Nelson, J.; Lui, K.; Sillence, D. O.: Barth

[123165]syndrome: clinical features and confirmation of gene localisation

[123166]to distal Xq28. Am. J. Med. Genet. 45: 327–334, 1993.

[123167]21858.

[123168]Barth, P. G.; Scholte, J. A.; Berden, J. A.; Van der Klei–Van Moorsel,

- [123169]J. M.; Luyt-Houwen, I. E. M.; Van't Veer-Korthof, E. T.; Van der Harten,
- [123170]J. J.; Sobotka-Plojhar, M. A.: An X-linked mitochondrial disease
- [123171]affecting cardiac muscle, skeletal muscle and neutrophil leucocytes. J.
- [123172]Neurol. Sci. 62: 327-355, 1983.
- [123173]21859.
- [123174]Barth, P. G.; Van't Veer-Korthof, E. T.; Van Delden, L.; Van Dam,
- [123175]K.; Van der Harten, J. J.; Kuipers, J. R. G.: An X-linked mitochondrial
- [123176]disease affecting cardiac muscle, skeletal muscle and neutrophil leukocytes.In:
- [123177]Busch, H. F. M.; Jennekens, F. G. I.; Schotte, H. R.: Mitochondria
- [123178]and Muscular Diseases. Beetsterzwaag, The Netherlands: Mefar (pub.)
- [123179]1981. Pp. 161-164.
- [123180]21860.
- [123181]Bione, S.; D'Adamo, P.; Maestrini, E.; Gedeon, A. K.; Bolhuis,

- [123182]P. A.; Toniolo, D.: A novel X-linked gene, G4.5.(sic) is responsible
for Barth syndrome. Nature Genet. 12: 385–389, 1996.
- [123184]21861.
- [123185]Bolhuis, P. A.; Hensels, G. W.; Hulsebos, T. J. M.; Baas, F.; Barth,
P. G.: Mapping of the locus for X-linked cardioskeletal myopathy
with neutropenia and abnormal mitochondria (Barth syndrome) to Xq28. Am. J. Hum. Genet. 48: 481–485, 1991.
- [123188]21862.
- [123190]Chen, Y.-T.; Rettig, W. J.; Yenamandra, A. K.; Kozak, C. A.; Chaganti,
R. S. K.; Posner, J. B.; Old, L. J.: Cerebellar degeneration-related antigen: a highly conserved neuroectodermal marker mapped to chromosomes
X in human and mouse. Proc. Nat. Acad. Sci. 87: 3077–3081, 1990.
- [123194]21863.

[123195]Dropcho, E. J.; Chen, Y. T.; Posner, J. B.; Old, L. J.: Cloning
 [123196]of a brain protein identified by autoantibodies from a pa-
 tient with
 [123197]paraneoplastic cerebellar degeneration. Proc. Nat. Acad.
 Sci. 84:
 [123198]4552–4556, 1987.
 [123199]21864.

[123200]Furneau, H. M.; Rosenblum, M. K.; Dalmau, J.; Wong, E.;
 Woodruff,
 [123201]P.; Graus, F.; Posner, J. B.: Selective expression of Purk-
 inje-cell
 [123202]antigens in tumor tissue from patients with paraneoplastic
 cerebellar
 [123203]degeneration. New Eng. J. Med. 322: 1844–1851, 1990.
 [123204]21865.

[123205]Furneau, H. M.; Wong, E.; Posner, J. B.: Isolation of cDNA
 clones
 [123206]encoding the major Yo paraneoplastic antigen. (Abstract)
 Neurology 40
 [123207](suppl. 1): 166 only, 1990.
 [123208]21866.

[123209]Hirst, M. C.; Bell, M. V.; MacKinnon, R. N.; Watson, J. E. V.;

[123210]Callen, D.; Sutherland, G.; Dahl, N.; Patterson, M. N.;

Schwartz,

[123211]C.; Ledbetter, D.; Ledbetter, S.; Davies, K. E.: Mapping of a
cerebellar

[123212]degeneration related protein and DXS304 around the
fragile site. Am.

[123213]J. Med. Genet. 38: 354–356, 1991.

[123214]21867.

[123215]Rettig, W. J.; Chen, Y. T.; Yenamandra, A. K.; Chaganti, R.
S.

[123216]K.; Dropcho, E. J.; Posner, J. B.; Old, L. J.: Assignment of
the

[123217]cerebellar degeneration–related autoantigen CDR to hu-
man chromosome

[123218]X. (Abstract) Cytogenet. Cell Genet. 46: 681 only, 1987.

[123219]21868.

[123220]Siniscalco, M.; Oberle, I.; Melis, P.; Alhadeff, B.; Murray, J.;

[123221]Filippi, G.; Mattioni, T.; Chen, Y.–T.; Furneaux, H.; Old, L.
J.;

[123222]Posner, J. B.: Physical and genetic mapping of the CDR

gene with

[123223]particular reference to its position with respect to the
FRAXA site. Am.

[123224]J. Med. Genet. 38: 357-362, 1991.

[123225]21869.

[123226]Siniscalco, M.; Alhadeff, B.; Kekish, O.; Melis, P.; Maltioni,

[123227]T.; Filippi, G.; Murray, J. C.; Oberle, I.; Chen, Y. T.; Old, L.
J.;

[123228]Posner, J. B.: The CDR gene, located at the upper bound-
ary of the

[123229]FRAXA, is tightly linked to DXS98 and DXS105 RFLPs,
moderately to

[123230]the FRA(X)-syndrome and loosely to the cluster of
DXS15-DXS52 RFLPs.

[123231](Abstract) Cytogenet. Cell Genet. 51: 1081 only, 1989.

[123232]21870.

[123233]Knight, J. C.; Renwick, P. J.; Downing, J. R.; Okuda, T.:
Physical

[123234]linkage of the cdc2-related gene (PCTK1) and the ubiqui-
tin-activating

[123235]enzyme E1 gene (UBE1) on human Xp11.3. Cytogenet. Cell

Genet. 71:

[123236]155–157, 1995.

[123237]21871.

[123238]Aarskog, D.: A familial syndrome of short stature associated with

[123239]facial dysplasia and genital anomalies. J. Pediat. 77:
856–861,

[123240]1970.

[123241]21872.

[123242]Baldellou, A.; Galve, L.; Bassecourt, M.: Risk of medullary
damage

[123243]in Aarskog–Scott syndrome. (Abstract) Clin. Genet. 23:
225 only,

[123244]1983.

[123245]21873.

[123246]Bawle, E.; Tyrkus, M.; Lipman, S.; Bozimowski, D.: Aarskog
syndrome:

[123247]full male and female expression associated with an X–
autosome translocation. Am.

[123248]J. Med. Genet. 17: 595–602, 1984.

[123249]21874.

[123250]Berman, P. A.; Desjardins, C.; Fraser, F. C.: Inheritance of the

[123251]Aarskog syndrome. Birth Defects Orig. Art. Ser. X(7): 151-159, 1974.

[123252]21875.

[123253]Buechner, M.; Hall, D. H.; Bhatt, H.; Hedgecock, E. M.: Cystic

[123254]canal mutants in *Caenorhabditis elegans* are defective in the apical

[123255]membrane domain of the renal (excretory) cell. Dev. Biol. 214: 227-241,

[123256]1999.

[123257]21876.

[123258]Casteels, M.; Samain, H.; Penninckx, F.; Coremans, G.; Beirinckx,

[123259]J.; Fryns, J. P.: Megadolichosigmoid in a young male with Aarskog

[123260]syndrome. Genet. Counsel. 5: 81-83, 1994.

[123261]21877.

[123262]Estrada, L.; Caron, E.; Gorski, J. L.: Fgd1, the Cdc42 guanine

- [123263]nucleotide exchange factor responsible for faciogenital dysplasia,
- [123264]is localized to the subcortical actin cytoskeleton and Golgi membrane. Hum.
- [123265]Molec. Genet. 10: 485–495, 2001.
- [123266]21878.
- [123267]Fernandez, I.; Tsukahara, M.; Mito, H.; Yoshii, H.; Uchida, M.;
- [123268]Matsuo, K.; Kajii, T.: Congenital heart defects in Aarskog syndrome. Am.
- [123269]J. Med. Genet. 50: 318–322, 1994.
- [123270]21879.
- [123271]Fryns, J.–P.: Dolichomegasigmoid in Aarskog syndrome. (Letter) Am.
- [123272]J. Med. Genet. 45: 122 only, 1993.
- [123273]21880.
- [123274]Fryns, J. P.: Aarskog syndrome: the changing phenotype with age. Am.
- [123275]J. Med. Genet. 43: 420–427, 1992.
- [123276]21881.

[123277]Fryns, J. P.; Macken, J.; Vinken, L.; Igodt-Ameye, L.; van
den

[123278]Berghe, H.: The Aarskog syndrome. Hum. Genet. 42:
129-135, 1978.

[123279]21882.

[123280]Funderburk, S. J.; Crandall, B. F.: The Aarskog syndrome in
three

[123281]brothers. Clin. Genet. 6: 119-124, 1974.

[123282]21883.

[123283]Furukawa, C. T.; Hall, B. D.; Smith, D. W.: The Aarskog
syndrome. J.

[123284]Pediat. 81: 1117-1122, 1972.

[123285]21884.

[123286]Gao, J.; Estrada, L.; Cho, S.; Ellis, R. E.; Gorski, J. L.: The

[123287]Caenorhabditis elegans homolog of FGD1, the human
Cdc42 GEF gene responsible

[123288]for faciogenital dysplasia, is critical for excretory cell mor-
phogenesis. Hum.

[123289]Molec. Genet. 10: 3049-3062, 2001.

[123290]21885.

- [123291]Glover, T. W.; Verga, V.; Rafael, J.; Barcroft, C.; Gorski, J.
[123292]L.; Bawle, E. V.; Higgins, J. V.: Translocation breakpoint in
Aarskog
[123293]syndrome maps to Xp11.21 between ALAS2 and DXS323.
Hum. Molec. Genet. 2:
[123294]1717–1718, 1993.
[123295]21886.
[123296]Grier, R. E.; Farrington, F. H.; Kendig, R.; Mamunes, P.:
Autosomal
[123297]dominant inheritance of the Aarskog phenotype.
(Abstract) Am. J.
[123298]Hum. Genet. 33: 64A only, 1981.
[123299]21887.
[123300]Hoo, J. J.: The Aarskog (facio–digito–genital) syndrome.
Clin.
[123301]Genet. 16: 269–276, 1979.
[123302]21888.
[123303]Kodama, M.; Fujimoto, S.; Namikawa, T.; Matsuda, I.:
Aarskog
[123304]syndrome with isolated growth hormone deficiency. Eu-
rop. J. Pediat. 135:

[123305]273–276, 1981.

[123306]21889.

[123307]Lebel, R. R.; May, M.; Pouls, S.; Lubs, H. A.; Stevenson, R. E.;

[123308]Schwartz, C. E.: Non-syndromic X-linked mental retardation associated

[123309]with a missense mutation (P312L) in the FGD1 gene. Clin. Genet. 61:

[123310]139–145, 2002.

[123311]21890.

[123312]Logie, L. J.; Porteous, M. E. M.: Intelligence and development

[123313]in Aarskog syndrome. Arch. Dis. Child. 79: 359–360, 1998.

[123314]21891.

[123315] Mikelsaar, R. V.-A.; Lurie, I. W.: Atypical case of Aarskog syndrome. J.

[123316] Med. Genet. 29: 349-350, 1992.

[123317] 21892.

[123318] Nielsen, K. B.: Aarskog syndrome in a Danish family: an illustration

[123319] of the need for dysmorphology in paediatrics. Clin. Genet. 33: 315-317,

[123320] 1988.

[123321] 21893.

[123322] Oberiter, V.; Lovrencic, M. K.; Schmutzer, L.; Kraus, O.: The

[123323] Aarskog syndrome. Acta Paediat. Scand. 69: 567-570, 1980.

[123324] 21894.

[123325] Orrico, A.; Galli, L.; Falciani, M.; Bracci, M.; Cavaliere, M.

[123326] L.; Rinaldi, M. M.; Musacchio, A.; Sorrentino, V.: A mutation in

[123327] the pleckstrin homology (PH) domain of the FGD1 gene in an Italian

[123328] family with faciogenital dysplasia (Aarskog-Scott syn-

drome). FEBS

[123329]Lett. 478: 216–220, 2000.

[123330]21895.

[123331]Pasteris, N. G.; Cadle, A.; Logie, L. J.; Porteous, M. E. M.;

[123332]Schwartz, C. E.; Stevenson, R. E.; Glover, T. W.; Wilroy, R.

S.; Gorski,

[123333]J. L.: Isolation and analysis of the faciogenital dysplasia

(Aarskog–Scott

[123334]syndrome) gene: a putative, rho/rac guanine nucleotide

exchange factor. Cell 79:

[123335]669–678, 1994.

[123336]21896.

[123337]Pasteris, N. G.; de Gouyon, B.; Cadle A. B.; Campbell, K.;

Herman,

[123338]G. E.; Gorski, J. L.: Cloning and regional localization of the

mouse

[123339]faciogenital dysplasia (Fgd1) gene. Mammalian Genome 6:

658–661,

[123340]1995.

[123341]21897.

[123342]Pedersen, J. C.; Fryns, J. P.; Bracke, P.; Geeraert, M.; Van

Den

[123343]Berghe, H.: The Aarskog syndrome. Ann. Genet. 23: 108–110, 1980.

[123344]21898.

[123345]Porteous, M. E. M.; Goudie, D. R.: Aarskog syndrome. J. Med.

[123346]Genet. 28: 44–47, 1991.

[123347]21899.

[123348]Rafael, J.; Verga, V.; Hall, B.; Burright, E.; Gorski, J.; Bawle,

[123349]J.; Higgins, J. V.; Glover, T. W.: Assignment of the translocation

[123350]breakpoint in a patient with Aarskog syndrome to Xp11.21. (Abstract) Am.

[123351]J. Hum. Genet. 51 (suppl.): A116 only, 1992.

[123352]21900.

[123353]Schwartz, C. E.; Gillessen-Kaesbach, G.; May, M.; Cappa, M.; Gorski,

[123354]J.; Steindl, K.; Neri, G.: Two novel mutations confirm FGD1 is responsible

[123355]for the Aarskog syndrome. Europ. J. Hum. Genet. 8: 869–874, 2000.

[123356]21901.

[123357]Scott, C. I., Jr.: Unusual facies, joint hypermobility, genital
[123358]anomaly and short stature: a new dysmorphic syndrome.

Birth Defects

[123359]Orig. Art. Ser. VII(6): 240–246, 1971.

[123360]21902.

[123361]Asghar, M.; Nevin, N. C.; Beattie, E. D.; McManus, D.;
Roberts,

[123362]G. M. A.; Phillips, J. A.: Contiguous gene deletion syn-
drome involving

[123363]glycerol kinase and Duchenne muscular dystrophy loci. J.
Inherit.

[123364]Metab. Dis. 22: 933–935, 1999.

[123365]21903.

[123366]Balducci, R.; Municchi, G.; Toscano, V.; Mangiantini, A.;
Sabatini,

[123367]R.; Caiola, S.; Tedeschi, B.; Orlandi, L.; Boscherini, B.:
Complex

[123368]glycerol kinase deficiency: an unusual cause of salt-
wasting in males. Clin.

[123369]Endocr. 42: 437–439, 1995.

[123370]21904.

[123371]Bartley, J. A.; Patil, S.; Davenport, S.; Goldstein, D.; Pick-
ens,

[123372]J.: Duchenne muscular dystrophy, glycerol kinase defi-
ciency, and

[123373]adrenal insufficiency associated with Xp21 interstitial
deletion. J.

[123374]Pediat. 108: 189-192, 1986.

[123375]21905.

[123376]Borresen, A.-L.; Hellerud, C.; Moller, P.; Sovik, O.; Berg, K.

[123377]: Prenatal diagnosis of glycerol-kinase deficiency associ-
ated with

[123378]a DNA deletion on the short arm of the X chromosome.
Clin. Genet. 32:

[123379]254-259, 1987.

[123380]21906.

[123381]Darras, B. T.; Francke, U.: Myopathy in complex glycerol
kinase

[123382]deficiency patients is due to 3-prime deletions of the dys-
trophin

[123383]gene. Am. J. Hum. Genet. 43: 126-130, 1988.

[123384]21907.

[123385]Davies, K. E.; Patterson, M. N.; Kenwrick, S. J.; Bell, M. V.;

[123386]Sloan, H. R.; Westman, J. A.; Elsas, L. J., II; Mahan, J.: Fine
mapping

[123387]of glycerol kinase deficiency and congenital adrenal hy-
poplasia within

[123388]Xp21 on the short arm of the human X chromosome. Am.
J. Med. Genet. 29:

[123389]557-564, 1988.

[123390]21908.

[123391]Dipple, K. M.; Zhang, Y.-H.; Huang, B.-L.; McCabe, L. L.;
Dallongeville,

[123392]J.; Inokuchi, T.; Kimura, M.; Marx, H. J.; Roederer, G. O.;
Shih,

[123393]Y.; Yamaguchi, S.; Yoshida, I.; McCabe, E. R. B.: Glycerol
kinase

[123394]deficiency: evidence for complexity in a single gene disor-
der. Hum.

[123395]Genet. 109: 55-62, 2001.

[123396]21909.

[123397]Arias, J.; Alberts, A. S.; Brindle, P.; Claret, F. X.; Smeal, T.;

[123398]Karin, M.; Feramisco, J.; Montminy, M.: Activation of cAMP and mitogen

[123399]responsive genes relies on a common nuclear factor. Nature 370:

[123400]226–229, 1994.

[123401]21910.

[123402]Blough, R. I.; Petrij, F.; Dauwerse, J. G.; Milatovich–Cherry,

[123403]A.; Weiss, L.; Saal, H. M.; Rubinstein, J. H.: Variation in microdeletions

[123404]of the cyclic AMP–responsive element–binding protein gene at chromosome

[123405]band 16p13.3 in the Rubinstein–Taybi syndrome. Am. J. Med. Genet. 90:

[123406]29–34, 2000.

[123407]21911.

[123408]Chen, X.–N.; Korenberg, J. R.: Localization of human CREBBP (CREB

[123409]binding protein) to 16p13.3 by fluorescence in situ hybridization. Cytogenet.

[123410]Cell Genet. 71: 56–57, 1995.

[123411]21912.

- [123412]Chrivia, J. C.; Kwok, R. P. S.; Lamb, N.; Hagiwara, M.;
Montminy,
[123413]M. R.; Goodman, R. H.: Phosphorylated CREB binds specif-
ically to
[123414]the nuclear protein CBP. *Nature* 365: 855–859, 1993.
[123415]21913.
- [123416]Borrow, J.; Stanton, V. P., Jr.; Andresen, J. M.; Becher, R.;
Behm,
[123417]F. G.; Chaganti, R. S. K.; Civin, C. I.; Distech, C.; Dube, I.;
Frischauf,
[123418]A. M.; Horsman, D.; Mitelman, F.; Volinia, S.; Watmore, A.
E.; Housman,
[123419]D. E.: The translocation t(8;16)(p11;p13) of acute myeloid
leukaemia
[123420]fuses a putative acetyltransferase to the CREB-binding
protein. *Nature*
[123421]*Genet.* 14: 33–41, 1996.
[123422]21914.
- [123423]Demarest, S. J.; Martinez–Yamout, M.; Chung, J.; Chen, H.;
Xu,
[123424]W.; Dyson, H. J.; Evans, R. M.; Wright, P. E.: Mutual syner-
gistic

- [123425]folding in recruitment of CBP/p300 by p160 nuclear receptor coactivators. *Nature* 415: 549–553, 2002.
- [123426]21915.
- [123427]Eckner, R.; Ewen, M. E.; Newsome, D.; Gerdes, M.; Decaprio, J. A.; Lawrence, J. B.; Livingston, D. M.: Molecular cloning and functional analysis of the adenovirus E1A-associated 300-kD protein (p300) reveals a protein with properties of a transcriptional adaptor. *Genes Dev.* 15: 869–884, 1994.
- [123432]21916.
- [123433]Giles, R. H.; Dauwerse, H. G.; van Ommen, G.-J. B.; Breuning, M. H.: Do human chromosomal bands 16p13 and 22q11–13 share ancestral origins? (Letter) *Am. J. Hum. Genet.* 63: 1240–1242, 1998.
- [123437]21917.
- [123438]Giles, R. H.; Petrij, F.; Dauwerse, H. G.; den Hollander, A.

I.;

[123439]Lushnikova, T.; van Ommen, G.-J. B.; Goodman, R. H.;
Deaven, L. L.;

[123440]Doggett, N. A.; Peters, D. J.; Breuning, M. H.: Construction
of a

[123441]1.2-Mb contig surrounding, and molecular analysis of, the
human CREB-binding

[123442]protein (CBP/CREBBP) gene on chromosome 16p13.3. Ge-
nomics 42: 96-114,

[123443]1997.

[123444]21918.

[123445]Gonzalez, G. A.; Montminy, M. R.: Cyclic AMP stimulates
somatostatin

[123446]gene transcription by phosphorylation of CREB at serine
133. Cell 59:

[123447]675-680, 1989.

[123448]21919.

[123449]Kung, A. L.; Rebel, V. I.; Bronson, R. T.; Ch'ng, L.-E.; Sieff,

[123450]C. A.; Livingston, D. M.; Yao, T.-P.: Gene dose-dependent
control

[123451]of hematopoiesis and hematologic tumor suppression by

CBP. Genes

[123452]Dev. 14: 272–277, 2000.

[123453]21920.

[123454]Kwok, R. P. S.; Lundblad, J. R.; Chrivia, J. C.; Richards, J.

[123455]P.; Bachinger, H. P.; Brennan, R. G.; Roberts, S. G. E.;

Green, M.

[123456]R.; Goodman, R. H.: Nuclear protein CBP is a coactivator
for the

[123457]transcription factor CREB. Nature 370: 223–226, 1994.

[123458]21921.

[123459]Lin, C. H.; Hare, B. J.; Wagner, G.; Harrison, S. C.; Maniatis,

[123460]T.; Fraenkel, E.: A small domain of CBP/p300 binds di-
verse proteins:

[123461]solution structure and functional studies. Molec. Cell 8:
581–590,

[123462]2001.

[123463]21922.

[123464]Lundblad, J. R.; Kwok, R. P. S.; Lurance, M. E.; Harter, M.

L.;

[123465]Goodman, R. H.: Adenoviral E1A-associated protein p300

as a functional

[123466]homologue of the transcriptional co-activator CBP. Nature
374: 85–88,

[123467]1995.

[123468]21923.

[123469]Murata, T.; Kurokawa, R.; Krones, A.; Tatsumi, K.; Ishii, M.;

[123470]Taki, T.; Masuno, M.; Ohashi, H.; Yanagisawa, M.; Rosen-
feld, M. G.;

[123471]Glass, C. K.; Hayashi, Y.: Defect of histone acetyltrans-
ferase activity

[123472]of the nuclear transcriptional coactivator CBP in Rubin-
stein–Taybi

[123473]syndrome. Hum. Molec. Genet. 10: 1071–1076, 2001.

[123474]21924.

[123475]Nordheim, A.: CREB takes CBP to tango. Nature 370:
177–178,

[123476]1994.

[123477]21925.

[123478]Ohno, S.: Patterns in genome evolution. Curr. Opin. Genet.
Dev. 3:

[123479]911–914, 1993.

[123480]21926.

[123481]Oike, Y.; Hata, A.; Mamiya, T.; Kaname, T.; Noda, Y.;
Suzuki,

[123482]M.; Yasue, H.; Nabeshima, T.; Araki, K.; Yamamura, K.:
Truncated

[123483]CBP protein leads to classical Rubinstein–Taybi syndrome
phenotypes

[123484]in mice: implications for a dominant–negative mechanism.
Hum. Molec.

[123485]Genet. 8: 387–396, 1999.

[123486]21927.

[123487]Kamb, A.; Shattuck–Eidens, D.; Eeles, R.; Liu, Q.; Gruis, N.
A.;

[123488]Ding, W.; Hussey, C.; Tran, T.; Miki, Y.; Weaver–Feldhaus,
J.; McClure,

[123489]M.; Aitken, J. F.; Anderson, D. E.; Bergman, W.; Frants, R.;
Goldgar,

[123490]D. E.; Green, A.; MacLennan, R.; Martin, N. G.; Meyer, L. J.;
Youl,

[123491]P.; Zone, J. J.; Skolnick, M. H.; Cannon–Albright, L. A.:
Analysis

[123492]of the p16 gene (CDKN2) as a candidate for the chromo–

some 9p melanoma

[123493]susceptibility locus. Nature Genet. 8: 22–26, 1994.

[123494]21928.

[123495]Riesewijk, A. M.; Blagitko, N.; Schinzel, A. A.; Hu, L.;
Schulz,

[123496]U.; Hamel, B. C. J.; Ropers, H.–H.; Kalscheuer, V. M.: Evi-
dence against

[123497]a major role of PEG1/MEST Silver–Russell syndrome. Eu-
rop. J. Hum.

[123498]Genet. 6: 114–120, 1998.

[123499]21929.

[123500]Riesewijk, A. M.; Hu, L.; Schulz, U.; Tariverdian, G.;
Hoglund,

[123501]P.; Kere, J.; Ropers, H.–H.; Kalscheuer, V. M.: Monoallelic
expression

[123502]of human PEG1/MEST is paralleled by parent–specific
methylation in

[123503]fetuses. Genomics 42: 236–244, 1997.

[123504]21930.

[123505]Sado, T.; Nakajima, N.; Tada, M.; Takagi, N.: A novel
mesoderm–specific

[123506]cDNA isolated from a mouse embryonal carcinoma cell
line. Dev. Growth

[123507]Differ. 35: 551–560, 1993.

[123508]21931.

[123509]Kamijo, T.; Zindy, F.; Roussel, M. F.; Quelle, D. E.; Down-
ing,

[123510]J. R.; Ashmun, R. A.; Grosveld, G.; Sherr, C. J.: Tumor sup-
pression

[123511]at the mouse INK4a locus mediated by the alternative
reading frame

[123512]product p19(ARF). Cell 91: 649–659, 1997.

[123513]21932.

[123514]Koh, J.; Enders, G. H.; Dynlacht, B. D.; Harlow, E.: Tumour-
derived

[123515]p16 alleles encoding proteins defective in cell-cycle inhi-
bition. Nature 375:

[123516]506–510, 1995.

[123517]21933.

[123518]Komiya, A.; Suzuki, H.; Aida, S.; Yatani, R.; Shimazaki, J.:

[123519]Mutational analysis of CDKN2 (CDK4I/MTS1) gene in tis-
sues and cell

[123520]lines of human prostate cancer. Jpn. J. Cancer Res. 86:
622–625,

[123521]1995.

[123522]21934.

[123523]Krimpenfort, P.; Quon, K. C.; Mooi, W. J.; Loonstra, A.;
Berns,

[123524]A.: Loss of p16(Ink4a) confers susceptibility to metastatic
melanoma

[123525]in mice. Nature 413: 83–85, 2001.

[123526]21935.

[123527]Labarriere, N.; Selvais, P. L.; Lemaigre, F. P.; Michel, A.;
Maiter,

[123528]D. M.; Rousseau, G. G.: A novel transcriptional activator
originating

[123529]from an upstream promoter in the human growth hor-
mone gene. J. Biol.

[123530]Chem. 270: 19205–19208, 1995.

[123531]21936.

[123532]Little, M.; Wainwright, B.: Methylation and p16: suppress-
ing

[123533]the suppressor. Nature Med. 1: 633–634, 1995.

[123534]21937.

[123535]Liu, L.; Dilworth, D.; Gao, L.; Monzon, J.; Summers, A.;
Lassam,

[123536]N.; Hogg, D.: Mutation of the CDKN2A 5-prime UTR cre-
ates an aberrant

[123537]initiation codon and predisposes to melanoma. Nature
Genet. 21:

[123538]128-132, 1999.

[123539]21938.

[123540]Liu, L.; Lassam, N. J.; Slingerland, J. M.; Bailey, D.; Cole,

[123541]D.; Jenkins, R.; Hogg, D.: Germline p16(INK4A) mutation
and protein

[123542]dysfunction in a family with inherited melanoma. Onco-
gene 11: 405-412,

[123543]1995.

[123544]21939.

[123545]Lukas, J.; Parry, D.; Aagaard, L.; Mann, D. J.; Bartkova, J.;

[123546]Strauss, M.; Peters, G.; Bartek, J.: Retinoblastoma-pro-
tein-dependent

[123547]cell-cycle inhibition by the tumour suppressor p16. Na-
ture 375:

[123548]503–506, 1995.

[123549]21940.

[123550]MacGeoch, C.; Bishop, J. A.; Bataille, V.; Bishop, D. T.;
Frischauf,

[123551]A. M.; Meloni, R.; Cuzick, J.; Pinney, E.; Spurr, N. K.: Ge-
netic

[123552]heterogeneity in familial malignant melanoma. Hum.
Molec. Genet. 3:

[123553]2195–2200, 1994.

[123554]21941.

[123555]MacKie, R. M.; Andrew, N.; Lanyon, W. G.; Connor, J. M.:
CDKN2A

[123556]germline mutations in U.K. patients with familial
melanoma and multiple

[123557]primary melanomas. J. Invest. Derm. 111: 269–272, 1998.

[123558]21942.

[123559]Mantelli, M.; Barile, M.; Ciotti, P.; Ghiorzo, P.; Lantieri, F.;

[123560]Pastorino, L.; Catricala, C.; Della Torre, G.; Folco, U.;
Grammatico,

[123561]P.; Padovani, L.; Pasini, B.; and 9 others: High prevalence
of

- [123562]the G101W germline mutation in the CDKN2A (P16ink4a) gene in 62 Italian
- [123563]malignant melanoma families. *Am. J. Med. Genet.* 107: 214–221, 2002.
- [123564]21943.
- [123565]Merlo, A.; Herman, J. G.; Mao, L.; Lee, D. J.; Gabrielson, E.;
- [123566]Burger, P. C.; Baylin, S. B.; Sidransky, D.: 5-prime CpG island methylation
- [123567]is associated with transcriptional silencing of the tumour suppressor
- [123568]p16/CDKN2/MTS1 in human cancers. *Nature Med.* 1: 686–692, 1995.
- [123569]21944.
- [123570]Monzon, J.; Liu, L.; Brill, H.; Goldstein, A. M.; Tucker, M. A.;
- [123571]From, L.; McLaughlin, J.; Hogg, D.; Lassam, N. J.: CDKN2A mutations
- [123572]in multiple primary melanomas. *New Eng. J. Med.* 338: 879–887, 1998.
- [123573]21945.
- [123574]Mori, T.; Miura, K.; Aoki, T.; Nishihira, T.; Mori, S.; Nakamura,

[123575]Y.: Frequent somatic mutation of the MTS1/CDK41
(multiple tumor suppressor/cyclin-dependent
[123576]kinase 4 inhibitor) gene in esophageal squamous cell car-
cinoma. Cancer
[123577]Res. 54: 3396-3397, 1994.
[123578]21946.

[123579]Nobori, T.; Miura, K.; Wu, D. J.; Lois, A.; Takabayashi, K.;
Carson,
[123580]D. A.: Deletions of the cyclin-dependent kinase-4 in-
hibitor gene
[123581]in multiple human cancers. Nature 368: 753-756, 1994.
[123582]21947.

[123583]Ogawa, S.; Hirano, N.; Sato, N.; Takahashi, T.; Hangaishi,
A.;
[123584]Tanaka, K.; Kurokawa, M.; Tanaka, T.; Mitani, K.; Yazaki,
Y.; Hirai,
[123585]H.: Homozygous loss of the cyclin-dependent kinase
4-inhibitor (p16)
[123586]gene in human leukemias. Blood 84: 2431-2435, 1994.
[123587]21948.

[123588]Ohhara, M.; Esumi, M.; Kurosu, Y.: Activation but not inac-

tivation

[123589] of the MTS1 gene is associated with primary colorectal carcinomas. Biochem.

[123590] Biophys. Res. Commun. 226: 791–795, 1996.

[123591] 21949.

[123592] Okuda, T.; Shurtleff, S. A.; Valentine, M. B.; Raimondi, S. C.;

[123593] Head, D. R.; Behm, F.; Curcio-Brint, A. M.; Liu, Q.; Pui, C.-H.; Sherr,

[123594] C. J.; Beach, D.; Look, A. T.; Downing, J. R.: Frequent deletion

[123595] of p16(INK4a)/MTS1 and p15(INK4b)/MTS2 in pediatric acute lymphoblastic

[123596] leukemia. Blood 85: 2321–2330, 1995.

[123597] 21950.

[123598] Petronzelli, F.; Sollima, D.; Coppola, G.; Martini-Neri, M. E.;

[123599] Neri, G.; Genuardi, M.: CDKN2A germline splicing mutation affecting

[123600] both P16(ink4) and P14(arf) RNA processing in a melanoma/neurofibroma

[123601] kindred. Genes Chromosomes Cancer 31: 398–401, 2001.

[123602]21951.

[123603]Petty, E. M.; Gibson, L. H.; Fountain, J. W.; Bolognia, J. L.;

[123604]Yang-Feng, T. L.; Housman, D. E.; Bale, A. E.: Molecular
definition

[123605]of a chromosome 9p21 germ-line deletion in a woman
with multiple melanomas

[123606]and a plexiform neurofibroma: implications for 9p tumour
suppressor

[123607]gene(s). Am. J. Hum. Genet. 53: 96–104, 1993.

[123608]21952.

[123609]Pilon, C.; Pistorello, M.; Moscon, A.; Altavilla, G.; Pagotto,

[123610]U.; Boscaro, M.; Fallo, F.: Inactivation of the p16 tumor
suppressor

[123611]gene in adrenocortical tumors. J. Clin. Endocr. Metab. 84:
2776–2779,

[123612]1999.

[123613]21953.

[123614]Pomerantz, J.; Schreiber-Agus, N.; Liegeois, N. J.; Silver-
man,

[123615]A.; Alland, L.; Chin, L.; Potes, J.; Chen, K.; Orlow, I.; Lee,
H.-W.;

[123616]Cordon-Cardo, C.; DePinho, R. A.: The Ink4a tumor suppressor gene

[123617]product, p19(Arf), interacts with MDM2 and neutralizes MDM2's inhibition

[123618]of p53. Cell 92: 713-723, 1998.

[123619]21954.

[123620]Jayakumar, A.; Chirala, S. S.; Chinault, A. C.; Baldini, A.; Abu-Elheiga,

[123621]L.; Wakil, S. J.: Isolation and chromosomal mapping of genomic clones

[123622]encoding the human fatty acid synthase gene. Genomics 23: 420-424,

[123623]1994.

[123624]21955.

[123625]Jayakumar, A.; Tai, M.-H.; Huang, W.-Y.; Al-Feel, W.; Hsu, M.;

[123626]Abu-Elheiga, L.; Chirala, S. S.; Wakil, S. J.: Human fatty acid synthase:

[123627]properties and molecular cloning. Proc. Nat. Acad. Sci. 92: 8695-8699,

[123628]1995.

[123629]21956.

[123630]Loftus, T. M.; Jaworsky, D. E.; Frehywot, G. L.; Townsend, C. A.;

[123631]Ronnett, G. V.; Lane, M. D.; Kuhajda, F. P.: Reduced food intake

[123632]and body weight in mice treated with fatty acid synthase inhibitors. Science 288:

[123633]2379–2381, 2000.

[123634]21957.

[123635]Wakil, S. J.: Fatty acid synthase, a proficient multifunctional

[123636]enzyme. Biochemistry 28: 4523–4530, 1989.

[123637]21958.

[123638]Ye, Q.; Chung, L. W. K.; Li, S.; Zhau, H. E.: Identification of

[123639]a novel FAS/ER- α fusion transcript expressed in human cancer cells. Biochim.

[123640]Biophys. Acta 1493: 373–377, 2000.

[123641]21959.

[123642]Argeson, A. C.; Druck, T.; Veronese, M. L.; Knopf, J. L.; Buchberg,

- [123643]A. M.; Huebner, K.; Siracusa, L. D.: Phospholipase C gamma-2 (Plcg2) and phospholipase C gamma-1 (Plcg1) map to distinct regions in the human and mouse genomes. *Genomics* 25: 29-35, 1995.
- [123646]21960.
- [123647]Hernandez, D.; Egan, S. E.; Yulug, I. G.; Fisher, E. M. C.: Mapping the gene that encodes phosphatidylinositol-specific phospholipase C-gamma-2 in the human and the mouse. *Genomics* 23: 504-507, 1994.
- [123650]21961.
- [123651]Kang, J. S.; Kohlhuber, F.; Hug, H.; Marme, D.; Eick, D.; Ueffing, M.: Cloning and functional analysis of the hematopoietic cell-specific phospholipase C-gamma-2 promoter. *FEBS Lett.* 399: 14-20, 1996.
- [123654]21962.
- [123655]Dumont, D. J.; Anderson, L.; Breitman, M. L.; Duncan, A. M. V.

- [123656] Assignment of the endothelial-specific protein receptor tyrosine kinase gene (TEK) to human chromosome 9p21. Genomics 23: 512-513, 1994.
- [123659] 1963.
- [123660] Dumont, D. J.; Yamaguchi, T. P.; Conlon, R. A.; Rossant, J.; Breitman, M. L.: tek, a novel tyrosine kinase gene located on mouse chromosome 4, is expressed in endothelial cells and their presumptive precursors. Oncogene 7: 1471-1480, 1992.
- [123664] 1964.
- [123665] Delot, E.; King, L. M.; Briggs, M. D.; Wilcox, W. R.; Cohn, D. H.: Trinucleotide expansion mutations in the cartilage oligomeric matrix protein (COMP) gene. Hum. Molec. Genet. 8: 123-128, 1999.
- [123668] 1965.
- [123669] Oliner, J. D.; Andresen, J. M.; Hansen, S. K.; Zhou, S.; Tjian,

[123670]R.: SREBP transcriptional activity is mediated through an interaction

[123671]with the CREB-binding protein. *Genes Dev.* 10: 2903–2911, 1996.

[123672]21966.

[123673]Panagopoulos, I.; Fioretos, T.; Isaksson, M.; Samuelsson, U.;

[123674]Billstrom, R.; Strombeck, B.; Mitelman, F.; Johansson, B.: Fusion

[123675]of the MORF and CBP genes in acute myeloid leukemia with the t(10;16)(q22;p13). *Hum.*

[123676]*Molec. Genet.* 10: 395–404, 2001.

[123677]21967.

[123678]Petrij, F.; Dorsman, J. C.; Dauwerse, H. G.; Giles, R. H.; Peeters,

[123679]T.; Hennekam, R. C. M.; Breuning, M. H.; Peters, D. J. M.: Rubinstein–Taybi

[123680]syndrome caused by a de novo reciprocal translocation t(2;16)(q36.3;p13.3). *Am.*

[123681]*J. Med. Genet.* 92: 47–52, 2000.

[123682]21968.

- [123683]Sobulo, O. M.; Borrow, J.; Tomek, R.; Reshmi, S.; Harden, A.;
- [123684]Schlegelberger, B.; Housman, D.; Doggett, N. A.; Rowley, J. D.; Zeleznik-Le, N. J.: MLL is fused to CBP, a histone acetyltransferase, in therapy-related acute myeloid leukemia with a t(11;16)(q23;p13.3). Proc. Nat. Acad. Sci. 94: 8732-8737, 1997.
- [123688]21969.
- [123689]Taine, L.; Goizet, C.; Wen, Z. Q.; Petrij, F.; Breuning, M. H.; Ayme, S.; Saura, R.; Arveiler, B.; Lacombe, D.: Submicroscopic deletion of chromosome 16p13.3 in patients with Rubinstein-Taybi syndrome. Am. J. Med. Genet. 78: 267-270, 1998.
- [123693]21970.
- [123694]Tini, M.; Benecke, A.; Um, S.-J.; Torchia, J.; Evans, R. M.; Chambon, P.: Association of CBP/p300 acetylase and thymine DNA glycosylase

[123696]links DNA repair and transcription. Molec. Cell 9:
265–277, 2002.

[123697]21971.

[123698]Weaver, B. K.; Kumar, K. P.; Reich, N. C.: Interferon regula-
tory

[123699]factor 3 and CREB-binding protein/p300 are subunits of
double-stranded

[123700]RNA-activated transcription factor DRAF1. Molec. Cell.
Biol. 18:

[123701]1359–1368, 1998.

[123702]21972.

[123703]Wydner, K. L.; Bhattacharya, S.; Eckner, R.; Lawrence, J. B.;

[123704]Livingston, D. M.: Localization of human CREB-binding
protein gene

[123705](CREBBP) to 16p13.2–p13.3 by fluorescence in situ hy-
bridization. Genomics 30:

[123706]395–396, 1995.

[123707]21973.

[123708]Yamauchi, T.; Oike, Y.; Kamon, J.; Waki, H.; Komeda, K.;
Tsuchida,

[123709]A.; Date, Y.; Li, M.-X.; Miki, H.; Akanuma, Y.; Nagai, R.;

Kimura,

[123710]S.; Saheki, T.; Nakazato, M.; Naitoh, T.; Yamamura, K.;
Kadowaki,

[123711]T.: Increased insulin sensitivity despite lipodystrophy in
Crebbp

[123712]heterozygous mice. Nature Genet. 30: 221–226, 2002.

[123713]21974.

[123714]Zanger, K.; Radovick, S.; Wondisford, F. E.: CREB binding
protein

[123715]recruitment to the transcription complex requires growth
factor-dependent

[123716]phosphorylation of its GF box. Molec. Cell 7: 551–558,
2001.

[123717]21975.

[123718]Hansen, J. J.; Durr, A.; Cournu-Rebeix, I.; Georgopoulos,
C.; Ang,

[123719]D.; Nielsen, M. N.; Davoine, C.-S.; Brice, A.; Fontaine, B.;
Gregersen,

[123720]N.; Bross, P.: Hereditary spastic paraplegia SPG13 is asso-
ciated

[123721]with a mutation in the gene encoding the mitochondrial
chaperonin

[123722]Hsp60. Am. J. Hum. Genet. 70: 1328–1332, 2002.

[123723]21976.

[123724]Todd, M. J.; Viitanen, P. V.; Lorimer, G. H.: Dynamics of the

[123725]chaperonin ATPase cycle: implications for facilitated protein folding. Science 265:

[123726]659–666, 1994.

[123727]21977.

[123728]Hellevuo, K.; Berry, R.; Sikela, J. M.; Tabakoff, B.: Localization

[123729]of the gene for a novel human adenylyl cyclase (ADCY7) to chromosome

[123730]16. Hum. Genet. 95: 197–200, 1995.

[123731]21978.

[123732]Hellevuo, K.; Yoshimura, M.; Kao, M.; Hoffman, P. L.; Cooper, D.

[123733]M. F.; Tabakoff, B.: A novel adenylyl cyclase sequence cloned from

[123734]the human erythroleukemia cell line. Biochem. Biophys. Res. Commun. 192:

[123735]311–318, 1993.

[123736]21979.

[123737]Moerman, P.; Fryns, J.-P.; Sastrowijoto, S. H.; Vandenberghe, K.;

[123738]Lauweryns, J. M.: Hereditary renal adysplasia: new observations and

[123739]hypotheses. *Pediat. Path.* 14: 405-410, 1994.

[123740]21980.

[123741]Kagawa, W.; Kurumizaka, H.; Ishitani, R.; Fukai, S.; Nureki, O.;

[123742]Shibata, T.; Yokoyama, S.: Crystal structure of the homologous-pairing

[123743]domain from the human Rad52 recombinase in the undecameric form. *Molec.*

[123744]*Cell* 10: 359-371, 2002.

[123745]21981.

[123746]Shen, Z.; Denison, K.; Lobb, R.; Gatewood, J. M.; Chen, D. J.:

[123747]The human and mouse homologs of the yeast RAD52 gene: cDNA cloning,

[123748]sequence analysis, assignment to human chromosome 12p12.2-p13, and

[123749]mRNA expression in mouse tissues. Genomics 25:
199–206, 1995.

[123750]21982.

[123751]Stapleton, D.; Woollatt, E.; Mitchelhill, K. I.; Nicholl, J. K.;

[123752]Fernandez, C. S.; Michell, B. J.; Witters, L. A.; Power, D. A.;
Sutherland,

[123753]G. R.; Kemp, B. E.: AMP-activated protein kinase isoen-
zyme family:

[123754]subunit structure and chromosomal location. FEBS Lett.
409: 452–456,

[123755]1997.

[123756]21983.

[123757]Cox, G. A.; Mahaffey, C. L.; Frankel, W. N.: Identification of

[123758]mouse neuromuscular degeneration gene and mapping of
a second site

[123759]suppressor allele. Neuron 21: 1327–1337, 1998.

[123760]21984.

[123761]Fukita, Y.; Mituza, T.; Shirozu, M.; Ozawa, K.; Shimuzu, A.;
Honjo,

[123762]T.: The human S-mu-bp2, a DNA-binding protein specific
to the single-stranded

[123763]guanine-rich sequence related to the immunoglobulin mu chain switch

[123764]region. J. Biol. Chem. 268: 17463-17470, 1993.

[123765]2 1985.

[123766]Grohmann, K.; Schuelke, M.; Diers, A.; Hoffmann, K.; Lucke, B.;

[123767]Adams, C.; Bertini, E.; Leonhardt-Horti, H.; Muntoni, F.; Ouvrier,

[123768]R.; Pfeufer, A.; Rossi, R.; Van Maldergem, L.; Wilmshurst, J. M.;

[123769]Wienker, T. F.; Sendtner, M.; Rudnik-Schoneborn, S.; Zerres, K.; Hubner,

[123770]C.: Mutations in the gene encoding immunoglobulin mu-binding protein

[123771]2 cause spinal muscular atrophy with respiratory distress type 1. Nature

[123772]Genet. 29: 75-77, 2001.

[123773]2 1986.

[123774]McBride, K.; Robitaille, L.; Tremblay, S.; Argentin, S.; Nemer,

[123775]M.: Fos/jun repression of cardiac specific transcription is targeted

[123776]at a tissue-specific cis element. Mol. Cell. Biol. 13:
600-612,
[123777]1993.
[123778]21987.
[123779]Sebastiani, G.; Durocher, D.; Gros, P.; Nemer, M.; Malo, D.:
Localization
[123780]of the Catf1 transcription factor gene to mouse chromo-
some 19. Mammalian
[123781]Genome 6: 147-148, 1995.
[123782]21988.
[123783]Zonana, J.; Jones, M.; Browne, D.; Litt, M.; Kramer, P.;
Becker,
[123784]H. W.; Brockdorff, N.; Rastan, S.; Davies, K. P.; Clarke, A.;
Thomas,
[123785]N. S. T.: High-resolution mapping of the X-linked hypo-
hidrotic ectodermal
[123786]dysplasia (EDA) locus. Am. J. Hum. Genet. 51: 1036-1046,
1992.
[123787]21989.
[123788]Zonana, J.; Roberts, S. H.; Thomas, N. S. T.; Harper, P. S.:
[123789]Recognition and reanalysis of a cell line from a manifest-

ing female

[123790]with X linked hypohidrotic ectodermal dysplasia and an
X;autosome

[123791]balanced translocation. J. Med. Genet. 25: 383-386, 1988.

[123792]21990.

[123793]Zonana, J.; Schnizel, A.; Upadhyaya, M.; Thomas, N. S. T.;
Anton-Lamprecht,

[123794]I.; Harper, P. S.: Prenatal diagnosis of X-linked hypo-
hidrotic ectodermal

[123795]dysplasia by linkage analysis. Am. J. Med. Genet. 35:
132-135, 1990.

[123796]21991.

[123797]Derry, J. M.; Barnard, P. J.: The gene for tissue inhibitor of

[123798]metalloproteinases (TIMP) is located within an intron of
the synapsin

[123799]I gene on the X chromosome. (Abstract) Cytogenet. Cell
Genet. 58:

[123800]2061-2062, 1991.

[123801]21992.

[123802]Gal, A.; Wieringa, B.; Smeets, D. F. C. M.; Bleeker-
Wagemakers,

[123803]L.; Ropers, H. H.: Submicroscopic interstitial deletion of the X chromosome explains a complex genetic syndrome dominated by Norrie disease. Cytogenet. Cell Genet. 42: 219–224, 1986.

[123806]21993.

[123807]Rugarli, E. I.; Ghezzi, C.; Valsecchi, V.; Ballabio, A.: The Kallmann syndrome gene product expressed in COS cells is cleaved on the cell surface to yield a diffusible component. Hum. Molec. Genet. 5: 1109–1115, 1996.

[123811]21994.

[123812]Schroffner, W. G.; Furth, E. D.: Hypogonadotropic hypogonadism with anosmia (Kallmann's syndrome) unresponsive to clomiphene citrate. J. Clin. Endocr. 31: 267–270, 1970.

[123815]21995.

[123816]Schwanzel-Fukuda, M.; Bick, D.; Pfaff, D. W.: Luteinizing hormone-releasing

[123817]hormone (LHRH)-expressing cells do not migrate normally
in an inherited

[123818]hypogonadal (Kallmann) syndrome. Brain Res. Molec. Brain
Res. 6:

[123819]311–326, 1989.

[123820]21996.

[123821]Soussi-Yanicostas, N.; de Castro, F.; Julliard, A. K.; Perfet-
tini,

[123822]I.; Chedotal, A.; Petit, C.: Anosmin-1, defective in the X-
linked

[123823]form of Kallmann syndrome, promotes axonal branch for-
mation from olfactory

[123824]bulb output neurons. Cell 109: 217–228, 2002.

[123825]21997.

[123826]Sparkes, R. S.; Simpson, R. W.; Paulsen, C. A.: Familial hy-
pogonadotropic

[123827]hypogonadism with anosmia. Arch. Intern. Med. 121:
534–538, 1968.

[123828]21998.

[123829]Wegenke, J. D.; Vehling, D. T.; Wear, J. B., Jr.; Gordon, E.
S.;

[123830]Bargman, G. J.; Deacon, J. S. R.; Herrmann, J. P. R.; Opitz, J. M.

[123831]: Familial Kallmann syndrome with unilateral renal aplasia. Clin.

[123832]Genet. 7: 368–381, 1975.

[123833]21999.

[123834]Soussi-Yanicostas, N.; Hardelin, J.-P.; del Mar Arroyo-Jimenez,

[123835]M.; Ardouin, O.; Legouis, R.; Levilliers, J.; Traincard, F.; Betton,

[123836]J.-M.; Cabanie, L.; Petit, C.: Initial characterization of anosmin-1,

[123837]a putative extracellular matrix protein synthesized by definite neuronal

[123838]cell populations in the central nervous system. J. Cell Sci. 109:

[123839]1749–1757, 1996.

[123840]22000.

[123841]Woods-Samuels, P.; Kazazian, H. H., Jr.; Antonarakis, S. E.:

[123842]Nonhomologous recombination in the human genome: deletions in the

[123843]human factor VIII gene. Genomics 10: 94–101, 1991.

[123844]22001.

[123845]Wray, S.; Grant, P.; Gainer, H.: Evidence that cells expressing
ing

[123846]lutinizing hormone–releasing hormone mRNA in the
mouse are derived

[123847]from progenitor cells in the olfactory placode. Proc. Nat.
Acad.

[123848]Sci. 86: 8132–8136, 1989.

[123849]22002.

[123850]Hata, A.; Tsuzuki, T.; Shimada, K.; Takiguchi, M.; Mori, M.;
Matsuda,

[123851]I.: Structure of the human ornithine transcarbamylase
gene. J. Biochem. 103:

[123852]302–308, 1988.

[123853]22003.

[123854]Hata, A.; Tsuzuki, T.; Shimada, K.; Takiguchi, M.; Mori, M.;
Matsuda,

[123855]I.: Isolation and characterization of the human ornithine
transcarbamylase

[123856]gene: structure of the 5–prime–end region. J. Biochem.

100: 717–725,

[123857]1986.

[123858]22004.

[123859]Hauser, E. R.; Finkelstein, J. E.; Valle, D.; Brusilow, S. W.

[123860]; Allopurinol-induced orotidinuria: a test for mutations at
the ornithine

[123861]carbamoyltransferase locus in women. New Eng. J. Med.
322: 1641–1645,

[123862]1990.

[123863]22005.

[123864]Hentzen, D.; Pelet, A.; Feldman, D.; Rabier, D.; Berthelot,
J.;

[123865]Munnich, A.: Fatal hyperammonemia resulting from a C-
to-T mutation

[123866]at a MspI site of the ornithine transcarbamylase gene.
Hum. Genet. 88:

[123867]153–156, 1991.

[123868]22006.

[123869]Herrin, J. T.; McCredie, D. A.: Peritoneal dialysis in the re-
duction

[123870]of blood ammonia levels in a case of hyperammonaemia.

Arch. Dis.

[123871]Child. 44: 149–151, 1969.

[123872]22007.

[123873]Hjelm, M.; de Silva, L. V. K.; Seakins, J. W. T.; Oberholzer,

[123874]V. G.; Rolles, C. J.: Evidence of inherited urea cycle defect
in

[123875]a case of fatal valproate toxicity. Brit. Med. J. 292: 23–24,
1986.

[123876]22008.

[123877]Hokanson, J. T.; O'Brien, W. E.; Idemoto, J.; Schafer, I. A.:

[123878]Carrier detection in ornithine transcarbamylase deficiency.
J. Pediat. 93:

[123879]75–78, 1978.

[123880]22009.

[123881]Holmes, A. K.; Fowler, B.; Sardharwalla, I. B.: Late-onset
ornithine

[123882]carbamyl transferase deficiency in a male patient: detailed
enzyme

[123883]studies. J. Inherit. Metab. Dis. 10 (suppl. 2): 299–301,
1987.

[123884]22010.

- [123885] Holzgreve, W.; Golbus, M. S.: Prenatal diagnosis of ornithine
- [123886] transcarbamylase deficiency utilizing fetal liver biopsy. Am. J.
- [123887] Hum. Genet. 36: 320–328, 1984.
- [123888] 22011.
- [123889] Honeycutt, D.; Callahan, K.; Rutledge, L.; Evans, B.: Heterozygote
- [123890] ornithine transcarbamylase deficiency presenting as symptomatic hyperammonemia
- [123891] during initiation of valproate therapy. Neurology 42: 666–668, 1992.
- [123892] 22012.
- [123893] Hoogenraad, N.; de Martinis, M. L.; Danks, D. M.: Immunological
- [123894] evidence for an ornithine transcarbamylase lesion resulting in the
- [123895] formation of enzyme with smaller protein subunits. J. Inher. Metab.
- [123896] Dis. 6: 149–152, 1983.
- [123897] 22013.

[123898]Hopkins, I. J.; Connelly, J. F.; Dawson, A. G.; Hird, F. J.;
Maddison,

[123899]T. G.: Hyperammonaemia due to ornithine transcarbamylase deficiency. Arch.

[123900]Dis. Child. 44: 143–148, 1969.

[123901]22014.

[123902]Horwich, A. L.; Fenton, W. A.; Williams, K. R.; Kalousek, F.;

[123903]Kraus, J. P.; Doolittle, R. F.; Konigsberg, W.; Rosenberg, L.
E.:

[123904]Structure and expression of a complementary DNA for the
nuclear coded

[123905]precursor of human mitochondrial ornithine transcarbamylase. Science 224:

[123906]1069–1074, 1984.

[123907]22015.

[123908]Horwich, A. L.; Kalousek, F.; Fenton, W. A.; Pollock, R. A.;
Rosenberg,

[123909]L. E.: Targeting of pre-ornithine transcarbamylase to mitochondria:

[123910]definition of critical regions and residues in the leader
peptide. Cell 44:

[123911]451–459, 1986.

[123912]22016.

[123913]Horwich, A. L.; Kalousek, F.; Mellman, I.; Rosenberg, L. E.:

[123914]A leader peptide is sufficient to direct mitochondrial import of a

[123915]chimeric protein. EMBO J. 4: 1129–1135, 1985.

[123916]22017.

[123917]Horwich, A. L.; Kraus, J. P.; Williams, K.; Kalousek, F.;
Konigsberg,

[123918]W.; Rosenberg, L. E.: Molecular cloning of the cDNA coding for rat

[123919]ornithine transcarbamoylase. Proc. Nat. Acad. Sci. 80:
4258–4262,

[123920]1983.

[123921]22018.

[123922]Hurt, E. C.; van Loon, A. P. G. M.: How proteins find mitochondria

[123923]and intramitochondrial compartments. Trends Biochem.
Sci. 11: 204–207,

[123924]1986.

[123925]22019.

[123926]Kang, E. S.; Solomon, S. S.; Gates, R. E.; Schaeffer, S. J.:

[123927]Red blood cell insulin binding studies in Reye's syndrome survivors

[123928]and families. Endocr. Res. Commun. 9: 121–133, 1982.

[123929]22020.

[123930]Kay, J. D. S.; Hilton-Jones, D.; Hyman, N.: Valproate toxicity

[123931]and ornithine carbamoyltransferase deficiency. (Letter)
Lancet II:

[123932]1283–1284, 1986.

[123933]22021.

[123934]Kodama, H.; Ohtake, A.; Mori, M.; Okabe, I.; Tatibana, M.;
Kamoshita,

[123935]S.: Ornithine transcarbamylase deficiency: a case with a
truncated

[123936]enzyme precursor and a case with undetectable mRNA activity. J. Inherit.

[123937]Metab. Dis. 9: 175–185, 1986.

[123938]22022.

[123939]Komaki, S.; Matsuura, T.; Oyanagi, K.; Hoshide, R.; Kiwaki, K.;

[123940]Endo, F.; Shimadzu, M.; Matsuda, I.: Familial lethal inheritance

[123941]of a mutated paternal gene in females causing X-linked ornithine transcarbamylase

[123942](OTC) deficiency. Am. J. Med. Genet. 69: 177–181, 1997.

[123943]2023.

[123944]Kornfeld, M.; Woodfin, B. M.; Papile, L.; Davis, L. E.; Bernard,

[123945]L. R.: Neuropathology of ornithine carbamyl transferase deficiency. Acta

[123946]Neuropath. 65: 261–264, 1985.

[123947]2024.

[123948]Korson, M. S.; Lillehei, C. W.; Vacanti, J. P.; Levy, H. L.:

[123949]Liver transplantation for ornithine transcarbamylase deficiency (OTCD).

[123950](Abstract) Am. J. Hum. Genet. 45 (suppl.): A8, 1989.

[123951]2025.

[123952]Krieger, I.; Snodgrass, P. J.; Roskamp, J.: Atypical clinical

[123953]course of ornithine transcarbamylase deficiency due to a new mutant

[123954](comparison with Reye's disease). J. Clin. Endocr. Metab. 48: 388–392,

[123955]1979.

[123956]22026.

[123957]Lee, J. T.; Nussbaum, R. L.: An arginine to glutamine mutation

[123958]in residue 109 of human ornithine transcarbamylase completely abolishes

[123959]enzymatic activity in COS1 cells. J. Clin. Invest. 84: 1762–1766,

[123960]1989.

[123961]22027.

[123962]Legius, E.; Baten, E.; Stul, M.; Marynen, P.; Cassiman, J.–J.

[123963]: Sporadic late onset ornithine transcarbamylase deficiency in a boy

[123964]with somatic mosaicism for an intragenic deletion. Clin. Genet. 38:

[123965]155–159, 1990.

[123966]22028.

[123967]Levin, B.; Abraham, J. M.; Oberholzer, V. G.; Burgess, E. A.:

[123968]Hyperammonaemia: a deficiency of liver ornithine transcarbamylase.

[123969]Occurrence in mother and child. Arch. Dis. Child. 44: 152-161, 1969.

[123970]22029.

[123971]Lindgren, V.; de Martinville, B.; Horwich, A. L.; Rosenberg, L.

[123972]E.; Francke, U.: Human ornithine transcarbamylase locus mapped to

[123973]band Xp21.1 near Duchenne muscular dystrophy locus. Science 226:

[123974]698-700, 1984.

[123975]22030.

[123976]Maddalena, A.; Sosnoski, D. M.; Berry, G. T.; Nussbaum, R. L.

[123977]: Mosaicism for an intragenic deletion in a boy with mild ornithine

[123978]transcarbamylase deficiency. New Eng. J. Med. 319: 999-1003, 1988.

[123979]22031.

[123980]Maddalena, A.; Spence, J. E.; O'Brien, W. E.; Nussbaum, R. L.

[123981]: Characterization of point mutations in the same arginine codon in

[123982]three unrelated patients with ornithine transcarbamylase deficiency. J.

[123983]Clin. Invest. 82: 1353-1358, 1988.

[123984]22032.

[123985]Maestri, N. E.; Brusilow, S. W.; Clissold, D. B.; Bassett, S.

[123986]S.: Long-term treatment of girls with ornithine transcarbamylase

[123987]deficiency. New Eng. J. Med. 335: 855-859, 1996.

[123988]22033.

[123989]Maestri, N. E.; Hauser, E. R.; Bartholomew, D.; Brusilow, S. W.

[123990]: Prospective treatment of urea cycle disorders. J. Pediat. 119:

[123991]923-928, 1991.

[123992]22034.

[123993]Alitalo, T.; Karna, J.; Forsius, H.; de la Chapelle, A.: X-linked

[123994]retinoschisis is closely linked to DXS41 and DXS16 but not DXS85. Clin.

[123995]Genet. 32: 192–195, 1987.

[123996]22035.

[123997]Alitalo, T.; Kruse, T. A.; de la Chapelle, A.: Refined localization

[123998]of the gene causing X-linked juvenile retinoschisis. Genomics 9:

[123999]505–510, 1991.

[124000]22036.

[124001]Arden, G. B.; Gorin, M. B.; Polkinghorne, P. J.; Jay, M.; Bird,

[124002]A. C.: Detection of the carrier state of X-linked retinoschisis. Am.

[124003]J. Ophthalm. 105: 590–595, 1988.

[124004]22037.

[124005]Byrne, J.; Warburton, D.: Male excess among anatomically normal

[124006]fetuses in spontaneous abortions. Am. J. Med. Genet. 26: 605–611,

[124007]1987.

[124008]22038.

[124009]Dahl, N.; Goonewardena, P.; Chotai, J.; Anvret, M.; Pettersson,

[124010]U.: DNA linkage of X-linked retinoschisis. Hum. Genet. 78: 228-232,

[124011]1988.

[124012]22039.

[124013]Dahl, N.; Goonewardena, P.; Chotai, J.; Wadelius, C.; Lindsten,

[124014]J.; Pettersson, U.: DNA linkage analysis of X-chromosome linked retinoschisis.

[124015](Abstract) Cytogenet. Cell Genet. 46: 602, 1987.

[124016]22040.

[124017]de Jong, P. T. V. M.; Zrenner, E.; van Meel, G. J.; Keunen, J.

[124018]E. E.; van Norren, D.: Mizuo phenomenon in X-linked retinoschisis:

[124019]pathogenesis of the Mizuo phenomenon. Arch. Ophthalmol. 109: 1104-1108,

[124020]1991.

[124021]22041.

[124022]de la Chapelle, A.; Alitalo, T.; Forsius, H.: X-linked juvenile
[124023]retinoschisis. In: Wright, A.; Jay, B. (eds.): Molecular Genet-
ics
[124024]of Inherited Eye Disorders. Chur: Harwood Academic Publ.
(1st
[124025]ed.): 1994.
[124026]22042.
[124027]Eksandh, L. C.; Ponjavic, V.; Ayyagari, R.; Bingham, E. L.;
Hiriyanna,
[124028]K. T.; Andreasson, S.; Ehinger, B.; Sieving, P. A.: Pheno-
typic expression
[124029]of juvenile X-linked retinoschisis in Swedish families with
different
[124030]mutations in the XLR51 gene. Arch. Ophthal. 118:
1098-1104, 2000.
[124031]22043.
[124032]Eriksson, A. W.; Vainio-Mattila, B.; Krause, U.; Fellman, J.;
[124033]Forsius, H.: Secondary sex ratio in families with X-
chromosomal disorders. Hereditas 57:
[124034]373-381, 1967.
[124035]22044.

- [124036] Forsius, H.; Eriksson, A.; Vainio-Mattila, B.: Geschlechts-
gebundene,
[124037] erbliche Retinoschisis in zwei Familien in Finnland. Klin.
Mbl. Augenheilk. 143:
[124038] 806–816, 1963.
[124039] 22045.
- [124040] Forsius, H.; Krause, U.; Helve, J.; Vuopala, V.; Mustonen,
E.;
[124041] Vainio-Mattila, B.; Fellman, J.; Eriksson, A. W.: Visual acu-
ity in
[124042] 183 cases of X-chromosomal retinoschisis. Canad. J.
Ophthal. 8:
[124043] 385–393, 1973.
[124044] 22046.
- [124045] Forsius, H.; Vainio-Mattila, B.; Eriksson, A.: X-linked
hereditary
[124046] retinoschisis. Brit. J. Ophthal. 46: 678–681, 1962.
[124047] 22047.
- [124048] Forsius, H. R.: Personal Communication. Oulu, Finland
1/1977.
[124049] 22048.

- [124050] Forsius, H. R.; Eriksson, A. W.: Retinoschisis X-chromosomalis. In:
- [124051] Eriksson, A. W.; Forsius, H. R.; Nevanlinna, H. R.; Workman, P. L.;
- [124052] Norio, R. K. (eds.): Population Structure and Genetic Disorders.
- [124053] New York: Academic Press (pub.) 1980. Pp. 673–676.
- [124054] 22049.
- [124055] Gehrig, A.; Weber, B. H. F.; Lorenz, B.; Andrassi, M.: First
- [124056] molecular evidence for a de novo mutation in RS1 (XLR1) associated
- [124057] with X-linked juvenile retinoschisis. J. Med. Genet. 36: 932–934,
- [124058] 1999.
- [124059] 22050.
- [124060] Gellert, G.; Petersen, J.; Krawczak, M.; Zoll, B.: Linkage relationship
- [124061] between retinoschisis and four marker loci. Hum. Genet. 79: 382–384,
- [124062] 1988.
- [124063] 22051.

[124064]George, N. D. L.; Yates, J. R. W.; Bradshaw, K.; Moore, A. T.

[124065]: Infantile presentation of X linked retinoschisis. Brit. J.

Ophthal 79:

[124066]653–657, 1995.

[124067]22052.

[124068]George, N. D. L.; Yates, J. R. W.; Moore, A. T.: X linked
retinoschisis. Brit.

[124069]J. Ophthal. 79: 697–702, 1995.

[124070]22053.

[124071]Gieser, E. P.; Falls, H. F.: Hereditary retinoschisis. Am. J.

[124072]Ophthal. 51: 1193–1200, 1961.

[124073]22054.

[124074]Grayson, C.; Reid, S. N. M.; Ellis, J. A.; Rutherford, A.; Sow-
den,

[124075]J. C.; Yates, J. R. W.; Farber, D. B.; Trump, D.:

Retinoschisin,

[124076]the X-linked retinoschisis protein, is a secreted photore-
ceptor protein,

[124077]and is expressed and released by Weri-Rb1 cells. Hum.
Molec. Genet. 9:

[124078]1873–1879, 2000.

[124079]22055.

[124080]Haas, J.: Ueber das Zusammenvorkommen von Veränderungen der Retina

[124081]und Chorioidea. Arch. Augenheilk. 37: 343–348, 1898.

[124082]22056.

[124083]Huopaniemi, L.; Rantala, A.; Forsius, H.; Somer, M.; de la Chapelle,

[124084]A.; Alitalo, T.: Three widespread founder mutations contribute to

[124085]high incidence of X-linked juvenile retinoschisis in Finland. Europ.

[124086]J. Hum. Genet. 7: 368–376, 1999.

[124087]22057.

[124088]Huopaniemi, L.; Rantala, A.; Tahvanainen, E.; de la Chapelle,

[124089]A.; Alitalo, T.: Linkage disequilibrium and physical mapping of X-linked

[124090]juvenile retinoschisis. Am. J. Hum. Genet. 60: 1139–1149, 1997.

[124091]22058.

- [124092]Inoue, Y.; Yamamoto, S.; Okada, M.; Tsujikawa, M.; Inoue, T.;
- [124093]Okada, A. A.; Kusaka, S.; Saito, Y.; Wakabayashi, K.; Miyake, Y.;
- [124094]Fujikado, T.; Tano, Y.: X-linked retinoschisis with point mutations
- [124095]in the XLR51 gene. Arch. Ophthal. 118: 93-96, 2000.
- [124096]22059.
- [124097]Ives, E. J.; Ewing, C. C.; Innes, R.: X-linked juvenile retinoschisis
- [124098]and Xg linkage in five families. (Abstract) Am. J. Hum. Genet. 22:
- [124099]17A-18A, 1970.
- [124100]22060.
- [124101]Kaplan, J.; Pelet, A.; Hentati, H.; Jeanpierre, M.; Briard, M.
- [124102]L.; Journal, H.; Munnich, A.; Dufier, J. L.: Contribution to carrier
- [124103]detection and genetic counselling in X linked retinoschisis. J. Med.
- [124104]Genet. 28: 383-388, 1991.
- [124105]22061.

[124106]Kato, K.; Miyake, Y.; Kachi, S.; Suzuki, T.; Terasaki, H.;
Kawase,

[124107]Y.; Kanda, T.: Axial length and refractive error in X-linked
retinoschisis. Am.

[124108]J. Ophthalm. 131: 812–814, 2001.

[124109]22062.

[124110]Kleinert, H.: Eine recessiv-geschlechtsgebundene Form
der idiopathischen

[124111]Netzhautspaltung bei nichtmyopen Jugendlichen. Albrecht
von Graefes

[124112]Arch. Klin. Exp. Ophthalm. 154: 295–305, 1953.

[124113]22063.

[124114]Mann, I.; MacRae, A.: Congenital vascular veils in the vit-
reous. Brit.

[124115]J. Ophthalm. 22: 1–10, 1938.

[124116]22064.

[124117]Manschot, W. A.: Pathology of hereditary juvenile
retinoschisis. Arch.

[124118]Ophthalm. 88: 131–138, 1972.

[124119]22065.

- [124120]Newton, M. S.; Collyer, S.; Phillips, C. I.: Male proband with
- [124121]X linked retinoschisis apparently inherited from his father's family.
- [124122](Letter) J. Med. Genet. 28: 213–214, 1991.
- [124123]22066.
- [124124]Pagenstecher, H. E.: Ueber eine unter dem Bilde der Netzhautablosung
- [124125]verlaufende, erbliche Erkrankung der Retina. Graefes Arch. Ophthal. 86:
- [124126]457–462, 1913.
- [124127]22067.
- [124128]Pawar, H.; Bingham, E. L.; Lunetta, K. L.; Segal, M.; Richards,
- [124129]J. E.; Boehnke, M.; Sieving, P. A.: Refined genetic mapping of juvenile
- [124130]X-linked retinoschisis. Hum. Hered. 45: 206–210, 1995.
- [124131]22068.
- [124132]Retinoschisis Consortium: Functional implications of the spectrum
- [124133]of mutations found in 234 cases with X-linked juvenile retinoschisis

[124134](XLRs). Hum. Molec. Genet. 7: 1185–1192, 1998.

[124135]22069.

[124136]Sauer, C. G.; Gehrig, A.; Warneke–Wittstock, R.; Marquardt, A.;

[124137]Ewing, C. C.; Gibson, A.; Lorenz, B.; Jurklies, B.; Weber, B. H. F.

[124138] Positional cloning of the gene associated with X-linked juvenile

[124139]retinoschisis. Nature Genet. 17: 164–170, 1997.

[124140]22070.

[124141]Sieving, P. A.; Bingham, E. L.; Roth, M. S.; Young, M. R.; Boehnke,

[124142]M.; Kuo, C.–Y.; Ginsburg, D.: Linkage relationship of X-linked juvenile

[124143]retinoschisis with Xp22.1–p22.3 probes. Am. J. Hum. Genet. 47: 616–621,

[124144]1990.

[124145]22071.

[124146]Sorsby, A.; Klein, M.; Gann, J. H.; Siggins, G.: Unusual retinal

[124147]detachment, possibly sex linked. Brit. J. Ophthal. 35:

1-10, 1951.

[124148]22072.

[124149]Springer, W. R.; Cooper, D. N.; Barondes, S. H.: Discoidin I

[124150]is implicated in cell-substratum attachment and ordered
cell migration

[124151]of Dictyostelium discoideum and resembles fibronectin.

Cell 39:

[124152]557-564, 1984.

[124153]22073.

[124154]Vainio-Mattila, B.; Eriksson, A. W.; Forsius, H.: X-
chromosomal

[124155]recessive retinoschisis in the region of Pori: an oph-
thalgo-genetical

[124156]analysis of 103 cases. Acta Ophthal. 47: 1135-1148,
1969.

[124157]22074.

[124158]Van de Vosse, E.; Bergen, A. A. B.; Meershoek, E. J.; Oost-
erwijk,

[124159]J. C.; Gregory, S.; Bakker, B.; Weissenbach, J.; Coffey, A. J.;
van

[124160]Ommen, G.-J. B.; Den Dunnen, J. T.: An Xp22.1-p22.2

YAC contig encompassing

[124161]the disease loci for RS, KFSD, CLS, HYP and RP15: refined
localization

[124162]of RS. *Europ. J. Hum. Genet.* 4: 101–104, 1996.

[124163]22075.

[124164]Weber, B. H. F.; Schrewe, H.; Molday, L. L.; Gehrig, A.;
White,

[124165]K. L.; Seeliger, M. W.; Jaissle, G. B.; Friedburg, C.; Tamm,
E.; Molday,

[124166]R. S.: Inactivation of the murine X-linked juvenile
retinoschisis

[124167]gene, Rs1h, suggests a role of retinoschisin in retinal cell
layer

[124168]organization and synaptic structure. *Proc. Nat. Acad. Sci.*
99: 6222–6227,

[124169]2002.

[124170]22076.

[124171]Wieacker, P.; Wienker, T. F.; Dallapiccola, B.; Bender, K.;
Davies,

[124172]K. E.; Ropers, H. H.: Linkage relationships between
retinoschisis,

[124173]Xg, and a cloned DNA sequence from the distal short arm

of the X chromosome. Hum.

[124174]Genet. 64: 143–145, 1983.

[124175]22077.

[124176]Yanoff, M.; Kertesz Rahn, E.; Zimmerman, L. E.:

Histopathology

[124177]of juvenile retinoschisis. Arch. Ophthal. 79: 49–53, 1968.

[124178]22078.

[124179]Hiraoka, M.; Trese, M. T.; Shastri, B. S.: X-linked juvenile

[124180]retinoschisis associated with a 4–base pair insertion at
codon 55

[124181]of the XLR51 gene. Biochem. Biophys. Res. Commun. 268:
370–372,

[124182]2000.

[124183]22079.

[124184]Hiriyanna, K. T.; Bingham, E. L.; Yashar, B. M.; Ayyagari,
R.;

[124185]Fishman, G.; Small, K. W.; Weinberg, D. V.; Weleber, R. G.;

Lewis,

[124186]R. A.; Andreasson, S.; Richards, J. E.; Sieving, P. A.: Novel
mutations

[124187]in XLR51 causing retinoschisis, including first evidence of

putative

[124188]leader sequence change. Hum. Mutat. 14: 423–427, 1999.

[124189]22080.

[124190]Newman, J. H.; Wheeler, L.; Lane, K. B.; Loyd, E.; Gaddipati, R.;

[124191]Phillips, J. A., III; Loyd, J. E.: Mutation in the gene for bone

[124192]morphogenetic protein receptor II as a cause of primary pulmonary

[124193]hypertension in a large kindred. New Eng. J. Med. 345: 319–324,

[124194]2001. Note: Erratum. New Eng. J. Med. 345: 1506 only, 2001.

[124195]22081.

[124196]Rosenzweig, B. L.; Imamura, T.; Okadome, T.; Cox, G. N.; Yamashita,

[124197]H.; ten Dijke, P.; Heldin, C.–H.; Miyazono, K.: Cloning and characterization

[124198]of a human type II receptor for bone morphogenetic proteins. Proc.

[124199]Nat. Acad. Sci. 92: 7632–7636, 1995.

[124200]22082.

- [124201]Di Paola, R.; Frittitta, L.; Miscio, G.; Bozzali, M.; Baratta, R.; Centra, M.; Spampinato, D.; Santagati, M. G.; Ercolino, T.; Cisternino, C.; Soccio, T. Mastroianno, S.; Tassi, V.; Almgren, P.; Pizuti, A.; Vigneri, R.; Trischitta, V.: A variation in 3-prime UTR of hPTP1B increases specific gene expression and associates with insulin resistance. Am. J. Hum. Genet. 70: 806–812, 2002.
- [124202]22083.
- [124203]Bradley, D. G.; Farrar, G. J.; Sharp, E. M.; Kenna, P.; Humphries, M. M.; McConnell, D. J.; Daiger, S. P.; McWilliam, P.; Humphries, P.: Autosomal dominant retinitis pigmentosa: exclusion of the gene from the short arm of chromosome 1 including the region surrounding the Rhesus locus. Am. J. Hum. Genet. 44: 570–576, 1989.
- [124204]22084.

- [124214]Mutt, V.; Jorpes, J. E.; Magnusson, S.: Structure of porcine secretin:
- [124215]the amino acid sequence. *Europ. J. Biochem.* 15: 513–519, 1970.
- [124216]22085.
- [124217]Narducci, M. G.; Fiorenza, M. T.; Kang, S.–M.; Bevilacqua, A.;
- [124218]Di Giacomo, M.; Remotti, D.; Picchio, M. C.; Fidanza, V.; Cooper,
- [124219]M. D.; Croce, C. M.; Mangia, F.; Russo, G.: TCL1 participates in
- [124220]early embryonic development and is overexpressed in human seminomas. *Proc.*
- [124221]*Nat. Acad. Sci.* 99: 11712–11717, 2002.
- [124222]22086.
- [124223]Casey, G.; Yamanaka, Y.; Freiss, H.; Kobrin, M. S.; Lopez, M.
- [124224]E.; Buchler, M.; Beger, H. G.; Korc, M.: p53 mutations are common
- [124225]in pancreatic cancer and are absent in chronic pancreatitis. *Cancer*
- [124226]*Lett.* 69: 151–160, 1993.

[124227]22087.

[124228]Castedo, M.; Ferri, K. F.; Blanco, J.; Roumier, T.; Larochette,

[124229]N.; Barretina, J.; Amendola, A.; Nardacci, R.; Metivier, D.; Este,

[124230]J. A.; Piacentini, M.; Kroemer, G.: Human immunodeficiency virus

[124231]1 envelope glycoprotein complex-induced apoptosis involves mammalian

[124232]target of rapamycin/FKBP12-rapamycin-associated protein-mediated p53

[124233]phosphorylation. J. Exp. Med. 194: 1097-1110, 2001.

[124234]22088.

[124235]Chen, L.-C.; Neubauer, A.; Kurisu, W.; Waldman, F. M.; Ljung,

[124236]B.-M.; Goodson, W., III; Goldman, E. S.; Moore, D., II; Balazs, M.;

[124237]Liu, E.; Mayall, B. H.; Smith, H. S.: Loss of heterozygosity on the

[124238]short arm of chromosome 17 is associated with high proliferative capacity

[124239]and DNA aneuploidy in primary human breast cancer.

Proc. Nat. Acad.

[124240]Sci. 88: 3847–3851, 1991.

[124241]22089.

[124242]Chen, P.-L.; Chen, Y.; Bookstein, R.; Lee, W.-H.: Genetic mechanisms

[124243]of tumor suppression by the human p53 gene. Science
250: 1576–1580,

[124244]1990.

[124245]22090.

[124246]Chiba, I.; Takahashi, T.; Nau, M. M.; D'Amico, D.; Curiel, D.

[124247]T.; Mitsudomi, T.; Buchhagen, D. L.; Carbone, D.; Pianta-
dosi, S.;

[124248]Koga, H.; Reissman, P.; Slamon, D. J.; Holmes, E. C.;
Minna, J. D.

[124249]: Mutations in the p53 gene are frequent in primary, re-
sected non-small

[124250]cell lung cancer. Oncogene 5: 1603–1610, 1990.

[124251]22091.

[124252]Cho, Y.; Gorina, S.; Jeffrey, P. D.; Pavletich, N. P.: Crystal

[124253]structure of a p53 tumor suppressor–DNA complex: un-

derstanding tumorigenic

[124254]mutations. Science 265: 346–355, 1994.

[124255]22092.

[124256]Chumakov, A. M.; Miller, C. W.; Chen, D. L.; Koeffler, H. P.:

[124257]Analysis of p53 transactivation through high-affinity
binding sites. Oncogene 8:

[124258]3005–3011, 1993.

[124259]22093.

[124260]Chung, R. Y.; Whaley, J. M.; Anderson, K. M.; Freiman, R.
N.;

[124261]Menon, A. G.; Seizinger, B. R.: p53 gene mutations in hu-
man glioblastomas

[124262]associated with early age onset and better survival.
(Abstract) Am.

[124263]J. Hum. Genet. 47 (suppl.): A4, 1990.

[124264]22094.

[124265]Conseiller, E.; Debussche, L.; Landais, D.; Venot, C.; Mara-
trat,

[124266]M.; Sierra, V.; Tocque, B.; Bracco, L.: CTS1: a p53-derived
chimeric

[124267]tumor suppressor gene with enhanced in vitro apoptotic

properties. J.

[124268]Clin. Invest. 101: 120–127, 1998.

[124269]22095.

[124270]Cooper, D. N.; Youssoufian, H.: The CpG dinucleotide and human

[124271]genetic disease. Hum. Genet. 78: 151–155, 1988.

[124272]22096.

[124273]Crawford, L.: Human p53 and human tumours. BioEssays
3: 117–120,

[124274]1985.

[124275]22097.

[124276]Crook, T.; Wrede, D.; Tidy, J. A.; Mason, W. P.; Evans, D. J.;

[124277]Vousden, K. H.: Clonal p53 mutation in primary cervical
cancer: association

[124278]with human–papillomavirus–negative tumours. Lancet
339: 1070–1073,

[124279]1992.

[124280]22098.

[124281]Culotta, E.; Koshland, D. E., Jr.: p53 sweeps through cancer

[124282]research. Science 262: 1958–1959, 1993.

[124283]22099.

[124284]Denissenko, M. F.; Chen, J. X.; Tang, M.; Pfeifer, G. P.: Cytosine

[124285]methylation determines hot spots of DNA damage in the human P53 gene. Proc.

[124286]Nat. Acad. Sci. 94: 3893–3898, 1997.

[124287]22100.

[124288]Denissenko, M. F.; Pao, A.; Tang, M.; Pfeifer, G. P.: Preferential

[124289]formation of benzo(a)pyrene adducts at lung cancer mutational hot

[124290]spots in P53. Science 274: 430–434, 1996.

[124291]22101.

[124292]Derry, W. B.; Putzke, A. P.; Rothman, J. H.: Caenorhabditis elegans

[124293]p53: role in apoptosis, meiosis, and stress resistance. Science 294:

[124294]591–595, 2001.

[124295]22102.

[124296]DiGiammarino, E. L.; Lee, A. S.; Cadwell, C.; Zhang, W.;
Bothner,
[124297]B.; Ribeiro, R. C.; Zambetti, G.; Kriwacki, R. W.: A novel
mechanism
[124298]of tumorigenesis involving pH-dependent destabilization
of a mutant
[124299]p53 tetramer. Nature Struct. Biol. 9: 12–16, 2002.
[124300]22103.

[124301]Donehower, L. A.; Harvey, M.; Slagle, B. L.; McArthur, M. J.;
[124302]Montgomery, C. A., Jr.; Butel, J. S.; Bradley, A.: Mice defi-
cient
[124303]for p53 are developmentally normal but susceptible to
spontaneous
[124304]tumours. Nature 356: 215–221, 1992.
[124305]22104.

[124306]Dumaz, N.; Drougard, C.; Sarasin, A.; Daya-Grosjean, L.:
Specific
[124307]UV-induced mutation spectrum in the p53 gene of skin
tumors from DNA-repair-deficient
[124308]xeroderma pigmentosum patients. Proc. Nat. Acad. Sci.
90: 10529–10533,

[124309]1993.

[124310]22105.

[124311]El-Deiry, W. S.; Kern, S. E.; Pietenpol, J. A.; Kinzler, K. W.;

[124312]Vogelstein, B.: Definition of a consensus binding site for
p53. Nature

[124313]Genet. 1: 45-49, 1992.

[124314]22106.

[124315]El-Deiry, W. S.; Tokino, T.; Velculescu, V. E.; Levy, D. B.;
Parsons,

[124316]R.; Trent, J. M.; Lin, D.; Mercer, E.; Kinzler, K. W.; Vogel-
stein,

[124317]B.: WAF1, a potential mediator of p53 tumor suppression.
Cell 75:

[124318]817-825, 1993.

[124319]22107.

[124320]Moore, K. J.; D'Amore-Bruno, M. A.; Korfhagen, T. R.;
Glasser,

[124321]S. W.; Whitsett, J. A.; Jenkins, N. A.; Copeland, N. G.: Chro-
mosomal

[124322]localization of three pulmonary surfactant protein genes
in the mouse. Genomics 12:

[124323]388–393, 1992.

[124324]22108.

[124325]Nogee, L. M.; Dunbar, A. E., III; Wert, S. E.; Askin, F.; Ham-
vas,

[124326]A.; Whitsett, J. A.: A mutation in the surfactant protein C
gene

[124327]associated with familial interstitial lung disease. New Eng.
J. Med. 344:

[124328]573–579, 2001.

[124329]22109.

[124330]Warr, R. G.; Hawgood, S.; Buckley, D. I.; Crisp, T. M.;
Schilling,

[124331]J.; Benson, B. J.; Ballard, P. L.; Clements, J. A.; White, R. T.:

[124332]Low molecular weight human pulmonary surfactant pro-
tein (SP5): isolation,

[124333]characterization, and cDNA and amino acid sequences.
Proc. Nat. Acad.

[124334]Sci. 84: 7915–7919, 1987.

[124335]22110.

[124336]Wood, S.; Yaremko, M. L.; Schertzer, M.; Kelemen, P. R.;
Minna,

[124337]J.; Westbrook, C. A.: Mapping of the pulmonary surfactant
SP5 (SFTP2)

[124338]locus to 8p21 and characterization of a microsatellite re-
peat marker

[124339]that shows frequent loss of heterozygosity in human car-
cinomas. Genomics 24:

[124340]597–600, 1994.

[124341]22111.

[124342]Kolble, K.; Lu, J.; Mole, S. E.; Kaluz, S.; Reid, K. B. M.: As-
signment

[124343]of the human pulmonary surfactant protein D gene
(SFTP4) to 10q22–q23

[124344]close to the surfactant protein A gene cluster. Genomics
17: 294–298,

[124345]1993.

[124346]22112.

[124347]Botas, C.; Poulain, F.; Akiyama, J.; Brown, C.; Allen, L.; Go-

erke,

[124348]J.; Clements, J.; Carlson, E.; Gillespie, A. M.; Epstein, C.;
Hawgood,

[124349]S.: Altered surfactant homeostasis and alveolar type II cell
morphology

[124350]in mice lacking surfactant protein D. Proc. Nat. Acad. Sci.
95:

[124351]11869–11874, 1998.

[124352]22113.

[124353]Crouch, E.; Rust, K.; Veile, R.; Donis-Keller, H.; Grosso, L.:

[124354]Genomic organization of human surfactant protein D
(SP-D): SP-D is

[124355]encoded on chromosome 10q22.2–23.1. J. Biol. Chem.
268: 2976–2983,

[124356]1993.

[124357]22114.

[124358]Holmskov, U.; Lawson, P.; Teisner, B.; Tornøe, I.; Willis, A.
C.;

[124359]Morgan, C.; Koch, C.; Reid, K. B. M.: Isolation and charac-
terization

[124360]of a new member of the scavenger receptor superfamily,

glycoprotein-340

[124361](gp-340), as a lung surfactant protein-D binding molecule. J. Biol.

[124362]Chem. 272: 13743-13749, 1997.

[124363]22115.

[124364]Holmskov, U.; Mollenhauer, J.; Madsen, J.; Vitved, L.; Gronlund,

[124365]J.; Tornoe, I.; Kliem, A.; Reid, K. B. M.; Poustka, A.; Skjodt, K.

[124366]: Cloning of gp-340, a putative opsonin receptor for lung surfactant

[124367]protein D. Proc. Nat. Acad. Sci. 96: 10794-10799, 1999.

[124368]22116.

[124369]LeVine, A. M.; Whitsett, J. A.; Hartshorn, K. L.; Crouch, E. C.;

[124370]Korfhagen, T. R.: Surfactant protein D enhances clearance of influenza

[124371]A virus from the lung in vivo. J. Immun. 167: 5868-5873, 2001.

[124372]22117.

[124373]Lu, J.; Willis, A. C.; Reid, K. B. M.: Purification, characteri-

zation

[124374]and cDNA cloning of human lung surfactant protein D.

Biochem. J. 284:

[124375]795–802, 1992.

[124376]22118.

[124377]Rust, K.; Grosso, L.; Zhang, V.; Chang, D.; Persson, A.;

Longmore,

[124378]W.; Cai, G.-Z.; Crouch, E.: Human surfactant protein D:

SP-D contains

[124379]a C-type lectin carbohydrate recognition domain. Arch.

Biochem. Biophys. 290:

[124380]116–126, 1991.

[124381]22119.

[124382]Wert, S. E.; Yoshida, M.; LeVine, A. M.; Ikegami, M.; Jones,

T.;

[124383]Ross, G. F.; Fisher, J. H.; Korfhagen, T. R.; Whitsett, J. A.:

Increased

[124384]metalloproteinase activity, oxidant production, and em-

physema in surfactant

[124385]protein D gene-inactivated mice. Proc. Nat. Acad. Sci. 97:

5972–5977,

[124386]2000.

[124387]22120.

[124388]Bunge, S.; Wedemann, H.; David, D.; Terwilliger, D. J.; van
den

[124389]Born, L. I.; Aulehla-Scholz, C.; Samanns, C.; Horn, M.; Ott,
J.; Schwinger,

[124390]E.; Schinzel, A.; Denton, M. J.; Gal, A.: Molecular analysis
and

[124391]genetic mapping of the rhodopsin gene in families with
autosomal dominant

[124392]retinitis pigmentosa. Genomics 17: 230–233, 1993.

[124393]22121.

[124394]Chang, H.-Y.; Ready, D. F.: Rescue of photoreceptor de-
generation

[124395]in rhodopsin-null *Drosophila* mutants by activated Rac1.
Science 290:

[124396]1978–1980, 2000.

[124397]22122.

[124398]Dikshit, M.; Agarwal, R.: Mutation analysis of codons 345
and

[124399]347 of rhodopsin gene in Indian retinitis pigmentosa pa-
tients. J.

[124400]Genet. 80: 111–116, 2001.

[124401]22123.

[124402]Dryja, T. P.; Hahn, L. B.; Cowley, G. S.; McGee, T. L.;
Berson,

[124403]E. L.: Mutation spectrum of the rhodopsin gene among
patients with

[124404]autosomal dominant retinitis pigmentosa. Proc. Nat. Acad.
Sci. 88:

[124405]9370–9374, 1991.

[124406]22124.

[124407]Dryja, T. P.; McGee, T. L.; Hahn, L. B.; Cowley, G. S.; Ols-
son,

[124408]J. E.; Reichel, E.; Sandberg, M. A.; Berson, E. L.: Mutations
within

[124409]the rhodopsin gene in patients with autosomal dominant
retinitis pigmentosa. New

[124410]Eng. J. Med. 323: 1302–1307, 1990.

[124411]22125.

[124412]Donis-Keller, H.; Green, P.; Helms, C.; Cartinhour, S.;
Weiffenbach,

[124413]B.; Stephens, K.; Keith, T. P.; Bowden, D. W.; Smith, D. R.;

Lander,

[124414]E. S.; Botstein, D.; Akots, G.; Rediker, K. S.; Gravius, T.:

Brown,

[124415]V. A.; Rising, M. B.; Parker, C.; Powers, J. A.; Watt, D. E.;

Kauffman,

[124416]E. R.; Bricker, A.; Phipps, P.; Muller-Kahle, H.; Fulton, T. R.;

Ng,

[124417]S.; Schumm, J. W.; Braman, J. C.; Knowlton, R. G.; Barker,

D. F.;

[124418]Crooks, S. M.; Lincoln, S. E.; Daly, M. J. and Abrahamson,

J.: A genetic

[124419]linkage map of the human genome. Cell 51: 319-337,

1987.

[124420]22126.

[124421]Dryja, T. P.; McGee, T. L.; Hahn, L. B.; Cowley, G. S.; Ols-

son,

[124422]J. E.; Reichel, E.; Sandberg, M. A.; Berson, E. L.: Missense

mutations

[124423]within the rhodopsin gene in patients with autosomal

dominant retinitis

[124424]pigmentosa. (Abstract) Am. J. Hum. Genet. 47 (suppl.):

A54, 1990.

[124425]22127.

[124426]Dryja, T. P.; McGee, T. L.; Reichel, E.; Hahn, L. B.; Cowley,
[124427]G. S.; Yandell, D. W.; Sandberg, M. A.; Berson, E. L.: A
point mutation
[124428]of the rhodopsin gene in one form of retinitis pigmentosa.
Nature 343:
[124429]364–366, 1990.
[124430]22128.
[124431]Elliott, R. W.; Sparkes, R. S.; Mohandas, T.; Grant, S. G.;
McGinnis,
[124432]J. F.: Localization of the rhodopsin gene to the distal half
of mouse
[124433]chromosome 6. Genomics 6: 635–644, 1990.
[124434]22129.
[124435]Farrar, G. J.; Findlay, J. B. C.; Kumar–Singh, R.; Kenna, P.;
[124436]Humphries, M. M.; Sharpe, E.; Humphries, P.: Autosomal
dominant retinitis
[124437]pigmentosa: a novel mutation in the rhodopsin gene in
the original
[124438]3q linked family. Hum. Molec. Genet. 1: 769–771, 1992.
[124439]22130.

- [124440]Farrar, G. J.; Kenna, P.; Redmond, R.; McWilliam, P.;
Bradley,
[124441]D. G.; Humphries, M. M.; Sharp, E. M.; Inglehearn, C. F.;
Bashir,
[124442]R.; Jay, M.; Watty, A.; Ludwig, M.; Schinzel, A.; Samanns,
C.; Gal,
[124443]A.; Bhattacharya, S.; Humphries, P.: Autosomal dominant
retinitis
[124444]pigmentosa: absence of the rhodopsin proline-
to-histidine substitution
[124445](codon 23) in pedigrees from Europe. Am. J. Hum. Genet.
47: 941-945,
[124446]1990.
[124447]22131.
[124448]Farrar, G. J.; Kenna, P.; Redmond, R.; Shiels, D.; McWilliam,
[124449]P.; Humphries, M. M.; Sharp, E. M.; Jordan, S.; Kumar-
Singh, R.; Humphries,
[124450]P.: Autosomal dominant retinitis pigmentosa: a mutation
in codon
[124451]178 of the rhodopsin gene in two families of Celtic origin.
Genomics 11:
[124452]1170-1171, 1991.

[124453]22132.

[124454]Fishman, G. A.; Stone, E. M.; Gilbert, L. D.; Kenna, P.;
Sheffield,

[124455]V. C.: Ocular findings associated with a rhodopsin gene
codon 58

[124456]transversion mutation in autosomal dominant retinitis
pigmentosa. Arch.

[124457]Ophthalmol. 109: 1387–1393, 1991.

[124458]22133.

[124459]Fishman, G. A.; Stone, E. M.; Gilbert, L. D.; Sheffield, V. C.

[124460]: Ocular findings associated with a rhodopsin gene codon
106 mutation:

[124461]glycine-to-arginine change in autosomal dominant retini-
tis pigmentosa. Arch.

[124462]Ophthalmol. 110: 646–653, 1992.

[124463]22134.

[124464]Fishman, G. A.; Vandenburgh, K.; Stone, E. M.; Gilbert, L.
D.;

[124465]Alexander, K. R.; Sheffield, V. C.: Ocular findings associ-
ated with

[124466]rhodopsin gene codon 267 and codon 190 mutations in

dominant retinitis

[124467]pigmentosa. Arch. Ophthalm. 110: 1582–1588, 1992.

[124468]22135.

[124469]Franke, R. R.; Konig, B.; Sakmar, T. P.; Khorana, H. G.;
Hofmann,

[124470]K. P.: Rhodopsin mutants that bind but fail to activate
transducin. Science 250:

[124471]123–125, 1990.

[124472]22136.

[124473]Gal, A.; Artlich, A.; Ludwig, M.; Niemeyer, G.; Olek, K.;
Schwinger,

[124474]E.; Schinzel, A.: Pro347-to-arg mutation of the rhodopsin
gene in

[124475]autosomal dominant retinitis pigmentosa. Genomics 11:
468–470, 1991.

[124476]22137.

[124477]Gal, A.; Samanns, C.; Watty, A.; Jimenez, J.; Ludwig, M.;
Chand,

[124478]A.; Pongratz, J.; Colantuoni, V.; Gussek, H.; Olsson, J.;
Neugebauer,

[124479]M.; Schinzel, A.; Denton, M.: A gene for autosomal domi-

nant retinitis

[124480]pigmentosa is closely linked to D3S20 on 3q. (Abstract)

Retinitis

[124481]Pigmentosa Meeting, Dublin July, 1990.

[124482]22138.

[124483]Hao, W.; Wenzel, A.; Obin, M. S.; Chen, C.-K.; Brill, E.;

Krasnoperova,

[124484]N. V.; Eversole-Cire, P.; Kleyner, Y.; Taylor, A.; Simon, M.

I.; Grimm,

[124485]C.; Reme, C. E.; Lem, J.: Evidence for two apoptotic path-

ways in

[124486]light-induced retinal degeneration. Nature Genet. 32:

254-260, 2002.

[124487]22139.

[124488]Heckenlively, J. R.; Rodriguez, J. A.; Daiger, S. P.: Autosomal

mal

[124489]dominant sectoral retinitis pigmentosa: two families with

transversion

[124490]mutation in codon 23 of rhodopsin. Arch. Ophthalmol. 109:

84-91, 1991.

[124491]22140.

[124492]Humphries, M. M.; Rancourt, D.; Farrar, G. J.; Kenna, P.;

Hazel,

[124493]M.; Bush, R. A.; Sieving, P. A.; Sheils, D. M.; McNally, N.;
Creighton,

[124494]P.; Erven, A.; Boros, A.; Gulya, K.; Capecchi, M. R.;
Humphries, P.

[124495]: Retinopathy induced in mice by targeted disruption of
the rhodopsin

[124496]gene. *Nature Genet.* 15: 216–219, 1997.

[124497]22141.

[124498]Humphries, P.: Personal Communication. Dublin, Ireland
2/1990.

[124499]22142.

[124500]Humphries, P.; Farrar, G. J.; Kenna, P.; McWilliam, P.: Re-
tinitis

[124501]pigmentosa: genetic mapping in X-linked and autosomal
forms of the

[124502]disease. *Clin. Genet.* 38: 1–13, 1990.

[124503]22143.

[124504]Humphries, P.; Kenna, P.; Farrar, G. J.: On the molecular
genetics

[124505]of retinitis pigmentosa. *Science* 256: 804–808, 1992.

[124506]22144.

[124507]Whitmore, T. E.; Holloway, J. L.; Lofton-Day, C. E.; Maurer, M.

[124508]F.; Chen, L.; Quinton, T. J.; Vincent, J. B.; Scherer, S. W.; Lok,

[124509]S.: Human secretin (SCT): gene structure, chromosome location, and

[124510]distribution of mRNA. Cytogenet. Cell Genet. 90: 47-52, 2000.

[124511]22145.

[124512]Domino, S. E.; Zhang, L.; Gillespie, P. J.; Saunders, T. L.; Lowe,

[124513]J. B.: Deficiency of reproductive tract alpha(1,2)fucosylated glycans

[124514]and normal fertility in mice with targeted deletions of the FUT1 or

[124515]FUT2 alpha(1,2)fucosyltransferase locus. Molec. Cell. Biol. 21:

[124516]8336-8345, 2001.

[124517]22146.

[124518]Gedde-Dahl, T., Jr.; Olaisen, B.; Teisberg, P.; Wilhelmy, M. C.;

[124519]Mevag, B.; Helland, R.: The locus for apolipoprotein E (apoE) is

[124520]close to the Lutheran (Lu) blood group locus on chromosome 19. Hum.

[124521]Genet. 67: 178–182, 1984.

[124522]22147.

[124523]Grubb, R.: Zur Genetik des Lewis-Systems. Naturwissenschaften 21:

[124524]560–561, 1953.

[124525]22148.

[124526]Le Pendu, J.; Lemieux, R. U.; Lambert, F.; Dalix, A.-M.; Oriol,

[124527]R.: Distribution of H type 1 and H type 2 antigenic determinants

[124528]in human sera and saliva. Am. J. Hum. Genet. 34: 402–415, 1982.

[124529]22149.

[124530]Oriol, R.; Danilovs, J.; Hawkins, B. R.: A new genetic model

[124531]proposing that the Se gene is a structural gene closely linked to

[124532]the H gene. Am. J. Hum. Genet. 33: 421–431, 1981.

[124533]22150.

[124534]Reguigne-Arnould, I.; Couillin, P.; Mollicone, R.; Faure, S.;

[124535]Fletcher, A.; Kelly, R. J.; Lowe, J. B.; Oriol, R.: Relative positions

[124536]of two clusters of human alpha-L-fucosyltransferases in 19q (FUT1-FUT2)

[124537]and 19p (FUT6-FUT3-FUT5) within the microsatellite genetic map of

[124538]chromosome 19. Cytogenet. Cell Genet. 71: 158-162, 1995.

[124539]22151.

[124540]Kolanus, W.; Nagel, W.; Schiller, B.; Zeitlmann, L.; Godar, S.;

[124541]Stockinger, H.; Seed, B.: Alpha-L-beta-2 integrin/LFA-1 binding to

[124542]ICAM-1 induced by cytohesin-1, a cytoplasmic regulatory molecule. Cell 86:

[124543]233-242, 1996.

[124544]22152.

[124545]Ogasawara, M.; Kim, S.-C.; Adamik, R.; Togawa, A.; Ferrans, V.

[124546]J.; Takeda, K.; Kirby, M.; Moss, J.; Vaughan, M.: Similarities
in
[124547]function and gene structure of cytohesin-4 and cyto-
hesin-1, guanine
[124548]nucleotide-exchange proteins for ADP-ribosylation fac-
tors. J. Biol.
[124549]Chem. 275: 3221-3230, 2000.
[124550]22153.
[124551]Findlay, D. M.; Fisher, L. W.; McQuillan, C. I.; Termine, J.
D.;
[124552]Young, M. F.: Isolation of the osteonectin gene: evidence
that a
[124553]variable region of the osteonectin molecule is encoded
within one
[124554]exon. Biochemistry 27: 1483-1489, 1988.
[124555]22154.
[124556]Gilmour, D. T.; Lyon, G. J.; Carlton, M. B. L.; Sanes, J. R.;
Cunningham,
[124557]J. M.; Anderson, J. R.; Hogan, B. L. M.; Evans, M. J.;
Colledge, W.
[124558]H.: Mice deficient for the secreted glycoprotein SPARC/
osteonectin/BM40

[124559]develop normally but show severe age-onset cataract formation and

[124560]disruption of the lens. EMBO J. 17: 1860–1870, 1998.

[124561]22155.

[124562]Sanyanusin, P.; McNoe, L. A.; Sullivan, M. J.; Weaver, R. G.;

[124563]Eccles, M. R.: Mutation of PAX2 in two siblings with renal-coloboma

[124564]syndrome. Hum. Molec. Genet. 4: 2183–2184, 1995.

[124565]22156.

[124566]Sanyanusin, P.; Norrish, J. H.; Ward, T. A.; Nebel, A.; McNoe,

[124567]L. A.; Eccles, M. R.: Genomic structure of the human PAX2 gene. Genomics 35:

[124568]258–261, 1996.

[124569]22157.

[124570]Sanyanusin, P.; Schimmenti, L. A.; McNoe, L. A.; Ward, T. A.;

[124571]Pierpont, M. E. M.; Sullivan, M. J.; Dobyns, W. B.; Eccles, M. R.

[124572]: Mutation of the PAX2 gene in a family with optic nerve colobomas,

[124573]renal anomalies and vesicoureteral reflux. Nature Genet.
9: 358–364,

[124574]1995.

[124575]22158.

[124576]Schimmenti, L. A.; Cunliffe, H. E.; McNoe, L. A.; Ward, T.
A.;

[124577]French, M. C.; Shim, H. H.; Zhang, Y.–H.; Proesmans, W.;
Leys, A.;

[124578]Byerly, K. A.; Braddock, S. R.; Masuno, M.; Imaizumi, K.;
Devriendt,

[124579]K.; Eccles, M. R.: Further delineation of renal–coloboma
syndrome

[124580]in patients with extreme variability of phenotype and
identical PAX2

[124581]mutations. Am. J. Hum. Genet. 60: 869–878, 1997.

[124582]22159.

[124583]Schimmenti, L. A.; Pierpont, M. E.; Carpenter, B. L. M.;
Kashtan,

[124584]C. E.; Johnson, M. R.; Dobyns, W. B.: Autosomal dominant
optic nerve

[124585]colobomas, vesicoureteral reflux, and renal anomalies.
Am. J. Med.

[124586]Genet. 59: 204–208, 1995.

[124587]22160.

[124588]Schimmenti, L. A.; Shim, H. H.; Wirtschafter, J. D.; Pan-
zarino,

[124589]V. A.; Kashtan, C. E.; Kirkpatrick, S. J.; Wargowski, D. S.;
France,

[124590]T. D.; Michel, E.; Dobyns, W. B.: Homonucleotide expan-
sion and contraction

[124591]mutations of PAX2 and inclusion of Chiari 1 malformation
as part of

[124592]renal–coloboma syndrome. Hum. Mutat. 14: 369–376,
1999.

[124593]22161.

[124594]Stapleton, P.; Weith, A.; Urbanek, P.; Kozmik, Z.; Bus-
slinger,

[124595]M.: Chromosomal localization of seven PAX genes and
cloning of a

[124596]novel family member, PAX–9. Nature Genet. 3: 292–298,
1993.

[124597]22162.

[124598]Tellier, A.–L.; Amiel, J.; Delezoide, A.–L.; Audollent, S.;
Auge,

[124599]J.; Esnault, D.; Encha-Razavi, F.; Munnich, A.; Lyonnet, S.; Vekemans, M.; Attie-Bitach, T.: Expression of the PAX2 gene in human embryos and exclusion in the CHARGE syndrome. Am. J. Med. Genet. 93: 85-88, 2000.

[124602]2000.

[124603]22163.

[124604]Tellier, A.-L.; Amiel, J.; Salomon, R.; Jolly, D.; Delezoide, A.-L.; Auge, J.; Gubler, M.-C.; Munnich, A.; Lyonnet, S.; Antignac, C.; Vekemans, M.; Broyer, M.; Attie-Bitach, T.: PAX2 expression during early human development and its mutations in renal hypoplasia with or without coloboma. (Abstract) Am. J. Hum. Genet. 63 (suppl.): A7 only, 1998.

[124609]A7 only, 1998.

[124610]22164.

[124611]Ward, T. A.; Nebel, A.; Reeve, A. E.; Eccles, M. R.: Alternative

[124612] messenger RNA forms and open reading frames within an additional conserved

[124613] region of the human PAX-2 gene. Cell Growth Differ. 5: 1015-1021,

[124614] 1994.

[124615] 22165.

[124616] Weaver, R. G.; Cashwell, L. F.; Lorentz, W.; Whiteman, D.; Geisinger,

[124617] K. R.; Ball, M.: Optic nerve coloboma associated with renal disease. Am.

[124618] J. Med. Genet. 29: 597-605, 1988.

[124619] 22166.

[124620] Barr, F. G.; Nauta, L. E.; Davis, R. J.; Schafer, B. W.; Nycum,

[124621] L. M.; Biegel, J. A.: In vivo amplification of the PAX3-FKHR and

[124622] PAX7-FKHR fusion genes in alveolar rhabdomyosarcoma. Hum. Molec.

[124623] Genet. 5: 15-21, 1996.

[124624] 22167.

[124625] Burri, M.; Tromvoukis, Y.; Bopp, D.; Frigerio, G.; Noll, M.: Conservation

[124626]of the paired domain in metazoans and its structure in
three isolated

[124627]human genes. EMBO J. 8: 1183–1190, 1989.

[124628]22168.

[124629]Gruss, P.; Walther, C.: Pax in development. Cell 69:
719–722,

[124630]1992.

[124631]22169.

[124632]Schafer, B. W.; Mattei, M. G.: The human paired domain
gene PAX7

[124633](Hup1) maps to chromosome 1p35–1p36.2. Genomics 17:
249–251, 1993.

[124634]22170.

[124635]Seale, P.; Sabourin, L. A.; Girgis–Gabardo, A.; Mansouri,
A.; Gruss,

[124636]P.; Rudnicki, M. A.: Pax7 is required for the specification
of myogenic

[124637]satellite cells. Cell 102: 777–786, 2000.

[124638]22171.

[124639]Shapiro, D. N.; Sublett, J. E.; Li, B.; Valentine, M. B.; Morris,

[124640]S. W.; Noll, M.: The gene for PAX7, a member of the
paired-box-containing
[124641]genes, is localized on human chromosome arm 1p36. Ge-
nomics 17: 767-769,
[124642]1993.
[124643]22172.
[124644]Helwig, U.; Imai, K.; Schmahl, W.; Thomas, B. E.; Varnum,
D. S.;
[124645]Nadeau, J. H.; Balling, R.: Interaction between undulated
and Patch
[124646]leads to an extreme form of spina bifida in double-mutant
mice. Nature
[124647]Genet. 11: 60-63, 1995.
[124648]22173.
[124649]Sun, J.; Rose, J. B.; Bird, P.: Gene structure, chromosomal
localization,
[124650]and expression of the murine homologue of human pro-
teinase inhibitor
[124651]6 (PI-6) suggests divergence of PI-6 from the ovalbumin
serpins. J.
[124652]Biol. Chem. 270: 16089-16096, 1995.

[124653]22174.

[124654]Cowley, C. M. E.; Simrak, D.; Spurr, N. K.; Arnemann, J.;
Buxton,

[124655]R. S.: The plakophilin 1 (PKP1) and plakoglobin (JUP) genes
map to

[124656]human chromosomes 1q and 17, respectively. Hum.
Genet. 100: 486–488,

[124657]1997.

[124658]22175.

[124659]Guenet, J.-L.; Simon-Chazottes, D.; Ringwald, M.; Kemler,
R.:

[124660]The genes coding for alpha and beta catenin (Catn1 and
Catn2) and

[124661]plakoglobin (Jup) map to mouse chromosomes 18, 9, and
11, respectively. Mammalian

[124662]Genome 6: 363–366, 1995.

[124663]22176.

[124664]Ruiz, P.; Brinkmann, V.; Ledermann, B.; Behrend, M.;
Grund, C.;

[124665]Thalhammer, C.; Vogel, F.; Birchmeier, C.; Gunthert, U.;
Franke, W.

[124666]W.; Birchmeier, W.: Targeted mutation of plakoglobin in mice reveals

[124667]essential functions of desmosomes in the embryonic heart. J. Cell

[124668]Biol. 135: 215–225, 1996.

[124669]22177.

[124670]Buckley, M. F.; Goding, J. W.: Plasma cell membrane glycoprotein

[124671]gene Pca-1 (alkaline phosphodiesterase I) is linked to the proto-oncogene

[124672]Myb on mouse chromosome 10. Immunogenetics 36: 199–201, 1992.

[124673]22178.

[124674]Buckley, M. F.; Loveland, K. A.; McKinstry, W. J.; Garson, O. M.;

[124675]Goding, J. W.: Plasma cell membrane glycoprotein PC-1: cDNA cloning

[124676]of the human molecule, amino acid sequence, and chromosomal location. J.

[124677]Biol. Chem. 265: 17506–17511, 1990.

[124678]22179.

[124679]Frittitta, L.; Baratta, R.; Spampinato, D.; Di Paola, R.; Piz-
zuti,

[124680]A.; Vigneri, R.; Trischitta, V.: The Q121 PC-1 variant and
obesity

[124681]have additive and independent effects in causing insulin
resistance. J.

[124682]Clin. Endocr. Metab. 86: 5888-5891, 2001.

[124683]22180.

[124684]Harahap, A. R.; Goding, J. W.: Distribution of the murine
plasma

[124685]cell antigen PC-1 in non-lymphoid tissues. J. Immun. 141:
2317-2320,

[124686]1988.

[124687]22181.

[124688]Huang, R.; Rosenbach, M.; Vaughn, R.; Provvedini, D.;
Rebbe, N.;

[124689]Hickman, S.; Goding, J.; Terkeltaub, R.: Expression of the
murine

[124690]plasma cell nucleotide pyrophosphohydrolase PC-1 is
shared by human

[124691]liver, bone, and cartilage cells: regulation of PC-1 expres-
sion in

[124692]osteosarcoma cells by transforming growth factor-beta. J.

Clin. Invest. 94:

[124693]560-567, 1994.

[124694]22182.

[124695]Kahn, C. R.: Causes of insulin resistance. Nature 373:

384-385,

[124696]1995.

[124697]22183.

[124698]Maddux, B. A.; Goldfine, I. D.: Membrane glycoprotein PC-
1 inhibition

[124699]of insulin receptor function occurs via direct interaction
with the

[124700]receptor alpha-subunit. Diabetes 49: 13-19, 2000.

[124701]22184.

[124702]Maddux, B. A.; Sbraccia, P.; Kumakura, S.; Sasson, S.;
Youngren,

[124703]J.; Fisher, A.; Spencer, S.; Grupe, A.; Henzel, W.; Stewart, T.
A.;

[124704]Reaven, G. M.; Goldfine, I. D.: Membrane glycoprotein PC-
1 and insulin

[124705]resistance in non-insulin-dependent diabetes mellitus.

Nature 373:

[124706]448–451, 1995.

[124707]22185.

[124708]Nakamura, I.; Ikegawa, S.; Okawa, A.; Okuda, S.;
Koshizuka, Y.;

[124709]Kawaguchi, H.; Nakamura, K.; Koyama, T.; Goto, S.;
Toguchida, J.;

[124710]Matsushita, M.; Ochi, T.; Takaoka, K.; Nakamura, Y.: Asso-
ciation

[124711]of the human NPPS gene with ossification of the posterior
longitudinal

[124712]ligament of the spine (OPLL). Hum. Genet. 104: 492–497,
1999.

[124713]22186.

[124714]Okawa, A.; Nakamura, I.; Goto, S.; Moriya, H.; Nakamura,
Y.; Ikegawa,

[124715]S.: Mutation in Npps in a mouse model of ossification of
the posterior

[124716]longitudinal ligament of the spine. Nature Genet. 19:
271–273, 1998.

[124717]22187.

[124718]Pizzuti, A.; Frittitta, L.; Argiolas, A.; Baratta, R.; Goldfine,

[124719]I. D.; Bozzali, M.; Ercolino, T.; Scarlato, G.; Iacoviello, L.;
Vigneri,
[124720]R.; Tassi, V.; Trischitta, V.: A polymorphism (K121Q) of
the human
[124721]glycoprotein PC-1 gene coding region is strongly associ-
ated with insulin
[124722]resistance. Diabetes 48: 1881-1884, 1999.
[124723]22188.
[124724]Rebbe, N. F.; Tong, B. D.; Finley, E. M.; Hickman, S.: Iden-
tification
[124725]of nucleotide pyrophosphatase/alkaline phosphodi-
esterase I activity
[124726]associated with the mouse plasma cell differentiation
antigen PC-1. Proc.
[124727]Nat. Acad. Sci. 88: 5192-5196, 1991.
[124728]22189.
[124729]Sakaguchi, A. Y.; Lalley, P. A.; Zabel, B. U.; Ellis, R. W.;
Scolnick,
[124730]E. M.; Naylor, S. L.: Chromosome assignments of four
mouse cellular
[124731]homologs of sarcoma and leukemia virus oncogenes.
Proc. Nat. Acad.

[124732]Sci. 81: 525–529, 1984.

[124733]22190.

[124734]Takahashi, T.; Old, L. J.; Boyse, E. A.: Surface alloantigens

[124735]of plasma cells. J. Exp. Med. 131: 1325–1341, 1970.

[124736]22191.

[124737]van Driel, I. R.; Goding, J. W.: Plasma cell membrane glycoprotein

[124738]PC-1: primary structure deduced from cDNA clones. J. Biol. Chem. 262:

[124739]4882–4887, 1987.

[124740]22192.

[124741]Frank, S. L.; Klisak, I.; Sparkes, R. S.; Lusk, A. J.: A gene

[124742]homologous to plasminogen located on human chromosome 2q11–p11. Genomics 4:

[124743]449–451, 1989.

[124744]22193.

[124745]Hughes, A. E.; Ralston, S. H.; Marken, J.; Bell, C.; MacPherson,

[124746]H.; Wallace, R. G. H.; van Hul, W.; Whyte, M. P.; Nakatsuka,

K.; Hovy,

[124747]L.; Anderson, D. M.: Mutations in TNFRSF11A, affecting
the signal

[124748]peptide of RANK, cause familial expansile osteolysis. Na-
ture Genet. 24:

[124749]45-48, 2000.

[124750]22194.

[124751]Hughes, A. E.; Shearman, A. M.; Weber, J. L.; Barr, R. J.;
Wallace,

[124752]R. G. H.; Osterberg, P. H.; Nevin, N. C.; Mollan, R. A. B.:
Genetic

[124753]linkage of familial expansile osteolysis to chromosome
18q. Hum.

[124754]Molec. Genet. 3: 359-361, 1994.

[124755]22195.

[124756]Brandriff, B. F.; Gordon, L. A.; Tynan, K. T.; Olsen, A. S.;
Mohrenweiser,

[124757]H. W.; Fertitta, A.; Carrano, A. V.; Trask, B. J.: Order and
genomic

[124758]distances among members of the carcinoembryonic anti-
gen (CEA) gene

[124759]family determined by fluorescence in situ hybridization.

Genomics 12:

[124760]773–779, 1992.

[124761]22196.

[124762]Teglund, S.; Olsen, A.; Khan, W. N.; Frangsmyr, L.; Hammarstrom,

[124763]S.: The pregnancy-specific glycoprotein (PSG) gene cluster on human

[124764]chromosome 19: fine structure of the 11 PSG genes and identification

[124765]of 6 new genes forming a third subgroup within the carcinoembryonic

[124766]antigen (CEA) family. Genomics 23: 669–684, 1994.

[124767]22197.

[124768]Thompson, J.; Koumari, R.; Wagner, K.; Barnert, S.; Schleussner,

[124769]C.; Schrewe, H.; Zimmermann, W.; Muller, G.; Schempp, W.; Zaninetta,

[124770]D.; Ammaturo, D.; Hardman, N.: The human pregnancy-specific glycoprotein

[124771]genes are tightly linked on the long arm of chromosome 19 and are

[124772]coordinately expressed. Biochem. Biophys. Res. Commun.

167: 848–859,
[124773]1990.
[124774]22198.
[124775]Olsen, A.; Teglund, S.; Nelson, D.; Gordon, L.; Copeland,
A.; Georgescu,
[124776]A.; Carrano, A.; Hammarstrom, S.: Gene organization of
the pregnancy-specific
[124777]glycoprotein region on human chromosome 19: assembly
and analysis
[124778]of a 700-kb cosmid contig spanning the region. Genomics
23: 659–668,
[124779]1994.
[124780]22199.
[124781]Thompson, J.; Zimmermann, W.; Osthus–Bugat, P.;
Schleussner, C.;
[124782]Eades–Perner, A.–M.; Barnert, S.; von Kleist, S.; Willcocks,
T.; Craig,
[124783]I.; Tynan, K.; Olsen, A.; Mohrenweiser, H.: Long-range
chromosomal
[124784]mapping of the carcinoembryonic antigen (CEA) gene
family cluster. Genomics 12:
[124785]761–772, 1992.

[124786]22200.

[124787]Tynan, K.; Olsen, A.; Trask, B.; de Jong, P.; Thompson, J.; Zimmermann,

[124788]W.; Carrano, A.; Mohrenweiser, H.: Assembly and analysis of cosmid

[124789]contigs in the CEA-gene family region of human chromosome 19. Nucleic

[124790]Acids Res. 20: 1629–1636, 1992.

[124791]22201.

[124792]Sheer, D.; Sheppard, D. M.; Le Beau, M.; Rowley, J. D.; San Roman,

[124793]C.; Solomon, E.: Localization of the oncogene c-erbA1 immediately

[124794]proximal to the acute promyelocytic leukaemia breakpoint on chromosome

[124795]17. Ann. Hum. Genet. 49: 167–171, 1985.

[124796]22202.

[124797]Yoshitake, H.; Rittling, S. R.; Denhardt, D. T.; Noda, M.: Osteopontin-deficient

[124798]mice are resistant to ovariectomy-induced bone resorption. Proc.

[124799]Nat. Acad. Sci. 96: 8156–8160, 1999.

[124800]22203.

[124801]Edelmann, W.; Zervas, M.; Costello, P.; Roback. L.; Fischer, I.;

[124802]Hammarback, J. A.; Cowan, N.; Davies, P.; Wainer, B.; Kucherlapati,

[124803]R.: Neuronal abnormalities in microtubule-associated protein 1B mutant

[124804]mice. Proc. Nat. Acad. Sci. 93: 1270–1275, 1996.

[124805]22204.

[124806]Hammarback, J. A.; Obar, R. A.; Hughes, S. M.; Vallee, R. B.:

[124807]MAP1B is encoded as a polyprotein that is processed to form a complex

[124808]N-terminal microtubule-binding domain. Neuron 7: 129–139, 1991.

[124809]22205.

[124810]Lien, L. L.; Boyce, F. M.; Kleyn, P.; Brzustowicz, L. M.; Menninger,

[124811]J.; Ward, D. C.; Gilliam, T. C.; Kunkel, L. M.: Mapping of human

[124812]microtubule-associated protein 1B in proximity to the
spinal muscular

[124813]atrophy locus at 5q13. Proc. Nat. Acad. Sci. 88:
7873–7876, 1991.

[124814]22206.

[124815]Lien, L. L.; Feener, C. A.; Fischbach, N.; Kunkel, L. M.:
Cloning

[124816]of human microtubule-associated protein 1B and the
identification

[124817]of a related gene on chromosome 15. Genomics 22:
273–280, 1994.

[124818]22207.

[124819]Wirth, B.; Voosen, B.; Rohrig, D.; Knapp, M.; Piechaczek,
B.; Rudnik-Schoneborn,

[124820]S.; Zerres, K.: Fine mapping and narrowing of the genetic
interval

[124821]of the spinal muscular atrophy region by linkage studies.
Genomics 15:

[124822]113–118, 1993.

[124823]22208.

[124824]Zhang, Y. Q.; Bailey, A. M.; Matthies, H. J. G.; Renden, R.
B.;

[124825]Smith, M. A.; Speese, S. D.; Rubin, G. M.; Broadie, K.:

Drosophila

[124826]fragile X-related gene regulates the MAP1B homolog

Futsch to control

[124827]synaptic structure and function. *Cell* 107: 591–603, 2001.

[124828]22209.

[124829]Garner, C. C.; Tucker, R. P.; Matus, A.: Selective localization

[124830]of messenger RNA for cytoskeletal protein MAP2 in dendrites. *Nature* 336:

[124831]674–677, 1988.

[124832]22210.

[124833]Kalcheva, N.; Albala, J.; O'Guin, K.; Rubino, H.; Garner, C.; Shafit-Zagardo,

[124834]B.: Genomic structure of human microtubule-associated protein 2 (MAP-2)

[124835]and characterization of additional MAP-2 isoforms. *Proc. Nat. Acad.*

[124836]*Sci.* 92: 10894–10898, 1995.

[124837]22211.

[124838]Kindler, S.; Garner, C. C.: Four repeat MAP2 isoforms in

human

[124839]and rat brain. Molec. Brain Res. 26: 218–224, 1994.

[124840]22212.

[124841]Marsden, K. M.; Doll, T.; Ferralli, J.; Botteri, F.; Matus, A.

[124842]: Transgenic expression of embryonic MAP2 in adult
mouse brain: implications

[124843]for neuronal polarization. J. Neurosci. 16: 3265–3273,
1996.

[124844]22213.

[124845]Neve, R. L.; Harris, P.; Kosik, K. S.; Kurnit, D. M.; Donlon,
T.

[124846]A.: Identification of cDNA clones for the human micro-
tubule-associated

[124847]protein tau and chromosomal localization of the genes for
tau and

[124848]microtubule-associated protein 2. Molec. Brain Res. 1:
271–280,

[124849]1986.

[124850]22214.

[124851]Chapin, S. J.; Bulinski, J. C.: Non-neuronal 210 x 10(3) M(r)

[124852]microtubule-associated protein (MAP4) contains a domain

homologous

[124853]to the microtubule-binding domains of neuronal MAP2
and tau. J. Cell

[124854]Sci. 98: 27-36, 1991.

[124855]22215.

[124856]Adriaansen, H. J.; Geurts Van Kessel, A. H. M.; Wijdenes-
De Bresser,

[124857]J. H. F. M.; Van Drunen-Schoenmaker, E.; Van Dongen, J. J.
M.: Expression

[124858]of the myeloid differentiation antigen CD33 depends on
the presence

[124859]of human chromosome 19 in human-mouse hybrids. Ann.
Hum. Genet. 54:

[124860]115-119, 1990.

[124861]22216.

[124862]Bonthron, D. T.; Dunlop, N.; Barr, D. G. D.; El Sanousi, A.
A.;

[124863]Al-Gazali, L. I.: Organisation of the human PAX4 gene and
its exclusion

[124864]as a candidate for the Wolcott-Rallison syndrome. J. Med.
Genet. 35:

[124865]288-292, 1998.

[124866]22217.

[124867]Inoue, H.; Nomiyama, J.; Nakai, K.; Matsutani, A.;
Tanizawa, Y.;

[124868]Oka, Y.: Isolation of full-length cDNA of mouse PAX4 gene
and identification

[124869]of its human homologue. Biochem. Biophys. Res. Com-
mun. 243: 628-633,

[124870]1998.

[124871]22218.

[124872]Mansouri, A.; St-Onge, L.; Gruss, P.: Role of Pax genes in
endoderm-derived

[124873]organs. Trends Endocr. Metab. 10: 164-167, 1999.

[124874]22219.

[124875]Matsushita, T.; Yamaoka, T.; Otsuka, S.; Moritani, M.; Mat-
sumoto,

[124876]T.; Itakura, M.: Molecular cloning of mouse paired-
box-containing

[124877]gene (Pax)-4 from an islet beta cell line and deduced se-
quence of

[124878]human Pax-4. Biochem. Biophys. Res. Commun. 242:
176-180, 1998.

[124879]22220.

[124880]Sosa-Pineda, B.; Chowdhury, K.; Torres, M.; Oliver, G.;
Gruss,

[124881]P.: The Pax4 gene is essential for differentiation of in-
sulin-producing

[124882]beta cells in the mammalian pancreas. Nature 386:
399-402, 1997.

[124883]22221.

[124884]St-Onge, L.; Sosa-Pineda, B.; Chowdhury, K.; Mansouri, A.;
Gruss,

[124885]P.: Pax6 is required for differentiation of glucagon-pro-
ducing alpha-cells

[124886]in mouse pancreas. Nature 387: 406-409, 1997.

[124887]22222.

[124888]Tamura, T.; Izumikawa, Y.; Kishino, T.; Soejima, H.; Jinno,
Y.;

[124889]Niikawa, N.: Assignment of the human PAX4 gene to chro-
mosome band

[124890]7q32 by fluorescence in situ hybridization. Cytogenet. Cell
Genet. 66:

[124891]132-134, 1994.

[124892]22223.

[124893]Adams, B.; Dorfler, P.; Aguzzi, A.; Kozmik, Z.; Urbanek, P.;
Maurer-Fogy,

[124894]I.; Busslinger, M.: Pax-5 encodes the transcription factor
BSAP and

[124895]is expressed in B lymphocytes, the developing CNS, and
adult testis. Genes

[124896]Dev. 6: 1589-1607, 1992.

[124897]22224.

[124898]Busslinger, M.; Klix, N.; Pfeffer, P.; Graninger, P. G.;
Kozmik,

[124899]Z.: Deregulation of PAX-5 by translocation of the E- μ
enhancer of

[124900]the IgH locus adjacent to two alternative PAX-5 promoters
in a diffuse

[124901]large-cell lymphoma. Proc. Nat. Acad. Sci. 93:
6129-6134, 1996.

[124902]22225.

[124903]Mikkola, I.; Heavey, B.; Horcher, M.; Busslinger, M.: Rever-
sion

[124904]of B cell commitment upon loss of Pax5 expression. Sci-
ence 297:

[124905]110–113, 2002.

[124906]22226.

[124907]Nutt, S. L.; Heavey, B.; Rolink, A. G.; Busslinger, M.: Com-
mitment

[124908]to the B-lymphoid lineage depends on the transcription
factor Pax5. Nature 401:

[124909]556–562, 1999.

[124910]22227.

[124911]Nutt, S. L.; Vambrie, S.; Steinlein, P.; Kozmik, Z.; Rolink,
A.;

[124912]Weith, A.; Busslinger, M.: Independent regulation of the
two Pax5

[124913]alleles during B-cell development. Nature Genet. 21:
390–395, 1999.

[124914]22228.

[124915]Ohno, H.; Furukawa, T.; Fukuhara, S.; Zong, S. Q.; Kame-
saki, H.;

[124916]Shows, T. B.; Le Beau, M. M.; McKeithan, T. W.; Kawakami,
T.; Honjo,

[124917]T.: Molecular analysis of a chromosomal translocation,
t(9;14)(p13;q32),

[124918]in a diffuse large-cell lymphoma cell line expressing the
Ki-1 antigen. Proc.

[124919]Nat. Acad. Sci. 87: 628-632, 1990.

[124920]22229.

[124921]Rolink, A. G.; Nutt, S. L.; Melchers, F.; Busslinger, M.:
Long-term

[124922]in vivo reconstitution of T-cell development by
Pax5-deficient B-cell

[124923]progenitors. Nature 401: 603-606, 1999.

[124924]22230.

[124925]Vorechovsky, I.; Koskinen, S.; Paganelli, R.; Smith, C. I. E.;

[124926]Busslinger, M.; Hammarstrom, L.: The PAX5 gene: a link-
age and mutation

[124927]analysis in candidate human primary immunodeficiencies.
Immunogenetics 42:

[124928]149-152, 1995.

[124929]22231.

[124930]Walther, C.; Guenet, J.-L.; Simon, D.; Deutsch, U.; Jostes,
B.;

[124931]Goulding, M. D.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a
murine

[124932]multigene family of paired box containing genes. Genomics 11: 424-434,

[124933]1991.

[124934]22232.

[124935]Koseki, H.; Zachgo, J.; Mizutani, Y.; Simon-Chazottes, D.; Guenet,

[124936]J.-L.; Balling, R.; Gossler, A.: Fine genetic mapping of the proximal

[124937]part of mouse chromosome 2 excludes Pax-8 as a candidate gene for

[124938]Danforth's short tail (Sd). Mammalian Genome 4: 324-327, 1993.

[124939]22233.

[124940]Kroll, T. G.; Sarraf, P.; Pecciarini, L.; Chen, C.-J.; Mueller,

[124941]E.; Splegeman, B. M.; Fletcher, J. A.:

PAX8-PPAR-gamma-1 fusion

[124942]in oncogene human thyroid carcinoma. Science 289: 1357-1360, 2000.

[124943]22234.

[124944]Macchia, P. E.; Lapi, P.; Krude, H.; Pirro, M. T.; Missero, C.;

[124945]Chiovato, L.; Souabni, A.; Baserga, M.; Tassi, V.; Pinchera,

A.; Fenzi,

[124946]G.; Gruters, A.; Busslinger, M.; Di Lauro, R.: PAX8 mutations associated

[124947]with congenital hypothyroidism caused by thyroid dysgenesis. Nature

[124948]Genet. 19: 83–86, 1998.

[124949]22235.

[124950]Mansouri, A.; Chowdhury, K.; Gruss, P.: Follicular cells of the

[124951]thyroid gland require Pax8 gene function. Nature Genet. 19: 87–90,

[124952]1998.

[124953]22236.

[124954]Pasca di Magliano, M.; Di Lauro, R.; Zannini, M.: Pax8 has a key

[124955]role in thyroid cell differentiation. Proc. Nat. Acad. Sci. 97:

[124956]13144–13149, 2000.

[124957]22237.

[124958]Plachov, D.; Chowdhury, K.; Walther, C.; Simon, D.; Guenet, J.–L.;

[124959]Gruss, P.: Pax–8, a murine paired box gene expressed in

the developing

[124960]excretory system and thyroid gland. Development 110:
643–651, 1990.

[124961]22238.

[124962]Tell, G.; Pellizzari, L.; Esposito, G.; Pucillo, C.; Macchia,

[124963]P. E.; Di Lauro, R.; Damante, G.: Structural defects of a
Pax8 mutant

[124964]that give rise to congenital hypothyroidism. Biochem. J.
341: 89–93,

[124965]1999.

[124966]22239.

[124967]Bongarzone, I.; Vigano, E.; Alberti, L.; Borrello, M. G.;

Pasini,

[124968]B.; Greco, A.; Mondellini, P.; Smith, D. P.; Ponder, B. A. J.;

Romeo,

[124969]G.; Pierotti, M. A.: Full activation of MEN2B mutant RET by
an additional

[124970]MEN2A mutation or by ligand GDNF stimulation. Onco-
gene 16: 2295–2301,

[124971]1998.

[124972]22240.

[124973]Carlson, K. M.; Bracamontes, J.; Jackson, C. E.; Clark, R.;
Lacroix,
[124974]A.; Wells, S. A., Jr.; Goodfellow, P. J.: Parent-of-origin ef-
fects
[124975]in multiple endocrine neoplasia type 2B. Am. J. Hum.
Genet. 55:
[124976]1076–1082, 1994.
[124977]22241.
[124978]Carlson, K. M.; Dou, S.; Chi, D.; Scavarda, N.; Toshima, K.;
Jackson,
[124979]C. E.; Wells, S. A., Jr.; Goodfellow, P. J.; Donis-Keller, H.:
Single
[124980]missense mutation in the tyrosine kinase catalytic domain
of the RET
[124981]protooncogene is associated with multiple endocrine neo-
plasia type
[124982]2B. Proc. Nat. Acad. Sci. 91: 1579–1583, 1994.
[124983]22242.
[124984]Mulligan, L. M.; Eng, C.; Healey, C. S.; Clayton, D.; Kwok, J.
[124985]B. J.; Gardner, E.; Ponder, M. A.; Frilling, A.; Jackson, C. E.;
Lehnert,

[124986]H.; Neumann, H. P. H.; Thibodeau, S. N.; Ponder, B. A. J.:

Specific

[124987]mutations of the RET proto-oncogene are related to disease phenotype

[124988]in MEN 2A and FMTC. Nature Genet. 6: 70–74, 1994.

[124989]22243.

[124990]Tartaglia, M.; Mehler, E. L.; Goldberg, R.; Zampino, G.; Brunner,

[124991]H. G.; Kremer, H.; van der Burgt, I.; Crosby, A. H.; Ion, A.; Jeffery,

[124992]S.; Kalidas, K.; Patton, M. A.; Kucherlapati, R. S.; Gelb, B. D.:

[124993]Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2,

[124994]cause Noonan syndrome. Nature Genet. 29: 465–468, 2001. Note: Errata:

[124995]Nature Genet. 29: 491 only, 2001; Nature Genet. 30: 123 only, 2002.

[124996]22244.

[124997]Bruss, M.; Kunz, J.; Lingen, B.; Bonisch, H.: Chromosomal mapping

[124998]of the human gene for the tricyclic antidepressant-sensi-

tive noradrenaline

[124999] transporter. Hum. Genet. 91: 278–280, 1993.

[125000] 22245.

[125001] Esler, M.; Jennings, G.; Lambert, G.; Meredith, I.; Horne, M.;

[125002] Eisenhofer, G.: Overflow of catecholamine neurotransmitters to the

[125003] circulation: source, fate, and functions. Physiol. Rev. 70: 963–985,

[125004] 1990.

[125005] 22246.

[125006] Fritz, J. D.; Jayanthi, L. D.; Thoreson, M. A.; Blakely, R. D.

[125007]: Cloning and chromosomal mapping of the murine norepinephrine transporter. J.

[125008] Neurochem. 70: 2241–2251, 1998.

[125009] 22247.

[125010] Gelernter, J.; Kruger, S.; Kidd, K. K.; Amara, S.: TaqI RFLP at

[125011] norepinephrine transporter protein (NET) locus. Hum. Molec. Genet. 2:

[125012] 820 only, 1993.

[125013]22248.

[125014]Gelernter, J.; Kruger, S.; Pakstis, A. J.; Pacholczyk, T.;
Sparkes,

[125015]R. S.; Kidd, K. K.; Amara, S.: Assignment of the nore-
pinephrine transporter

[125016]protein (NET1) locus to chromosome 16. Genomics 18:
690-692, 1993.

[125017]22249.

[125018]Kaye, W. H.; Jimerson, D. C.; Lake, C. R.; Ebert, M. H.: Al-
tered

[125019]norepinephrine metabolism following long-term weight
recovery in patients

[125020]with anorexia nervosa. Psychiat. Res. 14: 333-342, 1985.

[125021]22250.

[125022]Pacholczyk, T.; Blakely, R. D.; Amara, S. G.: Expression
cloning

[125023]of a cocaine- and antidepressant-sensitive human nora-
drenaline transporter. Nature 350:

[125024]350-354, 1991.

[125025]22251.

[125026]Paczkowski, F. A.; Bonisch, H.; Bryan-Lluka, L. J.: Pharma-

cological

[125027]properties of the naturally occurring ala457pro variant of the human

[125028]norepinephrine transporter. Pharmacogenetics 12: 165–173, 2002.

[125029]22252.

[125030]Pirke, K. M.; Kellner, M.; Philipp, E.; Laessle, R.; Krieg, J.

[125031]C.; Fichter, M. M.: Plasma norepinephrine after a standardized test

[125032]meal in acute and remitted patients with anorexia nervosa and in healthy

[125033]controls. Biol. Psychiat. 31: 1074–1077, 1992.

[125034]22253.

[125035]Porzgen, P.; Bonisch, H.; Bruss, M.: Molecular cloning and organization

[125036]of the coding region of the human norepinephrine transporter gene. Biochem.

[125037]Biophys. Res. Commun. 215: 1145–1150, 1995. Notes: Erratum: Biochem.

[125038]Biophys. Res. Commun. 227: 642–643, 1996.

[125039]22254.

[125040]Shannon, J. R.; Flattem, N. L.; Jordan, J.; Jacob, G.; Black,

[125041]B. K.; Biaggioni, I.; Blakely, R. D.; Robertson, D.: Ortho-
static

[125042]intolerance and tachycardia associated with nore-
pinephrine-transporter

[125043]deficiency. New Eng. J. Med. 342: 541-549, 2000.

[125044]22255.

[125045]Urwin, R. E.; Bennetts, B.; Wilcken, B.; Lampropoulos, B.;
Beumont,

[125046]P.; Clarke, S.; Russell, J.; Tanner, S.; Nunn, K. P.: Anorexia
nervosa

[125047](restrictive subtype) is associated with a polymorphism in
the novel

[125048]norepinephrine transporter gene promoter polymorphic
region. Molec.

[125049]Psychiat. 7: 652-657, 2002.

[125050]22256.

[125051]Tomilin, A.; Remenyi, A.; Lins, K.; Bak, H.; Leidel, S.;
Vriend,

[125052]G.; Wilmanns, M.; Scholer, H. R.: Synergism with the coac-
tivator

[125053]OBF-1 (OCA-B, BOB-1) is mediated by a specific POU dimer configuration. Cell 103:

[125054]853-864, 2000.

[125055]22257.

[125056]Ko, H.-S.; Fast, P.; McBride, W.; Staudt, L. M.: A human protein

[125057]specific for the immunoglobulin octamer DNA motif contains a functional

[125058]homeobox domain. Cell 55: 135-144, 1988.

[125059]22258.

[125060]Schubart, K.; Massa, S.; Schubart, D.; Corcoran, L. M.; Rolink,

[125061]A. G.; Matthias, P.: B cell development and immunoglobulin gene transcription

[125062]in the absence of Oct-2 and OBF-1. Nature Immun. 2: 69-74, 2001.

[125063]22259.

[125064]Staudt, L. M.; Clerc, R. G.; Singh, H.; LeBowitz, J. H.; Sharp,

[125065]P. A.; Baltimore, D.: Cloning of a lymphoid-specific cDNA encoding

[125066]a protein binding the regulatory octamer DNA motif. Sci-

ence 241:

[125067]577–580, 1988.

[125068]22260.

[125069]Parker, R. C.; Mardon, G.; Lebo, R. V.; Varmus, H. E.;
Bishop,

[125070]J. M.: Isolation of duplicated human c-src genes located
on chromosomes

[125071]1 and 20. Molec. Cell. Biol. 5: 831–838, 1985.

[125072]22261.

[125073]Tronick, S. R.; Popescu, N. C.; Cheah, M. S. C.; Swan, D. C.;
Amsbaugh,

[125074]S. C.; Lengel, C. R.; DiPaolo, J. A.; Robbins, K. C.: Isolation
and

[125075]chromosomal localization of the human fgr protoonco-
gene, a distinct

[125076]member of the tyrosine kinase gene family. Proc. Nat.
Acad. Sci. 82:

[125077]6595–6599, 1985.

[125078]22262.

[125079]Bork, P.: The modular architecture of a new family of
growth regulators

[125080]related to connective tissue growth factor. FEBS Lett. 327:
125-130,

[125081]1993.

[125082]22263.

[125083]Sugaya, K.; Fukagawa, T.; Matsumoto, K.; Mita, K.; Takahashi, E.;

[125084]Ando, A.; Inoko, H.; Ikemura, T.: Three genes in the human MHC class

[125085]III region near the junction with the class II: gene for receptor

[125086]of advanced glycosylation end products, PBX2 homeobox gene and a Notch

[125087]homolog, human counterpart of mouse mammary tumor gene int-3. Genomics 23:

[125088]408-419, 1994.

[125089]22264.

[125090]Joliot, V.; Martinerie, C.; Dambrine, G.; Plassiart, G.; Brisac,

[125091]M.; Crochet, J.; Perbal, B.: Proviral rearrangements and overexpression

[125092]of a new cellular gene (nov) in myeloblastosis-associated virus type

[125093]1-induced nephroblastomas. *Molec. Cell. Biol.* 12: 10-21, 1992.

[125094]22265.

[125095]Kim, H.-S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T., Jr.; Rosenfeld, R. G.: Identification of a family of low-affinity

[125097]insulin-like growth factor binding proteins (IFGBPs): characterization

[125098]of connective tissue growth factor as a member of the IGFBP superfamily. *Proc.*

[125099]*Nat. Acad. Sci.* 94: 12981-12986, 1997.

[125100]22266.

[125101]Martinerie, C.; Perbal, B.: Expression of a gene encoding a novel

[125102]potential IGF binding protein in human tissues. *C. R. Acad. Sci.*

[125103](Paris) 313 (ser. 3): 345-351, 1991.

[125104]22267.

[125105]Snaith, M. R.; Natarajan, D.; Taylor, L. B.; Choi, C.-P.; Martinerie,

[125106]C.; Perbal, B.; Schofield, P. N.; Boulter, C. A.: Genomic

structure

[125107]and chromosomal mapping of the mouse nov gene. Genomics 38: 425–428,

[125108]1996.

[125109]22268.

[125110]Soret, J.; Dambrine, G.; Perbal, B.: Induction of nephroblastoma

[125111]by myeloblastosis-associated virus type 1: state of proviral DNAs

[125112]in tumor cells. J. Virol. 63: 1803–1807, 1989.

[125113]22269.

[125114]Amson, R.; Sigaux, F.; Przedborski, S.; Flandrin, G.; Givol, D.;

[125115]Telerman, A.: The human protooncogene product p33pim is expressed

[125116]during fetal hematopoiesis and in diverse leukemias. Proc. Nat. Acad.

[125117]Sci. 86: 8857–8861, 1989.

[125118]22270.

[125119]Ark, B.; Gummere, G.; Bennett, D.; Artzt, K.: Mapping of the Pim-1

[125120] oncogene in mouse t-haplotypes and its use to define the relative

[125121] map positions of the tcl loci t-0(t-6) and t-w12 and the marker tf

[125122] (tufted). Genomics 10: 385-389, 1991.

[125123] 22271.

[125124] Cuypers, H. T.; Selten, G.; Berns, A.; Geurts van Kessel, A. H.

[125125] M.: Assignment of the human homologue of Pim-1, a mouse gene implicated

[125126] in leukemogenesis, to the pter-q12 region of chromosome 6. Hum. Genet. 72:

[125127] 262-265, 1986.

[125128] 22272.

[125129] Domen, J.; von Lindern, M.; Hermans, A.; Breuer, M.; Grosveld,

[125130] G.; Berns, A.: Comparison of the human and mouse PIM-1 cDNAs: nucleotide

[125131] sequence and immunological identification of the in vitro synthesized

[125132] PIM-1 protein. Oncogene Res. 1: 103-112, 1987.

[125133]22273.

[125134]Dunger, D. B.; Davies, K. E.; Pembrey, M.; Lake, B.; Pearson,
son,

[125135]P.; Williams, D.; Whitfield, A.; Dillon, M. J. D.: Deletion on
the

[125136]X chromosome detected by direct DNA analysis in one of
two unrelated

[125137]boys with glycerol kinase deficiency, adrenal hypoplasia,
and Duchenne

[125138]muscular dystrophy. Lancet I: 585-587, 1986.

[125139]22274.

[125140]Eriksson, A.; Lindstedt, S.; Ransnas, L.; von Wendt, L.: De-
ficiency

[125141]of glycerol kinase (EC 2.7.1.30). Clin. Chem. 29: 718-722,
1983.

[125142]22275.

[125143]Francke, U.; Harper, J. F.; Darras, B. T.; Cowan, J. M.; Mc-
Cabe,

[125144]E. R. B.; Kohlschutter, A.; Seltzer, W. K.; Saito, F.; Goto, J.;
Harpey,

[125145]J.-P.; Wise, J. E.: Congenital adrenal hypoplasia, myopathy,
and

[125146]glycerol kinase deficiency: molecular genetic evidence for deletions. Am.

[125147]J. Hum. Genet. 40: 212–227, 1987.

[125148]22276.

[125149]Gaudet, D.; Arsenault, S.; Perusse, L.; Vohl, M.–C.;
St.–Pierre,

[125150]J.; Bergeron, J.; Despres, J.–P.; Dewar, K.; Daly, M. J.; Hud-
son,

[125151]T.; Rioux, J. D.: Glycerol as a correlate of impaired glucose
tolerance:

[125152]dissection of a complex system by use of a simple genetic
trait. Am.

[125153]J. Hum. Genet. 66: 1558–1568, 2000.

[125154]22277.

[125155]Ginns, E. I.; Barranger, J. A.; McClean, S. W.; Sliva, C.;
Young,

[125156]R.; Schaefer, I.; Goodman, S. I.; McCabe, E. R. B.: A juvenile
form

[125157]of glycerol kinase deficiency with episodic vomiting,
acidemia, and

[125158]stupor. J. Pediat. 104: 736–739, 1984.

[125159]22278.

[125160]Goonewardena, P.; Dahl, N.; Ritzen, M.; Johnsen, E.; Petersson,

[125161]U.: Deletion in Xp associated with glycerol kinase deficiency, adrenal

[125162]aplasia and hypogonadotropic hypogonadism. (Abstract) Cytogenet.

[125163]Cell Genet. 46: 621 only, 1987.

[125164]22279.

[125165]Goussault, Y.; Turpin, E.; Neel, D.; Dreux, C.; Chanu, B.; Bakir,

[125166]R.; Rouffy, J.: 'Pseudohypertriglyceridemia' caused by hyperglycerolemia

[125167]due to congenital enzyme deficiency. Clin. Chim. Acta 123: 269-274,

[125168]1982.

[125169]22280.

[125170]Grier, R. E.; Howell, R. R.; Wu, D.; McCabe, E. R. B.: Isolated

[125171]glycerol kinase deficiency: nutritional and molecular genetic studies.

[125172](Abstract) Am. J. Hum. Genet. 45 (suppl.): A6 only, 1989.

[125173]22281.

[125174]Guggenheim, M. A.; McCabe, E. R. B.; Roig, M.; Goodman, S. I.;

[125175]Lum, G. M.; Bullen, W. W.; Ringel, S. P.: Glycerol kinase deficiency

[125176]with neuromuscular, skeletal, and adrenal abnormalities. Ann. Neurol. 7:

[125177]441–449, 1980.

[125178]22282.

[125179]Guo, W.; Worley, K.; Adams, V.; Mason, J.; Sylvester–Jackson,

[125180]D.; Zhang, Y.–H.; Towbin, J. A.; Fogt, D. D.; Madu, S.; Wheeler, D.

[125181]A.; McCabe, E. R. B.: Genomic scanning for expressed sequences in

[125182]Xp21 identifies the glycerol kinase gene. Nature Genet. 4: 367–371,

[125183]1993.

[125184]22283.

[125185]Hammond, J.; Howard, N. J.; Brookwell, R.; Purvis–Smith, S.; Wilcken,

- [125186]B.; Hoogenraad, N.: Proposed assignment of loci for X-linked adrenal hypoplasia and glycerol kinase genes. (Letter) Lancet I: 54 only, 1985.
- [125189]22284.
- [125190]Love, D. R.; Bloomfield, J. F.; Kenwrick, S. J.; Yates, J. R. W.; Davies, K. E.: Physical mapping distal to the DMD locus. Genomics 8: 106–112, 1990.
- [125193]22285.
- [125194]Mahbubul Huq, A. H. M.; Lovell, R. S.; Ou, C.-N.; Beaudet, A. L.; Craigen, W. J.: X-linked glycerol kinase deficiency in the mouse leads to growth retardation, altered fat metabolism, autonomous glucocorticoid secretion and neonatal death. Hum. Molec. Genet. 6: 1803–1809, 1997.
- [125198]22286.
- [125199]Mahbubul Huq, A. H. M.; Lovell, R. S.; Sampson, M. J.;

Decker,

[125200]W. K.; Dinulos, M. B.; Disteche, C. M.; Craigen, W. J.: Isolation,

[125201]mapping, and functional expression of the mouse X chromosome glycerol

[125202]kinase gene. Genomics 36: 530–534, 1996.

[125203]22287.

[125204]Nishimura, J.; Murakami, Y.; Kinoshita, T.: Paroxysmal nocturnal

[125205]hemoglobinuria: an acquired genetic disease. Am. J. Hemat. 62: 175–182,

[125206]1999.

[125207]22288.

[125208]Kosho, T.; Muroya, K.; Nagai, T.; Fujimoto, M.; Yokoya, S.; Sakamoto,

[125209]H.; Hirano, T.; Terasaki, H.; Ohashi, H.; Nishimura, G.; Sato, S.;

[125210]Matsuo, N.; Ogata, T.: Skeletal features and growth patterns in 14

[125211]patients with haploinsufficiency of SHOX: implications for the development

[125212]of Turner syndrome. J. Clin. Endocr. Metab. 84:

4613–4621, 1999.

[125213]22289.

[125214]Kuznetzova, T.; Baranov, A.; Ivaschenko, T.; Savitsky, G.
A.;

[125215]Lanceva, O. E.; Wang, M. R.; Giollant, M.; Malet, P.;
Kascheeva, T.;

[125216]Vakharlovsky, V.; Baranov, V. S.: X;Y translocation in a girl
with

[125217]short stature and some features of Turner's syndrome: cy-
togenetic

[125218]and molecular studies. J. Med. Genet. 31: 649–651, 1994.

[125219]22290.

[125220]May, C. A.; Shone, A. C.; Kalaydjieva, L.; Sajantila, A.; Jef-
freys,

[125221]A. J.: Crossover clustering and rapid decay of linkage dis-
equilibrium

[125222]in the Xp/Yp pseudoautosomal gene SHOX. Nature Genet.
31: 272–275,

[125223]2002.

[125224]22291.

[125225]Ogata, T.; Goodfellow, P.; Petit, C.; Aya, M.; Matsuo, N.:

Short

[125226] stature in a girl with a terminal Xp deletion distal to

DXYS15: localisation

[125227] of a growth gene(s) in the pseudoautosomal region. J.

Med. Genet. 29:

[125228] 455–459, 1992.

[125229] 22292.

[125230] Ogata, T.; Kosho, T.; Wakui, K.; Fukushima, Y.; Yoshimoto,

M.;

[125231] Miharu, N.: Short stature homeobox-containing gene du-

plication on

[125232] the der(X) chromosome in a female with 45,X/46,X, der(X),

gonadal

[125233] dysgenesis, and tall stature. J. Clin. Endocr. Metab. 85:

2927–2930,

[125234] 2000.

[125235] 22293.

[125236] Ogata, T.; Matsuo, N.; Nishimura, G.: SHOX haploinsuffi-

ciency

[125237] and overdosage: impact of gonadal function status. J.

Med. Genet. 38:

[125238] 1–6, 2001.

[125239]22294.

[125240]Ogata, T.; Muroya, K.; Sasaki, G.; Nishimura, G.; Kitoh, H.;
Hattori,

[125241]T.: SHOX nullizygosity and haploinsufficiency in a
Japanese family:

[125242]implication for the development of Turner skeletal fea-
tures. J. Clin.

[125243]Endocr. Metab. 87: 1390–1394, 2002.

[125244]22295.

[125245]Ogata, T.; Petit, C.; Rappold, G.; Matsuo, N.; Matsumoto,
T.;

[125246]Goodfellow, P.: Chromosomal localisation of a pseudoau-
tosomal growth

[125247]gene(s). J. Med. Genet. 29: 624–628, 1992.

[125248]22296.

[125249]Ogata, T.; Yoshizawa, A.; Muroya, K.; Matsuo, N.;
Fukushima, Y.;

[125250]Rappold, G.; Yokoya, S.: Short stature in a girl with partial
monosomy

[125251]of the pseudoautosomal region distal to DXYS15: further
evidence for

[125252]the assignment of the critical region for a pseudoautosomal growth

[125253]gene(s). J. Med. Genet. 32: 831–834, 1995.

[125254]22297.

[125255]Pfeiffer, R. A.: Observations in a case of an X/Y translocation,

[125256]t(X;Y)(p22;q11), in a mother and son. Cytogenet. Cell Genet. 26:

[125257]150–157, 1980.

[125258]22298.

[125259]Rao, E.; Blaschke, R. J.; Marchini, A.; Niesler, B.; Burnett,

[125260]M.; Rappold, G. A.: The Leri-Weill and Turner syndrome homeobox gene

[125261]SHOX encodes a cell-type specific transcriptional activator. Hum.

[125262]Molec. Genet. 10: 3083–3091, 2001.

[125263]22299.

[125264]Rao, E.; Weiss, B.; Fukami, M.; Mertz, A.; Meder, J.; Ogata, T.;

[125265]Heinrich, U.; Garcia-Heras, J.; Schiebel, K.; Rappold, G. A.: FISH-deletion

[125266]mapping defines a 270-kb short stature critical interval in the pseudoautosomal

[125267]region PAR1 on human sex chromosomes. Hum. Genet. 100: 236–239,

[125268]1997.

[125269]22300.

[125270]Rao, E.; Weiss, B.; Fukami, M.; Rump, A.; Niesler, B.; Mertz,

[125271]A.; Muroya, K.; Binder, G.; Kirsch, S.; Winkelmann, M.; Nordsiek,

[125272]G.; Heinrich, U.; Breuning, M. H.; Ranke, M. B.; Rosenthal, A.; Ogata,

[125273]T.; Rappold, G. A.: Pseudoautosomal deletions encompassing a novel

[125274]homeobox gene cause growth failure in idiopathic short stature and

[125275]Turner syndrome. Nature Genet. 16: 54–63, 1997.

[125276]22301.

[125277]Rappold, G. A.; Fukami, M.; Niesler, B.; Schiller, S.; Zumkeller,

[125278]W.; Bettendorf, M.; Heinrich, U.; Vlachopapadoupoulou, E.; Reinehr,

- [125279]T.; Onigata, K.; Ogata, T.: Deletions of the homeobox gene SHOX (short stature homeobox) are an important cause of growth failure in children with short stature. J. Clin. Endocr. Metab. 87: 1402–1406, 2002.
- [125282]22302.
- [125283]Schiller, S.; Spranger, S.; Schechinger, B.; Fukami, M.; Merker, S.; Drop, S. L. S.; Troger, J.; Knoblauch, H.; Kunze, J.; Seidel, J.; Rappold, G. A.: Phenotypic variation and genetic heterogeneity in Leri–Weill syndrome. Europ. J. Hum. Genet. 8: 54–62, 2000.
- [125287]22303.
- [125288]Shears, D. J.; Guillen–Navarro, E.; Sempere–Miralles, M.; Domingo–Jimenez, R.; Scambler, P. J.; Winter, R. M.: Pseudodominant inheritance of Langer mesomelic dysplasia caused by a SHOX homeobox missense mutation. Am.

[125291]J. Med. Genet. 110: 153–157, 2002.

[125292]22304.

[125293]Zinn, A. R.; Wei, F.; Zhang, L.; Elder, F. F.; Scott, C. I., Jr.;

[125294]Marttila, P.; Ross, J. L.: Complete SHOX deficiency causes
Langer

[125295]mesomelic dysplasia. Am. J. Med. Genet. 110: 158–163,
2002.

[125296]22305.

[125297]Zuffardi, O.; Maraschio, P.; Lo Curto, F.; Muller, U.; Gia-
rola,

[125298]A.; Perotti, L.: The role of Yp in sex determination: new
evidence

[125299]from X/Y translocations. Am. J. Med. Genet. 12: 175–184,
1982.

[125300]22306.

[125301]Acker, K. J.; Roels, H.; Beelaerts, W.; Pasternack, A.; Val-
cke,

[125302]R.: The histologic lesions of the kidney in the oculo-
cerebro-renal

[125303]syndrome of Lowe. Nephron 4: 193–214, 1967.

[125304]22307.

[125305]Attree, O.; Olivos, I. M.; Okabe, I.; Bailey, L. C.; Nelson, D.

[125306]L.; Lewis, R. A.; McInnes, R. R.; Nussbaum, R. L.: The
Lowe's oculocerebrorenal

[125307]syndrome gene encodes a protein highly homologous to
inositol polyphosphate-5-phosphatase. Nature 358:

[125308]239-242, 1992.

[125309]22308.

[125310]Auricchio, S.; Frischknecht, W.; Shmerling, D. H.: Primare
Tubulopathien.

[125311]III. Ein Fall von oculo-cerebro-renalem Syndrom
(Lowe-Syndrome). Helv.

[125312]Paediat. Acta 16: 647-655, 1961.

[125313]22309.

[125314]Bailey, L. C., Jr.; Olivos, I. M.; Leahey, A. M.; Attree, O. F.;

[125315]Okabe, I.; Lewis, R. A.; MacInnes, R. R.; Spinner, N. B.; Nel-
son,

[125316]D. L.; Nussbaum, R. L.: Characterization of a candidate
gene for

[125317]OCRL. (Abstract) Am. J. Hum. Genet. 51 (suppl.): A4 only,
1992.

[125318]22310.

[125319]Charnas, L. R.; Bernardini, I.; Rader, D.; Hoeg, J. M.; Gahl, W.

[125320]A.: Clinical and laboratory findings in the oculocerebrorenal syndrome

[125321]of Lowe, with special reference to growth and renal function. New

[125322]Eng. J. Med. 324: 1318–1325, 1991.

[125323]22311.

[125324]Chutorian, A.; Rowland, L. P.: Lowe's syndrome. Neurology 16:

[125325]115–122, 1966.

[125326]22312.

[125327]Delleman, J. W.; Bleeker-Wagemakers, E. M.; van Veelen, A. W. C.

[125328]: Opacities of the lens indicating carrier status in the oculo-cerebro-renal

[125329](Lowe) syndrome. J. Pediat. Ophthal. 14: 205–212, 1977.

[125330]22313.

[125331]Duden, R.; Griffiths, G.; Frank, R.; Argos, P.; Kreis, T. E.:

[125332]Beta-COP, a 110 kd protein associated with non-clathrin-coated vesicles

[125333]and the Golgi complex, shows homology to beta-adaptin.

Cell 64:

[125334]649–665, 1991.

[125335]22314.

[125336]Gardner, R. J. M.; Brown, N.: Lowe's syndrome: identification

[125337]of carriers by lens examination. J. Med. Genet. 13:
449–464, 1976.

[125338]22315.

[125339]Gropman, A.; Levin, S.; Yao, L.; Lin, T.; Suchy, S.; Sabnis,
S.;

[125340]Hadley, D.; Nussbaum, R.: Unusual renal features of Lowe
syndrome

[125341]in a mildly affected boy. Am. J. Med. Genet. 95: 461–466,
2000.

[125342]22316.

[125343]Harris, L. S.; Gitter, K. A.; Galin, M. A.; Plechaty, G. P.:

[125344]Oculo–cerebro–renal syndrome: report of a case in a baby
girl. Brit.

[125345]J. Ophthalm. 54: 278–280, 1970.

[125346]22317.

[125347]Hittner, H. M.; Carroll, A. J.; Prchal, J. T.: Linkage studies
[125348]in carriers of Lowe oculo-cerebro-renal syndrome. Am. J.
Hum. Genet. 34:
[125349]966-971, 1982.
[125350]22318.
[125351]Hodgson, S. V.; Heckmatt, J. Z.; Hughes, E.; Crolla, J. A.;
Dubowitz,
[125352]V.; Bobrow, M.: A balanced de novo X/autosome translo-
cation in a

[125353]girl with manifestations of Lowe syndrome. Am. J. Med.
Genet. 23:

[125354]837–847, 1986.

[125355]22319.

[125356]Kawano, T.; Indo, Y.; Nakazato, H.; Shimadzu, M.; Matsuda, I.

[125357]: Oculocerebrorenal syndrome of Lowe: three mutations in
the OCRL1

[125358]gene derived from three patients with different phenotypes. Am. J.

[125359]Med. Genet. 77: 348–355, 1998.

[125360]22320.

[125361]Kenworthy, L.; Charnas, L.: Evidence for a discrete behavioral

[125362]phenotype in the oculocerebrorenal syndrome of Lowe.
Am. J. Med.

[125363]Genet. 59: 283–290, 1995.

[125364]22321.

[125365]Kenworthy, L.; Park, T.; Charnas, L. R.: Cognitive and behavioral

[125366]profile of the oculocerebrorenal syndrome of Lowe. Am. J.

Med. Genet. 46:

[125367]297–303, 1993.

[125368]22322.

[125369]Leahey, A.–M.; Charnas, L. R.; Nussbaum, R. L.: Nonsense mutations

[125370]in the OCRL–1 gene in patients with the oculocerebrorenal syndrome

[125371]of Lowe. Hum. Molec. Genet. 2: 461–463, 1993.

[125372]22323.

[125373]Leahey, A. M.; Olivos, I. M.; Bailey, L. C.; Nussbaum, R. L.:

[125374]Characterization of the molecular defects in Lowe's syndrome. (Abstract) Am.

[125375]J. Hum. Genet. 51 (suppl.): A220 only, 1992.

[125376]22324.

[125377]Lin, T.; Lewis, R. A.; Nussbaum, R. L.: Molecular confirmation

[125378]of carriers for Lowe syndrome. Ophthalmology 106: 119–122, 1999.

[125379]22325.

[125380]Lin, T.; Orrison, B. M.; Leahey, A.–M.; Suchy, S. F.; Bernard,

[125381]D. J.; Lewis, R. A.; Nussbaum, R. L.: Spectrum of mutations
in the

[125382]OCRL1 gene in the Lowe oculocerebrorenal syndrome.
Am. J. Hum. Genet. 60:

[125383]1384–1388, 1997.

[125384]22326.

[125385]Kutsche, K.; Yntema, H.; Brandt, A.; Jantke, I.; Nothwang,
H. G.;

[125386]Orth, U.; Boavida, M. G.; David, D.; Chelly, J.; Fryns, J.-P.;
Moraine,

[125387]C.; Ropers, H.-H.; Hamel, B. C. J.; van Bokhoven, H.; Gal,
A.: Mutations

[125388]in ARHGEF6, encoding a guanine nucleotide exchange
factor for Rho

[125389]GTPases, in patients with X-linked mental retardation. Na-
ture Genet. 26:

[125390]247–250, 2000.

[125391]22327.

[125392]Lower, K. M.; Gecz, J.: Characterization of ARHGEF6, a
guanine

[125393]nucleotide exchange factor for Rho GTPases and a candi-
date gene for

[125394]X-linked mental retardation: mutation screening in Borjeson-Forssman-Lehmann

[125395]syndrome and MRX27. Am. J. Med. Genet. 100: 43-48, 2001.

[125396]22328.

[125397]Manser, E.; Loo, T.-H.; Koh, C.-G.; Zhao, Z.-S.; Chen, X.-Q.; Tan,

[125398]L.; Tan, I.; Leung, T.; Lim, L.: PAK kinases are directly coupled

[125399]to the PIX family of nucleotide exchange factors. Molec. Cell 1:

[125400]183-192, 1998.

[125401]22329.

[125402]Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Kawarabayashi,

[125403]Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.: Prediction

[125404]of the coding sequences of unidentified human genes. I. The coding

[125405]sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis

[125406]of randomly sampled cDNA clones from human immature

myeloid cell line,

[125407]KG-1. DNA Res. 1: 27-35, 1994.

[125408]22330.

[125409]von Ballestrem, C. L.; Boavida, M. G.; Zuther, C.; Carreiro, M.

[125410]H.; David, D.; Gal, A.; Schwinger, E.: Jumping translocation in a

[125411]phenotypically normal female. Clin. Genet. 49: 156-159, 1996.

[125412]22331.

[125413]Yntema, H. G.; Hamel, B. C.; Smits, A. P.; van Roosmalen, T.; van

[125414]den Helm, B.; Kremer, H.; Ropers, H. H.; Smeets, D. F.; van Bokhoven,

[125415]H.: Localisation of a gene for non-specific X linked mental retardation

[125416](MRX46) to Xq25-q26. J. Med. Genet. 35: 801-805, 1998.

[125417]22332.

[125418]Grozingier, C. M.; Hassig, C. A.; Schreiber, S. L.: Three proteins

[125419]define a class of human histone deacetylases related to

yeast Hda1p. Proc.

[125420]Nat. Acad. Sci. 96: 4868–4873, 1999.

[125421]22333.

[125422]Hubbert, C.; Guardiola, A.; Shao, R.; Kawaguchi, Y.; Ito, A.;
Nixon,

[125423]A.; Yoshida, M.; Wang, X.-F.; Yao, T.-P.: HDAC6 is a mi-
crotubule-associated

[125424]deacetylase. Nature 417: 455–458, 2002.

[125425]22334.

[125426]Mahlknecht, U.; Schnittger, S.; Landgraf, F.; Schoch, C.;
Ottmann,

[125427]O. G.; Hiddemann, W.; Hoelzer, D.: Assignment of the hu-
man histone

[125428]deacetylase 6 gene (HDAC6) to X chromosome p11.23 by
in situ hybridization. Cytogenet.

[125429]Cell Genet. 93: 135–136, 2001.

[125430]22335.

[125431]Nagase, T.; Isikawa, K.; Suyama, M.; Kikuno, R.; Hirose, A.;
M.;

[125432]Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara,
O.: Prediction

[125433]of the coding sequences of unidentified human genes. XII.

The complete

[125434]sequences of 100 new cDNA clones from brain which code
for large proteins

[125435]in vitro. DNA Res. 5: 355–364, 1998.

[125436]22336.

[125437]Ichida, F.; Tsubata, S.; Bowles, K. R.; Haneda, N.; Uese, K.;

[125438]Miyawaki, T.; Dreyer, W. J.; Messina, J.; Li, H.; Bowles, N.
E.; Towbin,

[125439]J. A.: Novel gene mutations in patients with left ventricular
noncompaction

[125440]or Barth syndrome. Circulation 103: 1256–1263, 2001.

[125441]22337.

[125442]Johnston, J.; Kelley, R. I.; Feigenbaum, A.; Cox, G. F.; Iyer,

[125443]G. S.; Funanage, V. L.; Proujansky, R.: Mutation characteri-
zation

[125444]and genotype–phenotype correlation in Barth syndrome.
Am. J. Hum.

[125445]Genet. 61: 1053–1058, 1997.

[125446]22338.

[125447]Kelley, R. I.; Cheatham, J. P.; Clark, B. J.; Nigro, M. A.;
Powell,
[125448]B. R.; Sherwood, G. W.; Sladky, J. T.; Swisher, W. P.: X-
linked dilated
[125449]cardiomyopathy with neutropenia, growth retardation, and
3-methylglutaconic
[125450]aciduria. J. Pediatr. 119: 738-747, 1991.
[125451]22339.
[125452]Sakamoto, O.; Kitoh, T.; Ohura, T.; Ohya, N.; Iinuma, K.:
Novel
[125453]missense mutation (R94S) in the TAZ (G4.5) gene in a
Japanese patient
[125454]with Barth syndrome. J. Hum. Genet. 47: 229-231, 2002.
[125455]22340.
[125456]Sakamoto, O.; Ohura, T.; Katsushima, Y.; Fujiwara, I.;
Ogawa,
[125457]E.; Miyabayashi, S.; Iinuma, K.: A novel intronic mutation
of the
[125458]TAZ (G4.5) gene in a patient with Barth syndrome: cre-
ation of a 5-prime
[125459]splice donor site with variant GC consensus and elonga-
tion of the

[125460]upstream exon. Hum. Genet. 109: 559–563, 2001.

[125461]22341.

[125462]Strasser, K.; Masuda, S.; Mason, P.; Pfannstiel, J.; Oppizzi, M.;

[125463]Rodriguez–Navarro, S.; Rondon, A. G.; Aguilera, A.; Struhl, K.; Reed,

[125464]R.; Hurt, E.: TREX is a conserved complex coupling transcription

[125465]with messenger RNA export. Nature 417: 304–308, 2002.

[125466]22342.

[125467]Fortenberry, Y.; Hwang, J.–R.; Apletalina, E. V.; Lindberg, I.

[125468]: Functional characterization of proSAAS: similarities and differences

[125469]with 7B2. J. Biol. Chem. 277: 5175–5186, 2002.

[125470]22343.

[125471]Fricker, L. D.; McKinzie, A. A.; Sun, J.; Curran, E.; Qian, Y.;

[125472]Yan, L.; Patterson, S. D.; Courchesne, P. L.; Richards, B.; Levin,

[125473]N.; Mzhavia, N.; Devi, L. A.; Douglass, J.: Identification and characterization

[125474]of proSAAS, a granin-like neuroendocrine peptide precursor that inhibits
[125475]prohormone processing. J. Neurosci. 20: 639–648, 2000.
[125476]22344.
[125477]Omori, Y.; Mesnil, M.; Yamasaki, H.: Connexin 32 mutations from
[125478]X-linked Charcot–Marie–Tooth disease patients: functional defects
[125479]and dominant negative effects. Molec. Biol. Cell 7: 907–916, 1996.
[125480]22345.
[125481]Tabaraud, F.; Lagrange, E.; Sindou, P.; Vandenberghe, A.; Levy,
[125482]N.; Vallat, J. M.: Demyelinating X-linked Charcot–Marie–Tooth disease:
[125483]unusual electrophysiological findings. Muscle Nerve 22: 1442–1447,
[125484]1999.
[125485]22346.
[125486]Moser, A. B.; Moser, H. W.: The prenatal diagnosis of X-linked

[125487]adrenoleukodystrophy. Prenatal Diag. 19: 46–48, 1999.

[125488]22347.

[125489]Sugarman, G. I.; Rimoin, D. L.; Lachman, R. S.: The facial–digital–genital

[125490](Aarskog) syndrome. Am. J. Dis. Child. 126: 248–252, 1973.

[125491]22348.

[125492]Tsukahara, M.; Fernandez, G. I.: Umbilical findings in Aarskog

[125493]syndrome. Clin. Genet. 45: 260–265, 1994.

[125494]22349.

[125495]Tyrkus, M.; Bawle, E.; Lipman, S.; Bozimowski, D.; Woolley, P.

[125496]V., Jr.: Aarskog–Scott syndrome inherited as an X-linked dominant

[125497]with full male–female expression. (Abstract) Am. J. Hum. Genet. 32:

[125498]134A only, 1980.

[125499]22350.

[125500]van den Bergh, P.; Fryns, J. P.; Wilms, G.; Piot, R.; Dra–lands,

- [125501]G.; van den Bergh, R.: Anomalous cerebral venous drainage in Aarskog
- [125502]syndrome. Clin. Genet. 25: 288–294, 1984.
- [125503]22351.
- [125504]Zheng, Y.; Fischer, D. J.; Santos, M. F.; Tigyi, G.; Pasteris,
- [125505]N. G.; Gorski, J. L.; Xu, Y.: The faciogenital dysplasia gene product
- [125506]FGD1 functions as a Cdc42Hs–specific guanine–nucleotide exchange factor. J.
- [125507]Biol. Chem. 271: 33169–33172, 1996.
- [125508]22352.
- [125509]Inaba, H.; Fujimaki, M.; Kazazian, H. H., Jr.; Antonarakis, S.
- [125510]E.: MspI polymorphic site in intron 22 of the factor VIII gene in
- [125511]the Japanese population. Hum. Genet. 84: 214–215, 1990.
- [125512]22353.
- [125513]Levinson, B.; Bermingham, J. R., Jr.; Metzenberg, A.; Ken-
- wrick,
- [125514]S.; Chapman, V.; Gitschier, J.: Sequence of the human factor VIII–associated

[125515]gene is conserved in mouse. Genomics 13: 862–865,
1992.

[125516]22354.

[125517]Levinson, B.; Kenwrick, S.; Lakich, D.; Hammonds, G.;
Gitschier,

[125518]J.: A transcribed gene in an intron of the human factor VIII
gene. Genomics 7:

[125519]1–11, 1990.

[125520]22355.

[125521]Peters, M. F.; Ross, C. A.: Isolation of a 40–kDa hunt-
ington-associated

[125522]protein. J. Biol. Chem. 276: 3188–3194, 2001.

[125523]22356.

[125524]Berra, B.; Gornati, R.; Rapelli, S.; Gatti, R.; Mancini, G. M.

[125525]S.; Ciana, G.; Bembi, B.: Infantile sialic acid storage dis-
ease:

[125526]biochemical studies. Am. J. Med. Genet. 58: 24–31, 1995.

[125527]22357.

[125528]Cameron, P. D.; Dubowitz, V.; Besley, G. T. N.; Fensom, A.
H.:

[125529]Sialic acid storage disease. Arch. Dis. Child. 65: 314–315, 1990.

[125530]22358.

[125531]Haataja, L.; Schleutker, J.; Laine, A.–P.; Renlund, M.; Savontaus,

[125532]M.–L.; Dib, C.; Weissenbach, J.; Peltonen, L.; Aula, P.: The genetic

[125533]locus for free sialic acid storage disease maps to the long arm of

[125534]chromosome 6. Am. J. Hum. Genet. 54: 1042–1049, 1994.

[125535]22359.

[125536]Mules, E. H.; Hayflick, S.; Miller, C. S.; Reynolds, L. W.; Thomas,

[125537]G. H.: Six novel deleterious and three neutral mutations in the gene

[125538]encoding the alpha–subunit of hexosaminidase A in non–Jewish individuals. Am.

[125539]J. Hum. Genet. 50: 834–841, 1992.

[125540]22360.

[125541]Myerowitz, R.: Tay–Sachs disease–causing mutations and neutral

[125542]polymorphisms in the Hex A gene. Hum. Mutat. 9:
195–208, 1997.

[125543]22361.

[125544]Myerowitz, R.: Splice junction mutation in some Ashkenazi
Jews

[125545]with Tay–Sachs disease: evidence against a single defect
within this

[125546]ethnic group. Proc. Nat. Acad. Sci. 85: 3955–3959, 1988.

[125547]22362.

[125548]Myerowitz, R.; Costigan, F. C.: The major defect in Ashke-
nazi

[125549]Jews with Tay–Sachs disease is an insertion in the gene for
the alpha–chain

[125550]of beta–hexosaminidase. J. Biol. Chem. 263:
18587–18589, 1988.

[125551]22363.

[125552]Myerowitz, R.; Hogikyan, N. D.: Different mutations in
Ashkenazi

[125553]Jewish and non–Jewish French Canadians with Tay–Sachs
disease. Science 232:

[125554]1646–1648, 1986.

[125555]22364.

[125556]Myerowitz, R.; Hogikyan, N. D.: A deletion involving Alu sequences

[125557]in the beta-hexosaminidase alpha-chain gene of French Canadians with

[125558]Tay-Sachs disease. J. Biol. Chem. 262: 15396-15399, 1987.

[125559]22365.

[125560]Myerowitz, R.; Piekarz, R.; Neufeld, E. F.; Shows, T. B.; Suzuki,

[125561]K.: Human beta-hexosaminidase alpha chain: coding sequence and homology

[125562]with the beta chain. Proc. Nat. Acad. Sci. 82: 7830-7834, 1985.

[125563]22366.

[125564]Myerowitz, R.; Proia, R. L.: cDNA clone for the alpha-chain of

[125565]human beta-hexosaminidase: deficiency of alpha-chain mRNA in Ashkenazi

[125566]Tay-Sachs fibroblasts. Proc. Nat. Acad. Sci. 81: 5394-5398, 1984.

[125567]22367.

[125568]Nakai, H.; Byers, M. G.; Nowak, N. J.; Shows, T. B.: Assignment

[125569]of beta-hexosaminidase A alpha-subunit to human chromosomal region

[125570]15q23-q24. Cytogenet. Cell Genet. 56: 164, 1991.

[125571]22368.

[125572]Nakano, T.; Muscillo, M.; Ohno, K.; Hoffman, A. J.; Suzuki, K.

[125573]: A point mutation in the coding sequence of the beta-hexosaminidase

[125574]alpha gene results in defective processing of the enzyme protein in

[125575]an unusual GM2-gangliosidosis variant. J. Neurochem. 51: 984-987,

[125576]1988.

[125577]22369.

[125578]Nakano, T.; Nanba, E.; Tanaka, A.; Ohno, K.; Suzuki, Y.; Suzuki,

[125579]K.: A new point mutation within exon 5 of beta-hexosaminidase alpha

[125580]gene in a Japanese infant with Tay-Sachs disease. Ann. Neurol. 27:

[125581]465–473, 1990.

[125582]22370.

[125583]Navon, R.; Kolodny, E. H.; Mitsumoto, H.; Thomas, G. H.;
Proia,

[125584]R. L.: Ashkenazi–Jewish and non–Jewish adult G(M2) gan-
gliosidosis

[125585]patients share a common genetic defect. Am. J. Hum.
Genet. 46: 817–821,

[125586]1990.

[125587]22371.

[125588]Navon, R.; Proia, R. L.: The mutations in Ashkenazi Jews
with

[125589]adult G(M2) gangliosidosis, the adult form of Tay–Sachs
disease. Science 243:

[125590]1471–1474, 1989.

[125591]22372.

[125592]Navon, R.; Proia, R. L.: Tay–Sachs disease in Moroccan
Jews:

[125593]deletion of a phenylalanine in the alpha–subunit of beta–
hexosaminidase. Am.

[125594]J. Hum. Genet. 48: 412–419, 1991.

[125595]22373.

[125596]Nishimoto, J.; Tanaka, A.; Nanba, E.; Suzuki, K.: Expression

[125597]of the beta-hexosaminidase alpha subunit gene with the four-base insertion

[125598]of infantile Jewish Tay-Sachs disease. J. Biol. Chem. 266: 14306-14309,

[125599]1991.

[125600]22374.

[125601]Koch, M. C.; Ricker, K.; Otto, M.; Grimm, T.; Hoffman, E. P.;

[125602]Rudel, R.; Bender, K.; Zoll, B.; Harper, P. S.; Lehmann-Horn, F.:

[125603]Confirmation of linkage of hyperkalaemic periodic paralysis to chromosome

[125604]17. J. Med. Genet. 28: 583-586, 1991.

[125605]22375.

[125606]Lehmann-Horn, F.; Rudel, R.; Ricker, K.: Non-dystrophic myotonias

[125607]and periodic paralyses. Neuromusc. Disord. 3: 161-168, 1993.

[125608]22376.

[125609]McClatchey, A. I.: Personal Communication. Boston, Mass.
3/16/1993.

[125610]22377.

[125611]McClatchey, A. I.; McKenna-Yasek, D.; Cros, D.; Worthen,
H. G.;

[125612]Kuncl, R. W.; DeSilva, S. M.; Cornblath, D. R.; Gusella, J. F.;
Brown,

[125613]R. H., Jr.: Novel mutations in families with unusual and
variable

[125614]disorders of the skeletal muscle sodium channel. Nature
Genet. 2:

[125615]148–152, 1992.

[125616]22378.

[125617]McClatchey, A. I.; Trofatter, J.; McKenna-Yasek, D.;
Raskind,

[125618]W.; Bird, T.; Pericak-Vance, M.; Gilchrist, J.; Arahata, K.;
Radosavljevic,

[125619]D.; Worthen, H. G.; Van den Bergh, P.; Haines, J. L.;
Gusella, J.

[125620]F.; Brown, R. H., Jr.: Dinucleotide repeat polymorphisms at
the SCN4A

[125621]locus suggest allelic heterogeneity of hyperkalemic periodic paralysis

[125622]and paramyotonia congenita. Am. J. Hum. Genet. 50: 896–901, 1992.

[125623]22379.

[125624]McClatchey, A. I.; Van den Bergh, P.; Pericak–Vance, M. A.; Raskind,

[125625]W.; Verellen, C.; McKenna–Yasek, D.; Rao, K.; Haines, J. L.; Bird,

[125626]T.; Brown, R. H., Jr.; Gusella, J. F.: Temperature–sensitive mutations

[125627]in the III–IV cytoplasmic loop region of the skeletal muscle sodium

[125628]channel gene in paramyotonia congenita. Cell 68: 769–774, 1992.

[125629]22380.

[125630]Moslehi, R.; Langlois, S.; Yam, I.; Friedman, J. M.: Linkage

[125631]of malignant hyperthermia and hyperkalemic periodic paralysis to the

[125632]adult skeletal muscle sodium channel (SCN4A) gene in a large pedigree. Am.

[125633]J. Med. Genet. 76: 21–27, 1998.

[125634]22381.

[125635]Ptacek, L. J.; George, A. L., Jr.; Barchi, R. L.; Griggs, R. C.;

[125636]Riggs, J. E.; Robertson, M.; Leppert, M. F.: Mutations in an
S4 segment

[125637]of the adult skeletal muscle sodium channel cause
paramyotonia congenita. Neuron 8:

[125638]891–897, 1992.

[125639]22382.

[125640]Ptacek, L. J.; George, A. L., Jr.; Griggs, R. C.; Tawil, R.;
Kallen,

[125641]R. G.; Barchi, R. L.; Robertson, M.; Leppert, M. F.: Identifi-
cation

[125642]of a mutation in the gene causing hyperkalemic periodic
paralysis. Cell 67:

[125643]1021–1027, 1991.

[125644]22383.

[125645]Ptacek, L. J.; Gouw, L.; Kwieninski, H.; McManis, P.;
Mendell,

[125646]J. R.; Barohn, A. L., Jr.; Robertson, M.; Leppert, M. F.:
Sodium

[125647]channel mutations in paramyotonia congenita and hyper-

kalemic periodic

[125648]paralysis. Ann. Neurol. 33: 300–307, 1993.

[125649]22384.

[125650]Ptacek, L. J.; Tawil, R.; Griggs, R. C.; Meola, G.; McManis, P.;

[125651]Barohn, R. J.; Mendell, J. R.; Harris, C.; Spitzer, R.; Santiago,

[125652]F.; Leppert, M. F.: Sodium channel mutations in acetazolamide-responsive

[125653]myotonia congenita, paramyotonia congenita, and hyperkalemic periodic

[125654]paralysis. Neurology 44: 1500–1503, 1994.

[125655]22385.

[125656]Ptacek, L. J.; Tawil, R.; Griggs, R. C.; Storvick, D.; Leppert,

[125657]M.: Linkage of atypical myotonia congenita to a sodium channel locus. Neurology 42:

[125658]431–433, 1992.

[125659]22386.

[125660]Ptacek, L. J.; Tyler, F.; Trimmer, J. S.; Agnew, W. S.; Leppert,

[125661]M.: Analysis in a large hyperkalemic periodic paralysis

pedigree

[125662]supports tight linkage to a sodium channel locus. Am. J. Hum. Genet. 49:

[125663]378–382, 1991.

[125664]22387.

[125665]Ricker, K.; Moxley, R. T., III; Heine, R.; Lehmann–Horn, F.:

[125666]Myotonia fluctuans: a third type of muscle sodium channel disease. Arch.

[125667]Neurol. 51: 1095–1102, 1994.

[125668]22388.

[125669]Rudolph, J. A.; Spier, S. J.; Byrns, G.; Hoffman, E. P.: Linkage

[125670]of hyperkalaemic periodic paralysis in Quarter horses to the horse

[125671]adult skeletal muscle sodium channel gene. Anim. Genet. 23: 241–250,

[125672]1992.

[125673]22389.

[125674]Rudolph, J. A.; Spier, S. J.; Byrns, G.; Rojas, C. V.; Bernoco,

[125675]D.; Hoffman, E. P.: Periodic paralysis in Quarter horses: a

sodium

[125676]channel mutation disseminated by selective breeding. *Nature Genet.* 2:

[125677]144–147, 1992.

[125678]22390.

[125679]Trudell, R. G.; Kaiser, K. K.; Griggs, R. C.: Acetazolamide responsive

[125680]myotonia congenita. *Neurology* 37: 488–491, 1987.

[125681]22391.

[125682]Tyler, F. H.; Stephens, F. E.; Gunn, F. D.; Perkoff, G. T.: Studies

[125683]in disorders of muscle. VII. Clinical manifestations and inheritance

[125684]of a type of periodic paralysis without hypokalemia. *J. Clin.*

[125685]Invest. 30: 492–502, 1951.

[125686]22392.

[125687]European Polycystic Kidney Disease Consortium: The polycystic

[125688]kidney disease 1 gene encodes a 14 kb transcript and lies within a

[125689] duplicated region on chromosome 16. Cell 77: 881–894, 1994.

[125690] 22393.

[125691] Gabow, P. A.: Autosomal dominant polycystic kidney disease. New

[125692] Eng. J. Med. 329: 332–342, 1993.

[125693] 22394.

[125694] Gal, A.; Wirth, B.; Kaariainen, H.; Lucotte, G.; Landais, P.;

[125695] Gillessen–Kaesbach, G.; Muller–Wiefel, D. E.; Zerres, K.: Childhood

[125696] manifestation of autosomal dominant polycystic kidney disease: no

[125697] evidence for genetic heterogeneity. Clin. Genet. 35: 13–19, 1989.

[125698] 22395.

[125699] Germino, G. G.; Barton, N. J.; Lamb, J.; Higgs, D. R.; Harris,

[125700] P.; Xiao, G. H.; Scherer, G.; Nakamura, Y.; Reeders, S. T.: Identification

[125701] of a locus which shows no genetic recombination with the autosomal

[125702] dominant polycystic kidney disease gene on chromosome

16. Am. J.

[125703]Hum. Genet. 46: 925–933, 1990.

[125704]22396.

[125705]Germino, G. G.; Weinstat–Saslow, D.; Himmelbauer, H.;
Gillespie,

[125706]G. A. J.; Somlo, S.; Wirth, B.; Barton, N.; Harris, K. L.;
Frischauf,

[125707]A.–M.; Reeders, S. T.: The gene for autosomal dominant
polycystic

[125708]kidney disease lies in a 750–kb CpG-rich region. Ge-
nomics 13: 144–151,

[125709]1992.

[125710]22397.

[125711]Gillespie, G. A. J.; Somlo, S.; Germino, G. G.; Weinstat–
Saslow,

[125712]D.; Reeders, S. T.: CpG island in the region of an autoso-
mal dominant

[125713]polycystic kidney disease locus defines the 5–prime end
of a gene

[125714]encoding a putative proton channel. Proc. Nat. Acad. Sci.
88: 4289–4293,

[125715]1991.

[125716]22398.

[125717]Harris, P. C.; Barton, N. J.; Higgs, D. R.; Reeders, S. T.;
Wilkie,

[125718]A. O. M.: A long-range restriction map between the alpha-
globin complex

[125719]and a marker closely linked to the polycystic kidney dis-
ease 1 (PKD1)

[125720]locus. Genomics 7: 195-206, 1990.

[125721]22399.

[125722]Harris, P. C.; Thomas, S.; Ratcliffe, P. J.; Breuning, M. H.;

[125723]Coto, E.; Lopez-Larrea, C.: Rapid genetic analysis of fami-
lies with

[125724]polycystic kidney disease 1 by means of a microsatellite
marker. Lancet 338:

[125725]1484-1487, 1991.

[125726]22400.

[125727]Himmelbauer, H.; Pohlschmidt, M.; Snarey, A.; Germino, G.
G.;

[125728]Weinstat-Saslow, D.; Somlo, S.; Reeders, S. T.; Frischauf,
A.-M.:

[125729]Human-mouse homologies in the region of the polycystic

kidney disease

[125730]gene (PKD1). Genomics 13: 35–38, 1992.

[125731]22401.

[125732]Himmelbauer, H.; Pohlschmidt, M.; Snarey, A.; Somlo, S.;
Weinstat-Saslow,

[125733]D.; Germino, G.; Reeders, S.; Frischauf, A.–M.: A syntenic
region

[125734]between human 16p13.3 and mouse chromosome 17.
(Abstract) Cytogenet.

[125735]Cell Genet. 58: 2136, 1991.

[125736]22402.

[125737]Hughes, J.; Ward, C. J.; Peral, B.; Aspinwall, R.; Clark, K.;

[125738]San Millan, J. L.; Gamble, V.; Harris, P. C.: The polycystic
kidney

[125739]disease 1 (PKD1) gene encodes a novel protein with multi-
ple cell recognition

[125740]domains. Nature Genet. 10: 151–160, 1995.

[125741]22403.

[125742]International Polycystic Kidney Disease Consortium: Poly-
cystic

[125743]kidney disease: the complete structure of the PKD1 gene

and its protein. Cell 81:

[125744]289–298, 1995.

[125745]22404.

[125746]Keith, T.; Reeders, S. T.; Brown, V. A.; Green, P.; Phipps, P.;

[125747]Bricker, A.; Barton, N. J.; Collier, P. S.; Knowlton, R.; Nelson,
son,

[125748]C.; Helms, C.; Donis–Keller, H.: Linkage studies of autosomal dominant

[125749]polycystic kidney disease (PKD1) and a genetic map of chromosome 16

[125750]based on 40 RFLP markers. (Abstract) Am. J. Hum. Genet.
41: A172,

[125751]1987.

[125752]22405.

[125753]Fujii, J.; Zarain–Herzberg, A.; Willard, H. F.; Tada, M.; MacLennan,

[125754]D. H.: Structure of the rabbit phospholamban gene, cloning of the

[125755]human cDNA, and assignment of the gene to human chromosome 6. J.

[125756]Biol. Chem. 266: 11669–11675, 1991.

[125757]22406.

[125758]McTiernan, C. F.; Frye, C. S.; Lemster, B. H.; Kinder, E. A.;
Ogletree-Hughes,

[125759]M. L.; Moravec, C. S.; Feldman, A. M.: The human phos-
pholamban gene:

[125760]structure and expression. J. Molec. Cell Cardiol. 31:
679-692, 1999.

[125761]22407.

[125762]Otsu, K.; Fujii, J.; Periasamy, M.; Difilippantonio, M.; Up-
pender,

[125763]M.; Ward, D. C.; MacLennan, D. H.: Chromosome mapping
of five human

[125764]cardiac and skeletal muscle sarcoplasmic reticulum pro-
tein genes. Genomics 17:

[125765]507-509, 1993.

[125766]22408.

[125767]Barhanin, J.; Lesage, F.; Guillemare, E.; Fink, M.; Lazdun-
ski,

[125768]M.; Romey, G.: K(v)LQT1 and Isk (mink) proteins associate
to form

[125769]the I(Ks) cardiac potassium current. Nature 384: 78-80,
1996.

[125770]22409.

[125771]Marx, S. O.; Kurokawa, J.; Reiken, S.; Motoike, H.;

D'Armiento,

[125772]J.; Marks, A. R.; Kass, R. S.: Requirement of a macromolecular signaling

[125773]complex for beta adrenergic receptor modulation of the KCNQ1-KCNE1

[125774]potassium channel. Science 295: 496-499, 2002.

[125775]22410.

[125776]Sanguinetti, M. C.; Curran, M. E.; Zou, A.; Shen, J.; Spector,

[125777]P. S.; Atkinson, D. L.; Keating, M. T.: Coassembly of K(v)LQT1 and

[125778]minK (IsK) proteins to form cardiac I(Ks) potassium channel. Nature 384:

[125779]80-83, 1996.

[125780]22411.

[125781]Coleman, R. A.; Smith, W. L.; Narumiya, S.: VIII. International

[125782]union of pharmacology classification of prostanoid receptors: properties,

[125783]distribution, and structure of the receptors and their sub-

types. Pharm.

[125784]Rev. 46: 205–229, 1994.

[125785]22412.

[125786]Duncan, A. M. V.; Anderson, L. L.; Funk, C. D.;
Abramovitz, M.;

[125787]Adam, M.: Chromosomal localization of the human
prostanoid receptor

[125788]gene family. Genomics 25: 740–742, 1995.

[125789]22413.

[125790]White, D. M.; Mikol, D. D.; Espinosa, R.; Weimer, B.; Le
Beau,

[125791]M. M.; Stefansson, K.: Structure and chromosomal local-
ization of

[125792]the human gene for a brain form of prostaglandin D-2
synthase. J.

[125793]Biol. Chem. 267: 23202–23208, 1992.

[125794]22414.

[125795]Sales, K. J.; Katz, A. A.; Davis, M.; Hinz, S.; Soeters, R. P.;

[125796]Hofmeyr, M. D.; Millar, R. P.; Jabbour, H. N.: Cyclooxyge-
nase-2 expression

[125797]and prostaglandin E2 synthesis are up-regulated in carci-

nomas of the

[125798]cervix: a possible autocrine/paracrine regulation of neoplastic cell

[125799]function via EP2/EP4 receptors. J. Clin. Endocr. Metab. 86: 2243–2249,

[125800]2001.

[125801]22415.

[125802]Taketo, M.; Rochelle, J. M.; Sugimoto, Y.; Namba, T.; Honda, A.;

[125803]Negishi, M.; Ichikawa, A.; Narumiya, S.; Seldin, M. F.: Mapping of

[125804]the genes encoding mouse thromboxane A2 receptor and prostaglandin

[125805]E receptor subtypes EP2 and EP3. Genomics 19: 585–588, 1994.

[125806]22416.

[125807]Catella–Lawson, F.; Reilly, M. P.; Kapoor, S. C.; Cucchiara, A.

[125808]J.; DeMarco, S.; Tournier, B.; Vyas, S. N.; FitzGerald, G. A.: Cyclooxygenase

[125809]inhibitors and the antiplatelet effects of aspirin. New Eng. J. Med. 345:

[125810]1809–1817, 2001.

[125811]22417.

[125812]Dube, J.-N.; Drouin, J.; Aminian, M.; Plant, M. H.; La-neuville,

[125813]O.: Characterization of a partial prostaglandin endoperox-
ide H synthase-1

[125814]deficiency in a patient with a bleeding disorder. Brit. J.
Haemat. 113:

[125815]878–885, 2001.

[125816]22418.

[125817]Funk, C. D.; Funk, L. B.; Kennedy, M. E.; Pong, A. S.;
Fitzgerald,

[125818]G. A.: Human platelet/erythroleukemia cell prostaglandin
G/H synthase:

[125819]cDNA cloning, expression, and gene chromosomal as-
signment. FASEB

[125820]J. 5: 2304–2312, 1991.

[125821]22419.

[125822]Gavett, S. H.; Madison, S. L.; Chulada, P. C.; Scarborough,
P.

[125823]E.; Qu, W.; Boyle, J. E.; Tiano, H. F.; Lee, C. A.; Langenbach,

R.;

[125824]Roggli, V. L.; Zeldin, D. C.: Allergic lung responses are increased

[125825]in prostaglandin H synthase-deficient mice. J. Clin. Invest. 104:

[125826]721-732, 1999.

[125827]22420.

[125828]Kirschenbaum, A.; Liotta, D. R.; Yao, S.; Liu, X.-H.; Klausner,

[125829]A. P.; Unger, P.; Shapiro, E.; Leav, I.; Levine, A. C.: Immunohistochemical

[125830]localization of cyclooxygenase-1 and cyclooxygenase-2 in the human

[125831]fetal and adult male reproductive tracts. J. Clin. Endocr. Metab. 85:

[125832]3436-3441, 2000.

[125833]22421.

[125834]Langenbach, R.; Morham, S. G.; Tiano, H. F.; Loftin, C. D.; Ghanayem,

[125835]B. I.; Chulada, P. C.; Mahler, J. F.; Lee, C. A.; Goulding, E. H.;

[125836]Kluckman, K. D.; Kim, H. S.; Smithies, O.: Prostaglandin

synthase

[125837]1 gene disruption in mice reduces arachidonic acid-induced inflammation

[125838]and indomethacin-induced gastric ulceration. Cell 83: 483-492, 1995.

[125839]22422.

[125840]Malkowski, M. G.; Ginell, S. L.; Smith, W. L.; Garavito, R. M.

[125841]: The productive conformation of arachidonic acid bound to prostaglandin

[125842]synthase. Science 289: 1933-1937, 2000.

[125843]22423.

[125844]Pareti, F. I.; Manucci, P. M.; D'Angelo, A.; Smith, J. B.; Sautebin,

[125845]L.; Galli, G.: Congenital deficiency of thromboxane and prostacyclin. Lancet I:

[125846]898-900, 1980.

[125847]22424.

[125848]Picot, D.; Loll, P. J.; Garavito, R. M.: The x-ray crystal structure

[125849]of the membrane protein prostaglandin H-2 synthase-1. Nature 367:

[125850]243–249, 1994.

[125851]22425.

[125852]Pieke, S. A.; Kimberling, W. J.; Kenyon, J. B.; Gabow, P.:

Genetic

[125853]heterogeneity of polycystic kidney disease: an estimate of
the proportion

[125854]of families unlinked to chromosome 16. (Abstract) Am. J.
Hum. Genet. 45

[125855](suppl.): A58 only, 1989.

[125856]22426.

[125857]Qian, F.; Germino, F. J.; Cai, Y.; Zhang, X.; Somlo, S.; Ger-
mino,

[125858]G. G.: PKD1 interacts with PKD2 through a probable
coiled–coil domain. Nature

[125859]Genet. 16: 179–183, 1997.

[125860]22427.

[125861]Ravine, D.; Walker, R. G.; Gibson, R. N.; Forrest, S. M.;
Richards,

[125862]R. I.; Friend, K.; Sheffield, L. J.; Kincaid–Smith, P.; Danks,
D.

[125863]M.: Phenotype and genotype heterogeneity in autosomal

dominant polycystic

[125864]kidney disease. Lancet 340: 1330–1333, 1992.

[125865]22428.

[125866]Reynolds, D. M.; Hayashi, T.; Cai, Y.; Veldhuisen, B.; Wat-
nick,

[125867]T. J.; Lens, X. M.; Mochizuki, T.; Qian, F.; Maeda, Y.; Li, L.;
Fossdal,

[125868]R.; Coto, E.; Wu, G.; Breuning, M. H.; Germino, G. G.; Pe-
ters, D.

[125869]J. M.; Somlo, S.: Aberrant splicing in the PKD2 gene as a
cause of

[125870]polycystic kidney disease. J. Am. Soc. Nephrol. 10:
2342–2351, 1999.

[125871]22429.

[125872]Romeo, G.; Devoto, M.; Costa, G.; Roncuzzi, L.; Catizone,
L.;

[125873]Zucchelli, P.; Germino, G.-G.; Keith, T.; Weatherall, D. J.;
Reeders,

[125874]S. T.: A second genetic locus for autosomal dominant
polycystic kidney

[125875]disease. Lancet II: 8–11, 1988.

[125876]22430.

[125877]San Millan, J. L.; Viribay, M.; Peral, B.; Martinez, I.; Weissenbach,
[125878]J.; Moreno, F.: Refining the localization of the PKD2 locus
on chromosome
[125879]4q by linkage analysis in Spanish families with autosomal
dominant
[125880]polycystic kidney disease type 2. Am. J. Hum. Genet. 56:
248-253,
[125881]1995.
[125882]22431.
[125883]Scheffers, M. S.; Le, H.; van der Bent, P.; Leonhard, W.;
Prins,
[125884]F.; Spruit, L.; Breuning, M. H.; de Heer, E.; Peters, D. J. M.:
Distinct
[125885]subcellular expression of endogenous polycystin-2 in the
plasma membrane
[125886]and Golgi apparatus of MDCK cells. Hum. Molec. Genet.
11: 59-67,
[125887]2002.
[125888]22432.
[125889]Schneider, M. C.; Rodriguez, A. M.; Nomura, H.; Zhou, J.;
Morton,

- [125890]C. C.; Reeders, S. T.; Weremowicz, S.: A gene similar to PKD1 maps to chromosome 4q22: a candidate gene for PKD2. *Genomics* 38: 1-4, 1996.
- [125891]22433.
- [125892]Torra, R.; Badenas, C.; San Millan, J. L.; Perez-Oller, L.; Estivill, X.; Darnell, A.: A loss-of-function model for cystogenesis in human autosomal dominant polycystic kidney disease type 2. *Am. J. Hum. Genet.* 65: 345-352, 1999.
- [125893]22434.
- [125894]Tsiokas, L.; Kim, E.; Arnould, T.; Sukhatme, V. P.; Walz, G.: Homo- and heterodimeric interactions between the gene products of PKD1 and PKD2. *Proc. Nat. Acad. Sci.* 94: 6965-6970, 1997.
- [125895]22435.
- [125896]Veldhuisen, B.; Saris, J. J.; de Haij, S.; Hayashi, T.;

Reynolds,

[125904]D. M.; Mochizuki, T.; Elles, R.; Fossdal, R.; Bogdanova, N.;
van Dijk,

[125905]M. A.; Coto, E.; Ravine, D.; Norby, S.; Verellen-Dumoulin,
C.; Breuning,

[125906]M. H.; Somlo, S.; Peters, D. J. M.: A spectrum of mutations
in the

[125907]second gene for autosomal dominant polycystic kidney
disease (PKD2). Am.

[125908]J. Hum. Genet. 61: 547-555, 1997.

[125909]22436.

[125910]Viribay, M.; Hayashi, T.; Telleria, D.; Mochizuki, T.;
Reynolds,

[125911]D. M.; Alonso, R.; Lens, X. M.; Moreno, F.; Harris, P. C.;
Somlo,

[125912]S.; San Millan, J. L.: Novel stop and frameshifting muta-
tions in

[125913]the autosomal dominant polycystic kidney disease 2
(PKD2) gene. Hum.

[125914]Genet. 101: 229-234, 1997.

[125915]22437.

[125916]Watnick, T.; He, N.; Wang, K.; Liang, Y.; Parfrey, P.; Heffer-

ton,

[125917]D.; St. George-Hyslop, P.; Germino, G.; Pei, Y.: Mutations
of PKD1

[125918]in ADPKD2 cysts suggest a pathogenic effect of trans-
heterozygous

[125919]mutations. Nature Genet. 25: 143–144, 2000.

[125920]22438.

[125921]Wu, G.; D'Agati, V.; Cai, Y.; Markowitz, G.; Park, J. H.;
Reynolds,

[125922]D. M.; Maeda, Y.; Le, T. C.; Hou, H., Jr.; Kucherlapati, R.;
Edelmann,

[125923]W.; Somlo, S.: Somatic inactivation of Pkd2 results in poly-
cystic

[125924]kidney disease. Cell 98: 177–188, 1998.

[125925]22439.

[125926]Wu, G.; Markowitz, G. S.; Li, L.; D'Agati, V. D.; Factor, S.
M.;

[125927]Geng, L.; Tibara, S.; Tuchman, J.; Cai, Y.; Park, J. H.; van
Adelsberg,

[125928]J.; Hou, H., Jr.; Kucherlapati, R.; Edelmann, W.; Somlo, S.:
Cardiac

[125929]defects and renal failure in mice with targeted mutations

in Pkd2. Nature

[125930]Genet. 24: 75–78, 2000.

[125931]22440.

[125932]Wu, G.; Mochizuki, T.; Le, T. C.; Cai, Y.; Hayashi, T.;
Reynolds,

[125933]D. M.; Somlo, S.: Molecular cloning, cDNA sequence analysis, and

[125934]chromosomal localization of mouse Pkd2. Genomics 45:
220–223, 1997.

[125935]22441.

[125936]Xenophontos, S.; Constantinides, R.; Hayashi, T.;
Mochizuki, T.;

[125937]Somlo, S.; Pierides, A.; Constantinou Deltas, C.: A transition frameshift

[125938]mutation induced by a cytosine insertion in the polycystic
kidney

[125939]disease 2 gene (PKD2). Hum. Molec. Genet. 6: 949–952,
1997.

[125940]22442.

[125941]Zerres, K.; Mucher, G.; Rudnik–Schoneborn, S.: Autosomal
recessive

[125942]polycystic kidney disease does not map to the second

gene locus for

[125943]autosomal dominant polycystic kidney disease on chromosome 4. Hum.

[125944]Genet. 93: 697–698, 1994.

[125945]22443.

[125946]Zerres, K.; Volpel, M. C.; Weiss, H.: Cystic kidneys: genetics,

[125947]pathologic anatomy, clinical picture, and prenatal diagnosis. Hum.

[125948]Genet. 68: 104–135, 1984.

[125949]22444.

[125950]Ala-Mello, S.; Koskimies, O.; Rapola, J.; Kaariainen, H.:
Nephronophthisis

[125951]in Finland: epidemiology and comparison of genetically
classified

[125952]subgroups. Europ. J. Hum. Genet. 7: 205–211, 1999.

[125953]22445.

[125954]Chamberlin, B. C.; Hagge, W. W.; Stickler, G. B.: Juvenile
nephronophthisis

[125955]and medullary cystic disease. Mayo Clin. Proc. 52:
485–491, 1977.

[125956]22446.

[125957]Hussels-Maumenee, I.; Pierce, E. R.; Bias, W. B.; Schleuter-mann,

[125958]D. A.: Linkage studies of typical retinitis pigmentosa and common

[125959]markers. Am. J. Hum. Genet. 27: 505-508, 1975.

[125960]22447.

[125961]Olsson, J. E.; Samanns, C.; Jimenez, J.; Pongratz, J.; Chand,

[125962]A.; Watty, A.; Seuchter, S. A.; Denton, M.; Gal, A.: Gene of type

[125963]II autosomal dominant retinitis pigmentosa maps on the long arm of

[125964]chromosome 3. Am. J. Med. Genet. 35: 595-599, 1990.

[125965]22448.

[125966]Pierce, E. A.; Quinn, T.; Meehan, T.; McGee, T. L.; Berson, E.

[125967]L.; Dryja, T. P.: Mutations in a gene encoding a new oxygen-regulated

[125968]photoreceptor protein cause dominant retinitis pigmentosa. Nature

[125969]Genet. 22: 248-254, 1999.

[125970]22449.

[125971]Sullivan, L. S.; Heckenlively, J. R.; Bowne, S. J.; Zuo, J.;
Hide,

[125972]W. A.; Gal, A.; Denton, M.; Inglehearn, C. F.; Blanton, S. H.;
Daiger,

[125973]S. P.: Mutations in a novel retina-specific gene cause au-
tosomal

[125974]dominant retinitis pigmentosa. Nature Genet. 22:
255-259, 1999.

[125975]22450.

[125976]Inglehearn, C.; Farrar, J.; Denton, M.; Gal, A.; Humphries,
P.;

[125977]Bhattacharya, S.: Evidence against a second autosomal
dominant retinitis

[125978]pigmentosa locus close to rhodopsin on chromosome 3q.
(Letter) Am.

[125979]J. Hum. Genet. 53: 536-537, 1993.

[125980]22451.

[125981]McInnes, R. R.; Bascom, R. A.: Retinal genetics: a nullifying

[125982]effect for rhodopsin. Nature Genet. 1: 155-157, 1992.

[125983]22452.

- [125984]McWilliam, P.; Farrar, G. J.; Kenna, P.; Bradley, D. G.;
Humphries,
[125985]M. M.; Sharp, E. M.; McConnell, D. J.; Lawler, M.; Sheils, D.;
Ryan,
[125986]C.; Stevens, K.; Daiger, S. P.; Humphries, P.: Autosomal
dominant
[125987]retinitis pigmentosa (ADRP): localization of an ADRP gene
to the long
[125988]arm of chromosome 3. Genomics 5: 619–622, 1989.
[125989]22453.
[125990]Chautard–Freire–Maia, E. A.: Probable assignment of the
serum
[125991]cholinesterase (E1) and transferrin (Tf) loci to chromo-
some 1 in man. Hum.
[125992]Hered. 27: 134–142, 1977.
[125993]22454.
[125994]Chautard–Freire–Maia, E. A.: Probable assignment of the
E1 and
[125995]Tf loci to chromosome 1 in man. Ciencia e Cultura (Brazil)
28 (suppl.):
[125996]309–310, 1976.

[125997]22455.

[125998]Niswender, K. D.; Morton, G. J.; Stearns, W. H.; Rhodes, C. J.;

[125999]Myers, M. G., Jr.; Schwartz, M. W.: Key enzyme in leptin-induced

[126000]anorexia. Nature 413: 794-795, 2001.

[126001]22456.

[126002]Ogawa, Y.; Masuzaki, H.; Isse, N.; Okazaki, T.; Mori, K.; Shigemoto,

[126003]M.; Satoh, N.; Tamura, N.; Hosoda, K.; Yoshimasa, Y.; Jingami, H.;

[126004]Kawada, T.; Nakao, K.: Molecular cloning of rat obese cDNA and augmented

[126005]gene expression in genetically obese Zucker fatty (fa/fa) rats. J.

[126006]Clin. Invest. 96: 1647-1652, 1995.

[126007]22457.

[126008]Pasco, J. A.; Henry, M. J.; Kotowicz, M. A.; Collier, G. R.; Ball,

[126009]M. J.; Ugoni, A. M.; Nicholson, G. C.: Serum leptin levels are associated

- [126010] with bone mass in nonobese women. J. Clin. Endocr. Metab. 86: 1884–1887, 2001.
- [126011] 22458.
- [126012] Pellemounter, M. A.; Cullen, M. J.; Baker, M. B.; Hecht, R.; Winters, D.; Boone, T.; Collins, F.: Effects of the obese gene product on body weight regulation in ob/ob mice. Science 269: 540–542, 1995.
- [126013] 22459.
- [126014] Rink, T. J.: In search of a satiety factor. Nature 372: 408–409, 1994.
- [126015] 22460.
- [126016] Rosenbaum, M.; Leibel, R. L.: The role of leptin in human physiology. (Editorial) New Eng. J. Med. 341: 913–915, 1999.
- [126017] 22461.
- [126018] Saad, M. F.; Damani, S.; Gingerich, R. L.; Riad-Gabriel, M. G.;

[126024]Khan, A.; Boyadjian, R.; Jinagouda, S. D.; El-Tawil, K.;
Rude, R.

[126025]K.; Kamdar, V.: Sexual dimorphism in plasma leptin concentration. J.

[126026]Clin. Endocr. Metab. 82: 579–584, 1997.

[126027]22462.

[126028]Schubring, C.; Kiess, W.; Englaro, P.; Rascher, W.; Dotsch, J.;

[126029]Hanitsch, S.; Attanasio, A.; Blum, W. F.: Levels of leptin in maternal

[126030]serum, amniotic fluid, and arterial and venous cord blood: relation

[126031]to neonatal and placental weight. J. Clin. Endocr. Metab. 82: 1480–1483,

[126032]1997.

[126033]22463.

[126034]Shanley, L. J.; Irving, A. J.; Rae, M. G.; Ashford, M. L. J.;

[126035]Harvey, J.: Leptin inhibits rat hippocampal neurons via activation

[126036]of large conductance calcium-activated K(+) channels. Nature Neurosci. 5:

[126037]299–300, 2002.

[126038]22464.

[126039]Shimomura, I.; Hammer, R. E.; Ikemoto, S.; Brown, M. S.;
Goldstein,

[126040]J. L.: Leptin reverses insulin resistance and diabetes mellitus in

[126041]mice with congenital lipodystrophy. *Nature* 401: 73–76,
1999.

[126042]22465.

[126043]Shimomura, I.; Matsuda, M.; Hammer, R. E.; Bashmakov,
Y.; Brown,

[126044]M. S.; Goldstein, J. L.: Decreased IRS–2 and increased
SREBP–1c lead

[126045]to mixed insulin resistance and sensitivity in livers of
lipodystrophic

[126046]and ob/ob mice. *Molec. Cell* 6: 77–86, 2000.

[126047]22466.

[126048]Sierra–Honigsmann, M. R.; Nath, A. K.; Murakami, C.; Gar-
cia–Cardena,

[126049]G.; Papapetropoulos, A.; Sessa, W. C.; Madge, L. A.;
Schechner, J.

[126050]S.; Schwabb, M. B.; Polverini, P. J.; Flores-Riveros, J. R.: Biological

[126051]action of leptin as an angiogenic factor. Science 281: 1583–1586,

[126052]1998.

[126053]22467.

[126054]Sooranna, S. R.; Ward, S.; Bajoria, R.: Fetal leptin influences

[126055]birth weight in twins with discordant growth. Pediat. Res. 49: 667–672,

[126056]2001.

[126057]22468.

[126058]Stattin, P.; Soderberg, S.; Hallmans, G.; Bylund, A.; Kaaks, R.;

[126059]Stenman, U.-H.; Bergh, A.; Olsson, T.: Leptin is associated with

[126060]increased prostate cancer risk: a nested case-referent study. J.

[126061]Clin. Endocr. Metab. 86: 1341–1345, 2001.

[126062]22469.

[126063]Strobel, A.; Issad, T.; Camoin, L.; Ozata, M.; Strosberg, A.

D.

[126064]: A leptin missense mutation associated with hypogonadism and morbid

[126065]obesity. (Letter) Nature Genet. 18: 213–215, 1998.

[126066]22470.

[126067]Szczypka, M. S.; Rainey, M. A.; Palmiter, R. D.: Dopamine is

[126068]required for hyperphagia in Lep(ob/ob) mice. Nature Genet. 25: 102–104,

[126069]2000.

[126070]22471.

[126071]Trayhurn, P.; Thurlby, P. L.; James, W. P. T.: Thermogenic defect

[126072]in pre-obese ob/ob mice. (Letter) Nature 266: 60–62, 1977.

[126073]22472.

[126074]Verploegen, S. A. B. W.; Plaetinck, G.; Devos, R.; Van der Heyden,

[126075]J.; Guisez, Y.: A human leptin mutant induces weight gain in normal

[126076]mice. FEBS Lett. 405: 237–240, 1997.

[126077]22473.

[126078]Weigle, D. S.; Ganter, S. L.; Kuijper, J. L.; Leonetti, D. L.;

[126079]Boyko, E. J.; Fujimoto, W. Y.: Effect of regional fat distribution

[126080]and Prader–Willi syndrome on plasma leptin levels. J. Clin. Endocr.

[126081]Metab. 82: 566–570, 1997.

[126082]22474.

[126083]Wellhoener, P.; Fruehwald–Schultes, B.; Kern, W.; Dantz, D.; Kerner,

[126084]W.; Born, J.; Fehm, H. L.; Peters, A.: Glucose metabolism rather

[126085]than insulin is a main determinant of leptin secretion in humans. J.

[126086]Clin. Endocr. Metab. 85: 1267–1271, 2000.

[126087]22475.

[126088]Zhang, Y.; Proenca, R.; Maffei, M.; Barone, M.; Leopold, L.; Friedman,

[126089]J. M.: Positional cloning of the mouse obese gene and its human homologue. Nature 372:

[126090]425–432, 1994.

[126091]22476.

[126092]Zhao, A. Z.; Huan, J.-N.; Gupta, S.; Pal, R.; Sahu, A.: A
phosphatidylinositol

[126093]3-kinase-phosphodiesterase 3B-cyclic AMP pathway in
hypothalamic action

[126094]of leptin on feeding. *Nature Neurosci.* 5: 727-728, 2002.

[126095]22477.

[126096]Siracusa, L. D.; Rosner, M. H.; Vigano, M. A.; Gilbert, D. J.;

[126097]Staudt, L. M.; Copeland, N. G.; Jenkins, N. A.: Chromoso-
mal location

[126098]of the octamer transcription factors, Otf-1, Otf-2, and
Otf-3, defines

[126099]multiple Otf-3-related sequences dispersed in the mouse
genome. *Genomics* 10:

[126100]313-326, 1991.

[126101]22478.

[126102]Sturm, R. A.; Eyre, H. J.; Baker, E.; Sutherland, G. R.: The
human

[126103]OTF1 locus which overlaps the CD3Z gene is located at
1q22-q23. *Cytogenet.*

[126104]*Cell Genet.* 68: 231-232, 1995.

[126105]22479.

[126106]Marshall, J. B.; Diaz-Arias, A. A.; Bochna, G. S.; Vogele, K.

[126107]A.: Achalasia due to diffuse esophageal leiomyomatosis
and inherited

[126108]as an autosomal dominant disorder. Gastroenterology 98:
1358-1365,

[126109]1990.

[126110]22480.

[126111]Mattei, M. G.; d'Auriol, L.; Andre, C.; Passage, E.; Mattei, J.

[126112]F.; Galibert, F.: Assignment of the human c-kit proto-
oncogene to

[126113]the q11-q12 region of chromosome 4, using in situ hy-
bridization. (Abstract) Cytogenet.

[126114]Cell Genet. 46: 657, 1987.

[126115]22481.

[126116]Nagata, H.; Worobec, A. S.; Oh, C. K.; Chowdhury, B. A.;
Tannenbaum,

[126117]S.; Suzuki, Y.; Metcalfe, D. D.: Identification of a point mu-
tation

[126118]in the catalytic domain of the protooncogene c-kit in pe-
ripheral blood

[126119]mononuclear cells of patients who have mastocytosis with
an associated

[126120]hematologic disorder. Proc. Nat. Acad. Sci. 92:
10560–10564, 1995.

[126121]22482.

[126122]Nishida, T.; Hirota, S.; Taniguchi, M.; Hashimoto, K.;
Isozaki,

[126123]K.; Nakamura, H.; Kanakura, Y.; Tanaka, T.; Takabayashi,
A.; Matsuda,

[126124]H.; Kitamura, Y.: Familial gastrointestinal stromal tumours
with

[126125]germline mutation of the KIT gene. (Letter) Nature Genet.
19: 323–324,

[126126]1998.

[126127]22483.

[126128]Nocka, K.; Tan, J. C.; Chiu, E.; Chu, T. Y.; Ray, P.; Trakt-
man,

[126129]P.; Besmer, P.: Molecular bases of dominant negative and
loss of

[126130]function mutations at the murine c-kit/white spotting lo-
cus: W-37,

[126131]W-v, W-41 and W. EMBO J. 9: 1805–1813, 1990.

[126132]22484.

[126133]Nomura, K.; Hatayama, I.; Narita, T.; Kaneko, T.; Shiraishi, M.

[126134]: A novel KIT gene missense mutation in a Japanese family with piebaldism.

[126135](Letter) J. Invest. Derm. 111: 337–338, 1998.

[126136]22485.

[126137]Pignon, J.–M.; Giraudier, S.; Duquesnoy, P.; Jouault, H.; Imbert,

[126138]M.; Vainchenker, W.; Vernant, J.–P.; Tulliez, M.: A new c-kit mutation

[126139]in a case of aggressive mast cell disease. Brit. J. Haemat. 96:

[126140]374–376, 1997.

[126141]22486.

[126142]Reinsch, N.; Thomsen, H.; Xu, N.; Brink, M.; Looft, C.; Kalm,

[126143]E.; Brockmann, G. A.; Grupe, S.; Kuhn, C.; Schwerin, M.; Leyhe, B.;

[126144]Hiendleder, S.; Erhardt, G.; Medjugorac, I.; Russ, I.; Forster, M.;

[126145]Reents, R.; Averdunk, G.: A QTL for the degree of spotting
in cattle

[126146]shows synteny with the KIT locus on chromosome 6. J.
Hered. 90:

[126147]629–634, 1999.

[126148]22487.

[126149]Selmanowitz, V. J.; Rabinowitz, A. D.; Orentreich, N.;
Wenk, E.

[126150]: Pigmentary correction of piebaldism by autografts. I.
Procedures

[126151]and clinical findings. J. Derm. Surg. Oncol. 3: 615–622,
1977.

[126152]22488.

[126153]Spritz, R. A.; Beighton, P.: Piebaldism with deafness:
molecular

[126154]evidence for an expanded syndrome. Am. J. Med. Genet.
75: 101–103,

[126155]1998.

[126156]22489.

[126157]Spritz, R. A.; Droetto, S.; Fukushima, Y.: Deletion of the
KIT

[126158]and PDGFRA genes in a patient with piebaldism. Am. J.

Med. Genet. 44:

[126159]492–495, 1992.

[126160]22490.

[126161]Spritz, R. A.; Giebel, L. B.: Mutation of the c-kit

(mast/stem

[126162]cell growth factor receptor) proto-oncogene in human
piebaldism. (Abstract) Am.

[126163]J. Hum. Genet. 49 (suppl.): 38, 1991.

[126164]22491.

[126165]Spritz, R. A.; Giebel, L. B.; Holmes, S. A.: Dominant nega-
tive

[126166]and loss of function mutations of the c-kit (mast/stem
cell growth

[126167]factor receptor) proto-oncogene in human piebaldism.
Am. J. Hum.

[126168]Genet. 50: 261–269, 1992.

[126169]22492.

[126170]Spritz, R. A.; Holmes, S. A.; Ramesar, R.; Greenberg, J.;
Curtis,

[126171]D.; Beighton, P.: Mutations of the KIT (mast/stem cell

growth factor

[126172]receptor) proto-oncogene account for a continuous range
of phenotypes

[126173]in human piebaldism. Am. J. Hum. Genet. 51: 1058–1065,
1992.

[126174]22493.

[126175]Syrris, P.; Malik, N. M.; Murday, V. A.; Patton, M. A.; Carter,

[126176]N. D.; Hughes, H. E.; Metcalfe, K.: Three novel mutations
of the

[126177]proto-oncogene KIT cause human piebaldism. (Letter) Am.
J. Med. Genet. 95:

[126178]79–81, 2000.

[126179]22494.

[126180]Tan, J. C.; Nocka, K.; Ray, P.; Traktman, P.; Besmer, P.: The

[126181]dominant W42 spotting phenotype results from a mis-
sense mutation in

[126182]the c-kit receptor kinase. Science 247: 209–212, 1990.

[126183]22495.

[126184]Taylor, M. L.; Dastych, J.; Sehgal, D.; Sundstrom, M.; Nils-
son,

[126185]G.; Akin, C.; Mage, R. G.; Metcalfe, D. D.: The Kit-

activating mutation

[126186]D816V enhances stem cell factor-dependent chemotaxis.

Blood 98:

[126187]1195–1199, 2001.

[126188]22496.

[126189]Thomsen, L.; Robinson, T. L.; Lee, J. C. F.; Farraway, L. A.;

[126190]Hughes, M. J. G.; Andrews, D. W.; Huizinga, J. D.: Interstitial cells
tial cells

[126191]of Cajal generate a rhythmic pacemaker current. Nature
Med. 4: 848–851,

[126192]1998.

[126193]22497.

[126194]Tian, Q.; Frierson, H. F., Jr.; Krystal, G. W.; Moskaluk, C. A.

[126195]: Activating c-kit gene mutations in human germ cell tumors. Am.

[126196]J. Path. 154: 1643–1647, 1999.

[126197]22498.

[126198]Tsujimura, T.; Furitsu, T.; Morimoto, M.; Isozaki, K.; Nomura,

[126199]S.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Ligand-

independent

[126200]activation of c-kit receptor tyrosine kinase in a murine mastocytoma

[126201]cell line P-815 generated by a point mutation. Blood 83: 2619-2626,

[126202]1994.

[126203]22499.

[126204]Vandenbark, G. R.; deCastro, C. M.; Taylor, H.; Dew-Knight, S.;

[126205]Kaufman, R. E.: Cloning and structural analysis of the human c-kit

[126206]gene. Oncogene 7: 1259-1266, 1992.

[126207]22500.

[126208]Worobec, A. S.; Semere, T.; Nagata, H.; Metcalfe, D. D.: Clinical

[126209]correlates of the presence of the asp816val c-kit mutation in the

[126210]peripheral blood mononuclear cells of patients with mastocytosis. Cancer 83:

[126211]2120-2129, 1998.

[126212]22501.

[126213]Yarden, Y.; Kuang, W.-J.; Yang-Feng, T.; Coussens, L.;
Munemitsu,

[126214]S.; Dull, T. J.; Chen, E.; Schlessinger, J.; Francke, U.; Ull-
rich,

[126215]A.: Human proto-oncogene c-kit: a new cell surface re-
ceptor tyrosine

[126216]kinase for an unidentified ligand. EMBO J. 6: 3341-3351,
1987.

[126217]22502.

[126218]Cheah, M. S. C.; Ley, T. J.; Tronick, S. R.; Robbins, K. C.:
Fgr

[126219]proto-oncogene mRNA induced in B lymphocytes by Ep-
stein-Barr virus

[126220]infection. Nature 319: 238-240, 1986.

[126221]22503.

[126222]Dracopoli, N. C.; Stanger, B. Z.; Lager, M.; Housman, D. E.:
Localization

[126223]of the FGR protooncogene on the genetic linkage map of
human chromosome

[126224]1p. Genomics 3: 124-128, 1988.

[126225]22504.

[126226]Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J.
D.:

[126227]Evidence for two distinct c-src loci on human chromo-
somes 1 and 20. Nature 312:

[126228]70-71, 1984.

[126229]22505.

[126230]Lebo, R. V.; Cheung, M.-C.; Bruce, B. D.: Rapid gene map-
ping by

[126231]dual laser chromosome sorting and spot blot DNA analy-
sis. (Abstract) Am.

[126232]J. Hum. Genet. 36: 101S, 1984.

[126233]22506.

[126234]Nishizawa, M.; Semba, K.; Yoshida, M. C.; Yamamoto, T.;
Sasaki,

[126235]M.; Toyoshima, K.: Structure, expression, and chromoso-
mal location

[126236]of the human c-fgr gene. Molec. Cell. Biol. 6: 511-517,
1986.

[126237]22507.

[126238]Kaiser, U. B.; Halvorsen, L. M.; Chen, M. T.: SP1, steroido-
genic

- [126239] factor 1 (SF-1), and early growth response protein 1 (Egr-1) binding
- [126240] sites form a tripartite gonadotropin-releasing hormone response element
- [126241] in the rat luteinizing hormone-beta gene promoter: an integral role
- [126242] for SF-1. *Molec. Endocr.* 14: 1235-1245, 2000.
- [126243] 22508.
- [126244] Liao, W. X.; Goh, H. H.; Roy, A. C.: Functional characterization
- [126245] of a natural variant of luteinizing hormone. *Hum. Genet.* 111: 219-224,
- [126246] 2002.
- [126247] 22509.
- [126248] Liao, W. X.; Roy, a. C.; Chan, C.; Arulkumaran, S.; Ratnam, S.
- [126249] S.: A new molecular variant of luteinizing hormone associated with
- [126250] female infertility. *Fertil. Steril.* 69: 102-106, 1998.
- [126251] 22510.
- [126252] Manna, P. R.; Joshi, L.; Reinhold, V. N.; Aubert, M. L.; Sug-

anuma,

[126253]N.; Pettersson, K.; Huhtaniemi, I. T.: Synthesis, purification
and

[126254]structural and functional characterization of recombinant
form of

[126255]a common genetic variant of human luteinizing hormone.
Hum. Molec.

[126256]Genet. 11: 301–315, 2002.

[126257]22511.

[126258]Mohrenweiser, H. W.; Tynan, K. M.; Branscomb, E. W.; de
Jong,

[126259]P. J.; Olsen, A.; Trask, B.; Carrano, A. V.: Development of
an integrated

[126260]genetic, functional and physical map of human chromo-
some 19. (Abstract) Cytogenet.

[126261]Cell Genet. 58: 2021 only, 1991.

[126262]22512.

[126263]Muyan, M.; Furuhashi, M.; Sugahara, T.; Boime, I.: The car-
boxy-terminal

[126264]region of the beta-subunits of luteinizing hormone and
chorionic gonadotropin

[126265]differentially influence secretion and assembly of the het-

erodimers. Molec.

[126266]Endocr. 10: 1678–1687, 1996.

[126267]22513.

[126268]Park, I. J.; Burnett, L. S.; Jones, H. W., Jr.; Migeon, C. J.;

[126269]Blizzard, R. M.: A case of male pseudohermaphroditism
associated

[126270]with elevated LH, normal FSH and low testosterone possibly due to

[126271]the secretion of an abnormal LH molecule. Acta Endocr.
83: 173–181,

[126272]1976.

[126273]22514.

[126274]Pettersson, K.; Ding, Y.-Q.; Huhtaniemi, I.: An immunologically

[126275]anomalous luteinizing hormone variant in a healthy
woman. J. Clin.

[126276]Endocr. Metab. 74: 164–171, 1992.

[126277]22515.

[126278]Quirk, C. C.; Lozada, K. L.; Keri, R. A.; Nilson, J. H.: A single

[126279]Pitx1 binding site is essential for activity of the LH-beta

promoter

[126280]in transgenic mice. *Molec. Endocr.* 15: 734–746, 2001.

[126281]22516.

[126282]Raivio, T.; Huhtaniemi, I.; Anttila, R.; Siimes, M. A.; Hage-
nas,

[126283]L.; Nilsson, C.; Pettersson, K.; Dunkel, L.: The role of
luteinizing

[126284]hormone-beta gene polymorphism in the onset and pro-
gression of puberty

[126285]in healthy boys. *J. Clin. Endocr. Metab.* 81: 3278–3282,
1996.

[126286]22517.

[126287]Ramanujam, L. N.; Liao, W. X.; Roy, A. C.; Loganath, A.;
Goh,

[126288]H. H.; Ng, S. C.: Association of molecular variants of
luteinizing

[126289]hormone with menstrual disorders. *Clin. Endocrin.* 51:
243–246, 1999.

[126290]22518.

[126291]Ramanujam, L. N.; Liao, W. X.; Roy, A. C.; Ng, S. C.: Asso-
ciation

[126292]of molecular variants of luteinizing hormone with male in-

fertility. Hum.

[126293]Reprod. 15: 925–928, 2000.

[126294]22519.

[126295]Roy, A. C.; Liao, W. X.; Chen, Y.; Arulkumaran, S.; Ratnam, S.

[126296]S.: Identification of 7 novel mutations in LH beta-subunit gene by

[126297]SSCP. Molec. Cell Biochem. 165: 151–153, 1996.

[126298]22520.

[126299]Shome, B.; Parlow, A. F.: The primary structure of the hormone-specific,

[126300]beta subunit of human pituitary luteinizing hormone (hLH). J. Clin.

[126301]Endocr. 36: 618–621, 1973.

[126302]22521.

[126303]Tapanainen, J. S.; Koivunen, R.; Fauser, B. C. J. M.; Taylor,

[126304]A. E.; Clayton, R. N.; Rajkova, M.; White, D.; Franks, S.; Anttila,

[126305]L.; Pettersson, K. S. I.; Huhtaniemi, I. T.: A new contributing factor

[126306]to polycystic ovary syndrome: the genetic variant of

luteinizing hormone. J.

[126307]Clin. Endocr. Metab. 84: 1711–1715, 1999.

[126308]22522.

[126309]van den Beld, A. W.; Huhtaniemi, I. T.; Pettersson, K. S. L.;

[126310]Pols, H. A. P.; Grobbee, D. E.; de Jong, F. H.; Lamberts, S.
W. J.

[126311]: Luteinizing hormone and different genetic variants, as
indicators

[126312]of frailty in healthy elderly men. J. Clin. Endocr. Metab. 84:
1334–1339,

[126313]1999.

[126314]22523.

[126315]Weck, J.; Anderson, A. C.; Jenkins, S.; Fallest, P. C.; Shup-
nik,

[126316]M. A.: Divergent, composite gonadotropin–releasing hor-
mone–responsive

[126317]elements in the rat luteinizing hormone subunit genes.
Molec. Endocr. 14:

[126318]472–485, 2000.

[126319]22524.

[126320]Weiss, J.; Axelrod, L.; Whitcomb, R. W.; Harris, P. E.; Crow-

ley,

[126321]W. F.; Jameson, J. L.: Hypogonadism caused by a single amino acid

[126322]substitution in the beta subunit of luteinizing hormone.
New Eng.

[126323]J. Med. 326: 179–183, 1992.

[126324]22525.

[126325]Atger, M.; Misrahi, M.; Sar, S.; Le Flem, L.; Dessen, P.; Milgrom,

[126326]E.: Structure of the human luteinizing hormone–
choriogonadotropin

[126327]receptor gene: unusual promoter and 5–prime non–
coding regions. Molec.

[126328]Cell. Endocr. 111: 113–123, 1995.

[126329]22526.

[126330]Bardin, C. W.; Bullock, I. P.; Sherins, R. J.; Mowszowicz, I.;

[126331]Blackburn, W. R.: Androgen metabolism and mechanism of
action in

[126332]male pseudohermaphroditism: a study of testicular feminization. Recent

[126333]Prog. Horm. Res. 29: 65–109, 1973.

[126334]22527.

[126335]Berthezene, F.; Forest, M. G.; Grimaud, J. A.; Claustrat, B.;
Mornex,

[126336]R.: Leydig-cell agenesis: a cause of male pseudo-
hermaphroditism. New

[126337]Eng. J. Med. 295: 969-972, 1976.

[126338]22528.

[126339]Brown, D. M.; Markland, C.; Dehner, L. P.: Leydig cell hy-
poplasia--a

[126340]cause of male pseudohermaphroditism. J. Clin. Endocr.
Metab. 46:

[126341]1-7, 1978.

[126342]22529.

[126343]Canto, P.; Soderlund, D.; Ramon, G.; Nishimura, E.;
Mendez, J.

[126344]P.: Mutational analysis of the luteinizing hormone receptor
gene

[126345]in two individuals with Leydig cell tumors. Am. J. Med.
Genet. 108:

[126346]148-152, 2002.

[126347]22530.

[126348]Eblen, A.; Bao, S.; Lei, Z. M.; Nakajima, S. T.; Rao, C. V.:

The

[126349]presence of functional luteinizing hormone/chorionic gonadotropin

[126350]receptors in human sperm. J. Clin. Endocr. Metab. 86: 2643–2648,

[126351]2001.

[126352]22531.

[126353]El-Awady, M. K.; Temtamy, S. A.; Salam, M. A.; Gad, Y. Z.:

Familial

[126354]Leydig cell hypoplasia as a cause of male pseudohermaphroditism. Hum.

[126355]Hered. 37: 36–40, 1987.

[126356]22532.

[126357]Evans, B. A. J.; Bowen, D. J.; Smith, P. J.; Clayton, P. E.;

Gregory,

[126358]J. W.: A new point mutation in the luteinising hormone receptor gene

[126359]in familial and sporadic male limited precocious puberty: genotype

[126360]does not always correlate with phenotype. J. Med. Genet. 33: 143–147,

[126361]1996.

[126362]22533.

[126363]Gospodarowicz, D.: Properties of the luteinizing hormone
receptor

[126364]of isolated bovine corpus luteum plasma membrane. J.
Biol. Chem. 248:

[126365]5042–5049, 1973.

[126366]22534.

[126367]Gromoll, J.; Eiholzer, U.; Nieschlag, E.; Simoni, M.: Male
hypogonadism

[126368]caused by homozygous deletion of exon 10 of the
luteinizing hormone

[126369](LH) receptor: differential action of human chorionic go-
nadotropin

[126370]and LH. J. Clin. Endocr. Metab. 85: 2281–2286, 2000.

[126371]22535.

- [126372]Robert, B.; Barton, P.; Minty, A.; Daubas, P.; Weydert, A.;
Bonhomme,
[126373]F.; Catalan, J.; Chazottes, D.; Guenet, J. L.; Gros, F.; Buck-
ingham,
[126374]M. E.: Investigation of genetic linkage between myosin and
actin
[126375]genes using an interspecific mouse back-cross. Nature
314: 181-183,
[126376]1985.
[126377]22536.
[126378]Saez, L. J.; Gianola, K. M.; McNally, E. M.; Feghali, R.; Eddy,
[126379]R.; Shows, T. B.; Leinwand, L. A.: Human cardiac myosin
heavy chain
[126380]genes and their linkage in the genome. Nucleic Acids Res.
15: 5443-5459,
[126381]1987.
[126382]22537.
[126383]Ye, K.; Aghdasi, B.; Luo, H. R.; Moriarity, J. L.; Wu, F. Y.;
Hong,
[126384]J. J.; Hurt, K. J.; Bae, S. S.; Suh, P.-G.; Snyder, S. H.: Phos-
pholipase

[126385]C-gamma-1 is a physiological guanine nucleotide exchange factor for

[126386]the nuclear GTPase PIKE. Nature 415: 541-544, 2002.

[126387]22538.

[126388]Lazarou, L. P.; Davies, F.; Sarfarazi, M.; Coles, G. A.; Harper,

[126389]P. S.: Adult polycystic kidney disease and linked RFLPs at the alpha-globin

[126390]locus: a genetic study in the South Wales population. J. Med. Genet. 24:

[126391]466-473, 1987.

[126392]22539.

[126393]Burn, J.; Chapman, P.; Delhanty, J.; Wood, C.; Laloo, F.; Cachon-Gonzalez,

[126394]M. B.; Tsioupra, K.; Church, W.; Rhodes, M.; Gunn, A.: The UK northern

[126395]region genetic register for familial adenomatous polyposis coli: use

[126396]of age of onset, congenital hypertrophy of the retinal pigment epithelium,

[126397]and DNA markers in risk calculations. J. Med. Genet. 28: 289-296,

[126398]1991.

[126399]22540.

[126400]Tsuji, M.; Kawano, S.; Tsuji, S.; Sawaoka, H.; Hori, M.;
DuBois,

[126401]R. N.: Cyclooxygenase regulates angiogenesis induced by
colon cancer

[126402]cells. Cell 93: 705–716, 1998.

[126403]22541.

[126404]Vane, J. R.; Mitchell, J. A.; Appleton, I.; Tomlinson, A.;
Bishop-Bailey,

[126405]D.; Croxtall, J.; Willoughby, D. A.: Inducible isoforms of
cyclooxygenase

[126406]and nitric-oxide synthase in inflammation. Proc. Nat.
Acad. Sci. 91:

[126407]2046–2050, 1994.

[126408]22542.

[126409]Yokoyama, C.; Tanabe, T.: Cloning of human gene encod-
ing prostaglandin

[126410]endoperoxide synthase and primary structure of the en-
zyme. Biochem.

[126411]Biophys. Res. Commun. 165: 888–894, 1989.

[126412]22543.

[126413]Schwarz, H. P.; Heeb, M. J.; Lottenberg, R.; Roberts, H.;
Griffin,

[126414]J. H.: Familial protein S deficiency with a variant protein S
molecule

[126415]in plasma and platelets. Blood 74: 213–221, 1989.

[126416]22544.

[126417]Stanislovitis, P.; Watkins, P. C.; Eddy, R.; Cohen, E. H.;
Dackowski,

[126418]W. R.; Wydro, R. M.; Shows, T. B.: Regional assignment of
the gene

[126419]for protein S to the human chromosome region
3p21–3q21. (Abstract) Am.

[126420]J. Hum. Genet. 41: A187 only, 1987.

[126421]22545.

[126422]Watkins, P. C.; Eddy, R.; Fukushima, Y.; Byers, M. G.; Co-
hen,

[126423]E. H.; Dackowski, W. R.; Wydro, R. M.; Shows, T. B.: The
gene for

[126424]protein S maps near the centromere of human chromo-
some 3. Blood 71:

[126425]238-241, 1988.

[126426]22546.

[126427]Watkins, P. C.; Eddy, R.; Fukushima, Y.; Byers, M. G.; Cohen,

[126428]E. H.; Dackowski, W. R.; Wydro, R. M.; Shows, T. B.: Regional assignment

[126429]of the gene for protein S (PROS) to the human chromosome region 3p11.1-q11.2.

[126430](Abstract) Cytogenet. Cell Genet. 46: 712 only, 1987.

[126431]22547.

[126432]Zoller, B.; Garcia de Frutos, P.; Dahlback, B.: Evaluation of

[126433]the relationship between protein S and C4b-binding protein isoforms

[126434]in hereditary protein S deficiency demonstrating type I and type III

[126435]deficiencies to be phenotypic variants of the same genetic disease. Blood 85:

[126436]3524-3531, 1995.

[126437]22548.

[126438]Kaplan, R.; Morse, B.; Huebner, K.; Croce, C.; Howk, R.; Ravera,

- [126439]M.; Ricca, G.; Jaye, M.; Schlessinger, J.: Cloning of three human
- [126440]tyrosine phosphatases reveals a multigene family of receptor-linked
- [126441]protein-tyrosine-phosphatases expressed in brain. Proc. Nat. Acad.
- [126442]Sci. 87: 7000-7004, 1990.
- [126443]22549.
- [126444]Jirik, F. R.; Anderson, L. L.; Duncan, A. M. V.: The human protein-tyrosine
- [126445]phosphatase PTP-alpha/LRP gene (PTPA) is assigned to chromosome 20p13. Cytogenet.
- [126446]Cell Genet. 60: 117-118, 1992.
- [126447]22550.
- [126448]Jirik, F. R.; Janzen, N. M.; Melhado, I. G.; Harder, K. W.: Cloning
- [126449]and chromosomal assignment of a widely expressed human receptor-like
- [126450]protein-tyrosine phosphatase. FEBS Lett. 273: 239-242, 1990.
- [126451]22551.

[126452]Matthews, R. J.; Cahir, E. D.; Thomas, M. L.: Identification
of

[126453]an additional member of the protein-tyro-
sine-phosphatase family: evidence

[126454]for alternative splicing in the tyrosine phosphatase do-
main. Proc.

[126455]Nat. Acad. Sci. 87: 4444-4448, 1990.

[126456]22552.

[126457]Rao, V. V. N. G.; Loffler, C.; Sap, J.; Schlessinger, J.; Hans-
mann,

[126458]I.: The gene for receptor-linked protein-tyro-
sine-phosphatase (PTPA)

[126459]is assigned to human chromosome 20p12-pter by in situ
hybridization

[126460](ISH and FISH). Genomics 13: 906-907, 1992.

[126461]22553.

[126462]Schnittger, S.; Rao, V. V. N. G.; Deutsch, U.; Gruss, P.;
Balling,

[126463]R.; Hansmann, I.: PAX1, a member of the paired box-
containing class

[126464]of developmental control genes, is mapped to human
chromosome 20p11.2

[126465]by in situ hybridization (ISH and FISH). Genomics 14:
740–744, 1992.

[126466]22554.

[126467]Ahmad, F.; Azevedo, J. L., Jr.; Cortright, R.; Dohm, G. L.;
Goldstein,

[126468]B. J.: Alternations in skeletal muscle protein-tyrosine
phosphatase

[126469]activity and expression in insulin-resistant human obesity
and diabetes. J.

[126470]Clin. Invest. 100: 449–458, 1997.

[126471]22555.

[126472]Brown-Shimer, S.; Johnson, K. A.; Lawrence, J. B.; Johnson,
C.;

[126473]Bruskin, A.; Green, N. R.; Hill, D. E.: Molecular cloning and
chromosome

[126474]mapping of the human gene encoding protein phosphoty-
rosyl phosphatase

[126475]1B. Proc. Nat. Acad. Sci. 87: 5148–5152, 1990.

[126476]22556.

[126477]Charbonneau, H.; Tonks, N. K.; Kumar, S.; Diltz, C. D.;
Harrylock,

- [126478]M.; Cool, D. E.; Krebs, E. G.; Fischer, E. H.; Walsh, K. A.:
Human
[126479]placenta protein-tyrosine-phosphatase: amino acid se-
quence and relationship
[126480]to a family of receptor-like proteins. Proc. Nat. Acad. Sci.
86:
[126481]5252-5256, 1989.
[126482]22557.
[126483]Chernoff, J.; Schievella, A. R.; Jost, C. A.; Erikson, R. L.;
Neel,
[126484]B. G.: Cloning of a cDNA for a major human protein-tyro-
sine-phosphatase. Proc.
[126485]Nat. Acad. Sci. 87: 2735-2739, 1990.
[126486]22558.
[126487]Kerr, N. C.; Wilroy, R. S., Jr.; Kaufman, R. A.: Type 3 Pfeif-
fer
[126488]syndrome with normal thumbs. Am. J. Med. Genet. 66:
138-143, 1996.
[126489]22559.
[126490]Lajeunie, E.; Cameron, R.; El Ghouzzi, V.; de Parseval, N.;
Journeau,

- [126491]P.; Gonzales, M.; Delezoide, A.-L.; Bonaventure, J.; Le
Merrer, M.;
- [126492]Renier, D.: Clinical variability in patients with Apert's syn-
drome. J.
- [126493]Neurosurg. 90: 443-447, 1999.
- [126494]22560.
- [126495]Lajeunie, E.; Ma, H. W.; Bonaventure, J.; Munnich, A.; Le
Merrer,
- [126496]M.; Renier, D.: FGFR2 mutations in Pfeiffer syndrome.
(Letter) Nature
- [126497]Genet. 9: 108, 1995.
- [126498]22561.
- [126499]Ma, H. W.; Lajeunie, E.; Le\Merrer, M.; de\Parseval, N.;
Serville,
- [126500]F.; Weissenbach, J.; Munnich, A.; Renier, D.: No evidence
of genetic
- [126501]heterogeneity in Crouzon craniofacial dysostosis. Hum.
Genet. 96:
- [126502]731-735, 1995.
- [126503]22562.
- [126504]Mattei, M.-G.; Moreau, A.; Gesnel, M.-C.; Houssaint, E.;

Breathnach,

[126505]R.: Assignment by in situ hybridization of a fibroblast growth factor

[126506]receptor gene to human chromosome band 10q26. Hum. Genet. 87: 84–86,

[126507]1991.

[126508]22563.

[126509]Meyers, G. A.; Day, D.; Goldberg, R.; Daentl, D. L.; Przylepa,

[126510]K. A.; Abrams, L. J.; Graham, J. M., Jr.; Feingold, M.; Moeschler,

[126511]J. B.; Rawnsley, E.; Scott, A. F.; Jabs, E. W.: FGFR2 exon IIIa and

[126512]IIIc mutations in Crouzon, Jackson–Weiss, and Pfeiffer syndromes:

[126513]evidence for missense changes, insertions, and a deletion due to alternative

[126514]RNA splicing. Am. J. Hum. Genet. 58: 491–498, 1996.

[126515]22564.

[126516]Miki, T.; Fleming, T. P.; Bottaro, D. P.; Rubin, J. S.; Ron, D.;

[126517]Aaronson, S. A.: Expression cDNA cloning of the KGF re–

ceptor by creation

[126518]of a transforming autocrine loop. Science 251: 72–75,
1991.

[126519]22565.

[126520]Mulvihill, J. J.: Craniofacial syndromes: no such thing as a

[126521]single gene disease. Nature Genet. 9: 101–103, 1995.

[126522]22566.

[126523]Munro, C. S.; Wilkie, A. O. M.: Epidermal mosaicism pro-
ducing

[126524]localised acne: somatic mutation in FGFR2. Lancet 352:
704–705,

[126525]1998.

[126526]22567.

[126527]Okajima, K.; Robinson, L. K.; Hart, M. A.; Abuelo, D. N.;
Cowan,

[126528]L. S.; Hasegawa, T.; Maumenee, I. H.; Jabs, E. W.: Ocular
anterior

[126529]chamber dysgenesis in craniosynostosis syndromes with a
fibroblast

[126530]growth factor receptor 2 mutation. Am. J. Med. Genet. 85:
160–170,

[126531]1999.

[126532]22568.

[126533]Oldridge, M.; Lunt, P. W.; Zackai, E. H.; McDonald-McGinn, D.

[126534]M.; Muenke, M.; Moloney, D. M.; Twigg, S. R. F.; Heath, J. K.; Howard,

[126535]T. D.; Hoganson, G.; Gagnon, D. M.; Jabs, E. W.; Wilkie, A. O. M.

[126536]: Genotype-phenotype correlation for nucleotide substitutions in the

[126537]IgII-IgIII linker of FGFR2. Hum. Molec. Genet. 6: 137-143, 1997.

[126538]22569.

[126539]Oldridge, M.; Zackai, E. H.; McDonald-McGinn, D. M.; Iseki, S.;

[126540]Morriss-Kay, G. M.; Twigg, S. R. F.; Johnson, D.; Wall, S. A.; Jiang,

[126541]W.; Theda, C.; Jabs, E. W.; Wilkie, A. O. M.: De novo Alu-element

[126542]insertions in FGFR2 identify a distinct pathological basis for Apert

[126543]syndrome. Am. J. Hum. Genet. 64: 446-461, 1999.

[126544]22570.

[126545]Park, W.-J.; Meyers, G. A.; Li, X.; Theda, C.; Day, D.; Orlow,

[126546]S. J.; Jones, M. C.; Jabs, E. W.: Novel FGFR2 mutations in
Crouzon

[126547]and Jackson-Weiss syndromes show allelic heterogeneity
and phenotypic

[126548]variability. Hum. Molec. Genet. 4: 1229-1233, 1995.

[126549]22571.

[126550]Passo-Buenos, M. R.; Richieri-Costa, A.; Sertie, A. L.;
Kneppers,

[126551]A.: Presence of the Apert canonical S252W FGFR2 muta-
tion in a patient

[126552]without severe syndactyly. J. Med. Genet. 35: 677-679,
1998.

[126553]22572.

[126554]Priolo, M.; Lerone, M.; Baffico, M.; Baldi, M.; Ravazzolo, R.;

[126555]Cama, A.; Capra, V.; Silengo, M.: Pfeiffer syndrome type 2
associated

[126556]with a single amino acid deletion in the FGFR2 gene.
(Letter) Clin.

[126557]Genet. 58: 81-83, 2000.

[126558]22573.

[126559]Rees, J.: FGFR2 mutations and acne. (Letter) Lancet 352:
668–669,

[126560]1998.

[126561]22574.

[126562]Rutland, P.; Pulleyn, L. J.; Reardon, W.; Baraitser, M.; Hay-
ward,

[126563]R.; Jones, B.; Malcolm, S.; Winter, R. M.; Oldridge, M.;
Slaney, S.

[126564]F.; Poole, M. D.; Wilkie, A. O. M.: Identical mutations in the
FGFR2

[126565]gene cause both Pfeiffer and Crouzon syndrome pheno-
types. Nature

[126566]Genet. 9: 173–176, 1995.

[126567]22575.

[126568]Schaefer, F.; Anderson, C.; Can, B.; Say, B.: Novel mutation

[126569]in the FGFR2 gene at the same codon as the Crouzon syn-
drome mutations

[126570]in a severe Pfeiffer syndrome type 2 case. Am. J. Med.
Genet. 75:

[126571]252–255, 1998.

[126572]22576.

[126573]Shuster, S.: Dermatology in Clinical Medicine. Oxford: Oxford
ford

[126574]Univ. Press , 1978.

[126575]22577.

[126576]Slaney, S. F.; Oldridge, M.; Hurst, J. A.; Morriss-Kay, G. M.;

[126577]Hall, C. M.; Poole, M. D.; Wilkie, A. O. M.: Differential ef-
fects

[126578]of FGFR2 mutations on syndactyly and cleft palate in Apert
syndrome. Am.

[126579]J. Hum. Genet. 58: 923-932, 1996.

[126580]22578.

[126581]Steinberger, D.; Collmann, H.; Schmalenberger, B.; Muller,
U.

[126582]: A novel mutation (A886G) in exon 5 of FGFR2 in mem-
bers of a family

[126583]with Crouzon phenotype and plagiocephaly. J. Med. Genet.
34: 420-422,

[126584]1997.

[126585]22579.

[126586]Steinberger, D.; Mulliken, J. B.; Muller, U.: Predisposition
[126587]for cysteine substitutions in the immunoglobulin-like
chain of FGFR2
[126588]in Crouzon syndrome. Hum. Genet. 96: 113–115, 1995.
[126589]22580.

[126590]Vaughan, K. T.; Weber, F. E.; Ried, T.; Ward, D. C.;
Reinach, F.
[126591]C.; Fischman, D. A.: Human myosin-binding protein H
(MyBP-H): complete
[126592]primary sequence, genomic organization, and chromoso-
mal localization. Genomics 16:
[126593]34–40, 1993.
[126594]22581.

[126595]Torbergesen, T.: A family with dominant hereditary myoto-
nia, muscular
[126596]hypertrophy, and increased muscular irritability, distinct
from myotonia
[126597]congenita Thomsen. Acta Neurol. Scand. 51: 225–232,
1975.
[126598]22582.

[126599]Harding, A. E.: The clinical features and classification of

the

[126600]late onset autosomal dominant cerebellar ataxias: a study
of 11 families,

[126601]including descendants of 'the Drew family of Walworth'.

Brain 105:

[126602]1–28, 1982.

[126603]22583.

[126604]Havener, W. H.: Cerebellar–macular abiotrophy. Arch.

Ophthal. 45:

[126605]40–43, 1951.

[126606]22584.

[126607]Holmberg, M.; Johansson, J.; Forsgren, L.; Heijbel, J.;

Sandgren,

[126608]O.; Holmgren, G.: Localization of autosomal dominant
cerebellar ataxia

[126609]associated with retinal degeneration and anticipation to
chromosome

[126610]3p12–p21.1. Hum. Molec. Genet. 4: 1441–1445, 1995.

[126611]22585.

[126612]Jampel, R. S.; Okazaki, H.; Bernstein, H.: Ophthalmoplegia
and

- [126613]retinal degeneration associated with spinocerebellar ataxia. Arch.
- [126614]Ophthalmol. 66: 247–259, 1961.
- [126615]22586.
- [126616]Jonasson, J.; Juvonen, V.; Sistonen, P.; Ignatius, J.; Johansson,
- [126617]D.; Bjorck, E. J.; Wahlstrom, J.; Melberg, A.; Holmgren, G.; Forsgren,
- [126618]L.; Holmberg, M.: Evidence for a common spinocerebellar ataxia type
- [126619]7 (SCA7) founder mutation in Scandinavia. Europ. J. Hum. Genet. 8:
- [126620]918–922, 2000.
- [126621]22587.
- [126622]Kaytor, M. D.; Duvick, L. A.; Skinner, P. J.; Koob, M. D.; Ranum,
- [126623]L. P. W.; Orr, H. T.: Nuclear localization of the spinocerebellar
- [126624]ataxia type 7 protein, ataxin-7. Hum. Molec. Genet. 8: 1657–1664,
- [126625]1999.

[126626]22588.

[126627]Koob, M. D.; Benzow, K. A.; Bird, T. D.; Day, J. W.; Moseley,

[126628]M. L.; Ranum, L. P. W.: Rapid cloning of expanded trinucleotide repeat

[126629]sequences from genomic DNA. *Nature Genet.* 18: 72–75, 1998.

[126630]22589.

[126631]Krols, L.; Martin, J.-J.; David, G.; Van Regemorter, N.; Benomar,

[126632]A.; Lofgren, A.; Stevanin, G.; Durr, A.; Brice, A.; Van Broeckhoven,

[126633]C.: Refinement of the locus for autosomal dominant cerebellar ataxia

[126634]type II to chromosome 3p21.1–14.1. *Hum. Genet.* 99: 225–232, 1997.

[126635]22590.

[126636]Lebre, A.-S.; Jamot, L.; Takahashi, J.; Spassky, N.; Lepage,

[126637]C.; Ravise, N.; Zander, C.; Fujigasaki, H.; Kussel-Andermann, P.;

[126638]Duyckaerts, C.; Camonis, J. H.; Brice, A.: Ataxin-7 interacts with

[126639]a Cbl-associated protein that it recruits into neuronal in-
tranuclear

[126640]inclusions. Hum. Molec. Genet. 10: 1201-1213, 2001.

[126641]22591.

[126642]Lindblad, K.; Savontaus, M.-L.; Stevanin, G.; Holmberg, M.;
Digre,

[126643]K.; Zander, C.; Ehrsson, H.; David, G.; Benomar, A.;
Nikoskelainen,

[126644]E.; Trottier, Y.; Holmgren, G.; Ptacek, L. J.; Anttinen, A.;
Brice,

[126645]A.; Schalling, M.: An expanded CAG repeat sequence in
spinocerebellar

[126646]ataxia type 7. Genome Res. 6: 965-971, 1996.

[126647]22592.

[126648]Matilla, A.; Gorbea, C.; Einum, D. D.; Townsend, J.; Micha-
lik,

[126649]A.; van Broeckhoven, C.; Jensen, C. C.; Murphy, K. J.;
Ptacek, L.

[126650]J.; Fu, Y.-H.: Association of ataxin-7 with the proteasome
subunit

[126651]S4 of the 19S regulatory complex. Hum. Molec. Genet. 10:
2821-2831,

[126652]2001.

[126653]22593.

[126654]Michalik, A.; Del-Favero, J.; Mauger, C.; Lofgren, A.; Van Broeckhoven,

[126655]C.: Genomic organisation of the spinocerebellar ataxia type 7 (SCA7)

[126656]gene responsible for autosomal dominant cerebellar ataxia with retinal

[126657]degeneration. Hum. Genet. 105: 410-417, 1999.

[126658]22594.

[126659]Stevanin, G.; Giunti, P.; David, G.; Belal, S.; Durr, A.; Ruberg,

[126660]M.; Wood, N.; Brice, A.: De novo expansion of intermediate alleles

[126661]in spinocerebellar ataxia 7. Hum. Molec. Genet. 7: 1809-1813, 1998.

[126662]22595.

[126663]Trottier, Y.; Lutz, Y.; Stevanin, G.; Imbert, G.; Devys, D.; Cancel,

[126664]G.; Saudou, F.; Weber, C.; David, G.; Tora, L.; Agid, Y.; Brice, A.;

- [126665]Mandel, J.-L.: Polyglutamine expansion as a pathological epitope
in Huntington's disease and four dominant cerebellar ataxias. *Nature* 378: 403–406, 1995.
- [126668]22596.
- [126669]Wadia, N. H.; Swami, R. K.: A new form of heredo-familial spinocerebellar degeneration with slow eye movements (nine families). *Brain* 94: 359–374, 1971.
- [126672]22597.
- [126673]Weiner, L. P.; Konigsmark, B. W.; Stoll, J., Jr.; Magladery, J. W.: Hereditary olivopontocerebellar atrophy with retinal degeneration: report of a family through six generations. *Arch. Neurol.* 16: 364–376, 1967.
- [126676]1967.
- [126677]22598.
- [126678]Woodworth, J. A.; Beckett, R. S.; Netsky, M. G.: A composite

[126679] of hereditary ataxias: a familial disorder with features of olivopontocerebellar

[126680] atrophy, Leber's optic atrophy and Friedreich's ataxia.
Arch. Intern.

[126681] Med. 104: 594–606, 1959.

[126682] 22599.

[126683] Yvert, G.; Lindenberg, K. S.; Picaud, S.; Landwehrmeyer, G.
B.;

[126684] Sahel, J. –A.; Mandel, J.–L.: Expanded polyglutamines induce neurodegeneration

[126685] and trans-neuronal alterations in cerebellum and retina of SCA7 transgenic

[126686] mice. Hum. Molec. Genet. 9: 2491–2506, 2000.

[126687] 22600.

[126688] Zander, C.; Takahashi, J.; El Hachimi, K. H.; Fujigasaki, H.;

[126689] Albanese, V.; Lebre, A. S.; Stevanin, G.; Duyckaerts, C.;
Brice, A.

[126690]: Similarities between spinocerebellar ataxia type 7 (SCA7)
cell models

[126691] and human brain: proteins recruited in inclusions and activation of

[126692]caspase-3. Hum. Molec. Genet. 10: 2569-2579, 2001.

[126693]22601.

[126694]Cazzaniga, G.; Tosi, S.; Aloisi, A.; Giudici, G.; Daniotti, M.;

[126695]Pioltelli, P.; Kearney, L.; Biondi, A.: The tyrosine kinase
Abl-related

[126696]gene ARG is fused to ETV6 in an AML-M4Eo patient with a
t(1;12)(q25;p13):

[126697]molecular cloning of both reciprocal transcripts. Blood 94:
4370-4373,

[126698]1999.

[126699]22602.

[126700]Iijima, Y.; Ito, T.; Oikawa, T.; Eguchi, M.; Eguchi-Ishimae,
M.;

[126701]Kamada, N.; Kishi, K.; Asano, S.; Sakaki, Y.; Sato, Y.: A new
ETV6/TEL

[126702]partner gene, ARG (ABL-related gene or ABL2), identified
in an AML-M3

[126703]cell line with a t(1;12)(q25;p13) translocation. Blood 95:
2126-2131,

[126704]2000.

[126705]22603.

[126706] Mastick, C. C.; Brady, M. J.; Saltiel, A. R.: Insulin stimulates
[126707] the tyrosine phosphorylation of caveolin. *J. Cell Biol.* 129:
1523–1531,
[126708] 1995.
[126709] 22604.
[126710] Petrelli, A.; Gilestro, G. F.; Lanzardo, S.; Comoglio, P. M.;
Migone,
[126711] N.; Giordano, S.: The endophilin–CIN85–Cbl complex me-
diates ligand–dependent
[126712] downregulation of c–Met. *Nature* 416: 187–190, 2002.
[126713] 22605.
[126714] Soubeyran, P.; Kowanetz, K.; Szymkiewicz, I.; Langdon, W.
Y.; Dikic,
[126715] J.: Cbl–CIN85–endophilin complex mediates ligand–in-
duced downregulation
[126716] of EGF receptors. *Nature* 416: 183–187, 2002.
[126717] 22606.
[126718] Cannizzaro, L. A.; Madaule, P.; Hecht, F.; Axel, R.; Croce,
C.
[126719] M.; Huebner, K.: Chromosome localization of human ARH
genes, a ras–related

[126720]gene family. Genomics 6: 197–203, 1990.

[126721]22607.

[126722]Chardin, P.; Madaule, P.; Tavitian, A.: Coding sequence of human

[126723]rho cDNAs clone 6 and clone 9. Nucleic Acid Res. 16: 2717 only,

[126724]1988.

[126725]22608.

[126726]Madaule, P.; Axel, R.: A novel ras-related gene family. Cell 41:

[126727]31–40, 1985.

[126728]22609.

[126729]Maekawa, M.; Ishizaki, T.; Boku, S.; Watanabe, N.; Fujita, A.;

[126730]Iwamatsu, A.; Obinata, T.; Ohashi, K.; Mizuno, K.; Narumiya, S.:

[126731]Signaling from Rho to the actin cytoskeleton through protein kinases

[126732]ROCK and LIM-kinase. Science 285: 895–898, 1999.

[126733]22610.

[126734]Ridley, A. J.; Hall, A.: The small GTP-binding protein rho regulates

[126735]the assembly of focal adhesions and actin stress fibers in response

[126736]to growth factors. Cell 70: 389-399, 1992.

[126737]22611.

[126738]Fagan, K. P.; Oliveira, L.; Pittler, S. J.: Sequence of rho small

[126739]GTP-binding protein cDNAs from human retina and identification of

[126740]novel 5-prime end cloning artifacts. Exp. Eye Res. 59: 235-237,

[126741]1994.

[126742]22612.

[126743]Morris, S. W.; Valentine, M. B.; Kirstein, M. N.; Huebner, K.:

[126744]Reassignment of the human ARH9 RAS-related gene to chromosome 1p13-p21. Genomics 15:

[126745]677-679, 1993.

[126746]22613.

[126747]Maesaki, R.; Ihara, K.; Shimizu, T.; Kuroda, S.; Kaibuchi, K.;

- [126748]Hakoshima, T.: The structural basis of Rho effector recognition revealed
by the crystal structure of human RhoA complexed with the effector domain of PKN/PRK1. Molec. Cell 4: 793–803, 1999.
22614.
- [126752]Nakamura, M.; Nagano, T.; Chikama, T.; Nishida, T.: Role of the
small GTP-binding protein Rho in epithelial cell migration in the
rabbit cornea. Invest. Ophthalmol. Vis. Sci. 42: 941–947, 2001.
22615.
- [126756]Rao, P. V.; Deng, P.-F.; Kumar, J.; Epstein, D. L.: Modulation
of aqueous humor outflow facility by the Rho kinase-specific inhibitor
Y-27632. Invest. Ophthalmol. Vis. Sci. 42: 1029–1037, 2001.
22616.
- [126760]Alexander, C.; Votruba, M.; Pesch, U. E. A.; Thiselton, D. L.;

[126761]Mayer, S.; Moore, A.; Rodriguez, M.; Kellner, U.; Leo-Kottler, B.;

[126762]Auburger, G.; Bhattacharya, S. S.; Wissinger, B.: OPA1, encoding

[126763]a dynamin-related GTPase, is mutated in autosomal dominant optic atrophy

[126764]linked to chromosome 3q28. Nature Genet. 26: 211–215, 2000.

[126765]22617.

[126766]Rosenfeld, P. J.; Cowley, G. S.; McGee, T. L.; Sandberg, M. A.;

[126767]Berson, E. L.; Dryja, T. P.: A null mutation in the rhodopsin gene

[126768]causes rod photoreceptor dysfunction and autosomal recessive retinitis

[126769]pigmentosa. Nature Genet. 1: 209–213, 1992.

[126770]22618.

[126771]Pulst, S.-M.; Nechiporuk, A.; Nechiporuk, T.; Gispert, S.; Chen,

[126772]X.-N.; Lopes-Cendes, I.; Pearlman, S.; Starkman, S.; Orozco-Diaz,

[126773]G.; Lunkes, A.; DeJong, P.; Rouleau, G. A.; Auburger, G.;

Korenberg,

[126774]J. R.; Figueroa, C.; Sahba, S.: Moderate expansion of a normally

[126775]biallelic trinucleotide repeat in spinocerebellar ataxia type

2. Nature

[126776]Genet. 14: 269–276, 1996.

[126777]22619.

[126778]Sanpei, K.; Takano, H.; Igarashi, S.; Sato, T.; Oyake, M.;

Sasaki,

[126779]H.; Wakisaka, A.; Tashiro, K.; Ishida, Y.; Ikeuchi, T.; Koide, R.;

[126780]Saito, M.; Sato, A.; Tanaka, T.; Hanyu, S.; Takiyama, Y.; Nishizawa,

[126781]M.; Shimizu, N.; Nomura, Y.; Segawa, M.; Iwabuchi, K.; Eguchi, I.;

[126782]Tanaka, H.; Takahashi, H.; Tsuji, S.: Identification of the spinocerebellar

[126783]ataxia type 2 gene using a direct identification of repeat expansion

[126784]and cloning technique, DIRECT. Nature Genet. 14: 277–284, 1996.

[126785]22620.

[126786]Chen, K.; Yang, W.; Grimsby, J.; Shih, J. C.: The human
5-HT2

[126787]receptor is encoded by a multiple intron-exon gene. Brain
Res. Molec.

[126788]Brain Res. 14: 20-26, 1992.

[126789]22621.

[126790]Holmes, C.; Arranz, M. J.; Powell, J. F.; Collier, D. A.; Love-
stone,

[126791]S.: 5-HT-2A and 5-HT-2C receptor polymorphisms and
psychopathology

[126792]in late onset Alzheimer's disease. Hum. Molec. Genet. 7:
1507-1509,

[126793]1998.

[126794]22622.

[126795]Hsieh, C.-L.; Bowcock, A. M.; Farrer, L. A.; Hebert, J. M.;
Huang,

[126796]K. N.; Cavalli-Sforza, L. L.; Julius, D.; Francke, U.: The
serotonin

[126797]receptor subtype 2 locus HTR2 is on human chromosome
13 near genes

[126798]for esterase D and retinoblastoma-1 and on mouse chro-
mosome 14. Somat.

[126799]Cell Molec. Genet. 16: 567–574, 1990.

[126800]22623.

[126801]Kato, M. V.; Shimizu, T.; Nagayoshi, M.; Kaneko, A.;
Sasaki, M.

[126802]S.; Ikawa, Y.: Genomic imprinting of the human serotonin-
receptor

[126803](HTR2) gene involved in development of retinoblastoma.
Am. J. Hum.

[126804]Genet. 59: 1084–1090, 1996.

[126805]22624.

[126806]Liu, J.; Chen, Y.; Kozak, C. A.; Yu, L.: The 5-HT(2) sero-
tonin

[126807]receptor gene Htr-2 is tightly linked to Es-10 on mouse
chromosome

[126808]14. Genomics 11: 231–234, 1991.

[126809]22625.

[126810]Peroutka, S. J.: 5-Hydroxytryptamine receptor subtypes.
Ann.

[126811]Rev. Neurosci. 11: 45–60, 1988.

[126812]22626.

[126813]Sparkes, R. S.; Lan, N.; Klisak, I.; Mohandas, T.; Diep, A.;
Kojis,

[126814]T.; Heinzmann, C.; Shih, J. C.: Assignment of a serotonin
5HT-2 receptor

[126815]gene (HTR2) to human chromosome 13q14-q21 and
mouse chromosome 14. Genomics 9:

[126816]461-465, 1991.

[126817]22627.

[126818]Williams, J.; McGuffin, P.; Nothen, M.; Owen, M. J.; EMAS
Collaborative

[126819]Group: Meta-analysis of association between the 5-HT
(2A) receptor

[126820]T102C polymorphism and schizophrenia. (Letter) Lancet
349: 1221

[126821]only, 1997.

[126822]22628.

[126823]Williams, J.; Spurlock, G.; McGuffin, P.; Mallet, J.; Nothen,
M.

[126824]M.; Gill, M.; Aschauer, H.; Nylander, P. O.; Macciardi, F.;
Owen,

[126825]M. J.: Association between schizophrenia and T102C poly-
morphism of

- [126826]the 5-hydroxytryptophan type 2a-receptor gene. Euro-
pean Multicentre
[126827]Association Study of Schizophrenia (EMASS) Group. Lancet
347: 1294-1296,
[126828]1996.
[126829]22629.
- [126830]Wulff, C.; Wilson, H.; Rudge, J. S.; Wiegand, S. J.; Lunn, S.
F.;
- [126831]Fraser, H. M.: Luteal angiogenesis: prevention and inter-
vention by
[126832]treatment with vascular endothelial growth factor trapA40.
J. Clin.
[126833]Endocr. Metab. 86: 3377-3386, 2001.
[126834]22630.
- [126835]Philipson, L. H.; Hice, R. E.; Schaefer, K.; LaMendola, J.;
Bell,
- [126836]G. I.; Neldon, D. J.; Steiner, D. F.: Sequence and functional
expression
[126837]in *Xenopus* oocytes of a human insulinoma and islet
potassium channel. Proc.
[126838]Nat. Acad. Sci. 88: 53-57, 1991.

[126839]22631.

[126840]McPherson, J. D.; Wasmuth, J. J.; Chandy, K. G.; Swanson, R.; Dethlefs,

[126841]B.; Chandy, G.; Wymore, R.; Ghanshani, S.: Chromosomal localization

[126842]of 7 potassium channel genes. (Abstract) Cytogenet. Cell Genet. 58:

[126843]1979 only, 1991.

[126844]22632.

[126845]Tamkun, M. M.; Knoth, K. M.; Walbridge, J. A.; Kroemer, H.; Roden,

[126846]D. M.; Glover, D. M.: Molecular cloning and characterization of two

[126847]voltage-gated K⁺ channel cDNAs from human ventricle. FASEB J. 5:

[126848]331-337, 1991.

[126849]22633.

[126850]Phromchotikul, T.; Browne, D. L.; Curran, M. E.; Keating, M. T.;

[126851]Litt, M.: Dinucleotide repeat polymorphism at the KCNA5 locus. Hum.

[126852]Molec. Genet. 2: 1512 only, 1993.

[126853]22634.

[126854]Bardien-Kruger, S.; Wulff, H.; Arieff, Z.; Brink, P.; Chandy, K.

[126855]G.; Corfield, V.: Characterisation of the human voltage-gated potassium

[126856]channel gene, KCNA7, a candidate gene for inherited cardiac disorders,

[126857]and its exclusion as cause of progressive familial heart block I (PFHBI). Europ.

[126858]J. Hum. Genet. 10: 36-43, 2002.

[126859]22635.

[126860]Barton, J. W.; Hart, I. M.; Patterson, D.: Mapping of a locus

[126861]correcting lack of phosphoribosylaminoimidazole carboxylase activity

[126862]in Chinese hamster ovary cell Ade(-)D mutants to human chromosome

[126863]4. Genomics 9: 314-321, 1991.

[126864]22636.

[126865]Brayton, K. A.; Chen, Z.; Zhou, G.; Nagy, P. L.; Gavalas, A.; Trent,

[126866]J. M.; Deaven, L. L.; Dixon, J. E.; Zalkin, H.: Two genes for
de

[126867]novo purine nucleotide synthesis on human chromosome
4 are closely

[126868]linked and divergently transcribed. J. Biol. Chem. 269:
5313–5321,

[126869]1994.

[126870]22637.

[126871]Chu, E. H. Y.; Sun, N. C.; Chang, C. C.: Induction of aux-
otrophic

[126872]mutants by treatment of Chinese hamster cells with
5–bromodeoxyuridine

[126873]and black light. Proc. Nat. Acad. Sci. 68: 3459–3463,
1972.

[126874]22638.

[126875]Feldman, R. I.; Taylor, M. W.: Purine mutants of mam-
malian cell

[126876]lines. II. Identification of a phosphoribosylpyrophosphate
amidotransferase–deficient

[126877]mutant of Chinese hamster lung cells. Biochem. Genet.
13: 227–234,

[126878]1975.

[126879]22639.

[126880]Cavalas, A.; Dixon, J. E.; Brayton, K. A.; Zalkin, H.: Coexpression

[126881]of two closely linked avian genes for purine nucleotide synthesis

[126882]from a bidirectional promoter. Molec. Cell. Biol. 13: 4784-4792,

[126883]1993.

[126884]22640.

[126885]Jones, C.; Morse, H. G.; Palmer, D. K.: Comparative gene mapping

[126886]of man and higher primates involving genes assigned to human chromosomes

[126887]3, 4, 11, 21. (Abstract) Cytogenet. Cell Genet. 40: 662-663, 1985.

[126888]22641.

[126889]Stanley, W.; Chu, E. H. Y.: Assignment of the gene for phosphoribosylpyrophosphate

[126890]amidotransferase to the pter-q21 region of human chromosome 4. Cytogenet.

[126891]Cell Genet. 22: 228-231, 1978.

[126892]22642.

[126893]Zalkin, H.; Dixon, J. E.: De novo purine nucleotide biosynthesis. Prog.

[126894]Nucleic Acids Res. Molec. Biol. 42: 259–287, 1992.

[126895]22643.

[126896]Parfrey, P. S.; Bear, J. C.; Morgan, J.; Cramer, B. C.; McManamon,

[126897]P. J.; Gault, M. H.; Churchill, D. N.; Singh, M.; Hewitt, R.; Somlo,

[126898]S.; Reeders, S. T.: The diagnosis and prognosis of autosomal dominant

[126899]polycystic kidney disease. New Eng. J. Med. 323: 1085–1090, 1990.

[126900]22644.

[126901]Peral, B.; Gamble, V.; San Millan, J. L.; Strong, C.; Sloane-Stanley,

[126902]J.; Moreno, F.; Harris, P. C.: Splicing mutations of the polycystic

[126903]kidney disease 1 (PKD1) gene induced by intronic deletion. Hum. Molec.

[126904]Genet. 4: 569–574, 1995.

[126905]22645.

[126906]Peral, B.; Ong, A. C. M.; San Millan, J. L.; Gamble, V.; Rees,

[126907]L.; Harris, P. C.: A stable, nonsense mutation associated
with a

[126908]case of infantile onset polycystic kidney disease 1 (PKD1).
Hum.

[126909]Molec. Genet. 5: 539–542, 1996.

[126910]22646.

[126911]Peral, B.; San Millan, J. L.; Hernandez, C.; Valero, A.; Lath-
rop,

[126912]G. M.; Beckmann, J. S.; Moreno, F.: Estimating locus het-
erogeneity

[126913]in autosomal dominant polycystic kidney disease (ADPKD)
in the Spanish

[126914]population. J. Med. Genet. 30: 910–913, 1993.

[126915]22647.

[126916]Peral, B.; San Millan, J. L.; Ong, A. C. M.; Gamble, V.; Ward,

[126917]C. J.; Strong, C.; Harris, P. C.: Screening the 3–prime re-
gion of

[126918]the polycystic kidney disease 1 (PKD1) gene reveals six
novel mutations. Am.

[126919]J. Hum. Genet. 58: 86–96, 1996.

[126920]22648.

[126921]Pound, S. E.; Carothers, A. D.; Pignatelli, P. M.; Macnicol, A.

[126922]M.; Watson, M. L.; Wright, A. F.: Evidence for linkage disequilibrium

[126923]between D16S94 and the adult onset polycystic kidney disease (PKD1)

[126924]gene. J. Med. Genet. 29: 247–248, 1992.

[126925]22649.

[126926]Qian, F.; Watnick, T. J.; Onuchic, L. F.; Germino, G. G.: The

[126927]molecular basis of focal cyst formation in human autosomal dominant

[126928]polycystic kidney disease type I. Cell 87: 979–987, 1996.

[126929]22650.

[126930]Reeders, S. T.: Personal Communication. Oxford, England 1986.

[126931]22651.

[126932]Reeders, S. T.: Multilocus polycystic disease. Nature Genet. 1:

[126933]235–237, 1992.

[126934]22652.

[126935]Reeders, S. T.: Personal Communication. Oxford, England
10/11/1985.

[126936]22653.

[126937]Reeders, S. T.; Breuning, M. H.; Davies, K. E.; Nicholls, R.
D.;

[126938]Jarman, A. P.; Higgs, D. R.; Pearson, P. L.; Weatherall, D. J.:
A

[126939]highly polymorphic DNA marker linked to adult polycystic
kidney disease

[126940]on chromosome 16. Nature 317: 542–544, 1985.

[126941]22654.

[126942]Reeders, S. T.; Breuning, M. H.; Ryynanen, M. A.; Wright,
A. F.;

[126943]Davies, K. E.; King, A. W.; Watson, M. L.; Weatherall, D. J.:
A study

[126944]of genetic linkage heterogeneity in adult polycystic kidney
disease. Hum.

[126945]Genet. 76: 348–351, 1987.

[126946]22655.

[126947]Reeders, S. T.; Keith, T.; Green, P.; Germino, G. G.; Barton,

[126948]N. J.; Lehmann, O. J.; Brown, V. A.; Phipps, P.; Morgan, J.;
Bear,

[126949]J. C.; Parfrey, P.: Regional localization of the autosomal
dominant

[126950]polycystic kidney disease locus. Genomics 3: 150–155,
1988.

[126951]22656.

[126952]Ryynanen, M.; Dolata, M. M.; Lampainen, E.; Reeders, S. T.:
Localisation

[126953]of a mutation producing autosomal dominant polycystic
kidney disease

[126954]without renal failure. J. Med. Genet. 24: 462–465, 1987.

[126955]22657.

[126956]Liu, S. C.; Palek, J.; Prchal, J. T.: Defective spectrin dimer-
dimer

[126957]association in hereditary elliptocytosis. Proc. Nat. Acad.
Sci. 79:

[126958]2072–2076, 1982.

[126959]22658.

[126960]Torban, E.; Pelletier, J.; Goodyer, P.: F329L polymorphism

in

[126961]the human PAX8 gene. *Am. J. Med. Genet.* 72: 186–187, 1997.

[126962]22659.

[126963]Gartner, J.; Kearns, W.; Pearson, P.; Valle, D.: Characterization

[126964]and localization of the human 70-kD peroxisomal membrane protein (PMP70)

[126965]gene. (Abstract) *Am. J. Hum. Genet.* 51 (suppl.): A168, 1992.

[126966]22660.

[126967]Gartner, J.; Kearns, W.; Rosenberg, C.; Pearson, P.; Copeland,

[126968]N. G.; Gilbert, D. J.; Jenkins, N. A.; Valle, D.: Localization of

[126969]the 70-kDa peroxisomal membrane protein to human 1p21-p22 and mouse 22661.

[126970]*Genomics* 15: 412–414, 1993.

[126971]

[126972]3. Gartner, J.; Moser, H.; Valle, D.: Mutations in the 70K peroxisomal

[126973]membrane protein gene in Zellweger syndrome. *Nature*

Genet. 1: 16–23,

[126974]1992.

[126975]22662.

[126976]Vilain, C.; Rydlewski, C.; Duprez, L.; Heinrichs, C.;

Abramowicz,

[126977]M.; Malvaux, P.; Renneboog, B.; Parma, J.; Costagliola, S.;

Vassart,

[126978]G.: Autosomal dominant transmission of congenital thyroid hypoplasia

[126979]due to loss-of-function mutation of PAX8. J. Clin. Endocr. Metab. 86:

[126980]234–238, 2001.

[126981]22663.

[126982]Walther, C.; Guenet, J.–L.; Simon, D.; Deutsch, U.; Jostes, B.;

[126983]Goulding, M.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a murine

[126984]multigene family of paired box-containing genes. Genomics 11: 424–434,

[126985]1991.

[126986]22664.

[126987] Grueneberg, D. A.; Natesan, S.; Alexandre, C.; Gilman, M.
Z.:

[126988] Human and Drosophila homeodomain proteins that en-
hance the DNA-binding

[126989] activity of serum response factor. Science 257:
1089-1095, 1992.

[126990] 22665.

[126991] Kern, M. J.; Argao, E. A.; Birkenmeier, E. H.; Rowe, L. B.;
Potter,

[126992] S. S.: Genomic organization and chromosome localization
of the murine

[126993] homeobox gene Pmx. Genomics 19: 334-340, 1994.

[126994] 22666.

[126995] Nakamura, T.; Yamazaki, Y.; Hatano, Y.; Miura, I.: NUP98
is fused

[126996] to PMX1 homeobox gene in human acute myelogenous
leukemia with chromosome

[126997] translocation t(1;11)(q23;p15). Blood 94: 741-747, 1999.

[126998] 22667.

[126999] Norris, R. A.; Scott, K. K.; Moore, C. S.; Stetten, G.; Brown,

[127000] C. R.; Jabs, E. W.; Wulfsberg, E. A.; Yu, J.; Kern, M. J.: Hu-

man

[127001]PRRX1 and PRRX2 genes: cloning, expression, genomic localization,

[127002]and exclusion as disease genes for Nager syndrome.

Mammalian Genome 11:

[127003]1000–1005, 2000.

[127004]22668.

[127005]Riazuddin, S.; Castelein, C. M.; Ahmed, Z. M.; Lalwani, A. K.;

[127006]Mastroianni, M. A.; Naz, S.; Smith, T. N.; Liburd, N. A.; Friedman,

[127007]T. B.; Griffith, A. J.; Riazuddin, S.; Wilcox, E. R.: Dominant modifier

[127008]DFNM1 suppresses recessive deafness DFNB26. Nature Genet. 26: 431–434,

[127009]2000.

[127010]22669.

[127011]Schaller, J.: Illness resembling lupus erythematosus in mothers

[127012]of boys with chronic granulomatous disease. Ann. Intern. Med. 76:

[127013]747–750, 1972.

[127014]22670.

[127015]Theofilopoulos, A. N.; Dixon, F. J.: Murine models of systemic

[127016]lupus erythematosus. Adv. Immun. 37: 269–390, 1985.

[127017]22671.

[127018]Eng, C.; Peacocke, M.: PTEN and inherited hamartoma-cancer syndromes.

[127019](Letter) Nature Genet. 19: 223 only, 1998.

[127020]22672.

[127021]Friedl, W.; Uhlhaas, S.; Schulmann, K.; Stolte, M.; Loff, S.; Back,

[127022]W.; Mangold, E.; Stern, M.; Knaebel, H. P.; Sutter, C.; Weber, R.

[127023]G.; Pistorius, S.; Burger, B.; Propping, P.: Juvenile polyposis:

[127024]massive gastric polyposis is more common in MADH4 mutation carriers

[127025]than in BMPR1A mutation carriers. Hum. Genet. 111: 108–111, 2002.

[127026]22673.

[127027]Howe, J. R.; Bair, J. L.; Sayed, M. G.; Anderson, M. E.;

Mitros,

[127028]F. A.; Petersen, G. M.; Velculescu, V. E.; Traverso, G.; Vogelstein,

[127029]B.: Germline mutations of the gene encoding bone morphogenetic protein

[127030]receptor 1A in juvenile polyposis. *Nature Genet.* 28: 184–187, 2001.

[127031]22674.

[127032]Houlston, R.; Bevan, S.; Williams, A.; Young, J.; Dunlop, M.; Rozen,

[127033]P.; Eng, C.; Markie, D.; Woodford–Richens, K.; Rodriguez–Bigas, M.

[127034]A.; Leggett, B.; Neale, K.; Phillips, R.; Sheridan, E.; Hodgson, S.;

[127035]Iwama, T.; Eccles, D.; Bodmer, W.; Tomlinson, I.: Mutations in DPC4

[127036](SMAD4) cause juvenile polyposis syndrome, but only account for a

[127037]minority of cases. *Hum. Molec. Genet.* 7: 1907–1912, 1998.

[127038]22675.

[127039]Howe, J. R.; Ringold, J. C.; Summers, R. W.; Mitros, F. A.;

Nishimura,

[127040]D. Y.; Stone, E. M.: A gene for familial juvenile polyposis maps

[127041]to chromosome 18q21.1. Am. J. Hum. Genet. 62: 1129–1136, 1998.

[127042]22676.

[127043]Howe, J. R.; Roth, S.; Ringold, J. C.; Summers, R. W.; Jarvinen,

[127044]H. J.; Sistonen, P.; Tomlinson, I. P. M.; Houlston, R. S.; Bevan,

[127045]S.; Mitros, F. A.; Stone, E. M.; Aaltonen, L. A.: Mutations in the

[127046]SMAD4/DPC4 gene in juvenile polyposis. Science 280: 1086–1088, 1998.

[127047]22677.

[127048]Kurose, K.; Araki, T.; Matsunaka, T.; Takada, Y.; Emi, M.: Variant

[127049]manifestation of Cowden disease in Japan: hamartomatous polyposis

[127050]of the digestive tract with mutation of the PTEN gene. (Letter) Am.

[127051]J. Hum. Genet. 64: 308–310, 1999.

[127052]22678.

[127053]Nelen, M. R.; Padberg, G. W.; Peeters, E. A.; Lin, A. Y.; van

[127054]den Helm, B.; Frants, R. R.; Coulon, V.; Goldstein, A. M.;
van Reen,

[127055]M. M.; Easton, D. F.; Eeles, R. A.; Hodgson, S.; and 10 others:
ers:

[127056]Localization of the gene for Cowden disease to chromosome 10q22-23. Nature

[127057]Genet. 13: 114-116, 1996.

[127058]22679.

[127059]Olschwang, S.; Serova-Sinilnikova, O. M.; Lenoir, G. M.;
Thomas,

[127060]G.: PTEN germ-line mutations in juvenile polyposis coli.
(Letter) Nature

[127061]Genet. 18: 12-14, 1998.

[127062]22680.

[127063]Meulenbelt, I.; Williams, C. J.; Te Koppele, J. M.; Van De
Giessen,

[127064]G. C.; Slagboom, P. E.: Population haplotype analysis and
evolutionary

[127065]relations of the COI2A1 gene. Ann. Hum. Genet. 60:

189-199, 1996.

[127066]22681.

[127067]Berry, C.; Cree, J.; Mann, T.: Aarskog's syndrome. Arch.
Dis.

[127068]Child. 55: 706-710, 1980.

[127069]22682.

[127070]Escobar, V.; Weaver, D. D.: Aarskog syndrome: new find-
ings and

[127071]genetic analysis. J.A.M.A. 240: 2638-2641, 1978.

[127072]22683.

[127073]Hanley, W. B.; McKusick, V. A.; Barranco, F. T.: Osteochon-
dritis

[127074]dissecans and associated malformations in brothers: a re-
view of familial

[127075]aspects. J. Bone Joint Surg. 49A: 925-937, 1967.

[127076]22684.

[127077]O'Reilly, M. S.; Holmgren, L.; Shing, Y.; Chen, C.; Rosen-
thal,

[127078]R. A.; Moses, M.; Lane, W. S.; Cao, Y.; Sage, E. H.; Folkman,
J.:

[127079]Angiostatin: a novel angiogenesis inhibitor that mediates

the suppression

[127080]of metastases by a Lewis lung carcinoma. Cell 79:
315–328, 1994.

[127081]22685.

[127082]Waite, K. A.; Eng, C.: Protean PTEN: form and function.
Am.

[127083]J. Hum. Genet. 70: 829–844, 2002.

[127084]22686.

[127085]Armstrong, J. G.; Davies, D. R.; Guy, S. P.; Frayling, I. M.;
Evans,

[127086]D. G. R.: APC mutations in familial adenomatous polyposis
families

[127087]in the northwest of England. Hum. Mutat. 10: 376–380,
1997.

[127088]22687.

[127089]Ashton–Rickardt, P. G.; Dunlop, M. G.; Nakamura, Y.; Mor-
ris, R.

[127090]G.; Purdie, C. A.; Steel, C. M.; Evans, H. J.; Bird, C. C.; Wyl-
lie,

[127091]A. H.: High frequency of APC loss in sporadic colorectal
carcinoma

[127092]due to breaks clustered in 5q21–22. Oncogene 4:

1169-1174, 1989.

[127093]22688.

[127094]Asman, H. B.; Pierce, E. R.: Familial multiple polyposis: a statistical

[127095]study of a large Kentucky kindred. Cancer 25: 972-981, 1970.

[127096]22689.

[127097]Baker, R. H.; Heinemann, M.-H.; Miller, H. H.; DeCosse, J. J.:

[127098]Hyperpigmented lesions of the retinal pigment epithelium in familial

[127099]adenomatous polyposis. Am. J. Med. Genet. 31: 427-435, 1988.

[127100]22690.

[127101]Bapat, B.; Odze, R.; Mitri, A.; Berk, T.; Ward, M.; Gallinger,

[127102]S.: Identification of somatic APC gene mutations in peri-ampullary

[127103]adenomas in a patient with familial adenomatous polyposis (FAP). Hum.

[127104]Molec. Genet. 2: 1957-1959, 1993.

[127105]22691.

[127106]Barber, J. C. K.; Ellis, K. H.; Bowles, L. V.; Delhanty, J. D.

[127107]A.; Ede, R. F.; Male, B. M.; Eccles, D. M.: Adenomatous polyposis

[127108]coli and a cytogenetic deletion of chromosome 5 resulting from a maternal

[127109]intrachromosomal insertion. J. Med. Genet. 31: 312-316, 1994.

[127110]22692.

[127111]Bell, B.; Mazzaferri, E. L.: Familial adenomatous polyposis (Gardner's

[127112]syndrome) and thyroid carcinoma: a case report and review of the literature. Digest.

[127113]Dis. Sci. 38: 185-190, 1993.

[127114]22693.

[127115]Benecke, E.: Ueber Epitheliome auf Atheromen (Epidermoide) und

[127116]Dermoidcysten der Haut. Frankf. Z. Pathol. 42: 502-515, 1931.

[127117]22694.

[127118]Berk, T.; Cohen, Z.; Cullen, J. B.: Familial polyposis and the

[127119]role of the preventive registry. Canad. Med. Assoc. J. 124:

1427-1428,

[127120]1981.

[127121]22695.

[127122]Beroud, C.; Soussi, T.: APC gene: database of germline
and somatic

[127123]mutations in human tumors and cell lines. Nucleic Acids
Res. 24:

[127124]121-124, 1996.

[127125]22696.

[127126]Bisgaard, M. L.; Fenger, K.; Bulow, S.; Niebuhr, E.; Mohr, J.

[127127]: Familial adenomatous polyposis (FAP): frequency, pene-
trance, and

[127128]mutation rate. Hum. Mutat. 3: 121-125, 1994.

[127129]22697.

[127130]Blair, N. P.; Trempe, C. L.: Hypertrophy of the retinal pig-
ment

[127131]epithelium associated with Gardner's syndrome. Am. J.
Ophthalmol. 90:

[127132]661-667, 1980.

[127133]22698.

- [127134]Bodmer, W. F.; Bailey, C. J.; Bodmer, J.; Bussey, H. J. R.; Ellis,
lis,
[127135]A.; Gorman, P.; Lucibello, F. C.; Murday, V. A.; Rider, S. H.;
Scambler,
[127136]P.; Sheer, D.; Solomon, E.; Spurr, N. K.: Localization of the
gene
[127137]for familial adenomatous polyposis on chromosome 5.
Nature 328:
[127138]614–616, 1987.
[127139]22699.
[127140]Boland, C. R.; Sato, J.; Appelman, H. D.; Bresalier, R. S.;
Feinberg,
[127141]A. P.: Microallelotyping defines the sequence and tempo
of allelic
[127142]losses at tumour suppressor gene loci during colorectal
cancer progression. Nature
[127143]Med. 1: 902–909, 1995.
[127144]22700.
[127145]Boolbol, S. K.; Dannenberg, A. J.; Chadburn, A.; Martucci,
C.;
[127146]Guo, X.-J.; Ramonetti, J. T.; Abreu-Goris, M.; Newmark, H.
L.; Lipkin,

[127147]M. L.; DeCosse, J. J.; Bertagnolli, M. M.: Cyclooxygenase-2 overexpression

[127148]and tumor formation are blocked by sulindac in a murine model of familial

[127149]adenomatous polyposis. Cancer Res. 56: 2556-2560, 1996.

[127150]22701.

[127151]Brensinger, J. D.; Laken, S. J.; Luce, M. C.; Powell, S. M.; Vance,

[127152]G. H.; Ahnen, D. J.; Petersen, G. M.; Hamilton, S. R.; Giardiello,

[127153]F. M.: Variable phenotype of familial adenomatous polyposis in pedigrees

[127154]with 3-prime mutation in the APC gene. Gut 43: 548-552, 1998.

[127155]22702.

[127156]Bull, M. J.; Ellis, F. D.; Sato, S.; Weaver, D. D.: Hypertrophy

[127157]of retinal pigment epithelium in Gardner syndrome.

(Abstract) Proc.

[127158]Greenwood Genet. Center 4: 136 only, 1985.

[127159]22703.

[127160]Bunyan, D. J.; Shea-Simonds, J.; Reck, A. C.; Finnis, D.; Eccles,

[127161]D. M.: Genotype-phenotype correlations of new causative APC gene

[127162]mutations in patients with familial adenomatous polyposis. J. Med.

[127163]Genet. 32: 728-731, 1995.

[127164]22704.

[127165]Awata, H.; Endo, F.; Matsuda, I.: Structure of the human 4-hydroxyphenylpyruvic

[127166]acid dioxygenase gene (HPD). Genomics 23: 534-539, 1994.

[127167]22705.

[127168]Nagase, T.; Seki, N.; Tanaka, A.; Ishikawa, K.; Nomura, N.: Prediction

[127169]of the coding sequences of unidentified human genes. IV. The coding

[127170]sequences of 40 new genes (KIAA0121-KIAA0160) deduced by analysis

[127171]of cDNA clones from human cell line KG-1. DNA Res. 2: 167-174, 1995.

[127172]22706.

[127173]Parrish, J. E.; Ciccodicola, A.; Wehnert, M.; Cox, G. F.;
Chen,
[127174]E.; Nelson, D. L.: A muscle-specific DNase I-like gene in
human Xq28. Hum.
[127175]Molec. Genet. 4: 1557-1564, 1995.
[127176]22707.
[127177]Pergolizzi, R.; Appierto, V.; Bosetti, A.; DeBellis, G. L.;
Rovida,
[127178]E.; Biunno, I.: Cloning of a gene encoding a DNase I-like
endonuclease
[127179]in the human Xq28 region. Gene 168: 267-270, 1996.
[127180]22708.
[127181]Klink, A.; Schiebel, K.; Winkelmann, M.; Rao, E.; Hors-
themke, B.;
[127182]Ludecke, H.-J.; Claussen, U.; Scherer, G.; Rappold, G.: The
human
[127183]protein kinase gene PKX1 on Xp22.3 displays Xp/Yp ho-
mology and is
[127184]a site of chromosomal instability. Hum. Molec. Genet. 4:
869-878,
[127185]1995.

[127186]22709.

[127187]Schiebel, K.; Mertz, A.; Winkelmann, B.; Glaser, B.;
Schempp, W.;

[127188]Rappold, G.: FISH localization of the human Y-homolog of
protein

[127189]kinase PRKX (PRKY) to Yp11.2 and two pseudogenes to
15q26 and Xq12-q13. Cytogenet.

[127190]Cell Genet. 76: 49-52, 1997.

[127191]22710.

[127192]Brown, C. J.; Carrel, L.; Willard, H. F.: Expression of genes

[127193]from the human active and inactive X chromosomes. Am.
J. Hum. Genet. 60:

[127194]1333-1343, 1997.

[127195]22711.

[127196]Dong, B.; Horowitz, D. S.; Kobayashi, R.; Krainer, A. R.:
Purification

[127197]and cDNA cloning of HeLa cell p54(nrb), a nuclear protein
with two

[127198]RNA recognition motifs and extensive homology to human
splicing factor

[127199]PSF and Drosophila NONA/BJ6. Nucleic Acids Res. 21:

4085–4092, 1993.

[127200]22712.

[127201]Nelson, D. L.; Ballabio, A.; Cremers, F.; Monaco, A. P.;
Schlessinger,

[127202]D.: Report of the sixth international workshop on X chromosome mapping

[127203]1995. Cytogenet. Cell Genet. 71: 308–342, 1995.

[127204]22713.

[127205]Janssens, R.; Boeynaems, J.-M.; Godart, M.; Communi, D.:
Cloning

[127206]of a human heptahelical receptor closely related to the
P2Y(5) receptor. Biochem.

[127207]Biophys. Res. Commun. 236: 106–112, 1997.

[127208]22714.

[127209]O'Dowd, B. F.; Nguyen, T.; Jung, B. P.; Marchese, A.;
Cheng, R.;

[127210]Heng, H. H. Q.; Kolakowski, L. F., Jr.; Lynch, K. R.; George,
S. R.

[127211]: Cloning and chromosomal mapping of four putative
novel human G-protein-coupled

[127212]receptor genes. Gene 187: 75–81, 1997.

[127213]22715.

[127214]Bienvenu, T.; Der-Sarkissian, H.; Billuart, P.; Tissot, M.;
Des

[127215]Portes, V.; Bruls, T.; Chabrolle, J. P.; Chauveau, P.; Cherry,
M.;

[127216]Kahn, A.; Cohen, D.; Beldjord, C.; Chelly, J.; Cherif, D.:
Mapping

[127217]of the X-breakpoint involved in a balanced X;12 translo-
cation in a

[127218]female with mild mental retardation. Europ. J. Hum. Genet.
5: 105-109,

[127219]1997.

[127220]22716.

[127221]Billuart, P.; Bienvenu, T.; Ronce, N.; des Portes, V.; Vinet,
M.

[127222]C.; Zemni, R.; Crollius, H. R.; Carrie, A.; Fauchereau, F.;
Cherry,

[127223]M.; Briault, S.; Hamel, B.; Fryns, J.-P.; Beldjord, C.; Kahn,
A.;

[127224]Moraine, C.; Chelly, J.: Oligophrenin-1 encodes a rhoGAP
protein

[127225]involved in X-linked mental retardation. Nature 392:

823–826, 1998.

[127226]22717.

[127227]Kozak, M.: Point mutations define a sequence flanking the
AUG

[127228]initiator codon that modulates translation by eukaryotic
ribosomes. Cell 44:

[127229]283–292, 1986.

[127230]22718.

[127231]Tentler, D.; Gustavsson, P.; Leisti, J.; Schueler, M.; Chelly,

[127232]J.; Timonen, E.; Anneren, G.; Willard, H. F.; Dahl, N.: Dele-
tion

[127233]including the oligophrenin-1 gene associated with en-
larged cerebral

[127234]ventricles, cerebellar hypoplasia, seizures and ataxia. Eu-
rop. J.

[127235]Hum. Genet. 7: 541–548, 1999.

[127236]22719.

[127237]Greenfield, A.; Carrel, L.; Pennisi, D.; Philippe, C.; Quaderi,

[127238]N.; Siggers, P.; Steiner, K.; Tam, P. P. L.; Monaco, A. P.;
Willard,

[127239]H. F.; Koopman, P.: The UTX gene escapes X inactivation

in mice and

[127240]humans. Hum. Molec. Genet. 7: 737-742, 1998.

[127241]22720.

[127242]Greenfield, A. J.; Scott, D.; Pennisi, D.; Ehrmann, I.; Ellis,

[127243]P.; Cooper, L.; Simpson, E.; Koopman, P.: An H-YDb epitope is encoded

[127244]by a novel mouse Y chromosome gene. Nature Genet. 14: 474-478, 1996.

[127245]22721.

[127246]Lin, C.-S.; Aebersold, R. H.; Leavitt, J.: Correction of the N-terminal

[127247]sequences of the human plastin isoforms by using anchored polymerase

[127248]chain reaction: identification of a potential calcium-binding domain. Molec.

[127249]Cell. Biol. 10: 1818-1821, 1990.

[127250]22722.

[127251]Fukuda, M. N.; Sato, T.; Nakayama, J.; Klier, G.; Mikami, M.; Aoki,

[127252]D.; Nozawa, S.: Trophinin and tastin, a novel cell adhesion molecule

[127253]complex with potential involvement in embryo implantation. Genes

[127254]Dev. 9: 1199–1210, 1995.

[127255]22723.

[127256]Pack, S. D.; Tanigami, A.; Ledbetter, D. H.; Sato, T.; Fukuda,

[127257]M. N.: Assignment of trophoblast/endometrial epithelium cell adhesion

[127258]molecule trophinin gene TRO to human chromosome bands Xp11.22–p11.21

[127259]by in situ hybridization. Cytogenet. Cell Genet. 79: 123–124, 1997.

[127260]22724.

[127261]Suzuki, N.; Nakayama, J.; Shih, I. M.; Aoki, D.; Nozawa, S.; Fukuda,

[127262]M. N.: Expression of trophinin, tastin, and bystin by trophoblast

[127263]and endometrial cells in human placenta. Biol. Reprod. 60: 621–627,

[127264]1999.

[127265]22725.

[127266]Yan, M.; Wang, L.-C.; Hymowitz, S. G.; Schilbach, S.; Lee, J.;

[127267]Goddard, A.; de Vos, A. M.; Gao, W.-Q.; Dixit, V. M.: Two-amino acid

[127268]molecular switch in an epithelial morphogen that regulates binding

[127269]to two distinct receptors. Science 290: 523–527, 2000.

[127270]22726.

[127271]Sana, T. R.; Debets, R.; Timans, J. C.; Bazan, J. F.; Kastelein,

[127272]R. A.: Computational identification, cloning, and characterization

[127273]of IL1R9, a novel interleukin-1 receptor-like gene encoded over an

[127274]unusually large interval of human chromosome Xq22.2–q22.3. Genomics 69:

[127275]252–262, 2000.

[127276]22727.

[127277]Bech-Hansen, N. T.; Naylor, M. J.; Maybaum, T. A.; Sparkes, R.

[127278]L.; Koop, B.; Birch, D. G.; Bergen, A. A. B.; Prinsen, C. F. M.; Polomeno,

[127279]R. C.; Gal, A.; Drack, A. V.; Musarella, M. A.; Jacobson, S. G.; Young,

[127280]R. S. L.; Weleber, R. G.: Mutations in NYX, encoding the leucine-rich

[127281]proteoglycan nyctalopin, cause X-linked complete congenital stationary

[127282]night blindness. Nature Genet. 26: 319–323, 2000.

[127283]22728.

[127284]Pusch, C. M.; Zeitz, C.; Brandau, O.; Pesch, K.; Achatz, H.; Feil,

[127285]S.; Scharfe, C.; Maurer, J.; Jacobi, F. K.; Pinckers, A.; Andreasson,

[127286]S.; Hardcastle, A.; Wissinger, B.; Berger, W.; Meindl, A.: The complete

[127287]form of X-linked congenital stationary night blindness is caused by

[127288]mutations in a gene encoding a leucine-rich repeat protein. Nature

[127289]Genet. 26: 324–327, 2000.

[127290]22729.

[127291]Isbrandt, D.; Leicher, T.; Waldschutz, R.; Zhu, X.; Luhmann, U.;

[127292]Michel, U.; Sauter, K.; Pongs, O.: Gene structures and expression

[127293]profiles of three human KCND (Kv4) potassium channels mediating A-type

[127294]currents I(to) and I(sa). Genomics 64: 144–154, 2000.

[127295]22730.

[127296]Cao, Z.; Henzel, W.; Gao, X.: IRAK: a kinase associated with the

[127297]interleukin-1 receptor. Science 271: 1128–1131, 1996.

[127298]22731.

[127299]Kanakaraj, P.; Ngo, K.; Wu, Y.; Angulo, A.; Ghazal, P.; Harris,

[127300]C.A.; Siekierka, J. J.; Peterson, P. A.; Fung-Leung, W. P.: Defective

[127301]interleukin (IL)-18-mediated natural killer and T helper cell type

[127302]1 responses in IL-1 receptor-associated kinase (IRAK)-deficient mice. J.

[127303]Exp. Med. 189: 1129–1138, 1999.

[127304]22732.

[127305]Thomas, J. A.; Allen, J. L.; Tsen, M.; Dubnicoff, T.; Danao,

J.;

[127306]Liao, X. C.; Cao, Z.; Wasserman, S. A.: Impaired cytokine signaling

[127307]in mice lacking the IL-1 receptor-associated kinase. J. Immun. 163:

[127308]978-984, 1999.

[127309]22733.

[127310]van Vliet, J.; Turner, J.; Crossley, M.: Human Kruppel-like factor

[127311]8: a CACCC-box binding protein that associates with CtBP and represses

[127312]transcription. Nucleic Acids Res. 28: 1955-1962, 2000.

[127313]22734.

[127314]Lurquin, C.; De Smet, C.; Brasseur, F.; Muscatelli, F.; Martelange,

[127315]V.; De Plaen, E.; Brasseur, R.; Monaco, A. P.; Boon, T.: Two members

[127316]of the human MAGEB gene family located in Xp21.3 are expressed in

[127317]tumors of various histological origins. Genomics 46: 397-408, 1997.

[127318]22735.

[127319]Aguilar, M. J.; O'Brien, J. S.; Taber, P.: The syndrome of familial

[127320]leukodystrophy, adrenal insufficiency and cutaneous melanosis. In:

[127321]Aronson, S. M.; Volk, B. W.: Inborn Disorders of Sphingolipid Metabolism.

[127322]Oxford: Pergamon Press (pub.) 1967. Pp. 149–166.

[127323]22736.

[127324]Alpern, M.; Sack, G. H., Jr.; Krantz, D. H.; Jenness, J.; Zhang,

[127325]H.; Moser, H. W.: Chromosomal rearrangement segregating with adrenoleukodystrophy:

[127326]associated changes in color vision. Proc. Nat. Acad. Sci. 90: 9494–9498,

[127327]1993.

[127328]22737.

[127329]ASHG statement: Professional disclosure of familial genetic

[127330]information: the American Society of Human Genetics Social Issues

[127331]Subcommittee on Familial Disclosure. Am. J. Hum. Genet. 62: 474–483,

[127332]1998.

[127333]22738.

[127334]Aubourg, P.; Adamsbaum, C.; Lavallard-Rousseau, M.-C.;
Rocchiccioli,

[127335]F.; Cartier, N.; Jambaque, I.; Jakobezak, C.; Lemaitre, A.;
Boureau,

[127336]F.; Wolf, C.; Bougneres, P.-F.: A two-year trial of oleic and
erucic

[127337]acids ('Lorenzo's oil') as treatment for adrenomyeloneu-
ropathy. New

[127338]Eng. J. Med. 329: 745-752, 1993.

[127339]22739.

[127340]Aubourg, P.; Blanche, S.; Jambaque, I.; Rocchiccioli, F.;
Kalifa,

[127341]G.; Naud-Saudreau, C.; Rolland, M.-O.; Debre, M.; Chaus-
sain, J.-L.;

[127342]Griscelli, C.; Fischer, A.; Bougneres, P.-F.: Reversal of early
neurologic

[127343]and neuroradiologic manifestations of X-linked
adrenoleukodystrophy

[127344]by bone marrow transplantation. New Eng. J. Med. 322:
1860-1866,

[127345]1990.

[127346]22740.

[127347]Aubourg, P.; Feil, R.; Guidoux, S.; Kaplan, J.-C.; Moser, H.;
Kahn,

[127348]A.; Mandel, J.-L.: The red-green visual pigment gene re-
gion in adrenoleukodystrophy. Am.

[127349]J. Hum. Genet. 46: 459-469, 1990.

[127350]22741.

[127351]Aubourg, P.; Mosser, J.; Douar, A. M.; Sarde, C. O.; Lopez,
J.;

[127352]Mandel, J. L.: Adrenoleukodystrophy gene: unexpected
homology to

[127353]a protein involved in peroxisome biogenesis. Biochimie
75: 293-302,

[127354]1993.

[127355]22742.

[127356]Aubourg, P. R.; Sack, G. H., Jr.; Meyers, D. A.; Lease, J. J.;

[127357]Moser, H. W.: Linkage of adrenoleukodystrophy to a poly-
morphic DNA

[127358]probe. Ann. Neurol. 21: 349-352, 1987.

[127359]22743.

[127360]Aubourg, P. R.; Sack, G. H., Jr.; Moser, H.: Frequent alterations

[127361]of visual pigment genes in adrenoleukodystrophy. Am. J. Hum. Genet. 42:

[127362]408-413, 1988.

[127363]22744.

[127364]Barcelo, A.; Giros, M.; Martinez-Bermejo, A.; Sarde, C.-O.; Coll,

[127365]M. J.; Moser, J.; Mandel, J. L.; Estivill, X.; Pampols, T.: A frame

[127366]shift mutation in the candidate gene of X-linked adrenoleukodystrophy. VI.

[127367]International Congress of Inborn Errors of Metabolism. Milano, Italy May-27-31,

[127368]1994. Pp. 228 only.

[127369]22745.

[127370]Barcelo, A.; Giros, M.; Sarde, C. O.; Martinez-Bermejo, A.; Mandel,

[127371]J. L.; Pampols, T.; Estivill, X.: Identification of a new frameshift

[127372]mutation (1801delAG) in the ALD gene. Hum. Molec.

Genet. 3: 1889–1890,

[127373]1994.

[127374]22746.

[127375]Berg, K. A.; Beaty, T. H.; Raven, M. B.; Moser, A. B.; Moser,

[127376]H. W.: X-linked adrenoleukodystrophy in a 362 member
kindred. (Abstract) Am.

[127377]J. Hum. Genet. 45 (suppl.): A39, 1989.

[127378]22747.

[127379]Berger, J.; Molzer, B.; Fae, I.; Bernheimer, H.: X-linked
adrenoleukodystrophy

[127380](ALD): a novel mutation of the ALD gene in 6 members of
a family presenting

[127381]with 5 different phenotypes. Biochem. Biophys. Res. Com-
mun. 205:

[127382]1638–1643, 1994.

[127383]22748.

[127384]Bezman, L.; Moser, A. B.; Raymond, G. V.; Rinaldo, P.;
Watkins,

[127385]P. A.; Smith, K. D.; Kass, N. E.; Moser, H. W.:
Adrenoleukodystrophy:

[127386] incidence, new mutation rate, and results of extended family screening. Ann.

[127387] Neurol. 49: 512–517, 2001.

[127388] 22749.

[127389] Bezman, L.; Moser, H. W.: Incidence of X-linked adrenoleukodystrophy

[127390] and the relative frequency of its phenotypes. (Editorial) Am. J.

[127391]Med. Genet. 76: 415–419, 1998.

[127392]22750.

[127393]Blaw, M. E.: Melanodermic type leucodystrophy
(adrenoleukodystrophy).In:

[127394]Vinken, P. J.; Bruyn, G. W. (eds.): Handbook of Clinical
Neurology.

[127395]Vol. 10. Amsterdam: North Holland 1970. Pp. 128–133.

[127396]22751.

[127397]Boue, J.; Oberle, I.; Heilig, R.; Mandel, J. L.; Moser, A.;
Moser,

[127398]H.; Larsen, J. W., Jr.; Dumez, Y.; Boue, A.: First trimester
prenatal

[127399]diagnosis of adrenoleukodystrophy by determination of
very long chain

[127400]fatty acid levels and by linkage analysis to a DNA probe.
Hum. Genet. 69:

[127401]272–274, 1985.

[127402]22752.

[127403]Brady, R. O.: Inherited metabolic diseases of the nervous
system. Science 193:

[127404]733–739, 1976.

[127405]22753.

[127406]Braun, A.; Ambach, H.; Kammerer, S.; Rolinski, B.; Stockler, S.;

[127407]Rabl, W.; Gartner, J.; Zierz, S.; Roscher, A. A.: Mutations in the

[127408]gene for X-linked adrenoleukodystrophy in patients with different

[127409]clinical phenotypes. Am. J. Hum. Genet. 56: 854-861, 1995.

[127410]22754.

[127411]Jonsson, J. J.; Renieri, A.; Gallagher, P. G.; Kashtan, C. E.;

[127412]Cherniske, E. M.; Bruttini, M.; Piccini, M.; Vitelli, F.; Bal-labio,

[127413]A.; Pober, B. R.: Alport syndrome, mental retardation, midface hypoplasia,

[127414]and elliptocytosis: a new X linked contiguous gene deletion syndrome? J.

[127415]Med. Genet. 35: 273-278, 1998.

[127416]22755.

[127417]Kang, M. J.; Fujino, T.; Sasano, H.; Minekura, H.; Yabuki, N.;

[127418]Nagura, H.; Iijima, H.; Yamamoto, T. T.: A novel arachido-
nate-preferring

[127419]acyl-CoA synthetase is present in steroidogenic cells of
the rat adrenal,

[127420]ovary, and testis. Proc. Nat. Acad. Sci. 94: 2880-2884,
1997.

[127421]22756.

[127422]Meloni, I.; Muscettola, M.; Raynaud, M.; Longo, I.; Bruttini,
M.;

[127423]Moizard, M.-P.; Gomot, M.; Chelly, J.; des Portes, V.;
Fryns, J.-P.;

[127424]Ropers, H.-H.; Magi, B.; Bellan, C.; Volpi, N.; Yntema, H.
G.; Lewis,

[127425]S. E.; Schaffer, J. E.; Renieri, A.: FAHL4, encoding fatty
acid-CoA

[127426]ligase 4, is mutated in nonspecific X-linked mental retar-
dation. Nature

[127427]Genet. 30: 436-440, 2002.

[127428]22757.

[127429]Minekura, H.; Kang, M.-J.; Inagaki, Y.; Cho, Y.-Y.; Suzuki,
H.;

[127430]Fujino, T.; Yamamoto, T. T.: Exon/intron organization and

transcription

[127431]units of the human acyl-CoA synthetase 4 gene. Biochem. Biophys.

[127432]Res. Commun. 286: 80-86, 2001.

[127433]22758.

[127434]Piccini, M.; Vitelli, F.; Bruttini, M.; Pober, B. R.; Jonsson,

[127435]J. J.; Villanova, M.; Zollo, M.; Borsani, G.; Ballabio, A.; Renieri,

[127436]A.: *FACL4*, a new gene encoding long-chain acyl-CoA synthetase 4,

[127437]is deleted in a family with Alport syndrome, elliptocytosis, and mental

[127438]retardation. Genomics 47: 350-358, 1998.

[127439]22759.

[127440]Vitelli, F.; Meloni, I.; Fineschi, S.; Favara, F.; Tiziana Storlazzi,

[127441]C.; Rocchi, M.; Renieri, A.: Identification and characterization

[127442]of mouse orthologs of the *AMMECR1* and *FACL4* genes deleted in AMME

[127443]syndrome: orthology of Xq22.3 and MmuXF1-F3. Cyto-

genet. Cell Genet. 88:

[127444]259–263, 2000.

[127445]22760.

[127446]Lahn, B. T.; Page, D. C.: Functional coherence of the human Y

[127447]chromosome. Science 278: 675–680, 1997.

[127448]22761.

[127449]Chung, J.; Lee, S.–G.; Song, K.: Identification of a human homolog

[127450]of a putative RNA helicase gene (mDEAD3) expressed in mouse erythroid

[127451]cells. Korean J. Biochem. 27: 193–197, 1995.

[127452]22762.

[127453]Park, S. H.; Lee, S.–G.; Kim, Y.; Song, K.: Assignment of a human

[127454]putative RNA helicase gene, DDX3, to human X chromosome bands p11.3–p11.23. Cytogenet.

[127455]Cell Genet. 81: 178–179, 1998.

[127456]22763.

[127457]Gasper, N. J.; Kinzy, T. G.; Scherer, B. J.; Humbelin, M.;

Hershey,

[127458]J. W. B.; Merrick, W. C.: Translation initiation factor eIF-2:
cloning

[127459]and expression of the human cDNA encoding the gamma-
subunit. J. Biol.

[127460]Chem. 269: 3415-3422, 1994.

[127461]22764.

[127462]Ehrmann, I. E.; Ellis, P. S.; Mazeyrat, S.; Duthie, S.; Brock-
dorff,

[127463]N.; Mattei, M. G.; Gavin, M. A.; Affara, N. A.; Brown, G. M.;
Simpson,

[127464]E.; Mitchell, M. J.; Scott, D. M.: Characterization of genes
encoding

[127465]translation initiation factor eIF-2-gamma in mouse and
human: sex

[127466]chromosome localization, escape from X-inactivation and
evolution. Hum.

[127467]Molec. Genet. 7: 1725-1737, 1998.

[127468]22765.

[127469]Lee, S. M. Y.; Tsui, S. K. W.; Chan, K. K.; Garcia-Barcelo,
M.;

[127470]Waye, M. M. Y.; Fung, K. P.; Liew, C. C.; Lee, C. Y.: Chro-

mosomal

[127471]mapping, tissue distribution and cDNA sequence of four-
and-a-half

[127472]LIM domain protein 1 (FHL1). Gene 216: 163–170, 1998.

[127473]22766.

[127474]Morgan, M. J.; Madgwick, A. J.; Charleston, B.; Pell, J. M.;
Loughna,

[127475]P. T.: The developmental regulation of a novel muscle
LIM-protein. Biochem.

[127476]Biophys. Res. Commun. 212: 840–846, 1995.

[127477]22767.

[127478]Morgan, M. J.; Madgwick, A. J. A.: Slim defines a novel
family

[127479]of LIM-proteins expressed in skeletal muscle. Biochem.
Biophys. Res.

[127480]Commun. 225: 632–638, 1996.

[127481]22768.

[127482]Blanchet, P.; Wellemeyer, M. L.; Burton, G. V.: Retinitis
pigmentosa

[127483]following cytotoxic chemotherapy in Usher's syndrome.
Am. J. Med.

[127484]Sci. 303: 319–320, 1992.

[127485]22769.

[127486]Connor, W. E.; Weleber, R. G.; DeFrancesco, C.; Lin, D. S.;
Wolf,

[127487]D. P.: Sperm abnormalities in retinitis pigmentosa. Invest.
Ophthalm.

[127488]Vis. Sci. 38: 2619–2628, 1997.

[127489]22770.

[127490]Barnard, P. J.; Derry, J. M. J.; Ryder–Cook, A. S.; Barnard, E.

[127491]A.: Localization of the GABA(A) receptor alpha–3 subunit
gene on

[127492]the mouse X chromosome. (Abstract) Cytogenet. Cell
Genet. 51: 958

[127493]only, 1989.

[127494]22771.

[127495]Bell, M. V.; Bloomfield, J.; McKinley, M.; Patterson, M. N.;
Darlison,

[127496]M. G.; Barnard, E. A.; Davies, K. E.: Physical linkage of a
GABA–A

[127497]receptor subunit gene to the DXS374 locus in human
Xq28. Am. J. Hum.

[127498]Genet. 45: 883–888, 1989.

[127499]22772.

[127500]Buckle, V. J.; Fujita, N.; Bateson, A. N.; Darlison, M. G.;
Barnard,

[127501]E. A.: Localization of the human GABA-A3 receptor sub-
unit gene to

[127502]Xq28: a candidate gene for X-linked manic depression.
(Abstract) Cytogenet.

[127503]Cell Genet. 51: 972 only, 1989.

[127504]22773.

[127505]Buckle, V. J.; Fujita, N.; Ryder-Cook, A. S.; Derry, J. M. J.;

[127506]Barnard, P. J.; Lebo, R. V.; Schofield, P. R.; Seeburg, P. H.;
Bateson,

[127507]A. N.; Darlison, M. G.; Barnard, E. A.: Chromosomal local-
ization

[127508]of GABA-A receptor subunit genes: relationship to human
genetic disease. Neuron 3:

[127509]647–654, 1989.

[127510]22774.

[127511]Derry, J. M. J.; Barnard, P. J.: Mapping of the glycine recep-
tor

[127512]alpha-2-subunit gene and the GABA-A alpha-3-subunit gene on the mouse

[127513]X chromosome. Genomics 10: 593-597, 1991.

[127514]22775.

[127515]Buckle, V. J.; Edwards, J. H.; Evans, E. P.; Jonasson, J. A.; Lyon,

[127516]M. F.; Peters, J.; Searle, A. G.: Comparative maps of human and mouse

[127517]X chromosomes. (Abstract) Cytogenet. Cell Genet. 40: 594-595, 1985.

[127518]22776.

[127519]Mansour, I.; Delague, V.; Cazeneuve, C.; Dode, C.; Chouery, E.;

[127520]Pecheux, C.; Medlej-Hashim, M.; Salem, N.; El Zein, L.; Levan-Petit,

[127521]J.; Lefranc, G.; Goossens, M.; Delpech, M.; Amselem, S.; Loiselet,

[127522]J.; Grateau, G.; Megarbane, A.; Naman, R.: Familial Mediterranean

[127523]fever in Lebanon: mutation spectrum, evidence for cases in Maronites,

[127524]Greek Orthodoxes, Greek Catholics, Syriacs and Chiites

and for an

[127525]association between amyloidosis and M694V and M694I mutations. *Europ.*

[127526]*J. Hum. Genet.* 9: 51–55, 2001.

[127527]22777.

[127528]Matzner, Y.; Brzezinski, A.: C5a-inhibitor deficiency in peritoneal

[127529]fluids from patients with familial Mediterranean fever. *New Eng.*

[127530]*J. Med.* 311: 287–290, 1984.

[127531]22778.

[127532]Matzner, Y.; Partridge, R. E. H.; Levy, M.; Babior, B. M.: Diminished

[127533]activity of a chemotactic inhibitor in synovial fluids from patients

[127534]with familial Mediterranean fever. *Blood* 63: 629–633, 1984.

[127535]22779.

[127536]Medlej-Hashim, M.; Petit, I.; Adib, S.; Chouery, E.; Salem, N.;

[127537]Delague, V.; Rawashdeh, M.; Mansour, I.; Lefranc, G.; Na-

man, R.; Loiselet,
 [127538]J.; Lecron, J.-C.; Serre, J.-L.; Megarbane, A.: Familial
 Mediterranean
 [127539]fever: association of elevated IgD plasma levels with spe-
 cific MEFV
 [127540]mutations. *Europ. J. Hum. Genet.* 9: 849–854, 2001.
 [127541]22780.
 [127542]Medlej-Hashim, M.; Salem, N.; Chouery, E.; Rawashdeh,
 M.; Delague,
 [127543]V.; Haffar, M.; Mansour, I.; Naman, R.; Lefranc, G.; Loise-
 let, J.;
 [127544]Megarbane, A.: Familial Mediterranean fever: the potential
 for misdiagnosis
 [127545]of E148V using the E148Q usual RFLP detection method.
 (Letter) *Clin.*
 [127546]*Genet.* 61: 71–73, 2002.
 [127547]22781.
 [127548]Meyerhoff, J.: Personal Communication. Salt Lake City,
 Utah
 [127549]and Baltimore, Md. 1978.
 [127550]22782.

[127551]Meyerhoff, J.: Familial Mediterranean fever: report of a large

[127552]family, review of the literature, and discussion of the frequency

[127553]of amyloidosis. *Medicine* 59: 66–77, 1980.

[127554]22783.

[127555]Mollaret, P.: La meningite endothelio-leucocytaire multi-recurrente

[127556]benign: syndrome nouveau du maladie nouvelle? *Bull. Soc. Med. Hop.*

[127557]Paris 60: 121–122, 1944.

[127558]22784.

[127559]Ozdemir, A. I.; Sokmen, C.: Familial Mediterranean fever among

[127560]the Turkish people. *Am. J. Gastroent.* 51: 311–316, 1969.

[127561]22785.

[127562]Ozyilkan, E.; Simsek, H.; Telatar, H.: Absence of asthma in patients

[127563]with familial Mediterranean fever. *Israel J. Med. Sci.* 30: 237–238,

[127564]1994.

[127565]22786.

[127566]Papin, S.; Duquesnoy, P.; Cazeneuve, C.; Pantel, J.;
Coppey-Moisan,

[127567]M.; Dargemont, C.; Amselem, S.: Alternative splicing at the
MEFV

[127568]locus involved in familial Mediterranean fever regulates
translocation

[127569]of the marenostriin/pyrin protein to the nucleus. Hum.
Molec. Genet. 9:

[127570]3001–3009, 2000.

[127571]22787.

[127572]Pras, E.; Aksentijevich, I.; Gruberg, L.; Balow, J. E., Jr.;
Prosen,

[127573]L.; Dean, M.; Steinberg, A. D.; Pras, M.; Kastner, D. L.:
Mapping

[127574]of a gene causing familial Mediterranean fever to the short
arm of

[127575]chromosome 16. New Eng. J. Med. 326: 1509–1513,
1992.

[127576]22788.

[127577]Pras, E.; Aksentijevich, I.; Gruberg, L.; Shen, Y.; Prosen, L.;

[127578]Balow, J. E., Jr.; Richards, R. I.; Pras, M.; Kastner, D. L.: Lo-
cus

[127579]homogeneity in familial Mediterranean fever. (Abstract)
Am. J. Hum.

[127580]Genet. 51 (suppl.): A371, 1992.

[127581]22789.

[127582]Pras, E.; Aksentijevich, I.; Levy, E.; Gruberg, L.; Prosen, L.;

[127583]Dean, M.; Pras, M.; Kastner, D. L.: The gene causing famil-
ial Mediterranean

[127584]fever maps to the short arm of chromosome 16 in Druze
and Moslem Arab

[127585]families. Hum. Genet. 94: 576-577, 1994.

[127586]22790.

[127587]Pras, E.; Livneh, A.; Balow, J. E., Jr.; Pras, E.; Kastner, D.

[127588]L.; Pras, M.; Langevitz, P.: Clinical differences between
North African

[127589]and Iraqi Jews with familial Mediterranean fever. Am. J.
Med. Genet. 75:

[127590]216-219, 1998.

[127591]22791.

[127592]Pras, M.; Bronshpigel, N.; Zemer, D.; Gafni, J.: Variable incidence
of amyloidosis in familial Mediterranean fever among different ethnic
groups. Johns Hopkins Med. J. 150: 22–26, 1982.
22792.

[127596]Ravid, M.; Robson, M.; Kedar, I.: Prolonged colchicine
treatment
in four patients with amyloidosis. Ann. Intern. Med. 87:
568–570,
1977.

[127599]22793.

[127600]Rawashdeh, M. O.; Majeed, H. A.: Familial Mediterranean
fever
in Arab children: the high prevalence and gene frequency.
Europ.
J. Pediat. 155: 540–544, 1996.

[127603]22794.

[127604]Reimann, H. A.; Moadie, J.; Semerdijian, S.; Sahyoun, P. F.:
Periodic peritonitis--heredity and pathology: report of
seventy-two

[127606]cases. J.A.M.A. 154: 1254–1259, 1954.

[127607]22795.

[127608]Rogers, D. B.; Shohat, M.; Petersen, G. M.; Bickal, J.; Congleton,

[127609]J.; Schwabe, A. D.; Rotter, J. I.: Familial Mediterranean fever in

[127610]Armenians: autosomal recessive inheritance with high gene frequency. Am.

[127611]J. Med. Genet. 34: 168–172, 1989.

[127612]22796.

[127613]Rubinger, D.; Friedlaender, M. M.; Popovtzer, M. M.: Amelioration

[127614]of familial Mediterranean fever during hemodialysis. New Eng. J.

[127615]Med. 301: 142–144, 1979.

[127616]22797.

[127617]Saatci, U.; Ozen, S.; Ozdemir, S.; Bakkaloglu, A.; Besbas, N.;

[127618]Topaloglu, R.; Arslan, S.: Familial Mediterranean fever in children:

[127619]report of a large series and discussion of the risk and

prognostic

[127620]factors of amyloidosis. *Europ. J. Pediat.* 156: 619–623, 1997.

[127621]22798.

[127622]Sack, G. H., Jr.; Talbot, C. C., Jr.; McCarthy, B. G.; Harris,

[127623]E. L.; Kastner, D.; Gruberg, L.; Pras, M.: Exclusion of linkage between

[127624]familial Mediterranean fever and the human serum amyloid A (SAA) gene

[127625]cluster. *Hum. Genet.* 87: 506–508, 1991.

[127626]22799.

[127627]Schaner, P.; Richards, N.; Wadhwa, A.; Aksentijevich, I.; Kastner,

[127628]D.; Tucker, P.; Gumucio, D.: Episodic evolution of pyrin in primates:

[127629]human mutations recapitulate ancestral amino acid states. *Nature*

[127630]*Genet.* 27: 318–321, 2001.

[127631]22800.

[127632]Schlesinger, M.; Ilfeld, D. N.; Zamir, R.; Brautbar, C.: Familial

- [127633]Mediterranean fever: no linkage with HLA. Tissue Antigens
24: 65-66,
[127634]1984.
[127635]22801.
- [127636]Schwabe, A. D.; Lehman, T. J. A.: C5a-inhibitor deficiency-
-a
[127637]role in familial Mediterranean fever? (Editorial) New Eng. J.
Med. 311:
[127638]325-326, 1984.
[127639]22802.
- [127640]Schwabe, A. D.; Monroe, J. B.: Meningitis in familial
Mediterranean
[127641]fever. Am. J. Med. 85: 715-717, 1988.
[127642]22803.
- [127643]Schwabe, A. D.; Nishizawa, A.: Recurrent polyserositis
(familial
[127644]Mediterranean fever) in a Japanese. Jpn. J. Med. 26:
370-372, 1987.
[127645]22804.
- [127646]Schwabe, A. D.; Peters, R. S.: Familial Mediterranean fever
in

[127647]Armenians: analysis of 100 cases. *Medicine* 53: 453-462, 1974.

[127648]22805.

[127649]Schwabe, A. D.; Terasaki, P. I.; Barnett, E. V.; Territo, M. C.;

[127650]Klinenberg, J. R.; Peters, R. S.: Familial Mediterranean fever--recent

[127651]advances in pathogenesis and management. *West. J. Med.* 127: 15-23,

[127652]1977.

[127653]22806.

[127654]Chern, C. J.; Beutler, E.: Biochemical and electrophoretic studies

[127655]of erythrocyte pyridoxine kinase in white and black Americans. *Am.*

[127656]*J. Hum. Genet.* 28: 9-17, 1976.

[127657]22807.

[127658]Hanna, M. C.; Turner, A. J.; Kirkness, E. F.: Human pyridoxal

[127659]kinase: cDNA cloning, expression, and modulation by ligands of the

[127660]benzodiazepine receptor. J. Biol. Chem. 272:
10756–10760, 1997.

[127661]22808.

[127662]Dougherty, K. M.; Brandriss, M. C.; Valle, D.: Cloning hu-
man pyrroline–5–carboxylate

[127663]reductase cDNA by complementation in *Saccharomyces*
cerevisiae. J.

[127664]Biol. Chem. 267: 871–875, 1992.

[127665]22809.

[127666]Merrill, M. J.; Yeh, G. C.; Phang, J. M.: Purified human ery-
throcyte

[127667]pyrroline–5–carboxylate reductase: preferential oxidation
of NADPH. J.

[127668]Biol. Chem. 264: 9352–9358, 1989.

[127669]22810.

[127670]Bigley, R. H.; Stenzel, P.; Jones, R. T.; Campos, J. O.; Koler,

[127671]R. D.: Tissue distribution of human pyruvate kinase
isozymes. Enzym.

[127672]Biol. Clin. 9: 10–20, 1968.

[127673]22811.

[127674]Bunn, H. F.: Personal Communication. Boston, Mass.
4/30/1981.

[127675]22812.

[127676]Chern, C. J.; Kennett, R.; Engel, L. E.; Mellman, W. J.;
Croce,

[127677]C. M.: Assignment of the structural genes for the alpha
subunit of

[127678]hexosaminidase A, mannosephosphate isomerase and
pyruvate kinase to

[127679]the region q22-qter of human chromosome 15. Somat.
Cell Genet. 3:

[127680]553-560, 1977.

[127681]22813.

[127682]Junien, C.; Robinson-Skala, H.; Dreyfus, J. C.; Ravise, N.;
Boue,

[127683]J.; Boue, A.; Kaplan, J. C.: PK3: a new chromosome enzyme
marker

[127684]for gene dosage studies in chromosome 15 imbalance.
Hum. Genet. 54:

[127685]191-196, 1980.

[127686]22814.

- [127687]Kahn, A.; Marie, J.; Garreau, H.; Sprengers, E. D.: Subunit structure,
[127688]interrelations and kinetic characteristics of the pyruvate kinase
[127689]from erythrocytes and liver. Biochim. Biophys. Acta 523: 58–74,
[127690]1978.
[127691]22815.
- [127692]Levine, M.; Muirhead, H.; Stammers, D. K.; Stuart, D. I.: Structure
[127693]of pyruvate kinase and similarities with other enzymes: possible implications
[127694]for protein taxonomy and evolution. Nature 271: 626–630, 1978.
[127695]22816.
- [127696]Nakashima, K.; Miwa, S.; Fujii, H.; Shinohara, K.; Yamauchi, K.;
[127697]Tsuji, Y.; Yanai, M.: Characterization of pyruvate kinase, PK Nagasaki. J.
[127698]Lab. Clin. Med. 90: 1012–1020, 1977.
[127699]22817.

- [127700]Ritter, H.; Friedrichson, U.; Schmitt, J.: Genetic variation of
[127701]mannose phosphate isomerase in man. Humangenetik 22:
261 only, 1974.
[127702]22818.
- [127703]Sheffield, V. C.; Fishman, G. A.; Beck, J. S.; Kimura, A. E.;
[127704]Stone, E. M.: Identification of novel rhodopsin mutations
associated
[127705]with retinitis pigmentosa by GC-clamped denaturing gra-
dient gel electrophoresis. Am.
[127706]J. Hum. Genet. 49: 699-706, 1991.
[127707]22819.
- [127708]Sieving, P. A.; Richards, J. E.; Bingham, E. L.; Naarendorp,
F.
[127709]: Dominant congenital complete nyctalopia and
gly90-to-asn rhodopsin
[127710]mutation. (Abstract) Invest. Ophthalm. Vis. Sci. 33: 1397,
1992.
[127711]22820.
- [127712]Sieving, P. A.; Richards, J. E.; Naarendorp, F.; Bingham, E.
L.;
[127713]Scott, K.; Alpern, M.: Dark-light: model for nightblindness

from

[127714]the human rhodopsin gly90-to-asp mutation. Proc. Nat.
Acad. Sci. 92:

[127715]880-884, 1995.

[127716]22821.

[127717]Sohocki, M. M.; Daiger, S. P.; Bowne, S. J.; Rodriquez, J. A.;

[127718]Northrup, H.; Heckenlively, J. R.; Birch, D. G.; Mintz-Hit-
tner, H.;

[127719]Ruiz, R. S.; Lewis, R. A.; Saperstein, D. A.; Sullivan, L. S.:
Prevalence

[127720]of mutations causing retinitis pigmentosa and other in-
herited retinopathies. Hum.

[127721]Mutat. 17: 42-51, 2001.

[127722]22822.

[127723]Souied, E.; Soubrane, G.; Benlian, P.; Coscas, G. J.; Gerber,

[127724]S.; Munnich, A.; Kaplan, J.: Retinitis punctata albescens
associated

[127725]with the arg135-to-trp mutation in the rhodopsin gene.
Am. J. Ophthal. 121:

[127726]19-25, 1996.

[127727]22823.

[127728]Sullivan, L. J.; Makris, G. S.; Dickinson, P.; Mulhall, L. E.

[127729]M.; App, D.; Forrest, S.; Cotton, R. G. H.; Loughnan, M. S.:

A new

[127730]codon 15 rhodopsin gene mutation in autosomal dominant
retinitis pigmentosa

[127731]is associated with sectorial disease. Arch. Ophthalmol. 111:
1512-1517,

[127732]1993.

[127733]22824.

[127734]Sung, C.-H.; Davenport, C. M.; Hennessey, J. C.; Maumenee, I.

[127735]H.; Jacobson, S. G.; Heckenlively, J. R.; Nowakowski, R.;
Fishman,

[127736]G.; Gouras, P.; Nathans, J.: Rhodopsin mutations in autosomal dominant

[127737]retinitis pigmentosa. Proc. Nat. Acad. Sci. 88: 6481-6485,
1991.

[127738]22825.

[127739]Sung, C.-H.; Davenport, C. M.; Nathans, J.: Rhodopsin
mutations

[127740]responsible for autosomal dominant retinitis pigmentosa:
clustering

[127741]of functional classes along the polypeptide chain. J. Biol.
Chem. 268:

[127742]26645–26649, 1993.

[127743]22826.

[127744]Sung, C.–H.; Makino, C.; Baylor, D.; Nathans, J.: A
rhodopsin

[127745]gene mutation responsible for autosomal dominant retini-
tis pigmentosa

[127746]results in a protein that is defective in localization to the
photoreceptor

[127747]outer segment. J. Neurosci. 14: 5818–5833, 1994.

[127748]22827.

[127749]Sung, C.–H.; Schneider, B. G.; Agarwal, N.; Papermaster, D.
S.;

[127750]Nathans, J.: Functional heterogeneity of mutant
rhodopsins responsible

[127751]for autosomal dominant retinitis pigmentosa. Proc. Nat.
Acad. Sci. 88:

[127752]8840–8844, 1991.

[127753]22828.

[127754]Vaithinathan, R.; Berson, E. L.; Dryja, T. P.: Further screen-

ing

[127755] of the rhodopsin gene in patients with autosomal dominant retinitis

[127756] pigmentosa. Genomics 21: 461–463, 1994.

[127757] 22829.

[127758] Weleber, R. G.; Pillers, D. M.; Powell, B. R.; Hanna, C. E.; Magenis,

[127759] R. E.; Buist, N. R. M.: Aland Island disease (Forsius–Eriksson syndrome)

[127760] associated with contiguous gene syndrome at Xp21: similarity to incomplete

[127761] congenital stationary night blindness. Arch. Ophthal. 107: 1170–1179,

[127762] 1989.

[127763] 22830.

[127764] Boehm, T.; Foroni, L.; Kaneko, Y.; Perutz, M. F.; Rabbitts, T.

[127765] H.: The rhombotin family of cysteine-rich LIM-domain oncogenes: distinct

[127766] members are involved in T-cell translocations to human chromosomes

[127767] 11p15 and 11p13. Proc. Nat. Acad. Sci. 88: 4367–4371,

1991.

[127768]22831.

[127769]Royer-Pokora, B.; Loos, U.; Ludwig, W.-D.: TTG-2, a new
gene encoding

[127770]a cysteine-rich protein with the LIM motif, is overex-
pressed in acute

[127771]T-cell leukaemia with the t(11;14)(p13;q11). Oncogene 6:
1887-1893,

[127772]1991.

[127773]22832.

[127774]Yamada, Y.; Pannell, R.; Forster, A.; Rabbitts, T. H.: The
oncogenic

[127775]LIM-only transcription factor Lmo2 regulates angiogenesis
but not

[127776]vasculogenesis in mice. Proc. Nat. Acad. Sci. 97: 320-324,
2000.

[127777]22833.

[127778]Yamada, Y.; Warren, A. J.; Dobson, C.; Forster, A.; Pannell,
R.;

[127779]Rabbitts, T. H.: The T cell leukemia LIM protein Lmo2 is
necessary

[127780]for adult mouse hematopoiesis. Proc. Nat. Acad. Sci. 95:

3890-3895,

[127781]1998.

[127782]22834.

[127783]Yang-Feng, T. L.; Thelander, L.; Lewis, W. H.; Srinivasan,
P. R.;

[127784]Francke, U.: Gene localization of the ribonucleotide reduc-
tase M2

[127785]subunit and of related and co-amplified sequences on hu-
man and mouse

[127786]chromosomes. (Abstract) Am. J. Hum. Genet. 39: A174
only, 1986.

[127787]22835.

[127788]Lindgren, V.; Bryke, C. R.; Ozcelik, T.; Yang-Feng, T. L.;
Francke,

[127789]U.: Phenotypic, cytogenetic, and molecular studies of
three patients

[127790]with constitutional deletions of chromosome 5 in the re-
gion of the

[127791]gene for familial adenomatous polyposis. Am. J. Hum.
Genet. 50:

[127792]988-997, 1992.

[127793]22836.

[127794]Barletta, C.; Druck, T.; LaForgia, S.; Calabretta, B.;
Drabkin,
[127795]H.; Patterson, D.; Croce, C. M.; Huebner, K.: Chromosome
locations
[127796]of the MYB related genes, AMYB and BMYB. Cancer Res.
51: 3821-3824,
[127797]1991.
[127798]22837.
[127799]Nomura, N.; Takahashi, M.; Matsui, M.; Ishii, S.; Date, T.;
Sasamoto,
[127800]S.; Ishizaki, R.: Isolation of human cDNA clones of MYB-
related genes,
[127801]A-MYB and B-MYB. Nucleic Acids Res. 16: 11075-11089,
1988.
[127802]22838.
[127803]Takahashi, T.; Nakagoshi, H.; Sarai, A.; Nomura, N.; Ya-
mamoto,
[127804]T.; Ishii, S.: Human A-myb gene encodes a transcriptional
activator
[127805]containing the negative regulatory domains. FEBS Lett.
358: 89-96,
[127806]1995.

[127807]22839.

[127808]Boylan, K. B.; Takahashi, N.; Diamond, M.; Hood, L. E.;
Prusiner,

[127809]S. B.: DNA length polymorphism located 5-prime to the
human myelin

[127810]basic protein gene. Am. J. Hum. Genet. 40: 387-400,
1987.

[127811]22840.

[127812]Dayhoff, M. O.: Atlas of Protein Sequence and Structure.
Myelin

[127813]membrane encephalitogenic protein. Washington: National
Biomedical

[127814]Research Foundation (pub.) 5: 1972. Pp. D324 only.

[127815]22841.

[127816]Eylar, E. H.; Brostoff, S.; Hashim, G.; Westall, F. C.: Basic

[127817]A1 protein of the myelin membrane: the complete amino
acid sequence. J.

[127818]Biol. Chem. 246: 5770-5784, 1971.

[127819]22842.

[127820]Gomez, C. M.; Muggleton-Harris, A. L.; Whittingham, D.
G.; Hood,

[127821]L. E.; Readhead, C.: Rapid preimplantation detection of mutant (shiverer) and normal alleles of the mouse myelin basic protein gene allowing selective implantation and birth of live young. Proc. Nat. Acad. Sci. 87: 4481–4484, 1990.

[127822]22843.

[127823]Kamholz, J.; de Ferra, F.; Puckett, C.; Lazzarini, R.: Identification of three forms of human myelin basic protein by cDNA cloning. Proc. Nat. Acad. Sci. 83: 4962–4966, 1986.

[127824]22844.

[127825]Kamholz, J.; Spielman, R.; Gogolin, K.; Modi, W.; O'Brien, S.; Lazzarini, R.: The human myelin–basic–protein gene: chromosomal localization and RFLP analysis. Am. J. Hum. Genet. 40: 365–373, 1987.

[127826]22845.

[127827]Lalley, P. A.; McKusick, V. A.: Report of the committee on

comparative

[127835]mapping (HGM8). Cytogenet. Cell Genet. 40: 536–566,
1985.

[127836]22846.

[127837]Martensen, R. E.: Myelin basic protein speciation. Prog.
Clin.

[127838]Biol. Res. 146: 511–521, 1984.

[127839]22847.

[127840]Marty, M. C.; Alliot, F.; Rutin, J.; Fritz, R.; Trisler, D.; Pes-
sac,

[127841]B.: The myelin basic protein gene is expressed in differen-
tiated

[127842]blood cell lineages and in hemopoietic progenitors. Proc.
Nat. Acad.

[127843]Sci. 99: 8856–8861, 2002.

[127844]22848.

[127845]Molineaux, S. M.; Engh, H.; de Ferra, F.; Hudson, L.; Laz-
zarini,

[127846]R. A.: Recombination within the myelin basic protein gene
created

[127847]the dysmyelinating shiverer mouse mutation. Proc. Nat.

Acad. Sci. 83:

[127848]7542–7546, 1986.

[127849]22849.

[127850]Popko, B.; Puckett, C.; Lai, E.; Shine, H. D.; Readhead, C.;
Takahashi,

[127851]N.; Hunt, S. W., III; Sidman, R. L.; Hood, L.: Myelin deficient
mice:

[127852]expression of myelin basic protein and generation of mice
with varying

[127853]levels of myelin. Cell 48: 713–721, 1987.

[127854]22850.

[127855]Readhead, C.; Popko, B.; Takahashi, N.; Shine, H. D.;
Saavedra,

[127856]R. A.; Sidman, R. L.; Hood, L.: Expression of a myelin basic
protein

[127857]gene in transgenic shiverer mice: correction of the dys-
myelinating

[127858]phenotype. Cell 48: 703–712, 1987.

[127859]22851.

[127860]Roach, A.; Boylan, K.; Horvath, S.; Prusiner, S. B.; Hood, L.

[127861]E.: Characterization of cloned cDNA representing rat

myelin basic

[127862]protein: absence of expression in shiverer mutant mice.

Cell 34:

[127863]799–806, 1983.

[127864]22852.

[127865]Roach, A.; Takahashi, N.; Pravtcheva, D.; Ruddle, F.; Hood, L.

[127866]: Chromosomal mapping of mouse myelin basic protein gene and structure

[127867]and transcription of the partially deleted gene in shiverer mutant

[127868]mice. Cell 42: 149–155, 1985.

[127869]22853.

[127870]Saxe, D. F.; Takahashi, N.; Hood, L.; Simon, M. I.: Localization

[127871]of the human myelin basic protein gene (MBP) to region 18q22–qter

[127872]by in situ hybridization. Cytogenet. Cell Genet. 39: 246–249, 1985.

[127873]22854.

[127874]Sheremata, W.; Cosgrove, J. B. R.; Hylar, E. H.: Cellular hy-

persensitivity

[127875]to basic myelin (A1) protein and clinical multiple sclerosis.

New

[127876]Eng. J. Med. 291: 14-17, 1974.

[127877]22855.

[127878]Sidman, R.: Personal Communication. Boston, Mass. 1983.

[127879]22856.

[127880]Sidman, R. L.; Conover, C. S.; Carson, J. H.: Shiverer gene
maps

[127881]near the distal end of chromosome 18 in the house
mouse. Cytogenet.

[127882]Cell Genet. 39: 241-245, 1985.

[127883]22857.

[127884]Sparkes, R. S.; Mohandas, T.; Heinzmann, C.; Roth, H. J.;
Klisak,

[127885]I.; Campagnoni, A. T.: Assignment of the myelin basic
protein gene

[127886]to human chromosome 18q22-qter. Hum. Genet. 75:
147-150, 1987.

[127887]22858.

[127888]Takahashi, N.; Roach, A.; Teplow, D. B.; Prusiner, S. B.;

Hood,

[127889]L.: Cloning and characterization of the myelin basic protein gene

[127890]from mouse: one gene can encode both 14 kd and 18.5 kd MBPs by alternate

[127891]use of exons. Cell 42: 139–148, 1985.

[127892]22859.

[127893]Eiberg, H.; Mohr, J.; Nielsen, L. S.; Simonsen, N.: Genetics

[127894]and linkage relationships of the C3 polymorphism: discovery of C3–Se

[127895]linkage and assignment of LES–C3–DM–Se–PEPD–Lu synteny to chromosome

[127896]19. Clin. Genet. 24: 159–170, 1983.

[127897]22860.

[127898]Barouch, L. A.; Harrison, R. W.; Skaf, M. W.; Rosas, G. O.; Cappola,

[127899]T. P.; Kobeissi, Z. A.; Hobai, I. A.; Lemmon, C. A.; Burnett, A. L.;

[127900]O'Rourke, B.; Rodriguez, E. R.; Huang, P. L.; Lima, J. A. C.; Berkowitz,

[127901]D. E.; Hare, J. M.: Nitric oxide regulates the heart by spa-

tial confinement

[127902]of nitric oxide synthase isoforms. Nature 416: 337–340,
2002.

[127903]22861.

[127904]Dimmeler, S.; Fleming, I.; Fisslthaler, B.; Hermann, C.;
Busse,

[127905]R.; Zeiher, A. M.: Activation of nitric oxide synthase in en-
dothelial

[127906]cells by Akt-dependent phosphorylation. Nature 399:
601–605, 1999.

[127907]22862.

[127908]Fulton, D.; Gratton, J.-P.; McCabe, T. J.; Fontana, J.; Fujio,

[127909]Y.; Walsh, K.; Franke, T. F.; Papapetropoulos, A.; Sessa, W.
C.:

[127910]Regulation of endothelium-derived nitric oxide production
by the protein

[127911]kinase Akt. Nature 399: 597–601, 1999.

[127912]22863.

[127913]Kruh, G. D.; King, C. R.; Kraus, M. H.; Popescu, N. C.;
Amsbaugh,

[127914]S. C.; McBride, W. O.; Aaronson, S. A.: A novel human gene

closely

[127915]related to the abl proto-oncogene. Science 234:
1545-1548, 1986.

[127916]22864.

[127917]Audrezet, M.-P.; Chen, J.-M.; Le Marechal, C.;
Ruszniewski, P.;

[127918]Robaszkiewicz, M.; Raguene, O.; Quere, I.; Scotet, V.;
Ferec, C.

[127919]: Determination of the relative contribution of three
genes--the cystic

[127920]fibrosis transmembrane conductance regulator gene, the
cationic trypsinogen

[127921]gene, and the pancreatic secretory trypsin inhibitor gene--
to the

[127922]etiology of idiopathic chronic pancreatitis. Europ. J. Hum.
Genet. 10:

[127923]100-106, 2002.

[127924]22865.

[127925]Arnaud, J.; Vavrusa, B.; Sevin, J.; Constans, J.: Human red-
cell

[127926]acid phosphatase (ACP1): a new mutant (ACP1*KUK) de-
tected by isoelectric

[127927] focusing, kinetics of thermostability and substrate activity.

Hum.

[127928] Hered. 39: 288–293, 1989.

[127929] 22866.

[127930] Arnaud, J.; Vavrusa, B.; Wiederanders, G.; Constans, J.:

Human

[127931] red-cell acid phosphatase (ACP1): kinetic and thermodynamic characterization

[127932] of the KUK variant. Hum. Hered. 42: 140–142, 1992.

[127933] 22867.

[127934] Beemer, F. A.; van der Heiden, C.; Van Hemel, J. O.;

Jansen, M.

[127935]: Letter to the editors. (Letter) Clin. Genet. 24: 151, 1983.

[127936] 22868.

[127937] Berg, K.: Close linkage between APOB and ACP1 excluded.

(Abstract) Cytogenet.

[127938] Cell Genet. 46: 580, 1987.

[127939] 22869.

[127940] Bottini, E.; Carapella, E.; Orzalesi, M.; Lucarelli, P.; Pas-

cone,

[127941]R.; Gloria-Bottini, F.; Coccia, M.: Is there a role of erythrocyte

[127942]acid phosphatase polymorphism in intrauterine development? (Letter) Am.

[127943]J. Hum. Genet. 32: 764-767, 1980.

[127944]22870.

[127945]Seldin, M. F.; Kruh, G. D.: Mapping of Abll within a conserved

[127946]linkage group on distal mouse chromosome 1 syntenic with human chromosome

[127947]1 using an interspecific cross. Genomics 4: 221-223, 1989.

[127948]22871.

[127949]Bello, M. J.; Salagnon, N.; Rey, J. A.; Guichaoua, M. R.; Berge-Lefranc,

[127950]J. L.; Jordan, B. R.; Luciani, J. M.: Precise in situ localization

[127951]of NCAM, ETS1, and D11S29 on human meiotic chromosomes. Cytogenet.

[127952]Cell Genet. 52: 7-10, 1989.

[127953]22872.

[127954]Concannon, P.; Malhotra, U.; Charmley, P.; Reynolds, J.;

Lange,

[127955]K.; Gatti, R. A.: The ataxia-telangiectasia gene (ATA) on chromosome

[127956]11 is distinct from the ETS-1 gene. Am. J. Hum. Genet. 46: 789-794,

[127957]1990.

[127958]22873.

[127959]Gold, D. P.; van Dongen, J. J. M.; Morton, C. C.; Bruns, G. A.

[127960]P.; van den Elsen, P.; Geurts van Kessel, A. H. M.; Terhorst, C.:

[127961]The gene encoding the epsilon subunit of the T3/T-cell receptor complex

[127962]maps to chromosome 11 in humans and to chromosome 9 in mice. Proc.

[127963]Nat. Acad. Sci. 84: 1664-1668, 1987.

[127964]22874.

[127965]Ohtani, N.; Zebedee, Z.; Huot, T. J. G.; Stinson, J. A.; Sugimoto,

[127966]M.; Ohashi, Y.; Sharrocks, A. D.; Peters, G.; Hara, E.: Opposing

[127967]effects of Ets and Id proteins on p16(INK4A) expression

during cellular

[127968]senescence. Nature 409: 1067–1070, 2001.

[127969]22875.

[127970]Sacchi, N.; Watson, D. K.; Geurts van Kessel, A. H. M.;
Hagemeijer,

[127971]A.; Kersey, J.; Drabkin, H. D.; Patterson, D.; Papas, T. S.:
Hu-ets-1

[127972]and Hu-ets-2 genes are transposed in acute leukemias
with (4;11) and

[127973](8;21) translocations. Science 231: 379–382, 1986.

[127974]22876.

[127975]Watson, D. K.; McWilliams-Smith, M. J.; Kozak, C.; Reeves,
R.;

[127976]Gearhart, J.; Nunn, M. F.; Nash, W.; Fowle, J. R., III; Dues-
berg,

[127977]P.; Papas, T. S.; O'Brien, S. J.: Conserved chromosomal po-
sitions

[127978]of dual domains of the ets protooncogene in cats, mice,
and humans. Proc.

[127979]Nat. Acad. Sci. 83: 1792–1796, 1986.

[127980]22877.

[127981]Watson, D. K.; McWilliams-Smith, M. J.; Nunn, M. F.; Duesberg,

[127982]P. H.; O'Brien, S. J.; Papas, T. S.: The ets sequence from the transforming

[127983]gene of avian erythroblastosis virus, E26, has unique domains on human

[127984]chromosomes 11 and 21: both loci are transcriptionally active. Proc.

[127985]Nat. Acad. Sci. 82: 7294-7298, 1985.

[127986]22878.

[127987]Bellacosa, A.; Franke, T. F.; Gonzalez-Portal, M. E.; Datta, K.;

[127988]Taguchi, T.; Gardner, J.; Cheng, J. Q.; Testa, J. R.; Tsichlis, P.

[127989]N.: Structure, expression and chromosomal mapping of c-akt: relationship

[127990]to v-akt and its implications. Oncogene 8: 745-754, 1993.

[127991]22879.

[127992]Brunet, A.; Bonni, A.; Zigmund, M. J.; Lin, M. Z.; Juo, P.; Hu,

[127993]L. S.; Anderson, M. J.; Arden, K. C.; Blenis, J.; Greenberg,

M. E.

[127994]: Akt promotes cell survival by phosphorylating and inhibiting a Forkhead

[127995]transcription factor. Cell 96: 857–868, 1999.

[127996]22880.

[127997]Chen, W. S.; Xu, P.-Z.; Gottlob, K.; Chen, M.-L.; Sokol, K.; Shiyanova,

[127998]T.; Roninson, I.; Weng, W.; Suzuki, R.; Tobe, K.; Kadowaki, T.; Hay,

[127999]N.: Growth retardation and increased apoptosis in mice with homozygous

[128000]disruption of the akt1 gene. Genes Dev. 15: 2203–2208, 2001.

[128001]22881.

[128002]Condorelli, G.; Drusco, A.; Stassi, G.; Bellacosa, A.; Roncarati,

[128003]R.; Iaccarino, G.; Russo, M. A.; Gu, Y.; Dalton, N.; Chung, C.; Latronico,

[128004]M. V. G.; Napoli, C.; Sadoshima, J.; Croce, C. M.; Ross, J., Jr.:

[128005]Akt induces enhanced myocardial contractility and cell size in vivo

[128006]in transgenic mice. Proc. Nat. Acad. Sci. 99:
12333–12338, 2002.

[128007]22882.

[128008]Delettre, C.; Lenaers, G.; Griffoin, J.–M.; Gigarel, N.;
Lorenzo,
[128009]C.; Belenguer, P.; Pelloquin, L.; Grosgeorge, J.; Turc–Carel,
C.;

[128010]Perret, E.; Astarie–Dequeker, C.; Lasquelllec, L.; Arnaud, B.;
Ducommun,
[128011]B.; Kaplan, J.; Hamel, C. P.: Nuclear gene OPA1, encoding a
mitochondrial
[128012]dynamin–related protein, is mutated in dominant optic at–
rophy. Nature
[128013]Genet. 26: 207–210, 2000.

[128014]22883.

[128015]Curry, C. J. R.; Magenis, R. E.; Brown, M.; Lanman, J. T., Jr.;

[128016]Tsai, J.; O'Lague, P.; Goodfellow, P.; Mohandas, T.;

Bergner, E. A.;

[128017]Shapiro, L. J.: Inherited chondrodysplasia punctata due to
a deletion
[128018]of the terminal short arm of an X chromosome. New Eng.
J. Med. 311:

[128019]1010–1015, 1984.

[128020]22884.

[128021]Bick, D.; Curry, C. J. R.; McGill, J. R.; Schorderet, D. F.; Bux,

[128022]R. C.; Moore, C. M.: Male infant with ichthyosis, Kallmann syndrome,

[128023]chondrodysplasia punctata, and an Xp chromosome deletion. Am. J.

[128024]Med. Genet. 33: 100–107, 1989.

[128025]22885.

[128026]Marlhens, F.; Chelly, J.; Kaplan, J. C.; Lefrancois, D.; Harpey,

[128027]J. P.; Dutrillaux, B.: Familial deletion of Xp21.2 with glycerol

[128028]kinase deficiency and congenital adrenal hypoplasia. Hum. Genet. 77:

[128029]379–383, 1987.

[128030]22886.

[128031]Lin, T.; Orrison, B. M.; Suchy, S. F.; Lewis, R. A.; Nussbaum,

[128032]R. L.: Mutations are not uniformly distributed throughout the OCRL1

[128033]gene in Lowe syndrome patients. Molec. Genet. Metab. 64:
58–61,

[128034]1998.

[128035]22887.

[128036]Poustka, A.; Dietrich, A.; Langenstein, G.; Toniolo, D.;
Warren,

[128037]S. T.; Lehrach, H.: Physical map of human Xq27–qter: lo-
calizing the

[128038]region of the fragile X mutation. Proc. Nat. Acad. Sci. 88:
8302–8306,

[128039]1991.

[128040]22888.

[128041]Ma, K.; Sharkey, A.; Kirsch, S.; Vogt, P.; Keil, R.; Hargreave,

[128042]T. B.; McBeath, S.; Chandley, A. C.: Towards the molecular
localisation

[128043]of the AZF locus: mapping of microdeletions in azoosper-
mic men within

[128044]14 subintervals of interval 6 of the human Y chromosome.
Hum. Molec.

[128045]Genet. 1: 29–33, 1992.

[128046]22889.

[128047]Adam, A.; Tippett, P.; Gavin, J.; Noades, J.; Sanger, R.;
Race,

[128048]R. R.: The linkage relation of Xg to G6PD in Israelis: the
evidence

[128049]of a second series of families. Am. J. Hum. Genet. 30:
211–218,

[128050]1966.

[128051]22890.

[128052]Ahluwalia, A.; Corcoran, C. M.; Vulliamy, T. J.; Ishwad, C.
S.;

[128053]Naidu, J. M.; Argusti, A.; Stevens, D. J.; Mason, P. J.; Luz-
zatto,

[128054]L.: G6PD Kalyan and G6PD Kerala; two deficient variants in
India

[128055]caused by the same 317 glu-to-lys mutation. Hum. Molec.
Genet. 3:

[128056]209–210, 1992.

[128057]22891.

[128058]Aksoy, K.; Yuregir, G. T.; Dikmen, N.; Unlukurt, I.: Three
new

[128059]G6PD variants, G6PD Adana, G6PD Samandag, and G6PD
Balcali in Cukurova,

[128060]Turkey. Hum. Genet. 76: 199–201, 1987.

[128061]22892.

[128062]Alfinito, F.; Cimmino, A.; Ferraro, F.; Cubellis, M. V.;
Vitagliano,

[128063]L.; Francese, M.; Zagari, A.; Rotoli, B.; Filosa, S.; Martini, G.

[128064]: Molecular characterization of G6PD deficiency in South-
ern Italy:

[128065]heterogeneity, correlation genotype–phenotype and de-
scription of a

[128066]new variant (G6PD Neapolis). Brit. J. Haemat. 98: 41–46,
1997.

[128067]22893.

[128068]Alperin, J. B.; Mills, G. C.: New variants of glucose-
6-phosphate

[128069]dehydrogenase (G6PD). Clin. Res. 20: 76, 1972.

[128070]22894.

[128071]Azevedo, E.; Kirkman, H. N.; Morrow, A. C.; Motulsky, A.
G.: Variants

[128072]of red cell glucose–6-phosphate dehydrogenase among
Asiatic Indians. Ann.

[128073]Hum. Genet. 31: 373–379, 1968.

[128074]22895.

[128075]Azevedo, E. S.; Yoshida, A.: Brazilian variant of glucose-6-phosphate

[128076]dehydrogenase (GD Minas Gerais). Nature 222: 380-382, 1969.

[128077]22896.

[128078]Babalola, A. O. G.; Beetlestone, J. G.; Luzzatto, L.: Genetic

[128079]variants of human erythrocyte glucose-6-phosphate dehydrogenase: kinetic

[128080]and thermodynamic parameters of variants A, B, and A- in relation

[128081]to quaternary structure. J. Biol. Chem. 251: 2993-3002, 1976.

[128082]22897.

[128083]Balinsky, D.; Cayanis, E.; Carter, G.; Jenkins, T.; Bersohn, I.

[128084]: A new variant of human erythrocyte glucose-6-phosphate dehydrogenase:

[128085]G6PD Port Elizabeth. Int. J. Biochem. 4: 235-244, 1973.

[128086]22898.

[128087]Balinsky, D.; Gomperts, E.; Cayanis, E.; Jenkins, T.; Bryer, D.;

[128088]Bersohn, I.; Metz, J.: Glucose 6-phosphate dehydrogenase
Johannesburg:

[128089]a new variant with reduced activity in a patient with con-
genital non-spherocytic

[128090]haemolytic anaemia. Brit. J. Haemat. 25: 385-392, 1973.

[128091]22899.

[128092]Araten, D. J.; Nafa, K.; Pakdeesuwan, K.; Luzzatto, L.:
Clonal

[128093]populations of hematopoietic cells with paroxysmal noc-
turnal hemoglobinuria

[128094]genotype and phenotype are present in normal individu-
als. Proc. Nat.

[128095]Acad. Sci. 96: 5209-5214, 1999.

[128096]22900.

[128097]Bessler, M.; Mason, P.; Hillmen, P.; Luzzatto, L.: Somatic
mutations

[128098]and cellular selection in paroxysmal nocturnal
haemoglobinuria. Lancet 343:

[128099]951-953, 1994.

[128100]22901.

[128101]Bessler, M.; Mason, P. J.; Hillmen, P.; Miyata, T.; Yamada,

N.;

[128102]Takeda, J.; Luzzatto, L.; Kinoshita, T.: Paroxysmal nocturnal haemoglobinuria

[128103](PNH) is caused by somatic mutations in the PIG-A gene.

EMBO J. 13:

[128104]110–117, 1994.

[128105]22902.

[128106]Brodsky, R. A.; Vala, M. S.; Barber, J. P.; Medof, M. E.;

Jones,

[128107]R. J.: Resistance to apoptosis caused by PIG-A gene mutations in

[128108]paroxysmal nocturnal hemoglobinuria. Proc. Nat. Acad.

Sci. 94: 8756–8760,

[128109]1997.

[128110]22903.

[128111]Hillmen, P.; Lewis, S. M.; Bessler, M.; Luzzatto, L.; Dacie, J.

[128112]V.: Natural history of paroxysmal nocturnal hemoglobinuria. New

[128113]Eng. J. Med. 333: 1253–1258, 1995.

[128114]22904.

[128115]Hirose, S.; Mohny, R. P.; Mutka, S. C.; Ravi, L.; Singleton,

D.

[128116]R.; Perry, G.; Tartakoff, A. M.; Medof, M. E.: Derivation and characterization

[128117]of glycoinositol-phospholipid anchor-defective human K562 cell clones. J.

[128118]Biol. Chem. 267: 5272-5278, 1992.

[128119]22905.

[128120]Hirose, S.; Ravi, L.; Prince, G. M.; Rosenfeld, M. G.; Silber,

[128121]R.; Andresen, S. W.; Hazra, S. V.; Medof, M. E.: Synthesis of mannosylglucosaminylinositol

[128122]phospholipids in normal but not paroxysmal nocturnal hemoglobinuria

[128123]cells. Proc. Nat. Acad. Sci. 89: 6025-6029, 1992.

[128124]22906.

[128125]Horikawa, K.; Kawaguchi, T.; Ishihara, S.; Nagakura, S.; Hida,

[128126]M.; Kagimoto, T.; Mitsuya, H.; Nakakuma, H.: Frequent detection of

[128127]T cells with mutations of the hypoxanthine-guanine phosphoribosyltransferase

[128128]gene in patients with paroxysmal nocturnal hemoglobin-

uria. Blood 99:

[128129]24–29, 2002.

[128130]22907.

[128131]Iida, Y.; Takeda, J.; Miyata, T.; Inoue, N.; Nishimura, J.; Kitanishi, T.; Maeda, K.; Kinoshita, T.: Characterization of genomic

[128132]PIG-A gene:
a gene for glycosylphosphatidylinositol-anchor biosynthesis and paroxysmal

[128133]nocturnal hemoglobinuria. Blood 83: 3126–3131, 1994.

[128134]22908.

[128135]Kawagoe, K.; Takeda, J.; Endo, Y.; Kinoshita, T.: Molecular

[128136]cloning
of murine Pig-a, a gene for GPI-anchor biosynthesis, and

[128137]demonstration
of interspecies conservation of its structure, function, and

[128138]genetic
locus. Genomics 23: 566–574, 1994.

[128139]22909.

[128140]Keller, P.; Tremml, G.; Rosti, V.; Bessler, M.: X inactivation

[128141]and somatic cell selection rescue female mice carrying a

[128142]

Piga-null

[128143]mutation. Proc. Nat. Acad. Sci. 96: 7479–7483, 1999.

[128144]22910.

[128145]Luzzatto, L.; Bessler, M.: The dual pathogenesis of paroxysmal

[128146]nocturnal hemoglobinuria. Curr. Opin. Hemat. 3: 101–110, 1996.

[128147]22911.

[128148]Luzzatto, L.; Bessler, M.; Rotoli, B.: Somatic mutations in paroxysmal

[128149]nocturnal hemoglobinuria: a blessing in disguise? Cell 88: 1–4,

[128150]1997.

[128151]22912.

[128152]Maugard, C.; Margueritte, G.; Tuffery, S.; Rabesandratana, H.;

[128153]Demaille, J.; Claustres, M.: Recurrent PIG-A mutation (IVS5+1G-A)

[128154]in a paediatric case of paroxysmal nocturnal haemoglobinuria: detection

[128155]by the protein truncation test. Brit. J. Haemat. 98: 21–24,

1997.

[128156]22913.

[128157]Miyata, T.; Takeda, J.; Iida, Y.; Yamada, N.; Inoue, N.;
Takahashi,

[128158]M.; Maeda, K.; Kitani, T.; Kinoshita, T.: The cloning of PIG-
A, a

[128159]component in the early step of GPI-anchor biosynthesis.
Science 259:

[128160]1318-1320, 1993.

[128161]22914.

[128162]Miyata, T.; Yamada, N.; Iida, Y.; Nishimura, J.; Takeda, J.;
Kitani,

[128163]T.; Kinoshita, T.: Abnormalities of PIG-A transcripts in
granulocytes

[128164]from patients with paroxysmal nocturnal hemoglobinuria.
New Eng.

[128165]J. Med. 330: 249-255, 1994.

[128166]22915.

[128167]Nafa, K.; Bessler, M.; Castro-Malaspina, H.; Jhanwar, S.;
Luzzatto,

[128168]L.: The spectrum of somatic mutations in the PIG-A gene

in paroxysmal

[128169]nocturnal hemoglobinuria includes large deletions and small duplications. Blood

[128170]Cells Molec. Dis. 24: 370–384, 1998.

[128171]22916.

[128172]Nafa, K.; Bessler, M.; Deeg, H. J.; Luzzatto, L.: New somatic

[128173]mutation in the PIG-A gene emerges at relapse of paroxysmal nocturnal

[128174]hemoglobinuria. Blood 92: 3422–3427, 1998.

[128175]22917.

[128176]Nafa, K.; Mason, P. J.; Hillmen, P.; Luzzatto, L.; Bessler, M.

[128177]: Mutations in the PIG-A gene causing paroxysmal nocturnal hemoglobinuria

[128178]are mainly of the frameshift type. Blood 86: 4650–4655, 1995.

[128179]22918.

[128180]Nagarajan, S.; Brodsky, R. A.; Young, N. S.; Medof, M. E.: Genetic

[128181]defects underlying paroxysmal nocturnal hemoglobinuria that arises

[128182]out of aplastic anemia. Blood 86: 4656–4661, 1995.

[128183]22919.

[128184]Nagarajan, S.; Brown, C. J.; Medof, M. E.: Identification of

[128185]a PIG-A related processed gene on chromosome 12. Hum.
Genet. 95:

[128186]691-697, 1995.

[128187]22920.

[128188]McCabe, E. R. B.: Personal Communication. Denver, Colo.
10/9/1985.

[128189]22921.

[128190]McCabe, E. R. B.; Fennessey, P. V.; Guggenheim, M. A.;
Miles,

[128191]B. S.; Bullen, W. W.; Sceats, D. J.; Goodman, S. I.: Human
glycerol

[128192]kinase deficiency with hyperglycerolemia and glyceroluria.
Biochem.

[128193]Biophys. Res. Commun. 78: 1327-1333, 1977.

[128194]22922.

[128195]Renier, W. O.; Nabben, F. A. E.; Hustinx, T. W. J.;
Veerkamp,

[128196]J. H.; Otten, B. J.; ter Laak, H. J.; ter Haar, B. G. A.;
Gabreels,

- [128197]F. J. M.: Congenital adrenal hypoplasia, progressive muscular dystrophy
- [128198]and severe mental retardation, in association with glycerol kinase
- [128199]deficiency, in male sibs. Clin. Genet. 24: 449–457, 1983.
- [128200]22923.
- [128201]Rose, C. I.; Haines, D. S. M.: Familial hyperglycerolemia. J.
- [128202]Clin. Invest. 61: 163–170, 1978.
- [128203]22924.
- [128204]Saito, F.; Goto, J.; Kakinuma, H.; Nakamura, F.; Murayama, S.;
- [128205]Nakano, I.; Tonomura, A.: Inherited Xp21 deletion in a boy with complex
- [128206]glycerol kinase deficiency syndrome. (Letter) Clin. Genet. 29: 92–93,
- [128207]1986.
- [128208]22925.
- [128209]Sargent, C. A.; Affara, N. A.; Bentley, E.; Pelmeur, A.; Bailey,
- [128210]D. M. D.; Davey, P.; Dow, D.; Leversha, M.; Aplin, H.; Besley, G.

[128211]T. N.; Ferguson-Smith, M. A.: Cloning of the X-linked
glycerol kinase

[128212]deficiency gene and its identification by sequence com-
parison to the

[128213]*Bacillus subtilis* homologue. *Hum. Molec. Genet.* 2:
97-106, 1993.

[128214]22926.

[128215]Sargent, C. A.; Kidd, A.; Moore, S.; Dean, J.; Besley, G. T.
N.;

[128216]Affara, N. A.: Five cases of isolated glycerol kinase defi-
ciency,

[128217]including two families: failure to find genotype:phenotype
correlation. *J.*

[128218]*Med. Genet.* 37: 434-441, 2000.

[128219]22927.

[128220]Sargent, C. A.; Young, C.; Marsh, S.; Ferguson-Smith, M.
A.; Affara,

[128221]N. A.: The glycerol kinase gene family: structure of the Xp
gene,

[128222]and related intronless retroposons. *Hum. Molec. Genet.* 3:
1317-1324,

[128223]1994.

[128224]22928.

[128225]Sjarif, D. R.; Ploos van Amstel, J. K.; Duran, M.; Beemer, F.

[128226]A.; Poll-The, B. T.: Isolated and contiguous glycerol kinase
gene

[128227]disorders: a review. J. Inherit. Metab. Dis. 23: 529-547,
2000.

[128228]22929.

[128229]Sjarif, D. R.; Sinke, R. J.; Duran, M.; Beemer, F. A.; Kleijer,

[128230]W. J.; Ploos van Amstel, J. K.; Poll-The, B. T.: Clinical het-
erogeneity

[128231]and novel mutations in the glycerol kinase gene in three
families

[128232]with isolated glycerol kinase deficiency. J. Med. Genet. 35:
650-656,

[128233]1998.

[128234]22930.

[128235]von Petrykowski, W.; Beckmann, R.; Bohm, N.; Ketelsen,
U.-P.;

[128236]Ropers, H. H.; Sauer, M.: Adrenal insufficiency, myopathic
hypotonia,

[128237]severe psychomotor retardation, failure to thrive, consti-

pation and

[128238]bladder ectasia in 2 brothers: adrenomyodystrophy. *Helv. Paediat.*

[128239]*Acta* 37: 387–400, 1982.

[128240]22931.

[128241]Walker, A. P.; Muscatelli, F.; Monaco, A. P.: Isolation of the
[128242]human Xp21 glycerol kinase gene by positional cloning.
Hum. Molec.

[128243]*Genet.* 2: 107–114, 1993.

[128244]22932.

[128245]Walker, A. P.; Muscatelli, F.; Stafford, A. N.; Chelly, J.;
Dahl,

[128246]N.; Blomquist, H. K.; Delanghe, J.; Willems, P. J.; Stein-
mann, B.;

[128247]Monaco, A. P.: Mutations and phenotype in isolated glyc-
erol kinase

[128248]deficiency. *Am. J. Hum. Genet.* 58: 1205–1211, 1996.

[128249]22933.

[128250]Wieringa, B.; Hustinx, T.; Scheres, J.; Hofker, M.; Schepens,

[128251]J.; Ropers, H. H.; ter Haar, B.: Glycerol kinase deficiency

syndrome

[128252]explained as X-chromosomal deletion. (Abstract) Cytogenet. Cell Genet. 40:

[128253]777 only, 1985.

[128254]22934.

[128255]Wieringa, B.; Hustinx, T.; Scheres, J.; Renier, W.; ter Haar,

[128256]B.: Complex glycerol kinase deficiency syndrome explained as X-chromosomal

[128257]deletion. (Letter) Clin. Genet. 27: 522-523, 1985.

[128258]22935.

[128259]Willard, H. F.: Cloning of the X-linked glycerol kinase gene. Hum.

[128260]Molec. Genet. 2: 95-96, 1993.

[128261]22936.

[128262]Zhang, Y.-H.; Dipple, K. M.; Vilain, E.; Huang, B.-L.; Finlayson,

[128263]G.; Therrell, B. L.; Worley, K.; Deininger, P.; McCabe, E. R. B.:

[128264]AluY insertion (IVS4-52ins316alu) in the glycerol kinase gene from

[128265]an individual with benign glycerol kinase deficiency. Hum.

Mutat. 15:

[128266]316–323, 2000.

[128267]22937.

[128268]Balinsky, D.; Rootman, A. J.; Nurse, G. T.; Cayanis, E.;
Lane,

[128269]A.; Jenkins, T.; Bersohn, I.: G6PD Kuanyama: a new variant
of human

[128270]erythrocyte glucose 6–phosphate dehydrogenase showing
slower than

[128271]normal electrophoretic mobility. S. Afr. J. Med. Sci. 39:
5–13,

[128272]1974.

[128273]22938.

[128274]Barretto, O. C.; Nonoyama, K.: Gd(+) Cuiaba, a new rare
glucose–6–phosphate

[128275]dehydrogenase variant presenting normal activity. Hum.
Genet. 77:

[128276]201–202, 1987.

[128277]22939.

[128278]Kornak, U.; Kasper, D.; Bosl, M. R.; Kaiser, E.; Schweizer,
M.;

[128279]Schulz, A.; Friedrich, W.; Delling, G.; Jentsch, T. J.: Loss of the

[128280]ClC-7 chloride channel leads to osteopetrosis in mice and man. Cell 104:

[128281]205–215, 2001.

[128282]22940.

[128283]Kornak, U.; Schulz, A.; Friedrich, W.; Uhlhaas, S.; Kremens, B.;

[128284]Voit, T.; Hasan, C.; Bode, U.; Jentsch, T. J.; Kubisch, C.: Mutations

[128285]in the $\alpha 3$ subunit of the vacuolar H(+)-ATPase cause infantile malignant

[128286]osteopetrosis. Hum. Molec. Genet. 9: 2059–2063, 2000.

[128287]22941.

[128288]Lowe, C. U.: Oculo-cerebral-renal syndrome. Maandschr. Kindergeneesk. 28:

[128289]77–80, 1960.

[128290]22942.

[128291]Lowe, C. U.; Terrey, M.; MacLachlan, E. A.: Organic-aciduria,

[128292]decreased renal ammonia production, hydrophthalmos,

and mental retardation. Am.

[128293]J. Dis. Child. 83: 164–184, 1952.

[128294]22943.

[128295]Martin, V. A. F.; Carson, N. A. J.: Inborn metabolic disorders

[128296]with associated ocular lesions in Northern Ireland. Trans. Ophthal.

[128297]Soc. U.K. 87: 847–870, 1967.

[128298]22944.

[128299]Matsuda, I.; Sugai, M.; Kajii, T.: Ornithine loading test in

[128300]Lowe's syndrome. J. Pediat. 77: 127–129, 1970.

[128301]22945.

[128302]Matsuda, I.; Takeda, T.; Sugai, M.; Matsuura, N.: Oculo-cerebrorenal

[128303]syndrome. Am. J. Dis. Child. 117: 205–212, 1969.

[128304]22946.

[128305]McCance, R. A.; Matheson, W. J.; Gresham, G. A.; Elkinton, J.

[128306]R.: The cerebro-ocular-renal dystrophies: a new variant. Arch. Dis.

[128307]Child. 35: 240–249, 1960.

[128308]22947.

[128309]Monnier, N.; Satre, V.; Lerouge, E.; Berthoin, F.; Lunardi, J.

[128310]: OCRL1 mutation analysis in French Lowe syndrome patients: implications

[128311]for molecular diagnosis strategy and genetic counseling.

Hum. Mutat. 16:

[128312]157–165, 2000.

[128313]22948.

[128314]Mueller, O. T.; Hartsfield, J. K., Jr.; Gallardo, L. A.; Essig,

[128315]Y.-P.; Miller, K. L.; Papenhausen, P. R.; Tedesco, T. A.:

Lowe oculocerebrorenal

[128316]syndrome in a female with a balanced X;20 translocation: mapping of

[128317]the X chromosome breakpoint. Am. J. Hum. Genet. 49: 804–810, 1991.

[128318]22949.

[128319]Nelson, D. L.; Ballabio, A.; Victoria, M. F.; Pieretti, M.; Bies,

[128320]R. D.; Gibbs, R. A.; Maley, J. A.; Chinault, A. C.; Webster, T. D.;

[128321]Caskey, C. T.: Alu-primed polymerase chain reaction for regional

[128322]assignment of 110 yeast artificial chromosome clones from the human

[128323]X chromosome: identification of clones associated with a disease locus. Proc.

[128324]Nat. Acad. Sci. 88: 6157-6161, 1991.

[128325]22950.

[128326]Oetliker, O.; Rossi, E.: The influence of extracellular fluid

[128327]volume on the renal bicarbonate threshold: a study of two children

[128328]with Lowe's syndrome. Pediat. Res. 3: 140-148, 1969.

[128329]22951.

[128330]Olivos-Glander, I. M.; Janne, P. A.; Nussbaum, R. L.: The oculocerebrorenal

[128331]syndrome gene product is a 105-kD protein localized to the Golgi complex. Am.

[128332]J. Hum. Genet. 57: 817-823, 1995.

[128333]22952.

[128334]Pallisgaard, G.; Goldschmidt, E.: The oculo-cerebro-renal syndrome

[128335] of Lowe in four generations of one family. Acta Paediat.

Scand. 60:

[128336] 146–148, 1971.

[128337] 22953.

[128338] Reilly, D. S.; Lewis, R. A.; Ledbetter, D. H.; Nussbaum, R. L.

[128339] Tightly linked flanking markers for the Lowe oculocerebrorenal syndrome,

[128340] with application to carrier assessment. Am. J. Hum. Genet. 42: 748–755,

[128341] 1988.

[128342] 22954.

[128343] Reilly, D. S.; Lewis, R. A.; Nussbaum, R. L.: Genetic and physical

[128344] mapping of Xq24–q26 markers flanking the Lowe oculocerebrorenal syndrome. Genomics 8:

[128345] 62–70, 1990.

[128346] 22955.

[128347] Richards, W.; Donnell, G. N.; Wilson, W. A.; Stowens, D.; Perry,

[128348] T.: The oculo-cerebro-renal syndrome of Lowe. Am. J. Dis. Child. 109:

[128349]185–203, 1965.

[128350]22956.

[128351]Roschinger, W.; Muntau, A. C.; Rudolph, G.; Roscher, A. A.;
Kammerer,

[128352]S.: Carrier assessment in families with Lowe oculocere-
brorenal syndrome:

[128353]novel mutations in the OCRL1 gene and correlation of di-
rect DNA diagnosis

[128354]with ocular examination. Molec. Genet. Metab. 69:
213–222, 2000.

[128355]22957.

[128356]Satre, V.; Monnier, N.; Berthoin, F.; Ayuso, C.; Joannard,
A.;

[128357]Jouk, P.-S.; Lopez-Pajares, I.; Megabarne, A.; Philippe, H.
J.; Plauchu,

[128358]H.; Torres, M. L.; Lunardi, J.: Characterization of a
germline mosaicism

[128359]in families with Lowe syndrome, and identification of
seven novel

[128360]mutations in the OCRL1 gene. Am. J. Hum. Genet. 65:
68–76, 1999.

[128361]22958.

[128362]Silver, D. N.; Lewis, R. A.; Ledbetter, D. H.; Bobrow, M.;
Nussbaum,

[128363]R. L.: Identification of tightly-linked flanking markers for
Lowe

[128364]syndrome. (Abstract) Cytogenet. Cell Genet. 46: 692 only,
1987.

[128365]22959.

[128366]Silver, D. N.; Lewis, R. A.; Nussbaum, R. L.: Mapping the
Lowe

[128367]oculocerebrorenal syndrome to Xq24-q26 by use of re-
striction fragment

[128368]length polymorphisms. J. Clin. Invest. 79: 282-285, 1987.

[128369]22960.

[128370]Streiff, E. B.; Straub, W.; Golay, L.: Les manifestations ocu-
laires

[128371]du syndrome de Lowe. Ophthalmologica 135: 632-639,
1958.

[128372]22961.

[128373]Suchy, S. F.; Lin, T.; Horwitz, J. A.; O'Brien, W. E.; Nuss-
baum,

[128374]R. L.: First report of prenatal biochemical diagnosis of
Lowe syndrome. Prenatal

[128375]Diag. 18: 1117–1121, 1998.

[128376]22962.

[128377]Svorc, J.; Masopust, J.; Komarkova, A.; Macek, M.; Hyanek, J.

[128378]: Oculocerebrorenal syndrome in a female child. Am. J. Dis. Child. 114:

[128379]186–190, 1967.

[128380]22963.

[128381]Tripathi, R.; Cibis, G. W.; Harris, D. J.; Tripathi, B.: Lowe's

[128382]syndrome. Birth Defects Orig. Art. Ser. 18(6): 629–644, 1982.

[128383]22964.

[128384]Wadelius, C.; Fagerholm, P.; Pettersson, U.; Anneren, G.: Lowe

[128385]oculocerebrorenal syndrome: DNA-based linkage of the gene to Xq24–q26,

[128386]using tightly linked flanking markers and the correlation to lens

[128387]examination in carrier diagnosis. Am. J. Hum. Genet. 44: 241–247,

[128388]1989.

[128389]22965.

[128390]Wilson, W. A.; Richards, W.; Donnell, G. N.: Oculo-cere-bral-renal

[128391]syndrome of Lowe: a review of eight cases noting the ge-netic inheritance. Arch.

[128392]Ophthal. 70: 5-11, 1963.

[128393]22966.

[128394]Witzleben, C. L.; Schoen, E. J.; Tu, W. H.; McDonald, L. W.:

[128395]Progressive morphologic renal changes in the oculo-cerebro-renal syndrome

[128396]of Lowe. Am. J. Med. 44: 319-324, 1968.

[128397]22967.

[128398]Zhang, X.; Jefferson, A. B.; Auethavekiat, V.; Majerus, P. W.

[128399]: The protein deficient in Lowe syndrome is a phos-phatidylinositol-4,5-bisphosphate

[128400]5-phosphatase. Proc. Nat. Acad. Sci. 92: 4853-4856, 1995.

[128401]22968.

[128402]Schram, A. W.; Goldfischer, S.; van Roermund, C. W. T.; Brouwer-Kelder,

[128403]E. M.; Collins, J.; Hashimoto, T.; Heymans, H. S. A.; van
den Bosch,

[128404]H.; Schutgens, R. B. H.; Tager, J. M.; Wanders, R. J. A.: Hu-
man peroxisomal

[128405]3-oxoacyl-coenzyme A thiolase deficiency. Proc. Nat.
Acad. Sci. 84:

[128406]2494-2496, 1987.

[128407]22969.

[128408]Suzuki, Y.; Shimozawa, N.; Yajima, S.; Tomatsu, S.; Kondo,
N.;

[128409]Nakada, Y.; Akaboshi, S.; Iai, M.; Tanabe, Y.; Hashimoto,
T.; Wanders,

[128410]R. J. A.; Schutgens, R. B. H.; Moser, H. W.; Orii, T.: Novel
subtype
[128411]of peroxisomal acyl-CoA oxidase deficiency and bifunc-
tional enzyme
[128412]deficiency with detectable enzyme protein: identification
by means
[128413]of complementation analysis. Am. J. Hum. Genet. 54:
36-43, 1994.
[128414]22970.

[128415]Rosenberg, C. L.; Wong, E.; Petty, E. M.; Bale, A. E.; Tsuji-
moto,
[128416]Y.; Harris, N. L.; Arnold, A.: PRAD1, a candidate BCL1
oncogene:
[128417]mapping and expression in centrocytic lymphoma. Proc.
Nat. Acad.
[128418]Sci. 88: 9638-9642, 1991.
[128419]22971.

[128420]Tsujimoto, Y.; Jaffe, E.; Cossman, J.; Gorham, J.; Nowell, P.
[128421]C.; Croce, C. M.: Clustering of breakpoints on chromo-
some 11 in human
[128422]B-cell neoplasms with the t(11;14) chromosome translo-
cation. Nature 315:

[128423]340–343, 1985.

[128424]22972.

[128425]Tsujiimoto, Y.; Yunis, J.; Onorato–Showe, L.; Erikson, J.;
Nowell,

[128426]P. C.; Croce, C. M.: Molecular cloning of the chromosomal
breakpoint

[128427]of B–cell lymphomas and leukemias with the t(11;14)
chromosome translocation. Science 224:

[128428]1403–1406, 1984.

[128429]22973.

[128430]Barbany, G.; Hoglund, M.; Simonsson, B.: Complete
molecular remission

[128431]in chronic myelogenous leukemia after imatinib therapy.
(Letter) New

[128432]Eng. J. Med. 347: 539–540, 2002.

[128433]22974.

[128434]Budarf, M.; Canaani, E.; Emanuel, B. S.: Linear order of the
four

[128435]BCR–related loci in 22q11. Genomics 3: 168–171, 1988.

[128436]22975.

[128437]Castellanos, A.; Pintado, B.; Weruaga, E.; Arevalo, R.;
Lopez,
[128438]A.; Orfao, A.; Sanchez-Garcia, I.: A BCR-ABL(p190) fusion
gene made
[128439]by homologous recombination causes B-cell acute lym-
phoblastic leukemias
[128440]in chimeric mice with independence of the endogenous
bcr product. Blood 90:
[128441]2168-2174, 1997. 1. Brunning, R. D.: Philadelphia chro-
mosome positive
[128442]leukemia. Hum. Path. 11: 307-309, 1980.
[128443]22976.
[128444]Chisoe, S. L.; Bodenteich, A.; Wang, Y.-F.; Wang, Y.-P.;
Burian,
[128445]D.; Clifton, S. W.; Crabtree, J.; Freeman, A.; Iyer, K.; Jian,
L.;
[128446]Ma, Y.; McLaury, H.-J.; Pan, H.-Q.; Sarhan, O. H.; Toth, S.;
Wang,
[128447]Z.; Zhang, G.; Heisterkamp, N.; Groffen, J.; Roe, B. A.: Se-
quence
[128448]and analysis of the human ABL gene, the BCR gene, and
regions involved

[128449]in the Philadelphia chromosomal translocation. Genomics
27: 67-82,

[128450]1995.

[128451]22977.

[128452]Court Brown, W. M.; Doll, R.: Mortality from cancer and
other

[128453]causes after radiotherapy for ankylosing spondylitis. Brit.
Med.

[128454]J. 2: 1327-1332, 1965.

[128455]22978.

[128456]Croce, C. M.; Huebner, K.; Isobe, M.; Fainstain, E.; Lifshitz,

[128457]B.; Shtivelman, E.; Canaani, E.: Mapping of four distinct
BCR-related

[128458]loci to chromosome region 22q11: order of BCR loci rela-
tive to chronic

[128459]myelogenous leukemia and acute lymphoblastic leukemia
breakpoints. Proc.

[128460]Nat. Acad. Sci. 84: 7174-7178, 1987.

[128461]22979.

[128462]Daley, G. Q.; Van Etten, R. A.; Baltimore, D.: Induction of
chronic

[128463]myelogenous leukemia in mice by the P210(bcr/abl) gene
of the Philadelphia

[128464]chromosome. Science 247: 824–830, 1990.

[128465]22980.

[128466]de Klein, A.; Geurts van Kessel, A.; Grosveld, G.; Bartram,
C.

[128467]R.; Hagemeijer, A.; Bootsma, D.; Spurr, N. K.; Heisterkamp,
N.; Groffen,

[128468]J.; Stephenson, J. R.: A cellular oncogene is translocated to
the

[128469]Philadelphia chromosome in chronic myelocytic
leukaemia. Nature 300:

[128470]765–767, 1982.

[128471]22981.

[128472]Diekmann, D.; Brill, S.; Garrett, M. D.; Totty, N.; Hsuan, J.;

[128473]Monfries, C.; Hall, C.; Lim, L.; Hall, A.: Bcr encodes a GT–
Pase–activating

[128474]protein for p21(rac). Nature 351: 400–402, 1991.

[128475]22982.

[128476]Druker, B. J.; Sawyers, C. L.; Kantarjian, H.; Resta, D. J.;
Reese,

[128477]S. F.; Ford, J. M.; Capdeville, R.; Talpaz, M.: Activity of a specific

[128478]inhibitor of the BCR-ABL tyrosine kinase in the blast crisis of chronic

[128479]myeloid leukemia and acute lymphoblastic leukemia with the Philadelphia

[128480]chromosome. New Eng. J. Med. 344: 1038-1042, 2001.

[128481]22983.

[128482]Druker, B. J.; Talpaz, M.; Resta, D. J.; Peng, B.; Buchdunger,

[128483]E.; Ford, J. M.; Lydon, N. B.; Kantarjian, H.; Capdeville, R.; Ohno-Jones,

[128484]S.; Sawyers, C. L.: Efficacy and safety of a specific inhibitor of

[128485]the BCR-ABL tyrosine kinase in chronic myeloid leukemia. New Eng.

[128486]J. Med. 344: 1031-1037, 2001.

[128487]22984.

[128488]Emanuel, B. S.; Selden, J. R.; Wang, E.; Nowell, P. C.; Croce,

[128489]C. M.: In situ hybridization and translocation breakpoint mapping.

[128490]J. Non-identical 22q11 breakpoints for the t(9;22) of CML
and the

[128491]t(8;22) of Burkitt lymphoma. Cytogenet. Cell Genet. 38:
127-131,

[128492]1984.

[128493]22985.

[128494]Misrahi, M.; Meduri, G.; Pissard, S.; Bouvattier, C.; Beau, I.;

[128495]Loosfelt, H.; Jolivet, A.; Rappaport, R.; Milgrom, E.;
Bougnères,

[128496]P.: Comparison of immunocytochemical and molecular
features with

[128497]the phenotype in a case of incomplete male pseudo-
hermaphroditism associated

[128498]with a mutation of the luteinizing hormone receptor. J.
Clin. Endocr.

[128499]Metab. 82: 2159-2165, 1997.

[128500]22986.

[128501]Perez-Palacios, G.; Scaglia, H. E.; Kofman, S.; Saavedra, D.;

[128502]Ochoa, S.; Laraza, O.; Perez, A. E.: Inherited deficiency of
gonadotropin

[128503]receptor in Leydig cells: a new form of male pseudo-

hermaphroditism.

[128504](Abstract) Am. J. Hum. Genet. 27: 71A, 1975.

[128505]22987.

[128506]Perez-Palacios, G.; Scaglia, H. E.; Kofman-Alfaro, S.;
Saavedra,

[128507]D.; Ochoa, S.; Larraza, O.; Perez, A. E.: Inherited male
pseudohermaphroditism

[128508]due to gonadotrophin unresponsiveness. Acta Endocr. 98:
148-155,

[128509]1981.

[128510]22988.

[128511]Rodien, P.; Cetani, F.; Costagliola, S.; Tonacchera, M.;
Duprez,

[128512]L.; Minegishi, T.; Govaerts, C.; Vassart, G.: Evidences for
an allelic

[128513]variant of the human LC/CG receptor rather than a gene
duplication:

[128514]functional comparison of wild-type and variant receptors.
J. Clin.

[128515]Endocr. Metab. 83: 4431-4434, 1998.

[128516]22989.

[128517]Rousseau-Merck, M. F.; Misrahi, M.; Atger, M.; Loosfelt, H.; Milgrom,

[128518]E.; Berger, R.: Localization of the human luteinizing hormone/choriogonadotropin

[128519]receptor gene (LHCGR) to chromosome 2p21. Cytogenet. Cell Genet. 54:

[128520]77-79, 1990.

[128521]22990.

[128522]Saldanha, P. H.; Arnhold, I. J. P.; Mendonca, B. B.; Bloise, W.;

[128523]Toledo, S. P. A.: A clinico-genetic investigation of Leydig cell

[128524]hypoplasia. Am. J. Med. Genet. 26: 337-344, 1987.

[128525]22991.

[128526]Schwartz, M.; Imperato-McGinley, J.; Peterson, R. E.; Cooper,

[128527]G.; Morris, P. L.; MacGillivray, M.; Hensle, T.: Male pseudo-hermaphroditism

[128528]secondary to an abnormality in Leydig cell differentiation. J. Clin.

[128529]Endocr. Metab. 53: 123-127, 1981.

[128530]22992.

[128531]Segaloff, D. L.; Ascoli, M.: The lutropin/chori-
gonadotropin

[128532]receptor: 4 years later. Endocr. Rev. 14: 324-347, 1993.

[128533]22993.

[128534]Shenker, A.; Laue, L.; Kosugi, S.; Merendino, J. J., Jr.;
Minegishi,

[128535]T.; Cutler, G. B., Jr.: A constitutively activating mutation of
the

[128536]lutinizing hormone receptor in familial male precocious
puberty. Nature 365:

[128537]652-654, 1993.

[128538]22994.

[128539]Rosenthal, I. M.; Refetoff, S.; Rich, B.; Barnes, R. B.; Sun-
thornthepvarakul,

[128540]T.; Parma, J.; Rosenfield, R. L.: Response to challenge with
gonadotropin-releasing

[128541]hormone agonist in a mother and her two sons with a
constitutively

[128542]activating mutation of the luteinizing hormone receptor-
-a clinical

[128543]research center study. J. Clin. Endocr. Metab. 81:
3802–3806, 1996.

[128544]22995.

[128545]Stavrou, S. S.; Zhu, Y.-S.; Cai, L.-Q.; Katz, M. D.; Herrera,
[128546]C.; Defillo-Ricart, M.; Imperato-McGinley, J.: A novel mu-
tation of

[128547]the human luteinizing hormone receptor in 46XY and
46XX sisters. J.

[128548]Clin. Endocr. Metab. 83: 2091–2098, 1998.

[128549]22996.

[128550]Toledo, S. P. A.; Brunner, H. G.; Kraaij, R.; Post, M.; Dahia,
[128551]P. L. M.; Hayashida, C. Y.; Kremer, H.; Themmen, A. P. N.:
An inactivating

[128552]mutation of the luteinizing hormone receptor causes
amenorrhea in

[128553]a 46,XX female. J. Clin. Endocr. Metab. 81: 3850–3854,
1996.

[128554]22997.

[128555]Tsai-Morris, C-H.; Geng, Y.; Buczko, E.; Dufau, M. L.: A
novel

[128556]human luteinizing hormone receptor gene. J. Clin. Endocr.

Metab. 83:

[128557]288–291, 1998.

[128558]22998.

[128559]Wu, S.–M.; Chan, W.–Y.: Male pseudohermaphroditism due to inactivating

[128560]luteinizing hormone receptor mutations. Arch. Med. Res. 30: 495–500,

[128561]1999.

[128562]22999.

[128563]Wu, S.–M.; Hallermeier, K. M.; Laue, L.; Brain, C.; Berry, A.

[128564]C.; Grant, D. B.; Griffin, J. E.; Wilson, J. D.; Cutler, G. B., Jr.;

[128565]Chan, W.–Y.: Inactivation of the luteinizing hormone/chorionic gonadotropin

[128566]receptor by an insertional mutation in Leydig cell hypoplasia. Molec.

[128567]Endocr. 12: 1651–1660, 1998.

[128568]23000.

[128569]Yano, K.; Hidaka, A.; Saji, M.; Polymeropoulos, M. H.; Okuno,

[128570]A.; Kohn, L. D.; Cutler, G. B., Jr.: A sporadic case of male–

limited

[128571]precocious puberty has the same constitutively activating point mutation

[128572]in luteinizing hormone/choriogonadotropin receptor gene as familial

[128573]cases. J. Clin. Endocr. Metab. 79: 1818–1823, 1994.

[128574]23001.

[128575]Yano, K.; Kohn, L. D.; Saji, M.; Okuno, A.; Cutler, G. B., Jr.

[128576]: Phe576 plays an important role in the secondary structure and intracellular

[128577]signaling of the human luteinizing hormone/chorionic gonadotropin

[128578]receptor. J. Clin. Endocr. Metab. 82: 2586–2591, 1997.

[128579]23002.

[128580]Yano, K.; Saji, M.; Hidaka, A.; Moriya, N.; Okuno, A.; Kohn, L.

[128581]D.; Cutler, G. B.: A new constitutively activating point mutation

[128582]in the luteinizing hormone/choriogonadotropin receptor gene in cases

[128583]of male-limited precocious puberty. J. Clin. Endocr. 80:

1162–1168,

[128584]1995.

[128585]23003.

[128586]Zenteno, J. C.; Canto, P.; Kofman–Alfaro, S.; Mendez, J. P.:

[128587]Evidence for genetic heterogeneity in male pseudo–
hermaphroditism due

[128588]to Leydig cell hypoplasia. J. Clin. Endocr. Metab. 84:
3803–3806,

[128589]1999.

[128590]23004.

[128591]Grandori, C.; Mac, J.; Siebelt, F.; Ayer, D. E.; Eisenman, R.
N.

[128592]: Myc–Max heterodimers activate a DEAD box gene and
interact with

[128593]multiple E box–related sites in vivo. EMBO J. 15:
4344–4357, 1996.

[128594]23005.

[128595]Deltas, C. C.: Mutations of the human polycystic kidney
disease

[128596]2 (PKD2) gene. Hum. Mutat. 18: 13–24, 2001.

[128597]23006.

- [128598]Elles, R. G.; Read, A. P.; Hodgkinson, K. A.; Watters, A.; Harris,
R.: Recombination or heterogeneity: is there a second locus for adult polycystic kidney disease? J. Med. Genet. 27: 413–417, 1990.
- [128601]23007.
- [128602]Fossdal, R.; Boovarsson, M.; Asmundsson, P.; Ragnarsson, J.; Peters, D.; Breuning, M. H.; Jensson, O.: Icelandic families with autosomal dominant polycystic kidney disease: families unlinked to chromosome 16p13.3 revealed by linkage analysis. Hum. Genet. 91: 609–613, 1993.
- [128606]23008.
- [128607]Gallagher, A. R.; Cedzich, A.; Gretz, N.; Somlo, S.; Witzgall, R.: The polycystic kidney disease protein PKD2 interacts with Hax-1, a protein associated with the actin cytoskeleton. Proc. Nat. Acad.
- [128610]Sci. 97: 4017–4022, 2000.

[128611]23009.

[128612]Gonzalez-Perrett, S.; Kim, K.; Ibarra, C.; Damiano, A. E.;
Zotta,

[128613]E.; Batelli, M.; Harris, P. C.; Reisin, I. L.; Arnaout, M. A.;
Cantiello,

[128614]H. F.: Polycystin-2, the protein mutated in autosomal
dominant polycystic

[128615]kidney disease (ADPKD), is a Ca^{2+} -permeable nonselective
cation

[128616]channel. Proc. Nat. Acad. Sci. 98: 1182-1187, 2001.

[128617]23010.

[128618]Grantham, J. J.; Calvet, J. P.: Polycystic kidney disease: in

[128619]danger of being X-rated? Proc. Nat. Acad. Sci. 98:
790-792, 2001.

[128620]23011.

[128621]Hanaoka, K.; Qian, F.; Boletta, A.; Bhunia, A. K.; Piontek,
K.;

[128622]Tsiokas, L.; Sukhatme, V. P.; Guggino, W. B.; Germino, G.
G.: Co-assembly

[128623]of polycystin-1 and -2 produces unique cation-permeable
currents. Nature 408:

[128624]990–994, 2000.

[128625]23012.

[128626]Hateboer, N.; van Dijk, M. A.; Bogdanova, N.; Coto, E.;
Saggar–Malik,

[128627]A. K.; San Millan, J. L.; Torra, R.; Breuning, M.; Ravine, D.:
Comparison

[128628]of phenotypes of polycystic kidney disease types 1 and 2.
Lancet 353:

[128629]103–107, 1999.

[128630]23013.

[128631]Hayashi, T.; Mochizuki, T.; Reynolds, D. M.; Wu, G.; Cai,
Y.;

[128632]Somlo, S.: Characterization of the exon structure of the
polycystic

[128633]kidney disease 2 gene (PKD2). Genomics 44: 131–136,
1997.

[128634]23014.

[128635]Jeffery, S.; Saggar–Malik, A. K.; Morgan, S.; MacGregor, G.
A.

[128636]: A family with autosomal dominant polycystic kidney dis-
ease not linked

[128637]to chromosome 16p13.3. Clin. Genet. 44: 173–176, 1993.

[128638]23015.

[128639]Kimberling, W. J.; Fain, P. R.; Kenyon, J. B.; Goldgar, D.;
Sujansky,

[128640]E.; Gabow, P. A.: Linkage heterogeneity of autosomal
dominant polycystic

[128641]kidney disease. New Eng. J. Med. 319: 913–918, 1988.

[128642]23016.

[128643]Kimberling, W. J.; Kumar, S.; Gabow, P. A.; Kenyon, J. B.;
Connolly,

[128644]C. J.; Somlo, S.: Autosomal dominant polycystic kidney
disease: localization

[128645]of the second gene to chromosome 4q13–q23. Genomics
18: 467–472,

[128646]1993.

[128647]23017.

[128648]Koptides, M.; Hadjimichael, C.; Koupepidou, P.; Pierides,
A.;

[128649]Deltas, C. C.: Germinal and somatic mutations in the PKD2
gene of

[128650]renal cysts in autosomal dominant polycystic kidney dis–

ease. Hum.

[128651]Molec. Genet. 8: 509–513, 1999.

[128652]23018.

[128653]Koptides, M.; Mean, R.; Demetriou, K.; Pierides, A.; Deltas, C.

[128654]C.: Genetic evidence for a trans–heterozygous model for cystogenesis

[128655]in autosomal dominant polycystic kidney disease. Hum. Molec. Genet. 9:

[128656]447–452, 2000.

[128657]23019.

[128658]Koulén, P.; Cai, Y.; Geng, L.; Maeda, Y.; Nishimura, S.; Witzgall,

[128659]R.; Ehrlich, B. E.; Somlo, S.: Polycystin–2 is an intracellular calcium

[128660]release channel. Nature Cell Biol. 4: 191–197, 2002.

[128661]23020.

[128662]Kumar, S.; Kimberling, W. J.; Gabow, P. A.; Kenyon, J. B.: Genetic

[128663]linkage studies of autosomal dominant polycystic kidney disease: search

[128664]for the second gene in a large Sicilian family. Hum. Genet.

87:

[128665]129–133, 1991.

[128666]23021.

[128667]Kumar, S.; Kimberling, W. J.; Gabow, P. A.; Shugart, Y. Y.;

Pieke–Dahl,

[128668]S.: Exclusion of autosomal dominant polycystic kidney
disease type

[128669]II (ADPKD2) from 160 cM of chromosome 1. J. Med. Genet.

27: 697–700,

[128670]1990.

[128671]23022.

[128672]Mochizuki, T.; Wu, G.; Hayashi, T.; Xenophontos, S. L.;

Veldhuisen,

[128673]B.; Saris, J. J.; Reynolds, D. M.; Cai, Y.; Gabow, P. A.;

Pierides,

[128674]A.; Kimberling, W. J.; Breuning, M. H.; Constantinou

Deltas, C.; Peters,

[128675]D. J. M.; Somlo, S.: PKD2, a gene for polycystic kidney dis-
ease that

[128676]encodes an integral membrane protein. Science 272:

1339–1342, 1996.

[128677]23023.

[128678]Newby, L. J.; Streets, A. J.; Zhao, Y.; Harris, P. C.; Ward, C.

[128679]J.; Ong, A. C.: Identification, characterization, and localization

[128680]of a novel kidney polycystin-1-polycystin-2 complex. J. Biol. Chem. 277:

[128681]20763-20773, 2002.

[128682]23024.

[128683]Norby, S.; Schwartz, M.: Possible locus for polycystic kidney

[128684]disease on chromosome 2. (Letter) Lancet 336: 323-324, 1990.

[128685]23025.

[128686]Parfrey, P. S. N.; Bear, J. C.; Morgan, J.; Cramer, B. C.; McManamon,

[128687]P. J.; Gault, M. H.; Churchill, D.; Singh, M.; Hewitt, R.; Somlo,

[128688]S.; Reeders, S.: The diagnosis and prognosis of autosomal dominant

[128689]polycystic kidney disease. New Eng. J. Med. 323: 1085-1090, 1990.

[128690]23026.

[128691]Pei, Y.; He, N.; Wang, K.; Kasenda, M.; Paterson, A. D.;
Chan,

[128692]G.; Liang, Y.; Roscoe, J.; Brissenden, J.; Hefferton, D.; Par-
frey,

[128693]P.; Somlo, S.; St. George-Hyslop, P.: A spectrum of muta-
tions in

[128694]the polycystic kidney disease-2 (PKD2) gene from eight
Canadian kindreds. J.

[128695]Am. Soc. Nephrol. 9: 1853-1860, 1998.

[128696]23027.

[128697]Pei, Y.; Paterson, A. D.; Wang, K. R.; He, N.; Hefferton, D.;

[128698]Watnick, T.; Germino, G. G.; Parfrey, P.; Somlo, S.; St.
George-Hyslop,

[128699]P.: Bilineal disease and trans-heterozygotes in autosomal
dominant

[128700]polycystic kidney disease. Am. J. Hum. Genet. 68:
355-363, 2001.

[128701]23028.

[128702]Pei, Y.; Wang, K.; Kasenda, M.; Paterson, A. D.; Liang, Y.;
Huang,

[128703]E.; Lian, J.; Rogovea, E.; Somlo, S.; St. George-Hyslop, P.: A novel

[128704]frameshift mutation induced by an adenosine insertion in the polycystic

[128705]kidney disease 2 (PKD2) gene. *Kidney Int.* 53: 1127–1132, 1998.

[128706]23029.

[128707]Cleutjens, K. B. J. M.; van Eekelen, C. C. E. M.; van der Ko-rput,

[128708]H. A. G. M.; Brinkmann, A. O.; Trapman, J.: Two androgen response

[128709]regions cooperate in steroid hormone regulated activity of the prostate-specific

[128710]antigen promoter. *J. Biol. Chem.* 271: 6379–6388, 1996.

[128711]23030.

[128712]Evans, B. A.; Drinkwater, C. C.; Richards, R. I.: Mouse glandular

[128713]kallikrein genes: structure and partial sequence analysis of the kallikrein

[128714]gene locus. *J. Biol. Chem.* 262: 8027–8034, 1987.

[128715]23031.

- [128716]Lundwall, A.; Lilja, H.: Molecular cloning of human prostate specific antigen cDNA. FEBS Lett. 214: 317–322, 1987.
- [128718]23032.
- [128719]Melegos, D. N.; Yu, H.; Ashok, M.; Wang, C.; Stanczyk, F.; Diamandis, E. P.: Prostate-specific antigen in female serum, a potential new marker of androgen excess. J. Clin. Endocr. Metab. 82: 777–780, 1997.
- [128722]1997.
- [128723]23033.
- [128724]Prendergast, G. C.; Lawe, D.; Ziff, E. B.: Association of Myn, the murine homolog of Max, with c-Myc stimulates methylation-sensitive DNA binding and Ras cotransformation. Cell 65: 395–408, 1991.
- [128727]23034.
- [128728]Wagner, A. J.; Le Beau, M. M.; Diaz, M. O.; Hay, N.: Expression,

[128729]regulation, and chromosomal localization of the Max gene. Proc. Nat.

[128730]Acad. Sci. 89: 3111–3115, 1992.

[128731]23035.

[128732]Zervos, A. S.; Faccio, L.; Gatto, J. P.; Kyriakis, J. M.; Brent,

[128733]R.: Mxi2, a mitogen-activated protein kinase that recognizes and

[128734]phosphorylates Max protein. Proc. Nat. Acad. Sci. 92: 10531–10534,

[128735]1995.

[128736]23036.

[128737]Emi, M.; Katagiri, T.; Harada, Y.; Saito, H.; Inazawa, J.; Ito,

[128738]I.; Kasumi, F.; Nakamura, Y.: A novel metalloprotease/disintegrin-like

[128739]gene at 17q21.3 is somatically rearranged in two primary breast cancers. Nature

[128740]Genet. 5: 151–157, 1993.

[128741]23037.

[128742]Katagiri, T.; Harada, Y.; Emi, M.; Nakamura, Y.: Human metalloprotease/disintegrin-like

[128743](MDC) gene: exon-intron organization and alternative splicing. Cytogenet.

[128744]Cell Genet. 68: 39-44, 1995.

[128745]23038.

[128746]Begley, C. G.; Lipkowitz, S.; Gobel, V.; Mahon, K. A.; Bertness,

[128747]V.; Green, A. R.; Gough, N. M.; Kirsch, I. R.: Molecular characterization

[128748]of NSCL, a gene encoding a helix-loop-helix protein expressed in the

[128749]developing nervous system. Proc. Nat. Acad. Sci. 89: 38-42, 1992.

[128750]23039.

[128751]Brown, L.; Espinosa, R., III; Le Beau, M. M.; Siciliano, M. J.;

[128752]Baer, R.: HEN1 and HEN2: a subgroup of basic helix-loop-helix genes

[128753]that are coexpressed in a human neuroblastoma. Proc. Nat. Acad. Sci. 89:

[128754]8492-8496, 1992.

[128755]23040.

[128756]Cogliati, T.; Good, D. J.; Haigney, M.; Delgado-Romero, P.;

Eckhaus,

[128757]M. A.; Koch, W. J.; Kirsch, I. R.: Predisposition to arrhythmia and

[128758]autonomic dysfunction in Nhlh1-deficient mice. Molec. Cell. Biol. 22:

[128759]4977-4983, 2002.

[128760]23041.

[128761]Lipkowitz, S.; Gobel, V.; Varterasian, M. L.; Nakahara, K.; Tchorz,

[128762]K.; Kirsch, I. R.: A comparative structural characterization of the

[128763]human NSCL-1 and NSCL-2 genes: two basic helix-loop-helix genes expressed

[128764]in the developing nervous system. J. Biol. Chem. 267: 21065-21071,

[128765]1992.

[128766]23042.

[128767]Mullick, A.; Groulx, N.; Trasler, D.; Gros, P.: Nhlh1, a basic

[128768]helix-loop-helix transcription factor, is very tightly linked to the

[128769]mouse looptail (Lp) mutation. Mammalian Genome 6:

700–704, 1995.

[128770]23043.

[128771]Meeker, T. C.; Nagarajan, L.; ar-Rushdi, A.; Rovera, G.; Huebner,

[128772]K.; Croce, C. M.: Characterization of the human PIM-1 gene: a putative

[128773]proto-oncogene coding for a tissue specific member of the protein

[128774]kinase family. *Oncogene Res.* 1: 87–101, 1987.

[128775]23044.

[128776]Pasqualucci, L.; Neumeister, P.; Goossens, T.; Nanjangud, G.; Chaganti,

[128777]R. S. K.; Kuppers, R.; Dalla-Favera, R.: Hypermutation of multiple

[128778]proto-oncogenes in B-cell diffuse large-cell lymphomas. *Nature* 412:

[128779]341–346, 2001.

[128780]23045.

[128781]Ragoussis, J.; Senger, G.; Mockridge, I.; Sanseau, P.; Ruddy, S.;

[128782]Dudley, K.; Sheer, D.; Trowsdale, J.: A testis-expressed Zn

finger

[128783] gene (ZNF76) in human 6p21.3 centromeric to the MHC is
closely linked

[128784] to the human homolog of the t-complex gene tcp-11.

Genomics 14:

[128785] 673-679, 1992.

[128786] 23046.

[128787] Saris, C. J. M.; Domen, J.; Berns, A.: The pim-1 oncogene
encodes

[128788] two related protein-serine/threonine kinases by alterna-
tive initiation

[128789] at AUG and CUG. EMBO J. 10: 655-664, 1991.

[128790] 23047.

[128791] Selten, G.; Cuypers, H. T.; Boelens, W.; Robanus-Maandag,
E.;

[128792] Verbeek, J.; Domen, J.; van Beveren, C.; Berns, A.: The pri-
mary structure

[128793] of the putative oncogene pim-1 shows extensive homol-
ogy with protein

[128794] kinases. Cell 46: 603-611, 1986.

[128795] 23048.

[128796]Zoghbi, H. Y.; Ballantyne, C. M.; O'Brien, W. E.; McCall, A. E.;

[128797]Kwiatkowski, T. J., Jr.; Ledbetter, S. A.; Beaudet, A. L.:
Deletion

[128798]and linkage mapping of eight markers from the proximal
short arm of

[128799]chromosome 6. Genomics 6: 352–357, 1990.

[128800]23049.

[128801]Knebelmann, B.; Forestier, L.; Drouot, L.; Quinones, S.;
Chuet,

[128802]C.; Benessy, F.; Saus, J.; Antignac, C.: Splice-mediated in-
sertion

[128803]of an Alu sequence in the COL4A3 mRNA causing autosomal
recessive

[128804]Alport syndrome. Hum. Molec. Genet. 4: 675–679, 1995.

[128805]23050.

[128806]Gromoll, J.; Partsch, C.-J.; Simoni, M.; Nordhoff, V.; Sip-
pell,

[128807]W. G.; Nieschlag, E.; Saxena, B. B.: A mutation in the first
transmembrane

[128808]domain of the lutropin receptor causes male precocious
puberty. J.

[128809]Clin. Endocr. Metab. 83: 476–480, 1998.

[128810]23051.

[128811]Kawate, N.; Kletter, G. B.; Wilson, B. E.; Netzloff, M. L.;
Menon,

[128812]K. M. J.: Identification of constitutively activating mutation
of

[128813]the luteinising hormone receptor in a family with male
limited gonadotrophin

[128814]independent precocious puberty (testotoxicosis). J. Med.
Genet. 32:

[128815]553–554, 1995.

[128816]23052.

[128817]Kosugi, S.; Van Dop, C.; Geffner, M. E.; Rabl, W.; Carel,
J.–C.;

[128818]Chaussain, J.–L.; Mori, T.; Merendino, J. J., Jr.; Shenker, A.:
Characterization

[128819]of heterogeneous mutations causing constitutive activa-
tion of the

[128820]luteinizing hormone receptor in familial male precocious
puberty. Hum.

[128821]Molec. Genet. 4: 183–188, 1995.

[128822]23053.

[128823]Kremer, H.; Kraaij, R.; Toledo, S. P. A.; Post, M.; Fridman, J.

[128824]B.; Hayashida, C. Y.; van Reen, M.; Milgrom, E.; Ropers,
H.-H.; Mariman,

[128825]E.; Themmen, A. P. N.; Brunner, H. G.: Male pseudo-
hermaphroditism

[128826]due to a homozygous missense mutation of the luteiniz-
ing hormone receptor

[128827]gene. *Nature Genet.* 9: 160-164, 1995.

[128828]23054.

[128829]Kremer, H.; Mariman, E.; Otten, B. J.; Moll, G. W., Jr.;
Stoelinga,

[128830]G. B. A.; Wit, J. M.; Jansen, M.; Drop, S. L.; Faas, B.; Ropers,
H.-H.;

[128831]Brunner, H. G.: Cosegregation of missense mutations of
the luteinizing

[128832]hormone receptor gene with familial male-limited preco-
cious puberty. *Hum.*

[128833]*Molec. Genet.* 2: 1779-1783, 1993.

[128834]23055.

[128835]Kremer, H.; Martens, J. W. M.; van Reen, M.; Verhoef-Post,

M.;

[128836]Wit, J. M.; Otten, B. J.; Drop, S. L. S.; Delemarre-van de
Waal, H.

[128837]A.; Pombo-Arias, M.; De Luca, F.; Potau, N.; Buckler, J. M.
H.; and

[128838]9 others: A limited repertoire of mutations of the luteiniz-
ing hormone

[128839](LH) receptor gene in familial and sporadic patients with
male LH-independent

[128840]precocious puberty. J. Clin. Endocr. Metab. 84:
1136-1140, 1999.

[128841]23056.

[128842]Latronico, A. C.; Abell, A. N.; Arnhold, I. J.; Liu, X.; Lins,

[128843]T. S. S.; Brito, V. N.; Billerbeck, A. E.; Segaloff, D. L.; Men-
donca,

[128844]B. B.: A unique constitutively activating mutation in third
transmembrane

[128845]helix of luteinizing hormone receptor causes sporadic
male gonadotropin-independent

[128846]precocious puberty. J. Clin. Endocr. Metab. 83:
2435-2440, 1998.

[128847]23057.

[128848]Latronico, A. C.; Anasti, J.; Arnhold, I. J. P.; Mendonca, B.

[128849]B.; Domenice, S.; Albano, M. C.; Zachman, K.; Wajchenberg, B. L.;

[128850]Tsigos, C.: A novel mutation of the luteinizing hormone receptor

[128851]gene causing male gonadotropin-independent precocious puberty. J.

[128852]Clin. Endocr. Metab. 80: 2490–2494, 1995.

[128853]23058.

[128854]Latronico, A. C.; Anasti, J.; Arnhold, I. J. P.; Rapaport, R.;

[128855]Mendonca, B. B.; Bloise, W.; Castro, M.; Tsigos, C.; Chrousos, G.

[128856]P.: Testicular and ovarian resistance to luteinizing hormone caused

[128857]by inactivating mutations of the luteinizing hormone-receptor gene. New

[128858]Eng. J. Med. 334: 507–512, 1996.

[128859]23059.

[128860]Laue, L.; Chan, W.-Y.; Hsueh, A. J. W.; Kudo, M.; Hsu, S. Y.;

[128861]Wu, S.-M.; Blomberg, L.; Cutler, G. B., Jr.: Genetic heterogeneity

[128862]of constitutively activating mutations of the human
luteinizing hormone
[128863]receptor in familial male-limited precocious puberty. Proc.
Nat.
[128864]Acad. Sci. 92: 1906-1910, 1995.
[128865]23060.
[128866]Laue, L.; Wu, S.-M.; Kudo, M.; Hsueh, A. J. W.; Cutler, G.
B.,
[128867]Jr.; Griffin, J. E.; Wilson, J. D.; Brain, C.; Berry, A. C.; Grant,
[128868]D. B.; Chan, W.-Y.: A nonsense mutation of the human
luteinizing
[128869]hormone receptor gene in Leydig cell hypoplasia. Hum.
Molec. Genet. 4:
[128870]1429-1433, 1995.
[128871]23061.
[128872]Lee, C. Y.; Ryan, R. J.: The uptake of human luteinizing
hormone
[128873](NLH) by slices of luteinized rat ovaries. Endocrinology 89:
1515-1523,
[128874]1971.
[128875]23062.

[128876]Latronico, A. C.; Chai, Y.; Arnhold, I. J. P.; Liu, X.; Mendonca,
 [128877]B. B.; Segaloff, D. L.: A homozygous microdeletion in helix
 7 of
 [128878]the luteinizing hormone receptor associated with familial
 testicular
 [128879]and ovarian resistance is due to both decreased cell sur-
 face expression
 [128880]and impaired effector activation by the cell surface recep-
 tor. Molec.
 [128881]Endocr. 12: 442-450, 1998.
 [128882]23063.
 [128883]Latronico, A. C.; Shinozaki, H.; Guerra, G., Jr.; Pereira, M.
 [128884]A. A.; Helena, S.; Lemos Marini, V.; Baptista, M. T. M.; Arn-
 hold,
 [128885]I. J. P.; Fanelli, F.; Mendonca, B. B.; Segaloff, D. L.: Go-
 nadotropin-independent
 [128886]precocious puberty due to luteinizing hormone receptor
 mutations in
 [128887]Brazilian boys: a novel constitutively activating mutation
 in the
 [128888]first transmembrane helix. J. Clin. Endocr. Metab. 85:

4799–4805,

[128889]2000.

[128890]23064.

[128891]Lee, C. Y.; Ryan, R. J.: Luteinizing hormone receptors: specific

[128892]binding of human luteinizing hormone to homogenates of
luteinized

[128893]rat ovaries. Proc. Nat. Acad. Sci. 69: 3520–3523, 1972.

[128894]23065.

[128895]Liu, G.; Duranteau, L.; Carel, J.-C.; Monroe, J.; Doyle, D. A.;

[128896]Shenker, A.: Leydig-cell tumors caused by an activating
mutation

[128897]of the gene encoding the luteinizing hormone receptor.
New Eng. J.

[128898]Med. 341: 1731–1736, 1999.

[128899]23066.

[128900]Loosfelt, H.; Misrahi, M.; Atger, M.; Salesse, R.; Vu Hai-Luu

[128901]Thi, M. T.; Jolivet, A.; Guiochon-Mantel, A.; Sar, S.; Jallal,
B.;

[128902]Garnier, J.; Milgrom, E.: Cloning and sequencing of

porcine LH-hCG

[128903]receptor cDNA: variants lacking transmembrane domain.

Science 245:

[128904]525–528, 1989.

[128905]23067.

[128906]Martens, J. W. M.; Verhoef-Post, M.; Abelin, N.; Erzabella,
M.;

[128907]Toledo, S. P. A.; Brunner, H. G.; Themmen, A. P. N.: A ho-
mozygous

[128908]mutation in the luteinizing hormone receptor causes par-
tial Leydig

[128909]cell hypoplasia: correlation between receptor activity and
phenotype. Molec.

[128910]Endocr. 12: 775–784, 1998.

[128911]23068.

[128912]McFarland, K. C.; Sprengel, R.; Phillips, H. S.; Kohler, M.;
Rosembliit,

[128913]N.; Nikolics, K.; Segaloff, D. L.; Seeburg, P. H.: Lutropin-
choriogonadotropin

[128914]receptor: an unusual member of the G protein-coupled
receptor family. Science 245:

[128915]494–499, 1989.

[128916]23069.

[128917]Min, L.; Ascoli, M.: Effect of activating and inactivating mutations

[128918]on the phosphorylation and trafficking of the human lutropin/choriogonadotropin

[128919]receptor. *Molec. Endocr.* 14: 1797–1810, 2000.

[128920]23070.

[128921]Minegishi, T.; Nakamura, K.; Takakura, Y.; Miyamoto, K.; Hasegawa,

[128922]Y.; Ibuki, Y.; Igarashi, M.: Cloning and sequencing of human LH/hCG

[128923]receptor cDNA. *Biochem. Biophys. Res. Commun.* 172: 1049–1054, 1990.

[128924]23071.

[128925]Chapin, S. J.; Lue, C.-M.; Yu, M. T.; Bulinski, J. C.: Differential

[128926]expression of alternatively spliced forms of MAP4: a repertoire of

[128927]structurally different microtubule-binding domains. *Biochemistry* 34:

[128928]2289–2301, 1995.

[128929]23072.

[128930]Laird, P. W.; van der Lugt, N. M. T.; Clarke, A.; Domen, J.;
Linders,

[128931]K.; McWhir, J.; Berns, A.; Hooper, M.: In vivo analysis of
Pim-1

[128932]deficiency. Nucleic Acids Res. 21: 4750-4755, 1993.

[128933]23073.

[128934]Gartner, J.; Obie, C.; Watkins, P.; Valle, D.: Restoration of

[128935]peroxisome biogenesis in a peroxisome-deficient mam-
malian cell line

[128936]by expression of either the 35 kDa or the 70 kDa peroxi-
somal membrane

[128937]proteins. J. Inherit. Metab. Dis. 17: 327-329, 1994.

[128938]23074.

[128939]Coto, E.; Sanz de Castro, S.; Aguado, S.; Alvarez, J.; Arias,
M.;

[128940]Menendez, M. J.; Lopez-Larrea, C.: DNA microsatellite
analysis of

[128941]families with autosomal dominant polycystic kidney dis-
ease types 1

[128942]and 2: evaluation of clinical heterogeneity between both

forms of

[128943]the disease. J. Med. Genet. 32: 442–445, 1995.

[128944]23075.

[128945]Maric, S. C.; Crozat, A.; Janne, O. A.: Structure and organization

[128946]of the human S-adenosylmethionine decarboxylase gene. J. Biol. Chem. 267:

[128947]18915–18923, 1992.

[128948]23076.

[128949]Maric, S. C.; Crozat, A.; Louhimo, J.; Knuutila, S.; Janne, O.

[128950]A.: The human S-adenosylmethionine decarboxylase gene: nucleotide

[128951]sequence of a pseudogene and chromosomal localization of the active

[128952]gene (AMD1) and the pseudogene (AMD2). Cytogenet. Cell Genet. 70:

[128953]195–199, 1995.

[128954]23077.

[128955]Mangan, M. E.; Olmsted, J. B.: The gene for microtubule-associated

[128956]protein 4 (Mtap4) maps to the distal region of mouse

chromosome 9. Mammalian

[128957]Genome 7: 918–925, 1996.

[128958]23078.

[128959]West, R. R.; Tenbarger, K. M.; Olmsted, J. B.: A model for
microtubule-associated

[128960]protein 4 structure: domains defined by comparisons of
human, mouse,

[128961]and bovine sequences. J. Biol. Chem. 266: 21886–21896,
1991.

[128962]23079.

[128963]Abel, K. J.; Boehnke, M.; Prahalad, M.; Ho, P.; Flejter, W. L.;

[128964]Watkins, M.; VanderStoep, J.; Chandrasekharappa, S. C.;
Collins, F.

[128965]S.; Glover, T. W.; Weber, B. L.: A radiation hybrid map of
the BRCA1

[128966]region of chromosome 17q12–q21. Genomics 17:
632–641, 1993.

[128967]23080.

[128968]Alonso, A. D. C.; Grundke-Iqbal, I.; Iqbal, K.: Alzheimer's
disease

[128969]hyperphosphorylated tau sequesters normal tau into tan–

gles of filaments

[128970]and disassembles microtubules. Nature Med. 2: 783–787, 1996.

[128971]23081.

[128972]Arima, K.; Kowalska, A.; Hasegawa, M.; Mukoyama, M.; Watanabe,

[128973]R.; Kawai, M.; Takahashi, K.; Iwatsubo, T.; Tabira, T.; Sunohara,

[128974]N.: Two brothers with frontotemporal dementia and parkinsonism with

[128975]an N279K mutation of the tau gene. Neurology 54: 1787–1795, 2000.

[128976]23082.

[128977]Baker, M.; Litvan, I.; Houlden, H.; Adamson, J.; Dickson, D.; Perez–Tur,

[128978]J.; Hardy, J.; Lynch, T.; Bigio, E.; Hutton, M.: Association of an

[128979]extended haplotype in the tau gene with progressive supranuclear palsy. Hum.

[128980]Molec. Genet. 8: 711–715, 1999.

[128981]23083.

[128982]Brown, J.; Lantos, P. L.; Roques, P.; Fidani, L.; Rossor, M.

N.

[128983]: Familial dementia with swollen achromatic neurons and corticobasal

[128984]inclusion bodies: a clinical and pathological study. J. Neurol. Sci. 135:

[128985]21–30, 1996.

[128986]23084.

[128987]Clark, L. N.; Poorkaj, P.; Wszolek, Z.; Geschwind, D. H.; Nasreddine,

[128988]Z. S.; Miller, B.; Li, D.; Payami, H.; Awert, F.; Markopoulou, K.;

[128989]Andreadis, A.; D'Souza, I.; Lee, V. M.-Y.; Reed, L.; Trojanowski,

[128990]J. Q.; Zhukareva, V.; Bird, T.; Schellenberg, G.; Wilhelmsen, K. C.

[128991]: Pathogenic implications of mutations in the tau gene in pallido-ponto-nigral

[128992]degeneration and related neurodegenerative disorders linked to chromosome

[128993]17. Proc. Nat. Acad. Sci. 95: 13103–13107, 1998.

[128994]23085.

[128995]Connell, J. W.; Gibb, G. M.; Betts, J. C.; Blackstock, W. P.;

Gallo,

[128996]J.-M.; Lovestone, S.; Hutton, M.; Anderton, B. H.: Effects of
FTDP-17

[128997]mutations on the in vitro phosphorylation of tau by glyco-
gen synthase

[128998]kinase 3-beta identified by mass spectrometry demon-
strate certain

[128999]mutations exert long-range conformational changes. FEBS
Lett. 493:

[129000]40-44, 2001.

[129001]23086.

[129002]Conrad, C.; Andreadis, A.; Trojanowski, J. Q.; Dickson, D.
W.; Kang,

[129003]D.; Chen, X.; Weiderholt, W.; Hansen, L.; Masliah, E.; Thal,
L. J.;

[129004]Katzman, R.; Xia, Y.; Saitoh, T.: Genetic evidence for the
involvement

[129005]of tau in progressive supranuclear palsy. Ann. Neurol. 41:
277-281,

[129006]1997.

[129007]23087.

[129008]Conrad, C.; Vianna, C.; Freeman, M.; Davies, P.: A poly-

morphic

[129009] gene nested within an intron of the tau gene: implications
for Alzheimer's

[129010] disease. Proc. Nat. Acad. Sci. 99: 7751–7756, 2002.

[129011] 23088.

[129012] Delisle, M. B.; Murrell, J. R.; Richardson, R.; Trofatter, J.

[129013] A.; Rascol, O.; Soulages, X.; Mohr, M.; Calvas, P.; Ghetti,
B.: A

[129014] mutation at codon 279 (N279K) in exon 10 of the tau
gene causes a

[129015] tauopathy with dementia and supranuclear palsy. Acta
Neuropath. 98:

[129016] 62–77, 1999.

[129017] 23089.

[129018] Donlon, T. A.; Harris, P.; Neve, R. L.: Localization of mi-
crotubule-associated

[129019] protein tau (MTBT1) to chromosome 17q21. (Abstract) Cy-
togenet. Cell

[129020] Genet. 46: 607, 1987.

[129021] 23090.

[129022] Goedert, M.; Crowther, R. A.; Spillantini, M. G.: Tau muta-

tions

[129023]cause frontotemporal dementias. Neuron 21: 955–958, 1998.

[129024]23091.

[129025]Goedert, M.; Spillantini, M. G.; Potier, M. C.; Ulrich, J.; Crowther,

[129026]R. A.: Cloning and sequencing of the cDNA encoding an isoform of

[129027]microtubule-associated protein tau containing four tandem repeats:

[129028]differential expression of tau protein mRNAs in human brain. EMBO

[129029]J. 8: 393–399, 1989.

[129030]23092.

[129031]Goedert, M.; Wischik, C. M.; Crowther, R. A.; Walker, J. E.; Klug,

[129032]A.: Cloning and sequencing of the cDNA encoding a core protein of

[129033]the paired helical filament of Alzheimer disease: identification as

[129034]the microtubule-associated protein tau. Proc. Nat. Acad. Sci. 85:

[129035]4051–4055, 1988.

[129036]23093.

[129037]Goode, B. L.; Chau, M.; Denis, P. E.; Feinstein, S. C.: Structural

[129038]and functional differences between 3-repeat and 4-repeat tau isoforms:

[129039]implications for normal tau function and the onset of neurodegenerative

[129040]disease. J. Biol. Chem. 275: 38182–38189, 2000.

[129041]23094.

[129042]Heutink, P.: Untangling tau-related dementia. Hum. Molec. Genet. 9:

[129043]979–986, 2000.

[129044]23095.

[129045]Heutink, P.; Stevens, M.; Rizzu, P.; Bakker, E.; Kros, J. M.;

[129046]Tibben, A.; Niermeijer, M. F.; van Duijn, C. M.; Oostra, B. A.; van

[129047]Swieten, J. C.: Hereditary frontotemporal dementia is linked to chromosome

[129048]17q21–q22: a genetic and clinicopathological study of three Dutch

[129049]families. Ann. Neurol. 41: 150–159, 1997.

[129050]23096.

[129051]Hiesberger, T.; Trommsdorff, M.; Howell, B. W.; Goffinet, A.;

[129052]Mumby, M. C.; Cooper, J. A.; Herz, J.: Direct binding of reelin to

[129053]VLDL receptor and apoE receptor 2 induces tyrosine phosphorylation

[129054]of disabled-1 and modulates tau phosphorylation. Neuron 24: 481–489,

[129055]1999.

[129056]23097.

[129057]Hong, M.; Zhukareva, V.; Vogelsberg–Ragaglia, V.; Ws–zolek, Z.;

[129058]Reed, L.; Miller, B. I.; Geschwind, D. H.; Bird, T. D.; McKeel, D.;

[129059]Goate, A.; Morris, J. C.; Wilhelmsen, K. C.; Schellenberg, G. D.;

[129060]Trojanowski, J. Q.; Lee, V. M.–Y.: Mutation–specific functional impairments

[129061]in distinct tau isoforms of hereditary FTDP–17. Science 282: 1914–1917,

[129062]1998.

[129063]23098.

[129064]Hutton, M.: Missense and splice site mutations in tau associated

[129065]with FTDP-17: multiple pathogenic mechanisms. Neurology 56 (suppl.

[129066]4): S21-S25, 2001.

[129067]23099.

[129068]Hutton, M.; Lendon, C. L.; Rizzu, P.; Baker, M.; Froelich, S.;

[129069]Houlden, H.; Pickering-Brown, S.; Chakraverty, S.; Isaacs, A.; Grover,

[129070]A.; Hackett, J.; Adamson, J.; and 39 others: Association of missense

[129071]and 5-prime-splice-site mutations in tau with the inherited dementia

[129072]FTDP-17. Nature 393: 702-705, 1998.

[129073]23100.

[129074]Iijima, M.; Tabira, T.; Poorkaj, P.; Schellenberg, G. D.; Trojanowski,

[129075]J. Q.; Lee, V. M.; Schmidt, M. L.; Takahashi, K.; Nabika, T.; Matsumoto,

[129076]T.; Yamashita, Y.; Yoshioka, S.; Ishino, H.: A distinct familial
 [129077]presenile dementia with a novel missense mutation in the
 tau gene. Neuroreport 10:
 [129078]497–501, 1999.
 [129079]23101.

[129080]Ishihara, T.; Hong, M.; Zhang, B.; Nakagawa, Y.; Lee, M. K.;
 Trojanowski,
 [129081]J. Q.; Lee, V. M.-Y.: Age-dependent emergence and progression of
 [129082]a tauopathy in transgenic mice overexpressing the shortest human tau
 [129083]isoform. Neuron 24: 751–762, 1999.
 [129084]23102.

[129085]Janssen, J. C.; Warrington, E. K.; Morris, H. R.; Lantos, P.;
 [129086]Brown, J.; Revesz, T.; Wood, N.; Khan, M. N.; Cipolotti, L.;
 Fox,
 [129087]N. C.; Rossor, M. N.: Clinical features of frontotemporal
 dementia
 [129088]due to the intronic tau 10 +16 mutation. Neurology 58:
 1161–1168,

[129089]2002.

[129090]23103.

[129091]Braun, T.; Grzeschik, K.-H.; Bober, E.; Arnold, H.-H.: The
MYF

[129092]genes, a group of human muscle determining factors, are
localized

[129093]on different human chromosomes. (Abstract) Cytogenet.
Cell Genet. 51:

[129094]969 only, 1989.

[129095]23104.

[129096]Henry, I.; Puech, A.; Antignac, C.; Couillin, P.; Jeanpierre,
M.;

[129097]Ahnine, L.; Barichard, F.; Boehm, T.; Augereau, P.; Scrable,
H.; Rabbitts,

[129098]T. H.; Rochefort, H.; Cavenee, W.; Junien, C.: Subregional
mapping

[129099]of BWS, CTSD, MYOD1, and a T-ALL breakpoint in 11p15.
(Abstract) Cytogenet.

[129100]Cell Genet. 51: 1013 only, 1989.

[129101]23105.

[129102]Kim, Y.-J.; Noguchi, S.; Hayashi, Y. K.; Tsukahara, T.;

Shimizu,

[129103]T.; Arahata, K.: The product of an oculopharyngeal muscular dystrophy

[129104]gene, poly(A)-binding protein 2, interacts with SKIP and stimulates

[129105]muscle-specific gene expression. Hum. Molec. Genet. 10: 1129-1139,

[129106]2001.

[129107]23106.

[129108]Olson, E. N.: MyoD family: a paradigm for development. Genes

[129109]Dev. 4: 1454-1461, 1990.

[129110]23107.

[129111]Troger, J.; Neyer, S.; Heufler, C.; Huemer, H.; Schmid, E.; Griesser,

[129112]U.; Kralinger, M.; Kremser, B.; Baldissera, I.; Kieselbach, G.: Substance

[129113]P and vasoactive intestinal polypeptide in the streptozotocin-induced

[129114]diabetic rat retina. Invest. Ophthalm. Vis. Sci. 42: 1045-1050, 2001.

[129115]23108.

[129116]Castagliuolo, I.; Riegler, M.; Pasha, A.; Nikulasson, S.; Lu, B.;

[129117]Gerard, C.; Gerard, N. P.; Pothoulakis, C.: Neurokinin-1 (NK-1) receptor

[129118]is required in Clostridium difficile-induced enteritis. J. Clin.

[129119]Invest. 101: 1547-1550, 1998.

[129120]23109.

[129121]De Felipe, C.; Herrero, J. F.; O'Brien, J. A.; Palmer, J. A.; Doyle,

[129122]C. A.; Smith, A. J. H.; Laird, J. M. A.; Belmonte, C.; Cervero, F.;

[129123]Hunt, S. P.: Altered nociception, analgesia and aggression in mice

[129124]lacking the receptor for substance P. Nature 392: 394-397, 1998.

[129125]23110.

[129126]Gerard, N. P.; Garraway, L. A.; Eddy, R. L., Jr.; Shows, T. B.;

[129127]Iijima, H.; Paquet, J. L.; Gerard, C.: Human substance P receptor

[129128](NK-1): organization of the gene, chromosome localization, and functional

[129129]expression of cDNA clones. Biochemistry 30:
10640–10646, 1991.

[129130]23111.

[129131]Hopkins, B.; Powell, S. J.; Danks, P.; Briggs, I.; Graham, A.:

[129132]Isolation and characterisation of the human lung NK-1 re-
ceptor cDNA. Biochem.

[129133]Biophys. Res. Commun. 180: 1110–1117, 1991.

[129134]23112.

[129135]Murtra, P.; Sheasby, A. M.; Hunt, S. P.; De Felipe, C.: Re-
warding

[129136]effects of opiates are absent in mice lacking the receptor
for substance

[129137]P. Nature 45: 180–183, 2000.

[129138]23113.

[129139]Page, N. M.; Woods, R. J.; Gardiner, S. M.; Lomthaisong, K.;
Gladwell,

[129140]R. T.; Butlin, D. J.; Manyonda, I. T.; Lowry, P. J.: Excessive
placental

[129141]secretion of neurokinin B during the third trimester causes
pre-eclampsia. Nature 405:

[129142]797–800, 2000.

[129143]23114.

[129144]Ishikawa, T.; Umesono, K.; Mangelsdorf, D. J.; Aburatani, H.; Stanger,

[129145]B. Z.; Shibasaki, Y.; Imawari, M.; Evans, R. M.; Takaku, F.:
A functional

[129146]retinoic acid receptor encoded by the gene on human
chromosome 12.

[129147]Molec. Endocr. 4: 837-844, 1990.

[129148]23115.

[129149]Krust, A.; Kastner, P.; Petkovich, M.; Zelent, A.; Chambon,
P.

[129150]: A third human retinoic acid receptor, hRAR-gamma.
Proc. Nat. Acad.

[129151]Sci. 86: 5310-5314, 1989.

[129152]23116.

[129153]Mattei, M.-G.; Riviere, M.; Krust, A.; Ingvarsson, S.;
Vennstrom,

[129154]B.; Islam, M. Q.; Levan, G.; Kautner, P.; Zelent, A.; Cham-
bon, P.;

[129155]Szpirer, J.; Szpirer, C.: Chromosomal assignment of
retinoic acid

[129156]receptor (RAR) genes in the human, mouse, and rat
genomes. Genomics 10:

[129157]1061–1069, 1991.

[129158]23117.

[129159]Ryseck, R.-P.; Macdonald-Bravo, H.; Mattei, M. G.;
Siegfried, R.

[129160]L.; Bravo, R.: Structure, mapping and expression of a
growth factor

[129161]inducible gene encoding a putative nuclear hormonal
binding receptor.

[129162]EMBO J. 8: 3327–3335, 1989.

[129163]23118.

[129164]Abramson, D. H.; Ellsworth, R. M.; Zimmerman, L. E.:
Monocular

[129165]cancer in retinoblastoma survivors. Trans. Am. Acad.
Ophthalmol. Otolaryng. 81:

[129166]454–457, 1976.

[129167]23119.

[129168]Aherne, G. E. S.; Roberts, D. F.: Retinoblastoma--a clinical
survey

[129169]and its genetic implications. Clin. Genet. 8: 275–290,

1975.

[129170]23120.

[129171]Alonso, J.; Garcia-Miguel, P.; Abelairas, J.; Mendiola, M.;
Sarret,

[129172]E.; Vendrell, M. T.; Navajas, A.; Pestana, A.: Spectrum of
germline

[129173]RB1 gene mutations in Spanish retinoblastoma patients:
phenotypic

[129174]and molecular epidemiological implications. Hum. Mutat.
17: 412-422,

[129175]2001.

[129176]23121.

[129177]Amoaku, W. M. K.; Willshaw, H. E.; Parkes, S. E.; Shah, K. J.;

[129178]Mann, J. R.: Trilateral retinoblastoma: a report of five pa-
tients. Cancer 78:

[129179]858-863, 1996.

[129180]23122.

[129181]Bader, J. L.; Meadows, A. T.; Zimmerman, L. E.; Rorke, L.
B.; Voute,

[129182]P. A.; Champion, L. A. A.; Miller, R. W.: Bilateral
retinoblastoma

- [129183]with ectopic intracranial retinoblastoma: trilateral retinoblastoma. Cancer
- [129184]Genet. Cytogenet. 5: 203–213, 1982.
- [129185]23123.
- [129186]Balaban–Malenbaum, G.; Gilbert, F.; Nichols, W. W.; Hill, R.; Shields, J.; Meadows, A. T.: A deleted chromosome no. 13 in human retinoblastoma cells: relevance to tumorigenesis. Cancer Genet. Cytogenet. 3: 243–250, 1981.
- [129189]1981.
- [129190]23124.
- [129191]Bandara, L. R.; Adamczewski, J. P.; Hunt, T.; La Thangue, N. B.
- [129192]: Cyclin A and the retinoblastoma gene product complex with a common transcription factor. Nature 352: 249–251, 1991.
- [129194]23125.
- [129195]Benedict, W. F.; Murphree, A. L.; Banerjee, A.; Spina, C. A.; Sparkes, M. C.; Sparkes, R. S.: Patient with 13 chromosome dele-

tion: evidence

[129197] that the retinoblastoma gene is a recessive cancer gene.

Science 219:

[129198] 973–975, 1983.

[129199] 23126.

[129200] Benedict, W. F.; Xu, H.-J.; Hu, S.-X.; Takahashi, R.: Role of

[129201] the retinoblastoma gene in the initiation and progression
of human

[129202] cancer. J. Clin. Invest. 85: 988–993, 1990.

[129203] 23127.

[129204] Bia, B.; Cowell, J. K.: Independent constitutional germline
mutations

[129205] occurring in the RB1 gene in cousins with bilateral
retinoblastoma. Oncogene 11:

[129206] 977–979, 1995.

[129207] 23128.

[129208] Blanquet, V.; Turleau, C.; de Grouchy, J.; Creau-Goldberg,
N.

[129209]: Physical map around the retinoblastoma gene: possible
genomic imprinting

[129210] suggested by Nrul digestion. Genomics 10: 350–355,

1991.

[129211]23129.

[129212]Blanquet, V.; Turleau, C.; Gross-Morand, M. S.; Sena-
maud-Beaufort,

[129213]C.; Doz, F.; Besmond, C.: Spectrum of germline mutations
in the RB1

[129214]gene: a study of 232 patients with hereditary and non
hereditary retinoblastoma. Hum.

[129215]Molec. Genet. 4: 383-388, 1995.

[129216]23130.

[129217]Jiang, Z.; Cote, J.; Kwon, J. M.; Goate, A. M.; Wu, J. Y.:
Aberrant

[129218]splicing of tau pre-mRNA caused by intronic mutations
associated with

[129219]the inherited dementia frontotemporal dementia with
Parkinsonism linked

[129220]to chromosome 17. Molec. Cell. Biol. 20: 4036-4048,
2000.

[129221]23131.

[129222]Magenis, R. E.; Smith, L.; Nadeau, J. H.; Johnson, K. R.;
Mountjoy,

[129223]K. G.; Cone, R. D.: Mapping of the ACTH, MSH, and neural

(MC3 and

[129224]MC4) melanocortin receptors in the mouse and human.

Mammalian Genome 5:

[129225]503–508, 1994.

[129226]23132.

[129227]Futterweit, W.; Ritch, R.; Teekhasaene, C.; Nelson, E. S.:

Coexistence

[129228]of Prader–Willi syndrome, congenital ectropion uveae with
glaucoma,

[129229]and factor XI deficiency. J.A.M.A. 255: 3280–3282, 1986.

[129230]23133.

[129231]Chow, D.; He, X.; Snow, A. L.; Rose–John, S.; Garcia, K. C.:

Structure

[129232]of an extracellular gp130 cytokine receptor signaling
complex. Science 291:

[129233]2150–2155, 2001.

[129234]23134.

[129235]Funatsu, H.; Yamashita, H.; Noma, H.; Mimura, T.; Ya–
mashita, T.;

[129236]Hori, S.: Increased levels of vascular endothelial growth
factor

[129237]and interleukin-6 in the aqueous humor of diabetics with
macular edema. Am.

[129238]J. Ophthalm. 133: 70-77, 2002.

[129239]23135.

[129240]Kelsell, D. P.; Risk, J. M.; Leigh, I. M.; Stevens, H. P.; Ellis,

[129241]A.; Hennies, H. C.; Reis, A.; Weissenbach, J.; Bishop, D. T.;
Spurr,

[129242]N. K.; Field, J. K.: Close mapping of the focal non-
epidermolytic

[129243]palmoplantar keratoderma (PPK) locus associated with oe-
sophageal cancer

[129244](TOC). Hum. Molec. Genet. 5: 857-860, 1996.

[129245]23136.

[129246]McLean, W. H. I.: Personal Communication. Philadelphia,
Pa.

[129247]2/26/1997.

[129248]23137.

[129249]McLean, W. H. I.; Rugg, E. L.; Lunny, D. P.; Morley, S. M.;
Lane,

[129250]E. B.; Swensson, O.; Dopping-Hepenstal, P. J. C.; Griffiths,
W. A.

- [129251]D.; Eady, R. A. J.; Higgins, C.; Navsaria, H. A.; Leigh, I. M.;
Strachan,
- [129252]T.; Kunkeler, L.; Munro, C. S.: Keratin 16 and keratin 17
mutations
- [129253]cause pachyonychia congenita. *Nature Genet.* 9: 273–278,
1995.
- [129254]23138.
- [129255]Shamsheer, M. K.; Navsaria, H. A.; Stevens, H. P.; Ratnavel,
R.
- [129256]C.; Purkis, P. E.; McLean, W. H.; Cook, L. J.; Griffiths, W. A.
D.;
- [129257]Geschmeissner, S.; Spurr, N.; Leigh, I. M.: Novel mutations
in keratin
- [129258]16 gene underly focal nonepidermolytic palmoplantar
keratoderma (NEPPK)
- [129259]in two families. *Hum. Mol. Genet.* 4: 1875–1881, 1995.
- [129260]23139.
- [129261]Smith, F. J. D.; Del Monaco, M.; Steijlen, P. M.; Munro, C.
S.;
- [129262]Morvay, M.; Coleman, C. M.; Rietveld, F. J. R.; Uitto, J.;
McLean,
- [129263]W. H. I.: Novel proline substitution mutations in keratin 16

in two

[129264]cases of pachyonychia congenita type 1. Brit. J. Derm.

141: 1010–1016,

[129265]1999.

[129266]23140.

[129267]Smith, F. J. D.; McKusick, V. A.; Nielsen, K.; Pfendner, E.;

Uitto,

[129268]J.; McLean, W. H. I.: Cloning of multiple keratin 16 genes

facilitates

[129269]prenatal diagnosis of pachyonychia congenita type 1. Pre-

natal Diag. 19:

[129270]941–946, 1999.

[129271]23141.

[129272]Stevens, H. P.; Kelsell, D. P.; Spurr, N. K.; Bishop, D. T.;

Purkis,

[129273]P. E.; Griffiths, W. A. D.; Rustin, M. H. A.; Leigh, I. M.: Ker-

atin

[129274]staining and linkage of non-epidermolytic focal palmo-

plantar keratodermas

[129275](PPK) to 17q. (Abstract) Brit. J. Derm. 131: 425, 1994.

[129276]23142.

- [129277]Terrinoni, A.; Puddu, P.; Didona, B.; De Laurenzi, V.;
Candi, E.;
- [129278]Smith, F. J. D.; McLean, W. H. I.; Melino, G.: A mutation in
the
- [129279]V1 domain of K16 is responsible for unilateral palmoplan-
tar verrucous
- [129280]nevus. J. Invest. Derm. 114: 1136–1140, 2000.
- [129281]23143.
- [129282]Lewis, J.; McGowan, E.; Rockwood, J.; Melrose, H.;
Nacharaju,
- [129283]P.; Van Slegtenhorst, M.; Gwinn-Hardy, K.; Murphy, M. P.;
Baker, M.;
- [129284]Yu, X.; Duff, K.; Hardy, J.; Corral, A.; Lin, W.-L.; Yen, S.-H.;
Dickson,
- [129285]D. W.; Davies, P.; Hutton, M.: Neurofibrillary tangles, amy-
otrophy
- [129286]and progressive motor disturbance in mice expressing
mutant (P301L)
- [129287]tau protein. Nature Genet. 25: 402–405, 2000. Note: Erra-
tum: Nature
- [129288]Genet. 26: 127 only, 2000.
- [129289]23144.

[129290]Lippa, C. F.; Zhukareva, V.; Kawarai, T.; Uryu, K.; Shafiq, M.;

[129291]Nee, L. E.; Grafman, J.; Liang, Y.; St George-Hyslop, P. H.; Trojanowski,

[129292]J. Q.; Lee, V. M.-Y.: Frontotemporal dementia with novel tau pathology

[129293]and a glu342val tau mutation. *Ann. Neurol.* 48: 850–858, 2000.

[129294]23145.

[129295]Litvan, I.; Baker, M.; Hutton, M.: Tau genotype: no effect on

[129296]onset, symptom severity, or survival in progressive supranuclear palsy. *Neurology* 57:

[129297]138–140, 2001.

[129298]23146.

[129299]Martin, E. R.; Scott, W. K.; Nance, M. A.; Watts, R. L.; Hubble,

[129300]J. P.; Koller, W. C.; Lyons, K.; Pahwa, R.; Stern, M. B.; Colcher,

[129301]A.; Hiner, B. C.; Jankovic, J.; and 20 others: Association of single-nucleotide

[129302]polymorphisms of the Tau gene with late-onset Parkinson

disease. J.A.M.A. 286:

[129303]2245–2250, 2001.

[129304]23147.

[129305]Moser, A. B.; Rasmussen, M.; Naidu, S.; Watkins, P. A.;
McGuinness,

[129306]M.; Hajra, A. K.; Chen, G.; Raymond, G.; Liu, A.; Gordon,
D.; Garnaas,

[129307]K.; Walton, D. S.; Skjedal, O. H.; Guggenheim, M. A.; Jack-
son, L.

[129308]G.; Elias, E. R.; Moser, H. W.: Phenotype of patients with
peroxisomal

[129309]disorders subdivided into sixteen complementation
groups. J. Pediat. 127:

[129310]13–22, 1995.

[129311]23148.

[129312]Paton, B. C.; Heron, S. E.; Nelson, P. V.; Morris, C. P.; Pou-
los,

[129313]A.: Absence of mutations raises doubts about the role of
the 70-kD

[129314]peroxisomal membrane protein in Zellweger syndrome.
(Letter) Am.

[129315]J. Hum. Genet. 60: 1535–1539, 1997.

[129316]23149.

[129317]Aljada, A.; Ghanim, H.; Friedman, J.; Garg, R.; Mohanty, P.;
Dandona,

[129318]P.: Troglitazone reduces the expression of PPAR-gamma
while stimulating

[129319]that of PPAR-alpha in mononuclear cells in obese sub-
jects. J. Clin.

[129320]Endocr. Metab. 86: 3130-3133, 2001.

[129321]23150.

[129322]Costet, P.; Legendre, C.; More, J.; Edgar, A.; Galtier, P.;
Pineau,

[129323]T.: Peroxisome proliferator-activated receptor alpha-
isoform deficiency

[129324]leads to progressive dyslipidemia with sexually dimorphic
obesity

[129325]and steatosis. J. Biol. Chem. 273: 29577-29585, 1998.

[129326]23151.

[129327]Djouadi, F.; Weinheimer, C. J.; Saffitz, J. E.; Pitchford, C.;

[129328]Bastin, J.; Gonzalez, F. J.; Kelly, D. P.: A gender-related
defect

[129329]in lipid metabolism and glucose homeostasis in peroxi-

some proliferator-activated
 [129330]receptor alpha-deficient mice. J. Clin. Invest. 102:
 1083-1091,
 [129331]1998.
 [129332]23152.
 [129333]Kersten, S.; Desvergne, B.; Wahli, W.: Roles of PPARs in
 health
 [129334]and disease. Nature 405: 421-424, 2000.
 [129335]23153.
 [129336]Kersten, S.; Seydoux, J.; Peters, J. M.; Gonzalez, F. J.;
 Desvergne,
 [129337]B.; Wahli, W.: Peroxisome proliferator-activated receptor
 alpha mediates
 [129338]the adaptive response to fasting. J. Clin. Invest. 103:
 1489-1498,
 [129339]1999.
 [129340]23154.
 [129341]Lee, S. S.-T.; Pineau, T.; Drago, J.; Lee, E. J.; Owens, J. W.;
 [129342]Kroetz, D. L.; Fernandez-Salguero, P. M.; Westphal, H.;
 Gonzalez,
 [129343]F. J.: Targeted disruption of the alpha isoform of the per-

oxisome

[129344]proliferator-activated receptor gene in mice results in
abolishment

[129345]of the pleiotropic effects of peroxisome proliferators.

Molec. Cell.

[129346]Biol. 15: 3012-3022, 1995.

[129347]23155.

[129348]Lee, Y.; Yu, X.; Gonzales, F.; Mangelsdorf, D. J.; Wang,
M.-Y.;

[129349]Richardson, C.; Witters, L. A.; Unger, R. H.: PPAR-alpha is
necessary

[129350]for the lipopenic action of hyperleptinemia on white adi-
pose and liver

[129351]tissue. Proc. Nat. Acad. Sci. 99: 11848-11853, 2002.

[129352]23156.

[129353]Leone, T. C.; Weinheimer, C. J.; Kelly, D. P.: A critical role

[129354]for the peroxisome proliferator-activated receptor alpha
(PPAR-alpha)

[129355]in the cellular fasting response: the PPAR-alpha-null
mouse as a model

[129356]of fatty acid oxidation disorders. Proc. Nat. Acad. Sci. 96:

7473–7478,

[129357]1999.

[129358]23157.

[129359]Michalik, L.; Desvergne, B.; Tan, N. S.; Basu–Modak, S.; Escher,

[129360]P.; Rieusset, J.; Peters, J. M.; Kaya, G.; Gonzalez, F. J.; Zakany,

[129361]J.; Metzger, D.; Chambon, P.; Duboule, D.; Wahli, W.: Impaired skin

[129362]wound healing in peroxisome proliferator–activated receptor (PPAR)–alpha

[129363]and PPAR–beta mutant mice. J. Cell Biol. 154: 799–814, 2001.

[129364]23158.

[129365]Sher, T.; Yi, H.–F.; McBride, O. W.; Gonzalez, F. J.: cDNA cloning,

[129366]chromosomal mapping, and functional characterization of the human

[129367]peroxisome proliferator activated receptor. Biochemistry 32: 5598–5604,

[129368]1993.

[129369]23159.

[129370] Xu, H. E.; Stanley, T. B.; Montana, V. G.; Lambert, M. H.;
Shearer,

[129371] B. G.; Cobb, J. E.; McKee, D. D.; Galardi, C. M.; Plunket, K.
D.;

[129372] Nolte, R. T.; Parks, D. J.; Moore, J. T.; Kliewer, S. A.; Will-
son,

[129373] T. M.; Stimmel, J. B.: Structural basis for antagonist-mediated recruitment

[129374] of nuclear co-repressors by PPAR-alpha. Nature 415:
813-817, 2002.

[129375] 23160.

[129376] Cameron, P.; Limjuco, G.; Rodkey, J.; Bennett, C.; Schmidt,
J.

[129377] A.: Amino acid sequence analysis of human interleukin 1
(IL-1): evidence

[129378] for biochemically distinct forms of IL-1. J. Exp. Med. 162:
790-801,

[129379] 1985.

[129380] 23161.

[129381] D'Eustachio, P.; Jadidi, S.; Fuhlbrigge, R. C.; Gray, P. W.;
Chaplin,

[129382] D. D.: Interleukin-1 alpha and beta genes: linkage on

chromosome

[129383]2 in the mouse. Immunogenetics 26: 339-343, 1987.

[129384]23162.

[129385]Dinarello, C. A.: An update on human interleukin-1: from molecular

[129386]biology to clinical relevance. J. Clin. Immun. 5: 287-297, 1985.

[129387]23163.

[129388]El-Omar, E. M.; Carrington, M.; Chow, W.-H.; McColl, K. E. L.;

[129389]Bream, J. H.; Young, H. A.; Herrera, J.; Lissowska, J.; Yuan, C.-C.;

[129390]Rothman, N.; Lanyon, G.; Martin, M.; Fraumeni, J. F., Jr.; Rabkin,

[129391]C. S.: Interleukin-1 polymorphisms associated with increased risk

[129392]of gastric cancer. Nature 404: 398-402, 2000. Note: Erratum: Nature

[129393]412: 99 only, 2001.

[129394]23164.

[129395]Furutani, Y.; Notake, M.; Yamayoshi, M.; Yamagishi, J.;

Nomura,

[129396]H.; Ohue, M.; Furuta, R.; Fukui, T.; Yamada, M.; Nakamura,
S.: Cloning

[129397]and characterization of the cDNAs for human and rabbit
interleukin-1

[129398]precursor. Nucleic Acids Res. 13: 5869-5882, 1985.

[129399]23165.

[129400]Hamajima, N.; Matsuo, K.; Saito, T.; Tajima, K.; Okuma, K.;
Yamao,

[129401]K.; Tominaga, S.: Interleukin 1 polymorphisms, lifestyle
factors,

[129402]and helicobacter pylori infection. Jpn. J. Cancer Res. 92:
383-389,

[129403]2001.

[129404]23166.

[129405]Le Beau, M. M.; Rowley, J. D.: Personal Communication.
Chicago,

[129406]Ill. 6/18/1986.

[129407]23167.

[129408]March, C. J.; Mosley, B.; Larsen, A.; Cerretti, D. P.; Braedt,

[129409]G.; Price, V.; Gillis, S.; Henney, C. S.; Kronheim, S. R.;

Grabstein,

[129410]K.; Conlon, P. J.; Hopp, T. P.; Cosman, D.: Cloning, sequence and

[129411]expression of two distinct human interleukin-1 complementary DNAs. *Nature* 315:

[129412]641-647, 1985.

[129413]23168.

[129414]Nicklin, M. J. H.; Weith, A.; Duff, G. W.: A physical map of

[129415]the region encompassing the human interleukin-1-alpha, interleukin-1-beta,

[129416]and interleukin-1 receptor antagonist genes. *Genomics* 19: 382-384,

[129417]1994.

[129418]23169.

[129419]Samad, T. A.; Moore, K. A.; Sapirstein, A.; Billet, S.; Allchorne,

[129420]A.; Poole, S.; Bonventre, J. V.; Woolf, C. J.: Interleukin-1-beta-mediated

[129421]induction of Cox-2 in the CNS contributes to inflammatory pain hypersensitivity. *Nature* 410:

[129422]471-475, 2001.

[129423]23170.

[129424]Vidal-Vanaclocha, F.; Fantuzzi, G.; Mendoza, L.; Fuentes,
A. M.;

[129425]Anasagasti, M. J.; Martin, J.; Carrascal, T.; Walsh, P.;
Reznikov,

[129426]L. L.; Kim, S.-H.; Novick, D.; Rubinstein, M.; Dinarello, C.
A.:

[129427]IL-18 regulates IL-1-beta-dependent hepatic melanoma
metastasis via

[129428]vascular cell adhesion molecule-1. Proc. Nat. Acad. Sci.
97: 734-739,

[129429]2000.

[129430]23171.

[129431]Webb, A.; Collins, K.; Auron, P.; Eddy, R.; Nakai, H.; Byers,

[129432]M.; Shows, T. B.: Genetics of acute phase response: a gene
for interleukin-1

[129433]is on chromosome 2. (Abstract) Am. J. Hum. Genet. 37:
A142 only,

[129434]1985.

[129435]23172.

[129436]Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai,

[129437]H.; Byers, M.; Shows, T. B.: The gene for interleukin-1 (IL1)
is

[129438]on human chromosome 2. (Abstract) Cytogenet. Cell
Genet. 40: 774

[129439]only, 1985.

[129440]23173.

[129441]Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai,

[129442]H.; Byers, M. G.; Haley, L. L.; Henry, W. M.; Shows, T. B.:
Interleukin-1

[129443]gene (IL1) assigned to long arm of human chromosome 2.

Lymphokine

[129444]Res. 5: 77–85, 1986.

[129445]23174.

[129446]Ansano, M.; Toda, M.; Sakaguchi, N.; Sakaguchi, S.: Autoimmune

[129447]disease as a consequence of developmental abnormality of a T cell

[129448]subpopulation. J. Exp. Med. 184: 387–396, 1996.

[129449]23175.

[129450]Cosman, D.; Cerretti, D. P.; Larsen, A.; Park, L.; March, C.; Dower,

[129451]S.; Gillis, S.; Urdal, D.: Cloning, sequence and expression of human

[129452]interleukin-2 receptor. Nature 312: 768–771, 1984.

[129453]23176.

[129454]Ferrari, S.; Cannizzaro, L. A.; Battini, R.; Huebner, K.; Baserga,

[129455]R.: The gene encoding human vimentin is located on the short arm

[129456]of chromosome 10. Am. J. Hum. Genet. 41: 616–626, 1987.

[129457]23177.

[129458]Greene, W. C.; Leonard, W. J.; Depper, J. M.; Nelson, D. L.;
Waldmann,

[129459]T. A.: The human interleukin-2 receptor: normal and ab-
normal expression

[129460]in T cells and in leukemias induced by the human T-
lymphotropic retroviruses. Ann.

[129461]Intern. Med. 105: 560-572, 1986.

[129462]23178.

[129463]Hatakeyama, M.; Minamoto, S.; Taniguchi, T.: Intracyto-
plasmic

[129464]phosphorylation sites of Tac antigen (p55) are not essen-
tial for the

[129465]conformation, function, and regulation of the human in-
terleukin 2

[129466]receptor. Proc. Nat. Acad. Sci. 83: 9650-9654, 1986.

[129467]23179.

[129468]Hatakeyama, M.; Minamoto, S.; Uchiyama, T.; Hardy, R. R.;
Yamada,

[129469]G.; Taniguchi, T.: Reconstitution of functional receptor for
human

[129470]interleukin-2 in mouse cells. Nature 318: 467-470, 1985.

[129471]23180.

[129472]Ihle, J. N.; Kerr, I. M.: Jaks and Stats in signaling by the cytokine

[129473]receptor superfamily. Trends Genet. 11: 69-74, 1995.

[129474]23181.

[129475]Ishida, N.; Kanamori, H.; Noma, T.; Nikaido, T.; Sabe, H.; Suzuki,

[129476]N.; Shimizu, A.; Honjo, T.: Molecular cloning and structure of the

[129477]human interleukin 2 receptor gene. Nucleic Acids Res. 13: 7579-7589,

[129478]1985.

[129479]23182.

[129480]Kondo, S.; Shimizu, A.; Maeda, M.; Tagaya, Y.; Yodoi, J.; Honjo,

[129481]T.: Expression of functional human interleukin-2 receptor in mouse

[129482]T cells by cDNA transfection. Nature 320: 75-77, 1986.

[129483]23183.

[129484]Leonard, W. J.; Depper, J. M.; Crabtree, G. R.; Rudikoff, S.;

[129485]Pumphrey, J.; Robb, R. J.; Kronke, M.; Svetlik, P. B.; Pepper,
N.

[129486]J.; Waldmann, T. A.; Greene, W. C.: Molecular cloning and
expression

[129487]of cDNAs for the human interleukin-2 receptor. Nature
311: 626-631,

[129488]1984.

[129489]23184.

[129490]Lekanne Deprez, R. H.; Riegman, P. H. J.; Groen, N. A.;
Warringa,

[129491]U. L.; van Biezen, N. A.; Molijn, A. C.; Bootsma, D.; de
Jong, P.

[129492]J.; Menon, A. G.; Kley, N. A.; Seizinger, B. R.; Zwarthoff, E.
C.

[129493]: Cloning and characterization of MN1, a gene from chro-
mosome 22q11,

[129494]which is disrupted by a balanced translocation in a
meningioma. Oncogene 10:

[129495]1521-1528, 1995.

[129496]23185.

[129497]Dahia, P. L. M.; Aguiar, R. C. T.; Alberta, J.; Kum, J. B.;
 Caron,
 [129498]S.; Sill, H.; Marsh, D. J.; Ritz, J.; Freedman, A.; Stiles, C.;
 Eng,
 [129499]C.: PTEN is inversely correlated with the cell survival factor
 Akt/PKB
 [129500]and is inactivated via multiple mechanisms in haemato-
 logical malignancies. Hum.
 [129501]Molec. Genet. 8: 185-193, 1999.
 [129502]23186.
 [129503]Bryson, G. L. M.; Massa, H.; Trask, B. J.; Van Etten, R. L.:
 Gene
 [129504]structure, sequence, and chromosomal localization of the
 human red
 [129505]cell-type low molecular weight acid phosphotyrosyl phos-
 phatase gene,
 [129506]ACP1. Genomics 30: 133-140, 1995.
 [129507]23187.
 [129508]Chen, T.-R.; McMorris, F. A.; Creagan, R.; Ricciuti, F. C.;
 Tischfield,
 [129509]J.; Ruddle, F. H.: Assignment of the genes for malate oxi-
 doreductase

[129510]decarboxylating to chromosome 6 and peptidase B and lactate dehydrogenase

[129511]B to chromosome 12 in man. Am. J. Hum. Genet. 25: 200–207, 1973.

[129512]23188.

[129513]Hamerton, J. L.; Mohandas, T.; McAlpine, P. J.; Douglas, G. R.

[129514]; Localization of human gene loci using spontaneous chromosome rearrangements

[129515]in human–Chinese hamster somatic cell hybrids. Am. J. Hum. Genet. 27:

[129516]595–608, 1975.

[129517]23189.

[129518]Herbschleb–Voogt, E.; Meera Khan, P.: Defining the locus of origin

[129519]of a genetically determined electrophoretic variant of a multilocus

[129520]enzyme system; the Calcutta–1 of human LDH system is a B–locus variant. Hum.

[129521]Genet. 57: 290–295, 1981.

[129522]23190.

- [129523] Houki, N.; Matsushima, Y.; Kitamura, M.; Tukada, T.;
Nishina, T.;
- [129524] Nakayama, T.: A case of deficiency of lactate dehydroge-
nase H-subunit. Jpn.
- [129525] J. Clin. Chem. 15: 85-90, 1986.
- [129526] 23191.
- [129527] Kitamura, M.; Iijima, N.; Hashimoto, F.; Hiratsuka, A.:
Hereditary
- [129528] deficiency of subunit H of lactate dehydrogenase. Clin.
Chim. Acta 34:
- [129529] 419-423, 1971.
- [129530] 23192.
- [129531] Maekawa, M.; Sudo, K.; Kitajima, M.; Matsuura, Y.; Li, S.
S.-L.;
- [129532] Kanno, T.: Analysis of a genetic mutation in an elec-
trophoretic variant
- [129533] of the human lactate dehydrogenase-B(H) subunit. Hum.
Genet. 91:
- [129534] 423-426, 1993.
- [129535] 23193.
- [129536] Maekawa, M.; Sudo, K.; Nagura, K.; Li, S. S.-L.; Kanno, T.:

Population

- [129537]screening of lactate dehydrogenase deficiencies in
Fukuoka Prefecture
- [129538]in Japan and molecular characterization of three independent mutations
- [129539]in the lactate dehydrogenase-B(H) gene. Hum. Genet. 93:
74-76, 1994.
- [129540]23194.
- [129541]Malpuech, G.; Kaplan, J. C.; Rethore, M. O.; Junien, C.;
Geneix,
- [129542]A.: Une observation de deletion partielle du bras court du
chromosome
- [129543]12: localisation du gene de la lacticodeshydrogenase B.
Lyon Med. 233:
- [129544]275-279, 1975.
- [129545]23195.
- [129546]Mayeda, K.; Weiss, L.; Lindahl, R.; Dully, M.: Localization of
- [129547]the human lactate dehydrogenase B gene on the short arm
of chromosome
- [129548]12. Am. J. Hum. Genet. 26: 59-64, 1974.
- [129549]23196.

- [129550]Miwa, S.; Nishima, T.; Kanehashi, Y.; Kitamura, M.; Hiratsuka,
A.; Shizume, K.: Studies on erythrocyte metabolism in a
case with
hereditary deficiency of H-subunit of lactate dehydroge-
nase. Acta
Haemat. Jpn. 34: 228-232, 1971.
23197.
- [129555]Bell, D. W.; Varley, J. M.; Szydlo, T. E.; Kang, D. H.; Wahrer,
D. C. R.; Shannon, K. E.; Lubratovich, M.; Verselis, S. J.; Is-
selbacher,
K. J.; Fraumeni, J. F.; Birch, J. M.; Li, F. P.; Garber, J. E.;
Haber,
D. A.: Heterozygous germ line hCHK2 mutations in Li-
Fraumeni syndrome. Science 286:
2528-2531, 1999.
23198.
- [129561]Birch, J. M.; Hartley, A. L.; Tricker, K. J.; Prosser, J.; Condie,
A.; Kelsey, A. M.; Harris, M.; Jones, P. H. M.; Binchy, A.;
Crowther,
D.; Craft, A. W.; Eden, O. B.; Evans, D. G. R.; Thompson, E.;

Mann,

[129564]J. R.; Martin, J.; Mitchell, E. L. D.; Santibanez-Koref, M. F.:

Prevalence

[129565]and diversity of constitutional mutations in the p53 gene
among 21

[129566]Li-Fraumeni families. Cancer Res. 54: 1298-1304, 1994.

[129567]23199.

[129568]Goy, A.; Gilles, F.; Remache, Y.; Zelenetz, A. D.: Physical
linkage

[129569]of the lysyl oxidase-like (LOXL1) gene to the PML gene on
human chromosome

[129570]15q22. Cytogenet. Cell Genet. 88: 22-24, 2000.

[129571]23200.

[129572]Kenyon, K.; Modi, W. S.; Contente, S.; Friedman, R. M.: A
novel

[129573]human cDNA with a predicted protein similar to lysyl ox-
idase maps

[129574]to chromosome 15q24-q25. J. Biol. Chem. 268:
18435-18437, 1993.

[129575]23201.

[129576]Kim, Y.; Boyd, C. D.; Csiszar, K.: A new gene with se-

quence and

[129577] structural similarity to the gene encoding human lysyl oxidase. J.

[129578] Biol. Chem. 270: 7176–7182, 1995.

[129579] 23202.

[129580] Wydner, K. S.; Kim, Y.; Csiszar, K.; Boyd, C. D.; Passmore, H.

[129581] C.: An intron capture strategy used to identify and map a lysyl oxidase-like

[129582] gene on chromosome 9 in the mouse. Genomics 40: 342–345, 1997.

[129583] 23203.

[129584] Schmidt, B.; Servidei, S.; Gabbai, A. A.; Silva, A. C.; de Sousa

[129585] Bulle de Oliveira, A.; DiMauro, S.: McArdle's disease in two generations:

[129586] autosomal recessive transmission with manifesting heterozygote. Neurology 37:

[129587] 1558–1561, 1987.

[129588] 23204.

[129589] Arch, E. M.; Goodman, B. K.; Van Wesep, R. A.; Liaw, D.;

Clarke,

[129590]K.; Parsons, R.; McKusick, V. A.; Geraghty, M. T.: Deletion
of PTEN

[129591]in a patient with Bannayan–Riley–Ruvalcaba syndrome
suggests allelism

[129592]with Cowden disease. Am. J. Med. Genet. 71: 489–493,
1997.

[129593]23205.

[129594]Celebi, J. T.; Tsou, H. C.; Chen, F. F.; Zhang, H.; Ping, X. L.;

[129595]Lebwohl, M. G.; Kezis, J.; Peacocke, M.: Phenotypic find-
ings of Cowden

[129596]syndrome and Bannayan–Zonana syndrome in a family as-
sociated with

[129597]a single germline mutation in PTEN. J. Med. Genet. 36:
360–364,

[129598]1999.

[129599]23206.

[129600]Dudek, H.; Datta, S. R.; Franke, T. F.; Birnbaum, M. J.; Yao,
R.;

[129601]Cooper, G. M.; Segal, R. A.; Kaplan, D. R.; Greenberg, M.
E.: Regulation

[129602]of neuronal survival by the serine–threonine protein ki-

nase Akt. Science 275:

[129603]661–663, 1997.

[129604]23207.

[129605]Franke, T. F.; Kaplan, D. R.; Cantley, L. C.; Toker, A.: Direct

[129606]regulation of the Akt proto-oncogene product by phosphatidylinositol-3,4-bisphosphate. Science 275:

[129607]665–667, 1997.

[129608]23208.

[129609]Franke, T. F.; Yang, S.-I.; Chan, T. O.; Datta, K.; Kazanietz, A.; Morrison, D. K.; Kaplan, D. R.; Tsichlis, P. N.: The protein

[129610]kinase encoded by the Akt proto-oncogene is a target of the PDGF-activated

[129611]phosphatidylinositol 3-kinase. Cell 81: 727–736, 1995.

[129612]23209.

[129613]Furnari, F. B.; Huang, H. J.; Cavenee, W. K.: The phospho-

[129614]inositol phosphatase activity of PTEN mediates a serum-sensitive G1 growth

[129615]arrest in glioma cells. Cancer Res. 58: 5002–5008, 1998.

[129616]

[129617]23210.

[129618]Hajdуч, E.; Litherland, G. J.; Hundal, H. S.: Protein kinase

[129619]B (PKB/Akt)--a key regulator of glucose transport? FEBS
Lett. 492:

[129620]199-203, 2001.

[129621]23211.

[129622]Hemmings, B. A.: Akt signaling: linking membrane events
to life

[129623]and death decisions. Science 275: 628-630, 1997.

[129624]23212.

[129625]Jones, P. F.; Jakubowicz, T.; Pitossi, F. J.; Maurer, F.; Hem-
mings,

[129626]B. A.: Molecular cloning and identification of a serine/
threonine

[129627]protein kinase of the second-messenger subfamily. Proc.
Nat. Acad.

[129628]Sci. 88: 4171-4175, 1991.

[129629]23213.

[129630]Lee, M.-J.; Thangada, S.; Paik, J.-H.; Sapkota, G. P.; An-
cellin,

- [129631]N.; Chae, S.-S.; Wu, M.; Morales-Ruiz, M.; Sessa, W. C.; Alessi, D.
- [129632]R.; Hla, T.: Akt-mediated phosphorylation of the G protein-coupled
- [129633]receptor EDG-1 is required for endothelial cell chemotaxis. *Molec.*
- [129634]*Cell* 8: 693-704, 2001.
- [129635]23214.
- [129636]Maira, S.-M.; Galetic, I.; Brazil, D. P.; Kaeck, S.; Ingley, E.;
- [129637]Thelen, M.; Hemmings, B. A.: Carboxyl-terminal modulator protein
- [129638](CTMP), a negative regulator of PKB/Akt and v-Akt at the plasma membrane. *Scien*
- [129639]*ce* 294: 374-380, 2001.
- [129640]23215.
- [129641]Powell, D. W.; Rane, M. J.; Chen, Q.; Singh, S.; McLeish, K. R.
- [129642]: Identification of 14-3-3-zeta as a protein kinase B/Akt substrate. *J.*
- [129643]*Biol. Chem.* 277: 21639-21642, 2002.
- [129644]23216.

[129645]Staal, S. P.: Molecular cloning of the akt oncogene and its human

[129646]homologues AKT1 and AKT2: amplification of AKT1 in a primary human

[129647]gastric adenocarcinoma. Proc. Nat. Acad. Sci. 84: 5034–5037, 1987.

[129648]23217.

[129649]Staal, S. P.; Huebner, K.; Croce, C. M.; Parsa, N. Z.; Testa,

[129650]J. R.: The AKT1 proto-oncogene maps to human chromosome 14, band

[129651]q32. Genomics 2: 96–98, 1988.

[129652]23218.

[129653]Vanhaesebroeck, B.; Alessi, D. R.: The PI3K–PDK1 connection:

[129654]more than just a road to PKB. Biochem. J. 346: 561–576, 2000.

[129655]23219.

[129656]Weng, L.–P.; Brown, J. L.; Eng, C.: PTEN induces apoptosis and

[129657]cell cycle arrest through phosphoinositol–3–kinase/Akt–dependent and

[129658]—independent pathways. Hum. Molec. Genet. 10:
237–242, 2001.

[129659]23220.

[129660]Yang, J.; Cron, P.; Thompson, V.; Good, V. M.; Hess, D.;
Hemmings,

[129661]B. A.; Barford, D.: Molecular mechanism for the regulation
of protein

[129662]kinase B/Akt by hydrophobic motif phosphorylation.
Molec. Cell 9:

[129663]1227–1240, 2002.

[129664]23221.

[129665]Le Beau, M. M.; Rowley, J. D.; Sacchi, N.; Watson, D. K.; Pa-
pas,

[129666]T. S.; Diaz, M. O.: Hu-ets-2 is translocated to chromo-
some 8 in the

[129667]t(8;21) in acute myelogenous leukemia. Cancer Genet. Cy-
togenet. 23:

[129668]269–274, 1986.

[129669]23222.

[129670]Mavrothalassitis, G. J.; Watson, D. K.; Papas, T. S.: Molecu-
lar

- [129671]and functional characterization of the promoter of ETS2,
the human
- [129672]c-ets-2 gene. Proc. Nat. Acad. Sci. 87: 1047–1051, 1990.
- [129673]23223.
- [129674]Sacchi, N.; Cheng, S. V.; Tanzi, R. E.; Gusella, J. F.;
Drabkin,
- [129675]H. A.; Patterson, D.; Haines, J. H.; Papas, T. S.: The ETS
genes
- [129676]on chromosome 21 are distal to the breakpoint of the
acute myelogenous
- [129677]leukemia translocation (8;21). Genomics 3: 110–116,
1988.
- [129678]23224.
- [129679]Sumarsono, S. H.; Wilson, T. J.; Tymms, M. J.; Venter, D. J.;
Corrick,
- [129680]C. M.; Kola, R.; Lahoud, M. H.; Papas, T. S.; Seth, A.; Kola,
I.:
- [129681]Down's syndrome-like skeletal abnormalities in Ets2
transgenic mice. Nature 379:
- [129682]534–540, 1996.
- [129683]23225.

[129684]Chu, E. H. Y.; Chang, C. C.; Sun, N. C.: Synteny of the human

[129685]genes for gal-1-PT, ACP-1, MDH-1, and gal-plus-activator and assignment

[129686]to chromosome 2. Birth Defects Orig. Art. Ser. XI(3): 103-106, 1975.

[129687]Note: Alternate: Cytogenet. Cell Genet. 14: 273-276, 1975...

[129688]23226.

[129689]Rasmussen, S. K.; Lautier, C.; Hansen, L.; Echwald, S. M.; Hansen,

[129690]T.; Ekstrom, C. T.; Urhammer, S. A.; Borch-Johnsen, K.; Grigorescu,

[129691]F.; Smith, R. J.; Pedersen, O.: Studies of the variability of the

[129692]genes encoding the insulin-like growth factor I receptor and its ligand

[129693]in relation to type 2 diabetes mellitus. J. Clin. Endocr. Metab. 85:

[129694]1606-1610, 2000.

[129695]23227.

[129696]Dissing, J.; Johnsen, A. H.: Human red cell acid phos-

phatase (ACP1):

[129697]the primary structure of the two pairs of isozymes encoded by the

[129698]ACP1*A and ACP1*C alleles. Biochim. Biophys. Acta 1121: 261-268,

[129699]1992.

[129700]23228.

[129701]Dissing, J.; Johnsen, A. H.; Sensabaugh, G. F.: Human red cell

[129702]acid phosphatase (ACP1): the amino acid sequence of the two isozymes

[129703]Bf and Bs encoded by the ACP1*B allele. J. Biol. Chem. 266: 20619-20625,

[129704]1991.

[129705]23229.

[129706]Emanuel, B. S.; Zackai, E. H.; Van Dyke, D. C.; Swallow, D. M.;

[129707]Allen, F. H.; Mellman, W. J.: Deletion mapping: further evidence

[129708]for the location of acid phosphatase (ACP-1) within 2p23. Am. J.

[129709]Med. Genet. 4: 167-172, 1979.

[129710]23230.

[129711]Ferguson-Smith, M. A.; Newman, B. F.; Ellis, P. M.; Thomson, D.

[129712]M. G.; Riley, I. D.: Assignment by deletion of human red cell acid

[129713]phosphatase gene locus to the short arm of chromosome 2. Nature 243:

[129714]271-273, 1973.

[129715]23231.

[129716]Fisher, R. A.; Harris, H.: Studies on the separate isoenzymes

[129717]of red cell acid phosphatase phenotypes A and B: chromatographic separation

[129718]of the isoenzymes. Ann. Hum. Genet. 34: 431-438, 1971.

[129719]23232.

[129720]Fuhrmann, W.; Lichte, K. H.: Human red cell acid phosphatase

[129721]polymorphism: a study on gene frequency and forensic use of the system

[129722]in cases of disputed paternity. Humangenetik 3: 121-126, 1966.

[129723]23233.

[129724]Giblett, E. R.; Scott, N. M.: Red cell acid phosphatase:
racial

[129725]distribution and report of a new phenotype. Am. J. Hum.
Genet. 17:

[129726]425-432, 1965.

[129727]23234.

[129728]Herbich, J.; Fisher, R. A.; Hopkinson, D. A.: Atypical segre-
gation

[129729]of human red cell acid phosphatase phenotypes: evidence
for a rare

[129730]'silent' allele P(O). Ann. Hum. Genet. 34: 145-152, 1970.

[129731]23235.

[129732]Herbich, J.; Meinhardt, K.: The rare 'silent' allele P(O) or P(V)

[129733](P Vienna) of human red cell acid phosphatase, typed in a
second family. Humangenetik 15:

[129734]345-348, 1972.

[129735]23236.

[129736]Hopkinson, D. A.; Spencer, N.; Harris, H.: Red cell acid
phosphatase

[129737]variants: a new human polymorphism. Nature 199:
969-971, 1963.

[129738]23237.

[129739]Hulten, M.; Lindsten, J.; Pen-Ming, L. M.; Fraccaro, M.;
Mannini,

[129740]A.; Tiepolo, L.; Robson, E. B.; Heiken, A.; Tillinger, K. G.:
Possible

[129741]localization of the genes for the Kidd blood group on an
autosome

[129742]involved in a reciprocal translocation. Nature 211:
1067-1068, 1966.

[129743]23238.

[129744]Junien, C.; Kaplan, J.-C.; Bernheim, A.; Berger, R.: Regional

[129745]assignment of red cell acid phosphatase locus to band
2p25. Hum.

[129746]Genet. 48: 17-21, 1979.

[129747]23239.

[129748]Karp, G. W., Jr.; Sutton, H. E.: Some new phenotypes of
human

[129749]red cell acid phosphatase. Am. J. Hum. Genet. 19: 54-62,
1967.

[129750]23240.

[129751]Lothe, R. A.; Gedde-Dahl, T.; Olaisen, B.; Bakker, E.; Pearson,
son,

[129752]P.: Very close linkage between D2S1 and ACP1 on chromosome 2p. Ann.

[129753]Hum. Genet. 50: 361-367, 1986.

[129754]23241.

[129755]Mace, M. A.; Cook, P. J. L.; Robson, E. B.: Linkage data on red

[129756]cell acid phosphatase from family studies. Ann. Hum. Genet. 38:

[129757]471-477, 1975.

[129758]23242.

[129759]Mace, M. A.; Robson, E. B.: Linkage data on ACP-1 and MNSs. Cytogenet.

[129760]Cell Genet. 13: 123-125, 1974.

[129761]23243.

[129762]Magenis, R. E.; Koler, R. D.; Lovrien, E. W.; Bigley, R. H.; Duval,

[129763]M. C.; Overton, K. M.: Gene dosage: evidence for assignment of erythrocyte

[129764]acid phosphatase locus to chromosome 2. Proc. Nat.
Acad. Sci. 72:

[129765]4526–4530, 1975.

[129766]23244.

[129767]Mayr, W. R.: No close linkage between MNSs and red cell
acid

[129768]phosphatase. Hum. Hered. 26: 1–3, 1976.

[129769]23245.

[129770]Miller, S. A.; Nelson, M. S.; Dykes, D. D.; Polesky, H. F.:
Comparison

[129771]of acid phosphatase ACP1 variants by isoelectric focusing
and conventional

[129772]electrophoresis: identification of three new alleles,
ACP1*N, ACP1*P

[129773]and ACP1*S. Hum. Hered. 37: 371–375, 1987.

[129774]23246.

[129775]Mohrenweiser, H. W.; Novotny, J. E.: ACP-1-GUA-1: a low-
activity

[129776]variant of human erythrocyte acid phosphatase: associa-
tion with increased

[129777]glutathione reductase activity. Am. J. Hum. Genet. 34:

425-433,

[129778]1982.

[129779]23247.

[129780]Nezbeda, P.: Occurrence of the ACP-1 null allele in
Czechoslovakia. Hum.

[129781]Genet. 46: 227-229, 1979.

[129782]23248.

[129783]Palmarino, R.; Agostino, R.; Gloria, F.; Lucarelli, P.; Bus-
inco,

[129784]L.; Antognoni, G.; Maggioni, G.; Workman, P. L.; Bottini, E.:
Red

[129785]cell acid phosphatase: another polymorphism correlated
with malaria? Am.

[129786]J. Phys. Anthropol. 43: 177-186, 1975.

[129787]23249.

[129788]Povey, S.; Swallow, D. M.; Bobrow, M.; Craig, I.; Van
Heyningen,

[129789]V.: Probable assignment of the locus determining human
red cell acid

[129790]phosphatase ACP(1) to chromosome 2 using somatic cell
hybrids. Ann.

[129791]Hum. Genet. 38: 1-5, 1974.

[129792]23250.

[129793]Renwick, J. H.: Assignment and map-positioning of human
loci

[129794]using chromosomal variation. Ann. Hum. Genet. 35:
79-97, 1971.

[129795]23251.

[129796]Sensabaugh, G. F.; Golden, V. L.: Phenotype dependence
in the

[129797]inhibition of red cell acid phosphatase (ACP) by folates.
Am. J.

[129798]Hum. Genet. 30: 553-560, 1978.

[129799]23252.

[129800]Swallow, D. M.; Povey, S.; Harris, H.: Activity of the 'red
cell'

[129801]acid phosphatase locus in other tissues. Ann. Hum. Genet.
37: 31-38,

[129802]1973.

[129803]23253.

[129804]Wu, R. M.; Cheng, C. W.; Chen, K. H.; Lu, S. L.; Shan, D. E.;

- [129805]Ho, Y. F.; Chern, H. D.: The COMT L allele modifies the association
[129806]between MAOB polymorphism and PD in Taiwanese. Neurology 56: 375–382,
[129807]2001.
[129808]23254.
- [129809]Leonard, W. J.; Depper, J. M.; Kanehisa, M.; Kronke, M.;
Peffer,
[129810]N. J.; Svetlik, P. B.; Sullivan, M.; Greene, W. C.: Structure of
[129811]the human interleukin–2 receptor gene. Science 230:
633–639, 1985.
[129812]23255.
- [129813]Leonard, W. J.; Depper, J. M.; Robb, R. J.; Waldmann, T. A.;
Greene,
[129814]W. C.: Characterization of the human receptor for T-cell
growth factor. Proc.
[129815]Nat. Acad. Sci. 80: 6957–6961, 1983.
[129816]23256.
- [129817]Marx, J. L.: The interleukin–2 receptor gene is cloned. Science 226:
[129818]1064–1065, 1985.

[129819]23257.

[129820]Nikaido, T.; Shimizu, A.; Ishida, N.; Sabe, H.; Teshigawara, K.;

[129821]Maeda, M.; Uchiyama, T.; Yodoi, J.; Honjo, T.: Molecular cloning

[129822]of a cDNA encoding human interleukin-2 receptor. Nature 311: 631-635,

[129823]1984.

[129824]23258.

[129825]Robb, R. J.; Rusk, C. M.; Neeper, M. P.: Structure-function relationships

[129826]for the interleukin 2 receptor: location of ligand and anti-body binding

[129827]sites on the Tac receptor chain by mutational analysis. Proc. Nat.

[129828]Acad. Sci. 85: 5654-5658, 1988.

[129829]23259.

[129830]Sharfe, N.; Dadi, H. K.; Shahar, M.; Roifman, C. M.: Human immune

[129831]disorder arising from mutation of the alpha chain of the interleukin-2

[129832]receptor. Proc. Nat. Acad. Sci. 94: 3168–3171, 1997.

[129833]23260.

[129834]Shevach, E. M.: Certified professionals: CD4(+)CD25(+) suppressor

[129835]T cells. J. Exp. Med. 193: F41–F45, 2001.

[129836]23261.

[129837]Tsudo, M.; Kozak, R. W.; Goldman, C. K.; Waldmann, T. A.: Demonstration

[129838]of a non-Tac peptide that binds interleukin 2: a potential participant

[129839]in a multichain interleukin 2 receptor complex. Proc. Nat. Acad.

[129840]Sci. 83: 9694–9698, 1986.

[129841]23262.

[129842]Urdal, D. L.; March, C. J.; Gillis, S.; Larsen, A.; Dower, S.

[129843]K.: Purification and chemical characterization of the receptor for

[129844]interleukin 2 from activated human T lymphocytes and from a human

[129845]T-cell lymphoma cell line. Proc. Nat. Acad. Sci. 81: 6481–6485,

[129846]1984.

[129847]23263.

[129848]Mohrenweiser, H. W.; Neel, J. V.: Frequency of thermostability

[129849]variants: estimation of total 'rare' variant frequency in human populations. Proc.

[129850]Nat. Acad. Sci. 78: 5729-5733, 1981.

[129851]23264.

[129852]Farrer, L. A.; Conneally, P. M.: A genetic model for age at onset

[129853]in Huntington disease. Am. J. Hum. Genet. 37: 350-357, 1985.

[129854]23265.

[129855]Farrer, L. A.; Conneally, P. M.; Yu, P.: The natural history

[129856]of Huntington disease: possible role of 'aging genes.'. Am. J. Med.

[129857]Genet. 18: 115-123, 1984.

[129858]23266.

[129859]Farrer, L. A.; Cupples, L. A.; Wiater, P.; Conneally, P. M.; Gusella,

[129860]J. F.; Myers, R. H.: The normal Huntington disease (HD) allele, or
[129861]a closely linked gene, influences age at onset of HD. Am.
J. Hum.
[129862]Genet. 53: 125–130, 1993.
[129863]23267.
[129864]Ferrante, R. J.; Kowall, N. W.; Beal, M. F.; Richardson, E. P.,
[129865]Jr.; Bird, E. D.; Martin, J. B.: Selective sparing of a class of
[129866]striatal neurons in Huntington's disease. Science 230:
561–563,
[129867]1985.
[129868]23268.
[129869]Folstein, S.; Abbott, M.; Moser, R.; Parhad, I.; Clark, A.;
Folstein,
[129870]M.: A phenocopy of Huntington's disease: lacunar infarcts
of the
[129871]corpus striatum. Johns Hopkins Med. J. 148: 104–108,
1981.
[129872]23269.
[129873]Folstein, S. E.; Abbott, M. H.; Franz, M. L.; Huang, S.;
Chase,

- [129874]G. A.; Folstein, M. F.: Phenotypic heterogeneity in Huntington disease. J. Neurogenet. 1: 175–184, 1984.
- [129875]23270.
- [129876]Folstein, S. E.; Chase, G. A.; Wahl, W. E.; McDonnell, A. M.;
- [129877]Folstein, M. F.: Huntington disease in Maryland: clinical aspects of racial variation. Am. J. Hum. Genet. 41: 168–179, 1987.
- [129878]23271.
- [129879]Folstein, S. E.; Phillips, J. A., III; Meyers, D. A.; Chase, G. A.; Abbott, M. H.; Franz, M. L.; Waber, P. G.; Kazazian, H. H., Jr.;
- [129880]Conneally, P. M.; Hobbs, W.; Tanzi, R.; Faryniarz, A.; Gibbons, K.;
- [129881]Gusella, J.: Huntington's disease: two families with differing clinical features show linkage to the G8 probe. Science 229: 776–779, 1985.
- [129882]23272.
- [129883]Freeman, T. B.; Cicchetti, F.; Hauser, R. A.; Deacon, T. W.; Li,

[129888]X.-J.; Hersch, S. M.; Nauert, G. M.; Sanberg, P. R.; Kordower, J.

[129889]H.; Saporta, S.; Isacson, O.: Transplanted fetal striatum in Huntington's

[129890]disease: phenotypic development and lack of pathology. Proc. Nat.

[129891]Acad. Sci. 97: 13877-13882, 2000.

[129892]23273.

[129893]Froster-Iskenius, U. G.; Hayden, M. R.; Wang, H. S.; Kalousek,

[129894]D. K.; Horsman, D.; Pfeiffer, R. A.; Schottky, A.; Schwinger, E.:

[129895]A family with Huntington disease and reciprocal translocation 4;5. Am.

[129896]J. Hum. Genet. 38: 759-767, 1986.

[129897]23274.

[129898]Gellera, C.; Meoni, C.; Castellotti, B.; Zappacosta, B.; Girotti,

[129899]F.; Taroni, F.; DiDonato, S.: Errors in Huntington disease diagnostic

[129900]test caused by trinucleotide deletion in the IT15 gene. (Letter) Am.

[129901]J. Hum. Genet. 59: 475–477, 1996.

[129902]23275.

[129903]Gervais, F. G.; Singaraja, R.; Xanthoudakis, S.; Gutekunst, C.-A.;

[129904]Leavitt, B. R.; Metzler, M.; Hackam, A. S.; Tam, J.; Vaillancourt,

[129905]J. P.; Houtzager, V.; Rasper, D. M.; Roy, S.; Hayden, M. R.; Nicholson,

[129906]D. W.: Recruitment and activation of caspase-8 by the huntingtin-interacting

[129907]protein Hip-1 and a novel partner Hippi. Nature Cell Biol. 4: 95–105,

[129908]2002.

[129909]23276.

[129910]Gilliam, T. C.; Bucan, M.; MacDonald, M. E.; Zimmer, M.; Haines,

[129911]J. L.; Cheng, S. V.; Pohl, T. M.; Meyers, R. H.; Whaley, W. L.; Allitto,

[129912]B. A.; Faryniarz, A.; Wasmuth, J. J.; Frischauf, A.-M.; Conneally,

[129913]P. M.; Lehrach, H.; Gusella, J. F.: A DNA segment encoding two genes

- [129914]very tightly linked to Huntington's disease. Science 238:
950-952,
- [129915]1987.
- [129916]23277.
- [129917]Gilliam, T. C.; Tanzi, R. E.; Haines, J. L.; Bonner, T. I.; Fary-
niarz,
- [129918]A. G.; Hobbs, W. J.; MacDonald, M. E.; Cheng, S. V.; Fol-
stein, S.
- [129919]E.; Conneally, P. M.; Wexler, N. S.; Gusella, J. F.: Localiza-
tion
- [129920]of the Huntington's disease gene to a small segment of
chromosome
- [129921]4 flanked by D4S10 and the telomere. Cell 50: 565-571,
1987.
- [129922]23278.
- [129923]Giordani, B.; Berent, S.; Boivin, M. J.; Penney, J. B.; Lehti-
nen,
- [129924]S.; Markel, D. S.; Hollingsworth, Z.; Butterbaugh, G.;
Hichwa, R.
- [129925]D.; Gusella, J. F.; Young, A. B.: Longitudinal neuropsychol-
ogical
- [129926]and genetic linkage analysis of persons at risk for Hunt-

ington's disease. Arch.

[129927]Neurol. 52: 59-64, 1995.

[129928]23279.

[129929]Goetz, I.; Roberts, E.; Comings, D. E.: Fibroblasts in Huntington's

[129930]disease. New Eng. J. Med. 293: 1225-1227, 1975.

[129931]23280.

[129932]Goldberg, Y. P.; Andrew, S. E.; Theilmann, J.; Kremer, B.; Squitieri,

[129933]F.; Telenius, H.; Brown, J. D.; Hayden, M. R.: Familial predisposition

[129934]to recurrent mutations causing Huntington's disease: genetic risk

[129935]to sibs of sporadic cases. J. Med. Genet. 30: 987-990, 1993.

[129936]23281.

[129937]Goldberg, Y. P.; Kalchman, M. A.; Metzler, M.; Nasir, J.; Zeisler,

[129938]J.; Graham, R.; Koide, H. B.; O'Kusky, J.; Sharp, A. H.; Ross, C.

[129939]A.; Jirik, F.; Hayden, M. R.: Absence of disease phenotype

and intergenerational

[129940]stability of the CAG repeat in transgenic mice expressing the human

[129941]Huntington disease transcript. Hum. Molec. Genet. 5: 177-185, 1996.

[129942]23282.

[129943]Goldberg, Y. P.; Nicholson, D. W.; Rasper, D. M.; Kalchman, M.

[129944]A.; Koide, H. B.; Graham, R. K.; Bromm, M.; Kazemi-Esfarjani, P.;

[129945]Thornberry, N. A.; Vaillancourt, J. P.; Hayden, M. R.: Cleavage of

[129946]huntingtin by apopain, a proapoptotic cysteine protease, is modulated

[129947]by the polyglutamine tract. Nature Genet. 13: 442-449, 1996.

[129948]23283.

[129949]Goldberg, Y. P.; Rommens, J. M.; Andrew, S. E.; Hutchinson, G.

[129950]B.; Lin, B.; Theilmann, J.; Graham, R.; Graves, M. L.; Starr, E.;

[129951]McDonald, H.; Nasir, J.; Schappert, K.; Kalchman, M. A.;

Clarke, L.

[129952]A.; Hayden, M. R.: Identification of an Alu retrotransposi-
tion event

[129953]in close proximity to a strong candidate gene for Hunting-
ton's disease. Nature 362:

[129954]370-373, 1993.

[129955]23284.

[129956]Green, H.: Human genetic diseases due to codon reitera-
tion: relationship

[129957]to an evolutionary mechanism. Cell 74: 955-956, 1993.

[129958]23285.

[129959]Greenberg, L. J.; Martell, R. W.; Theilman, J.; Hayden, M.
R.;

[129960]Joubert, J.: Genetic linkage between Huntington disease
and the D4S10

[129961]locus in South African families: further evidence against
non-allelic

[129962]heterogeneity. Hum. Genet. 87: 701-708, 1991.

[129963]23286.

[129964]Gu, M.; Gash, M. T.; Mann, V. M.; Javoy-Agid, F.; Cooper, J.
M.;

[129965]Schapira, A. H. V.: Mitochondrial defect in Huntington's disease

[129966]caudate nucleus. Ann. Neurol. 39: 385–389, 1996.

[129967]23287.

[129968]Gusella, J.; Tanzi, R. E.; Bader, P. I.; Phelan, M. C.; Stevenson,

[129969]R.; Hayden, M. R.; Hofman, K. J.; Faryniarz, A. G.; Gibbons, K.:

[129970]Deletion of Huntington's disease linked G8 (D4S10) locus in Wolf–Hirschhorn

[129971]syndrome. Nature 318: 75–78, 1985.

[129972]23288.

[129973]Gusella, J. F.; Gibbons, K.; Hobbs, W.; Heft, R.; Anderson, M.;

[129974]Rashtchian, R.; Folstein, S.; Wallace, P.; Conneally, P. M.; Tanzi,

[129975]R.: The G8 locus linked to Huntington's disease. (Abstract) Am.

[129976]J. Hum. Genet. 36: 139S, 1984.

[129977]23289.

[129978]Gusella, J. F.; McNeil, S.; Persichetti, F.; Srinidhi, J.; Novel–

letto,

[129979]A.; Bird, E.; Faber, P.; Vonsattel, J.-P.; Myers, R. H.; Mac-Donald,

[129980]M. E.: Huntington's disease. Cold Spring Harbor Symp. Quant. Biol. 61:

[129981]615-626, 1996.

[129982]23290.

[129983]Dey, B. R.; Furlanetto, R. W.; Nissley, S. P.: Cloning of human

[129984]p55-gamma, a regulatory subunit of phosphatidylinositol 3-kinase,

[129985]by a yeast two-hybrid library screen with the insulin-like growth

[129986]factor-I receptor. Gene 209: 175-183, 1998.

[129987]23291.

[129988]Francke, U.; Yang-Feng, T. L.; Brissenden, J. E.; Ullrich, A.:

[129989]Chromosomal mapping of genes involved in growth control. Cold Spring

[129990]Harbor Symp. Quant. Biol. 51: 855-866, 1986.

[129991]23292.

[129992]Maor, S. B.; Abramovitch, S.; Erdos, M. R.; Brody, L. C.;

Werner,

[129993]H.: BRCA1 suppresses insulin-like growth factor-I receptor promoter

[129994]activity: potential interaction between BRCA1 and Sp1.
Molec. Genet.

[129995]Metab. 69: 130-136, 2000.

[129996]23293.

[129997]Poduslo, S. E.; Dean, M.; Kolch, U.; O'Brien, S. J.: Detecting
[129998]high-resolution polymorphisms in human coding loci by
combining PCR

[129999]and single-strand conformation polymorphism (SSCP)
analysis. Am.

[130000]J. Hum. Genet. 49: 106-111, 1991.

[130001]23294.

[130002]Okumura, N.; Terasawa, F.; Ueno, I.; Oki, K.; Yamauchi, K.;
Hidaka,

[130003]H.; Tozuka, M.; Okura, M.; Katsuyama, T.: Genetic analyses in homozygous

[130004]and heterozygous variants of lactate dehydrogenase-B (H)
subunit:

[130005]LD-B Matsumoto I and II (LD-B W323R). Clin. Chim. Acta

287: 163-171,

[130006]1999.

[130007]23295.

[130008]Rethore, M.-O.; Junien, C.; Malpuech, G.; Baccichetti, C.;
Tenconi,

[130009]R.; Kaplan, J. C.; de Romeuf, J.; Lejeune, J.: Localisation du
gene

[130010]de la glyceraldehyde 3-phosphate dehydrogenase (G3PD)
sur le segment

[130011]distal du bras court du chromosome 12. Ann. Genet. 19:
140-142,

[130012]1976.

[130013]23296.

[130014]Rethore, M.-O.; Kaplan, J.-C.; Junien, C.; Cruveiller, J.;
Dutrillaux,

[130015]B.; Aurias, A.; Carpentier, S.; Lafourcade, J.; Lejeune, J.:
Augmentation

[130016]de l'activite de la LDH-B chez un garçon trisomique 12p
par malsegregation

[130017]d'une translocation maternelle t(12;14)(q12;p11). Ann.
Genet. 18:

[130018]81-87, 1975.

[130019]23297.

[130020]Sakai, I.; Sharief, F. S.; Pan, Y.-C. E.; Li, S. S.-L.: The cDNA

[130021]and protein sequences of human lactate dehydrogenase B.

Biochem.

[130022]J. 248: 933-936, 1987.

[130023]23298.

[130024]Santachiara, A. S.; Nabholz, M.; Miggiano, V.; Darlington,

A.

[130025]J.; Bodmer, W. F.: Linkage between human lactate dehy-

drogenase B

[130026]and peptidase B genes. Nature 227: 248-251, 1970.

[130027]23299.

[130028]Steinbach, P.; Rehder, H.: Tetrasomy for the short arm of

chromosome

[130029]12 with accessory isochromosome (+i[12p]) and a marked

LDH-B gene

[130030]dosage effect. Clin. Genet. 32: 1-4, 1987.

[130031]23300.

[130032]Sudo, K.: Personal Communication. Komae City, Japan

7/12/1993.

[130033]23301.

[130034]Sudo, K.; Maekawa, M.; Ikawa, S.; Machida, K.; Kitamura,
M.; Li,

[130035]S. S.-L.: A missense mutation found in human lactate de-
hydrogenase-B

[130036](H) variant gene. Biochem. Biophys. Res. Commun. 168:
672-676, 1990.

[130037]23302.

[130038]Sudo, K.; Maekawa, M.; Luedemann, M. M.; Deaven, L. L.;
Li, S.

[130039]S.-L.: Human lactate dehydrogenase-B processed pseudo-
gene: nucleotide

[130040]sequence analysis and assignment to the X-chromosome.
Biochem. Biophys.

[130041]Res. Commun. 171: 67-74, 1990.

[130042]23303.

[130043]Sudo, K.; Maekawa, M.; Tomonaga, A.; Tsukada, T.;
Nakayama, T.;

[130044]Kitamura, M.; Li, S. S.-L.; Kanno, T.; Toriumi, J.: Molecular
characterization

[130045]of genetic mutations in human lactate dehydrogenase
(LDH)B (H) variant. Hum.

[130046]Genet. 89: 158-162, 1992.

[130047]23304.

[130048]Van Someren, H.; Meera Khan, P.; Westerveld, A.; Bootsma,
D.:

[130049]Human genetics--two new linkage groups carrying different loci for

[130050]LDH and glutamic-pyruvic transaminase found. Nature
240: 221-222,

[130051]1972.

[130052]23305.

[130053]Weiss, L.; Mayeda, K.; Lindahl, R.; Dully, M.: Localization of

[130054]human LDH-B gene of the short arm of chromosome 12.
(Abstract) Am.

[130055]J. Hum. Genet. 25: 85A only, 1973.

[130056]23306.

[130057]Askew, D. S.; Bartholomew, C.; Buchberg, A. M.; Valentine,
M. B.;

[130058]Jenkins, N. A.; Copeland, N. G.; Ihle, J. N.: His-1 and His-
2: identification

[130059]and chromosomal mapping of two commonly rearranged
sites of viral

[130060]integration in a myeloid leukemia. Oncogene 6:
2041–2047, 1991.

[130061]23307.

[130062]Zhong, S.; Delva, L.; Rachez, C.; Cenciarelli, C.; Gandini,
D.;

[130063]Zhang, H.; Kalantry, S.; Freedman, L. P.; Pandolfi, P. P.: A
RA-dependent,

[130064]tumour–growth suppressive transcription complex is the
target of the

[130065]PML–RAR–alpha and T18 oncoproteins. Nature Genet. 23:
287–295, 1999.

[130066]23308.

[130067]Kamb, A.; Gruis, N. A.; Weaver–Feldhaus, J.; Liu, Q.;
Harshman,

[130068]K.; Tavtigian, S. V.; Stockert, E.; Day, R. S., III; Johnson, B.
E.;

[130069]Skolnick, M. H.: A cell cycle regulator potentially involved
in genesis

[130070]of many tumor types. Science 264: 436–440, 1994.

[130071]23309.

[130072]Bonner, T.; O'Brien, S. J.; Nash, W. G.; Rapp, U. R.; Morton,
C.

[130073]C.; Leder, P.: The human homologs of the raf (mil) oncogene are located
[130074]on human chromosomes 3 and 4. Science 223: 71-74,
1984.
[130075]23310.

[130076]Bonner, T. I.; Kerby, S. B.; Suttrave, P.; Gunnell, M. A.; Mark,
[130077]G.; Rapp, U. R.: Structure and biological activity of human
homologs
[130078]of the raf/mil oncogene. Molec. Cell. Biol. 5: 1400-1407,
1985.
[130079]23311.

[130080]Bonner, T. I.; Oppermann, H.; Seeburg, P.; Kerby, S. B.;
Gunnell,
[130081]M. A.; Young, A. C.; Rapp, U. R.: The complete coding se-
quence of
[130082]the human raf oncogene and the corresponding structure
of the c-raf-1
[130083]gene. Nucleic Acids Res. 14: 1009-1015, 1986.
[130084]23312.

[130085]Bonner, T. I.; Rapp, U. R.; Nash, W. G.; O'Brien, S. J.: Two
human

[130086] homologues to a new retroviral oncogene raf-1 and raf-2
are assigned

[130087] to human chromosomes 3 and 4, respectively. (Abstract)
Cytogenet.

[130088] Cell Genet. 37: 424, 1984.

[130089] 23313.

[130090] Fukui, M.; Yamamoto, T.; Kawai, S.; Maruo, K.; Toyoshima,
K.:

[130091] Detection of a raf-related and two other transforming
DNA sequences

[130092] in human tumors maintained in nude mice. Proc. Nat.
Acad. Sci. 82:

[130093] 5954-5958, 1985.

[130094] 23314.

[130095] Hiroshige, S.; Carlock, L.; Smith, M.: Regional assignment
of

[130096] the RAF2 gene to the region 4pter-4p15. (Abstract) Am. J.
Hum. Genet. 39:

[130097] A157, 1986.

[130098] 23315.

[130099] Hopyan, S.; Gokgoz, N.; Poon, R.; Gensure, R. C.; Yu, C.;

Cole,

[130100]W. G.; Bell, R. S.; Juppner, H.; Andrulis, I. L.; Wunder, J. S.;

Alman,

[130101]B. A.: A mutant PTH/PTHrP type I receptor in enchondro-
matosis. Nature

[130102]Genet. 30: 306–310, 2002.

[130103]23316.

[130104]Sakamoto, O.; Iwama, A.; Amitani, R.; Takehara, T.; Yam-
aguchi,

[130105]N.; Yamamoto, T.; Masuyama, K.; Yamanaka, T.; Ando, M.;
Suda, T.:

[130106]Role of macrophage–stimulating protein and its receptor,
RON tyrosine

[130107]kinase, in ciliary motility. J. Clin. Invest. 99: 701–709,
1997.

[130108]23317.

[130109]Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura, N.;
Shimizu,

[130110]N.: Assignment of hepatocyte growth factor (HGF) gene to
chromosome

[130111]7q21.1. (Abstract) Cytogenet. Cell Genet. 58: 1921 only,
1991.

[130112]23318.

[130113]Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura, N.;
Shimizu,

[130114]N.: Regional localization of the hepatocyte growth factor
(HGF) gene

[130115]to human chromosome 7 band q21.1. Genomics 11:
410-415, 1991.

[130116]23319.

[130117]Gherardi, E.; Stoker, M.: Hepatocytes and scatter factor.
Nature 346:

[130118]228 only, 1990.

[130119]23320.

[130120]Gohda, E.; Tsubouchi, H.; Nakayama, H.; Hirono, S.;
Sakiyama, O.;

[130121]Takahashi, K.; Miyazaki, H.; Hashimoto, S.; Daikuhara, Y.:
Purification

[130122]and partial characterization of hepatocyte growth factor
from plasma

[130123]of a patient with fulminant hepatic failure. J. Clin. Invest.
81:

[130124]414-419, 1988.

[130125]23321.

[130126]Kilby, M. D.; Afford, S.; Li, X. F.; Strain, A. J.; Ahmed, A.;

[130127]Whittle, M. J.: Localisation of hepatocyte growth factor and
its

[130128]receptor (c-met) protein and mRNA in human term pla-
centa. Growth

[130129]Factors 13: 133-139, 1996.

[130130]23322.

[130131]Lai, L.; Goldschneider, I.: Cutting edge: identification of a

[130132]hybrid cytokine consisting of IL-7 and the beta-chain of
the hepatocyte

[130133]growth factor/scatter factor. J. Immun. 167: 3550-3554,
2001.

[130134]23323.

[130135]Maina, F.; Casagrande, F.; Audero, E.; Simeone, A.; Co-
moglio, P.

[130136]M.; Klein, R.; Ponzetto, C.: Uncoupling of Grb2 from the
Met receptor

[130137]in vivo reveals complex roles in muscle development. Cell
87: 531-542,

[130138]1996.

[130139]23324.

[130140]Miyazawa, K.; Tsubouchi, H.; Naka, D.; Takahashi, K.; Okigaki,

[130141]M.; Arakaki, N.; Nakayama, H.; Hirono, S.; Sakiyama, O.; Takahashi,

[130142]K.; Gohda, E.; Daikuhara, Y.; Kitamura, N.: Molecular cloning and

[130143]sequence analysis of cDNA for human hepatocyte growth factor. Biochem.

[130144]Biophys. Res. Commun. 163: 967–973, 1989.

[130145]23325.

[130146]Nakamura, T.; Nishizawa, T.; Hagiya, M.; Seki, T.; Shimonishi,

[130147]M.; Sugimura, A.; Tashiro, K.; Shimizu, S.: Molecular cloning and

[130148]expression of human hepatocyte growth factor. Nature 342: 440–443,

[130149]1989.

[130150]23326.

[130151]Noonan, F. P.; Recio, J. A.; Takayama, H.; Duray, P.; Anver, M.

[130152]R.; Rush, W. L.; De Fabo, E. C.; Merlino, G.: Neonatal sunburn and

[130153]melanoma in mice: severe sunburn in newborn, but not adult, mice is

[130154]linked with melanoma in later life. Nature 413: 271–272, 2001.

[130155]23327.

[130156]Powell, E. M.; Mars, W. M.; Levitt, P.: Hepatocyte growth factor/scatter

[130157]factor is a motogen for interneurons migrating from the ventral to

[130158]dorsal telencephalon. Neuron 30: 79–89, 2001.

[130159]23328.

[130160]Rubin, J. S.; Chan, A. M.–L.; Bottaro, D. P.; Burgess, W. H.;

[130161]Taylor, W. G.; Cech, A. C.; Hirschfield, D. W.; Wong, J.; Miki, T.;

[130162]Finch, P. W.; Aaronson, S. A.: A broad-spectrum human lung fibroblast-derived

[130163]mitogen is a variant of hepatocyte growth factor. Proc. Nat. Acad.

[130164]Sci. 88: 415–419, 1991.

[130165]23329.

[130166]Saccone, S.; Narsimhan, R. P.; Gaudino, G.; Dalpra, L.; Co-
moglio,

[130167]P. M.; Della Valle, G.: Regional mapping of the human
hepatocyte

[130168]growth factor (HGF)-scatter factor gene to chromosome
7q21.1. Genomics 13:

[130169]912-914, 1992.

[130170]23330.

[130171]Schmidt, C.; Blatt, F.; Goedecke, S.; Brinkmann, V.; Zschi-
esche,

[130172]W.; Sharpe, M.; Gherardi, E.; Birchmeier, C.: Scatter factor/
hepatocyte

[130173]growth factor is essential for liver development. Nature
373: 699-702,

[130174]1995.

[130175]23331.

[130176]Uehara, Y.; Minowa, O.; Mori, C.; Shiota, K.; Kuno, J.;
Noda,

[130177]T.; Kitamura, N.: Placental defect and embryonic lethality
in mice

[130178]lacking hepatocyte growth factor/scatter factor. Nature
373: 702–705,

[130179]1995.

[130180]23332.

[130181]Weidner, K. M.; Arakaki, N.; Hartmann, G.; Vandekerck-
hove, J.;

[130182]Weingart, S.; Rieder, H.; Fonatsch, C.; Tsubouchi, H.;
Hishida, T.;

[130183]Daikuhara, Y.; Birchmeier, W.: Evidence for the identity of
human

[130184]scatter factor and human hepatocyte growth factor. Proc.
Nat. Acad.

[130185]Sci. 88: 7001–7005, 1991.

[130186]23333.

[130187]Zarnegar, R.; Petersen, B.; DeFrances, M. C.; Michalopou-
los, G.

[130188]: Localization of hepatocyte growth factor (HGF) gene on
human chromosome

[130189]7. Genomics 12: 147–150, 1992.

[130190]23334.

[130191]Gusella, J. F.; Tanzi, R.; Anderson, M. A.; Ottina, K.; Wal-

lace,

[130192]M.; Conneally, P. M.: Linkage analysis of Huntington's disease using

[130193]RFLPs. (Abstract) Cytogenet. Cell Genet. 37: 484–485, 1984.

[130194]23335.

[130195]Gusella, J. F.; Tanzi, R. E.; Anderson, M. A.; Hobbs, W.; Gibbons,

[130196]K.; Raschtchian, R.; Gilliam, T. C.; Wallace, M. R.; Wexler, N. S.;

[130197]Conneally, P. M.: DNA markers for nervous system diseases. Science 225:

[130198]1320–1326, 1984.

[130199]23336.

[130200]Gusella, J. F.; Wexler, N. S.; Conneally, P. M.; Naylor, S. L.;

[130201]Anderson, M. A.; Tanzi, R. E.; Watkins, P. C.; Ottina, K.; Wallace,

[130202]M. R.; Sakaguchi, A. Y.; Young, A. B.; Shoulson, I.; Bonilla, E.;

[130203]Martin, J. B.: A polymorphic DNA marker genetically linked to Huntington's

[130204]disease. Nature 306: 234–238, 1983.

[130205]23337.

[130206]Gutekunst, C.-A.; Levey, A. I.; Heilman, C. J.; Whaley, W. L.;

[130207]Yi, H.; Nash, N. R.; Rees, H. D.; Madden, J. J.; Hersch, S. M.: Identification

[130208]and localization of huntingtin in brain and human lymphoblastoid cell

[130209]lines with anti-fusion protein antibodies. Proc. Nat. Acad. Sci. 92:

[130210]8710-8714, 1995.

[130211]

[130212]100. Haines, J.; Tanzi, R.; Wexler, N.; Harper, P.; Folstein, S.;

[130213]Cassiman, J.; Meyers, R.; Young, A.; Hayden, M.; Falek, A.; Tolosa,

[130214]E.; Crespi, S.; Campanella, G.; Holmgren, G.; Anvret, M.; Kanazawa,

[130215]I.; Gusella, J.; Conneally, M.: No evidence of linkage heterogeneity

[130216]between Huntington disease (HD) and G8 (D4S10). (Abstract) Am. J.

[130217]Hum. Genet. 39: A156, 1986.

[130218]

[130219]101. Hansson, O.; Petersen, A.; Leist, M.; Nicotera, P.;
Castilho,

[130220]R. F.; Brundin, P.: Transgenic mice expressing a Hunting-
ton's disease

[130221]mutation are resistant to quinolinic acid-induced striatal
excitotoxicity. Proc.

[130222]Nat. Acad. Sci. 96: 8727-8732, 1999.

[130223]

[130224]102. Harding, A. E.: Genetic aspects of autosomal domi-
nant late onset

[130225]cerebellar ataxia. J. Med. Genet. 18: 436-441, 1981.

[130226]

[130227]103. Harper, P. S.: The epidemiology of Huntington's dis-
ease. Hum.

[130228]Genet. 89: 365-376, 1992.

[130229]

[130230]104. Harper, P. S.: Localization of the gene for Hunting-
ton's chorea. Trends

[130231]Neurosci. 7: 1-2, 1984.

[130232]

[130233]105. Harper, P. S.; Lim, C.; Craufurd, D.: Ten years of
presymptomatic

[130234]testing for Huntington's disease: the experience of the UK
Huntington's

[130235]Disease Prediction Consortium. J. Med. Genet. 37:
567-571, 2000.

[130236]

[130237]106. Harper, P. S.; Sarfarazi, M.: Genetic prediction and
family

[130238]structure in Huntington's chorea. Brit. Med. J. 290:
1929-1931,

[130239]1985.

[130240]

[130241]107. Harper, P. S.; Tyler, A.; Walker, D. A.; Newcombe, R.
G.; Davies,

[130242]K.: Huntington's chorea: the basis for long-term preven-
tion. Lancet II:

[130243]346-349, 1979.

[130244]

[130245]108. Harper, P. S.; Youngman, S.; Anderson, M. A.; Sar-

farazi, M.;

[130246]Quarrell, O.; Tanzi, R.; Shaw, D.; Wallace, P.; Conneally, P.
M.;

[130247]Gusella, J. F.: Genetic linkage between Huntington's dis-
ease and

[130248]the DNA polymorphism G8 in South Wales families. J. Med.
Genet. 22:

[130249]447-450, 1985.

[130250]

[130251]109. Hayden, M. R.: Huntington's Chorea. Berlin and New
York: Springer-Verlag

[130252](pub.) 1981.

[130253]

[130254]110. Hayden, M. R.; Beighton, P.: Genetic aspects of Hunt-
ington's

[130255]chorea: results of a national survey. Am. J. Med. Genet.
11: 135-141,

[130256]1982.

[130257]

[130258]111. Hayden, M. R.; Hewitt, J.; Wasmuth, J. J.; Kastelein, J.
J.;

[130259]Langlois, S.; Conneally, M.; Haines, J.; Smith, B.; Hilbert, C.; Allard,

[130260]D.: A polymorphic DNA marker that represents a conserved expressed

[130261]sequence in the region of the Huntington disease gene. Am. J. Hum.

[130262]Genet. 42: 125–131, 1988.

[130263]

[130264]112. Hayden, M. R.; Martin, W. R. W.; Stoessl, A. J.; Clark, C.; Hollenberg,

[130265]S.; Adam, M. J.; Ammann, W.; Harrop, R.; Rogers, J.; Ruth, T.; Sayre,

[130266]C.; Pate, B. D.: Positron emission tomography in the early diagnosis

[130267]of Huntington's disease. Neurology 36: 888–894, 1986.

[130268]

[130269]113. Hayes, C. V.: Genetic testing for Huntington's disease—a family

[130270]issue. (Editorial) New Eng. J. Med. 327: 1449–1451, 1992.

[130271]

[130272]114. Heiser, V.; Scherzinger, E.; Boeddrich, A.; Nordhoff,

E.; Lurz,

[130273]R.; Schugardt, N.; Lehrach, H.; Wanker, E. E.: Inhibition of huntingtin

[130274]fibrillogenesis by specific antibodies and small molecules: implications

[130275]for Huntington's disease therapy. Proc. Nat. Acad. Sci. 97: 6739–6744,

[130276]2000.

[130277]

[130278]115. Hilditch–Maguire, P.; Trettel, F.; Passani, L. A.; Auerbach,

[130279]A.; Persichetti, F.; MacDonald, M. E.: Huntingtin: an iron-regulated

[130280]protein essential for normal nuclear and perinuclear organelles. Hum.

[130281]Molec. Genet. 9: 2789–2797, 2000.

[130282]

[130283]116. Hodge, S. E.; Spence, M. A.; Crandall, B. F.; Sparkes, R. S.;

[130284]Sparkes, M. C.; Crist, M.; Tideman, S.: Huntington disease: linkage

[130285]analysis with age-of-onset corrections. Am. J. Med. Genet.

5: 247–254,

[130286]1980.

[130287]

[130288]117. Hodgson, J. G.; Agopyan, N.; Gutekunst, C.–A.; Leavitt, B. R.;

[130289]LePiane, F.; Singaraja, R.; Smith, D. J.; Bissada, N.; McCutcheon,

[130290]K.; Nasir, J.; Jamot, L.; Li, X.–J.; Stevens, M. E.; Rosemond, E.;

[130291]Roder, J. C.; Phillips, A. G.; Rubin, E. M.; Hersch, S. M.; Hayden,

[130292]M. R.: A YAC mouse model for Huntington's disease with full-length

[130293]mutant huntingtin, cytoplasmic toxicity, and selective striatal neurodegeneration. *Neuron* 23:

[130294]181–192, 1999.

[130295]

[130296]118. Hodgson, J. G.; Smith, D. J.; McCutcheon, K.; Koide, H. B.; Nishiyama,

[130297]K.; Dinulos, M. B.; Stevens, M. E.; Bissada, N.; Nasir, J.; Kanazawa,

[130298]J.; Disteche, C. M.; Rubin, E. M.; Hayden, M. R.: Human

huntingtin

[130299] derived from YAC transgenes compensates for loss of
murine huntingtin

[130300] by rescue of the embryonic lethal phenotype. Hum. Molec.
Genet. 5:

[130301] 1875–1885, 1996.

[130302]

[130303] 119. Holmgren, G.; Winnberg Almqvist, E.; Anvret, M.;
Conneally, M.;

[130304] Hobbs, W.; Mattsson, B.; Wahlstrom, J.; Winblad, B.;
Gusella, J. F.

[130305]: Linkage of G8 (D4S10) in two Swedish families with
Huntington's

[130306] disease. Clin. Genet. 32: 289–294, 1987.

[130307]

[130308] 120. Hoogeveen, A. T.; Willemsen, R.; Meyer, N.; de Rooij,
K. E.;

[130309] Roos, R. A. C.; van Ommen, G.–J. B.; Galjaard, H.: Charac-
terization

[130310] and localization of the Huntington disease gene product.
Hum. Molec.

[130311] Genet. 2: 2069–2073, 1993.

[130312]

[130313]121. Horton, T. M.; Graham, B. H.; Corral-Debrinski, M.; Shoffner,

[130314]J. M.; Kaufman, A. E.; Beal, M. F.; Wallace, D. C.: Marked increase

[130315]in mitochondrial DNA deletion levels in the cerebral cortex of Huntington's

[130316]disease patients. *Neurology* 45: 1879–1883, 1995.

[130317]

[130318]122. Huntington's Disease Collaborative Research Group: A novel

[130319]gene containing a trinucleotide repeat that is expanded and unstable

[130320]on Huntington's disease chromosomes. *Cell* 72: 971–983, 1993.

[130321]

[130322]123. Huntington, G.: On chorea. *Med. Surg. Reporter* 26: 317–321,

[130323]1872.

[130324]

[130325]124. Ikeda, H.; Yamaguchi, M.; Sugai, S.; Aze, Y.; Naru-

miya, S.; Kakizuka,

[130326]A.: Expanded polyglutamine in the Machado–Joseph disease protein

[130327]induces cell death in vitro and in vivo. *Nature Genet.* 13: 196–202,

[130328]1996.

[130329]

[130330]125. Ikonen, E.; Ignatius, J.; Norio, R.; Palo, J.; Peltonen, L.:

[130331]Huntington disease in Finland: a molecular and genealogical study. *Hum.*

[130332]*Genet.* 89: 275–280, 1992.

[130333]

[130334]126. Illarioshkin, S. N.; Igarashi, S.; Onodera, O.; Markova, E. D.;

[130335]Nikolskaya, N. N.; Tanaka, H.; Chabrashwili, T. Z.; Insarova, N. G.;

[130336]Endo, K.; Ivanova–Smolenskaya, I. A.; Tsuji, S.: Trinucleotide repeat

[130337]length and rate of progression of Huntington's disease. *Ann. Neurol.* 36:

[130338]630–635, 1994.

[130339]

[130340]127. Jana, N. R.; Zemskov, E. A.; Wang, G.; Nukina, N.: Altered proteasomal

[130341]function due to the expression of polyglutamine-expanded truncated

[130342]N-terminal huntingtin induces apoptosis by caspase activation through

[130343]mitochondrial cytochrome c release. Hum. Molec. Genet. 10: 1049-1059,

[130344]2001.

[130345]

[130346]128. Kahlem, P.; Green, H.; Djian, P.: Transglutaminase action imitates

[130347]Huntington's disease: selective polymerization of huntingtin containing

[130348]expanded polyglutamine. Molec. Cell 1: 595-601, 1998.

[130349]

[130350]129. Kanazawa, I.; Kondo, I.; Ikeda, J.-E.; Ikeda, T.; Shizu, Y.;

[130351]Yoshida, M.; Narabayashi, H.; Kuroda, S.; Tsunoda, H.; Mizuta, E.;

- [130352] Okuno, Y.; Sugawara, K.; Murata, M.; Takahashi, M.;
Gusella, J. F.
- [130353] Studies on DNA markers (D4S10 and D4S43/S127) ge-
netically linked
to Huntington's disease in Japanese families. Hum. Genet.
85: 257-260,
1990.
- [130356]
- [130357] 130. Karpuj, M. V.; Becher, M. W.; Springer, J. E.; Chabas,
D.; Youssef,
S.; Pedotti, R.; Mitchell, D.; Steinman, L.: Prolonged sur-
vival and
decreased abnormal movements in transgenic model of
Huntington disease,
with administration of the transglutaminase inhibitor cys-
tamine. Nature
Med. 8: 143-149, 2002. Note: Erratum: Nature Med. 8:
303 only, 2002.
- [130362]
- [130363] 131. Kazantsev, A.; Walker, H. A.; Slepko, N.; Bear, J. E.;
Preisinger,
E.; Steffan, J. S.; Zhu, Y.-Z.; Gertler, F. B.; Housman, D. E.;

Marsh,

[130365]J. L.; Thompson, L. M.: A bivalent Huntingtin binding peptide suppresses

[130366]polyglutamine aggregation and pathogenesis in *Drosophila*. *Nature*

[130367]*Genet.* 30: 367–376, 2002.

[130368]

[130369]132. Kazemi-Esfarjani, P.; Benzer, S.: Genetic suppression of polyglutamine

[130370]toxicity in *Drosophila*. *Science* 287: 1837–1840, 2000.

[130371]

[130372]133. Keene, C. D.; Rodrigues, C. M. P.; Eich, T.; Chhabra, M. S.;

[130373]Steer, C. J.; Low, W. C.: Tauroursodeoxycholic acid, a bile acid,

[130374]is neuroprotective in a transgenic animal model of Huntington's disease. *Proc.*

[130375]*Nat. Acad. Sci.* 99: 10671–10676, 2002.

[130376]

[130377]134. Keene, C. D.; Rodrigues, C. M. P.; Eich, T.; Linehan-Stieers,

[130378]C.; Abt, A.; Kren, B. T.; Steer, C. J.; Low, W. C.: A bile acid protects

[130379]against motor and cognitive deficits and reduces striatal degeneration

[130380]in the 3-nitropropionic acid model of Huntington's disease. Exp.

[130381]Neurol. 171: 351-360, 2001.

[130382]

[130383]135. Kehoe, P.; Krawczak, M.; Harper, P. S.; Owen, M. J.; Jones, A.

[130384]L.: Age of onset in Huntington disease: sex specific influence of

[130385]apolipoprotein E genotype and normal CAG repeat length. J. Med. Genet. 36:

[130386]108-111, 1999.

[130387]

[130388]136. Kennedy, L.; Shelbourne, P. F.: Dramatic mutation instability

[130389]in HD mouse striatum: does polyglutamine load contribute to cell-specific

[130390]vulnerability in Huntington's disease? Hum. Molec. Genet. 9: 2539-2544,

[130391]2000.

[130392]

[130393]137. Kerbeshian, J.; Burd, L.; Leech, C.; Rorabaugh, A.:
Huntington

[130394]disease and childhood-onset Tourette syndrome. Am. J.
Med. Genet. 39:

[130395]1-3, 1991.

[130396]

[130397]138. Khoshnan, A.; Ko, J.; Patterson, P. H.: Effects of intra-
cellular

[130398]expression of anti-huntingtin antibodies of various speci-
ficities

[130399]on mutant huntingtin aggregation and toxicity. Proc. Nat.
Acad. Sci. 99:

[130400]1002-1007, 2002.

[130401]

[130402]139. Klawans, H. L., Jr.; Paulson, G. W.; Ringel, S. P.; Bar-
beau,

[130403]A.: L-dopa in the detection of presymptomatic Hunting-
ton's chorea. New

[130404]Eng. J. Med. 286: 1332-1334, 1972.

[130405]

[130406]140. Ko, J.; Ou, S.; Patterson, P. H.: New anti-huntingtin
monoclonal

[130407]antibodies: implications for huntingtin conformation and
its binding

[130408]proteins. Brain Res. Bull. 56: 319–329, 2001.

[130409]

[130410]141. Kovtun, I. V.; McMurray, C. T.: Trinucleotide expansion in haploid

[130411]germ cells by gap repair. Nature Genet. 27: 407–411,
2001.

[130412]

[130413]142. Kovtun, I. V.; Therneau, T. M.; McMurray, C. T.: Gender of the

[130414]embryo contributes to CAG instability in transgenic mice
containing

[130415]a Huntington's disease gene. Hum. Molec. Genet. 9:
2767–2775, 2000.

[130416]

[130417]143. Kremer, B.; Goldberg, P.; Andrew, S. E.; Theilmann, J.;
Telenius,

[130418]H.; Zeisler, J.; Squitieri, F.; Lin, B.; Bassett, A.; Almqvist, E.;

[130419]Bird, T. D.; Hayden, M. R.: A worldwide study of the Huntington's

[130420]disease mutation: the sensitivity and specificity of measuring CAG

[130421]repeats. New Eng. J. Med. 330: 1401–1406, 1994.

[130422]

[130423]144. Laccone, F.; Christian, W.: A recurrent expansion of a maternal

[130424]allele with 36 CAG repeats causes Huntington disease in two sisters. Am.

[130425]J. Hum. Genet. 66: 1145–1148, 2000.

[130426]

[130427]145. Landegent, J. E.; Jansen in de Wal, N.; Fisser–Groen, Y. M.;

[130428]Bakker, E.; van der Ploeg, M.; Pearson, P. L.: Fine mapping of the

[130429]Huntington disease linked D4S10 locus by non-radioactive in situ hybridization. Hum.

[130430]Genet. 73: 354–357, 1986.

[130431]

[130432]146. Lanska, D. J.; Lavine, L.; Lanska, M. J.; Schoenberg, B. S.:

[130433]Huntington's disease mortality in the United States. Neu-
rology 38:

[130434]769-772, 1988.

[130435]

[130436]147. Lazzarini, A.; McCormack, M. K.; Lepore, F.: Maternal
transmission

[130437]of juvenile Huntington's disease in U.S. black families.
(Abstract) Am.

[130438]J. Hum. Genet. 36: 62S, 1984.

[130439]

[130440]148. Leavitt, B. R.; Guttman, J. A.; Hodgson, J. G.; Kimel,
G. H.;

[130441]Singaraja, R.; Vogl, A. W.; Hayden, M. R.: Wild-type hunt-
ingtin reduces

[130442]the cellular toxicity of mutant huntingtin in vivo. Am. J.
Hum. Genet. 68:

[130443]313-324, 2001.

[130444]

[130445]149. Leeflang, E. P.; Tavare, S.; Marjoram, P.; Neal, C. O.

S.; Srinidhi,

[130446]J.; MacDonald, M. E.; de Young, M.; Wexler, N. S.; Gusella,
J. F.;

[130447]Arnhem, N.: Analysis of germline mutation spectra at the
Huntington's

[130448]disease locus supports a mitotic mutation mechanism.

Hum. Molec.

[130449]Genet. 8: 173–183, 1999.

[130450]

[130451]150. Leeflang, E. P.; Zhang, L.; Tavaré, S.; Hubert, R.;

Srinidhi,

[130452]J.; MacDonald, M. E.; Myers, R. H.; de Young, M.; Wexler,

N. S.; Gusella,

[130453]J. F.; Arnheim, N.: Single sperm analysis of the trinucleotide repeats

in the Huntington's disease gene: quantification of the

[130454]mutation frequency

spectrum. Hum. Molec. Genet. 4: 1519–1526, 1995.

[130455]

[130456]

[130457]151. Leung, C. M.; Chan, Y. W.; Chang, C. M.; Yu, Y. L.;

Chen, C.

[130458]N.: Huntington's disease in Chinese: a hypothesis of its

origin. J.

[130459]Neurol. Neurosurg. Psychiat. 55: 681–684, 1992.

[130460]

[130461]152. Li, H.; Li, S.-H.; Johnston, H.; Shelbourne, P. F.; Li,

X.-J.

[130462]: Amino-terminal fragments of mutant huntingtin show selective accumulation

[130463]in striatal neurons and synaptic toxicity. Nature Genet. 25: 385-389,

[130464]2000.

[130465]

[130466]153. Li, S.-H.; Lam, S.; Cheng, A. L.; Li, X.-J.: Intracellular huntingtin

[130467]increases the expression of caspase-1 and induces apoptosis. Hum.

[130468]Molec. Genet. 9: 2859-2867, 2000.

[130469]

[130470]154. Li, X.-J.; Li, S.-H.; Sharp, A. H.; Nucifora, F. C., Jr.; Schilling,

[130471]G.; Lanahan, A.; Worley, P.; Snyder, S. H.; Ross, C. A.: A huntingtin-associated

[130472]protein enriched in brain and implications for pathology. Nature 378:

[130473]398-402, 1995.

[130474]

- [130475]155. Lin, B.; Nasir, J.; Kalchman, M. A.; McDonald, H.; Zeisler, J.;
- [130476]Goldberg, Y. P.; Hayden, M. R.: Structural analysis of the 5-prime
- [130477]region of mouse and human Huntington disease genes reveals conservation
- [130478]of putative promoter region and di- and trinucleotide polymorphisms. *Genomics* 25:
- [130479]707-715, 1995.
- [130480]
- [130481]156. Lin, C.-H.; Tallaksen-Greene, S.; Chien, W.-M.; Cearley, J. A.;
- [130482]Jackson, W. S.; Crouse, A. B.; Ren, S.; Li, X.-J.; Albin, R. L.; Detloff,
- [130483]P. J.: Neurological abnormalities in a knock-in mouse model of Huntington's
- [130484]disease. *Hum. Molec. Genet.* 10: 137-144, 2001.
- [130485]
- [130486]157. Lindblad, A. N.: To test or not to test: an ethical conflict
- [130487]with presymptomatic testing of individuals at 25% risk for Huntington's

[130488]disorder. Clin. Genet. 60: 442–446, 2001.

[130489]

[130490]158. Lovestone, S.; Hodgson, S.; Sham, P.; Differ, A.–M.;
Levy, R.

[130491]: Familial psychiatric presentation of Huntington's disease.
J. Med.

[130492]Genet. 33: 128–131, 1996.

[130493]

[130494]159. Lunkes, A.; Lindenberg, K. S.; Ben–Haiem, L.; Weber,
C.; Devys,

[130495]D.; Landwehrmeyer, G. B.; Mandel, J.–L.; Trottier, Y.: Pro-
teases

[130496]acting on mutant huntingtin generate cleaved products
that differentially

[130497]build up cytoplasmic and nuclear inclusions. Molec. Cell
10: 259–269,

[130498]2002.

[130499]

[130500]160. Lunkes, A.; Mandel, J.–L.: A cellular model that reca-
pitulates

[130501]major pathogenic steps of Huntington's disease. Hum.

Molec. Genet. 7:

[130502]1355–1361, 1998.

[130503]

[130504]161. Luthi–Carter, R.; Strand, A.; Peters, N. L.; Solano, S.
M.; Hollingsworth,

[130505]Z. R.; Menon, A. S.; Frey, A. S.; Spektor, B. S.; Penney, E. B.;
Schilling,

[130506]G.; Ross, C. A.; Borchelt, D. R.; Tapscott, S. J.; Young, A.
B.; Cha,

[130507]J.–H. J.; Olson, J. M.: Decreased expression of striatal sig-
naling

[130508]genes in a mouse model of Huntington's disease. Hum.
Molec. Genet. 9:

[130509]1259–1271, 2000.

[130510]

[130511]162. Lyon, R. L.: Huntington's chorea in the Moray Firth
area. Brit.

[130512]Med. J. 1: 1301–1306, 1962.

[130513]

[130514]163. MacDonald, M. E.; Barnes, G.; Srinidhi, J.; Duyao, M.
P.; Ambrose,

- [130515]C. M.; Myers, R. H.; Gray, J.; Conneally, P. M.; Young, A.; Penney,
- [130516]J.; Shoulson, I.; Hollingsworth, Z.; Koroshetz, W.; Bird, E.; Vonsattel,
- [130517]J. P.; Bonilla, E.; Moscovitz, C.; Penchaszadeh, G.; Brzus-towicz,
- [130518]L.; Alvir, J.; Bickham Conde, J.; Cha, J.-H.; Dure, L.; Gomez, F.;
- [130519]Ramos-Arroyo, M.; Sanchez-Ramos, J.; Snodgrass, S. R.; de Young, M.;
- [130520]Wexler, N. S.; MacFarlane, H.; Anderson, M. A.; Jenkins, B.; Gusella,
- [130521]J. F.: Gametic but not somatic instability of CAG repeat length in
- [130522]Huntington's disease. *J. Med. Genet.* 30: 982-986, 1993.
- [130523]
- [130524]164. MacDonald, M. E.; Cheng, S. V.; Zimmer, M.; Haines, J. L.; Poustka,
- [130525]A.; Allitto, B.; Smith, B.; Whaley, W. L.; Romano, D. M.; Jagadeesh,
- [130526]J.; Myers, R. H.; Lehrach, H.; Wasmuth, J. J.; Frischauf, A.-M.; Gusella,

[130527]J. F.: Clustering of multiallele DNA markers near the Huntington's

[130528]disease gene. J. Clin. Invest. 84: 1013–1016, 1989.

[130529]

[130530]165. MacDonald, M. E.; Duyao, M.; Calzonetti, T.; Auerbach, A.; Ryan,

[130531]A.; Barnes, G.; White, J. K.; Auerbach, W.; Vonsattel, J.-P.; Gusella,

[130532]J. F.; Joyner, A. L.: Targeted inactivation of the mouse Huntington's

[130533]disease gene homolog Hdh. Cold Spring Harbor Symp. Quant. Biol. 61:

[130534]627–638, 1996.

[130535]

[130536]166. MacDonald, M. E.; Haines, J. L.; Zimmer, M.; Cheng, S. V.; Youngman,

[130537]S.; Whaley, W. L.; Wexler, N.; Bucan, M.; Allitto, B. A.; Smith, B.;

[130538]Leavitt, J.; Poustka, A.; Harper, P.; Lehrach, H.; Wasmuth, J. J.;

[130539]Frischauf, A.-M.; Gusella, J. F.: Recombination events suggest potential

[130540]sites for the Huntington's disease gene. Neuron 3:
183–190, 1989.

[130541]

[130542]167. MacDonald, M. E.; Novelletto, A.; Lin, C.; Tagle, D.;
Barnes,

[130543]G.; Bates, G.; Taylor, S.; Allitto, B.; Altherr, M.; Myers, R.;
Lehrach,

[130544]H.; Collins, F. S.; Wasmuth, J. J.; Frontali, M.; Gusella, J. F.:

[130545]The Huntington's disease candidate region exhibits many
different

[130546]haplotypes. Nature Genet. 1: 99–103, 1992.

[130547]

[130548]168. MacDonald, M. E.; Vonsattel, J. P.; Shrinidhi, J.;
Couropmitree,

[130549]N. N.; Cupples, L. A.; Bird, E. D.; Gusella, J. F.; Myers, R.
H.:

[130550]Evidence for the GluR6 gene associated with younger on-
set age of Huntington's

[130551]disease. Neurology 53: 1330–1332, 1999.

[130552]

[130553]169. Magenis, E.; Gusella, J.; Weliky, K.; Haight, G.;

Sheehy, B.

[130554] Huntington disease-linked (HD) restriction fragment polymorphism

[130555] localized to band p16 of chromosome 4 by in situ hybridization. (Abstract) Cytogenet.

[130556] Cell Genet. 40: 685, 1985.

[130557]

[130558] 170. Magenis, R. E.; Gusella, J.; Weliky, K.; Olson, S.; Haight, G.;

[130559] Toth-Fejel, S.; Sheehy, R.: Huntington disease-linked restriction

[130560] fragment length polymorphism localized within band p16.1 of chromosome

[130561] 4 by in situ hybridization. Am. J. Hum. Genet. 39: 383-391, 1986.

[130562]

[130563] 171. Maltzberger, J. T.: Even unto the twelfth generation-Huntington's

[130564] chorea. J. Hist. Med. Allied Sci. 16: 1-17, 1961.

[130565]

[130566] 172. Mangiarini, L.; Sathasivam, K.; Mahal, A.; Mott, R.;

Seller,

[130567]M.; Bates, G. P.: Instability of highly expanded CAG repeats in mice

[130568]transgenic for the Huntington's disease mutation. Nature Genet. 15:

[130569]197–200, 1997.

[130570]

[130571]173. Mangiarini, L.; Sathasivam, K.; Seller, M.; Cozens, B.; Harper,

[130572]A.; Hetherington, C.; Lawton, M.; Trottier, Y.; Lehrach, H.; Davies,

[130573]S. W.; Bates, G. P.: Exon 1 of the HD gene with an expanded CAG repeat

[130574]is sufficient to cause a progressive neurological phenotype in transgenic

[130575]mice. Cell 87: 493–506, 1996.

[130576]

[130577]174. Martin, J. B.; Gusella, J. F.: Huntington's disease: pathogenesis

[130578]and management. New Eng. J. Med. 315: 1267–1276, 1986.

[130579]

[130580]175. Marx, J. L.: A parent's sex may affect gene expression. Science 239:

[130581]352–353, 1988.

[130582]

[130583]176. Mastromauro, C. A.; Meissen, G. J.; Cupples, L. A.; Kiely, D.

[130584]K.; Berkman, B.; Myers, R. H.: Estimation of fertility and fitness

[130585]in Huntington disease in New England. Am. J. Med. Genet. 33: 248–254,

[130586]1989.

[130587]

[130588]177. Masuda, N.; Goto, J.; Murayama, N.; Watanabe, M.; Kondo, I.;

[130589]Kanazawa, I.: Analysis of triplet repeats in the huntingtin gene

[130590]in Japanese families affected with Huntington's disease. J. Med.

[130591]Genet. 32: 701–705, 1995.

[130592]

[130593]178. Mazziotta, J. C.; Phelps, M. E.; Pahl, J. J.; Huang,

S.-C.; Baxter,

[130594]L. R.; Riege, W. H.; Hoffman, J. M.; Kuhl, D. E.; Lanto, A. B.;

Wapenski,

[130595]J. A.; Markham, C. H.: Reduced cerebral glucose

metabolism in asymptomatic

[130596]subjects at risk for Huntington's disease. New Eng. J. Med.

316:

[130597]357-362, 1987.

[130598]

[130599]179. McKeown, C.; Read, A. P.; Dodge, A.; Stecko, O.;

Mercer, A.;

[130600]Harris, R.: Wolf-Hirschhorn locus is distal to D4S10 on

short arm

[130601]of chromosome 4. J. Med. Genet. 24: 410-412, 1987.

[130602]

[130603]180. McKusick, V. A.: Personal Communication. Baltimore,

Md. 1977.

[130604]

[130605]181. McLaughlin, B. A.; Spencer, C.; Eberwine, J.: CAG

trinucleotide

[130606]RNA repeats interact with RNA-binding proteins. Am. J.

Hum. Genet. 59:

[130607]561–569, 1996.

[130608]

[130609]182. Meissen, G. J.; Myers, R. H.; Mastromauro, C. A.; Koro-

roshetz,

[130610]W. J.; Klinger, K. W.; Farrer, L. A.; Watkins, P. A.; Gusella, J.

[130611]F.; Bird, E. D.; Martin, J. B.: Predictive testing for Hunting-

ton's

[130612]disease with use of a linked DNA marker. New Eng. J. Med.

318: 535–542,

[130613]1988.

[130614]

[130615]183. Metzler, M.; Helgason, C. D.; Dragatsis, I.; Zhang, T.;

Gan,

[130616]L.; Pineault, N.; Zeitlin, S. O.; Humphries, R. K.; Hayden, M.

R.

[130617]: Huntingtin is required for normal hematopoiesis. Hum.

Molec. Genet. 9:

[130618]387–394, 2000.

[130619]

[130620]184. Millan, F. A.; Curits, A.; Mennie, M.; Holloway, S.;

Boxer, M.;

[130621]Faed, M. J. W.; Crawford, J. W.; Liston, W. A.; Brock, D. J.
H.:

[130622]Prenatal exclusion testing for Huntington's disease: a
problem of

[130623]too much information. J. Med. Genet. 26: 83–85, 1989.

[130624]

[130625]185. Mochizuki, H.; Kamakura, K.; Kumada, M.; Goto, J.;
Kanazawa,

[130626]I.; Motoyoshi, K.: A patient with Huntington's disease pre-
senting

[130627]with laryngeal chorea. Europ. Neurol. 41: 119–120, 1999.

[130628]

[130629]186. Morris, M. J.; Tyler, A.; Lazarou, L.; Meredith, L.;
Harper,

[130630]P. S.: Problems in genetic prediction for Huntington's dis-
ease. Lancet II:

[130631]601–603, 1989.

[130632]

[130633]187. Morrison, P. J.; Johnston, W. P.; Nevin, N. C.: The epi-
demiology

[130634]of Huntington's disease in Northern Ireland. J. Med. Genet.

32:

[130635]524–530, 1995.

[130636]

[130637]188. Mouchiroud, D.; D'Onofrio, G.; Aissani, B.; Macaya, G.; Gautier,

[130638]C.; Bernardi, G.: The distribution of genes in the human genome. Gene 100:

[130639]181–187, 1991.

[130640]

[130641]189. Muchowski, P. J.; Ning, K.; D'Souza-Schorey, C.; Fields, S.:

[130642]Requirement of an intact microtubule cytoskeleton for aggregation

[130643]and inclusion body formation by a mutant huntingtin fragment. Proc.

[130644]Nat. Acad. Sci. 99: 727–732, 2002.

[130645]

[130646]190. Myers, R. H.; Cupples, L. A.; Schoenfeld, M.; D'Agostino, R.

[130647]B.; Terrin, N. C.; Goldmakher, N.; Wolf, P. A.: Maternal fac-

tors

[130648]in onset of Huntington disease. Am. J. Hum. Genet. 37:
511–523,

[130649]1985.

[130650]

[130651]191. Myers, R. H.; Goldman, D.; Bird, E. D.; Sax, D. S.;
Merril, C.

[130652]R.; Schoenfeld, M.; Wolf, P. A.: Maternal transmission in
Huntington's

[130653]disease. Lancet I: 208–210, 1983.

[130654]

[130655]192. Myers, R. H.; Leavitt, J.; Farrer, L. A.; Jagadeesh, J.;
McFarlane,

[130656]H.; Mastromauro, C. A.; Mark, R. J.; Gusella, J. F.: Homozy-
gote for

[130657]Huntington disease. Am. J. Hum. Genet. 45: 615–618,
1989.

[130658]

[130659]193. Myers, R. H.; Madden, J. J.; Teague, J. L.; Falek, A.:
Factors

[130660]related to onset age of Huntington disease. Am. J. Hum.

Genet. 34:

[130661]481–488, 1982.

[130662]

[130663]194. Myrianthopoulos, N. C.: Huntington's chorea. J. Med.

Genet. 3:

[130664]298–314, 1966.

[130665]

[130666]195. Narain, Y.; Wyttenbach, A.; Rankin, J.; Furlong, R. A.;

Rubinsztein,

[130667]D. C.: A molecular investigation of true dominance in

Huntington's

[130668]disease. J. Med. Genet. 36: 739–746, 1999.

[130669]

[130670]196. Nasir, J.; Floresco, S. B.; O'Kusky, J. R.; Diewert, V. M.;

Richman,

[130671]J. M.; Zeisler, J.; Borowski, A.; Marth, J. D.; Phillips, A. G.;

Hayden,

[130672]M. R.: Targeted disruption of the Huntington's disease

gene results

[130673]in embryonic lethality and behavioral and morphological

changes in

[130674]heterozygotes. Cell 81: 811–823, 1995.

[130675]

[130676]197. Nasir, J.; Lin, B.; Bucan, M.; Koizumi, T.; Nadeau, J. H.; Hayden,

[130677]M. R.: The murine homologues of the Huntington disease gene (Hdh)

[130678]and the alpha-adducin gene (Add1) map to mouse chromosome 5 within

[130679]a region of conserved synteny with human chromosome 4p16.3. Genomics 22:

[130680]198–201, 1994.

[130681]

[130682]198. Navarrete, C.; Martinez, I.; Salamanca, F.: Paternal line of

[130683]transmission in chorea of Huntington with very early onset. Genet.

[130684]Counsel. 5: 175–178, 1994.

[130685]

[130686]199. Ona, V. O.; Li, M.; Vonsattel, J. P. G.; Andrews, L. J.; Khan,

[130687]S. Q.; Chung, W. M.; Frey, A. S.; Menon, A. S.; Li, X.-J.;

Stieg,

[130688]P. E.; Yuan, J.; Penney, J. B.; Young, A. B.; Cha, J.-H. J.;

Friedlander,

[130689]R. M.: Inhibition of caspase-1 slows disease progression
in a mouse

[130690]model of Huntington's disease. Nature 399: 263-267,
1999.

[130691]

[130692]200. Ordway, J. M.; Tallaksen-Greene, S.; Gutekunst,
C.-A.; Bernstein,

[130693]E. M.; Cearley, J. A.; Wiener, H. W.; Dure, L. S., IV; Lindsey,
R.;

[130694]Hersch, S. M.; Jope, R. S.; Albin, R. L.; Detloff, P. J.: Ectopi-
cally

[130695]expressed CAG repeats cause intranuclear inclusions and
a progressive

[130696]late onset neurological phenotype in the mouse. Cell 91:
753-763,

[130697]1997.

[130698]

[130699]201. Orr, H. T.; Zoghbi, H. Y.: Reversing neurodegenera-
tion: a promise

[130700]unfolds. Cell 101: 1–4, 2000.

[130701]

[130702]202. Osler, W.: Remarks on the varieties of chronic chorea, and a

[130703]report upon two families of the hereditary form, with one autopsy. J.

[130704]Nerv. Ment. Dis. 18: 97–111, 1893.

[130705]

[130706]203. Palo, J.; Somer, H.; Ikonen, E.; Karila, L.; Peltonen, L.: Low

[130707]prevalence of Huntington's disease in Finland. Lancet I: 805–806,

[130708]1987.

[130709]

[130710]204. Panov, A. V.; Gutekunst, C.–A.; Leavitt, B. R.; Hayden, M. R.;

[130711]Burke, J. R.; Strittmatter, W. J.; Greenamyre, J. T.: Early mitochondrial

[130712]calcium defects in Huntington's disease are a direct effect of polyglutamines. Nature

[130713]Neurosci. 5: 731–736, 2002.

[130714]

[130715]205. Parrish, J. E.; Nelson, D. L.: Methods for finding
genes: a

[130716]major rate-limiting step in positional cloning. Genet. Anal.
Tech.

[130717]Appl. 10: 29-41, 1993.

[130718]

[130719]206. Paulson, H. L.; Bonini, N. M.; Roth, K. A.: Polyglu-
tamine disease

[130720]and neuronal cell death. Proc. Nat. Acad. Sci. 97:
12957-12958,

[130721]2000.

[130722]

[130723]207. Peel, A. L.; Rao, R. V.; Cottrell, B. A.; Hayden, M. R.;
Ellerby,

[130724]L. M.; Bredesen, D. E.: Double-stranded RNA-dependent
protein kinase,

[130725]PKR, binds preferentially to Huntington's disease (HD)
transcripts

[130726]and is activated in HD tissue. Hum. Molec. Genet. 10:
1531-1538,

[130727]2001.

[130728]

[130729]208. Pericak-Vance, M. A.; Conneally, P. M.; Merritt, A. D.;
Roos,
[130730]R.; Norton, J. A., Jr.; Vance, J. M.: Genetic linkage studies
in
[130731]Huntington disease. Cytogenet. Cell Genet. 22: 640-645,
1978.

[130732]

[130733]209. Perry, T. L.; Hansen, S.; Kloster, M.: Huntington's
chorea:
[130734]deficiency of gamma-aminobutyric acid in brain. New Eng.
J. Med. 288:
[130735]337-342, 1973.

[130736]

[130737]210. Petersen, A.; Larsen, K. E.; Behr, G. G.; Romero, N.;
Przedborski,
[130738]S.; Brundin, P.; Sulzer, D.: Expanded CAG repeats in exon
1 of the
[130739]Huntington's disease gene stimulate dopamine-mediated
striatal neuron

[130740]autophagy and degeneration. Hum. Molec. Genet. 10:
1243–1254, 2001.

[130741]

[130742]211. Peyser, C. E.; Folstein, M.; Chase, G. A.; Starkstein, S.;
Brandt,

[130743]J.; Cockrell, J. R.; Bylsma, F.; Coyle, J. T.; McHugh, P. R.;
Folstein,

[130744]S. E.: Trial of d-alpha-tocopherol in Huntington's disease.
Am.

[130745]J. Psychiat. 152: 1771–1775, 1995.

[130746]

[130747]212. Portera-Cailliau, C.; Hedreen, J. C.; Price, D. L.; Koli-
atsos,

[130748]V. E.: Evidence for apoptotic cell death in Huntington dis-
ease and

[130749]excitotoxic animal models. J. Neurosci. 15: 3775–3787,
1995.

[130750]

[130751]213. Pridmore, S. A.: The large Huntington's disease fam-
ily of Tasmania. Med.

[130752]J. Aust. 153: 593–595, 1990.

[130753]

[130754]214. Quarrell, O. W. J.; Meredith, A. L.; Tyler, A.; Youngman, S.;

[130755]Upadhyaya, M.; Harper, P. S.: Exclusion testing for Huntington's

[130756]disease in pregnancy with a closely linked DNA marker.

Lancet I:

[130757]1281-1283, 1987.

[130758]

[130759]215. Quarrell, O. W. J.; Tyler, A.; Cole, G.; Harper, P. S.:
The

[130760]problem of isolated cases of Huntington's disease in South
Wales 1974-1984. Clin.

[130761]Genet. 30: 433-439, 1986.

[130762]

[130763]216. Quarrell, O. W. J.; Tyler, A.; Jones, M. P.; Nordin, M.;
Harper,

[130764]P. S.: Population studies of Huntington's disease in Wales.
Clin.

[130765]Genet. 33: 189-195, 1988.

[130766]

[130767]217. Ranen, N. G.; Stine, O. C.; Abbott, M. H.; Sherr, M.;
Codori,

[130768]A.-M.; Franz, M. L.; Chao, N. I.; Chung, A. S.; Pleasant, N.;
Callahan,

[130769]C.; Kasch, L. M.; Ghaffari, M.; Chase, G. A.; Kazazian, H.
H.; Brandt,

[130770]J.; Folstein, S. E.; Ross, C. A.: Anticipation and instability of
[130771]IT-15 (CAG) n repeats in parent-offspring pairs with Hunt-
ington disease. Am.

[130772]J. Hum. Genet. 57: 593-602, 1995.

[130773]

[130774]218. Read, A. P.: Huntington's disease: testing the test.
Nature

[130775]Genet. 4: 329-330, 1993.

[130776]

[130777]219. Reddy, P. H.; Williams, M.; Charles, V.; Garrett, L.;
Pike-Buchanan,

[130778]L.; Whetsell, W. O., Jr.; Miller, G.; Tagle, D. A.: Behavioral
abnormalities

[130779]and selective neuronal loss in HD transgenic mice ex-
pressing mutated

[130780]full-length HD cDNA. Nature Genet. 20: 198–202, 1998.

[130781]

[130782]220. Reed, T. E.; Chandler, J. H.: Huntington's chorea in Michigan.

[130783]I. Demography and genetics. Am. J. Hum. Genet. 10: 203–225, 1958.

[130784]

[130785]221. Reed, T. E.; Neel, J. V.: Huntington's chorea in Michigan. II.

[130786]Selection and mutation. Am. J. Hum. Genet. 11: 107–136, 1959.

[130787]

[130788]222. Reik, W.: Genomic imprinting: a possible mechanism for the parental

[130789]origin effect in Huntington's chorea. J. Med. Genet. 25: 805–808,

[130790]1988.

[130791]

[130792]223. Reik, W.: Genomic imprinting and genetic disorders in man. Trends

[130793]Genet. 5: 331–336, 1989.

[130794]

[130795]224. Reik, W.; Collick, A.; Norris, M. L.; Barton, S. C.;
Surani,

[130796]M. A.: Genomic imprinting determines methylation of
parental alleles

[130797]in transgenic mice. *Nature* 328: 248–251, 1987.

[130798]

[130799]225. Reiner, A.; Albin, R. L.; Anderson, K. D.; D'Amato, C.
J.; Penney,

[130800]J. B.; Young, A. B.: Differential loss of striatal projection
neurons

[130801]in Huntington disease. *Proc. Nat. Acad. Sci.* 85:
5733–5737, 1988.

[130802]

[130803]226. Richfield, E. K.; Herkenham, M.: Selective vulnerabil-
ity in

[130804]Huntington's disease: preferential loss of cannabinoid re-
ceptors in

[130805]lateral globus pallidus. *Ann. Neurol.* 36: 577–584, 1994.

[130806]

[130807]227. Ridley, R. M.; Frith, C. D.; Crow, T. J.; Conneally, P.

M.:

[130808]Anticipation in Huntington's disease is inherited through the male

[130809]line but may originate in the female. J. Med. Genet. 25: 589-595,

[130810]1988.

[130811]

[130812]228. Ridley, R. M.; Frith, C. D.; Farrer, L. A.; Conneally, P. M.

[130813]: Patterns of inheritance of the symptoms of Huntington's disease

[130814]suggestive of an effect of genomic imprinting. J. Med. Genet. 28:

[130815]224-231, 1991.

[130816]

[130817]229. Robbins, C.; Theilmann, J.; Youngman, S.; Haines, J.; Altherr,

[130818]M. J.; Harper, P. S.; Payne, C.; Junker, A.; Wasmuth, J.; Hayden,

[130819]M. R.: Evidence from family studies that the gene causing Huntington

[130820]disease is telomeric to D4S95 and D4S90. Am. J. Hum.

Genet. 44:

[130821]422-425, 1989.

[130822]

[130823]230. Rosenberg, N. K.; Sorensen, S. A.; Christensen, A.-L.:
Neuropsychological

[130824]characteristics of Huntington's disease carriers: a double
blind study. J.

[130825]Med. Genet. 32: 600-604, 1995.

[130826]

[130827]231. Rosenblatt, A.; Brinkman, R. R.; Liang, K. Y.;
Almqvist, E. W.;

[130828]Margolis, R. L.; Huang, C. Y.; Sherr, M.; Franz, M. L.; Ab-
bott, M.

[130829]H.; Hayden, M. R.; Ross, C. A.: Familial influence on age of
onset

[130830]among siblings with Huntington disease. Am. J. Med.
Genet. 105:

[130831]399-403, 2001.

[130832]

[130833]232. Roses, A. D.: From genes to mechanisms to thera-
pies: lessons

[130834]to be learned from neurological disorders. Nature Med. 2:
267–269,

[130835]1996.

[130836]

[130837]233. Rubinsztein, D. C.; Amos, W.; Leggo, J.; Goodburn,
S.; Ramesar,

[130838]R. S.; Old, J.; Bontrop, R.; McMahon, R.; Barton, D. E.; Fer-
guson-Smith,

[130839]M. A.: Mutational bias provides a model for the evolution
of Huntington's

[130840]disease and predicts a general increase in disease preva-
lence. Nature

[130841]Genet. 7: 525–530, 1994.

[130842]

[130843]234. Rubinsztein, D. C.; Leggo, J.; Coles, R.; et al.: Pheno-
typic

[130844]characterization of individuals with 30–40 CAG repeats in
the Huntington

[130845]disease (HD) gene reveals HD cases with 36 repeats and
apparently

[130846]normal elderly individuals with 36–39 repeats. Am. J.
Hum. Genet. 59:

[130847]16–22, 1996.

[130848]

[130849]235. Sabl, J. F.; Laird, C. D.: Epigene conversion: a proposal with

[130850]implications for gene mapping in humans. *Am. J. Hum. Genet.* 50:

[130851]1171–1177, 1992.

[130852]

[130853]236. Saccone, S.; De Sario, A.; Della Valle, G.; Bernardi, G.:
The

[130854]highest gene concentrations in the human genome are in
telomeric bands

[130855]of metaphase chromosomes. *Proc. Nat. Acad. Sci.* 89:
4913–4917, 1992.

[130856]

[130857]237. Sapienza, C.; Peterson, A. C.; Rossant, J.; Balling, R.:
Degree

[130858]of methylation of transgenes is independent of gamete of
origin. *Nature* 328:

[130859]251–254, 1987.

[130860]

- [130861]238. Sapp, E.; Ge, P.; Aizawa, H.; Bird, E.; Penney, J.;
Young, A.
[130862]B.; Vonsattel, J.-P.; DiFiglia, M.: Evidence for a preferential
loss
[130863]of enkephalin immunoreactivity in the external globus
pallidus in
[130864]low grade Huntington's disease using high resolution im-
age analysis. Neuroscience 64:
[130865]397-404, 1995.
[130866]
[130867]239. Sathasivam, K.; Hobbs, C.; Turmaine, M.; Mangiarini,
L.; Mahal,
[130868]A.; Bertaux, F.; Wanker, E. E.; Doherty, P.; Davies, S. W.;
Bates,
[130869]G. P.: Formation of polyglutamine inclusions in non-CNS
tissue. Hum.
[130870]Molec. Genet. 8: 813-822, 1999.
[130871]
[130872]240. Sathasivam, K.; Woodman, B.; Mahal, A.; Bertaux, F.;
Wanker,
[130873]E. E.; Shima, D. T.; Bates, G. P.: Centrosome disorganiza-
tion in

[130874] fibroblast cultures derived from R6/2 Huntington's disease (HD) transgenic

[130875] mice and HD patients. Hum. Molec. Genet. 10: 2425–2435, 2001.

[130876]

[130877] 241. Saudou, F.; Finkbeiner, S.; Devys, D.; Greenberg, M. E.: Huntingtin

[130878] acts in the nucleus to induce apoptosis but death does not correlate

[130879] with the formation of intranuclear inclusions. Cell 95: 55–66, 1998.

[130880]

[130881] 242. Sax, D. S.; Bird, E. D.; Gusella, J. F.; Myers, R. H.: Phenotypic

[130882] variation in 2 Huntington's disease families with linkage to chromosome

[130883] 4. Neurology 39: 1332–1336, 1989.

[130884]

[130885] 243. Scherzinger, E.; Lurz, R.; Turmaine, M.; Mangiarini, L.; Hollenbach,

[130886] B.; Hasenbank, R.; Bates, G. P.; Davies, S. W.; Lehrach, H.; Wanker,

[130887]E. E.: Huntingtin-encoded polyglutamine expansions form amyloid-like

[130888]protein aggregates in vitro and in vivo. Cell 90: 549-558, 1997.

[130889]

[130890]244. Scherzinger, E.; Sittler, A.; Schweiger, K.; Heiser, V.; Lurz,

[130891]R.; Hasenbank, R.; Bates, G. P.; Lehrach, H.; Wanker, E. E.: Self-assembly

[130892]of polyglutamine-containing huntingtin fragments into amyloid-like

[130893]fibrils: implications for Huntington's disease pathology. Proc. Nat.

[130894]Acad. Sci. 96: 4604-4609, 1999.

[130895]

[130896]245. Schilling, G.; Becher, M. W.; Sharp, A. H.; Jinnah, H. A.; Duan,

[130897]K.; Kotzuk, J. A.; Slunt, H. H.; Ratovitski, T.; Cooper, J. K.; Jenkins,

[130898]N. A.; Copeland, N. G.; Price, D. L.; Ross, C. A.; Borchelt, D. R.

[130899]: Intranuclear inclusions and neuritic aggregates in trans-

genic mice

[130900]expressing a mutant N-terminal fragment of huntingtin.

Hum. Molec.

[130901]Genet. 8: 397-407, 1999.

[130902]

[130903]246. Schwarcz, R.; Okuno, E.; White, R. J.; Bird, E. D.;

Whetsell,

[130904]W. O., Jr.: 3-Hydroxyanthranilate oxygenase activity is increased

[130905]in the brains of Huntington disease victims. Proc. Nat.

Acad. Sci. 85:

[130906]4079-4081, 1988.

[130907]

[130908]247. Scrimgeour, E. M.: Possible introduction of Huntington's chorea

[130909]into Pacific Islands by New England whalers. Am. J. Med.

Genet. 15:

[130910]607-613, 1983.

[130911]

[130912]248. Scrimgeour, E. M.; Samman, Y.; Brock, D. J. H.: Huntington's

[130913]disease in a Sudanese family from Khartoum. Hum. Genet.
96: 624–625,
[130914]1995.

[130915]

[130916]249. Shelbourne, P. F.; Killeen, N.; Hevner, R. F.; Johnston,
H. M.;

[130917]Tecott, L.; Lewandoski, M.; Ennis, M.; Ramirez, L.; Li, Z.;
Iannicola,

[130918]C.; Littman, D. R.; Myers, R. M.: A Huntington's disease
CAG expansion

[130919]at the murine Hdh locus is unstable and associated with
behavioural

[130920]abnormalities in mice. Hum. Molec. Genet. 8: 763–774,
1999.

[130921]

[130922]250. Shiwach, R.: Psychopathology in Huntington's disease
patients. Acta

[130923]Psychiat. Scand. 90: 241–246, 1994.

[130924]

[130925]251. Shiwach, R. S.; Norbury, C. G.: A controlled psychi-
atric study

[130926] of individuals at risk for Huntington's disease. Brit. J. Psychiatry. 165:

[130927] 500–505, 1994.

[130928]

[130929] 252. Silber, E.; Kromberg, J.; Temlett, J. A.; Krause, A.; Saffer,

[130930] D.: Huntington's disease confirmed by genetic testing in five African

[130931] families. Mov. Disord. 13: 726–730, 1998.

[130932]

[130933] 253. Simpson, S. A.; Johnston, A. W.: The prevalence and patterns

[130934] of care of Huntington's chorea in Grampian. Brit. J. Psychiatry. 155:

[130935] 799–804, 1989.

[130936]

[130937] 254. Sisodia, S. S.: Nuclear inclusions in glutamine repeat disorders:

[130938] are they pernicious, coincidental, or beneficial? Cell 95: 1–4,

[130939] 1998.

[130940]

[130941]255. Sittler, A.; Lurz, R.; Lueder, G.; Priller, J.; Hayer-Hartl,

[130942]M. K.; Hartl, F. U.; Lehrach, H.; Wanker, E. E.: Gel-

danamycin activates

[130943]a heat shock response and inhibits huntingtin aggregation

in a cell

[130944]culture model of Huntington's disease. Hum. Molec.

Genet. 10: 1307-1315,

[130945]2001.

[130946]

[130947]256. Skraastad, M. I.; Van de Vosse, E.; Belfroid, R.; Hold,

K.; Vegter-van

[130948]der Vlis, M.; Sandkuijl, L. A.; Bakker, E.; van Ommen, G. J.

B.:

[130949]Significant linkage disequilibrium between the Huntington

disease

[130950]gene and the loci D4S10 and D4S95 in the Dutch popula-

tion. Am. J.

[130951]Hum. Genet. 51: 730-735, 1992.

[130952]

[130953]257. Snell, R. G.; MacMillan, J. C.; Cheadle, J. P.; Fenton, I.;

Lazarou,

[130954]L. P.; Davies, P.; MacDonald, M. E.; Gusella, J. F.; Harper, P. S.;

[130955]Shaw, D. J.: Relationship between trinucleotide repeat expansion

[130956]and phenotypic variation in Huntington's disease. Nature Genet. 4:

[130957]393–397, 1993.

[130958]

[130959]258. Snell, R. G.; Youngman, S.; Lehrach, H.; Sarfarazi, M.; Harper,

[130960]P. S.; Shaw, D. J.: A new probe (2R3) in the region of Huntington's

[130961]disease. (Abstract) Cytogenet. Cell Genet. 51: 1083, 1989.

[130962]

[130963]259. Stebbins, C. E.; Russo, A. A.; Schneider, C.; Rosen, N.; Hartl,

[130964]F. U.; Pavletich, N. P.: Crystal structure of an Hsp90–geldanamycin

[130965]complex: targeting of a protein chaperone by an antitumor agent. Cell 89:

[130966]239–250, 1997.

[130967]

[130968]260. Steffan, J. S.; Bodal, L.; Pallos, J.; Poelman, M.; Mc-Campbell,

[130969]A.; Apostol, B. L.; Kazantsev, A.; Schmidt, E.; Zhu, Y.-Z.; Greenwald,

[130970]M.; Kurokawa, R.; Housman, D. E.; Jackson, G. R.; Marsh, J. L.; Thompson,

[130971]L. M.: Histone deacetylase inhibitors arrest polyglutamine-dependent

[130972]neurodegeneration in *Drosophila*. *Nature* 413: 739-743, 2001.

[130973]

[130974]261. Steffan, J. S.; Kazantsev, A.; Spasic-Boskovic, O.; Greenwald,

[130975]M.; Zhu, Y.-Z.; Gohler, H.; Wanker, E. E.; Bates, G. P.; Housman,

[130976]D. E.; Thompson, L. M.: The Huntington's disease protein interacts

[130977]with p53 and CREB-binding protein and represses transcription. *Proc.*

[130978]*Nat. Acad. Sci.* 97: 6763-6768, 2000.

[130979]

[130980]262. Stine, O. C.; Smith, K. D.: The estimation of selection coefficients

[130981]in Afrikaners: Huntington disease, porphyria variegata, and lipoid

[130982]proteinosis. Am. J. Hum. Genet. 46: 452–458, 1990.

[130983]

[130984]263. Strobel, S. A.; Doucette–Stamm, L. A.; Riba, L.; Housman, D.

[130985]E.; Dervan, P. B.: Site–specific cleavage of human chromosome 4 mediated

[130986]by triple–helix formation. Science 254: 1639–1642, 1991.

[130987]

[130988]264. Tolmie, J. L.; Davidson, H. R.; May, H. M.; McIntosh, K.; Paterson,

[130989]J. S.; Smith, B.: The prenatal exclusion test for Huntington's disease:

[130990]experience in the West of Scotland, 1986–1993. J. Med. Genet. 32:

[130991]97–101, 1995.

[130992]

[130993]265. Tranebjaerg, L.; Petersen, A.; Hove, K.; Rehder, H.;

Mikkelsen,

[130994]M.: Clinical and cytogenetic studies in a large (4;8)
translocation

[130995]family with pre- and postnatal Wolf syndrome. Ann.
Genet. 27: 224-229,

[130996]1984.

[130997]

[130998]266. Trettel, F.; Rigamonti, D.; Hilditch-Maguire, P.;
Wheeler, V.

[130999]C.; Sharp, A. H.; Persichetti, F.; Cattaneo, E.; MacDonald,
M. E.

[131000]: Dominant phenotypes produced by the HD mutation in
STHdh(Q111) striatal

[131001]cells. Hum. Molec. Genet. 9: 2799-2809, 2000.

[131002]

[131003]267. Trottier, Y.; Biancalana, V.; Mandel, J.-L.: Instability
of

[131004]CAG repeats in Huntington's disease: relation to parental
transmission

[131005]and age of onset. J. Med. Genet. 31: 377-382, 1994.

[131006]

- [131007]268. Trottier, Y.; Devys, D.; Imbert, G.; Saudou, F.; An, I.;
Lutz,
[131008]Y.; Weber, C.; Agid, Y.; Hirsch, E. C.; Mandel, J.-L.: Cellular
localization
[131009]of the Huntington's disease protein and discrimination of
the normal
[131010]and mutated form. Nature Genet. 10: 104–110, 1995.
[131011]
[131012]269. Tyler, A.; Quarrell, O. W. J.; Lazarou, L. P.; Meredith,
A. L.;
[131013]Harper, P. S.: Exclusion testing in pregnancy for Hunting-
ton's disease. J.
[131014]Med. Genet. 27: 488–495, 1990.
[131015]
[131016]270. van Dellen, A.; Blakemore, C.; Deacon, R.; York, D.;
Hannan,
[131017]A. J.: Delaying the onset of Huntington's in mice: this un-
remitting
[131018]disease develops later in animals stimulated by their envi-
ronment.
[131019](Letter) Nature 404: 721–722, 2000.

[131020]

[131021]271. van der Weiden, R. M. F.: George Huntington and
George Sumner

[131022]Huntington: a tale of two doctors. (Letter) Hist. Phil. Life
Sci. 11:

[131023]297–304, 1989.

[131024]

[131025]272. van Dijk, J. G.; van der Velde, E. A.; Roos, R. A. C.;
Bruyn,

[131026]G. W.: Juvenile Huntington disease. Hum. Genet. 73:
235–239, 1986.

[131027]

[131028]273. Vessie, P. R.: Original article on the transmission of
Huntington's

[131029]chorea for 300 years--the Bures family group. J. Nerv.
Ment. Dis. 76:

[131030]553–573, 1932.

[131031]

[131032]274. Volkers, W. S.; Went, L. N.; Vegter–van der Vlis, M.;
Harper,

[131033]P. S.; Caro, A.: Genetic linkage studies in Huntington's

chorea. Ann.

[131034]Hum. Genet. 44: 75–79, 1980.

[131035]

[131036]275. Walker, D. A.; Harper, P. S.; Wells, C. E. C.; Tyler, A.;
Davies,

[131037]K.; Newcombe, R. G.: Huntington's chorea in South Wales:
a genetic

[131038]and epidemiological study. Clin. Genet. 19: 213–221,
1981.

[131039]

[131040]276. Wallace, D. C.; Hall, A. C.: Evidence of genetic het-
erogeneity

[131041]in Huntington's chorea. J. Neurol. Neurosurg. Psychiat. 35:
789–800,

[131042]1972.

[131043]

[131044]277. Wang, H. S.; Greenberg, C. R.; Hewitt, J.; Kalousek,
D.; Hayden,

[131045]M. R.: Subregional assignment of the linked marker G8
(D4S10) for

[131046]Huntington disease to chromosome 4p16.1–16.3. Am. J.

Hum. Genet. 39:

[131047]392-396, 1986.

[131048]

[131049]278. Wang, H. S.; Greenberg, C. R.; Kalousek, D.; Gusella,
J.; Horsman,

[131050]D.; Hayden, M. R.: Subregional assignment of the linked
marker D4S10

[131051](G8) for Huntington disease by in situ hybridization.
(Abstract) Cytogenet.

[131052]Cell Genet. 40: 772, 1985.

[131053]

[131054]279. Warner, J.; Barron, L.; St Clair, D.; Brock, D.: Reliabil-
ity

[131055]of clinical diagnosis of Huntington's disease. (Letter) J.
Neurol.

[131056]Neurosurg. Psychiat. 63: 1277, 1994.

[131057]

[131058]280. Wasmuth, J. J.; Hewitt, J.; Smith, B.; Allard, D.; Haines,
J.

[131059]L.; Skarecky, D.; Partlow, E.; Hayden, M. R.: A highly poly-
morphic

[131060] locus very tightly linked to the Huntington's disease gene.

Nature 332:

[131061] 734–736, 1988.

[131062]

[131063] 281. Went, L. N.; Vegter–van der Vlis, M.; Bruyn, G. W.:

Parental

[131064] transmission in Huntington's disease. Lancet I:

1100–1102, 1984.

[131065]

[131066] 282. Wexler, N. S.: The Tiresias complex: Huntington's disease as

[131067] a paradigm of testing for late–onset disorders. FASEB J. 6: 2820–2825,

[131068] 1992.

[131069]

[131070] 283. Wexler, N. S.; Bonilla, E.; Young, A. B.; Shoulson, I.; Gomez,

[131071] F.; Starosta, S.; Travers, H.; Villalobas, M.; de Quiroz, I.; Erbe,

[131072] R.; Penney, J. B.; Uzzell, R. S.; Burnham, F. A.; Daugherty, L.; Jones,

- [131073]B.; Mapstone, C.; Rivas, M.; Messer, E.; Wexler, A.; Snodgrass, R.;
- [131074]Rosenzweig, G.; Esteves, J.; Marsol, N.; Bailey, S.; Brinley, F. J.;
- [131075]Goldstein, E.; Greene, A. E.; Kidd, J. R.; Kidd, K. K.; Gusella, J.
- [131076]F.; Conneally, P. M.; Moreno, H.: Huntington's disease in Venezuela
- [131077]and gene linkage. (Abstract) Cytogenet. Cell Genet. 37: 605, 1984.
- [131078]
- [131079]284. Wexler, N. S.; Young, A.; Tanzi, R.; Starosta, S.; Gomez, F.;
- [131080]Travers, H.; Snodgrass, S. R.; Moreno, H.; Shoulson, I.; Penney, J.;
- [131081]Conneally, P. M.; Gusella, J.: Huntington's disease homozygotes identified.
- [131082](Abstract) Am. J. Hum. Genet. 37: A82, 1985.
- [131083]
- [131084]285. Wexler, N. S.; Young, A. B.; Tanzi, R. E.; Travers, H.; Starosta-Rubinstein,
- [131085]S.; Penney, J. B.; Snodgrass, S. R.; Shoulson, I.; Gomez, F.;

Ramos

[131086]Arroyo, M. A.; Penchaszadeh, G. K.; Moreno, H.; Gibbons, K.; Faryniarz,

[131087]A.; Hobbs, W.; Anderson, M. A.; Bonilla, E.; Conneally, P. M.; Gusella,

[131088]J. F.: Homozygotes for Huntington's disease. *Nature* 326: 194–197,

[131089]1987.

[131090]

[131091]286. Wheeler, V. C.; Gutekunst, C.–A.; Vrbanc, V.; Lebel, L.–A.;

[131092]Schilling, G.; Hersch, S.; Friedlander, R. M.; Gusella, J. F.; Vonsattel,

[131093]J.–P.; Borchelt, D. R.; MacDonald, M. E.: Early phenotypes that presage

[131094]late-onset neurodegenerative disease allow testing of modifiers in

[131095]Hdh CAG knock-in mice. *Hum. Molec. Genet.* 11: 633–640, 2002.

[131096]

[131097]287. Wheeler, V. C.; White, J. K.; Gutekunst, C.–A.; Vrbanc, V.;

- [131098]Weaver, M.; Li, X. J.; Li, S.-H.; Yi, H.; Vonsattel, J.-P.;
Gusella,
[131099]J. F.; Hersch, S.; Auerbach, W.; Joyner, A. L.; MacDonald,
M. E.:
[131100]Long glutamine tracts cause nuclear localization of a novel
form of
[131101]huntingtin in medium spiny striatal neurons in Hdh-Q92
and Hdh-Q111
[131102]knock-in mice. Hum. Molec. Genet. 9: 503-513, 2000.
[131103]
[131104]288. Wiggins, S.; Whyte, P.; Huggins, M.; Adam, S.; Theil-
mann, J.;
[131105]Bloch, M.; Sheps, S. B.; Schechter, M. T.; Hayden, M. R.:
The psychological
[131106]consequences of predictive testing for Huntington's dis-
ease. New
[131107]Eng. J. Med. 327: 1401-1405, 1992.
[131108]
[131109]289. Wolff, G.; Deuschl, G.; Wienker, T. F.; Hummel, K.;
Bender, K.;
[131110]Lucking, C. H.; Schumacher, M.; Hammer, J.; Oepen, G.:
New mutation

[131111]to Huntington's disease. J. Med. Genet. 26: 18-27, 1989.

[131112]

[131113]290. Wright, H. H.; Still, C. N.; Abramson, R. K.: Hunting-
ton's disease

[131114]in black kindreds in South Carolina. Arch. Neurol. 38:
412-414,

[131115]1981.

[131116]

[131117]291. Xiang, F.; Almqvist, E. W.; Huq, M.; Lundin, A.; Hay-
den, M. R.;

[131118]Edstrom, L.; Anvret, M.; Zhang, Z.: A Huntington disease-
like neurodegenerative

[131119]disorder maps to chromosome 20p. Am. J. Hum. Genet.
63: 1431-1438,

[131120]1998.

[131121]

[131122]292. Yamamoto, A.; Lucas, J. J.; Hen, R.: Reversal of neu-
ropathology

[131123]and motor dysfunction in a conditional model of Hunting-
ton's disease. Cell 101:

[131124]57-66, 2000.

[131125]

[131126]293. Youngman, S.; Sarfarazi, M.; Quarrell, O. W. J.; Conneally, P.

[131127]M.; Gibbons, K.; Harper, P. S.; Shaw, D. J.; Tanzi, R. E.; Wallace,

[131128]M. R.; Gusella, J. F.: Studies of a DNA marker (G8) genetically linked

[131129]to Huntington disease in British families. Hum. Genet. 73: 333-339,

[131130]1986.

[131131]

[131132]294. Zabel, B. U.; Naylor, S. L.; Sakaguchi, A. Y.; Gusella, J. F.

[131133]: Regional localization of a DNA polymorphism (D4S10) linked to Huntington's

[131134]disease at 4p16-p15. (Abstract) Cytogenet. Cell Genet. 40: 787,

[131135]1985.

[131136]

[131137]295. Zabel, B. U.; Naylor, S. L.; Sakaguchi, A. Y.; Gusella, J. F.

[131138]: Mapping of the DNA locus D4S10 and the linked Huntington's disease

[131139] gene to 4p16-p15. Cytogenet. Cell Genet. 42: 187-190, 1986.

[131140]

[131141] 296. Zeitlin, S.; Liu, J.-P.; Chapman, D. L.; Papaioannou, V. E.;

[131142] Efstratiadis, A.: Increased apoptosis and early embryonic lethality

[131143] in mice nullizygous for the Huntington's disease gene homologue. Nature

[131144] Genet. 11: 155-163, 1995.

[131145]

[131146] 297. Zlotogora, J.: Dominance and homozygosity. Am. J. Med. Genet. 68:

[131147] 412-416, 1997.

[131148]

[131149] 298. Zuccato, C.; Ciammola, A.; Rigamonti, D.; Leavitt, B. R.; Goffredo,

[131150] D.; Conti, L.; MacDonald, M. E.; Friedlander, R. M.; Silani, V.; Hayden,

[131151]M. R.; Timmusk, T.; Sipione, S.; Cattaneo, E.: Loss of huntingtin-mediated

[131152]BDNF gene transcription in Huntington's disease. Science 293: 493–498,

[131153]2001.

[131154]

[131155]299. Zuhlke, C.; Riess, O.; Bockel, B.; Lange, H.; Thies, U.: Mitotic

[131156]stability and meiotic variability of the (CAG)_n repeat in the Huntington

[131157]disease gene. Hum. Molec. Genet. 2: 2063–2067, 1993.

[131158]

[131159]300. Zuo, J.; Robbins, C.; Taillon-Miller, P.; Cox, D. R.; Myers,

[131160]R. M.: Cloning of the Huntington disease region in yeast artificial

[131161]chromosomes. Hum. Molec. Genet. 1: 149–159, 1992.

[131162]23338.

[131163]Calandro, L. M.; Baer, D. M.; Sensabaugh, G. F.: Characterization

[131164]of a recombinant that locates the hereditary hemochro-

matosis gene

[131165]telomeric to HLA-F. Hum. Genet. 96: 339-342, 1995.

[131166]23339.

[131167]Campbell, K.; Flavin, N.; Ivens, A.; Robert, B.; Buckingham, M.;

[131168]Williamson, R.: The human homeobox gene HOX7 maps to 4p16.1 and is

[131169]deleted in Wolf-Hirschhorn syndrome patients. (Abstract) Am. J. Hum.

[131170]Genet. 45 (suppl.): A179 only, 1989.

[131171]23340.

[131172]Gehring, W. J.; Qian, Y. Q.; Billeter, M.; Furukubo-Tokunaga, K.;

[131173]Schier, A. F.; Resendez-Perez, D.; Affolter, M.; Otting, G.; Wuthrich,

[131174]K.: Homeodomain-DNA recognition. Cell 78: 211-223, 1994.

[131175]23341.

[131176]Greenberg, F.; Elder, F. F. B.; Ledbetter, D. H.; Altherr, M.;

[131177]Wasmuth, J. J.: Molecular confirmation of Wolf-Hirschhorn syndrome

[131178]with apparently normal chromosomes. (Abstract) Am. J.
Hum. Genet. 45

[131179](suppl.): A76 only, 1989.

[131180]23342.

[131181]Hewitt, J. E.; Clark, L. N.; Ivens, A.; Williamson, R.: Struc-
ture

[131182]and sequence of the human homeobox gene HOX7. Ge-
nomics 11: 670-678,

[131183]1991.

[131184]23343.

[131185]Hirschhorn, K.; Cooper, H. L.; Firschein, I. L.: Deletion of
short

[131186]arms of chromosome 4-5 in a child with defects of mid-
line fusion. Humangenetik 1:

[131187]479-482, 1965.

[131188]23344.

[131189]Hu, G.; Vastardis, H.; Bendall, A. J.; Wang, Z.; Logan, M.;
Zhang,

[131190]H.; Nelson, C.; Stein, S.; Greenfield, N.; Seidman, C. E.;
Seidman,

[131191]J. G.; Abate-Shen, C.: Haploinsufficiency of MSX1: a mech-

anism for

[131192]selective tooth agenesis. *Molec. Cell. Biol.* 18: 6044–6051, 1998.

[131193]23345.

[131194]Hwang, S.-J.; Beaty, T. H.; McIntosh, I.; Hefferon, T.; Panny,

[131195]S. R.: Association between homeobox-containing gene MSX1 and the

[131196]occurrence of limb deficiency. *Am. J. Med. Genet.* 75: 419–423, 1998.

[131197]23346.

[131198]Ivens, A.; Flavin, N.; Williamson, R.; Dixon, M.; Bates, G.; Buckingham,

[131199]M.; Robert, B.: The human homeobox gene HOX7 maps to chromosome 4p16.1

[131200]and may be implicated in Wolf–Hirschhorn syndrome. *Hum. Genet.* 84:

[131201]473–476, 1990.

[131202]23347.

[131203]Jumlongras, D.; Bei, M.; Stimson, J. M.; Wang, W.-F.; De-Palma,

[131204]S. R.; Seidman, C. E.; Felbor, U.; Maas, R.; Seidman, J. G.;

Olsen,

[131205]B. R.: A nonsense mutation in MSX1 causes Witkop syndrome. Am. J.

[131206]Hum. Genet. 69: 67–74, 2001.

[131207]23348.

[131208]Odelberg, S. J.; Kollhoff, A.; Keating, M. T.: Dedifferentiation

[131209]of mammalian myotubes induced by msx1. Cell 103: 1099–1109, 2000.

[131210]23349.

[131211]Robert, B.; Sassoon, D.; Jacq, B.; Gehring, W.; Buckingham, M.

[131212]: Hox-7, a mouse homeobox gene with a novel pattern of expression

[131213]during embryogenesis. EMBO J. 8: 91–100, 1989.

[131214]23350.

[131215]Satokata, I.; Maas, R.: Msx1 deficient mice exhibited cleft palate

[131216]and abnormalities of craniofacial and tooth development. Nature Genet. 6:

[131217]348–355, 1994.

[131218]23351.

[131219]Scarel, R. M.; Trevilatto, P. C.; Di Hipolito, O., Jr.; Ca-
margo,

[131220]L. E. A.; Line, S. R. P.: Absence of mutations in the home-
odomain

[131221]of the MSX1 gene in patients with hypodontia. Am. J. Med.
Genet. 92:

[131222]346–349, 2000.

[131223]23352.

[131224]Stimson, J. M.; Sivers, J. E.; Hlava, G. L.: Features of
oligodontia

[131225]in three generations. J. Clin. Pediat. Dent. 21: 269–275,
1997.

[131226]23353.

[131227]van den Boogaard, M.–J. H.; Dorland, M.; Beemer, F. A.;
Ploos

[131228]van Amstel, H. K.: MSX1 mutation is associated with oro-
facial clefting

[131229]and tooth agenesis in humans. (Letter) Nature Genet. 24:
342–343,

[131230]2000. Note: Erratum: Nature Genet. 25: 125 only, 2000.

[131231]23354.

[131232]Vastardis, H.; Karimbux, N.; Guthua, S. W.; Seidman, J. G.;
Seidman,

[131233]C. E.: A human MSX1 homeodomain missense mutation
causes evidence

[131234]tooth agenesis. Nature Genet. 13: 417–421, 1996.

[131235]23355.

[131236]Wolf, U.; Reinwein, H.; Porsch, R.; Schroter, R.; Baitsch, H.

[131237]: Defizienz an den kurzen Armen eines Chromosoms Nr.
4. Humangenetik 1:

[131238]397–413, 1965.

[131239]23356.

[131240]Del Campo, M.; Jones, M. C.; Veraksa, A. N.; Curry, C. J.;
Jones,

[131241]K. L.; Mascarello, J. T.; Ali-Kahn-Catts, Z.; Drumheller, T.;
McGinnis,

[131242]W.: Monodactylous limbs and abnormal genitalia are asso-
ciated with

[131243]hemizygosity for the human 2q31 region that includes the
HOXD cluster. Am.

[131244]J. Hum. Genet. 65: 104–110, 1999.

[131245]23357.

[131246]Zakany, J.; Kmita, M.; Alarcon, P.; de la Pompa, J.-L.;
Duboule,

[131247]D.: Localized and transient transcription of Hox genes
suggests a

[131248]link between patterning and the segmentation clock. Cell
106: 207–217,

[131249]2001.

[131250]23358.

[131251]Lafuse, W. P.; Zwilling, B. S.: Localization of the inhibin
beta-B

[131252]gene on mouse chromosome 1. Mammalian Genome 4:
399–400, 1993.

[131253]23359.

[131254]Salmenkivi, K.; Arola, J.; Voutilainen, R.; Ilvesmaki, V.;
Haglund,

[131255]C.; Kahri, A. I.; Heikkila, P.; Liu, J.: Inhibin/activin beta-
B-subunit

[131256]expression in pheochromocytomas favors benign diagnosis. J. Clin.

[131257]Endocr. Metab. 86: 2231–2235, 2001.

[131258]23360.

[131259]Schrewe, H.; Gendron-Maguire, M.; Harbison, M. L.; Gridley, T.

[131260]: Mice homozygous for a null mutation of activin beta B are viable

[131261]and fertile. Mech. Dev. 47: 43-51, 1994.

[131262]23361.

[131263]Kotanides, H.; Reich, N. C.: Interleukin-4-induced STAT6 recognizes

[131264]and activates a target site in the promoter of the interleukin-4 receptor

[131265]gene. J. Biol. Chem. 271: 25555-25561, 1996.

[131266]23362.

[131267]Caggana, M.; Walker, K.; Reilly, A. A.; Conroy, J. M.; Duva, S.;

[131268]Walsh, A. C.: Population-based studies reveal differences in the

[131269]allelic frequencies of two functionally significant human interleukin-4

[131270]receptor polymorphisms in several ethnic groups. Genet. Med. 1:

[131271]267-271, 1999.

[131272]23363.

[131273]Deichmann, K. A.; Heinzmann, A.; Forster, J.; Dischinger, S.; Mehl,

[131274]C.; Brueggelnte, E.; Hildebrandt, F.; Moseler, M.; Kuehr, J.: Linkage

[131275]and allelic association of atopy and markers flanking the IL4-receptor

[131276]gene. Clin. Exp. Allergy 28: 151-155, 1998.

[131277]23364.

[131278]MacLeod, J. N.; Lee, A. K.; Liebhaber, S. A.; Cooke, N. E.: Developmental

[131279]control and alternative splicing of the placentally expressed transcripts

[131280]from the human growth hormone gene cluster. J. Biol. Chem. 267:

[131281]14219-14226, 1992.

[131282]23365.

[131283]Wurzel, J. M.; Parks, J. S.; Herd, J. E.; Nielsen, P. V.: A gene

[131284]deletion is responsible for absence of human chorionic somatomammotropin. DNA 1:

[131285]251-257, 1982.

[131286]23366.

[131287]Gallin, J. I.: Personal Communication. Bethesda, Md.
10/12/1990.

[131288]23367.

[131289]McCombs, J. L.; Teng, C. T.; Pentecost, B. T.; Magnuson, V.
L.;

[131290]Moore, C. M.; McGill, J. R.: Chromosomal localization of
human lactotransferrin

[131291]gene (LTF) by in situ hybridization. Cytogenet. Cell Genet.
47:

[131292]16-17, 1988.

[131293]23368.

[131294]Moriuchi, M.; Moriuchi, H.: A milk protein lactoferrin en-
hances

[131295]human T cell leukemia virus type I and suppresses HIV-1
infection. J.

[131296]Immun. 166: 4231-4236, 2001.

[131297]23369.

[131298]Naylor, S. L.; Marshall, A.; Solomon, A.; McGill, J. R.; Mc-
Combs,

[131299]J.; Magnuson, V. L.; Moore, C. M.; Lalley, P. A.; Pentecost, B. T.;

[131300]Teng, C.: Lactoferrin maps to human chromosome 3(q21-q23) and mouse

[131301]chromosome 9. (Abstract) Cytogenet. Cell Genet. 46: 669 only, 1987.

[131302]23370.

[131303]Powell, M. J.; Ogden, J. E.: Nucleotide sequence of human lactoferrin

[131304]cDNA. Nucleic Acids Res. 18: 4013 only, 1990.

[131305]23371.

[131306]Qiu, J.; Hendrixson, D. R.; Baker, E. N.; Murphy, T. F.; St. Geme,

[131307]J. W., III; Plaut, A. G.: Human milk lactoferrin inactivates two

[131308]putative colonization factors expressed by Haemophilus influenzae. Proc.

[131309]Nat. Acad. Sci. 95: 12641-12646, 1998.

[131310]23372.

[131311]Rado, T. A.; Wei, X.; Benz, E. J., Jr.: Isolation of lactoferrin

[131312]cDNA from a human myeloid library and expression of

mRNA during normal

[131313]and leukemic myelopoiesis. Blood 70: 989–993, 1987.

[131314]23373.

[131315]Singh, P. K.; Parsek, M. R.; Greenberg, E. P.; Welsh, M. J.: A

[131316]component of innate immunity prevents bacterial biofilm
development. Nature 417:

[131317]552–555, 2002.

[131318]23374.

[131319]Teng, C. T.; Pentecost, B. T.; Marshall, A.; Solomon, A.;
Bowman,

[131320]B. H.; Lalley, P. A.; Naylor, S. L.: Assignment of the lacto-
transferrin

[131321]gene to human chromosome 3 and to mouse chromosome
9. Somat. Cell

[131322]Molec. Genet. 13: 689–693, 1987.

[131323]23375.

[131324]Yang, F.; Lum, J.; Baldwin, W. D.; Brune, J. L.; van Bragt, P.;

[131325]Bowman, B. H.: Genetic analysis of human iron binding
glycoproteins.

[131326](Abstract) Am. J. Hum. Genet. 35: 184A only, 1983.

[131327]23376.

[131328]Tanaka, A. R.; Ikeda, Y.; Abe-Dohmae, S.; Arakawa, R.;
Sadanami,

[131329]K.; Kidera, A.; Nakagawa, S.; Nagase, T.; Aoki, R.; Kioka,
N.; Amachi,

[131330]T.; Yokoyama, S.; Ueda, K.: Human ABCA1 contains a large
amino terminal

[131331]extracellular domain homologous to an epitope of Sjo-
gren's syndrome. Biochem.

[131332]Biophys. Res. Commun. 283: 1019-1025, 2001.

[131333]23377.

[131334]Aoki, N.; Suzuki, H.; Ito, K.; Ito, M.: A novel point mutation

[131335]of the FALDH gene in a Japanese family with Sjogren-
Larsson syndrome. J.

[131336]Invest. Derm. 114: 1065-1066, 2000.

[131337]23378.

[131338]Blumel, J.; Watkins, M.; Eggers, G. W. N.: Spastic
quadriplegia

[131339]combined with congenital ichthyosiform erythroderma
and oligophrenia. Am.

[131340]J. Dis. Child. 96: 724-726, 1958.

[131341]23379.

[131342]Chang, C.; Yoshida, A.: Human fatty aldehyde dehydrogenase gene

[131343](ALDH10): organization and tissue-dependent expression. Genomics 40:

[131344]80–85, 1997.

[131345]23380.

[131346]De Laurenzi, V.; Rogers, G. R.; Hamrock, D. J.; Marekov, L. N.;

[131347]Steinert, P. M.; Compton, J. G.; Markova, N.; Rizzo, W. B.: Sjogren–Larsson

[131348]syndrome is caused by mutations in the fatty aldehyde dehydrogenase

[131349]gene. Nature Genet. 12: 52–57, 1996.

[131350]23381.

[131351]De Laurenzi, V.; Rogers, G. R.; Tarcsa, E.; Carney, G.; Marekov,

[131352]L.; Bale, S. J.; Compton, J. G.; Markova, N.; Steinert, P. M.; Rizzo,

[131353]W. B.: Sjogren–Larsson syndrome is caused by a common mutation in

[131354]northern European and Swedish patients. J. Invest. Derm.
109: 79–83,

[131355]1997.

[131356]23382.

[131357]Gedde–Dahl, T., Jr.; Rajka, G.; Larsen, T. E.; Jellum, E.: Au-
tosomal

[131358]recessive ichthyosis in Norway: II. Sjogren–Larsson–like
ichthyosis

[131359]without CNS or eye involvement. Clin. Genet. 26:
242–244, 1984.

[131360]23383.

[131361]Gilbert, W. R., Jr.; Smith, J. L.; Nyhan, W. L.: The Sjogren–
Larsson

[131362]syndrome. Arch. Ophthal. 80: 308–316, 1968.

[131363]23384.

[131364]Goldsmith, L. A.; Baden, H. P.; Canty, T. G.: Sjogren–Lars-
son

[131365]syndrome. Acta Derm. Venerol. 51: 374–378, 1971.

[131366]23385.

[131367]Gustavson, K. H.; Jagell, S.: Dermatoglyphic patterns in the
Sjogren–Larsson

[131368]syndrome. Clin. Genet. 17: 120–124, 1980.

[131369]23386.

[131370]Heijer, A.; Reed, W. B.: Sjogren–Larsson syndrome: congenital

[131371]ichthyosis, spastic paralysis, and oligophrenia. Arch. Derm. 92:

[131372]545–552, 1965.

[131373]23387.

[131374]Jl, L.; Oostheim, W.; van Werkhoven, M.; Willemsen, M. A. A.

[131375]P.; Wanders, R. J. A.: Molecular basis of Sjogren–Larsson syndrome:

[131376]frequency of the 1297–1298 del GA and 943C–T mutation in 29 patients. J.

[131377]Inherit. Metab. Dis. 22: 319–321, 1999.

[131378]23388.

[131379]Jagell, S.; Gustavson, K.–H.; Holmgren, G.: Sjogren–Larsson syndrome

[131380]in Sweden: a clinical, genetic and epidemiological study. Clin. Genet. 19:

[131381]233–256, 1981.

[131382]23389.

[131383]Jagell, S.; Linden, S.: Ichthyosis in the Sjogren-Larsson syndrome. Clin.

[131384]Genet. 21: 243-252, 1982.

[131385]23390.

[131386]Jagell, S.; Polland, W.; Sandgren, O.: Specific changes in the

[131387]fundus typical for the Sjogren-Larsson syndrome: an ophthalmological

[131388]study of 35 patients. Acta Ophthal. 58: 321-330, 1980.

[131389]23391.

[131390]Shohat, M.; Bu, X.; Shohat, T.; Fischel-Ghodsian, N.; Magal, N.;

[131391]Nakamura, Y.; Schwabe, A. D.; Schlezinger, M.; Danon, Y.; Rotter,

[131392]J. I.: The gene for familial Mediterranean fever in both Armenians

[131393]and non-Ashkenazi Jews is linked to the alpha-globin complex on 16p:

[131394]evidence for locus homogeneity. Am. J. Hum. Genet. 51: 1349-1354,

[131395]1992.

[131396]23392.

[131397]Shohat, M.; Korenberg, J. R.; Schlesinger, M.; Shohat, T.;
Petersen,

[131398]G. M.; Schwabe, A.; Rotter, J. I.: The gene for serum amy-
loid P component

[131399](APCS) is not a genetic marker for amyloidosis in familial
Mediterranean

[131400]fever. (Abstract) Cytogenet. Cell Genet. 51: 1079, 1989.

[131401]23393.

[131402]Shohat, M.; Korenberg, J. R.; Schwabe, A. D.; Rotter, J. I.:

[131403]Hypothesis: familial Mediterranean fever--a genetic disorder of the

[131404]lipocortin family? Am. J. Med. Genet. 34: 163-167, 1989.

[131405]23394.

[131406]Shohat, M.; Korenberg, J. R.; Shohat, T.; Sack, G.; Schwabe,
A.;

[131407]Rotter, J. I.: Serum amyloid A (SAA) locus is not associated
with

[131408]familial Mediterranean fever (FMF) or FMF-amyloidosis.
(Abstract) Cytogenet.

[131409]Cell Genet. 51: 1079, 1989.

[131410]23395.

[131411]Shohat, M.; Livneh, A.; Zemer, D.; Pras, M.; Sohar, E.: Twin

[131412]studies in familial Mediterranean fever. Am. J. Med. Genet.
44:

[131413]179–182, 1992.

[131414]23396.

[131415]Shohat, M.; Shohat, T.; Rotter, J. I.; Schlesinger, M.; Pe-
tersen,

[131416]G. M.; Pribyl, T.; Sack, G.; Schwabe, A. D.; Korenberg, J. R.:
Serum

[131417]amyloid A and P protein genes in familial Mediterranean
fever. Genomics 8:

[131418]83–89, 1990.

[131419]

[131420]100. Shohat, T.; Shohat, M.; Petersen, G. M.; Sparkes, R.
S.; Langfield,

[131421]D.; Bickal, J.; Korenberg, J. R.; Schwabe, A. D.; Rotter, J. I.:

[131422]Genetic marker family studies in familial Mediterranean
fever (FMF)

[131423]in Armenians. Clin. Genet. 38: 332–339, 1990.

[131424]

[131425]101. Shohat, T.; Shohat, M.; Tyan, D. B.; Wang, S. J.;
Sparkes, R.

[131426]S.; Schwabe, A. D.; Rotter, J. I.: Familial Mediterranean
fever--linkage

[131427]studies with genetic markers on chromosome 6. Tissue
Antigens 36:

[131428]103–107, 1990.

[131429]

[131430]102. Siegal, S.: Benign paroxysmal peritonitis. Ann. Intern.
Med. 23:

[131431]1–22, 1945.

[131432]

[131433]103. Siegal, S.: Familial paroxysmal polyserositis: analysis
of fifty

[131434]cases. Am. J. Med. 36: 893–918, 1964.

[131435]

[131436]104. Sohar, E.; Gafni, J.; Pras, M.; Heller, H.: Familial
Mediterranean

[131437]fever: a survey of 470 cases and review of literature. Am.
J. Med. 43:

[131438]227-253, 1967.

[131439]

[131440]105. Sokmen, C.: Comment. Trans. Assoc. Am. Phys. 72:
120-121,

[131441]1959.

[131442]

[131443]106. Sood, R.; Blake, T.; Aksentijevich, I.; Wood, G.; Chen,
X.; Gardner,

[131444]D.; Shelton, D. A.; Mangelsdorf, M.; Orsborn, A.; Pras, E.;
Balow,

[131445]J. E. Jr.; Centola, M. and 16 others: Construction of a 1-Mb
restriction-mapped

[131446]cosmid contig containing the candidate region for the fa-
milial Mediterranean

[131447]fever locus (MEFV) on chromosome 16p13.3. Genomics
42: 83-95, 1997.

[131448]

[131449]107. Tamir, N.; Langevitz, P.; Zemer, D.; Pras, E.; Shinar,
Y.; Padeh,

[131450]S.; Zaks, N.; Pras, M.; Livneh, A.: Late-onset familial
Mediterranean

[131451]fever (FMF): a subset with distinct clinical, demographic,
and molecular

[131452]genetic characteristics. Am. J. Med. Genet. 87: 30–35,
1999.

[131453]

[131454]108. Touitou, I.: The spectrum of familial Mediterranean
fever (FMF)

[131455]mutations. Europ. J. Hum. Genet. 9: 473–483, 2001.

[131456]

[131457]109. Tutar, H. E.; Imamoglu, A.; Atalay, S.: Recurrent peri-
carditis

[131458]in familial Mediterranean fever. (Letter) Acta Pediat. 88:
1045–1046,

[131459]1999.

[131460]

[131461]110. Wolff, S. M.; Dinarello, C. A.; Dale, D. C.; Goldfinger,
S. E.;

[131462]Alling, D. W.: Colchicine therapy of familial Mediterranean
fever.

[131463](Abstract) Clin. Res. 22: 567A, 1974.

[131464]

[131465]111. Yazigi, A.; Abou-Charaf, L. C.: Colchicine for recurrent pericarditis

[131466]in children. Acta Paediat. 87: 603-604, 1998.

[131467]

[131468]112. Yilmaz, E.; Ozen, S.; Balci, B.; Duzova, A.; Topaloglu, R.; Besbas,

[131469]N.; Saatci, U.; Bakkaloglu, A.; Ozguc, M.: Mutation frequency of

[131470]familial Mediterranean fever and evidence for a high carrier rate

[131471]in the Turkish population. *Europ. J. Hum. Genet.* 9: 553–555, 2001.

[131472]

[131473]113. Yuval, Y.; Hemo-Zisser, M.; Zemer, D.; Sohar, E.; Pras, M.:

[131474]Dominant inheritance in two families with familial Mediterranean fever

[131475](FMF). *Am. J. Med. Genet.* 57: 455–457, 1995.

[131476]

[131477]114. Zemer, D.; Livneh, A.; Pras, M.; Sohar, E.: Familial Mediterranean

[131478]fever in the colchicine era: the fate of one family. *Am. J. Med.*

[131479]*Genet.* 45: 340–344, 1993.

[131480]

[131481]115. Zemer, D.; Pras, M.; Sohar, E.; Modan, M.; Cabili, S.;
Gafni,

[131482]J.: Colchicine in the prevention and treatment of the amy-
loidosis

[131483]of familial Mediterranean fever. New Eng. J. Med. 314:
1001-1005,

[131484]1986.

[131485]

[131486]116. Zemer, D.; Revach, M.; Pras, M.; Modan, B.; Schor, S.;
Sohar,

[131487]E.; Gafni, J.: A controlled trial of colchicine in preventing
attacks

[131488]of familial Mediterranean fever. New Eng. J. Med. 291:
932-934,

[131489]1974.

[131490]23397.

[131491]Labay, V.; Raz, T.; Baron, D.; Mandel, H.; Williams, H.; Bar-
rett,

[131492]T.; Szargel, R.; McDonald, L.; Shalata, A.; Nosaka, K.; Gre-
gory, S.;

[131493]Cohen, N.: Mutations in SLC19A2 cause thiamine-responsive megaloblastic

[131494]anaemia associated with diabetes mellitus and deafness. Nature Genet. 22:

[131495]300–304, 1999.

[131496]23398.

[131497]Millat, G.; Marcais, C.; Rafi, M. A.; Yamamoto, T.; Morris, J.

[131498]A.; Pentchev, P. G.; Ohno, K.; Wenger, D. A.; Vanier, M. T.: Niemann–Pick

[131499]C1 disease: the I1061T substitution is a frequent mutant allele in

[131500]patients of western European descent and correlates with a classic

[131501]juvenile phenotype. Am. J. Hum. Genet. 65: 1321–1329, 1999.

[131502]23399.

[131503]Agostoni, C.; Massetto, N.; Biasucci, G.; Rottoli, A.; Bonvissuto,

[131504]M.; Bruzzese, M.; Giovannini, M.; Riva, E.: Effects of long-chain

[131505]polyunsaturated fatty acid supplementation on fatty acid status and

- [131506]visual function in treated children with hyperphenylalaninemia. J.
- [131507]Pediat. 137: 504–509, 2000.
- [131508]23400.
- [131509]Eisensmith, R. C.; Okano, Y.; Dasovich, M.; Wang, T.; Guttler,
- [131510]F.; Lou, H.; Guldberg, P.; Lichter–Konecki, U.; Konecki, D. S.; Svensson,
- [131511]E.; Hagenfeldt, L.; Rey, F.; Munnich, A.; Lyonnet, S.; Cockburn, F.;
- [131512]Connor, J. M.; Pembrey, M. E.; Smith, I.; Gitzelmann, R.; Steinmann,
- [131513]B.; Apold, J.; Eiken, H. G.; Giovannini, M.; Riva, E.; Longhi, R.;
- [131514]Romano, C.; Cerone, R.; Naughten, E. R.; Mullins, C.; Cahalane, S.;
- [131515]Ozalp, I.; Fekete, G.; Schuler, D.; Berencsi, G. Y.; Nasz, I.; Brdicka,
- [131516]R.; Kamaryt, J.; Pijackova, A.; Cabalska, B.; Boszkowa, K.; Schwartz,
- [131517]E.; Kalinin, V. N.; Jin, L.; Chakraborty, R.; Woo, S. L. C.: Multiple

[131518]origins for phenylketonuria in Europe. Am. J. Hum. Genet.
51: 1355–1365,
[131519]1992.
[131520]23401.
[131521]Kirk, E. P. E.; Fletcher, J. M.; Sharp, P.; Carey, B.; Poulos,
[131522]A.: X-linked adrenoleukodystrophy: the Australasian ex-
perience. Am.
[131523]J. Med. Genet. 76: 420–423, 1998.
[131524]23402.
[131525]Kobayashi, T.; Noda, S.; Umezaki, H.; Goto, I.; Suzuki, S.;
Kitaguchi,
[131526]T.; Kuroiwa, Y.: Familial spinocerebellar degeneration as
an expression
[131527]of adrenoleukodystrophy. J. Neurol. Neurosurg. Psychiat.
49: 1438–1440,
[131528]1986.
[131529]23403.
[131530]Kobayashi, T.; Yamada, T.; Yasutake, T.; Shinnoh, N.;
Goto, I.;
[131531]Iwaki, T.: Adrenoleukodystrophy gene encodes an 80 kDa
membrane protein. Biochem.

[131532]Biophys. Res. Commun. 201: 1029–1034, 1994.

[131533]23404.

[131534]Kok, F.; Neumann, S.; Sarde, C.–O.; Zheng, S.; Wu, K.–H.;
Wei,

[131535]H.–M.; Bergin, J.; Watkins, P. A.; Gould, S.; Sack, G.; Moser,
H.;

[131536]Mandel, J.–L.; Smith, K. D.: Mutational analysis of patients
with

[131537]X-linked adrenoleukodystrophy. Hum. Mutat. 6: 104–115,
1995.

[131538]23405.

[131539]Kolodny, E. H.: The adrenoleukodystrophy–adreno–
myeloneuropathy

[131540]complex: is it treatable? (Editorial) Ann. Neurol. 21:
230–231,

[131541]1987.

[131542]23406.

[131543]Korenke, G. C.; Fuchs, S.; Krasemann, E.; et al.: Cerebral
adrenoleukodystrophy

[131544](ALD) in only one of monozygous twins with an identical
ALD genotype. Ann.

[131545]Neurol. 40: 254–257, 1996.

[131546]23407.

[131547]Krasemann, E. W.; Meier, V.; Korenke, G. C.; Hunneman, D.
H.;

[131548]Hanefeld, F.: Identification of mutations in the ALD–gene
of 20 families

[131549]with adrenoleukodystrophy/adrenomyeloneuropathy.
Hum. Genet. 97:

[131550]194–197, 1996.

[131551]23408.

[131552]Kruse, B.; Barker, P. B.; van Zijl, P. C. M.; Duyn, J. H.; Moo-
nen,

[131553]C. T. W.; Moser, H. W.: Multislice proton magnetic reso-
nance spectroscopic

[131554]imaging in X–linked adrenoleukodystrophy. Ann. Neurol.
36: 595–608,

[131555]1994.

[131556]23409.

[131557]Lachtermacher, M. B. R.; Seuanez, H. N.; Moser, A. B.;
Moser,

[131558]H. W.; Smith, K. D.: Determination of 30 X–linked

adrenoleukodystrophy

[131559] mutations, including 15 not previously described. Hum.

Mutat. 15:

[131560] 348–353, 2000.

[131561] 23410.

[131562] Laureti, S.; Casucci, G.; Santeusano, F.; Angeletti, G.;

Aubourg,

[131563] P.; Brunetti, P.: X-linked adrenoleukodystrophy is a fre-

quent cause

[131564] of idiopathic Addison's disease in young adult male pa-

tients. J.

[131565] Clin. Endocr. Metab. 81: 470–474, 1996.

[131566] 23411.

[131567] Lazo, O.; Contreras, M.; Hashmi, M.; Stanley, W.; Irazu, C.;

Singh,

[131568] J.: Peroxisomal lignoceroyl-CoA ligase deficiency in child-

hood adrenoleukodystrophy

[131569] and adrenomyeloneuropathy. Proc. Nat. Acad. Sci. 85:

7647–7651,

[131570] 1988.

[131571] 23412.

[131572]Ligtenberg, M. J. L.; Kemp, S.; Sarde, C.-O.; van Geel, B.
M.;

[131573]Kleijer, W. J.; Barth, P. G.; Mandel, J.-L.; van Oost, B. A.;
Bolhuis,

[131574]P. A.: Spectrum of mutations in the gene encoding the
adrenoleukodystrophy

[131575]protein. Am. J. Hum. Genet. 56: 44-50, 1995.

[131576]23413.

[131577]Luciani, G. B.; Pessotto, R.; Mazzucco, A.:
Adrenoleukodystrophy

[131578]presenting as postperfusion syndrome. (Letter) New Eng.
J. Med. 336:

[131579]731-732, 1997.

[131580]23414.

[131581]Maestri, N. E.; Beaty, T. H.: Predictions of a 2-locus model

[131582]for disease heterogeneity: application to adrenoleukodys-
trophy. Am.

[131583]J. Med. Genet. 44: 576-582, 1992.

[131584]23415.

[131585]Malm, G.; Ringden, O.; Anvret, M.; von Döbeln, U.; Hagen-
feldt,

[131586]L.; Isberg, B.; Knuutila, S.; Nennesmo, I.; Winiarski, J.; Marcus,

[131587]C.: Treatment of adrenoleukodystrophy with bone marrow transplantation. Acta

[131588]Paediat. 86: 484-492, 1997.

[131589]23416.

[131590]Manz, H. J.; Schuelein, M.; McCullough, D. C.; Kishimoto, Y.;

[131591]Eiben, R. M.: New phenotypic variant of adrenoleukodystrophy: pathologic,

[131592]ultrastructural, and biochemical study in two brothers. J. Neurol.

[131593]Sci. 45: 245-260, 1980.

[131594]23417.

[131595]Martin, J. J.; Dompas, B.; Ceuterick, C.; Jacobs, K.: Adrenomyeloneuropathy

[131596]and adrenoleukodystrophy in two brothers. Europ. Neurol. 19: 281-287,

[131597]1980.

[131598]23418.

[131599]Menkes, J. H.; Corbo, L. M.: Adrenoleukodystrophy: accu-

mutation

[131600]of cholesterol esters with very long chain fatty acids. Neu-
rology 27:

[131601]928-932, 1977.

[131602]23419.

[131603]Migeon, B. R.; Moser, H. W.; Moser, A. B.; Axelman, J.; Sil-
lence,

[131604]D.; Norum, R. A.: Adrenoleukodystrophy: evidence for X
linkage inactivation,

[131605]and selection favoring the mutant allele in heterozygous
cells. Proc.

[131606]Nat. Acad. Sci. 78: 5066-5070, 1981.

[131607]23420.

[131608]Millat, G.; Marcais, C.; Tomasetto, C.; Chikh, K.; Fensom,
A.

[131609]H.; Harzer, K.; Wenger, D. A.; Ohno, K.; Vanier, M. T.: Nie-
mann-Pick

[131610]C1 disease: correlations between NPC1 mutations, levels
of NPC1 protein,

[131611]and phenotypes emphasize the functional significance of
the putative

[131612]sterol-sensing domain and of the cysteine-rich luminal

loop. Am.

[131613]J. Hum. Genet. 68: 1373–1385, 2001.

[131614]23421.

[131615]Morris, J. A.; Zhang, D.; Coleman, K. G.; Nagle, J.;

Pentchev,

[131616]P. G.; Carstea, E. D.: The genomic organization and poly-

morphism

[131617]analysis of the human Niemann–Pick C1 gene. Biochem.

Biophys. Res.

[131618]Commun. 261: 493–498, 1999.

[131619]23422.

[131620]Omura, K.; Suzuki, Y.; Norose, N.; Sato, M.; Maruyama, K.;

Koeda,

[131621]T.: Type C Niemann–Pick disease: clinical and biochemical

studies

[131622]on 6 cases. Brain Dev. 11: 57–61, 1989.

[131623]23423.

[131624]Palmer, M.; Green, W. R.; Maumenee, I. H.; Valle, D. L.;

Singer,

[131625]H. S.; Morton, S. J.; Moser, H. W.: Niemann–Pick disease–

–type C:

[131626]ocular histopathologic and electron microscopic studies.

Arch. Ophthalmol. 103:

[131627]817–822, 1985.

[131628]23424.

[131629]Patel, S. C.; Suresh, S.; Kumar, U.; Hu, C. Y.; Cooney, A.;

Blanchette–Mackie,

[131630]E. J.; Neufeld, E. B.; Patel, R. C.; Brady, R. O.; Patel, Y. C.;

Pentchev,

[131631]P. G.; Ong, W.–Y.: Localization of Niemann–Pick C1 protein

in astrocytes:

[131632]implications for neuronal degeneration in Niemann–Pick

type C disease. Proc.

[131633]Nat. Acad. Sci. 96: 1657–1662, 1999.

[131634]23425.

[131635]Pentchev, P. G.; Boothe, A. D.; Kruth, H. S.; Weintraub, H.;

Stivers,

[131636]J.; Brady, R. O.: A genetic storage disorder in BALB/C mice

with

[131637]a metabolic block in esterification of exogenous chole-

sterol. J.

[131638]Biol. Chem. 259: 5784–5791, 1984.

[131639]23426.

[131640]Sun, X.-H.: Constitutive expression of the Id1 gene impairs mouse

[131641]B cell development. Cell 79: 893–900, 1994.

[131642]23427.

[131643]Nowell, P. C.; Hungerford, D. A.: Chromosome studies on normal

[131644]and leukemic human leukocytes. J. Nat. Cancer Inst. 25: 85–109,

[131645]1960.

[131646]23428.

[131647]Dowbenko, D. J.; Diep, A.; Taylor, B. A.; Lysis, A. J.; Lasky,

[131648]L. A.: Characterization of the murine homing receptor gene reveals

[131649]correspondence between protein domains and coding exons. Genomics 9:

[131650]270–277, 1991.

[131651]23429.

[131652]Aarskog, N. K.; Vedeler, C. A.: Real-time quantitative polymerase

[131653]chain reaction: a new method that detects both the peripheral myelin

[131654]protein 22 duplication in Charcot-Marie-Tooth type 1A disease and

[131655]the peripheral myelin protein 22 deletion in hereditary neuropathy

[131656]with liability to pressure palsies. Hum. Genet. 107: 494-498, 2000.

[131657]23430.

[131658]Chance, P. F.; Alderson, M. K.; Leppig, K. A.; Lensch, M. W.; Matsunami,

[131659]N.; Smith, B.; Swanson, P. D.; Odelberg, S. J.; Distèche, C. M.; Bird,

[131660]T. D.: DNA deletion associated with hereditary neuropathy with liability

[131661]to pressure palsies. Cell 72: 143-151, 1993.

[131662]23431.

[131663]Lehmann, A. R.; Bootsma, D.; Clarkson, S. G.; Cleaver, J. E.;

[131664]McAlpine, P. J.; Tanaka, K.; Thompson, L. H.; Wood, R. D.: Nomenclature

[131665]of human DNA repair genes. Mutat. Res. 315: 41-42,

1994.

[131666]23432.

[131667]Abruzzo, L. V.; Jaffe, E. S.; Cotelingam, J. D.; Whang-Peng, J.;

[131668]Del Duca, V., Jr.; Medeiros, L. J.: T-cell lymphoblastic lymphoma

[131669]with eosinophilia associated with subsequent myeloid malignancy. Am.

[131670]J. Surg. Path. 16: 236-245, 1992.

[131671]23433.

[131672]Chesi, M.; Nardini, E.; Brents, L. A.; Schrock, E.; Ried, T.; Kuehl,

[131673]W. M.; Bergsagel, P. L.: Frequent translocation t(4;14)(p16.3;q32.3)

[131674]in multiple myeloma is associated with increased expression and activating

[131675]mutations of fibroblast growth factor receptor 3. Nature Genet. 16:

[131676]260-264, 1997.

[131677]23434.

[131678]Demiroglu, A.; Steer, E. J.; Heath, C.; Taylor, K.; Bentley,

M.;

[131679] Allen, S. L.; Koduru, P.; Brody, J. P.; Hawson, G.; Rodwell, R.; Doody,

[131680] M.-L.; Carnicero, F.; Reiter, A.; Goldman, J. M.; Melo, J. V.; Cross,

[131681] N. C. P.: The t(8;22) in chronic myeloid leukemia fuses BCR to FGFR1:

[131682] transforming activity and specific inhibition of FGFR1 fusion proteins. Blood 98:

[131683] 3778-3783, 2001.

[131684] 23435.

[131685] Guasch, G.; Mack, G. J.; Popovici, C.; Dastugue, N.; Birnbaum,

[131686] D.; Rattner, J. B.; Pebusque, M.-J.: FGFR1 is fused to the centrosome-associated

[131687] protein CEP110 in the 8p12 stem cell myeloproliferative disorder with

[131688] t(8;9)(p12;q33). Blood 95: 1788-1796, 2000.

[131689] 23436.

[131690] Inhorn, R. C.; Aster, J. C.; Roach, S. A.; Slapak, C. A.; Soifer,

[131691] R.; Tantravahi, R.; Stone, R. M.: A syndrome of lym-

phoblastic lymphoma,
[131692]eosinophilia, and myeloid hyperplasia/malignancy associated with t(8;13)(p11;q11):
[131693]description of a distinctive clinicopathologic entity. Blood
85:
[131694]1881-1887, 1995.
[131695]23437.
[131696]Lee, P. L.; Johnson, D. E.; Cousens, L. S.; Fried, V. A.;
Williams,
[131697]L. T.: Purification and complementary DNA cloning of a
receptor for
[131698]basic fibroblast growth factor. Science 245: 57-60, 1989.
[131699]23438.
[131700]Mavrothalassitis, G.; Tzimagiorgis, G.; Mitsialis, A.; Zannis,
[131701]V.; Plaitakis, A.; Papamatheakis, J.; Moschonas, N.: Isolation and
[131702]characterization of cDNA clones encoding human liver
glutamate dehydrogenase:
[131703]evidence for a small gene family. Proc. Nat. Acad. Sci. 85:
3494-3498,
[131704]1988.

[131705]23439.

[131706]Michaelidis, T. M.; Tzimagiorgis, G.; Moschonas, N. K.; Papamatheakis,

[131707]J.: The human glutamate dehydrogenase gene family:
gene organization

[131708]and structural characterization. Genomics 16: 150–160,
1993.

[131709]23440.

[131710]Miki, Y.; Taki, T.; Ohura, T.; Kato, H.; Yanagisawa, M.;
Hayashi,

[131711]Y.: Novel missense mutations in the glutamate dehydro-
genase gene

[131712]in the congenital hyperinsulinism–hyperammonemia syn-
drome. J. Pediatr. 136:

[131713]69–72, 2000.

[131714]23441.

[131715]Nakatani, Y.; Schneider, M.; Banner, C.; Freese, E.: Com-
plete

[131716]nucleotide sequence of human glutamate dehydrogenase
cDNA. Nucleic

[131717]Acids Res. 16: 6237 only, 1988.

[131718]23442.

[131719]Nelson, R. L.; Povey, M. S.; Hopkinson, D. A.; Harris, H.:
Electrophoresis

[131720]of human L-glutamate dehydrogenase: tissue distribution
and preliminary

[131721]population survey. Biochem. Genet. 15: 87-91, 1977.

[131722]23443.

[131723]Plaitakis, A.; Berl, S.; Yahr, M. D.: Abnormal glutamate
metabolism

[131724]in an adult-onset degenerative neurological disorder. Sci-
ence 216:

[131725]193-196, 1982.

[131726]23444.

[131727]Plaitakis, A.; Berl, S.; Yahr, M. D.: Neurological disorders

[131728]associated with deficiency of glutamate dehydrogenase.
Ann. Neurol. 15:

[131729]144-153, 1984.

[131730]23445.

[131731]Plaitakis, A.; Nicklas, W. J.; Desnick, R. J.: Glutamate dehy-
drogenase

[131732]deficiency in three patients with spinocerebellar syndrome. Ann.

[131733]Neurol. 7: 297–303, 1980.

[131734]23446.

[131735]Santer, R.; Kinner, M.; Passarge, M.; Superti-Furga, A.; Mayatepek,

[131736]E.; Meissner, T.; Schneppenheim, R.; Schaub, J.: Novel missense mutations

[131737]outside the allosteric domain of glutamate dehydrogenase are prevalent

[131738]in European patients with the congenital hyperinsulinism–hyperammonemia

[131739]syndrome. Hum. Genet. 108: 66–71, 2001.

[131740]23447.

[131741]Shaughnessy, J., Jr.; Mock, B.; Duncan, R.; Potter, M.; Banner,

[131742]C.: A restriction fragment length polymorphism at murine Glud locus

[131743] cosegregates with Rib-1, Es-10, and Tcra on chromosome 14. Nucleic

[131744]Acids Res. 17: 2881 only, 1989.

[131745]23448.

[131746]Smith, T. J.; Peterson, P. E.; Schmidt, T.; Fang, J.; Stanley,

[131747]C. A.: Structures of bovine glutamate dehydrogenase
complexes elucidate

[131748]the mechanism of purine regulation. J. Molec. Biol. 307:
707–720,

[131749]2001.

[131750]23449.

[131751]Sorbi, S.; Tonini, S.; Giannini, E.; Piacentini, S.; Marini, P.;

[131752]Amaducci, L.: Abnormal platelet glutamate dehydrogenase
activity

[131753]and activation in dominant and nondominant olivoponto-
cerebellar atrophy. Ann.

[131754]Neurol. 19: 239–245, 1986.

[131755]23450.

[131756]Stanley, C. A.; Lieu, Y.; Hsu, B.; Poncz, M.: Hypoglycemia
in

[131757]infants with hyperinsulinism and hyperammonemia: gain
of function

[131758]mutations in the pathway of leucine-mediated insulin se-
cretion. (Abstract) Diabetes 46

[131759](suppl. 1): 217A only, 1997.

[131760]23451.

[131761]Stanley, C. A.; Lieu, Y. K.; Hsu, B. Y. L.; Burlina, A. B.;
Greenberg,

[131762]C. R.; Hopwood, N. J.; Perlman, K.; Rich, B. H.; Zammarchi,
E.; Poncz,

[131763]M.: Hyperinsulinism and hyperammonemia in infants with
regulatory

[131764]mutations of the glutamate dehydrogenase gene. New
Eng. J. Med. 338:

[131765]1352–1357, 1998.

[131766]23452.

[131767]Tzimagiorgis, G.; Adamson, M. C.; Kozak, C. A.;
Moschonas, N.

[131768]K.: Chromosomal mapping of glutamate dehydrogenase
gene sequences

[131769]to mouse chromosomes 7 and 14. Genomics 10: 83–88,
1991.

[131770]23453.

[131771]Tzimagiorgis, G.; Leversha, M. A.; Chroniary, K.; Gouliel-
mos,

[131772]G.; Sargent, C. A.; Ferguson-Smith, M.; Moschonas, N. K.:

Structure

[131773]and expression analysis of a member of the human glutamate dehydrogenase

[131774](GLUD) gene family mapped to chromosome 10p11.2.

Hum. Genet. 91:

[131775]433-438, 1993.

[131776]23454.

[131777]Yamaguchi, T.; Hayashi, K.; Murakami, H.; Ota, K.;

Maruyama, S.

[131778]: Glutamate dehydrogenase deficiency in spinocerebellar degeneration. Neurochem.

[131779]Res. 7: 627-636, 1982.

[131780]23455.

[131781]Agus, D. B.; Gambhir, S. S.; Pardridge, W. M.; Spielholz, C.;

Baselga,

[131782]J.; Vera, J. C.; Golde, D. W.: Vitamin C crosses the blood-brain

[131783]barrier in the oxidized form through the glucose transporters. J.

[131784]Clin. Invest. 100: 2842-2848, 1997.

[131785]23456.

[131786]Ardinger, R. H., Jr.; Buetow, K. H.; Weiss, M. J.; Nemer, M.;
DeHaven,

[131787]C. R.; Murray, J. C.: Multipoint linkage relationships of 6
loci

[131788]on 1p (ALPL, GLUT, PGD, PGM1, PND, RH). (Abstract) Am.
J. Hum. Genet. 41:

[131789]A154 only, 1987.

[131790]23457.

[131791]Baroni, M. G.; Oelbaum, R. S.; Pozzilli, P.; Stocks, J.; Li,
S.-R.;

[131792]Fiore, V.; Galton, D. J.: Polymorphisms at the GLUT1
(HepG2) and

[131793]GLUT4 (muscle/adipocyte) glucose transporter genes and
non-insulin-dependent

[131794]diabetes mellitus (NIDDM). Hum. Genet. 88: 557-561,
1992.

[131795]23458.

[131796]Brockmann, K.; Wang, D.; Korenke, C. G.; von Moers, A.;
Ho, Y.-Y.;

[131797]Pascual, J. M.; Kuang, K.; Yang, H.; Ma, L.; Kranz-Eble, P.;
Fischbarg,

[131798]J.; Hanefeld, F.; De Vivo, D. C.: Autosomal dominant Glut-1 deficiency

[131799]syndrome and familial epilepsy. Ann. Neurol. 50: 476-485, 2001.

[131800]23459.

[131801]Klepper, J.; Willemsen, M.; Verrips, A.; Guertsen, E.; Hermann,

[131802]R.; Kutzick, C.; Florcken, A.; Voit, T.: Autosomal dominant transmission

[131803]of GLUT1 deficiency. Hum. Molec. Genet. 10: 63-68, 2001.

[131804]23460.

[131805]Lazar, V.; Bidart, J.-M.; Caillou, B.; Mahe, C.; Lacroix, L.; Filetti,

[131806]S.; Schlumberger, M.: Expression of the Na(+)/I(-) symporter gene

[131807]in human thyroid tumors: a comparison study with other thyroid-specific

[131808]genes. J. Clin. Endocr. Metab. 84: 3228-3234, 1999.

[131809]23461.

[131810]Li, S. R.; Baroni, M. G.; Oelbaum, R. S.; Stock, J.; Galton, D.

[131811]J.: Association of genetic variant of the glucose transporter with

[131812]non-insulin-dependent diabetes mellitus. Lancet II: 368–370, 1988.

[131813]23462.

[131814]Mueckler, M.; Caruso, C.; Baldwin, S. A.; Panico, M.; Blench, I.;

[131815]Morris, H. R.; Allard, W. J.; Lienhard, G. E.; Lodish, H. F.: Sequence

[131816]and structure of a human glucose transporter. Science 229: 941–945,

[131817]1985.

[131818]23463.

[131819]Sarkar, H. K.; Thorens, B.; Lodish, H. F.; Kaback, H. R.: Expression

[131820]of the human erythrocyte glucose transporter in Escherichia coli. Proc.

[131821]Nat. Acad. Sci. 85: 5463–5467, 1988.

[131822]23464.

[131823]Seidner, G.; Alvarez, M. G.; Yeh, J.-I.; O'Driscoll, K. R.; Klepper,

[131824]J.; Stump, T. S.; Wang, D.; Spinner, N. B.; Birnbaum, M. J.;
De Vivo,

[131825]D. C.: GLUT-1 deficiency syndrome caused by haploinsuf-
ficiency of

[131826]the blood-brain barrier hexose carrier. *Nature Genet.* 18:
188-191,

[131827]1998.

[131828]23465.

[131829]Shepherd, P. R.; Kahn, B. B.: Glucose transporters and in-
sulin

[131830]action: implications for insulin resistance and diabetes
mellitus. *New*

[131831]*Eng. J. Med.* 341: 248-257, 1999.

[131832]23466.

[131833]Shows, T. B.; Eddy, R. L.; Byers, M. G.; Fukushima, Y.; De-
haven,

[131834]C. R.; Murray, J. C.; Bell, G. I.: Polymorphic human glucose
transporter

[131835]gene (GLUT) is on chromosome 1p31.3-p35. *Diabetes* 36:
546-549, 1987.

[131836]23467.

[131837]Kelley, M. J.; Pech, M.; Seuanez, H. N.; Rubin, J. S.; O'Brien,
[131838]S. J.; Aaronson, S. A.: Emergence of the keratinocyte
growth factor
[131839]multigene family during the great ape radiation. Proc. Nat.
Acad.
[131840]Sci. 89: 9287–9291, 1992.
[131841]23468.
[131842]Mattei, M.–G.; deLapeyriere, O.; Bresnick, J.; Dickson, C.;
Birnbaum,
[131843]D.; Mason, I.: Mouse Fgf7 (fibroblast growth factor 7) and
Fgf8 (fibroblast
[131844]growth factor 8) genes map to chromosomes 2 and 19 re-
spectively. Mammalian
[131845]Genome 6: 196–197, 1995.
[131846]23469.
[131847]Rubin, J. S.; Osada, H.; Finch, P. W.; Taylor, W. G.;
Rudikoff,
[131848]S.; Aaronson, S. A.: Purification and characterization of a
newly
[131849]identified growth factor specific for epithelial cells. Proc.
Nat.

[131850]Acad. Sci. 86: 802–806, 1989.

[131851]23470.

[131852]Werner, S.; Smola, H.; Liao, X.; Longaker, M. T.; Krieg, T.;
Hofschneider,

[131853]P. H.; Williams, L. T.: The function of KGF in morphogene-
sis of epithelium

[131854]and reepithelialization of wounds. Science 266: 819–822,
1994.

[131855]23471.

[131856]Zimonjic, D. B.; Kelley, M. J.; Rubin, J. S.; Aaronson, S. A.;

[131857]Popescu, N. C.: Fluorescence in situ hybridization analysis
of keratinocyte

[131858]growth factor gene amplification and dispersion in evolu-
tion of great

[131859]apes and humans. Proc. Nat. Acad. Sci. 94: 11461–11465,
1997.

[131860]23472.

[131861]Hoeck, H. C.; Vestergaard, P.; Jakobsen, P. E.; Falhof, J.;

Laurberg,

[131862]P.: Diagnosis of growth hormone (GH) deficiency in adults
with hypothalamic–pituitary

[131863] disorders: comparison of test results using pyridostigmine plus GH-releasing

[131864] hormone (GHRH), clonidine plus GHRH, and insulin-induced hypoglycemia

[131865] as GH secretagogues. J. Clin. Endocr. Metab. 85: 1467-1472, 2000.

[131866] 23473.

[131867] Igarashi, Y.; Ogawa, M.; Kamijo, T.; Iwatani, N.; Nishi, Y.; Kohno,

[131868] H.; Masumura, T.; Koga, J.: A new mutation causing inherited growth

[131869] hormone deficiency: a compound heterozygote of a 6.7 kb deletion and

[131870] a two base deletion in the third exon of the GH-1 gene. Hum. Molec.

[131871] Genet. 2: 1073-1074, 1993.

[131872] 23474.

[131873] Johannsson, G.; Marin, P.; Lonn, L.; Ottosson, M.; Stenlof, K.;

[131874] Bjorntorp, P.; Sjostrom, L.; Bengtsson, B.-A.: Growth hormone treatment

[131875] of abdominally obese men reduces abdominal fat mass,

improves glucose

[131876]and lipoprotein metabolism, and reduces diastolic blood pressure. J.

[131877]Clin. Endocr. Metab. 82: 727–734, 1997.

[131878]23475.

[131879]Kamijo, T.; Hayashi, Y.; Shimatsu, A.; Kinoshita, E.; Yoshimoto,

[131880]M.; Ogawa, M.; Seo, H.: Mutations in intron 3 of GH-1 gene associated

[131881]with isolated GH deficiency type II in three Japanese families. Clin.

[131882]Endocr. 51: 355–360, 1999.

[131883]23476.

[131884]Kowarski, A. A.; Schneider J.; Ben-Galim, E.; Weldon, V. V.; Daughaday,

[131885]W. H.: Growth failure with normal serum RIA-GH and low somatomedin

[131886]activity: somatomedin restoration and growth acceleration after exogenous

[131887]GH. J. Clin. Endocr. Metab. 47: 461–464, 1978.

[131888]23477.

[131889]Lebo, R. V.: Personal Communication. San Francisco, Calif.

[131890]7/31/1980.

[131891]23478.

[131892]Leiberman, E.; Pesler, D.; Parvari, R.; Elbedour, K.; Abdul-Latif,

[131893]H.; Brown, M. R.; Parks, J. S.; Carmi, R.: Short stature in carriers

[131894]of recessive mutation causing familial isolated growth hormone deficiency. Am.

[131895]J. Med. Genet. 90: 188-192, 2000.

[131896]23479.

[131897]Maghnie, M.; Strigazzi, C.; Tinelli, C.; Autelli, M.; Cisternino,

[131898]M.; Loche, S.; Severi, F.: Growth hormone (GH) deficiency (GHD) of

[131899]childhood onset: reassessment of GH status and evaluation of the predictive

[131900]criteria for permanent GHD in young adults. J. Clin. Endocr. Metab. 84:

[131901]1324-1328, 1999.

[131902]23480.

[131903]Mahajan, T.; Lightman, S. L.: A simple test for growth hormone

[131904]deficiency in adults. J. Clin. Endocr. Metab. 85:
1473–1476, 2000.

[131905]23481.

[131906]Martial, J. A.; Hallewell, R. A.; Baxter, J. D.; Goodman, H.
M.

[131907]: Human growth hormone: complementary DNA cloning
and expression in

[131908]bacteria. Science 205: 602–607, 1979.

[131909]23482.

[131910]Martinez, A. S.; Domene, H. M.; Ropelato, M. G.; Jasper, H.
G.;

[131911]Pennisi, P. A.; Escobar, M. E.; Heinrich, J. J.: Estrogen
priming

[131912]effect on growth hormone (GH) provocative test: a useful
tool for

[131913]the diagnosis of GH deficiency. J. Clin. Endocr. Metab. 85:
4168–4172,

[131914]2000.

[131915]23483.

[131916]Masuda, N.; Watahiki, M.; Tanaka, M.; Yamakawa, M.; Shimizu, K.;

[131917]Nagai, J.; Nakashima, K.: Molecular cloning of cDNA encoding 20 kDa

[131918]variant human growth hormone and the alternative splicing mechanism. Biochim.

[131919]Biophys. Acta 949: 125–131, 1988.

[131920]23484.

[131921]McCarthy, E. M. S.; Phillips, J. A., III: Characterization of

[131922]an intron splice enhancer that regulates alternative splicing of human

[131923]GH pre-mRNA. Hum. Molec. Genet. 7: 1491–1496, 1998.

[131924]23485.

[131925]Mendlewicz, J.; Linkowski, P.; Kerkhofs, M.; Leproult, R.; Copinschi,

[131926]G.; Van Cauter, E.: Genetic control of 24-hour growth hormone secretion

[131927]in man: a twin study. J. Clin. Endocr. Metab. 84: 856–862, 1999.

[131928]23486.

[131929]Mericq, V.; Cassorla, F.; Salazar, T.; Avila, A.; Iniguez, G.;

- [131930] Bowers, C. Y.; Merriam, G. R.: Effects of eight months treatment
- [131931] with graded doses of a growth hormone (GH)-releasing peptide in GH-deficient
- [131932] children. J. Clin. Endocr. Metab. 83: 2355–2360, 1998.
- [131933] 23487.
- [131934] Morgan, J. R.; Barrandon, Y.; Green, H.; Mulligan, R. C.: Expression
- [131935] of an exogenous growth hormone gene by transplantable human epidermal
- [131936] cells. Science 237: 1476–1479, 1987.
- [131937] 23488.
- [131938] Moseley, C. T.; Mullis, P. E.; Prince, M. A.; Phillips, J. A.,
- [131939] III: An exon splice enhancer mutation causes autosomal dominant GH
- [131940] deficiency. J. Clin. Endocr. Metab. 87: 847–852, 2002.
- [131941] 23489.
- [131942] Mullis, P. E.; Akinci, A.; Kanaka, C.; Eble, A.; Brook, C. G.
- [131943] D.: Prevalence of human growth hormone-1 gene deletions among patients

[131944]with isolated growth hormone deficiency from different populations. *Pediat.*

[131945]*Res.* 31: 532–534, 1992.

[131946]23490.

[131947]Niall, H. D.; Hogan, M. L.; Sauer, R.; Rosenblum, I. Y.; Greenwood,

[131948]F. C.: Sequence of pituitary and placental lactogenic and growth

[131949]hormones: evolution from a primordial peptide by gene reduplication. *Proc.*

[131950]*Nat. Acad. Sci.* 68: 866–869, 1971.

[131951]23491.

[131952]Nishi, Y.; Tanaka, T.; Takano, K.; Fujieda, K.; Igarashi, Y.;

[131953]Hanew, K.; Hirano, T.; Yokoya, S; Tachibana, K.; Saito, T.; Watanabe,

[131954]S.: Recent status in the occurrence of leukemia in growth hormone-treated

[131955]patients in Japan. *J. Clin. Endocr. Metab.* 84: 1961–1965, 1999.

[131956]23492.

[131957]Paladini, A. C.; Pena, C.; Retegui, L. A.: The intriguing na-

ture

[131958]of the multiple actions of growth hormone. Trends
Biochem. Sci. 4:

[131959]256–260, 1979.

[131960]23493.

[131961]Pena-Almazan, S.; Buchlis, J.; Miller, S.; Shine, B.;
Macgillivray,

[131962]M.: Linear growth characteristics of congenitally GH-
deficient infants

[131963]from birth to one year of age. J. Clin. Endocr. Metab. 86:
5691–5694,

[131964]2001.

[131965]23494.

[131966]Pfeifer, M.; Verhovec, R.; Zizek, B.; Prezelj, J.; Poredos, P.;

[131967]Clayton, R. N.: Growth hormone (GH) treatment reverses
early atherosclerotic

[131968]changes in GH-deficient adults. J. Clin. Endocr. Metab. 84:
453–457,

[131969]1999.

[131970]23495.

[131971]Phillips, J. A., III: Inherited defects in growth hormone syn–

thesis

[131972]and action.In: Scriver, C. R.; Beaudet, A. L.; Sly, W. S.;
Valle,

[131973]D. (eds.): The Metabolic and Molecular Bases of Inherited
Disease.

[131974]Vol. II. New York: McGraw-Hill (7th ed.): 1995. Pp.
3023-3044.

[131975]23496.

[131976]Phillips, J. A., III: Personal Communication. Baltimore, Md.

[131977]1/17/1983.

[131978]23497.

[131979]Phillips, J. A., III; Cogan, J. D.: Genetic basis of endocrine

[131980]disease 6: molecular basis of familial human growth hor-
mone deficiency. J.

[131981]Clin. Endocr. Metab. 78: 11-16, 1994.

[131982]23498.

[131983]Phillips, J. A., III; Hjelle, B. L.; Seeburg, P. H.; Plotnick,

[131984]L. P.; Migeon, C. J.; Zachmann, M.: Heterogeneity in the
molecular

[131985]basis of familial growth hormone deficiency (IGHD).

(Abstract) Am.

[131986]J. Hum. Genet. 33: 52A, 1981.

[131987]23499.

[131988]Rappaport, R.; Mugnier, E.; Limoni, C.; Crosnier, H.; Czernichow,

[131989]P.; Leger, J.; Limal, J.-M.; Rochiccioli, P.; Soskin, S.;
Brauner,

[131990]R.; Pinto, G.; Pawels, C.; and 27 others: A 5-year prospective
study of growth hormone (GH)-deficient children treated

[131991]with GH before
the age of 3 years. J. Clin. Endocr. Metab. 82: 452-456,

[131992]1997.

[131993]23500.

[131994]Buckingham, M.; Alonso, S.; Barton, P.; Cohen, A.; Daubas,
P.;

[131995]Garner, I.; Robert, B.; Weydert, A.: Actin and myosin multi-
gene families:

[131996]their expression during the formation and maturation of
striated muscle. Am.

[131997]J. Med. Genet. 25: 623-634, 1986.

[131998]23501.

- [131999]Edwards, Y. H.; Parkar, M.; Povey, S.; West, L. F.; Parrington,
ton,
- [132000]J. M.; Solomon, E.: Human myosin heavy chain genes assigned to chromosome
- [132001]17 using a human cDNA clone as probe. *Ann. Hum. Genet.*
49: 101–109,
- [132002]1985.
- [132003]23502.
- [132004]Epp, T. A.; Dixon, I. M. C.; Wang, H.–Y.; Sole, M. J.; Liew,
C.–C.
- [132005]: Structural organization of the human cardiac alpha–
myosin heavy
- [132006]chain gene (MYH6). *Genomics* 18: 505–509, 1993.
- [132007]23503.
- [132008]Geisterfer–Lowrance, A. A. T.; Kass, S.; Tanigawa, G.; Vos–
berg,
- [132009]H.–P.; McKenna, W.; Seidman, C. E.; Seidman, J. G.: A
molecular basis
- [132010]for familial hypertrophic cardiomyopathy: a beta cardiac
myosin heavy
- [132011]chain gene missense mutation. *Cell* 62: 999–1006, 1990.

[132012]23504.

[132013]Jaenicke, T.; Diederich, K. W.; Haas, W.; Schleich, J.;
Lichter,

[132014]P.; Pfordt, M.; Bach, A.; Vosberg, H.-P.: The complete se-
quence of

[132015]the human beta-myosin heavy chain gene and a compara-
tive analysis

[132016]of its product. Genomics 8: 194-206, 1990.

[132017]23505.

[132018]Kurabayashi, M.; Tsuchimochi, H.; Komuro, I.; Takaku, F.;
Yazaki,

[132019]Y.: Molecular cloning and characterization of human car-
diac alpha-

[132020]and beta-form myosin heavy chain complementary DNA
clones: regulation

[132021]of expression during development and pressure overload
in human atrium. J.

[132022]Clin. Invest. 82: 524-531, 1988.

[132023]23506.

[132024]Lichter, P.; Umeda, P. K.; Levin, J. E.; Vosberg, H.-P.: Par-
tial

[132025] characterization of the human beta-myosin heavy-chain gene which is
[132026] expressed in heart and skeletal muscle. *Europ. J. Biochem.*
160:
[132027] 419-426, 1986.
[132028] 23507.
[132029] Mahdavi, V.; Chambers, A. P.; Nadal-Ginard, B.: Cardiac
alpha-
[132030] and beta-myosin heavy chain genes are organized in tan-
dem. *Proc.*
[132031] *Nat. Acad. Sci.* 81: 2626-2630, 1984.
[132032] 23508.
[132033] Mahdavi, V.; Periasamy, M.; Nadal-Ginard, B.: Molecular
characterization
[132034] of two myosin heavy chain genes expressed in the adult
heart. *Nature* 297:
[132035] 659-664, 1982.
[132036] 23509.
[132037] Matsuoka, R.; Chambers, A.; Kimura, M.; Kanda, N.; Bruns,
G.;
[132038] Yoshida, M.; Takao, A.: Molecular cloning and chromoso-

mal localization

[132039] of a gene coding for human cardiac myosin heavy-chain.

Am. J. Med.

[132040] Genet. 29: 369–376, 1988.

[132041] 23510.

[132042] Matsuoka, R.; Yoshida, M. C.; Kanda, N.; Kimura, M.;

Ozasa, H.;

[132043] Takao, A.: Human cardiac myosin heavy chain gene

mapped within chromosome

[132044] region 14q11.2–q13. Am. J. Med. Genet. 32: 279–284,

1989.

[132045] 23511.

[132046] Nadal-Ginard, B.; Mahdavi, V.: Molecular basis of cardiac

performance:

[132047] plasticity of the myocardium generated through protein

isoform switches. J.

[132048] Clin. Invest. 84: 1693–1700, 1989.

[132049] 23512.

[132050] Niimura, H.; Patton, K. K.; McKenna, W. J.; Soultis, J.;

Maron,

[132051] B. J.; Seidman, J. G.; Seidman, C. E.: Sarcomere protein

gene mutations

[132052]in hypertrophic cardiomyopathy of the elderly. Circulation
105:

[132053]446-451, 2002.

[132054]23513.

[132055]Rappold, G. A.; Vosberg, H.-P.: Chromosomal localization
of a

[132056]human myosin heavy-chain gene by in situ hybridization.
Hum. Genet. 65:

[132057]195-197, 1983.

[132058]23514.

[132059]Adams, J. G., III; Morrison, W. T.; Steinberg, M. H.: A dou-
ble

[132060]crossover within a single human gene: Hb Parchman
(NH-2-delta-beta-delta-COOH).

[132061](Abstract) Am. J. Hum. Genet. 33: 34A only, 1981.

[132062]23515.

[132063]Adams, J. G., III; Morrison, W. T.; Steinberg, M. H.:
Hemoglobin

[132064]Parchman: double crossover within a single human gene.
Science 218:

[132065]291-293, 1982.

[132066]23516.

[132067]Ahern, E. J.; Ahern, V. N.; Aarons, G. H.; Jones, R. T.;
Brimhall,

[132068]B.: Hemoglobin Lepore Washington in two Jamaican families: interaction

[132069]with beta chain variants. Blood 40: 246-256, 1972.

[132070]23517.

[132071]Baglioni, C.: The fusion of two peptide chains in
hemoglobin Lepore

[132072]and its interpretation as a genetic deletion. Proc. Nat.
Acad. Sci. 48:

[132073]1880-1886, 1962.

[132074]23518.

[132075]Baglioni, C.; Ventruto, V.: Human abnormal hemoglobins.
II. A

[132076]chemical study of hemoglobin Lepore from a homozygote
individual. Europ.

[132077]J. Biochem. 5: 29-32, 1968.

[132078]23519.

[132079]McLean, R. H.; Hoefnagel, D.: Partial lipodystrophy and familial
milial

[132080]C3 deficiency. Hum. Hered. 30: 149–154, 1980.

[132081]23520.

[132082]Reaven, G. M.: Role of insulin resistance in human disease.
Diabetes 37:

[132083]1595–1607, 1988.

[132084]23521.

[132085]Sparkes, R. S.; Zollman, S.; Klisak, I.; Kirchgessner, T. G.;

[132086]Komaromy, M. C.; Mohandas, T.; Schotz, M. C.; Lusic, A. J.:
Human

[132087]genes involved in lipolysis of plasma lipoproteins: mapping of loci

[132088]for lipoprotein lipase to 8p22 and hepatic lipase to 15q21.
Genomics 1:

[132089]138–144, 1987.

[132090]23522.

[132091]Nurden, A. T.; Didry, D.; Kieffer, N.; McEver, R. P.: Residual

[132092]amounts of glycoproteins IIb and IIIa may be present in
the platelets

[132093] of most patients with Glanzmann's thrombasthenia. Blood
65: 1021-1024,

[132094] 1985.

[132095] 23523.

[132096] Nurden, A. T.; Rosa, J.-P.; Fournier, D.; Legrand, C.; Didry,

[132097] D.; Parquet, A.; Pidard, D.: A variant of Glanzmann's
thrombasthenia

[132098] with abnormal glycoprotein IIb-IIIa complexes in the
platelet membrane. J.

[132099] Clin. Invest. 79: 962-969, 1987.

[132100] 23524.

[132101] Budka, H.; Sluga, E.; Heiss, W.-D.: Spastic paraplegia as-
sociated

[132102] with Addison's disease: adult variant of adreno-
leukodystrophy. J.

[132103] Neurol. 213: 237-250, 1976.

[132104] 23525.

[132105] Cremers, F. P. M.; Pfeiffer, R. A.; van de Pol, T. J. R.;
Hofker,

[132106] M. H.; Kruse, T. A.; Wieringa, B.; Ropers, H. H.: An intersti-
tial

[132107]duplication of the X chromosome in a male allows physical
fine mapping

[132108]of probes from the Xq13-q22 region. Hum. Genet. 77:
23-27, 1987.

[132109]23526.

[132110]Bauer, P.; Knoblich, R.; Bauer, C.; Finckh, U.; Hufen, A.;
Kropp,

[132111]J.; Braun, S.; Kustermann-Kuhn, B.; Schmidt, D.; Harzer,
K.; Rolfs,

[132112]A.: NPC1: complete genomic sequence, mutation analysis,
and characterization

[132113]of haplotypes. Hum. Mutat. 19: 30-38, 2002.

[132114]23527.

[132115]Besley, G. T. N.; Hoogeboom, A. J. M.; Hoogeveen, A.;
Kleijer,

[132116]W. J.; Galjaard, H.: Somatic cell hybridisation studies
showing different

[132117]gene mutations in Niemann-Pick variants. Hum. Genet.
54: 409-412,

[132118]1980.

[132119]23528.

[132120]Blanchette-Mackie, E. J.; Dwyer, N. K.; Amende, L. M.;
Kruth, H.

[132121]S.; Butler, J. D.; Sokol, J.; Comly, M. E.; Vanier, M. T.; Au-
gust,

[132122]J. T.; Brady, R. O.; Pentchev, P. G.: Type-C Niemann-Pick
disease:

[132123]low density lipoprotein uptake is associated with prema-
ture cholesterol

[132124]accumulation in the Golgi complex and excessive choles-
terol storage

[132125]in lysosomes. Proc. Nat. Acad. Sci. 85: 8022-8026, 1988.

[132126]23529.

[132127]Boustany, R.-M. N.; Kaye, E.; Alroy, J.: Ultrastructural find-
ings

[132128]in skin from patients with Niemann-Pick disease, type C.
Pediat.

[132129]Neurol. 6: 177-183, 1990.

[132130]23530.

[132131]Brady, R. O.: Personal Communication. Bethesda, Md.
3/7/1986.

[132132]23531.

- [132133]Brady, R. O.: Sphingomyelin lipidoses: Niemann–Pick disease.In:
- [132134]Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.; Goldstein,
- [132135]J. L.; Brown, M. S.: The Metabolic Basis of Inherited Disease.
- [132136]New York: McGraw–Hill (pub.) (5th ed.): 1983. Pp. 731 only.
- [132137]23532.
- [132138]Brown, D. E.; Thrall, M. A.; Walkley, S. U.; Wenger, D. A.; Mitchell,
- [132139]T. W.; Smith, M. O.; Royals, K. L.; March, P. A.; Allison, R. W.:
- [132140]Feline Niemann–Pick disease type C. Am. J. Path. 144: 1412–1415,
- [132141]1994.
- [132142]23533.
- [132143]Carstea, E. D.; Morris, J. A.; Coleman, K. G.; Loftus, S. K.;
- [132144]Zhang, D.; Cummings, C.; Gu, J.; Rosenfeld, M. A.; Pavan, W. J.; Krizman,
- [132145]D. B.; Nagle, J.; Polymeropoulos, M. H.; and 26 others: Niemann–Pick

[132146]C1 disease gene: homology to mediators of cholesterol homeostasis. Science 277:

[132147]228–231, 1997.

[132148]23534.

[132149]Carstea, E. D.; Parker, C. C.; Fandino, L. B.; Vanier, M. T.;

[132150]Overhauser, J.; Weissenbach, J.; Pentchev, P. G.; Brady, R. O.; Polymeropoulos,

[132151]M. H.: Localizing the human Niemann–Pick C gene to 18q11–12. (Abstract) Am.

[132152]J. Hum. Genet. 55 (suppl.): A182, 1994.

[132153]23535.

[132154]Carstea, E. D.; Polymeropoulos, M. H.; Parker, C. C.; Detera–Wadleigh,

[132155]S. D.; O'Neill, R. R.; Patterson, M. C.; Goldin, E.; Xiao, H.; Straub,

[132156]R. E.; Vanier, M. T.; Brady, R. O.; Pentchev, P. G.: Linkage of Niemann–Pick

[132157]disease type C to human chromosome 18. Proc. Nat. Acad. Sci. 90:

[132158]2002–2004, 1993.

[132159]23536.

[132160]Ceuterick, C.; Martin, J. J.: Nerve biopsy findings in Niemann-Pick

[132161]type II (NPC). (Letter) Acta Neuropath. 88: 602-603, 1994.

[132162]23537.

[132163]Christomanou, H.: Niemann-Pick disease, type C: evidence for

[132164]the deficiency of an activating factor stimulating sphingomyelin and

[132165]glucocerebroside degradation. Hoppe Seylers Z. Physiol. Chem. 361:

[132166]1489-1502, 1980.

[132167]23538.

[132168]Davies, J. P.; Chen, F. W.; Ioannou, Y. A.: Transmembrane molecular

[132169]pump activity of Niemann-Pick C1 protein. Science 290: 2295-2298,

[132170]2000.

[132171]23539.

[132172]Erickson, R. P.; Garver, W. S.; Camargo, F.; Hossain, G. S.; Heidenreich,

[132173]R. A.: Pharmacological and genetic modifications of so-

matic cholesterol

[132174]do not substantially alter the course of CNS disease in
Niemann–Pick

[132175]C mice. J. Inherit. Metab. Dis. 23: 54–62, 2000.

[132176]23540.

[132177]Fink, J. K.; Filling–Katz, M. R.; Sokol, J.; Cogan, D. G.;
Pikus,

[132178]A.; Sonies, B.; Soong, B.; Pentchev, P. G.; Comly, M. E.;
Brady, R.

[132179]O.; Barton, N. W.: Clinical spectrum of Niemann–Pick dis-
ease type

[132180]C. Neurology 39: 1040–1049, 1989.

[132181]23541.

[132182]Gilbert, E. F.; Callahan, J.; Viseskul, C.; Opitz, J. M.: Nie-
mann–Pick

[132183]disease type C: pathological, histochemical, ultrastructural
and biochemical

[132184]studies. Europ. J. Pediat. 136: 263–274, 1981.

[132185]23542.

[132186]Greer, W. L.; Dobson, M. J.; Girouard, G. S.; Byers, D. M.;
Riddell,

- [132187]D. C.; Neumann, P. E.: Mutations in NPC1 highlight a conserved NPC1-specific
cysteine-rich domain. *Am. J. Hum. Genet.* 65: 1252–1260,
1999.
- [132189]23543.
- [132190]Greer, W. L.; Riddell, D. C.; Byers, D. M.; Welch, J. P.;
Girouard,
[132191]G. S.; Sparrow, S. M.; Gillan, T. L.; Neumann, P. E.: Linkage
of
[132192]Niemann–Pick disease type D to the same region of human chromosome
[132193]18 as Niemann–Pick disease type C. *Am. J. Hum. Genet.*
61: 139–142,
[132194]1997.
- [132195]23544.
- [132196]Greer, W. L.; Riddell, D. C.; Gillan, T. L.; Girouard, G. S.;
[132197]Sparrow, S. M.; Byers, D. M.; Dobson, M. J.; Neumann, P.
E.: The
[132198]Nova Scotia (type D) form of Niemann–Pick disease is
caused by a 3097G–T
[132199]transversion in NPC1. *Am. J. Hum. Genet.* 63: 52–54,
1998.

[132200]23545.

[132201]Imrie, J.; Wraith, J. E.: Niemann–Pick disease type C. Arch.

[132202]Dis. Child. 84: 427–429, 2001.

[132203]23546.

[132204]Ioannou, Y. A.: The structure and function of the Niemann–Pick

[132205]C1 protein. Molec. Genet. Metab. 71: 175–181, 2000.

[132206]23547.

[132207]Kruth, H. S.; Comly, M. E.; Butler, J. D.; Vanier, M. T.; Fink,

[132208]J. K.; Wenger, D. A.; Patel, S.; Pentchev, P. G.: Type C Niemann–Pick

[132209]disease: abnormal metabolism of low density lipoprotein in homozygous

[132210]and heterozygous fibroblasts. J. Biol. Chem. 261: 16769–16774, 1986.

[132211]23548.

[132212]Liscum, L.; Faust, J. R.: Low density lipoprotein (LDL)–mediated

[132213]suppression of cholesterol synthesis and LDL uptake is defective in

[132214]Niemann–Pick type C fibroblasts. J. Biol. Chem. 262:
17002–17008,

[132215]1987.

[132216]23549.

[132217]Liu, Y.; Wu, Y.–P.; Wada, R.; Neufeld, E. B.; Mullin, K. A.;
Howard,

[132218]A. C.; Pentchev, P. G.; Vanier, M. T.; Suzuki, K.; Proia, R. L.:

[132219]Alleviation of neuronal ganglioside storage does not im-
prove the clinical

[132220]course of the Niemann–Pick C disease mouse. Hum.
Molec. Genet. 9:

[132221]1087–1092, 2000.

[132222]23550.

[132223]Loftus, S. K.; Morris, J. A.; Carstea, E. D.; Gu, J. Z.; Cum-
mings,

[132224]C.; Brown, A.; Ellison, J.; Ohno, K.; Rosenfeld, M. A.; Tagle,
D.

[132225]A.; Pentchev, P. G.; Pavan, W. J.: Murine model of Nie-
mann–Pick C

[132226]disease: mutation in a cholesterol homeostasis gene. Sci-
ence 277:

[132227]232–235, 1997.

[132228]23551.

[132229]Lowenthal, A. C.; Cummings, J. F.; Wenger, D. A.; Thrall, M. A.;

[132230]Wood, P. A.; de Lahunta, A.: Feline sphingolipidosis re-
sembling Niemann–Pick

[132231]disease type C. Acta Neuropath. 81: 189–197, 1990.

[132232]23552.

[132233]Meiner, V.; Shpitzen, S.; Mandel, H.; Klar, A.; Ben–Neriah, Z.;

[132234]Zlotogora, J.; Sagi, M.; Lossos, A.; Bargal, R.; Sury, V.; Carmi,

[132235]R.; Leitersdorf, E.; Zeigler, M.: Clinical–biochemical corre-
lation

[132236]in molecularly characterized patients with Niemann–Pick
type C. Genet.

[132237]Med. 3: 343–348, 2001.

[132238]23553.

[132239]Cappa, M.; Bertini, E.; del Balzo, P.; Cambiaso, P.; Di Biase,

[132240]A.; Salvati, S.: High dose immunoglobulin IV treatment in
adrenoleukodystrophy. J.

[132241]Neurol. Neurosurg. Psychiat. 57 (suppl.): 69–70, 1994.

[132242]23554.

[132243]Cartier, N.; Sarde, C.-O.; Douar, A.-M.; Mosser, J.; Mandel, J.-L.;

[132244]Aubourg, P.: Abnormal messenger RNA expression and a missense mutation

[132245]in patients with X-linked adrenoleukodystrophy. Hum. Molec. Genet. 2:

[132246]1949–1951, 1993.

[132247]23555.

[132248]Cotrufo, R.; Melone, M. A. B.; Monsurro, M. R.; Di Iorio, G.;

[132249]Carella, C.; Moser, H. W.: Phenotype heterogeneity among hemizygotes

[132250]in a family biochemically screened for adrenoleukodystrophy. Am.

[132251]J. Med. Genet. 26: 833–838, 1987.

[132252]23556.

[132253]Davis, L. E.; Snyder, R. D.; Orth, D. N.; Nicholson, W. E.; Kornfeld,

[132254]M.; Seelinger, D. F.: Adrenoleukodystrophy and adrenomyeloneuropathy

[132255]associated with partial adrenal insufficiency in three generations

[132256]of a kindred. Am. J. Med. 66: 342–347, 1979.

[132257]23557.

[132258]Di Chiro, G.; Eiben, R. M.; Manz, H. J.; Jacobs, I. B.; Schellinger,

[132259]D.: A new CT pattern in adrenoleukodystrophy. Radiology 137: 687–692,

[132260]1980.

[132261]23558.

[132262]Di Rocco, M.; Doria-Lamba, L.; Caruso, U.: Monozygotic twins

[132263]with X-linked adrenoleukodystrophy and different phenotypes. Ann.

[132264]Neurol. 50: 424 only, 2001.

[132265]23559.

[132266]Dodd, A.; Rowland, S. A.; Hawkes, S. L. J.; Kennedy, M. A.; Love,

[132267]D. R.: Mutations in the adrenoleukodystrophy gene. Hum. Mutat. 9:

[132268]500–511, 1997.

[132269]23560.

[132270]Dvorakova, L.; Storkanova, G.; Unterrainer, G.; Hujova, J.;
Kmoch,

[132271]S.; Zeman, J.; Hrebicek, M.; Berger, J.: Eight novel ABCD1
gene mutations

[132272]and three polymorphisms in patients with X-linked
adrenoleukodystrophy:

[132273]the first polymorphism causing an amino acid exchange.
Hum. Mutat. 18:

[132274]52–60, 2001.

[132275]23561.

[132276]Eichler, E. E.; Budarf, M. L.; Rocchi, M.; Deaven, L. L.;
Doggett,

[132277]N. A.; Baldini, A.; Nelson, D. L.; Mohrenweiser, H. W.: In-
terchromosomal

[132278]duplications of the adrenoleukodystrophy locus: a phe-
nomenon of pericentromeric

[132279]plasticity. Hum. Molec. Genet. 6: 991–1002, 1997.

[132280]23562.

[132281]El-Deiry, S. S.; Naidu, S.; Blevins, L. S.; Ladenson, P. W.:

[132282]Assessment of adrenal function in women heterozygous

for adrenoleukodystrophy. J.

[132283]Clin. Endocr. Metab. 82: 856–860, 1997.

[132284]23563.

[132285]Fanconi, A.; Prader, A.; Isler, W.; Luthy, F.; Siebenmann, R.

[132286]E.: Morbus Addison mit Hirnsklerose im Kindesalter. Ein
hereditaeres

[132287]Syndrom mit X-chromosomaler Vererbung? Helv. Paediat.
Acta 18: 480–501,

[132288]1963.

[132289]23564.

[132290]Fanen, P.; Guidoux, S.; Sarde, C.-O.; Mandel, J.-L.;
Goossens,

[132291]M.; Aubourg, P.: Identification of mutations in the putative
ATP-binding

[132292]domain of the adrenoleukodystrophy gene. J. Clin. Invest.
94: 516–520,

[132293]1994.

[132294]23565.

[132295]Federico, A.; Dotti, M. T.; Annunziata, P.; Bonuccelli, U.;
Fenzi,

[132296]G.; Ciacci, G.; Malandrini, A.; Meucci, G.; Guazzi, G. C.:

Adrenomyeloneurodystrophy

[132297] with late cerebral involvement and evidence of a multiple autoimmune

[132298] disorder. J. Inherit. Metab. Dis. 11 (suppl. 2): 169–172, 1988.

[132299] 23566.

[132300] Feil, R.; Aubourg, P.; Mosser, J.; Douar, A.–M.; Le Paslier, D.;

[132301] Philippe, C.; Mandel, J.–L.: Adrenoleukodystrophy: a complex chromosomal

[132302] rearrangement in the Xq28 red/green–color–pigment gene region indicates

[132303] two possible gene locations. Am. J. Hum. Genet. 49: 1361–1371, 1991.

[132304] 23567.

[132305] Forss–Petter, S.; Werner, H.; Berger, J.; Lassmann, H.; Molzer,

[132306] B.; Schwab, M. H.; Bernheimer, H.; Zimmermann, F.; Nave, K.–A.: Targeted

[132307] inactivation of the X–linked adrenoleukodystrophy gene in mice. J.

[132308] Neurosci. Res. 50: 829–843, 1997.

[132309]23568.

[132310]Fuchs, S.; Sarde, C. O.; Wedemann, H.; Schwinger, E.;
Mandel,

[132311]J. L.; Gal, A.: Missense mutations are frequent in the gene
for X-chromosomal

[132312]adrenoleukodystrophy (ALD). Hum. Molec. Genet. 3:
1903-1905, 1994.

[132313]23569.

[132314]Griffin, J. W.; Goren, E.; Schaumburg, H. H.; Engel, W. K.;
Loriaux,

[132315]L.: Adrenomyeloneuropathy: a probable variant of
adrenoleukodystrophy.

[132316]J. Clinical and endocrinologic aspects. Neurology 27:
1107-1113,

[132317]1977.

[132318]23570.

[132319]Guimaraes, C. P.; Lemos, M.; Menezes, I.; Coelho, T.; Sa-
Miranda,

[132320]C.; Azevedo, J. E.: Characterisation of two mutations in the
ABCD1

[132321]gene leading to low levels of normal ALDP. Hum. Genet.
109: 616-622,

[132322]2001.

[132323]23571.

[132324]Gumbinas, M.; Liu, H. M.; Dawson, G.; Larsen, M.; Green,
O.:

[132325]Progressive spastic paraparesis and adrenal insufficiency.
Arch.

[132326]Neurol. 33: 678-680, 1976.

[132327]23572.

[132328]Haberfeld, W.; Spieler, F.: Zur diffusen Hirn-
Ruckenmarksklerose

[132329]im Kindesalter. Dt. Z. Nervheilk 40: 436-463, 1910.

[132330]23573.

[132331]Harris-Jones, J. N.; Nixon, P. G. F.: Familial Addison's dis-
ease

[132332]with spastic paraplegia. J. Clin. Endocr. 15: 739-744,
1955.

[132333]23574.

[132334]Hashmi, M.; Stanley, W.; Singh, I.: Lignoceroyl-CoASH lig-
ase:

[132335]enzyme defect in fatty acid beta-oxidation system in X-
linked childhood

[132336]adrenoleukodystrophy. FEBS Lett. 196: 247–250, 1986.

[132337]23575.

[132338]Hecaen, H.; De Ajuriaguerra, J.: Balint's syndrome (psychic paralysis

[132339]of visual fixation) and its minor forms. Brain 77: 373–400, 1954.

[132340]23576.

[132341]Heffungs, W.; Hameister, H.; Ropers, H. H.: Addison disease and

[132342]cerebral sclerosis in an apparently heterozygous girl: evidence for

[132343]inactivation of the adrenoleukodystrophy locus. Clin. Genet. 18:

[132344]184–188, 1980.

[132345]23577.

[132346]Ho, J. K.; Moser, H.; Kishimoto, Y.; Hamilton, J. A.: Interactions

[132347]of a very long chain fatty acid with model membranes and serum albumin:

[132348]implications for the pathogenesis of adrenoleukodystrophy. J. Clin.

[132349]Invest. 96: 1455–1463, 1995.

[132350]23578.

[132351]Hoefnagel, D.; Brun, A.; Ingbar, S. H.; Goldman, H.: Addison's

[132352]disease and diffuse cerebral sclerosis. J. Neurol. Neurosurg. Psychiat. 30:

[132353]56–60, 1967.

[132354]23579.

[132355]Hoefnagel, D.; Van den Noort, S.; Ingbar, S. H.: Diffuse cerebral

[132356]sclerosis with endocrine abnormalities in young males. Brain 85:

[132357]553–568, 1962.

[132358]23580.

[132359]Holmberg, B. H.; Hagg, E.; Hagenfeldt, L.: Adrenomyeloneuropathy--report

[132360]on a family. J. Intern. Med. 230: 535–538, 1991.

[132361]23581.

[132362]Igarashi, M.; Schaumburg, H. H.; Powers, J.; Kishimoto, Y.; Kolodny,

[132363]E. H.; Suzuki, K.: Fatty acid abnormality in
adrenoleukodystrophy. J.

[132364]Neurochem. 26: 851–860, 1976.

[132365]23582.

[132366]Kano, S.; Watanabe, M.; Kanai, M.; Koike, R.; Onodera, O.;
Tsuji,

[132367]S.; Okamoto, K.; Shoji, M.: A Japanese family with
adrenoleukodystrophy

[132368]with a codon 291 deletion: a clinical, biochemical, patho-
logical,

[132369]and genetic report. J. Neurol. Sci. 158: 187–192, 1998.

[132370]23583.

[132371]Kemp, S.; Ligtenberg, M. J. L.; van Geel, B. M.; Barth, P. G.;

[132372]Sarde, C.-O.; van Oost, B. A.; Bolhuis, P. A.: Two intronic
mutations

[132373]in the adrenoleukodystrophy gene. Hum. Mutat. 6:
272–273, 1995.

[132374]23584.

[132375]Kemp, S.; Pujol, A.; Waterham, H. R.; van Geel, B. M.;
Boehm,

[132376]C. D.; Raymond, G. V.; Cutting, G. R.; Wanders, R. J. A.;

Moser, H.

[132377]W.: ABCD1 mutations and the X-linked adrenoleukodys-
trophy mutation

[132378]database: role in diagnosis and clinical correlations. Hum.
Mutat. 18:

[132379]499–515, 2001.

[132380]23585.

[132381]Ito, M.; Yuan, C.-X.; Malik, S.; Gu, W.; Fondell, J. D.; Yama-
mura,

[132382]S.; Fu, Z.-Y.; Zhang, X.; Qin, J.; Roeder, R. G.: Identity be-
tween

[132383]TRAP and SMCC complexes indicates novel pathways for
the function

[132384]of nuclear receptors and diverse mammalian activators.
Molec. Cell 3:

[132385]361–370, 1999.

[132386]23586.

[132387]Nagase, T.; Seki, N.; Ishikawa, K.; Tanaka, A.; Nomura, N.:
Prediction

[132388]of the coding sequences of unidentified human genes. V.
The coding

[132389]sequences of 40 new genes (KIAA0161–KIAA0200) de-

duced by analysis

[132390]of cDNA clones from human cell line KG-1. DNA Res. 3:
17-24, 1996.

[132391]23587.

[132392]Maeda, K.; Matsuhashi, S.; Hori, K.; Xin, Z.; Mukai, T.;
Tabuchi,

[132393]K.; Egashira, M.; Niikawa, N.: Cloning and characterization
of a

[132394]novel human gene, TM4SF6, encoding a protein belonging
to the transmembrane

[132395]4 superfamily, and mapped to Xq22. Genomics 52:
240-242, 1998.

[132396]23588.

[132397]Todd, S. C.; Doctor, V. S.; Levy, S.: Sequences and expres-
sion

[132398]of six new members of the tetraspanin/TM4SF family.
Biochim. Biophys.

[132399]Acta 1399: 101-104, 1998.

[132400]23589.

[132401]Vaccari, T.; Beltrame, M.; Ferrari, S.; Bianchi, M. E.: Hmg4,

[132402]a new member of the Hmg1/2 gene family. Genomics 49:

247–252, 1998.

[132403]23590.

[132404]Wilke, K.; Wiemann, S.; Gaul, R.; Gong, W.; Poustka, A.:

Isolation

[132405]of human and mouse HMG2a cDNAs: evidence for an
HMG2a-specific 3-prime

[132406]untranslated region. *Gene* 198: 269–274, 1997.

[132407]23591.

[132408]Bird, A. P.: HTF islands as gene markers in the vertebrate
nucleus. *Trends*

[132409]*Genet.* 3: 342–347, 1987.

[132410]23592.

[132411]Anson, D. S.; Blake, D. J.; Winship, P. R.; Birnbaum, D.;
Brownlee,

[132412]G. G.: Nullisomic deletion of the mcf.2 transforming gene
in two

[132413]haemophilia B patients. *EMBO J.* 7: 2795–2799, 1988.

[132414]23593.

[132415]Gong, Y.; Slee, R. B.; Fukai, N.; Rawadi, G.; Roman–Roman,
S.;

[132416]Reginato, A. M.; Wang, H.; Cundy, T.; Glorieux, F. H.; Lev,

D.; Zacharin,
[132417]M.; Oexle, K.; and 50 others: LDL receptor-related protein
5 (LRP5)
[132418]affects bone accrual and eye development. Cell 107:
513–523, 2001.
[132419]23594.

[132420]Abadie, V.; Jaruzelska, J.; Lyonnet, S.; Millasseau, P.;
Berthelon,
[132421]M.; Rey, F.; Munnich, A.; Rey, J.: Illegitimate transcription
of
[132422]the phenylalanine hydroxylase gene in lymphocytes for
identification
[132423]of mutations in phenylketonuria. Hum. Molec. Genet. 2:
31–34, 1993.
[132424]23595.

[132425]Abadie, V.; Lyonnet, S.; Maurin, N.; Berthelon, M.; Caillaud,
C.;
[132426]Giraud, F.; Mattei, J.-F.; Rey, J.; Rey, F.; Munnich, A.: CpG
dinucleotides
[132427]are mutation hot spots in phenylketonuria. Genomics 5:
936–939,
[132428]1989.

[132429]23596.

[132430]Abadie, V.; Lyonnet, S.; Maurin, N.; Berthelon, M.; Caillaud, C.;

[132431]Giraud, F.; Mattei, J.-F.; Rey, J.; Rey, F.; Munnich, A.: CpG dinucleotides

[132432]are mutation hot spots in phenylketonuria. (Abstract) Am. J. Hum.

[132433]Genet. 45 (suppl.): A169 only, 1989.

[132434]23597.

[132435]Weiss, P.; Tietze, F.; Gahl, W. A.; Seppala, R.; Ashwell, G.:

[132436]Identification of the metabolic defect in sialuria. J. Biol. Chem. 264:

[132437]17635-17636, 1989.

[132438]23598.

[132439]Kennedy, M. A.; Rowland, S. A.; Miller, A. L.; Morris, C. M.;

[132440]Neville, L. A.; Dodd, A.; Fifield, W. J.; Love, D. R.: Structure

[132441]and location of the murine adrenoleukodystrophy gene. Genomics 32:

[132442]395-400, 1996.

[132443]23599.

- [132444] Bassi, M. T.; Ramesar, R. S.; Caciotti, B.; Winship, I. M.; De
- [132445] Grandi, A.; Riboni, M.; Townes, P. L.; Beighton, P.; Ballabio, A.;
- [132446] Borsani, G.: X-linked late-onset sensorineural deafness caused by
- [132447] a deletion involving OA1 and a novel gene containing WD-40 repeats. Am.
- [132448] J. Hum. Genet. 64: 1604-1616, 1999.
- [132449] 23600.
- [132450] Converse, P. J.: Personal Communication. Baltimore, Md. 8/24/2001.
- [132451] 23601.
- [132452] Disteche, C. M.; Dinulos, M. B.; Bassi, M. T.; Elliott, R. W.;
- [132453] Rugarli, E. I.: Mapping of the murine Tbl1 gene reveals a new rearrangement
- [132454] between mouse and human X chromosomes. Mammalian Genome 9: 1062-1064,
- [132455] 1998.
- [132456] 23602.
- [132457] Dong, X.; Tsuda, L.; Zavitz, K. H.; Lin, M.; Li, S.; Carthew, R.

[132458]W.; Zipursky, S. L.: ebi regulates epidermal growth factor receptor

[132459]signaling pathways in Drosophila. Genes Dev. 13: 954–965, 1999.

[132460]23603.

[132461]Matsuzawa, S.; Reed, J. C.: Siah–1, SIP, and Ebi collaborate in

[132462]a novel pathway for beta–catenin degradation linked to p53 responses. Molec.

[132463]Cell 7: 915–926, 2001.

[132464]23604.

[132465]Zhang, J.; Kalkum, M.; Chait, B. T.; Roeder, R. G.: The N–CoR–HDAC3

[132466]nuclear receptor corepressor complex inhibits the JNK pathway through

[132467]the integral subunit GPS2. Molec. Cell 9: 611–623, 2002.

[132468]23605.

[132469]Mu, J.; Skurat, A. V.; Roach, P. J.: Glycogenin–2, a novel self–glucosylating

[132470]protein involved in liver glycogen biosynthesis. J. Biol. Chem. 272:

[132471]27589–27597, 1997.

[132472]23606.

[132473]Bartley, J. A.; Miller, D. K.; Hayford, J. T.; McCabe, E. R. B.

[132474]: Concordance of X-linked glycerol kinase deficiency with
X-linked

[132475]congenital adrenal hypoplasia. *Lancet* II: 733–736, 1982.

[132476]23607.

[132477]Jansen, G. A.; Ofman, R.; Ferdinandusse, S.; IJlst, L.; Mui-
jsers,

[132478]A. O.; Skjeldal, O. H.; Stokke, O.; Jakobs, C.; Besley, G. T.
N.;

[132479]Wraith, J. E.; Wanders, R. J. A.: Refsum disease is caused
by mutations

[132480]in the phytanoyl-CoA hydroxylase gene. *Nature Genet.* 17:
190–193,

[132481]1997.

[132482]23608.

[132483]Mihalik, S. J.; Morrell, J. C.; Kim, D.; Sacksteder, K. A.;
Watkins,

[132484]P. A.; Gould, S. J.: Identification of PAHX, a Refsum disease
gene. *Nature*

[132485]Genet. 17: 185–189, 1997.

[132486]23609.

[132487]Wilcken, B.; Don, N.; Greenaway, R.; Hammond, J.; Sosula, L.:

[132488]Sialuria: a second case. J. Inherit. Metab. Dis. 10: 97–102, 1987.

[132489]23610.

[132490]Bickel, H.; Gerrard, J.; Hickmans, E. M.: Influence of phenylalanine

[132491]intake on phenylketonuria. Lancet 2: 812–813, 1953.

[132492]23611.

[132493]Bowden, J. A.; McArthur, C. L., III: Possible biochemical model

[132494]for phenylketonuria. Nature 235: 230, 1972.

[132495]23612.

[132496]Brenton, D. P.; Lilburn, M.: Maternal phenylketonuria: a study

[132497]from the United Kingdom. Europ. J. Pediat. 155 (suppl. 1): 177–180,

[132498]1996.

[132499]23613.

[132500]Burgard, P.; Rupp, A.; Konecki, D. S.; Trefz, F. K.; Schmidt,
[132501]H.; Lichter-Konecki, U.: Phenylalanine hydroxylase geno-
types, predicted
[132502]residual enzyme activity and phenotypic parameters of di-
agnosis and
[132503]treatment of phenylketonuria. *Europ. J. Pediat.* 155
(suppl.): S11-S15,
[132504]1996.
[132505]23614.

[132506]Byck, S.; Tyfield, L.; Carter, K.; Scriver, C. R.: Prediction
[132507]of multiple hypermutable codons in the human PAH gene:
codon 280 contains
[132508]recurrent mutations in Quebec and other populations.
Hum. Mutat. 9:
[132509]316-321, 1997.
[132510]23615.

[132511]Cabalska, B.: Personal Communication. Warsaw, Poland
1980.
[132512]23616.

[132513]Caillaud, C.; Lyonnet, S.; Rey, F.; Melle, D.; Frebourg, T.;
Berthelon,

[132514]M.; Vilarinho, L.; Vaz Osorio, R.; Rey, J.; Munnich, A.: A
3-base

[132515]pair in-frame deletion of the phenylalanine hydroxylase
gene results

[132516]in a kinetic variant of phenylketonuria. J. Biol. Chem. 266:
9351-9354,

[132517]1991.

[132518]23617.

[132519]Carter, C. O.; Woolf, L. I.: The birthplaces of parents and
grandparents

[132520]of a series of patients with phenylketonuria in southeast
England. Ann.

[132521]Hum. Genet. 25: 57-64, 1961.

[132522]23618.

[132523]Carter, K. C.; Byck, S.; Waters, P. J.; Richards, B.; Nowacki,

[132524]P. M.; Laframboise, R.; Lambert, M.; Treacy, E.; Sriver, C.
R.:

[132525]Mutation at the phenylalanine hydroxylase gene (PAH) and
its use to

[132526]document population genetic variation: the Quebec expe-
rience. Europ.

[132527]J. Hum. Genet. 6: 61–70, 1998.

[132528]23619.

[132529]Centerwall, W. R.; Neff, C. A.: Phenylketonuria: a case report

[132530]of children of Jewish ancestry. Arch. Pediat. 78: 379–384, 1961.

[132531]23620.

[132532]Chao, H.-K.; Hsiao, K.-J.; Su, T.-S.: A silent mutation induces

[132533]exon skipping in the phenylalanine hydroxylase gene in phenylketonuria. Hum.

[132534]Genet. 108: 14–19, 2001.

[132535]23621.

[132536]Chen, K.-J.; Chao, H.-K.; Hsiao, K.-J.; Su, T.-S.: Identification

[132537]and characterization of a novel liver-specific enhancer of the human

[132538]phenylalanine hydroxylase gene. Hum. Genet. 110: 235–243, 2002.

[132539]23622.

[132540]Chen, S.-H.; Hsiao, K.-J.; Lin, L.-H.; Liu, T.-T.; Tang,

R.-B.;

[132541]Su, T.-S.: Study of restriction fragment length polymorphisms at

[132542]the human phenylalanine hydroxylase locus and evaluation of its potential

[132543]application in prenatal diagnosis of phenylketonuria in Chinese. Hum.

[132544]Genet. 81: 226-230, 1989.

[132545]23623.

[132546]Cipcic-Schmidt, S.; Trefz, F. K.; Funders, B.; Seidlitz, G.; Ullrich,

[132547]K.: German maternal phenylketonuria study. Europ. J. Pediat. 155

[132548](suppl. 1): 173-176, 1996.

[132549]23624.

[132550]Cohen, B. E.; Bodonyi, E.; Szeinberg, A.: Phenylketonuria in

[132551]Jews. Lancet I: 344-345, 1961.

[132552]23625.

[132553]Corsello, G.; Bosco, P.; Cali, F.; Greco, D.; Cammarata, M.; Ciaccio,

[132554]M.; Piccione, M.; Romano, V.: Maternal phenylketonuria in

two Sicilian

[132555]families identified by maternal blood phenylalanine level
screening

[132556]and identification of a new phenylalanine hydroxylase
gene mutation

[132557](P407L). (Letter) Europ. J. Pediat. 158: 83–88, 1999.

[132558]23626.

[132559]Coskun, T.; Ozalp, I.; Kale, G.; Gogus, S.: Scleroderma-like

[132560]skin lesions in two patients with phenylketonuria. Europ. J.
Pediat. 150:

[132561]109–110, 1990.

[132562]23627.

[132563]Cotton, R. G. H.: Heterogeneity of phenylketonuria at the
clinical,

[132564]protein and DNA levels. J. Inherit. Metab. Dis. 13:
739–750, 1990.

[132565]23628.

[132566]Cunningham, G. C.; Day, R. W.; Berman, J. L.; Hsia, D.
Y.–Y.:

[132567]Phenylalanine tolerance tests in families with phenylke-
tonuria and

- [132568]hyperphenylalaninemia. Am. J. Dis. Child. 117: 626-635, 1969.
- [132569]23629.
- [132570]Daiger, S. P.; Chakraborty, R.; Reed, L.; Fekete, G.; Schuler, D.; Berenssi, G.; Nasz, I.; Brdicka, R.; Kamaryt, J.; Pijackova, A.;
- [132572]Moore, S.; Sullivan, S.; Woo, S. L. C.: Polymorphic DNA haplotypes
- [132573]at the phenylalanine hydroxylase (PAH) locus in European families
- [132574]with phenylketonuria (PKU). Am. J. Hum. Genet. 45: 310-318, 1989.
- [132575]23630.
- [132576]Gaedigk, R.; Duncan, A. M. V.; Miyazaki, I.; Robinson, B. H.; Dosch, H.-M.: ICA1 encoding p69, a protein linked to the development of
- [132578]type 1 diabetes, maps to human chromosome 7p22. Cytogenet. Cell Genet. 66:
- [132579]274-276, 1994.
- [132580]23631.

[132581]Gaedigk, R.; Karges, W.; Hui, M. F.; Scherer, S. W.; Dosch, H.-M.

[132582]Genomic organization and transcript analysis of ICAp69, a target

[132583]antigen in diabetic autoimmunity. Genomics 38: 382–391, 1996.

[132584]23632.

[132585]Pietropaolo, M.; Castano, L.; Babu, S.; Buelow, R.; Kuo, Y.-L.

[132586]S.; Martin, S.; Martin, A.; Powers, A. C.; Prochazka, M.; Naggert,

[132587]J.; Leiter, E. H.; Eisenbarth, G. S.: Islet cell autoantigen 69 kD

[132588](ICA69): molecular cloning and characterization of a novel diabetes-associated

[132589]autoantigen. J. Clin. Invest. 92: 359–371, 1993.

[132590]23633.

[132591]Diaz, M. O.; Le Beau, M. M.; Pitha, P.; Rowley, J. D.: Interferon

[132592]and c-ets-1 genes in the translocation (9;11)(p22;q23) in human acute

[132593]monocytic leukemia. Science 231: 265–267, 1986.

[132594]23634.

[132595]Ohlsson, M.; Feder, J.; Cavalli-Sforza, L. L.; von Gabain, A.

[132596]: Close linkage of alpha and beta interferons and infrequent duplication

[132597]of beta interferon in humans. Proc. Nat. Acad. Sci. 82: 4473-4476,

[132598]1985.

[132599]23635.

[132600]Owerbach, D.; Rutter, W. J.; Shows, T. B.; Gray, P.; Goeddel,

[132601]D. V.; Lawn, R. M.: Leukocyte and fibroblast interferon genes are

[132602]located on human chromosome 9. Proc. Nat. Acad. Sci. 78: 3123-3127,

[132603]1981.

[132604]23636.

[132605]Siegal, F. P.; Kadowaki, N.; Shodell, M.; Fitzgerald-Bocarsly,

[132606]P. A.; Shah, K.; Ho, S.; Antonenko, S.; Liu, Y.-J.: The nature of

[132607]the principal type 1 interferon-producing cells in human

blood. Science 284:

[132608]1835–1837, 1999.

[132609]23637.

[132610]Heath, P.; Elvin, P.; Jenner, D.; Gammack, A.; Morten, J.;
Markham,

[132611]A.: Localisation of a cDNA clone for human cytokeratin 18
to chromosome

[132612]17p11–p12 by in situ hybridisation. Hum. Genet. 85:
669–670, 1990.

[132613]23638.

[132614]Ku, N.–O.; Michie, S.; Oshima, R. G.; Omary, M. B.: Chronic
hepatitis,

[132615]hepatocyte fragility, and increased soluble phosphogly-
cokeratins in

[132616]transgenic mice expressing a keratin 18 conserved argi-
nine mutant. J.

[132617]Cell Biol. 131: 1303–1314, 1995.

[132618]23639.

[132619]Ku, N.–O.; Michie, S. A.; Soetikno, R. M.; Resurreccion, E.
Z.;

[132620]Broome, R. L.; Oshima, R. G.; Omary, M. B.: Susceptibility

to hepatotoxicity

[132621]in transgenic mice that express a dominant-negative human keratin

[132622]18 mutant. J. Clin. Invest. 98: 1034–1046, 1996.

[132623]23640.

[132624]Kulesh, D. A.; Oshima, R. G.: Complete structure of the gene for

[132625]human keratin 18. Genomics 4: 339–347, 1989.

[132626]23641.

[132627]Lendahl, U.; Zimmerman, L. B.; McKay, R. D. G.: CNS stem cells

[132628]express a new class of intermediate filament protein. Cell 60: 585–595,

[132629]1990.

[132630]23642.

[132631]Steinert, P. M.; Roop, D. R.: Molecular and cellular biology of

[132632]intermediate filaments. Ann. Rev. Biochem. 57: 593–625, 1988.

[132633]23643.

[132634]Waseem, A.; Gough, A. C.; Spurr, N. K.; Lane, E. B.: Local–

ization

[132635]of the gene for human simple epithelial keratin 18 to
chromosome 12

[132636]using polymerase chain reaction. Genomics 7: 188–194,
1990.

[132637]23644.

[132638]Erikson, J.; Griffin, C.; ar-Rushdi, A.; Valtieri, M.; Hoxie,

[132639]J.; Finan, J.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.;
Croce, C.

[132640]M.: Heterogeneity of chromosome 22 breakpoint in
Philadelphia-positive

[132641](Ph+) acute lymphocytic leukemia. Proc. Nat. Acad. Sci.
83: 1807–1811,

[132642]1986.

[132643]23645.

[132644]Fioretos, T.; Heisterkamp, N.; Groffen, J.: No evidence for
genomic

[132645]imprinting of the human BCR gene. Blood 83: 3441–3444,
1994.

[132646]23646.

[132647]Fitzgerald, P. H.: Evidence that chromosome band 22q12

is concerned

[132648]with cell proliferation in chronic myeloid leukemia. Hum.

Genet. 33:

[132649]269–274, 1976.

[132650]23647.

[132651]Ganesan, T. S.; Rassool, F.; Guo, A.-P.; Th'ng, K. H.;

Dowding,

[132652]C.; Hibbin, J. A.; Young, B. D.; White, H.; Kumaran, T. O.;

Galton,

[132653]D. A. G.; Goldman, J. M.: Rearrangement of the bcr gene in

Philadelphia

[132654]chromosome-negative chronic myeloid leukemia. Blood

68: 957–960,

[132655]1986.

[132656]23648.

[132657]Goldman, J. M.; Melo, J. V.: Targeting the BCR-ABL tyro-

sine kinase

[132658]in chronic myeloid leukemia. (Editorial) New Eng. J. Med.

344: 1084–1086,

[132659]2001.

[132660]23649.

[132661] Gorre, M. E.; Mohammed, M.; Ellwood, K.; Hsu, N.; Paquette, R.;

[132662] Rao, P. N.; Sawyers, C. L.: Clinical resistance to STI-571 cancer

[132663] therapy caused by BCR-ABL gene mutation or amplification. Science 293:

[132664] 876-883, 2001.

[132665] 23650.

[132666] Groffen, J.; Stephenson, J. R.; Heisterkamp, N.; de Klein, A.;

[132667] Bartram, C. R.; Grosveld, G.: Philadelphia chromosomal breakpoints

[132668] are clustered within a limited region, bcr, on chromosome 22. Cell 36:

[132669] 93-99, 1984.

[132670] 23651.

[132671] Grossman, A.; Silver, R. T.; Arlin, Z.; Coleman, M.; Camposano,

[132672] E.; Gascon, P.; Benn, P. A.: Fine mapping of chromosome 22 breakpoints

[132673] within the breakpoint cluster region (bcr) implies a role for bcr

[132674]lexon 3 in determining disease duration in chronic myeloid leukemia. Am.

[132675]J. Hum. Genet. 45: 729–738, 1989.

[132676]23652.

[132677]Grosveld, G.; Verwoerd, T.; van Agthoven, T.; de Klein, A.; Ramachandran,

[132678]K. L.; Heisterkamp, N.; Stam, K.; Groffen, J.: The chronic myelocytic

[132679]cell line K562 contains a breakpoint in bcr and produces a chimeric

[132680]bcr/c-abl transcript. Molec. Cell. Biol. 6: 607–616, 1986.

[132681]23653.

[132682]Haas, O. A.: Are ABL and BCR imprinted? No definitive answers,

[132683]but more questions. Leukemia 9: 740–745, 1995.

[132684]23654.

[132685]Haas, O. A.; Argyriou-Tirita, A.; Lion, T.: Parental origin of

[132686]chromosomes involved in the translocation t(9;22). Nature 359: 414–416,

[132687]1992.

[132688]23655.

[132689]Hariharan, I. K.; Adams, J. M.: cDNA sequence for human
bcr,

[132690]the gene that translocates to the abl oncogene in chronic
myeloid

[132691]leukaemia. EMBO J. 6: 115-119, 1987.

[132692]23656.

[132693]Heisterkamp, N.; Jenster, G.; ten Hoeve, J.; Zovich, D.; Pat-
tengale,

[132694]P. K.; Groffen, J.: Acute leukaemia in bcr/abl transgenic
mice. Nature 344:

[132695]251-253, 1990.

[132696]23657.

[132697]Heisterkamp, N.; Stam, K.; Groffen, J.; de Klein, A.;
Grosveld,

[132698]G.: Structural organization of the bcr gene and its role in
the Ph-1

[132699]translocation. Nature 315: 758-761, 1985.

[132700]23658.

[132701]Hermans, A.; Heisterkamp, N.; von Lindern, M.; van Baal,
S.; Meijer,

- [132702]D.; van der Plas, D.; Wiedemann, L. M.; Groffen, J.;
Bootsma, D.;
- [132703]Grosveld, G.: Unique fusion of bcr and c-abl genes in
Philadelphia
- [132704]chromosome positive acute lymphoblastic leukemia. Cell
51: 33-40,
- [132705]1987.
- [132706]23659.
- [132707]Huettner, C. S.; Zhang, P.; Van Etten, R. A.; Tenen, D. G.:
Reversibility
- [132708]of acute B-cell leukaemia induced by BCR-ABL1. Nature
Genet. 24:
- [132709]57-60, 2000.
- [132710]23660.
- [132711]Jacobs, A.: Benzene and leukemia. Brit. J. Haemat. 72:
119-121,
- [132712]1989.
- [132713]23661.
- [132714]Klein, G.: Specific chromosomal translocations and the
genesis
- [132715]of the B-cell-derived tumors in mice and men. Cell 32:

311-315,

[132716]1983.

[132717]23662.

[132718]Koeffler, H. P.; Golde, D. W.: Chronic myelogenous
leukemia--new

[132719]concepts. New Eng. J. Med. 304: 1201-1209 and
1269-1274, 1981.

[132720]23663.

[132721]Kohno, S.-I.; Sandberg, A. A.: Chromosomes and causa-
tion of human

[132722]cancer and leukemia. XXXIX. Usual and unusual findings in
Ph(1)-positive

[132723]CML. Cancer 46: 2227-2237, 1980.

[132724]23664.

[132725]Laurent, E.; Talpaz, M.; Kantarjian, H.; Kurzrock, R.: The
BCR

[132726]gene and Philadelphia chromosome-positive leukemogen-
esis. Cancer

[132727]Res. 61: 2343-2355, 2001.

[132728]23665.

[132729]Lillicrap, D. A.; Sterndale, H.: Familial chronic myeloid

leukaemia.

[132730](Letter) Lancet II: 699, 1984.

[132731]23666.

[132732]Lim, Y.-M.; Wong, S.; Lau, G.; Witte, O. N.; Colicelli, J.:

BCR/ABL

[132733]inhibition by an escort/phosphatase fusion protein. Proc.

Nat. Acad.

[132734]Sci. 97: 12233-12238, 2000.

[132735]23667.

[132736]Litz, C. E.; Copenhaver, C. M.: Paternal origin of the rear-

ranged

[132737]major breakpoint cluster region in chronic myeloid

leukemia. Blood 83:

[132738]3445-3448, 1994.

[132739]23668.

[132740]Maru, Y.; Witte, O. N.: The BCR gene encodes a novel ser-

ine/threonine

[132741]kinase activity within a single exon. Cell 67: 459-468,

1991.

[132742]23669.

[132743]Melo, J. V.; Yan, X.-H.; Diamond, J.; Goldman, J. M.: Lack

of

[132744]imprinting of the ABL gene. (Letter) Nature Genet. 8:
318–319, 1994.

[132745]23670.

[132746]Melo, J. V.; Yan, X.-H.; Diamond, J.; Goldman, J. M.: Bal-
anced

[132747]parental contribution to the ABL component of the BCR-
ABL gene in

[132748]chronic myeloid leukemia. Leukemia 9: 734–745, 1995.

[132749]23671.

[132750]Mes-Masson, A.-M.; McLaughlin, J.; Daley, G. Q.; Paskind,
M.;

[132751]Witte, O. N.: Overlapping cDNA clones define the complete
coding

[132752]region for the P210(c-abl) gene product associated with
chronic myelogeneous

[132753]leukemia cells containing the Philadelphia chromosome.
Proc. Nat.

[132754]Acad. Sci. 83: 9768–9772, 1986.

[132755]23672.

[132756]Mills, K. I.; MacKenzie, E. D.; Birnie, G. D.: The site of the

[132757]breakpoint within the bcr is a prognostic factor in
Philadelphia-positive

[132758]CML patients. Blood 72: 1237-1241, 1988.

[132759]23673.

[132760]Mittelman, F.; Levan, G.: Clustering of aberrations to specific

[132761]chromosomes in human neoplasms. III. Incidence and geographic distribution

[132762]of chromosome aberrations in 856 cases. Hereditas 89:
207-232, 1978.

[132763]23674.

[132764]Nowell, P. C.; Hungerford, D. A.: A minute chromosome in
human

[132765]chronic granulocytic leukemia. Science 132: 1497, 1960.

[132766]23675.

[132767]Kusari, J.; Takata, Y.; Hatada, E.; Freidenberg, G.; Kolterman,

[132768]O.; Olefsky, J. M.: Insulin resistance and diabetes due to
different

[132769]mutations in the tyrosine kinase domain of both insulin
receptor gene

[132770]alleles. J. Biol. Chem. 266: 5260–5267, 1991.

[132771]23676.

[132772]Lander, E. S.; Botstein, D.: Homozygosity mapping: a way
to map

[132773]human recessive traits with the DNA of inbred children.
Science 236:

[132774]1567–1570, 1987.

[132775]23677.

[132776]Sagane, K.; Ohya, Y.; Hasegawa, Y.; Tanaka, I.: Metallo-
proteinase-like,

[132777]disintegrin-like, cysteine-rich proteins MDC2 and MDC3:
novel human

[132778]cellular disintegrins highly expressed in the brain.
Biochem. J. 334:

[132779]93–98, 1998.

[132780]23678.

[132781]Splendore, A.; Passos-Bueno, M. R.; Jabs, E. W.; Van
Maldergem,

[132782]L.; Wulfsberg, E. A.: TCOF1 mutations excluded from a
role in other

[132783]first and second branchial arch-related disorders. (Letter)

Am. J.

[132784]Med. Genet. 111: 324–327, 2002.

[132785]23679.

[132786]Alsobrook, J. P., II; Zohar, A. H.; Leboyer, M.; Chabane, N.;
Ebstein,

[132787]R. P.; Pauls, D. L.: Association between the COMT locus
and obsessive–compulsive

[132788]disorder in females but not males. Am. J. Med. Genet.
(Neuropsychiat.

[132789]Genet.) 114: 116–120, 2002.

[132790]23680.

[132791]Karayiorgou, M.; Sobin, C.; Blundell, M. L.; Galke, B. L.;
Malinova,

[132792]L.; Goldberg, P.; Ott, J.; Gogos, J. A.: Family–based associ-
ation

[132793]studies support a sexually dimorphic effect of COMT and
MAOA on genetic

[132794]susceptibility to obsessive–compulsive disorder. Biol. Psy-
chiat. 45:

[132795]1178–1189, 1999.

[132796]23681.

[132797]Taira, M.; Yoshida, T.; Miyagawa, K.; Sakamoto, H.; Terada, M.;

[132798]Sugimura, T.: cDNA sequence of human transforming gene hst and identification

[132799]of the coding sequence required for transforming activity. Proc.

[132800]Nat. Acad. Sci. 84: 2980–2984, 1987.

[132801]23682.

[132802]Allikmets, R.; Shroyer, N. F.; Singh, N.; Seddon, J. M.; Lewis,

[132803]R. A.; Bernstein, P. S.; Peiffer, A.; Zabriskie, N. A.; Hutchinson,

[132804]A.; Dean, M.; Lupski, J. R.; Leppert, M.: Mutation of the Stargardt

[132805]disease gene (ABCR) in age-related macular degeneration. Science 277:

[132806]1805–1807, 1997.

[132807]23683.

[132808]De La Paz, M. A.; Guy, V. K.; Abou-Donia, S.; Heinis, R.; Bracken,

[132809]B.; Vance, J. M.; Gilbert, J. R.; Gass, J. D. M.; Haines, J. L.; Pericak-Vance,

- [132810]M. A.: Analysis of the Stargardt disease gene (ABCR) in age-related
macular degeneration. Ophthalmology 106: 1531–1536, 1999.
- [132811]23684.
- [132812]Farndon, J. R.; Leight, G. S.; Dilley, W. G.; Baylin, S. B.; Smallridge,
R. C.; Harrison, T. S.; Wells, S. A., Jr.: Familial medullary thyroid
carcinoma without associated endocrinopathies: a distinct clinical
entity. Brit. J. Surg. 73: 278–281, 1986.
- [132813]23685.
- [132814]Taylor, M. D.; Liu, L.; Raffel, C.; Hui, C.; Mainprize, T. G.; Zhang, X.; Agatep, R.; Chiappa, S.; Gao, L.; Lowrance, A.; Hao, A.; Goldstein, A. M.; Stavrou, T.; Scherer, S. W.; Dura, W. T.; Wainwright,
B.; Squire, J. A.; Rutka, J. T.; Hogg, D.: Mutations in SUFU predispose
to medulloblastoma. Nature Genet. 31: 306–310, 2002.

[132823]23686.

[132824]Ball, E. W.; Meynell, M. J.; Beale, D.; Kynoch, P.; Lehmann, H.;

[132825]Strelton, A. O. W.: Haemoglobin alpha(2) prime: alpha 2 gamma 2 (16

[132826]glycine to arginine). Nature 209: 1217–1218, 1966.

[132827]23687.

[132828]Barnabas, J.; Muller, C. J.: Haemoglobin Lepore (Hollandia). Nature 194:

[132829]931–932, 1962.

[132830]23688.

[132831]Beaven, G. H.; Gratzner, W. B.; Stevens, B. L.; Shooter, E. M.;

[132832]Ellis, M. J.; White, J. C.; Gillespie, J. E. O.: An abnormal haemoglobin

[132833](Lepore–Cyprus) resembling haemoglobin Lepore and its interaction

[132834]with thalassaemia. Brit. J. Haemat. 10: 159–170, 1964.

[132835]23689.

[132836]Brennan, S. O.; Williamson, D.; Smith, M. B.; Cauchi, M. N.; Macphee,

[132837]A.; Carrell, R. W.: HbA(2) Victoria, delta24 (B6) gly-to-asp:
a new

[132838]delta chain variant occurring with beta-thalassemia.
Hemoglobin 8:

[132839]163-168, 1984.

[132840]23690.

[132841]Camaschella, C.; Alfarano, A.; Gottardi, E.; Travi, M.; Prim-
ignani,

[132842]P.; Cappio, F. C.; Saglio, G.: Prenatal diagnosis of fetal
hemoglobin

[132843]Lepore-Boston disease on maternal peripheral blood.
Blood 75: 2102-2106,

[132844]1990.

[132845]23691.

[132846]Codrington, J. F.; Kutlar, F.; Harris, H. F.; Wilson, J. B.;
Stoming,

[132847]T. A.; Huisman, T. H. J.: Hb A(2)-Wrens or
delta98(FG5)val-to-met,

[132848]an unstable delta-chain variant identified by sequence
analysis of

[132849]amplified DNA. Biochim. Biophys. Acta 1009: 87-89,
1989.

[132850]23692.

[132851]Curtain, C. C.: A structural study of abnormal
haemoglobins occurring

[132852]in New Guinea. Aust. J. Exp. Biol. Med. Sci. 42: 89–97,
1964.

[132853]23693.

[132854]De Angioletti, M.; Di Girgenti, C.; Messineo, R.; Capra, M.;
Carestia,

[132855]C.: Hb A2–Monreale (delta–146 (HC3) his–to–arg), a novel
delta chain

[132856]variant detected in west Sicily. Hemoglobin 26: 1–5, 2002.

[132857]23694.

[132858]De Jong, W. W.; Went, L. N.: Hemoglobin A(2)–NYU in the
Netherlands:

[132859]incidence of delta–chain variants in human populations.
Hum. Hered. 24:

[132860]32–39, 1974.

[132861]23695.

[132862]De Jong, W. W. W.; Bernini, L. F.: Haemoglobin Babinga
(delta

[132863]136 glycine–aspartic acid): a new delta chain variant. Na–

ture 219:

[132864]1360–1362, 1968.

[132865]23696.

[132866]Efremov, G. D.: Hemoglobins Lepore and anti-Lepore.

Hemoglobin 2:

[132867]197–233, 1978.

[132868]23697.

[132869]McMichael, A.; Makgoba, W.: Complexity in human histocompatibility

[132870]loci. Nature 293: 701–702, 1981.

[132871]23698.

[132872]Moen, T.; Albrechtsen, D.; Flatmark, A.; Jakobsen, A.; Jervell,

[132873]J.; Halvorsen, S.; Solheim, B. G.; Thorsby, E.: Importance of HLA-DR

[132874]matching in cadaveric renal transplantation: a prospective one-center

[132875]study of 170 transplants. New Eng. J. Med. 303: 850–854, 1980.

[132876]23699.

[132877]Olerup, O.; Troye-Blomberg, M.; Schreuder, G. M. T.; Ri-

ley, E.

[132878]M.: HLA-DR and -DQ gene polymorphism in West Africans
is twice as

[132879]extensive as in North European Caucasians: evolutionary
implications. Proc.

[132880]Nat. Acad. Sci. 88: 8480-8484, 1991.

[132881]23700.

[132882]Park, M. S.; Terasaki, P. I.; Bernoco, D.: Relationship be-
tween

[132883]MT and DR antigens.In: Terasaki, P. I.: Histocompatibility
Testing

[132884]1980. Los Angeles: UCLA Press (pub.) 1980.

[132885]23701.

[132886]Park, M. S.; Terasaki, P. I.; Bernoco, D.; Iwaki, Y.: Evidence

[132887]for a second B-cell locus separate from the DR locus.
Transplant.

[132888]Proc. 10: 823-828, 1978.

[132889]23702.

[132890]Pisella, P.-J.; Brignole, F.; Debbasch, C.; Lozato, P.-A.;
Creuzot-Garcher,

[132891]C.; Bara, J.; Saiag, P.; Warnet, J.-M.; Baudouin, C.: Flow cy-

tometric

[132892]analysis of conjunctival epithelium in ocular rosacea and
keratoconjunctivitis

[132893]sicca. Ophthalmology 107: 1841-1849, 2000.

[132894]23703.

[132895]Rollini, P.; Mach, B.; Gorski, J.: Linkage map of three HLA-
DR

[132896]beta-chain genes: evidence for a recent duplication event.
Proc.

[132897]Nat. Acad. Sci. 82: 7197-7201, 1985.

[132898]23704.

[132899]Sachs, J. A.; Jaraquemada, D.; Festenstein, H.: Intra HLA-D
region

[132900]recombinant maps HLA-DR between HLA-B and HLA-D.
Tissue Antigens 17:

[132901]43-56, 1981.

[132902]23705.

[132903]Schwartz, B. D.: Personal Communication. St. Louis, Mo.
2/22/1983.

[132904]23706.

[132905]Shackelford, D. A.; Mann, D. L.; van Rood, J. J.; Ferrara, G.

[132906]B.; Strominger, J. L.: Human B-cell alloantigens DC1, MT1,
and LB12

[132907]are identical to each other but distinct from the HLA-DR
antigen. Proc.

[132908]Nat. Acad. Sci. 78: 4566-4570, 1981.

[132909]23707.

[132910]Strominger, J. L.: Biology of the human histocompatibility
leukocyte

[132911]antigen (HLA) system and a hypothesis regarding the gen-
eration of

[132912]autoimmune diseases. J. Clin. Invest. 77: 1411-1415,
1986.

[132913]23708.

[132914]Suciu-Foca, N.; Weiner, J.; Rohowsky, C.; McKiernan, P.;
Susinno,

[132915]E.; Rubinstein, P.: Indications that Dw determinants are
controlled

[132916]by distinct (but closely linked) genes. Transplant. Proc. 10:
799-804,

[132917]1978.

[132918]23709.

[132919]Thursz, M. R.; Thomas, H. C.; Greenwood, B. M.; Hill, A. V.
S.

[132920]Heterozygote advantage for HLA class-II type in hepatitis
B virus

[132921]infection. (Letter) Nature Genet. 17: 11-12, 1997.

[132922]23710.

[132923]Tiercy, J.-M.; Gorski, J.; Jeannet, M.; Mach, B.: Identifica-
tion

[132924]and distribution of three serologically undetected alleles
of HLA-DR

[132925]by oligonucleotide-DNA typing analysis. Proc. Nat. Acad.
Sci. 85:

[132926]198-202, 1988.

[132927]23711.

[132928]Tiwari, J. L.; Terasaki, P. I.: HLA and Disease Associations.

[132929]New York: Springer-Verlag (pub.) 1985.

[132930]23712.

[132931]Trowsdale, J.: Genetics and polymorphism: class II anti-
gens. Brit.

[132932]Med. Bull. 43: 15-36, 1987.

[132933]23713.

[132934]Tsubota, K.; Fukagawa, K.; Fujihara, T.; Shimmura, S.;
Saito,

[132935]I.; Saito, K.; Takeuchi, T.: Regulation of human leukocyte
antigen

[132936]expression in human conjunctival epithelium. Invest.
Ophthalm. Vis.

[132937]Sci. 40: 28–34, 1999.

[132938]23714.

[132939]Walker, L. E.; Hewick, R.; Hunkapiller, M. W.; Hood, L. E.;
Dreyer,

[132940]W. J.; Reisfeld, R. A.: N-terminal amino acid sequences of
the alpha

[132941]and beta chains of HLA-DR1 and HLA-DR2 antigens. Bio-
chemistry 23:

[132942]185–188, 1983.

[132943]23715.

[132944]Spelbrink, J. N.; Li, F.-Y.; Tiranti, V.; Nikali, K.; Yuan,
Q.-P.;

[132945]Tariq, M.; Wanrooij, S.; Garrido, N.; Comi, G.; Morandi, L.;
Santoro,

[132946]L.; Toscano, A.; and 9 others: Human mitochondrial DNA deletions

[132947]associated with mutations in the gene encoding twinkle, a phage T7

[132948]gene 4-like protein localized in mitochondria. Nature Genet. 28:

[132949]223-231, 2001.

[132950]23716.

[132951]Farfel, Z.; Iiri, T.; Shapira, H.; Roitman, A.; Mouallem, M.;

[132952]Bourne, H. R.: Pseudohypoparathyroidism, a novel mutation in the

[132953]beta/gamma- contact region of Gs-alpha impairs receptor stimulation. J.

[132954]Biol. Chem. 271: 19653-19655, 1996.

[132955]23717.

[132956]Heisler, L. K.; Cowley, M. A.; Tecott, L. H.; Fan, W.; Low, M.

[132957]J.; Smart, J. L.; Rubinstein, M.; Tatro, J. B.; Marcus, J. N.; Holstege,

[132958]H.; Lee, C. E.; Cone, R. D.; Elmquist, J. K.: Activation of central

[132959]melanocortin pathways by fenfluramine. Science 297:

609–611, 2002.

[132960]23718.

[132961]Folster–Holst, R.; Moises, H. W.; Yang, L.; Fritsch, W.;
Weissenbach,

[132962]J.; Christophers, E.: Linkage between atopy and the IgE
high–affinity

[132963]receptor gene at 11q13 in atopic dermatitis families.
Hum. Genet. 102:

[132964]236–239, 1998.

[132965]23719.

[132966]Hill, M. R.; Cookson, W. O. C. M.: A new variant of the beta
subunit

[132967]of the high–affinity receptor for immunoglobulin E
(Fc–epsilon–RI–beta

[132968]E237G): associations with measures of atopy and
bronchial hyper–responsiveness. Hum.

[132969]Molec. Genet. 5: 959–962, 1996.

[132970]23720.

[132971]Hupp, K.; Siwarski, D.; Mock, B. A.; Kinet, J.–P.: Gene map–
ping

[132972]of the three subunits of the high affinity FcR for IgE to

mouse chromosomes

[132973]1 and 19. J. Immun. 143: 3787–3791, 1989.

[132974]23721.

[132975]Kuster, H.; Zhang, L.; Brini, A. T.; MacGlashan, D. W. J.;
Kinet,

[132976]J.-P.: The gene and cDNA for the human high affinity im-
munoglobulin

[132977]E receptor beta chain and expression of the complete hu-
man receptor. J.

[132978]Biol. Chem. 267: 12782–12787, 1992.

[132979]23722.

[132980]Nagata, H.; Mutoh, H.; Kumahara, K.; Arimoto, Y.;
Tomemori, T.;

[132981]Sakurai, D.; Arase, K.; Ohno, K.; Yamakoshi, T.; Nakano,
K.; Okawa,

[132982]T.; Numata, T.; Konno, A.: Association between nasal al-
lergy and

[132983]a coding variant of the Fc-epsilon-RI-beta gene
Glu237Gly in a Japanese

[132984]population. Hum. Genet. 109: 262–266, 2001.

[132985]23723.

[132986] Shirakawa, T.; Li, A.; Dubowitz, M.; Dekker, J. W.; Shaw, A. E.;

[132987] Faux, J. A.; Ra, C.; Cookson, W. O. C. M.; Hopkin, J. M.: Association

[132988] between atopy and variants of the beta subunit of the high-affinity

[132989] immunoglobulin E receptor. *Nature Genet.* 7: 125–130, 1994.

[132990] 23724.

[132991] Shirakawa, T.; Mao, X.-Q.; Sasaki, S.; Enomoto, T.; Kawai, M.;

[132992] Morimoto, K.; Hopkin, J.: Association between atopic asthma and a

[132993] coding variant of Fc-epsilon-RI-beta in a Japanese population. *Hum.*

[132994] *Molec. Genet.* 5: 1129–1130, 1996.

[132995] 23725.

[132996] Szepietowski, P.; Gaudray, P.: FCER1B, a candidate gene for atopy,

[132997] is located in 11q13 between CD20 and TCN1. *Genomics* 19: 399–400,

[132998] 1994.

[132999]23726.

[133000]Allen, G.; Fantes, K. H.: A family of structural genes for
human

[133001]lymphoblastoid (leucocyte-type) interferon. Nature 287:
408–411,

[133002]1980.

[133003]23727.

[133004]Diaz, M. O.; Bohlander, S.; Allen, G.: Nomenclature of the
human

[133005]interferon genes. J. Interferon Cytokine Res. 16: 179–180,
1996.

[133006]23728.

[133007]Diaz, M. O.; Pomykala, H. M.; Bohlander, S. K.; Maltepe, E.;
Malik,

[133008]K.; Brownstein, B.; Olopade, O. I.: Structure of the human
type-I

[133009]interferon gene cluster determined from a YAC clone con-
tig. Genomics 22:

[133010]540–552, 1994.

[133011]23729.

[133012]Douglas, R. M.; Moore, B. W.; Miles, H. B.; Davies, L. M.;

Graham,

[133013]N. M. H.; Ryan, P.; Worswick, D. A.; Albrecht, J. K.: Prophylactic

[133014]efficacy of intranasal alpha-2-interferon against rhinovirus infections

[133015]in the family setting. New Eng. J. Med. 314: 65-70, 1986.

[133016]23730.

[133017]Edge, M. D.; Green, A. R.; Heathcliffe, G. R.; Meacock, P. A.;

[133018]Schuch, W.; Scanlon, D. B.; Atkinson, T. C.; Newton, C. R.; Markham,

[133019]A. F.: Total synthesis of a human leukocyte interferon gene. Nature 292:

[133020]756-762, 1981.

[133021]23731.

[133022]Fountain, J. W.; Karayiorgou, M.; Taruscio, D.; Graw, S. L.; Buckler,

[133023]A. J.; Ward, D. C.; Dracopoli, N. C.; Housman, D. E.: Genetic and

[133024]physical map of the interferon region on chromosome 9p. Genomics 14:

[133025]105-112, 1992.

[133026]23732.

[133027]Gillespie, D.; Carter, W.: Concerted evolution of human interferon

[133028]alpha genes. J. Interferon Res. 3: 83-88, 1983.

[133029]23733.

[133030]Hayden, F. G.; Albrecht, J. K.; Kaiser, D. L.; Gwaltney, J. M.,

[133031]Jr.: Prevention of natural colds by contact prophylaxis with intranasal

[133032]alpha-2-interferon. New Eng. J. Med. 314: 71-75, 1986.

[133033]23734.

[133034]Hitzeman, R. A.; Hagie, F. E.; Levine, H. L.; Goeddel, D. V.;

[133035]Ammerer, G.; Hall, B. D.: Expression of a human gene for interferon

[133036]in yeast. Nature 293: 717-722, 1981.

[133037]23735.

[133038]Imai, M.; Sano, T.; Yanase, Y.; Miyamoto, K.; Yonehara, S.; Mori,

[133039]H.; Honda, T.; Fukuda, S.; Nakamura, T.; Miyakawa, Y.; Mayumi, M.

[133040]: Demonstration of two subtypes of human leukocyte in-

terferon (IFN- α)

[133041]by monoclonal antibodies. J. Immun. 128: 2824-2825,
1982.

[133042]23736.

[133043]Isaacs, D.; Clarke, J. R.; Tyrrell, D. A. J.; Webster, A. D. B.;

[133044]Valman, H. B.: Deficient production of leucocyte interferon
(interferon- α)

[133045]in vitro and in vivo in children with recurrent respiratory
tract

[133046]infections. Lancet II: 950-952, 1981.

[133047]23737.

[133048]Lawn, R. M.; Adelman, J.; Dull, T. J.; Gross, M.; Goeddel,
D.;

[133049]Ullrich, A.: DNA sequence of two closely linked human
leukocyte interferon

[133050]genes. Science 212: 1159-1162, 1981.

[133051]23738.

[133052]Lawn, R. M.; Goeddel, D. V.; Ullrich, A.: The human inter-
feron

[133053]gene family.(Abstract) Sixth Int. Cong. Hum. Genet.,
Jerusalem 55

[133054]only, 1981.

[133055]23739.

[133056]Miyata, T.; Hayashida, H.: Recent divergence from a common ancestor

[133057]of human IFN- α genes. Nature 295: 165-168, 1982.

[133058]23740.

[133059]Mory, Y.; Chernajovsky, Y.; Feinstein, S. I.; Chen, L.; Weissenbach,

[133060]J.; Revel, M.: Expression of the cloned human interferon beta-1 gene

[133061]in E. coli.(Abstract) Sixth Int. Cong. Hum. Genet., Jerusalem 56

[133062]only, 1981.

[133063]23741.

[133064]Reis, A.: Personal Communication. Berlin, Germany 3/28/1994.

[133065]23742.

[133066]Blanchet-Bardon, C.; Nazzaro, V.; Chevrant-Breton, J.; Espie, M.;

[133067]Kerbrat, P.; Le Marec, B.: Hereditary epidermolytic palmo-plantar

[133068]keratoderma associated with breast and ovarian cancer in a large kindred. Brit.

[133069]J. Derm. 117: 363-370, 1987.

[133070]23743.

[133071]Appel, S.; Filter, M.; Reis, A.; Hennies, H. C.; Bergheim, A.;

[133072]Ogilvie, E.; Arndt, S.; Simmons, A.; Lovett, M.; Hide, W.; Ramsay,

[133073]M.; Reichwald, K.; Zimmermann, W.; Rosenthal, A.: Physiological and transcriptional

[133074]map of the critical region for keratolytic winter erythema (KWE) on

[133075]chromosome 8p22-p23 between D8S550 and D8S1759. Europ. J. Hum. Genet. 10:

[133076]17-25, 2002.

[133077]23744.

[133078]Clarke, C. A.; McConnell, R. B.: Six cases of carcinoma of the

[133079]oesophagus occurring in one family. Brit. Med. J. 2: 1137-1138,

[133080]1954.

[133081]23745.

[133082]Ellis, A.; Field, J. K.; Field, E. A.; Friedmann, P. S.; Fryer,
[133083]A.; Howard, P.; Leigh, I. M.; Risk, J.; Shaw, J. M.; Whittaker,
J.
[133084]: Tylosis associated with carcinoma of the oesophagus
and oral leukoplakia
[133085]in a large Liverpool family--a review of six generations.
Europ.
[133086]J. Cancer 30B: 102-112, 1994.
[133087]23746.
[133088]Harper, P. S.; Harper, R. M. J.; Howel-Evans, A. W.: Carci-
noma
[133089]of the oesophagus with tylosis. Quart. J. Med. 39:
317-333, 1970.
[133090]23747.
[133091]Olavarria, E.; Craddock, C.; Dazzi, F.; Marin, D.; Marktel,
S.;
[133092]Apperley, J. F.; Goldman, J. M.: Imatinib mesylate (STI571)
in the
[133093]treatment of relapse of chronic myeloid leukemia after al-
logeneic
[133094]stem cell transplantation. Blood 99: 3861-3862, 2002.

[133095]23748.

[133096]Pegoraro, L.; Matera, L.; Ritz, J.; Levis, A.; Palumbo, A.; Biagini,

[133097]G.: Establishment of a Ph(1)-positive human cell line (BV173). J.

[133098]Nat. Cancer Inst. 70: 447-451, 1983.

[133099]23749.

[133100]Perrotti, D.; Cesi, V.; Trotta, R.; Guerzoni, C.; Santilli, G.;

[133101]Campbell, K.; Iervolino, A.; Condorelli, F.; Gambacorti-Passerini,

[133102]C.; Caligiuri, M. A.; Calabretta, B.: BCR-ABL suppresses C/EBP-alpha

[133103]expression through inhibitory action of hnRNP E2. Nature Genet. 30:

[133104]48-58, 2002.

[133105]23750.

[133106]Prakash, O.; Yunis, J. J.: High resolution chromosomes of the

[133107]t(9;22) positive leukemias. Cancer Genet. Cytogenet. 11: 361-367,

[133108]1984.

[133109]23751.

[133110]Priest, J. R.; Robison, L. L.; McKenna, R. W.; Lindquist, L.
L.;

[133111]Warkentin, P. I.; LeBien, T. W.; Woods, W. G.; Kersey, J. H.;
Coccia,

[133112]P. F.; Nesbit, M. E., Jr.: Philadelphia chromosome positive
childhood

[133113]acute lymphoblastic leukemia. Blood 56: 15-22, 1980.

[133114]23752.

[133115]Rowley, J. D.: A new consistent chromosomal abnormality
in chronic

[133116]myelogenous leukemia identified by quinacrine fluores-
cence and Giemsa

[133117]staining. Nature 243: 290-293, 1973.

[133118]23753.

[133119]Rubin, C. M.; Carrino, J. J.; Dickler, M. N.; Leibowitz, D.;
Smith,

[133120]S. D.; Westbrook, C. A.: Heterogeneity of genomic fusion
of BCR and

[133121]ABL in Philadelphia chromosome-positive acute lym-
phoblastic leukemia. Proc.

[133122]Nat. Acad. Sci. 85: 2795–2799, 1988.

[133123]23754.

[133124]Saglio, G.; Storlazzi, C. T.; Giugliano, E.; Surace, C.; Anelli,

[133125]L.; Rege-Cambrin, G.; Zagaria, A.; Velasco, A. J.; Heiniger, A.; Scaravaglio,

[133126]P.; Gomez, A. T.; Gomez, J. R.; Archidiacono, N.; Banfi, S.; Rocchi,

[133127]M.: A 76-kb duplicon maps close to the BCR gene on chromosome 22

[133128]and the ABL gene on chromosome 9: possible involvement in the genesis

[133129]of the Philadelphia chromosome translocation. Proc. Nat. Acad. Sci. 99:

[133130]9882–9887, 2002.

[133131]23755.

[133132]Savage, D. G.; Antman, K. H.: Imatinib mesylate—a new oral targeted

[133133]therapy. New Eng. J. Med. 346: 683–693, 2002.

[133134]23756.

[133135]Sawyers, C. L.: Chronic myeloid leukemia. New Eng. J. Med. 340:

[133136]1330–1340, 1999.

[133137]23757.

[133138]Schaefer–Rego, K.; Dudek, H.; Popenoe, D.; Arlin, Z.;
Mears, J.

[133139]G.; Bank, A.; Leibowitz, D.: CML patients in blast crisis
have breakpoints

[133140]localized to a specific region of the BCR. Blood 70:
448–455, 1987.

[133141]23758.

[133142]Shtivelman, E.; Gale, R. P.; Drazan, O.; Berrebi, A.; Zaizov,

[133143]R.; Kubonishi, I.; Miyoshi, I.; Canaani, E.: bcr–abl RNA in
patients

[133144]with chronic myelogenous leukemia. Blood 69: 971–973,
1987.

[133145]23759.

[133146]Shtivelman, E.; Lifshitz, B.; Gale, R. P.; Canaani, E.: Fused

[133147]transcript of abl and bcr genes in chronic myelogenous
leukaemia. Nature 315:

[133148]550–554, 1985.

[133149]23760.

[133150]Skorski, T.; Nieborowska-Skorska, M.; Nicolaides, N. C.;
 Szczalik,
 [133151]C.; Iversen, P.; Iozzo, R. V.; Zon, G.; Calabretta, B.: Sup-
 pression
 [133152]of Philadelphia-1 leukemia cell growth in mice by BCR-
 ABL antisense
 [133153]oligodeoxynucleotide. Proc. Nat. Acad. Sci. 91:
 4504-4508, 1994.
 [133154]23761.

[133155]Stam, K.; Heisterkamp, N.; Grosveld, G.; de Klein, A.;
 Verma,
 [133156]R. S.; Coleman, M.; Dosik, H.; Groffen, J.: Evidence of a
 new chimeric
 [133157]bcr/c-abl mRNA in patients with chronic myelocytic
 leukemia and the
 [133158]Philadelphia chromosome. New Eng. J. Med. 313:
 1429-1433, 1985.
 [133159]23762.

[133160]Stam, K.; Heisterkamp, N.; Reynolds, F. H., Jr.; Groffen, J.:
 [133161]Evidence that the phl gene encodes a 160,000-dalton
 phosphoprotein
 [133162]with associated kinase activity. Molec. Cell. Biol. 7:

1955–1960,

[133163]1987.

[133164]23763.

[133165]Swan, D. C.; McBride, O. W.; Robbins, K. C.; Keithley, D. A.;

[133166]Reddy, E. P.; Aaronson, S. A.: Chromosomal mapping of
the simian

[133167]sarcoma virus onc gene analogue in human cells. Proc.
Nat. Acad.

[133168]Sci. 79: 4691–4695, 1982.

[133169]23764.

[133170]Tanabe, T.; Kuwabara, T.; Warashina, M.; Tani, K.; Taira,
K.;

[133171]Asano, S.: Oncogene inactivation in a mouse model: tissue
invasion

[133172]by leukaemic cells is stalled by loading them with a de-
signer ribozyme. Nature 406:

[133173]473–474, 2000.

[133174]23765.

[133175]Teyssier, J. R.; Bartram, C. R.; Deville, J.; Potron, G.; Pi-
geon,

[133176]F.: C-abl oncogene and chromosome 22 'bcr' juxtaposi-

tion in chronic

[133177]myelogenous leukemia. New Eng. J. Med. 312:

1393–1394, 1985.

[133178]23766.

[133179]Tkachuk, D. C.; Westbrook, C. A.; Andreeff, M.; Donlon, T.

A.;

[133180]Cleary, M. L.; Suryanarayan, K.; Homge, M.; Redner, A.;

Gray, J.;

[133181]Pinkel, D.: Detection of bcr–abl fusion in chronic myeloge-
neous leukemia

[133182]by in situ hybridization. Science 250: 559–562, 1990.

[133183]23767.

[133184]Verhest, A.; Monsieur, R.: Philadelphia chromosome–posi-
tive thrombocythemia

[133185]with leukemic transformation. (Letter) New Eng. J. Med.

308: 1603,

[133186]1983.

[133187]23768.

[133188]Verma, R. S.; Dosik, H.: Heteromorphisms of the Philadel-
phia

[133189](Ph–1) chromosome in patients with chronic myelogenous

leukaemia (CML).

[133190]l. Classification and clinical significance. Brit. J. Haemat.

45:

[133191]215-222, 1980.

[133192]23769.

[133193]Aisenberg, A. C.; Wilkes, B. M.; Jacobson, J. O.: The bcl-2
gene

[133194]is rearranged in many diffuse B-cell lymphomas. Blood
71: 969-972,

[133195]1988.

[133196]23770.

[133197]Bakhshi, A.; Jensen, J. P.; Goldman, P.; Wright, J. J.;
McBride,

[133198]O. W.; Epstein, A. L.; Korsmeyer, S. J.: Cloning the chro-
mosomal

[133199]breakpoint of t(14;18) human lymphomas: clustering
around J(H) on

[133200]chromosome 14 and near a transcriptional unit on 18. Cell
41: 899-906,

[133201]1985.

[133202]23771.

[133203]Bakhshi, A.; Wright, J. J.; Graninger, W.; Seto, M.; Owens, J.;

[133204]Cossman, J.; Jensen, J. P.; Goldman, P.; Korsmeyer, S. J.:
Mechanism

[133205]of the t(14;18) chromosomal translocation: structural
analysis of

[133206]both derivative 14 and 18 reciprocal partners. Proc. Nat.
Acad. Sci. 84:

[133207]2396–2400, 1987.

[133208]23772.

[133209]Chen, J.; Flannery, J. G.; LaVail, M. M.; Steinberg, R. H.; Xu,

[133210]J.; Simon, M. I.: Bcl-2 overexpression reduces apoptotic
photoreceptor

[133211]cell death in three different retinal degenerations. Proc.
Nat. Acad.

[133212]Sci. 93: 7042–7047, 1996.

[133213]23773.

[133214]Cleary, M. L.; Sklar, J.: Nucleotide sequence of a t(14;18)
chromosomal

[133215]breakpoint in follicular lymphoma and demonstration of a
breakpoint–cluster

[133216]region near a transcriptionally active locus on chromosome 18. Proc.

[133217]Nat. Acad. Sci. 82: 7439-7443, 1985.

[133218]23774.

[133219]Cleary, M. L.; Smith, S. D.; Sklar, J.: Cloning and structural

[133220]analysis of cDNAs for bcl-2 and a hybrid bcl-2/immunoglobulin transcript

[133221]resulting from the t(14;18) translocation. Cell 47: 19-28, 1986.

[133222]23775.

[133223]Cockerham, G. C.; Hidayat, A. A.; Bijwaard, K. E.; Sheng, Z.-M.

[133224]: Re-evaluation of 'reactive lymphoid hyperplasia of the uvea': an

[133225]immunohistochemical and molecular analysis of 10 cases. Ophthalmology 107:

[133226]151-158, 2000.

[133227]23776.

[133228]Croce, C. M.: Personal Communication. Philadelphia, Pa. 1989.

[133229]23777.

[133230]Efremov, G. D.; Rudivic, R.; Niazi, G. A.; Hunter, E., Jr.;
Huisman,

[133231]T. H. J.; Schroeder, W. A.: An individual with Hb-Lep-
ore-Baltimore-delta

[133232]beta-thalassaemia in a Yugoslavian family. Scand. J.
Haemat. 16:

[133233]81-89, 1976.

[133234]23778.

[133235]Fessas, P.; Stamatoyannopoulos, G.; Karaklis, A.:
Hemoglobin

[133236]'Pylos': study of a hemoglobinopathy resembling tha-
lassemia in the

[133237]heterozygous, homozygous and double heterozygous
state. Blood 19:

[133238]1-22, 1962.

[133239]23779.

[133240]Fioretti, G.; De Angioletti, M.; Masciangelo, F.; Lacerra, G.;

[133241]Scarallo, A.; de Bonis, C.; Pagano, L.; Guarino, E.; De Rosa,
L.;

[133242]Salvati, F.; Carestia, C.: Origin heterogeneity of Hb Lep-
ore-Boston

[133243]gene in Italy. Am. J. Hum. Genet. 50: 781–786, 1992.

[133244]23780.

[133245]Fujita, S.; Ohta, Y.; Saito, S.; Kobayashi, Y.; Naritomi, Y.;

[133246]Kawaguchi, T.; Imamura, T.; Wada, Y.; Hayashi, A.:

Hemoglobin A(2)

[133247]Honai (delta90(F6) glu-to-val): a new delta chain variant.

Hemoglobin 9:

[133248]597–607, 1985.

[133249]23781.

[133250]Galanello, R.; Gasperini, D.; Perseu, L.; Barella, S.; Ideo, A.;

[133251]Cao, A.: Hb A(2)–Sant' Antioco (delta93(F9) cys-to-gly): a
new delta

[133252]chain variant identified by sequencing of amplified DNA.

Hemoglobin 18:

[133253]437–439, 1994.

[133254]23782.

[133255]Gasperini, D.; Perseu, L.; Cossu, P.; Podda, R.; Cao, A.;

Galanello,

[133256]R.: A novel delta-0-thalassemia mutation: TCG-to-TAG
(trp-to-stop)

[133257]at codon 37. Hum. Mutat. 3: 71–72, 1994.

[133258]23783.

[133259]Gerald, P. S.; Diamond, L. K.: A new hereditary
hemoglobinopathy

[133260](the Lepore trait) and its interaction with thalassemia trait.
Blood 12:

[133261]835–844, 1958.

[133262]23784.

[133263]Harano, T.; Harano, K.; Kushida, Y.; Ueda, S.; Kawakami,
H.:

[133264]Hb A(2)–Niigata [δ –1(NA1)val–to–ala]: a new delta
chain variant

[133265]found in the Japanese population. Hemoglobin 15:
335–339, 1991.

[133266]23785.

[133267]Horton, B.; Payne, R. A.; Bridges, M. T.; Huisman, T. H. J.:

[133268]Studies on an abnormal minor hemoglobin component
Hb–beta(2). Clin.

[133269]Chim. Acta 6: 246–253, 1961.

[133270]23786.

[133271] Huehns, E. R.: A third haemoglobin abnormality in two individuals

[133272] with Hb-H disease. In: Lehmann, H.; Betke, K.:
Haemoglobin-Colloquium.

[133273] Stuttgart: Georg Thieme Verlag (pub.) 1962. Pp. 76 only.
[133274] 23787.

[133275] Huehns, E. R.; Dance, N.; Beaven, G. H.; Stevens, B. L.:
Further

[133276] investigations in haemoglobin H disease. (Abstract) Proc.
9th Cong.

[133277] Int. Soc. Hemat., Mexico City 7-9, 1962.

[133278] 23788.

[133279] Huisman, T. H. J.; Sydenstricker, V. P.: Haemoglobin: difference

[133280] in gross structure of two electrophoretically identical minor haemoglobin

[133281] components. Nature 193: 489-491, 1962.

[133282] 23789.

[133283] Jeffreys, A. J.: DNA sequence variants in the G-gamma, A-gamma,

[133284] delta and beta globin genes of man. Cell 18: 1-10, 1979.

[133285]23790.

[133286]Jones, R. T.; Brimhall, B.; Huehns, E. R.; Barnicot, N. A.:
Hemoglobin

[133287]Sphakia: a delta-chain variant of hemoglobin A2 from
Crete. Science 151:

[133288]1406–1408, 1966.

[133289]23791.

[133290]Jones, R. T.; Brimhall, B.; Huisman, T. H. J.: Structural
characterization

[133291]of two delta chain variants: hemoglobin A-prime-2 (B2)
and hemoglobin

[133292]Flatbush. J. Biol. Chem. 242: 5141–5145, 1967.

[133293]23792.

[133294]Juricic, D.; Crepinko, I.; Efremov, G. D.; Lam, H.; Webber,
B.

[133295]B.; Headlee, M. G.; Huisman, T. H. J.: Hb A(2)–Zagreb or
delta125(H3)gln-to-glu,

[133296]a new delta chain variant in association with delta-
beta-thalassemia. Hemoglobin 7:

[133297]443–448, 1983.

[133298]23793.

[133299]Klein, H. L.; Petes, T. D.: Intrachromosomal gene conversion

[133300]in yeast. Nature 289: 144–148, 1981.

[133301]23794.

[133302]Labie, D.; Schroeder, W. A.; Huisman, T. H. J.: The amino acid

[133303]sequence of the delta–beta chains of haemoglobin Lepore Augusta =

[133304]Lepore Washington. Biochim. Biophys. Acta 127: 428–437, 1966.

[133305]23795.

[133306]Lanclos, K. D.; Patterson, J.; Efremov, G. D.; Wong, S. C.; Villegas,

[133307]A.; Ojwang, P. J.; Wilson, J. B.; Kutlar, F.; Huisman, T. H. J.:

[133308]Characterization of chromosomes with hybrid genes for Hb Lepore–Washington,

[133309]Hb Lepore–Baltimore, Hb P–Nilotic, and Hb Kenya. Hum. Genet. 77:

[133310]40–45, 1987.

[133311]23796.

[133312]Lee, R. C.; Huisman, T. H. J.: A variant of hemoglobin A–2

found

[133313]in a Negro family. Blood 24: 495–501, 1964.

[133314]23797.

[133315]Lehmann, H.; Jenkins, T.; Plowman, D.; Nurse, G. T.: Ho-
mozygosity

[133316]for the delta-chain variant haemoglobin A(2)-prime
(HbB-2) (delta16

[133317]gly-to-arg). Hemoglobin 9: 363–372, 1985.

[133318]23798.

[133319]Leung, H.; Gilbert, A. T.; Fleming, P. J.; Wong, J.; Hughes,
W.

[133320]G.; Hussein, S.; Nash, A. R.: Hb A(2)-Parkville or
delta47(CD6) asp-to-val,

[133321]a new delta chain variant. Hemoglobin 15: 407–416,
1991.

[133322]23799.

[133323]Lie-Injo, L. E.; Pribadi, W.; Westendorp-Boerma, F.; Efre-
mov,

[133324]G. D.; Wilson, J. B.; Reynolds, C. A.; Huisman, T. H. J.:
Hemoglobin

[133325]A(2)-Indonesia or alpha(2) beta(2) 69(E13)gly to arg.

Biochim. Biophys.

[133326]Acta 229: 335–342, 1971.

[133327]23800.

[133328]Losekoot, M.; Fodde, R.; Giordano, P. C.; Bernini, L. F.: A novel

[133329]delta-0-thalassemia arising from a frameshift insertion, detected

[133330]by direct sequencing of enzymatically amplified DNA. Hum. Genet. 83:

[133331]75–78, 1989.

[133332]23801.

[133333]Loudianos, G.; Cao, A.; Pirastu, M.; Vassilopoulos, G.; Kol-
lia,

[133334]P.; Loukopoulos, D.: Molecular basis of the delta tha-
lassemia in

[133335]cis to hemoglobin Knossos variant. (Letter) Blood 77:
2087–2088,

[133336]1991.

[133337]23802.

[133338]Loudianos, G.; Murru, S.; Kanavakis, E.; Metaxotou-Mavro-
mati,

[133339]A.; Theodoropoulou, D.; Kattamis, C.; Cao, A.; Pirastu, M.:

A new

[133340]delta chain variant hemoglobin A(2)-Corfu or al-

pha(2)delta(2) 116

[133341]arg-to-cys (G18), detected by delta-globin gene analysis

in a Greek

[133342]family. Hum. Genet. 87: 237-238, 1991.

[133343]23803.

[133344]Loudianos, G.; Murru, S.; Ristaldi, M. S.; Cossu, P.; Pilia,

G.;

[133345]Porcu, S.; Sciarratta, G. V.; Parodi, M. I.; Cao, A.; Pirastu,

M.

[133346]: A novel delta-thalassemia mutation: a G-to-C substitu-

tion at codon

[133347]30 of the delta-globin gene in a person of southern Italian

origin. Hum.

[133348]Mutat. 1: 169-171, 1992.

[133349]23804.

[133350]Loudianos, G.; Porcu, S.; Cossu, P.; Tannoia, N.; Vitucci,

A.;

[133351]Campanale, D.; Cao, A.; Pirastu, M.: A new delta-chain

variant hemoglobin

[133352]A(2)-Puglia or delta26 glu-to-asp (B8), detected by DNA analysis in

[133353]a family of southern Italian extraction. Hum. Mutat. 2: 327-329,

[133354]1993.

[133355]23805.

[133356]Martin, S. L.; Zimmer, E. A.; Kan, Y. W.; Wilson, A. C.: Silent

[133357]delta-globin gene in Old World monkeys. Proc. Nat. Acad. Sci. 77:

[133358]3563-3566, 1980.

[133359]23806.

[133360]Matsuda, M.; Sakamoto, N.; Fukumaki, Y.: Delta-thalassemia caused

[133361]by disruption of the site for an erythroid-specific transcription

[133362]factor, GATA-1, in the delta-globin gene promoter. Blood 80: 1347-1351,

[133363]1992.

[133364]23807.

[133365]Moi, P.; Loudianos, G.; Lavinha, J.; Murru, S.; Cossu, P.; Casu,

[133366]R.; Oggiano, L.; Longinotti, M.; Cao, A.; Pirastu, M.: Delta-thalassemia

[133367]due to a mutation in an erythroid-specific binding protein sequence

[133368]3-prime to the delta-globin gene. Blood 79: 512-516, 1992.

[133369]23808.

[133370]Moi, P.; Paglietti, E.; Sanna, A.; Brancati, C.; Tagarelli, A.;

[133371]Galanello, R.; Cao, A.; Pirastu, M.: Delineation of the molecular

[133372]basis of delta- and normal HbA2 beta-thalassemia. Blood 72: 530-533,

[133373]1988.

[133374]23809.

[133375]Guida, P. M.; Todd, J. E.; Moore, S. W.; Beal, J. M.: Zollinger-Ellison

[133376]syndrome with interesting variations: report of twelve cases including

[133377]one of carcinoid of the duodenum. Am. J. Surg. 112: 807-817, 1966.

[133378]23810.

- [133379]Guo, S.S.; Sawicki, M. P.: Molecular and genetic mechanisms of
tumorigenesis in multiple endocrine neoplasia type-1.
Molec. Endocr. 15:
1653-1664, 2001.
23811.
- [133383]Guru, S. C.; Agarwal, S. K.; Manickam, P.; Olufemi, S.-E.;
Crabtree,
J. S.; Weisemann, J. M.; Kester, M. B.; Kim, Y. S.; Wang, Y.;
Emmert-Buck,
M. R.; Liotta, L. A.; Spiegel, A. M.; Boguski, M. S.; Roe, B.
A.;
Collins, F. S.; Marx, S. J.; Burns, L.; Chandrasekharappa, S.
C.:
A transcript map for the 2.8-Mb region containing the
multiple endocrine
neoplasia type 1 locus. Genome Res. 7: 725-735, 1997.
23812.
- [133390]Guru, S. C.; Crabtree, J. S.; Brown, K. D.; Dunn, K. J.; Man-
ickam,
P.; Prasad, N. B.; Wangsa, D.; Burns, A. L.; Spiegel, A. M.;
Marx,

[133392]S. J.; Pavan, W. J.; Collins, F. S.; Chandrasekharappa, S. C.:
Isolation,
[133393]genomic organization, and expression analysis of Men1,
the murine
[133394]homolog of the MEN1 gene. Mammalian Genome 10:
592–596, 1999.
[133395]23813.

[133396]Guru, S. C.; Goldsmith, P. K.; Burns, A. L.; Marx, S. J.;
Spiegel,
[133397]A. M.; Collins, F. S.; Chandrasekharappa, S. C.: Menin, the
product
[133398]of the MEN1 gene, is a nuclear protein. Proc. Nat. Acad.
Sci. 95:
[133399]1630–1634, 1998.
[133400]23814.

[133401]Heppner, C.; Kester, M. B.; Agarwal, S. K.; Debelenko, L.
V.;

[133402]Emmert–Buck, M. R.; Guru, S. C.; Manickam, P.; Olufemi,
S.–E.; Skarulis,

[133403]M. C.; Doppman, J. L.; Alexander, R. H.; Kim, Y. S.; Saggar,
S. K.;

[133404]Lubensky, I. A.; Zhuang, Z.; Liotta, L. A.; Chan–

drasekharappa, S.

[133405]C.; Collins, F. S.; Spiegel, A. M.; Burns, A. L.; Marx, S. J.:

Somatic

[133406]mutation of the MEN1 gene in parathyroid tumours. Na-

ture Genet. 16:

[133407]375-378, 1997.

[133408]23815.

[133409]Heppner, C.; Reincke, M.; Agarwal, S. K.; Mora, P.; Allolio,

B.;

[133410]Burns, A. L.; Spiegel, A. M.; Marx, S. J.: MEN1 gene analysis

in

[133411]sporadic adrenocortical neoplasms. J. Clin. Endocr. Metab.

84: 216-219,

[133412]1999.

[133413]23816.

[133414]Hershon, K. S.; Kelly, W. A.; Shaw, C. M.; Schwartz, R.;

Bierman,

[133415]E. L.: Prolactinomas as part of the multiple endocrine neo-

plastic

[133416]syndrome type 1. Am. J. Med. 74: 713-720, 1983.

[133417]23817.

[133418]Hessman, O.; Skogseid, B.; Westin, G.; Akerstrom, G.: Multiple

[133419]allelic deletions and intratumoral genetic heterogeneity in MEN1 pancreatic

[133420]tumors. J. Clin. Endocr. Metab. 86: 1355–1361, 2001.

[133421]23818.

[133422]Johnson, G. J.; Summerskill, W. H. J.; Anderson, V. E.; Keating,

[133423]F. R.: Clinical and genetic investigation of a large kindred with

[133424]multiple endocrine adenomatosis. New Eng. J. Med. 277: 1379–1385,

[133425]1967.

[133426]23819.

[133427]Jones, B. S.; O'Hagan, J. J.; Phear, D. N.; Sheville, E.: A case

[133428]of the Zollinger–Ellison syndrome associated with hyperplasia of salivary

[133429]and Brunner's glands. Gut 11: 837–839, 1970.

[133430]23820.

[133431]Kaji, H.; Canaff, L.; Lebrun, J.-J.; Goltzman, D.; Hendy, G. N.

- [133432]: Inactivation of menin, a Smad3-interacting protein,
blocks transforming
- [133433]growth factor type beta signaling. Proc. Nat. Acad. Sci. 98:
3837–3842,
- [133434]2001.
- [133435]23821.
- [133436]Kjellman, M.; Roshani, L.; Teh, B. T.; Kallioniemi, O.-P.;
Hoog,
- [133437]A.; Gray, S.; Farnebo, L.-O.; Holst, M.; Backdahl, M.; Lars-
son, C.
- [133438]: Genotyping of adrenocortical tumors: very frequent
deletions of
- [133439]the MEN1 locus in 11q13 and of a 1-centimorgan region
in 2p16. J.
- [133440]Clin. Endocr. Metab. 84: 730–735, 1999.
- [133441]23822.
- [133442]Lamers, C. B. H. W.; Froeling, P. G. A. M.: Clinical signifi-
cance
- [133443]of hyperparathyroidism in familial multiple endocrine ade-
nomatosis. Am.
- [133444]J. Med. 66: 422–424, 1979.

[133445]23823.

[133446]Lamers, C. B. H. W.; Stadil, F.; van Tongeren, J. H.: Prevalence

[133447]of endocrine abnormalities in patients with the Zollinger–Ellison

[133448]syndrome in their families. Am. J. Med. 64: 607–612, 1978.

[133449]23824.

[133450]Larsson, C.; Shepherd, J.; Nakamura, Y.; Blomberg, C.; Weber,

[133451]G.; Werelius, B.; Hayward, N.; Teh, B.; Tokino, T.; Seizinger, B.;

[133452]Skogseid, B.; Oberg, K.; Nordenskjold, M.: Predictive testing for

[133453]multiple endocrine neoplasia type 1 using DNA polymorphisms. J. Clin.

[133454]Invest. 89: 1344–1349, 1992.

[133455]23825.

[133456]Larsson, C.; Skogseid, B.; Oberg, K.; Nakamura, Y.; Nordenskjold,

[133457]M.: Multiple endocrine neoplasia type 1 gene maps to chromosome 11

[133458]and is lost in insulinoma. Nature 332: 85–87, 1988.

[133459]23826.

[133460]Larsson, C.; Skogseid, B.; Oberg, K.; Nordenskjold, M.:

Linkage

[133461]of multiple endocrine neoplasia type 1 (MEN1) to markers
at 11q13.

[133462](Abstract) Cytogenet. Cell Genet. 51: 1029, 1989.

[133463]23827.

[133464]Lemmens, I.; Van de Ven, W. J. M.; Kas, K.; Zhang, C. X.;

Giraud,

[133465]S.; Wautot, V.; Buisson, N.; De Witte, K.; Salandre, J.;

Lenoir, G.;

[133466]Pugeat, M.; Calender, A.; and 19 others: Identification of
the

[133467]multiple endocrine neoplasia type 1 (MEN1) gene. Hum.
Molec. Genet. 6:

[133468]1177–1183, 1997.

[133469]23828.

[133470]Long, T. T.; Barton, T. K.; Draffin, R.; Reeves, W. J.; Mc-

Carty,

[133471]K. S., Jr.: Conservative management of the Zollinger-Elli-

son syndrome:

[133472] ectopic gastrin production by an ovarian cystadenoma.

J.A.M.A. 243:

[133473] 1837–1839, 1980.

[133474] 23829.

[133475] Lulu, D. J.; Corcoran, T. E.; Andre, M.: Familial endocrine
adenomatosis

[133476] with associated Zollinger–Ellison syndrome: Wermer's
syndrome. Am.

[133477] J. Surg. 115: 695–701, 1968.

[133478] 23830.

[133479] Mallette, L. E.; Bilezikian, J. P.; Ketcham, A. S.; Aurbach, G.

[133480] D.: Parathyroid carcinoma in familial hyperparathyroidism.
Am. J.

[133481] Med. 57: 642–648, 1974.

[133482] 23831.

[133483] Maton, P. N.; Gardner, J. D.; Jensen, R. T.: Cushing's syn-
drome

[133484] in patients with the Zollinger–Ellison syndrome. New Eng.
J. Med. 315:

[133485] 1–5, 1986.

[133486]23832.

[133487]McCarthy, D. M.: Zollinger–Ellison syndrome. Ann. Rev.
Med. 33:

[133488]197–215, 1982.

[133489]23833.

[133490]McCarthy, D. M.; Olinger, E. J.; May, R. J.; Long, B. W.;
Gardner,

[133491]J. D.: H2–histamine receptor blocking agents in the
Zollinger–Ellison

[133492]syndrome: experience in seven cases and implications for
long-term

[133493]therapy. Ann. Intern. Med. 87: 668–675, 1977.

[133494]23834.

[133495]Mee, A. S.; Ismail, S.; Bornman, P. C.; Marks, I. N.: Chang-
ing

[133496]concepts in the presentation, diagnosis and management
of the Zollinger–Ellison

[133497]syndrome. Quart. J. Med. 206: 256–267, 1983.

[133498]23835.

[133499]Nakamura, Y.; Larsson, C.; Julier, C.; Bystrom, C.; Skog-
seid,

[133500]B.; Wells, S.; Oberg, K.; Carlson, M.; Taggart, T.; O'Connell, P.;

[133501]Leppert, M.; Lalouel, J.-M.; Nordenskjold, M.; White, R.:
Localization

[133502]of the genetic defect in multiple endocrine neoplasia type
I within

[133503]a small region of chromosome 11. Am. J. Hum. Genet. 44:
751-755,

[133504]1989.

[133505]23836.

[133506]Nordenskjold, M.; Jones, C. A.; Glaser, T.; Werelius, B.;
Larsson,

[133507]C.; Jansson, M.: Mapping of chromosome 11q probes on
somatic cell

[133508]hybrids. (Abstract) Cytogenet. Cell Genet. 51: 1054, 1989.

[133509]23837.

[133510]Basson, C. T.; Bachinsky, D. R.; Lin, R. C.; Levi, T.; Elkins,

[133511]J. A.; Soultz, J.; Grayzel, D.; Kroumpouzou, E.; Traill, T. A.;
Leblanc-Straceski,

[133512]J.; Renault, B.; Kucherlapati, R.; Seidman, J. G.; Seidman, C.
E.

[133513]: Mutations in human cause limb and cardiac malforma-
tion in Holt-Oram

[133514]syndrome. Nature Genet. 15: 30-35, 1997.

[133515]23838.

[133516]Basson, C. T.; Huang, T.; Lin, R. C.; Bachinsky, D. R.; Wer-
emowicz,

[133517]S.; Vaglio, A.; Bruzzone, R.; Quadrelli, R.; Lerone, M.;
Romeo, G.;

[133518]Silengo, M.; Pereira, A.; Krieger, J.; Mesquita, S. F.; Kamis-

ago,

[133519]M.; Morton, C. C.; Pierpont, M. E. M.; Muller, C. W.; Seidman, J.

[133520]G.; Seidman, C. E.: Different TBX5 interactions in heart and limb

[133521]defined by Holt–Oram syndrome mutations. Proc. Nat. Acad. Sci. 96:

[133522]2919–2924, 1999.

[133523]23839.

[133524]Glenn, C. L.; Wang, W. Y. S.; Benjafield, A. V.; Morris, B. J.

[133525]: Linkage and association of tumor necrosis factor receptor 2 locus

[133526]with hypertension, hypercholesterolemia and plasma shed receptor. Hum.

[133527]Molec. Genet. 9: 1943–1949, 2000.

[133528]23840.

[133529]Jones, C.: Synteny between the Pro(plus) marker and human glutamate

[133530]oxaloacetate transaminase. Somat. Cell Genet. 1: 345–354, 1975.

[133531]23841.

[133532]Lasky, L. A.; Singer, M. S.; Dowbenko, D.; Imai, Y.; Henzel, W.

[133533]J.; Grimley, C.; Fennie, C.; Gillett, N.; Watson, S. R.; Rosen, S.

[133534]D.: An endothelial ligand for L-selectin is a novel mucin-like molecule. Cell 69:

[133535]927-938, 1992.

[133536]23842.

[133537]Lasky, L. A.; Singer, M. S.; Yednock, T. A.; Dowbenko, D.; Fennie,

[133538]C.; Rodriguez, H.; Nguyen, T.; Stachel, S.; Rosen, S. D.: Cloning

[133539]of a lymphocyte homing receptor reveals a lectin domain. Cell 56:

[133540]1045-1055, 1989.

[133541]23843.

[133542]Ord, D. C.; Ernst, T. J.; Zhou, L.-J.; Rambaldi, A.; Spertini,

[133543]O.; Griffin, J.; Tedder, T. F.: Structure of the gene encoding the

[133544]human leukocyte adhesion molecule-1 (TQ1, Leu-8) of lymphocytes and

[133545]neutrophils. J. Biol. Chem. 265: 7760–7767, 1990.

[133546]23844.

[133547]Siegelman, M. H.; Weissman, I. L.: Human homologue of mouse lymph

[133548]node homing receptor: evolutionary conservation at tandem cell interaction

[133549]domains. Proc. Nat. Acad. Sci. 86: 5562–5566, 1989.

[133550]23845.

[133551]Sitrin, R. G.; Pan, P. M.; Blackwood, R. A.; Huang, J.; Petty,

[133552]H. R.: Cutting edge: evidence for a signaling partnership between

[133553]urokinase receptors (CD87) and L-selectin (CD62L) in human polymorphonuclear

[133554]neutrophils. J. Immun. 166: 4822–4825, 2001.

[133555]23846.

[133556]Tedder, T. F.; Isaacs, C. M.; Ernst, T. J.; Demetri, G. D.; Adler,

[133557]D. A.; Disteche, C. M.: Isolation and chromosomal localization of

[133558]cDNAs encoding a novel human lymphocyte cell surface molecule, LAM-1:

[133559]homology with the mouse lymphocyte homing receptor
and other human

[133560]adhesion proteins. J. Exp. Med. 170: 123–133, 1989.

[133561]23847.

[133562]Aizawa, S.; Nakano, H.; Ishida, T.; Horie, R.; Nagai, M.; Ito,

[133563]K.; Yagita, H.; Okumura, K.; Inoue, J.; Watanabe, T.: Tumor
necrosis

[133564]factor receptor-associated factor (TRAF) 5 and TRAF2 are
involved

[133565]in CD30-mediated NF-kappa-B activation. J. Biol. Chem.
272: 2042–2045,

[133566]1997.

[133567]23848.

[133568]Durkop, H.; Latza, U.; Hummel, M.; Eitelbach, F.; Seed, B.;
Stein,

[133569]H.: Molecular cloning and expression of a new member of
the nerve

[133570]growth factor receptor family that is characteristic for
Hodgkin's

[133571]disease. Cell 68: 421–427, 1992.

[133572]23849.

[133573]Kurts, C.; Carbone, F. R.; Krummel, M. F.; Koch, K. M.;
Miller,

[133574]J. F. A. P.; Heath, W. R.: Signalling through CD30 protects
against

[133575]autoimmune diabetes mediated by CD8 T cells. Nature
398: 341–344,

[133576]1999.

[133577]23850.

[133578]McClive, P. J.; Morahan, G.: Assignment of the mouse ho-
mologues

[133579]of 6 loci from HSA1p to chromosomes 3 and 4. Genomics
23: 243–246,

[133580]1994.

[133581]23851.

[133582]Stein, H.; Gerdes, J.; Schwab, U.; Lemke, H.; Mason, D. Y.;
Ziegler,

[133583]A.; Schienle, W.; Diehl, V.: Identification of Hodgkin and
Sternberg–Reed

[133584]cells as a unique cell type derived from a newly detected
small cell

[133585]population. Int. J. Cancer 30: 445–449, 1982.

[133586]23852.

[133587]de Lau, W.; Clevers, H.: LEF1 turns over a new leaf. *Nature Genet.* 28:

[133588]3–5, 2001.

[133589]23853.

[133590]Hovanes, K.; Li, T. W. H.; Munguia, J. E.; Truong, T.; Milovanovic,

[133591]T.; Marsh, J. L.; Holcombe, R. F.; Waterman, M. L.: Beta-catenin-sensitive

[133592]isoforms of lymphoid enhancer factor-1 are selectively expressed in

[133593]colon cancer. *Nature Genet.* 28: 53–57, 2001.

[133594]23854.

[133595]Love, J. J.; Li, X.; Case, D. A.; Giese, K.; Grosschedl, R.; Wright,

[133596]P. E.: Structural basis for DNA bending by the architectural transcription

[133597]factor LEF-1. *Nature* 376: 791–795, 1995.

[133598]23855.

[133599]Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.: LEF1,

[133600]the gene for lymphoid enhancer-binding factor 1, mapped
to human chromosome
[133601]4 (q23-q25) and distal mouse chromosome 3 by Southern
blot analysis
[133602]and fluorescence in situ hybridization. (Abstract) Cyto-
genet. Cell
[133603]Genet. 58: 1888 only, 1991.
[133604]23856.
[133605]Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.: Gene
[133606]for lymphoid enhancer-binding factor 1 (LEF1) mapped to
human chromosome
[133607]4 (q23-q25) and mouse chromosome 3 near Egf. Ge-
nomics 11: 1040-1048,
[133608]1991.
[133609]23857.
[133610]van Genderen, C.; Okamura, R. M.; Farinas, I.; Quo, R. G.;
Parslow,
[133611]T. G.; Bruhn, L.; Grosschedl, R.: Development of several
organs that
[133612]require inductive epithelial-mesenchymal interactions is
impaired

[133613]in LEF-1-deficient mice. Genes Dev. 8: 2691-2703, 1994.

[133614]23858.

[133615]Waterman, M. L.; Fischer, W. H.; Jones, K. A.: A thymus-specific

[133616]member of the HMG protein family regulates the human T cell receptor

[133617]C alpha enhancer. Genes Dev. 5: 656-669, 1991.

[133618]23859.

[133619]Zhou, P.; Byrne, C.; Jacobs, J.; Fuchs, E.: Lymphoid enhancer

[133620]factor 1 directs hair follicle patterning and epithelial cell fate. Genes

[133621]Dev. 9: 700-713, 1995.

[133622]23860.

[133623]White, P. S.; Jensen, S. J.; Rajalingam, V.; Stairs, D.; Sulman,

[133624]E. P.; Maris, J. M.; Biegel, J. A.; Wooster, R.; Brodeur, G. M.:

[133625]Physical mapping of the CA6, ENO1, and SLC2A5 (GLUT5) genes and reassignment

[133626]of SLC2A5 to 1p36.2. Cytogenet. Cell Genet. 81: 60-64, 1998.

[133627]23861.

[133628]Puranam, R. S.; Eubanks, J. H.; Heinemann, S. F.; McNamara, J.

[133629]O.: Chromosomal localization of gene for human glutamate receptor

[133630]subunit-7. Somat. Cell Molec. Genet. 19: 581-588, 1993.

[133631]23862.

[133632]Paschen, W.; Hedreen, J. C.; Ross, C. A.: RNA editing of the glutamate

[133633]receptor subunits GluR2 and GluR6 in human brain tissue. J. Neurochem. 63:

[133634]1596-1602, 1994.

[133635]23863.

[133636]Gregor, P.; Reeves, R. H.; Jabs, E. W.; Yang, X.; Dackowski, W.;

[133637]Rochelle, J. M.; Brown, R. H., Jr.; Haines, J. L.; O'Hara, B. F.;

[133638]Uhl, G. R.; Seldin, M. F.: Chromosomal localization of glutamate

[133639]receptor genes: relationship to familial amyotrophic lateral sclerosis

[133640]and other neurological disorders of mice and humans.

Proc. Nat. Acad.

[133641]Sci. 90: 3053–3057, 1993.

[133642]23864.

[133643]McNamara, J. O.; Eubanks, J. H.; McPherson, J. D.; Wasmuth, J.

[133644]J.; Evans, G. A.; Heinemann, S. F.: Chromosomal localization of human

[133645]glutamate receptor genes. J. Neurosci. 12: 2555–2562, 1992.

[133646]23865.

[133647]Armstrong, N.; Gouaux, E.: Mechanisms for activation and antagonism

[133648]of an AMPA-sensitive glutamate receptor: crystal structures of the

[133649]GluR2 ligand binding core. Neuron 28: 165–181, 2000.

[133650]23866.

[133651]Brusa, R.; Zimmermann, F.; Koh, D.-S.; Feldmeyer, D.; Gass, P.;

[133652]Seeburg, P. H.; Sprengel, R.: Early-onset epilepsy and postnatal

[133653]lethality associated with an editing-deficient GluR-B allele

in mice. Science 270:

[133654]1677–1680, 1995.

[133655]23867.

[133656]Hollmann, M.; Hartley, M.; Heinemann, S.: Ca(2+) permeability

[133657]of KA–AMPA–gated glutamate receptor channels depends on subunit composition. Science 252:

[133658]851–853, 1991.

[133659]23868.

[133660]Kohler, M.; Kornau, H. C.; Seeburg, P. H.: The organization of

[133661]the gene for the functionally dominant alpha-amino-3-hydroxy-5-methylisoxazole-4-propionic

[133662]acid receptor subunit Glu4-B. J. Biol. Chem. 269: 17367–17370, 1994.

[133663]23869.

[133664]Sun, W.; Ferrer-Montiel, A. V.; Schinder, A. F.; McPherson, J.

[133665]P.; Evans, G. A.; Montal, M.: Molecular cloning, chromosomal mapping,

[133666]and functional expression of human brain glutamate re-

ceptors. Proc.

[133667]Nat. Acad. Sci. 89: 1443–1447, 1992.

[133668]23870.

[133669]Sun, Y.; Olson, R.; Horning, M.; Armstrong, N.; Mayer, M.;
Gouaux,

[133670]E.: Mechanism of glutamate receptor desensitization. Na-
ture 417:

[133671]245–253, 2002.

[133672]23871.

[133673]Shows, T. B.: Linkage of loci for human pyruvate kinase
and mannosephosphate

[133674]isomerase in somatic cell hybrids. (Abstract) Am. J. Hum.
Genet. 24:

[133675]13A only, 1972.

[133676]23872.

[133677]Shows, T. B.: Somatic cell genetics of enzyme markers as-
sociated

[133678]with three human linkage groups.In: Davidson, R. L.: Proc.
Conf.

[133679]Somatic Cell Hybridization, Orlando, Fla. (pub.) 1973.

[133680]23873.

- [133681]Champion, M. J.; Brown, J. A.; Shows, T. B.: Assignment of cytoplasmic
- [133682]alpha-mannosidase (MAN-A) and confirmation of the mitochondrial isocitrate
- [133683]dehydrogenase (IDH-M) genes to the q11--qter region of chromosome
- [133684]15 in man. Cytogenet. Cell Genet. 22: 498-502, 1978.
- [133685]23874.
- [133686]Ferguson-Smith, M. A.; Westerveld, A.: Report of the committee
- [133687]on the genetic constitution of chromosomes 13, 14, 15, 16, 17, 18,
- [133688]19, 20, 21, and 22 (HGM5). Cytogenet. Cell Genet. 25: 59-73, 1979.
- [133689]23875.
- [133690]Neri, G.; Ricci, R.; Pelino, A.; Bova, R.; Tedeschi, B.; Serra, A.: A boy with ring chromosome 15 derived from a
- t(15q;15q) Robertsonian
- [133692]translocation in the mother: cytogenetic and biochemical findings. Am.
- [133693]J. Med. Genet. 14: 307-314, 1983.

[133694]23876.

[133695]Chui, D.; Oh-Eda, M.; Liao, Y.-F.; Panneerselvam, K.; Lai, A.;

[133696]Marek, K. W.; Freeze, H. H.; Moremen, K. W.; Fukuda, M. N.; Marth,

[133697]J. D.: Alpha-mannosidase-II deficiency results in dyserythropoiesis

[133698]and unveils an alternate pathway in oligosaccharide biosynthesis. Cell 90:

[133699]157-167, 1997.

[133700]23877.

[133701]Chui, D.; Sellakumar, G.; Green, R. S.; Sutton-Smith, M.; McQuistan,

[133702]T.; Marek, K. W.; Morris, H. R.; Dell, A.; Marth, J. D.: Genetic

[133703]remodeling of protein glycosylation in vivo induces autoimmune disease. Proc.

[133704]Nat. Acad. Sci. 98: 1142-1147, 2001.

[133705]23878.

[133706]Gasparini, P.; del Giudice, E. M.; Delaunay, J.; Totaro, A.; Granatiero,

- [133707]M.; Melchionda, S.; Zelante, L.; Iolascon, A.: Localization of the
[133708]congenital dyserythropoietic anemia II locus to chromosome 20q11.2
[133709]by genomewide search. Am. J. Hum. Genet. 61:
1112-1116, 1997.
[133710]23879.
- [133711]Misago, M.; Liao, Y.-F.; Kudo, S.; Eto, S.; Mattei, M.-G.;
Moremen,
[133712]K. W.; Fukuda, M. N.: Molecular cloning and expression of
cDNAs encoding
[133713]human alpha-mannosidase II and a previously unrecognized alpha-mannosidase
[133714]II(X) isozyme. Proc. Nat. Acad. Sci. 92: 11766-11770,
1995.
[133715]23880.
- [133716]Moremen, K. W.; Robbins, P. W.: Isolation, characterization, and
[133717]expression of cDNAs encoding murine alpha-mannosidase II, a Golgi
[133718]enzyme that controls conversion of high mannose to complex N-glycans. J.

[133719]Cell Biol. 115: 1521–1534, 1991.

[133720]23881.

[133721]Wada, A.; Sakamoto, H.; Katoh, O.; Yoshida, T.; Yokota, J.;
Little,

[133722]P. F. R.; Sugimura, T.; Terada, M.: Two homologous onco-
genes, HST1

[133723]and INT2, are closely located in human genome. Biochem.
Biophys.

[133724]Res. Commun. 157: 828–835, 1988.

[133725]23882.

[133726]Yoshida, T.; Tsutsumi, M.; Sakamoto, H.; Miyagawa, K.;
Teshima,

[133727]S.; Sugimura, T.; Terada, M.: Expression of the HST1
oncogene in

[133728]human germ cell tumors. Biochem. Biophys. Res. Com-
mun. 155: 1324–1329,

[133729]1988.

[133730]23883.

[133731]Birchmeier, C.; Birnbaum, D.; Waitches, G.; Fasano, O.;
Wigler,

[133732]M.: Characterization of an activated human ROS gene.

Molec. Cell.

[133733]Biol. 6: 3109–3116, 1986.

[133734]23884.

[133735]Birchmeier, C.; Sharma, S.; Wigler, M.: Expression and re-
arrangement

[133736]of the ROS1 gene in human glioblastoma cells. Proc. Nat.
Acad. Sci. 84:

[133737]9270–9274, 1987.

[133738]23885.

[133739]Nagarajan, L.; Louie, E.; Tsujimoto, Y.; Balduzzi, P. C.;
Huebner,

[133740]K.; Croce, C. M.: The human c–ros gene (ROS) is located at
chromosome

[133741]region 6q16–6q22. Proc. Nat. Acad. Sci. 83: 6568–6572,
1986.

[133742]23886.

[133743]Rabin, M.; Birnbaum, D.; Wigler, M.; Ruddle, F. H.: MCF3
oncogene

[133744]mapped to region on chromosome 6 associated with ma-
lignant transformation.

[133745](Abstract) Am. J. Hum. Genet. 37 only, 1985.

[133746]23887.

[133747]Rabin, M.; Birnbaum, D.; Young, D.; Birchmeier, C.; Wigler, M.;

[133748]Ruddle, F. H.: Human ROS1 and MAS1 oncogenes located in regions of

[133749]chromosome 6 associated with tumor-specific rearrangements. Oncogene

[133750]Res. 1: 169-178, 1987.

[133751]23888.

[133752]Satoh, H.; Yoshida, M. C.; Matsushime, H.; Shibuya, M.; Sasaki,

[133753]M.: Regional localization of the human c-ros-1 on 6q22 and flt on

[133754]13q12. Jpn. J. Cancer Res. 78: 772-775, 1987.

[133755]23889.

[133756]Yoshida, M. C.; Satoh, H.; Matsushime, H.; Shibuya, M.; Sasaki,

[133757]M.: Two ros-related protooncogenes, c-ros-1 and flt, are regionally

[133758]mapped on human chromosomes 6q22 and 13q12, respectively. (Abstract) Cytogenet.

[133759]Cell Genet. 46: 724 only, 1987.

[133760]23890.

[133761]Fong, G.-H.; Rossant, J.; Breitman, M. L.: Role of the Flt-1
receptor

[133762]tyrosine kinase in regulating the assembly of vascular en-
dothelium. Nature 376:

[133763]65-69, 1995.

[133764]23891.

[133765]He, Y.; Smith, S. K.; Day, K. A.; Clark, D. E.; Licence, D. R.;

[133766]Charnock-Jones, D. S.: Alternative splicing of vascular en-
dothelial

[133767]growth factor (VEGF)-R1 (FLT-1) pre-mRNA is important
for the regulation

[133768]of VEGF activity. Molec. Endocr. 13: 537-545, 1999.

[133769]23892.

[133770]Imbert, A.; Rosnet, O.; Marchetto, S.; Ollendorff, V.; Birn-
baum,

[133771]D.; Pebusque, M.-J.: Characterization of a yeast artificial
chromosome

[133772]from human chromosome band 13q12 containing the
FLT1 and FLT3 receptor-type

[133773] tyrosine kinase genes. Cytogenet. Cell Genet. 67:
175–177, 1994.

[133774] 23893.

[133775] Kendall, R. L.; Wang, G.; Thomas, K. A.: Identification of a
natural

[133776] soluble form of the vascular endothelial growth factor re-
ceptor, FLT-1,

[133777] and its heterodimerization with KDR. Biochem. Biophys.
Res. Commun. 226:

[133778] 324–328, 1996.

[133779] 23894.

[133780] Shalaby, F.; Rossant, J.; Yamaguchi, T. P.; Gertsenstein, M.;
Wu,

[133781] X.-F.; Breitman, M. L.; Schuh, A. C.: Failure of blood-is-
land formation

[133782] and vasculogenesis in Flk-1-deficient mice. Nature 376:
62–65, 1995.

[133783] 23895.

[133784] Wiesmann, C.; Fuh, G.; Christinger, H. W.; Eigenbrot, C.;
Wells,

[133785] J. A.; de Vos, A. M.: Crystal structure at 1.7 Angstrom res-
olution

[133786]of VEGF in complex with domain 2 of the Flt-1 receptor.

Cell 91:

[133787]695-704, 1997. Cell 91:695-704, 1997

[133788]23896.

[133789]Dabiri, G. A.; Young, C. L.; Rosenbloom, J.; Southwick, F.

S.:

[133790]Molecular cloning of human macrophage capping protein

cDNA: a unique

[133791]member of the gelsolin/villin family expressed primarily in
macrophages. J.

[133792]Biol. Chem. 267: 16545-16552, 1992.

[133793]23897.

[133794]Mishra, V. S.; Henske, E. P.; Kwiatkowski, D. J.; Southwick,

F.

[133795]S.: The human actin-regulatory protein cap G: gene struc-
ture and

[133796]chromosome location. Genomics 23: 560-565, 1994.

[133797]23898.

[133798]Southwick, F. S.; DiNubile, M. J.: Rabbit alveolar

macrophages

[133799]contain a Ca(2+)-sensitive, 41,000-Dalton protein which

reversibly

[133800]blocks the 'barbed' ends of actin filaments but does not sever them. J.

[133801]Biol. Chem. 261: 14191-14195, 1986.

[133802]23899.

[133803]Fukai, K.; Oh, J.; Frenk, E.; Almodovar, C.; Spritz, R. A.:

Linkage

[133804]disequilibrium mapping of the gene for Hermansky-Pudlak syndrome to

[133805]chromosome 10q23.1-q23.3. Hum. Molec. Genet. 4: 1665-1669, 1995.

[133806]23900.

[133807]Le Beau, M. M.; Diaz, M. O.; Plowman, G. D.; Brown, J. P.; Rowley,

[133808]J. D.: Chromosomal sublocalization of the human p97 melanoma antigen. Hum.

[133809]Genet. 72: 294-296, 1986.

[133810]23901.

[133811]Plowman, G. D.; Brown, J. P.; Enns, C. A.; Schroder, J.; Nikinmaa,

[133812]B.; Sussman, H. H.; Hellstrom, K. E.; Hellstrom, I.: Assign-

ment of

[133813]the gene for human melanoma-associated antigen p97 to
chromosome 3. Nature 303:

[133814]70–72, 1983.

[133815]23902.

[133816]Rose, T. M.; Plowman, G. D.; Teplow, D. B.; Dreyer, W. J.;
Hellstrom,

[133817]K. E.; Brown, J. P.: Primary structure of the human
melanoma-associated

[133818]antigen p97 (melanotransferrin) deduced from the mRNA
sequence. Proc.

[133819]Nat. Acad. Sci. 83: 1261–1265, 1986.

[133820]23903.

[133821]Seligman, P. A.; Butler, C. D.; Massey, E. J.; Kaur, J. A.;
Brown,

[133822]J. P.; Plowman, G. D.; Miller, Y.; Jones, C.: The p97 antigen
is

[133823]mapped to the q24–qter region of chromosome 3, the
same region as

[133824]the transferrin receptor. Am. J. Hum. Genet. 38: 540–548,
1986.

[133825]23904.

[133826]Yang, F.; Lum, J. B.; McGill, J. R.; Moore, C. M.; Naylor, S. L.;

[133827]van Bragt, P. H.; Baldwin, W. D.; Bowman, B. H.: Human transferrin:

[133828]cDNA characterization and chromosomal localization. Proc. Nat. Acad.

[133829]Sci. 81: 2752–2756, 1984.

[133830]23905.

[133831]Randerson–Moor, J. A.; Harland, M.; Williams, S.; Cuthbert–Heavens,

[133832]D.; Sheridan, E.; Aveyard, J.; Sibley, K.; Whitaker, L.; Knowles,

[133833]M.; Bishop, J. N.; Bishop, D. T.: A germline deletion of p14(ARF)

[133834]but not CDKN2A in a melanoma–neural system tumour syndrome family. Hum.

[133835]Molec. Genet. 10: 55–62, 2001.

[133836]23906.

[133837]Fong, L. Y. Y.; Fidanza, V.; Zanesi, N.; Lock, L. F.; Siracusa,

[133838]L. D.; Mancini, R.; Siprashvili, Z.; Ottey, M.; Martin, S. E.; Druck,

[133839]T.; McCue, P. A.; Croce, C. M.; Huebner, K.: Muir-Torre-like syndrome
 [133840]in Fhit-deficient mice. Proc. Nat. Acad. Sci. 97:
 4742-4747, 2000.
 [133841]23907.

[133842]Chemelli, R. M.; Willie, J. T.; Sinton, C. M.; Elmquist, J. K.;
 [133843]Scammell, T.; Lee, C.; Richardson, J. A.; Williams, S. C.;
 Xiong,
 [133844]Y.; Kisanuki, Y.; Fitch, T. E.; Nakazato, M.; Hammer, R. E.;
 Saper,
 [133845]C. B.; Yanagisawa, M.: Narcolepsy in orexin knockout
 mice: molecular
 [133846]genetics of sleep regulation. Cell 437-451, 1999.
 [133847]23908.

[133848]Adkins, S.; Gan, K. N.; Mody, M.; La Du, B. N.: Molecular
 basis
 [133849]for the polymorphic forms of human serum paraoxonase/
 arylesterase:
 [133850]glutamine or arginine at position 191, for the respective A
 and B
 [133851]allozymes. Am. J. Hum. Genet. 52: 598-608, 1993.

[133852]23909.

[133853]Antikainen, M.; Murtomaki, S.; Syvanne, M.; Pahlman, R.;
Tahvanainen,

[133854]E.; Jauhiainen, M.; Frick, M. H.; Ehnholm, C.: The gln-
arg191 polymorphism

[133855]of the human paraoxonase gene (HUMPONA) is not asso-
ciated with the

[133856]risk of coronary artery disease in Finns. J. Clin. Invest. 98:
883-885,

[133857]1996.

[133858]23910.

[133859]Augustinsson, K.-B.; Henricson, B.: A genetically con-
trolled esterase

[133860]in rat plasma. Biochim. Biophys. Acta 124: 323-331,
1966.

[133861]23911.

[133862]Barbieri, M.; Bonafe, M.; Marfella, R.; Ragno, E.; Giugliano,
D.;

[133863]Franceschi, C.; Paolisso, G.: LL-paraoxonase genotype is
associated

[133864]with a more severe degree of homeostasis model assess-
ment IR in healthy

[133865]subjects. J. Clin. Endocr. Metab. 87: 222–225, 2002.

[133866]23912.

[133867]Brophy, V. H.; Jampsa, R. L.; Clendenning, J. B.; McKinstry, L.

[133868]A.; Jarvik, G. P.; Furlong, C. E.: Effects of 5–prime regulatory-region

[133869]polymorphisms on paraoxonase–gene (PON1) expression. Am. J. Hum.

[133870]Genet. 68: 1428–1436, 2001.

[133871]23913.

[133872]Cherry, N.; Mackness, M.; Durrington, P.; Povey, A.; Dippnall,

[133873]M.; Smith, T.; Mackness, B.: Paraoxonase (PON1) polymorphisms in

[133874]farmers attributing ill health to sheep dip. Lancet 359: 763–764,

[133875]2002.

[133876]23914.

[133877]Clendenning, J. B.; Humbert, R.; Green, E. D.; Wood, C.; Traver,

[133878]D.; Furlong, C. E.: Structural organization of the human

PON1 gene. Genomics 35:

[133879]586–589, 1996.

[133880]23915.

[133881]Coates, P. M.; Mestriner, M. A.; Hopkinson, D. A.: A preliminary

[133882]interpretation of the esterase isozymes of human tissues.
Ann. Hum.

[133883]Genet. 39: 1–20, 1975.

[133884]23916.

[133885]Davies, H. G.; Richter, R. J.; Keifer, M.; Broomfield, C. A.;
Sowalla,

[133886]J.; Furlong, C. E.: The effect of the human serum paraoxonase polymorphism

[133887]is reversed with diazoxon, soman and sarin. Nature Genet.
14: 334–336,

[133888]1996.

[133889]23917.

[133890]Deakin, S.; Leviev, I.; Nicaud, V.; Meynet, M.–C. B.; Tired,
L.;

[133891]James, R. W.: Paraoxonase–1 L55M polymorphism is associated with

[133892]an abnormal oral glucose tolerance test and differentiates
high risk

[133893]coronary disease families. J. Clin. Endocr. Metab. 87:
1268–1273,

[133894]2002.

[133895]23918.

[133896]Eckerson, H. W.; Romson, J.; Wyte, C.; La Du, B. N.: The
human

[133897]serum paraoxonase polymorphism: identification of phe-
notypes by their

[133898]response to salts. Am. J. Hum. Genet. 35: 214–227, 1983.

[133899]23919.

[133900]Eckerson, H. W.; Wyte, C. M.; La Du, B. N.: The human
serum paraoxonase/arylesterase

[133901]polymorphism. Am. J. Hum. Genet. 35: 1126–1136, 1983.

[133902]23920.

[133903]Eiberg, H.; Mohr, J.: Linkage relations of the paraoxonase
polymorphism

[133904]with 43 marker systems. (Abstract) Cytogenet. Cell Genet.
25: 150,

[133905]1979.

[133906]23921.

[133907]Rey-Campos, J.; Chouard, T.; Yaniv, M.; Cereghini, S.:
vHNF1

[133908]is a homeoprotein that activates transcription and forms
heterodimers

[133909]with HNF1. EMBO J. 10: 1445-1457, 1991.

[133910]23922.

[133911]Shih, D. Q.; Bussen, M.; Sehayek, E.; Ananthanarayanan,
M.; Shneider,

[133912]B. L.; Suchy, F. J.; Shefer, S.; Bollileni, J. S.; Gonzalez, F. J.;

[133913]Breslow, J. L.; Stoffel, M.: Hepatocyte nuclear factor-
1-alpha is

[133914]an essential regulator of bile acid and plasma cholesterol
metabolism. Nature

[133915]Genet. 27: 375-382, 2001.

[133916]23923.

[133917]Urhammer, S. A.; Hansen, T.; Ekstrom, C. T.; Eiberg, H.;
Pederson,

[133918]O.: The Ala/Val98 polymorphism of the hepatocyte nu-
clear factor-1-alpha

[133919]gene contributes to the interindividual variation in serum

C-peptide

[133920]response during an oral glucose tolerance test: evidence from studies

[133921]of 231 glucose-tolerant first degree relatives of type 2 diabetic

[133922]probands. J. Clin. Endocr. Metab. 83: 4506-4509, 1998.

[133923]23924.

[133924]Urhammer, S. A.; Moller, A. M.; Nyholm, B.; Ekstrom, C. T.; Eiberg,

[133925]H.; Clausen, J. O.; Hansen, T.; Pedersen, O.; Schmitz, O.: The effect

[133926]of two frequent amino acid variants of the hepatocyte nuclear factor-1-alpha

[133927]gene on estimates of the pancreatic beta-cell function in Caucasian

[133928]glucose-tolerant first-degree relatives of type 2 diabetic patients. J.

[133929]Clin. Endocr. Metab. 83: 3992-3995, 1998.

[133930]23925.

[133931]Urhammer, S. A.; Rasmussen, S. K.; Kaisaki, P. J.; Oda, N.; Yamagata,

[133932]K.; Moller, A. M.; Fridberg, M.; Hansen, L.; Hansen, T.; Bell,

G.

[133933]J.; Pedersen, O.: Genetic variation in the hepatocyte nuclear factor-1-alpha

[133934]gene in Danish Caucasians with late-onset NIDDM. Diabetesologia 40:

[133935]473-475, 1997.

[133936]23926.

[133937]Vaxillaire, M.; Rouard, M.; Yamagata, K.; Oda, N.; Kaisaki, P.

[133938]J.; Boriraj, V. V.; Chevre, J.-C.; Boccio, V.; Cox, R. D.; Lathrop,

[133939]G. M.; Dussoix, P.; Philippe, J.; Timsit, J.; Charpentier, G.; Velho,

[133940]G.; Bell, G. I.; Froguel, P.: Identification of nine novel mutations

[133941]in the hepatocyte nuclear factor 1 alpha gene associated with maturity-onset

[133942]diabetes of the young (MODY3). Hum. Molec. Genet. 6: 583-586, 1997.

[133943]23927.

[133944]Yamada, S.; Nishigori, H.; Onda, H.; Utsugi, T.; Yanagawa, T.;

[133945]Maruyama, T.; Onigata, K.; Nagashima, K.; Nagai, R.;
Morikawa, A.;

[133946]Takeuchi, T.; Takeda, J.: Identification of mutations in the
hepatocyte

[133947]nuclear factor (HNF)-1-alpha gene in Japanese subjects
with IDDM. Diabetes 46:

[133948]1643-1647, 1997.

[133949]23928.

[133950]Yamagata, K.; Oda, N.; Kalsaki, P. J.; Menzel, S.; Furuta, H.;

[133951]Vaxillaire, M.; Southam, L.; Cox, R. D.; Lathrop, G. M.;
Borhaj, V.

[133952]V.; Chen, X.; Cox, N. J.; Oda, Y.; Yano, H.; Le Beau, M. M.;
Yamada,

[133953]S.; Nishigori, H.; Takeda, J.; Fajans, S. S.; Hattersley, A. T.;
Iwasaki,

[133954]N.; Hansen, T.; Pedersen, O.; Polonsky, K. S.; Turner, R. C.;
Velho,

[133955]G.; Chevre, J.-C.; Froguel, P.; Bell, G. I.: Mutations in the
hepatocyte

[133956]nuclear factor-1-alpha gene in maturity-onset diabetes of
the young

[133957](MODY3). Nature 384: 455-457, 1996.

[133958]23929.

[133959]Yoshiuchi, I.; Yamagata, K.; Yoshimoto, M.; Zhu, Q.; Yang, Q.;

[133960]Nammo, T.; Uenaka, R.; Kinoshita, E.; Hanafusa, T.; Miyagawa, J.;

[133961]Matsuzawa, Y.: Analysis of a non-functional HNF-1-alpha (TCF1) mutation

[133962]in Japanese subjects with familial type 1 diabetes. Hum. Mutat. 18:

[133963]345-351, 2001.

[133964]23930.

[133965]Dhanasekaran, S. M.; Barrette, T. R.; Ghosh, D.; Shah, R.; Varambally,

[133966]S.; Kurachi, K.; Pienta, K. J.; Rubin, M. A.; Chinnaiyan, A. M.:

[133967]Delineation of prognostic biomarkers in prostate cancer. Nature 412:

[133968]822-826, 2001.

[133969]23931.

[133970]Leytus, S. P.; Loeb, K. R.; Hagen, F. S.; Kurachi, K.; Davie, E.

[133971]W.: A novel trypsin-like serine protease (hepsin) with a putative

[133972]transmembrane domain expressed by human liver and hepatoma cells. *Biochemistry* 27:

[133973]1067-1074, 1988.

[133974]23932.

[133975]Tsuji, A.; Torres-Rosado, A.; Arai, T.; Le Beau, M. M.; Lemons,

[133976]R. S.; Chou, S.-H.; Kurachi, K.: Hepsin, a cell membrane-associated

[133977]protease: characterization, tissue distribution, and gene localization. *J.*

[133978]*Biol. Chem.* 266: 16948-16953, 1991.

[133979]23933.

[133980]Wu, Q.; Yu, D.; Post, J.; Halks-Miller, M.; Sadler, J. E.; Morser,

[133981]J.: Generation and characterization of mice deficient in hepsin,

[133982]a hepatic transmembrane serine protease. *J. Clin. Invest.* 101: 321-326,

[133983]1998.

[133984]23934.

[133985]Cannella, B.; Hoban, C. J.; Gao, Y.-L.; Garcia-Arenas, R.;
Lawson,

[133986]D.; Marchionni, M.; Gwynne, D.; Raine, C. S.: The neureg-
ulin, glial

[133987]growth factor 2, diminishes autoimmune demyelination
and enhances

[133988]remyelination in a chronic relapsing model for multiple
sclerosis. Proc.

[133989]Nat. Acad. Sci. 95: 10100-10105, 1998.

[133990]23935.

[133991]Falls, D. L.; Rosen, K. M.; Corfas, G.; Lane, W. S.; Fis-
chbach,

[133992]G. D.: ARIA, a protein that stimulates acetylcholine recep-
tor synthesis,

[133993]is a member of the Neu ligand family. Cell 72: 801-815,
1993.

[133994]23936.

[133995]Fernandez, P.-A.; Tang, D. G.; Cheng, L.; Prochiantz, A.;
Mudge,

[133996]A. W.; Raff, M. C.: Evidence that axon-derived neuregulin
promotes

[133997]oligodendrocyte survival in the developing rat optic nerve.

Neuron 28:

[133998]81–90, 2000.

[133999]23937.

[134000]Holmes, W. E.; Sliwkowski, M. X.; Akita, R. W.; Henzel, W.

J.;

[134001]Lee, J.; Park, J. W.; Yansura, D.; Abadi, N.; Raab, H.; Lewis,

G.

[134002]D.; Shepard, H. M.; Kuang, W.-J.; Wood, W. I.; Goeddel, D.

V.; Vandlen,

[134003]R. L.: Identification of heregulin, a specific activator of
p185(erbB2). Science 256:

[134004]1205–1210, 1992.

[134005]23938.

[134006]Huang, Y. Z.; Won, S.; Ali, D. W.; Wang, Q.; Tanowitz, M.;

Du,

[134007]Q. S.; Pelkey, K. A.; Yang, D. J.; Xiong, W. C.; Salter, M. W.;

Mei,

[134008]L.: Regulation of neuregulin signaling by PSD-95 interact-
ing with

[134009]ErbB4 at CNS synapses. Neuron 26: 443–455, 2000.

[134010]23939.

[134011]Lee, J.; Wood, W. I.: Assignment of heregulin (HGL) to human chromosome

[134012]8p22-p11 by PCR analysis of somatic cell hybrid DNA. Genomics 16:

[134013]790-791, 1993.

[134014]23940.

[134015]Martinou, J.-C.; Falls, D. L.; Fischbach, G. D.; Merlie, J. P.

[134016]: Acetylcholine receptor-inducing activity stimulates expression of

[134017]the epsilon-subunit gene of the muscle acetylcholine receptor. Proc.

[134018]Nat. Acad. Sci. 88: 7669-7673, 1991.

[134019]23941.

[134020]Ni, C.-Y.; Murphy, M. P.; Golde, T. E.; Carpenter, G.: Gamma-secretase

[134021]cleavage and nuclear localization of ErbB-4 receptor tyrosine kinase. Science 294:

[134022]2179-2181, 2001.

[134023]23942.

[134024]Orr-Urtreger, A.; Trakhtenbrot, L.; Ben-Levy, R.; Wen, D.;
Rechavi,
[134025]G.; Lonai, P.; Yarden, Y.: Neural expression and chromo-
somal mapping
[134026]of NEU differentiation factor to 8p12-p21. Proc. Nat.
Acad. Sci. 90:
[134027]1867-1871, 1993.
[134028]23943.
[134029]Boyd, D.; Jain, S. K.; Crampton, J.; Barrett, K. J.; Drysdale,
[134030]J.: Isolation and characterization of a cDNA clone for hu-
man ferritin
[134031]heavy chain. Proc. Nat. Acad. Sci. 81: 4751-4755, 1984.
[134032]23944.
[134033]Caskey, J. H.; Jones, C.; Miller, Y. E.; Seligman, P. A.: Hu-
man
[134034]ferritin gene is assigned to chromosome 19. Proc. Nat.
Acad. Sci. 80:
[134035]482-486, 1983.
[134036]23945.
[134037]Costanzo, F.; Colombo, M.; Staempfli, S.; Santoro, C.;
Marone,

[134038]M.; Frank, R.; Delius, H.; Cortese, R.: Structure of gene and pseudogenes

[134039]of human apoferritin H. Nucleic Acids Res. 14: 721–736, 1986.

[134040]23946.

[134041]Cragg, S. J.; Drysdale, J.; Worwood, M.: Genes for the 'H' subunit

[134042]of human ferritin are present on a number of human chromosomes. Hum.

[134043]Genet. 71: 108–112, 1985.

[134044]23947.

[134045]Eisenstein, R. S.: Iron regulatory proteins and the molecular

[134046]control of mammalian iron metabolism. Annu. Rev. Nutr. 20: 627–662,

[134047]2000.

[134048]23948.

[134049]Ferreira, C.; Bucchini, D.; Martin, M.–E.; Levi, S.; Arosio, P.;

[134050]Grandchamp, B.; Beaumont, C.: Early embryonic lethality of H ferritin

[134051]gene deletion in mice. J. Biol. Chem. 275: 3021–3024,

2000.

[134052]23949.

[134053]Ferreira, C.; Santambrogio, P.; Martin, M.-E.; Andrieu, V.;
Feldmann,

[134054]G.; Henin, D.; Beaumont, C.: H ferritin knockout mice: a
model of

[134055]hyperferritinemia in the absence of iron overload. Blood
98: 525-532,

[134056]2001.

[134057]23950.

[134058]Gailani, M. R.; Petty, E. M.; Horsthemke, B.; Arnold, A.;
Marx,

[134059]S. J.; Bale, A. E.: Physical mapping of chromosome
11q12-13 by pulsed

[134060]field gel electrophoresis (PFGE). (Abstract) Cytogenet. Cell
Genet. 58:

[134061]1959 only, 1991.

[134062]23951.

[134063]Gatti, R. A.; Shaked, R.; Mohandas, T. K.; Salser, W.: Hu-
man

[134064]ferritin genes: chromosomal assignments and polymor-

phisms. Am. J.

[134065]Hum. Genet. 41: 654–667, 1987.

[134066]23952.

[134067]Harrison, P. M.; Arosio, P.: The ferritins: molecular properties,

[134068]iron storage function and cellular regulation. Biochim. Biophys.

[134069]Acta 1275: 161–203, 1996.

[134070]23953.

[134071]Hentze, M. W.; Keim, S.; Papadopoulos, P.; O'Brien, S.; Modi,

[134072]W.; Drysdale, J.; Leonard, W. J.; Harford, J. B.; Klausner, R. D.

[134073]: Cloning, characterization, expression, and chromosomal localization

[134074]of a human ferritin heavy-chain gene. Proc. Nat. Acad. Sci. 83:

[134075]7226–7230, 1986.

[134076]23954.

[134077]Kato, J.; Fujikawa, K.; Kanda, M.; Fukuda, N.; Sasaki, K.; Takayama,

[134078]T.; Kobune, M.; Takada, K.; Takimoto, R.; Hamada, H.; Ikeda, T.; Niitsu, Y.: A mutation, in the iron-responsive element of H ferritin mRNA, causing autosomal dominant iron overload. *Am. J. Hum. Genet.* 69: 191–197, 2001.

[134082]23955.

[134083]Leibold, E. A.; Munro, H. N.: Cytoplasmic protein binds in vitro to a highly conserved sequence in the 5-prime untranslated region of ferritin heavy- and light-subunit mRNAs. *Proc. Nat. Acad. Sci.* 85: 2171–2175, 1988.

[134087]23956.

[134088]Lauren, P.: The two histological main types of gastric carcinoma: diffuse and so-called intestinal-type carcinoma. An attempt at a histo-clinical classification. *Acta Path. Microbiol. Scand.* 64: 31–49, 1965.

[134091]23957.

[134092]La Vecchia, C.; Negri, E.; Franceschi, S.; Gentile, A.: Family

[134093]history and the risk of stomach and colorectal cancer.

Cancer 70:

[134094]50–55, 1992.

[134095]23958.

[134096]Zanghieri, G.; Di Gregorio, C.; Sacchetti, C.; Fante, R.; Sas-
satelli,

[134097]R.; Cannizzo, G.; Carriero, A.; Ponz de Leon, M.: Familial
occurrence

[134098]of gastric cancer in the 2-year experience of a popula-
tion-based registry. Cancer 66:

[134099]2047–2051, 1990.

[134100]23959.

[134101]Canfield, V. A.; Okamoto, C. T.; Chow, D.; Dorfman, J.;
Gros, P.;

[134102]Forte, J. G.; Levenson, R.: Cloning of the H,K-ATPase beta
subunit:

[134103]tissue-specific expression, chromosomal assignment, and
relationship

[134104]to Na,K-ATPase beta subunits. J. Biol. Chem. 265:

19878–19884, 1990.

[134105]23960.

[134106]Maeda, M.; Oshiman, K.-I.; Tamura, S.; Futai, M.: Human gastric

[134107](H(+) + K(+))-ATPase gene: similarity to (Na(+) + K(+))-ATPase genes

[134108]in exon/intron organization but difference in control region. J.

[134109]Biol. Chem. 265: 9027–9032, 1990.

[134110]23961.

[134111]Song, I.; Yamada, T.; Trent, J. M.: Mapping of the gene encoding

[134112]the alpha-subunit of the human H(+),K(+)-ATPase to chromosome 19q13.1

[134113]by fluorescent in situ hybridization. Genomics 14: 547–548, 1992.

[134114]23962.

[134115]Anderson, L. A.; Friedman, L.; Osborne-Lawrence, S.; Lynch, E.;

[134116]Weissenbach, J.; Bowcock, A.; King, M.-C.: High-density genetic map

[134117]of the BRCA1 region of chromosome 17q12–q21. Ge-

nomics 17: 618–623,

[134118]1993.

[134119]23963.

[134120]Cheung, A. T.; Dayanandan, B.; Lewis, J. T.; Korbitt, G. S.;
Rajotte,

[134121]R. V.; Bryer–Ash, M.; Boylan, M. O.; Wolfe, M. M.; Kieffer,
T. J.

[134122]: Glucose–dependent insulin release from genetically en-
gineered K

[134123]cells. Science 290: 1959–1962, 2000.

[134124]23964.

[134125]Fukushima, Y.; Byers, M. G.; Eddy, R. L.; Haley, L. L.;
Henry,

[134126]W. M.; Inagaki, N.; Seino, Y.; Takeda, J.; Yano, H.; Yamada,
Y.; Imura,

[134127]H.; Bell, G. I.; Shows, T. B.: Localization of the human gas-
tric

[134128]inhibitory polypeptide gene (GIP) to chromosome
17q21.3–q22. (Abstract) Cytogenet.

[134129]Cell Genet. 51: 1001 only, 1989.

[134130]23965.

[134131]Inagaki, N.; Seino, Y.; Takeda, J.; Yano, H.; Yamada, Y.;
Bell,

[134132]G. I.; Eddy, R. L.; Fukushima, Y.; Byers, M. G.; Shows, T. B.;
Imura,

[134133]H.: Gastric inhibitory polypeptide: structure and chromo-
somal localization

[134134]of the human gene. Molec. Endocr. 3: 1014-1021, 1989.

[134135]23966.

[134136]Lacroix, A.; Bolte, E.; Tremblay, J.; Dupre, J.; Poitras, P.;
Fournier,

[134137]H.; Garon, J.; Garrel, D.; Bayard, F.; Taillefer, R.; Flanagan,
R.

[134138]J.; Hamet, P.: Gastric inhibitory polypeptide-dependent
cortisol

[134139]hypersecretion--a new cause of Cushing's syndrome. New
Eng. J. Med. 327:

[134140]974-980, 1992.

[134141]23967.

[134142]Lewis, T. B.; O'Connell, P.; Leach, R. J.: Localization of glu-
cose-dependent

[134143]insulinotropic polypeptide (GIP) to a gene cluster on chro-
mosome 17q. Genomics 19:

[134144]589–591, 1994.

[134145]23968.

[134146]Mohlke, K. L.; Nichols, W. C.; Westrick, R. J.; Novak, E. K.;
Cooney,

[134147]K. A.; Swank, R. T.; Ginsburg, D.: A novel modifier gene
for plasma

[134148]von Willebrand factor level maps to distal mouse chromo-
some 11. Proc.

[134149]Nat. Acad. Sci. 93: 15352–15357, 1996.

[134150]23969.

[134151]Reznik, Y.; Allali-Zerah, V.; Chayvialle, J. A.; Leroyer, R.;
Leymarie,

[134152]P.; Travert, G.; Lebrethon, M.-C.; Budi, I.; Balliere, A.-M.;
Mahoudeau,

[134153]J.: Food-dependent Cushing's syndrome mediated by
aberrant adrenal

[134154]sensitivity to gastric inhibitory polypeptide. New Eng. J.
Med. 327:

[134155]981–986, 1992.

[134156]23970.

[134157]Ambrose, C. M.; Duyao, M. P.; Barnes, G.; Bates, G. P.; Lin,

C.

[134158]S.; Srinidhi, J.; Baxendale, S.; Hummerich, H.; Lehrach, H.;
Altherr,

[134159]M.; Wasmuth, J.; Buckler, A.; Church, D.; Housman, D.;
Berks, M.;

[134160]Micklem, G.; Durbin, R.; Dodge, A.; Read, A.; Gusella, J.;
MacDonald,

[134161]M. E.: Structure and expression of the Huntington's dis-
ease gene:

[134162]evidence against simple inactivation due to an expanded
CAG repeat. Somat.

[134163]Cell Molec. Genet. 20: 27-38, 1994.

[134164]23971.

[134165]Andrew, S. E.; Goldberg, Y. P.; Kremer, B.; Squitieri, F.;
Theilmann,

[134166]J.; Zeisler, J.; Telenius, H.; Adam, S.; Almquist, E.; Anvret,
M.;

[134167]Lucotte, G.; Stoessl, A. J.; Campanella, G.; Hayden, M. R.:
Huntington

[134168]disease without CAG expansion: phenocopies or errors in
assignment? Am.

[134169]J. Hum. Genet. 54: 852-863, 1994.

[134170]23972.

[134171]Andrew, S. E.; Goldberg, Y. P.; Kremer, B.; Telenius, H.;
Theilmann,

[134172]J.; Adam, S.; Starr, E.; Squitieri, F.; Lin, B.; Kalchman, M. A.;

[134173]Graham, R. K.; Hayden, M. R.: The relationship between
trinucleotide

[134174](CAG) repeat length and clinical features of Huntington's
disease. Nature

[134175]Genet. 4: 398-403, 1993.

[134176]23973.

[134177]Aronin, N.; Chase, K.; Young, C.; Sapp, E.; Schwarz, C.;
Matta,

[134178]N.; Kornreich, R.; Landwehrmeyer, B.; Bird, E.; Beal, M. F.;
Vonsattel,

[134179]J.-P.; Smith, T.; Carraway, R.; Boyce, F. M.; Young, A. B.;
Penney,

[134180]J. B.; DiFiglia, M.: CAG expansion affects the expression of
mutant

[134181]huntingtin in Huntington's disease brain. Neuron 15:
1193-1201,

[134182]1995.

[134183]23974.

[134184]Auerbach, W.; Hurlbert, M. S.; Hilditch-Maguire, P.;
Wadghiri,

[134185]Y. Z.; Wheeler, V. C.; Cohen, S. I.; Joyner, A. L.; MacDonald,
M.

[134186]E.; Turbull, D. H.: The HD mutation causes progressive
lethal neurological

[134187]disease in mice expressing reduced levels of huntingtin.
Hum. Molec.

[134188]Genet. 10: 2515–2523, 2001.

[134189]23975.

[134190]Bamford, K. A.; Caine, E. D.; Kido, D. K.; Cox, C.; Shoul-
son,

[134191]J.: A prospective evaluation of cognitive decline in early
Huntington's

[134192]disease: functional and radiographic correlates. Neurology
45: 1867–1873,

[134193]1995.

[134194]23976.

[134195]Barbeau, A.: Parental ascent in the juvenile form of Hunt-
ington's

[134196]chorea. (Letter) Lancet II: 937 only, 1970.

[134197]23977.

[134198]Barkley, D. S.; Hardiwidjaja, S.; Menkes, J. H.: Abnormalities
ties

[134199]in growth of skin fibroblasts of patients with Huntington's
disease. Ann.

[134200]Neurol. 1: 426-430, 1977.

[134201]23978.

[134202]Barnes, G. T.; Duyao, M. P.; Ambrose, C. M.; McNeil, S.;
Perischetti,

[134203]F.; Srinidhi, J.; Gusella, J. F.; MacDonald, M. E.: Mouse
Huntington's

[134204]disease gene homolog (Hdh). Somat. Cell Molec. Genet.
20: 87-97,

[134205]1994.

[134206]23979.

[134207]Bates, G. P.; Mangiarini, L.; Mahal, A.; Davies, S. W.: Trans-
genic

[134208]models of Huntington's disease. Hum. Molec. Genet. 6:
1633-1637,

[134209]1997.

[134210]23980.

[134211]Bates, G. P.; Valdes, J.; Hummerich, H.; Baxendale, S.; Le
Paslier,

[134212]D. L.; Monaco, A. P.; Tagle, D.; MacDonald, M. E.; Altherr,
M.; Ross,

[134213]M.; Brownstein, B. H.; Bentley, D.; Wasmuth, J. J.; Gusella,
J. F.;

[134214]Cohen, D.; Collins, F.; Lehrach, H.: Characterization of a
yeast

[134215]artificial chromosome contig spanning the Huntington's
disease gene

[134216]candidate region. Nature Genet. 1: 180–187, 1992.

[134217]23981.

[134218]Baxendale, S.; Abdulla, S.; Elgar, G.; Buck, D.; Berks, M.;
Micklem,

[134219]G.; Durbin, R.; Bates, G.; Brenner, S.; Beck, S.; Lehrach, H.:
Comparative

[134220]sequence analysis of the human and pufferfish Hunting-
ton's disease

[134221]genes. Nature Genet. 10: 67–76, 1995.

[134222]23982.

- [134223]Behan, P. O.; Bone, I.: Hereditary chorea without dementia.
J.
- [134224]Neurol. Neurosurg. Psychiat. 40: 687–691, 1977.
- [134225]23983.
- [134226]Bernardi, G.: The isochore organization of the human
genome. Annu.
- [134227]Rev. Genet. 23: 637–661, 1989.
- [134228]23984.
- [134229]Bibb, J. A.; Yan, Z.; Svenningsson, P.; Snyder, G. L.; Pieri-
bone,
- [134230]V. A.; Horiuchi, A.; Nairn, A. C.; Messer, A.; Greengard, P.:
Severe
- [134231]deficiencies in dopamine signaling in presymptomatic
Huntington's
- [134232]disease mice. Proc. Nat. Acad. Sci. 97: 6809–6814, 2000.
- [134233]23985.
- [134234]Bird, E. D.; Caro, A. J.; Pilling, J. B.: A sex related factor
- [134235]in the inheritance of Huntington's chorea. Ann. Hum.
Genet. 37:
- [134236]255–260, 1974.

[134237]23986.

[134238]Bloch, M.; Hayden, M. R.: Predictive testing for Huntington disease

[134239]in childhood: challenges and implications. Am. J. Hum. Genet. 46:

[134240]1–4, 1990.

[134241]23987.

[134242]Bloch, M.; Hayden, M. R.: Preclinical testing in Huntington disease.

[134243](Letter) Am. J. Med. Genet. 27: 733–734, 1987.

[134244]23988.

[134245]Boehnke, M.; Conneally, P. M.; Lange, K.: Two models for a maternal

[134246]factor in the inheritance of Huntington disease. Am. J. Hum. Genet. 35:

[134247]845–860, 1983.

[134248]23989.

[134249]Brackenridge, C. J.: The relation of type of initial symptoms

[134250]and line of transmission to ages at onset and death in Huntington's

[134251]disease. Clin. Genet. 2: 287–297, 1971.

[134252]23990.

[134253]Brackenridge, C. J.: Familial correlations for age at onset
and

[134254]age at death in Huntington's disease. J. Med. Genet. 9:
23–32, 1972.

[134255]23991.

[134256]Brackenridge, C. J.: Relationship of parental age of rigidity
[134257]in Huntington's disease. J. Med. Genet. 11: 136–140,
1974.

[134258]23992.

[134259]Brackenridge, C. J.; Case, J.; Chiu, E.; Probert, D. N.;
Teltscher,

[134260]B.; Wallace, D. C.: A linkage study of the loci for Hunting-
ton's

[134261]disease and some common polymorphic markers. Ann.
Hum. Genet. 42:

[134262]201–211, 1978.

[134263]23993.

[134264]Brinkman, R. R.; Mezei, M. M.; Theilmann, J.; Almqvist, E.;
Hayden,

[134265]M. R.: The likelihood of being affected with Huntington disease by

[134266]a particular age, for a specific CAG size. Am. J. Hum. Genet. 60:

[134267]1202–1210, 1997.

[134268]23994.

[134269]Brothers, C. R. D.: The history and incidence of Huntington's

[134270]chorea in Tasmania. Proc. Roy. Aust. Coll. Physicians 4: 48–50,

[134271]1949.

[134272]23995.

[134273]Buetow, K. H.; Shiang, R.; Yang, P.; Nakamura, Y.; Lathrop, G.

[134274]M.; White, R.; Wasmuth, J. J.; Wood, S.; Berdahl, L. D.; Lysens,

[134275]N. J.; Ritty, T. M.; Wise, M. E.; Murray, J. C.: A detailed multipoint

[134276]map of human chromosome 4 provides evidence for linkage heterogeneity

[134277]and position-specific recombination rates. Am. J. Hum. Genet. 48:

[134278]911–925, 1991.

[134279]23996.

[134280]Bundey, S.: New mutations in Huntington's chorea. (Letter)
J.

[134281]Med. Genet. 20: 76–77, 1983.

[134282]23997.

[134283]Byers, R. K.; Dodge, J. A.: Huntington's chorea in children:

[134284]report of four cases. Neurology 17: 587–596, 1967.

[134285]23998.

[134286]Campbell, A. M. G.; Corner, B. D.; Norman, R. M.; Urich,
H.:

[134287]The rigid form of Huntington's disease. J. Neurol. Neuro-
surg. Psychiat. 24:

[134288]71–77, 1961.

[134289]23999.

[134290]Cariello, L.; de Cristofaro, T.; Zanetti, L.; Cuomo, T.; Di
Maio,

[134291]L.; Campanella, G.; Rinaldi, S.; Zanetti, P.; Di Lauro, R.;
Varrone,

[134292]S.: Transglutaminase activity is related to CAG repeat

length in

[134293]patients with Huntington's disease. Hum. Genet. 98:
633–635, 1996.

[134294]24000.

[134295]Marshall, D.: Ectodermal dysplasia: report of kindred with
ocular

[134296]abnormalities and hearing defect. Am. J. Ophthalm. 45:
143–156, 1958.

[134297]24001.

[134298]Korkko, J.; Ritvaniemi, P.; Haataja, L.; Kaariainen, H.;
Kivirikko,

[134299]K. I.; Prockop, D. J.; Ala-Kokko, L.: Mutation in type II pro-
collagen

[134300](COL2A1) that substitutes aspartate for glycine alpha-I-67
and that

[134301]causes cataracts and retinal detachment: evidence for
molecular heterogeneity

[134302]in the Wagner syndrome and the Stickler syndrome
(arthro-ophthalmopathy). Am.

[134303]J. Hum. Genet. 53: 55–61, 1993.

[134304]24002.

[134305]Fryns, J. P.; Kleczkowska, A.; Moerman, P.; Vandenberghe,

K.:

[134306]Hereditary hydronephrosis and the short arm of chromosome 6. (Letter) Hum.

[134307]Genet. 91: 514–515, 1993.

[134308]24003.

[134309]Groenen, P. M. A.; Garcia, E.; Debeer, P.; Devriendt, K.; Fryns,

[134310]J. P.; Van de Ven, W. J. M.: Structure, sequence, and chromosome

[134311]19 localization of human USF2 and its rearrangement in a patient with

[134312]multicystic renal dysplasia. Genomics 38: 141–148, 1996.

[134313]24004.

[134314]Groenen, P. M. A.; Vanderlinden, G.; Devriendt, K.; Fryns, J.-P;

[134315]Van de Ven, W. J. M.: Rearrangement of the human CDC5L gene by a

[134316]t(6;19)(p21;q13.1) in a patient with multicystic renal dysplasia. Genomics 49:

[134317]218–229, 1998.

[134318]24005.

[134319]Sepehrnia, B.; Kamboh, M. I.; Adams–Campbell, L. L.;
Nwankwo,

[134320]M.; Ferrell, R. E.: Genetic studies of human apolipoproteins. VII.

[134321]Population distribution of polymorphisms of apolipoproteins A–I, A–II,

[134322]A–IV, C–II, E, and H in Nigeria. Am. J. Hum. Genet. 43:
847–853,

[134323]1988.

[134324]24006.

[134325]Bost, F.; Bourguignon, J.; Martin, J.–P.; Sesboue, R.;
Thiberville,

[134326]L.; Diarra–Mehrpour, M.: Isolation and characterization of
the human

[134327]inter–alpha–trypsin inhibitor heavy–chain H1 gene. Europ.
J. Biochem. 218:

[134328]283–291, 1993.

[134329]24007.

[134330]Galizzi, J.–P.; Zuber, C. E.; Harada, N.; Gorman, D. M.;
Kastelein,

[134331]R.; Banchereau, J.; Howard, M.; Miyajima, A.: Molecular
cloning of

[134332]a cDNA encoding the human interleukin 4 receptor. Int. Immun. 2:

[134333]669–675, 1990.

[134334]24008.

[134335]Grimbacher, B.; Holland, S. M.; Puck, J. M.: The interleukin-4

[134336]receptor variant Q576R in hyper-IgE syndrome. (Letter) New Eng. J.

[134337]Med. 338: 1073–1074, 1998.

[134338]24009.

[134339]Idzerda, R. L.; March, C. J.; Mosley, B.; Lyman, S. D.; VandenBos,

[134340]T.; Gimpel, S. D.; Din, W. S.; Grabstein, K. H.; Widmer, M. B.; Park,

[134341]L. S.; Cosman, D.; Beckmann, M. P.: Human interleukin-4 receptor

[134342]confers biological responsiveness and defines a novel receptor superfamily. J.

[134343]Exp. Med. 171: 861–873, 1990.

[134344]24010.

[134345]Khurana Hershey, G. K.; Friedrich, M. F.; Esswein, L. A.;

Thomas,

[134346]M. L.; Chatila, T. A.: The association of atopy with a gain-of-function

[134347]mutation in the alpha subunit of the interleukin-4 receptor. New

[134348]Eng. J. Med. 337: 1720-1725, 1997.

[134349]24011.

[134350]Kruse, S.; Japha, T.; Tedner, M.; Sparholt, S. H.; Forster, J.;

[134351]Kuehr, J.; Deichmann, K. A.; Abello, A.: The polymorphisms S503P

[134352]and Q576R in the interleukin-4 receptor alpha gene are associated

[134353]with atopy and influence the signal transduction. Immunology 96:

[134354]365-371, 1999.

[134355]24012.

[134356]Mitsuyasu, H.; Izuhara, K.; Mao, X.-Q.; Gao, P.-S.; Arinobu, Y.;

[134357]Enomoto, T.; Kawai, M.; Sasaki, S.; Dake, Y.; Hamasaki, N.; Shirakawa,

[134358]T.; Hopkin, J. M.: Ile50val variant of IL4R-alpha upregu-

lates IgE

[134359]synthesis and associates with atopic asthma. (Letter) Nature Genet. 19:

[134360]119–120, 1998.

[134361]24013.

[134362]Corcoran, A. E.; Riddell, A.; Krooshoop, D.; Venkitaraman, A. R.

[134363]: Impaired immunoglobulin gene rearrangement in mice lacking the IL-7

[134364]receptor. Nature 391: 904–907, 1998.

[134365]24014.

[134366]Ober, C.; Leavitt, S. A.; Tsalenko, A.; Howard, T. D.; Hoki, D.

[134367]M.; Daniel, R.; Newman, D. L.; Wu, X.; Parry, R.; Lester, L. A.; Solway,

[134368]J.; Blumenthal, M.; King, R. A.; Xu, J.; Meyers, D. A.; Bleecker,

[134369]E. R.; Cox, N. J.: Variation in the interleukin 4–receptor alpha

[134370]gene confers susceptibility to asthma and atopy in ethnically diverse

[134371]populations. Am. J. Hum. Genet. 66: 517–526, 2000.

[134372]24015.

[134373]Ober, C.; Tsalenko, A.; Willadsen, S.; Newman, D.; Daniel, R.;

[134374]Wu, X.; Andal, J.; Hoki, D.; Schneider, D.; True, K.; Schou, C.; Parry,

[134375]R.; Cox, N.: Genome-wide screen for atopy susceptibility alleles

[134376]in the Hutterites. Clin. Exp. Allergy 29 (suppl. 4): 11–15, 1999.

[134377]24016.

[134378]Patuzzo, C.; Trabetti, E.; Malerba, G.; Martinati, L. C.; Boner,

[134379]A. L.; Pescolliderungg, L.; Zanoni, G.; Pignatti, P. F.: No linkage

[134380]or association of the IL-4R-alpha gene Q576R mutation with atopic

[134381]asthma in Italian families. (Letter) J. Med. Genet. 37: 382–384,

[134382]2000.

[134383]24017.

[134384]Pritchard, M. A.; Baker, E.; Whitmore, S. A.; Sutherland, G. R.;

[134385] Idzerda, R. L.; Park, L. S.; Cosman, D.; Jenkins, N. A.;

Gilbert,

[134386] D. J.; Copeland, N. G.; Beckmann, M. P.: The interleukin-4
receptor

[134387] gene (IL4R) maps to 16p11.2-16p12.1 in human and to
the distal region

[134388] of mouse chromosome 7. Genomics 10: 801-806, 1991.

[134389] 24018.

[134390] Suzuki, H.; Chung, F.; Palmer, E.; Sasaki, T.; Ohara, N.;
Taylor,

[134391] B. A.; Ohara, J.-I.: Gene mapping of mouse IL-4 receptor:
the loci

[134392] of interleukin 4 (IL-4) receptor gene and lymphocyte func-
tion associated

[134393] antigen 1 (LFA-1) gene are closely linked on chromosome
7. Immunogenetics 34:

[134394] 252-256, 1991.

[134395] 24019.

[134396] Zurawski, S. M.; Vega, F., Jr.; Huyghe, B.; Zurawski, G.: Re-
ceptors

[134397] for interleukin-13 and interleukin-4 are complex and
share a novel

[134398] component that functions in signal transduction. EMBO J.
12: 2663–2670,

[134399] 1993.

[134400] 24020.

[134401] Koshland, M. E.: The coming of age of the immunoglobulin J chain. Annu.

[134402] Rev. Immun. 3: 425–453, 1985.

[134403] 24021.

[134404] Max, E. E.; McBride, O. W.; Morton, C. C.; Robinson, M. A.:
Human

[134405] J chain gene: chromosomal localization and associated restriction

[134406] fragment length polymorphisms. Proc. Nat. Acad. Sci. 83:
5592–5596,

[134407] 1986.

[134408] 24022.

[134409] Diarra-Mehrpour, M.; Bourguignon, J.; Sesboue, R.; Mattei,
M.-G.;

[134410] Passage, E.; Salier, J.-P.; Martin, J.-P.: Human plasma inter-alpha-trypsin

[134411] inhibitor is encoded by four genes on three chromosomes.

Europ. J.

[134412]Biochem. 179: 147–154, 1989.

[134413]24023.

[134414]Engchild, J. J.; Thogersen, I. B.; Pizzo, S. V.; Salvesen, G.:

[134415]Analysis of inter- α -trypsin inhibitor and a novel
trypsin inhibitor,

[134416]pre- α -trypsin inhibitor, from human plasma:
polypeptide chain

[134417]stoichiometry and assembly by glycan. J. Biol. Chem. 264:
15975–15981,

[134418]1989.

[134419]24024.

[134420]Salier, J. P.; Diarra-Mehrpour, M.; Sesboue, R.; Bour-
guignon, J.;

[134421]Benarous, R.; Ohkubo, I.; Kurachi, S.; Kurachi, K.; Martin, J.
P.

[134422]: Isolation and characterization of cDNAs encoding the
heavy chain

[134423]of human inter- α -trypsin inhibitor (I- α -TI): un-
ambiguous evidence

[134424]for multipolypeptide chain structure of I- α -TI. Proc.

Nat. Acad.

[134425]Sci. 84: 8272–8276, 1987.

[134426]24025.

[134427]Salier, J. P.; Simon, D.; Rouet, P.; Raguenes, G.; Muscatelli,

[134428]F.; Gebhard, W.; Guenet, J. L.; Mattei, M. G.: Homologous
chromosomal

[134429]locations of the four genes for inter-alpha-inhibitor and
pre-alpha-inhibitor

[134430]family in human and mouse: assignment of the ancestral
gene for the

[134431]lipocalin superfamily. Genomics 14: 83–88, 1992.

[134432]24026.

[134433]Diarra-Mehrpour, M.; Sarafan, N.; Bourguignon, J.; Bonnet,
F.;

[134434]Bost, F.; Martin, J.-P.: Human inter-alpha-trypsin inhibitor
heavy

[134435]chain H3 gene: genomic organization, promoter analysis,
and gene linkage. J.

[134436]Biol. Chem. 273: 26809–26819, 1998.

[134437]24027.

[134438]Goodwin, R. G.; Friend, D.; Ziegler, S. F.; Jerzy, R.; Falk, B.

[134439]A.; Gimpel, S.; Cosman, D.; Dower, S. K.; March, C. J.; Narmen, A.

[134440]E.; Park, L. S.: Cloning of the human and murine interleukin-7 receptors:

[134441]demonstration of a soluble form and homology to a new receptor superfamily. Cell 60:

[134442]941-951, 1990.

[134443]24028.

[134444]Kondo, M.; Takeshita, T.; Higuchi, M.; Nakamura, M.; Sudo, T.;

[134445]Nishikawa, S.-I.; Sugamura, K.: Functional participation of the IL-2

[134446]receptor gamma chain in IL-7 receptor complexes. Science 263: 1453-1454,

[134447]1994.

[134448]24029.

[134449]Lai, S. Y.; Molden, J.; Goldsmith, M. A.: Shared gamma(c) subunit

[134450]within the human interleukin-7 receptor complex: a molecular basis

[134451]for the pathogenesis of X-linked severe combined immunodeficiency. J.

[134452]Clin. Invest. 99: 169–177, 1997.

[134453]24030.

[134454]Lynch, M.; Baker, E.; Park, L. S.; Sutherland, G. R.; Goodwin,

[134455]R. G.: The interleukin–7 receptor gene is at 5p13. Hum. Genet. 89:

[134456]566–568, 1992.

[134457]24031.

[134458]Maki, K.; Sunaga, S.; Komagata, Y.; Kodaira, Y.; Mabuchi, A.; Karasuyama,

[134459]H.; Yokomuro, K.; Miyazaki, J.; Ikuta, K.: Interleukin 7 receptor–deficient

[134460]mice lack gamma/delta T cells. Proc. Nat. Acad. Sci. 93: 7172–7177,

[134461]1996.

[134462]24032.

[134463]Noguchi, M.; Nakamura, Y.; Russell, S. M.; Ziegler, S. F.; Tsang,

[134464]M.; Cao, X.; Leonard, W. J.: Interleukin–2 receptor gamma chain:

[134465]a functional component of the interleukin–7 receptor. Sci-

ence 262:

[134466]1877–1880, 1993.

[134467]24033.

[134468]Peschon, J. J.; Morrissey, P. J.; Grabstein, K. H. et al.: Early

[134469]lymphocyte expansion is severely impaired in interleukin
7 receptor-deficient

[134470]mice. J. Exp. Med. 180: 1955–1960, 1994.

[134471]24034.

[134472]Puel, A.; Ziegler, S. F.; Buckley, R. H.; Leonard, W. J.: De-
fective

[134473]IL7R expression in T-B+NK+ severe combined immunod-
efficiency. Nature

[134474]Genet. 20: 394–397, 1998.

[134475]24035.

[134476]Mezei, M. M.; Mankodi, A.; Brais, B.; Marineau, C.; Thorn-
ton, C.

[134477]A.; Rouleau, G. A.; Karpati, G.: Minimal expansion of the
GCG repeat

[134478]in the PABP2 gene does not predispose to sporadic inclu-
sion body myositis. Neurology 52:

[134479]669–670, 1999.

[134480]24036.

[134481]Aleman, A.; Verhaar, H. J. J.; De Haan, E. H. F.; De Vries, W.

[134482]R.; Samson, M. M.; Drent, M. L.; Van Der Veen, E. A.;

Koppeschaar,

[134483]H. P. F.: Insulin-like growth factor-I and cognitive function
in

[134484]healthy older men. *J. Clin. Endocr. Metab.* 84: 471-475,
1999.

[134485]24037.

[134486]Baker, J.; Liu, J.-P.; Robertson, E. J.; Efstratiadis, A.: Role

[134487]of insulin-like growth factors in embryonic and postnatal
growth. *Cell* 75:

[134488]73-82, 1993.

[134489]24038.

[134490]Bowcock, A.; Sartorelli, V.: Polymorphism and mapping of
the IGF1

[134491]gene, and absence of association with stature among
African Pygmies. *Hum.*

[134492]*Genet.* 85: 349-354, 1990.

[134493]24039.

- [134494]Camacho–Hubner, C.; Woods, K. A.; Miraki–Moud, F.; Hindmarsh, P.
- [134495]C.; Clark, A. J.; Hansson, Y.; Johnston, A.; Baxter, R. C.; Savage,
- [134496]M. O.: Effects of recombinant human insulin–like growth factor I
- [134497](IGF–I) therapy on the growth hormone–IGF system of a patient with
- [134498]a partial IGF–I gene deletion. J. Clin. Endocr. Metab. 84: 1611–1616,
- [134499]1999.
- [134500]24040.
- [134501]Goddard, A. D.; Covello, R.; Luoh, S.–M.; Clackson, T.; Attie,
- [134502]K. M.; Gesundheit, N.; Rundle, A. C.; Wells, J. A.; Carlsson, L. M.
- [134503]S.: Mutations of the growth hormone receptor in children with idiopathic
- [134504]short stature. New Eng. J. Med. 333: 1093–1098, 1995.
- [134505]24041.
- [134506]Guler, H.–P.; Binz, K.; Eigenmann, E.; Jaggi, S.; Zimmermann, D.;

- [134507]Zapf, J.; Froesch, E. R.: Small stature and insulin-like growth factors:
- [134508]prolonged treatment of mini-poodles with recombinant human insulin-like
- [134509]growth factor I. Acta Endocr. 121: 456–464, 1989.
- [134510]24042.
- [134511]Schmale, H.; Ivell, R.; Breindl, M.; Darmer, D.; Richter, D.:
- [134512]The mutant vasopressin gene from diabetes insipidus (Brattleboro)
- [134513]rats is transcribed but the message is not efficiently translated. EMBO
- [134514]J. 3: 3289–3293, 1984.
- [134515]24043.
- [134516]Willcutts, M. D.; Felner, E.; White, P. C.: Autosomal recessive
- [134517]familial neurohypophyseal diabetes insipidus with continued secretion
- [134518]of mutant weakly active vasopressin. Hum. Molec. Genet. 8: 1303–1307,
- [134519]1999.
- [134520]24044.

[134521]Bichet, D. G.; Hendy, G. N.; Lonergan, M.; Arthus, M.-F.;
Ligier,

[134522]S.; Pausova, Z.; Kluge, R.; Zingg, H.; Saenger, P.; Oppen-
heimer, E.;

[134523]Hirsch, D. J.; Gilgenkrantz, S.; Salles, J.-P.; Oberle, I.; Man-
del,

[134524]J.-L.; Gregory, M. C.; Fujiwara, T. M.; Morgan, K.; Scriver, C. R.

[134525]: X-linked nephrogenic diabetes insipidus: from the ship Hopewell

[134526]to RFLP studies. Am. J. Hum. Genet. 51: 1089–1102, 1992.

[134527]24045.

[134528]Ten Bessel, R. W.; Peters, E. R.: Progressive hydronephrosis,

[134529]hydroureter, and dilatation of the bladder in siblings with congenital

[134530]nephrogenic diabetes insipidus. J. Pediatr. 77: 439–443, 1970.

[134531]24046.

[134532]Fajans, S. S.: Maturity-onset diabetes of the young (MODY). Diabetes

[134533]Metab. Rev. 5: 579–606, 1989.

[134534]24047.

[134535]Harrison, G. S.; Drabkin, H. A.; Kao, F.-T.; Hartz, J.; Hart, I.

[134536]M.; Chu, E. H. Y.; Wu, B. J.; Morimoto, R. I.: Chromosomal location

[134537] of human genes encoding major heat-shock protein
HSP70. Somat. Cell

[134538] Molec. Genet. 13: 119–130, 1987.

[134539] 24048.

[134540] Imai, Y.; Soda, M.; Hatakeyama, S.; Akagi, T.; Hashikawa,
T.; Nakayama,

[134541] K.; Takahashi, R.: CHIP is associated with Parkin, a gene
responsible

[134542] for familial Parkinson's disease, and enhances its ubiquitin
ligase

[134543] activity. Molec. Cell 10: 55–67, 2002.

[134544] 24049.

[134545] Laroia, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.: Control

[134546] of mRNA decay by heat shock-ubiquitin-proteasome
pathway. Science 284:

[134547] 499–502, 1999.

[134548] 24050.

[134549] Milner, C. M.; Campbell, R. D.: Polymorphic analysis of the
three

[134550] MHC-linked HSP70 genes. Immunogenetics 36: 357–362,
1992.

[134551]24051.

[134552]Milner, C. M.; Campbell, R. D.: Structure and expression of
the

[134553]three MHC-linked HSP70 genes. Immunogenetics 32:
242-251, 1990.

[134554]24052.

[134555]Sargent, C. A.; Dunham, I.; Trowsdale, J.; Campbell, R. D.:
Human

[134556]major histocompatibility complex contains genes for the
major heat

[134557]shock protein HSP70. Proc. Nat. Acad. Sci. 86:
1968-1972, 1989.

[134558]24053.

[134559]Bonnycastle, L. L. C.; Yu, C.-E.; Hunt, C. R.; Trask, B. J.;
Clancy,

[134560]K. P.; Weber, J. L.; Patterson, D.; Schellenberg, G. D.:
Cloning,

[134561]sequencing, and mapping of the human chromosome 14
heat shock protein

[134562]gene (HSPA2). Genomics 23: 85-93, 1994.

[134563]24054.

[134564]Dix, D. J.; Allen, J. W.; Collins, B. W.; Mori, C.; Nakamura, N.;

[134565]Poorman–Allen, P.; Goulding, E. H.; Eddy, E. M.: Targeted gene disruption

[134566]of Hsp70–2 results in failed meiosis, germ cell apoptosis, and male

[134567]infertility. Proc. Nat. Acad. Sci. 93: 3264–3268, 1996.

[134568]24055.

[134569]Hunt, C. R.; Gasser, D. L.; Chaplin, D. D.; Pierce, J. C.; Kozak,

[134570]C. A.: Chromosomal localization of five murine HSP70 gene family

[134571]members: Hsp70–1, Hsp70–2, Hsp70–3, Hsc70t and Grp78. Genomics 16:

[134572]193–198, 1993.

[134573]24056.

[134574]Roux, A.–F.; Nguyen, V. T. T.; Squire, J. A.; Cox, D. W.: A heat

[134575]shock gene at 14q22: mapping and expression. Hum. Molec. Genet. 3:

[134576]1819–1822, 1994.

[134577]24057.

[134578]Wolpowitz, D.; Mason, T. B. A.; Dietrich, P.; Mendelsohn, M.;

[134579]Talmage, D. A.; Role, L. W.: Cysteine-rich domain isoforms of the

[134580]neuregulin-1 gene are required for maintenance of peripheral synapses. Neuron 25:

[134581]79-91, 2000.

[134582]24058.

[134583]Bolton, P.; Powell, J.; Rutter, M.; Buckle, V.; Yates, J. R. W.;

[134584]Ishikawa-Brush, Y.; Monaco, A. P.: Autism, mental retardation, multiple

[134585]exostoses and short stature in a female with 46,X,t(X;8)(p22.13;q22.1). Psychiat.

[134586]Genet. 5: 51-55, 1995.

[134587]24059.

[134588]David, G.; van der Schueren, B.; Marynen, P.; Cassiman, J.-J.;

[134589]van den Berghe, H.: Molecular cloning of amphiglycan, a novel integral

[134590]membrane heparan sulfate proteoglycan expressed by ep-

ithelial and

[134591]fibroblastic cells. J. Cell Biol. 118: 961–969, 1992.

[134592]24060.

[134593]Ishikawa–Brush, Y.; Powell, J. F.; Bolton, P.; Miller, A. P.;
Francis,

[134594]F.; Willard, H. F.; Lehrach, H.; Monaco, A. P.: Autism and
multiple

[134595]exostoses associated with an X;8 translocation occurring
within the

[134596]GRPR gene and 3–prime to the SDC2 gene. Hum. Molec.
Genet. 6: 1241–1250,

[134597]1997.

[134598]24061.

[134599]Lories, V.; De Broeck, H.; David, G.; Cassiman, J. J.; Van
den

[134600]Berghe, H.: Heparan sulfate proteoglycans of human lung
fibroblasts:

[134601]structural heterogeneity of the core proteins of the hy-
drophobic cell–associated

[134602]forms. J. Biol. Chem. 262: 854–859, 1987.

[134603]24062.

[134604] Marynen, P.; Zhang, J.; Cassiman, J.-J.; David, G.: The gene
for
[134605] the 48kDa and the 90kDa core proteins of cell surface-
associated heparan
[134606] sulphate proteoglycan of human lung fibroblast maps to
chromosome
[134607] 8q2. (Abstract) Cytogenet. Cell Genet. 51: 1040, 1989.
[134608] 24063.

[134609] Marynen, P.; Zhang, J.; Cassiman, J.-J.; Van den Berghe,
H.; David,
[134610] G.: Partial primary structure of the 48- and 90-kilodalton
core proteins
[134611] of cell surface-associated heparan sulfate proteoglycans
of lung fibroblasts:
[134612] prediction of an integral membrane domain and evidence
for multiple
[134613] distinct core proteins at the cell surface of human lung fi-
broblasts. J.
[134614] Biol. Chem. 264: 7017-7024, 1989.
[134615] 24064.

[134616] Spring, J.; Goldberger, O. A.; Jenkins, N. A.; Gilbert, D. J.;

[134617] Copeland, N. G.; Bernfield, M.: Mapping of the syndecan genes in
[134618] the mouse: linkage with members of the Myc gene family.
Genomics 21:
[134619] 597–601, 1994.
[134620] 24065.

[134621] Glaser, T.; Matthews, K. E.; Hudson, J. W.; Seth, P.; Hous-
man,
[134622] D. E.; Crerar, M. M.: Localization of the muscle, liver and
brain
[134623] glycogen phosphorylase genes on linkage maps of mouse
chromosomes
[134624] 19, 12 and 2, respectively. Genomics 5: 510–521, 1989.
[134625] 24066.

[134626] Newgard, C. B.; Littman, D. R.; van Genderen, C.; Smith,
M.; Fletterick,
[134627] R. J.: Human brain glycogen phosphorylase: cloning, se-
quence analysis,
[134628] chromosomal mapping, tissue expression, and compari-
son with the human
[134629] liver and muscle isozymes. J. Biol. Chem. 263:
3850–3857, 1988.

[134630]24067.

[134631]Yasuda, K.; Espinosa, R., III; Davis, E. M.; Le Beau, M. M.;
Bell,

[134632]G. I.: Human somatostatin receptor genes: localization of
SSTR5 to

[134633]human chromosome 20p11.2. Genomics 17: 785-786,
1993.

[134634]24068.

[134635]Barclay, A. N.: Personal Communication. Oxford, England
1/9/1989.

[134636]24069.

[134637]Douglas, J.; Albertson, D. G.; Barclay, A. N.; Davis, M.;
Rabbitts,

[134638]P. H.: RFLP and mapping of human MOX-1 gene on chro-
mosome 3. Nucleic

[134639]Acids Res. 16: 9067 only, 1988.

[134640]24070.

[134641]Hoek, R. M.; Ruuls, S. R.; Murphy, C. A.; Wright, G. J.; God-
dard,

[134642]R.; Zurawski, S. M.; Blom, B.; Homola, M. E.; Streit, W. J.;
Brown,

[134643]M. H.; Barclay, A. N.; Sedgwick, J. D.: Down-regulation of the macrophage

[134644]lineage through interaction with OX2 (CD200). Science 290: 1768-1771,

[134645]2000.

[134646]24071.

[134647]McCaughan, G. W.; Clark, M. J.; Barclay, A. N.: Characterization

[134648]of the human homolog of the rat MRC OX-2 membrane glycoprotein. Immunogenetics 25:

[134649]329-335, 1987.

[134650]24072.

[134651]McCaughan, G. W.; Clark, M. J.; Hurst, J.; Grosveld, F.; Barclay,

[134652]A. N.: The gene for MRC OX-2 membrane glycoprotein is localized on

[134653]human chromosome 3. Immunogenetics 25: 133-135, 1987.

[134654]24073.

[134655]Wright, G. J.; Jones, M.; Puklavec, M. J.; Brown, M. H.; Barclay,

[134656]A. N.: The unusual distribution of the neuronal/lymphoid cell surface

[134657]CD200 (OX2) glycoprotein is conserved in humans. Immunology 102:

[134658]173–179, 2001.

[134659]24074.

[134660]Buijs, A.; Sherr, S.; van Baal, S.; van Bezouw, S.; van der Plas,

[134661]D.; Geurts van Kessel, A.; Riegman, P.; Lekanne Deprez, R.; Zwarthoff,

[134662]E.; Hagemeijer, A.; Grosveld, G.: Translocation (12;22)(p13;q11)

[134663]in myeloproliferative disorders results in fusion of the ETS-like

[134664]TEL gene on 12q13 to the MN1 gene on 22q11. Oncogene 10: 1511–1519,

[134665]1995.

[134666]24075.

[134667]Lekanne Deprez, R. H.; Groen, N. A.; van Biezen, N. A.; Hagemeijer,

[134668]A.; van Drunen, E.; Koper, J. W.; Avezaat, C. J. J.; Bootsma, D.;

[134669]Zwarthoff, E. C.: A t(4;22) in a meningioma points to the
 localization
 [134670]of a putative tumor-suppressor gene. Am. J. Hum. Genet.
 48: 783-790,
 [134671]1991.
 [134672]24076.
 [134673]Reche, P. A.; Soumelis, V.; Gorman, D. M.; Clifford, T.; Liu,
 [134674]M.; Travis, M.; Zurawski, S. M.; Johnston, J.; Liu, Y.-J.;
 Spits,
 [134675]H.; de Waal Malefyt, R.; Kastelein, R. A.; Bazan, J. F.: Hu-
 man thymic
 [134676]stromal lymphopoietin preferentially stimulates myeloid
 cells. J.
 [134677]Immun. 167: 336-343, 2001.
 [134678]24077.
 [134679]Chrousos, G. P.: Personal Communication. Bethesda, Md.
 10/21/2002.
 [134680]24078.
 [134681]Chrousos, G. P.; MacLusky, N. J.; Brandon, D. D.; Tomita,
 M.; Renquist,
 [134682]D. M.; Loriaux, D. L.; Lipsett, M. B.: Progesterone resis-

tance.In:

[134683]Chrousos, G. P.; Loriaux, D. L.; Lipsett, M. B.: Steroid Hormone

[134684]Resistance: Mechanisms and Clinical Aspects. New York: Plenum Press

[134685](pub.) 1986.

[134686]24079.

[134687]Keller, D. W.; Wiest, W. G.; Askin, F. B.; Johnson, L. W.; Strickler,

[134688]R. C.: Pseudocorpus luteum insufficiency: a local defect of progesterone

[134689]action on endometrial stroma. J. Clin. Endocr. Metab. 48: 127-132,

[134690]1979.

[134691]24080.

[134692]Daiger, S. P.; Reed, L.; Huang, S.-S.; Zeng, Y.-T.; Wang, T.;

[134693]Lo, W. H. Y.; Okano, Y.; Hase, Y.; Fukuda, Y.; Oura, T.; Tada, K.;

[134694]Woo, S. L. C.: Polymorphic DNA haplotypes at the phenylalanine hydroxylase

[134695](PAH) locus in Asian families with phenylketonuria (PKU).

Am. J.

[134696]Hum. Genet. 45: 319–324, 1989.

[134697]24081.

[134698]Desviat, L. R.; Perez, B.; De Lucca, M.; Cornejo, V.;

Schmidt,

[134699]B.; Ugarte, M.: Evidence in Latin America of recurrence of

V388M,

[134700]a phenylketonuria mutation with high in vitro residual ac-

tivity. Am.

[134701]J. Hum. Genet. 57: 337–342, 1995.

[134702]24082.

[134703]Desviat, L. R.; Perez, B.; Ugarte, M.: A new PKU mutation

associated

[134704]with haplotype 12. Hum. Molec. Genet. 1: 765–766, 1992.

[134705]24083.

[134706]Desviat, L. R.; Perez, B.; Ugarte, M.: Phenylketonuria in

Spanish

[134707]Gypsies: prevalence of the IVS10nt546 mutation on haplo-

type 34. Hum.

[134708]Mutat. 9: 66–68, 1997.

[134709]24084.

[134710]Dianzani, I.; Camaschella, C.; Saglio, G.; Ferrero, G. B.; Ramus,

[134711]S.; Ponzzone, A.; Cotton, R. G. H.: Molecular analysis of contiguous

[134712]exons of the phenylalanine hydroxylase gene: identification of a new

[134713]PKU mutation. J. Med. Genet. 30: 228–231, 1993.

[134714]24085.

[134715]Dianzani, I.; Devoto, M.; Camaschella, C.; Saglio, G.; Ferrero,

[134716]G. B.; Cerone, R.; Romano, C.; Romeo, G.; Giovannini, M.; Riva, E.;

[134717]Angeneydt, F.; Trefz, F. K.; Okano, Y.; Woo, S. L. C.: Haplotype

[134718]distribution and molecular defects at the phenylalanine hydroxylase

[134719]locus in Italy. Hum. Genet. 86: 69–72, 1990.

[134720]24086.

[134721]Dianzani, I.; Forrest, S. M.; Camaschella, C.; Saglio, G.; Ponzzone,

[134722]A.; Cotton, R. G. H.: Screening for mutations in the phenylalanine

- [134723]hydroxylase gene from Italian patients with phenylketonuria by using
- [134724]the chemical cleavage method: a new splice mutation. Am. J. Hum. Genet. 48: 631–635, 1991.
- [134725]24087.
- [134726]DiLella, A. G.; Huang, W.-M.; Woo, S. L. C.: Screening for phenylketonuria mutations by DNA amplification with the polymerase chain reaction. Lancet I: 497–499, 1988.
- [134727]24088.
- [134728]DiLella, A. G.; Kwok, S. C. M.; Ledley, F. D.; Marvit, J.; Woo, S. L. C.: Molecular structure and polymorphic map of the human phenylalanine hydroxylase gene. Biochemistry 25: 743–749, 1986.
- [134729]24089.
- [134730]DiLella, A. G.; Marvit, J.; Brayton, K.; Woo, S. L. C.: An amino-acid substitution involved in phenylketonuria is in linkage disequilibrium

[134737]with DNA haplotype 2. Nature 327: 333–336, 1987.

[134738]24090.

[134739]DiLella, A. G.; Marvit, J.; Guttler, F.; Woo, S. L. C.: Molecular

[134740]genetics of phenylketonuria. (Abstract) 7th Int. Cong. Hum. Genet.,

[134741]Berlin 665–666, 1986.

[134742]24091.

[134743]DiLella, A. G.; Marvit, J.; Lidsky, A. S.; Guttler, F.; Woo, S.

[134744]L. C.: Tight linkage between a splicing mutation and a specific DNA

[134745]haplotype in phenylketonuria. Nature 322: 799–803, 1986.

[134746]24092.

[134747]Drogari, E.; Smith, I.; Beasley, M.; Lloyd, J. K.: Timing of

[134748]strict diet in relation to fetal damage in maternal phenylketonuria:

[134749]an international collaborative study by the MRC/DHSS phenylketonuria

[134750]register. Lancet II: 927–930, 1987.

[134751]24093.

[134752]Dworniczak, B.; Aulehla-Scholz, C.; Horst, J.: Phenylalanine

[134753]hydroxylase gene: silent mutation uncovers evolutionary origin of

[134754]different alleles. Clin. Genet. 38: 270-273, 1990.

[134755]24094.

[134756]Dworniczak, B.; Aulehla-Scholz, C.; Horst, J.: Phenylketonuria:

[134757]detection of a frequent haplotype 4 allele mutation. Hum. Genet. 84:

[134758]95-96, 1989.

[134759]24095.

[134760]Dworniczak, B.; Aulehla-Scholz, C.; Kalaydjieva, L.; Bartholome,

[134761]K.; Grudza, K.; Horst, J.: Aberrant splicing of phenylalanine hydroxylase

[134762]mRNA: the major cause for phenylketonuria in parts of southern Europe. Genomics 11:

[134763]242-246, 1991.

[134764]24096.

[134765]Dworniczak, B.; Grudda, K.; Stumper, J.; Bartholome, K.;
Aulehla-Scholz,

[134766]C.; Horst, J.: Phenylalanine hydroxylase gene: novel mis-
sense mutation

[134767]in exon 7 causing severe phenylketonuria. Genomics 9:
193-199, 1991.

[134768]24097.

[134769]Dworniczak, B.; Kalaydjieva, L.; Aulehla-Scholz, C.; Ullrich,

[134770]K.; Kremensky, I.; Radeva, B.; Horst, J.: Recurrent non-
sense mutation

[134771]in exon 7 of the phenylalanine hydroxylase gene. Hum.
Genet. 87:

[134772]731-733, 1991.

[134773]24098.

[134774]Dworniczak, B.; Kalaydjieva, L.; Pankoke, S.; Aulehla-
Scholz,

[134775]C.; Allen, G.; Horst, J.: Analysis of exon 7 of the human
phenylalanine

[134776]hydroxylase gene: a mutation hot spot? Hum. Mutat. 1:
138-146, 1992.

[134777]24099.

[134778]Economou-Petersen, E.; Henriksen, K. F.; Guldberg, P.;
Guttler,

[134779]F.: Molecular basis for nonphenylketonuria hyperphenyl-
alaninemia. Genomics 14:

[134780]1-5, 1992.

[134781]24100.

[134782]Eigel, A.; Dworniczak, B.; Kalaydjieva, L.; Horst, J.: A
frameshift

[134783]mutation in exon 2 of the phenylalanine hydroxylase gene
linked to

[134784]RFLP haplotype 1. Hum. Genet. 87: 739-741, 1991.

[134785]24101.

[134786]Eiken, H. G.; Knappskog, P. M.; Apold, J.; Flatmark, T.: PKU

[134787]mutation G46S is associated with increased aggregation
and degradation

[134788]of the phenylalanine hydroxylase enzyme. Hum. Mutat. 7:
228-238,

[134789]1996.

[134790]24102.

[134791]Eiken, H. G.; Knappskog, P. M.; Apold, J.; Skjelkvale, L.;
Boman,

- [134792]H.: A de novo phenylketonuria mutation: ATG (met) to ATA (ile) in the start codon of the phenylalanine hydroxylase gene. Hum. Mutat. 1: 388–391, 1992.
- [134793]24103.
- [134794]Eiken, H. G.; Knappskog, P. M.; Boman, H.; Thune, K. S.; Kaada, G.; Motzfeldt, K.; Apold, J.: Relative frequency, heterogeneity and geographic clustering of PKU mutations in Norway. Europ. J. Hum. Genet. 4: 205–213, 1996.
- [134795]24104.
- [134796]Eiken, H. G.; Knappskog, P. M.; Guldberg, P.; Apold, J.: DGGE analysis as supplement to SSCP analysis of the phenylalanine hydroxylase gene: detection of eight (one de novo, seven inherited) of nine remaining Norwegian PKU mutations. Hum. Mutat. 8: 19–22, 1996.

[134805]24105.

[134806]Eiken, H. G.; Stangeland, K.; Skjelkvale, L.; Knappskog, P.
M.;

[134807]Boman, H.; Apol, J.: PKU mutations R408Q and F299C in
Norway: haplotype

[134808]associations, geographic distributions and phenotype
characteristics. Hum.

[134809]Genet. 88: 608–612, 1992.

[134810]24106.

[134811]Dautigny, A.; Bernier, I.; Colombani, J.; Jolles, P.: Human
platelets

[134812]as a source of HL-A antigens: a study of various solubi-
lization techniques. Biochimie 57:

[134813]1197–1201, 1975.

[134814]24107.

[134815]Moser, A. B.; Kreiter, N.; Bezman, L.; Lu, S.; Raymond, G.
V.;

[134816]Naidu, S.; Moser, H. W.: Plasma very long chain fatty acids
in 3,000

[134817]peroxisome disease patients and 29,000 controls. Ann.
Neurol. 45:

[134818]100–110, 1999.

[134819]24108.

[134820]Hoskins, J. A.; Jack, G.; Wade, H. E.; Peiris, R. J. D.; Wright,

[134821]E. C.; Starr, D. J. T.; Stern, J.: Enzymatic control of phenyl-
alanine

[134822]intake in phenylketonuria. *Lancet* I: 392–394, 1980.

[134823]

[134824]100. Howell, R. R.; Stevenson, R. E.: The offspring of
phenylketonuric

[134825]women. *Soc. Biol.* 18 (suppl.): S19–S29, 1971.

[134826]

[134827]101. Hsia, D. Y.–Y.: Phenylketonuria and its variants. *Prog.*
Med.

[134828]*Genet.* 7: 29–68, 1970.

[134829]

[134830]102. Huang, S.; Ren, Z.; Zeng, Y.; Woo, S. L. C.: PAH 399
GTA(val)–to–GTT(val),

[134831]a new silent mutation found in the Chinese. *Hum. Genet.*
86: 305–306,

[134832]1991.

[134833]

[134834]103. Huang, S.-Z.; Zhou, X.-D.; Ren, Z.-R.; Zeng, Y.-T.;
Woo, S. L.

[134835]C.: Prenatal detection of an arg-to-ter mutation at codon
111 of

[134836]the PAH gene using DNA amplification. Prenatal Diag. 10:
289-293,

[134837]1990.

[134838]

[134839]104. Huntley, C. C.; Stevenson, R. E.: Maternal phenylke-
tonuria:

[134840]course of two pregnancies. Obstet. Gynec. 34: 694-700,
1969.

[134841]

[134842]105. Hutchesson, A. C. J.; Hall, S. K.; Preece, M. A.; Green,
A.:

[134843]Screening for tyrosinaemia type I. Arch. Child. Dis. 74:
F191-F194,

[134844]1996.

[134845]

[134846]106. Ivaschenko, T.; Baranov, V. S.: Rapid and efficient

PCR/Styl

[134847]test for identification of common mutation R408W in
phenylketonuria

[134848]patients. J. Med. Genet. 30: 153–154, 1993.

[134849]

[134850]107. Jaruzelska, J.; Henriksen, K. F.; Guttler, F.; Riess, O.;
Borski,

[134851]K.; Blin, N.; Slomski, R.: The codon 408 mutation associ-
ated with

[134852]haplotype 2 is predominant in Polish families with
phenylketonuria. Hum.

[134853]Genet. 86: 247–250, 1991.

[134854]

[134855]108. Jaruzelska, J.; Melle, D.; Matuszak, R.; Borski, K.;
Munnich,

[134856]A.: A new 15 bp deletion in exon 11 of the phenylalanine
hydroxylase

[134857]gene in phenylketonuria. Hum. Molec. Genet. 1: 763–764,
1992.

[134858]

[134859]109. Jervis, G. A.: Phenylpyruvic oligophrenia deficiency of

phenylalanine-oxidizing

[134860]system. Proc. Soc. Exper. Biol. Med. 82: 514-515, 1953.

[134861]

[134862]110. Jervis, G. A.: Studies on phenylpyruvic oligophrenia:
the position

[134863]of the metabolic error. J. Biol. Chem. 169: 651-656, 1947.

[134864]

[134865]111. John, S. W. M.; Rozen, R.; Laframboise, R.; Laberge,
C.; Sriver,

[134866]C. R.: Five mutations at the PAH locus account for almost
90% of

[134867]PKU mutations in French-Canadians from eastern Quebec.
Hum. Mutat. 1:

[134868]72-74, 1992.

[134869]

[134870]112. John, S. W. M.; Rozen, R.; Laframboise, R.; Laberge,
C.; Sriver,

[134871]C. R.: Novel PKU mutation on haplotype 2 in French-
Canadians. Am.

[134872]J. Hum. Genet. 45: 905-909, 1989.

[134873]

[134874]113. John, S. W. M.; Rozen, R.; Scriver, C. R.; Laframboise, R.; Laberge, C.: Recurrent mutation, gene conversion, or recombination at the human phenylalanine hydroxylase locus: evidence in French-Canadians and a catalog of mutations. Am. J. Hum. Genet. 46: 970-974, 1990.

[134878]

[134879]114. John, S. W. M.; Scriver, C. R.; Laframboise, R.; Rozen, R.:

[134880]In vitro and in vivo correlations for I65T and M1V mutations at the

[134881]phenylalanine hydroxylase locus. Hum. Mutat. 1: 147-153, 1992.

[134882]

[134883]115. Justice, M. J.; Siracusa, L. D.; Gilbert, D. J.; Heisterkamp,

[134884]N.; Groffen, J.; Chada, K.; Silan, C. M.; Copeland, N. G.; Jenkins,

[134885]N. A.: A genetic linkage map of mouse chromosome 10: localization

[134886]of eighteen molecular markers using a single interspecific
backcross. Genetics 125:

[134887]855–866, 1990.

[134888]

[134889]116. Kalanin, J.; Takarada, Y.; Kagawa, S.; Yamashita, K.;
Ohtsuka,

[134890]N.; Matsuoka, A.: Gypsy phenylketonuria: a point mutation
of the

[134891]phenylalanine hydroxylase gene in gypsy families from
Slovakia. Am.

[134892]J. Med. Genet. 49: 235–239, 1994.

[134893]

[134894]117. Kalaydjieva, L.; Dworniczak, B.; Aulehla–Scholz, C.;
Devoto,

[134895]M.; Romeo, G.; Sturhmann, M.; Kucinskas, V.; Yurgelyavi-
cius, V.; Horst,

[134896]J.: Silent mutations in the phenylalanine hydroxylase gene
as an

[134897]aid to the diagnosis of phenylketonuria. J. Med. Genet. 28:
686–690,

[134898]1991.

[134899]

[134900]118. Kalaydjieva, L.; Dworniczak, B.; Aulehla-Scholz, C.;
Kremensky,

[134901]I.; Bronzova, J.; Eigel, A.; Horst, J.: Classical phenylke-
tonuria

[134902]in Bulgaria: RFLP haplotypes and frequency of the major
mutations. J.

[134903]Med. Genet. 27: 742-745, 1990.

[134904]

[134905]119. Kalaydjieva, L.; Dworniczak, B.; Kucinskas, V.;
Yurgeliavicius,

[134906]V.; Kunert, E.; Horst, J.: Geographical distribution gradi-
ents of

[134907]the major PKU mutations and the linked haplotypes. Hum.
Genet. 86:

[134908]411-413, 1991.

[134909]

[134910]120. Kamaryt, J.; Mrskos, A.; Podhradská, O.; Kolcova, V.;
Cabalska,

[134911]B.; Duzynska, N.; Borzymowska, J.: PKU locus: genetic
linkage with

[134912]human amylase (AMY) loci and assignment to linkage group 1. Hum.

[134913]Genet. 43: 205–210, 1978.

[134914]

[134915]121. Kaufman, S.: Phenylketonuria: biochemical mechanisms. Adv.

[134916]Neurochem. 2: 1–132, 1976.

[134917]

[134918]122. Kaufman, S.: A model of human phenylalanine metabolism in normal

[134919]subjects and in phenylketonuric patients. Proc. Nat. Acad. Sci. 96:

[134920]3160–3164, 1999.

[134921]

[134922]123. Kaufman, S.: Differential diagnosis of variant forms of hyperphenylalaninemia. Pediatrics 65:

[134923]840–842, 1980.

[134924]

[134925]124. Kaufman, S.; Fisher, D. B.: Purification and some physical properties

[134926]of phenylalanine hydroxylase from rat liver. J. Biol. Chem.
245:

[134927]4745–4750, 1970.

[134928]

[134929]125. Kaufman, S.; Max, E. E.; Kang, E. S.: Phenylalanine
hydroxylase

[134930]activity in liver biopsies from hyperphenylalaninemia het-
erozygotes:

[134931]deviation from proportionality with gene dosage. Pediat.
Res. 9:

[134932]632–634, 1975.

[134933]

[134934]126. Kawashima, H.; Kawano, M.; Masaki, A.; Sato, T.:
Three cases

[134935]of untreated classical PKU: a report on cataracts and brain
calcification. Am.

[134936]J. Med. Genet. 29: 89–93, 1988.

[134937]

[134938]127. Kerr, G. R.; Chamove, A. S.; Harlow, H. F.; Waisman,
H. A.:

[134939]'Fetal PKU': the effect of maternal hyperphenylalaninemia

during pregnancy

[134940]in the rhesus monkey (*Macaca mulatta*). *Pediatrics* 42:
27–36, 1968.

[134941]

[134942]128. Kidd, K. K.: Phenylketonuria: population genetics of a
disease. *Nature* 327:

[134943]282–283, 1987.

[134944]

[134945]129. Kleiman, S.; Vanagaite, L.; Bernstein, J.; Schwartz, G.;
Brand,

[134946]N.; Elitzur, A.; Woo, S. L. C.; Shiloh, Y.: Phenylketonuria:
variable

[134947]phenotypic outcomes of the R261Q mutation and mater-
nal PKU in the

[134948]offspring of a healthy homozygote. *J. Med. Genet.* 30:
284–288, 1993.

[134949]

[134950]130. Knapp, A.; Tintschewa, R.; Scheibe, E.; Schiebe, E.;
Jager, B.;

[134951]Biebler, K. E.: The genetic linkage between the PKU locus
and the

[134952]loci for amylase–1, amylase–2, Fy, PGM–1, and Rh and the

question

[134953] of assignment of the PKU locus to chromosome no. 1.

Hum. Genet. 60:

[134954] 122–125, 1982.

[134955]

[134956] 131. Knappskog, P. M.; Eiken, H. G.; Martinez, A.; Flatmark, T.; Apold,

[134957] J.: The PKU mutation S349P causes complete loss of catalytic activity

[134958] in the recombinant phenylalanine hydroxylase enzyme.

Hum. Genet. 95:

[134959] 171–173, 1995.

[134960]

[134961] 132. Koch, R.; Moats, R.; Guttler, F.; Guldberg, P.; Nelson, M., Jr.

[134962] Blood–brain phenylalanine relationships in persons with phenylketonuria. Pediatrics 106:

[134963] 1093–1096, 2000.

[134964]

[134965] 133. Komrower, G. M.; Sardharwalla, I. B.; Coutts, J. M. J.; Ingham,

[134966]D.: Management of maternal phenylketonuria: an emerging clinical

[134967]problem. Brit. Med. J. I: 1383–1387, 1979.

[134968]

[134969]134. Konecki, D. S.; Lichter-Konecki, U.: The phenylketonuria locus:

[134970]current knowledge about alleles and mutations of the phenylalanine

[134971]hydroxylase gene in various populations. Hum. Genet. 87: 377–388,

[134972]1991.

[134973]

[134974]135. Konecki, D. S.; Schlotter, M.; Trefz, F. K.; Lichter-Konecki,

[134975]U.: The identification of two mis-sense mutations at the PAH gene

[134976]locus in a Turkish patient with phenylketonuria. Hum. Genet. 87:

[134977]389–393, 1991.

[134978]

[134979]136. Kwok, S. C. M.; Ledley, F. D.; DiLella, A. G.; Robson,

K. J.

[134980]H.; Woo, S. L. C.: Nucleotide sequence of a full-length complementary

[134981]DNA clone and amino acid sequence of human phenylalanine hydroxylase. *Biochemistry* 24:

[134982]556–561, 1985.

[134983]

[134984]137. Labrune, P.; Melle, D.; Rey, F.; Berthelon, M.; Cailaud, C.;

[134985]Rey, J.; Munnich, A.; Lyonnet, S.: Single-strand conformation polymorphism

[134986]for detection of mutations and base substitutions in phenylketonuria. *Am.*

[134987]*J. Hum. Genet.* 48: 1115–1120, 1991.

[134988]

[134989]138. Lasala, J. M.; Coscia, C. J.: Accumulation of a tetrahydroisoquinoline

[134990]in phenylketonuria. *Science* 203: 283–284, 1979.

[134991]

[134992]139. Leandro, P.; Rivera, I.; Ribeiro, V.; de Almeida, I. T.; da Silva,

[134993]C.; Lechner, M. C.: Mutation analysis of phenylketonuria in
South

[134994]and Central Portugal: prevalence of V388M mutation.
Hum. Mutat. 6:

[134995]192–194, 1995.

[134996]

[134997]140. Ledbetter, S. A.; Ledbetter, D. H.; Ledley, F. D.; Woo,
S.:

[134998]Localization of phenylalanine hydroxylase (PAH) and al-
pha-1 antitrypsin

[134999](AAT) loci in mouse genome by synteny and in situ hy-
bridization. (Abstract) Am.

[135000]J. Hum. Genet. 41: A173 only, 1987.

[135001]

[135002]141. Ledley, F. D.; DiLella, A. G.; Woo, S. L. C.: Molecular
biology

[135003]of phenylalanine hydroxylase and phenylketonuria. Trends
Genet. 1:

[135004]309–313, 1985.

[135005]

[135006]142. Ledley, F. D.; Grenett, H. E.; DiLella, A. G.; Kwok, S. C.

M.;

[135007]Woo, S. L. C.: Gene transfer and expression of human
phenylalanine

[135008]hydroxylase. Science 228: 77-79, 1985.

[135009]

[135010]143. Ledley, F. D.; Grenett, H. E.; McGinnis-Shelnutt, M.;
Woo, S.

[135011]L. C.: Retroviral-mediated gene transfer of human pheny-
lalanine hydroxylase

[135012]into NIH 3T3 and hepatoma cells. Proc. Nat. Acad. Sci. 83:
409-413,

[135013]1986.

[135014]

[135015]144. Ledley, F. D.; Koch, R.; Jew, K.; Beaudet, A.; O'Brien,
W. E.;

[135016]Bartos, D. P.; Woo, S. L. C.: Phenylalanine hydroxylase ex-
pression

[135017]in liver of a fetus with phenylketonuria. J. Pediat. 113:
463-468,

[135018]1988.

[135019]

[135020]145. Ledley, F. D.; Levy, H. L.; Woo, S. L. C.: Molecular analysis

[135021]of the inheritance of phenylketonuria and mild hyperphenylalaninemia

[135022]in families with both disorders. New Eng. J. Med. 314: 1276–1280,

[135023]1986.

[135024]

[135025]146. Lenke, R. R.; Levy, H. L.: Maternal phenylketonuria and hyperphenylalaninemia:

[135026]an international survey of the outcome of untreated and treated pregnancies. New

[135027]Eng. J. Med. 303: 1202–1208, 1980.

[135028]

[135029]147. Leuzzi, V.; Bianchi, M. C.; Tosetti, M.; Carducci, C.; Carducci,

[135030]C.; Antonozzi, I.: Clinical significance of brain phenylalanine concentration

[135031]assessed by in vivo proton magnetic resonance spectroscopy in phenylketonuria. J.

[135032]Inherit. Metab. Dis. 23: 563–570, 2000.

[135033]

[135034]148. Levy, H. L.: Molecular genetics of phenylketonuria and its implications.

[135035](Editorial) Am. J. Hum. Genet. 45: 667–670, 1989.

[135036]

[135037]149. Levy, H. L.; Guldberg, P.; Guttler, F.; Hanley, W. B.; Matalon,

[135038]R.; Rouse, B. M.; Trefz, F.; Azen, C.; Allred, E. N.; de la Cruz,

[135039]F.; Koch, R.: Congenital heart disease in maternal phenylketonuria:

[135040]report from the Maternal PKU Collaborative Study. Pediat. Res. 49:

[135041]636–642, 2001.

[135042]

[135043]150. Levy, H. L.; Karolkewicz, V.; Houghton, S. A.; Mac-Cready, R.

[135044]A.: Screening the 'normal' population in Massachusetts for phenylketonuria. New

[135045]Eng. J. Med. 282: 1455–1458, 1970.

[135046]

[135047]151. Levy, H. L.; Lobbregt, D.; Barnes, P. D.; Poussaint, T.
Y.:

[135048]Maternal phenylketonuria: magnetic resonance imaging of
the brain

[135049]in offspring. J. Pediat. 128: 770–775, 1996.

[135050]

[135051]152. Levy, H. L.; Waisbren, S. E.: Effects of untreated ma-
ternal

[135052]phenylketonuria and hyperphenylalaninemia on the fetus.
New Eng.

[135053]J. Med. 309: 1269–1274, 1983.

[135054]

[135055]153. Li, J.; Eisensmith, R. C.; Wang, T.; Lo, W. H. Y.;
Huang, S.-Z.;

[135056]Zeng, Y.-T.; Yuan, L.-F.; Liu, S.-R.; Woo, S. L. C.: Identifi-
cation

[135057]of three novel missense PKU mutations among Chinese.
Genomics 13:

[135058]894–895, 1992.

[135059]

[135060]154. Lichter-Konecki, U.; Konecki, D. S.; DiLella, A. G.;

Brayton,

[135061]K.; Marvit, J.; Hahn, T. M.; Trefz, F. K.; Woo, S. L. C.:

Phenylalanine

[135062]hydroxylase deficiency caused by a single base substitution in an

[135063]exon of the human phenylalanine hydroxylase gene. Biochemistry 27:

[135064]2881-2885, 1988.

[135065]

[135066]155. Lidsky, A. S.; Guttler, F.; Woo, S. L. C.: Prenatal diagnosis

[135067]of classic phenylketonuria by DNA analysis. Lancet I: 549-551, 1985.

[135068]

[135069]156. Lidsky, A. S.; Law, M. L.; Morse, H. G.; Kao, F.-T.; Rabin, M.;

[135070]Ruddle, F. H.; Woo, S. L. C.: Regional mapping of the phenylalanine

[135071]hydroxylase gene and the phenylketonuria locus in the human genome. Proc.

[135072]Nat. Acad. Sci. 82: 6221-6225, 1985.

[135073]

[135074]157. Lidsky, A. S.; Ledley, F. D.; DiLella, A. G.; Kwok, S. C. M.;

[135075]Daiger, S. P.; Robson, K. J. H.; Woo, S. L. C.: Extensive restriction

[135076]site polymorphism at the human phenylalanine hydroxylase locus and

[135077]application in prenatal diagnosis of phenylketonuria. Am. J. Hum.

[135078]Genet. 37: 619–634, 1985.

[135079]

[135080]158. Lidsky, A. S.; Robson, K. J. H.; Thirumalachary, C.; Barker,

[135081]P. E.; Ruddle, F. H.; Woo, S. L. C.: The PKU locus in man is on chromosome

[135082]12. Am. J. Hum. Genet. 36: 527–533, 1984.

[135083]

[135084]159. Lin, C.-H.; Hsiao, K.-J.; Tsai, T.-F.; Chao, H.-K.; Su, T.-S.

[135085]: Identification of a missense phenylketonuria mutation at codon 408

[135086]in Chinese. Hum. Genet. 89: 593–596, 1992.

[135087]

[135088]160. Liu, S. R.; Zuo, Q. H.: Newborn screening for
phenylketonuria

[135089]in eleven districts. Chinese Med. J. 99: 113–118, 1986.

[135090]

[135091]161. Lyonnet, S.; Caillaud, C.; Rey, F.; Berthelon, M.;
Frezal, J.;

[135092]Rey, J.; Munnich, A.: Molecular genetics of phenylke-
tonuria in Mediterranean

[135093]countries: a mutation associated with partial phenylala-
nine hydroxylase

[135094]deficiency. Am. J. Hum. Genet. 44: 511–517, 1989.

[135095]

[135096]162. Lyonnet, S.; Melle, D.; de Braekeleer, M.; Lafram-
boise, R.; Rey,

[135097]F.; John, S. W. M.; Berthelon, M.; Berthelot, J.; Journal, H.;
Le

[135098]Marec, B.; Parent, P.; de Parscau, L.; Saudubray, J.-M.;
Rozen, R.;

[135099]Rey, J.; Munnich, A.; Scriver, C. R.: Time and space clusters
of

[135100]the French-Canadian M1V phenylketonuria mutation in
France. Am. J.

[135101]Hum. Genet. 51: 191-196, 1992.

[135102]

[135103]163. Marvit, J.; DiLella, A. G.; Brayton, K.; Ledley, F. D.;
Robson,

[135104]K. J. H.; Woo, S. L. C.: GT to AT transition at a splice donor
site

[135105]causes skipping of the preceding exon in phenylketonuria.
Nucleic

[135106]Acids Res. 15: 5613-5628, 1987.

[135107]

[135108]164. Matalon, R.; Justice, P.; Deanching, M. N.: Phenylala-
nine hydroxylase

[135109]in human placenta: novel system for study of phenylke-
tonuria. (Letter) Lancet I:

[135110]853-854, 1977.

[135111]

[135112]165. McDonald, J. D.; Bode, V. C.; Dove, W. F.; Shedlovsky,
A.: Pah(hph-5):

[135113]a mouse mutant deficient in phenylalanine hydroxylase.

Proc. Nat.

[135114] Acad. Sci. 87: 1965–1967, 1990.

[135115]

[135116] 166. McDonald, J. D.; Charlton, C. K.: Characterization of mutations

[135117] at the mouse phenylalanine hydroxylase locus. Genomics 39: 402–405,

[135118] 1997.

[135119]

[135120] 167. Melle, D.; Verelst, P.; Rey, F.; Berthelon, M.; Francois, B.;

[135121] Munnich, A.; Lyonnet, S.: Two distinct mutations at a single BamHI

[135122] site in phenylketonuria. J. Med. Genet. 28: 38–40, 1991.

[135123]

[135124] 168. Menkes, J. H.; Aeberhard, E.: Maternal phenylketonuria. J.

[135125] Pediat. 74: 924–931, 1969.

[135126]

[135127] 169. Murphey, R. M.: Phenylketonuria (PKU) and the single

gene: an

[135128]old story retold. Behav. Genet. 13: 141–157, 1983.

[135129]

[135130]170. Nowacki, P.; Byck, S.; Prevost, L.; Scriver, C. R.: The
PAH

[135131]mutation analysis consortium database: update 1996. Nu-
cleic Acids

[135132]Res. 25: 139–142, 1997.

[135133]

[135134]171. Nyhan, W. L.: Fifty years ago: Asbjorn Folling and
phenylketonuria. Trends

[135135]Biochem. Sci. 9: 71–72, 1984.

[135136]

[135137]172. O'Connell, P.; Leppert, M.; Hoff, M.; Kumlin, E.;
Thomas, W.;

[135138]Cai, G.; Law, M.; White, R.: A linkage map for human chro-
mosome 12.

[135139](Abstract) Am. J. Hum. Genet. 37: A169 only, 1985.

[135140]

[135141]173. O'Flynn, M. E.; Tillman, P.; Hsia, D. Y.-Y.: Hyper-

phenylalaninemia

[135142]without phenylketonuria. Am. J. Dis. Child. 113: 22-30,
1967.

[135143]

[135144]174. Okano, Y.; Asada, M.; Kang, Y.; Nishi, Y.; Hase, Y.;
Oura, T.;

[135145]Isshiki, G.: Molecular characterization of phenylketonuria
in Japanese

[135146]patients. Hum. Genet. 103: 613-618, 1998.

[135147]

[135148]175. Okano, Y.; Eisensmith, R. C.; Dasovich, M.; Wang, T.;
Guttler,

[135149]F.; Woo, S. L. C.: A prevalent missense mutation in North-
ern Europe

[135150]associated with hyperphenylalaninaemia. Europ. J. Pediat.
150: 347-352,

[135151]1991.

[135152]

[135153]176. Okano, Y.; Wang, T.; Eisensmith, R. C.; Guttler, F.;
Woo, S.

[135154]L. C.: Recurrent mutation in the human phenylalanine hy-

droxylase

[135155]gene. Am. J. Hum. Genet. 46: 919–924, 1990.

[135156]

[135157]177. Okano, Y.; Wang, T.; Eisensmith, R. C.; Longhi, R.;
Riva, E.;

[135158]Giovannini, M.; Cerone, R.; Romano, C.; Woo, S. L. C.:
Phenylketonuria

[135159]missense mutations in the Mediterranean. Genomics 9:
96–103, 1991.

[135160]

[135161]178. Okano, Y.; Wang, T.; Eisensmith, R. C.; Steinmann, B.;
Gitzelmann,

[135162]R.; Woo, S. L. C.: Missense mutations associated with RFLP
haplotypes

[135163]1 and 4 of the human phenylalanine hydroxylase gene.
Am. J. Hum.

[135164]Genet. 46: 18–25, 1990.

[135165]

[135166]179. Okano, Y.; Wang, T.; Eisensmith, R. C.; Woo, S. L. C.:
PKU mutations

[135167]among Caucasians. (Abstract) Am. J. Hum. Genet. 45:

A211 only, 1989.

[135168]

[135169]180. Paine, R. S.: The variability in manifestations of untreated

[135170]patients with phenylketonuria (phenylpyruvic aciduria).
Pediatrics 20:

[135171]290-302, 1957.

[135172]

[135173]181. Paul, T. D.; Brandt, I. K.; Elsas, L. J.; Jackson, C. E.;
Mamunes,

[135174]P.; Nance, C. S.; Nance, W. E.: Phenylketonuria heterozygote detection

[135175]in families with affected children. Am. J. Hum. Genet. 30:
293-301,

[135176]1978.

[135177]

[135178]182. Paul, T. D.; Brandt, I. K.; Elsas, L. J.; Jackson, C. E.;
Nance,

[135179]C. S.; Nance, W. E.: Linkage analysis using heterozygote detection

[135180]in phenylketonuria. Clin. Genet. 16: 217-232, 1979.

[135181]

[135182]183. Perez, B.; Desviat, L. R.; Die, M.; Cornejo, V.;
Chamoles, N.

[135183]A.; Nicolini, H.; Ugarte, M.: Presence of the Mediterranean
PKU mutation

[135184]IVS10 in Latin America. Hum. Molec. Genet. 2:
1289–1290, 1993.

[135185]

[135186]184. Perez, B.; Desviat, L. R.; Die, M.; Ugarte, M.: Mutation
analysis

[135187]of phenylketonuria in Spain: prevalence of two Mediter-
ranean mutations. Hum.

[135188]Genet. 89: 341–342, 1992.

[135189]

[135190]185. Perry, T. L.; Hansen, S.; Tischler, B.; Bunting, R.; Dia-
mond,

[135191]S.: Glutamine depletion in phenylketonuria: possible cause
of the

[135192]mental defect. New Eng. J. Med. 282: 761–766, 1970.

[135193]

[135194]186. Pitt, D. B.; O'Day, J.: Phenylketonuria does not cause

cataracts. Europ.

[135195]J. Pediat. 150: 661–664, 1991.

[135196]

[135197]187. Ramus, S. J.; Forrest, S. M.; Cotton, R. G. H.: Illegitimate

[135198]transcription of phenylalanine hydroxylase for detection of mutations

[135199]in patients with phenylketonuria. Hum. Mutat. 1: 154–158, 1992.

[135200]

[135201]188. Ramus, S. J.; Forrest, S. M.; Saleeba, J. A.; Cotton, R. G. H.

[135202]: CpG hotspot causes second mutation in codon 408 of the phenylalanine

[135203]hydroxylase gene. Hum. Genet. 90: 147–148, 1992.

[135204]

[135205]189. Rao, D. C.; Keats, B. J.; Lalouel, J. M.; Morton, N. E.; Yee,

[135206]S.: A maximum likelihood map of chromosome 1. Am. J. Hum. Genet. 31:

[135207]680–696, 1979.

[135208]

[135209]190. Rey, F.; Berthelon, M.; Caillaud, C.; Lyonnet, S.;
Abadie, V.;

[135210]Blandin-Savoja, F.; Feingold, J.; Saudubray, J.-M.; Frezal,
J.; Munnich,

[135211]A.; Rey, J.: Clinical and molecular heterogeneity of phenyl-
alanine

[135212]hydroxylase deficiencies in France. Am. J. Hum. Genet. 43:
914-921,

[135213]1988.

[135214]

[135215]191. Riess, O.; Michel, A.; Speer, A.; Cobet, G.; Coutelle,
C.: Introduction

[135216]of genomic diagnosis of classical phenylketonuria to the
health care

[135217]system of the German Democratic Republic. Clin. Genet.
32: 209-215,

[135218]1987.

[135219]

[135220]192. Rohr, F. J.; Doherty, L. B.; Waisbren, S. E.; Bailey, I. V.;

[135221]Ampola, M. G.; Benacerraf, B.; Levy, H. L.: New England

maternal

[135222]PKU project: prospective study of untreated and treated pregnancies

[135223]and their outcomes. J. Pediat. 110: 391–398, 1987.

[135224]

[135225]193. Romeo, G.; Menozzi, P.; Ferlini, A.; Prosperi, L.; Cerone, R.;

[135226]Scalisi, S.; Romano, C.; Antonozzi, I.; Riva, E.; Piceni Sereni, L.;

[135227]Zammarchi, E.; Lenzi, G.; Sartorio, R.; Andria, G.; Cioni, M.; Fois,

[135228]A.; Burroni, M.; Burlina, A. B.; Carnevale, F.: Incidence of classic

[135229]PKU in Italy estimated from consanguineous marriages and from neonatal

[135230]screening. Clin. Genet. 24: 339–345, 1983.

[135231]

[135232]194. Rosenblatt, D. S.; Scriver, C. R.: Heterogeneity in genetic

[135233]control of phenylalanine metabolism in man. Nature 218: 677–678,

[135234]1968.

[135235]

[135236]195. Rouse, B.; Azen, C.; Koch, R.; Matalon, R.; Hanley, W.;
de la

[135237]Cruz, F.; Trefz, F.; Friedman, E.; Shifrin, H.: Maternal
phenylketonuria

[135238]collaborative study (MPKUCS) offspring: facial anomalies,
malformations,

[135239]and early neurological sequelae. Am. J. Med. Genet. 69:
89–95, 1997.

[135240]

[135241]196. Rouse, B.; Matalon, R.; Koch, R.; Azen, C.; Levy, H.;
Hanley,

[135242]W.; Trefz, F.; de la Cruz, F.: Maternal phenylketonuria syn-
drome:

[135243]congenital heart defects, microcephaly, and developmen-
tal outcomes. J.

[135244]Pediat. 136: 57–61, 2000.

[135245]

[135246]197. Sarkissian, C. N.; Shao, Z.; Blain, F.; Peevers, R.; Su,
H.;

[135247]Heft, R.; Chang, T. M. S.; Scriver, C. R.: A different ap-
proach to

[135248] treatment of phenylketonuria: phenylalanine degradation
with recombinant

[135249] phenylalanine ammonia lyase. Proc. Nat. Acad. Sci. 96:
2339–2344,

[135250] 1999.

[135251]

[135252] 198. Saugstad, L. F.: Frequency of phenylketonuria in Nor-
way. Clin.

[135253] Genet. 7: 40–51, 1975.

[135254]

[135255] 199. Saugstad, L. F.: Anthropological significance of
phenylketonuria. Clin.

[135256] Genet. 7: 52–61, 1975.

[135257]

[135258] 200. Scott, T. M.; Fyfe, W. M.; Hart, D. M.: Maternal
phenylketonuria:

[135259] abnormal baby despite low phenylalanine diet during
pregnancy. Arch.

[135260] Dis. Child. 55: 634–649, 1980.

[135261]

[135262]201. Scriver, C. R.: Personal Communication. Montreal,
Quebec, Canada

[135263]1974.

[135264]

[135265]202. Scriver, C. R.; Clow, C. L.: Phenylketonuria and other
phenylalanine

[135266]hydroxylation mutants in man. Ann. Rev. Genet. 14:
179–202, 1980.

[135267]

[135268]203. Scriver, C. R.; Clow, C. L.: Phenylketonuria: epitome
of human

[135269]biochemical genetics. New Eng. J. Med. 303: 1336–1342
and 1394–1400,

[135270]1980.

[135271]

[135272]204. Scriver, C. R.; Eisensmith, R. C.; Woo, S. L. C.; Kauf-
man, S.

[135273]: The hyperphenylalaninemias of man and mouse. Annu.
Rev. Genet. 28:

[135274]141–165, 1994.

[135275]

[135276]205. Scriver, C. R.; Gregory, D. M.; Sovetts, D.; Clow, C. L.;
Tissenbaum,

[135277]G.: Normal plasma free amino acid values in adults: the
influence

[135278]of some common physiological variables. Metabolism 34:
868–873,

[135279]1985.

[135280]

[135281]206. Scriver, C. R.; Kaufman, S.; Woo, S. L. C.: The hyper-
phenylalaninemias.In:

[135282]Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D.: The
Metabolic

[135283]Basis of Inherited Disease. New York: McGraw–Hill (pub.)
(6th

[135284]ed.) I: 1989. Pp. 495–546.

[135285]

[135286]207. Shimizu, A.; Sakai, Y.; Ohno, K.; Masaki, S.; Kuwano,
R.; Takahashi,

[135287]Y.; Miyashita, N.; Watanabe, T.: A molecular genetic link-
age map

[135288]of mouse chromosome 10, including the Myb, S100b, Pah,
Sl, and Ifg

[135289]genes. Biochem. Genet. 30: 529–535, 1992.

[135290]

[135291]208. Smith, C. B.; Kang, J.: Cerebral protein synthesis in a genetic

[135292]mouse model of phenylketonuria. Proc. Nat. Acad. Sci. 97: 11014–11019,

[135293]2000.

[135294]

[135295]209. Smith, I.; Macartney, F. J.; Erdohazi, M.; Pincott, J. R.; Wolff,

[135296]O. H.; Brenton, D. P.; Biddle, S. A.; Fairweather, D. V. I.; Dobbing,

[135297]J.: Fetal damage despite low-phenylalanine diet after conception

[135298]in a phenylketonuric woman. Lancet I: 17–19, 1979.

[135299]

[135300]210. Steffens, C.: No difference in dermatoglyphics of fingers and

[135301]palms between phenylketonuria patients and controls. (Letter) Hum.

[135302]Genet. 69: 195 only, 1985.

[135303]

[135304]211. Stegink, L. D.; Filer, L. J., Jr.; Baker, G. L.; Bell, E. F.;

[135305]Ziegler, E. E.; Brummel, M. C.; Krause, W. L.: Repeated in-
gestion

[135306]of aspartame-sweetened beverage: effect on plasma
amino acid concentrations

[135307]in individuals heterozygous for phenylketonuria.

Metabolism 38:

[135308]78-84, 1989.

[135309]

[135310]212. Stuhmann, M.; Riess, O.; Monch, E.; Kurdoglu, G.:

Haplotype

[135311]analysis of the phenylalanine hydroxylase gene in Turkish
phenylketonuria

[135312]families. Clin. Genet. 36: 117-121, 1989.

[135313]

[135314]213. Superti-Furga, A.; Steinmann, B.; Duc, G.; Gitzel-
mann, R.: Maternal

[135315]phenylketonuria syndrome in cousins caused by mild, un-
recognized phenylketonuria

[135316]in their mothers homozygous for the phenylalanine hy-

droxylase arg261-to-gln

[135317]mutation. *Europ. J. Pediat.* 150: 493–497, 1991.

[135318]

[135319]214. Svensson, E.; Andersson, B.; Hagenfeldt, L.: Two mutations within

[135320]the coding sequence of the phenylalanine hydroxylase gene. *Hum. Genet.* 85:

[135321]300–304, 1990.

[135322]

[135323]215. Svensson, E.; Eisensmith, R. C.; Dworniczak, B.; von Dobeln,

[135324]U.; Hagenfeldt, L.; Horst, J.; Woo, S. L. C.: Two missense mutations

[135325]causing mild hyperphenylalaninemia associated with DNA haplotype 12. *Hum.*

[135326]*Mutat.* 1: 129–137, 1992.

[135327]

[135328]216. Takahashi, K.; Kure, S.; Matsubara, Y.; Narisawa, K.: Novel

[135329]phenylketonuria mutation detected by analysis of ectopically transcribed

[135330]phenylalanine hydroxylase mRNA from lymphoblast.

(Letter) Lancet 340:

[135331]1473 only, 1992.

[135332]

[135333]217. Teebi, A. S.; Al-Awadi, S. A.; Farag, T. I.; Naguib, K.
K.; El-Khalifa,

[135334]M. Y.: Phenylketonuria in Kuwait and Arab countries. Eu-
rop. J. Pediat. 146:

[135335]59-60, 1987.

[135336]

[135337]218. Thompson, A. J.; Tillotson, S.; Smith, I.; Kendall, B.;
Moore,

[135338]S. G.; Brenton, D. P.: Brain MRI changes in phenylke-
tonuria: associations

[135339]with dietary status. Brain 116: 811-821, 1993.

[135340]

[135341]219. Tourian, A. Y.; Sidbury, J. B., Jr.: Phenylketonuria.In:
Stanbury,

[135342]J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: Metabolic Basis
of

[135343]Inherited Disease. New York: McGraw-Hill (pub.) (4th ed.):

1978.

[135344]Pp. 240–255.

[135345]

[135346]220. Tsai, T.–F.; Hsiao, K.–J.; Su, T.–S.: Phenylketonuria
mutation

[135347]in Chinese haplotype 44 identical with haplotype 2 muta-
tion in northern–European

[135348]Caucasians. Hum. Genet. 84: 409–411, 1990.

[135349]

[135350]221. Tyfield, L. A.; Meredith, A. L.; Osborn, M. J.; Harper,
P. S.

[135351]: Identification of the haplotype pattern associated with
the mutant

[135352]PKU allele in the Gypsy population of Wales. J. Med. Genet.
26:

[135353]499–503, 1989.

[135354]

[135355]222. Tyfield, L. A.; Stephenson, A.; Cockburn, F.; Harvie,
A.; Bidwell,

[135356]J. L.; Wood, N. A. P.; Pilz, D. T.; Harper, P.; Smith, I.: Se-
quence

[135357]variation at the phenylalanine hydroxylase gene in the
British Isles. Am.

[135358]J. Hum. Genet. 60: 388–396, 1997.

[135359]

[135360]223. Ullrich, K.; Weglage, J.; Schuierer, G.; Funders, B.;
Pietsch,

[135361]M.; Koch, H. G.; Hahn-Ullrich, H.: Cranial MRI in PKU: eval-
uation

[135362]of a critical threshold for blood phenylalanine. (Letter)
Neuropediatrics 25:

[135363]278–279, 1994.

[135364]

[135365]224. Usha, R.; Uma, R.; Farag, T. I.; Girish, Y.; Al-Ghanim,
M. M.

[135366]A.; Al-Najdi, K.; Al-Awadi, S. A.; El-Badramany, M. H.: Late
diagnosis

[135367]of phenylketonuria in a Bedouin mother. Am. J. Med.
Genet. 44: 713–715,

[135368]1992.

[135369]

[135370]225. Vajro, P.; Strisciuglio, P.; Houssin, D.; Huault, G.;

Laurent,

[135371]J.; Alvarez, F.; Bernard, O.: Correction of phenylketonuria after

[135372]liver transplantation in a child with cirrhosis. (Letter) New Eng.

[135373]J. Med. 329: 363 only, 1993.

[135374]

[135375]226. van Spronsen, F. J.; Reijngoud, D.-J.; Smit, G. P. A.; Nagel,

[135376]G. T.; Stellaard, F.; Berger, R.; Heymans, H. S. A.: Phenylketonuria:

[135377]the in vivo hydroxylation rate of phenylalanine into tyrosine is decreased. J.

[135378]Clin. Invest. 101: 2875-2880, 1998.

[135379]

[135380]227. Waisbren, S. E.; Hanley, W.; Levy, H. L.; Shifrin, H.; Allred,

[135381]E.; Azen, C.; Chang, P.-N.; Cipic-Schmidt, S.; de la Cruz, F.; Hall,

[135382]R.; Matalon, R.; Nanson, J.; Rouse, B.; Trefz, F.; Koch, R.: Outcome

[135383]at age 4 years in offspring of women with maternal

phenylketonuria:

[135384]the maternal PKU collaborative study. J.A.M.A. 283:
756-762, 2000.

[135385]

[135386]228. Wang, T.; Okano, Y.; Eisensmith, R.; Huang, S.-Z.;
Zeng, Y.-T.;

[135387]Lo, W. H. Y.; Woo, S. L. C.: Molecular genetics of phenylke-
tonuria

[135388]in Orientals: linkage disequilibrium between a termination
mutation

[135389]and haplotype 4 of the phenylalanine hydroxylase gene.
Am. J. Hum.

[135390]Genet. 45: 675-680, 1989.

[135391]

[135392]229. Wang, T.; Okano, Y.; Eisensmith, R. C.; Fekete, G.;
Schuler,

[135393]D.; Berencsi, G.; Nasz, I.; Woo, S. L. C.: Molecular genetics
of

[135394]PKU in Eastern Europe: a nonsense mutation associated
with haplotype

[135395]4 of the phenylalanine hydroxylase gene. Somat. Cell
Molec. Genet. 16:

[135396]85–89, 1990.

[135397]

[135398]230. Wang, T.; Okano, Y.; Eisensmith, R. C.; Harvey, M. L.;
Lo, W.

[135399]H. Y.; Huang, S.-Z.; Zeng, Y.-T.; Yuan, L.-F.; Furuyama, J.;
Oura,

[135400]T.; Sommer, S. S.; Woo, S. L. C.: Founder effect of a preva-
lent phenylketonuria

[135401]mutation in the Oriental population. Proc. Nat. Acad. Sci.
88: 2146–2150,

[135402]1991.

[135403]

[135404]231. Wang, T.; Okano, Y.; Eisensmith, R. C.; Lo, W. H. Y.;
Huang,

[135405]S.-Z.; Zeng, Y.-T.; Woo, S. L. C.: Identification of a novel
phenylketonuria

[135406](PKU) mutation in the Chinese: further evidence for multi-
ple origins

[135407]of PKU in Asia. Am. J. Hum. Genet. 48: 628–630, 1991.

[135408]

[135409]232. Wang, T.; Okano, Y.; Eisensmith, R. C.; Lo, W. H. Y.;

Huang,

[135410]S.-Z.; Zeng, Y.-T.; Yuan, L.-F.; Liu, S.-R.; Woo, S. L. C.:

Identification

[135411]of three novel PKU mutations among Chinese: evidence

for recombination

[135412]or recurrent mutation at the PAH locus. Genomics 13:

230-231, 1992.

[135413]

[135414]233. Wang, T.; Okano, Y.; Eisensmith, R. C.; Lo, W. H. Y.;

Huang,

[135415]S.-Z.; Zeng, Y.-T.; Yuan, L.-F.; Liu, S.-R.; Woo, S. L. C.:

Missense

[135416]mutations prevalent in Orientals with phenylketonuria:

molecular characterization

[135417]and clinical implications. Genomics 10: 449-456, 1991.

[135418]

[135419]234. Wang, T.; Woo, S. L. C.: Personal Communication.

Houston, Tex.

[135420]1/20/1990.

[135421]

[135422]235. Waters, P. J.; Parniak, M. A.; Akerman, B. R.; Scriver,

C. R.

[135423]; Characterization of phenylketonuria missense substitutions, distant

[135424]from the phenylalanine hydroxylase active site, illustrates a paradigm

[135425]for mechanism and potential modulation of phenotype. Molec. Genet.

[135426]Metab. 69: 101–110, 2000.

[135427]

[135428]236. Weglage, J.; Grenzebach, M.; Pietsch, M.; Feldmann, R.; Linnenbank,

[135429]R.; Denecke, J.; Koch, H. G.: Behavioural and emotional problems

[135430]in early-treated adolescents with phenylketonuria in comparison with

[135431]diabetic patients and healthy controls. J. Inherit. Metab. Dis. 23:

[135432]487–496, 2000.

[135433]

[135434]237. Weglage, J.; Pietsch, M.; Denecke, J.; Sprinz, A.; Feldmann,

[135435]R.; Grenzebach, M.; Ullrich, K.: Regression of neuropsych-

chological

[135436]deficits in early-treated phenylketonurics during adolescence. J.

[135437]Inherit. Metab. Dis. 22: 693–705, 1999.

[135438]

[135439]238. Weinstein, M.; Eisensmith, R. C.; Abadie, V.; Avigad, S.; Lyonnet,

[135440]S.; Schwartz, G.; Munnich, A.; Woo, S. L. C.; Shiloh, Y.: A missense

[135441]mutation, S349P, completely inactivates phenylalanine hydroxylase

[135442]in North African Jews with phenylketonuria. Hum. Genet. 90: 645–649,

[135443]1993.

[135444]

[135445]239. Woo, S. L. C.: Personal Communication. Houston, Tex. 1/11/1983.

[135446]

[135447]240. Woo, S. L. C.: Collation of RFLP haplotypes at the human phenylalanine

[135448]hydroxylase (PAH) locus. (Letter) Am. J. Hum. Genet. 43:

781-783,

[135449]1988.

[135450]

[135451]241. Woo, S. L. C.: Molecular basis and population genetics of phenylketonuria. Biochemistry 28:

[135452]1-7, 1989.

[135453]

[135454]242. Woo, S. L. C.; Chandra, T.; Stackhouse, R.; Robson, K. J. H.

[135455]: Analysis of phenylketonuria (PKU) by cloning and characterization

[135456]of the human phenylalanine hydroxylase gene. (Abstract) Am. J. Hum.

[135457]Genet. 34: 168A only, 1982.

[135458]

[135459]243. Woo, S. L. C.; Lidsky, A.; Law, M.; Kao, F. T.: Regional mapping

[135460]of the human phenylalanine hydroxylase gene and PKU locus to 12q21-qter.

[135461](Abstract) Am. J. Hum. Genet. 36: 210S only, 1984.

[135462]

[135463]244. Woo, S. L. C.; Lidsky, A. S.; Guttler, F.; Chandra, T.;
Robson,

[135464]K. J. H.: Cloned human phenylalanine hydroxylase gene
permits prenatal

[135465]diagnosis and carrier detection of classical phenylke-
tonuria. Nature 306:

[135466]151-155, 1983.

[135467]

[135468]245. Woo, S. L. C.; Lidsky, A. S.; Guttler, F.; Thiru-
malachary, C.;

[135469]Robson, K. J. H.: Prenatal diagnosis of classical phenylke-
tonuria

[135470]by gene mapping. J.A.M.A. 251: 1998-2002, 1984.

[135471]

[135472]246. Woolf, L. I.: The heterozygote advantage in
phenylketonuria.

[135473](Letter) Am. J. Hum. Genet. 38: 773-775, 1986.

[135474]

[135475]247. Woolf, L. I.; Cranston, W. I.; Goodwin, B. L.: Genetics
of phenylketonuria.

[135476]I. Heterozygosity for phenylketonuria. II. Third allele at the

phenylalanine

[135477]hydroxylase locus in man. Nature 213: 882–885, 1967.

[135478]

[135479]248. Woolf, L. I.; Goodwin, B. L.; Cranston, W. I.; Wade, D. N.; Woolf,

[135480]F.; Hudson, F. P.; McBean, M. S.: A third allele at the phenylalanine–hydroxylase

[135481]locus in mild phenylketonuria (hyperphenylalaninaemia). Lancet I:

[135482]114–117, 1968.

[135483]

[135484]249. Woolf, L. I.; McBean, M. S.; Woolf, F. M.; Cahalane, S. F.:

[135485]Phenylketonuria as a balanced polymorphism: the nature of the heterozygote

[135486]advantage. Ann. Hum. Genet. 38: 461–469, 1975.

[135487]

[135488]250. Yu, J. S.; O'Halloran, M. T.: Atypical phenylketonuria in a

[135489]family with a phenylketonuric mother. Pediatrics 46: 707–711, 1970.

[135490]

[135491]251. Zschocke, J.; Graham, C. A.; Stewart, F. J.; Carson, D. J.; Nevin,

[135492]N. C.: Non-phenylketonuria hyperphenylalaninaemia in Northern Ireland:

[135493]frequent mutation allows screening and early diagnosis. Hum. Mutat. 4:

[135494]114–118, 1994.

[135495]

[135496]252. Zschocke, J.; Mallory, J. P.; Eiken, H. G.; Nevin, N. C.: Phenylketonuria

[135497]and the peoples of Northern Ireland. Hum. Genet. 100: 189–194, 1997.

[135498]

[135499]253. Zschocke, J.; Quak, E.; Knauer, A.; Fritz, B.; Aslan, M.; Hoffmann,

[135500]G. F.: Large heterozygous deletion masquerading as homozygous missense

[135501]mutation: a pitfall in diagnostic mutation analysis. J. Inherit.

[135502]Metab. Dis. 22: 687–692, 1999.

[135503]

[135504]254. Zygulska, M.; Eigel, A.; Aulehla-Scholz, C.; Pietrzyk, J. J.;

[135505]Horst, J.: Molecular analysis of PKU haplotypes in the population

[135506]of southern Poland. Hum. Genet. 86: 292-294, 1991.

[135507]

[135508]255. Zygulska, M.; Eigel, A.; Dworniczak, B.; Sutkowska, A.; Pietrzyk,

[135509]J. J.; Horst, J.: Phenylketonuria in Poland: 66% of PKU alleles are

[135510]caused by three mutations. Hum. Genet. 88: 91-94, 1991.

[135511]24109.

[135512]Bessant, D. A. R.; Payne, A. M.; Plant, C.; Bird, A. C.; Bhat-tacharya,

[135513]S. S.: Further refinement of the Usher 2A locus at 1q41. J. Med.

[135514]Genet. 35: 773-774, 1998.

[135515]24110.

[135516]Konrad, M.; Saunier, S.; Heidet, L.; Silbermann, F.; Bennisy,

[135517]F.; Calado, J.; Le Paslier, D.; Broyer, M.; Gubler, M.-C.;
Antignac,

[135518]C.: Large homozygous deletions of the 2q13 region are a
major cause

[135519]of juvenile nephronophthisis. Hum. Molec. Genet. 5:
367-371, 1996.

[135520]24111.

[135521]Degos, L.; Dautigny, A.; Brouet, J. C.; Colombani, M.; Ar-
daillou,

[135522]N.; Caen, J. P.; Colombani, J.: A molecular defect in throm-
basthenic

[135523]platelets. J. Clin. Invest. 56: 236-240, 1975.

[135524]24112.

[135525]Djaffar, I.; Vilette, D.; Pidard, D.; Wautier, J.-L.; Rosa, J.-P.

[135526]: Human platelet antigen 3 (HPA-3): localization of the de-
terminant

[135527]of the alloantibody Lek(a)(HPA-3a) to the C-terminus of
platelet glycoprotein

[135528]IIb heavy chain and contribution of O-linked carbohy-
drates. Thromb.

[135529]Haemost. 69: 485-489, 1993.

[135530]24113.

[135531]Friedman, L. L.; Bowie, E. J. W.; Thompson, J. H., Jr.;
Brown,

[135532]A. L., Jr.; Owen, C. A., Jr.: Familial Glanzmann's thrombas-
thenia. Mayo

[135533]Clin. Proc. 39: 908-918, 1964.

[135534]24114.

[135535]Giltay, J. C.; Leeksa, O. C.; Breederveld, C.; van Mourik, J.

[135536]A.: Normal synthesis and expression of endothelial IIb/IIIa
in Glanzmann's

[135537]thrombasthenia. Blood 69: 809-812, 1987.

[135538]24115.

[135539]Grimaldi, C. M.; Chen, F.; Wu, C.; Weiss, H. J.; Coller, B. S.;

[135540]French, D. L.: Glycoprotein IIb leu214pro mutation pro-
duces Glanzmann

[135541]thrombasthenia with both quantitative and qualitative ab-
normalities

[135542]in GPIIb/IIIa. Blood 91: 1562-1571, 1998.

[135543]24116.

[135544]Gross, R.; Gerok, W.; Lohr, G. W.; Vogell, W.; Waller, H. D.;

[135545]Theopold, W.: Ueber die Natur der Thrombasthenie.

Thrombopathie Glanzmann-Naegeli. Klin.

[135546]Wschr. 38: 193-206, 1960.

[135547]24117.

[135548]Hathaway, W. E.: Bleeding disorders due to platelet dys-

function. Am.

[135549]J. Dis. Child. 121: 127-134, 1971.

[135550]24118.

[135551]Herrmann, F. H.; Meyer, M.; Gogstad, G. O.; Solum, N. O.:

Glycoprotein

[135552]IIb-IIIa complex in platelets of patients and heterozygotes

of Glanzmann's

[135553]thrombasthenia. Thromb. Res. 32: 615-622, 1983.

[135554]24119.

[135555]Herrmann, F. H.; Meyer, M.; Ihle, E.: Protein and glycopro-

tein

[135556]abnormalities in an unusual subtype of Glanzmann's

thrombasthenia. Haemostasis 12:

[135557]337-344, 1982.

[135558]24120.

[135559]Kanska, B.; Niewiarowski, S.; Ostrowski, L.; Poplawski, A.;
Prokopowicz,

[135560]J.: Macrothrombocytic thrombopathia. Clinical, coagula-
tion and hereditary

[135561]aspects. Thromb. Diath. Haemorrh. 10: 88-100, 1963.

[135562]24121.

[135563]Karpatkin, S.; Weiss, H. J.: Deficiency of glutathione per-
oxidase

[135564]associated with high levels of reduced glutathione in
Glanzmann's

[135565]thrombasthenia. New Eng. J. Med. 287: 1062-1066, 1972.

[135566]24122.

[135567]Kato, A.; Yamamoto, K.; Miyazaki, S.; Jung, S. M.; Moroi,
M.;

[135568]Aoki, N.: Molecular basis for Glanzmann's thrombasthenia
(GT) in

[135569]a compound heterozygote with glycoprotein IIb gene: a
proposal for

[135570]the classification of GT based on the biosynthetic pathway

of glycoprotein

[135571]Iib-IIIa complex. Blood 79: 3212-3218, 1992.

[135572]24123.

[135573]Khanduri, U.; Pulimood, R.; Sudarsanam, A.; Carman, R.
H.; Jadhav,

[135574]M.; Pereira, S.: Glanzmann's thrombasthenia: a review and
report

[135575]of 42 cases from South India. Thromb. Haemost. 46:
717-721, 1981.

[135576]24124.

[135577]Kunicki, T. J.; Aster, R. H.: Deletion of the platelet-specific

[135578]alloantigen PI(A1) from platelets in Glanzmann's throm-
basthenia. J.

[135579]Clin. Invest. 61: 1225-1231, 1978.

[135580]24125.

[135581]Lelong, J. C.: La thrombopathie de Glanzmann-Naegeli.
Paris:

[135582]R. Foulon et Cie. (pub.) 1960.

[135583]24126.

[135584]Levy, J. M.; Mayer, G.; Sacrez, R.; Ruff, R.; Francfort, J. J.;

[135585]Rodier, L.: Thrombasthenie de Glanzmann-Naegeli: etude
d'un groupe

[135586]ethnique a forte endogamie. Ann. Pediat. 18: 129-137,
1971.

[135587]24127.

[135588]Luo, X. Y.; Bray, P. F.; Magenis, R. E.: Precise localization

[135589]of the gene for platelet membrane glycoprotein IIb to
17q21.32 using

[135590]structural rearrangements of chromosome 17. (Abstract)
Cytogenet.

[135591]Cell Genet. 51: 1036 only, 1989.

[135592]24128.

[135593]Marx, R.; Jean, G.: Studien zur Pathogenese der Throm-
basthenie

[135594]Glanzmann-Naegeli. Klin. Wschr. 40: 942-953, 1962.

[135595]24129.

[135596]McEver, R. P.; Baenziger, J. U.; Majerus, P. W.: Isolation and

[135597]structural characterization of the polypeptide subunits of
membrane

[135598]glycoprotein IIb-IIIa from human platelets. Blood 59:
80-85, 1982.

[135599]24130.

[135600]McEver, R. P.; Baenziger, N. L.; Majerus, P. W.: Isolation
and

[135601]quantitation of the platelet membrane glycoprotein defi-
cient in thrombasthenia

[135602]using a monoclonal hybridoma antibody. J. Clin. Invest.
66: 1311-1318,

[135603]1980.

[135604]24131.

[135605]Meyer, M.; Herrmann, F. H.: Diversity of glycoprotein defi-
ciencies

[135606]in Glanzmann's thrombasthenia. Thromb. Haemost. 54:
626-629, 1985.

[135607]24132.

[135608]Moser, K.; Lechner, K.; Vinazzer, H.: A hitherto not de-
scribed

[135609]enzyme defect in thrombasthenia: glutathione reductase
deficiency. Thromb.

[135610]Diath. Haemorrh. 19: 46-52, 1968.

[135611]24133.

[135612]Nachman, R. L.: Thrombasthenia: immunologic evidence

of a platelet

[135613]protein abnormality. J. Lab. Clin. Med. 67: 411-419, 1966.

[135614]24134.

[135615]Newman, P. J.: Platelet GPIIb-IIIa: molecular variations and

[135616]alloantigens. Thromb. Haemost. 66: 111-118, 1991.

[135617]24135.

[135618]Nurden, A. T.; Caen, J. P.: An abnormal platelet glycoprotein

[135619]pattern in three cases of Glanzmann's thrombasthenia.
Brit. J. Haemat. 28:

[135620]253-260, 1974.

[135621]24136.

[135622]Eisensmith, R. C.; Woo, S. L. C.: Molecular basis of
phenylketonuria

[135623]and related hyperphenylalaninemias: mutations and polymorphisms in

[135624]the human phenylalanine hydroxylase gene. Hum. Mutat.
1: 13-23,

[135625]1992.

[135626]24137.

[135627]Eisensmith, R. C.; Woo, S. L. C.: Updated listing of haplo-
types

[135628]at the human phenylalanine hydroxylase (PAH) locus.
(Letter) Am.

[135629]J. Hum. Genet. 51: 1445-1448, 1992.

[135630]24138.

[135631]Eisensmith, R. C.; Woo, S. L. C.: Gene therapy for
phenylketonuria. Europ.

[135632]J. Pediat. 155 (suppl. 1): S16-S19, 1996.

[135633]24139.

[135634]Fisch, R. O.; Tagatz, G.; Stassart, J. P.: Gestational carrier-
-a

[135635]reproductive haven for offspring of mothers with
phenylketonuria (PKU):

[135636]an alternative therapy for maternal PKU. J. Inherit. Metab.
Dis. 16:

[135637]957-961, 1993.

[135638]24140.

[135639]Flatz, G.; Oelbe, M.; Herrmann, H.: Ethnic distribution of
phenylketonuria

[135640]in the north German population. Hum. Genet. 65:

396–399, 1984.

[135641]24141.

[135642]Folling, A.: Ueber Ausscheidung von Phenylbrenztrauben-
saeure

[135643]in den Harn als Stoffwechselanomalie in Verbindung mit
Imbezillitaet. Ztschr.

[135644]Physiol. Chem. 227: 169–176, 1934.

[135645]24142.

[135646]Forrest, S. M.; Dahl, H. H.; Howells, D. W.; Dianzani, I.;
Cotton,

[135647]R. G. H.: Mutation detection in phenylketonuria by using
chemical

[135648]cleavage of mismatch: importance of using probes from
both normal

[135649]and patient samples. Am. J. Hum. Genet. 49: 175–183,
1991.

[135650]24143.

[135651]Frankenburg, W. K.; Duncan, B. R.; Coffelt, R. W.; Koch, R.;
Coldwell,

[135652]J. G.; Son, C. D.: Maternal phenylketonuria: implications
for growth

[135653]and development. J. Pediat. 73: 560–570, 1968.

[135654]24144.

[135655]Friedman, P. A.; Fisher, D. B.; Kang, E. S.; Kaufman, S.: De-
tection

[135656]of hepatic phenylalanine 4-hydroxylase in classical
phenylketonuria. Proc.

[135657]Nat. Acad. Sci. 70: 552-556, 1973.

[135658]24145.

[135659]Gjetting, T.; Petersen, M.; Guldberg, P.; Guttler, F.: Mis-
sense

[135660]mutations in the N-terminal domain of human phenylala-
nine hydroxylase

[135661]interfere with binding of regulatory phenylalanine. Am. J.
Hum. Genet. 68:

[135662]1353-1360, 2001.

[135663]24146.

[135664]Greeves, L. G.; Patterson, C. C.; Carson, D. J.; Thom, R.;
Wolfenden,

[135665]M. C.; Zshock, J.; Graham, C. A.; Nevin, N. C.; Trimble, E.
R.:

[135666]Effect of genotype on changes in intelligence quotient af-
ter dietary

- [135667]relaxation in phenylketonuria and hyperphenylalani-
naemia. Arch. Dis.
- [135668]Child. 82: 216–221, 2000.
- [135669]24147.
- [135670]Gregory, D. M.; Sovetts, D.; Clow, C. L.; Sriver, C. R.:
Plasma
- [135671]free amino acid values in normal children and adoles-
cents. Metabolism 35:
- [135672]967–969, 1986.
- [135673]24148.
- [135674]Griffiths, P. V.; Demellweek, C.; Fay, N.; Robinson, P. H.;
Davidson,
- [135675]D. C.: Wechsler subscale IQ and subtest profile in early
treated
- [135676]phenylketonuria. Arch. Dis. Child. 82: 209–215, 2000.
- [135677]24149.
- [135678]Guldborg, P.; Henriksen, K. F.; Sipila, I.; Guttler, F.; de la
- [135679]Chapelle, A.: Phenylketonuria in a low incidence popula-
tion: molecular
- [135680]characterization of mutations in Finland. J. Med. Genet.
32: 976–978,

[135681]1995.

[135682]24150.

[135683]Guldberg, P.; Henriksen, K. F.; Thony, B.; Blau, N.; Guttler,

[135684]F.: Molecular heterogeneity of nonphenylketonuria hyper-
phenylalaninemia

[135685]in 25 Danish patients. Genomics 21: 453–455, 1994.

[135686]24151.

[135687]Guldberg, P.; Lou, H. C.; Henriksen, K. F.; Mikkelsen, I.;
Olsen,

[135688]B.; Holck, B.; Guttler, F.: A novel missense mutation in the
phenylalanine

[135689]hydroxylase gene of a homozygous Pakistani patient with
non-PKU hyperphenylalaninemia. Hum.

[135690]Molec. Genet. 2: 1061–1062, 1993.

[135691]24152.

[135692]Guldberg, P.; Zschocke, J.; Dagbjartsson, A.; Henriksen, K.
F.;

[135693]Guttler, F.: A molecular survey of phenylketonuria in Ice-
land: identification

[135694]of a founding mutation and evidence of predominant
Norse settlement. Europ.

[135695]J. Hum. Genet. 5: 376–381, 1997.

[135696]24153.

[135697]Guthrie, R.: The introduction of newborn screening for phenylketonuria:

[135698]a personal history. Europ. J. Pediat. 155 (suppl. 1): 4–5, 1996.

[135699]24154.

[135700]Guthrie, R.; Susi, A.: A simple phenylalanine method for detecting

[135701]phenylketonuria in large populations of newborn infants. Pediatrics 32:

[135702]338–343, 1963.

[135703]24155.

[135704]Guldborg, P.; Rey, F.; Zschocke, J.; Romano, V.; Francois, B.;

[135705]Michiels, L.; Ullrich, K.; Hoffmann, G. F.; Burgard, P.; Schmidt,

[135706]H.; Meli, C.; Riva, E.; Dianzani, I.; Ponzone, A.; Rey, J.; Gütler,

[135707]F.: A European multicenter study of phenylalanine hydroxylase deficiency:

- [135708]classification of 105 mutations and a general system for
genotype-based
- [135709]prediction of metabolic phenotype. Am. J. Hum. Genet. 63:
71-79,
- [135710]1998.
- [135711]24156.
- [135712]Guttler, F.: Hyperphenylalaninemia: diagnosis and classifi-
cation
- [135713]of the various types of phenylalanine hydroxylase defi-
ciency in childhood. Acta
- [135714]Paediat. Scand. 280 (suppl.): 1-80, 1980.
- [135715]24157.
- [135716]Guttler, F.; Azen, C.; Guldberg, P.; Romstad, A.; Hanley, W.
B.;
- [135717]Levy, H. L.; Matalon, R.; Rouse, B. M.; Trefz, F.; de la Cruz,
F.;
- [135718]Koch, R.: Relationship among genotype, biochemical phe-
notype, and
- [135719]cognitive performance in females with phenylalanine hy-
droxylase deficiency:
- [135720]report from the maternal phenylketonuria collaborative
study. Pediatrics 104:

[135721]258–262, 1999.

[135722]24158.

[135723]Guttler, F.; Hansen, G.: Heterozygote detection in phenylketonuria. Clin.

[135724]Genet. 11: 137–146, 1977.

[135725]24159.

[135726]Guttler, F.; Ledley, F. D.; Lidsky, A. S.; DiLella, A. G.; Sullivan,

[135727]S. E.; Woo, S. L. C.: Correlation between polymorphic DNA haplotypes

[135728]at phenylalanine hydroxylase locus and clinical phenotypes of phenylketonuria. J.

[135729]Pediat. 110: 68–71, 1987.

[135730]24160.

[135731]Guttler, F.; Woo, S. L. C.: Molecular genetics of PKU. J. Inherit.

[135732]Metab. Dis. 9 (suppl. 1): 58–68, 1986.

[135733]24161.

[135734]Hanley, W. B.; Clarke, J. T. R.; Schoonheydt, W.: Maternal phenylketonuria

[135735](PKU)--a review. Clin. Biochem. 20: 149-156, 1987.

[135736]24162.

[135737]Hanley, W. B.; Koch, R.; Levy, H. L.; Matalon, R.; Rouse, B.;

[135738]Azen, C.; de la Cruz, F.: The North American Maternal
Phenylketonuria

[135739]Collaborative Study, developmental assessment of the off-
spring: preliminary

[135740]report. Europ. J. Pediat. 155 (suppl. 1): 169-172, 1996.

[135741]24163.

[135742]Hanley, W. B.; Lee, A. W.; Hanley, A. J. G.; Lehotay, D. C.;
Austin,

[135743]V. J.; Schoonheydt, W. E.; Platt, B. A.; Clarke, J. T. R.: 'Hypo-
tyrosinemia'

[135744]in phenylketonuria. Molec. Genet. Metab. 69: 286-294,
2000.

[135745]24164.

[135746]Hertzberg, M.; Jahromi, K.; Ferguson, V.; Dahl, H. H. M.;
Mercer,

[135747]J.; Mickleson, K. N. P.; Trent, R. J.: Phenylalanine hydroxy-
lase

[135748]gene haplotypes in Polynesians: evolutionary origins and

absence of

[135749]alleles associated with severe phenylketonuria. Am. J.

Hum. Genet. 44:

[135750]382–387, 1989.

[135751]24165.

[135752]Hoang, L.; Byck, S.; Prevost, L.; Scriver, C. R.: PAH Muta-

tion

[135753]Analysis Consortium Database: a database for disease-

producing and

[135754]other allelic variation at the human PAH locus. Nucleic

Acids Res. 24:

[135755]127–131, 1996.

[135756]24166.

[135757]Hofman, K. J.; Steel, G.; Kazazian, H. H.; Valle, D.:

Phenylketonuria

[135758]in U.S. blacks: molecular analysis of the phenylalanine hy-

droxylase

[135759]gene. Am. J. Hum. Genet. 48: 791–798, 1991.

[135760]24167.

[135761]Garland, D.; Rao, P. V.; Del Corso, A.; Mura, U.; Zigler, J.

S.,

[135762]Jr.: Zeta-crystallin is a major protein in the lens of
Camelus dromedarius.

[135763]Arch. Biochem. Biophys. 285: 134-136, 1991.

[135764]24168.

[135765]Gonzalez, P.; Rao, P. V.; Zigler, J. S., Jr.: Organization of
[135766]the human zeta-crystallin/quinone reductase gene
(CRYZ). Genomics 21:
[135767]317-324, 1994.

[135768]24169.

[135769]Heinzmann, C.; Kojis, T. L.; Gonzalez, P.; Rao, P. V.; Zigler,
[135770]J. S., Jr.; Polymeropoulos, M. H.; Klisak, I.; Sparkes, R. S.;
Mohandas,
[135771]T.; Bateman, J. B.: Assignment of the zeta-crystallin gene
(CRYZ)
[135772]to human chromosome 1p22-p31 and identification of
restriction fragment
[135773]length polymorphisms. Genomics 23: 403-407, 1994.

[135774]24170.

[135775]Huang, Q.-L.; Russell, P.; Stone, S. H.; Zigler, J. S., Jr.:
Zeta-crystallin,

[135776]a novel lens protein from the guinea pig. Curr. Eye Res. 6:
725-732,

[135777]1987.

[135778]24171.

[135779]Rodriguez, I. R.; Gonzalez, P.; Zigler, J. S., Jr.; Borrás, T.

[135780]; A guinea-pig hereditary cataract contains a splice site
deletion

[135781]in a crystallin gene. Biochim. Biophys. Acta 1180: 44-52,
1992.

[135782]24172.

[135783]Morrison, N. A.; Yeoman, R.; Kelly, P. J.; Eisman, J. A.:
Contribution

[135784]of trans-acting factor alleles to normal physiological vari-
ability:

[135785]vitamin D receptor gene polymorphisms and circulating
osteocalcin. Proc.

[135786]Nat. Acad. Sci. 89: 6665-6669, 1992.

[135787]24173.

[135788]Pan, L. C.; Price, P. A.: The propeptide of rat bone
gamma-carboxyglutamic

[135789]acid protein shares homology with other vitamin K-

dependent protein

[135790]precursors. Proc. Nat. Acad. Sci. 82: 6109–6113, 1985.

[135791]24174.

[135792]Puchacz, E.; Lian, J. B.; Stein, G. S.; Wozney, J.; Huebner, K.;

[135793]Croce, C.: Chromosomal localization of the human osteocalcin gene. Endocrinology 124:

[135794]2648–2650, 1989.

[135795]24175.

[135796]Tabas, J. A.; Zasloff, M.; Wasmuth, J. J.; Emanuel, B. S.; Altherr,

[135797]M. R.; McPherson, J. D.; Wozney, J. M.; Kaplan, F. S.: Bone morphogenetic

[135798]protein: chromosomal localization of human genes for BMP1, BMP2A,

[135799]and BMP3. Genomics 9: 283–289, 1991.

[135800]24176.

[135801]Wozney, J. M.; Rosen, V.; Celeste, A. J.; Mitsock, L. M.; Whitters,

[135802]M. J.; Kriz, R. W.; Hewick, R. M.; Wang, E. A.: Novel regulators

[135803]of bone formation: molecular clones and activities. Science 242:

[135804]1528–1534, 1988.

[135805]24177.

[135806]Connor, J. M.: Fibrodysplasia ossificans progressiva: lessons

[135807]from rare maladies. (Editorial) New Eng. J. Med. 335: 591–593, 1996.

[135808]24178.

[135809]Dooley, C. A.; Attia, G. R.; Rainey, W. E.; Moore, D. R.; Carr,

[135810]B. R.: Bone morphogenetic protein inhibits ovarian androgen production. J.

[135811]Clin. Endocr. Metab. 85: 3331–3337, 2000.

[135812]24179.

[135813]McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba,

[135814]Canada 7/15/1992.

[135815]24180.

[135816]Monsoro–Burq, A.–H.; le Douarin, N. M.: BMP4 plays a key role

[135817]in left-right patterning in chick embryos by maintaining
Sonic Hedgehog

[135818]asymmetry. *Molec. Cell* 7: 789-799, 2001.

[135819]24181.

[135820]Shafritz, A. B.; Shore, E. M.; Gannon, F. H.; Zasloff, M. A.;
Taub,

[135821]R.; Muenke, M.; Kaplan, F. S.: Overexpression of an os-
teogenic morphogen

[135822]in fibrodysplasia ossificans progressiva. *New Eng. J. Med.*
335:

[135823]555-561, 1996.

[135824]24182.

[135825]Tabas, J. A.; Hahn, G. V.; Cohen, R. B.; Seaunez, H. N.;
Modi,

[135826]W. S.; Wozney, J. M.; Zasloff, M.; Kaplan, F. S.: Chromoso-
mal assignment

[135827]of the human gene for bone morphogenetic protein 4.
Clin. Orthop.

[135828]*Rel. Res.* 293: 310-316, 1993.

[135829]24183.

[135830]Tucker, A. S.; Matthews, K. L.; Sharpe, P. T.: Transforma-

tion

[135831] of tooth type induced by inhibition of BMP signaling. Science 282:

[135832] 1136–1138, 1998.

[135833] 24184.

[135834] van den Wijngaard, A.; Olde Weghuis, D.; Boersma, C. J. C.; van

[135835] Zoelen, E. J. J.; Geurts van Kessel, A.; Olijve, W.: Fine mapping

[135836] of the human bone morphogenetic protein–4 gene (BMP4) to chromosome

[135837] 14q22–q23 by in situ hybridization. Genomics 27: 559–560, 1995.

[135838] 24185.

[135839] van den Wijngaard, A.; van Kraay, M.; van Zoelen, E. J. J.; Olijve,

[135840] W.; Boersma, C. J. C.: Genomic organization of the human bone morphogenetic

[135841] protein–4 gene: molecular basis for multiple transcripts. Biochem.

[135842] Biophys. Res. Commun. 219: 789–794, 1996.

[135843] 24186.

- [135844]Amano, S.; Scott, I. C.; Takahara, K.; Koch, M.; Champli-
aud, M.-F.;
- [135845]Gerecke, D. R.; Keene, D. R.; Hudson, D. L.; Nishiyama, T.;
Lee, S.;
- [135846]Greenspan, D. S.; Burgeson, R. E.: Bone morphogenetic
protein 1 is
- [135847]an extracellular processing enzyme of the laminin 5
gamma-2 chain. J.
- [135848]Biol. Chem. 275: 22728-22735, 2000.
- [135849]24187.
- [135850]Bond, J. S.; Beynon, R. J.: The astacin family of metalloen-
dopeptidases. Protein
- [135851]Sci. 4: 1247-1261, 1995.
- [135852]24188.
- [135853]Ceci, J. D.; Kingsley, D. M.; Silan, C. M.; Copeland, N. G.;
- Jenkins,
- [135854]N. A.: An interspecific backcross linkage map of the prox-
imal half
- [135855]of mouse chromosome 14. Genomics 6: 673-678, 1990.
- [135856]24189.
- [135857]Kessler, E.; Takahara, K.; Biniaminov, L.; Brusel, M.;

Greenspan,

[135858]D.: Bone morphogenic protein-1: the type I procollagen
C-proteinase. Science 271:

[135859]360-362, 1996.

[135860]24190.

[135861]Kremmidiotis, G.; Baker, E.; Crawford, J.; Eyre, H. J.; Nah-
mias,

[135862]J.; Callen, D. F.: Localization of human cadherin genes to
chromosome

[135863]regions exhibiting cancer-related loss of heterozygosity.
Genomics 49:

[135864]467-471, 1998.

[135865]24191.

[135866]Herrenknecht, K.; Ozawa, M.; Eckerskorn, C.; Lottspeich,
F.; Lenter,

[135867]M.; Kemler, R.: The uvomorulin-anchorage protein alpha-
catenin is

[135868]a vinculin homologue. Proc. Nat. Acad. Sci. 88:
9156-9160, 1991.

[135869]24192.

[135870]Blomback, M.; Blomback, B.; Mammen, E. F.; Prasad, A. S.:

Fibrinogen

[135871]Detroit--a molecular defect in the N-terminal disulphide knot of human

[135872]fibrinogen? Nature 218: 134-137, 1968.

[135873]24193.

[135874]Magnuson, V. L.; McCombs, J. L.; Lee, C.-C.; Yang, F.; Bowman,

[135875]B. H.; McGill, J. R.: Human alpha-2-HS-glycoprotein localized to

[135876]3q27-q29 by in situ hybridization. Cytogenet. Cell Genet. 47: 72-74,

[135877]1988.

[135878]24194.

[135879]Hromas, R.; Collins, S.; Raskind, W.; Deaven, L.; Kaushansky, K.

[135880]: Hem-1, a potential membrane protein, with expression restricted

[135881]to blood cells. Biochim. Biophys. Acta 1090: 241-244, 1991.

[135882]24195.

[135883]Chen, Y.-H.; Lin, S.-J.; Lin, M.-W.; Tsai, H.-L.; Kuo, S.-S.;

Chen,

[135884]J.-W.; Charng, M.-J.; Wu, T.-C.; Chen, L.-C.; Ding, P. Y.-A.;

Pan,

[135885]W.-H.; Jou, Y.-S.; Chau, L.-Y.: Microsatellite polymorphism

in promoter

[135886]of heme oxygenase-1 gene is associated with susceptibil-

ity to coronary

[135887]artery disease in type 2 diabetic patients. Hum. Genet.

111: 1-8,

[135888]2002.

[135889]24196.

[135890]Fujita, T.; Toda, K.; Karimova, A.; Yan, S.-F.; Naka, Y.; Yet,

[135891]S.-F.; Pinsky, D. J.: Paradoxical rescue from ischemic lung

injury

[135892]by inhaled carbon monoxide driven by derepression of

fibrinolysis. Nature

[135893]Med. 7: 598-604, 2001.

[135894]24197.

[135895]Kutty, R. K.; Kutty, G.; Rodriguez, I. R.; Chader, G. J.; Wig-

gart,

[135896]B.: Chromosomal localization of the human oxygenase

genes: heme oxygenase-1

[135897](HMOX1) maps to chromosome 22q12 and heme oxygenase-2 (HMOX2) maps

[135898]to chromosome 16p13.3. Genomics 20: 513-516, 1994.

[135899]24198.

[135900]Poss, K. D.; Tonegawa, S.: Reduced stress defense in heme oxygenase

[135901]1-deficient cells. Proc. Nat. Acad. Sci. 94: 10925-10930, 1997.

[135902]24199.

[135903]Saito-Ohara, F.; Ikeuchi, T.; Matsumoto, M.; Kurata, S.: Assignment

[135904]of the mouse heme oxygenase genes: heme oxygenase-1 (Hmox1) to chromosome

[135905]10 band C1 and heme oxygenase-2 (Hmox2) to chromosome 16 band B1. Cytogenet.

[135906]Cell Genet. 77: 180-181, 1997.

[135907]24200.

[135908]Seroussi, E.; Kedra, D.; Kost-Alimova, M.; Sandberg-Nordqvist,

[135909]A.-C.; Fransson, I.; Jacobs, J. F. M.; Fu, Y.; Pan, H.-Q.; Roe,

B.

- [135910]A.; Imreh, S.; Dumanski, J. P.: TOM1 genes map to human chromosome 22q13.1 and mouse chromosome 8C1 and encode proteins similar to the endosomal proteins HGS and STAM. Genomics 57: 380–388, 1999.
- [135913]24201.
- [135914]Yachie, A.; Niida, Y.; Wada, T.; Igarashi, N.; Kaneda, H.; Toma, T.; Ohta, K.; Kasahara, Y.; Koizumi, S.: Oxidative stress causes enhanced endothelial cell injury in human heme oxygenase–1 deficiency. J. Clin. Invest. 103: 129–135, 1999.
- [135918]24202.
- [135919]Yamada, N.; Yamaya, M.; Okinaga, S.; Nakayama, K.; Sekizawa, K.; Shibahara, S.; Sasaki, H.: Microsatellite polymorphism in the heme oxygenase–1 gene promoter is associated with susceptibility to emphysema. Am.

[135922]J. Hum. Genet. 66: 187–195, 2000. Note: Erratum: Am. J.

Hum. Genet.

[135923]68: 1542 only, 2001.

[135924]24203.

[135925]Yoshida, T.; Biro, P.; Cohen, T.; Muller, R. M.; Shibahara, S.

[135926]; Human heme oxygenase cDNA and induction of its
mRNA by hemin. Europ.

[135927]J. Biochem. 171: 457–461, 1988.

[135928]24204.

[135929]Stevenson, R. E.; Schroer, R. J.; Taylor, H. A.; Compton, J.
D.;

[135930]Livingston, R. E., III: Dislocated lens, dolichostenomelia,
and joint

[135931]stiffness. Proc. Greenwood Genet. Center 1: 16–22, 1982.

[135932]24205.

[135933]Celli, J.; Duijf, P.; Hamel, B. C. J.; Bamshad, M.; Kramer, B.;

[135934]Smits, A. P. T.; Newbury-Ecob, R.; Hennekam, R. C. M.;

Van Buggenhout,

[135935]G.; van Haeringen, A.; Woods, C. G.; van Essen, A. J.; de
Waal, R.;

- [135936]Vriend, G.; Haber, D. A.; Yang, A.; McKeon, F.; Brunner, H. G.; van Bokhoven, H.: Heterozygous germline mutations in the p53 homolog p63 are the cause of EEC syndrome. Cell 99: 143–153, 1999.
- [135939]24206.
- [135940]O'Farrell, P. H.: High resolution two-dimensional electrophoresis of proteins. J. Biol. Chem. 250: 4007–4021, 1975.
- [135942]24207.
- [135943]Osawa, M.; Umetsu, K.; Ohki, T.; Nagasawa, T.; Suzuki, T.; Takeichi, S.: Molecular evidence for human alpha2-HS glycoprotein (AHSG) polymorphism. Hum. Genet. 99: 18–21, 1997.
- [135946]24208.
- [135947]Pedersen, K. O.: Fetuin, a new globulin isolated from serum.
- [135948](Letter) Nature 154: 575, 1944.
- [135949]24209.

- [135950]Rizzu, P.; Baldini, A.: Three members of the human cystatin gene
- [135951]superfamily, AHSG, HRG, and KNG, map within one megabase of genomic
- [135952]DNA at 3q27. Cytogenet. Cell Genet. 70: 26–28, 1995.
- [135953]24210.
- [135954]Terkelsen, O. B. F.; Jahnen–Dechent, W.; Nielsen, H.; Moos, T.;
- [135955]Fink, E.; Nawratil, P.; Muller–Esterl, W.; Mollgard, K.: Rat fetuin:
- [135956]distribution of protein and mRNA in embryonic and neonatal rat tissues. Anat.
- [135957]Embryol. 197: 125–133, 1998.
- [135958]24211.
- [135959]Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha-2-HS-glycoprotein
- [135960]typing by isoelectric focusing. Hum. Genet. 67: 70–71, 1984.
- [135961]24212.
- [135962]Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha-2-HS-glycoprotein

[135963]allele (AHS*5*) in two Japanese families. Hum. Genet. 68:
264–265,

[135964]1984.

[135965]24213.

[135966]Umetsu, K.; Yuasa, I.; Ikuta, M.; Suzuki, T.: Alpha-
2-HS-glycoprotein

[135967](A2HS) polymorphism in a Japanese population: existence
of two new

[135968]variants. Jpn. J. Hum. Genet. 32: 263–266, 1987.

[135969]24214.

[135970]Umetsu, K.; Yuasa, I.; Suzuki, T.: The polymorphism of
desialyzed

[135971]alpha-2-HS-glycoprotein (AHS): isoelectric focusing in
2.5M urea as

[135972]a method for identification of genetic variants. Hum.
Genet. 73:

[135973]372–373, 1986.

[135974]24215.

[135975]Yoshioka, Y.; Gejyo, F.; Marti, T.; Rickli, E. E.; Burgi, W.;

[135976]Offner, G. D.; Troxler, R. F.; Schmid, K.: The complete
amino acid

[135977]sequence of the A-chain of human plasma alpha-
2HS-glycoprotein. J.

[135978]Biol. Chem. 261: 1665-1676, 1986.

[135979]24216.

[135980]Zelinski, T.; Kaita, H.; Lewis, M.; Coghlan, G.; Craig, D.:
The

[135981]sequence of chromosome 3 loci AHS:TF:CHE1. Hum.
Hered. 37: 1-6,

[135982]1987.

[135983]24217.

[135984]Cleve, H.: Genetic studies on the deficiency of beta-
2-glycoprotein

[135985]I of human serum. Humangenetik 5: 294-304, 1968.

[135986]24218.

[135987]Cleve, H.; Rittner, C.: Further family studies on the genetic

[135988]control of beta-2-glycoprotein I concentration in human
serum. Humangenetik 7:

[135989]93-97, 1969.

[135990]24219.

[135991]Eiberg, H.; Mohr, J.; Nielsen, L. S.: The beta-

2-glycoprotein

[135992]I (BG): allele frequencies and linkage relationships.

(Abstract) Cytogenet.

[135993]Cell Genet. 37: 462 only, 1984.

[135994]24220.

[135995]Eiberg, H.; Nielsen, L. S.; Mohr, J.: Exclusion mapping of apolipoprotein

[135996]H (APOH) and relationship between electrophoretic and quantitative

[135997]polymorphism. (Abstract) Cytogenet. Cell Genet. 51: 994 only, 1989.

[135998]24221.

[135999]Haagerup, A.; Kristensen, T.; Kruse, T. A.: Polymorphism and genetic

[136000]mapping of the gene encoding human beta-2-glycoprotein I to chromosome

[136001]17. (Abstract) Cytogenet. Cell Genet. 58: 2005 only, 1991.

[136002]24222.

[136003]Haupt, H.; Schwick, H. G.; Storiko, K.: Ueber einen erblichen

[136004]beta-2-Glykoprotein I-Mangel. Humangenetik 5:

291-293, 1968.

[136005]24223.

[136006]Hoeg, J. M.; Segal, P.; Gregg, R. E.; Chang, Y. S.; Lindgren, F.

[136007]T.; Adamson, G. L.; Frank, M.; Brickman, C.; Brewer, H. B., Jr.:

[136008]Characterization of plasma lipids and lipoproteins in patients with

[136009]beta-2-glycoprotein I (apolipoprotein H) deficiency. Atherosclerosis 55:

[136010]25-34, 1985.

[136011]24224.

[136012]Kamboh, M. I.; Ferrell, R. E.; Sepehrnia, B.: Genetic studies of human apolipoproteins. IV. Structural heterogeneity of apolipoprotein

[136014]H (beta-2-glycoprotein I). Am. J. Hum. Genet. 42: 452-457, 1988.

[136015]24225.

[136016]Koppe, A. L.; Walter, H.; Chopra, V. P.; Bajatzadeh, M.: Investigations

[136017]on the genetics and population genetics of the beta-

2-glycoprotein

[136018] polymorphism. Humangenetik 9: 164–171, 1970.

[136019] 24226.

[136020] Lozier, J.; Takahashi, N.; Putnam, F. W.: Complete amino acid

[136021] sequence of human plasma beta-2-glycoprotein I. Proc. Nat. Acad.

[136022] Sci. 81: 3640–3644, 1984.

[136023] 24227.

[136024] Mehdi, H.; Nunn, M.; Steel, D. M.; Whitehead, A. S.; Perez, M.;

[136025] Walker, L.; Peeples, M. E.: Nucleotide sequence and expression of

[136026] the human gene encoding apolipoprotein H (beta-2-glycoprotein I). Gene 108:

[136027] 293–298, 1991.

[136028] 24228.

[136029] Nakaya, Y.; Schaefer, E. J.; Brewer, H. B.: Activation of human

[136030] post-heparin lipase by apolipoprotein H (beta-2-glycoprotein I). Biochem.

[136031]Biophys. Res. Commun. 95: 1168–1172, 1980.

[136032]24229.

[136033]Nonaka, M.; Matsuda, Y.; Shiroishi, T.; Moriwaki, K.; Nonaka,

[136034]M.; Natsuume–Sakai, S.: Molecular cloning of mouse beta-2-glycoprotein

[136035]and mapping of the gene to chromosome 11. Genomics 13: 1082–1087,

[136036]1992.

[136037]24230.

[136038]Rahimi, A. G.; Goedde, H. W.; Flatz, G.; Kaifie, S.; Benkmann,

[136039]H.–G.; Delbruck, H.: Serum protein polymorphisms in four populations

[136040]of Afghanistan. Am. J. Hum. Genet. 29: 356–360, 1977.

[136041]24231.

[136042]Richter, A.; Cleve, H.: Genetic variations of human serum beta-2-glycoprotein

[136043]demonstrated by isoelectric focusing. Electrophoresis 9: 317–322,

[136044]1988.

[136045]24232.

[136046]Sanghera, D. K.; Nestlerode, C. S.; Ferrell, R. E.; Kamboh, M.

[136047]I.: Chimpanzee apolipoprotein H (beta-2-glycoprotein I): report on

[136048]the gene structure, a common polymorphism, and a high prevalence of

[136049]antiphospholipid antibodies. Hum. Genet. 109: 63-72, 2001.

[136050]24233.

[136051]Sanghera, D. K.; Wagenknecht, D. R.; McIntyre, J. A.; Kamboh,

[136052]M. I.: Identification of structural mutations in the fifth domain

[136053]of apolipoprotein H (beta-2-glycoprotein I) which affect phospholipid

[136054]binding. Hum. Molec. Genet. 6: 311-316, 1997.

[136055]24234.

[136056]Sepehrnia, B.; Kamboh, M. I.; Adams-Campbell, L. L.; Bunker, C.

[136057]H.; Nwankwo, M.; Majumder, P. P.; Ferrell, R. E.: Genetic studies

[136058] of human apolipoproteins. VIII. Role of the apolipoprotein
H polymorphism

[136059] in relation to serum lipoprotein concentrations. Hum.
Genet. 82:

[136060] 118–122, 1989.

[136061] 24235.

[136062] Kost, G. J.; Verity, A.: A new variant of late-onset
myophosphorylase

[136063] deficiency. Muscle Nerve 3: 195–201, 1980.

[136064] 24236.

[136065] Rosenmann, A.; Scriver, C. R.; Clow, C. L.; Levy, H. L.: His-
tidinaemia.

[136066] Part II: Impact; a retrospective study. J. Inherit. Metab. Dis.
6:

[136067] 54–57, 1983.

[136068] 24237.

[136069] Johnson, J. L.; Coyne, K. E.; Rajagopalan, K. V.; Van Hove,
J.

[136070] L. K.; Mackay, M.; Pitt, J.; Boneh, A.: Molybdopter in syn-
thase mutations

[136071] in a mild case of molybdenum cofactor deficiency. Am. J.

Med. Genet. 104:

[136072]169–173, 2001.

[136073]24238.

[136074]Johnson, J. L.; Wuebbens, M. M.; Mandell, R.; Shih, V. E.:

Molybdenum

[136075]cofactor biosynthesis in humans: identification of two
complementation

[136076]groups of cofactor-deficient patients and preliminary
characterization

[136077]of a diffusible molybdopterin precursor. J. Clin. Invest. 83:
897–903,

[136078]1989.

[136079]24239.

[136080]Reiss, J.; Christensen, E.; Dorche, C.: Molybdenum cofac-
tor deficiency:

[136081]first prenatal genetic analysis. Prenatal Diag. 19:
386–388, 1999.

[136082]24240.

[136083]Reiss, J.; Christensen, E.; Kurlemann, G.; Zabet, M.-T.;
Dorche,

[136084]C.: Genomic structure and mutational spectrum of the bi-

cistronic

[136085]MOCS1 gene defective in molybdenum cofactor deficiency
type A. Hum.

[136086]Genet. 103: 639–644, 1998.

[136087]24241.

[136088]Reiss, J.; Cohen, N.; Dorche, C.; Mandel, H.; Mendel, R. R.;
Stallmeyer,

[136089]B.; Zobot, M.–T.; Dierks, T.: Mutations in a polycistronic
nuclear

[136090]gene associated with molybdenum cofactor deficiency.
Nature Genet. 20:

[136091]51–53, 1998.

[136092]24242.

[136093]Reiss, J.; Dorche, B.; Stallmeyer, B.; Mendel, R. R.; Cohen,
N.;

[136094]Zobot, M. T.: Human molybdopterin synthase gene: ge-
nomic structure

[136095]and mutations in molybdenum cofactor deficiency type B.
Am. J. Hum.

[136096]Genet. 64: 706–711, 1999.

[136097]24243.

- [136098]Inui, K.; Emmett, M.; Wenger, D. A.: Immunological evidence for
deficiency in an activator protein for sulfatide sulfatase in
a variant
form of metachromatic leukodystrophy. Proc. Nat. Acad.
Sci. 80:
3074–3077, 1983.
- [136102]24244.
- [136103]Austin, J.; Armstrong, D.; Fouch, S.; Mitchell, C.; Stumpf,
D.
A.; Shearer, L.; Briner, O.: Metachromatic leukodystrophy
(MLD).
VIII. MLD in adults: diagnosis and pathogenesis. Arch.
Neurol. 18:
225–240, 1968.
- [136107]24245.
- [136108]Austin, J.; McAfee, D.; Armstrong, D.; O'Rourke, M.;
Shearer, L.;
Bachhawat, B. K.: Abnormal sulphatase activities in two
human diseases
(metachromatic leukodystrophy and gargoylism).
Biochem. J. 93: 15C–17C,

[136111]1964.

[136112]24246.

[136113]Austin, J.; McAfee, D.; Shearer, L.: Metachromatic form of diffuse

[136114]cerebral sclerosis. IV. Low sulfatase activity in the urine of nine

[136115]living patients with metachromatic leukodystrophy (MLD). Arch. Neurol. 12:

[136116]447-455, 1965.

[136117]24247.

[136118]Austin, J. H.: Some recent findings in leukodystrophies and in

[136119]gargoylism.In: Aronson, S. M.; Volk, B. W.: Inborn Disorders of Sphingolipid

[136120]Metabolism. Oxford: Pergamon Press (pub.) 1967. Pp. 359-387.

[136121]24248.

[136122]Austin, J. H.: Metachromatic form of diffuse cerebral sclerosis.

[136123]III. Significance of sulfatide and other lipid abnormalities in white

[136124]matter and kidney. Neurology 10: 470–483, 1960.

[136125]24249.

[136126]Baldinger, S.; Pierpont, M. E.; Wenger, D. A.: Pseudodeficiency

[136127]of arylsulfatase A: a counseling dilemma. Clin. Genet. 31: 70–76,

[136128]1987.

[136129]24250.

[136130]Barth, M. L.; Fensom, A.; Harris, A.: Missense mutations in the

[136131]aryl sulphatase A genes of metachromatic leukodystrophy patients. Hum.

[136132]Molec. Gen. 2: 2117–2121, 1993.

[136133]24251.

[136134]Barth, M. L.; Fensom, A.; Harris, A.: Identification of seven

[136135]novel mutations associated with metachromatic leukodystrophy. Hum.

[136136]Mutat. 6: 170–176, 1995.

[136137]24252.

[136138]Lebo, R. V.; Anderson, L. A.; DiMauro, S.; Lynch, E.;

Hwang, P.;

[136139]Fletcher, R.: Rare McArdle disease locus polymorphic site on 11q13

[136140]contains CpG sequence. Hum. Genet. 86: 17–24, 1990.

[136141]24253.

[136142]Lebo, R. V.; Gorin, F.; Fletcher, R. J.; Kao, F.-T.; Cheung,

[136143]M.-C.; Bruce, B. D.; Kan, Y. W.: High-resolution chromosome sorting

[136144]and DNA spot-blot analysis assign McArdle's syndrome to chromosome

[136145]11. Science 225: 57–59, 1984.

[136146]24254.

[136147]Lehoczky, T.; Halasy, M.; Simon, G.; Harnos, G.: Glycogenic myopathy:

[136148]a case of skeletal muscle-glycogenosis in twins. J. Neurol. Sci. 2:

[136149]366–384, 1965.

[136150]24255.

[136151]Mantle, D.; Lauffart, B.; Atack, J.; Lane, R. J. M.: Absence

[136152]of biochemical heterogeneity in McArdle's disease: a high

resolution

[136153]SDS–polyacrylamide gel electrophoresis study. J. Neurol. Sci. 78:

[136154]63–70, 1987.

[136155]24256.

[136156]Martin, M. A.; Rubio, J. C.; Garcia, A.; Fernandez, M. A.; Campos,

[136157]Y.; Krawczak, M.; Cooper, D. N.; Arenas, J.: Resolution of a mispaired

[136158]secondary structure intermediate could account for a novel micro–insertion/deletion

[136159](387 insA/del 8 bp) in the PYGM gene causing McArdle's disease. Clin.

[136160]Genet. 59: 48–51, 2001.

[136161]24257.

[136162]Martinuzzi, A.; Vergani, L.; Carrozzo, R.; Fanin, M.; Bartoloni,

[136163]L.; Angelini, C.; Askanas, V.; Engel, W. K.: Expression of muscle–type

[136164]phosphorylase in innervated and aneural cultured muscle of patients

[136165]with myophosphorylase deficiency. J. Clin. Invest. 92:

1774–1780,

[136166]1993.

[136167]24258.

[136168]McArdle, B.: Myopathy due to a defect in muscle glycogen breakdown. Clin.

[136169]Sci. 10: 13–33, 1951.

[136170]24259.

[136171]Mellick, R. S.; Mahler, R. F.; Hughes, B. P.: McArdle's syndrome.

[136172]Phosphorylase-deficient myopathy. Lancet I: 1045–1048, 1962.

[136173]24260.

[136174]Miranda, A. F.; Nette, E. G.; Hartlage, P. L.; DiMauro, S.: Phosphorylase

[136175]isoenzymes in normal and myophosphorylase-deficient human heart. Neurology 29:

[136176]1538–1541, 1979.

[136177]24261.

[136178]Papadimitriou, A.; Manta, P.; Divari, R.; Karabetsos, A.; Papadimitriou,

[136179]E.; Bresolin, N.: McArdle's disease: two clinical expressions

in

[136180]the same pedigree. J. Neurol. 237: 267–270, 1990.

[136181]24262.

[136182]Pearson, C. M.; Rimer, D. G.; Mommaerts, W. F. H. M.: A
metabolic

[136183]myopathy due to absence of muscle phosphorylase. Am. J.
Med. 30:

[136184]502–517, 1961.

[136185]24263.

[136186]Ross, B. D.; Radda, G. K.; Gadian, D. G.; Rocker, G.; Esiri,
M.;

[136187]Falconer–Smith, J.: Examination of a case of suspected
McArdle's

[136188]syndrome by (31)P nuclear magnetic resonance. New Eng.
J. Med. 304:

[136189]1338–1342, 1981.

[136190]24264.

[136191]Rowland, L. P.; Lovelace, R. E.; Schotland, D. L.; Araki, S.;

[136192]Carmel, P.: The clinical diagnosis of McArdle's disease:
identification

[136193]of another family with deficiency of muscle phosphory–

lase. Neurology 16:

[136194]93–100, 1966.

[136195]24265.

[136196]Sarova–Pinhas, I.; Sadeh, M.: McArdle disease in a Druze family. Israel

[136197]J. Med. Sci. 25: 64–68, 1989.

[136198]24266.

[136199]Schmid, R.; Hammaker, L.: Hereditary absence of muscle phosphorylase

[136200](McArdle's syndrome). New Eng. J. Med. 264: 223–225, 1961.

[136201]24267.

[136202]Schmid, R.; Mahler, R.: Chronic progressive myopathy with myoglobinuria:

[136203]demonstration of a glycogenolytic defect in the muscle. J. Clin.

[136204]Invest. 38: 2044–2058, 1959.

[136205]24268.

[136206]Servidei, S.; Shanske, S.; Zeviani, M.; Lebo, R.; Fletterick,

[136207]R.; DiMauro, S.: McArdle's disease: biochemical and

molecular genetic

[136208] studies. Ann. Neurol. 24: 774-781, 1988.

[136209] 24269.

[136210] Slonim, A. E.; Goans, P. J.: Myopathy in McArdle's syndrome:

[136211] improvement with a high-protein diet. New Eng. J. Med. 312: 355-359,

[136212] 1985.

[136213] 24270.

[136214] Tabata, T.; Kikunami, K.; Matsushita, Y.; Inoue, T.; Inoue, T.;

[136215] Okamoto, T.; Kono, N.; Takahashi, M.; Tarui, S.; Morii, H.: Acute

[136216] renal failure in McArdle's disease. Nephron 44: 371-374, 1986.

[136217] 24271.

[136218] Tsujino, S.; Rubin, L. A.; Shanske, S.; DiMauro, S.: An A-to-C

[136219] substitution involving the translation initiation codon in a patient

[136220] with myophosphorylase deficiency (McArdle's disease).

Hum. Mutat. 4:

[136221]73–75, 1994.

[136222]24272.

[136223]Tsuchino, S.; Shanske, S.; DiMauro, S.: Molecular genetic heterogeneity

[136224]of myophosphorylase deficiency (McArdle's disease). New Eng. J. Med. 329:

[136225]241–245, 1993.

[136226]24273.

[136227]Tsuchino, S.; Shanske, S.; Martinuzzi, A.; Heiman–Patterson, T.;

[136228]DiMauro, S.: Two novel missense mutations (E654K, L396P) in Caucasian

[136229]patients with myophosphorylase deficiency (McArdle's disease). Hum.

[136230]Mutat. 6: 276–277, 1995.

[136231]24274.

[136232]Tsuchino, S.; Shanske, S.; Nonaka, I.; Eto, Y.; Mendell, J. R.;

[136233]Fenichel, G. M.; DiMauro, S.: Three new mutations in patients with

[136234]myophosphorylase deficiency (McArdle disease). Am. J.

Hum. Genet. 54:

[136235]44–52, 1994.

[136236]24275.

[136237]Vorgerd, M.; Kubisch, C.; Burwinkel, B.; Reichmann, H.;
Mortier,

[136238]W.; Tettenborn, B.; Pongratz, D.; Lindemuth, R.; Tegen-
thoff, M.; Malin,

[136239]J.-P.; Kilimann, M. W.: Mutation analysis in myophospho-
rylase deficiency

[136240](McArdle's disease). Ann. Neurol. 43: 326–331, 1998.

[136241]24276.

[136242]Hers, H. G.: Etudes enzymatiques sur fragments hepa-
tiques: application

[136243]a la classification des glycogenoses. Rev. Int. Hepat. 9:
35–55,

[136244]1959.

[136245]24277.

[136246]Scriver, C. R.; Levy, H. L.: Histidinaemia. Part I: Reconciling

[136247]retrospective and prospective findings. J. Inherit. Metab.
Dis. 6:

[136248]51–53, 1983.

[136249]24278.

[136250]Suchi, M.; Harada, N.; Wada, Y.; Takagi, Y.: Molecular cloning

[136251]of a cDNA encoding human histidase. Biochim. Biophys. Acta 1216:

[136252]293–295, 1993.

[136253]24279.

[136254]Suchi, M.; Sano, H.; Mizuno, H.; Wada, Y.: Molecular cloning

[136255]and structural characterization of the human histidase gene (HAL). Genomics 29:

[136256]98–104, 1995.

[136257]24280.

[136258]Taylor, R. G.; Garcia-Heras, J.; Sadler, S. J.; Lafreniere, R.

[136259]G.; Willard, H. F.; Ledbetter, D. H.; McInnes, R. R.: Localization

[136260]of histidase to human chromosome region 12q22–q24.1 and mouse chromosome

[136261]region 10C2–D1. Cytogenet. Cell Genet. 56: 178–181, 1991.

[136262]24281.

[136263]Taylor, R. G.; Garcia-Heras, J.; Sexsmith, E.; Lambert, M.;
Ledbetter,
[136264]D.; McInnes, R. R.: Molecular characterization of histidase
cDNAs,
[136265]and sublocalization of the human histidase gene to
12q22-23, and the
[136266]mouse gene to 10C2-D1.(Abstract) Am. J. Hum. Genet. 45
(suppl.):
[136267]A164, 1989.
[136268]24282.

[136269]Taylor, R. G.; Grieco, D.; Clarke, G. A.; McInnes, R. R.; Tay-
lor,
[136270]B. A.: Identification of the mutation in murine histidinemia
(his)
[136271]and genetic mapping of the murine histidase locus (Hal)
on chromosome
[136272]10. Genomics 16: 231-240, 1993.
[136273]24283.

[136274]Wadman, S. K.; Van Sprang, F. J.; Van Stekelenburg, G. J.;
de
[136275]Bree, P. K.: Three new cases of histidinemia: clinical and
biochemical

[136276]data. Acta Paediat. Scand. 56: 485–492, 1967.

[136277]24284.

[136278]Woody, N. C.; Snyder, C. H.; Harris, J. A.: Histidinemia.
Am.

[136279]J. Dis. Child. 110: 606–613, 1965.

[136280]24285.

[136281]Martin–Subero, J. I.; Gesk, S.; Harder, L.; Sonoki, T.;
Tucker,

[136282]P. W.; Schlegelberger, B.; Grote, W.; Novo, F. J.; Calasanz,
M. J.;

[136283]Hansmann, M. L.; Dyer, M. J. S.; Siebert, R.: Recurrent in-
volvement

[136284]of the REL and BCL11A loci in classical Hodgkin lym-
phoma. Blood 99:

[136285]1474–1477, 2002.

[136286]24286.

[136287]Hankinson, S. E.; Willett, W. C.; Colditz, G. A.; Hunter, D.
J.;

[136288]Michaud, D. S.; Deroo, B.; Rosner, B.; Speizer, F. E.; Pollak,
M.

[136289]: Circulating concentrations of insulin–like growth factor–I

and risk

[136290]of breast cancer. Lancet 351: 1393–1396, 1998.

[136291]24287.

[136292]Esumi, N.; Budarf, M.; Ciccarelli, L.; Sellinger, B.; Kozak, C.

[136293]A.; Wistow, G.: Conserved gene structure and genomic
linkage for

[136294]D-dopachrome tautomerase (DDT) and MIF. Mammalian
Genome 9: 753–757,

[136295]1998.

[136296]24288.

[136297]Roger, T.; David, J.; Glauser, M. P.; Calandra, T.: MIF regu-
lates

[136298]innate immune response through modulation of Toll-like
receptor 4. Nature 414:

[136299]920–924, 2001.

[136300]24289.

[136301]Emi, M.; Asaoka, H.; Matsumoto, A.; Itakura, H.; Kurihara,
Y.;

[136302]Wada, Y.; Kanamori, H.; Yazaki, Y.; Takahashi, E.; Lepert,
M.; Lalouel,

[136303]J.–M.; Kodama, T.; Mukai, T.: Structure, organization, and

chromosomal

[136304]mapping of the human macrophage scavenger receptor
gene. J. Biol.

[136305]Chem. 268: 2120–2125, 1993.

[136306]24290.

[136307]Latil, A.; Lidereau, R.: Genetic aspects of prostate cancer.
Virchows

[136308]Arch. 432: 389–406, 1998.

[136309]24291.

[136310]Matsumoto, A.; Naito, M.; Itakura, H.; Ikemoto, S.; Asaoka,
H.;

[136311]Hayakawa, I.; Kanamori, H.; Aburatani, H.; Takaku, F.;

Suzuki, H.;

[136312]Kobari, Y.; Miyai, T.; Takahashi, K.; Cohen, E. H.; Wydro,
R.; Housman,

[136313]D. E.; Kodama, T.: Human macrophage scavenger recep-
tors: primary

[136314]structure, expression, and localization in atherosclerotic
lesions. Proc.

[136315]Nat. Acad. Sci. 87: 9133–9137, 1990.

[136316]24292.

[136317] Xu, J.; Zheng, S. L.; Hawkins, G. A.; Faith, D. A.; Kelly, B.;

[136318] Isaacs, S. D.; Wiley, K. E.; Chang, B.; Ewing, C. M.; Bu-
jnovszky,

[136319] P.; Carpten, J. D.; Bleecker, E. R.; Walsh, P. C.; Trent, J. M.;
Meyers,

[136320] D. A.; Isaacs, W. B.: Linkage and association studies of
prostate

[136321] cancer susceptibility: evidence for linkage at 8p22-23.
Am. J. Hum.

[136322] Genet. 69: 341-350, 2001.

[136323] 24293.

[136324] Xu, J.; Zheng, S. L.; Komiya, A.; Mychaleckyj, J. C.; Isaacs,
S.

[136325] D.; Hu, J. J.; Sterling, D.; Lange, E. M.; Hawkins, G. A.;
Turner,

[136326] A.; Ewing, C. M.; Faith, D. A.; and 19 others: Germline mu-
tations

[136327] and sequence variants of the macrophage scavenger re-
ceptor 1 gene

[136328] are associated with prostate cancer risk. Nature Genet. 32:
321-325,

[136329] 2002.

[136330]24294.

[136331]Holness, C. L.; Simmons, D. L.: Molecular cloning of CD68,
a human

[136332]macrophage marker related to lysosomal glycoproteins.
Blood 81:

[136333]1607-1613, 1993.

[136334]24295.

[136335]Jones, E.; Quinn, C. M.; See, C. G.; Montgomery, D. S.;
Ford, M.

[136336]J., Kolble, K.; Gordon, S.; Greaves, D. R.: The linked human
elongation

[136337]initiation factor 4A1 (EIF4A1) and CD68 genes map to
chromosome 17p13. Genomics 53:

[136338]248-250, 1998.

[136339]24296.

[136340]Chavrier, P.; Janssen-Timmen, U.; Mattei, M.-G.; Zerial,
M.; Bravo,

[136341]R.; Charnay, P.: Structure, chromosome location, and ex-
pression of

[136342]the mouse zinc finger gene Krox-20: multiple gene prod-
ucts and coregulation

[136343]with the proto-oncogene c-fos. Molec. Cell. Biol. 9:
787-797, 1989.

[136344]24297.

[136345]Le Beau, M. M.; Espinosa, R., III; Neuman, W. L.; Stock, W.;
Roulston,

[136346]D.; Larson, R. A.; Keinanen, M.; Westbrook, C. A.: Cytoge-
netic and

[136347]molecular delineation of the smallest commonly deleted
region of chromosome

[136348]5 in malignant myeloid diseases. Proc. Nat. Acad. Sci. 90:
5484-5488,

[136349]1993.

[136350]24298.

[136351]Barrow, L. L.; Simin, K.; Jones, J. M.; Lee, D. C.; Meisler, M.

[136352]H.: Conserved linkage of early growth response 4, annexin
4, and

[136353]transforming growth factor alpha on mouse chromosome
6. Genomics 19:

[136354]388-390, 1994.

[136355]24299.

[136356]Crosby, S. D.; Veile, R. A.; Donis-Keller, H.; Baraban, J. M.;

[136357]Bhat, R. V.; Simburger, K. S.; Milbrandt, J.: Neural-specific expression,
[136358]genomic structure, and chromosomal localization of the gene encoding
[136359]the zinc-finger transcription factor NGFI-C. Proc. Nat. Acad. Sci. 89:
[136360]4739-4743, 1992.
[136361]24300.

[136362]Boerkoel, C. F.; Takashima, H.; Bacino, C. A.; Daentl, D.; Lupski,
[136363]J. R.: EGR2 mutation R359W causes a spectrum of Dejerine-Sottas neuropathy. Neurogenetics 3:
[136364]153-157, 2001.
[136365]24301.

[136366]Willcutt, E. G.; Pennington, B. F.; Smith, S. D.; Cardon, L. R.;

[136367]Gayan, J.; Knopik, V. S.; Olson, R. K.; DeFries, J. C.: Quantitative
[136368]trait locus for reading disability on chromosome 6p is pleiotropic
[136369]for attention-deficit/hyperactivity disorder. Am. J. Med. Genet. 114:

[136370]260–268, 2002.

[136371]24302.

[136372]Koiwai, O.; Nishizawa, M.; Hasada, K.; Aono, S.; Adachi, Y.;
Mamiya,

[136373]N.; Sato, H.: Gilbert's syndrome is caused by a heterozy-
gous missense

[136374]mutation in the gene for bilirubin UDP-
glucuronosyltransferase. Hum.

[136375]Molec. Genet. 4: 1183–1186, 1995.

[136376]24303.

[136377]Maruo, Y.; Wada, S.; Yamamoto, K.; Sato, H.; Yamano, T.;
Shimada,

[136378]M.: A case of anorexia nervosa with hyperbilirubinaemia in
a patient

[136379]homozygous for a mutation in the bilirubin UDP-
glucuronosyltransferase

[136380]gene. Europ. J. Pediat. 158: 547–549, 1999.

[136381]24304.

[136382]Andres, D. A.; Milatovich, A.; Ozcelik, T.; Wenzlau, J. M.;
Brown,

[136383]M. S.; Goldstein, J. L.; Francke, U.: cDNA cloning of the two

subunits

[136384]of human CAAX farnesyltransferase and chromosomal
mapping of FNTA

[136385]and FNTB loci and related sequences. Genomics 18:
105–112, 1993.

[136386]24305.

[136387]Long, S. B.; Casey, P. J.; Beese, L. S.: Reaction path of pro-
tein

[136388]farnesyltransferase at atomic resolution. Nature 419:
645–650, 2002.

[136389]24306.

[136390]Roifman, C. M.; Zhang, J.; Chitayat, D.; Sharfe, N.: A partial

[136391]deficiency of interleukin-7R-alpha is sufficient to abro-
gate T-cell

[136392]development and cause severe combined immunodeficiency. Blood 96:

[136393]2803–2807, 2000.

[136394]24307.

[136395]Bowne, S. J.; Sullivan, L. S.; Blanton, S. H.; Cepko, C. L.;
Blackshaw,

[136396]S.; Birch, D. G.; Houghbanks-Wheaton, D.; Heckenlively, J.

R.; Daiger,

[136397]S. P.: Mutations in the inosine monophosphate dehydrogenase 1 gene

[136398](IMPDH1) cause the RP10 form of autosomal dominant retinitis pigmentosa. Hum.

[136399]Molec. Genet. 11: 559-568, 2002.

[136400]24308.

[136401]Collart, F. R.; Huberman, E.: Cloning and sequence analysis of

[136402]the human and Chinese hamster inosine-5-prime-monophosphate dehydrogenase

[136403]cDNAs. J. Biol. Chem. 263: 15769-15772, 1988.

[136404]24309.

[136405]Doggett, N. A.; Callen, D. F.; Chen, Z. L.; Moore, S.; Tesmer,

[136406]J. G.; Duesing, L. A.; Stallings, R. L.: Identification and regional

[136407]localization of a human IMP dehydrogenase-like locus (IMPDHL1) at

[136408]16p13.13. Genomics 18: 687-689, 1993.

[136409]24310.

[136410]Gu, J. J.; Kaiser-Rogers, K.; Rao, K.; Mitchell, B. S.: Assignment
[136411]of the human type I IMP dehydrogenase gene (IMPDH1) to
[136412]7q31.3-q32. Genomics 24: 179-181, 1994.
[136413]24311.
[136414]Natsumeda, Y.; Ohno, S.; Kawasaki, H.; Konno, Y.; Weber,
G.; Suzuki,
[136415]K.: Two distinct cDNAs for human IMP dehydrogenase. J.
Biol. Chem. 265:
[136416]5292-5295, 1990.
[136417]24312.
[136418]Kennan, A.; Aherne, A.; Palfi, A.; Humphries, M.; McKee,
A.; Stitt,
[136419]A.; Simpson, D. A. C.; Demtroder, K.; Orntoft, T.; Ayuso,
C.; Kenna,
[136420]P. F.; Farrar, G. J.; Humphries, P.: Identification of an IM-
PDH1
[136421]mutation in autosomal dominant retinitis pigmentosa
(RP10) revealed
[136422]following comparative microarray analysis of transcripts
derived from

- [136423]retinas of wild-type and Rho-/- mice. Hum. Molec. Genet.
11: 547-558,
[136424]2002.
[136425]24313.
[136426]Crompton, M. R.; Moss, S. E.; Crompton, M. J.: Diversity in
the
[136427]lipocortin/calpactin family. Cell 55: 1-3, 1988.
[136428]24314.
[136429]Horlick, K. R.; Cheng, I. C.; Wong, W. T.; Wakeland, E. K.;
Nick,
[136430]H. S.: Mouse lipocortin I gene structure and chromosomal
assignment:
[136431]gene duplication and the origins of a gene family. Ge-
nomics 10:
[136432]365-374, 1991.
[136433]24315.
[136434]Huebner, K.; Cannizzaro, L. A.; Croce, C. M.; Frey, A. Z.;
Wallner,
[136435]B. P.; Hecht, B. K.; Hecht, F.: Chromosome localization of
the human
[136436]genes for lipocortin I and the lipocortin II family.

(Abstract) Cytogenet.

[136437]Cell Genet. 46: 631 only, 1987.

[136438]24316.

[136439]Huebner, K.; Cannizzaro, L. A.; Frey, A. Z.; Hecht, B. K.;
Hecht,

[136440]F.; Croce, C. M.; Wallner, B. P.: Chromosomal localization
of the

[136441]human genes for lipocortin I and lipocortin II. Oncogene
Res. 2:

[136442]299–310, 1988.

[136443]24317.

[136444]Pepinsky, R. B.; Tizard, R.; Mattaliano, R. J.; Sinclair, L. K.;

[136445]Miller, G. T.; Browning, J. L.; Chow, E. P.; Burne, C.; Huang,
K.-S.;

[136446]Pratt, D.; Wachter, L.; Hession, C.; Frey, A. Z.; Wallner, B.
P.:

[136447]Five distinct calcium and phospholipid binding proteins
share homology

[136448]with lipocortin I. J. Biol. Chem. 263: 10799–10811, 1988.

[136449]24318.

[136450]Shohat, M.; Korenberg, J. R.; Schwabe, A. D.; Rotter, J. I.:

Hypothesis:

[136451]familial Mediterranean fever—a genetic disorder of the lipocortin

[136452]family? Am. J. Med. Genet. 34: 163–167, 1989.

[136453]24319.

[136454]Wallner, B. P.; Mattaliano, R. J.; Hession, C.; Cate, R. L.; Tizard,

[136455]R.; Sinclair, L. K.; Foeller, C.; Chow, E. P.; Browning, J. L.; Ramachandran,

[136456]K. L.; Pepinsky, R. B.: Cloning and expression of human lipocortin,

[136457]a phospholipase A2 inhibitor with potential anti-inflammatory activity. Nature 320:

[136458]77–81, 1986.

[136459]24320.

[136460]Harman, S. M.; Metter, E. J.; Blackman, M. R.; Landis, P. K.;

[136461]Carter, H. B.: Serum levels of insulin-like growth factor I (IGF-I),

[136462]IGF-II, IGF-binding protein-3, and prostate-specific antigen as predictors

[136463]of clinical prostate cancer. J. Clin. Endocr. Metab. 85:

4258-4265,

[136464]2000.

[136465]24321.

[136466]Holly, J.: Insulin-like growth factor-I and new opportunities

[136467]for cancer prevention. Lancet 351: 1373-1375, 1998.

[136468]24322.

[136469]Hoppener, J. W. M.; de Pagter-Holthuisen, P.; Geurts van Kessel,

[136470]A. H. M.; Jansen, M.; Kittur, S. D.; Antonarakis, S. E.; Lips, C.

[136471]J. M.; Sussenbach, J. S.: The human gene encoding insulin-like growth

[136472]factor I is located on chromosome 12. Hum. Genet. 69: 157-160, 1985.

[136473]24323.

[136474]Le Bouc, Y.; Dreyer, D.; Jaeger, F.; Binoux, M.; Sondermeyer,

[136475]P.: Complete characterization of the human IGF-I nucleotide sequence

[136476]isolated from a newly constructed adult liver cDNA library.

FEBS

[136477]Lett. 196: 108–112, 1986.

[136478]24324.

[136479]Hellstrom, A.; Perruzzi, C.; Ju, M.; Engstrom, E.; Hard,
A.-L.;

[136480]Liu, J.-L.; Albertsson-Wikland, K.; Carlsson, B.; Niklasson,
A.; Sjodell,

[136481]L.; LeRoith, D.; Senger, D. R.; Smith, L. E. H.: Low IGF-I
suppresses

[136482]VEGF-survival signaling in retinal endothelial cells: direct
correlation

[136483]with clinical retinopathy of prematurity. Proc. Nat. Acad.
Sci. 98:

[136484]5804–5808, 2001.

[136485]24325.

[136486]Lembo, G.; Rockman, H. A.; Hunter, J. J.; Steinmetz, H.;
Koch,

[136487]W. J.; Ma, L.; Printz, M. P.; Ross, J., Jr.; Chien, K. R.; Powell–
Braxton,

[136488]L.: Elevated blood pressure and enhanced myocardial con-
tractility

[136489]in mice with severe IGF–1 deficiency. J. Clin. Invest. 98:

2648–2655,

[136490]1996.

[136491]24326.

[136492]Li, C. H.; Yamashiro, D.; Gospodarowicz, D.; Kaplan, S. L.;

Van

[136493]Vliet, G.: Total synthesis of insulin-like growth factor I

(somatomedin

[136494]C). Proc. Nat. Acad. Sci. 80: 2216–2220, 1983.

[136495]24327.

[136496]Liu, J.-P.; Baker, J.; Perkins, A. S.; Robertson, E. J.; Efstra-

tiadis,

[136497]A.: Mice carrying null mutations of the genes encoding in-

sulin-like

[136498]growth factor I (Igf-1) and type 1 IGF receptor (Igf1r). Cell

75:

[136499]59–72, 1993.

[136500]24328.

[136501]Mathews, L. S.; Norstedt, G.; Palmiter, R. D.: Regulation of

[136502]insulin-like growth factor I gene expression by growth

hormone. Proc.

[136503]Nat. Acad. Sci. 83: 9343–9347, 1986.

[136504]24329.

[136505]Morton, C.; Rall, L.; Bell, G.; Shows, T.: Human insulin-like

[136506]growth factor-1 (IGF1) is encoded at 12q22-q24.1, and
insulin-like

[136507]growth factor-2 (IGF2) is at 11p15. (Abstract) Cytogenet.
Cell Genet. 40:

[136508]703 only, 1985.

[136509]24330.

[136510]Musaro, A.; McCullagh, K.; Paul, A.; Houghton, L.; Do-
browolny,

[136511]G.; Molinaro, M.; Barton, E. R.; Sweeney, H. L.; Rosenthal,
N.: Localized

[136512]Igf-1 transgene expression sustains hypertrophy and re-
generation in

[136513]senescent skeletal muscle. Nature Genet. 27: 195-200,
2001.

[136514]24331.

[136515]Musaro, A.; McCullagh, K. J. A.; Naya, F. J.; Olson, E. N.;
Rosenthal,

[136516]N.: IGF-1 induces skeletal myocyte hypertrophy through
calcineurin

[136517]in association with GATA-2 and NF-ATc1. Nature 400:
581-585, 1999.

[136518]24332.

[136519]Playford, M. P.; Bicknell, D.; Bodmer, W. F.; Macaulay, V. M.

[136520]; Insulin-like growth factor 1 regulates the location, sta-
bility,

[136521]and transcriptional activity of beta-catenin. Proc. Nat.
Acad. Sci. 97:

[136522]12103-12108, 2000.

[136523]24333.

[136524]Powell-Braxton, L.; Hollingshead, P.; Warburton, C.; Dowd,
M.;

[136525]Pitts-Meek, S.; Dalton, D.; Gillett, N.; Stewart, T. A.: IGF-I
is

[136526]required for normal embryonic growth in mice. Genes
Dev. 7: 2609-2617,

[136527]1993.

[136528]24334.

[136529]Rapp, R.; Deger, A.; Blum, W.; Koch, R.; Weber, U.: Charac-
terization

[136530]of the protein which binds insulin-like growth factor in

human serum. Europ.

[136531]J. Biochem. 172: 421–425, 1988.

[136532]24335.

[136533]Rotwein, P.: Two insulin-like growth factor I messenger
RNAs

[136534]are expressed in human liver. Proc. Nat. Acad. Sci. 83:
77–81, 1986.

[136535]24336.

[136536]Schoenle, E. J.; Zenobi, P. D.; Torresani, T.; Werder, E. A.;

[136537]Zachmann, M.; Froesch, E. R.: Recombinant human in-
sulin-like growth

[136538]factor I (rhIGF I) reduces hyperglycaemia in patients with
extreme

[136539]insulin resistance. Diabetologia 34: 675–679, 1991.

[136540]24337.

[136541]Semsarian, C.; Wu, M.-J.; Ju, Y.-K.; Marciniak, T.; Yeoh, T.;

[136542]Allen, D. G.; Harvey, R. P.; Graham, R. M.: Skeletal muscle
hypertrophy

[136543]is mediated by a Ca^{2+} -dependent calcineurin signalling
pathway. Nature 400:

[136544]576–581, 1999.

[136545]24338.

[136546]Svoboda, M. E.; Van Wyk, J. J.; Klapper, D. G.; Fellows, R. E.;

[136547]Grissom, F. E.; Schleuter, R. J.: Purification of somatomedin-C from

[136548]human plasma: chemical and biological properties, partial sequence

[136549]analysis, and relationship to other somatomedins. Biochemistry 19:

[136550]790-797, 1980.

[136551]24339.

[136552]Taylor, B. A.; Grieco, D.: Localization of the gene encoding

[136553]insulin-like growth factor I on mouse chromosome 10. Cytogenet. Cell

[136554]Genet. 56: 57-58, 1991.

[136555]24340.

[136556]Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T.

[136557]B.: Insulin-like growth factor genes: chromosome organization and

[136558]association with disease. (Abstract) Am. J. Hum. Genet. 36: 121S

[136559]only, 1984.

[136560]24341.

[136561]Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T.

[136562]B.: Localization of insulin-like growth factor genes to human chromosomes

[136563]11 and 12. Nature 310: 784–786, 1984.

[136564]24342.

[136565]Ullrich, A.; Berman, C. H.; Dull, T. J.; Gray, A.; Lee, J. M.

[136566]: Isolation of the human insulin-like growth factor I gene using a

[136567]single synthetic DNA probe. EMBO J. 3: 361–364, 1984.

[136568]24343.

[136569]Gough, N. M.; Rakar, S.; Harpur, A.; Wilks, A. F.: Localization

[136570]of genes for two members of the JAK family of protein tyrosine kinases

[136571]to murine chromosomes 4 and 19. Mammalian Genome 6: 247–248, 1995.

[136572]24344.

[136573]Ihle, J. N.: Cytokine receptor signalling. Nature 377: 591–594,

[136574]1995.

[136575]24345.

- [136576]Pritchard, M. A.; Baker, E.; Callen, D. F.; Sutherland, G. R.;
[136577]Wilks, A. F.: Two members of the JAK family of protein ty-
rosine kinases
[136578]map to chromosomes 1p31.3 and 9p24. Mammalian
Genome 3: 36-38, 1992.
[136579]24346.
- [136580]Campbell, G. S.; Argetsinger, L. S.; Ihle, J. N.; Kelly, P. A.;
[136581]Rillema, J. A.; Carter-Su, C.: Activation of JAK2 tyrosine ki-
nase
[136582]by prolactin receptors in Nb2 cells and mouse mammary
gland explants. Proc.
[136583]Nat. Acad. Sci. 91: 5232-5236, 1994.
[136584]24347.
- [136585]Huang, L. J.; Constantinescu, S. N.; Lodish, H. F.: The N-
terminal
[136586]domain of Janus kinase 2 is required for Golgi processing
and cell
[136587]surface expression of erythropoietin receptor. Molec. Cell
8: 1327-1338,
[136588]2001.
[136589]24348.

[136590]Neubauer, H.; Cumano, A.; Muller, M.; Wu, H.; Huffstadt, U.; Pfeffer,

[136591]K.: Jak2 deficiency defines an essential developmental checkpoint

[136592]in definitive hematopoiesis. Cell 93: 397–409, 1998.

[136593]24349.

[136594]Parganas, E.; Wang, D.; Stravopodis, D.; Topham, D. J.; Marine,

[136595]J.–C.; Teglund, S.; Vanin, E. F.; Bodner, S.; Colamonici, O. R.; van

[136596]Deursen, J. M.; Grosveld, G.; Ihle, J. N.: Jak2 is essential for

[136597]signaling through a variety of cytokine receptors. Cell 93: 385–395,

[136598]1998.

[136599]24350.

[136600]Saltzman, A.; Stone, M.; Franks, C.; Searfoss, G.; Munro, R.; Jaye,

[136601]M.; Ivashchenko, Y.: Cloning and characterization of human Jak–2

[136602]kinase: high mRNA expression in immune cells and muscle tissue. Biochem.

[136603]Biophys. Res. Commun. 246: 627–633, 1998.

[136604]24351.

[136605]Schwaller, J.; Parganas, E.; Wang, D.; Cain, D.; Aster, J. C.;

[136606]Williams, I. R.; Lee, C.-K.; Gerthner, R.; Kitamura, T.;

Frantsve,

[136607]J.; Anastasiadou, E.; Loh, M. L.; Levy, D. E.; Ihle, J. N.;

Gilliland,

[136608]D. G.: Stat5 is essential for the myelo- and lymphoprolif-
erative

[136609]disease induced by TEL/JAK2. Molec. Cell 6: 693–704,
2000.

[136610]24352.

[136611]Watling, D.; Guschin, D.; Muller, M.; Silvennoinen, O.; Wit-
thuhn,

[136612]B. A.; Quelle, F. W.; Rogers, N. C.; Schindler, C.; Stark, G.
R.;

[136613]Ihle, J. N.; Kerr, I. M.: Complementation by the protein ty-
rosine

[136614]kinase JAK2 of a mutant cell line defective in the inter-
feron-gamma

[136615]signal transduction pathway. Nature 366: 166–170, 1993.

[136616]24353.

[136617]Cihakova, D.; Trebusak, K.; Heino, M.; Fadeyev, V.; Tiul-
pakov,

[136618]A.; Battelino, T.; Tar, A.; Halasz, Z.; Blumel, P.; Tawfik, S.;
Krohn,

[136619]K.; Lebl, J.; Peterson, P.; MEWPE-APECED Study Group:
Novel AIRE

[136620]mutations and P450 cytochrome autoantibodies in Central
and Eastern

[136621]European patients with APECED. Hum. Mutat. 18:
225–232, 2001.

[136622]24354.

[136623]Clemente, M. G.; Obermayer–Straub, P.; Meloni, A.; Strass-
burg,

[136624]C. P.; Arangino, V.; Tukey, R. H.; de Virgiliis, S.; Manns, M.
P.

[136625]: Cytochrome P450 1A2 is a hepatic autoantigen in au-
toimmune polyglandular

[136626]syndrome type 1. J. Clin. Endocr. Metab. 82: 1353–1361,
1997.

[136627]24355.

[136628]Craig, J. M.; Schiff, L. H.; Boone, J. E.: Chronic moniliasis

[136629] associated with Addison's disease. *Am. J. Dis. Child.* 89: 669–684, 1955.

[136630] 24356.

[136631] Ekwall, O.; Hedstrand, H.; Haavik, J.; Perheentupa, J.; Betterle, C.; Gustafsson, J.; Husebye, E.; Rorsman, F.; Kampe, O.: Pteridin-dependent hydroxylases as autoantigens in autoimmune polyendocrine syndrome type I. *J. Clin. Endocr. Metab.* 85: 2944–2950, 2000.

[136632] 24357.

[136633] Cetani, F.; Barbesino, G.; Borsari, S.; Pardi, E.; Cianferotti, L.; Pinchera, A.; Marcocci, C.: A novel mutation of the autoimmune regulator gene in an Italian kindred with autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy, acting in a dominant fashion and strongly cosegregating with hypothyroid autoimmune thyroiditis. *J. Clin. Endocr. Metab.* 86:

[136642]4747–4752, 2001.

[136643]24358.

[136644]Finnish–German APECED Consortium: An autoimmune disease, APECED,

[136645]caused by mutations in a novel gene featuring two PHD-type zinc-finger

[136646]domains. *Nature Genet.* 17: 399–403, 1997.

[136647]24359.

[136648]Foz, M.; Mirada, A.; Guardia, J.: Endocrine disorders in a family.

[136649](Letter) *Lancet* II: 269 only, 1970.

[136650]24360.

[136651]Gass, J. D. M.: The syndrome of keratoconjunctivitis, superficial

[136652]moniliiasis, idiopathic hypoparathyroidism and Addison's disease. *Am.*

[136653]*J. Ophthalm.* 54: 660–674, 1962.

[136654]24361.

[136655]Gylling, M.; Tuomi, T.; Bjorses, P.; Kontiainen, S.; Partanen,

[136656]J.; Christie, M. R.; Knip, M.; Perheentupa, J.; Miettinen, A.:

Beta-cell

[136657]autoantibodies, human leukocyte antigen II alleles, and
type 1 diabetes

[136658]in autoimmune polyendocrinopathy-candidia-
sis-ectodermal dystrophy. J.

[136659]Clin. Endocr. Metab. 85: 4434-4440, 2000.

[136660]24362.

[136661]Hedstrand, H.; Ekwall, O.; Haavik, J.; Landgren, E.; Betterle,

[136662]C.; Perheentupa, J.; Gustafsson, J.; Husebye, E.; Rorsman,
F.; Kampe,

[136663]O.: Identification of tyrosine hydroxylase as an autoanti-
gen in autoimmune

[136664]polyendocrine syndrome type I. Biochem. Biophys. Res.
Commun. 267:

[136665]456-461, 2000.

[136666]24363.

[136667]Heino, M.; Peterson, P.; Kudoh, J.; Shimizu, N.; An-
tonarakis,

[136668]S. E.; Scott, H. S.; Krohn, K.: APECED mutations in the au-
toimmune

[136669]regulator (AIRE) gene. Hum. Mutat. 18: 205–211, 2001.

[136670]24364.

[136671]Heino, M.; Scott, H. S.; Chen, Q.; Peterson, P.; Maenpaa, U.;

[136672]Papasavvas, M.–P.; Mittaz, L.; Barras, C.; Rossier, C.; Chrousos,

[136673]G. P.; Stratakis, C. A.; Nagamine, K.; Kudoh, J.; Shimizu, N.; Maclaren,

[136674]N.; Antonarakis, S. E.; Krohn, K.: Mutation analyses of North American

[136675]APS–1 patients. Hum. Mutat. 13: 69–74, 1999.

[136676]24365.

[136677]Hendrix, T. R.: Personal Communication. Baltimore, Md. 1/14/1985.

[136678]24366.

[136679]Hiekkala, H.: Idiopathic hypoparathyroidism, adrenal insufficiency

[136680]and moniliasis in children. Ann. Paediat. Fenn. 10: 213–222, 1964.

[136681]24367.

[136682]Hung, W.; Migeon, C. J.; Parrott, R. H.: A possible autoim–

mune

[136683]basis for Addison's disease in three siblings, one with idiopathic

[136684]hypoparathyroidism, pernicious anemia and superficial moniliasis. New

[136685]Eng. J. Med. 269: 658–663, 1963.

[136686]24368.

[136687]Husebye, E. S.; Gebre-Medhin, G.; Tuomi, T.; Perheentupa, J.;

[136688]Landin-Olsson, M.; Gustafsson, J.; Rorsman, F.; Kampe, O.: Autoantibodies

[136689]against aromatic L-amino acid decarboxylase in autoimmune polyendocrine

[136690]syndrome type I. J. Clin. Endocr. Metab. 82: 147–150, 1997.

[136691]24369.

[136692]Ishii, T.; Suzuki, Y.; Ando, N.; Matsuo, N.; Ogata, T.: Novel

[136693]mutations of the autoimmune regulator gene in two siblings with autoimmune

[136694]polyendocrinopathy–candidiasis–ectodermal dystrophy. J. Clin. Endocr.

[136695]Metab. 85: 2922–2926, 2000.

[136696]24370.

[136697]Kenny, F. M.; Holliday, M. D.: Hypoparathyroidism, moniliasis,

[136698]Addison's and Hashimoto's disease. Hypercalcemia treated with intravenously

[136699]administered sodium sulfate. New Eng. J. Med. 271: 708-713, 1964.

[136700]24371.

[136701]Krohn, K.; Uibo, R.; Aavik, E.; Peterson, P.; Savilahti, K.:

[136702]Identification by molecular cloning of an autoantigen associated with

[136703]Addison's disease as steroid 17-alpha-hydroxylase. Lancet 339: 770-773,

[136704]1992.

[136705]24372.

[136706]Kunin, A. S.; MacKay, B. R.; Burns, S. L.; Halberstam, M. J.:

[136707]The syndrome of hypoparathyroidism and adrenocortical insufficiency,

[136708]a possible sequel of hepatitis: case report and review of the literature. Am.

[136709]J. Med. 34: 856-866, 1963.

[136710]24373.

[136711]Louria, D. B.; Shannon, D. C.; Johnson, G.; Caroline, L.;
Okas,

[136712]A.; Taschdjian, C.: The susceptibility to moniliasis in chil-
dren

[136713]with endocrine hypofunction. Trans. Assoc. Am. Phys. 80:
236-249,

[136714]1967.

[136715]24374.

[136716]Maghnie, M.; Cosi, G.; Genovese, E.; Manca-Bitti, M. L.;
Cohen,

[136717]A.; Zecca, S.; Tinelli, C.; Gallucci, M.; Bernasconi, S.;
Boscherini,

[136718]B.; Severi, F.; Arico, M.: Central diabetes insipidus in chil-
dren

[136719]and young adults. New Eng. J. Med. 343: 998-1007, 2000.

[136720]24375.

[136721]Marieb, N. J.; Melby, J. C.; Lyall, S. S.: Isolated hypoaldos-
teronism

[136722]associated with idiopathic hypoparathyroidism. Arch. In-
tern. Med. 134:

[136723]424–429, 1974.

[136724]24376.

[136725]McKusick, V. A.: Personal Communication. Baltimore, Md.
1985.

[136726]24377.

[136727]Meloni, A.; Perniola, R.; Faa, V.; Corvaglia, E.; Cao, A.;
Rosatelli,

[136728]M. C.: Delineation of the molecular defects in the AIRE
gene in autoimmune

[136729]polyendocrinopathy–candidiasis–ectodermal dystrophy
patients from

[136730]Southern Italy. J. Clin. Endocr. Metab. 87: 841–846, 2002.

[136731]24378.

[136732]Nagamine, K.; Peterson, P.; Scott, H. S.; Kudoh, J.; Mi-
noshima,

[136733]S.; Heino, M.; Krohn, K. J. E.; Lalioti, M. D.; Mullis, P. E.;
Antonarakis,

[136734]S. E.; Kawasaki, K.; Asakawa, S.; Ito, F.; Shimizu, N.: Posi-
tional

[136735]cloning of the APECED gene. Nature Genet. 17: 393–398,
1997.

[136736]24379.

[136737]Neufeld, M.; Maclaren, N.; Blizzard, R.: Autoimmune
polyglandular

[136738]syndrome. *Pediat. Ann.* 9: 154–162, 1980.

[136739]24380.

[136740]Neufeld, M.; Maclaren, N. K.; Blizzard, R. M.: Two types of
autoimmune

[136741]Addison's disease associated with different polyglandular
autoimmune

[136742](PGA) syndromes. *Medicine* 60: 355–362, 1981.

[136743]24381.

[136744]Nithiyananthan, R.; Heward, J. M.; Allahabadia, A.; Barnett,
A.

[136745]H.; Franklyn, J. A.; Gough, S. C. L.: A heterozygous dele-
tion of

[136746]the autoimmune regulator (AIRE1) gene, autoimmune thy-
roid disease,

[136747]and type 1 diabetes: no evidence for association. *J. Clin.*
Endocr.

[136748]*Metab.* 85: 1320–1322, 2000.

[136749]24382.

- [136750]Okano, M.; Bell, D. W.; Haber, D. A.; Li, E.: DNA methyltransferases
- [136751]Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian
- [136752]development. Cell 99: 247–257, 1999.
- [136753]24383.
- [136754]Wijmenga, C.; van den Heuvel, L. P. W. J.; Strengman, E.; Luyten,
- [136755]J. A. F. M.; van der Burgt, I. J. A. M.; de Groot, R.; Smeets, D.
- [136756]F. C. M.; Draaisma, J. M. T.; van Dongen, J. J.; De Abreu, R. A.;
- [136757]Pearson, P. L.; Sandkuijl, L. A.; Weemaes, C. M. R.: Localization
- [136758]of the ICF syndrome to chromosome 20 by homozygosity mapping. Am.
- [136759]J. Hum. Genet. 63: 803–809, 1998.
- [136760]24384.
- [136761]Xu, G.–L.; Bestor, T. H.; Bourc'his, D.; Hsieh, C.–L.; Tommerup,
- [136762]N.; Bugge, M.; Hulten, M.; Qu, X.; Russo, J. J.; Viegas–Pequignot,

[136763]E.: Chromosome instability and immunodeficiency syndrome caused by

[136764]mutations in a DNA methyltransferase gene. Nature 402: 187-191,

[136765]1999.

[136766]24385.

[136767]Yamamoto, T.; Ninomiya, H.; Matsumoto, M.; Ohta, Y.; Nanba, E.;

[136768]Tsutsumi, Y.; Yamakawa, K.; Millat, G.; Vanier, M. T.; Pentchev, P.

[136769]G.; Ohno, K.: Genotype-phenotype relationship of Niemann-Pick disease

[136770]type C: a possible correlation between clinical onset and levels of

[136771]NPC1 protein in isolated skin fibroblasts. J. Med. Genet. 37: 707-711,

[136772]2000.

[136773]24386.

[136774]Alagille, D.; Josso, F.; Binet, J. L.; Blin, M. L.: La dystrophie

[136775]thrombocytaire hemorrhagique: discussion nosologique. Nouv. Rev.

[136776]Franc. Hemat. 4: 755–790, 1964.

[136777]24387.

[136778]Awidi, A. S.: Increased incidence of Glanzmann's throm-
basthenia

[136779]in Jordan as compared with Scandinavia. Scand. J. Haemat.
30: 218–222,

[136780]1983.

[136781]24388.

[136782]Basani, R .B.; Vilaire, G.; Shattil, S. J.; Kolodziej, M. A.;
Bennett,

[136783]J. S.; Poncz, M.: Glanzmann thrombasthenia due to a two
amino acid

[136784]deletion in the fourth calcium-binding domain of alphaIIb:
demonstration

[136785]of the importance of calcium-binding domains in the con-
formation of

[136786]alphaIIb/beta3. Blood 88: 167–173, 1996.

[136787]24389.

[136788]Bellucci, S.; Devergie, A.; Gluckman, E.; Tobelem, G.;
Lethielleux,

[136789]P.; Benbunan, M.; Schaison, G.; Boiron, M.: Complete cor-

rection of

[136790]Glanzmann's thrombasthenia by allogeneic bone-marrow transplantation. Brit.

[136791]J. Haemat. 59: 635-641, 1985.

[136792]24390.

[136793]Beutler, E.: Glanzmann's thrombasthenia and reduced glutathione. New

[136794]Eng. J. Med. 287: 1094-1095, 1972.

[136795]24391.

[136796]Bowie, E. J. W.; Owen, C. A., Jr.: Thrombopathy. Seminars Hemat. 5:

[136797]73-82, 1968.

[136798]24392.

[136799]Bowie, E. J. W.; Thompson, J. H., Jr.; Owen, C. A., Jr.: A new

[136800]abnormality of platelet function. Thromb. Diath. Haemorrh. 11: 195-203,

[136801]1964.

[136802]24393.

[136803]Bray, P. F.: Inherited diseases of platelet glycoproteins: considerations

[136804]for rapid molecular characterization. Thromb. Haemost.
72: 492–502,

[136805]1994.

[136806]24394.

[136807]Bray, P. F.; Barsh, G.; Rosa, J.-P.; Luo, X. Y.; Magenis, E.;

[136808]Shuman, M. A.: Physical linkage of the genes for platelet
membrane

[136809]glycoproteins IIb and IIIa. Proc. Nat. Acad. Sci. 85:
8683–8687,

[136810]1988.

[136811]24395.

[136812]Bray, P. F.; Rosa, J.-P.; Johnston, G. I.; Shiu, D. T.; Cook,

[136813]R. G.; Lau, C.; Kan, Y. W.; McEver, R. P.; Shuman, M. A.:
Platelet

[136814]glycoprotein IIb: chromosomal localization and tissue ex-
pression. J.

[136815]Clin. Invest. 80: 1812–1817, 1987.

[136816]24396.

[136817]Bray, P. F.; Rosa, J.-P.; Lingappa, V. R.; Kan, Y. W.; McEver,

[136818]R. P.; Shuman, M. A.: Biogenesis of the platelet receptor

for fibrinogen:

[136819]evidence for separate precursors for glycoproteins IIb and
IIla. Proc.

[136820]Nat. Acad. Sci. 83: 1480-1484, 1986.

[136821]24397.

[136822]Burk, C. D.; Newman, P. J.; Lyman, S.; Gill, J.; Collier, B. S.;

[136823]Poncz, M.: A deletion in the gene for glycoprotein IIb as-
sociated

[136824]with Glanzmann's thrombasthenia. J. Clin. Invest. 87:
270-276, 1991.

[136825]24398.

[136826]Collier, B. S.; Seligsohn, U.; Little, P. A.: Type I Glanzmann

[136827]thrombasthenia patients from the Iraqi-Jewish and Arab
populations

[136828]in Israel can be differentiated by platelet glycoprotein IIIa
immunoblot

[136829]analysis. Blood 69: 1696-1703, 1987.

[136830]24399.

[136831]Corby, D. G.; Zirbel, C. L.; Lindley, A.; Schulman, I.:
Thrombasthenia. Am.

[136832]J. Dis. Child. 121: 140-144, 1971.

[136833]24400.

[136834]Cronberg, S.; Nilsson, I. M.; Zetterqvist, E.: Investigation
[136835]of a family with members with both severe and mild de-
gree of thrombasthenia. Acta

[136836]Paediat. Scand. 56: 189–197, 1967.

[136837]24401.

[136838]Chuang, D. T.; Shih, V. E.: Disorders of branched chain
amino

[136839]acid and keto acid metabolism.In: Scriver, C. R.; Beaudet,
A. L.;

[136840]Sly, W. S.; Valle, D.: Metabolic and Molecular Bases of In-
herited

[136841]Disease. New York: McGraw–Hill (pub.) (7th ed.): 1995. Pp.
1239–1277.

[136842]24402.

[136843]Barth, M. L.; Fensom, A.; Harris, A.: Missense mutations in
the

[136844]arylsulphatase A genes of metachromatic leukodystrophy
patients. Hum.

[136845]Molec. Genet. 2: 2117–2121, 1993.

[136846]24403.

[136847]Barth, M. L.; Fensom, A.; Harris, A.: Prevalence of common mutations

[136848]in the arylsulphatase A gene in metachromatic leukodystrophy patients

[136849]diagnosed in Britain. Hum. Genet. 91: 73–77, 1993.

[136850]24404.

[136851]Lake, B. D.; Smith, V. V.; Judge, M. R.; Harper, J. I.; Besley,

[136852]G. T. N.: Hexanol dehydrogenase activity shown by enzyme histochemistry

[136853]on skin biopsies allows differentiation of Sjogren–Larsson syndrome

[136854]from other ichthyoses. J. Inherit. Metab. Dis. 14: 338–340, 1991.

[136855]24405.

[136856]Link, J. K.; Roldan, E. C.: Mental deficiency, spasticity, and

[136857]congenital ichthyosis: report of a case. J. Pediat. 52: 712–714,

[136858]1958.

[136859]24406.

[136860]Pigg, M.; Jagell, S.; Sillen, A.; Weissenbach, J.; Gustavson,

[136861]K.-H.; Wadelius, C.: The Sjogren-Larsson syndrome gene is close to

[136862]D17S805 as determined by linkage analysis and allelic association. Nature

[136863]Genet. 8: 361-364, 1994.

[136864]24407.

[136865]Rayner, A.; Lampert, R. P.; Rennert, O. M.: Familial ichthyosis,

[136866]dwarfism, mental retardation, and renal disease. J. Pediat. 92:

[136867]766-768, 1978.

[136868]24408.

[136869]Richards, B. W.: Congenital ichthyosis, spastic diplegia and

[136870]mental deficiency. (Letter) Brit. Med. J. 2: 714 only, 1960.

[136871]24409.

[136872]Rizzo, W. B.; Carney, G.; Lin, Z.: The molecular basis of Sjogren-Larsson

[136873]syndrome: mutation analysis of the fatty aldehyde dehydrogenase gene. Am.

[136874]J. Hum. Genet. 65: 1547-1560, 1999.

[136875]24410.

[136876]Rizzo, W. B.; Craft, D. A.: Sjogren-Larsson syndrome: deficient

[136877]activity of the fatty aldehyde dehydrogenase component of fatty alcohol:NAD+

[136878]oxidoreductase in cultured fibroblasts. J. Clin. Invest. 88: 1643-1648,

[136879]1991.

[136880]24411.

[136881]Rizzo, W. B.; Dammann, A. L.; Craft, D.; Black, S.; Henderson

[136882]Tilton, A.; Africk, D.; Chaves-Carballo, E.: Sjogren-Larsson syndrome:

[136883]deficient fatty alcohol:NAD+ oxidoreductase (FAO) activity in mixed

[136884]leukocytes. (Abstract) Am. J. Hum. Genet. 41: A16 only, 1987.

[136885]24412.

[136886]Rizzo, W. B.; Dammann, A. L.; Craft, D. A.: Sjogren-Larsson syndrome:

[136887]impaired fatty alcohol oxidation in cultured fibroblasts due to deficient

[136888]fatty alcohol:nicotinamide adenine dinucleotide oxidore-
ductase activity. J.

[136889]Clin. Invest. 81: 738-744, 1988.

[136890]24413.

[136891]Rizzo, W. B.; Dammann, A. L.; Craft, D. A.; Black, S. H.;
Henderson

[136892]Tilton, A.; Africk, D.; Chaves-Carballo, E.; Holmgren, G.;
Jagell,

[136893]S.: Sjogren-Larsson syndrome: inherited defect in the fatty
alcohol

[136894]cycle. J. Pediat. 115: 228-234, 1989.

[136895]24414.

[136896]Rogers, G. R.; Markova, N. G.; De Laurenzi, V.; Rizzo, W.
B.;

[136897]Compton, J. G.: Genomic organization and expression of
the human

[136898]fatty aldehyde dehydrogenase gene (FALDH). Genomics
39: 127-135,

[136899]1997.

[136900]24415.

[136901]Rogers, G. R.; Rizzo, W. B.; Zlotogorski, A.; Hashem, N.;

Lee,

[136902]M.; Bale, S. J.; Compton, J. G.: Sjogren-Larsson syndrome (SLS) and

[136903]fatty aldehyde dehydrogenase (FALDH) genes map to chromosome 17p.

[136904](Abstract) Am. J. Hum. Genet. 57: A202 only, 1995.

[136905]24416.

[136906]Rogers, G. R.; Rizzo, W. B.; Zlotogorski, A.; Hashem, N.; Lee,

[136907]M.; Compton, J. G.; Bale, S. J.: Genetic homogeneity in Sjogren-Larsson

[136908]syndrome: linkage to chromosome 17p in families of different non-Swedish

[136909]ethnic origins. Am. J. Hum. Genet. 57: 1123-1129, 1995.

[136910]24417.

[136911]Selmanowitz, V. J.; Porter, M. J.: The Sjogren-Larsson syndrome. Am.

[136912]J. Med. 42: 412-422, 1967.

[136913]24418.

[136914]Sillen, A.; Anton-Lamprecht, I.; Braun-Quentin, C.; Kraus, C.

[136915]S.; Sayli, B. S.; Ayuso, C.; Jagell, S.; Kuster, W.; Wadelius, C.

[136916]: Spectrum of mutations and sequence variants in the
FALDH gene in

[136917]patients with Sjogren–Larsson syndrome. Hum. Mutat. 12:
377–384,

[136918]1998.

[136919]24419.

[136920]Sillen, A.; Holmgren, G.; Wadelius, C.: First prenatal diag-
nosis

[136921]by mutation analysis in a family with Sjogren–Larsson
syndrome. Prenat.

[136922]Diag. 17: 1147–1149, 1997.

[136923]24420.

[136924]Sillen, A.; Jagell, S.; Wadelius, C.: A missense mutation in

[136925]the FALDH gene identified in Sjogren–Larsson syndrome
patients originating

[136926]from the northern part of Sweden. Hum. Genet. 100:
201–203, 1997.

[136927]24421.

[136928]Sjogren, T.: Oligophrenia combined with congenital
ichthyosiform

[136929]erythrodermia, spastic syndrome and macular retinal de-
generation:

[136930]a clinical and genetic study. Acta Genet. Statist. Med. 6:
80-91,

[136931]1956.

[136932]24422.

[136933]Sjogren, T.; Larsson, T.: Oligophrenia in combination with
congenital

[136934]ichthyosis and spastic disorders: a clinical and genetic
study. Acta

[136935]Psychiat. Neurol. Scand. 32 (suppl. 113): 1-112, 1957.

[136936]24423.

[136937]Tsukamoto, N.; Chang, C.; Yoshida, A.: Mutations associ-
ated with

[136938]Sjogren-Larsson syndrome. Ann. Hum. Genet. 61:
235-242, 1997.

[136939]24424.

[136940]Willemsen, M. A. A. P.; Cruysberg, J. R. M.; Rotteveel, J. J.;

[136941]Aandekerck, A. L.; Van Domburg, P. H. M. F.; Deutman, A.
F.: Juvenile

[136942]macular dystrophy associated with deficient activity of

fatty aldehyde

[136943]dehydrogenase in Sjogren–Larsson syndrome. Am. J. Ophthalmol. 130:

[136944]782–789, 2000.

[136945]24425.

[136946]Zaleski, W. A.: Congenital ichthyosis, mental retardation and

[136947]spasticity (Sjogren–Larsson syndrome). Canad. Med. Assoc. J. 86:

[136948]951–954, 1962.

[136949]24426.

[136950]Zhang, M.; Dwyer, N. K.; Love, D. C.; Cooney, A.; Comly, M.; Neufeld,

[136951]E.; Pentchev, P. G.; Blanchette–Mackie, E. J.; Hanover, J. A.: Cessation

[136952]of rapid late endosomal tubulovesicular trafficking in Niemann–Pick

[136953]type C1 disease. Proc. Nat. Acad. Sci. 98: 4466–4471, 2001.

[136954]24427.

[136955]Loeffen, J.; Elpeleg, O.; Smeitink, J.; Smeets, R.; Stockler–

Ipsiroglu,

[136956]S.; Mandel, H.; Sengers, R.; Trijbels, F.; van den Heuvel, L.:

Mutations

[136957]in the complex I NDUF52 gene of patients with cardiomyopathy and encephalomyopathy. Ann.

[136958]Neurol. 49: 195–201, 2001.

[136959]24428.

[136960]Smeitink, J.; van den Heuvel, L.: Human mitochondrial complex

[136961]I in health and disease. Am. J. Hum. Genet. 64: 1505–1510, 1999.

[136962]24429.

[136963]Baumgartner, E. R.; Suormala, T.; Wick, H.; Bausch, J.; Bonjour,

[136964]J.–P.: Biotinidase deficiency: factors responsible for the increased

[136965]biotin requirement. J. Inherit. Metab. Dis. 8 (suppl. 1): 59–64,

[136966]1985.

[136967]24430.

[136968]Burlina, A. B.; Sherwood, W. G.; Marchioro, M. V.; Dalla

Bernardina,

[136969]B.; Gaburro, D.: Neonatal screening for biotinidase deficiency in

[136970]north eastern Italy. *Europ. J. Pediat.* 147: 317–318, 1988.

[136971]24431.

[136972]Cole, H.; Reynolds, T. R.; Lockyer, J. M.; Buck, G. A.; Den-
son,

[136973]T.; Spence, J. E.; Hymes, J.; Wolf, B.: Human serum bio-
tinidase:

[136974]cDNA cloning, sequence, and characterization. *J. Biol.*
Chem. 269:

[136975]6566–6570, 1994.

[136976]24432.

[136977]Cole, H.; Weremowicz, S.; Morton, C. C.; Wolf, B.: Localiza-
tion

[136978]of serum biotinidase (BTD) to human chromosome 3 in
band p25. *Genomics* 22:

[136979]662–663, 1994.

[136980]24433.

[136981]Dionisi-Vici, C.; Bachmann, C.; Graziani, M. C.; Sabetta,
G.:

- [136982]Laryngeal stridor as a leading symptom in a biotinidase-deficient
patient. J. Inherit. Metab. Dis. 11: 312-313, 1988.
- [136983]24434.
- [136984]Dunkel, G.; Scriver, C. R.; Clow, C. L.; Melancon, S.;
Lemieux,
B.; Grenier, A.; Laberge, C.: Prospective ascertainment of
complete
and partial serum biotinidase deficiency in the newborn. J.
Inherit.
Metab. Dis. 12: 131-138, 1989.
- [136985]24435.
- [136986]Fischer, A.; Munnich, A.; Saudubray, J. M.; Mamas, S.;
Coude, F.
X.; Charpentier, C.; Dray, F.; Frezal, J.; Griscelli, C.: Biotin-
responsive
immunoregulatory dysfunction in multiple carboxylase
deficiency. J.
Clin. Immun. 2: 35-38, 1982.
- [136987]24436.
- [136988]Gaudry, M.; Munnich, A.; Saudubray, J. M.; Ogier, H.;

Mitchell,

[136996]G.; Marsac, C.; Causse, M.; Marquet, A.; Frezal, J.: Deficient liver

[136997]biotinidase activity in multiple carboxylase deficiency.

(Letter) Lancet II:

[136998]397, 1983.

[136999]24437.

[137000]Giardini, O.; D'Eufemia, P.; Cantani, A.; Ruberto, U.; Sabetta,

[137001]G.; Castro, M.; Cardi, E.: Congenital lactic acidosis associated

[137002]with laryngeal stridor: report of 2 cases. Riv. Ital. Pediat. 7:

[137003]139-142, 1981.

[137004]24438.

[137005]Gordon, A.: Biotinidase mutational 'hotspot'. (Letter) Nature

[137006]Genet. 13: 144-145, 1996.

[137007]24439.

[137008]Greter, J.; Holme, E.; Lindstedt, S.; Koivikko, M.: Biotin-responsive

[137009]3-methylcrotonylglycinuria with biotinidase deficiency. J.

Inherit.

[137010]Metab. Dis. 8: 103-104, 1985.

[137011]24440.

[137012]Hart, P. S.; Hymes, J.; Wolf, B.: Biochemical and immunological

[137013]characterization of serum biotinidase in profound biotinidase deficiency. Am.

[137014]J. Hum. Genet. 50: 126-136, 1992.

[137015]24441.

[137016]Heard, G. S.; Wolf, B.; Jefferson, L. G.; Weissbecker, K. A.;

[137017]Nance, W. E.; Secor McVoy, J. R.; Napolitano, A.; Mitchell, P. L.;

[137018]Lambert, F. W.; Linyear, A. S.: Neonatal screening for biotinidase

[137019]deficiency: results of a 1-year pilot study. J. Pediat. 108: 40-46,

[137020]1986.

[137021]24442.

[137022]Hymes, J.; Stanley, C. M.; Wolf, B.: Mutations in BTBD caus-

ing

[137023]biotinidase deficiency. Hum. Mutat. 18: 375–381, 2001.

[137024]24443.

[137025]Kalayci, O.; Coskun, T.; Tokatli, A.; Demir, E.; Erdem, G.;
Gungor,

[137026]C.; Yukselen, A.; Ozalp, I.: Infantile spasms as the initial
symptom

[137027]of biotinidase deficiency. J. Pediat. 124: 103–104, 1994.

[137028]24444.

[137029]Kennedy, R.; Girdwood, R. W. A.; King, M. D.: Neonatal
screening

[137030]for biotinidase deficiency: a pilot study in Scotland. J. In-
herit.

[137031]Metab. Dis. 12: 344–345, 1989.

[137032]24445.

[137033]Minns, R. A.; Kirk, J.: Biotinidase deficiency in Scotland.
(Letter) Dev.

[137034]Med. Child Neurol. 36: 748–749, 1994.

[137035]24446.

[137036]Muhl, A.; Moslinger, D.; Item, C. B.; Stockler-Ipsiroglu, S.:

[137037]Molecular characterisation of 34 patients with biotinidase

deficiency

[137038]ascertained by newborn screening and family investigation. Europ.

[137039]J. Hum. Genet. 9: 237-243, 2001.

[137040]24447.

[137041]Munnich, A.; Saudubray, J. M.; Carre, G.; Coude, F. X.; Ogier,

[137042]H.; Charpentier, C.; Frezal, J.: Defective biotin absorption in multiple

[137043]carboxylase deficiency. (Letter) Lancet II: 263 only, 1981.

[137044]24448.

[137045]Norrsgard, K. J.; Pomponio, R. J.; Hymes, J.; Wolf, B.: Mutations

[137046]causing profound biotinidase deficiency in children ascertained by

[137047]newborn screening in the United States occur at different frequencies

[137048]than in symptomatic children. Pediat. Res. 46: 20-27, 1999.

[137049]24449.

[137050]Norrsgard, K. J.; Swango, K. L.; Wolf, B.: Mutation (D444H)

in

[137051]the biotinidase gene causes approximately 50% loss of
enzyme activity

[137052]and is common in the general population. (Abstract) Am.
J. Hum. Genet.

[137053](Suppl.) 59: A275 only, 1996.

[137054]24450.

[137055]Pomponio, R. J.; Coskun, T.; Demirkol, M.; Tokatli, A.;
Ozalp,

[137056]I.; Huner, G.; Baykal, T.; Wolf, B.: Novel mutations cause
biotinidase

[137057]deficiency in Turkish children. J. Inherit. Metab. Dis. 23:
120-128,

[137058]2000.

[137059]24451.

[137060]Pomponio, R. J.; Hymes, J.; Reynolds, T. R.; Meyers, G. A.;
Fleischhauer,

[137061]K.; Buck, G. A.; Wolf, B.: Mutations in the human bio-
tinidase gene

[137062]that cause profound biotinidase deficiency in symptomatic
children:

[137063]molecular, biochemical, and clinical analysis. Pediat. Res.

42:

[137064]840–848, 1997.

[137065]24452.

[137066]Pomponio, R. J.; Narasimahan, V.; Reynolds, T. R.; Buck, G. A.;

[137067]Povirk, L. F.; Wolf, B.: Deletion/insertion mutation that causes

[137068]biotinidase deficiency may result from the formation of a quasipalindromic

[137069]structure. Hum. Molec. Genet. 5: 1657–1661, 1996.

[137070]24453.

[137071]Pomponio, R. J.; Norrgard, K. J.; Hymes, J.; Reynolds, T. R.;

[137072]Buck, G. A.; Baumgartner, R.; Suormala, T.; Wolf, B.: Arg538 to cys

[137073]mutation in a CpG dinucleotide of the human biotinidase gene is the

[137074]second most common cause of profound biotinidase deficiency in symptomatic

[137075]children. Hum. Genet. 99: 506–512, 1997.

[137076]24454.

[137077]Pomponio, R. J.; Reynolds, T. R.; Cole, H.; Buck, G. A.;

Wolf,

[137078]B.: Mutational hotspot in the human biotinidase gene
causes profound

[137079]biotinidase deficiency. Nature Genet. 11: 96–98, 1995.

[137080]24455.

[137081]Pomponio, R. J.; Reynolds, T. R.; Mandel, H.; Admoni, O.;
Melone,

[137082]P. D.; Buck, G. A.; Wolf, B.: Profound biotinidase deficiency
caused

[137083]by a point mutation that creates a downstream cryptic
3–prime splice

[137084]acceptor site within an exon of the human biotinidase
gene. Hum.

[137085]Molec. Genet. 6: 739–745, 1997.

[137086]24456.

[137087]Packman, S.; Sweetman, L.; Baker, H.; Wall, S.: The neona-
tal

[137088]form of biotin–responsive multiple carboxylase deficiency.
J. Pediat. 99:

[137089]418–420, 1981.

[137090]24457.

[137091]Packman, S.; Sweetman, L.; Yoshino, M.; Baker, H.; Cowan, M.:

[137092]Biotin-responsive multiple carboxylase deficiency of infantile onset. J.

[137093]Pediat. 99: 421-423, 1981.

[137094]24458.

[137095]Daiger, S. P.; Lidsky, A. S.; Chakraborty, R.; Koch, R.; Guttler,

[137096]F.; Woo, S. L. C.: Polymorphic DNA haplotypes at the phenylalanine

[137097]hydroxylase locus in prenatal diagnosis of phenylketonuria. Lancet I:

[137098]229-231, 1986.

[137099]24459.

[137100]Pizarro, T. T.; Michie, M. H.; Bentz, M.; Woraratanadham, J.;

[137101]Smith, M. F., Jr.; Foley, E.; Moskaluk, C. A.; Bickston, S. J.; Cominelli,

[137102]F.: IL-18, a novel immunoregulatory cytokine, is up-regulated in

[137103]Crohn's disease: expression and localization in intestinal mucosal

[137104]cells. J. Immun. 162: 6829–6835, 1999.

[137105]24460.

[137106]Oggiano, L.; Pirastu, M.; Moi, P.; Longinotti, M.; Perseu, L.;

[137107]Cao, A.: Molecular characterization of a normal HbA-2
beta-thalassemia

[137108]determinant in a Sardinian family. Brit. J. Haemat. 67:
225–229,

[137109]1987.

[137110]24461.

[137111]Ohba, Y.; Igarashi, M.; Tsukahara, M.; Nakashima, M.;
Sanada,

[137112]C.; Ami, M.; Arai, Y.; Miyaji, T.: Hb A(2) Yokoshima,
delta25(B7)

[137113]gly-to-asp, a new delta chain variant found in a Japanese
family. Hemoglobin 9:

[137114]613–615, 1985.

[137115]24462.

[137116]Ostertag, W.; Smith, E. W.: Hemoglobin–Lepore–Baltimore,
a third

[137117]type of a delta, beta crossover (delta 50, beta 86). Europ.
J. Biochem. 10:

[137118]371–376, 1969.

[137119]24463.

[137120]Papadakis, M.; Drakoulakou, O.; Papapanagiotou, E.;
Pessini, D.;

[137121]Loutradi–Anagnostou, A.: Hb A(2)–Agrinio
(delta-43(CD2)glu-to-gly

[137122](GAG-to-GGG)): a new delta-chain variant detected in a
Greek family. Hemoglobin 19:

[137123]295–299, 1995.

[137124]24464.

[137125]Petes, T. D.: Evidence that structural variants within the
human

[137126]delta-globin protein may reflect genetic interactions be-
tween the

[137127]delta- and beta-globin genes. (Letter) Am. J. Hum. Genet.
34: 820–823,

[137128]1982.

[137129]24465.

[137130]Ramirez, F.; Mears, J. G.; Nudel, U.; Bank, A.; Luzzatto, L.;

[137131]Di Prisco, G.; D'Avino, R.; Pepe, G.; Gambino, R.; Cimino,
R.; Quattrin,

[137132]N.: Defects in DNA and globin messenger RNA in homozygotes for hemoglobin

[137133]Lepore. J. Clin. Invest. 63: 736–742, 1979.

[137134]24466.

[137135]Ranney, H. M.; Jacobs, A. S.: Simultaneous occurrence of haemoglobins

[137136]C and Lepore in an Afro-American. Nature 204: 163–166, 1964.

[137137]24467.

[137138]Ranney, H. M.; Jacobs, A. S.; Ramot, B.; Bradley, T. B., Jr.:

[137139]Hemoglobin NYU, a delta chain variant, alpha 2 delta 2(12 lys). J.

[137140]Clin. Invest. 48: 2057–2062, 1969.

[137141]24468.

[137142]Renda, M.; Piazza, T.; Ciaccio, C.; Maggio, A.: Delta(+)-27 homozygosis

[137143]in a Sicilian family. Haematologica 77: 82–83, 1992.

[137144]24469.

[137145]Rieder, R. F.; Clegg, J. B.; Weiss, H. J.; Christy, N. P.; Rabinowitz,

[137146]R.: Hemoglobin A-2-Roosevelt: delta 20 val-to-glu.

Biochim. Biophys.

[137147]Acta 439: 501-504, 1976.

[137148]24470.

[137149]Romero Garcia, C.; Navarro, J. L.; Lam, H.; Webber, B. B.;

Headlee,

[137150]M. G.; Wilson, J. B.; Huisman, T. H. J.: Hb A(2)-Manzanares

or delta

[137151]121(GH4) glu-to-val, an unstable delta chain variant observed in a

[137152]Spanish family. Hemoglobin 7: 435-442, 1983.

[137153]24471.

[137154]Salkie, M. L.; Gordon, P. A.; Rigal, W. M.; Lam, H.; Wilson,

J.

[137155]B.; Headlee, M. E.; Huisman, T. H. J.: Hb A2-Canada or

delta 99(G1)

[137156]asp-to-asn, a newly discovered delta chain variant with increased

[137157]oxygen affinity occurring in cis to beta-thalassemia.

Hemoglobin 6:

[137158]223-231, 1982.

[137159]24472.

[137160]Schiliro, G.; Russo-Mancuso, G.; Dibenedetto, S. P.; Samperi,

[137161]P.; Di Cataldo, A.; Ragusa, R.; Testa, R.: Six rare hemoglobin variants

[137162]found in Sicily. Hemoglobin 15: 431-437, 1991.

[137163]24473.

[137164]Sharma, R. S.; Harding, D. L.; Wong, S. D.; Wilson, J. B.; Gravely,

[137165]M. E.; Huisman, T. H. J.: A new delta chain variant, hemoglobin-A2

[137166]Melbourne, or delta 43 glu-to-lys (CD2). Biochim. Biophys. Acta 359:

[137167]233-235, 1974.

[137168]24474.

[137169]Sharma, R. S.; Williams, L.; Wilson, J. B.; Huisman, T. H. J.

[137170]: Hemoglobin A(2) Coburg or alpha-2-delta-2 116 arg-to-his (G18). Biochim.

[137171]Biophys. Acta 393: 379-382, 1975.

[137172]24475.

[137173]Spritz, R. A.; DeRiel, J. K.; Forget, B. G.; Weissman, S. M.:

[137174]Complete nucleotide sequence of the human delta-globin gene. Cell 21:

[137175]639-646, 1980.

[137176]24476.

[137177]Trifillis, P.; Ioannou, P.; Schwartz, E.; Surrey, S.: Identification

[137178]of four novel delta-globin gene mutations in Greek Cypriots using

[137179]polymerase chain reaction and automated fluorescence-based DNA sequence

[137180]analysis. Blood 78: 3298-3305, 1991.

[137181]24477.

[137182]Vella, F.; Graham, B.: A variant of hemoglobin A(2) in Alberta

[137183]Indians. Clin. Biochem. 2: 455-460, 1969.

[137184]24478.

[137185]Williamson, D.; Brennan, S. O.; Strosberg, H.; Whitty, J.; Carell,

[137186]R. W.: Hemoglobin A(2) Fitzroy delta142 ala-to-asg: a new delta-chain

[137187]variant. Hemoglobin 8: 325–332, 1984.

[137188]24479.

[137189]Wilson, J. T.; Wilson, L. B.; Ohta, Y.: A case of homozygous

[137190]delta-thalassemia not due to a deletion of the delta-
globin structural

[137191]gene. Biochem. Biophys. Res. Commun. 99: 1035–1039,
1981.

[137192]24480.

[137193]Le Coniat, M.; Kinet, J. P.; Berger, R.: The human genes for
the

[137194]alpha and gamma subunits of the mast cell receptor for
immunoglobulin

[137195]E are located on human chromosome band 1q23. Im-
munogenetics 32:

[137196]183–186, 1990.

[137197]24481.

[137198]Pestka, S.: The human interferons--from protein purifica-
tion

[137199]and sequence to cloning and expression in bacteria: be-
fore, between,

[137200]and beyond. Arch. Biochem. Biophys. 221: 1–37, 1983.

[137201]24482.

[137202]Sehgal, P. B.; Sagar, A. D.; Braude, I. A.: Further heterogeneity

[137203]of human alpha-interferon mRNA species. Science 214: 803–805, 1981.

[137204]24483.

[137205]Shows, T. B.; Sakaguchi, A. Y.; Naylor, S. L.; Goeddel, D. V.;

[137206]Lawn, R. M.: Clustering of leukocyte and fibroblast interferon genes

[137207]on human chromosome 9. Science 218: 373–374, 1982.

[137208]24484.

[137209]Slate, D. L.; D'Eustachio, P.; Pravtcheva, D.; Cunningham, A.

[137210]C.; Nagata, S.; Weissmann, C.; Ruddle, F. H.: Chromosomal location

[137211]of a human alpha interferon gene family. J. Exp. Med. 155: 1019–1024,

[137212]1982.

[137213]24485.

[137214]Trent, J. M.; Olson, S.; Lawn, R. M.: Chromosomal localization

tion

[137215]of human leukocyte, fibroblast and immune interferon
genes by means

[137216]of in situ hybridization. Proc. Nat. Acad. Sci. 79:
7809-7813, 1982.

[137217]24486.

[137218]Ullrich, A.; Gray, A.; Goeddel, D. V.; Dull, T. J.: Nucleotide

[137219]sequence of a portion of human chromosome 9 contain-
ing a leukocyte

[137220]interferon gene cluster. J. Molec. Biol. 156: 467-486,
1982.

[137221]24487.

[137222]Virelizier, J. L.; Griscelli, C.: Defaut selectif de secretion

[137223]d'interferon associe a un deficit d'activite cytotoxique na-
turelle. Arch.

[137224]Franc. Pediat. 38: 77-81, 1981.

[137225]24488.

[137226]Virelizier, J. L.; Lenoir, G.; Griscelli, C.: Persistent Epstein-
Barr

[137227]virus infection in a child with hypergammaglobulinaemia
and immunoblastic

- [137228] proliferation associated with a selective defect in interferon secretion. Lancet II: 231–234, 1978.
- [137229] 24489.
- [137230] It Hart, L. M.; Stolk, R. P.; Heine, R. J.; Grobbee, D. E.; van der Does, F. E. E.; Maassen, J. A.: Association of the insulin-receptor variant met-985 with hyperglycemia and non-insulin-dependent diabetes mellitus in the Netherlands: a population-based study. Am. J. Hum. Genet. 59: 1119–1125, 1996.
- [137231] 24490.
- [137232] Accili, D.; Drago, J.; Lee, E. J.; Johnson, M. D.; Cool, M. H.; Salvatore, P.; Asico, L. D.; Jose, P. A.; Taylor, S. I.; Westphal, H.: Early neonatal death in mice homozygous for a null allele of the insulin receptor gene. Nature Genet. 12: 106–109, 1996.
- [137233] 24491.

[137242]Accili, D.; Frapier, C.; Mosthaf, L.; McKeon, C.; Elbein, S.
C.;

[137243]Permutt, M. A.; Ramos, E.; Lander, E.; Ullrich, A.; Taylor, S.
I.

[137244]: A mutation in the insulin receptor gene that impairs
transport of

[137245]the receptor to the plasma membrane and causes insulin-
resistant diabetes. EMBO

[137246]J. 8: 2509–2517, 1989.

[137247]24492.

[137248]Al-Gazali, L. I.; Khalil, M.; Devadas, K.: A syndrome of in-
sulin

[137249]resistance resembling leprechaunism in five sibs of con-
sanguineous

[137250]parents. J. Med. Genet. 30: 470–475, 1993.

[137251]24493.

[137252]Bar, R. S.; Muggeo, M.; Roth, J.; Kahn, C. R.; Havrankova,
J.;

[137253]Imperato-McGinley, J.: Insulin resistance, acanthosis ni-
gricans,

[137254]and normal insulin receptors in a young woman: evidence
for a postreceptor

[137255]defect. J. Clin. Endocr. Metab. 47: 620–625, 1978.

[137256]24494.

[137257]Barbetti, F.; Gejman, P. V.; Taylor, S. I.; Raben, N.; Cama, A.;

[137258]Bonora, E.; Pizzo, P.; Moghetti, P.; Muggeo, M.; Roth, J.:
Detection

[137259]of mutations in insulin receptor gene by denaturing gradient gel electrophoresis. Diabetes 41:

[137260]408–415, 1992.

[137261]24495.

[137262]Barnes, N. D.; Palumbo, P. J.; Hayles, A. B.; Folgar, H.: Insulin

[137263]resistance, skin changes, and virilization: a recessively inherited

[137264]syndrome possibly due to pineal gland dysfunction. Diabetologia 10:

[137265]285–289, 1974.

[137266]24496.

[137267]Belke, D. D.; Betuing, S.; Tuttle, M. J.; Graveleau, C.;
Young,

[137268]M. E.; Pham, M.; Zhang, D.; Cooksey, R. C.; McClain, D. A.;

Litwin,

[137269]S. E.; Taegtmeier, H.; Severson, D.; Kahn, C. R.; Abel, E.
D.: Insulin

[137270]signaling coordinately regulates cardiac size, metabolism,
and contractile

[137271]protein isoform expression. J. Clin. Invest. 109: 629–639,
2002.

[137272]24497.

[137273]Benecke, H.; Flier, J. S.; Moller, D. E.: Alternatively spliced

[137274]variants of the insulin receptor protein: expression in nor-
mal and

[137275]diabetic human tissues. J. Clin. Invest. 89: 2066–2070,
1992.

[137276]24498.

[137277]Bruning, J. C.; Gautam, D.; Burks, D. J.; Gillette, J.; Schu-
bert,

[137278]M.; Orban, P. C.; Klein, R.; Krone, W.; Muller-Wieland, D.;
Kahn,

[137279]C. R.: Role of brain insulin receptor in control of body
weight and

[137280]reproduction. Science 289: 2122–2125, 2000.

[137281]24499.

[137282]Bruning, J. C.; Michael, M. D.; Winnay, J. N.; Hayashi, T.;
Horsch,

[137283]D.; Accili, D.; Goodyear, L. J.; Kahn, C. R.: A muscle-spe-
cific insulin

[137284]receptor knockout exhibits features of the metabolic syn-
drome of NIDDM

[137285]without altering glucose tolerance. Molec. Cell 2:
559-569, 1998.

[137286]24500.

[137287]Cama, A.; de la Luz Sierra, M.; Ottini, L.; Kadowaki, T.;
Gorden,

[137288]P.; Imperato-McGinley, J.; Taylor, S. I.: A mutation in the
tyrosine

[137289]kinase domain of the insulin receptor associated with in-
sulin resistance

[137290]in an obese woman. J. Clin. Endocr. Metab. 73: 894-901,
1991.

[137291]24501.

[137292]Li, Q. Y.; Newbury-Ecob, R. A.; Terrett, J. A.; Wilson, D. I.;

[137293]Curtis, A. R. J.; Yi, C. H.; Gebuhr, T.; Bullen, P. J.; Robson,
S.

[137294]C.; Strachan, T.; Bonnet, D.; Lyonnet, S.; Young, I. D.; Rae-

burn,

[137295]J. A.; Buckler, A. J.; Law, D. J.; Brook, J. D.: Holt-Oram syndrome

[137296]is caused by mutations in TBX5, a member of the
Brachyury (T) gene

[137297]family. Nature Genet. 15: 21-29, 1997.

[137298]24502.

[137299]Terrett, J. A.; Newbury-Ecob, R.; Cross, G. S.; Fenton, I.;
Raeburn,

[137300]J. A.; Young, I. D.; Brook, J. D.: Holt-Oram syndrome is a
genetically

[137301]heterogeneous disease with one locus mapping to human
chromosome 12q. Nature

[137302]Genet. 6: 401-404, 1994.

[137303]24503.

[137304]Yang, J.; Hu, D.; Xia, J.; Yang, Y.; Ying, B.; Hu, J.; Zhou, X.

[137305]: Three novel TBX5 mutations in Chinese patients with
Holt-Oram syndrome. Am.

[137306]J. Med. Genet. 92: 237-240, 2000.

[137307]24504.

[137308]Henry, I.; Humphries, S. E.; Tata, F.; Barichard, F.; Holm,

M.;

[137309]Williamson, R.; Junien, C.: The gene for HMG CoA reductase (HMGCR)

[137310]is on human chromosome 5. (Abstract) Cytogenet. Cell Genet. 40:

[137311]649-650, 1985.

[137312]24505.

[137313]Humphries, S. E.; Tata, F.; Henry, I.; Barichard, F.; Holm, M.;

[137314]Junien, C.; Williamson, R.: The isolation, characterisation, and

[137315]chromosomal assignment of the gene for human 3-hydroxy-3-methylglutaryl

[137316]coenzyme A reductase, (HMG-CoA reductase). Hum. Genet. 71: 254-258,

[137317]1985.

[137318]24506.

[137319]Rutherford, S.; Johnson, M. P.; Curtain, R. P.; Griffiths, L.

[137320]R.: Chromosome 17 and the inducible nitric oxide synthase gene in

[137321]human essential hypertension. Hum. Genet. 109:

408-415, 2001.

[137322]24507.

[137323]Walter, J. W.; North, P. E.; Waner, M.; Mizeracki, A.; Blei, F.;

[137324]Walker, J. W. T.; Reinisch, J. F.; Marchuk, D. A.: Somatic
mutation

[137325]of vascular endothelial growth factor receptors in juvenile
hemangioma. Genes

[137326]Chromosomes Cancer 33: 295-303, 2002.

[137327]24508.

[137328]Ragoussis, J.; Senger, G.; Trowsdale, J.; Campbell, I. G.:
Genomic

[137329]organization of the human folate receptor genes on chro-
mosome 11q13. Genomics 14:

[137330]423-430, 1992.

[137331]24509.

[137332]Barber, R. C.; Shaw, G. M.; Lammer, E. J.; Greer, K. A.;
Biela,

[137333]T. A.; Lacey, S. W.; Wasserman, C. R.; Finnell, R. H.: Lack of
association

[137334]between mutations in the folate receptor-alpha gene and
spina bifida. Am.

[137335]J. Med. Genet. 76: 310–317, 1998.

[137336]24510.

[137337]Bowcock, A. M.; Lacey, S.; Saltman, D.; Mohandas, T. K.;
Kamen,

[137338]B. A.; Taggart, R. T.: Localization of the folate receptor
gene to

[137339]chromosome 11q13. (Abstract) Cytogenet. Cell Genet. 58:
1955 only,

[137340]1991.

[137341]24511.

[137342]Campbell, I. G.; Jones, T. A.; Foulkes, W. D.; Trowsdale, J.:

[137343]Folate-binding protein is a marker for ovarian cancer.
Cancer Res. 51:

[137344]5329–5338, 1991.

[137345]24512.

[137346]Kranes, A.; Balogh, K., Jr.: Liver disease in a patient with

[137347]von Hippel–Lindau disease. New Eng. J. Med. 275:
950–959, 1966.

[137348]24513.

[137349]Lamiell, J. M.: Personal Communication. San Francisco,

Calif.

[137350]9/1987.

[137351]24514.

[137352]Lamiell, J. M.; Salazar, F. G.; Hsia, Y. E.: Von Hippel–Lindau

[137353]disease affecting 43 members of a single kindred.

Medicine 68: 1–29,

[137354]1989.

[137355]24515.

[137356]Latif, F.; Tory, K.; Gnarr, J.; Yao, M.; Duh, F.–M.; Orcutt,

[137357]M. L.; Stackhouse, T.; Kuzmin, I.; Modi, W.; Geil, L.;

Schmidt, L.;

[137358]Zhou, F.; Li, H.; Wei, M. H.; Chen, F.; Glenn, G.; Choyke, P.;

Walther,

[137359]M. M.; Weng, Y.; Duan, D.–S. R.; Dean, M.; Glavac, D.;

Richards, F.

[137360]M.; Crossey, P. A.; Ferguson–Smith, M. A.; Le Paslier, D.;

Chumakov,

[137361]J.; Cohen, D.; Chinault, A. C.; Maher, E. R.; Linehan, W. M.;

Zbar,

[137362]B.; Lerman, M. I.: Identification of the von Hippel–Lindau

disease

[137363]tumor suppressor gene. Science 260: 1317–1320, 1993.

[137364]24516.

[137365]Lee, S.; Chen, D. Y. T.; Humphrey, J. S.; Gnarr, J. R.; Linehan,

[137366]W. M.; Klausner, R. D.: Nuclear/cytoplasmic localization of the von

[137367]Hippel–Lindau tumor suppressor gene product is determined by cell

[137368]density. Proc. Nat. Acad. Sci. 93: 1770–1775, 1996.

[137369]24517.

[137370]Lenz, T.; Thiede, H. M.; Nussberger, J.; Atlas, S. A.; Distler,

[137371]A.; Schulte, K. L.: Hyperreninemia and secondary hyperaldosteronism

[137372]in a patient with pheochromocytoma and von Hippel–Lindau disease. Nephron 62:

[137373]345–350, 1992.

[137374]24518.

[137375]Lindau, A.: Zur Frage der Angiomatosis Retinae und Ihrer Hirncomplication. Acta

[137376]Ophthal. 4: 193–226, 1927.

[137377]24519.

[137378]Loeb, D. B.; Pericak-Vance, M. A.; Stajich, J. M.; Vance, J. M.

[137379]: A novel mutation in the von Hippel-Lindau gene. Hum. Molec. Genet. 3:

[137380]1423-1424, 1994.

[137381]24520.

[137382]Lui, W. O.; Chen, J.; Glasker, S.; Bender, B. U.; Madura, C.;

[137383]Khoo, S. K.; Kort, E.; Larsson, C.; Neumann, H. P. H.; Teh, B. T.

[137384]: Selective loss of chromosome 11 in pheochromocytomas associated

[137385]with the VHL syndrome. Oncogene 21: 1117-1122, 2002.

[137386]24521.

[137387]Maddock, I. R.; Moran, A.; Maher, E. R.; Teare, M. D.; Norman,

[137388]A.; Payne, S. J.; Whitehouse, R.; Dodd, C.; Lavin, M.; Hartley, N.;

[137389]Super, M.; Evans, D. G. R.: A genetic register for von Hippel-Lindau

[137390]disease. J. Med. Genet. 33: 120-127, 1996.

[137391]24522.

[137392]Maher, E. R.; Bentley, E.; Yates, J. R. W.; Barton, D.; Jennings,

[137393]A.; Fellows, I. W.; Ponder, M. A.; Ponder, B. A. J.; Benjamin, C.;

[137394]Harris, R.; Ferguson-Smith, M. A.: Mapping of von Hippel-Lindau disease

[137395]to chromosome 3p confirmed by genetic linkage analysis. J. Neurol.

[137396]Sci. 100: 27-30, 1990.

[137397]24523.

[137398]Maher, E. R.; Bentley, E.; Yates, J. R. W.; Latif, F.; Lerman,

[137399]M.; Zbar, B.; Affara, N. A.; Ferguson-Smith, M. A.: Mapping of the

[137400]von Hippel-Lindau disease locus to a small region of chromosome 3p

[137401]by genetic linkage analysis. Genomics 10: 957-960, 1991.

[137402]24524.

[137403]Maher, E. R.; Iselius, L.; Yates, J. R. W.; Littler, M.; Benjamin,

[137404]C.; Harris, R.; Sampson, J.; Williams, A.; Ferguson-Smith,

M. A.;

[137405]Morton, N.: Von Hippel–Lindau disease: a genetic study. J. Med.

[137406]Genet. 28: 443–447, 1991.

[137407]24525.

[137408]Maher, E. R.; Yates, J. R. W.; Ferguson–Smith, M. A.: Statistical

[137409]analysis of the two stage mutation model in von Hippel–Lindau disease,

[137410]and in sporadic cerebellar haemangioblastoma and renal cell carcinoma. J.

[137411]Med. Genet. 27: 311–314, 1990.

[137412]24526.

[137413]Mahon, P. C.; Hirota, K.; Semenza, G. L.: FIH–1: a novel protein

[137414]that interacts with HIF–1–alpha and VHL to mediate repression of HIF–1

[137415]transcriptional activity. Genes Dev. 15: 2675–2686, 2001.

[137416]

[137417]100. Manski, T. J.; Heffner, D. K.; Glenn, G. M.; Patronas, N. J.;

[137418]Pikus, A. T.; Katz, D.; Lebovics, R.; Sledjeski, K.; Choyke, P. L.;

[137419]Zbar, B.; Linehan, W. M.; Oldfield, E. H.: Endolymphatic sac tumors:

[137420]a source of morbid hearing loss in von Hippel–Lindau disease. J.A.M.A. 277:

[137421]1461–1466, 1997.

[137422]

[137423]101. Maxwell, P. H.; Wiesener, M. S.; Chang, G.–W.; Clifford, S. C.;

[137424]Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.; Maher, E.

[137425]R.; Ratcliffe, P. J.: The tumour suppressor protein VHL targets hypoxia-inducible

[137426]factors for oxygen-dependent proteolysis. Nature 399: 271–275, 1999.

[137427]

[137428]102. McCabe, C. M.; Flynn, H. W., Jr.; Shields, C. L.; Shields, J.

[137429]A.; Regillo, C. D.; McDonald, H. R.; Berrocal, M. H.; Gass, J. D.

[137430]M.; Mieler, W. F.: Juxtapapillary capillary hemangiomas:

clinical

[137431] features and visual acuity outcomes. Ophthalmology 107:
2240–2249,

[137432] 2000.

[137433]

[137434] 103. McKusick, V. A.; Abbey, H.; Bettersby, E. J.; Borhani,
N. O.;

[137435] Boyer, S. H., IV; Cohen, B. H.; Ferguson–Smith, M. A.;
Franke, F.

[137436] R.; Gordon, H.; Handmaker, S. D.; Harris, W. S.; Hawkins,
M. R.;

[137437] and 13 others: Medical genetics 1960. J. Chronic Dis. 14:
1–198,

[137438] 1961. Fig. 71.

[137439]

[137440] 104. Melmon, K. L.; Rosen, S. W.: Lindau's disease: review
of the

[137441] literature and study of a large kindred. Am. J. Med. 36:
595–617,

[137442] 1964.

[137443]

- [137444]105. Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W. G., Jr.;
[137445]Pavletich, N. P.: Structure of an HIF-1- α -pVHL complex: hydroxyproline
[137446]recognition in signaling. Science 296: 1886-1889, 2002.
[137447]
[137448]106. Mukhopadhyay, D.; Knebelmann, B.; Cohen, H. T.;
Ananth, S.; Sukhatme,
[137449]V. P.: The von Hippel-Lindau tumor suppressor gene
product interacts
[137450]with Sp1 to repress vascular endothelial growth factor
promoter activity. Molec
[137451]. Cell. Biol. 17: 5629-5639, 1997.
[137452]
[137453]107. Neumann, H. P. H.; Berger, D. P.; Sigmund, G.; Blum,
U.; Schmidt,
[137454]D.; Parmer, R. J.; Volk, B.; Kirste, G.: Pheochromocytomas,
multiple
[137455]endocrine neoplasia type 2, and von Hippel-Lindau disease. New Eng.
[137456]J. Med. 329: 1531-1538, 1993.

[137457]

[137458]108. Neumann, H. P. H.; Eng, C.; Mulligan, L. M.; Glavac, D.; Zauner,

[137459]J.; Ponder, B. A. J.; Crossey, P. A.; Maher, E. R.; Brauch, H.:
Consequences

[137460]of direct genetic testing for germline mutations in the
clinical management

[137461]of families with multiple endocrine neoplasia, type II.

J.A.M.A. 274:

[137462]1149–1151, 1995.

[137463]

[137464]109. Neumann, H. P. H.; Wiestler, O. D.: Clustering of fea-
tures of

[137465]von Hippel–Lindau syndrome: evidence for a complex ge-
netic locus. Lancet 337:

[137466]1052–1054, 1991.

[137467]

[137468]110. Nibbelink, D. W.; Peters, B. H.; McCormick, W. F.: On
the association

[137469]of pheochromocytoma and cerebellar hemangioblastoma.
Neurology 19:

[137470]455–460, 1969.

[137471]

[137472]111. Oberstrass, J.; Reifenberger, G.; Reifenberger, J.;
Wechsler,

[137473]W.; Collins, V. P.: Mutation of the von Hippel–Lindau tu-
mour suppressor

[137474]gene in capillary haemangioblastomas of the central ner-
vous system. J.

[137475]Path. 179: 151–156, 1996.

[137476]

[137477]112. Ohh, M.; Yauch, R. L.; Lonergan, K. M.; Whaley, J. M.;
Stemmer–Rachamimov,

[137478]A. O.; Louis, D. N.; Gavin, B. J.; Kley, N.; Kaelin, W. G., Jr.;
Iliopoulos,

[137479]O.: The von Hippel–Lindau tumor suppressor protein is
required for

[137480]proper assembly of an extracellular fibronectin matrix.
Molec. Cell 1:

[137481]959–968, 1998.

[137482]

[137483]113. Olschwang, S.; Richard, S.; Boisson, C.; Giraud, S.;

Laurent-Puig,

[137484]P.; Resche, F.; Thomas, G.: Germline mutation profile of the VHL

[137485]gene in von Hippel-Lindau disease and in sporadic hemangioblastoma. Hum.

[137486]Mutat. 12: 424-430, 1998.

[137487]

[137488]114. Otenasek, F. J.; Silver, M. L.: Spinal hemangioma (hemangioblastoma)

[137489]in Lindau's disease: report of six cases in a single family. J. Neurosurg. 18:

[137490]295-300, 1961.

[137491]

[137492]115. Pack, S. D.; Zbar, B.; Pak, E.; Ault, D. O.; Humphrey, J. S.;

[137493]Pham, T.; Hurley, K.; Weil, R. J.; Park, W.-S.; Kuzmin, I.; Stolle,

[137494]C.; Glenn, G.; Liotta, L. A.; Lerman, M. I.; Klausner, R. D.; Linehan,

[137495]W. M.; Zhuang, Z.: Constitutional von Hippel-Lindau (VHL) gene deletions

[137496]detected in VHL families by fluorescence in situ hybridiza-

tion. Cancer

[137497]Res. 59: 5560–5564, 1999.

[137498]

[137499]116. Pause, A.; Lee, S.; Lonergan, K. M.; Klausner, R. D.:

The von

[137500]Hippel–Lindau tumor suppressor gene is required for cell
cycle exit

[137501]upon serum withdrawal. Proc. Nat. Acad. Sci. 95:
993–998, 1998.

[137502]

[137503]117. Price, E. B., Jr.: Papillary cystadenoma of the epi-
didymis:

[137504]a clinicopathologic analysis of 20 cases. Arch. Path. 91:
456–470,

[137505]1971.

[137506]

[137507]118. Probst, A.; Lotz, M.; Heitz, P.: Von Hippel–Lindau's
disease,

[137508]syringomyelia and multiple endocrine tumors: a complex
neuroendocrinopathy. Virchows

[137509]Arch. Path. Anat. Histol. 378: 265–272, 1978.

[137510]

[137511]119. Prowse, A. H.; Webster, A. R.; Richards, F. M.;
Richard, S.;

[137512]Olschwang, S.; Resche, F.; Affara, N. A.; Maher, E. R.: So-
matic inactivation

[137513]of the VHL gene in Von Hippel–Lindau disease tumors.
Am. J. Hum.

[137514]Genet. 60: 765–771, 1997.

[137515]

[137516]120. Rho, Y. M.: Von Hippel–Lindau's disease: a report of
5 cases. Canad.

[137517]Med. Assoc. J. 101: 135–142, 1969.

[137518]

[137519]121. Richard, S.; Croisille, L.; Yvart, J.; Casadeval, N.; Es-
chwege,

[137520]P.; Aghakhani, N.; David, P.; Gaudric, A.; Scigalla, P.; Her-
mine,

[137521]O.: Paradoxical secondary polycythemia in von Hippel–
Lindau patients

[137522]treated with anti–vascular endothelial growth factor re-
ceptor therapy. Blood 99:

[137523]3851–3853, 2002.

[137524]

[137525]122. Richards, F. M.; Crossey, P. A.; Phipps, M. E.; Foster, K.; Latif,

[137526]F.; Evans, G.; Sampson, J.; Lerman, M. I.; Zbar, B.; Affara, N. A.;

[137527]Ferguson-Smith, M. A.; Maher, E. R.: Detailed mapping of germline

[137528]deletions of the von Hippel–Lindau disease tumour suppressor gene. Hum.

[137529]Molec. Genet. 3: 595–598, 1994.

[137530]

[137531]123. Richards, F. M.; Maher, E. R.; Latif, F.; Phipps, M. E.; Tory,

[137532]K.; Lush, M.; Crossey, P. A.; Oostra, B.; Gustavson, K. H.; Green,

[137533]J.; Turner, G.; Yates, J. R. W.; Linehan, W. M.; Affara, N. A.; Lerman,

[137534]M.; Zbar, B.; Ferguson-Smith, M. A.: Detailed genetic mapping of

[137535]the von Hippel–Lindau disease tumour suppressor gene. J. Med. Genet. 30:

[137536]104–107, 1993.

[137537]

[137538]124. Richards, F. M.; Phipps, M. E.; Latif, F.; Yao, M.;
Crossey,

[137539]P. A.; Foster, K.; Linehan, W. M.; Affara, N. A.; Lerman, M.
I.; Zbar,

[137540]B.; Ferguson-Smith, M. A.; Maher, E. R.: Mapping the von
Hippel–Lindau

[137541]disease tumour suppressor gene: identification of
germline deletions

[137542]by pulsed field gel electrophoresis. Hum. Molec. Genet. 2:
879–882,

[137543]1993.

[137544]

[137545]125. Richards, F. M.; Schofield, P. N.; Fleming, S.; Maher,
E. R.

[137546]: Expression of the von Hippel–Lindau disease tumour
suppressor gene

[137547]during human embryogenesis. Hum. Molec. Genet. 5:
639–644, 1996.

[137548]

[137549]126. Rubenstein, J. L.; Yaari, H.: von Hippel–Lindau and
the genetics

[137550]of astrocytoma. J. Nat. Cancer Inst. 86: 142–143, 1994.

[137551]

[137552]127. Sander, S.; Normann, T.; Mathisen, W.: Pheochromo-
cytoma associated

[137553]with von–Hippel–Lindau's disease in a family. Scand. J.
Urol. Nephrol. 4:

[137554]259–263, 1970.

[137555]

[137556]128. Schechterman, L.: Lindau's disease: report of an un-
usual case

[137557]and two additional cases in a Negro family. Med. Ann.
D.C. 30: 64–76,

[137558]1961.

[137559]

[137560]129. Schimke, R. N.; Collins, D. L.; Rothberg, P. G.: Func-
tioning

[137561]carotid paraganglioma in the von Hippel–Lindau syn-
drome. (Letter) Am.

[137562]J. Med. Genet. 80: 533–534, 1998.

[137563]

[137564]130. Schoenfeld, A.; Davidowitz, E. J.; Burk, R. D.: A second major

[137565]native von Hippel–Lindau gene product, initiated from an internal

[137566]translation start site, functions as a tumor suppressor. Proc. Nat.

[137567]Acad. Sci. 95: 8817–8822, 1998.

[137568]

[137569]131. Seizinger, B. R.; Farmer, G.; Haines, J.; Anderson, K.; Whaley,

[137570]J.; Hettlich, C.; Decker, J.; Rouleau, G.; Smith, D.; Drabkin, H.;

[137571]Filling–Katz, M.; Neumann, H.; Collins, D.; Hsia, E.; Green, J.; Waziri,

[137572]M.; Gusella, J.; Li, F.: Isolating the gene(s) for von Hippel–Lindau

[137573]disease and renal cell carcinoma. (Abstract) Am. J. Hum. Genet. 45

[137574](suppl.): A32, 1989.

[137575]

[137576]132. Seizinger, B. R.; Rouleau, G. A.; Ozelius, L. J.; Lane,
[137577]A. H.; Farmer, G. E.; Lamiell, J. M.; Haines, J.; Yuen, J.
[137578]W.; Collins, D.; Majoor-Krakauer, D.; et al.: Von Hippel-
Lindau
[137579]disease maps to the region of chromosome 3 associated
with renal cell
[137580]carcinoma. Nature 332: 268-269, 1988.

[137581]

[137582]133. Seizinger, B. R.; Smith, D. I.; Filling-Katz, M. R.; Neumann,

[137583]H.; Green, J. S.; Choyke, P. L.; Anderson, K. M.; Freiman, R. N.;

[137584]Klauck, S. M.; Whaley, J.; Decker, H.-J. H.; Hsia, Y. E.; Collins,

[137585]D.; Halperin, J.; Lamiell, J. M.; Oostra, B.; Waziri, M. H.; Gorin,

[137586]M. B.; Scherer, G.; Drabkin, H. A.; Aronin, N.; Schinzel, A.; Martuza,

[137587]R. L.; Gusella, J. F.; Haines, J. L.: Genetic flanking markers refine

[137588]diagnostic criteria and provide insights into the genetics of Von

[137589]Hippel Lindau disease. Proc. Nat. Acad. Sci. 88: 2864-2868, 1991.

[137590]

[137591]134. Sgambati, M. T.; Stolle, C.; Choyke, P. L.; Walther, M. M.; Zbar,

[137592]B.; Linehan, W. M.; Glenn, G. M.: Mosaicism in von Hippel-Lindau

[137593]disease: lessons from kindreds with germline mutations
identified

[137594]in offspring with mosaic parents. Am. J. Hum. Genet. 66:
84–91,

[137595]2000.

[137596]

[137597]135. Sharp, W. V.; Platt, R. L.: Familial pheochromocytoma:
association

[137598]with von-Hippel-Lindau's disease. Angiology 22:
141–146, 1971.

[137599]

[137600]136. Shokeir, M. H. K.: Von Hippel–Lindau syndrome: a
report on three

[137601]kindreds. J. Med. Genet. 7: 155–157, 1970.

[137602]

[137603]137. Silver, M. L.: Hereditary vascular tumors of the ner-
vous system. J.A.M.A.

[137604]156: 1053–1056, 1954.

[137605]

[137606]138. Stolle, C.; Glenn, G.; Zbar, B.; Humphrey, J. S.;
Choyke, P.;

[137607]Walther, M.; Pack, S.; Hurley, K.; Audrey, C.; Klausner, R.;
Linehan,

[137608]W. M.: Improved detection of germline mutations in the
von Hippel–Lindau

[137609]disease tumor suppressor gene. Hum. Mutat. 12:
417–423, 1998.

[137610]

[137611]139. Thomas, M.; Burnside, R. M.: Von Hippel–Lindau dis-
ease. Am.

[137612]J. Ophthalm. 51: 140–146, 1961.

[137613]

[137614]140. Tisherman, S. E.; Gregg, F. J.; Danowski, T. S.: Famil-
ial pheochromocytoma

[137615]. J.A.M.A. 182: 152–156, 1962.

[137616]

[137617]141. Tisherman, S. E.; Tisherman, B. G.; Tisherman, S. A.;
Dunmire,

[137618]S.; Levey, G. S.; Mulvihill, J. J.: Three-decade investigation
of

[137619]familial pheochromocytoma: an allele of von Hippel–Lin-
dau disease? Arch.

[137620]Int. Med. 153: 2550–2556, 1993.

[137621]

[137622]142. Tory, K.; Brauch, H.; Linehan, M.; Barba, D.; Oldfield, E.; Filling-Katz,

[137623]M.; Seizinger, B.; Nakamura, Y.; White, R.; Marshall, F. F.; Lerman,

[137624]M. I.; Zbar, B.: Specific genetic change in tumors associated with

[137625]von Hippel–Lindau disease. J. Nat. Cancer Inst. 81: 1097–1101, 1989.

[137626]

[137627]143. Tsuda, H.; Fukushima, S.; Takahashi, M.; Hikosaka, Y.; Hayashi,

[137628]K.: Familial bilateral papillary cystadenoma of the epididymis: report

[137629]of three cases in siblings. Cancer 37: 1831–1839, 1976.

[137630]

[137631]144. Tyers, M.; Willems, A. R.: One ring to rule a super-family of

[137632]E3 ubiquitin ligases. Science 284: 602–604, 1999.

[137633]

- [137634]145. Vance, J. M.; Small, K. W.; Jones, M. A.; Stajich, J. M.; Yamaoka, L. H.; Roses, A. D.; Hung, W.-Y.; Pericak-Vance, M. A.: Confirmation of linkage in von Hippel-Lindau disease. Genomics 6: 565-567, 1990.
- [137637]
- [137638]146. van der Harst, E.; de Krijger, R. R.; Dinjens, W. N. M.; Weeks, L. E.; Bonjer, H. J.; Bruining, H. A.; Lamberts, S. W. J.; Koper, J. W.: Germline mutations in the vhl gene in patients presenting with phaeochromocytomas. Int. J. Cancer. 77: 337-340, 1998.
- [137642]
- [137643]147. Versteeg, R.: Aberrant methylation in cancer. (Editorial) Am. J. Hum. Genet. 60: 751-754, 1997.
- [137645]
- [137646]148. Vogelstein, B.: Personal Communication. Baltimore, Md. 1/6/1995.

[137647]

[137648]149. von Hippel, E.: Ueber eine sehr seltene Erkrankung
der Netzhaut. Albrecht

[137649]von Graefes Arch. Ophthal. 59: 83–106, 1904.

[137650]

[137651]150. Webster, A. R.; Maher, E. R.; Bird, A. C.; Moore, A. T.:
Risk

[137652]of multisystem disease in isolated ocular angioma
(haemangioblastoma). J.

[137653]Med. Genet. 37: 62–63, 2000.

[137654]

[137655]151. Webster, A. R.; Richards, F. M.; MacDonald, F. E.;
Moore, A.

[137656]T.; Maher, E. R.: An analysis of phenotypic variation in the
familial

[137657]cancer syndrome von Hippel–Lindau disease: evidence for
modifier effects. Am.

[137658]J. Hum. Genet. 63: 1025–1035, 1998.

[137659]

[137660]152. Wells, R. A.; Reenders, S. T.; Green, J.; Johnson, G. J.;
Robson,

- [137661]K. J. H.: Linkage studies of von Hippel–Lindau syndrome using multiple–locus hypervariable DNA probes. (Abstract) Cytogenet. Cell Genet. 46: 714, 1987.
- [137664]
- [137665]153. Wesolowski, D. P.; Ellwood, R. A.; Schwab, R. E.; Farah, J.: Hippel–Lindau syndrome in identical twins. Brit. J. Radiol. 54: 982–986, 1981.
- [137668]
- [137669]154. Wiesener, M. S.; Seyfarth, M.; Warnecke, C.; Jurgensen, J. S.; Rosenberger, C.; Morgan, N. V.; Maher, E. R.; Frei, U.; Eckardt, K.–U. Paraneoplastic erythrocytosis associated with an inactivating point mutation of the von Hippel–Lindau gene in a renal cell carcinoma. Blood 99: 3562–3565, 2002.

[137674]

[137675]155. Wise, K. S.; Gibson, J. A.: Von Hippel–Lindau's disease and

[137676]phaeochromocytoma. Brit. Med. J. 1: 441, 1971.

[137677]

[137678]156. Wyburn–Mason, R.: Arteriovenous aneurysm of mid-brain and retina,

[137679]facial naevi and mental changes. Brain 66: 163–203, 1943.

[137680]

[137681]157. Zatyka, M.; da Silva, N. F.; Clifford, S. C.; Morris, M. R.;

[137682]Wiesener, M. S.; Eckardt, K.–U.; Houlston, R. S.; Richards, F. M.;

[137683]Latif, F.; Maher, E. R.: Identification of cyclin D1 and other novel

[137684]targets for the von Hippel–Lindau tumor suppressor gene by expression

[137685]array analysis and investigation of cyclin D1 genotype as a modifier

[137686]in von Hippel–Lindau disease. Cancer Res. 62: 3803–3811, 2002.

[137687]

[137688]158. Zbar, B.; Kishida, T.; Chen, F.; Schmidt, L.; Maher, E. R.; Richards,

[137689]F. M.; Crossey, P. A.; Webster, A. R.; Affara, N. A.; Ferguson-Smith,

[137690]M. A.; Brauch, H.; Glavac, D.; and 14 others: Germline mutations

[137691]in the Von Hippel-Lindau disease (VHL) gene in families from North

[137692]America, Europe, and Japan. Hum. Mutat. 8: 348-357, 1996.

[137693]

[137694]159. Zhuang, Z.; Emmert-Buck, M. R.; Roth, M. J.; Gnarr, J.; Linehan,

[137695]W. M.; Liotta, L. A.; Lubensky, I. A.: Von Hippel-Lindau disease

[137696]gene deletion detected in microdissected sporadic human colon carcinoma

[137697]specimens. Hum. Path. 27: 152-156, 1996.

[137698]24527.

[137699]Devlin, R. H.; Deeb, S.; Brunsell, J.; Hayden, M. R.: Partial

[137700] gene duplication involving exon-Alu interchange results
in lipoprotein

[137701] lipase deficiency. *Am. J. Hum. Genet.* 46: 112-119, 1990.

[137702] 24528.

[137703] Dichek, H. L.; Fojo, S. S.; Beg, O. U.; Skarlatos, S. I.; Brun-
zell,

[137704] J. D.; Cutler, G. B., Jr.; Brewer, H. B., Jr.: Identification of

[137705] 2 separate allelic mutations in the lipoprotein lipase gene
of a patient

[137706] with the familial hyperchylomicronemia syndrome. *J. Biol.*
Chem. 266:

[137707] 473-477, 1991.

[137708] 24529.

[137709] Eckel, R. H.: Lipoprotein lipase: a multifunctional enzyme
relevant

[137710] to common metabolic diseases. *New Eng. J. Med.* 320:
1060-1068, 1989.

[137711] 24530.

[137712] Emi, M.; Hata, A.; Robertson, M.; Iverius, P.-H.; Hegele, R.;

[137713] Lalouel, J.-M.: Lipoprotein lipase deficiency resulting from
a nonsense

[137714]mutation in exon 3 of the lipoprotein lipase gene. Am. J. Hum. Genet. 47:

[137715]107–111, 1990.

[137716]24531.

[137717]Emi, M.; Wilson, D. E.; Iverius, P.–H.; Wu, L.; Hata, A.; Hegele,

[137718]R.; Williams, R. R.; Lalouel, J.–M.: Missense mutation (gly-to-glu188)

[137719]of human lipoprotein lipase imparting functional deficiency. J. Biol.

[137720]Chem. 265: 5910–5916, 1990.

[137721]24532.

[137722]Faustinella, F.; Chang, A.; Van Biervliet, J. P.; Rosseneu, M.;

[137723]Vinaimont, N.; Smith, L. C.; Chen, S.–H.; Chan, L.: Catalytic triad

[137724]residue mutation (asp156-to-gly) causing familial lipoprotein lipase

[137725]deficiency: co-inheritance with a nonsense mutation (ser447-to-ter)

[137726]in a Turkish family. J. Biol. Chem. 266: 14418–14424, 1991.

[137727]24533.

[137728]Feoli-Fonseca, J. C.; Levy, E.; Godard, M.; Lambert, M.: Familial

[137729]lipoprotein lipase deficiency in infancy: clinical, biochemical, and

[137730]molecular study. J. Pediat. 133: 417-423, 1998.

[137731]24534.

[137732]Fisher, R. M.; Humphries, S. E.; Talmud, P. J.: Common variation

[137733]in the lipoprotein lipase gene: effects on plasma lipids and risk

[137734]of atherosclerosis. Atherosclerosis 135: 145-159, 1997.

[137735]24535.

[137736]Franklin, S. M.: Splenomegaly with lipaemia. Proc. Roy. Soc.

[137737]Med. 30: 711, 1937.

[137738]24536.

[137739]Funke, H.; Klug, J.; Assmann, G.: Hind III RFLP in the lipoprotein

[137740]lipase gene, (LPL). Nucleic Acids Res. 15: 9102, 1987.

[137741]24537.

[137742]Funke, H.; Reckwerth, A.; Stapenhorst, D.; Schulze Beier-
ing, M.;

[137743]Jansen, M.; Assmann, G.: Bst NI (Eco RI) RFLP in the
lipoprotein

[137744]lipase gene (LPL). Nucleic Acids Res. 16: 2741, 1988.

[137745]24538.

[137746]Gagne, C.; Brun, L. D.; Julien, P.; Moorjani, S.; Lupien, P. J.

[137747]: Primary lipoprotein lipase activity deficiency: clinical in-
vestigation

[137748]of a French Canadian population. Canad. Med. Assoc. J.
140: 405-411,

[137749]1989.

[137750]24539.

[137751]Gilbert, B.; Rouis, M.; Griglio, S.; de Lumley, L.; Laplaud,
P.-M.

[137752]: Lipoprotein lipase (LPL) deficiency: a new patient ho-
mozygote for

[137753]the preponderant mutation gly88-to-glu in the human
LPL gene and review

[137754]of reported mutations: 75% are clustered in exons 5 and

6. Ann. Genet. 44:

[137755]25–32, 2001.

[137756]24540.

[137757]Ginzinger, D. G.; Lewis, M. E. S.; Ma, Y.; Jones, B. R.; Liu,

[137758]G.; Jones, S. D.; Hayden, M. R.: A mutation in the lipoprotein lipase

[137759]gene is the molecular basis of chylomicronemia in a colony of domestic

[137760]cats. J. Clin. Invest. 97: 1257–1266, 1996.

[137761]24541.

[137762]Gotoda, T.; Yamada, N.; Kawamura, M.; Kozaki, K.; Mori, N.; Ishibashi,

[137763]S.; Shimano, H.; Takaku, F.; Yazaki, Y.; Furuichi, Y.; Murase, T.

[137764]: Heterogeneous mutations in the human lipoprotein lipase gene in

[137765]patients with familial lipoprotein lipase deficiency. J. Clin. Invest. 88:

[137766]1856–1864, 1991.

[137767]24542.

[137768]Gotoda, T.; Yamada, N.; Murase, T.; Miyake, S.; Murakami,

R.;

[137769]Kawamura, M.; Kozaki, K.; Mori, N.; Shimano, H.; Shimada, M.; Yazaki,

[137770]Y.: A newly identified null allelic mutation in the human lipoprotein

[137771]lipase (LPL) gene of a compound heterozygote with familial LPL deficiency. *Biochim.*

[137772]*Biophys. Acta* 1138: 353-356, 1992.

[137773]24543.

[137774]Hata, A.; Emi, M.; Luc, G.; Basdevant, A.; Gambert, P.; Iverius,

[137775]P.-H.; Lalouel, J.-M.: Compound heterozygote for lipoprotein lipase

[137776]deficiency: ser-to-thr(244) and transition in 3-prime splice site

[137777]of intron 2 (AG-to-AA) in the lipoprotein lipase gene. *Am. J. Hum.*

[137778]*Genet.* 47: 721-726, 1990.

[137779]24544.

[137780]Hata, A.; Ridinger, D. N.; Sutherland, S. D.; Emi, M.; Kwong,

[137781]L. K.; Shuhua, J.; Lubbers, A.; Guy-Grand, B.; Basdevant,

A.; Iverius,

[137782]P. H.; Wilson, D. E.; Lalouel, J.-M.: Missense mutations in
exon

[137783]5 of the human lipoprotein lipase gene: inactivation cor-
relates with

[137784]loss of dimerization. J. Biol. Chem. 267: 20132-20139,
1992.

[137785]24545.

[137786]Haubenwallner, S.; Horl, G.; Shachter, N. S.; Presta, E.;
Fried,

[137787]S. K.; Hofler, G.; Kostner, G. M.; Breslow, J. L.; Zechner, R.:
A

[137788]novel missense mutation in the gene for lipoprotein lipase
resulting

[137789]in a highly conservative amino acid substitution
(asp180-to-glu) causes

[137790]familial chylomicronemia (type I hyperlipoproteinemia).
Genomics 18:

[137791]392-396, 1993.

[137792]24546.

[137793]Havel, R. J.; Gordon, R. S.: Idiopathic hyperlipemia:
metabolic

[137794]studies in an affected family. J. Clin. Invest. 39:
1777-1790, 1960.

[137795]24547.

[137796]Heaney, A. P.; Sharer, N.; Rameh, B.; Braganza, J. M.; Dur-
rington,

[137797]P. N.: Prevention of recurrent pancreatitis in familial
lipoprotein

[137798]lipase deficiency with high-dose antioxidant therapy. J.
Clin. Endocr.

[137799]Metab. 84: 1203-1205, 1999.

[137800]24548.

[137801]Heinzmann, C.; Ladas, J.; Antonarakis, S.; Kirchgessner,
T.;

[137802]Schotz, M.; Lasis, A. J.: RFLP for the human lipoprotein li-
pase (LPL)

[137803]gene: HindIII. Nucleic Acids Res. 15: 6763, 1987.

[137804]24549.

[137805]Heizmann, C.; Kirchgessner, T.; Kwiterovich, P. O.; Ladas,
J.

[137806]A.; Derby, C.; Antonarakis, S. E.; Lasis, A. J.: DNA poly-
morphism

[137807]haplotypes of the human lipoprotein lipase gene: possible association

[137808]with high density lipoprotein levels. Hum. Genet. 86: 578–584, 1991.

[137809]24550.

[137810]Henderson, H.; Ma, Y.; Kastelein, J.; Roederer, G.; Julien, P.;

[137811]Brunzell, J.; Hayden, M. R.: Identification of the molecular defects

[137812]underlying chylomicronemia in the majority of 75 separate probands

[137813]with LPL deficiency.(Abstract) Clin. Res. 39: 336A, 1991.

[137814]24551.

[137815]Henderson, H. E.; Bijvoet, S. M.; Mannens, M. A. M. M.; Bruin,

[137816]T.; Erkelens, D. W.; Hayden, M. R.; Kastelein, J. J. P.: Ile225-to-thr

[137817]loop mutation in the lipoprotein lipase (LPL) gene is a de novo event. Am.

[137818]J. Med. Genet. 78: 313–316, 1998.

[137819]24552.

[137820]Sjlander, A.; Birgander, R.; Rannug, A.; Alexandrie, A.-K.;
Tornling,

[137821]G.; Beckman, G.: Association between the p21 codon
31A1 (arg) allele

[137822]and lung cancer. Hum. Hered. 46: 221-225, 1996.

[137823]24553.

[137824]Fryns, J. P.; van den Berghe, K.; van Assche, A.; van den
Berghe,

[137825]H.: Prenatal diagnosis of campomelic dwarfism. Clin.
Genet. 19:

[137826]199-201, 1981.

[137827]24554.

[137828]Mellows, H. J.; Pryse-Davies, J.; Bennett, M. J.; Carter, C. O.

[137829]: The campomelic syndrome in two female siblings. Clin.
Genet. 18:

[137830]137-141, 1980.

[137831]24555.

[137832]Bjursell, C.; Wahlstrom, J.; Berg, K.; Stibler, H.; Kristians-
son,

[137833]B.; Matthijs, G.; Martinsson, T.: Detailed mapping of the
phosphomannomutase

[137834]2 (PMM2) gene and mutation detection enable improved analysis for

[137835]Scandinavian CDG type I families. *Europ. J. Hum. Genet.* 6: 603–611,

[137836]1998.

[137837]24556.

[137838]Satoyoshi, E.; Kowa, H.: A myopathy due to glycolytic abnormality. *Arch. Neurol.* 17: 248–256, 1967.

[137840]24557.

[137841]Servidei, S.; Bonilla, E.; Diedrich, R. G.; Kornfeld, M.; Oates,

[137842]J. D.; Davidson, M.; Vora, S.; DiMauro, S.: Fatal infantile form

[137843]of muscle phosphofructokinase deficiency. *Neurology* 36: 1465–1470,

[137844]1986.

[137845]24558.

[137846]Tani, K.; Fujii, H.; Takegawa, S.; Miwa, S.; Koyama, W.; Kanayama,

[137847]M.; Imanaka, A.; Imanaka, F.; Kuramoto, A.: Two cases of

phosphofructokinase

[137848]deficiency associated with congenital hemolytic anemia
found in Japan. Am.

[137849]J. Hemat. 14: 165-174, 1983.

[137850]24559.

[137851]Tarui, S.; Okuno, G.; Ikura, Y.; Tanaka, T.; Suda, M.;
Nishikawa,

[137852]M.: Phosphofructokinase deficiency in skeletal muscle: a
new type

[137853]of glycogenosis. Biochem. Biophys. Res. Commun. 19:
517-523, 1965.

[137854]24560.

[137855]Tsuji, S.; Servidei, S.; Tonin, P.; Shanske, S.; Azan, G.;
DiMauro,

[137856]S.: Identification of three novel mutations in non-
Ashkenazi Italian

[137857]patients with muscle phosphofructokinase deficiency. Am.
J. Hum.

[137858]Genet. 54: 812-819, 1994.

[137859]24561.

[137860]Matsumoto, K.; Yoshitomi, H.; Rossant, J.; Zaret, K. S.:

Liver

[137861]organogenesis promoted by endothelial cells prior to vascular function. Science 294:

[137862]559–563, 2001.

[137863]24562.

[137864]Matthews, W.; Jordan, C. T.; Gavin, M.; Jenkins, N. A.; Copeland,

[137865]N. G.; Lemischka, I. R.: A receptor tyrosine kinase cDNA isolated

[137866]from a population of enriched primitive hematopoietic cells and exhibiting

[137867]close genetic linkage to c-kit. Proc. Nat. Acad. Sci. 88: 9026–9030,

[137868]1991.

[137869]24563.

[137870]Millauer, B.; Shawver, L. K.; Plate, K. H.; Risau, W.; Ullrich,

[137871]A.: Glioblastoma growth inhibited in vivo by a dominant-negative

[137872]Flk-1 mutant. Nature 367: 576–579, 1994.

[137873]24564.

[137874]Millauer, B.; Witzigmann-Voos, S.; Schnurch, H.; Martinez,

R.;

[137875]Moller, N. P. H.; Risau, W.; Ullrich, A.: High affinity VEGF binding

[137876]and developmental expression suggest Flk-1 as a major regulator of

[137877]vasculogenesis and angiogenesis. Cell 72: 835-846, 1993.

[137878]24565.

[137879]Plate, K. H.; Breier, G.; Millauer, B.; Ullrich, A.; Risau, W.

[137880]: Up-regulation of vascular endothelial growth factor and its cognate

[137881]receptors in a rat glioma model of tumor angiogenesis. Cancer Res. 53:

[137882]5822-5827, 1993.

[137883]24566.

[137884]Quinn, T. P.; Peters, K. G.; De Vries, C.; Ferrara, N.; Williams,

[137885]L. T.: Fetal liver kinase 1 is a receptor for vascular endothelial

[137886]growth factor and is selectively expressed in vascular endothelium. Proc.

[137887]Nat. Acad. Sci. 90: 7533-7537, 1993.

[137888]24567.

[137889]Sait, S. N. J.; Dougher-Vermazen, M.; Shows, T. B.; Ter-
man, B.

[137890]J.: The kinase insert domain receptor gene (KDR) has been
relocated

[137891]to chromosome 4q11-q12. Cytogenet. Cell Genet. 70:
145-146, 1995.

[137892]24568.

[137893]Shalaby, F.; Rossant, J.; Yamaguchi, T. P.; Gertsenstein, M.;

[137894]Wu, X.-F.; Breitman, M. L.; Schuh, A. C.: Failure of blood-
island

[137895]formation and vasculogenesis in Flk-1-deficient mice. Na-
ture 376:

[137896]62-65, 1995.

[137897]24569.

[137898]Shay-Salit, A.; Shushy, M.; Wolfowitz, E.; Yahav, H.; Bre-
viario,

[137899]F.; Dejana, E.; Resnick, N.: VEGF receptor 2 and the ad-
herens junction

[137900]as a mechanical transducer in vascular endothelial cells.
Proc. Nat.

[137901]Acad. Sci. 99: 9462–9467, 2002.

[137902]24570.

[137903]Spritz, R. A.; Strunk, K. M.; Lee, S.-T.; Lu-Kuo, J. M.; Ward,

[137904]D. C.; Le Paslier, D.; Altherr, M. R.; Dorman, T. E.; Moir, D.
T.

[137905]; A YAC contig spanning a cluster of human type III recep-
tor protein

[137906]tyrosine kinase genes (PDGFRA-KIT-KDR) in chromosome
segment 4q12. Genomics 22:

[137907]431–436, 1994.

[137908]24571.

[137909]Terman, B.; Carrion, M.; Eddy, R. L.; Shows, T. B.: Tyrosine

[137910]kinase growth factor receptor (KDR) localized to chromo-
some 4 (Abstract) Cytogenet.

[137911]Cell Genet. 58: 1890 only, 1991.

[137912]24572.

[137913]Terman, B. I.; Carrion, M. E.; Kovacs, E.; Rasmussen, B. A.;
Eddy,

[137914]R. L.; Shows, T. B.: Identification of a new endothelial cell
growth

[137915]factor receptor tyrosine kinase. Oncogene 6: 1677–1683,
1991.

[137916]24573.

[137917]Terman, B. I.; Jani-Sait, S.; Carrion, M. E.; Shows, T. B.: The

[137918]KDR gene maps to human chromosome 4q31.2–q32, a lo-
cus which is distinct

[137919]from locations for other type III growth factor receptor ty-
rosine

[137920]kinases. Cytogenet. Cell Genet. 60: 214–215, 1992.

[137921]24574.

[137922]Yamashita, J.; Itoh, H.; Hirashima, M.; Ogawa, M.;
Nishikawa,

[137923]S.; Yurugi, T.; Naito, M.; Nakao, K.; Nishikawa, S.–I.:
Flk1–positive

[137924]cells derived from embryonic stem cells serve as vascular
progenitors. Nature 408:

[137925]92–96, 2000.

[137926]24575.

[137927]Ziegler, B. L.; Valtieri, M.; Porada, G. A.; De Maria, R.;
Muller,

[137928]R.; Masella, B.; Gabbianelli, M.; Casella, I.; Pelosi, E.; Bock,

T.;

[137929]Zanjani, E. D.; Peschle, C.: KDR receptor: a key marker
defining

[137930]hematopoietic stem cells. Science 285: 1553–1558, 1999.

[137931]24576.

[137932]Ala-Mello, S.; Sankila, E.-M.; Koskimies, O.; de la
Chapelle, A.;

[137933]Kaariainen, H.: Molecular studies in Finnish patients with
familial

[137934]juvenile nephronophthisis exclude a founder effect and
support a common

[137935]mutation causing mechanism. J. Med. Genet. 35:
279–283, 1998.

[137936]24577.

[137937]Alexander, F.; Campbell, S.: Familial uremic medullary
cystic

[137938]disease. Pediatrics 45: 1024–1028, 1970.

[137939]24578.

[137940]Antignac, C.; Arduy, C. H.; Beckmann, J. S.; Benessy, F.;

Gros,

[137941]F.; Medhioub, M.; Hildebrandt, F.; Dufier, J.-L.;

Kleinknecht, C.;

[137942]Broyer, M.; Weissenbach, J.; Habib, R.; Cohen, D.: A gene
for familial

[137943]juvenile nephronophthisis (recessive medullary cystic kidney disease)

[137944]maps to chromosome 2p. Nature Genet. 3: 342-345,
1993.

[137945]24579.

[137946]Benzing, T.; Gerke, P.; Hopker, K.; Hildebrandt, F.; Kim, E.;
Walz,

[137947]G.: Nephrocystin interacts with Pyk2, p130(Cas), and
tensin and triggers

[137948]phosphorylation of Pyk2. Proc. Nat. Acad. Sci. 98:
9784-9789, 2001.

[137949]24580.

[137950]Betz, R.; Rensing, C.; Otto, E.; Mincheva, A.; Zehnder, D.;
Lichter,

[137951]P.; Hildebrandt, F.: Children with ocular motor apraxia
type Cogan

[137952]carry deletions in the gene (NPHP1) for juvenile
nephronophthisis. J.

[137953]Pediat. 136: 828-831, 2000.

[137954]24581.

[137955]Broberger, O.; Winberg, J.; Zetterstrom, R.: Juvenile nephronophthisis.

[137956]I. A genetically determined nephropathy with hypotonic polyuria and

[137957]azotaemia. Acta Paediat. 49: 470–479, 1960.

[137958]24582.

[137959]Caridi, G.; Murer, L.; Bellantuono, R.; Sorino, P.; Caringella,

[137960]D. A.; Gusmano, R.; Ghiggeri, G. M.: Renal–retinal syndromes: association

[137961]of retinal anomalies and recessive nephronophthisis in patients with

[137962]homozygous deletion of the NPH1 locus. Am. J. Kidney Dis. 32: 1059–1062,

[137963]1998.

[137964]24583.

[137965]Cohen, A. H.; Hoyer, J. R.: Nephronophthisis: a primary tubular

[137966]basement membrane defect. Lab. Invest. 55: 564–572, 1986.

[137967]24584.

[137968]Collan, Y.; Sipponen, P.; Haapanen, E.; Lindahl, J.; Jokinen,
[137969]E. J.; Hjelt, L.: Hereditary nephronophthisis with a life span
of
[137970]3 decades--light and electron microscopical, immunohis-
tochemical,
[137971]clinical and family studies. Wilhelm Roux. Arch. Develop.
Biol. 376:
[137972]195-208, 1977.
[137973]24585.
[137974]Donaldson, J. C.; Dempsey, P. J.; Reddy, S.; Bouton, A. H.;
Coffey,
[137975]R. J.; Hanks, S. K.: Crk-associated substrate p130(Cas) in-
teracts
[137976]with nephrocystin and both proteins localize to cell-cell
contacts
[137977]of polarized epithelial cells. Exp. Cell Res. 256: 168-175,
2000.
[137978]24586.
[137979]Fanconi, G.; Hanhart, E.; Von Albertini, A.; Uehlinger, E.;
Dolivo,
[137980]G.; Prader, A.: Die familiaere juvenile Nephronophthise.
(Die idiopathische

[137981]Parenchymatose). *Helv. Paediat. Acta* 6: 1–49, 1951.

[137982]24587.

[137983]Gibson, A. A. M.; Arneil, G. C.: Nephronophthisis: report of

[137984]8 cases in Britain. *Arch. Dis. Child.* 47: 84–89, 1972.

[137985]24588.

[137986]Giselson, N.; Heinegard, D.; Holmberg, C. G.; Lindberg, L.
G.;

[137987]Lindstedt, E.; Lindstedt, G.; Schersten, B.: Renal medullary
cystic

[137988]disease or familial juvenile nephronophthisis: a renal
tubular disease. *Am.*

[137989]*J. Med.* 48: 174–184, 1970.

[137990]24589.

[137991]Hackzell, G.; Lundmark, C.: Familial juvenile nephronoph-
thisis. *Acta*

[137992]*Paediat.* 47: 428–440, 1958.

[137993]24590.

[137994]Herdman, R. C.; Good, R. A.; Vernier, R. L.: Medullary cys-
tic

[137995]disease in two siblings. *Am. J. Med.* 43: 335–344, 1967.

[137996]24591.

[137997]Hildebrandt, F.: Personal Communication. Freiburg, Germany

[137998]10/1/1997.

[137999]24592.

[138000]Hildebrandt, F.; Otto, E.; Rensing, C.; Nothwang, H. G.; Vollmer,

[138001]M.; Adolphs, J.; Hanusch, H.; Brandis, M.: A novel gene encoding

[138002]an SH3 domain protein is mutated in nephronophthisis type 1. Nature

[138003]Genet. 17: 149–153, 1997.

[138004]24593.

[138005]Hildebrandt, F.; Singh–Sawhney, I.; Schnieders, B.; Papenfuss,

[138006]T.; Brandis, M.; Members of the APN Study Group: Refined genetic

[138007]mapping of a gene for familial juvenile nephronophthisis (NPH1) and

[138008]physical mapping of linked markers. Genomics 25: 360–364, 1995.

[138009]24594.

[138010]Khoury, Z.; Brezis, M.; Mogle, P.: Familial medullary
sponge

[138011]kidney in association with congenital absence of teeth
(anodontia). Nephron 48:

[138012]231–233, 1988.

[138013]24595.

[138014]Cho, K. R.; Oliner, J. D.; Simons, J. W.; Hedrick, L.; Fearon,

[138015]E. R.; Preisinger, A. C.; Hedge, P.; Silverman, G. A.; Vogel-
stein,

[138016]B.: The DCC gene: structural analysis and mutations in
colorectal

[138017]carcinomas. Genomics 19: 525–531, 1994.

[138018]24596.

[138019]Fazeli, A.; Dickinson, S. L.; Hermiston, M. L.; Tighe, R. V.;
Steen,

[138020]R. G.; Small, C. G.; Stoeckli, E. T.; Keino-Masu, K.; Masu,
M.; Rayburn,

[138021]H.; Simons, J.; Bronson, R. T.; Gordon, J. I.; Tessier-Lavi-
gne, M.;

[138022]Weinberg, R. A.: Phenotype of mice lacking functional

Deleted in

[138023]colorectal cancer (Dcc) gene. Nature 386: 796–804, 1997.

[138024]24597.

[138025]Fearon, E. R.; Cho, K. R.; Nigro, J. M.; Kern, S. E.; Simons, J.

[138026]W.; Ruppert, J. M.; Hamilton, S. R.; Preisinger, A. C.;

Thomas, G.;

[138027]Kinzler, K. W.; Vogelstein, B.: Identification of a chromo-
some 18q

[138028]gene that is altered in colorectal cancers. Science 247:
49–56,

[138029]1990.

[138030]24598.

[138031]Gotley, D. C.; Reeder, J. A.; Fawcett, J.; Walsh, M. D.; Bates,

[138032]P.; Simmons, D. L.; Antalis, T. M.: The deleted in colon
cancer (DCC)

[138033]gene is consistently expressed in colorectal cancers and
metastases. Oncogene 13:

[138034]787–795, 1996.

[138035]24599.

[138036]Hohne, M. W.; Halatsch, M.–E.; Kahl, G. F.; Weinel, R. J.:

Frequent

[138037]loss of expression of the potential tumor suppressor gene
DCC in ductal

[138038]pancreatic adenocarcinoma. Cancer Res. 52: 2616–2619,
1992.

[138039]24600.

[138040]Jen, J.; Kim, H.; Piantadosi, S.; Liu, Z.-F.; Levitt, R. C.; Sis-
tonen,

[138041]P.; Kinzler, K. W.; Vogelstein, B.; Hamilton, S. R.: Allelic
loss

[138042]of chromosome 18q and prognosis in colorectal cancer.
New Eng. J.

[138043]Med. 331: 213–221, 1994.

[138044]24601.

[138045]Keino-Masu, K.; Masu, M.; Hinck, L.; Leonardo, E. D.;
Chan, S.

[138046]S.-Y.; Culotti, J. G.; Tessier-Lavigne, M.: Deleted in col-
orectal

[138047]cancer (DCC) encodes a netrin receptor. Cell 87: 175–185,
1996.

[138048]24602.

[138049]Keynes, R.; Cook, G. M. W.: Axon guidance molecules. Cell

[138050]161–169, 1995.

[138051]24603.

[138052]Kolodziej, P. A.; Timpe, L. C.; Mitchell, K. J.; Fried, S. R.;

[138053]Goodman, C. S.; Jan, L. Y.; Jan, Y. N.: Frazzled encodes a
Drosophila

[138054]member of the DCC immunoglobulin subfamily and is re-
quired for CNS

[138055]and motor axon guidance. Cell 87: 197–204, 1996.

[138056]24604.

[138057]Maesawa, C.; Tamura, G.; Ogasawara, S.; Suzuki, Y.;
Sakata, K.;

[138058]Sugimura, J.; Nishizuka, S.; Sato, N.; Ishida, K.; Saito, K.;
Satodate,

[138059]R.: Loss of heterozygosity at the DCC gene locus is not
crucial for

[138060]the acquisition of metastatic potential in oesophageal
squamous cell

[138061]carcinoma. Europ. J. Cancer. 32A: 896–898, 1996.

[138062]24605.

[138063]Mehlen, P.; Rabizadeh, S.; Snipas, S. J.; Assa-Munt, N.;

Salvesen,

[138064]G. S.; Bredesen, D. E.: The DCC gene product induces apoptosis by

[138065]a mechanism requiring receptor proteolysis. Nature 395: 801–804,

[138066]1998.

[138067]24606.

[138068]Miyake, S.; Nagai, K.; Yoshino, K.; Oto, M.; Endo, M.; Yuasa,

[138069]Y.: Point mutations and allelic deletion of tumor suppressor gene

[138070]DCC in human esophageal squamous cell carcinomas and their relation

[138071]to metastasis. Cancer Res. 54: 3007–3010, 1994.

[138072]24607.

[138073]Nigro, J. M.; Cho, K. R.; Fearon, E. R.; Kern, S. E.; Ruppert,

[138074]J. M.; Oliner, J. D.; Kinzler, K. W.; Vogelstein, B.: Scrambled exons. Cell 64:

[138075]607–613, 1991.

[138076]24608.

[138077]Shibata, D.; Reale, M. A.; Lavin, P.; Silverman, M.; Fearon,

E.

[138078]R.; Steele, G., Jr.; Jessup, J. M.; Loda, M.; Summerhayes, I.

C.:

[138079]The DCC protein and prognosis in colorectal cancer. New
Eng. J. Med. 335:

[138080]1727-1732, 1996.

[138081]24609.

[138082]Stein, E.; Tessier-Lavigne, M.: Hierarchical organization of

[138083]guidance receptors: silencing of netrin attraction by Slit
through

[138084]a Robo/DCC receptor complex. Science 291: 1928-1938,
2001.

[138085]24610.

[138086]Tanaka, K.; Oshimura, M.; Kikuchi, R.; Seki, M.; Hayashi,
T.;

[138087]Miyaki, M.: Suppression of tumorigenicity in human colon
carcinoma

[138088]cells by introduction of normal chromosome 5 or 18. Na-
ture 349:

[138089]340-342, 1991.

[138090]24611.

[138091]Uchino, S.; Tsuda, H.; Noguchi, M.; Yokota, J.; Terada, M.;
Saito,
[138092]T.; Kobayashi, M.; Sugimura, T.; Hirohashi, S.: Frequent
loss of
[138093]heterozygosity at the DCC locus in gastric cancer. Cancer
Res. 52:
[138094]3099–3102, 1992.
[138095]24612.
[138096]Vogelstein, B.: Personal Communication. Baltimore, Md.
11/30/1995.
[138097]24613.
[138098]Zetter, B. R.: Adhesion molecules in tumor metastasis.
Semin.
[138099]Cancer Biol. 4: 219–229, 1993.
[138100]24614.
[138101]Devor, E. J.; Grandy, D. K.; Civelli, O.; Litt, M.; Burgess, A.
[138102]K.; Isenberg, K. E.; van de Wetering, B. J. M.; Oostra, B.:
Genetic
[138103]linkage is excluded for the D2–dopamine receptor
lambda–HD2G1 and
[138104]flanking loci on chromosome 11q22–q23 in Tourette syn–

drome. Hum.

[138105]Hered. 40: 105–108, 1990.

[138106]24615.

[138107]Voutsinas, S.; Wynne–Davies, R.: The infrequency of malignant

[138108]disease in diaphyseal aclasis and neurofibromatosis. J. Med. Genet. 20:

[138109]345–349, 1983.

[138110]24616.

[138111]Freemantle, S. J.; Taylor, S. M.; Krystal, G.; Moran, R. G.: Upstream

[138112]organization of and multiple transcripts from the human folypoly–gamma–glutamate

[138113]synthetase gene. J. Biol. Chem. 270: 9579–9584, 1995.

[138114]24617.

[138115]Garrow, T. A.; Admon, A.; Shane, B.: Expression cloning of a human

[138116]cDNA encoding folypoly(gamma–glutamate) synthetase and determination

[138117]of its primary structure. Proc. Nat. Acad. Sci. 89: 9151–9155, 1992.

[138118]24618.

[138119]Jones, C.; Kao, F.-T.: Regional mapping of the folylpolyglutamate

[138120]synthetase gene (FPGS) to 9cen-q34. (Abstract) Cytogenet. Cell Genet. 37:

[138121]499-500, 1984.

[138122]24619.

[138123]Jones, C.; Kao, F.-T.; Taylor, R. T.: Chromosomal assignment of

[138124]the gene for folylpolyglutamate synthetase to human chromosome 9. Cytogenet.

[138125]Cell Genet. 28: 181-194, 1980.

[138126]24620.

[138127]Jones, C.; Kao, F. T.: Assignment of the human gene complementing

[138128]the auxotrophic marker GAT-minus to chromosome 9. (Abstract) Cytogenet.

[138129]Cell Genet. 25: 168, 1979.

[138130]24621.

[138131]Kao, F. T.; Puck, T. T.: Genetics of somatic mammalian cells.

[138132]VII. Induction and isolation of nutritional mutants in Chinese hamster

[138133]cells. Proc. Nat. Acad. Sci. 60: 1275–1281, 1968.

[138134]24622.

[138135]Sussman, D. J.; Milman, G.; Shane, B.: Characterization of human

[138136]folylpolyglutamate synthetase expressed in Chinese hamster ovary cells. Somat.

[138137]Cell Molec. Genet. 12: 531–540, 1986.

[138138]24623.

[138139]Taylor, R. T.; Hanna, M. L.: Folate-dependent enzymes in cultured

[138140]Chinese hamster cells: folylpolyglutamate synthetase and its absence

[138141]in mutants auxotrophic for glycine, adenosine and thymidine. Arch.

[138142]Biochem. Biophys. 181: 331–344, 1977.

[138143]24624.

[138144]Taylor, S. M.; Freemantle, S. J.; Moran, R. G.: Structural organization

[138145]of the human folylpoly- γ -glutamate synthetase

gene: evidence for

[138146]a single genomic locus. Cancer Res. 55: 6030–6034,
1995.

[138147]24625.

[138148]Buters, J. T. M.; Tang, B.-K.; Pineau, T.; Gelboin, H. V.;
Kimura,

[138149]S.; Gonzalez, F. J.: Role of CYP1A2 in caffeine pharmacoki-
netics

[138150]and metabolism: studies using mice deficient in CYP1A2.
Pharmacogenetics 6:

[138151]291–296, 1996.

[138152]24626.

[138153]Butler, M. A.; Iwasaki, M.; Guengerich, F. P.; Kadlubar, F. F.

[138154]: Human cytochrome P-450(PA) (P-450IA2), the
phenacetin O-deethylase,

[138155]is primarily responsible for the hepatic 3-demethylation
of caffeine

[138156]and N-oxidation of carcinogenic arylamines. Proc. Nat.
Acad. Sci. 86:

[138157]7696–7700, 1989.

[138158]24627.

[138159]Christiansen, L.; Bygum, A.; Jensen, A.; Thomsen, K.;
Brandrup,
[138160]F.; Horder, M.; Petersen, N. E.: Association between
CYP1A2 polymorphism
[138161]and susceptibility to porphyria cutanea tarda. Hum. Genet.
107:
[138162]612–614, 2000.
[138163]24628.
[138164]Devonshire, H. W.; Kong, I.; Cooper, M.; Sloan, T. P.; Idle,
J.
[138165]R.; Smith, R. L.: The contribution of genetically determined
oxidation
[138166]status to inter-individual variation in phenacetin disposi-
tion. Brit.
[138167]J. Clin. Pharm. 16: 157–166, 1983.
[138168]24629.
[138169]Ikeya, K.; Jaiswal, A. K.; Owens, R. A.; Jones, J. E.; Nebert,
[138170]D. W.; Kimura, S.: Human CYP1A2: sequence, gene struc-
ture, comparison
[138171]with the mouse and rat orthologous gene, and differences
in liver

- [138172]1A2 mRNA expression. Molec. Endocr. 3: 1399–1408, 1989.
- [138173]24630.
- [138174]Jaiswal, A. K.; Nebert, D. W.; McBride, O. W.; Gonzalez, F. J.
- [138175]: Human P(3)450: cDNA and complete protein sequence, repetitive Alu
- [138176]sequences in the 3–prime nontranslated region, and localization of
- [138177]gene to chromosome 15. J. Exp. Path. 3: 1–17, 1987.
- [138178]24631.
- [138179]Liang, H.–C. L.; Li, H.; McKinnon, R. A.; Duffy, J. J.; Potter,
- [138180]S. S.; Puga, A.; Nebert, D. W.: Cyp1a2(–/–) null mutant mice develop
- [138181]normally but show deficient drug metabolism. Proc. Nat. Acad. Sci. 93:
- [138182]1671–1676, 1996.
- [138183]24632.
- [138184]Nebert, D. W.: Personal Communication. Bethesda, Md. 2/3/1988.
- [138185]24633.

[138186]Sesardic, D.; Boobis, A. R.; Edwards, R. J.; Davies, D. S.: A

[138187]form of cytochrome P450 in man, orthologous to form d
in the rat,

[138188]catalyses the O-deethylation of phenacetin and is in-
ducible by cigarette

[138189]smoking. Brit. J. Clin. Pharm. 26: 363–372, 1988.

[138190]24634.

[138191]Tantcheva-Poor, I.; Zaigler, M.; Rietbrock, S.; Fuhr, U.: Es-
timation

[138192]of cytochrome P-450 CYP1A2 activity in 863 healthy Cau-
casians using

[138193]a saliva-based caffeine test. Pharmacogenetics 9:
131–144, 1999.

[138194]24635.

[138195]Wooding, S. P.; Watkins, W. S.; Bamshad, M. J.; Dunn, D.
M.; Weiss,

[138196]R. B.; Jorde, L. B.: DNA sequence variation in a 3.7-kb
noncoding

[138197]sequence 5-prime of the CYP1A2 gene: implications for
human population

[138198]history and natural selection. Am. J. Hum. Genet. 71:
528–542, 2002.

[138199]24636.

[138200]Bale, A. E.; Mitchell, A. L.; Gonzalez, F. J.; McBride, O. W.:

[138201]Localization of CYP2F1 by multipoint linkage analysis and
pulsed-field

[138202]gel electrophoresis. Genomics 10: 284-286, 1991.

[138203]24637.

[138204]Mitchell, A. L.; Bale, A. E.; Gonzalez, F.; McBride, O. W.:
Mapping

[138205]of cytochrome p450 IIA, IIB, and IIF subfamilies on chro-
mosome 19

[138206]by linkage analysis. (Abstract) Cytogenet. Cell Genet. 51:
1045,

[138207]1989.

[138208]24638.

[138209]Nhamburo, P. T.; Kimura, S.; McBride, O. W.; Kozak, C. A.;
Gelboin,

[138210]H. V.; Gonzalez, F. J.: The human CYP2F gene subfamily:
identification

[138211]of a cDNA encoding a new cytochrome P450, cDNA-di-
rected expression,

[138212]and chromosome mapping. Biochemistry 29: 5491-5499,

1990.

[138213]24639.

[138214]Thum, T.; Borlak, J.: Gene expression in distinct regions of
the

[138215]heart. Lancet 355: 979–983, 2000.

[138216]24640.

[138217]Wells, D. E.; Hill, A.; Lin, X.; Ahn, J.; Brown, N.; Wagner, M.

[138218]J.: Identification of novel mutations in the human EXT1 tu-
mor suppressor

[138219]gene. Hum. Genet. 99: 612–615, 1997.

[138220]24641.

[138221]Wicklund, C. L.; Pauli, R. M.; Johnston, D.; Hecht, J. T.:

Natural

[138222]history study of hereditary multiple exostoses. Am. J. Med.
Genet. 55:

[138223]43–46, 1995.

[138224]24642.

[138225]Wuyts, W.; Van Hul, W.: Molecular basis of multiple exos-
toses:

[138226]mutations in the EXT1 and EXT2 genes. Hum. Mutat. 15:

220–227, 2000.

[138227]24643.

[138228]Wuyts, W.; Van Hul, W.; De Boulle, K.; Hendrickx, J.;
Bakker,

[138229]E.; Vanhoenacker, F.; Mollica, F.; Ludecke, H.-J.; Sayli, B.
S.; Pazzaglia,

[138230]U. E.; Mortier, G.; Hamel, B.; Conrad, E. U.; Matsushita, M.;
Raskind,

[138231]W. H.; Willems, P. J.: Mutations in the EXT1 and EXT2
genes in hereditary

[138232]multiple exostoses. Am. J. Hum. Genet. 62: 346–354,
1998.

[138233]24644.

[138234]Xu, L.; Xia, J.; Jiang, H.; Zhou, J.; Li, H.; Wang, D.; Pan, Q.;

[138235]Long, Z.; Fan, C.; Deng, H.-X.: Mutation analysis of heredit-
itary multiple

[138236]exostoses in the Chinese. Hum. Genet. 105: 45–50, 1999.

[138237]24645.

[138238]Yoshiura, K.; Inazawa, J.; Koyama, K.; Nakamura, Y.; Ni-
ikawa,

[138239]N.: Mapping of the 8q23 translocation breakpoint of

t(8;13) observed

[138240]in a patient with multiple exostoses. Genes Chromosomes
Cancer 9:

[138241]57–61, 1994.

[138242]24646.

[138243]Bellaïche, Y.; The, I.; Perrimon, N.: Tout-velu is a
Drosophila

[138244]homologue of the putative tumour suppressor EXT-1 and
is needed for

[138245]Hh diffusion. Nature 394: 85–88, 1998.

[138246]24647.

[138247]Clines, G. A.; Ashley, J. A.; Shah, S.; Lovett, M.: The struc-
ture

[138248]of the human multiple exostoses 2 gene and characteriza-
tion of homologs

[138249]in mouse and Caenorhabditis elegans. Genome Res. 7:
359–367, 1997.

[138250]24648.

[138251]Francannet, C.; Cohen-Tanugi, A.; Le Merrer, M.; Munnich,
A.; Bonaventure,

[138252]J.; Legeai-Mallet, L.: Genotype–phenotype correlation in

hereditary

[138253]multiple exostoses. J. Med. Genet. 38: 430–434, 2001.

[138254]24649.

[138255]Ligon, A. H.; Potocki, L.; Shaffer, L. G.; Stickens, D.; Evans,

[138256]G. A.: Gene for multiple exostoses (EXT2) maps to

11(p11.2p12) and

[138257]is deleted in patients with a contiguous gene syndrome.

(Letter) Am.

[138258]J. Med. Genet. 75: 538–540, 1998.

[138259]24650.

[138260]Lind, T.; Tufaro, F.; McCormick, C.; Lindahl, U.; Lidholt, K.

[138261]: The putative tumor suppressors EXT1 and EXT2 are gly-

cosyltransferases

[138262]required for the biosynthesis of heparan sulfate. J. Biol.

Chem. 273:

[138263]26265–26268, 1998.

[138264]24651.

[138265]McGaughan, J. M.; Ward, H. B.; Evans, D. G. R.: WAGR

syndrome

[138266]and multiple exostoses in a patient with

del(11)(p11.2p14.2). J.

[138267]Med. Genet. 32: 823–824, 1995.

[138268]24652.

[138269]Potocki, L.; Greenberg, F.; Shaffer, L. G.: Interstitial deletion

[138270]of 11(p12p11.2): a rare chromosomal syndrome with mental retardation,

[138271]parietal foramina, and multiple exostoses. (Abstract) Am. J. Hum.

[138272]Genet. 57: A123, 1995.

[138273]24653.

[138274]Potocki, L.; Shaffer, L. G.: Interstitial deletion of 11(p11.2p12):

[138275]a newly described contiguous gene deletion syndrome involving the

[138276]gene for hereditary multiple exostoses (EXT2). Am. J. Med. Genet. 62:

[138277]319–325, 1996.

[138278]24654.

[138279]Quirini, G. E.; Meyer, J. R.; Herman, M.; Russell, E. J.: Osteochondroma

[138280]of the thoracic spine: an unusual cause of spinal cord compression. Am.

[138281]J. Neuroradiol. 17: 961-964, 1996.

[138282]24655.

[138283]Simmons, A. D.; Musy, M. M.; Lopes, C. S.; Hwang, L.-Y.; Yang,

[138284]Y.-P.; Lovett, M.: A direct interaction between EXT proteins and

[138285]glycosyltransferases is defective in hereditary multiple exostoses. Hum.

[138286]Molec. Genet. 8: 2155-2164, 1999.

[138287]24656.

[138288]Stickens, D.; Clines, G.; Burbee, D.; Ramos, P.; Thomas, S.; Hogue,

[138289]D.; Hecht, J. T.; Lovett, M.; Evans, G. A.: The EXT2 multiple exostoses

[138290]gene defines a family of putative tumour suppressor genes. Nature

[138291]Genet. 14: 25-32, 1996.

[138292]24657.

[138293]Stickens, D.; Evans, G. A.: Isolation and characterization of

- [138294]the murine homolog of the human EXT2 multiple exostoses gene. *Biochem. Molec. Med.* 61: 16–21, 1997.
- [138295]24658.
- [138296]24658.
- [138297]Wu, Y.-Q.; Heutink, P.; de Vries, B. B. A.; Sandkuijl, L. A.; van den Ouweland, A. M. W.; Niermeijer, M. F.; Galjaard, H.; Reyniers, E.; Willems, P. J.; Halley, D. J. J.: Assignment of a second locus for multiple exostoses to the pericentromeric region of chromosome 11. *Hum. Molec. Genet.* 3: 167–171, 1994.
- [138298]24659.
- [138299]Joutel, A.; Favrole, P.; Labauge, P.; Chabriat, H.; Lescoat, C.; Andreux, F.; Domenga, V.; Cecillon, M.; Vahedi, K.; Ducros, A.; Cave-Riant, F.; Bousser, M. G.; Tournier-Lasserre, E.: Skin biopsy immunostaining with a Notch3 monoclonal antibody for CADASIL diagnosis. *Lancet* 358:

[138307]2049–2051, 2001.

[138308]24660.

[138309]Aoki, M.; Abe, K.; Kameya, T.; Watanabe, M.; Itoyama, Y.:
Maternal

[138310]anticipation of DRPLA. Hum. Molec. Genet. 3: 1197–1198,
1994.

[138311]24661.

[138312]Barinaga, M.: An intriguing new lead on Huntington's dis-
ease. Science 271:

[138313]1233–1234, 1996.

[138314]24662.

[138315]Burke, J. R.; Enghild, J. J.; Martin, M. E.; Jou, Y.–S.; Myers,

[138316]R. M.; Roses, A. D.; Vance, J. M.; Strittmatter, W. J.: Hunt-
ingtin

[138317]and DRPLA proteins selectively interact with the enzyme
GAPDH. Nature

[138318]Med. 2: 347–350, 1996.

[138319]24663.

[138320]Burke, J. R.; Wingfield, M. S.; Lewis, K. E.; Roses, A. D.;
Lee,

[138321]J. E.; Hulette, C.; Pericak-Vance, M. A.; Vance, J. M.: The
Haw River

[138322]syndrome: dentatorubropallidoluysian atrophy (DRPLA) in
an African-American

[138323]family. Nature Genet. 7: 521-524, 1994.

[138324]24664.

[138325]Cancel, G.; Durr, A.; Stevanin, G.; Chneiweiss, H.; Duyck-
aerts,

[138326]C.; Serdaru, M.; de Toffol, B.; Agid, Y.; Brice, A.: Is DRPLA
also

[138327]linked to 14q? (Letter) Nature Genet. 6: 8, 1994.

[138328]24665.

[138329]Connarty, M.; Dennis, N. R.; Patch, C.; Macpherson, J. N.;
Harvey,

[138330]J. F.: Molecular re-investigation of patients with Hunting-
ton's disease

[138331]in Wessex reveals a family with dentatorubral and palli-
doluysian atrophy. Hum.

[138332]Genet. 97: 76-78, 1996.

[138333]24666.

[138334]Dubourg, O.; Durr, A.; Chneiweiss, H.; Stevanin, G.; Can-

cel, G.;

[138335]Penet, C.; Agid, Y.; Brice, A.: La forme ataxo-choreique de
DRPLA

[138336]existe-t-elle en Europe? Recherche de la mutation dans
120 familles. Rev.

[138337]Neurol. (Paris) 151: 657-660, 1995.

[138338]24667.

[138339]Hayashi, Y.; Kakita, A.; Yamada, M.; Koide, R.; Igarashi, S.;
Takano,

[138340]H.; Ikeuchi, T.; Wakabayashi, K.; Egawa, S.; Tsuji, S.; Taka-
hashi,

[138341]H.: Hereditary dentatorubral-pallidoluysian atrophy: de-
tection of

[138342]widespread ubiquitinated neuronal and glial intranuclear
inclusions

[138343]in the brain. Acta Neuropath. 96: 547-552, 1998.

[138344]24668.

[138345]Hirayama, K.; Iizuka, R.; Maehara, K.; Watanabe, T.: Clini-
copathological

[138346]study of dentatorubro-pallidoluysian atrophy. Part 1. Its
clinical

[138347]form and analysis of symptomatology. Shinkei Kenkyu no

Shinpo 25:

[138348]725–736, 1981.

[138349]24669.

[138350]Hirayama, K.; Takayanagi, T.; Nakamura, R.; Yanagisawa, N.; Hattori,

[138351]T.; Kita, K.; Yanagimoto, S.; Fujita, M.; Nagaoka, M.; Sato-mura, Y.;

[138352]Sobue, I.; Iizuka, R.; Toyokura, Y.; Satoyoshi, E.:
Spinocerebellar

[138353]degenerations in Japan: a nationwide epidemiological and
clinical

[138354]study. *Acta Neurol. Scand.* 89 (suppl. 153): 1–22, 1994.

[138355]24670.

[138356]Iizuka, R.; Hirayama, K.; Maehara, K.: Dentatorubro-palli-doluysian

[138357]atrophy: a clinico-pathological study. *J. Neurol. Neuro-surg. Psychiat.* 47:

[138358]1288–1298, 1984.

[138359]24671.

[138360]Ikeuchi, T.; Igarashi, S.; Takiyama, Y.; Onodera, O.; Oyake, M.;

[138361]Takano, H.; Koide, R.; Tanaka, H.; Tsuji, S.: Non-mendelian transmission

[138362]in dentatorubral-pallidoluysian atrophy and Machado-Joseph disease:

[138363]the mutant allele is preferentially transmitted in male meiosis. Am.

[138364]J. Hum. Genet. 58: 730-733, 1996.

[138365]24672.

[138366]Iwabuchi, K.; Amano, N.; Yokoi, S.; Nakano, T.; Yagishita, S.

[138367]: Two familial cases of dentato-rubro-pallidoluysian atrophy with

[138368]pseudo-Huntington's chorea (in Japanese). Clin. Neurol.

25: 1052-1060,

[138369]1985.

[138370]24673.

[138371]Eubanks, J. H.; Djabali, M.; Selleri, L.; Grandy, D. K.; Civelli,

[138372]O.; McElligott, D. L.; Evans, G. A.: Structure and linkage of the

[138373]D2 dopamine receptor and neural cell adhesion molecule genes on human

[138374]chromosome 11q23. Genomics 14: 1010-1018, 1992.

[138375]24674.

[138376]Gejman, P. V.; Ram, A.; Gelernter, J.; Friedman, E.; Cao, Q.;

[138377]Pickar, D.; Blum, K.; Noble, E. P.; Kranzler, H. R.; O'Malley, S.;

[138378]Hamer, D. H.; Whitsitt, F.; Rao, P.; DeLisi, L. E.; Virkkunen, M.;

[138379]Linnoila, M.; Goldman, D.; Gershon, E. S.: No structural mutation

[138380]in the dopamine D2 receptor gene in alcoholism or schizophrenia: analysis

[138381]using denaturing gradient gel electrophoresis. J.A.M.A. 271: 204-208,

[138382]1994.

[138383]24675.

[138384]Gelernter, J.; Grandy, D. K.; Bunzow, J.; Pakstis, A. J.; Civelli,

[138385]O.; Retief, A. E.; Litt, M.; Kidd, K. K.: D(2) dopamine receptor

[138386]locus (probe hD2G1) maps close to D11S29 (probe L7) and is also linked

[138387]to PBGD (probe PBGD0.0) and D11S84 (probe p-2-7-ID6)
on 11q. (Abstract) Cytogenet.

[138388]Cell Genet. 51: 1002 only, 1989.

[138389]24676.

[138390]Gelernter, J.; Pakstis, A. J.; Grandy, D.; Litt, M.; Retief, A.

[138391]E.; Kennedy, J. L.; Hing-Loh, A.; Schoolfield, G.; Civelli, O.;
Kidd,

[138392]K. K.: Linkage map of eight human chromosome 11q
markers, including

[138393]DRD2, spanning 60 cM. Cytogenet. Cell Genet. 60: 26-28,
1992.

[138394]24677.

[138395]Gelernter, J.; Pakstis, A. J.; Pauls, D. L.; Kurlan, R.;
Gancher,

[138396]S. T.; Civelli, O.; Grandy, D.; Kidd, K. K.: Gilles de la
Tourette

[138397]syndrome is not linked to D2-dopamine receptor. Arch.
Gen. Psychiat. 47:

[138398]1073-1077, 1990.

[138399]24678.

[138400]Giros, B.; Sokoloff, P.; Martres, M.-P.; Riou, J.-F.; Emorine,

[138401]L. J.; Schwartz, J.-C.: Alternative splicing directs the expression

[138402]of two D(2) dopamine receptor isoforms. Nature 342: 923–926, 1989.

[138403]24679.

[138404]Grandy, D. K.; Litt, M.; Allen, L.; Bunzow, J. R.; Marchionni,

[138405]M.; Makam, H.; Reed, L.; Magenis, R. E.; Civelli, O.: The human dopamine

[138406]D(2) receptor gene is located on chromosome 11 at q22–q23 and identifies

[138407]a TaqI RFLP. Am. J. Hum. Genet. 45: 778–785, 1989.

[138408]24680.

[138409]Grandy, D. K.; Marchionni, M. A.; Makam, H.; Stofko, R. E.; Alfano,

[138410]M.; Frothingham, L.; Fischer, J. B.; Burke–Howie, K. J.; Bunzow, J.

[138411]R.; Server, A. C.; Civelli, O.: Cloning of the cDNA and gene for

[138412]a human D2 dopamine receptor. Proc. Nat. Acad. Sci. 86: 9762–9766,

[138413]1989.

[138414]24681.

[138415]Klein, C.; Brin, M. F.; Kramer, P.; Sena-Esteves, M.; de
Leon,

[138416]D.; Doheny, D.; Bressman, S.; Fahn, S.; Breakefield, X. O.;
Ozelius,

[138417]L. J.: Association of a missense change in the D2
dopamine receptor

[138418]with myoclonus dystonia. Proc. Nat. Acad. Sci. 96:
5173–5176, 1999.

[138419]24682.

[138420]Maldonado, R.; Salardi, A.; Valverde, O.; Samad, T. A.;
Roques,

[138421]B. P.; Borrelli, E.: Absence of opiate rewarding effects in
mice

[138422]lacking dopamine D2 receptors. Nature 388: 586–589,
1997.

[138423]24683.

[138424]Milligan, G.: Receptors as kissing cousins. Science 288:
65–67,

[138425]2000.

[138426]24684.

[138427]Monsma, F. J., Jr.; McVittie, L. D.; Gerfen, C. R.; Mahan, L.
 [138428]C.; Sibley, D. R.: Multiple D(2) dopamine receptors pro-
 duced by alternative
 [138429]RNA splicing. *Nature* 342: 926–929, 1989.
 [138430]24685.
 [138431]Oliveri, R. L.; Annesi, G.; Zappia, M.; Civitelli, D.; Monte-
 santi,
 [138432]R.; Branca, D.; Nicoletti, G.; Spadafora, P.; Pasqua, A. A.;
 Cittadella,
 [138433]R.; Andreoli, V.; Gambardella, A.; Aguglia, U.; Quattrone,
 A.: Dopamine
 [138434]D2 receptor gene polymorphism and the risk of levodopa-
 induced dyskinesias
 [138435]in *PD. Neurology* 53: 1425–1430, 1999.
 [138436]24686.
 [138437]Rocheville, M.; Lange, D. C.; Kumar, U.; Patel, S. C.; Patel,
 [138438]R. C.; Patel, Y. C.: Receptors for dopamine and somato-
 statin: formation
 [138439]of hetero-oligomers with enhanced functional activity.
Science 288:
 [138440]154–157, 2000.

[138441]24687.

[138442]Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.;
Grandy,

[138443]D.; Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an
11q22.3-9p22

[138444]translocation with affective disorder: proximity of the
dopamine D2

[138445]receptor gene relative to the translocation breakpoint.
(Abstract) Am.

[138446]J. Hum. Genet. 45 (suppl.): A220 only, 1989.

[138447]24688.

[138448]St Clair, D.; Blackwood, D.; Muir, W.; Carothers, A.; Walker,

[138449]M.; Spowart, G.; Gosden, C.; Evans, H. J.: Association
within a family

[138450]of a balanced autosomal translocation with major mental
illness. Lancet 336:

[138451]13-16, 1990.

[138452]24689.

[138453]Suarez, B. K.; Parsian, A.; Hampe, C. L.; Todd, R. D.; Reich,

[138454]T.; Cloninger, C. R.: Linkage disequilibria at the D2
dopamine receptor

[138455]locus (DRD2) in alcoholics and controls. Genomics 19:
12–20, 1994.

[138456]24690.

[138457]Usiello, A.; Baik, J.–H.; Rouge–Pont, F.; Picetti, R.; Dierich,

[138458]A.; LeMeur, M.; Piazza, P. V.; Borrelli, E.: Distinct functions
of

[138459]the two isoforms of dopamine D2 receptors. Nature 408:
199–203,

[138460]2000.

[138461]24691.

[138462]Wang, J.; Liu, Z.–L.; Chen, B.: Association study of
dopamine

[138463]D2, D3 receptor gene polymorphisms with motor fluctua-
tions in PD. Neurology 56:

[138464]1757–1759, 2001.

[138465]24692.

[138466]Accili, D.; Fishburn, C. S.; Drago, J.; Steiner, H.; Lachowicz,

[138467]J. E.; Park, B.–H.; Gauda, E. B.; Lee, E. J.; Cool, M. H.; Sib-
ley,

[138468]D. R.; Gerfen, C. R.; Westphal, H.; Fuchs, S.: A targeted
mutation

[138469]of the D3 dopamine receptor gene is associated with hyperactivity

[138470]in mice. Proc. Nat. Acad. Sci. 93: 1945–1949, 1996.

[138471]24693.

[138472]Asico, L. D.; Ladines, C.; Fuchs, S.; Accili, D.; Carey, R. M.;

[138473]Semeraro, C.; Pocchiari, F.; Felder, R. A.; Eisner, G. M.; Jose, P.

[138474]A.: Disruption of the dopamine D3 receptor gene produces renin-dependent

[138475]hypertension. J. Clin. Invest. 102: 493–498, 1998.

[138476]24694.

[138477]Crocq, M.–A.; Mant, R.; Asherson, P.; Williams, J.; Hode, Y.; Mayerova,

[138478]A.; Collier, D.; Lannfelt, L.; Sokoloff, P.; Schwartz, J.–C.; Gill,

[138479]M.; Macher, J.–P.; McGuffin, P.; Owen, M. J.: Association between

[138480]schizophrenia and homozygosity at the dopamine D3 receptor gene. J.

[138481]Med. Genet. 29: 858–860, 1992.

[138482]24695.

- [138483]Guillin, O.; Diaz, J.; Carroll, P.; Griffon, N.; Schwartz, J.-C.;
- [138484]Sokoloff, P.: BDNF controls dopamine D3 receptor expression and triggers
- [138485]behavioural sensitization. *Nature* 411: 86–89, 2001.
- [138486]24696.
- [138487]Ilani, T.; Ben-Shachar, D.; Strous, R. D.; Mazor, M.; Sheinkman,
- [138488]A.; Kotler, M.; Fuchs, S.: A peripheral marker for schizophrenia:
- [138489]increased levels of D3 dopamine receptor mRNA in blood lymphocytes. *Proc.*
- [138490]*Nat. Acad. Sci.* 98: 625–628, 2001.
- [138491]24697.
- [138492]Le Coniat, M.; Sokoloff, P.; Hillion, J.; Martres, M.-P.; Giros,
- [138493]B.; Pilon, C.; Schwartz, J.-C.; Berger, R.: Chromosomal localization
- [138494]of the human D-3 dopamine receptor gene. *Hum. Genet.* 87: 618–620,
- [138495]1991.
- [138496]24698.

- [138497]Nanko, S.; Sasaki, T.; Fukuda, R.; Hattori, M.; Dai, X. Y.; Kazamatsuri, H.; Kuwata, S.; Juji, T.; Gill, M.: A study of the association between schizophrenia and the dopamine D3 receptor gene. Hum. Genet. 92: 336–338, 1993.
- [138501]24699.
- [138502]Nothen, M. M.; Cichon, S.; Propping, P.; Fimmers, R.; Schwab, S. G.; Wildenauer, D. B.: Excess of homozygosity at the dopamine D3 receptor gene in schizophrenia not confirmed. (Letter) J. Med. Genet. 30: 708–712, 1993.
- [138506]24700.
- [138507]Bevilacqua, M.; Butcher, E.; Furie, B.; Furie, B.; Gallatin, M.; Gimbrone, M.; Harlan, J.; Kishimoto, K.; Lasky, L.; McEver, R.; Paulson, J.; Rosen, S.; Seed, B.; Siegelman, M.; Springer, T.; Stoolman, L.;

- [138510] Tedder, T.; Varki, A.; Wagner, D.; Weissman, I.; Zimmerman, G.: Selectins:
[138511] a family of adhesion receptors. (Letter) Cell 67: 233 only,
1991.
[138512] 24701.
- [138513] Bevilacqua, M. P.; Stengelin, S.; Gimbrone, M. A., Jr.; Seed,
B.
[138514]: Endothelial leukocyte adhesion molecule 1: an inducible
receptor
[138515] for neutrophils related to complement regulatory proteins
and lectins. Science 243:
[138516] 1160–1165, 1989.
[138517] 24702.
- [138518] Collins, T.; Williams, A.; Johnston, G. I.; Kim, J.; Eddy, R.;
[138519] Shows, T.; Gimbrone, M. A., Jr.; Bevilacqua, M. P.: Struc-
ture and
[138520] chromosomal location of the gene for endothelial-leuko-
cyte adhesion
[138521] molecule 1. J. Biol. Chem. 266: 2466–2473, 1991.
[138522] 24703.
- [138523] DeLisser, H. M.; Christofidou-Solomidou, M.; Sun, J.;

Nakada, M.

[138524]T.; Sullivan, K. E.: Loss of endothelial surface expression
of E-selectin

[138525]in a patient with recurrent infections. Blood 94: 884–894,
1999.

[138526]24704.

[138527]Wang, N.; Chintala, S. K.; Fini, M. E.; Schuman, J. S.: Acti-
vation

[138528]of a tissue-specific stress response in the aqueous out-
flow pathway

[138529]of the eye defines the glaucoma disease phenotype. Na-
ture Med. 7:

[138530]304–309, 2001.

[138531]24705.

[138532]Watson, M. L.; Kingsmore, S. F.; Johnston, G. I.; Siegelman,
M.

[138533]H.; Le Beau, M. M.; Lemons, R. S.; Bora, N. S.; Howard, T.
A.; Weissman,

[138534]I. L.; McEver, R. P.; Seldin, M. F.: Genomic organization of
the

[138535]selectin family of leukocyte adhesion molecules on human
and mouse

[138536]chromosome 1. J. Exp. Med. 172: 263–272, 1990.

[138537]24706.

[138538]Wenzel, K.; Felix, S.; Kleber, F. X.; Brachold, R.; Menke, T.;

[138539]Schattke, S.; Schulte, K. L.; Glaser, C.; Rohde, K.; Baumann,
G.;

[138540]Speer, A.: E-selectin polymorphism and atherosclerosis:
an association

[138541]study. Hum. Molec. Genet. 3: 1935–1937, 1994.

[138542]24707.

[138543]Zheng, F.; Chevalier, J. A.; Zhang, L. Q.; Virgil, D.; Ye, S.
Q.;

[138544]Kwiterovich, P. O.: An HphI polymorphism in the E-
selectin gene is

[138545]associated with premature coronary artery disease. Clin.
Genet. 59:

[138546]58–64, 2001.

[138547]24708.

[138548]Burgess, W. H.; Mehlman, T.; Marshak, D. R.; Fraser, B. A.;
Maciag,

[138549]T.: Structural evidence that endothelial cell growth factor
beta

[138550] is the precursor of both endothelial cell growth factor al-
pha and

[138551] acidic fibroblast growth factor. Proc. Nat. Acad. Sci. 83:
7216-7220,

[138552] 1986.

[138553] 24709.

[138554] Eckenstein, F. P.: Fibroblast growth factors in the nervous
system. J.

[138555] Neurobiol. 25: 1467-1480, 1994.

[138556] 24710.

[138557] Gautschi-Sova, P.; Muller, T.; Bohlen, P.: Amino acid se-
quence

[138558] of human acidic fibroblast growth factor. Biochem. Bio-
phys. Res.

[138559] Commun. 140: 874-880, 1986.

[138560] 24711.

[138561] Jaye, M.; Howk, R.; Burgess, W.; Ricca, G. A.; Chiu, I.-M.;
Ravera,

[138562] M. W.; O'Brien, S. J.; Modi, W. S.; Maciag, T.; Drohan, W. N.:
Human

[138563] endothelial cell growth factor: cloning, nucleotide se-

quence, and

[138564]chromosome localization. Science 233: 541–545, 1986.

[138565]24712.

[138566]Jung, J.; Zheng, M.; Goldfarb, M.; Zaret, K. S.: Initiation of

[138567]mammalian liver development from endoderm by fibroblast growth factors. Science 284:

[138568]1998–2003, 1999.

[138569]24713.

[138570]Mergia, A.; Eddy, R.; Abraham, J. A.; Fiddes, J. C.; Shows, T.

[138571]B.: The genes for basic and acidic fibroblast growth factors are

[138572]on different human chromosomes. Biochem. Biophys. Res. Commun. 138:

[138573]644–651, 1986.

[138574]24714.

[138575]Pellegrini, L.; Burke, D. F.; von Delft, F.; Mulloy, B.; Blundell,

[138576]T. L.: Crystal structure of fibroblast growth factor receptor ectodomain

[138577]bound to ligand and heparin. Nature 407: 1029–1034,

2000.

[138578]24715.

[138579]Plotnikov, A. N.; Hubbard, S. R.; Schlessinger, J.; Moham-
madi,

[138580]M.: Crystal structures of two FGF-FGFR complexes reveal
the determinants

[138581]of ligand-receptor specificity. Cell 101: 413-424, 2000.

[138582]24716.

[138583]Wang, W.-P.; Lehtoma, K.; Varban, M. L.; Krishnan, I.;
Chiu, I.-M.

[138584]: Cloning of the gene coding for human class 1 heparin-
binding growth

[138585]factor and its expression in fetal tissues. Molec. Cell. Biol.
9:

[138586]2387-2395, 1989.

[138587]24717.

[138588]Kayes-Wandover, K.; White, P. C.: Steroidogenic enzyme
gene expression

[138589]in the human heart. J. Clin. Endocr. Metab. 85:
2519-2525, 2000.

[138590]24718.

[138591]Carrero-Valenzuela, R. D.; Quan, F.; Lightowlers, R.; Ken-
naway,

[138592]N. G.; Litt, M.; Forte, M.: Human cytochrome c oxidase
subunit VIb:

[138593]characterization and mapping of a multigene family. Gene
102: 229-236,

[138594]1991.

[138595]24719.

[138596]Taanman, J.-W.; Schrage, C.; Ponne, N.; Bolhuis, P.; de
Vries,

[138597]H.; Agsteribbe, E.: Nucleotide sequence of cDNA encoding
subunit

[138598]VIb of human cytochrome c oxidase. Nucleic Acids Res.
17: 1766

[138599]only, 1989.

[138600]24720.

[138601]Taanman, J.-W.; van der Veen, A. Y.; Schrage, C.; de Vries, H.;

[138602]Buys, C. H. C. M.: Assignment of the gene coding for human cytochrome

[138603]c oxidase subunit VIb to chromosome 19, band q13.1, by fluorescence

[138604]in situ hybridisation. Hum. Genet. 87: 325–327, 1991.

[138605]24721.

[138606]Freddi, S.; Savarirayan, R.; Bateman, J. F.: Molecular diagnosis

[138607]of Stickler syndrome: A COL2A1 stop codon mutation screening strategy

[138608]that is not compromised by mutant mRNA instability. Am. J. Med. Genet. 90:

[138609]398–406, 2000.

[138610]24722.

[138611]Griffith, A. J.; Sprunger, L. K.; Sirko-Osadsa, D. A.; Tiller,

[138612]G. E.; Meisler, M. H.; Warman, M. L.: Marshall syndrome associated

[138613]with a splicing defect at the COL11A1 gene. Am. J. Hum.

Genet. 62:

[138614]816-823, 1998.

[138615]24723.

[138616]Cox, D. R.; Epstein, L. B.; Epstein, C. J.: Genes coding for sensitivity

[138617]to interferon (IfRec) and soluble superoxide dismutase (SOD-1) are

[138618]linked in mouse and man and map to mouse chromosome 16. Proc. Nat.

[138619]Acad. Sci. 77: 2168-2172, 1980.

[138620]24724.

[138621]De Clercq, E.; Edy, V. G.; Cassiman, J.-J.: Chromosome 21 does

[138622]not code for an interferon receptor. Nature 264: 249-251, 1976.

[138623]24725.

[138624]Epstein, L. B.; Epstein, C. J.: Localization of the gene AVG for

[138625]the antiviral expression of immune and classical interferon to the

[138626]distal portion of the long arm of chromosome 21. J. Infect.

[138627](suppl.): A56-A62, 1976.

[138628]24726.

[138629]Faltynek, C. R.; Branca, A. A.; McCandless, S.; Baglioni, C.:

[138630]Characterization of an interferon receptor on human lymphoblastoid

[138631]cells. Proc. Nat. Acad. Sci. 80: 3269-3273, 1983.

[138632]24727.

[138633]Fournier, A.; Zhang, Z. Q.; Tan, Y. H.: Human beta:alpha but not

[138634]gamma interferon binding site is a product of the chromosome 21 interferon

[138635]action gene. Somat. Cell Molec. Genet. 11: 291-295, 1985.

[138636]24728.

[138637]Lutfalla, G.; Gardiner, K.; Proudhon, D.; Vielh, E.; Uze, G.:

[138638]The structure of the human interferon alpha/beta receptor gene. J.

[138639]Biol. Chem. 267: 2802-2809, 1992.

[138640]24729.

[138641]Lutfalla, G.; Roeckel, N.; Mogensen, K. E.; Mattei, M. G.;
Uze,

[138642]G.: Assignment of the human interferon- α receptor
gene to chromosome

[138643]21q22.1 by in situ hybridization. J. Interferon Res. 10:
515-517,

[138644]1990.

[138645]24730.

[138646]Maroun, L. E.: Interferon action and chromosome 21 tri-
somy. (Letter) J.

[138647]Theor. Biol. 86: 603-606, 1980.

[138648]24731.

[138649]Novick, D.; Cohen, B.; Rubinstein, M.: The human inter-
feron α/β

[138650]receptor: characterization and molecular cloning. Cell 77:
391-400,

[138651]1994.

[138652]24732.

[138653]Raziuddin, A.; Sarkar, F. H.; Dutkowski, R.; Shulman, L.;
Ruddle,

[138654]F. H.; Gupta, S. L.: Receptors for human α and β in-

terferon

[138655]but not for gamma interferon are specified by human chromosome 21. Proc.

[138656]Nat. Acad. Sci. 81: 5504-5508, 1984.

[138657]24733.

[138658]Revel, M.; Bash, D.; Ruddle, F. H.: Antibodies to a cell-surface

[138659]component coded by human chromosome 21 inhibit action of interferon. Nature 260:

[138660]139-141, 1976.

[138661]24734.

[138662]Calabrese, G.; Crescenzi, C.; Morizio, E.; Palka, G.; Guerra, E.;

[138663]Alberti, S.: Assignment of TACSTD1 (alias TROP1, M4S1) to human chromosome

[138664]2p21 and refinement of mapping of TACSTD2 (alias TROP2, M1S1) to human

[138665]chromosome 1p32 by in situ hybridization. Cytogenet. Cell Genet. 92:

[138666]164-165, 2001.

[138667]24735.

[138668]Fornaro, M.; Dell'Arciprete, R.; Stella, M.; Bucci, C.; Nutini,
[138669]M.; Capri, M. G.; Alberti, S.: Cloning of the gene encoding
Trop-2,
[138670]a cell-surface glycoprotein expressed by human carcino-
mas. Int. J.
[138671]Cancer 62: 610-618, 1995.
[138672]24736.
[138673]Linnenbach, A. J.; Seng, B. A.; Wu, S.; Robbins, S.; Scollon,
M.;
[138674]Pyr, J. J.; Druck, T.; Huebner, K.: Retroposition in a family
of
[138675]carcinoma-associated antigen genes. Molec. Cell Biol. 13:
1507-1515,
[138676]1993.
[138677]24737.
[138678]Linnenbach, A. J.; Wojcierowski, J.; Wu, S.; Pyr, J. J.; Ross,
[138679]A. H.; Dietzschold, B.; Speicher, D.; Koprowski, H.: Se-
quence investigation
[138680]of the major gastrointestinal tumor-associated antigen
gene family,
[138681]GA733. Proc. Nat. Acad. Sci. 86: 27-31, 1989.

[138682]24738.

[138683]Ren, Z.; Lin, P.-Y.; Klintworth, G. K.; Iwata, F.; Munier, F.
L.;

[138684]Schorderet, D. F.; El Matri, L.; Theendakara, V.; Basti, S.;
Reddy,

[138685]M.; Hejtmancik, J. F.: Allelic and locus heterogeneity in au-
tosomal

[138686]recessive gelatinous drop-like corneal dystrophy. Hum.
Genet. 110:

[138687]568-577, 2002.

[138688]24739.

[138689]TsujiKawa, M.; Kurahashi, H.; Tanaka, T.; Nishida, K.; Shi-
momura,

[138690]Y.; Tano, Y.; Nakamura, Y.: Identification of the gene re-
sponsible

[138691]for gelatinous drop-like corneal dystrophy. Nature Genet.
21: 420-423,

[138692]1999.

[138693]24740.

[138694]Briegel, K.; Lim, K.-C.; Plank, C.; Beug, H.; Engel, J. D.;
Zenke,

- [138695]M.: Ectopic expression of a conditional GATA-2/estrogen receptor
- [138696]chimera arrests erythroid differentiation in a hormone-dependent manner. *Genes Dev.* 7: 1097-1109, 1993.
- [138698]24741.
- [138699]Ciciotte, S. L.; Tsai, F.-Y.; Peters, L. L.: Gata2 maps to mouse chromosome 6. *Mammalian Genome* 8: 69-70, 1997.
- [138701]24742.
- [138702]Dasen, J. S.; O'Connell, S. M.; Flynn, S. E.; Treier, M.; Gleiberman, A. S.; Szeto, D. P.; Hooshmand, F.; Aggarwal, A. K.; Rosenfeld, M.
- [138704]G.: Reciprocal interactions of Pit1 and GATA2 mediate signaling gradient-induced determination of pituitary cell types. *Cell* 97: 587-598, 1999.
- [138706]24743.
- [138707]Dorfman, D. M.; Wilson, D. B.; Bruns, G. A.; Orkin, S. H.: Human

[138708]transcription factor GATA-2: evidence for regulation of
preproendothelin-1

[138709]gene expression in endothelial cells. J. Biol. Chem. 267:
1279-1285,

[138710]1992.

[138711]24744.

[138712]Lee, M.-E.; Temizer, D. H.; Clifford, J. A.; Quertermous, T.:

[138713]Cloning of the GATA-binding protein that regulates en-
dothelin-1 gene

[138714]expression in endothelial cells. J. Biol. Chem. 266:
16188-16192,

[138715]1991.

[138716]24745.

[138717]Tsai, F.-Y.; Keller, G.; Kuo, F. C.; Weiss, M.; Chen, J.;
Rosenblatt,

[138718]M.; Alt, F. W.; Orkin, S. H.: An early haematopoietic defect
in mice

[138719]lacking the transcription factor GATA2. Nature 371:
221-226, 1994.

[138720]24746.

[138721]Zhou, Y.; Lim, K.-C.; Onodera, K.; Takahashi, S.; Ohta, J.;

Minegishi,

[138722]N.; Tsai, F.-Y.; Orkin, S. H.; Yamamoto, M.; Engel, J. D.:

Rescue

[138723]of the embryonic lethal hematopoietic defect reveals a
critical role

[138724]for GATA-2 in urogenital development. EMBO J. 17:
6689-6700, 1998.

[138725]24747.

[138726]Becker-Follmann, J.; Gaa, A.; Bausch, E.; Natt, E.; Scherer,
G.;

[138727]von Deimling, O.: High-resolution mapping of a linkage
group on mouse

[138728]chromosome 8 conserved on human chromosome 16Q.
Mammalian Genome 8:

[138729]172-177, 1997.

[138730]24748.

[138731]Brooks, P. C.; Silletti, S.; von Schalscha, T. L.; Friedlander,

[138732]M.; Cheresh, D. A.: Disruption of angiogenesis by PEX, a
noncatalytic

[138733]metalloproteinase fragment with integrin binding activity.
Cell 92:

[138734]391-400, 1998.

[138735]24749.

[138736]Collier, I. E.; Bruns, G. A. P.; Goldberg, G. I.; Gerhard, D. S.

[138737]: On the structure and chromosome location of the 72-
and 92-kDa human

[138738]type IV collagenase genes. Genomics 9: 429-434, 1991.

[138739]24750.

[138740]Corry, D. B.; Rishi, K.; Kanellis, J.; Kiss, A.; Song, L.; Xu,

[138741]J.; Feng, L.; Werb, Z.; Kheradmand, F.: Decreased allergic
lung inflammatory

[138742]cell egression and increased susceptibility to asphyxiation
in MMP2-deficiency. Nature

[138743]Immun. 3: 347-353, 2002.

[138744]24751.

[138745]Devarajan, P.; Johnston, J. J.; Ginsberg, S. S.; Van Wart, H.
E.;

[138746]Berliner, N.: Structure and expression of neutrophil gelati-
nase cDNA:

[138747]identity with type IV collagenase from HT1080 cells. J.
Biol. Chem. 267:

[138748]25228-25232, 1992.

[138749]24752.

[138750]Fan, Y.-S.; Eddy, R. L.; Huhtala, P.; Byers, M. G.; Haley, L. L.;

[138751]Henry, W. M.; Tryggvason, K.; Shows, T. B.: Collagenase type IV (CLG4)

[138752]is mapped to human chromosome 16q21. (Abstract) Cytogenet. Cell Genet. 51:

[138753]996, 1989.

[138754]24753.

[138755]Huhtala, P.; Chow, L. T.; Tryggvason, K.: Structure of the human

[138756]type IV collagenase gene. J. Biol. Chem. 265: 11077-11082, 1990.

[138757]24754.

[138758]Huhtala, P.; Eddy, R. L.; Fan, Y. S.; Byers, M. G.; Shows, T.

[138759]B.; Tryggvason, K.: Completion of the primary structure of the human

[138760]type IV collagenase preproenzyme and assignment of the gene (CLG4)

[138761]to the q21 region of chromosome 16. Genomics 6: 554-559, 1990.

[138762]24755.

[138763]Irwin, J. C.; Kirk, D.; Gwatkin, R. B. L.; Navre, M.; Cannon,

[138764]P.; Giudice, L. C.: Human endometrial matrix metallopro-
teinase-2,

[138765]a putative menstrual proteinase: hormonal regulation in
cultured stromal

[138766]cells and messenger RNA expression during the menstrual
cycle. J.

[138767]Clin. Invest. 97: 438-447, 1996.

[138768]24756.

[138769]Itoh, T.; Tanioka, M.; Yoshida, H.; Yoshioka, T.; Nishimoto,
H.;

[138770]Itohara, S.: Reduced angiogenesis and tumor progression
in gelatinase

[138771]A-deficient mice. Cancer Res. 58: 1048-1051, 1998.

[138772]24757.

[138773]Kato, T.; Kure, T.; Chang, J.-H.; Gabison, E. E.; Itoh, T.;
Itohara,

[138774]S.; Azar, D. T.: Diminished corneal angiogenesis in gelati-
nase A-deficient

[138775]mice. FEBS Lett. 508: 187-190, 2001.

[138776]24758.

[138777]Morgunova, E.; Tuuttila, A.; Bergmann, U.; Isupov, M.;
Lindqvist,

[138778]Y.; Schneider, G.; Tryggvason, K.: Structure of human pro-
matrix

[138779]metalloproteinase-2: activation mechanism revealed. Sci-
ence 284:

[138780]1667-1670, 1999.

[138781]24759.

[138782]Vu, T. H.: Don't mess with the matrix. Nature Genet. 28:
202-204,

[138783]2001.

[138784]24760.

[138785]Willecke, K.; Jungbluth, S.; Dahl, E.; Hennemann, H.;
Heynkes,

[138786]R.; Grzeschik, K.-H.: Six genes of the human connexin
gene family

[138787]coding for gap junctional proteins are assigned to four
different

[138788]human chromosomes. Europ. J. Cell Biol. 53: 275-280,
1990.

[138789]24761.

[138790]Reed, K. E.; Westphale, E. M.; Larson, D. M.; Wang, H.-Z.;
Veenstra,

[138791]R. D.; Beyer, E. C.: Molecular cloning and functional ex-
pression

[138792]of human connexin37, an endothelial cell gap junction
protein. J.

[138793]Clin. Invest. 91: 997-1004, 1993.

[138794]24762.

[138795]Van Camp, G.; Coucke, P.; Speleman, F.; Van Roy, N.;
Beyer, E.

[138796]C.; Oostra, B. A.; Willems, P. J.: The gene for human gap
junction

[138797]protein connexin37 (GJA4) maps to chromosome 1p35.1,
in the vicinity

[138798]of D1S195. Genomics 30: 402-403, 1995.

[138799]24763.

[138800]Gelb, B. D.; Zhang, J.; Cotter, P. D.; Gershin, I. F.; Desnick,

[138801]R. J.: Physical mapping of the human connexin 40 (GJA5)
flavin-containing

[138802]monooxygenase 5, and natriuretic peptide receptor A

genes on 1q21. Genomics 39:

[138803]409–411, 1997.

[138804]24764.

[138805]Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Cardiac myocytes

[138806]express multiple gap junction proteins. Circ. Res. 70:

438–444,

[138807]1992.

[138808]24765.

[138809]Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Molecular cloning

[138810]of two human cardiac gap junction proteins, connexin40

and connexin45. J.

[138811]Molec. Cell. Cardiol. 26: 861–868, 1994.

[138812]24766.

[138813]Oviedo–Orta, E.; Hoy, T.; Evans, W. H.: Intercellular com-

munication

[138814]in the immune system: differential expression of con-

nexin40 and 43,

[138815]and perturbation of gap junction channel functions in pe-

ripheral blood

[138816]and tonsil human lymphocyte subpopulations. Immunol–

- ogy 99: 578–590,
[138817]2000.
- [138818]24767.
- [138819]Seul, K. H.; Tadros, P. N.; Beyer, E. C.: Mouse connexin40:
gene
[138820]structure and promoter analysis. Genomics 46: 120–126,
1997.
- [138821]24768.
- [138822]Britz–Cunningham, S. H.; Shah, M. M.; Zuppan, C. W.;
Fletcher,
[138823]W. H.: Mutations of the connexin43 gap–junction gene in
patients
[138824]with heart malformations and defects of laterality. New
Eng. J. Med. 332:
[138825]1323–1329, 1995.
- [138826]24769.
- [138827]Brueckner, M.; D'Eustachio, P.; Horwich, A. L.: Linkage
mapping
[138828]of a mouse gene, iv, that controls left–right asymmetry of
the heart
[138829]and viscera. Proc. Nat. Acad. Sci. 86: 5035–5038, 1989.

[138830]24770.

[138831]Burdine, R. D.; Schier, A. F.: Conserved and divergent mechanisms

[138832]in left-right axis formation. Genes Dev. 14: 763-776, 2000.

[138833]24771.

[138834]Casey, B.; Ballabio, A.: Connexin43 mutations in sporadic and

[138835]familial defects of laterality. (Letter) New Eng. J. Med. 333: 941,

[138836]1995.

[138837]24772.

[138838]Corcos, I. A.; Meese, E. U.; Loch-Carusio, R.: Human connexin 43

[138839]gene locus, GJA1, sublocalized to band 6q21-q23.2. Cytogenet. Cell

[138840]Genet. 64: 31-32, 1993.

[138841]24773.

[138842]Debrus, S.; Tuffery, S.; Matsuoka, R.; Galal, O.; Sarda, P.; Sauer,

[138843]U.; Bozio, A.; Tanman, B.; Toutain, A.; Claustres, M.; Le

Paslier,

[138844]D.; Bouvagnet, P.: Lack of evidence for connexin 43 gene mutations

[138845]in human autosomal recessive lateralization defects. J. Molec. Cell.

[138846]Cardiol. 29: 1423–1431, 1997.

[138847]24774.

[138848]Przylepa, K. A.; Paznekas, W.; Zhang, M.; Golabi, M.; Bias, W.;

[138849]Bamshad, M. J.; Carey, J. C.; Hall, B. D.; Stevenson, R.; Orlow, S.

[138850]J.; Cohen, M. M., Jr.; Jabs, E. W.: Fibroblast growth factor receptor

[138851]2 mutations in Beare–Stevenson cutis gyrata syndrome. Nature Genet. 13:

[138852]492–494, 1996.

[138853]24775.

[138854]Hai, T.; Liu, F.; Coukos, W. J.; Green, M. R.: Transcription factor

[138855]ATF cDNA clones: an extensive family of leucine zipper proteins able

[138856]to selectively form DNA-binding heterodimers. Genes Dev.

3: 2083–2090,

[138857]1989.

[138858]24776.

[138859]Herzig, S.; Long, F.; Jhala, U. S.; Hedrick, S.; Quinn, R.;
Bauer,

[138860]A.; Rudolph, D.; Schutz, G.; Yoon, C.; Puigserver, P.;
Spiegelman,

[138861]B.; Montminy, M.: CREB regulates hepatic gluconeogenesis
through

[138862]the coactivator PGC-1. *Nature* 413: 179–183, 2001.

[138863]24777.

[138864]Benson, A. M.; Hunkeler, M. J.; Talalay, P.: Increase of
NAD(P)H:quinone

[138865]reductase by dietary antioxidants: possible role in protec-
tion against

[138866]carcinogenesis and toxicity. *Proc. Nat. Acad. Sci.* 77:
5216–5220,

[138867]1980.

[138868]24778.

[138869]Chen, L. Z.; Harris, P. C.; Apostolou, S.; Baker, E.; Holman,
K.;

[138870] Lane, S. A.; Nancarrow, J. K.; Whitmore, S. A.; Stallings, R. L.;

[138871] Hildebrand, C. E.; Richards, R. I.; Sutherland, G. R.; Callen, D.

[138872] F.: A refined physical map of the long arm of human chromosome 16. Genomics 10:

[138873] 308–312, 1991.

[138874] 24779.

[138875] Edwards, Y. H.; Hopkinson, D. A.; Carritt, B.: A genetic characterization

[138876] of the human diaphorase–4 deficiency. Ann. Hum. Genet. 47: 97–105,

[138877] 1983.

[138878] 24780.

[138879] Grzeschik, K.–H.: Assignment of a structural gene for a fourth

[138880] human diaphorase (DIA–4) to chromosome 16 in man–mouse somatic cell

[138881] hybrids. Hum. Genet. 53: 189–193, 1980.

[138882] 24781.

[138883] Jaiswal, A. K.; McBride, O. W.; Adesnik, M.; Nebert, D. W.:

Human

[138884]dioxin-inducible cytosolic NAD(P)H:menadione oxidoreductase: cDNA

[138885]sequence and localization of gene to chromosome 16. J. Biol. Chem. 263:

[138886]13572-13578, 1988.

[138887]24782.

[138888]Kelsey, K. T.; Ross, D.; Traver, R. D.; Christiani, D. C.; Zuo,

[138889]Z. F.; Spitz, M. R.; Wang, M.; Xu, X.; Lee, B. K.; Schwartz, B. S.;

[138890]Wiencke, J. K.: Ethnic variation in the prevalence of a common NAD(P)H

[138891]quinone oxidoreductase polymorphism and its implications for anti-cancer

[138892]chemotherapy. Brit. J. Cancer 76: 852-854, 1997.

[138893]24783.

[138894]Lavinha, J.; Morrison, N.; Glasgow, L.; Ferguson-Smith, M. A.:

[138895]Further evidence for the regional localization of human APRT and DIA4

[138896]on chromosome 16. (Abstract) Cytogenet. Cell Genet. 37:

517 only,

[138897]1984.

[138898]24784.

[138899]Moran, J. L.; Siegel, D.; Ross, D.: A potential mechanism
underlying

[138900]the increased susceptibility of individuals with a polymor-
phism in

[138901]NAD(P)H:quinone oxidoreductase 1 (NQO1) to benzene
toxicity. Proc.

[138902]Nat. Acad. Sci. 96: 8150–8155, 1999.

[138903]24785.

[138904]Povey, S.; Wilson, D.; Edwards, Y. H.: Assignment of a hu-
man diaphorase

[138905](DIA-4) to chromosome 16. Ann. Hum. Genet. 43:
349–353, 1980.

[138906]24786.

[138907]Radjendirane, V.; Joseph, P.; Lee, Y.-H.; Kimura, S.; Klein-
Szanto,

[138908]A. J. P.; Gonzalez, F. J.; Jaiswal, A. K.: Disruption of the DT
diaphorase

[138909](NQO1) gene in mice leads to increased menadione toxic-

ity. J. Biol.

[138910]Chem. 273: 7382–7389, 1998.

[138911]24787.

[138912]Rothman, N.; Smith, M. T.; Hayes, R. B.; Traver, R. D.;
Hoener,

[138913]B.; Campleman, S.; Li, G. L.; Dosemeci, M.; Linet, M.;
Zhang, L.;

[138914]Xi, L.; Wacholder, S.; Lu, W.; Meyer, K. B.; Titenko–Holland,
N.;

[138915]Stewart, J. T.; Yin, S.; Ross, D.: Benzene poisoning, a risk
factor

[138916]for hematological malignancy, is associated with the
NQO1 609C–T mutation

[138917]and rapid fractional excretion of chlorzoxazone. Cancer
Res. 57:

[138918]2839–2842, 1997.

[138919]24788.

[138920]Smith, M. T.: Benzene, NQO1, and genetic susceptibility to
cancer.

[138921](Commentary) Proc. Nat. Acad. Sci. 96: 7624–7626, 1999.

[138922]24789.

[138923]Traver, R. D.; Horikoshi, T.; Danenberg, K. D.; Stadlbauer, T.

[138924]H.; Danenberg, P. V.; Ross, D.; Gibson, N. W.:

NAD(P)H:quinone oxidoreductase

[138925]gene expression in human colon carcinoma cells: characterization of

[138926]a mutation which modulates DT-diaphorase activity and mitomycin sensitivity. Cancer

[138927]Res. 52: 797-802, 1992.

[138928]24790.

[138929]Traver, R. D.; Siegel, D.; Beall, H. D.; Phillips, R. M.; Gibson,

[138930]N. W.; Franklin, W. A.; Ross, D.: Characterization of a polymorphism

[138931]in NAD(P)H: quinone oxidoreductase (DT-diaphorase). Brit. J. Cancer 75:

[138932]69-75, 1997.

[138933]24791.

[138934]Zhang, L.; Rothman, N.; Wang, Y.; Hayes, R. B.; Li, G.; Dosemeci,

[138935]M.; Yin, S.; Kolachana, P.; Titenko-Holland, N.; Smith, M. T.: Increased

[138936]aneusomy and long arm deletion of chromosomes 5 and 7
in the lymphocytes

[138937]of Chinese workers exposed to benzene. Carcinogenesis
19: 1955–1961,

[138938]1998.

[138939]24792.

[138940]Zhang, L.; Wang, Y.; Shang, N.; Smith, M. T.: Benzene
metabolites

[138941]induce the loss of long arm deletion of chromosomes 5
and 7 in human

[138942]lymphocytes. Leukemia Res. 22: 105–113, 1998.

[138943]24793.

[138944]Roychoudhury, A. K.; Nei, M.: Human Polymorphic Genes:
World Distribution.

[138945]New York: Oxford Univ. Press (pub.) 1988.

[138946]24794.

[138947]Hark, A. T.; Schoenherr, C. J.; Katz, D. J.; Ingram, R. S.;
Levrose,

[138948]J. M.; Tilghman, S. M.: CTCF mediates methylation–sensi-
tive enhancer–blocking

[138949]activity at the H19/Igf2 locus. Nature 405: 486–489,

2000.

[138950]24795.

[138951]Hark, A. T.; Tilghman, S. M.: Chromatin conformation of
the H19

[138952]epigenetic mark. Hum. Molec. Genet. 7: 1979–1985,
1998.

[138953]24796.

[138954]Jones, B. K.; Levorse, J.; Tilghman, S. M.: Deletion of a nu-
clease-sensitive

[138955]region between the Igf2 and H19 genes leads to Igf2 mis-
regulation

[138956]and increased adiposity. Hum. Molec. Genet. 10:
807–814, 2001.

[138957]24797.

[138958]Jones, J. M.; Meisler, M. H.; Seldin, M. F.; Lee, B. K.; Eicher,

[138959]E. M.: Localization of insulin-2 (Ins-2) and the obesity
mutant tubby

[138960](tub) to distinct regions of mouse chromosome 7. Ge-
nomics 14: 197–199,

[138961]1992.

[138962]24798.

[138963]Rainier, S.; Johnson, L. A.; Dobry, C. J.; Ping, A. J.; Grundy,
[138964]P. E.; Feinberg, A. P.: Relaxation of imprinted genes in hu-
man cancer. Nature 362:
[138965]747–749, 1993.
[138966]24799.
[138967]Srivastava, M.; Hsieh, S.; Grinberg, A.; Williams-Simons, L.;
[138968]Huang, S.-P.; Pfeifer, K.: H19 and Igf2 monoallelic expres-
sion is
[138969]regulated in two distinct ways by a shared cis acting regu-
latory region
[138970]upstream of H19. Genes Dev. 14: 1186–1195, 2000.
[138971]24800.
[138972]Thorvaldsen, J. L.; Duran, K. L.; Bartolomei, M. S.: Deletion
[138973]of the H19 differentially methylated domain results in loss
of imprinted
[138974]expression of H19 and Igf2. Genes Dev. 12: 3693–3702,
1998.
[138975]24801.
[138976]Webber, A.; Ingram, R. S.; Levorse, J. M.; Tilghman, S. M.:
Location

[138977]of enhancers is essential for imprinting of H19 and Igf2
genes. Nature 391:

[138978]711–715, 1998.

[138979]24802.

[138980]Zhang, Y.; Tycko, B.: Monoallelic expression of the human
H19

[138981]gene. Nature Genet. 1: 40–44, 1992.

[138982]24803.

[138983]Campanelli, J. T.; Hoch, W.; Rupp, F.; Kreiner, T.; Scheller,
R.

[138984]H.: Agrin mediates cell contact–induced acetylcholine re-
ceptor clustering. Cell 67:

[138985]909–916, 1991.

[138986]24804.

[138987]DeChiara, T. M.; Bowen, D. C.; Valenzuela, D. M.; Sim-
mons, M. V.;

[138988]Poueymirou, W. T.; Thomas, S.; Kinetz, E.; Compton, D. L.;
Rojas,

[138989]E.; Park, J. S.; Smith, C.; DiStefano, P. S.; Glass, D. J.; Bur-
den,

[138990]S. J.; Yancopoulos, G. D.: The receptor tyrosine kinase

MuSK is required

[138991]for neuromuscular junction formation in vivo. Cell 85:
501–512,

[138992]1996.

[138993]24805.

[138994]Chan, S. Y.; Empig, C. J.; Welte, F. J.; Speck, R. F.;
Schmaljohn,

[138995]A.; Kreisberg, J. F.; Goldsmith, M. A.: Folate receptor- α is
pha is

[138996]a cofactor for cellular entry by Marburg and Ebola viruses.
Cell 106:

[138997]117–126, 2001.

[138998]24806.

[138999]Czeizel, A. E.; Dudas, I.: Prevention of the first occurrence

[139000]of neural-tube defects by periconceptional vitamin sup-
plementation. New

[139001]Eng. J. Med. 327: 1832–1835, 1992.

[139002]24807.

[139003]De Marco, P.; Moroni, A.; Merello, E.; de Franchis, R.; An-
dreussi,

[139004]L.; Finnell, R. H.; Barber, R. C.; Cama, A.; Capra, V.: Folate

pathway

[139005]gene alterations in patients with neural tube defects. Am. J. Med.

[139006]Genet. 95: 216–223, 2000.

[139007]24808.

[139008]Elwood, P. C.: Molecular cloning and characterization of the human

[139009]folate-binding protein cDNA from placenta and malignant tissue culture

[139010](KB) cells. J. Biol. Chem. 264: 14893–14901, 1989.

[139011]24809.

[139012]Elwood, P. C.; Nachmanoff, K.; Saikawa, Y.; Page, S. T.; Pacheco,

[139013]P.; Roberts, S.; Chung, K.–N.: The divergent 5–prime termini of the

[139014]alpha human folate receptor (hFR) mRNAs originate from two tissue-specific

[139015]promoters and alternative splicing: characterization of the alpha

[139016]hFR gene structure. Biochemistry 36: 1467–1478, 1997.

[139017]24810.

- [139018]Boman, B. M.; Wildrick, D. M.; Alfaro, S. R.: Chromosome
18 allele
- [139019]Loss at the D18S6 locus in human colorectal carcinomas.
Biochem.
- [139020]Biophys. Res. Commun. 155: 463–469, 1988.
- [139021]24811.
- [139022]Chan, S. S.–Y.; Zheng, H.; Su, M.–W.; Wilk, R.; Killeen, M.
T.;
- [139023]Hedgecock, E. M.; Culotti, J. G.: UNC–40, a *C. elegans* ho–
molog of
- [139024]the DCC (deleted in colorectal cancer), is required in
motile cells
- [139025]responding to UNC–6 netrin cues. Cell 87: 187–195,
1996.
- [139026]24812.
- [139027]Sarmanova, J.; Benesova, K.; Gut, I.; Nedelcheva–Kris–
tensen, V.;
- [139028]Tynkova, L.; Soucek, P.: Genetic polymorphisms of bio–
transformation
- [139029]enzymes in patients with Hodgkin's and non–Hodgkin's
lymphomas. Hum.
- [139030]Molec. Genet. 10: 1265–1273, 2001.

[139031]24813.

[139032]Albrecht, F. E.; Drago, J.; Felder, R. A.; Printz, M. P.; Eisner,

[139033]G. M.; Robillard, J. E.; Sibley, D. R.; Westphal, H. J.; Jose, P.

[139034]A.: Role of the D-1A dopamine receptor in the pathogenesis of genetic

[139035]hypertension. J. Clin. Invest. 97: 2283-2288, 1996.

[139036]24814.

[139037]Bermak, J. C.; Li, M.; Bullock, C.; Zhou, Q.-Y.: Regulation of

[139038]transport of the dopamine D1 receptor by a new membrane-associated

[139039]ER protein. Nature Cell Biol. 3: 492-498, 2001.

[139040]24815.

[139041]Boulton, J.; Lewis, M. S.; Wainscoat, J. S.: Physical linkage

[139042]of glucocorticoid receptor (GRL) and dopamine D1 receptor (DRD1) on

[139043]the long arm of chromosome 5.(Abstract) Cytogenet. Cell Genet. 58:

[139044]1894 only, 1991.

[139045]24816.

[139046]Castner, S. A.; Williams, G. V.; Goldman-Rakic, P. S.: Re-
versal

[139047]of antipsychotic-induced working memory deficits by
short-term dopamine

[139048]D1 receptor stimulation. Science 287: 2020-2022, 2000.

[139049]24817.

[139050]Dearry, A.; Gingrich, J. A.; Falardeau, P.; Fremeau, R. T.,
Jr.;

[139051]Bates, M. D.; Caron, M. G.: Molecular cloning and expres-
sion of the

[139052]gene for a human D(1) dopamine receptor. Nature 347:
72-76, 1990.

[139053]24818.

[139054]Grandy, D. K.; Zhou, Q.-Y.; Allen, L.; Litt, R.; Magenis, R.
E.;

[139055]Civelli, O.; Litt, M.: A human D(1) dopamine receptor gene
is located

[139056]on chromosome 5 at q35.1 and identifies an EcoRI RFLP.
Am. J. Hum.

[139057]Genet. 47: 828-834, 1990.

[139058]24819.

[139059]Krushkal, J.; Xiong, M.; Ferrell, R.; Sing, C. F.; Turner, S. T.;

[139060]Boerwinkle, E.: Linkage and association of adrenergic and
dopamine

[139061]receptor genes in the distal portion of the long arm of
chromosome

[139062]5 with systolic blood pressure variation. Hum. Molec.
Genet. 7:

[139063]1379–1383, 1998.

[139064]24820.

[139065]Mayerhofer, A.; Hemmings, H. C., Jr.; Snyder, G. L.; Green-
gard,

[139066]P.; Boddien, S.; Berg, U.; Brucker, C.: Functional
dopamine-1 receptors

[139067]and DARPP-32 are expressed in human ovary and granu-
losa luteal cells

[139068]in vitro. J. Clin. Endocr. Metab. 84: 257–264, 1999.

[139069]24821.

[139070]Sunahara, R. K.; Niznik, H. B.; Weiner, D. M.; Stormann, T.
M.;

[139071]Brann, M. R.; Kennedy, J. L.; Gelernter, J. E.; Rozmahel, R.;

Yang,

[139072]Y.; Israel, Y.; Seeman, P.; O'Dowd, B. F.: Human dopamine
D1 receptor

[139073]encoded by an intronless gene on chromosome 5. Nature
347: 80–83,

[139074]1990.

[139075]24822.

[139076]Wilkie, T. M.; Chen, Y.; Gilbert, D. J.; Moore, K. J.; Yu, L.;

[139077]Simon, M. I.; Copeland, N. G.; Jenkins, N. A.: Identification,
chromosomal

[139078]location, and genome organization of mammalian G-
protein-coupled receptors. Genomics 18:

[139079]175–184, 1993.

[139080]24823.

[139081]Xu, M.; Moratalla, R.; Gold, L. H.; Hiroi, N.; Koob, G. F.;
Graybiel,

[139082]A. M.; Tonegawa, S.: Dopamine D1 receptor mutant mice
are deficient

[139083]in striatal expression of dynorphin and in dopamine-mediated behavioral

[139084]responses. Cell 79: 729–742, 1994.

[139085]24824.

[139086]Zhou, Q.-Y.; Grandy, D. K.; Thambi, L.; Kushner, J. A.; Van Tol,

[139087]H. H. M.; Cone, R.; Pribnow, D.; Salon, J.; Bunzow, J. R.; Civelli,

[139088]O.: Cloning and expression of human and rat D(1) dopamine receptors. *Nature* 347:

[139089]76–80, 1990.

[139090]24825.

[139091]Balk, J.-H.; Picetti, R.; Salardi, A.; Thirlet, G.; Dierich, A.;

[139092]Depaulls, A.; Le Meur, M.; Borrelli, E.: Parkinsonian-like locomotor

[139093]impairment in mice lacking dopamine D2 receptors. *Nature* 377: 424–428,

[139094]1995.

[139095]24826.

[139096]Basu, S.; Nagy, J. A.; Pal, S.; Vasile, E.; Eckelhoefer, I. A.;

[139097]Bliss, V. S.; Manseau, E. J.; Dasgupta, P. S.; Dvorak, H. F.; Mukhopadhyay,

[139098]D.: The neurotransmitter dopamine inhibits angiogenesis induced by

[139099]vascular permeability factor/vascular endothelial growth factor. Nature

[139100]Med. 7: 569–574, 2001.

[139101]24827.

[139102]Blum, K.; Noble, E. P.; Sheridan, P. J.; Montgomery, A.; Ritchie,

[139103]T.; Jagadeeswaran, P.; Nogami, H.; Briggs, A. H.; Cohn, J. B.: Allelic

[139104]association of human dopamine D(2) receptor gene in alcoholism. J.A.M.A. 263:

[139105]2055–2060, 1990.

[139106]24828.

[139107]Bolos, A. M.; Dean, M.; Lucas–Derse, S.; Ramsburg, M.; Brown, G.

[139108]L.; Goldman, D.: Population and pedigree studies reveal a lack of

[139109]association between the dopamine D(2) receptor gene and alcoholism. J.A.M.A. 264:

[139110]3156–3160, 1990.

[139111]24829.

[139112]Bunzow, J. R.; Van Tol, H. H.; Grandy, D. K.; Albert, P.; Sa–

lon,

[139113]J.; Christie, M.; Machida, C. A.; Neve, K. A.; Civelli, O.:

Cloning

[139114]and expression of a rat D2 dopamine receptor cDNA. Nature 336: 783–787,

[139115]1988.

[139116]24830.

[139117]Dal Toso, R.; Sommer, B.; Ewert, M.; Herb, A.; Pritchett, D. B.;

[139118]Bach, A.; Shivers, B. D.; Seeburg, P. H.: The dopamine D(2) receptor:

[139119]two molecular forms generated by alternative splicing.

EMBO J. 8:

[139120]4025–4034, 1989.

[139121]24831.

[139122]Matsutani, A.; Hing, A.; Steinbrueck, T.; Janssen, R.; Weber,

[139123]J.; Permutt, M. A.; Donis-Keller, H.: Mapping the human liver/islet

[139124]glucose transporter (GLUT2) gene within a genetic linkage map of chromosome

[139125]3q using a (CA)_n dinucleotide repeat polymorphism and

characterization

[139126]of the polymorphism in three racial groups. Genomics 13:
495–501,

[139127]1992.

[139128]24832.

[139129]Meera Khan, P.: Personal Communication. Leiden, The
Netherlands

[139130]1977.

[139131]24833.

[139132]Garman, S. C.; Kinet, J.-P.; Jardetzky, T. S.: Crystal struc-
ture

[139133]of the human high-affinity IgE receptor. Cell 95: 951–961,
1998.

[139134]24834.

[139135]Garman, S. C.; Wurzburg, B. A.; Tarchevskaya, S. S.; Kinet,
J.-P.;

[139136]Jardetzky, T. S.: Structure of the Fc fragment of human IgE
bound

[139137]to its high-affinity receptor Fc-epsilon-RI-alpha. Nature
406: 259–266,

[139138]2000.

[139139]24835.

[139140]Liu, F.-T.; Albrandt, K.; Robertson, M. W.: cDNA heterogeneity

[139141]suggests structural variants related to the high-affinity IgE receptor. Proc.

[139142]Nat. Acad. Sci. 85: 5639-5643, 1988.

[139143]24836.

[139144]Shimizu, A.; Tepler, I.; Benfey, P. N.; Berenstein, E. H.; Siraganian,

[139145]R. P.; Leder, P.: Human and rat mast cell high-affinity immunoglobulin

[139146]E receptors: characterization of putative alpha-chain gene products. Proc.

[139147]Nat. Acad. Sci. 85: 1907-1911, 1988.

[139148]24837.

[139149]Miyajima, I.; Dombrowicz, D.; Martin, T. R.; Ravetch, J. V.; Kinet,

[139150]J.-P.; Galli, S. J.: Systemic anaphylaxis in the mouse can be mediated

[139151]largely through IgG1 and Fc-gamma-RIII: assessment of the cardiopulmonary

[139152]changes, mast cell degranulation, and death associated with active

[139153]or IgE- or IgG1-dependent passive anaphylaxis. J. Clin. Invest. 99:

[139154]901-914, 1997.

[139155]24838.

[139156]Tepler, I.; Morton, C. C.; Shimizu, A.; Holcombe, R. F.; Eddy,

[139157]R.; Shows, T. B.; Leder, P.: The gene for the human mast cell high-affinity

[139158]IgE receptor alpha chain: chromosomal localization to 1q21-q23 and

[139159]RFLP analysis. Am. J. Hum. Genet. 45: 761-765, 1989.

[139160]24839.

[139161]Zhu, D.; Kepley, C. L.; Zhang, M.; Zhang, K.; Saxon, A.: A novel

[139162]human immunoglobulin Fc-gamma-Fc-epsilon bifunctional fusion protein

[139163]inhibits Fc-epsilon-RI-mediated degranulation. Nature Med. 8: 518-521,

[139164]2002.

[139165]24840.

[139166]Bain, G.; Robanus Maandag, E. C.; Izon, D. J.; Amsen, D.;
Kruisbeek,

[139167]A. M.; Weintraub, B. C.; Krop, I.; Schlissel, M. S.; Feeney, A.
J.;

[139168]van Roon, M.; van der Valk, M.; te Riele, H. P. J.; Berns, A.;
Murre,

[139169]C.: E2A proteins are required for proper B cell develop-
ment and initiation

[139170]of immunoglobulin gene rearrangements. Cell 79:
885–892, 1994.

[139171]24841.

[139172]El Ghouzzi, V.; Legeai-Mallet, L.; Aresta, S.; Benoist, C.;
Munnich,

[139173]A.; de Gunzburg, J.; Bonaventure, J.: Saethre–Chotzen mu-
tations cause

[139174]TWIST protein degradation or impaired nuclear location.
Hum. Molec.

[139175]Genet. 9: 813–819, 2000.

[139176]24842.

[139177]Henthorn, P.; Kiledjian, M.; Kadesch, T.: Two distinct tran-
scription

[139178]factors that bind the immunoglobulin enhancer mu-
E5/kappa-E2 motif. Science 247:

[139179]467-470, 1990.

[139180]24843.

[139181]Kamps, M. P.; Murre, C.; Sun, X.; Baltimore, D.: A new
homeobox

[139182]gene contributes the DNA binding domain of the t(1;19)
translocation

[139183]protein in pre-B ALL. Cell 60: 547-555, 1990.

[139184]24844.

[139185]Mellentin, J. D.; Murre, C.; Donlon, T. A.; McCaw, P. S.;
Smith,

[139186]S. D.; Carroll, A. J.; McDonald, M. E.; Baltimore, D.; Cleary,
M.

[139187]L.: The gene for enhancer binding proteins E12/E47 lies at
the t(1;19)

[139188]breakpoint in acute leukemias. Science 246: 379-382,
1989.

[139189]24845.

[139190]Murre, C.; McCaw, P. S.; Baltimore, D. :Cell 56: 777-783,
1989.

[139191]24846.

[139192]Nourse, J.; Mellentin, J. D.; Galili, N.; Wilkinson, J.; Stan-
bridge,

[139193]E.; Smith, S. D.; Cleary, M. L.: Chromosomal translocation
t(1;19)

[139194]results in synthesis of a homeobox fusion mRNA that
codes for a potential

[139195]chimeric transcription factor. Cell 60: 535-545, 1990.

[139196]24847.

[139197]Matsutani, A.; Koranyi, L.; Cox, N.; Permutt, M. A.: Poly-
morphisms

[139198]of GLUT2 and GLUT4 genes: use in evaluation of genetic
susceptibility

[139199]to NIDDM in Blacks. Diabetes 39: 1534-1542, 1990.

[139200]24848.

[139201]Moller, A. M.; Jensen, N. M.; Pildal, J.; Drivsholm, T.;
Borch-Johnsen,

[139202]K.; Urhammer, S. A.; Hansen, T.; Pedersen, O.: Studies of
genetic

[139203]variability of the glucose transporter 2 promoter in pa-
tients with

- [139204]type 2 diabetes mellitus. J. Clin. Endocr. Metab. 86:
2181–2186,
[139205]2001.
[139206]24849.
[139207]Mueckler, M.; Caruso, C.; Baldwin, S. A.; Panico, M.;
Blench,
[139208]J.; Morris, H. R.; Allard, W. J.; Lienhard, G. E.; Lodish, H. F.:
[139209]Sequence and structure of a human glucose transporter.
Science 229:
[139210]941–945, 1985.
[139211]24850.
[139212]Mueckler, M.; Kruse, M.; Strube, M.; Riggs, A. C.; Chiu, K.
C.;
[139213]Permutt, M. A.: A mutation in the Glut2 glucose trans-
porter gene
[139214]of a diabetic patient abolishes transport activity. J. Biol.
Chem. 269:
[139215]17765–17767, 1994.
[139216]24851.
[139217]Muller, D.; Santer, R.; Krawinkel, M.; Christiansen, B.;
Schaub,

- [139218]J.: Fanconi-Bickel syndrome presenting in neonatal screening for galactosemia. J. Inherit. Metab. Dis. 20: 607-608, 1997.
- [139219]24852.
- [139221]Oka, Y.; Asano, T.; Shibaskai, Y.; Lin, J.-L.; Tsukuda, K.; Katagiri, H.; Akanuma, Y.; Takaku, F.: C-terminal truncated glucose transporter is locked into an inward-facing form without transport activity. Nature 345: 550-553, 1990.
- [139222]24853.
- [139226]Orci, L.; Thorens, B.; Ravazzola, M.; Lodish, H. F.: Localization of the pancreatic beta cell glucose transporter to specific plasma membrane domains. Science 245: 295-297, 1989.
- [139227]24854.
- [139230]Permutt, M. A.; Koranyi, L.; Keller, K.; Lacy, P. E.; Scharp, D. W.; Mueckler, M.: Cloning and functional expression of a human

[139232]pancreatic islet glucose-transporter cDNA. Proc. Nat. Acad. Sci. 86:

[139233]8688-8692, 1989.

[139234]24855.

[139235]Sakamoto, O.; Ogawa, E.; Ohura, T.; Igarashi, Y.; Matsubara, Y.;

[139236]Narisawa, K.; Iinuma, K.: Mutation analysis of the GLUT2 gene in

[139237]patients with Fanconi-Bickel syndrome. Pediat. Res. 48: 586-589,

[139238]2000.

[139239]24856.

[139240]Sanjad, S. A.; Kaddoura, R. E.; Nazer, H. M.; Akhtar, M.; Sakati,

[139241]N. A.: Fanconi's syndrome with hepatorenal glycogenosis associated

[139242]with phosphorylase b kinase deficiency. Am. J. Dis. Child. 147:

[139243]957-959, 1993.

[139244]24857.

[139245]Santer, R.; Groth, S.; Kinner, M.; Dombrowski, A.; Berry, G.

T.;

[139246]Brodehl, J.; Leonard, J. V.; Moses, S.; Norgren, S.; Skovby, F.; Schneppenheim,

[139247]R.; Steinmann, B.; Schaub, J.: The mutation spectrum of the facilitative

[139248]glucose transporter gene SLC2A2 (GLUT2) in patients with Fanconi-Bickel

[139249]syndrome. Hum. Genet. 110: 21-29, 2002.

[139250]24858.

[139251]Santer, R.; Schneppenheim, R.; Dombrowski, A.; Gotze, H.; Steinmann,

[139252]B.; Schaub, J.: Fanconi-Bickel syndrome--a congenital defect of the

[139253]liver-type facilitative glucose transporter. J. Inherit. Metab. Dis. 21:

[139254]191-194, 1998.

[139255]24859.

[139256]Santer, R.; Schneppenheim, R.; Dombrowski, A.; Gotze, H.; Steinmann,

[139257]B.; Schaub, J.: Mutations in GLUT2, the gene for the liver-type glucose

[139258]transporter, in patients with Fanconi-Bickel syndrome.

Nature Genet. 17:

[139259]324–326, 1997. Note: Erratum: Nature Genet. 18: 298 only, 1998.

[139260]24860.

[139261]Steinmann, B.; Zeller, L.: Personal Communication. Zurich and

[139262]Santa Maria Calanca, Switzerland 8/2/1997.

[139263]24861.

[139264]Takeda, J.; Kayano, T.; Fukumoto, H.; Bell, G. I.: Organization

[139265]of the human GLUT2 (pancreatic beta-cell and hepatocyte) glucose transporter

[139266]gene. Diabetes 42: 773–777, 1993.

[139267]24862.

[139268]Tanizawa, Y.; Riggs, A. C.; Chiu, K. C.; Janssen, R. C.; Bell,

[139269]D. S. H.; Go, R. P. C.; Roseman, J. M.; Acton, R. T.; Permutt, M.

[139270]A.: Variability of the pancreatic islet beta cell/liver (GLUT 2)

[139271]glucose transporter gene in NIDDM patients. Diabetologia 37: 420–427,

[139272]1994.

[139273]24863.

[139274]Thorens, B.; Deriaz, N.; Bosco, D.; DeVos, A.; Pipeleers, D.;

[139275]Schuit, F.; Meda, P.; Porret, A.: Protein kinase A-
dependent phosphorylation

[139276]of GLUT2 in pancreatic beta cells. J. Biol. Chem. 271:
8075-8081,

[139277]1996.

[139278]24864.

[139279]Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.

[139280]M.; Kayano, T.; Shows, T. B.; Bell, G. I.: Assignment of
genes encoding

[139281]three human glucose transporter/transporter-like pro-
teins (GLUT4,

[139282]GLUT5 and GLUT6) to chromosomes 17, 1 and 5, respec-
tively. (Abstract) Cytogenet.

[139283]Cell Genet. 51: 997 only, 1989.

[139284]24865.

[139285]Gould, G. W.; Holman, G. D.: The glucose transporter fam-
ily: structure,

[139286]function and tissue-specific expression. Biochem. J. 295:
329-341,

[139287]1993.

[139288]24866.

[139289]Hauguel-de Mouzon, S.; Challier, J. C.; Kacemi, A.;
Cauzac, M.;

[139290]Malek, A.; Girard, J.: The GLUT3 glucose transporter iso-
form is differentially

[139291]expressed within human placental cell types. J. Clin. En-
docr. Metab. 82:

[139292]2689-2694, 1997.

[139293]24867.

[139294]Kayano, T.; Burant, C. F.; Fukumoto, H.; Gould, G. W.; Fan,
Y.

[139295]S.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Seino, S.; Bell, G.
I.

[139296]: Human facilitative glucose transporters: isolation, func-
tional characterization,

[139297]and gene localization of cDNAs encoding an isoform
(GLUT5) expressed

[139298]in small intestine, kidney, muscle, and adipose tissue and
an unusual

- [139299]glucose transporter pseudogene-like sequence (GLUT6). J. Biol. Chem. 265: 13276-13282, 1990.
- [139300]24868.
- [139301]Kayano, T.; Fukumoto, H.; Eddy, R. L.; Fan, Y.-S.; Byers, M. G.;
- [139302]Shows, T. B.; Bell, G. I.: Evidence for a family of human glucose transporter-like proteins: sequence and gene localization of a protein expressed in fetal skeletal muscle and other tissues. J. Biol. Chem. 263: 15245-15248, 1988.
- [139303]24869.
- [139304]Gitelman, B. J.; Tomkins, D. J.; Partington, M. W.; Roberts, M.;
- [139305]H.; Simpson, N. E.: Gene dosage studies of glutamic oxaloacetic transaminase (GOT) and hexokinase (HK) in two patients with possible partial trisomy 10q. (Abstract) Am. J. Hum. Genet. 32: 41A only, 1980.

[139312]24870.

[139313]Junien, C.; Despoisse, S.; Turleau, C.; de Grouchy, J.;
Bucher,

[139314]T.; Fundele, R.: Assignment of phosphoglycerate mutase
(PGAMA) to

[139315]human chromosome 10: regional mapping of GOT1 and
PGAMA to subbands

[139316]10q26.1 (or q25.3). Ann. Genet. 25: 25-27, 1982.

[139317]24871.

[139318]O'Brien, S. J.; Bonner, T. I.; Cohen, M.; O'Connell, C.; Nash,

[139319]W. G.: Mapping of an endogenous retroviral sequence to
human chromosome

[139320]18. Nature 303: 74-77, 1983.

[139321]24872.

[139322]Overhauser, J.; Mewar, R.; Rojas, K.; Lia, K.; Kline, A. D.;
Silverman,

[139323]G. A.: STS map of genes and anonymous DNA fragments
on human chromosome

[139324]18 using a panel of somatic cell hybrids. Genomics 15:
387-391,

[139325]1993.

[139326]24873.

[139327]Yunis, J. J.; Oken, M. M.; Kaplan, M. E.; Ensrud, K. M.;
Howe,

[139328]R. R.; Theologides, A.: Distinctive chromosomal abnormalities in

[139329]histologic subtypes of non-Hodgkin's lymphoma. New
Eng. J. Med. 307:

[139330]1231-1236, 1982.

[139331]24874.

[139332]Abdalla, S. A.; Pece-Barbara, N.; Vera, S.; Tapia, E.; Paez,
E.;

[139333]Bernabeu, C.; Letarte, M.: Analysis of ALK-1 and endoglin
in newborns

[139334]from families with hereditary hemorrhagic telangiectasia
type 2. Hum.

[139335]Molec. Genet. 9: 1227-1237, 2000.

[139336]24875.

[139337]Bellon, T.; Corbi, A.; Lastres, P.; Cales, C.; Cebrian, M.;
Vera,

[139338]S.; Cheifetz, S.; Massague, J.; Letarte, M.; Bernabeu, C.:
Identification

- [139339]and expression of two forms of the human transforming growth factor- β -binding
- [139340]protein endoglin with distinct cytoplasmic regions. *Europ. J. Immun.* 23:
- [139341]2340–2345, 1993.
- [139342]24876.
- [139343]Bourdeau, A.; Dumont, D. J.; Letarte, M.: A murine model of hereditary
- [139344]hemorrhagic telangiectasia. *J. Clin. Invest.* 104: 1343–1351, 1999.
- [139345]24877.
- [139346]Fernandez–Ruiz, E.; St. Jacques, S.; Bellon, T.; Letarte, M.; Bernabeu,
- [139347]C.: Assignment of the human endoglin gene (END) to 9q34–qter. *Cytogenet. Cell Genet.* 64: 204–207, 1993.
- [139348]24878.
- [139349]24878.
- [139350]Gallione, C. J.; Klaus, D. J.; Yeh, E. Y.; Stenzel, T. T.; Xue,
- [139351]Y.; Anthony, K. B.; McAllister, K. A.; Baldwin, M. A.; Berg, J. N.;
- [139352]Lux, A.; Smith, J. D.; Vary, C. P. H.; Craigen, W. J.; Wester–

mann,

[139353]C. J. J.; Warner, M. L.; Miller, Y. E.; Jackson, C. E.;

Guttmacher,

[139354]A. E.; Marchuk, D. A.: Mutation and expression analysis of
the endoglin

[139355]gene in hereditary hemorrhagic telangiectasia reveals null
alleles. Hum.

[139356]Mutat. 11: 286–294, 1998.

[139357]24879.

[139358]Gallione, C. J.; Scheessele, E. A.; Reinhardt, D.; Duits, A. J.;

[139359]Berg, J. N.; Westermann, C. J. J.; Marchuk, D. A.: Two com-
mon endoglin

[139360]mutations in families with hereditary hemorrhagic telang-
iectasia in

[139361]the Netherlands Antilles: evidence for a founder effect.
Hum. Genet. 107:

[139362]40–44, 2000.

[139363]24880.

[139364]Gougos, A.; Letarte, M.: Primary structure of endoglin, an
RGD-containing

[139365]glycoprotein of human endothelial cells. J. Biol. Chem.

265: 8361–8364,
[139366]1990.

[139367]24881.

[139368]Kozak, M.: Context effects and inefficient initiation at
non-AUG
[139369]codons in eucaryotic cell-free translation systems. Molec.
Cell.
[139370]Biol. 9: 5073–5080, 1989.

[139371]24882.

[139372]Li, D. Y.; Sorensen, L. K.; Brooke, B. S.; Urness, L. D.;
Davis,
[139373]E. C.; Taylor, D. G.; Boak, B. B.; Wendel, D. P.: Defective
angiogenesis
[139374]in mice lacking endoglin. Science 284: 1534–1537, 1999.

[139375]24883.

[139376]Lux, A.; Gallione, C. J.; Marchuk, D. A.: Expression analysis
[139377]of endoglin missense and truncation mutations: insights
into protein
[139378]structure and disease mechanisms. Hum. Molec. Genet. 9:
745–755,
[139379]2000.

[139380]24884.

[139381]McAllister, K. A.; Grogg, K. M.; Johnson, D. W.; Gallione, C.

[139382]J.; Baldwin, M. A.; Jackson, C. E.; Helmbold, E. A.; Markel,
D. S.;

[139383]McKinnon, W. C.; Murrell, J.; McCormick, M. K.; Pericak-
Vance, M.

[139384]A.; Heutink, P.; Oostra, B. A.; Haitjema, T.; Westerman, C.
J. J.;

[139385]Porteous, M. E.; Guttmacher, A. E.; Letarte, M.; Marchuk,
D. A.:

[139386]Endoglin, a TGF- β binding protein of endothelial cells,
is the

[139387]gene for hereditary haemorrhagic telangiectasia type 1.
Nature Genet. 8:

[139388]345-351, 1994.

[139389]24885.

[139390]Paquet, M.-E.; Pece-Barbara, N.; Vera, S.; Cymerman, U.;
Karabegovic,

[139391]A.; Shovlin, C.; Letarte, M.: Analysis of several endoglin
mutants

[139392]reveals no endogenous mature or secreted protein capa-
ble of interfering

[139393]with normal endoglin function. Hum. Molec. Genet. 10:
1347–1357,

[139394]2001.

[139395]24886.

[139396]Pece, N.; Vera, S.; Cymerman, U.; White, R. I., Jr.; Wrana, J.

[139397]L.; Letarte, M.: Mutant endoglin in hereditary hemorrhagic
telangiectasia

[139398]type 1 is transiently expressed intracellularly and is not a
dominant

[139399]negative. J. Clin. Invest. 100: 2568–2579, 1997.

[139400]24887.

[139401]Pece-Barbara, N.; Cymerman, U.; Vera, S.; Marchuk, D. A.;
Letarte,

[139402]M.: Expression analysis of four endoglin missense muta-
tions suggests

[139403]that haploinsufficiency is the predominant mechanism for
hereditary

[139404]hemorrhagic telangiectasia type 1. Hum. Molec. Genet. 8:
2171–2181,

[139405]1999.

[139406]24888.

[139407]Pilz, A.; Woodward, K.; Peters, J.; Povey, S.; Abbott, C.:

Comparative

[139408]mapping of 38 human chromosome 9 loci in the laboratory mouse. (Abstract) Ann.

[139409]Hum. Genet. 58: 231-232, 1994.

[139410]24889.

[139411]Qureshi, S. T.; Gros, P.; Letarte, M.; Malo, D.: The murine endoglin

[139412]gene (Eng) maps to chromosome 2. Genomics 26: 165-166, 1995.

[139413]24890.

[139414]Abbott, C.; Jackson, I. J.; Carritt, B.; Povey, S.: The human

[139415]homolog of the mouse brown gene maps to the short arm of chromosome

[139416]9 and extends the known region of homology with mouse chromosome 4. Genomics 11:

[139417]471-473, 1991.

[139418]24891.

[139419]Box, N. F.; Wyeth, J. R.; Mayne, C. J.; O'Gorman, L. E.; Martin,

[139420]N. G.; Sturm, R. A.: Complete sequence and polymorphism

study of

[139421]the human TYRP1 gene encoding tyrosinase-related protein 1. Mammalian

[139422]Genome 9: 50-53, 1998.

[139423]24892.

[139424]Chintamaneni, C. D.; Ramsay, M.; Colman, M.-A.; Fox, M. F.; Pickard,

[139425]R. T.; Kwon, B. S.: Mapping the human CAS2 gene, the homologue of

[139426]the mouse brown (b) locus, to human chromosome 9p22-pter. Biochem.

[139427]Biophys. Res. Commun. 178: 227-235, 1991.

[139428]24893.

[139429]Cohen, T.; Muller, R. M.; Tomita, Y.; Shibahara, S.: Nucleotide

[139430]sequence of the cDNA encoding human tyrosinase-related protein. Nucleic

[139431]Acids Res. 18: 2807-2808, 1990.

[139432]24894.

[139433]Halaban, R.; Moellmann, G.: Murine and human b locus pigmentation

- [139434]genes encode a glycoprotein (gp75) with catalase activity.
Proc.
- [139435]Nat. Acad. Sci. 87: 4809–4813, 1990.
- [139436]24895.
- [139437]Bode–Lesniewska, B.; Dours–Zimmermann, M. T.; Oder–
matt, B. F.;
- [139438]Briner, J.; Heitz, P. U.; Zimmermann, D. R.: Distribution of
the
- [139439]large aggregating proteoglycan versican in adult human
tissues. J.
- [139440]Histochem. Cytochem. 44: 303–312, 1996.
- [139441]24896.
- [139442]Dours–Zimmermann, M. T.; Zimmermann, D. R.: A novel
glycosaminoglycan
- [139443]attachment domain identified in two alternative splice
variants of
- [139444]human versican. J. Biol. Chem. 269: 32992–32998, 1994.
- [139445]24897.
- [139446]Iozzo, R. V.; Naso, M. F.; Cannizzaro, L. A.; Wasmuth, J. J.;
McPherson,
- [139447]J. D.: Mapping of the versican proteoglycan gene (CSPG2)

to the long

[139448]arm of human chromosome 5 (5q12–5q14). Genomics 14:
845–851, 1992.

[139449]24898.

[139450]Kjellen, L.; Lindahl, U.: Proteoglycans: structures and in-
teractions. Annu.

[139451]Rev. Biochem. 60: 443–475, 1991.

[139452]24899.

[139453]Naso, M. F.; Morgan, J. L.; Buchberg, A. M.; Siracusa, L. D.;
Iozzo,

[139454]R. V.: Expression pattern and mapping of the murine ver-
sican gene

[139455](Cspg2) to chromosome 13. Genomics 29: 297–300,
1995.

[139456]24900.

[139457]Naso, M. F.; Zimmermann, D. R.; Iozzo, R. V.: Characteri-
zation

[139458]of the complete genomic structure of the human versican
gene and functional

[139459]analysis of its promoter. J. Biol. Chem. 269:
32999–33008, 1994.

[139460]24901.

[139461]Zako, M.; Shinomura, T.; Ujita, M.; Ito, K.; Kimata, K.: Ex-
pression
[139462]of PG-M(V3), an alternatively spliced form of PG-M with-
out a chondroitin
[139463]sulfate attachment region in mouse and human tissues. J.
Biol. Chem. 270:
[139464]3914-3918, 1995.
[139465]24902.

[139466]Zimmermann, D. R.; Ruoslahti, E.: Multiple domains of the
large
[139467]fibroblast proteoglycan, versican. EMBO J. 8: 2975-2981,
1989.
[139468]24903.

[139469]Grunder, S.; Geisler, H.-S.; Rainer, S.; Fink, J. K.: Acid-
sensing
[139470]ion channel (ASIC) 4 gene: physical mapping, genomic or-
ganisation,
[139471]and evaluation as a candidate for paroxysmal dystonia.
Europ. J.
[139472]Hum. Genet. 9: 672-676, 2001.
[139473]24904.

[139474] de Vries, W. N.; de Wet, W. J.: The molecular defect in an autosomal

[139475] dominant form of osteogenesis imperfecta: synthesis of type I procollagen

[139476] containing cysteine in the triple-helical domain of pro-alpha-1(I)

[139477] chains. J. Biol. Chem. 261: 9056-9064, 1986.

[139478] 24905.

[139479] de Vries, W. N.; de Wet, W. J.: Development and diseases of cartilage

[139480] and bone matrix. In: Sen, A.; Thornhill, T.: UCLA Symposia on Molecular

[139481] and Cellular Biology--New Series. New York: Alan R. Liss (pub.)

[139482] 46: 1987.

[139483] 24906.

[139484] Deak, S. B.; Scholz, P. M.; Amenta, P. S.; Constantinou, C. D.;

[139485] Levi-Minzi, S. A.; Gonzalez-Lavin, L.; Mackenzie, J. W.: The substitution

[139486] of arginine for glycine 85 of the alpha-1(I) procollagen chain results

[139487]in mild osteogenesis imperfecta: the mutation provides direct evidence

[139488]for three discrete domains of cooperative melting of intact type I

[139489]collagen. J. Biol. Chem. 266: 21827–21832, 1991.

[139490]24907.

[139491]Diesel, A. J.; Schumacher, A. M.; Flavell, R. A.: A Hind III

[139492]restriction site polymorphism in the human collagen alpha-1(I)-like

[139493]gene on chromosome no. 7. Hum. Genet. 62: 175–176, 1982.

[139494]24908.

[139495]Forlino, A.; Zolezzi, F.; Valli, M.; Pignatti, P. F.; Cetta, G.;

[139496]Brunelli, P. C.; Mottes, M.: Severe (type III) osteogenesis imperfecta

[139497]due to glycine substitutions in the central domain of the collagen

[139498]triple helix. Hum. Molec. Genet. 3: 2201–2206, 1994.

[139499]24909.

[139500]Huerre, C.; Junien, C.; Weil, D.; Chu, M.-L.; Morabito, M.;
Van

- [139501] Cong, N.; Myers, J. C.; Foubert, C.; Gross, M.-S.; Prockop, D. J.;
- [139502] Boue, A.; Kaplan, J.-C.; de la Chapelle, A.; Ramirez, F.: Human type
- [139503] procollagen genes are located on different chromosomes. Proc. Nat.
- [139504] Acad. Sci. 79: 6627-6630, 1982.
- [139505] 24910.
- [139506] Izant, J. G.; Weintraub, H.: Inhibition of thymidine kinase gene
- [139507] expression by anti-sense RNA: a molecular approach to genetic analysis. Cell 36:
- [139508] 1007-1015, 1984.
- [139509] 24911.
- [139510] Khillan, J. S.; Li, S.-W.; Prockop, D. J.: Partial rescue of
- [139511] a lethal phenotype of fragile bones in transgenic mice with a chimeric
- [139512] antisense gene directed against a mutated collagen gene. Proc. Nat.
- [139513] Acad. Sci. 91: 6298-6302, 1994.
- [139514] 24912.

[139515]Korkko, J.; Ala-Kokko, L.; De Paepe, A.; Nuytinck, L.; Earley,

[139516]J.; Prockop, D. J.: Analysis of the COL1A1 and COL1A2 genes by PCR

[139517]amplification and scanning by conformation-sensitive gel electrophoresis

[139518]identifies only COL1A1 mutations in 15 patients with osteogenesis

[139519]imperfecta type I: identification of common sequences of null-allele

[139520]mutations. Am. J. Hum. Genet. 62: 98-110, 1998.

[139521]24913.

[139522]Korkko, J.; Kuivaniemi, H.; Paassilta, P.; Zhuang, J.; Tromp,

[139523]G.; DePaepe, A.; Prockop, D. J.; Ala-Kokko, L.: Two new recurrent

[139524]nucleotide mutations in the COL1A1 gene in four patients with osteogenesis

[139525]imperfecta: about one-fifth are recurrent. Hum. Mutat. 9: 148-156,

[139526]1997.

[139527]24914.

- [139528]Lazarides, E.; Lukens, L. N.: Collagen synthesis on polysomes
[139529]in vivo and in vitro. Nature N.B. 232: 37-40, 1971.
[139530]24915.
- [139531]Lund, A. M.; Skovby, F.; Schwartz, M.: (G586V) substitutions
[139532]in the alpha-1 and alpha-2 chains of collagen I: effect of alpha-chain
[139533]stoichiometry on the phenotype of osteogenesis imperfecta? Hum. Mutat. 9:
[139534]431-436, 1997.
[139535]24916.
- [139536]Mackay, K.; Lund, A. M.; Raghunath, M.; Steinmann, B.; Dagleish,
[139537]R.: SSCP detection of a gly565-to-val substitution in the pro-alpha-1(I)
[139538]collagen chain resulting in osteogenesis imperfecta type II. Hum.
[139539]Genet. 91: 439-444, 1993.
[139540]24917.
- [139541]Marini, J. C.; Grange, D. K.; Gottesman, G. S.; Lewis, M. B.;

[139542]Koeplin, D. A.: Osteogenesis imperfecta type IV: detection of a point

[139543]mutation in one alpha-1(I) collagen allele (COL1A1) by RNA/RNA hybrid

[139544]analysis. J. Biol. Chem. 264: 11893-11900, 1989.

[139545]24918.

[139546]Marini, J. C.; Lewis, M. B.; Chen, K.: Moderately severe osteogenesis

[139547]imperfecta associated with substitutions of serine for glycine in

[139548]the alpha-1(I) chain of type I collagen. Am. J. Med. Genet. 45:

[139549]241-245, 1993.

[139550]24919.

[139551]Mayer, S. A.; Rubin, B. S.; Starman, B. J.; Byers, P. H.: Spontaneous

[139552]multivessel cervical artery dissection in a patient with a substitution

[139553]of alanine for glycine (G13A) in the alpha-1(I) chain of type I collagen. Neurology 47:

[139554]552-556, 1996.

[139555]24920.

[139556]Mizuno, T.; Chou, M.; Inouye, M.: A unique mechanism
regulating

[139557]gene expression: translational inhibition by a complemen-
tary RNA transcript

[139558](micRNA). Proc. Nat. Acad. Sci. 81: 1966–1970, 1984.

[139559]24921.

[139560]Mottes, M.; Gomez Lira, M. M.; Valli, M.; Scarano, G.;
Lonardo,

[139561]F.; Forlino, A.; Cetta, G.; Pignatti, P. F.: Paternal mosaicism
for

[139562]a COL1A1 dominant mutation (alpha-1 ser-415) causes
recurrent osteogenesis

[139563]imperfecta. Hum. Mutat. 2: 196–204, 1993.

[139564]24922.

[139565]Mottes, M.; Sangalli, A.; Valli, M.; Gomez Lira, M.; Tenni,
R.;

[139566]Buttitta, P.; Pignatti, P. F.; Cetta, G.: Mild dominant osteo-
genesis

[139567]imperfecta with intrafamilial variability: the cause is a ser-
ine for

[139568]glycine alpha-1(I) 901 substitution in a type-I collagen gene. Hum.

[139569]Genet. 89: 480-484, 1992.

[139570]24923.

[139571]Munke, M.; Harbers, K.; Jaenisch, R.; Francke, U.: Chromosomal

[139572]mapping of four different integration sites of Moloney murine leukemia

[139573]virus including the locus for alpha-1(I) collagen in mouse. Cytogenet.

[139574]Cell Genet. 43: 140-149, 1986.

[139575]24924.

[139576]Namikawa, C.; Suzumori, K.; Fukushima, Y.; Sasaki, M.; Hata, A.

[139577]: Recurrence of osteogenesis imperfecta because of paternal mosaicism:

[139578]gly862-to-ser substitution in a type I collagen gene (COL1A1). Hum.

[139579]Genet. 95: 666-670, 1995.

[139580]24925.

[139581]Nicholls, A. C.; Oliver, J.; Renouf, D. V.; Keston, M.; Pope,

[139582]F. M.: Substitution of cysteine for glycine at residue 415 of
one

[139583]allele of the alpha-1(I) chain of type I procollagen in type
III/IV

[139584]osteogenesis imperfecta. J. Med. Genet. 28: 757-764,
1991.

[139585]24926.

[139586]Steinmann, B.; Rao, V. H.; Vogel, A.; Bruckner, P.; Gitzel-
mann,

[139587]R.; Byers, P. H.: Cysteine in the triple-helical domain of
one allelic

[139588]product of the alpha-1(I) gene of type I collagen produces
a lethal

[139589]form of osteogenesis imperfecta. J. Biol. Chem. 250:
11129-11139,

[139590]1984.

[139591]24927.

[139592]Albert, L. J.; Inman, R. D.: Molecular mimicry and autoim-
munity. New

[139593]Eng. J. Med. 341: 2068-2074, 1999.

[139594]24928.

[139595]Asada, H.; Kawamura, Y.; Maruyama, K.; Kume, H.; Ding, R.-G.; Ji,

[139596]F. Y.; Kanbara, N.; Kuzume, H.; Sanbo, M.; Yagi, T.; Obata, K.: Mice

[139597]lacking the 65 kDa isoform of glutamic acid decarboxylase (GAD65)

[139598]maintain normal levels of GAD67 and GABA in their brains but are susceptible

[139599]to seizures. Biochem. Biophys. Res. Commun. 229: 891–895, 1996.

[139600]24929.

[139601]Baekkeskov, S.; Aanstoot, H.-J.; Christgau, S.; Reetz, A.; Solimena,

[139602]M.; Cascalho, M.; Folli, F.; Richter-Olesen, H.; Camilli, P.-D.:

[139603]Identification of the 64K autoantigen in insulin-dependent diabetes

[139604]as the GABA-synthesizing enzyme glutamic acid decarboxylase. Nature 347:

[139605]151–156, 1990.

[139606]24930.

[139607]Bu, D.-F.; Erlander, M. G.; Hitz, B. C.; Tillakaratne, N. J. K.;

[139608]Kaufman, D. L.; Wagner-McPherson, C. B.; Evans, G. A.;
Tobin, A. J.

[139609]: Two human glutamate decarboxylases, 65-kDa GAD and
67-kDa GAD, are

[139610]each encoded by a single gene. Proc. Nat. Acad. Sci. 89:
2115-2119,

[139611]1992.

[139612]24931.

[139613]Cram, D. S.; Barnett, L. D.; Joseph, J. L.; Harrison, L. C.:
Cloning

[139614]and partial nucleotide sequence of human glutamic acid
decarboxylase

[139615]cDNA from brain and pancreatic islets. Biochem. Biophys.
Res. Commun. 176:

[139616]1239-1244, 1991.

[139617]24932.

[139618]De Aizpurua, H. J.; Wilson, Y. M.; Harrison, L. C.: Glutamic
acid

- [139619]decarboxylase autoantibodies in preclinical insulin-dependent diabetes. Proc.
- [139620]Nat. Acad. Sci. 89: 9841-9845, 1992.
- [139621]24933.
- [139622]Edelhoff, S.; Grubin, C. E.; Karlsen, A. E.; Adler, D. A.; Foster,
- [139623]D.; Distèche, C. M.; Lernmark, A.: Mapping of glutamic acid decarboxylase
- [139624](GAD) genes. Genomics 17: 93-97, 1993.
- [139625]24934.
- [139626]Jones, D. B.; Armstrong, N. W.: Cocksackie virus and diabetes revisited.
- [139627](Letter) Nature Med. 1: 284 only, 1995.
- [139628]24935.
- [139629]Karlsen, A. E.; Hagopian, W. A.; Grubin, C. E.; Dube, S.; Distèche,
- [139630]C. M.; Adler, D. A.; Barmeier, H.; Mathewes, S.; Grant, F. J.; Foster,
- [139631]D.; Lernmark, A.: Cloning and primary structure of a human islet
- [139632]isoform of glutamic acid decarboxylase from chromosome

10. Proc.

[139633]Nat. Acad. Sci. 88: 8337–8341, 1991.

[139634]24936.

[139635]Kash, S. F.; Johnson, R. S.; Tecott, L. H.; Noebels, J. L.;
Mayfield,

[139636]R. D.; Hanahan, D.; Baekkeskov, S.: Epilepsy in mice deficient in

[139637]the 65-kDa isoform of glutamic acid decarboxylase. Proc.
Nat. Acad.

[139638]Sci. 94: 14060–14065, 1997.

[139639]24937.

[139640]Kaufman, D. L.; Erlander, M. G.; Clare-Salzler, M.; Atkinson,
son,

[139641]M. A.; Maclaren, N. K.; Tobin, A. J.: Autoimmunity to two
forms of

[139642]glutamate decarboxylase in insulin-dependent diabetes
mellitus. J.

[139643]Clin. Invest. 89: 283–292, 1992.

[139644]24938.

[139645]Lohmann, T.; Hawa, M.; Leslie, R. D. G.; Lane, R.; Picard, J.;

[139646]Londei, M.: Immune reactivity to glutamic acid decarboxy-

lase 65 in

[139647]stiff-man syndrome and type 1 diabetes mellitus. Lancet
356: 31-35,

[139648]2000.

[139649]24939.

[139650]Richter, W.; Endl, J.; Eiermann, T. H.; Brandt, M.; Kientsch-En-
gel,

[139651]R.; Thivolet, C.; Jungfer, H.; Scherbaum, W. A.: Human
monoclonal

[139652]islet cell antibodies from a patient with insulin-dependent
diabetes

[139653]mellitus reveal glutamate decarboxylase as the target
antigen. Proc.

[139654]Nat. Acad. Sci. 89: 8467-8471, 1992.

[139655]24940.

[139656]Tian, J.; Lehmann, P. V.; Kaufman, D. L.: T cell cross-
reactivity

[139657]between coxsackievirus and glutamate decarboxylase is
associated with

[139658]a murine diabetes susceptibility allele. J. Exp. Med. 180:
1979-1984,

[139659]1994.

[139660]24941.

[139661]Chiang, S.-H.; Baumann, C. A.; Kanzaki, M.; Thurmond, D. C.; Watson,

[139662]R. T.; Neudauer, C. L.; Macara, I. G.; Pessin, J. E.; Saltiel, A.

[139663]R.: Insulin-stimulated GLUT4 translocation requires the CAP-dependent

[139664]activation of TC10. *Nature* 410: 944-948, 2001.

[139665]24942.

[139666]Ribon, V.; Printen, J. A.; Hoffman, N. G.; Kay, B. K.; Saltiel,

[139667]A. R.: A novel, multifunctional c-Cbl binding protein in insulin

[139668]receptor signaling in 3T3-L1 adipocytes. *Molec. Cell. Biol.* 18:

[139669]872-879, 1998.

[139670]24943.

[139671]Astrin, K. H.; Arredondo-Vega, F. X.; Desnick, R. J.; Smith, M.

[139672]: Assignment of the gene for cytosolic alanine amino-transferase (AAT1)

[139673]to human chromosome 8. *Ann. Hum. Genet.* 46: 125-133, 1982.

[139674]24944.

[139675]Chen, S.-H.; Giblett, E. R.: Polymorphism of soluble glutamic-pyruvic

[139676]transaminase: a new genetic marker in man. Science 173: 148-149,

[139677]1971.

[139678]24945.

[139679]Cook, P. J. L.; Jeremiah, S. J.; Buckton, K. E.: Exclusion mapping

[139680]of GPT. (Abstract) Cytogenet. Cell Genet. 32: 258 only, 1982.

[139681]24946.

[139682]Mombaerts, P.; Wang, F.; Dulac, C.; Vassar, R.; Chao, S. K.; Nemes,

[139683]A.; Mendelsohn, M.; Edmondson, J.; Axel, R.: The molecular biology

[139684]of olfactory perception. Cold Spring Harbor Symp. Quant. Biol. 61:

[139685]135-145, 1996.

[139686]24947.

[139687]Reed, R. R.: Genetic approaches to mammalian olfaction.

Cold

- [139688]Spring Harbor Symp. Quant. Biol. 61: 165–172, 1996.
- [139689]24948.
- [139690]Ronnett, G. V.; Snyder, S. H.: Molecular messengers of olfaction. Trends
- [139691]Neurosci. 15: 508–513, 1992.
- [139692]24949.
- [139693]Schwab, S. G.; Hallmayer, J.; Lerer, B.; Albus, M.; Borrmann, M.;
- [139694]Honig, S.; Strauss, M.; Segman, R.; Lichtermann, D.; Knapp, M.; Trixler,
- [139695]M.; Maier, W.; Wildenauer, D. B.: Support for a chromosome 18p locus
- [139696]conferring susceptibility to functional psychoses in families with
- [139697]schizophrenia, by association and linkage analysis. Am. J. Hum. Genet. 63:
- [139698]1139–1152, 1998.
- [139699]24950.
- [139700]Davignon, I.; Barnard, M.; Gavrilova, O.; Sweet, K.; Wilkie, T.

[139701]M.: Gene structure of murine Gna11 and Gna15: tandemly duplicated

[139702]Gq class G protein alpha subunit genes. Genomics 31: 359–366, 1996.

[139703]24951.

[139704]Jiang, M.; Pandey, S.; Tran, V. T.; Fong, H. K.: Guanine nucleotide-binding

[139705]regulatory proteins in retinal pigment epithelial cells. Proc. Nat.

[139706]Acad. Sci. 88: 3907–3911, 1991.

[139707]24952.

[139708]Offermanns, S.; Zhao, L.-P.; Gohla, A.; Sarosi, I.; Simon, M. I.;

[139709]Wilkie, T. M.: Embryonic cardiomyocyte hypoplasia and craniofacial

[139710]defects in G-alpha-q/G-alpha-11-mutant mice. EMBO J. 17: 4304–4312,

[139711]1998.

[139712]24953.

[139713]Strathmann, M. P.; Simon, M. I.: G-alpha-12 and G-alpha-13 subunits

[139714]define a fourth class of G protein alpha subunits. Proc.
Nat. Acad.

[139715]Sci. 88: 5582–5586, 1991.

[139716]24954.

[139717]Ahmed, S. F.; Barr, D. G. D.; Bonthron, D. T.: GNAS1 muta-
tions

[139718]and progressive osseous heteroplasia. (Letter) New Eng. J.
Med. 346:

[139719]1669–1670, 2002.

[139720]24955.

[139721]Ahmed, S. F.; Dixon, P. H.; Bonthron, D. T.; Stirling, H. F.;
Barr,

[139722]D. G. D.; Kelnar, C. J. H.; Thakker, R. V.: GNAS1 mutational
analysis

[139723]in pseudohypoparathyroidism. Clin. Endocr. 49: 525–531,
1998.

[139724]24956.

[139725]Ahrens, W.; Hiort, O.; Staedt, P.; Kirschner, T.; Marschke,
C.;

[139726]Kruse, K.: Analysis of the GNAS1 gene in Albright's hered-
itary osteodystrophy. J.

[139727]Clin. Endocr. Metab. 86: 4630–4634, 2001.

[139728]24957.

[139729]Aldred, M. A.; Trembath, R. C.: Activating and inactivating mutations

[139730]in the human GNAS1 gene. Hum. Mutat. 16: 183–189, 2000.

[139731]24958.

[139732]Ashley, P. L.; Ellison, J.; Sullivan, K. A.; Bourne, H. R.; Cox,

[139733]D. R.: Chromosomal assignment of the murine Gi and Gs genes. (Abstract) Am.

[139734]J. Hum. Genet. 41: A155 only, 1987.

[139735]24959.

[139736]Ballare, E.; Mantovani, S.; Lania, A.; Di Blasio, A. M.; Vallar,

[139737]L.; Spada, A.: Activating mutations of the GS- α gene are associated

[139738]with low levels of GS- α protein in growth hormone-secreting tumors. J.

[139739]Clin. Endocr. Metab. 83: 4386–4390, 1998.

[139740]24960.

[139741]Bastepe, M.; Pincus, J. E.; Sugimoto, T.; Tojo, K. Kanatani,

M.;

[139742]Azuma, Y.; Kruse, K.; Rosenbloom, A. L.; Koshiyama, H.; Juppner, H.

[139743]; Positional dissociation between the genetic mutation responsible

[139744]for pseudohypoparathyroidism type 1b and the associated methylation

[139745]defect at exon A/B: evidence for a long-range regulatory element within

[139746]the imprinted GNAS1 locus. Hum. Molec. Genet. 10: 1231-1241, 2001.

[139747]24961.

[139748]Bastepe, M.; Juppner, H.: GNAS1 mutations and progressive osseous

[139749]heteroplasia. (Letter) New Eng. J. Med. 346: 1671 only, 2002.

[139750]24962.

[139751]Bastepe, M.; Lane, A. H.; Juppner, H.: Parental uniparental isodisomy

[139752]of chromosome 20q--and the resulting changes in GNAS1 methylation--as

[139753]a plausible cause of pseudohypoparathyroidism. Am. J.

Hum. Genet. 68:

[139754]1283–1289, 2001.

[139755]24963.

[139756]Bianco, P.; Riminucci, M.; Majolagbe, A.; Kuznetsov, S. A.;
Collins,

[139757]M. T.; Mankani, M. H.; Corsi, A.; Bone, H. G.; Wientroub,
S.; Spiegel,

[139758]A. M.; Fisher, L. W.; Robey, P. G.: Mutations of the GNAS1
gene,

[139759]stromal cell dysfunction, and osteomalacic changes in
non-McCune-Albright

[139760]fibrous dysplasia of bone. J. Bone Miner. Res. 15:
120–128, 2000.

[139761]24964.

[139762]Billestrup, N.; Swanson, L. W.; Vale, W.: Growth hormone–
releasing

[139763]factor stimulates proliferation of somatotrophs in vitro.
Proc. Nat.

[139764]Acad. Sci. 83: 6854–6857, 1986.

[139765]24965.

[139766]Candelieri, G. A.; Glorieux, F. H.; Prud'Homme, J.;

St.-Arnaud,

[139767]R.: Increased expression of the c-fos proto-oncogene in bone from

[139768]patients with fibrous dysplasia. New Eng. J. Med. 332: 1546-1551,

[139769]1995.

[139770]24966.

[139771]Candelieri, G. A.; Roughley, P. J.; Glorieux, F. H.: Polymerase

[139772]chain reaction-based technique for the selective enrichment and analysis

[139773]of mosaic arg201 mutations in G alpha s from patients with fibrous

[139774]dysplasia of bone. Bone 21: 201-206, 1997.

[139775]24967.

[139776]Carel, J. C.; Le Stunff, C.; Condamine, L.; Mallet, E.; Chaus-sain,

[139777]J. L.; Adnot, P.; Garabedian, M.; Bougneres, P.: Resistance to the

[139778]lipolytic action of epinephrine: a new feature of protein GS deficiency. J.

[139779]Clin. Endocr. Metab. 84: 4127-4131, 1999.

[139780]24968.

[139781]Carter, A.; Bardin, C.; Collins, R.; Simons, C.; Bray, P.;
Spiegel,

[139782]A.: Reduced expression of multiple forms of the alpha
subunit of

[139783]the stimulatory GTP-binding protein in pseudohy-
poparathyroidism type

[139784]la. Proc. Nat. Acad. Sci. 84: 7266-7269, 1987.

[139785]24969.

[139786]Cattanach, B. M.; Kirk, M.: Differential activity of mater-
nally

[139787]and paternally derived chromosome regions in mice. Na-
ture 315: 496-498,

[139788]1985.

[139789]24970.

[139790]Coutant, R.; Lumbroso, S.; Rey, R.; Lahlou, N.; Venara, M.;
Rouleau,

[139791]S.; Sultan, C.; Limal, J.-M.: Macroorchidism due to au-
tonomous hyperfunction

[139792]of Sertoli cells and GS-alpha gene mutation: an unusual
expression

[139793]of McCune–Albright syndrome in a prepubertal boy. J.
Clin. Endocr.

[139794]Metab. 86: 1778–1781, 2001.

[139795]24971.

[139796]Kim, H.–S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T.,
[139797]Jr.; Rosenfeld, R. G.: Identification of a family of low–
affinity

[139798]insulin–like growth factor binding proteins (IGFBPs): char–
acterization

[139799]of connective tissue growth factor as a member of the
IGFBP superfamily. Proc.

[139800]Nat. Acad. Sci. 94: 12981–12986, 1997.

[139801]24972.

[139802]Martinerie, C.; Viegas–Pequignot, E.; Guenard, I.; Dutril–
laux,

[139803]B.; Nguyen, V. C.; Bernheim, A.; Perbal, B.: Physical map–
ping of

[139804]human loci homologous to the chicken nov proto–
oncogene. Oncogene 7:

[139805]2529–2534, 1992.

[139806]24973.

- [139807]Nakanishi, T.; Yamaai, T.; Asano, M.; Nawachi, K.; Suzuki, M.;
- [139808]Sugimoto, T.; Takigawa, M.: Overexpression of connective tissue growth
- [139809]factor/hypertrophic chondrocyte-specific gene product 24 decreases
- [139810]bone density in adult mice and induces dwarfism. Biochem. Biophys. Res. Commun. 281: 678-681, 2001.
- [139812]24974.
- [139813]O'Donovan, N.; Galvin, M.; Morgan, J. G.: Physical mapping of
- [139814]the CXC chemokine locus on human chromosome 4. Cytogenet. Cell Genet. 84:
- [139815]39-42, 1999.
- [139816]24975.
- [139817]McGill, J. R.; Boyd, D.; Barrett, K. J.; Drysdale, J. W.; Moore, C. M.: Localization of human ferritin H (heavy) and L (light) subunits
- [139819]by in situ hybridization. (Abstract) Am. J. Hum. Genet. 36: 146S

[139820]only, 1984.

[139821]24976.

[139822]Murray, M. T.; White, K.; Munro, H. N.: Conservation of
ferritin

[139823]heavy subunit gene structure: implications for the regula-
tion of ferritin

[139824]gene expression. Proc. Nat. Acad. Sci. 84: 7438-7442,
1987.

[139825]24977.

[139826]Papadopoulos, P.; Bhavsar, D.; Zappone, E.; David, V.;
Jones,

[139827]C.; Worwood, M.; Drysdale, J.: A second human ferritin H
locus on

[139828]chromosome 11. Cytogenet. Cell Genet. 61: 107-108,
1992.

[139829]24978.

[139830]Richard, C. W.; Withers, D. A.; Meeker, T. C.; Myers, R. M.:

[139831]A radiation hybrid map of the proximal long arm of hu-
man chromosome

[139832]11 containing the MEN-1 and bcl-1 disease locus.
(Abstract) Cytogenet.

[139833]Cell Genet. 58: 1970 only, 1991.

[139834]24979.

[139835]Worwood, M.; Brook, J. D.; Cragg, S. J.; Hellkuhl, B.; Jones,

[139836]B. M.; Perera, P.; Roberts, S. H.; Shaw, D. J.: Assignment of
human

[139837]ferritin genes to chromosomes 11 and 19q13.3-19qter.

Hum. Genet. 69:

[139838]371-374, 1985.

[139839]24980.

[139840]Wu, K.-J.; Polack, A.; Dalla-Favera, R.: Coordinated regu-
lation

[139841]of iron-controlling genes, H-ferritin and IRP2, by c-MYC.
Science 283:

[139842]676-679, 1999.

[139843]24981.

[139844]Yachou, A.; Mattei, M. G.; Roeckel, N.; Grandchamp, B.;
Beaumont,

[139845]C.: Mouse ferritin H sequences map to chromosomes 3, 6,
and 19. Genomics 9:

[139846]204-206, 1991.

[139847]24982.

[139848]Yachou, A.-K.; Renaudie, F.; Guenet, J.-L.; Simon-
Chazottes, D.;

[139849]Jones, R.; Grandchamp, B.; Beaumont, C.: Mouse ferritin H
multigene

[139850]family is polymorphic and contains a single multiallelic
functional

[139851]gene located on chromosome 19. Genomics 10: 531-538,
1991.

[139852]24983.

[139853]Youssofian, H.; Chance, P.; Tuck-Muller, C. M.; Jabs, E.
W.:

[139854]Association of a new chromosomal deletion
[del(1)(q32q42)] with diaphragmatic

[139855]hernia: assignment of a human ferritin gene. Hum. Genet.
78: 267-270,

[139856]1988.

[139857]24984.

[139858]Takeda, J.; Seino, Y.; Tanaka, K.-I.; Fukumoto, H.; Kayano,
T.;

[139859]Takahashi, H.; Mitani, T.; Kurono, M.; Suzuki, T.; Tobe, T.;
Imura,

- [139860]H.: Sequence of an intestinal cDNA encoding human gastric inhibitory polypeptide precursor. Proc. Nat. Acad. Sci. 84: 7005–7008, 1987.
- [139861]24985.
- [139862]Xu, W.; Gorman, P. A.; Rider, S. H.; Hedge, P. J.; Moore, G.; Prichard, C.; Sheer, D.; Solomon, E.: Construction of a genetic map of human chromosome 17 by use of chromosome-mediated gene transfer. Proc. Nat. Acad. Sci. 85: 8563–8567, 1988.
- [139863]24986.
- [139864]Fishman, G. I.; Eddy, R. L.; Shows, T. B.; Rosenthal, L.; Leinwand, L. A.: The human connexin gene family of gap junction proteins: distinct chromosomal locations but similar structures. Genomics 10: 250–256, 1991.
- [139865]24987.
- [139866]Campbell, H. D.; Webb, G. C.; Kono, T.; Taniguchi, T.;

Ford, J.

[139874]H.; Young, I. G.: Assignment of the interleukin-2 receptor
beta chain

[139875]gene (IL-2rb) to band E on mouse chromosome 15. Ge-
nomics 12: 179-180,

[139876]1992.

[139877]24988.

[139878]Burgeson, R. E.; Chiquet, M.; Deutzmann, R.; Ekblom, P.;
Engel,

[139879]J.; Kleinman, H.; Martin, G. R.; Meneguzzi, G.; Paulsson,
M.; Sanes,

[139880]J.; Timpl, R.; Tryggvason, K.; Yamada, Y.; Yurchenco, P. D.:
A new

[139881]nomenclature for the laminins. Matrix Biol. 14: 209-211,
1994.

[139882]24989.

[139883]Gremlich, S.; Porret, A.; Hani, E. H.; Cherif, D.; Vionnet, N.;

[139884]Froguel, P.; Thorens, B.: Cloning, functional expression,
and chromosomal

[139885]localization of the human pancreatic islet glucose-de-
pendent insulinitropic

[139886]polypeptide receptor. Diabetes 44: 1202-1208, 1995.

[139887]24990.

[139888]Miyawaki, K.; Yamada, Y.; Ban, N.; Ihara, Y.; Tsukiyama, K.;
Zhou,

[139889]H.; Fujimoto, S.; Oku, A.; Tsuda, K.; Toyokuni, S.; Hiai, H.;
Mizunoya,

[139890]W.; and 9 others: Inhibition of gastric inhibitory polypep-
tide

[139891]signaling prevents obesity. *Nature Med.* 8: 738–742,
2002.

[139892]24991.

[139893]Stoffel, M.; Fernald, A. A.; Le Beau, M. M.; Bell, G. I.: As-
signment

[139894]of the gastric inhibitory polypeptide receptor gene (GIPR)
to chromosome

[139895]bands 19q13.2–q13.3 by fluorescence in situ hybridiza-
tion. *Genomics* 28:

[139896]607–609, 1995.

[139897]24992.

[139898]Usdin, T. B.; Mezey, E.; Button, D. C.; Brownstein, M. J.;
Bonner,

[139899]T. I.: Gastric inhibitory polypeptide receptor, a member of
the secretin–vasoactive

[139900]intestinal peptide receptor family, is widely distributed in
peripheral

[139901]organs and the brain. *Endocrinology* 133: 2861–2870,
1993.

[139902]24993.

[139903]Volz, A.; Groke, R.; Lankat-Buttgereit, B.; Fehmann, H.-C.;
Bode,

[139904]H. P.; Goke, B.: Molecular cloning, functional expression,
and signal

[139905]transduction of the GIP-receptor cloned from a human in-
sulinoma. *FEBS*

[139906]*Lett.* 373: 23–29, 1995.

[139907]24994.

[139908]Yamada, Y.; Hayami, T.; Nakamura, K.; Kaisaki, P. J.;
Someya, Y.;

[139909]Wang, C.-Z.; Seino, S.; Seino, Y.: Human gastric inhibitory
polypeptide

[139910]receptor: cloning of the gene (GIPR) and cDNA. *Genomics*
29: 773–776,

[139911]1995.

[139912]24995.

- [139913]Miyawaki, K.; Yamada, Y.; Yano, H.; Niwa, H.; Ban, N.; Ihara, Y.;
- [139914]Kubota, A.; Fujimoto, S.; Kajikawa, M.; Kuroe, A.; Tsuda, K.; Hashimoto,
- [139915]H.; Yamashita, T.; Jomori, T.; Tashiro, F.; Miyazaki, J.; Seino, Y.
- [139916]Glucose intolerance caused by a defect in the entero-insular axis:
- [139917]a study in gastric inhibitory polypeptide receptor knock-out mice. Proc.
- [139918]Nat. Acad. Sci. 96: 14843–14847, 1999.
- [139919]24996.
- [139920]Bentley, P. H.; Kenner, G. W.; Sheppard, R. C.: Structure of human
- [139921]gastrins I and II. Nature 209: 583–585, 1966.
- [139922]24997.
- [139923]Boel, E.; Vuust, J.; Norris, F.; Norris, K.; Wind, A.; Rehfeld, J. F.; Marcker, K. A.: Molecular cloning of human gastrin cDNA: evidence
- [139925]for evolution of gastrin by gene duplication. Proc. Nat. Acad. Sci. 80:

[139926]2866–2869, 1983.

[139927]24998.

[139928]Flejter, W. L.; Barcroft, C. L.; Guo, S.–W.; Lynch, E. D.;
Boehnke,

[139929]M.; Chandrasekharappa, S.; Hayes, S.; Collins, F. S.; Weber,
B. L.;

[139930]Glover, T. W.: Multicolor FISH mapping with Alu–
PCR–amplified YAC

[139931]clone DNA determines the order of markers in the BRCA1
region on chromosome

[139932]17q12–q21. Genomics 17: 624–631, 1993.

[139933]24999.

[139934]Flejter, W. L.; Kukowska–Latallo, J. F.; Kiouisis, S.; Chan–
drasekharappa,

[139935]S. C.; King, S. E.; Chamberlain, J. S.: Tetranucleotide repeat
polymorphism

[139936]at D17S846 maps within 40 kb of GAS at 17q12–q22.
Hum. Molec. Genet. 2:

[139937]1080 only, 1993.

[139938]25000.

[139939]Friis–Hansen, L.; Rourke, I. J.; Bundgaard, J. R.; Rehfeld, J.

- [139940]F.; Samuelson, L. C.: Molecular structure and genetic mapping of
[139941]the mouse gastrin gene. FEBS Lett. 386: 128–132, 1996.
[139942]25001.
- [139943]Fukushige, S.; Murotsu, T.; Matsubara, K.: Chromosomal assignment
[139944]of human genes for gastrin, thyrotropin (TSH)–beta sub-unit and C-erb-2
[139945]by chromosome sorting combined with velocity sedimentation and southern
[139946]hybridization. Biochem. Biophys. Res. Commun. 134: 477–483, 1986.
[139947]25002.
- [139948]Gregory, R. A.; Tracy, H. J.; Agarwal, K. L.; Grossman, M. I.:
[139949]Amino acid constitution of two gastrins isolated from Zollinger–Ellison
[139950]tumor tissue. Gut 10: 603–608, 1969.
[139951]25003.
- [139952]Ito, R.; Sato, K.; Helmer, T.; Jay, G.; Agarwal, K.: Structural
[139953]analysis of the gene encoding human gastrin: the large intron contains

[139954]an Alu sequence. Proc. Nat. Acad. Sci. 81: 4662–4666,
1984.

[139955]25004.

[139956]Lund, T.; Geurts van Kessel, A. H. M.; Huan, S.; Dixon, J. E.

[139957]: The genes for human gastrin and cholecystokinin are lo-
cated on different

[139958]chromosomes. Hum. Genet. 73: 77–80, 1986.

[139959]25005.

[139960]Lund, T.; Geurts van Kessel, A. H. M.; Westerveld, A.: The
human

[139961]gastrin gene is located at human chromosome 17.
(Abstract) Cytogenet.

[139962]Cell Genet. 40: 683 only, 1985.

[139963]25006.

[139964]Wiborg, O.; Berglund, L.; Boel, E.; Norris, F.; Norris, K.; Re-
hfeld,

[139965]J. F.; Marcker, K. A.; Vuust, J.: Structure of a human gastrin
gene. Proc.

[139966]Nat. Acad. Sci. 81: 1067–1069, 1984.

[139967]25007.

- [139968]Justice, M. J.; Gilbert, D. J.; Kinzler, K. W.; Vogelstein, B.;
[139969]Buchberg, A. M.; Ceci, J. D.; Matsuda, Y.; Chapman, V. M.;
Patriotis,
[139970]C.; Makris, A.; Tsiichlis, P. N.; Jenkins, N. A.; Copeland, N.
G.:
[139971]A molecular genetic linkage map of mouse chromosome
18 reveals extensive
[139972]linkage conservation with human chromosomes 5 and 18.
Genomics 13:
[139973]1281–1288, 1992.
[139974]25008.
[139975]Cox, D. W.; Francke, U.: Direct assignment of orosomu-
coid to human
[139976]chromosome 9 and alpha-2-HS-glycoprotein to chromo-
some 3 using human
[139977]fetal liver x rat hepatoma hybrids. Hum. Genet. 70:
109–115, 1985.
[139978]25009.
[139979]Umetsu, K.; Yuasa, I.; Nishimura, H.; Sasaki, H.; Suzuki, T.:
[139980]Genetic polymorphisms of orosomucoid and alpha-
2-HS-glycoprotein in

[139981]a Philippine population. Hum. Hered. 38: 287–290, 1988.

[139982]25010.

[139983]Triantafilou, K.; Triantafilou, M.; Dedrick, R. L.: A
CD14-independent

[139984]LPS receptor cluster. Nature Immun. 2: 338–345, 2001.

[139985]25011.

[139986]Klinger, H. P.: Suppression of tumorigenicity in somatic
cell

[139987]hybrids. I. Suppression and reexpression of tumorigenicity
in diploid

[139988]human x D98/AH2 hybrids and independent segregation
of tumorigenicity

[139989]from other cell phenotypes. Cytogenet. Cell Genet. 27:
254–266,

[139990]1980.

[139991]25012.

[139992]Lichy, J. H.; Modi, W. S.; Seuanetz, H. N.; Howley, P. M.:
Identification

[139993]of a human chromosome 11 gene which is differentially
regulated in

[139994]tumorigenic and nontumorigenic somatic cell hybrids of

HeLa cells. Cell

[139995]Growth Differ. 3: 541–548, 1992.

[139996]25013.

[139997]Stanbridge, E. J.: Suppression of malignancy in human cells. Nature 260:

[139998]17–20, 1976.

[139999]25014.

[140000]Corrigall, V. M.; Arastu, M.; Khan, S.; Shah, C.; Fife, M.; Smeets,

[140001]T.; Tak, P.–P.; Panayi, G. S.: Functional IL–2 receptor beta (CD122)

[140002]and gamma (CD132) chains are expressed by fibroblast-like synoviocytes:

[140003]activation by IL–2 stimulates monocyte chemoattractant protein–1 production. J.

[140004]Immun. 166: 4141–4147, 2001.

[140005]25015.

[140006]Gnarra, J. R.; Otani, H.; Wang, M. G.; McBride, O. W.; Sharon,

[140007]M.; Leonard, W. J.: Human interleukin 2 receptor beta-chain gene:

[140008]chromosomal localization and identification of 5-prime regulatory

[140009]sequences. Proc. Nat. Acad. Sci. 87: 3440–3444, 1990.

[140010]25016.

[140011]Hatakeyama, M.; Tsudo, M.; Minamoto, S.; Kono, T.; Doi, T.; Miyata,

[140012]T.; Miyasaka, M.; Taniguchi, T.: Interleukin-2 receptor beta chain

[140013]gene: generation of three receptor forms by cloned human alpha and

[140014]beta chain cDNA's. Science 244: 551–556, 1989.

[140015]25017.

[140016]Lamaze, C.; Dujeancourt, A.; Baba, T.; Lo, C. G.; Benmerah, A.;

[140017]Dautry-Varsat, A.: Interleukin 2 receptors and detergent-resistant

[140018]membrane domains define a clathrin-independent endocytic pathway. Molec.

[140019]Cell 7: 661–671, 2001.

[140020]25018.

[140021]Shibuya, H.; Yoneyama, M.; Nakamura, Y.; Harada, H.;

Hatakeyama,

[140022]M.; Minamoto, S.; Kono, T.; Doi, T.; White, R.; Taniguchi, T.: The

[140023]human interleukin-2 receptor beta-chain gene: genomic organization,

[140024]promoter analysis and chromosomal assignment. Nucleic Acids Res. 18:

[140025]3697-3703, 1990.

[140026]25019.

[140027]Suzuki, H.; Kundig, T. M.; Furlonger, C.; Wakeham, A.; Timms, E.;

[140028]Matsuyama, T.; Schmits, R.; Simard, J. J. L.; Ohashi, P. S.; Griesser,

[140029]H.; Taniguchi, T.; Paige, C. J.; Mak, T. W.: Deregulated T cell activation

[140030]and autoimmunity in mice lacking interleukin-2 receptor beta. Science 268:

[140031]1472-1476, 1995.

[140032]25020.

[140033]Tsuda, M.; Kitamura, F.; Miyasaka, M.: Characterization of the

[140034]interleukin 2 receptor beta chain using three distinct

monoclonal

[140035]antibodies. Proc. Nat. Acad. Sci. 86: 1982–1986, 1989.

[140036]25021.

[140037]Ballard, F. J.; Baxter, R. C.; Binoux, M.; Clemmons, D. R.;
Drop,

[140038]S. L. S.; Hall, K.; Hintz, R. L.; Rechler, M. M.; Rutanen, E.
M.;

[140039]Schwander, J. C.: Report on the nomenclature of the IGF
binding proteins. J.

[140040]Clin. Endocr. Metab. 70: 817–818, 1990.

[140041]25022.

[140042]Popovici, R. M.; Lu, M.; Bhatia, S.; Faessen, G. H.; Giaccia,
A.

[140043]J.; Giudice, L. C.: Hypoxia regulates insulin-like growth
factor-binding

[140044]protein 1 in human fetal hepatocytes in primary culture:
suggestive

[140045]molecular mechanisms for in utero fetal growth restriction
caused

[140046]by uteroplacental insufficiency. J. Clin. Endocr. Metab. 86:
2653–2659,

[140047]2001.

[140048]25023.

[140049]Allander, S. V.; Larsson, C.; Ehrenborg, E.; Suwanichkul,
A.; Weber,

[140050]G.; Morris, S. L.; Bajalica, S.; Kiefer, M. C.; Luthman, H.;
Powell,

[140051]D. R.: Characterization of the chromosomal gene and pro-
moter for

[140052]human insulin-like growth factor binding protein-5. J.
Biol. Chem. 269:

[140053]10891-10898, 1994.

[140054]25024.

[140055]Bauer, M.; Dieterich, W.; Ehnis, T.; Schuppan, D.: Complete
primary

[140056]structure of human collagen type XIV (undulin). Biochim.
Biophys.

[140057]Acta 1354: 183-188, 1997.

[140058]25025.

[140059]Dublet, B.; van der Rest, M.: Type XIV collagen, a new ho-
motrimeric

[140060]molecule extracted from fetal bovine skin and tendon,
with a triple

[140061]helical disulfide-bonded domain homologous to type IX
and type XII

[140062]collagens. J. Biol. Chem. 266: 6853-6858, 1991.

[140063]25026.

[140064]Just, M.; Herbst, H.; Hummel, M.; Durkop, H.; Tripier, D.;
Stein,

[140065]H.; Schuppan, D.: Undulin is a novel member of the fi-
bronectin-tenascin

[140066]family of extracellular matrix glycoproteins. J. Biol. Chem.
266:

[140067]17326-17332, 1991.

[140068]25027.

[140069]Schnittger, S.; Herbst, H.; Schuppan, D.; Dannenberg, C.;
Bauer,

[140070]M.; Fonatsch, C.: Localization of the undulin gene (UND) to
human

[140071]chromosome band 8q23. Cytogenet. Cell Genet. 68:
233-234, 1995.

[140072]25028.

[140073]Schuppan, D.; Cantaluppi, M. C.; Becker, J.; Veit, A.; Bunte,
T.;

- [140074]Troyer, D.; Schuppan, F.; Schmid, M.; Ackermann, R.;
Hahn, E. G.:
- [140075]Undulin, an extracellular matrix glycoprotein associated
with collagen
- [140076]fibrils. J. Biol. Chem. 265: 8823–8832, 1990.
- [140077]25029.
- [140078]Eklund, L.; Piihola, J.; Komulainen, J.; Sormunen, R.; Ong-
varrasopone,
- [140079]C.; Fassler, R.; Muona, A.; Ilves, M.; Ruskoaho, H.; Takala,
T. E.
- [140080]S.; Pihlajaniemi, T.: Lack of type XV collagen causes a
skeletal
- [140081]myopathy and cardiovascular defects in mice. Proc. Nat.
Acad. Sci. 98:
- [140082]1194–1199, 2001.
- [140083]25030.
- [140084]Hagg, P. M.; Hagg, P. O.; Peltonen, S.; Autio-Harminen,
H.; Pihlajaniemi,
- [140085]T.: Location of type XV collagen in human tissues and its
accumulation
- [140086]in the interstitial matrix of the fibrotic kidney. Am. J. Path.
150:

[140087]2075–2086, 1997.

[140088]25031.

[140089]Hagg, P. M.; Horelli-Kuitunen, N.; Eklund, L.; Palotie, A.;
Pihlajaniemi,

[140090]T.: Cloning of mouse type XV collagen sequences and
mapping of the

[140091]corresponding gene to 4B1–3: comparison of mouse and
human alpha-1(XV)

[140092]collagen sequences indicates divergence in the number of
small collagenous

[140093]domains. Genomics 45: 31–41, 1997.

[140094]25032.

[140095]Hagg, P. M.; Muona, A.; Lietard, J.; Kivirikko, S.; Pihla-
janiemi,

[140096]T.: Complete exon–intron organization of the human gene
for the alpha-1

[140097]chain of type XV collagen (COL15A1) and comparison with
the homologous

[140098]Col18a1 gene. J. Biol. Chem. 273: 17824–17831, 1998.

[140099]25033.

[140100]Huebner, K.; Cannizzaro, L. A.; Jabs, E. W.; Kivirikko, S.;

Manzone,

[140101]H.; Pihlajaniemi, T.; Myers, J. C.: Chromosomal assignment
of a gene

[140102]encoding a new collagen type (COL15A1) to 9q21-q22.

Genomics 14:

[140103]220-224, 1992.

[140104]25034.

[140105]Kivirikko, S.; Heinamaki, P.; Rehn, M.; Honkanen, N.; My-
ers, J.

[140106]C.; Pihlajaniemi, T.: Primary structure of the alpha-1 chain
of human

[140107]type XV collagen and exon-intron organization in the
3-prime region

[140108]of the corresponding gene. J. Biol. Chem. 269:
4773-4779, 1994.

[140109]25035.

[140110]Muragaki, Y.; Abe, N.; Ninomiya, Y.; Olsen, B. R.; Ooshima,
A.

[140111]: The human alpha-1(XV) collagen chain contains a large
amino-terminal

[140112]non-triple helical domain with a tandem repeat structure
and homology

[140113]to alpha-1(XVIII) collagen. J. Biol. Chem. 269: 4042-4046, 1994.

[140114]25036.

[140115]Myers, J. C.; Dion, A. S.; Abraham, V.; Amenta, P. S.: Type XV

[140116]collagen exhibits a widespread distribution in human tissues but a

[140117]distinct localization in basement membrane zones. Cell Tissue Res. 286:

[140118]493-505, 1996.

[140119]25037.

[140120]Myers, J. C.; Kivirikko, S.; Gordon, M. K.; Dion, A. S.; Pihlajaniemi,

[140121]T.: Identification of a previously unknown human collagen chain,

[140122]alpha-1(XV), characterized by extensive interruptions in the triple-helical

[140123]region. Proc. Nat. Acad. Sci. 89: 10144-10148, 1992.

[140124]25038.

[140125]Ramchandran, R.; Dhanabal, M.; Volk, R.; Waterman, M. J. F.; Segal,

[140126]M.; Lu, H.; Knebelmann, B.; Sukhatme, V. P.: Antiangiogenic activity

[140127]of restin, NC10 domain of human collagen XV: comparison to endostatin. Biochem.

[140128]Biophys. Res. Commun. 255: 735–739, 1999.

[140129]25039.

[140130]Rehn, M.; Hintikka, E.; Pihlajaniemi, T.: Primary structure of

[140131]the alpha 1 chain of mouse type XVIII collagen, partial structure

[140132]of the corresponding gene, and comparison of the alpha 1(XVIII) chain

[140133]with its homologue, the alpha 1(XV) collagen chain. J. Biol. Chem. 269:

[140134]13929–13935, 1994.

[140135]25040.

[140136]Sasaki, T.; Larsson, H.; Tisi, D.; Claesson-Welsh, L.; Hohenester,

[140137]E.; Timpl, R.: Endostatins derived from collagens XV and XVIII differ

[140138]in structural and binding properties, tissue distribution and anti-angiogenic

[140139]activity. *J. Molec. Biol.* 301: 1179–1190, 2000.

[140140]25041.

[140141]Pan, T.-C.; Zhang, R.-Z.; Mattei, M.-G.; Timpl, R.; Chu, M.-L.

[140142]: Cloning and chromosomal location of human alpha-1(XVI) collagen.

[140143]*Proc. Nat. Acad. Sci.* 89: 6565–6569, 1992.

[140144]25042.

[140145]Yamaguchi, N.; Kimura, S.; McBride, O. W.; Hori, H.; Yamada, Y.;

[140146]Kanamori, T.; Yamakoshi, H.; Nagai, Y.: Molecular cloning and partial

[140147]characterization of a novel collagen chain, alpha-1(XVI), consisting

[140148]of repetitive collagenous domains and cysteine-containing non-collagenous

[140149]segments. *J. Biochem.* 112: 856–863, 1992.

[140150]25043.

[140151]O'Reilly, M. S.; Boehm, T.; Shing, Y.; Fukai, N.; Vasios, G.; Lane,

[140152]W. S.; Flynn, E.; Birkhead, J. R.; Olsen, B. R.; Folkman, J.:

Endostatin:

- [140153]an endogenous inhibitor of angiogenesis and tumor growth. Cell 88: 277–285, 1997.
- [140155]25044.
- [140156]Sakai, L. Y.; Keene, D. R.; Glanville, R. W.; Bachinger, H. P.
- [140157]: Purification and partial characterization of fibrillin, a cysteine-rich structural component of connective tissue microfibrils. J. Biol. Chem. 266: 14763–14770, 1991.
- [140160]25045.
- [140161]Eiberg, H.; Bisgaard, M. L.; Mohr, J.: Linkage between alpha-1-B-glycoprotein (A1BG) and Lutheran (LU) red blood group system: assignment to chromosome 19: new genetic variants of A1BG. Clin. Genet. 36: 415–418, 1989.
- [140164]25046.
- [140165]Eiberg, H.; Nielsen, L. S.; Gahne, B.; Juneja, R. K.; Mohr, J.
- [140166]: Exclusion data for the alpha-1-B glycoprotein (A1BG)

polymorphism.

[140167](Abstract) Cytogenet. Cell Genet. 51: 994 only, 1989.

[140168]25047.

[140169]Gahne, B.; Juneja, R. K.; Stratil, A.: Genetic polymorphism
of

[140170]human plasma alpha-1-B-glycoprotein: phenotyping by
immunoblotting

[140171]or by a simple method of 2-D electrophoresis. Hum.
Genet. 76: 111-115,

[140172]1987.

[140173]25048.

[140174]Ishioka, N.; Takahashi, N.; Putnam, F. W.: Amino acid se-
quence

[140175]of human plasma alpha-1B-glycoprotein: homology to the
immunoglobulin

[140176]supergene family. Proc. Nat. Acad. Sci. 83: 2363-2367,
1986.

[140177]25049.

[140178]Juneja, R. K.; Weitkamp, L. R.; Stratil, A.; Gahne, B.; Gut-
tormsen,

[140179]S. A.: Further studies of the plasma alpha-

1-B-glycoprotein polymorphism:

[140180]two new alleles and allele frequencies in Caucasians and
in American

[140181]blacks. Hum. Hered. 38: 267-272, 1988.

[140182]25050.

[140183]Anderson, L.; Anderson, N. G.: High resolution two-
dimensional

[140184]electrophoresis of human plasma proteins. Proc. Nat.
Acad. Sci. 74:

[140185]5421-5425, 1977.

[140186]25051.

[140187]Boutin, B.; Feng, S. H.; Arnaud, P.: The genetic polymor-
phism

[140188]of alpha(2)-HS glycoprotein: study by ultrathin-layer iso-
electric

[140189]focusing and immunoblot. Am. J. Hum. Genet. 37:
1098-1105, 1985.

[140190]25052.

[140191]Cox, D. W.; Andrews, B. J.: Silver stain immunofixation for
alpha-2-HS-glycoprotein:

[140192]a new method for detection of protein heterogeneity. In:

Stathakos,

[140193]D.: Electrophoresis '82. Berlin: Walter de Gruyter (pub.)
1983.

[140194]Pp. 243–247.

[140195]25053.

[140196]Cox, D. W.; Andrews, B. J.; Wills, D. E.: Genetic polymor-
phism

[140197]of alpha-2-HS-glycoprotein. Am. J. Hum. Genet. 38:
699–706, 1986.

[140198]25054.

[140199]Cox, D. W.; Francke, U.; Allderdice, P. W.; McAlpine, P. J.:
Gene

[140200]mapping of human serum proteins using hepatoma hy-
brids and human chromosome

[140201]deletions and duplications. Genetics 107 (suppl.):
s22–s23, 1984.

[140202]25055.

[140203]Eiberg, H.; Mohr, J.; Nielsen, L. S.: A2HS: new methods of
phenotyping

[140204]and analysis of linkage relations: assignment to chromo-
some 3. (Abstract) Cytogenet.

[140205]Cell Genet. 37: 461 only, 1984.

[140206]25056.

[140207]Gejyo, F.; Chang, J.-L.; Burgi, W.; Schmid, K.; Offner, G. D.;

[140208]Troxler, R. F.; Van Halbeek, H.; Dorland, L.; Gerwig, G. J.;
Vliegenthart,

[140209]J. F. G.: Characterization of the B-chain of human plasma
alpha(2)HS-glycoprotein:

[140210]the complete amino acid sequence and primary structure
of its heteroglycan. J.

[140211]Biol. Chem. 258: 4966-4971, 1983.

[140212]25057.

[140213]Jahnen-Dechent, W.: Personal Communication. Mainz,
Germany

[140214]7/22/1998.

[140215]25058.

[140216]Jahnen-Dechent, W.; Schinke, T.; Trindl, A.; Muller-Esterl,
W.;

[140217]Sablitzky, F.; Kaiser, S.; Blessing, M.: Cloning and targeted
deletion

[140218]of the mouse fetuin gene. J. Biol. Chem. 272:
31496-31503, 1997.

[140219]25059.

- [140220]Lee, C.-C.; Bowman, B. H.; Yang, F.: Human alpha-2-HS-glycoprotein: the A and B chains with a connecting sequence are encoded by a single mRNA transcript. Proc. Nat. Acad. Sci. 84: 4403-4407, 1987.
- [140221]25060.
- [140222]Fajans, S. S.; Bell, G. I.; Polonsky, K. S.: Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. New Eng. J. Med. 345: 971-980, 2001.
- [140223]25061.
- [140224]Lindner, T.; Gragnoli, C.; Furuta, H.; Cockburn, B. N.; Petzold, C.; Rietzsch, H.; Weiss, U.; Schulze, J.; Bell, G. I.: Hepatic function in a family with a nonsense mutation (R154X) in the hepatocyte nuclear factor-4-alpha/MODY1 gene. J. Clin. Invest. 100: 1400-1405, 1997.
- [140225]25062.

- [140233]Moller, A. M.; Dalgaard, L. T.; Ambye, L.; Hansen, L.; Schmitz, O.; Hansen, T.; Pedersen, O.: A novel Phe75fsdelT mutation in the hepatocyte nuclear factor-4-alpha gene in a Danish pedigree with maturity-onset diabetes of the young. J. Clin. Endocr. Metab. 84: 367-369, 1999.
- [140237]25063.
- [140238]Thomas, H.; Jaschowitz, K.; Bulman, M.; Frayling, T. M.; Mitchell, S. M. S.; Roosen, S.; Lingott-Frieg, A.; Tack, C. J.; Ellard, S.; Ryffel, G. U.; Hattersley, A. T.: A distant upstream promoter of the HNF-4-alpha gene connects the transcription factors involved in maturity-onset diabetes of the young. Hum. Molec. Genet. 10: 2089-2097, 2001.
- [140243]25064.
- [140245]Froguel, P.; Vaxillaire, M.; Sun, F.; Velho, G.; Zouali, H.; Butel,

- [140246]M. O.; Lesage, S.; Vionnet, N.; Clement, K.; Fougerousse, F.; Tanizawa,
- [140247]Y.; Weissenbach, J.; Beckmann, J. S.; Lathrop, G. M.; Passa, P.; Permutt,
- [140248]M. A.; Cohen, D.: Close linkage of glucokinase locus on chromosome
- [140249]7p to early-onset non-insulin-dependent diabetes mellitus. Nature 356:
- [140250]162-164, 1992.
- [140251]25065.
- [140252]Vionnet, N.; Stoffel, M.; Takeda, J.; Yasuda, K.; Bell, G. I.;
- [140253]Zouali, H.; Lesage, S.; Velho, G.; Iris, F.; Passa, P.; Froguel, P.;
- [140254]Cohen, D.: Nonsense mutation in the glucokinase gene causes early-onset
- [140255]non-insulin-dependent diabetes mellitus. Nature 356: 721-722, 1992.
- [140256]25066.
- [140257]Dunger, D. B.; Ong, K. K. L.; Huxtable, S. J.; Sherriff, A.; Woods,
- [140258]K. A.; Ahmed, M. L.; Golding, J.; Pembrey, M. E.; Ring, S.;
- ALSPAC

[140259]Study Team; Bennett, S. T.; Todd, J. A.: Association of the
INS

[140260]VNTR with size at birth. Nature Genet. 19: 98–100, 1998.

[140261]25067.

[140262]Joseph, L. J.; Le Beau, M. M.; Jamieson, G. A., Jr.; Acharya,
S.;

[140263]Shows, T. B.; Rowley, J. D.; Sukhatme, V. P.: Molecular
cloning,

[140264]sequencing, and mapping of EGR2, a human early growth
response gene

[140265]encoding a protein with 'zinc-binding finger' structure.
Proc. Nat.

[140266]Acad. Sci. 85: 7164–7168, 1988.

[140267]25068.

[140268]Nagarajan, R.; Svaren, J.; Le, N.; Araki, T.; Watson, M.; Mil-
brandt,

[140269]J.: EGR2 mutations in inherited neuropathies dominant-
negatively

[140270]inhibit myelin gene expression. Neuron 30: 355–368,
2001.

[140271]25069.

- [140272] Timmerman, V.; De Jonghe, P.; Ceuterick, C.; De Vriendt, E.; Lofgren, A.; Nelis, E.; Warner, L. E.; Lupski, J. R.; Martin, J.-J.; Van Broeckhoven, C.: Novel missense mutation in the early growth response 2 gene associated with Dejerine-Sottas syndrome phenotype. *Neurology* 52: 1827-1832, 1999.
- [140276] 25070.
- [140277] Warner, L. E.; Mancias, P.; Butler, I.; Lupski, J. R.: Mutation in the early growth response 2 (EGR2) transcription factor associated with recessive congenital hypomyelinating neuropathy (CHN). (Abstract) *Am. J. Hum. Genet.* 61 (suppl.): A350 only, 1997.
- [140281] 25071.
- [140282] Warner, L. E.; Mancias, P.; Butler, I. J.; McDonald, C. M.; Keppen, L.; Koob, K. G.; Lupski, J. R.: Mutations in the early growth response

[140285]2 (EGR2) gene are associated with hereditary myelinopathies. Nature

[140286]Genet. 18: 382–384, 1998.

[140287]25072.

[140288]Warner, L. E.; Svaren, J.; Milbrandt, J.; Lupski, J. R.: Functional

[140289]consequences of mutations in the early growth response 2 gene (EGR2)

[140290]correlate with severity of human myelinopathies. Hum. Molec. Genet. 8:

[140291]1245–1251, 1999.

[140292]25073.

[140293]Wu, J.; Joseph, L.; Sukhatme, V. P.; Kidd, K. K.: A HindIII polymorphism

[140294]identified by the human early growth response gene 2 (EGR2) on chromosome 25074.

[140295]Nucleic Acids Res. 16: 11855 only, 1988.

[140296]25075.

[140297]Kaneda, Y.; Hayes, H.; Uchida, T.; Yoshida, M. C.; Okada, Y.:

[140298]Regional assignment of five genes on human chromosome

19. Chromosoma 95:

[140299]8-12, 1987.

[140300]25076.

[140301]Boyle, J. M.; Hey, Y.; Fox, M.: Human ecto-5-prime nucleotidase

[140302]maps to chromosome 6q14-q21. (Abstract) Cytogenet. Cell Genet. 51:

[140303]968 only, 1989.

[140304]25077.

[140305]Boyle, J. M.; Hey, Y.; Geurts van Kessel, A.; Fox, M.: Assignment

[140306]of ecto-5-prime-nucleotidase to human chromosome 6. Hum. Genet. 81:

[140307]88-92, 1988.

[140308]25078.

[140309]Misumi, Y.; Ogata, S.; Ohkubo, K.; Hirose, S.; Ikehara, Y.: Primary

[140310]structure of human placental 5-prime-nucleotidase and identification

[140311]of the glycolipid anchor in the mature form. Europ. J. Biochem. 191:

[140312]563–569, 1990.

[140313]25079.

[140314]Monreal, A. W.; Ferguson, B. M.; Headon, D. J.; Street, S. L.;

[140315]Overbeek, P. A.; Zonana, J.: Mutations in the human homologue of

[140316]mouse dl cause autosomal recessive and dominant hypohidrotic ectodermal

[140317]dysplasia. *Nature Genet.* 22: 366–369, 1999.

[140318]25080.

[140319]Schrijver, I.; Liu, W.; Brenn, T.; Furthmayr, H.; Francke, U.

[140320]: Cysteine substitutions in epidermal growth factor-like domains of

[140321]fibrillin-1: distinct effects on biochemical and clinical phenotypes. *Am.*

[140322]*J. Hum. Genet.* 65: 1007–1020, 1999.

[140323]25081.

[140324]Schrijver, I.; Liu, W.; Francke, U.: The pathogenicity of the

[140325]Pro1148-to-ala substitution in the FBN1 gene: causing or predisposing

[140326]to Marfan syndrome and aortic aneurysm, or clinically innocent? Hum.

[140327]Genet. 99: 607–611, 1997.

[140328]25082.

[140329]Schrijver, I.; Liu, W.; Odom, R.; Brenn, T.; Oefner, P.; Furthmayr,

[140330]H.; Francke, U.: Premature termination mutations in FBN1: distinct

[140331]effects on differential allelic expression and on protein and clinical

[140332]phenotypes. Am. J. Hum. Genet. 71: 223–237, 2002.

[140333]25083.

[140334]Siracusa, L. D.; McGrath, R.; Ma, Q.; Moskow, J. J.; Manne, J.;

[140335]Christner, P. J.; Buchberg, A. M.; Jimenez, S. A.: A tandem duplication

[140336]within the fibrillin 1 gene is associated with the mouse tight skin

[140337]mutation. Genome Res. 6: 300–313, 1996.

[140338]25084.

[140339]Sood, S.; Eldadah, Z. A.; Krause, W. L.; McIntosh, I.; Dietz,

- [140340]H. C.: Mutation in fibrillin-1 and the Marfanoid-craniosynostosis
- [140341](Shprintzen-Goldberg) syndrome. Nature Genet. 12: 209-211, 1996.
- [140342]25085.
- [140343]Stahl-Hallengren, C.; Ukkonen, T.; Kainulainen, K.; Kristofersson,
- [140344]U.; Saxne, T.; Tornqvist, K.; Peltonen, L.: An extra cysteine in
- [140345]one of the non-calcium-binding epidermal growth factor-like motifs
- [140346]of the FBN1 polypeptide is connected to a novel variant of Marfan
- [140347]syndrome. J. Clin. Invest. 94: 709-713, 1994.
- [140348]25086.
- [140349]Tiecke, F.; Katzke, S.; Booms, P.; Robinson, P. N.; Neumann, L.;
- [140350]Godfrey, M.; Mathews, K. R.; Scheuner, M.; Hinkel, G. K.; Brenner,
- [140351]R. E.; Hovels-Gurich, H. H.; Hagemeier, C.; Fuchs, J.; Skovby, F.;
- [140352]Rosenberg, T.: Classic, atypically severe and neonatal

Marfan syndrome:

- [140353] twelve mutations and genotype-phenotype correlations in FBN1 exons
24-40. *Europ. J. Hum. Genet.* 9: 13-21, 2001.
- [140355] 25087.
- [140356] Trask, T. M.; Ritty, T. M.; Broekelmann, T.; Tisdale, C.; Mecham, R. P.: N-terminal domains of fibrillin 1 and fibrillin 2 direct the formation of homodimers: a possible first step in microfibril assembly. *Biochem. J.* 340: 693-701, 1999.
- [140360] 25088.
- [140361] Tynan, K.; Comeau, K.; Pearson, M.; Wilgenbus, P.; Levitt, D.; Gasner, C.; Berg, M. A.; Miller, D. C.; Francke, U.: Mutation screening of complete fibrillin-1 coding sequence: report of five new mutations, including two in 8-cysteine domains. *Hum. Molec. Genet.* 2: 1813-1821, 1993.

[140366]25089.

[140367]Velinov, M.; Sarfarazi, M.; Young, K.; Hodes, M. E.; Con-
neally,

[140368]P. M.; Jackson, C. E.; Tsipouras, P.: Limb-girdle muscular
dystrophy

[140369]is closely linked to the fibrillin locus on chromosome 15.
Connect.

[140370]Tissue Res. 29: 13-21, 1993.

[140371]25090.

[140372]Wang, M.; Mathews, K. R.; Imaizumi, K.; Beiraghi, S.;
Blumberg,

[140373]B.; Scheuner, M.; Graham, J. M., Jr.; Godfrey, M.: P1148A in
fibrillin-1

[140374]is not a mutation anymore. (Letter) Nature Genet. 15: 12
only, 1997.

[140375]25091.

[140376]Wang, M.; Wang, J.-Y.; Cisler, J.; Imaizumi, K.; Burton, B.
K.;

[140377]Jones, M. C.; Lamberti, J. J.; Godfrey, M.: Three novel fib-
rillin

[140378]mutations in exons 25 and 27: classic versus neonatal
Marfan syndrome. Hum.

[140379]Mutat. 9: 359–362, 1997.

[140380]25092.

[140381]Watanabe, Y.; Yano, S.; Koga, Y.; Yukizane, S.; Nishiyori, A.;

[140382]Yoshino, M.; Kato, H.; Ogata, T.; Adachi, M.: P1148A in fibrillin-1

[140383]is not a mutation leading to Shprintzen–Goldberg syndrome. (Letter) Hum.

[140384]Mutat. 10: 326–327, 1997.

[140385]25093.

[140386]Whiteman, P.; Smallridge, R. S.; Knott, V.; Cordle, J. J.; Downing,

[140387]A. K.; Handford, P. A.: A G1127S change in calcium-binding epidermal

[140388]growth factor-like domain 13 of human fibrillin-1 causes short range

[140389]conformational effects. J. Biol. Chem. 276: 17156–17162, 2001.

[140390]25094.

[140391]Alving, B. M.; Henschen, A. H.: Fibrinogen Giessen I: a congenital

[140392]homozygously expressed dysfibrinogenemia with A-alpha
arg-to-his substitution. Am.

[140393]J. Hemat. 25: 479-482, 1987.

[140394]25095.

[140395]Arocha-Pinango, C. L.; Rodriguez, S.; Nagy, H.; Perez Re-
quejo,

[140396]J. L.: Fibrinogen Lima: a new dysfibrinogenaemia with a
high-molecular

[140397]weight alpha-chain and effective polymerization. Blood
Coagul. Fibrinolysis 1:

[140398]561-565, 1990.

[140399]25096.

[140400]Aschbacher, A.; Buetow, K.; Chung, D.; Walsh, S.; Murray,
J.:

[140401]Linkage disequilibrium of RFLP's associated with alpha,
beta, and

[140402]gamma fibrinogen predict gene order on chromosome 4.
(Abstract) Am.

[140403]J. Hum. Genet. 37: A186 only, 1985.

[140404]25097.

[140405]Asl, L. H.; Liepnieks, J. J.; Uemichi, T.; Rebibou, J.-M.; Jus-

trabo,

[140406]E.; Droz, D.; Mousson, C.; Chalopin, J.-M.; Benson, M. D.;
Delpech,

[140407]M.; Grateau, G.: Renal amyloidosis with a frame shift mu-
tation in

[140408]fibrinogen A(alpha)-chain gene producing a novel amyloid
protein. Blood 90:

[140409]4799-4805, 1997.

[140410]25098.

[140411]Asselta, R.; Duga, S.; Spena, S.; Santagostino, E.; Peyvandi,
F.;

[140412]Piseddu, G.; Targhetta, R.; Malcovati, M.; Mannucci, P. M.;
Tenchini,

[140413]M. L.: Congenital afibrinogenemia: mutations leading to
premature

[140414]termination codons in fibrinogen A-alpha-chain gene are
not associated

[140415]with the decay of the mutant mRNAs. Blood 98:
3685-3692, 2001.

[140416]25099.

[140417]Aznar, J.; Fernandez-Pavon, A.; Reganon, E.; Vila, V.; Orel-
lana,

[140418]F.: Fibrinogen Valencia--a new case of congenital dysfibrinogenemia. Thromb.

[140419]Diath. Haemorrh. 32: 564-577, 1974.

[140420]25100.

[140421]Barthels, M.; Sandvoss, G.: Fibrinogen Hannover, another abnormal

[140422]fibrinogen. Blut 34: 99-106, 1977.

[140423]25101.

[140424]Beck, E. A.; Charache, P.; Jackson, D. P.: A new inherited coagulation

[140425]disorder caused by an abnormal fibrinogen ('fibrinogen Baltimore'). Nature 208:

[140426]143-145, 1965.

[140427]25102.

[140428]Beck, E. A.; Shainoff, J. R.; Vogel, A.; Jackson, D. P.: Functional

[140429]evaluation of an inherited abnormal fibrinogen: fibrinogen 'Baltimore.'. J.

[140430]Clin. Invest. 50: 1874-1884, 1971.

[140431]25103.

[140432]Benson, M. D.; Liepnieks, J.; Uemichi, T.; Wheeler, G.; Cor-
rea,

[140433]R.: Hereditary renal amyloidosis associated with a mutant
fibrinogen

[140434]alpha-chain. Nature Genet. 3: 252-255, 1993.

[140435]25104.

[140436]Blomback, B.; Blomback, M.: Molecular defects and vari-
ants of

[140437]fibrinogen. Nouv. Rev. Franc. Hemat. 10: 671-678, 1970.

[140438]25105.

[140439]Blomback, B.; Hessel, B.; Fields, R.; Procyk, R.: Fibrinogen

[140440]Aarhus: an abnormal fibrinogen with A alpha 19 arg-
to-gly substitution.In:

[140441]Mosesson, M. W.; Amrani, D. L.; Siebenlist, K. R.; Di Orio,
J. P.

[140442]: Fibrinogen 3: Biochemistry, Biological Functions, Gene
Regulation

[140443]and Expression.. Amsterdam: Elsevier (pub.) 1988. Pp.
263-266.

[140444]25106.

[140445]Ballo, R.; Beighton, P. H.; Ramesar, R. S.: Stickler-like syn-

drome

[140446]due to a dominant negative mutation in the COL2A1 gene.

Am. J. Med.

[140447]Genet. 80: 6–11, 1998.

[140448]25107.

[140449]Chan, E. F.; Gat, U.; McNiff, J. M.; Fuchs, E.: A common hu-

man

[140450]skin tumour is caused by activating mutations in beta-

catenin. Nature

[140451]Genet. 21: 410–413, 1999.

[140452]25108.

[140453]Oh, S. P.; Kamagata, Y.; Muragaki, Y.; Timmons, S.;

Ooshima, A.;

[140454]Olsen, B. R.: Isolation and sequencing of cDNAs for pro-

teins with

[140455]multiple domains of Gly-X-Y repeats identify a novel fam-

ily of collagenous

[140456]proteins. Proc. Nat. Acad. Sci. 91: 4229–4233, 1994.

[140457]25109.

[140458]Oh, S. P.; Warman, M. L.; Seldin, M. F.; Cheng, S.-D.; Knoll,

J.

[140459]H. M.; Timmons, S.; Olsen, B. R.: Cloning of cDNA and genomic DNA
[140460]encoding human type XVIII collagen and localization of the
alpha-1(XVIII)
[140461]collagen gene to mouse chromosome 10 and human
chromosome 21. Genomics 19:
[140462]494-499, 1994.
[140463]25110.
[140464]Rehn, M.; Hintikka, E.; Pihlajaniemi, T.: Characterization of
[140465]the mouse gene for the alpha-1 chain of type XVIII collagen (Col18a1)
[140466]reveals that the three variant N-terminal polypeptide
forms are transcribed
[140467]from two widely separated promoters. Genomics 32:
436-446, 1996.
[140468]25111.
[140469]Rehn, M.; Pihlajaniemi, T.: Alpha-1(XVIII), a collagen chain
with
[140470]frequent interruptions in the collagenous sequence, a distinct tissue
[140471]distribution, and homology with type XV collagen. Proc.
Nat. Acad.

[140472]Sci. 91: 4234–4238, 1994.

[140473]25112.

[140474]Saarela, J.; Ylikarppa, R.; Rehn, M.; Purmonen, S.; Pihlajaniemi,

[140475]T.: Complete primary structure of two variant forms of human type

[140476]XVIII collagen and tissue-specific differences in the expression of

[140477]the corresponding transcripts. Matrix Biol. 16: 319–328, 1998.

[140478]25113.

[140479]Horelli-Kuitunen, N.; Kvist, A.-P.; Helaakoski, T.; Kivirikko,

[140480]K.; Pihlajaniemi, T.; Palotie, A.: The order and transcriptional

[140481]orientation of the human COL13A1 and P4HA genes on chromosome 10 long

[140482]arm determined by high-resolution FISH. Genomics 46: 299–302, 1997.

[140483]25114.

[140484]Pajunen, L.; Tamminen, M.; Solomon, E.; Pihlajaniemi, T.: Assignment

[140485] of the gene coding for the alpha 1 chain of collagen type
XIII (COL13A1)

[140486] to human chromosome region 10q11-qter. Cytogenet.
Cell Genet. 52:

[140487] 190-193, 1989.

[140488] 25115.

[140489] Shows, T. B.; Tikka, L.; Byers, M. G.; Eddy, R. L.; Haley, L.
L.;

[140490] Henry, W. M.; Prockop, D. J.; Tryggvason, K.: Assignment
of the human

[140491] collagen alpha-1(XIII) chain gene (COL13A1) to the q22
region of chromosome

[140492] 10. Genomics 5: 128-133, 1989.

[140493] 25116.

[140494] Tikka, L.; Pihlajaniemi, T.; Henttu, P.; Prockop, D. J.; Tryg-
gvason,

[140495] K.: Gene structure for the alpha-1 chain of a human
short-chain collagen

[140496] (type XIII) with alternatively spliced transcripts and trans-
lation

[140497] termination codon at the 5-prime end of the last exon.
Proc. Nat.

[140498]Acad. Sci. 85: 7491–7495, 1988.

[140499]25117.

[140500]Cunningham, B. A.; Hemperly, J. J.; Murray, B. A.; Prediger, E.

[140501]A.; Brackenbury, R.; Edelman, G. M.: Neural cell adhesion molecule:

[140502]structure, immunoglobulin-like domains, cell surface modulation, and

[140503]alternative RNA splicing. Science 236: 799–806, 1987.

[140504]25118.

[140505]D'Eustachio, P.; Davisson, M. T.: Resolution of the staggerer

[140506](sg) mutation from the neural cell adhesion molecule locus (Ncam)

[140507]on mouse chromosome 9. Mammalian Genome 4: 278–280, 1993.

[140508]25119.

[140509]D'Eustachio, P.; Owens, G. C.; Edelman, G. M.; Cunningham, B. A.

[140510]: Chromosomal location of the gene encoding the neural cell adhesion

[140511]molecule (N-CAM) in the mouse. Proc. Nat. Acad. Sci. 82:
7631-7635,

[140512]1985.

[140513]25120.

[140514]Lin, D. M.; Fetter, R. D.; Kopczynski, C.; Grenningloh, G.;
Goodman,

[140515]C. S.: Genetic analysis of fasciclin II in Drosophila: defasci-
cultation,

[140516]refasciculation, and altered fasciculation. Neuron 13:
1055-1069,

[140517]1994.

[140518]25121.

[140519]Altare, F.; Jouanguy, E.; Lamhamedi-Cherradi, S.; Fon-
daneche, M.-C.;

[140520]Fizame, C.; Ribierre, F.; Merlin, G.; Dembic, Z.; Schreiber,
R.; Lisowska-Grospierre,

[140521]B.; Fischer, A.; Seboun, E.; Casanova, J.-L.: A causative re-
lationship

[140522]between mutant IFNGR1 alleles and impaired cellular re-
sponse to IFN-gamma

[140523]in a compound heterozygous child. (Letter) Am. J. Hum.
Genet. 62:

[140524]723–726, 1998.

[140525]25122.

[140526]Dalton, D. K.; Pitts–Meek, S.; Keshav, S.; Figari, I. S.;
Bradley,

[140527]A.; Stewart, T. A.: Multiple defects of immune cell function
in mice

[140528]with disrupted interferon–gamma genes. Science 259:
1739–1742, 1993.

[140529]25123.

[140530]Dessein, A. J.; Hillaire, D.; Elwali, N. E. M. A.; Marquet, S.;

[140531]Mohamed–Ali, Q.; Mirghani, A.; Henri, S.; Abdelhameed, A.
A.; Saeed,

[140532]O. K.; Magzoub, M. M. A.; Abel, L.: Severe hepatic fibrosis
in Schistosoma

[140533]mansoni infection is controlled by a major locus that is
closely linked

[140534]to the interferon–gamma receptor gene. Am. J. Hum.
Genet. 65: 709–721,

[140535]1999.

[140536]25124.

[140537]Fellous, M.: Personal Communication. Paris, France

10/24/1986.

[140538]25125.

[140539]Fellous, M.; Couillin, P.; Rosa, F.; Metezeau, P.; Foubert, C.;

[140540]Gross, M. S.; Frezal, J.; Van Cong, N.: Receptor for human gamma

[140541]interferon is specified by human chromosome 18.
(Abstract) Cytogenet.

[140542]Cell Genet. 40: 627-628, 1985.

[140543]25126.

[140544]Jouanguy, E.; Altare, F.; Lamhamedi, S.; Revy, P.; Emile, J.-F.;

[140545]Newport, M.; Levin, M.; Blanche, S.; Seboun, E.; Fischer, A.; Casanova,

[140546]J.-L.: Interferon-gamma-receptor deficiency in an infant with fatal

[140547]bacille Calmette-Guerin infection. New Eng. J. Med. 335: 1956-1961,

[140548]1996.

[140549]25127.

[140550]Jouanguy, E.; Dupuis, S.; Pallier, A.; Doffinger, R.; Fon-

daneche,

[140551]M.-C.; Fieschi, C.; Lamhamedi-Cherradi, S.; Altare, F.;
Emile, J.-F.;

[140552]Lutz, P.; Bordigoni, P.; Cokugras, H.; Akcakaya, N.; Land-
man-Parker,

[140553]J.; Donnadieu, J.; Camcioglu, Y.; Casanova, J.-L.: In a novel
form

[140554]of IFN-gamma receptor 1 deficiency, cell surface recep-
tors fail to

[140555]bind IFN-gamma. J. Clin. Invest. 105: 1429-1436, 2000.

[140556]25128.

[140557]Jouanguy, E.; Lamhamedi-Cherradi, S.; Lammas, D.; Dor-
man, S. E.;

[140558]Fondaneche, M.-C.; Dupuis, S.; Doffinger, R.; Altare, F.;
Girdlestone,

[140559]J.; Emile, J.-F.; Ducoulombier, H.; Edgar, D.; and 10 others:
A

[140560]human IFNGR1 small deletion hotspot associated with
dominant susceptibility

[140561]to mycobacterial infection. Nature Genet. 21: 370-378,
1999.

[140562]25129.

[140563]Kaplan, D. H.; Shankaran, V.; Dighe, A. S.; Stockert, E.;
 Aguet,
 [140564]M.; Old, L. J.; Schreiber, R. D.: Demonstration of an inter-
 feron
 [140565]gamma-dependent tumor surveillance system in immuno-
 competent mice. Proc.
 [140566]Nat. Acad. Sci. 95: 7556-7561, 1998.
 [140567]25130.
 [140568]Le Coniat, M.; Alcaide-Loridan, C.; Fellous, M.; Berger, R.:
 [140569]Human interferon gamma receptor 1 (IFNGR1) gene maps
 to chromosome
 [140570]region 6q23-6q24. Hum. Genet. 84: 92-94, 1989.
 [140571]25131.
 [140572]Lekstrom-Himes, J. A.; Gallin, J. I.: Immunodeficiency dis-
 eases
 [140573]caused by defects in phagocytes. New Eng. J. Med. 343:
 1703-1714,
 [140574]2000.
 [140575]25132.
 [140576]Levin, M.; Newport, M. J.; D'Souza, S.; Kalabalikis, P.;
 Brown,

- [140577]J. N.; Lenicker, H. M.; Agius, P. V.; Davies, E. G.; Thrasher, A.;
- [140578]Klein, N.; Blackwell, J. M.: Familial disseminated atypical mycobacterial
- [140579]infection in childhood: a human mycobacterial susceptibility gene? Lancet 345:
- [140580]79–83, 1995.
- [140581]25133.
- [140582]Mariano, T. M.; Kozak, C. A.; Langer, J. A.; Pestka, S.: The
- [140583]mouse immune interferon receptor gene is located on chromosome 10. J.
- [140584]Biol. Chem. 262: 5812–5814, 1987.
- [140585]25134.
- [140586]Deloukas, P.; Schuler, G. D.; Gyapay, G.; Beasley, E. M.; Soderlund,
- [140587]C.; Rodriguez–Tome, P.; Hui, L.; Matise, T. C.; McKusick, K. B.; Beckmann,
- [140588]J. S.; Bentolila, S.; Bihoreau, M.–T.; and 53 others: A physical
- [140589]map of 30,000 human genes. Science 282: 744–746, 1998.

[140590]25135.

[140591]Manser, E.; Leung, T.; Salihuddin, H.; Tan, L.; Lim, L.: A
non-receptor

[140592]tyrosine kinase that inhibits the GTPase activity of
p21cdc42. Nature 363:

[140593]364–367, 1993.

[140594]25136.

[140595]Wu, W. J.; Erickson, J. W.; Lin, R.; Cerione, R. A.: The
gamma-subunit

[140596]of the coatamer complex binds Cdc42 to mediate trans-
formation. Nature 405:

[140597]800–804, 2000.

[140598]25137.

[140599]Basu, S.; Binder, R. J.; Ramalingam, T.; Srivastava, P. K.:
CD91

[140600]is a common receptor for heat shock proteins gp96,
hsp90, hsp70, and

[140601]calreticulin. Immunity 14: 303–313, 2001.

[140602]25138.

[140603]Beisiegel, U.; Weber, W.; Ihrke, G.; Herz, J.; Stanley, K. K.:

[140604]The LDL-receptor-related protein, LRP, is an apolipoprotein E-binding

[140605]protein. *Nature* 341: 162-164, 1989.

[140606]25139.

[140607]Binder, R. J.; Han, D. K.; Srivastava, P. K.: CD91: a receptor

[140608]for heat shock protein gp96. *Nature Immun.* 1: 151-155, 2000.

[140609]25140.

[140610]Forus, A.; Maelandsmo, G. M.; Fodstad, Y.; Myklebost, O.: The

[140611]genes for the alpha-2-macroglobulin receptor/LDL receptor-related

[140612]protein and GLI are located within a chromosomal segment of about

[140613]300 kilobases and are coamplified in a rhabdomyosarcoma cell line.

[140614](Abstract) *Cytogenet. Cell Genet.* 58: 1977 only, 1991.

[140615]25141.

[140616]Forus, A.; Myklebost, O.: A physical map of a 1.3-Mb region on

[140617]the long arm of chromosome 12, spanning the GLI and

LRP loci. Genomics 14:

[140618]117–120, 1992.

[140619]25142.

[140620]Hilliker, C.; Van Leuven, F.; Van Den Berghe, H.: Assignment of

[140621]the gene coding for the alpha(2)-macroglobulin receptor to mouse chromosome

[140622]15 and to human chromosome 12q13–q14 by isotopic and nonisotopic in

[140623]situ hybridization. Genomics 13: 472–474, 1992.

[140624]25143.

[140625]Kristensen, T.; Moestrup, S. K.; Gliemann, J.; Bendtsen, L.; Sand,

[140626]O.; Sottrup-Jensen, L.: Evidence that the newly cloned low-density-lipoprotein

[140627]receptor related protein (LRP) is the alpha-2-macroglobulin receptor. FEBS

[140628]Lett. 276: 151–155, 1990.

[140629]25144.

[140630]Lendon, C. L.; Talbot, C. J.; Craddock, N. J.; Han, S. W.; Wragg,

- [140631] M.; Morris, J. C.; Goate, A. M.: Genetic association studies between
- [140632] dementia of the Alzheimer's type and three receptors for apolipoprotein
- [140633] E in a Caucasian population. *Neurosci. Lett.* 222: 187-190, 1997.
- [140634] 25145.
- [140635] Myklebost, O.; Arheden, K.; Rogne, S.; Geurts van Kessel, A.;
- [140636] Mandahl, N.; Herz, J.; Stanley, K.; Heim, S.; Mitelman, F.: The gene
- [140637] for the human putative apoE receptor is on chromosome 12 in the segment

[140638]q13–14. Genomics 5: 65–69, 1989.

[140639]25146.

[140640]Pericak–Vance, M. A.; Bass, M. P.; Yamaoka, L. H.; Gaskell, P.

[140641]C.; Scott, W. K.; Terwedow, H. A.; Menold, M. M.; Conneally, P. M.;

[140642]Small, G. W.; Vance, J. M.; Saunders, A. M.; Roses, A. D.; Haines,

[140643]J. L.: Complete genomic screen in late-onset familial Alzheimer disease:

[140644]evidence for a new locus on chromosome 12. J.A.M.A. 278: 1237–1241,

[140645]1997.

[140646]25147.

[140647]Scott, W. K.; Yamaoka, L. H.; Bass, M. P.; Gaskell, P. C.; Conneally,

[140648]P. M.; Small, G. W.; Farrer, L. A.; Auerbach, S. A.; Saunders, A.

[140649]M.; Roses, A. D.; Haines, J. L.; Pericak–Vance, M. A.: No genetic

[140650]association between the LRP receptor and sporadic or late-onset familial

[140651]Alzheimer disease. Neurogenetics 1: 179–183, 1998.

[140652]25148.

[140653]Strickland, D. K.; Ashcom, J. D.; Williams, S.; Burgess, W.
H.;

[140654]Migliorini, M.; Argraves, W. S.: Sequence identity between
the alpha-2-macroglobulin

[140655]receptor and low density lipoprotein receptor-related pro-
tein suggests

[140656]that this molecule is a multifunctional receptor. J. Biol.
Chem. 265:

[140657]17401–17404, 1990.

[140658]25149.

[140659]Yochem, J.; Greenwald, I.: A gene for a low density
lipoprotein

[140660]receptor-related protein in the nematode *Caenorhabditis*
elegans. Proc.

[140661]Nat. Acad. Sci. 90: 4572–4576, 1993.

[140662]25150.

[140663]Ladías, J. A. A.; Karathanasis, S. K.: Regulation of the
apolipoprotein

[140664]AI gene by ARP-1, a novel member of the steroid receptor

superfamily. Science 251:

[140665]561–565, 1991.

[140666]25151.

[140667]Modi, W. S.; Seuanes, H.; Mietus-Snyder, M.; O'Brien, S. J.;
Karathanasis,

[140668]S. K.: Chromosomal localization of the ARP-1 gene to
15q26. (Abstract) Cytogenet.

[140669]Cell Genet. 58: 1995 only, 1991.

[140670]25152.

[140671]Pereira, F. A.; Qiu, Y.; Zhou, G.; Tsai, M.-J.; Tsai, S. Y.: The
[140672]orphan nuclear receptor COUP-TFII is required for angio-
genesis and

[140673]heart development. Genes Dev. 13: 1037–1049, 1999.

[140674]25153.

[140675]Widom, R. L.; Ladas, J. A. A.; Kouidou, S.; Karathanasis, S.
K.

[140676]: Synergistic interactions between transcription factors
control expression

[140677]of the apolipoprotein AI gene in liver cells. Molec. Cell.
Biol. 11:

[140678]677–687, 1991.

[140679]25154.

[140680]Agre, P.; Smith, B. L.; Baumgarten, R.; Preston, G. M.;
Pressman,

[140681]E.; Wilson, P.; Illum, N.; Anstee, D. J.; Lande, M. B.; Zeidel,
M.

[140682]L.: Human red cell aquaporin CHIP. II. Expression during
normal fetal

[140683]development and in a novel form of congenital dysery-
thropoietic anemia. J.

[140684]Clin. Invest. 94: 1050-1058, 1994.

[140685]25155.

[140686]Ikeda, H.; Yamaguchi, M.; Sugai, S.; Aze, Y.; Narumiya, S.;
Kakizuka,

[140687]A.: Expanded polyglutamine in the Machado-Joseph dis-
ease protein

[140688]induces cell death in vitro and in vivo. Nature Genet. 13:
196-202,

[140689]1996.

[140690]25156.

[140691]Santisteban, I.; Arredondo-Vega, F. X.; Kelly, S.; Mary, A.;
Fischer,

[140692]A.; Hummell, D. S.; Lawton, A.; Sorensen, R. U.; Stiehm, E.
R.; Uribe,
[140693]L.; Weinberg, K.; Hershfield, M. S.: Novel splicing, mis-
sense, and
[140694]deletion mutations in seven adenosine deaminase-defi-
cient patients
[140695]with late/delayed onset of combined immunodeficiency
disease: contribution
[140696]of genotype to phenotype. J. Clin. Invest. 92: 2291-2302,
1993.
[140697]
[140698]100. Schmalstieg, F. C.; Mills, G. C.; Tsuda, H.; Goldman,
A. S.:
[140699]Severe combined immunodeficiency in a child with a
healthy adenosine
[140700]deaminase deficient mother. Pediat. Res. 17: 935-940,
1983.
[140701]
[140702]101. Schrader, W. P.; Pollara, B.; Meuwissen, H. J.: Charac-
terization
[140703]of the residual adenosine deaminating activity in the
spleen of a

[140704]patient with combined immunodeficiency disease and
adenosine deaminase

[140705]deficiency. Proc. Nat. Acad. Sci. 75: 446–450, 1978.

[140706]

[140707]102. Scott, C. R.; Chen, S.–H.; Giblett, E. R.: Deletion of
the carrier

[140708]state in combined immunodeficiency disease associated
with deaminase

[140709]deficiency. J. Clin. Invest. 53: 1194–1196, 1974.

[140710]

[140711]103. Shovlin, C. L.; Hughes, J. M. B.; Simmonds, H. A.;
Fairbanks,

[140712]L.; Deacock, S.; Lechler, R.; Roberts, I.; Webster, A. D. B.:
Adult

[140713]presentation of adenosine deaminase deficiency. (Letter)
Lancet 341:

[140714]1471, 1993.

[140715]

[140716]104. Shovlin, C. L.; Simmonds, H. A.; Fairbanks, L. D.;
Deacock, S.

[140717]J.; Hughes, J. M. B.; Lechler, R. I.; Webster, A. D. B.; Sun,

X.-M.;

- [140718]Webb, J. C.; Soutar, A. K.: Adult onset immunodeficiency caused by
[140719]inherited adenosine deaminase deficiency. J. Immun. 153:
2331-2339,
[140720]1994.
- [140721]
- [140722]105. Spencer, N.; Hopkinson, D. A.; Harris, H.: Adenosine
deaminase
[140723]polymorphism in man. Ann. Hum. Genet. 32: 9-14, 1968.
- [140724]
- [140725]106. Stephan, V.; Wahn, V.; Le Deist, F.; Dirksen, U.; Bro-
ker, B.;
- [140726]Muller-Fleckenstein, I.; Horneff, G.; Schroten, H.; Fischer,
A.; de
[140727]Saint Basile, G.: Atypical X-linked severe combined im-
munodeficiency
[140728]due to possible spontaneous reversion of the genetic de-
fect in T cells. New
[140729]Eng. J. Med. 335: 1563-1567, 1996.
- [140730]

[140731]107. Tariverdian, G.; Ritter, H.: Adenosine deaminase polymorphism

[140732](EC 3.5.4.4): formal genetics and linkage relations. Humangenetik 7:

[140733]176–178, 1969.

[140734]

[140735]108. Tischfield, J. A.; Creagan, R. P.; Nichols, E. A.; Ruddle, F.

[140736]H.: Assignment of a gene for adenosine deaminase to human chromosome

[140737]20. Hum. Hered. 24: 1–11, 1974.

[140738]

[140739]109. Tzall, S.; Ellenbogen, A.; Eng, F.; Hirschhorn, R.: Identification

[140740]and characterization of nine RFLPs at the adenosine deaminase (ADA)

[140741]locus. Am. J. Hum. Genet. 44: 864–875, 1989.

[140742]

[140743]110. Umetsu, D. T.; Schlossman, C. M.; Ochs, H. D.; Herschfield, M.

[140744]S.: Heterogeneity of phenotype in two siblings with

adenosine deaminase

[140745]deficiency. J. Allergy Clin. Immun. 93: 543-550, 1994.

[140746]

[140747]111. Valerio, D.; Dekker, B. M. M.; Duyvesteyn, M. G. C.;
van der

[140748]Voorn, L.; Berkvens, T. M.; van Ormondt, H.; van der Eb, A.
J.: One

[140749]adenosine deaminase allele in a patient with severe com-
bined immunodeficiency

[140750]contains a point mutation abolishing enzyme activity.
EMBO J. 5:

[140751]113-119, 1986.

[140752]

[140753]112. Valerio, D.; Duyvesteyn, M. G. C.; Dekker, B. M. M.;
Weeda, G.;

[140754]Berkvens, T. M.; van der Voorn, L.; van Ormondt, H.; van
der Eb, A.

[140755]J.: Adenosine deaminase: characterization and expression
of a gene

[140756]with a remarkable promoter. EMBO J. 4: 437-443, 1985.

[140757]

- [140758]113. Valerio, D.; Duyvesteyn, M. G. C.; Meera Khan, P.; Pearson, P.
- [140759]L.; Geurts van Kessel, A.; van Ormondt, H.: Direct assignment of
- [140760]ADA gene to chromosome 20. (Abstract) Cytogenet. Cell Genet. 37:
- [140761]599, 1984.
- [140762]
- [140763]114. Valerio, D.; Duyvesteyn, M. G. C.; van Ormondt, H.; Meera Khan,
- [140764]P.; van der Eb, A. J.: Adenosine deaminase (ADA) deficiency in cells
- [140765]derived from humans with severe combined immunodeficiency is due to
- [140766]an aberration of the ADA protein. Nucleic Acids Res. 12: 1015-1024,
- [140767]1984.
- [140768]
- [140769]115. Valerio, D.; McIvor, R. S.; Williams, S. R.; Duyvesteyn, M. G.
- [140770]C.; van Ormondt, H.; van der Eb, A. J.; Martin, D. W., Jr.: Cloning

[140771]of human adenosine deaminase cDNA and expression in mouse cells. Gene 31:

[140772]147–153, 1984.

[140773]

[140774]116. Van der Weyden, M. B.; Kelley, W. N.: Adenosine deaminase deficiency

[140775]in severe combined immunodeficiency: evidence for a posttranslational

[140776]defect. (Abstract) J. Clin. Invest. 53: 81A–82A, 1974.

[140777]

[140778]117. Wakamiya, M.; Blackburn, M. R.; Jurecic, R.; McArthur, M. J.;

[140779]Geske, R. S.; Cartwright, J., Jr.; Mitani, K.; Vaishnav, S.; Belmont,

[140780]J. W.; Kellems, R. E.; Finegold, M. J.; Montgomery, Jr., C. A.; Bradley,

[140781]A.; Caskey, C. T.: Disruption of the adenosine deaminase gene causes

[140782]hepatocellular impairment and perinatal lethality in mice. Proc.

[140783]Nat. Acad. Sci. 92: 3673–3677, 1995.

[140784]

[140785]118. Weitkamp, L. R.: Further data on the genetic linkage relations

[140786]of the adenosine deaminase locus. Hum. Hered. 21: 351-356, 1971.

[140787]

[140788]119. Weitkamp, L. R.: Genetic linkage relationships of the ADA and

[140789]6-PGD loci in 'Humangenetik.' (Letter) Humangenetik 15: 359-360,

[140790]1972.

[140791]

[140792]120. Wiginton, D. A.; Adrian, G. S.; Friedman, R. L.; Suttle, D. P.;

[140793]Hutton, J. J.: Cloning of cDNA sequences of human adenosine deaminase. Proc.

[140794]Nat. Acad. Sci. 80: 7481-7485, 1983.

[140795]

[140796]121. Wiginton, D. A.; Adrian, G. S.; Hutton, J. J.: Sequence of human

[140797]adenosine deaminase cDNA including the coding region

and a small intron. Nucleic

[140798]Acids Res. 12: 2439–2446, 1984.

[140799]

[140800]122. Wiginton, D. A.; Hutton, J. J.: Immunoreactive protein
in adenosine

[140801]deaminase deficient human lymphoblast cell lines. J. Biol.
Chem. 257:

[140802]3211–3217, 1982.

[140803]

[140804]123. Wiginton, D. A.; Kaplan, D. J.; States, J. C.; Akeson, A.
L.;

[140805]Perme, C. M.; Bilyk, I. J.; Vaughn, A. J.; Lattier, D. L.; Hut-
ton,

[140806]J. J.: Complete sequence and structure of the gene for hu-
man adenosine

[140807]deaminase. Biochemistry 25: 8234–8244, 1986.

[140808]

[140809]124. Yokoyama, S.; Hayashi, T.; Yoshimura, Y.; Irimada,
K.; Saito,

[140810]T.; Akiba, T.; Tsuchiya, S.: Severe combined immunodeficiency disease

[140811]with adenosine deaminase deficiency. Tohoku J. Exp. Med.
129: 197–202,

[140812]1979.

[140813]

[140814]125. Yount, J.; Nichols, P.; Ochs, H. D.; Hammar, S. P.;
Scott, C.

[140815]R.; Chen, S.–H.; Giblett, E. R.; Wedgwood, R. J.: Absence of
erythrocyte

[140816]adenosine deaminase associated with severe combined
immunodeficiency. J.

[140817]Pediat. 84: 173–177, 1974.

[140818]

[140819]126. Ziegler, J. B.; Lee, C. H.; Van Der Weyden, M. B.; Bag-
nara, A.

[140820]S.; Beveridge, J.: Severe combined immunodeficiency and
adenosine

[140821]deaminase deficiency: failure of enzyme replacement
therapy. Arch.

[140822]Dis. Child. 55: 452–457, 1980.

[140823]

[140824]127. Ziegler, J. B.; Van Der Weyden, M. B.; Lee, C. H.;

Daniel, A.

[140825] Prenatal diagnosis for adenosine deaminase deficiency. J. Med.

[140826] Genet. 18: 154–156, 1981.

[140827] 25157.

[140828] Dooley, T. P.: Personal Communication. San Antonio, Tex. 7/21/1992.

[140829] 25158.

[140830] Dooley, T. P.; Weiland, K. L.; Simon, M.: cDNA sequence of human

[140831] p11 calpactin I light chain. Genomics 13: 866–868, 1992.

[140832] 25159.

[140833] Harder, T.; Kube, E.; Gerke, V.: Cloning and characterization

[140834] of the human gene encoding p11: structural similarity to other members

[140835] of the S-100 gene family. Gene 113: 269–274, 1992.

[140836] 25160.

[140837] Kube, E.; Weber, K.; Gerke, V.: Primary structure of human, chicken,

[140838] and *Xenopus laevis* p11, a cellular ligand of the Src-kinase

substrate,

[140839]annexin II. Gene 102: 255–259, 1991.

[140840]25161.

[140841]Okuse, K.; Malik–Hall, M.; Baker, M. D.; Poon, W.–Y. L.;
Kong,

[140842]H.; Chao, M. V.; Wood, J. N.: Annexin II light chain regu-
lates sensory

[140843]neuron–specific sodium channel expression. Nature 417:
653–656,

[140844]2002.

[140845]25162.

[140846]Saris, C. J. M.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.;

[140847]Noonan, D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and
tissue

[140848]distribution of the mRNA for bovine and murine p11, the
S100–related

[140849]light chain of the protein–tyrosine kinase substrate p36
(calpactin

[140850]l). J. Biol. Chem. 262: 10663–10671, 1987.

[140851]25163.

[140852]Eikenboom, J. C. J.; Vink, T.; Briet, E.; Sixma, J. J.; Reitsma,

[140853]P. H.: Multiple substitutions in the von Willebrand factor gene that
[140854]mimic the pseudogene sequence. Proc. Nat. Acad. Sci. 91:
2221–2224,
[140855]1994.
[140856]25164.
[140857]Gralnick, H. R.; Coller, B. S.: Molecular defects in
haemophilia
[140858]A and von Willebrand's disease. Lancet I: 837–838, 1976.
[140859]25165.
[140860]Watanabe, H.; Orii, K. E.; Fukao, T.; Song, X.–Q.; Aoyama,
T.;
[140861]Jlst, L.; Ruiter, J.; Wanders, R. J. A.; Kondo, N.: Molecular
basis
[140862]of very long chain acyl–CoA dehydrogenase deficiency in
three Israeli
[140863]patients: identification of a complex mutant allele with
P65L and
[140864]K247Q mutations, the former being an exonic mutation
causing exon
[140865]3 skipping. Hum. Mutat. 15: 430–438, 2000.

[140866]25166.

[140867]Gavish, H.; dos Santos, C. C.; Buchwald, M.: Generation of
a non-functional

[140868]Fanconi anemia group C protein (FACC) by site-directed in
vitro mutagenesis.

[140869](Abstract) Am. J. Hum. Genet. 51 (suppl.): A128 only,
1992.

[140870]25167.

[140871]Gavish, H.; dos Santos, C. C.; Buchwald, M.: A
leu554-to-pro substitution

[140872]completely abolishes the functional complementing activ-
ity of the

[140873]Fanconi anemia (FACC) protein. Hum. Molec. Genet. 2:
123-126, 1993.

[140874]25168.

[140875]Ahmad, W.; Haque, M. F.; Brancolini, V.; Tsou, H. C.;
Haque, S.;

[140876]Lam, H.; Aita, V. M.; Owen, J.; deBlaquiere, M.; Frank, J.;
Cserhalmi-Friedman,

[140877]P. B.; Leask, A.; McGrath, J. A.; Peacocke, M.; Ahmad, M.;
Ott, J.;

[140878]Christiano, A. M.: Alopecia universalis associated with a

mutation

[140879]in the human hairless gene. Science 279: 720–724, 1998.

[140880]25169.

[140881]Matthijs, G.; Schollen, E.; Heykants, L.; Grunewald, S.:

Phosphomannomutase

[140882]deficiency: the molecular basis of the classical Jaeken syndrome (CDGS

[140883]type Ia). Molec. Genet. Metab. 68: 220–226, 1999.

[140884]25170.

[140885]Matthijs, G.; Schollen, E.; Pardon, E.; Veiga-Da-Cunha, M.;

Jaeken,

[140886]J.; Cassiman, J.-J.; Van Schaftingen, E.: Mutations in PMM2, a phosphomannomutase

[140887]gene on chromosome 16p13, in carbohydrate-deficient glycoprotein type

[140888]I syndrome (Jaeken syndrome). Nature Genet. 16: 88–92, 1997.

[140889]25171.

[140890]Matthijs, G.; Schollen, E.; Pirard, M.; Budarf, M. L.; Van Schaftingen,

[140891]E.; Cassiman, J.-J.: PMM (PMM1), the human homologue of

SEC53 or

[140892] yeast phosphomannomutase, is localized on chromosome
22q13. Genomics 40:

[140893] 41–47, 1997.

[140894] 25172.

[140895] Broeks, A.; de Klein, A.; Floore, A. N.; Muijtjens, M.; Klei-
jer,

[140896] W. J.; Jaspers, N. G.; van't Veer, L. J.: ATM germline muta-
tions

[140897] in classical ataxia–telangiectasia patients in the Dutch
population. Hum.

[140898] Mutat. 12: 330–337, 1998.

[140899] 25173.

[140900] Munoz, F.; Lestringant, G.; Sybert, V.; Frydman, M.; Al-
swaini,

[140901] A.; Frossard, P. M.; Jorgenson, R.; Zonana, J.: Definitive
evidence

[140902] for an autosomal recessive form of hypohidrotic ectoder-
mal dysplasia

[140903] clinically indistinguishable from the more common X-
linked disorder. Am.

[140904] J. Hum. Genet. 61: 94–100, 1997.

[140905]25174.

[140906]Van Schaftingen, E.; Jaeken, J.: Phosphomannomutase deficiency

[140907]is a cause of carbohydrate-deficient glycoprotein syndrome type I. FEBS

[140908]Lett. 377: 318-320, 1995.

[140909]25175.

[140910]Carritt, B.: Somatic cell genetic evidence for the presence of

[140911]a gene for citrullinemia on human chromosome 9. Cytogenet. Cell Genet. 19:

[140912]44-48, 1977.

[140913]25176.

[140914]Carritt, B.; Goldfarb, P. S. G.; Hooper, M. L.; Slack, C.: Chromosome

[140915]assignment of a human gene for argininosuccinate synthetase expression

[140916]in Chinese hamster-human somatic cell hybrids. Exp. Cell Res. 106:

[140917]71-78, 1977.

[140918]25177.

[140919]Daiger, S. P.; Hoffman, N. S.; Wildin, R. S.; Su, T.-S.: Multi-
ple,
[140920]independent restriction site polymorphisms in human
DNA detected with
[140921]a cDNA probe to argininosuccinate synthetase (AS). Am. J.
Hum. Genet. 36:
[140922]736-749, 1984.
[140923]25178.
[140924]Daiger, S. P.; Wildin, R. S.; Su, T.-S.: Polymorphic variants
[140925]of restriction fragments of human DNA detected with a
probe to argininosuccinate
[140926]synthetase. (Abstract) Am. J. Hum. Genet. 33: 136A only,
1981.
[140927]25179.
[140928]Gripp, K. W.; Stolle, C. A.; Celle, L.; McDonald-McGinn, D.
M.;
[140929]Whitaker, L. A.; Zackai, E. H.: TWIST gene mutation in a
patient
[140930]with radial aplasia and craniosynostosis: further evidence
for heterogeneity
[140931]of Baller-Gerold syndrome. Am. J. Med. Genet. 82:
170-176, 1999.

[140932]25180.

[140933]Seto, M. L.; Lee, S. J.; Sze, R. W.; Cunningham, M. L.: Another

[140934]TWIST on Baller-Gerold syndrome. Am. J. Med. Genet.
104: 323-330,

[140935]2001.

[140936]25181.

[140937]Karathanasis, S. K.: Apolipoprotein multigene family: tandem organization

[140938]of human apolipoprotein AI, CIII, and AIV genes. Proc. Nat.
Acad.

[140939]Sci. 82: 6374-6378, 1985.

[140940]25182.

[140941]Bailey, R.; Miller, D. A.: Mapping the human apolipoprotein
genes

[140942]CII and E to band 19q13.1 by in situ hybridization.
(Abstract) Am.

[140943]J. Hum. Genet. 41: A156 only, 1987.

[140944]25183.

[140945]Bartres-Faz, D.; Clemente, I. C.; Junque, C.; Valveny, N.;
Lopez-Alomar,

[140946]A.; Sanchez-Aldeguer, J.; Lopez-Guillen, A.; Moral, P.:
APOE and
[140947]APOC1 genetic polymorphisms in age-associated memory
impairment. Neurogenetics 3:
[140948]215–219, 2001.
[140949]25184.
[140950]Crook, T. H.; Bartus, R. T.; Ferris, S. H.; Whitehouse, P.;
Cohen,
[140951]G. D.; Gershon, S.: Age-associated memory impairment:
proposed diagnostic
[140952]criteria and measures of clinical change. Report of a Na-
tional Institute
[140953]of Mental Health work group. Dev. Neuropsychol. 2:
261–276, 1986.
[140954]25185.
[140955]Davison, P. J.; Norton, P.; Wallis, S. C.; Gill, L.; Cook, M.;
[140956]Williamson, R.; Humphries, S. E.: There are two gene se-
quences for
[140957]human apolipoprotein CI (APO CI) on chromosome 19,
one of which is
[140958]4 kb from the gene for APO E. Biochem. Biophys. Res.
Commun. 136:

[140959]876–884, 1986.

[140960]25186.

[140961]Gautier, T.; Masson, D.; Jong, M. C.; Duverneuil, L.; Le
Guern,

[140962]N.; Deckert, V.; Pais de Barros, J.–P.; Dumont, L.; Bataille,
A.;

[140963]Zak, Z.; Jiang, X.–C.; Tall, A. R.; Havekes, L. M.; Lagrost, L.:

[140964]Apolipoprotein CI deficiency markedly augments plasma
lipoprotein

[140965]changes mediated by human cholesteryl ester transfer
protein (CETP)

[140966]in CETP transgenic/ApoCI–knocked out mice. J. Biol.
Chem. 277: 31354–31363,

[140967]2002.

[140968]25187.

[140969]Lauer, S. J.; Walker, D.; Elshourbagy, N. A.; Reardon, C. A.;

Levy–Wilson,

[140970]B.; Taylor, J. M.: Two copies of the human apolipoprotein
C–I gene

[140971]are linked closely to the apolipoprotein E gene. J. Biol.
Chem. 263:

[140972]7277-7286, 1988.

[140973]25188.

[140974]Myklebost, O.; Rogne, S.: The gene for human apolipoprotein CII

[140975]is located 4.3 kilobases away from the apolipoprotein E gene on chromosome

[140976]19. Hum. Genet. 73: 286-289, 1986.

[140977]25189.

[140978]Myklebost, O.; Rogne, S.: A physical map of the apolipoprotein

[140979]gene cluster on human chromosome 19. Hum. Genet. 78: 244-247, 1988.

[140980]25190.

[140981]Richards, A. J.; Baguley, D. M.; Yates, J. R. W.; Lane, C.; Nicol,

[140982]M.; Harper, P. S.; Scott, J. D.; Snead, M. P.: Variation in the vitreous

[140983]phenotype of Stickler syndrome can be caused by different amino acid

[140984]substitutions in the X position of the type II collagen Gly-X-Y triple

[140985]helix. Am. J. Hum. Genet. 67: 1083–1094, 2000.

[140986]25191.

[140987]Stickler, G. B.; Belau, P. G.; Farrell, F. J.; Jones, J. D.; Pugh,

[140988]D. G.; Steinberg, A. G.; Ward, L. E.: Hereditary progressive
arthro–ophthalmopathy. Mayo

[140989]Clin. Proc. 40: 433–455, 1965.

[140990]25192.

[140991]Wilkin, D. J.; Liberfarb, R.; Davis, J.; Levy, H. P.; Cole, W.

[140992]G.; Francomano, C. A.; Cohn, D. H.: Rapid determination
of COL2A1

[140993]mutations in individuals with Stickler syndrome: analysis
of potential

[140994]premature termination codons. Am. J. Med. Genet. 94:
141–148, 2000.

[140995]25193.

[140996]Williams, C. J.; Ganguly, A.; Considine, E.; McCarron, S.;
Prockop,

[140997]D. J.; Walsh–Vockley, C.; Michels, V. V.: A(–2)–to–G transi-
tion at

[140998]the 3–prime acceptor splice site of IVS17 characterizes the
COL2A1

[140999]gene mutation in the original Stickler syndrome kindred.
Am. J. Med.

[141000]Genet. 63: 461–467, 1996.

[141001]25194.

[141002]Zlotogora, J.; Sagi, M.; Schuper, A.; Leiba, H.; Merin, S.:
Variability

[141003]of Stickler syndrome. Am. J. Med. Genet. 42: 337–339,
1992.

[141004]25195.

[141005]Brown, S.; Wiebel, F. J.; Gelboin, H. V.; Minna, J. D.: As-
signment

[141006]of a locus required for flavoprotein–linked monooxyge-
nase expression

[141007]to human chromosome 2. Proc. Nat. Acad. Sci. 73:
4628–4632, 1976.

[141008]25196.

[141009]Chen, Y. T.; Tukey, R. H.; Swan, D. C.; Negishi, N.; Nebert,
D.

[141010]W.: Characterization of the human P1–450 genomic gene.
(Abstract) Clin.

[141011]Res. 31: 456A, 1983.

[141012]25197.

[141013]Corchero, J.; Pimprale, S.; Kimura, S.; Gonzalez, F. J.: Organization

[141014]of the CYP1A cluster on human chromosome 15: implications for gene

[141015]regulation. Pharmacogenetics 11: 1–6, 2001.

[141016]25198.

[141017]Gorman, C.; Padmanabhan, R.; Howard, B. H.: High efficiency DNA-mediated

[141018]transformation of primate cells. Science 221: 551–553, 1983.

[141019]25199.

[141020]Gorman, C. M.: CAT: an easy assay for gene expression (citation

[141021]classic). Current Contents (Life Sciences) 36(22): 8, 1993.

[141022]25200.

[141023]Hildebrand, C. E.; Gonzalez, F. J.; Kozak, C. A.; Nebert, D. W.

[141024]: Regional linkage analysis of the dioxin-inducible P-450 gene family

[141025]on mouse chromosome 9. Biochem. Biophys. Res. Com-

mun. 130: 396–406,

[141026]1985.

[141027]25201.

[141028]Lee, S.; Wu, X.; Reid, M. E.; Zelinski, T.; Redman, C. M.:

Molecular

[141029]basis of the Kell (K1) phenotype. Blood 85: 912–916,

1995.

[141030]25202.

[141031]Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.: The
human

[141032]Kell blood group gene maps to chromosome 7q33 and its
expression is

[141033]restricted to erythroid cells. Blood 81: 2804–2809, 1993.

[141034]25203.

[141035]Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.:

Molecular

[141036]cloning and primary structure of Kell blood group protein.
Proc.

[141037]Nat. Acad. Sci. 88: 6353–6357, 1991.

[141038]25204.

[141039]Marsh, W. L.: Molecular biology of blood groups: cloning

of the

[141040]Kell gene. Transfusion 32: 98–101, 1992.

[141041]25205.

[141042]Morton, N. E.; Krieger, H.; Steinberg, A. G.; Rosenfield, R.
E.

[141043]: Genetic evidence confirming the localization of Sutter in
the Kell

[141044]blood-group system. Vox Sang. 10: 608–613, 1965.

[141045]25206.

[141046]Murphy, M. T.; Morrison, N.; Miles, J. S.; Fraser, R. H.;
Spurr,

[141047]N. K.; Boyd, E.: Regional chromosomal assignment of the
Kell blood

[141048]group locus (KEL) to chromosome 7q33–q35 by fluores-
cence in situ hybridization:

[141049]evidence for the polypeptide nature of antigenic variation.
Hum.

[141050]Genet. 91: 585–588, 1993.

[141051]25207.

[141052]Parke, J. T.; Riccardi, V. M.; Lewis, R. A.; Ferrell, R. E.:

[141053]A syndrome of microcephaly and retinal pigmentary ab-

normalities without

[141054]mental retardation in a family with coincidental autosomal dominant

[141055]hyperreflexia. Am. J. Med. Genet. 17: 585-594, 1984.

[141056]25208.

[141057]Purohit, K. R.; Weber, J. L.; Ward, L. J.; Keats, B. J. B.: The

[141058]Kell blood group locus is close to the cystic fibrosis locus on chromosome

[141059]7. Hum. Genet. 89: 457-458, 1992.

[141060]25209.

[141061]Russo, D.; Wu, X.; Redman, C. M.; Lee, S.: Expression of Kell

[141062]blood group protein in nonerythroid tissues Blood 96: 340-346, 2000.

[141063]25210.

[141064]Stroup, M.; MacIlroy, M.; Walker, R.; Aydelotte, J. V.: Evidence

[141065]that Sutter belongs to the Kell blood group system. Transfusion 5:

[141066]309-314, 1965.

[141067]25211.

[141068]Wagner, T.; Berer, A.; Lanzer, G.; Geissler, K.: Kell is not
[141069]restricted to the erythropoietic lineage but is also ex-
pressed on
[141070]myeloid progenitor cells. Brit. J. Haemat. 110: 409–411,
2000.
[141071]25212.

[141072]Weitkamp, L. R.; Townes, P. L.; Johnston, E.: Linkage data
on
[141073]urinary pepsinogen and the Kell blood group. Birth De-
fects Orig.
[141074]Art. Ser. 11(3): 281–282, 1975. Note: Alternate: Cyto-
genet. Cell
[141075]Genet. 14: 451–452, 1975.
[141076]25213.

[141077]Wimer, B. M.; Marsh, W. L.; Taswell, H. F.: Clinical charac-
teristics
[141078]of the McLeod blood group phenotype. (Abstract) 19th
Annual Meeting
[141079]of the Am. Soc. Hemat., Boston , 1976.
[141080]25214.

[141081]Yu, L.-C.; Twu, Y.-C.; Chang, C.-Y.; Lin, M.: Molecular ba-

sis

[141082]of the Kell-null phenotype: a mutation at the splice site of human

[141083]KEL gene abolishes the expression of Kell blood group antigens. J.

[141084]Biol. Chem. 276: 10247-10252, 2001.

[141085]25215.

[141086]Zelinski, T. A.; Coghlan, G. E.; Myal, Y.; White, L. J.; Philipps,

[141087]S. E.: Assignment of the Kell blood group locus to chromosome 7q.

[141088](Abstract) Cytogenet. Cell Genet. 58: 1927 only, 1991.

[141089]25216.

[141090]Field, L. L.; Marazita, M. L.; Spence, M. A.; Crandall, B. F.;

[141091]Sparkes, R. S.: Is JK linked to IGK on chromosome 2? (Abstract) Cytogenet.

[141092]Cell Genet. 40: 628-629, 1985.

[141093]25217.

[141094]Gedde-Dahl, T.: Personal Communication. Oslo, Norway 9/26/1986.

[141095]25218.

[141096]Geitvik, G. A.; Hoyheim, B.; Gedde-Dahl, T.; Grzeschik, K.
H.;

[141097]Lothe, R.; Tomter, H.; Olaisen, B.: The Kidd (JK) blood
group locus

[141098]assigned to chromosome 18 by close linkage to a DNA-
RFLP. Hum. Genet. 77:

[141099]205-209, 1987.

[141100]25219.

[141101]Hulten, M.; Lindsten, J.; Pen-Ming, L. M.; Fraccaro, M.;
Mannini,

[141102]A.; Trepolo, L.; Robson, E. B.; Heiken, A.; Tellingén, K. G.:
Possible

[141103]localization of the genes for the Kidd blood group on an
autosome

[141104]involved in a reciprocal translocation. Nature 211:
1067-1068, 1968.

[141105]25220.

[141106]Leppert, M.; Ferrell, R.; Kamboh, M. I.; Beasley, J.;
O'Connell,

[141107]P.; Lathrop, M.; Lalouel, J.-M.; White, R.: Linkage of the
polymorphic

[141108]protein markers F13B, C1S, C1R, and blood group antigen

Kidd in CEPH

[141109]reference families. (Abstract) Cytogenet. Cell Genet. 46:
647, 1987.

[141110]25221.

[141111]Lucien, N.; Chiaroni, J.; Cartron, J.-P.; Bailly, P.: Partial

[141112]deletion in the JK locus causing a Jk(null) phenotype.

Blood 99:

[141113]1079–1081, 2002.

[141114]25222.

[141115]Lucien, N.; Sidoux-Walter, F.; Olives, B.; Moulds, J.; Le

Pennec,

[141116]P.-Y.; Cartron, J.-P.; Bailly, P.: Characterization of the gene

encoding

[141117]the human Kidd blood group/urea transporter protein:

evidence for

[141118]splice site mutations in Jk-null individuals. J. Biol. Chem.

273:

[141119]12973–12980, 1998.

[141120]25223.

[141121]Olives, B.; Mattei, M.-G.; Huet, M.; Neau, P.; Martial, S.;

Cartron,

[141122]J.-P.; Bailly, P.: Kidd blood group and urea transport function of
[141123]human erythrocytes are carried by the same protein. J.
Biol. Chem. 270:
[141124]15607–15610, 1995.
[141125]25224.
[141126]Albertsen, H.; Plaetke, R.; Ballard, L.; Fujimoto, E.; Connolly,
[141127]J.; Lawrence, E.; Rodriguez, P.; Robertson, M.; Bradley, P.;
Milner,
[141128]B.; Fuhrman, D.; Marks, A.; Sargent, R.; Cartwright, P.;
Matsunami,
[141129]N.; White, R.: Genetic mapping of the BRCA1 region on
chromosome
[141130]17q21. Am. J. Hum. Genet. 54: 516–525, 1994.
[141131]25225.
[141132]Andersen, T. I.; Borresen, A.-L.; Moller, P.: A common
BRCA1 mutation
[141133]in Norwegian breast and ovarian cancer families? Am. J.
Hum. Genet. 59:
[141134]486–487, 1996.

[141135]25226.

[141136]Anderson, S. F.; Schlegel, B. P.; Nakajima, T.; Wolpin, E. S.;

[141137]Parvin, J. D.: BRCA1 protein is linked to the RNA polymerase II holoenzyme

[141138]complex via RNA helicase A. *Nature Genet.* 19: 254–256, 1998.

[141139]25227.

[141140]Arason, A.; Barkardottir, R. B.; Egilsson, V.: Linkage analysis

[141141]of chromosome 17q markers and breast–ovarian cancer in Icelandic families,

[141142]and possible relationship to prostatic cancer. *Am. J. Hum. Genet.* 52:

[141143]711–717, 1993.

[141144]25228.

[141145]Womack, J. E.: Personal Communication. College Station, Texas

[141146]2/26/1990.

[141147]25229.

[141148]Hiraoka, L. R.; Hsu, L.; Hsieh, C.–L.: Assignment of ALDH3 to

[141149]human chromosome 17p11.2 and ALDH5 to human chromosome 9p13. Genomics 25:

[141150]323–325, 1995.

[141151]25230.

[141152]Vasiliou, V.; Bairoch, A.; Tipton, K. F.; Nebert, D. W.: Eukaryotic

[141153]aldehyde dehydrogenase (ALDH) genes: human polymorphisms, and recommended

[141154]nomenclature based on divergent evolution and chromosomal mapping. Pharmacogenetics 9:

[141155]421–434, 1999.

[141156]25231.

[141157]Harada, S.; Agarwal, D. P.; Goedde, H. W.: Electrophoretic and

[141158]biochemical studies of human aldehyde dehydrogenase isozymes in various

[141159]tissues. Life Sci. 26: 1773–1780, 1980.

[141160]25232.

[141161]Hsu, L. C.; Chang, W.–C.; Yoshida, A.: Cloning of a new human

[141162]aldehyde dehydrogenase gene and comparison with liver

cytosolic ALDH1

[141163]and mitochondrial ALDH2 genes. (Abstract) Am. J. Hum. Genet. 45

[141164](suppl.): A196 only, 1989.

[141165]25233.

[141166]Hsu, L. C.; Chang, W. C.: Cloning and characterization of a new

[141167]functional human aldehyde dehydrogenase gene. J. Biol. Chem. 266:

[141168]12257-12265, 1991.

[141169]25234.

[141170]Ashworth, A.: Two acetyl-CoA acetyltransferase genes located in

[141171]the t-complex region of mouse chromosome 17 partially overlap the

[141172]Tcp-1 and Tcp-1x genes. Genomics 18: 195-198, 1993.

[141173]25235.

[141174]Bennett, M. J.; Hosking, G. P.; Smith, M. F.; Gray, R. G. F.; Middleton,

[141175]B.: Biochemical investigations on a patient with a defect in cytosolic

[141176]acetoacetyl-CoA thiolase, associated with mental retardation. J.

[141177]Inherit. Metab. Dis. 7: 125-128, 1984.

[141178]25236.

[141179]Wunderle, V. M.; Critcher, R.; Hastie, N.; Goodfellow, P. N.;

[141180]Schedl, A.: Deletion of long-range regulatory elements upstream of

[141181]SOX9 causes campomelic dysplasia. Proc. Nat. Acad. Sci. 95: 10649-10654,

[141182]1998.

[141183]25237.

[141184]Young, I. D.; Zuccollo, J. M.; Maltby, E. L.; Broderick, N. J.

[141185]: Campomelic dysplasia associated with a de novo 2q;17q reciprocal

[141186]translocation. J. Med. Genet. 29: 251-252, 1992.

[141187]25238.

[141188]Lynch, H. T.; Schuelke, G. S.; Kimberling, W. J.; Albano, W. A.;

[141189]Lynch, J. F.; Biscione, K. A.; Lipkin, M. L.; Deschner, E. E.; Mikol,

- [141190]Y. B.; Sandberg, A. A.; Elston, R. C.; Bailey-Wilson, J. E.;
Danes,
[141191]B. S.: Hereditary nonpolyposis colorectal cancer (Lynch
syndromes
[141192]I and II). II. Biomarker studies. Cancer 56: 939-951, 1985.
[141193]25239.
[141194]Bourn, D.; Carter, S. A.; Mason, S.; Evans, D. G. R.; Stra-
chan,
[141195]T.: Germline mutations in the neurofibromatosis type 2
tumor suppressor
[141196]gene. Hum. Molec. Genet. 3: 813-816, 1994.
[141197]25240.
[141198]Bouzas, E. A.; Freidlin, V.; Parry, D. M.; Eldridge, R.;
Kaiser-Kupfer,
[141199]M. I.: Lens opacities in neurofibromatosis 2: further sig-
nificant
[141200]correlations. Brit. J. Ophthal. 77: 354-357, 1993.
[141201]25241.
[141202]Bouzas, E. A.; Parry, D. M.; Eldridge, R.; Kaiser-Kupfer, M.
I.
[141203]: Visual impairment in patients with neurofibromatosis 2.

Neurology 43:

[141204]622–623, 1993.

[141205]25242.

[141206]Bruder, C. E. G.; Hirvela, C.; Tapia-Paez, I.; Fransson, I.;
Segraves,

[141207]R.; Hamilton, G.; Zhang, X. X.; Evans, D. G.; Wallace, A. J.;
Baser,

[141208]M. E.; Zucman-Rossi, J.; Hergersberg, M.; and 22 others:
High resolution

[141209]deletion analysis of constitutional DNA from neurofibro-
matosis type

[141210]2 (NF2) patients using microarray-CGH. Hum. Molec.
Genet. 10: 271–282,

[141211]2001.

[141212]25243.

[141213]Claudio, J. O.; Malo, D.; Rouleau, G. A.: The mouse neu-
rofibromatosis

[141214]type 2 gene maps to chromosome 11. Genomics 21:
437–439, 1994.

[141215]25244.

[141216]Egan, R. A.; Thompson, C. R.; MacCollin, M.; Lessell, S.:

Monocular

[141217]levator paresis in neurofibromatosis type 2. Neurology
56: 1222–1224,

[141218]2001.

[141219]25245.

[141220]Eldridge, R.: Central neurofibromatosis with bilateral
acoustic

[141221]neuroma. Adv. Neurol. 29: 57–65, 1981.

[141222]25246.

[141223]Eldridge, R.; Parry, D. M.; Kaiser–Kupfer, M. I.: Neurofibro-
matosis

[141224]2 (NF2): clinical heterogeneity and natural history based
on 39 individuals

[141225]in 9 families and 16 sporadic cases. (Abstract) Am. J. Hum.
Genet. 49

[141226](suppl.): 133, 1991.

[141227]25247.

[141228]Evans, D. G. R.; Birch, J. M.; Ramsden, R. T.: Paediatric
presentation

[141229]of type 2 neurofibromatosis. Arch. Dis. Child. 81:
496–499, 1999.

[141230]25248.

[141231]Evans, D. G. R.; Bourn, D.; Wallace, A.; Ramsden, R. T.;
Mitchell,

[141232]J. D.; Strachan, T.: Diagnostic issues in a family with late
onset

[141233]type 2 neurofibromatosis. J. Med. Genet. 32: 470-474,
1995.

[141234]25249.

[141235]Evans, D. G. R.; Huson, S. M.; Donnai, D.; Neary, W.; Blair,
V.;

[141236]Newton, V.; Strachan, T.; Harris, R.: A genetic study of
type 2 neurofibromatosis

[141237]in the United Kingdom. II. Guidelines for genetic coun-
selling. J.

[141238]Med. Genet. 29: 847-852, 1992.

[141239]25250.

[141240]Evans, D. G. R.; Huson, S. M.; Donnai, D.; Neary, W.; Blair,
V.;

[141241]Teare, D.; Newton, V.; Strachan, T.; Ramsden, R.; Harris,
R.: A genetic

[141242]study of type 2 neurofibromatosis in the United Kingdom.
I. Prevalence,

[141243]mutation rate, fitness, and confirmation of maternal transmission

[141244]effect on severity. J. Med. Genet. 29: 841–846, 1992.

[141245]25251.

[141246]Evans, D. G. R.; Trueman, L.; Wallace, A.; Collins, S.; Strachan,

[141247]T.: Genotype/phenotype correlations in type 2 neurofibromatosis (NF2):

[141248]evidence for more severe disease associated with truncating mutations. J.

[141249]Med. Genet. 35: 450–455, 1998.

[141250]25252.

[141251]Evans, D. G. R.; Wallace, A. J.; Wu, C. L.; Trueman, L.; Ramsden,

[141252]R. T.; Strachan, T.: Somatic mosaicism: a common cause of classic

[141253]disease in tumor-prone syndromes? Lessons from type 2 neurofibromatosis. Am.

[141254]J. Hum. Genet. 63: 727–736, 1998.

[141255]25253.

[141256]Fabricant, R. N.; Todaro, G. J.; Eldridge, R.: Increased lev-

els

[141257]of a nerve-growth factor cross-reacting protein in 'central'
neurofibromatosis. Lancet I:

[141258]4-7, 1979.

[141259]25254.

[141260]Feiling, A.; Ward, E.: A familial form of acoustic tumour.
Brit.

[141261]Med. J. 1: 496-497, 1920.

[141262]25255.

[141263]Fernandez-Valle, C.; Tang, Y.; Ricard, J.; Rodenas-Ruano,
A.;

[141264]Taylor, A.; Hackler, E.; Biggerstaff, J.; Iacovelli, J.: Paxillin

[141265]binds schwannomin and regulates its density-dependent
localization

[141266]and effect on cell morphology. Nature Genet. 31:
354-362, 2002.

[141267]25256.

[141268]Gardner, W. J.; Frazier, C. H.: Hereditary bilateral acoustic

[141269]tumors. J. Hered. 22: 7-8, 1933.

[141270]25257.

[141271]Gardner, W. J.; Frazier, C. H.: Bilateral acoustic neurofibromas.

[141272]A clinical study and survey of a family of five generations with bilateral

[141273]deafness in 38 members. Arch. Neurol. Psychiat. 23: 266–302, 1930.

[141274]25258.

[141275]Gardner, W. J.; Turner, O.: Bilateral acoustic neurofibromas:

[141276]further clinical and pathologic data on hereditary deafness and Recklinghausen's

[141277]disease. Arch. Neurol. Psychiat. 44: 76–99, 1940.

[141278]25259.

[141279]Giovannini, M.; Robanus-Maandag, E.; van der Valk, M.; Niwa-Kawakita,

[141280]M.; Abramowski, V.; Goutebroze, L.; Woodruff, J. M.; Berns, A.; Thomas,

[141281]G.: Conditional biallelic Nf2 mutation in the mouse promotes manifestations

[141282]of human neurofibromatosis type 2. Genes Dev. 14: 1617–1630, 2000.

[141283]25260.

[141284]Gutmann, D. H.: The neurofibromatoses: when less is more. Hum.

[141285]Molec. Genet. 10: 747-755, 2001.

[141286]25261.

[141287]Gutmann, D. H.; Haipek, C. A.; Burke, S. P.; Sun, C.-X.;
Scoles,

[141288]D. R.; Pulst, S. M.: The NF2 interactor, hepatocyte growth
factor-regulated

[141289]tyrosine kinase substrate (HRS), associates with merlin in
the 'open'

[141290]conformation and suppresses cell growth and motility.
Hum. Molec.

[141291]Genet. 10: 825-834, 2001.

[141292]25262.

[141293]Gutmann, D. H.; Hirbe, A. C.; Haipek, C. A.: Functional
analysis

[141294]of neurofibromatosis 2 (NF2) missense mutations. Hum.
Molec. Genet. 10:

[141295]1519-1529, 2001.

[141296]25263.

[141297]Gutmann, D. H.; Sherman, L.; Seftor, L.; Haipek, C.; Lu, K.

H.;

[141298]Hendrix, M.: Increased expression of the NF2 tumor suppressor gene

[141299]product, merlin, impairs cell motility, adhesion and spreading. Hum.

[141300]Molec. Genet. 8: 267-275, 1999.

[141301]25264.

[141302]Jacoby, L. B.; Jones, D.; Davis, K.; Kronn, D.; Short, M. P.;

[141303]Gusella, J.; MacCollin, M.: Molecular analysis of the NF2 tumor-suppressor

[141304]gene in schwannomatosis. Am. J. Hum. Genet. 61: 1293-1302, 1997.

[141305]25265.

[141306]Kaiser-Kupfer, M. I.; Freidlin, V.; Datiles, M. B.; Edwards, P.

[141307]A.; Sherman, J. L.; Parry, D.; McCain, L. M.; Eldridge, R.:

The association

[141308]of posterior capsular lens opacities with bilateral acoustic neuromas

[141309]in patients with neurofibromatosis type 2. Arch. Ophthalmol. 107: 541-544,

[141310]1989.

[141311]25266.

[141312]Kanter, W. R.; Eldridge, R.; Fabricant, R.; Allen, J. C.; Koerber,

[141313]T.: Central neurofibromatosis with bilateral acoustic neuroma: genetic,

[141314]clinical and biochemical distinctions from peripheral neurofibromatosis. *Neurology* 30:

[141315]851–859, 1980.

[141316]25267.

[141317]Kehrer-Sawatzki, H.; Udart, M.; Krone, W.; Baden, R.; Fahsold,

[141318]R.; Thomas, G.; Schmucker, B.; Assum, G.: Mutational analysis and

[141319]expression studies of the neurofibromatosis type 2 (NF2) gene in a

[141320]patient with a ring chromosome 22 and NF2. *Hum. Genet.* 100: 67–74,

[141321]1997.

[141322]25268.

[141323]Kedes, L.; Ng, S.-Y.; Lin, C.-S.; Gunning, P.; Eddy, R.; Shows,

[141324]T.; Leavitt, J.: The human beta-actin multigene family.
Trans. Assoc.

[141325]Am. Phys. 98: 42-46, 1985.

[141326]25269.

[141327]Leavitt, J.; Bushar, G.; Kakunaga, T.; Hamada, H.; Hirakawa, T.;

[141328]Goldman, D.; Merrill, C.: Variations in expression of mutant beta-actin

[141329]accompanying incremental increases in human fibroblast tumorigenicity. Cell 28:

[141330]259-268, 1982.

[141331]25270.

[141332]Nakajima-Iijima, S.; Hamada, H.; Reddy, P.; Kakunaga, T.:
Molecular

[141333]structure of the human cytoplasmic beta-actin gene; interspecies homology

[141334]of sequences in the introns. Proc. Nat. Acad. Sci. 82:
6133-6137,

[141335]1985.

[141336]25271.

[141337]Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.;

Kedes,

[141338]L.; Shows, T.: Chromosome 7 assignment of the human
beta-actin functional

[141339]gene (ACTB) and the chromosomal dispersion of pseudo-
genes. (Abstract) Cytogenet.

[141340]Cell Genet. 40: 712 only, 1985.

[141341]25272.

[141342]Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.;
Shows,

[141343]T.; Kedes, L.: Evolution of the functional human beta-actin
gene

[141344]and its multi-pseudogene family: conservation of the
noncoding regions

[141345]and chromosomal dispersion of pseudogenes. Molec. Cell.
Biol. 5:

[141346]2720-2732, 1985.

[141347]25273.

[141348]Toyama, S.; Toyama, S.: A variant form of beta-actin in a
mutant

[141349]of KB cells resistant to cytochalasin B. Cell 37: 609-614,
1984.

[141350]25274.

[141351] Vandekerckhove, J.; Weber, K.: Mammalian cytoplasmic
actins are
[141352] the products of at least two genes and differ in primary
structure
[141353] in at least 25 identified positions from skeletal muscle
actins. Proc.
[141354] Nat. Acad. Sci. 75: 1106–1110, 1978.
[141355] 25275.
[141356] Alderdice, P. W.; Kaita, H.; Lewis, M.; McAlpine, P. J.;
Wong,
[141357] P.; Anderson, J.; Giblett, E. R.: Segregation of marker loci
in families
[141358] with an inherited paracentric insertion of chromosome 9.
Am. J. Hum.
[141359] Genet. 39: 612–617, 1986.
[141360] 25276.
[141361] Beutler, E.; Carson, D.; Dannawi, H.; Forman, L.; Kuhl, W.;
West,
[141362] C.; Westwood, B.: Metabolic compensation for profound
erythrocyte
[141363] adenylate kinase deficiency: a hereditary enzyme defect
without hemolytic

[141364]anemia. J. Clin. Invest. 72: 648–655, 1983.

[141365]25277.

[141366]Beutler, E.; Carson, D. A.; Dannawi, H.; Forman, L.; Kuhl, W.;

[141367]West, C.; Westwood, B.: Red cell adenylate kinase deficiency: another

[141368]non-disease? (Abstract) Blood 60: 33A only, 1982.

[141369]25278.

[141370]Bianchi, P.; Zappa, M.; Bredi, E.; Vercellati, C.; Pelissero, G.;

[141371]Barraco, F.; Zanella, A.: A case of complete adenylate kinase deficiency

[141372]due to a nonsense mutation in AK-1 gene (arg107-to-stop, CGA-to-TGA)

[141373]associated with chronic haemolytic anaemia. Brit. J. Haemat. 105:

[141374]75–79, 1999.

[141375]25279.

[141376]Boivin, P.; Galand, C.; Hakim, J.; Simony, D.; Seligman, M.: Une

[141377]nouvelle erythroenzymopathie: anemie hemolytique con-

genitale non spherocytaire

[141378]et deficit hereditaire en adenylate-kinase erythrocytaire.

Presse

[141379]Med. 79: 215-218, 1971.

[141380]25280.

[141381]Boivin, P.; Galand, C.; Hakim, J.; Simony, D.; Seligman, M.:

Deficit

[141382]congenital en adenylate-kinase erythrocytaire. (Letter)

Presse Med. 78:

[141383]1443 only, 1970.

[141384]25281.

[141385]Bowman, J. E.; Frischer, H.; Ajmar, F.; Carson, P. E.; Gower,

M.

[141386]K.: Population, family and biochemical investigation of
human adenylate

[141387]kinase polymorphism. Nature 214: 1156-1158, 1967.

[141388]25282.

[141389]Brock, D. J. H.: Evidence against a common subunit in
adenylate

[141390]kinase and pyruvate kinase. Humangenetik 10: 30-34,
1970.

[141391]25283.

[141392]Cavalli-Sforza, L. L.; King, M. C.; Go, R. C. P.; Namboodiri, K.

[141393]K.; Lynch, H. T.; Wong, L.; Kaplan, E. B.; Elston, R. C.: Possible

[141394]linkage between transcobalamin II (TC II) and adenylate kinase (AK).

[141395](Abstract) Cytogenet. Cell Genet. 25: 140-141, 1979.

[141396]25284.

[141397]Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Slaughter, C. A.;

[141398]Gray, J. E.; Blank, C. E.; James, F. E.; Ridler, M. A. C.; Insley,

[141399]J.; Hulten, M.: Segregation of ABO, AK(1) and ACONs in families with

[141400]abnormalities of chromosome 9. Ann. Hum. Genet. 41: 365-377, 1978.

[141401]25285.

[141402]Fenger, K.; Sorensen, S. A.: Evaluation of a possible sex difference

[141403]in recombination for the ABO-AK linkage. Am. J. Hum. Genet. 27:

[141404]784–788, 1975.

[141405]25286.

[141406]Ferguson-Smith, M. A.; Aitken, D. A.; Turleau, C.; de
Grouchy,

[141407]J.: Localisation of the human ABO: Np-1: AK-1 linkage
group by regional

[141408]assignment of AK-1 to 9q34. Hum. Genet. 34: 35–43,
1976.

[141409]25287.

[141410]Fildes, R. A.; Harris, H.: Genetically determined variation
of

[141411]adenylate kinase in man. Nature 209: 261–262, 1966.

[141412]25288.

[141413]Geurts van Kessel, A. H. M.; Hagemeijer, A.; Westerveld,
A.; Meera

[141414]Khan, P.; de Groot, P. G.; Pearson, P. L.: Characterization
of chromosomal

[141415]abnormalities in chronic myeloid leukemia using somatic
cell hybrids.

[141416](Abstract) Cytogenet. Cell Genet. 32: 280 only, 1982.

[141417]25289.

- [141418]Lachant, N. A.; Zerez, C. R.; Barredo, J.; Lee, D. W.; Savely,
[141419]S. M.; Tanaka, K. R.: Hereditary erythrocyte adenylate ki-
nase deficiency:
[141420]a defect of multiple phosphotransferases? Blood 77:
2774-2784, 1991.
[141421]25290.
- [141422]Matsuura, S.; Igarashi, M.; Tanizawa, Y.; Yamada, M.;
Kishi, F.;
[141423]Kajii, T.; Fujii, H.; Miwa, S.; Sakurai, M.; Nakazawa, A.: Hu-
man
[141424]adenylate kinase deficiency associated with hemolytic
anemia: a single
[141425]base substitution affecting solubility and catalytic activity
of the
[141426]cytosolic adenylate kinase. J. Biol. Chem. 264:
10148-10155, 1989.
[141427]25291.
- [141428]Miwa, S.; Fujii, H.; Tani, K.; Takahashi, K.; Takizawa, T.;
Igarashi,
[141429]T.: Red cell adenylate kinase deficiency associated with
hereditary
[141430]nonspherocytic hemolytic anemia: clinical and biochemical

studies. Am.

[141431]J. Hemat. 14: 325-333, 1983.

[141432]25292.

[141433]Mohandas, T.; Sparkes, R. S.; Sparkes, M. C.; Shulkin, J. D.;

[141434]Toomey, K. E.; Funderburk, S. J.: Regional localization of
human

[141435]gene loci on chromosome 9: studies of somatic cell hy-
brids containing

[141436]human translocations. Am. J. Hum. Genet. 31: 586-600,
1979.

[141437]25293.

[141438]Povey, S.; Slaughter, C. A.; Wilson, D. E.; Gormley, I. P.;
Buckton,

[141439]K. E.; Perry, P.; Bobrow, M.: Evidence for the assignment of
loci

[141440]AK 1, AK 3 and ACON to chromosome 9 in man. Ann.
Hum. Genet. 39:

[141441]413-422, 1976.

[141442]25294.

[141443]Qualtieri, A.; Pedace, V.; Bisconte, M. G.; Bria, M.; Gulino,

[141444]B.; Andreoli, V.; Brancati, C.: Severe erythrocyte adenylate

kinase

[141445]deficiency due to homozygous A-to-G substitution at
codon 164 of human

[141446]AK1 gene associated with chronic haemolytic anaemia.
Brit. J. Haemat. 99:

[141447]770-776, 1997.

[141448]25295.

[141449]Rapley, S.; Robson, E. B.; Harris, H.; Smith, S. M.: Data on
[141450]the incidence, segregation and linkage relations of the
adenylate

[141451]kinase (AK) polymorphism. Ann. Hum. Genet. 31:
237-242, 1967.

[141452]25296.

[141453]Ohno, S.: Original domain for the serum albumin family
arose

[141454]from repeated sequences. Proc. Nat. Acad. Sci. 78:
7657-7661, 1981.

[141455]25297.

[141456]Ott, H.: Analbuminemia. In: Linneweh, F.: Erbliche Stoff-
wechselkrankheiten.

[141457]Munich: Urban und Schwarzenberg (pub.) 1962. Pp. 44.

[141458]25298.

[141459]Peach, R. J.; Brennan, S. O.: Structural characterization of

[141460]a glycoprotein variant of human serum albumin: albumin

Casebrook (494

[141461]asp-to-asn). Biochim. Biophys. Acta 1097: 49-54, 1991.

[141462]25299.

[141463]Peach, R. J.; Fellowes, A. P.; Brennan, S. O.; George, P. M.:

[141464]Albumin Rugby Park: a truncated albumin variant caused

by a G-to-C

[141465]splice-site mutation in intron 13. Biochim. Biophys. Acta

1180:

[141466]107-110, 1992.

[141467]25300.

[141468]Petersen, C. E.; Scottolini, A. G.; Cody, L. R.; Mandel, M.;

Reimer,

[141469]N.; Bhagavan, N. V.: A point mutation in the human serum

albumin

[141470]gene results in familial dysalbuminaemic hyperthyroxi-

naemia. J. Med.

[141471]Genet. 31: 355-359, 1994.

[141472]25301.

[141473]Pinkert, C. A.; Ornitz, D. M.; Brinster, R. L.; Palmiter, R. D.

[141474]: An albumin enhancer located 10 kb upstream functions
along with

[141475]its promoter to direct efficient, liver-specific expression in
transgenic

[141476]mice. Genes Dev. 1: 268-276, 1987.

[141477]25302.

[141478]Pohlenz, J.; Sadow, P. M.; Koffler, T.; Schonberger, W.;
Weiss,

[141479]R. E.; Refetoff, S.: Congenital hypothyroidism in a child
with unsuspected

[141480]familial dysalbuminemic hyperthyroxinemia caused by a
mutation (R218H)

[141481]in the human albumin gene. J. Pediatr. 139: 887-891,
2001.

[141482]25303.

[141483]Prager, E. M.; Wilson, A. C.; Lowenstein, J. M.; Sarich, V. M.

[141484]: Mammoth albumin. Science 209: 287-289, 1980.

[141485]25304.

[141486]Premachandra, B. N.; Wolfe, B.; Williams, I. K.: Coexistence
[141487]of familial dysalbuminemic hyperthyroxinemia with famil-
ial hypercholesterolemia
[141488]and multiple lipoprotein type hyperlipidemia. Am. J. Med.
84: 345-351,
[141489]1988.
[141490]25305.
[141491]Putnam, F. W.: Personal Communication. Bloomington,
Ind. 8/4/1993.
[141492]25306.
[141493]Rajatanavin, R.; Fournier, L.; DeCosimo, D.; Abreau, C.;
Braverman,
[141494]L. E.: Elevated serum free thyroxine by thyroxine analog
radioimmunoassays
[141495]in euthyroid patients with familial dysalbuminemic hyper-
thyroxinemia. Ann.
[141496]Intern. Med. 97: 865-866, 1982.
[141497]25307.
[141498]Rajatanavin, R.; Young, R. A.; Braverman, L. E.: Effect of
chloride
[141499]on serum thyroxine binding in familial dysalbuminemic

hyperthyroxinemia. J.

[141500]Clin. Endocr. Metab. 58: 388–391, 1984.

[141501]25308.

[141502]Rochu, D.; Fine, J. M.: New method for identifying genetic variants

[141503]of human proalbumin. Clin. Chem. 32: 2063–2065, 1986.

[141504]25309.

[141505]Ruffner, D. E.; Dugaiczky, A.: Splicing mutation in human hereditary

[141506]analbuminemia. Proc. Nat. Acad. Sci. 85: 2125–2129, 1988.

[141507]25310.

[141508]Ruiz, M.; Rajatanavin, R.; Young, R. A.; Taylor, C.; Brown, R.;

[141509]Braverman, L. E.; Ingbar, S. H.: Familial dysalbuminemic hyperthyroxinemia:

[141510]a syndrome that can be confused with thyrotoxicosis. New Eng. J.

[141511]Med. 306: 635–639, 1982.

[141512]

[141513]100. Sakamoto, Y.; Davis, E.; Madison, J.; Watkins, S.;

McLaughlin,

[141514]H.; Leahy, D. T.; Putnam, F. W.: Purification and structural study

[141515]of two albumin variants in an Irish population. Clin. Chim. Acta 204:

[141516]179–188, 1991.

[141517]

[141518]101. Sanders, G. T. B.; Tarnoky, A. L.: Albumin Amsterdam: a new

[141519]European albumin variant. IRCS Med. Sci. 7: 581 only, 1979.

[141520]

[141521]102. Sarcione, E. J.; Aungst, C. W.: Studies in bisalbuminemia: binding

[141522]properties of the two albumins. Blood 20: 156–164, 1962.

[141523]

[141524]103. Sargent, T. D.; Wu, J.-R.; Sala-Trepat, J. M.; Wallace, R. B.;

[141525]Reyes, A. A.; Bonner, J.: The rat serum albumin gene: analysis of

[141526]cloned sequences. Proc. Nat. Acad. Sci. 76: 3256–3260,

1979.

[141527]

[141528]104. Sarich, V. M.: Generation time and albumin evolution.
Biochem.

[141529]Genet. 7: 205–212, 1972.

[141530]

[141531]105. Schell, L. M.; Agarwal, S. S.; Blumberg, B. S.; Levy, H.;
Bennett,

[141532]H.; Laughlin, W. S.; Martin, J. P.: Distribution of albumin
variants

[141533]Naskapi and Mexico among Aleuts, Frobisher Bay Eski-
mos, and Micmac,

[141534]Naskapi, Mohawk, Omaha and Apache Indians. Am. J.
Phys. Anthrop. 49:

[141535]111–118, 1978.

[141536]

[141537]106. Schell, L. M.; Blumberg, B. S.: The genetics of human
serum

[141538]albumin.In: Rosenoer, V. M.; Oratz, M.; Rothschild, M. A.:
Albumin

[141539]Structure, Function and Uses. Oxford: Pergamon Press

(pub.) 1977.

[141540]Pp. 113–141.

[141541]

[141542]107. Shalaby, F.; Shafritz, D. A.: Exon skipping during
splicing

[141543]of albumin mRNA precursors in Nagase analbuminemic
rats. Proc. Nat.

[141544]Acad. Sci. 87: 2652–2656, 1990.

[141545]

[141546]108. Shashaty, G.; Atamer, M.: Acquired bisalbuminemia
with hyperamylasemia. Digest.

[141547]Dis. 17: 59–67, 1972.

[141548]

[141549]109. Shibata, T.; Abe, T.: Linkage between the loci for
serum albumin

[141550]and vitamin D binding protein (GC) in the Japanese quail.
Animal

[141551]Genet. 27: 195–197, 1996.

[141552]

[141553]110. Silverberg, J. D. H.; Premachandra, B. N.: Familial hy-

perthyroxinemia

[141554] due to abnormal thyroid hormone binding. Ann. Intern. Med. 96: 183–186,

[141555] 1982.

[141556]

[141557] 111. Sunthornthepvarakul, T.; Likitmaskul, S.; Ngown-garmratana, S.;

[141558] Angsusingha, K.; Kitvitayasak, S.; Scherberg, N. H.; Refetoff, S.

[141559]: Familial dysalbuminemic hypertriiodothyroninemia: a new, dominantly

[141560] inherited albumin defect. J. Clin. Endocr. Metab. 83: 1448–1454,

[141561] 1998.

[141562]

[141563] 112. Swain, B. K.; Talukder, G.; Sharma, A.: Bisalbuminaemia: reports

[141564] from Calcutta. Biomedicine 33: 172–173, 1980.

[141565]

[141566] 113. Takahashi, N.; Takahashi, Y.; Blumberg, B. S.; Putnam, F. W.

[141567]: Amino acid substitutions in genetic variants of human serum albumin

[141568]and in sequences inferred from molecular cloning. Proc. Nat. Acad.

[141569]Sci. 84: 4413–4417, 1987.

[141570]

[141571]114. Takahashi, N.; Takahashi, Y.; Isobe, T.; Putnam, F. W.; Fujita,

[141572]M.; Satoh, C.; Neel, J. V.: Amino acid substitutions in inherited

[141573]albumin variants from Amerindian and Japanese populations. Proc.

[141574]Nat. Acad. Sci. 84: 8001–8005, 1987.

[141575]

[141576]115. Takahashi, N.; Takahashi, Y.; Putnam, F. W.: Structural changes

[141577]and metal binding by proalbumins and other amino-terminal genetic

[141578]variants of human serum albumin. Proc. Nat. Acad. Sci. 84: 7403–7407,

[141579]1987.

[141580]

[141581]116. Tarnoky, A. L.; Lestas, A. N.: A new type of bisalbuminaemia. Clin.

[141582]Chim. Acta 9: 551-558, 1964.

[141583]

[141584]117. Urano, Y.; Sakai, M.; Watanabe, K.; Tamaoki, T.: Tandem arrangement

[141585]of the albumin and alpha-fetoprotein genes in the human genome. Gene 32:

[141586]255-261, 1984.

[141587]

[141588]118. Vanzetti, G.; Porta, F.; Prencipe, L.; Scherini, A.; Fraccaro,

[141589]M.: A homozygote for a serum albumin variant of the fast type. Hum.

[141590]Genet. 46: 5-9, 1979.

[141591]

[141592]119. Vaysse, J.; Pilardeau, P.; Garnier, M.: Trisalbuminemia. (Letter) New

[141593]Eng. J. Med. 305: 833-834, 1981.

[141594]

[141595]120. Wada, N.; Chiba, H.; Shimizu, C.; Kijima, H.; Kubo, M.; Koike

[141596]T.: A novel missense mutation in codon 218 of the albumin gene in

[141597]a distinct phenotype of familial dysalbuminemic hyperthyroxinemia

[141598]in a Japanese kindred. J. Clin. Endocr. Metab. 82: 3246–3250, 1997.

[141599]

[141600]121. Watkins, S.; Madison, J.; Galliano, M.; Minchiotti, L.; Putnam,

[141601]F. W.: A nucleotide insertion and frameshift cause analbuminemia

[141602]in an Italian family. Proc. Nat. Acad. Sci. 91: 2275–2279, 1994.

[141603]

[141604]122. Weitkamp, L. R.: Comparative gene mapping: linkage between the

[141605]albumin and Gc loci in the horse. (Abstract) Am. J. Hum. Genet. 30:

[141606]128A only, 1978.

[141607]

[141608]123. Weitkamp, L. R.; Buck, A. A.: Phenotype frequencies
for four

[141609]serum proteins in Afghanistan: two 'new' albumin variants.
Humangenetik 15:

[141610]335-340, 1972.

[141611]

[141612]124. Weitkamp, L. R.; Chagnon, N. A.: Albumin Maku: a
new variant

[141613]of human serum albumin. Nature 217: 759-760, 1968.

[141614]

[141615]125. Weitkamp, L. R.; Franglen, G.; Rokala, D. A.; Polesky,
H. F.;

[141616]Simpson, N. E.; Sunderman, F. W., Jr.; Bell, H. E.; Saave, J.;
Lisker,

[141617]R.; Bohls, S. W.: An electrophoretic comparison of human
serum albumin

[141618]variants: eight distinguishable types. Hum. Hered. 19:
159-169,

[141619]1969.

[141620]

- [141621]126. Weitkamp, L. R.; Renwick, J. H.; Berger, J. P.; Shreffler, D.
C.; Drachmann, O.; Wuhrmann, F.; Braend, M.; Franglen, G.: Additional
data and summary for albumin-GC linkage in man. Hum. Hered. 20:
1-7, 1970.
- [141625]
- [141626]127. Weitkamp, L. R.; Robson, E. B.; Shreffler, D. C.; Corney, G.
An unusual human serum albumin variant: further data
on genetic
linkage between loci for human serum albumin and
group-specific component
(GC). Am. J. Hum. Genet. 20: 392-397, 1968.
- [141630]
- [141631]128. Weitkamp, L. R.; Rucknagel, D. L.; Gershowitz, H.:
Genetic linkage
between structural loci for albumin and group specific
component (GC). Am.
J. Hum. Genet. 18: 559-571, 1966.

[141634]

[141635]129. Weitkamp, L. R.; Salzano, F. M.; Neel, J. V.; Porta, F.;
Geerdink,

[141636]R. A.; Tarnoky, A. L.: Human serum albumin: twenty-three
genetic

[141637]variants and their population distribution. Ann. Hum.
Genet. 36:

[141638]381–392, 1973.

[141639]

[141640]130. Weitkamp, L. R.; Shreffler, D. C.; Robbins, J. L.;
Drachmann,

[141641]O.; Adner, P. L.; Weime, R. J.; Simon, N. M.; Cooke, K. B.;
Sandor,

[141642]G.; Wuhrmann, F.; Braend, M.; Tarnoky, A. L.: An elec-
trophoretic

[141643]comparison of serum albumin variants from nineteen un-
related families. Acta

[141644]Genet. Statist. Med. 17: 399–405, 1967.

[141645]

[141646]131. Wieme, R. J.: On the presence of two albumins in
certain normal

[141647]human sera and its genetic determination. Clin. Chim.
Acta 5: 443–445,

[141648]1960.

[141649]

[141650]132. Williams, D. I.; Martin, N. H.: Bisalbuminemia with
curious

[141651]acrocyanotic skin changes (two cases). Proc. Roy. Soc.
Med. 53:

[141652]566–568, 1960.

[141653]

[141654]133. Yabu, Y.; Amir, S. M.; Ruiz, M.; Braverman, L. E.; Ing-
bar, S.

[141655]H.: Heterogeneity of thyroxine binding by serum albumins
in normal

[141656]subjects and patients with familial dysalbuminemic hyper-
thyroxinemia. J.

[141657]Clin. Endocr. Metab. 60: 451–459, 1985.

[141658]

[141659]134. Yabu, Y.; Miyai, K.; Kobayashi, A.; Miki, K.; Doi, K.;
Takamatsu,

[141660]J.; Mozai, T.; Matsuzuka, F.; Kuma, K.: A new type of albumin with

[141661]predominantly increased binding affinity for
3,3–prime,5–triiodothyronine

[141662]in a patient with Graves' disease. J. Endocr. Invest. 10:
163–169,

[141663]1987.

[141664]

[141665]135. Yeo, P. P. B.; Yabu, Y.; Etzkorn, J. R.; Rajatanavin, R.;
Braverman,

[141666]L. E.; Ingbar, S. H.: A four generation study of dysalbuminemic hyperthyroxinemia:

[141667]diagnosis in the presence of an acquired excess of thyroxine–binding

[141668]globulin. J. Endocr. Invest. 10: 33–38, 1987.

[141669]

[141670]136. Ying, Q.; Liang, Z.; Wu, H.; Wang, L.: The gene fre–

quency of

[141671]serum albumin variants in Chinese and the electrophoretic
characterization

[141672]of several serum albumin variants. Scientia Sinica 24:
1597–1602,

[141673]1981.

[141674]25311.

[141675]Osier, M. V.; Pakstis, A. J.; Soodyall, H.; Comas, D.; Gold-
man,

[141676]D.; Odunsi, A.; Okonofua, F.; Parnas, J.; Schulz, L. O.;
Bertranpetit,

[141677]J.; Bonne–Tamir, B.; Lu, R.–B.; Kidd, J. R.; Kidd, K. K.: A
global

[141678]perspective on genetic variation at the ADH genes reveals
unusual

[141679]patterns of linkage disequilibrium and diversity. Am. J.
Hum. Genet. 71:

[141680]84–99, 2002.

[141681]25312.

[141682]Comings, D. E.; Gade–Andavolu, R.; Gonzalez, N.; Blake,
H.; Wu,

[141683]S.; MacMurray, J. P.: Additive effect of three noradrenergic

genes

[141684](ADRA2A, ADRA2C, DBH) on attention-deficit hyperactivity disorder

[141685]and learning disabilities in Tourette syndrome subjects.

Clin. Genet. 55:

[141686]160–172, 1999.

[141687]25313.

[141688]Halperin, J. M.; Newcorn, J. H.; Koda, V. H.; Pick, L.; McKay,

[141689]K. E.; Knott, P.: Noradrenergic mechanisms in ADHD children with

[141690]and without reading disabilities: a replication and extension. J.

[141691]Am. Acad. Child Adolesc. Psychiat. 36: 1688–1697, 1997.

[141692]25314.

[141693]Hein, L.; Altman, J. D.; Kobilka, B. K.: Two functionally distinct

[141694]alpha-2-adrenergic receptors regulate sympathetic neurotransmission. Nature 402:

[141695]181–184, 1999.

[141696]25315.

[141697]Hoehe, M.; Berrettini, W.; Leppert, M.; Lalouel, J.-M.; Byer-

ley,

[141698]W.; Gershon, E.; White, R.: Genetic mapping of adrenergic receptor

[141699]genes. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A143 only, 1989.

[141700]25316.

[141701]Hoehe, M. R.; Berrettini, W. H.; Lentes, K.-U.: Dra I identifies

[141702]a two allele DNA polymorphism in the human alpha-2-adrenergic receptor

[141703]gene (ADRAR), using a 5.5 kb probe (p ADRAR). Nucleic Acids Res. 16:

[141704]9070 only, 1988.

[141705]25317.

[141706]Kobilka, B. K.; Matsui, H.; Kobilka, T. S.; Yang-Feng, T. L.; Francke,

[141707]U.; Caron, M. G.; Lefkowitz, R. J.; Regan, J. W.: Cloning, sequencing,

[141708]and expression of the gene coding for the human platelet alpha-2-adrenergic

[141709]receptor. Science 238: 650-656, 1987.

[141710]25318.

[141711]Oakey, R. J.; Caron, M. G.; Lefkowitz, R. J.; Seldin, M. F.:
Genomic

[141712]organization of adrenergic and serotonin receptors in the
mouse: linkage

[141713]mapping of sequence-related genes provides a method
for examining

[141714]mammalian chromosome evolution. Genomics 10:
338-344, 1991.

[141715]25319.

[141716]Philipp, M.; Brede, M. E.; Hadamek, K.; Gessler, M.; Lohse,
M.

[141717]J.; Hein, L.: Placental alpha-2-adrenoceptors control vas-
cular development

[141718]at the interface between mother and embryo. Nature
Genet. 31: 311-315,

[141719]2002.

[141720]25320.

[141721]Surprenant, A.; Horstman, D. A.; Akbarali, H.; Limbird, L.
E.:

[141722]A point mutation of the alpha-2-adrenoceptor that blocks
coupling

[141723]to potassium but not calcium currents. Science 257:

977-980, 1992.

[141724]25321.

[141725]Yang-Feng, T. L.; Kobilka, B. K.; Caron, M. G.; Lefkowitz, R.

[141726]J.; Francke, U.: Chromosomal assignment of genes for an alpha-adrenergic

[141727]receptor (ADRAR) and for another member of this receptor family coupled

[141728]to guanine nucleotide regulatory proteins (RG21). (Abstract) Cytogenet.

[141729]Cell Genet. 46: 722-723, 1987.

[141730]25322.

[141731]Bruno, J. F.; Whittaker, J.; Song, J.; Berelowitz, M.: Molecular

[141732]cloning and sequencing of a cDNA encoding a human alpha-1A adrenergic

[141733]receptor. Biochem. Biophys. Res. Commun. 179: 1485-1490, 1991.

[141734]25323.

[141735]Esbenshade, T. A.; Hirasawa, A.; Tsujimoto, G.; Tanaka, T.; Yano,

[141736]J.; Minneman, K. P.; Murphy, T. J.: Cloning of the human

alpha-1D-adrenergic

[141737]receptor and inducible expression of three subtypes in
SK-N-MC cells. Molec.

[141738]Pharm. 47: 977-985, 1995.

[141739]25324.

[141740]Forray, C.; Bard, J. A.; Wetzel, J. M.; Chiu, G.; Shapiro, E.;

[141741]Tang, R.; Lepor, H.; Hartig, P. R.; Weinshank, R. L.;
Branchek, T.

[141742]A.; Gluchowski, C.: The alpha-1-adrenergic receptor that
mediates

[141743]smooth muscle contraction in human prostate has the
pharmacological

[141744]properties of the cloned human alpha-1C subtype. Molec.
Pharm. 45:

[141745]703-708, 1994.

[141746]25325.

[141747]Loftus, S. K.; Shiang, R.; Warrington, J. A.; Bengtsson, U.;

McPherson,

[141748]J. D.; Wasmuth, J. J.: Genes encoding adrenergic receptors
are not

[141749]clustered on the long arm of human chromosome 5. Cyto-

genet. Cell

[141750]Genet. 67: 69–74, 1994.

[141751]25326.

[141752]Lomasney, J. W.; Cotecchia, S.; Lorenz, W.; Leung, W.-Y.;
Schwinn,

[141753]D. A.; Yang-Feng, T. L.; Brownstein, M.; Lefkowitz, R. J.;
Caron,

[141754]M. G.: Molecular cloning and expression of the cDNA for
the alpha-1A-adrenergic

[141755]receptor: the gene for which is located on human chromo-
some 5. J.

[141756]Biol. Chem. 266: 6365–6369, 1991.

[141757]25327.

[141758]Schwinn, D. A.; Johnston, G. I.; Page, S. O.; Mosley, M. J.;
Wilson,

[141759]K. H.; Worman, N. P.; Campbell, S.; Fidock, M. D.; Furness,
L. M.;

[141760]Parry-Smith, D. J.; Peter, B.; Bailey, D. S.: Cloning and
pharmacological

[141761]characterization of human alpha-1 adrenergic receptors:
sequence corrections

[141762]and direct comparison with other species homologues. J.

Pharm. Exp.

[141763]Ther. 272: 134–142, 1995.

[141764]25328.

[141765]Schwinn, D. A.; Lomasney, J. W.: Pharmacologic characterization

[141766]of cloned alpha-1-adrenoceptor subtypes: selective antagonists suggest

[141767]the existence of a fourth subtype. Europ. J. Pharm. 227: 433–436,

[141768]1992.

[141769]25329.

[141770]Weinberg, D. H.; Trivedi, P.; Tan, C. P.; Mitra, S.; Perkins-Barrow,

[141771]A.; Borkowski, D.; Strader, C. D.; Bayne, M.: Cloning, expression

[141772]and characterization of human alpha adrenergic receptors alpha-1A,

[141773]alpha-1B, and alpha-1C. Biochem. Biophys. Res. Commun. 201: 1296–1304,

[141774]1994.

[141775]25330.

[141776]Yang-Feng, T. L.; Han, H.; Lomasney, J. W.; Caron, M. G.:
Localization
[141777]of the cDNA for an alpha-1-adrenergic receptor subtype
(ADRA1D) to
[141778]chromosome band 20p13. Cytogenet. Cell Genet. 66:
170-171, 1994.
[141779]25331.

[141780]Markert, M. L.; Hershfield, M. S.; Wiginton, D. A.; States, J.
[141781]C.; Ward, F. E.; Bigner, S. H.; Buckley, R. H.; Kaufman, R.
E.; Hutton,
[141782]J. J.: Identification of a deletion in the adenosine deami-
nase gene
[141783]in a child with severe combined immunodeficiency. J. Im-
mun. 138:
[141784]3203-3206, 1987.
[141785]25332.

[141786]Markert, M. L.; Hutton, J. J.; Wiginton, D. A.; States, J. C.;
[141787]Kaufman, R. E.: Adenosine deaminase (ADA) deficiency
due to deletion
[141788]of the ADA gene promoter and first exon by homologous
recombination

[141789]between two Alu elements. J. Clin. Invest. 81: 1323–1327, 1988.

[141790]25333.

[141791]Markert, M. L.; Norby–Slycord, C.; Ward, F. E.: A high proportion

[141792]of ADA point mutations associated with a specific alanine-to-valine

[141793]substitution. Am. J. Hum. Genet. 45: 354–361, 1989.

[141794]25334.

[141795]Meuwissen, H. J.; Pollara, B.; Pickering, R. J.: Combined immunodeficiency

[141796]disease associated with adenosine deaminase deficiency (report on

[141797]a workshop held in Albany, New York, October 1, 1973). J. Pediat. 86:

[141798]169–181, 1975.

[141799]25335.

[141800]Migchielsen, A. A. J.; Breuer, M. L.; van Roon, M. A.; te Riele,

[141801]H.; Zurcher, C.; Ossendorp, F.; Toutain, S.; Hershfield, M. S.; Berns,

[141802]A.; Valerio, D.: Adenosine-deaminase-deficient mice die perinatally

[141803]and exhibit liver-cell degeneration, atelectasis and small intestinal

[141804]cell death. Nature Genet. 10: 279-287, 1995.

[141805]25336.

[141806]Mitchell, B. S.; Mejias, E.; Daddona, P. E.; Kelley, W. N.: Purinogenic

[141807]immunodeficiency disease: selective toxicity of deoxyribo-
nucleosides

[141808]for T-cells. Proc. Nat. Acad. Sci. 75: 5011-5014, 1978.

[141809]25337.

[141810]Moen, R. C.; Horowitz, S. D.; Sondel, P. M.; Borchering, W. R.;

[141811]Trigg, M. E.; Billing, R.; Hong, R.: Immunologic reconstitution after

[141812]haploidentical bone marrow transplantation for immune deficiency disorders:

[141813]treatment of bone marrow cells with monoclonal antibody CT-2 and complement. Blood 70:

[141814]664-669, 1987.

[141815]25338.

[141816]Mohandas, T.; Sparkes, R. S.; Suh, E. J.; Hershfield, M. S.:

[141817]Regional localization of the human genes for S-adenosylhomocysteine

[141818]hydrolase (cen-q131) and adenosine deaminase (q131-qter) on chromosome

[141819]20. Hum. Genet. 66: 292-295, 1984.

[141820]25339.

[141821]Nielsen, K. B.; Tommerup, N.; Jespersen, B.; Nygaard, P.; Kleif,

[141822]L.: Segregation of a t(3;20) translocation through three generations

[141823]resulting in unbalanced karyotypes in six persons. J. Med. Genet. 23:

[141824]446-451, 1986.

[141825]25340.

[141826]Orkin, S. H.; Daddona, P. E.; Shewach, D. S.; Markham, A. F.;

[141827]Bruns, G. A.; Goff, S. C.; Kelley, W. N.: Molecular cloning of human

[141828]adenosine deaminase gene sequences. J. Biol. Chem. 258:

12753–12756,

[141829]1983.

[141830]25341.

[141831]Ozsahin, H.; Arredondo–Vega, F. X.; Santisteban, I.;

Fuhrer, H.;

[141832]Tuchs Schmid, P.; Jochum, W.; Aguzzi, A.; Lederman, H. M.;

Fleischman,

[141833]A.; Winkelstein, J. A.; Seger, R. A.; Hershfield, M. S.:

Adenosine

[141834]deaminase deficiency in adults. Blood 89: 2849–2855,

1997.

[141835]25342.

[141836]Onodera, M.; Ariga, T.; Kawamura, N.; Kobayashi, I.;

Ohtsu, M.;

[141837]Yamada, M.; Tame, A.; Furuta, H.; Okano, M.; Matsumoto,

S.; Kotani,

[141838]H.; McGarrity, G. J.; Blaese, R. M.; Sakiyama, Y.: Successful

peripheral

[141839]T–lymphocyte–directed gene transfer for a patient with

severe combined

[141840]immune deficiency caused by adenosine deaminase defi-

ciency. Blood 91:

[141841]30–36, 1998.

[141842]25343.

[141843]Palmer, T. D.; Hock, R. A.; Osborne, W. R. A.; Miller, A. D.:

[141844]Efficient retrovirus-mediated transfer and expression of a
human adenosine

[141845]deaminase gene in diploid skin fibroblasts from an
adenosine deaminase-deficient

[141846]human. Proc. Nat. Acad. Sci. 84: 1055–1059, 1987.

[141847]25344.

[141848]Parkman, R.; Gelfand, E. W.; Rosen, F. S.; Sanderson, A.;
Hirschhorn,

[141849]R.: Severe combined immunodeficiency and adenosine
deaminase deficiency. New

[141850]Eng. J. Med. 292: 714–719, 1975.

[141851]25345.

[141852]Petersen, M. B.; Tranebjaerg, L.; Tommerup, N.; Nygaard,
P.; Edwards,

[141853]H.: New assignment of the adenosine deaminase gene lo-
cus to chromosome

[141854]20q13.11 by study of a patient with interstitial deletion
20q. J.

[141855]Med. Genet. 24: 93–96, 1987.

[141856]25346.

[141857]Polmar, S. H.; Stern, R. C.; Schwartz, A. L.; Wetzler, E. M.;

[141858]Chase, P. A.; Hirschhorn, R.: Enzyme replacement therapy
for adenosine

[141859]deaminase deficiency and severe combined immunodeficiency. New Eng.

[141860]J. Med. 295: 1337–1343, 1976.

[141861]25347.

[141862]Ratech, H.; Greco, M. A.; Gallo, G.; Rimoin, D. L.; Kamino,
H.;

[141863]Hirschhorn, R.: Pathologic findings in adenosine–deami-
nase–deficient

[141864]severe combined immunodeficiency. I. Kidney, adrenal,
and chondro–osseous

[141865]tissue alterations. Am. J. Path. 120: 157–169, 1985.

[141866]25348.

[141867]Ritter, H.; Wendt, G. G.; Tariverdian, G.; Zelch, J.; Rube, M.;

[141868]Kirchberg, G.: Genetics and linkage analysis of adenosine
deaminase. Humangenetik 14:

[141869]69–71, 1971.

[141870]25349.

[141871]Rogers, M. H.; Lwin, R.; Fairbanks, L.; Gerritsen, B.; Gaspar,

[141872]H. B.: Cognitive and behavioral abnormalities in adenosine deaminase

[141873]deficient severe combined immunodeficiency. *J. Pediat.* 139: 44–50,

[141874]2001.

[141875]25350.

[141876]Rothschild, C. B.; Akots, G.; Hayworth, R.; Pettenati, M. J.;

[141877]Rao, P. N.; Wood, P.; Stolz, F.–M.; Hansmann, I.; Serino, K.; Keith,

[141878]T. P.; Fajans, S. S.; Bowden, D. W.: A genetic map of chromosome

[141879]20q12–q13.1: multiple highly polymorphic microsatellite and RFLP markers

[141880]linked to the maturity-onset diabetes of the young (MODY) locus. *Am.*

[141881]*J. Hum. Genet.* 52: 110–123, 1993.

[141882]25351.

[141883]Rubinstein, A.; Hirschhorn, R.; Sicklick, M.; Murphy, R. A.:

[141884]In vivo and in vitro effects of thymosin and adenosine
deaminase on

[141885]adenosine–deaminase–deficient lymphocytes. New Eng. J.
Med. 300:

[141886]387–392, 1979.

[141887]25352.

[141888]Rudd, N. L.; Bain, H. W.; Giblett, E.; Chen, S.–H.; Worton, R.

[141889]G.: Partial trisomy 20 confirmed by gene dosage studies.
Am. J.

[141890]Med. Genet. 4: 357–364, 1979.

[141891]25353.

[141892]Koch, G.; Shows, T. B.: Somatic cell genetics of adenosine
deaminase

[141893]expression and severe combined immune deficiency dis-
ease in man. Proc.

[141894]Nat. Acad. Sci. 77: 4211–4215, 1980.

[141895]25354.

[141896]Gautam, M.; Noakes, P. G.; Moscoso, L.; Rupp, F.; Scheller,
R.

[141897]H.; Merlie, J. P.; Sanes, J. R.: Defective neuromuscular synaptogenesis

[141898]in agrin-deficient mutant mice. Cell 85: 525–535, 1996.

[141899]25355.

[141900]Glass, D. J.; Bowen, D. C.; Stitt, T. N.; Radziejewski, C.; Bruno,

[141901]J.; Ryan, T. E.; Gies, D. R.; Shah, S.; Mattsson, K.; Burden, S. J.;

[141902]DiStefano, P. S.; Valenzuela, D. M.; DeChiara, T. M.; Yancopoulos,

[141903]G. D.: Agrin acts via a MuSK receptor complex. Cell 85: 513–523,

[141904]1996.

[141905]25356.

[141906]Khan, A. A.; Bose, C.; Yam, L. S.; Soloski, M. J.; Rupp, F.: Physiological

[141907]regulation of the immunological synapse by agrin. Science 292: 1681–1686,

[141908]2001.

[141909]25357.

[141910]Lin, W.; Burgess, R. W.; Dominguez, B.; Pfaff, S. L.; Sanes,

J.

[141911]R.; Lee, K.-F.: Distinct roles of nerve and muscle in post-synaptic

[141912]differentiation of the neuromuscular synapse. Nature 410: 1057-1064,

[141913]2001.

[141914]25358.

[141915]McMahan, U. J.: The agrin hypothesis Cold Spring Harb. Symp.

[141916]Quant. Biol. 50: 407-418, 1990.

[141917]25359.

[141918]Rupp, F.; Ozcelik, T.; Linial, M.; Peterson, K.; Francke, U.; Scheller,

[141919]R.: Structure and chromosomal localization of the mammalian agrin

[141920]gene. J. Neurosci. 12: 3535-3544, 1992.

[141921]25360.

[141922]Rupp, F.; Payan, D. G.; Magill-Solc, C.; Cowan, D. M.; Scheller,

[141923]R. H.: Structure and expression of a rat agrin. Neuron 6: 811-823,

[141924]1991.

[141925]25361.

[141926]D'Alfonso, S.; Richiardi, P. M.: A polymorphic variation in a

[141927]putative regulation box of the TNFA promoter region. Immunogenetics 39:

[141928]150–154, 1994.

[141929]25362.

[141930]Mietus-Snyder, M.; Charmley, P.; Korf, B.; Ladas, J. A. A.:
Gatti,

[141931]R. A. and Karathanasis, S. K.: Genetic linkage of the human apolipoprotein

[141932]AI-CIII-AIV gene cluster and the neural cell adhesion molecule (NCAM)

[141933]gene. Genomics 7: 633–637, 1990.

[141934]25363.

[141935]Mietus-Snyder, M.; Korf, B.; Ladas, J. A.; Karathanasis, S.
K.

[141936]: Linkage of the human apolipoproteins A1, C3, A4 and the neural cell

[141937]adhesion molecule (NCAM) genes. (Abstract) Cytogenet. Cell Genet. 51:

[141938]1044 only, 1989.

[141939]25364.

[141940]Nguyen, C.; Mattei, M. G.; Goridis, C.; Mattei, J. F.; Jordan,

[141941]B. R.: Localization of the human N-CAM gene to chromosome 11 by in

[141942]situ hybridization with a murine N-CAM cDNA probe.
(Abstract) Cytogenet.

[141943]Cell Genet. 40: 713 only, 1985.

[141944]25365.

[141945]Nguyen, C.; Mattei, M. G.; Mattei, J.-F.; Santoni, M.-J.;
Goridis,

[141946]C.; Jordan, B. R.: Localization of the human NCAM gene to
band q23

[141947]of chromosome 11: the third gene coding for a cell interaction molecule

[141948]mapped to the distal portion of the long arm of chromosome 11. J.

[141949]Cell Biol. 102: 711-715, 1986.

[141950]25366.

[141951]Rabinowitz, J. E.; Rutishauser, U.; Magnuson, T.: Targeted
mutation

[141952]of Ncam to produce a secreted molecule results in a dominant embryonic

[141953]lethality. Proc. Nat. Acad. Sci. 93: 6421–6424, 1996.

[141954]25367.

[141955]Rutishauser, U.; Acheson, A.; Hall, A. K.; Mann, D. M.; Sunshine,

[141956]J.: The neural cell adhesion molecule (NCAM) as a regulator of cell-cell

[141957]interactions. Science 240: 53–57, 1988.

[141958]25368.

[141959]Rutishauser, U.; Goridis, C.: NCAM: the molecule and its genetics. Trends

[141960]Genet. 2: 72–76, 1986.

[141961]25369.

[141962]Telatar, M.; Lange, E.; Uhrhammer, N.; Gatti, R. A.: New localization

[141963]of NCAM, proximal to DRD2 at chromosome 11q23. Mammalian Genome 6:

[141964]59–60, 1995.

[141965]25370.

- [141966]Durbin, H.; Novelli, M.; Bodmer, W.: Detection of a 4-bp insertion
- [141967](CACA) functional polymorphism at nucleotide 241 of the cellular adhesion
- [141968]regulatory molecule CMAR (formerly CAR). Genomics 19: 181–182, 1994.
- [141969]25371.
- [141970]Koyama, K.; Emi, M.; Nakamura, Y.: The cell adhesion regulator
- [141971](CAR) gene, TaqI and insertion/deletion polymorphisms, and regional
- [141972]assignment to the peritelomeric region of 16q by linkage analysis. Genomics 16:
- [141973]264–265, 1993.
- [141974]25372.
- [141975]Pullman, W. E.; Bodmer, W. F.: Cloning and characterization of
- [141976]a gene that regulates cell adhesion. Nature 356: 529–532, 1992.
- [141977]25373.
- [141978]Agellon, L. B.; Drover, V. A. B.; Cheema, S. K.; Gbaguidi, G. F.;

[141979]Walsh, A.: Dietary cholesterol fails to stimulate the human cholesterol

[141980]7-alpha-hydroxylase gene (CYP7A1) in transgenic mice. J. Biol. Chem. 277:

[141981]20131-20134, 2002.

[141982]25374.

[141983]Angelin, B.; Einarsson, K.; Hellstrom, K.; Leijid, B.: Bile acid

[141984]kinetics in relation to endogenous triglyceride metabolism in various

[141985]types of hyperlipoproteinemia. J. Lipid Res. 19: 1004-1016, 1978.

[141986]25375.

[141987]Angelin, B.; Hershon, K. S.; Brunzell, J. D.: Bile acid metabolism

[141988]in hereditary forms of hypertriglyceridemia: evidence for an increased

[141989]synthesis rate in monogenic familial hypertriglyceridemia. Proc.

[141990]Nat. Acad. Sci. 84: 5434-5438, 1987.

[141991]25376.

[141992]Cohen, J. C.; Cali, J. J.; Jelinek, D. F.; Mehrabian, M.;

Sparkes,

[141993]R. S.; Lusi, A. J.; Russell, D. W.; Hobbs, H. H.: Cloning of
the

[141994]human cholesterol 7- α -hydroxylase gene (CYP7) and
localization

[141995]to chromosome 8q11-q12. Genomics 14: 153-161, 1992.

[141996]25377.

[141997]Drover, V. A. B.; Wong, N. C. W.; Agellon, L. B.: A distinct
thyroid

[141998]hormone response element mediates repression of the
human cholesterol

[141999]7- α -hydroxylase (CYP7A1) gene promoter. Molec. En-
docr. 16: 14-23,

[142000]2002.

[142001]25378.

[142002]Goodwin, B.; Jones, S. A.; Price, R. R.; Watson, M. A.; Mc-
Kee,

[142003]D. D.; Moore, L. B.; Galardi, C.; Wilson, J. G.; Lewis, M. C.;
Roth,

[142004]M. E.; Maloney, P. R.; Willson, T. M.; Kliewer, S. A.: A regu-
latory

[142005]cascade of the nuclear receptors FXR, SHP-1, and LXR-1

represses bile

[142006]acid biosynthesis. Molec. Cell 6: 517–526, 2000.

[142007]25379.

[142008]Molowa, D. T.; Chen, W. S.; Cimis, G. M.; Tan, C. P.: Transcriptional

[142009]regulation of the human cholesterol 7 alpha-hydroxylase gene. Biochemistry 31:

[142010]2539–2544, 1992.

[142011]25380.

[142012]Nitta, M.; Ku, S.; Brown, C.; Okamoto, A. Y.; Shan, B.: CPF: an

[142013]orphan nuclear receptor that regulates liver-specific expression of

[142014]the human cholesterol 7-alpha-hydroxylase gene. Proc. Nat. Acad.

[142015]Sci. 96: 6660–6665, 1999.

[142016]25381.

[142017]Noshiro, M.; Okuda, K.: Molecular cloning and sequence analysis

[142018]of cDNA encoding human cholesterol 7 alpha-hydroxylase. FEBS Lett. 268:

[142019]137–140, 1990.

[142020]25382.

[142021]Paumgartner, G.; Sauerbruch, T.: Gallstones: pathogenesis. *Lancet* 338:

[142022]1117–1121, 1991.

[142023]25383.

[142024]Wang, J.; Freeman, D. J.; Grundy, S. M.; Levine, D. M.;
Guerra,

[142025]R.; Cohen, J. C.: Linkage between cholesterol
7- α -hydroxylase

[142026]and high plasma low-density lipoprotein cholesterol concentrations. *J.*

[142027]*Clin. Invest.* 101: 1283–1291, 1998.

[142028]25384.

[142029]Gress, C. J.; Jacenko, O.: Growth plate compressions and
altered

[142030]hematopoiesis in collagen X null mice. *J. Cell Biol.* 149:
983–993,

[142031]2000.

[142032]25385.

[142033]Hasegawa, T.; Kozlowski, K.; Nishimura, G.; Hara, H.;
Hasegawa,

[142034]Y.; Aso, T.; Koto, S.; Nagai, T.; Tsuchiya, Y.: Japanese type
of

[142035]spondylo-metaphyseal dysplasia. *Pediat. Radiol.* 24:
194-197, 1994.

[142036]25386.

[142037]Ikegawa, S.; Nakamura, K.; Nagano, A.; Haga, N.; Naka-
mura, Y.:

[142038]Mutations in the N-terminal globular domain of the type X
collagen

[142039]gene (COL10A1) in patients with Schmid metaphyseal
chondrodysplasia. *Hum.*

[142040]*Mutat.* 9: 131-135, 1997.

[142041]25387.

[142042]Ikegawa, S.; Nishimura, G.; Nagai, T.; Hasegawa, T.;
Ohashi, H.;

[142043]Nakamura, Y.: Mutation of the type X collagen gene
(COL10A1) causes

[142044]spondylometaphyseal dysplasia. *Am. J. Hum. Genet.* 63:
1659-1662,

[142045]1998.

[142046]25388.

[142047]Kirsch, T.; von der Mark, K.: Isolation of human type X collagen

[142048]and immunolocalization in fetal human cartilage. *Europ. J. Biochem.* 196:

[142049]575–580, 1991.

[142050]25389.

[142051]Kuivaniemi, H.; Tromp, G.; Prockop, D. J.: Mutations in fibrillar

[142052]collagens (types I, II, III, and XI), fibril-associated collagen (type

[142053]IX), and network-forming collagen (type X) cause a spectrum of diseases

[142054]of bone, cartilage, and blood vessels. *Hum. Mutat.* 9: 300–315, 1997.

[142055]25390.

[142056]McIntosh, I.; Abbot, M. H.; Francomano, C. A.: Concentration

[142057]of mutations causing Schmid metaphyseal chondrodysplasia in the C-terminal

[142058]noncollagenous domain of type X collagen. *Hum. Mut.* 5: 121–125,

[142059]1995.

[142060]25391.

[142061]McIntosh, I.; Abbot, M. H.; Warman, M. L.; Olsen, B. R.;
Francomano,

[142062]C. A.: Additional mutations of type X collagen confirm
COL10A1 as

[142063]the Schmid metaphyseal chondrodysplasia locus. Hum.
Molec. Genet. 3:

[142064]303–307, 1994.

[142065]25392.

[142066]Nielsen, V. H.; Bendixen, C.; Arnbjerg, J.; Sorensen, C. M.;
Jensen,

[142067]H. E.; Shukri, N. M.; Thomsen, B.: Abnormal growth plate
function

[142068]in pigs carrying a dominant mutation in type X collagen.
Mammalian

[142069]Genome 11: 1087–1092, 2000.

[142070]25393.

[142071]Pokharel, R. K.; Alimsardjono, H.; Uno, K.; Fujii, S.; Shiba,

[142072]R.; Matsuo, M.: A novel mutation substituting tryptophan
with arginine

[142073]in the carboxyl-terminal, noncollagenous domain of collagen X in a
[142074]case of Schmid metaphyseal chondrodysplasia. Biochem. Biophys. Res. Commun. 217: 1157-1162, 1995.
[142075]25394.
[142076]Rosati, R.; Horan, G. S. B.; Pinero, G. J.; Garofalo, S.; Keene, D. R.; Horton, W. A.; Vuorio, E.; de Crombrughe, B.; Behringer, R. R.: Normal long bone growth and development in type X collagen-null mice. Nature Genet. 8: 129-135, 1994.
[142080]25395.
[142082]Sawai, H.; Ida, A.; Nakata, Y.; Koyama, K.: Novel missense mutation resulting in the substitution of tyrosine by cysteine at codon 597 of the type X collagen gene associated with Schmid metaphyseal chondrodysplasia. J. Hum. Genet. 43: 259-261, 1998.

[142086]25396.

[142087]Schmid, T. M.; Linsenmayer, T. F.: Immunohistochemical
localization

[142088]of short chain cartilage collagen (type X) in avian tissues.
J. Cell

[142089]Biol. 100: 598-605, 1985.

[142090]25397.

[142091]Stratakis, C. A.; Orban, Z.; Burns, A. L.; Vottero, A.; Mitsi-
ades,

[142092]C. S.; Marx, S. J.; Abbassi, V.; Chrousos, G. P.: Dideoxyfin-
gerprinting

[142093](ddF) analysis of the type X collagen gene (COL10A1) and
identification

[142094]of a novel mutation (S671P) in a kindred with Schmid
metaphyseal chondrodysplasia. Biochem.

[142095]Molec. Med. 59: 112-117, 1996.

[142096]25398.

[142097]Sweetman, W. A.; Rash, B.; Sykes, B.; Beighton, P.; Hecht, J.

[142098]T.; Zabel, B.; Thomas, J. T.; Boot-Handford, R.; Grant, M.
E.; Wallis,

[142099]G. A.: SSCP and segregation analysis of the human type X

collagen

[142100]gene (COL10A1) in heritable forms of chondrodysplasia.

Am. J. Hum.

[142101]Genet. 51: 841–849, 1992.

[142102]25399.

[142103]Thomas, J. T.; Cresswell, C. J.; Rash, B.; Hoyland, J.;

Freemont,

[142104]A. J.; Grant, M. E.; Boot-Handford, R. P.: The human colla-

gen X gene:

[142105]complete primary sequence and reexpression in os-

teoarthritis. Biochem.

[142106]Soc. Trans. 19: 804–808, 1991.

[142107]25400.

[142108]Thomas, J. T.; Cresswell, C. J.; Rash, B.; Nicolai, H.; Jones,

[142109]T.; Solomon, E.; Grant, M. E.; Boot-Handford, R. P.: The

human collagen

[142110]X gene: complete primary translated sequence and chro-

mosomal localization. Biochem.

[142111]J. 280: 617–623, 1991.

[142112]25401.

[142113]Wallis, G. A.; Rash, B.; Sweetman, W. A.; Thomas, J. T.; Su-

per,

[142114]M.; Evans, G.; Grant, M. E.; Boot-Handford, R. P.: Amino acid substitutions

[142115]of conserved residues in the carboxyl-terminal domain of the alpha-I(X)

[142116]chain of type X collagen occur in two unrelated families with metaphyseal

[142117]chondrodysplasia type Schmid. Am. J. Hum. Genet. 54: 169-178, 1994.

[142118]25402.

[142119]Haeffliger, J.-A.; Bruzzone, R.; Jenkins, N. A.; Gilbert, D. J.;

[142120]Copeland, N. G.; Paul, D. L.: Four novel members of the connexin

[142121]family of gap junction proteins: molecular cloning, expression, and

[142122]chromosome mapping. J. Biol. Chem. 267: 2057-2064, 1992.

[142123]25403.

[142124]Hsieh, C.-L.; Kumar, N. M.; Gilula, N. B.; Francke, U.: Distribution

[142125]of genes for gap junction membrane channel proteins on human and mouse

[142126]chromosomes. Somat. Cell Molec. Genet. 17: 191–200,
1991.

[142127]25404.

[142128]Hattersley, A. T.; Beards, F.; Ballantyne, E.; Appleton, M.;
Harvey,

[142129]R.; Ellard, S.: Mutations in the glucokinase gene of the fe-
tus result

[142130]in reduced birth weight. Nature Genet. 19: 268–270,
1998.

[142131]25405.

[142132]McCarthy, M.: Weighing in on diabetes risk. Nature Genet.
19:

[142133]209–210, 1998.

[142134]25406.

[142135]Neel, J. V.: Diabetes mellitus: a 'thrifty' genotype rendered

[142136]detrimental by 'progress'? Am. J. Hum. Genet. 14:
353–362, 1962.

[142137]25407.

[142138]Spielman, R. S.; McGinnis, R. E.; Ewens, W. J.: Transmission

[142139]test for linkage disequilibrium: the insulin gene region
and insulin-dependent

[142140]diabetes mellitus (IDDM). Am. J. Hum. Genet. 52:
506–516, 1993.

[142141]25408.

[142142]Altshuler, D.; Hirschhorn, J. N.; Klannemark, M.; Lindgren,
C.

[142143]M.; Vohl, M.–C.; Nemesh, J.; Lane, C. R.; Schaffner, S. F.;
Bolk,

[142144]S.; Brewer, C.; Tuomi, T.; Gaudet, D.; Hudson, T. J.; Daly,
M.; Groop,

[142145]L.; Lander, E. S.: The common PPAR–gamma pro12ala
polymorphism is

[142146]associated with decreased risk of type 2 diabetes. Nature
Genet. 76–80,

[142147]2000.

[142148]25409.

[142149]DeStefano, A. L.; Baldwin, C. T.; Burzstyn, M.; Gavras, I.;
Handy,

[142150]D. E.; Joost, O.; Martel, T.; Nicolaou, M.; Schwartz, F.;
Streeten,

[142151]D. H. P.; Farrer, L. A.; Gavras, H.: Autosomal dominant or–
thostatic

[142152]hypotensive disorder maps to chromosome 18q. Am. J.

Hum. Genet. 63:

[142153]1425–1430, 1998.

[142154]25410.

[142155]Zneimer, S. M.; Womack, J. E.: Regional localization of the
fibronectin

[142156]and gamma crystallin genes to mouse chromosome 1 by
in situ hybridization. Cytogenet.

[142157]Cell Genet. 48: 238–241, 1988.

[142158]25411.

[142159]den Dunnen, J. T.; Moormann, R. J. M.; Cremers, F. P. M.;
Schoenmakers,

[142160]J. G. G.: Two human gamma–crystallin genes are linked
and riddled

[142161]with Alu–repeats. Gene 38: 197–204, 1985.

[142162]25412.

[142163]Daniels, G.; King, M.–J.; Avent, N. D.; Khalid, G.; Reid, M.;

[142164]Mallinson, G.; Symthe, J.; Cedergren, B.: A point mutation
in the

[142165]GYPC gene results in the expression of the blood group
An(a) antigen

[142166]on glycophorin D but not on glycophorin C: further evi–

dence that glycophorin

[142167]D is a product of the GYPC gene. Blood 82: 3198–3203, 1993.

[142168]25413.

[142169]El-Maliki, B.; Blanchard, D.; Dahr, W.; Beyreuther, K.; Cartron,

[142170]J.-P.: Structural homology between glycophorins C and D of human

[142171]erythrocytes. Europ. J. Biochem. 183: 639–643, 1989.

[142172]25414.

[142173]Jorgensen, J.; Drachmann, O.; Gavin, J.: Duch, Dh(a), a low frequency

[142174]red cell antigen. Hum. Hered. 32: 73–75, 1982.

[142175]25415.

[142176]King, M. J.; Avent, N. D.; Mallinson, G.; Reid, M. E.: Point

[142177]mutation in the glycophorin C gene results in the expression of the

[142178]blood group antigen Dh(a). Vox Sang. 63: 56–58, 1992.

[142179]25416.

[142180]Le Van Kim, C.; Colin, Y.; Blanchard, D.; Dahr, W.; London,

J.;

[142181]Cartron, J.-P.: Gerbich blood group deficiency of the
Ge:-1,-2,-3

[142182]and Ge:-1,-2,3 types: immunochemical study and ge-
nomic analysis with

[142183]cDNA probes. *Europ. J. Biochem.* 165: 571-579, 1987.

[142184]25417.

[142185]Mattei, M. G.; Colin, Y.; Le Van Kim, C.; Mattei, J. F.;
Cartron,

[142186]J. P.: Localization of the gene for human erythrocyte gly-
cophorin

[142187]C to chromosome 2, q14-q21. *Hum. Genet.* 74: 420-422,
1986.

[142188]25418.

[142189]Pasvol, G.; Anstee, D. J.; Tanner, M. J. A.: Glycophorin C
and

[142190]the invasion of red cells by *Plasmodium falciparum*. *Lancet*
I: 907-908,

[142191]1984.

[142192]25419.

[142193]Race, R. R.; Sanger, R.: *Blood Groups in Man*. Oxford:

Blackwell

[142194]Sci. Publ. (pub.) (6th ed.): 1975. Pp. 416–421.

[142195]25420.

[142196]Reid, M. E.: The Gerbich blood group antigens: a review.
Med.

[142197]Lab. Sci. 43: 177–182, 1972.

[142198]25421.

[142199]Reid, M. E.; Sullivan, C.; Taylor, M.; Anstee, D. J.: Inheri-
tance

[142200]of human-erythrocyte Gerbich blood group antigens. Am.
J. Hum. Genet. 41:

[142201]1117–1123, 1987.

[142202]25422.

[142203]Simmons, R. T.; Albrey, J. A.: A 'new' blood group antigen
Webb

[142204](Wb) of low frequency found in two Australian families.
Med. J. Aust. I:

[142205]8–10, 1963.

[142206]25423.

[142207]Sondag, D.; Alloisio, N.; Blanchard, D.; Ducluzeau, M.–T.;

Colonna,

[142208]P.; Bachir, D.; Bloy, C.; Cartron, J.-P.; Delaunay, J.: Gerbich reactivity

[142209]in 4.1(-) hereditary elliptocytosis and protein 4.1 level in blood

[142210]group Gerbich deficiency. Brit. J. Haemat. 65: 43-50, 1987.

[142211]25424.

[142212]Spring, F. A.: Immunochemical characterisation of the low-incidence

[142213]antigen, Dh(a). Vox Sang. 61: 65-68, 1991.

[142214]25425.

[142215]Telen, M. J.; Le Van Kim, C.; Guizzo, M. L.; Cartron, J.-P.; Colin,

[142216]Y.: Erythrocyte Webb-type glycophorin C variant lacks N-glycosylation

[142217]due to an asparagine to serine substitution. Am. J. Hemat. 37: 51-52,

[142218]1991.

[142219]25426.

[142220]Winardi, R.; Reid, M.; Conboy, J.; Mohandas, N.: Molecular

analysis

[142221]of glycophorin C deficiency in human erythrocytes. Blood
81: 2799-2803,

[142222]1993.

[142223]25427.

[142224]Bierhuizen, M. F. A.; Mattei, M.-G.; Fukuda, M.: Expression
of

[142225]the developmental I antigen by a cloned human cDNA en-
coding a member

[142226]of a beta-1,6-N-acetylglucosaminyltransferase gene fam-
ily. Genes

[142227]Dev. 7: 468-478, 1993.

[142228]25428.

[142229]Lin-Chu, M.; Broadberry, R. E.; Okubo, Y.; Tanaka, M.: The
i phenotype

[142230]and congenital cataracts among Chinese in Taiwan (Letter)
Transfusion 31:

[142231]676-677, 1991.

[142232]25429.

[142233]Ogata, H.; Okubo, Y.; Akabane, T.: Phenotype i associated
with

[142234]congenital cataract in Japanese. Transfusion 19: 166–168,
1979.

[142235]25430.

[142236]Yeh, J.-C.; Ong, E.; Fukuda, M.: Molecular cloning and ex-
pression

[142237]of a novel beta-1,6-N-acetylglucosaminyltransferase that
forms core

[142238]2, core 4, and I branches. J. Biol. Chem. 274: 3215–3221,
1999.

[142239]25431.

[142240]Yu, L.-C.; Twu, Y.-C.; Chang, C.-Y.; Lin, M.: Molecular ba-
sis

[142241]of the adult i phenotype and the gene responsible for the
expression

[142242]of the human blood group I antigen. Blood 98:
3840–3845, 2001.

[142243]25432.

[142244]Allen, F. H., Jr.; Krabbe, S. M.; Corcoran, P. A.: A new phe-
notype

[142245](McLeod) in the Kell blood-group system. Vox Sang. 6:
555–560, 1961.

[142246]25433.

[142247]Avent, N. D.; Martin, P. G.: Kell typing by allele-specific
PCR

[142248](ASP). Brit. J. Haemat. 93: 728–730, 1996.

[142249]25434.

[142250]Chown, B.; Lewis, M.; Kaita, K.: A 'new' Kell blood-group
phenotype. Nature 180:

[142251]711 only, 1957.

[142252]25435.

[142253]Conneally, P. M.; Nance, W. E.; Huntzinger, R. S.: Linkage
analysis

[142254]of Kell–Sutter and PTC loci. (Abstract) Am. J. Hum. Genet.
26: 22A

[142255]only, 1974.

[142256]25436.

[142257]Abdelhak, S.; Kalatzis, V.; Heilig, R.; Compain, S.; Samson,
D.;

[142258]Vincent, C.; Weil, D.; Cruaud, C.; Sahly, I.; Leibovici, M.;
Bitner-Grindzicz,

[142259]M.; Francis, M.; Lacombe, D.; Vigneron, J.; Charachon, R.;
Boven,

[142260]K.; Bedbeder, P.; Van Regemorter, N.; Weissenbach, J.; Pe-

tit, C.:

[142261]A human homologue of the *Drosophila* eyes absent gene underlies branchio-oto-renal

[142262](BOR) syndrome and identifies a novel gene family. *Nature Genet.* 15:

[142263]157-164, 1997.

[142264]25437.

[142265]Tommerup, N.; Schempp, W.; Meinecke, P.; Pedersen, S.; Bolund,

[142266]L.; Brandt, C.; Goodpasture, C.; Guldberg, P.; Held, K.; Reinwein,

[142267]H.; Saugstad, O. D.; Scherer, G.; Skjeldal, O.; Toder, R.; Westvik,

[142268]J.; van der Hagen, C. B.; Wolf, U.: Assignment of an autosomal sex

[142269]reversal locus (SRA1) and campomelic dysplasia (CMPD1) to 17q24.3-q25.1. *Nature*

[142270]*Genet.* 4: 170-174, 1993.

[142271]25438.

[142272]Phillips, D. J.; Albertsson-Wikland, K.; Eriksson, K.; Wide, L.

[142273]: Changes in the isoforms of luteinizing hormone and fol-

lile-stimulating

[142274]hormone during puberty in normal children. J. Clin. Endocr. Metab. 82:

[142275]3103-3106, 1997.

[142276]25439.

[142277]Anderson, M. J.; Shelton, G. D.; Cavenee, W. K.; Arden, K. C.:

[142278]Embryonic expression of the tumor-associated PAX3-FKHR fusion protein

[142279]interferes with the developmental functions of Pax3. Proc. Nat. Acad.

[142280]Sci. 98: 1589-1594, 2001.

[142281]25440.

[142282]Anderson, M. J.; Viars, C. S.; Czekay, S.; Cavenee, W. K.; Arden,

[142283]K. C.: Cloning and characterization of three human fork-head genes

[142284]that comprise an FKHR-like gene subfamily. Genomics 47: 187-199,

[142285]1998.

[142286]25441.

[142287] Barr, F. G.; Galili, N.; Holick, J.; Biegel, J. A.; Rovera, G.; Emanuel, B. S.: Rearrangement of the PAX3 paired box gene in the paediatric solid tumor alveolar rhabdomyosarcoma. Nature Genet. 3: 113-117, 1993.

[142290] 25442.

[142291] Davis, R. J.; Barr, F. G.: Fusion genes resulting from alternative chromosomal translocations are overexpressed by gene-specific mechanisms in alveolar rhabdomyosarcoma. Proc. Nat. Acad. Sci. 94: 8047-8051, 1997.

[142295] 25443.

[142296] Fredericks, W. J.; Galili, N.; Mukhopadhyay, S.; Rovera, G.; Bennicelli, J.; Barr, F. G.; Rauscher, F. J., III: The PAX3-FKHR fusion protein created by the t(2;13) translocation in alveolar rhabdomyosarcomas

[142300] is a more potent transcriptional activator than PAX3.

Molec. Cell.

[142301] Biol. 15: 1522–1535, 1995.

[142302] 25444.

[142303] Galili, N.; Davis, R. J.; Fredericks, W. J.; Mukhopadhyay, S.;

[142304] Rauscher, F. J., III; Emanuel, B. S.; Rovera, G.; Barr, F. G.:

Fusion

[142305] of a fork head domain gene to PAX3 in the solid tumour
alveolar rhabdomyosarcoma. Nature

[142306] Genet. 5: 230–235, 1993.

[142307] 25445.

[142308] Khan, J.; Bittner, M. L.; Saal, L. H.; Teichmann, U.; Azorsa,
D.

[142309] O.; Gooden, G. C.; Pavan, W. J.; Trent, J. M.; Meltzer, P. S.:
cDNA

[142310] microarrays detect activation of a myogenic transcription
program

[142311] by the PAX3–FKHR fusion oncogene. Proc. Nat. Acad. Sci.
96: 13264–13269,

[142312] 1999.

[142313] 25446.

[142314] Lahat, H.; Pras, E.; Olender, T.; Avidan, N.; Ben-Asher, E.;
Man,

[142315] O.; Levy-Nissenbaum, E.; Khoury, A.; Lorber, A.; Goldman,
B.; Lancet,

[142316] D.; Eldar, M.: A missense mutation in a highly conserved
region of

[142317] CASQ2 is associated with autosomal recessive cate-
cholamine-induced

[142318] polymorphic ventricular tachycardia in Bedouin families
from Israel. Am.

[142319] J. Hum. Genet. 69: 1378-1384, 2001.

[142320] 25447.

[142321] Viskin, S.; Belhassen, B.: Polymorphic ventricular tach-
ycardias

[142322] in the absence of organic heart disease: classification, dif-
ferential

[142323] diagnosis, and implication for therapy. Prog. Cardiovas.
Dis. 41:

[142324] 17-34, 1998.

[142325] 25448.

[142326] Angle, C. R.: Congenital bowing and angulation of the
long bones. Pediatrics 13:

[142327]257–268, 1954.

[142328]25449.

[142329]Bain, A. D.; Barrett, H. S.: Congenital bowing of the long bones:

[142330]report of a case. Arch. Dis. Child. 34: 516–524, 1959.

[142331]25450.

[142332]Bell, D. M.; Leung, K. K. H.; Wheatley, S. C.; Ng, L. J.; Zhou,

[142333]S.; Ling, K. W.; Sham, M. H.; Koopman, P.; Tam, P. P. L.; Cheah, K.

[142334]S. E.: SOX9 directly regulates the type-II collagen gene. Nature

[142335]Genet. 16: 174–178, 1997.

[142336]25451.

[142337]Bi, W.; Huang, W.; Whitworth, D. J.; Deng, J. M.; Zhang, Z.; Behringer,

[142338]R. R.; de Crombrughe, B.: Haploinsufficiency of Sox9 results in

[142339]defective cartilage primordia and premature skeletal mineralization. Proc.

[142340]Nat. Acad. Sci. 98: 6698–6703, 2001.

[142341]25452.

[142342]Bishop, C. E.; Whitworth, D. J.; Qin, Y.; Agoulnik, A. I.;
Agoulnik,

[142343]J. U.; Harrison, W. R.; Behringer, R. R.; Overbeek, P. A.: A
transgenic

[142344]insertion upstream of Sox9 is associated with dominant
XX sex reversal

[142345]in the mouse. Nature Genet. 26: 490–494, 2000.

[142346]25453.

[142347]Caffey, J. P.: Prenatal bowing and thickening of tubular
bones,

[142348]with multiple cutaneous dimples in arms and legs: a con-
genital syndrome

[142349]of mechanical origin. Am. J. Dis. Child. 74: 543–562,
1947.

[142350]25454.

[142351]Cameron, F. J.; Hageman, R. M.; Cooke–Yarborough, C.;
Kwok, C.;

[142352]Goodwin, L. L.; Sillence, D. O.; Sinclair, A. H.: A novel germ
line

[142353]mutation in SOX9 causes familial campomelic dysplasia
and sex reversal. Hum.

[142354]Molec. Genet. 5: 1625–1630, 1996.

[142355]25455.

[142356]Cameron, F. J.; Sinclair, A. H.: Mutations in SRY and SOX9:
testis-determining

[142357]genes. Hum. Mutat. 9: 388–395, 1997.

[142358]25456.

[142359]Cooke, C. T.; Mulcahy, M. T.; Cullity, G. J.; Watson, M.;
Sprague,

[142360]P.: Campomelic dysplasia with sex reversal: morphological
and cytogenetic

[142361]studies of a case. Pathology 17: 526–529, 1985.

[142362]25457.

[142363]Cremin, B. J.; Orsmond, G.; Beighton, P.: Autosomal reces-
sive

[142364]inheritance in campomelic dwarfism.(Letter) Lancet I:
488–489,

[142365]1973.

[142366]25458.

[142367]Dagna Bricarelli, F.; Fraccaro, M.; Lindsten, J.; Muller, U.;

[142368]Baggio, P.; Carbone, L. D. L.; Hjerpe, A.; Lindgren, F.;

Mayerova,

[142369]A.; Ringertz, H.; Ritzen, E. M.; Rovetta, D. C.; Sicchero, C.;
Wolf,

[142370]U.: Sex-reversed XY females with campomelic dysplasia
are H-Y negative. Hum.

[142371]Genet. 57: 15-22, 1981.

[142372]25459.

[142373]Ebensperger, C.; Jager, R. J.; Lattermann, U.; Dagna
Bricarelli,

[142374]F.; Keutel, J.; Lindsten, J.; Rehder, H.; Muller, U.; Wolf, U.:
No

[142375]evidence of mutations in four candidate genes for male
sex determination/differentiation

[142376]in sex-reversed XY females with campomelic dysplasia.
Ann. Genet. 34:

[142377]233-238, 1991.

[142378]25460.

[142379]Fontaine, G.; Walbaum, R.; Farriaux, J. P.; Tilmont, P.;
Peuzin,

[142380]F.; Delecour, M.: Le conseil genetique dans la dysplasie
campomelique

[142381](a propos de deux observations). J. Genet. Hum. 28:

267–279, 1980.

[142382]25461.

[142383]Foster, J. W.; Dominguez–Steglich, M. A.; Guioli, S.; Kwok, C.;

[142384]Weller, P. A.; Stevanovic, M.; Weissenbach, J.; Mansour, S.; Young,

[142385]J. D.; Goodfellow, P. N.; Brook, J. D.; Schafer, A. J.: Campomelic

[142386]dysplasia and autosomal sex reversal caused by mutations in an SRY–related

[142387]gene. *Nature* 372: 525–530, 1994.

[142388]25462.

[142389]Friedrich, U.; Schaefer, E.; Meinecke, P.: Campomelic dysplasia

[142390]without overt campomelia. *Clin. Dysmorph.* 1: 172–178, 1992.

[142391]25463.

[142392]Gasca, S.; Canizares, J.; de Santa Barbara, P.; Mejean, C.; Poulat,

[142393]F.; Berta, P.; Boizet–Bonhoure, B.: A nuclear export signal within

[142394]the high mobility group domain regulates the nucleocyto–

plasmic translocation

[142395]of SOX9 during sexual determination. Proc. Nat. Acad. Sci.
99: 11199–11204,

[142396]2002.

[142397]25464.

[142398]Glass, R. B. J.; Rosenbaum, K. N.: Acampomelic cam-
pomelic dysplasia:

[142399]further radiographic variations. Am. J. Med. Genet. 69:
29–32, 1997.

[142400]25465.

[142401]Hall, B.; Spranger, J. W.: Campomelic dysplasia: further
elucidation

[142402]of a distinct entity. Am. J. Dis. Child. 134: 285–289, 1980.

[142403]25466.

[142404]Hoefnagel, D.; Wurster-Hill, D. H.; Dupree, W. B.;
Benirschke,

[142405]K.; Fuld, G. L.: Camptomelic dwarfism associated with XY-
gonadal

[142406]dysgenesis and chromosome anomalies. Clin. Genet. 13:
489–499, 1978.

[142407]25467.

[142408]Houston, C. S.; Opitz, J. M.; Spranger, J. W.; Macpherson, R.

[142409]J.; Reed, M. H.; Gilbert, E. F.; Herrmann, J.; Schinzel, A.:
The

[142410]campomelic syndrome: review, report of 17 cases, and
follow-up on

[142411]the currently 17-year-old boy first reported by Maroteaux
et al in

[142412]1971. Am. J. Med. Genet. 15: 3–28, 1983.

[142413]25468.

[142414]Hovmoller, M. L.; Osuna, A.; Eklof, O.; Fredga, K.; Hjerpe, A.;

[142415]Lindsten, J.; Ritzen, M.; Stanescu, V.; Svenningsen, N.:
Camptomelic

[142416]dwarfism. A genetically determined mesenchymal disorder
combined with

[142417]sex reversal. Hereditas 86: 51–62, 1977.

[142418]25469.

[142419]Huang, W.; Chung, U.; Kronenberg, H. M.; de Crombrughe, B.:

[142420]The chondrogenic transcription factor Sox9 is a target of
signaling

[142421]by the parathyroid hormone-related peptide in the growth plate of

[142422]endochondral bones. Proc. Nat. Acad. Sci. 98: 160-165, 2001.

[142423]25470.

[142424]Kanai, Y.; Koopman, P.: Structural and functional characterization

[142425]of the mouse Sox9 promoter: implications for campomelic dysplasia. Hum.

[142426]Molec. Genet. 8: 691-696, 1999.

[142427]25471.

[142428]Medema, R. H.; Kops, G. J. P. L.; Bos, J. L.; Burgering, B. M.

[142429]T.: AFX-like forkhead transcription factors mediate cell-cycle regulation

[142430]by Ras and PKB through p27(kip1). Nature 404: 782-787, 2000.

[142431]25472.

[142432]Eicher, E. M.; Womack, J. E.: Chromosomal location of soluble

[142433]glutamic-pyruvic transaminase-1 (Gpt-1) in the mouse. Biochem. Genet. 15:

[142434]1-8, 1977.

[142435]25473.

[142436]Falk, C. T.; Huss, J.: Linkage data on the chromosome 16 markers

[142437]HP and PGP: no additional support for or against the mapping of GPT.

[142438](Abstract) Cytogenet. Cell Genet. 40: 626 only, 1985.

[142439]25474.

[142440]Ferrell, R. E.: Personal Communication. Pittsburgh, Pa. 1/22/1988.

[142441]25475.

[142442]Jeremiah, S. J.; Kielty, C.; Povey, S.; McLellan, T.: Immunological

[142443]characterization of human GPT in hybrids supports assignment to chromosome 25476.

[142444](Abstract) Cytogenet. Cell Genet. 37: 498 only, 1984.

[142445]

[142446]8. Kalimanovska, V.; Majkic-Singh, N.; Stojanov, M.; Grozdanic, V.;

[142447]Vucetic, G.; Andelic, M.; Gligorovic, V.; Tomasevic, R.: Human red

[142448]cell glutamic-pyruvic transaminase polymorphism in Serbia, Yugoslavia. Hum.

[142449]Hered. 33: 319-321, 1983.

[142450]25477.

[142451]Kielty, C. M.; Povey, S.; Hopkinson, D. A.: Regulation of expression

[142452]of liver-specific enzymes: II. Activation and chromosomal localization

[142453]of soluble glutamate-pyruvate transaminase. Ann. Hum. Genet. 46:

[142454]135-143, 1982.

[142455]25478.

[142456]King, M.-C.; Go, R. C. P.; Elston, R. C.; Lynch, H. T.; Petrakis,

[142457]N. L.: Allele increasing susceptibility to human breast cancer may

[142458]be linked to the glutamate-pyruvate transaminase locus. Science 208:

[142459]406-408, 1980.

[142460]25479.

[142461]Kompf, J.: Population genetics of soluble glutamic-pyruvic

transaminase

[142462](EC:2.6.1.2): gene frequencies in southwestern Germany.

Humangenetik 14:

[142463]76–77, 1971.

[142464]25480.

[142465]Lahav, M.; Szeinberg, A.: A red-cell glutamic-pyruvic
transaminase

[142466]polymorphism in several population groups in Israel.

Hum. Hered. 22:

[142467]533–538, 1972.

[142468]25481.

[142469]Marazita, M. L.; Spence, M. A.; Sparkes, R. S.; Field, L. L.;

[142470]Crandall, B. F.; Sparkes, M. C.; Crist, M.: Linkage relations
of

[142471]GPT (glutamic-pyruvate transaminase). (Abstract) Cyto-
genet. Cell

[142472]Genet. 40: 690 only, 1985.

[142473]25482.

[142474]McLellan, T.: Two previously undetected variants of glu-
tamic-pyruvic

[142475]transaminase found by acidic polyacrylamide gel elec-

trophoresis. Am.

[142476]J. Hum. Genet. 34: 623-628, 1982.

[142477]25483.

[142478]McLellan, T.; Cannon, L. A.; Bishop, D. T.; Skolnick, M. H.:

[142479]The cumulative lod score between a breast cancer susceptibility locus

[142480]and GPT is -3.86. (Abstract) Cytogenet. Cell Genet. 37: 536-537,

[142481]1984.

[142482]25484.

[142483]Mithal, Y.; Lane, A. B.; Jenkins, T.: Absence of red cell glutamic-pyruvate

[142484]transaminase: discovery of a 'silent' allele homozygote. Am. J. Hum.

[142485]Genet. 32: 42-46, 1980.

[142486]25485.

[142487]O'Connell, P.; Nakamura, Y.; Lathrop, G. M.; Leppert, M.; Cartwright,

[142488]P.; Lalouel, J.-M.; White, R.: Three genetic linkage groups on chromosome

[142489]8. (Abstract) Cytogenet. Cell Genet. 46: 672 only, 1987.

[142490]25486.

[142491]Olaisen, B.: Genetics of the GPT system: family, mother-child

[142492]and association studies. Clin. Genet. 7: 245-254, 1975.

[142493]25487.

[142494]Pelzer, C. F.; Norum, R. A.: Identification of human red cell
[142495]glutamate-pyruvate transaminase (GPT) phenotypes by
isoelectric focusing. Am.

[142496]J. Hum. Genet. 37: 147-152, 1985.

[142497]25488.

[142498]Rocha, J.; Amorim, A.; Almeida, V. M.; Oliveira, J. P.; Leao,
[142499]M.; Tavares, M. C.; Pereira, M. S.; Vidal-Pinheiro, L.: Gene
dosage

[142500]evidence for the regional assignment of GPT
(glutamate-pyruvate transaminase;

[142501]E.C. 2.6.1.2) locus to 8q24.2-8qter. Hum. Genet. 80:
299-300, 1988.

[142502]25489.

[142503]Sanders, M. F.; King, M. C.; Lattanzio, D.; Crandall, J.; Le-
ung,

[142504]R.: Absence of linkage between HP and GPT. (Abstract)
Cytogenet.

[142505]Cell Genet. 37: 536–537, 1984.

[142506]25490.

[142507]Santachiara Benerecetti, A. S.; Beretta, M.; Pampiglione, S.:

[142508]Red cell glutamic–pyruvic transaminase polymorphism in
a sample of

[142509]the Italian population: a new variant allele: GPT(8). Hum.
Hered. 25:

[142510]276–278, 1975.

[142511]25491.

[142512]Sohocki, M. M.; Sullivan, L. S.; Harrison, W. R.; Sodergren,
E.

[142513]J.; Elder, F. F. B.; Weinstock, G.; Tanase, S.; Daiger, S. P.:
Human

[142514]glutamate pyruvate transaminase (GPT): localization to
8q24.3, cDNA

[142515]and genomic sequences, and polymorphic sites. Genomics
40: 247–252,

[142516]1997.

[142517]25492.

- [142518]Sparkes, M. C.; Crist, M.; Sparkes, R. S.: Glutamate pyruvate
transaminase null allele in seven new families. Hum.
Genet. 65:
147–148, 1983.
- [142521]25493.
- [142522]Wijnen, L. M. M.; Meera Khan, P.: Assignment of GPT to
human
chromosome 16. (Abstract) Cytogenet. Cell Genet. 32: 327
only, 1982.
- [142524]25494.
- [142525]Cremonini, N.; Graziano, E.; Chiarini, V.; Sforza, A.;
Zampa,
G. A.: Atypical McCune–Albright syndrome associated with
growth hormone–prolactin
pituitary adenoma: natural history, long–term follow–up,
and SMS 201–995--bromocriptine
combined treatment results. J. Clin. Endocr. Metab. 75:
1166–1169,
1992.
- [142530]25495.

[142531]DeChiara, T.; Robertson, E. J.; Efstratiadis, A.: Parental imprinting
[142532]of the mouse insulin-like growth factor II gene. Cell 64:
849–859,
[142533]1991.
[142534]25496.
[142535]Eddy, M. C.; Jan de Beur, S. M.; Yandow, S. M.; McAlister, W.
[142536]H.; Shore, E. M.; Kaplan, F. S.; Whyte, M. P.; Levine, M. A.:
Deficiency
[142537]of the alpha-subunit of the stimulatory G protein and severe extraskeletal
[142538]ossification. J. Bone Miner. Res. 15: 2074–2083, 2000.
[142539]25497.
[142540]Falconer, M. A.; Cope, C. L.; Robb-Smith, A. H. T.: Fibrous dysplasia
[142541]of bone with endocrine disorders and cutaneous pigmentation (Albright's
[142542]disease). Quart. J. Med. 11: 121–154, 1942.
[142543]25498.
[142544]Jackson, I. J.: A cDNA encoding tyrosinase-related protein

maps

[142545]to the brown locus in mouse. Proc. Nat. Acad. Sci. 85:
4392–4396,

[142546]1988. Note: Erratum: Proc. Nat. Acad. Sci. 86: 997 only,
1989.

[142547]25499.

[142548]Albertsen, H. M.; Smith, S. A.; Mazoyer, S.; Fujimoto, E.;
Stevens,

[142549]J.; Williams, B.; Rodriguez, P.; Cropp, C. S.; Slijepcevic, P.;
Carlson,

[142550]M.; Robertson, M.; Bradley, P.; Lawrence, E.; Harrington,
T.; Mei

[142551]Sheng, Z.; Hoopes, R.; Sternberg, N.; Brothman, A.; Calla-
han, R.;

[142552]Ponder, B. A. J.; White, R.: A physical map and candidate
genes in

[142553]the BRCA1 region on chromosome 17q12–21. Nature
Genet. 7: 472–479,

[142554]1994.

[142555]25500.

[142556]Ninomiya, Y.; Olsen, B. R.: Synthesis and characterization
of

- [142557]cDNA encoding a cartilage-specific short collagen. Proc. Nat. Acad.
- [142558]Sci. 81: 3014-3018, 1984.
- [142559]25501.
- [142560]Khatib, Z. A.; Inaba, T.; Valentine, M.; Look, A. T.: Chromosomal
- [142561]localization and cDNA cloning of the human DBP and TEF genes. Genomics 23:
- [142562]344-351, 1994.
- [142563]25502.
- [142564]Pilla, M.; Perachon, S.; Sautel, F.; Garrido, F.; Mann, A.; Wermuth,
- [142565]C. G.; Schwartz, J.-C.; Everitt, B. J.; Sokoloff, P.: Selective inhibition
- [142566]of cocaine-seeking behaviour by a partial dopamine D3 receptor agonist. Nature 400:
- [142567]371-375, 1999.
- [142568]25503.
- [142569]Kaplan, R.; Jaye, M.; Burgess, W. H.; Schlaepfer, D. D.; Haigler,
- [142570]H. T.: Cloning and expression of cDNA for human en-

donexin II, a Ca^{2+}

[142571]and phospholipid binding protein. J. Biol. Chem. 263:
8037–8043,

[142572]1988.

[142573]25504.

[142574]Arinami, T.; Ishikawa, M.; Inoue, A.; Yanagisawa, M.;
Masaki, T.;

[142575]Yoshida, M. C.; Hamaguchi, H.: Chromosomal assign-
ments of the human

[142576]endothelin family genes: the endothelin-1 gene (EDN1) to
6p23–p24,

[142577]the endothelin-2 gene (EDN2) to 1p34, and the endothe-
lin-3 gene (EDN3)

[142578]to 20q13.2–q13.3. Am. J. Hum. Genet. 48: 990–996,
1991.

[142579]25505.

[142580]Bourgeois, C.; Robert, B.; Rebourcet, R.; Mondon, F.;
Mignot, T.–M.;

[142581]Duc–Goiran, P.; Ferre, F.: Endothelin-1 and ET(A) receptor
expression

[142582]in vascular smooth muscle cells from human placenta: a
new ET(A) receptor

[142583] messenger ribonucleic acid is generated by alternative splicing of

[142584] exon 3. J. Clin. Endocr. Metab. 82: 3116–3123, 1997.

[142585] 25506.

[142586] Inoue, A.; Yanagisawa, M.; Kimura, S.; Kasuya, Y.;

Miyauchi, T.;

[142587] Goto, K.; Masaki, T.: The human endothelin family: three structurally

[142588] and pharmacologically distinct iso-peptides predicted by three separate

[142589] genes. Proc. Nat. Acad. Sci. 86: 2863–2867, 1989.

[142590] 25507.

[142591] Muller, G.; Ruppert, S.; Schmid, E.; Schutz, G.: Functional analysis

[142592] of alternatively spliced tyrosinase gene transcripts. EMBO J. 7:

[142593] 2723–2730, 1988.

[142594] 25508.

[142595] Murty, V. V. V. S.; Bouchard, B.; Mathew, S.; Vijayasaradhi, S.;

[142596] Houghton, A. N.: Assignment of the human TYRP (brown)

locus to chromosome

[142597]region 9p23 by nonradioactive in situ hybridization. Genomics 13:

[142598]227-229, 1992.

[142599]25509.

[142600]Ramsay, M.; Colman, M. A.; Jenkins, T.; Fox, M.; Chintamani,

[142601]C.; Pickard, R.; Kwon, B.: The human CAS2 locus (homologous to the

[142602]mouse b locus) maps to 9p22-pter. (Abstract) Cytogenet. Cell Genet. 58:

[142603]1943 only, 1991.

[142604]25510.

[142605]Shibahara, S.; Tomita, Y.; Yoshizawa, M.; Shibata, K.; Tagami,

[142606]H.: Identification of mutation in the pigment cell-specific gene

[142607]located at the brown locus in mouse. Pigment Cell Res. Suppl. 2:

[142608]90-95, 1992.

[142609]25511.

- [142610]Shaikh, S.; Ball, D.; Craddock, N.; Castle, D.; Hunt, N.;
Mant,
[142611]R.; Owen, M.; Collier, D.; Gill, M.: The dopamine D3 recep-
tor gene:
[142612]no association with bipolar affective disorder. J. Med.
Genet. 30:
[142613]308-309, 1993.
[142614]25512.
- [142615]Sokoloff, P.; Giros, B.; Martres, M.-P.; Bouthenet, M.-L.;
Schwartz,
[142616]J.-C.: Molecular cloning and characterization of a novel
dopamine
[142617]receptor (D-3) as a target for neuroleptics. Nature 347:
146-151,
[142618]1990.
[142619]25513.
- [142620]Spurlock, G.; Williams, J.; McGuffin, P.; Aschauer, H. N.;
Lenzinger,
[142621]E.; Fuchs, K.; Sieghart, W. C.; Meszaros, K.; Fathi, N.; Lau-
rent,
[142622]C.; Mallet, J.; Macciardi, F.; Pedrini, S.; Gill, M.; Hawi, Z.;
Gibson,

[142623]S.; Jazin, E. E.; Yang, H.-T.; Adolfsson, R.; Pato, C. N.;
Dourado,

[142624]A. M.; Owen, M. J.: European Multicentre Association Study
of Schizophrenia:

[142625]a study of the DRD2 ser311-to-cys and DRD3 ser9-to-gly
polymorphisms. Am.

[142626]J. Med. Genet. 81: 24-28, 1998.

[142627]25514.

[142628]Barsh, G. S.; Seeburg, P. H.; Gelinas, R. E.: The human
growth

[142629]hormone gene family:structure and evolution of the chro-
mosomal locus. Nucleic

[142630]Acids Res. 11: 3939-3958, 1983.

[142631]25515.

[142632]Halford, S.; Freedman, M. S.; Bellingham, J.; Inglis, S. L.;
Poopalasundaram,

[142633]S.; Soni, B. G.; Foster, R. G.; Hunt, D. M.: Characterization
of

[142634]a novel human opsin gene with wide tissue expression
and identification

[142635]of embedded and flanking genes on chromosome 1q43.
Genomics 72:

[142636]203–208, 2001.

[142637]25516.

[142638]Haneda, M.; Chan, S. J.; Kwok, S. C. M.; Rubenstein, A. H.;
Steiner,

[142639]D. F.: Studies on mutant human insulin genes: identifica-
tion and

[142640]sequence analysis of a gene encoding (Ser-B24) insulin.
Proc. Nat.

[142641]Acad. Sci. 80: 6366–6370, 1983.

[142642]25517.

[142643]Hart, A. W.; Baeza, N.; Apelqvist, A.; Edlund, H.: Attenua-
tion

[142644]of FGF signalling in mouse beta-cells leads to diabetes.
Nature 408:

[142645]864–868, 2000.

[142646]25518.

[142647]Mammarella, S.; Romano, F.; Di Valerio, A.; Creati, B.; Es-
posito,

[142648]D. L.; Palmirotta, R.; Capani, F.; Vitullo, P.; Volpe, G.; Bat-
tista,

[142649]P.; Della Loggia, F.; Mariani–Costantini, R.; Cama, A.: In–

teraction

[142650]between the G1057D variant of IRS-2 and overweight in
the pathogenesis

[142651]of type 2 diabetes. Hum. Molec. Genet. 9: 2517-2521,
2000.

[142652]25519.

[142653]Triggs-Raine, B. L.; Kirkpatrick, R. D.; Kelly, S. L.; Norquay,

[142654]L. D.; Cattini, P. A.; Yamagata, K.; Hanley, A. J. G.; Zinman,
B.;

[142655]Harris, S. B.; Barrett, P. H.; Hegele, R. A.: HNF1-alpha
G319S, a

[142656]transactivation-deficient mutant, is associated with al-
tered dynamics

[142657]of diabetes onset in an Oji-Cree community. Proc. Nat.
Acad. Sci. 99:

[142658]4614-4619, 2002.

[142659]25520.

[142660]Tuomi, T.; Carlsson, A.; Li, H.; Isomaa, B.; Miettinen, A.;
Nilsson,

[142661]A.; Nissen, M.; Ehrnstrom, B.-O.; Forsen, B.; Snickars, B.;
Lahti,

[142662]K.; Forsblom, C.; Saloranta, C.; Taskinen, M.-R.; Groop, L.

C.: Clinical

[142663]and genetic characteristics of type 2 diabetes with and without GAD

[142664]antibodies. Diabetes 48: 150–157, 1999.

[142665]25521.

[142666]Yuan, M.; Konstantopoulos, N.; Lee, J.; Hansen, L.; Li, Z.-W.;

[142667]Karin, M.; Shoelson, S. E.: Reversal of obesity- and diet-induced

[142668]insulin resistance with salicylates or targeted disruption of Ikk-beta. Science 293:

[142669]1673–1677, 2001.

[142670]25522.

[142671]Nuytinck, L.; Dalgleish, R.; Spotila, L.; Renard, J.-P.; Van Regemorter,

[142672]N.; De Paepe, A.: Substitution of glycine-661 by serine in the alpha-1(I)

[142673]and alpha-2(I) chains of type I collagen results in different clinical

[142674]and biochemical phenotypes. Hum. Genet. 97: 324–329, 1996.

[142675]25523.

[142676]Nuytinck, L.; Freund, M.; Lagae, L.; Pierard, G. E.; Hermanns-Le,
manns-Le,
[142677]T.; De Paepe, A.: Classical Ehlers-Danlos syndrome caused
by a mutation
[142678]in type I collagen. Am. J. Hum. Genet. 66: 1398-1402,
2000.
[142679]25524.
[142680]Oliver, J. E.; Thompson, E. M.; Pope, F. M.; Nicholls, A. C.:
[142681]Mutation in the carboxy-terminal propeptide of the pro-
alpha1(I) chain
[142682]of type I collagen in a child with severe osteogenesis im-
perfecta
[142683](OI type III): possible implications for protein folding.
Hum. Mutat. 7:
[142684]318-326, 1996.
[142685]25525.
[142686]Pace, J. M.; Atkinson, M.; Willing, M. C.; Wallis, G.; Byers,
[142687]P. H.: Deletions and duplications of Gly-Xaa-Yaa triplet
repeats
[142688]in the triple helical domains of type I collagen chains dis-
rupt helix

[142689]formation and result in several types of osteogenesis im-
perfecta. Hum.

[142690]Mutat. 18: 319-326, 2001.

[142691]25526.

[142692]Pack, M.; Constantinou, C. D.; Kalia, K.; Nielsen, K. B.;
Prockop,

[142693]D. J.: Substitution of serine for alpha-1(I)-glycine 844 in a
severe

[142694]variant of osteogenesis imperfecta minimally destabilizes
the triple

[142695]helix of type I procollagen: the effects of glycine substitu-
tions

[142696]on thermal stability are either position or amino acid spe-
cific. J.

[142697]Biol. Chem. 264: 19694-19699, 1989.

[142698]25527.

[142699]Pereira, R.; Halford, K.; Sokolov, B. P.; Khillan, J. S.;
Prockop,

[142700]D. J.: Phenotypic variability and incomplete penetrance of
spontaneous

[142701]fractures in an inbred strain of transgenic mice expressing
a mutated

[142702]collagen gene (COL1A1). J. Clin. Invest. 93: 1765–1769,
1994.

[142703]25528.

[142704]Pereira, R.; Khillan, J. S.; Helminen, H. J.; Hume, E. L.;
Prockop,

[142705]D. J.: Transgenic mice expressing a partially deleted gene
for type

[142706]I procollagen (COL1A1): a breeding line with a phenotype
of spontaneous

[142707]fractures and decreased bone collagen and mineral. J.
Clin. Invest. 91:

[142708]709–716, 1993.

[142709]25529.

[142710]Pereira, R. F.; O'Hara, M. D.; Laptev, A. V.; Halford, K. W.;

[142711]Pollard, M. D.; Class, R.; Simon, D.; Livezey, K.; Prockop,
D. J.

[142712]: Marrow stromal cells as a source of progenitor cells for
nonhematopoietic

[142713]tissues in transgenic mice with a phenotype of osteogene-
sis imperfecta. Proc.

[142714]Nat. Acad. Sci. 95: 1142–1147, 1998.

[142715]25530.

[142716]Pestka, S.; Daugherty, B. L.; Jung, V.; Hotta, K.; Pestka, R.

[142717]K.: Anti-mRNA: specific inhibition of translation of single
mRNA

[142718]molecules. Proc. Nat. Acad. Sci. 81: 7525-7528, 1984.

[142719]25531.

[142720]Raghunath, M.; Steinmann, B.; Delozier-Blanchet, C.; Ex-
termann,

[142721]P.; Superti-Furga, A.: Prenatal diagnosis of collagen disor-
ders by

[142722]direct biochemical analysis of chorionic villus biopsies.
Pediat.

[142723]Res. 36: 441-448, 1994.

[142724]25532.

[142725]Redford-Badwal, D. A.; Stover, M. L.; Valli, M.; McKinstry,
M.

[142726]B.; Rowe, D. W.: Nuclear retention of COL1A1 messenger
RNA identifies

[142727]null alleles causing mild osteogenesis imperfecta. J. Clin.
Invest. 97:

[142728]1035-1040, 1996.

[142729]25533.

[142730]Retief, E.; Parker, M. I.; Retief, A. E.: Regional chromosome

[142731]mapping of human collagen genes alpha 2(I) and alpha
1(I) (COLIA2

[142732]and COLIA1). Hum. Genet. 69: 304–308, 1985.

[142733]25534.

[142734]Sainz, J.; Van Tornout, J. M.; Sayre, J.; Kaufman, F.;
Gilsanz,

[142735]V.: Association of collagen type 1 alpha-1 gene polymor-
phism with

[142736]bone density in early childhood. J. Clin. Endocr. Metab. 84:
853–855,

[142737]1999.

[142738]25535.

[142739]Schnieke, A.; Harbers, K.; Jaenisch, R.: Embryonic lethal
mutation

[142740]in mice induced by retrovirus insertion into the alpha-1 (I)
collagen

[142741]gene. Nature 304: 315–320, 1983.

[142742]25536.

[142743]Schwarze, U.; Starman, B. J.; Byers, P. H.: Redefinition of
exon
[142744]7 in the COL1A1 gene of type I collagen by an intron 8
splice-donor-site
[142745]mutation in a form of osteogenesis imperfecta: influence
of intron
[142746]splice order on outcome of splice-site mutation. Am. J.
Hum. Genet. 65:
[142747]336-344, 1999.
[142748]25537.
[142749]Sippola-Thiele, M.; Tromp, G. C.; Prockop, D. J.; Ramirez,
F.
[142750]: Assessment of small polymorphisms in defined human
collagen gene
[142751]segments. Hum. Genet. 72: 245-247, 1986.
[142752]25538.
[142753]Solomon, E.; Hiorns, L.; Sheer, D.; Rowe, D.: Confirmation
that
[142754]the type I collagen gene on chromosome 17 is
COL1A1(alpha-1(I)), using
[142755]a human genomic probe. Ann. Hum. Genet. 48: 39-42,
1984.

[142756]25539.

[142757]Solomon, E.; Hiorns, L. R.; Cheah, K. S. E.; Parkar, M.;
Weiss,

[142758]E.; Flavell, R. A.: Assignment of a human alpha-1(I)-like
collagen

[142759]gene to chromosome 12, by molecular hybridiza-
tion.(Abstract) Cytogenet.

[142760]Cell Genet. 37: 588-589, 1984.

[142761]25540.

[142762]Solomon, E.; Sykes, B.: Assignment of alpha-1 (I), alpha-2,
and

[142763]possibly alpha-1 (III), chains of human collagen to chro-
mosome 7.(Abstract) Cytogenet.

[142764]Cell Genet. 25: 205, 1979.

[142765]25541.

[142766]Steinmann, B.: Personal Communication. Zurich, Switzer-
land

[142767]11/28/1994.

[142768]25542.

[142769]Barker, P. E.; Shipp, M. A.; D'Adamio, L.; Masteller, E. L.;
Reinherz,

[142770]E. L.: The common acute lymphoblastic leukemia antigen gene maps

[142771]to chromosomal region 3(q21-q27). J. Immun. 142: 283-287, 1989.

[142772]25543.

[142773]D'Adamio, L.; Shipp, M. A.; Masteller, E. L.; Reinherz, E. L.:

[142774]Organization of the gene encoding common acute lymphoblastic leukemia

[142775]antigen (neutral endopeptidase 24.11): multiple miniexons and separate

[142776]5'-prime untranslated regions. Proc. Nat. Acad. Sci. 86: 7103-7107,

[142777]1989.

[142778]25544.

[142779]Debiec, H.; Guigonis, V.; Mougenot, B.; Decobert, F.; Haymann,

[142780]J.-P.; Bensman, A.; Deschenes, G.; Ronco, P. M.: Antenatal membranous

[142781]glomerulonephritis due to anti-neutral endopeptidase antibodies. New

[142782]Eng. J. Med. 346: 2053-2060, 2002.

[142783]25545.

[142784]Letarte, M.; Vera, S.; Tran, R.; Addis, J. B. L.; Onizuka, R. J.;

[142785]Quackenbush, E. J.; Jongeneel, C. V.; McInnes, R. R.: Com-
mon acute

[142786]lymphocytic leukemia antigen is identical to neutral en-
dopeptidase. J.

[142787]Exp. Med. 168: 1247-1253, 1988.

[142788]25546.

[142789]Shipp, M. A.; Vijayaraghavan, J.; Schmidt, E. V.; Masteller,
E.

[142790]L.; D'Adamio, L.; Hersh, L. B.; Reinherz, E. L.: Common
acute lymphoblastic

[142791]leukemia antigen (CALLA) is active neutral endopeptidase
24.11 ('enkephalinase'):

[142792]direct evidence by cDNA transfection analysis. Proc. Nat.
Acad. Sci. 86:

[142793]297-301, 1989.

[142794]25547.

[142795]Tran-Paterson, R.; Willard, H. F.; Letarte, M.: The common
acute

[142796]lymphoblastic leukemia antigen (neutral endopeptidase-

-3.4.24.11)

[142797] gene is located on human chromosome 3. Cancer Genet.

Cytogenet. 42:

[142798] 129-134, 1989.

[142799] 25548.

[142800] Boyd, Y.; Cockburn, D.; Holt, S.; Munro, E.; van Ommen, G.

J.;

[142801] Gillard, B.; Affara, N.; Ferguson-Smith, M.; Craig, I.: Map-

ping of

[142802] 12 translocation breakpoints in the Xp21 region with re-

spect to the

[142803] locus for Duchenne muscular dystrophy. Cytogenet. Cell

Genet. 48:

[142804] 28-34, 1988.

[142805] 25549.

[142806] Sellar, G. C.; Blake, D. J.; Reid, K. B.: Characterization and

[142807] organization of the genes encoding the A-, B-, and C-

chains of human

[142808] complement subcomponent C1q: the complete derived

amino acid sequence

[142809] of human C1q. Biochem. J. 274: 481-490, 1991.

[142810]25550.

[142811]Topaloglu, R.; Bakkaloglu, A.; Slingsby, J. H.; Mihatsch, M. J.;

[142812]Pascual, M.; Norsworthy, P.; Morley, B. J.; Saatci, U.; Schif-ferli,

[142813]J. A.; Walport, M. J.: Molecular basis of hereditary C1q de-ficiency

[142814]associated with SLE and IgA nephropathy in a Turkish family. Kidney

[142815]Int. 50: 635-642, 1996.

[142816]25551.

[142817]Skok, J.; Solomon, E.; Reid, K. B. M.; Thompson, R. A.: Dis-tinct

[142818]genes for fibroblast and serum C1q. Nature 292: 549-551, 1981.

[142819]25552.

[142820]Hannema, A. J.; Kluin-Nelemans, J. C.; Hack, C. E.; Eeren-berg-Belmer,

[142821]A. J. M.; Mallee, C.; van Helden, H. P. T.: SLE like syndrome and

[142822]functional deficiency of C1q in members of a large family. Clin.

[142823]Exp. Immun. 55: 106–114, 1984.

[142824]25553.

[142825]McAdam, R. A.; Goundis, D.; Reid, K. B. M.: A homozygous point

[142826]mutation results in a stop codon in the C1q deficient individual. Immunogenetics 27:

[142827]259–264, 1988.

[142828]25554.

[142829]Sellar, G. C.; Cockburn, D.; Reid, K. B. M.: Localization of the

[142830]gene cluster encoding the A, B, and C chains of human C1q to 1p34.1–1p36.3. Immunogenetics 35:

[142831]214–216, 1992.

[142832]25555.

[142833]Solomon, E.; Skok, J.; Griffin, J.; Reid, K. B. M.: Human C1q

[142834]B chain (C1QB) is on chromosome 1p. (Abstract) Cytogenet. Cell Genet. 40:

[142835]749 only, 1985.

[142836]25556.

[142837]Thompson, R. A.; Haeney, M.; Reid, K. B. M.; Davis, J. G.;

White,

[142838]R. H.; Cameron, A. H.: A genetic defect of the C1q sub-component of

[142839]complement associated with childhood (immune complex) nephritis. New

[142840]Eng. J. Med. 303: 22-24, 1980.

[142841]25557.

[142842]Lahn, B. T.; Page, D. C.: Four evolutionary strata on the human

[142843]X chromosome. Science 286: 964-967, 1999.

[142844]25558.

[142845]Lee, F. A.; Issacs, H.; Strauss, J.: The 'campptomelic' syndrome.

[142846]Short life-span dwarfism with respiratory distress, hypotonia, peculiar

[142847]facies, and multiple skeletal and cartilaginous deformities. Am.

[142848]J. Dis. Child. 124: 485-496, 1972.

[142849]25559.

[142850]Lynch, S. A.; Gaunt, M. L.; Minford, A. M. B.: Campomelic dysplasia:

[142851]evidence of autosomal dominant inheritance. J. Med.

Genet. 30: 683–686,

[142852]1993.

[142853]25560.

[142854]Macpherson, R. I.; Skinner, S. A.; Donnenfeld, A. E.: Acam-
pomelic

[142855]campomelic dysplasia. Pediat. Radiol. 20: 90–93, 1989.

[142856]25561.

[142857]Mansour, S.; Hall, C. M.; Pembrey, M. E.; Young, I. D.: A
clinical

[142858]and genetic study of campomelic dysplasia. J. Med. Genet.
32: 415–420,

[142859]1995.

[142860]25562.

[142861]Maraia, R.; Saal, H. M.; Wangsa, D.: A chromosome 17q de
novo

[142862]paracentric inversion in a patient with campomelic dyspla-
sia; case

[142863]report and etiologic hypothesis. Clin. Genet. 39: 401–408,
1991.

[142864]25563.

- [142865]Maroteaux, P.; Spranger, J. W.; Opitz, J. M.; Kucera, J.;
Lowry,
[142866]R. B.; Schimke, R. N.; Kagan, S. M.: Le syndrome cam-
pomelique. Presse
[142867]Med. 22: 1157-1162, 1971.
[142868]25564.
[142869]Meyer, J.; Sudbeck, P.; Held, M.; Wagner, T.; Schmitz, M.
L.;
[142870]Bricarelli, F. D.; Eggermont, E.; Friedrich, U.; Haas, O. A.;
Kobelt,
[142871]A.; Leroy, J. G.; van Maldergem, L.; Michel, E.; Mitulla, B.;
Pfeiffer,
[142872]R. A.; Schinzel, A.; Schmidt, H.; Scherer, G.: Mutational
analysis
[142873]of the SOX9 gene in campomelic dysplasia and autosomal
sex reversal:
[142874]lack of genotype/phenotype correlations. Hum. Molec.
Genet. 6: 91-98,
[142875]1997.
[142876]25565.
[142877]Moedjono, S. J.; Crandall, B. F.; Sparkes, R. S.; Feldman, G.

[142878]M.; Austin, G. E.; Perry, S.: The campomelic syndrome in a
singleton

[142879]and monozygotic twins. Clin. Genet. 18: 397–401, 1980.

[142880]25566.

[142881]Moog, U.; Jansen, N. J. G.; Scherer, G.; Schrander–Stumpel,
C.

[142882]T. R. M.: Acampomelic campomelic syndrome. Am. J. Med.
Genet. 104:

[142883]239–245, 2001.

[142884]25567.

[142885]Morais da Silva, S.; Hacker, A.; Harley, V.; Goodfellow, P.;
Swain,

[142886]A.; Lovell–Badge, R.: Sox9 expression during gonadal de-
velopment

[142887]implies a conserved role for the gene in testis differentia-
tion in

[142888]mammals and birds. Nature Genet. 14: 62–68, 1996.

[142889]25568.

[142890]Murakami, S.; Kan, M.; McKeehan, W. L.; de Crombrughe,
B.: Up–regulation

[142891]of the chondrogenic Sox9 gene by fibroblast growth fac-

tors is mediated

[142892]by the mitogen-activated protein kinase pathway. Proc.
Nat. Acad.

[142893]Sci. 97: 1113-1118, 2000.

[142894]25569.

[142895]Ninomiya, S.; Isomura, M.; Narahara, K.; Seino, Y.; Naka-
mura,

[142896]Y.: Isolation of a testis-specific cDNA on chromosome 17q
from a

[142897]region adjacent to the breakpoint of t(12;17) observed in
a patient

[142898]with acampomelic campomelic dysplasia and sex reversal.
Hum. Molec.

[142899]Genet. 5: 69-72, 1996.

[142900]25570.

[142901]Ninomiya, S.; Yokoyama, Y.; Teraoka, M.; Mori, R.; Inoue,
C.;

[142902]Yamashita, S.; Tamai, H.; Funato, M.; Seino, Y.: A novel
mutation

[142903](296 del G) of the SOX9 gene in a patient with campomelic
syndrome

[142904]and sex reversal. Clin. Genet. 58: 224-227, 2000.

[142905]25571.

[142906]Olney, P. N.; Kean, L. S.; Graham, D.; Elsas, L. J.; May, K.
M.

[142907]: Campomelic syndrome and deletion of SOX9. Am. J. Med.
Genet. 84:

[142908]20–24, 1999.

[142909]25572.

[142910]Ozkilic, A.; Seven, M.; Yuksel, A.: A case of acampomelic
campomelic

[142911]dysplasia. Genet. Counsel. 13: 23–28, 2002.

[142912]25573.

[142913]Patel, M.; Dorman, K. S.; Zhang, Y.–H.; Huang, B.–L.;
Arnold,

[142914]A. P.; Sinsheimer, J. S.; Vilain, E.; McCabe, E. R. B.: Primate
DAX1,

[142915]SRY, and SOX9: evolutionary stratification of sex–
determination pathway. Am.

[142916]J. Hum. Genet. 68: 275–280, 2001.

[142917]25574.

[142918]Pfeifer, D.; Kist, R.; Dewar, K.; Devon, K.; Lander, E. S.; Bir–
ren,

- [142919]B.; Korniszewski, L.; Back, E.; Scherer, G.: Campomelic dysplasia translocation breakpoints are scattered over 1 Mb proximal to SOX9: evidence for an extended control region. Am. J. Hum. Genet. 65: 111–124, 1999.
- [142922]25575.
- [142923]Puck, S. M.; Haseltine, F. P.; Francke, U.: Absence of H-Y antigen in an XY female with campomelic dysplasia. Hum. Genet. 57: 23–27, 1981.
- [142926]25576.
- [142927]Rimoin, D. L.: Personal Communication. Torrance, Calif. 8/12/1976.
- [142928]25577.
- [142929]Rodriguez, J. I.: Vascular anomalies in campomelic syndrome. Am. J. Med. Genet. 46: 185–192, 1993.
- [142932]25578.

- [142933]Savarirayan, R.; Bankier, A.: Acampomelic campomelic dysplasia
[142934]with de novo 5q;17q reciprocal translocation and severe phenotype. J.
[142935]Med. Genet. 35: 597–599, 1998.
[142936]25579.
- [142937]Schimke, R. N.: XY sex–reversed campomelia--possibly an X–linked
[142938]disorder? (Letter) Clin. Genet. 16: 62–63, 1979.
[142939]25580.
- [142940]Shafai, T.; Schwartz, L.: Camptomelic syndrome in siblings. J.
[142941]Pediat. 89: 512–513, 1976.
[142942]25581.
- [142943]Spranger, J.: Advances in bone dysplasias. (Abstract) Sixth
[142944]Int. Cong. Hum. Genet. Jerusalem , 1981.
[142945]25582.
- [142946]Stuve, A.; Wiedemann, H.–R.: Congenital bowing of the long bones
[142947]in two sisters. (Letter) Lancet I: 495, 1971.

[142948]25583.

[142949]Sudbeck, P.; Schmitz, M. L.; Baeuerle, P. A.; Scherer, G.:
Sex

[142950]reversal by loss of the C-terminal transactivation domain
of human

[142951]SOX9. Nature Genet. 13: 230-232, 1996.

[142952]25584.

[142953]Thong, M.-K.; Scherer, G.; Kozlowski, K.; Haan, E.; Morris,
L.

[142954]: Acampomelic campomelic dysplasia with SOX9 mutation.
Am. J. Med.

[142955]Genet. 93: 421-425, 2000.

[142956]25585.

[142957]Thurmon, T. F.; De Fraites, E. B.; Anderson, E. E.: Familial

[142958]campomelic dwarfism. J. Pediat. 83: 841-843, 1973.

[142959]25586.

[142960]Vidal, V. P. I.; Chaboissier, M.-C.; de Rooij, D. G.; Schedl,

[142961]A.: Sox9 induces testis development in XX transgenic
mice. Nature

[142962]Genet. 28: 216-217, 2001.

[142963]25587.

[142964]Wagner, T.; Wirth, J.; Meyer, J.; Zabel, B.; Held, M.; Zimmer,

[142965]J.; Pasantes, J.; Dagna Bricarelli, F.; Keutel, J.; Hustert, E.; Wolf,

[142966]U.; Tommerup, N.; Schempp, W.; Scherer, G.: Autosomal sex reversal

[142967]and campomelic dysplasia are caused by mutations in and around the

[142968]SRY-related gene SOX9. Cell 79: 1111-1120, 1994.

[142969]25588.

[142970]Weller, S. D. V.: Hypophosphatasia with congenital dimples. Proc.

[142971]Roy. Soc. Med. 52: 637, 1959.

[142972]25589.

[142973]Wirth, J.; Wagner, T.; Meyer, J.; Pfeiffer, R. A.; Tietze, H.-U.;

[142974]Schempp, W.; Scherer, G.: Translocation breakpoints in three patients

[142975]with campomelic dysplasia and autosomal sex reversal map more than

[142976]130 kb from SOX9. Hum. Genet. 97: 186–193, 1996.

[142977]25590.

[142978]Wright, E.; Hargrave, M. R.; Christiansen, J.; Cooper, L.;
Kun,

[142979]J.; Evans, T.; Gangadharan, U.; Greenfield, A.; Koopman,
P.: The

[142980]Sry-related gene Sox9 is expressed during chondrogenesis in mouse

[142981]embryos. Nature Genet. 9: 15–20, 1995.

[142982]25591.

[142983]Albritton, L. M.; Bowcock, A. M.; Eddy, R. L.; Morton, C. C.;
Tseng,

[142984]L.; Farrer, L. A.; Cavalli-Sforza, L. L.; Shows, T. B.; Cunningham,

[142985]J. M.: The human cationic amino acid transporter (ATRC1):
physical

[142986]and genetic mapping to 13q12–q14. Genomics 12:
430–434, 1992.

[142987]25592.

[142988]Bowcock, A. M.; Gerken, S. C.; Barnes, R. I.; Shiang, R.;
Jabs,

- [142989]E. W.; Warren, A. C.; Antonarakis, S.; Retief, A. E.; Vergnaud, G.;
- [142990]Leppert, M.; Lalouel, J.-M.; White, R. L.; Cavalli-Sforza, L. L.;
- [142991]The CEPH consortium linkage map of human chromosome 13. *Genomics* 16: 486–496, 1993.
- [142992]25593.
- [142993]Kim, J. W.; Closs, E. I.; Albritton, L. M.; Cunningham, J. M.:
- [142995]Transport of cationic amino acids by the mouse ecotropic retrovirus receptor. *Nature* 352: 725–728, 1991.
- [142996]25594.
- [142998]Kozak, C. A.; Albritton, L. M.; Cunningham, J. M.: Genetic mapping of a cloned sequence responsible for susceptibility to leukemia viruses. *J. Virol.* 64: 3119–3121, 1990.
- [143001]25595.
- [143002]Oie, H. K.; Gazdar, A. F.; Lalley, P. A.; Russell, E. K.; Minna,

- [143003]J. D.; DeLarco, J.; Todaro, G. J.; Francke, U.: Mouse chromosome
- [143004]5 codes for ecotropic murine leukaemia virus cell-surface receptor. *Nature* 274:
- [143005]60-62, 1978.
- [143006]25596.
- [143007]Ruddle, N. H.; Conta, B. S.; Leinwand, L.; Kozak, C.; Ruddle, F.;
- [143008]Besmer, P.; Baltimore, D.: Assignment of the receptor for ecotropic
- [143009]murine leukemia virus to mouse chromosome 5. *J. Exp. Med.* 148: 451-465,
- [143010]1978.
- [143011]25597.
- [143012]Cook, P. W.; Piepkorn, M.; Clegg, C. H.; Plowman, G. D.; DeMay,
- [143013]J. M.; Brown, J. R.; Pittelkow, M. R.: Transgenic expression of the
- [143014]human amphiregulin gene induces a psoriasis-like phenotype. *J. Clin.*
- [143015]*Invest.* 100: 2286-2294, 1997.

[143016]25598.

[143017]Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly, J.;

[143018]Bowen-Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of the amphiregulin

[143019]and the platelet-growth factor receptor alpha genes to the proximal

[143020]long arm of chromosome 4. (Abstract) Cytogenet. Cell Genet. 51:

[143021]990 only, 1989.

[143022]25599.

[143023]Kimura, H.; Fischer, W. H.; Schubert, D.: Structure, expression

[143024]and function of a schwannoma-derived growth factor. Nature 348:

[143025]257-260, 1990.

[143026]25600.

[143027]Gillessen-Kaesbach, G.; Demuth, S.; Thiele, H.; Theile, U.; Lich,

[143028]C.; Horsthemke, B.: A previously unrecognised phenotype characterised

- [143029]by obesity, muscular hypotonia, and ability to speak in patients with
- [143030]Angelman syndrome caused by an imprinting defect. *Eur. J. Hum. Genet.* 7: 638–644, 1999.
- [143031]25601.
- [143032]25601.
- [143033]Acosta, J.; Hettinga, J.; Fluckiger, R.; Krumrei, N.; Goldfine, A.; Angarita, L.; Halperin, J.: Molecular basis for a link between
- [143034]complement and the vascular complications of diabetes. *Proc. Nat. Acad. Sci.* 97: 5450–5455, 2000.
- [143035]25602.
- [143036]25602.
- [143037]Bickmore, W. A.; Longbottom, D.; Oghene, K.; Fletcher, J. M.; van
- [143038]Heyningen, V.: Colocalization of the human CD59 gene to 11p13 with
- [143039]the MIC11 cell surface antigen. *Genomics* 17: 129–135, 1993.
- [143040]25603.
- [143041]25603.
- [143042]Davies, A.; Simmons, D. L.; Hale, G.; Harrison, R. A.; Tighe,

H.;

[143043]Lachmann, P. J.; Waldmann, H.: CD59, an LY-6-like protein expressed

[143044]in human lymphoid cells, regulates the action of the complement membrane

[143045]attack complex on homologous cells. J. Exp. Med. 170: 637-654, 1989.

[143046]25604.

[143047]Forsberg, U. H.; Bazil, V.; Stefanova, I.; Schroder, J.: Gene

[143048]for human CD59 (likely Ly-6 homologue) is located on the short arm

[143049]of chromosome 11. Immunogenetics 30: 188-193, 1989.

[143050]25605.

[143051]Harada, R.; Okada, N.; Fujita, T.; Okada, H.: Purification of

[143052]1F5 antigen that prevents complement attack on homologous cell membranes. J.

[143053]Immun. 144: 1823-1828, 1990.

[143054]25606.

[143055]Heckl-Ostreicher, B.; Ragg, S.; Drechsler, M.; Scherthan, H.; Royer-Pokora,

[143056]B.: Localization of the human CD59 gene by fluorescence

in situ hybridization

[143057]and pulsed-field gel electrophoresis. Cytogenet. Cell
Genet. 63:

[143058]144–146, 1993.

[143059]25607.

[143060]Holt, D. S.; Botto, M.; Bygrave, A. E.; Hanna, S. M.; Walport,

[143061]M. J.; Morgan, B. P.: Targeted deletion of the CD59 gene
causes spontaneous

[143062]intravascular hemolysis and hemoglobinuria. Blood 98:
442–449, 2001.

[143063]25608.

[143064]Holt, D. S.; Powell, M. B.; Rushmere, N. K.; Morgan, B. P.:
Genomic

[143065]structure and chromosome location of the gene encoding
mouse CD59. Cytogenet.

[143066]Cell Genet. 89: 264–267, 2000.

[143067]25609.

[143068]Huppi, K.; Duncan, R.; Potter, M.: Myc-1 is centromeric to
the

[143069]linkage group Ly-6-Sis-Gdc-1 on mouse chromosome 15.
Immunogenetics 27:

[143070]215–219, 1988.

[143071]25610.

[143072]Kamiura, S.; Nolan, C. M.; Meruelo, D.: Long-range physical map

[143073]of the Ly-6 complex: mapping the Ly-6 multigene family by field-inversion

[143074]and two-dimensional gel electrophoresis. Genomics 12: 89–105, 1992.

[143075]25611.

[143076]Low, M. G.; Saltiel, A. R.: Structural and functional roles of

[143077]glycosyl-phosphatidylinositol in membranes. Science 239: 268–275,

[143078]1988.

[143079]25612.

[143080]Mahoney, J. F.; Urakaze, M.; Hall, S.; DeGasperi, R.; Chang, H.-M.;

[143081]Sugiyama, E.; Warren, C. D.; Borowitz, M.; Nicholson-Weller, A.; Rosse,

[143082]W. F.; Yeh, E. T. H.: Defective glycosylphosphatidylinositol anchor

[143083]synthesis in paroxysmal nocturnal hemoglobinuria granu-

locytes. Blood 79:

[143084]1400-1403, 1992.

[143085]25613.

[143086]Mao, M.; Yu, M.; Tong, J.-H.; Ye, J.; Zhu, J.; Huang, Q.-H.;

Fu,

[143087]G.; Yu, L.; Zhao, S.-Y.; Waxman, S.; Lanotte, M.; Wang,

Z.-Y.; Tan,

[143088]J.-Z.; Chan, S.-J.; Chen, Z.: RIG-E, a human homolog of

the murine

[143089]Ly-6 family, is induced by retinoic acid during the differ-

entiation

[143090]of acute promyelocytic leukemia cell. Proc. Nat. Acad. Sci.

93:

[143091]5910-5914, 1996.

[143092]25614.

[143093]Meri, S.; Morgan, B. P.; Davies, A.; Daniels, R. H.;

Olavesen,

[143094]M. G.; Waldmann, H.; Lachmann, P. J.: Human protectin

(CD59), an

[143095]18,000-20,000 MW complement lysis restricting factor,

inhibits C5b-8

[143096]catalysed insertion of C9 into lipid bilayers. Immunology

71: 1-9,

[143097]1990.

[143098]25615.

[143099]Meri, S.; Morgan, B. P.; Wing, M.; Jones, J.; Davies, A.; Po-
dack,

[143100]E.; Lachmann, P. J.: Human protectin (CD59), an 18-20-kD
homologous

[143101]complement restriction factor, does not restrict perforin-
mediated

[143102]lysis. J. Exp. Med. 172: 367-370, 1990.

[143103]25616.

[143104]Motoyama, N.; Okada, N.; Yamashina, M.; Okada, H.:
Paroxysmal

[143105]nocturnal hemoglobinuria due to hereditary nucleotide
deletion in

[143106]the HRF20 (CD59) gene. Europ. J. Immun. 22: 2669-2673,
1992.

[143107]25617.

[143108]Okada, N.; Harada, R.; Fujiita, T.; Okada, H.: A novel
membrane

[143109]glycoprotein capable of inhibiting membrane attack by

homologous complement. Int.

[143110]Immun. 1: 205–208, 1989.

[143111]25618.

[143112]Stubbs, L.; Carver, E.; Ashworth, L.; Lopez–Molina, L.: Location

[143113]of the DBP transcription factor gene in human and mouse.
Mammalian

[143114]Genome 7: 65–67, 1996.

[143115]25619.

[143116]Szpirer, C.; Riviere, M.; Cortese, R.; Nakamura, T.; Islam, M.

[143117]Q.; Levan, G.; Szpirer, J.: Chromosomal localization in man and rat

[143118]of the genes encoding the liver–enriched transcription factors C/EBP,

[143119]DBP, and HNF1/LFB–1 (CEBP, DBP, and transcription factor 1, TCF1,

[143120]respectively) and of the hepatocyte growth factor/scatter factor gene

[143121](HGF). Genomics 13: 293–300, 1992.

[143122]25620.

- [143123]Gersten, K. M.; Natsuka, S.; Trinchera, M.; Petryniak, B.;
Kelly,
- [143124]R. J.; Hiraiwa, N.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N.
G.;
- [143125]Lowe, J. B.: Molecular cloning, expression, chromosomal
assignment,
- [143126]and tissue-specific expression of a murine alpha-
(1,3)-fucosyltransferase
- [143127]locus corresponding to the human ELAM-1 ligand fucosyl
transferase. J.
- [143128]Biol. Chem. 270: 25047-25056, 1995.
- [143129]25621.
- [143130]Angeletti, R. H.: Chromogranins and neuroendocrine se-
cretion.
- [143131](Editorial) Lab. Invest. 55: 387-390, 1986.
- [143132]25622.
- [143133]Bhargava, G.; Russell, J.; Sherwood, L. M.: Phosphorylation
of
- [143134]parathyroid secretory protein. Proc. Nat. Acad. Sci. 80:
878-881,
- [143135]1983.

[143136]25623.

[143137]Cetin, Y.; Aunis, D.; Bader, M.-F.; Galindo, E.; Jorns, A.;
Bargsten,

[143138]G.; Grube, D.: Chromostatin, a chromogranin A-derived
bioactive peptide,

[143139]is present in human pancreatic insulin (beta) cells. Proc.
Nat. Acad.

[143140]Sci. 90: 2360–2364, 1993.

[143141]25624.

[143142]Janssen, J. W. G.; Schulz, A. S.; Steenvoorden, A. C. M.;
Schmidberger,

[143143]M.; Strehl, S.; Ambros, P. F.; Bartram, C. R.: A novel puta-
tive tyrosine

[143144]kinase receptor with oncogenic potential. Oncogene 6:
2113–2120,

[143145]1991.

[143146]25625.

[143147]Liu, E.; Hjelle, B.; Bishop, J. M.: Transforming genes in
chronic

[143148]myelogenous leukemia. Proc. Nat. Acad. Sci. 85:
1952–1956, 1988.

[143149]25626.

[143150]Lu, Q.; Lemke, G.: Homeostatic regulation of the immune system

[143151]by receptor tyrosine kinases of the Tyro 3 family. Science 293:

[143152]306–311, 2001.

[143153]25627.

[143154]Manfioletti, G.; Brancolini, C.; Avanzi, G.; Schneider, C.: The

[143155]protein encoded by a growth arrest–specific gene (gas6) is a new member

[143156]of the vitamin K–dependent proteins related to protein S, a negative

[143157]coregulator in the blood coagulation cascade. Molec. Cell. Biol. 13:

[143158]4976–4985, 1993.

[143159]25628.

[143160]O'Bryan, J. P.; Frye, R. A.; Cogswell, P. C.; Neubauer, A.; Kitch,

[143161]B.; Prokop, C.; Espinosa, R., III; Le Beau, M. M.; Earp, H. S.; Liu,

- [143162]E. T.: Axl, a transforming gene isolated from primary human myeloid leukemia cells, encodes a novel receptor tyrosine kinase. Molec. Cell. Biol. 11: 5016–5031, 1991.
- [143163]25629.
- [143164]Varnum, B. C.; Young, C.; Elliott, G.; Garcia, A.; Bartley, T. D.; Fridell, Y.-W.; Hunt, R. W.; Trail, G.; Clogston, C.; Toso, R.
- [143165]J.; Yanagihara, D.; Bennett, L.; Sylber, M.; Merewether, L. A.; Tseng, A.; Escobar, E.; Liu, E. T.; Yamane, H. K.: Axl receptor tyrosine kinase stimulated by the vitamin K-dependent protein encoded by growth-arrest-specific gene 6. Nature 373: 623–626, 1995.
- [143166]25630.
- [143167]Galbraith, G. M. P.; Pandey, J. P.: Tumor necrosis factor alpha (TNF-alpha) gene polymorphism in alopecia areata. Hum. Genet. 96:

[143175]433–436, 1995.

[143176]25631.

[143177]Tazi–Ahnini, R.; di Giovine, F. S.; McDonagh, A. J. G.; Messenger,

[143178]A. G.; Amadou, C.; Cox, A.; Duff, G. W.; Cork, M. J.: Structure and

[143179]polymorphism of the human gene for the interferon–induced p78 protein

[143180](MX1): evidence of association with alopecia areata in the Down syndrome

[143181]region. Hum. Genet. 106: 639–645, 2000.

[143182]25632.

[143183]Wilson, A. G.; di Giovine, F. S.; Blakemore, A. I. F.; Duff, G.

[143184]W.: Single base polymorphism in the human tumour necrosis factor

[143185]alpha (TNF–alpha) gene detectable by NcoI restriction of PCR product. Hum.

[143186]Molec. Genet. 1: 353 only, 1992.

[143187]25633.

[143188]Belanger, L.; Roy, S.; Allard, D.: New albumin gene 3–prime adjacent

[143189]to the alpha-1-fetoprotein locus. J. Biol. Chem. 269:

5481-5484,

[143190]1994.

[143191]25634.

[143192]Lichenstein, H. S.; Lyons, D. E.; Wurfel, M. M.; Johnson, D.

A.;

[143193]McGinley, M. D.; Leidli, J. C.; Trollinger, D. B.; Mayer, J. P.;

Wright,

[143194]S. D.; Zukowski, M. M.: Afamin is a new member of the al-

bumin, alpha-fetoprotein,

[143195]and vitamin D-binding protein gene family. J. Biol. Chem.

269: 18149-18154,

[143196]1994.

[143197]25635.

[143198]Nishio, H.; Dugaiczky, A.: Complete structure of the hu-

man alpha-albumin

[143199]gene, a new member of the serum albumin multigene

family. Proc. Nat.

[143200]Acad. Sci. 93: 7557-7561, 1996.

[143201]25636.

[143202]Song, Y.-H.; Naumova, A. K.; Liebhaber, S. A.; Cooke, N.

E.: Physical

[143203]and meiotic mapping of the region of human chromosome
4q11-q13 encompassing

[143204]the vitamin D binding protein DBP/Gc-globulin and albu-
min multigene

[143205]cluster. Genome Res. 9: 581-587, 1999.

[143206]25637.

[143207]Ettinger, S.; Weksler, M. E.; Zhou, X.; Blass, J.; Szabo, P.:

[143208]Chromosomal fragility associated with familial Alzheimer's
disease. Ann.

[143209]Neurol. 36: 190-199, 1994.

[143210]25638.

[143211]Fox, N. C.; Kennedy, A. M.; Harvey, R. J.; Lantos, P. L.;
Roques,

[143212]P. K.; Collinge, J.; Hardy, J.; Hutton, M.; Stevens, J. M.;
Warrington,

[143213]E. K.; Rossor, M. N.: Clinicopathological features of famil-
ial Alzheimer's

[143214]disease associated with the M139V mutation in the prese-
nilin 1 gene:

[143215]pedigree but not mutation specific age at onset provides

evidence

[143216]for a further genetic factor. Brain 120: 491–501, 1997.

[143217]25639.

[143218]Seri, M.; Celli, I.; Betsos, N.; Claudiani, F.; Camera, G.;

Romeo,

[143219]G.: A cys634gly substitution of the RET proto-oncogene in
a family

[143220]with recurrence of multiple endocrine neoplasia type 2A
and cutaneous

[143221]lichen amyloidosis. Clin. Genet. 51: 86–90, 1997.

[143222]25640.

[143223]Georgakopoulos, A.; Marambaud, P.; Efthimiopoulos, S.;

Shioi,

[143224]J.; Cui, W.; Li, H.-C.; Schutte, M.; Gordon, R.; Holstein, G.

R.;

[143225]Martinelli, G.; Mehta, P.; Friedrich, V. L., Jr.; Robakis, N. K.:

[143226]Presenilin-1 forms complexes with the cadherin/catenin
cell-cell adhesion

[143227]system and is recruited to intercellular and synaptic con-
tacts. Molec.

[143228]Cell 4: 893–902, 1999.

[143229]25641.

[143230]Grilli, M.; Diodato, E.; Lozza, G.; Brusa, R.; Casarini, M.;
Uberti,

[143231]D.; Rozmahel, R.; Westaway, D.; St George-Hyslop, P.;
Memo, M.; Ongini,

[143232]E.: Presenilin-1 regulates the neuronal threshold to exci-
totoxicity

[143233]both physiologically and pathologically. Proc. Nat. Acad.
Sci. 97:

[143234]12822-12827, 2000.

[143235]25642.

[143236]Gustafson, L.; Brun, A.; Englund, E.; Hagnell, O.; Nilsson,
K.;

[143237]Stensmyr, M.; Ohlin, A.-K.; Abrahamson, M.: A 50-year
perspective

[143238]of a family with chromosome-14-linked Alzheimer's dis-
ease. Hum. Genet. 102:

[143239]253-257, 1998.

[143240]25643.

[143241]Handler, M.; Yang, X.; Shen, J.: Presenilin-1 regulates neu-
ronal

[143242]differentiation during neurogenesis. Development 127:
2593–2606,

[143243]2000.

[143244]25644.

[143245]Hartmann, D.: From Alzheimer's disease to skin tumors:
the catenin

[143246]connection. Proc. Nat. Acad. Sci. 98: 10522–10523, 2001.

[143247]25645.

[143248]Harvey, R. J.; Ellison, D.; Hardy, J.; Hutton, M.; Roques, P.

[143249]K.; Collinge, J.; Fox, N. C.; Rossor, M. N.: Chromosome 14
familial

[143250]Alzheimer's disease: the clinical and neuropathological
characteristics

[143251]of a family with a leucine-to-serine (L250S) substitution at
codon

[143252]250 of the presenilin 1 gene. J. Neurol. Neurosurg. Psy-
chiat. 64:

[143253]44–49, 1998.

[143254]25646.

[143255]Hsiao, K.; Chapman, P.; Nilsen, S.; Eckman, C.; Harigaya,
Y.;

[143256]Younkin, S.; Yang, F.; Cole, G.: Correlative memory deficits, A- β elevation, and amyloid plaques in transgenic mice. Science 274: 99–103, 1996.

[143259]25647.

[143260]Hull, M.; Fiebich, B. L.; Dykieriek, P.; Schmidtke, K.; Nitzsche, E.; Orszagh, M.; Deuschl, G.; Moser, E.; Schumacher, M.; Lucking, C.; Berger, M.; Bauer, J.: Early-onset Alzheimer's disease due to mutations of the presenilin-1 gene on chromosome 14: a 7-year follow-up of a patient with a mutation at codon 139. Europ. Arch. Psychiat. Clin. Neurosci. 248: 123–129, 1998.

[143266]25648.

[143267]Jarrett, J. T.; Berger, E. P.; Lansbury, P. T.: The carboxy terminus of the β amyloid protein is critical for the seeding of amyloid

[143269]formation: implications for the pathogenesis of
Alzheimer's disease. Biochemistry 32:

[143270]4693–4697, 1993.

[143271]25649.

[143272]Johnson, K. A.; Lopera, F.; Jones, K.; Becker, A.; Sperling,
R.;

[143273]Hilson, J.; Londono, J.; Siegert, I.; Arcos, M.; Moreno, S.;
Madrigal,

[143274]L.; Ossa, J.; Pineda, N.; Ardila, A.; Roselli, M.; Albert, M. S.;

[143275]Kosik, K. S.; Rios, A.: Presenilin-1-associated abnormali-
ties in

[143276]regional cerebral perfusion. Neurology 56: 1545–1551,
2001.

[143277]25650.

[143278]Kamal, A.; Almenar-Queral, A.; LeBlanc, J. F.; Roberts, E.
A.;

[143279]Goldstein, L. S. B.: Kinesin-mediated axonal transport of a
membrane

[143280]compartment containing beta-secretase and presenilin-1
requires APP. Nature 414:

[143281]643–648, 2001.

[143282]25651.

[143283]Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.; Goldstein, L. S.

[143284]: Axonal transport of amyloid precursor protein is mediated by direct

[143285]binding to the kinesin light chain subunit of kinesin-I.
Neuron 28:

[143286]449–459, 2000.

[143287]25652.

[143288]Kang, D. E.; Soriano, S.; Xia, X.; Eberhart, C. G.; De
Strooper,

[143289]B.; Zheng, H.; Koo, E. H.: Presenilin couples the paired
phosphorylation

[143290]of beta-catenin independent of Axin: implications for
beta-catenin

[143291]activation in tumorigenesis. Cell 110: 751–762, 2002.

[143292]25653.

[143293]Kopan, R.; Goate, A.: A common enzyme connects Notch
signaling

[143294]and Alzheimer's disease. Genes Dev. 14: 2799–2806,
2000.

[143295]25654.

[143296]Kovacs, D. M.; Fausett, H. J.; Page, K. J.; Kim, T.-W.; Moir,
[143297]R. D.; Merriam, D. E.; Hollister, R. D.; Hallmark, O. G.;
Mancini,
[143298]R.; Felsenstein, K. M.; Hyman, B. T.; Tanzi, R. E.; Wasco,
W.: Alzheimer-associated
[143299]presenilins 1 and 2: neuronal expression in brain and lo-
calization
[143300]to intracellular membranes in mammalian cells. Nature
Med. 2: 224-229,
[143301]1996.
[143302]25655.
[143303]Kwok, J. B. J.; Taddei, K.; Hallupp, M.; Fisher, C.; Brooks,
W.
[143304]S.; Broe, G. A.; Hardy, J.; Fulham, M. J.; Nicholson, G. A.;
Stell,
[143305]R.; St. George Hyslop, P. H.; Fraser, P. E.; and 6 others:
Two
[143306]novel (M233T and R278T) presenilin-1 mutations in early-
onset Alzheimer's
[143307]disease pedigrees and preliminary evidence for associa-
tion of presenilin-1
[143308]mutations with a novel phenotype. Neuroreport 8:

1537–1542, 1997.

[143309]25656.

[143310]Lambert, J.-C.; Mann, D. M. A.; Harris, J. M.; Chartier-Harlin,

[143311]M.-C.; Cumming, A.; Coates, J.; Lemmon, H.; StClair, D.; Iwatsubo,

[143312]T.; Lendon, C.: The -48 C/T polymorphism in the presenilin 1 promoter

[143313]is associated with an increased risk of developing Alzheimer's disease

[143314]and an increased A-beta load in brain. J. Med. Genet. 38: 353–355,

[143315]2001.

[143316]25657.

[143317]Leissring, M. A.; Akbari, Y.; Fanger, C. M.; Cahalin, M. D.; Mattson,

[143318]M. P.; LaFerla, F. M.: Capacitative calcium entry deficits and elevated

[143319]luminal calcium content in mutant presenilin-1 knockin mice. J. Cell

[143320]Biol. 149: 793–797, 2000.

[143321]25658.

[143322]Lemere, C. A.; Lopera, F.; Kosik, K. S.; Lendon, C. L.; Ossa,
 [143323]J.; Saido, T. C.; Yamaguchi, H.; Ruiz, A.; Martinez, A.;
 Madrigal,
 [143324]L.; Hincapie, L.; Arango, J. C.; Anthony, D. C.; Koo, E. H.;
 Goate,
 [143325]A. M.; Selkoe, D. J.; Arango, J. C.: The E280A presenilin 1
 Alzheimer
 [143326]mutation produces increased A-beta-42 deposition and
 severe cerebellar
 [143327]pathology. Nature Med. 2: 1146-1150, 1996.
 [143328]25659.
 [143329]Lewis, P. A.; Perez-Tur, J.; Golde, T. E.; Hardy, J.: The pre-
 senilin
 [143330]1 C92S mutation increases A-beta-42 production.
 Biochem. Biophys.
 [143331]Res. Commun. 277: 261-263, 2000.
 [143332]25660.
 [143333]Li, J.; Xu, M.; Zhou, H.; Ma, J.; Potter, H.: Alzheimer prese-
 nilins
 [143334]in the nuclear membrane, interphase kinetochores, and
 centrosomes

[143335] suggest a role in chromosome segregation. Cell 90:
917–927, 1997.

[143336] 25661.

[143337] Li, Y.–M.; Lai, M.–T.; Xu, M.; Huang, Q.; DiMuzio–Mower,
J.; Sardana,

[143338] M. K.; Shi, X.–P.; Yin, K.–C.; Shafer, J. A.; Gardell, S. J.: Pre–
senilin

[143339] 1 is linked with gamma–secretase activity in the detergent
solubilized

[143340] state. Proc. Nat. Acad. Sci. 97: 6138–6143, 2000.

[143341] 25662.

[143342] Matsubara–Tsutsui, M.; Yasuda, M.; Yamagata, H.; No–
mura, T.; Taguchi,

[143343] K.; Kohara, K.; Miyoshi, K.; Miki, T.: Molecular evidence of
presenilin

[143344] 1 mutation in familial early onset dementia. Am. J. Med.
Genet. 114:

[143345] 292–298, 2002.

[143346] 25663.

[143347] Hofstra, R. M. W.; Sijmons, R. H.; Stelwagen, T.; Stulp, R.
P.;

[143348]Kousseff, B. G.; Lips, C. J. M.; Steijlen, P. M.; Van Voorst
Vader,

[143349]P. C.; Buys, C. H. C. M.: RET mutation screening in familial
cutaneous

[143350]lichen amyloidosis and in skin amyloidosis associated with
multiple

[143351]endocrine neoplasia. J. Invest. Derm. 107: 215–218, 1996.
[143352]25664.

[143353]Bordet, T.; Lesbordes, J.-C.; Rouhani, S.; Castelnau-
Ptakhine,

[143354]L.; Schmalbruch, H.; Haase, G.; Kahn, A.: Protective effects
of cardiotrophin-1

[143355]adenoviral gene transfer on neuromuscular degeneration
in transgenic

[143356]ALS mice. Hum. Molec. Genet. 10: 1925–1933, 2001.
[143357]25665.

[143358]Narahara, K.; Kikkawa, K.; Kimira, S.; Kimoto, H.; Ogata,
M.;

[143359]Kasai, R.; Hamawaki, M.; Matsuoka, K.: Regional mapping
of catalase

[143360]and Wilms tumor--aniridia, genitourinary abnormalities,
and mental

- [143361]retardation triad loci to the chromosome segment
11p1305–p1306. Hum.
- [143362]Genet. 66: 181–185, 1984.
- [143363]25666.
- [143364]Turleau, C.; de Grouchy, J.; Nihoul–Fekete, C.; Dufier, J. L.;
- [143365]Chavin–Colin, F.; Junien, C.: Del11p13/nephroblastoma
without aniridia. Hum.
- [143366]Genet. 67: 455–456, 1984.
- [143367]25667.
- [143368]Newport, M. J.; Huxley, C. M.; Huston, S.; Hawrylowicz, C.
M.;
- [143369]Oostra, B. A.; Williamson, R.; Levin, M.: A mutation in the
interferon–gamma–receptor
- [143370]gene and susceptibility to mycobacterial infection. New
Eng. J. Med. 335:
- [143371]1941–1949, 1996.
- [143372]25668.
- [143373]Cowan, C. A.; Yokoyama, N.; Bianchi, L. M.; Henkemeyer,
M.; Frittsch,
- [143374]B.: EphB2 guides axons at the midline and is necessary for
normal

[143375]vestibular function. Neuron 26: 417–430, 2000.

[143376]25669.

[143377]Novick, D.; Orchansky, P.; Revel, M.; Rubinstein, M.: The human

[143378]interferon–gamma receptor: purification, characterization, and preparation

[143379]of antibodies. J. Biol. Chem. 262: 8483–8487, 1987.

[143380]25670.

[143381]Orchansky, P.; Rubinstein, M.; Fischer, D. G.: The interferon–gamma

[143382]receptor in human monocytes is different from the one in nonhematopoietic

[143383]cells. J. Immun. 136: 169–173, 1986.

[143384]25671.

[143385]Papanicolaou, G. J.; Parsa, N. Z.; Meltzer, P. S.; Trent, J. M.

[143386]: Assignment of interferon gamma receptor (IFNGR1) to human chromosome

[143387]bands 6q24.1–q24.2 by in situ hybridization. Cytogenet. Cell Genet. 76:

[143388]181–182, 1997. Note: Erratum: Cytogenet. Cell Genet. 78: 132 only,

[143389]1997.

[143390]25672.

[143391]Pierre-Audigier, C.; Jouanguy, E.; Lamhamedi, S.; Altare, F.;

[143392]Rauzier, J.; Vincent, V.; Canioni, D.; Emile, J. F.; Fischer, A.;

[143393]Blanche, S.; Gaillard, J. L.; Casanova, J. L.: Fatal disseminated

[143394]Mycobacterium smegmatis infection in a child with inherited interferon

[143395]gamma receptor deficiency. Clin. Infect. Dis. 24: 982-984, 1997.

[143396]25673.

[143397]Rashidbaigi, A.; Langer, J. A.; Jung, V.; Jones, C.; Morse, H.

[143398]G.; Tischfield, J. A.; Trill, J. J.; Kung, H.-F.; Pestka, S.: The

[143399]gene for the human immune interferon receptor is located on chromosome

[143400]6. Proc. Nat. Acad. Sci. 83: 384-388, 1986.

[143401]25674.

[143402]Rettig, W. J.; Grzeschik, K.-H.; Yenamandra, A. K.; Garcia, E.;

[143403]Old, L. J.: Definition of selectable cell surface markers for human

[143404]chromosomes and chromosome segments in rodent-human hybrids. Somat.

[143405]Cell Molec. Genet. 14: 223-231, 1988.

[143406]25675.

[143407]Shankaran, V.; Ikeda, H.; Bruce, A. T.; White, J. M.; Swanson,

[143408]P. E.; Old, L. J.; Schreiber, R. D.: IFN-gamma and lymphocytes prevent

[143409]primary tumour development and shape tumour immunogenicity. Nature 410:

[143410]1107-1111, 2001.

[143411]25676.

[143412]Bauer, R.; Imhof, A.; Pscherer, A.; Kopp, H.; Moser, M.; Seegers,

[143413]S.; Kerscher, M.; Tainsky, M. A.; Hofstaedter, F.; Buettner, R.:

[143414]The genomic structure of the human AP-2 transcription factor. Nucleic

[143415]Acids Res. 22: 1413-1420, 1994.

[143416]25677.

[143417]Buettner, R.; Kannan, P.; Imhof, A.; Bauer, R.; Yim, S. O.;
Glockshuber,

[143418]R.; Van Dyke, M. W.; Tainsky, M. A.: An alternatively
spliced mRNA

[143419]from the AP-2 gene encodes a negative regulator of tran-
scriptional

[143420]activation by AP-2. Molec. Cell. Biol. 13: 4174-4185,
1993.

[143421]25678.

[143422]Ahmad, I.: Mash-1 is expressed during ROD photorecep-
tor differentiation

[143423]and binds an E-box, E(opsin-1), in the rat opsin gene.
Brain Res.

[143424]Dev. Brain Res. 90: 184-189, 1995.

[143425]25679.

[143426]Ball, D. W.; Azzoli, C. G.; Baylin, S. B.; Chi, D.; Dou, S.; Do-
nis-Keller,

[143427]H.; Cumaraswamy, A.; Borges, M.; Nelkin, B. D.: Identifica-
tion of

[143428]a human achaete-scute homolog highly expressed in neu-
roendocrine tumors. Proc.

[143429]Nat. Acad. Sci. 90: 5648–5652, 1993.

[143430]25680.

[143431]Guillemot, F.; Lo, L.-C.; Johnson, J. E.; Auerbach, A.; Anderson,

[143432]D. J.; Joyner, A. L.: Mammalian achaete–scute homolog 1 is required

[143433]for the early development of olfactory and autonomic neurons. Cell 75:

[143434]463–476, 1993.

[143435]25681.

[143436]Renault, B.; Lieman, J.; Ward, D.; Krauter, K.; Kucherlapati, R.

[143437]: Localization of the human achaete–scute homolog gene (ASCL1) distal

[143438]to phenylalanine hydroxylase (PAH) and proximal to tumor rejection

[143439]antigen (TRA1) on chromosome 12q22–q23. Genomics 30: 81–83, 1995.

[143440]25682.

[143441]Deen, P. M. T.; Weghuis, D. O.; Geurts van Kessel, A.; Wieringa,

[143442]B.; van Os, C. H.: The human gene for water channel
aquaporin 1 (AQP1)

[143443]is localized on chromosome 7p15–p14. Cytogenet. Cell
Genet. 65:

[143444]243–246, 1994.

[143445]25683.

[143446]de Groot, B. L.; Grubmuller, H.: Water permeation across
biological

[143447]membranes: mechanism and dynamics of aquaporin–1
and Glpf. Science 294:

[143448]2353–2357, 2001.

[143449]25684.

[143450]de la Chapelle, A.; Vuopio, P.; Sanger, R.; Teesdale, P.:
Monosomy–7

[143451]and the Colton blood–groups. (Letter) Lancet II: 817 only,
1975.

[143452]25685.

[143453]Denker, B. M.; Smith, B. L.; Kuhajda, F. P.; Agre, P.: Identi-
fication,

[143454]purification, and partial characterization of a novel M(r)
28,000

[143455]integral membrane protein from erythrocytes and renal tubules. J.

[143456]Biol. Chem. 263: 15634–15642, 1988.

[143457]25686.

[143458]Fushimi, K.; Uchida, S.; Hara, Y.; Hirata, Y.; Marumo, F.; Sasaki,

[143459]S.: Cloning and expression of apical membrane water channel of rat

[143460]kidney collecting tubule. Nature 361: 549–552, 1993.

[143461]25687.

[143462]Hoffman, J. F.: Aquaporin: a wee burn runs through it. (Editorial) J.

[143463]Clin. Invest. 92: 1604–1605, 1993.

[143464]25688.

[143465]Keen, T. J.; Inglehearn, C. F.; Patel, R. J.; Green, E. D.; Peluso,

[143466]D. C.; Bhattacharya, S. S.: Localization of the aquaporin 1 (AQP1)

[143467]gene within a YAC contig containing the polymorphic markers D7S632

[143468]and D7S526. Genomics 25: 599–600, 1995.

[143469]25689.

[143470]King, L. S.; Choi, M.; Fernandez, P. C.; Cartron, J.-P.; Agre,

[143471]P.: Defective urinary concentrating ability due to a complete deficiency

[143472]of aquaporin-1. New Eng. J. Med. 345: 175-179, 2001.

[143473]25690.

[143474]King, L. S.; Nielsen, S.; Agre, P.; Brown, R. H.: Decreased pulmonary

[143475]vascular permeability in aquaporin-1-null humans. Proc. Nat. Acad.

[143476]Sci. 99: 1059-1063, 2002.

[143477]25691.

[143478]Knepper, M. A.: The aquaporin family of molecular water channels. Proc.

[143479]Nat. Acad. Sci. 91: 6255-6258, 1994.

[143480]25692.

[143481]Ma, T.; Yang, B.; Gillespie, A.; Carlson, E. J.; Epstein, C. J.;

[143482]Verkman, A. S.: Severely impaired urinary concentrating ability in

[143483]transgenic mice lacking aquaporin-1 water channels. J.

Biol. Chem. 273:

[143484]4296–4299, 1998.

[143485]25693.

[143486]Maurel, C.; Reizer, J.; Schroeder, J. I.; Chrispeels, M.: The

[143487]vacuolar membrane protein gamma-TIP creates water
specific channels

[143488]in *Xenopus* oocytes. EMBO J. 12: 2241–2247, 1993.

[143489]25694.

[143490]Moon, C.; Preston, G. M.; Griffin, C. A.; Jabs, E. W.; Agre, P.

[143491]: The human aquaporin-CHIP gene: structure, organiza-
tion, and chromosomal

[143492]localization. J. Biol. Chem. 268: 15772–15778, 1993.

[143493]25695.

[143494]Moon, C.; Williams, J. B.; Preston, G. M.; Copeland, N. G.;
Gilbert,

[143495]D. J.; Nathans, D.; Jenkins, N. A.; Agre, P.: The mouse
Aquaporin-1

[143496]gene. Genomics 30: 354–357, 1995.

[143497]25696.

[143498]Murata, K.; Mitsuoka, K.; Hirai, T.; Walz, T.; Agre, P.; Heymann,

[143499]J. B.; Engel, A.; Fujiyoshi, Y.: Structural determinants of water

[143500]permeation through aquaporin-1. Nature 407: 599-605, 2000.

[143501]25697.

[143502]Parsons, S. F.; Jones, J.; Anstee, D. J.; Judson, P. A.; Gardner,

[143503]B.; Wiener, E.; Poole, J.; Illum, N.; Wickramasinghe, S. N.: A novel

[143504]form of congenital dyserythropoietic anemia associated with deficiency

[143505]of erythroid CD44 and a unique blood group phenotype [In(a-b-), Co(a-b-)]. Blood 83:

[143506]860-868, 1994.

[143507]25698.

[143508]Pasquali, F.; Bernasconi, P.; Casalone, R.; Fraccaro, M.; Bernasconi,

[143509]C.; Lazzarino, M.; Morra, E.; Alessandrino, E. P.; Marchi, M. A.;

[143510]Sanger, R.: Pathogenic significance of 'pure' monosomy 7

in myeloproliferative

[143511]disorders: analysis of 14 cases.. Hum. Genet. 62: 40–51,
1982.

[143512]25699.

[143513]Preston, G. M.; Agre, P.: Isolation of the cDNA for erythro-
cyte

[143514]integral membrane protein of 28 kilodaltons: member of
an ancient

[143515]channel family. Proc. Nat. Acad. Sci. 88: 11110–11114,
1991.

[143516]25700.

[143517]Preston, G. M.; Smith, B. L.; Zeidel, M. L.; Moulds, J. J.;
Agre,

[143518]P.: Mutations in aquaporin–1 in phenotypically normal hu-
mans without

[143519]functional CHIP water channels. Science 265: 1585–1587,
1994.

[143520]25701.

[143521]Smith, B. L.; Baumgarten, R.; Nielsen, S.; Raben, D.; Zeidel,

[143522]M. L.; Agre, P.: Concurrent expression of erythroid and re-
nal aquaporin

[143523]CHIP and appearance of water channel activity in perinatal

rats. J.

[143524]Clin. Invest. 92: 2035–2041, 1993.

[143525]25702.

[143526]Smith, B. L.; Preston, G. M.; Spring, F.; Anstee, D. J.; Agre,

[143527]P.: Human red blood cell aquaporin CHIP: I. Molecular
characterization

[143528]of ABH and Colton blood group antigens. J. Clin. Invest.
94: 1043–1049,

[143529]1994.

[143530]25703.

[143531]Sui, H.; Han, B.-G.; Lee, J. K.; Walian, P.; Jap, B. K.: Struc-
tural

[143532]basis of water-specific transport through the AQP1 water
channel. Nature 414:

[143533]872–878, 2001.

[143534]25704.

[143535]Tang, W.; Cai, S.-P.; Eng, B.; Poon, M.-C.; Waye, J. S.; Il-
lum,

[143536]N.; Chui, D. H. K.: Expression of embryonic zeta-globin
and epsilon-globin

[143537]chains in a 10-year-old girl with congenital anemia. Blood

- 81: 1636–1640,
[143538]1993.
- [143539]25705.
- [143540]Wickramasinghe, S. N.; Illum, N.; Wimberley, P. D.: Con-
genital
[143541]dyserythropoietic anaemia with novel intra-erythroblastic
and intra-erythrocytic
[143542]inclusions. *Brit. J. Haemat.* 79: 322–330, 1991.
- [143543]25706.
- [143544]Yang, B.; Ma, T.; Verkman, A. S.: Erythrocyte water perme-
ability
[143545]and renal function in double knockout mice lacking aqua-
porin-1 and
[143546]aquaporin-3. *J. Biol. Chem.* 276: 624–628, 2001.
- [143547]25707.
- [143548]de The, H.; Chomienne, C.; Lanotte, M.; Degos, L.; Dejean,
A.:
[143549]The t(15;17) translocation of acute promyelocytic
leukaemia fuses
[143550]the retinoic acid receptor alpha gene to a novel tran-
scribed locus. *Nature* 347:

[143551]558–561, 1990.

[143552]25708.

[143553]de The, H.; Lavau, C.; Marchio, A.; Chomienne, C.; Degos, L.; Dejean,

[143554]A.: The PML–RAR–alpha fusion mRNA generated by the t(15;17) translocation

[143555]in acute promyelocytic leukemia encodes a functionally altered RAR. Cell 66:

[143556]675–684, 1991.

[143557]25709.

[143558]Di Croce, L.; Raker, V. A.; Corsaro, M.; Fazi, F.; Fanelli, M.;

[143559]Faretta, M.; Fuks, F.; Lo Coco, F.; Kouzarides, T.; Nervi, C.; Minucci,

[143560]S.; Pelicci, P. G.: Methyltransferase recruitment and DNA hypermethylation

[143561]of target promoters by an oncogenic transcription factor. Science 295:

[143562]1079–1082, 2002.

[143563]25710.

[143564]Dyck, J. A.; Maul, G. G.; Miller, W. H., Jr.; Chen, J. D.; Kakizuka,

[143565]A.; Evans, R. M.: A novel macromolecular structure is a
target of
[143566]the promyelocyte-retinoic acid receptor oncoprotein. Cell
76: 333-343,
[143567]1994.
[143568]25711.
[143569]Goddard, A. D.; Borrow, J.; Freemont, P. S.; Solomon, E.:
Characterization
[143570]of a zinc finger gene disrupted by the t(15;17) in acute
promyelocytic
[143571]leukemia. Science 254: 1371-1374, 1991.
[143572]25712.
[143573]Goddard, A. D.; Yuan, J. Q.; Fairbairn, L.; Dexter, M.; Bor-
row,
[143574]J.; Kozak, C.; Solomon, E.: Cloning of the murine homolog
of the
[143575]leukemia-associated PML gene. Mammalian Genome 6:
732-737, 1995.
[143576]25713.
[143577]Grignani, F.; Ferrucci, P. F.; Testa, U.; Talamo, G.; Fagioli,
[143578]M.; Alcalay, M.; Mencarelli, A.; Grignani, F.; Peschle, C.;

Nicoletti,

[143579]I.; Pelicci, P. G.: The acute promyelocytic leukemia-specific
PML-RAR-alpha

[143580]fusion protein inhibits differentiation and promotes survival of myeloid

[143581]precursor cells. Cell 74: 423-431, 1993.

[143582]25714.

[143583]Diverio, D.; Lo Coco, F.; D'Adamo, F.; Biondi, A.; Fagioli,
M.;

[143584]Grignani, F.; Rambaldi, A.; Rossi, V.; Avvisati, G.; Petti, M.
C.;

[143585]Testi, A. M.; Liso, V.; Specchia, G.; Fioritoni, G.; Recchia,
A.;

[143586]Frassoni, F.; Ciolli, S.; Pelicci, P. G.: Identification of DNA
rearrangements

[143587]at the retinoic acid receptor-alpha (RAR-alpha) locus in all
patients

[143588]with acute promyelocytic leukemia and mapping of APL
breakpoints within

[143589]the RAR-alpha second intron. Blood 79: 3331-3336,
1992.

[143590]25715.

[143591]Khan, M. M.; Nomura, T.; Kim, H.; Kaul, S. C.; Wadhwa, R.;
Shinagawa,

[143592]T.; Ichikawa-Iwata, E.; Zhong, S.; Pandolfi, P. P.; Ishii, S.:
Role

[143593]of PML and PML-RAR-alpha in Mad-mediated transcrip-
tional repression. Molec.

[143594]Cell 7: 1233-1243, 2001.

[143595]25716.

[143596]Lo Coco, F.; Diverio, D.; Falini, B.; Biondi, A.; Nervi, C.;
Pelicci,

[143597]P. G.: Genetic diagnosis and molecular monitoring in the
management

[143598]of acute promyelocytic leukemia. Blood 94: 12-22, 1999.

[143599]25717.

[143600]Pearson, M.; Carbone, R.; Sebastiani, C.; Cioce, M.; Fagioli,

[143601]M.; Saito, S.; Higashimoto, Y.; Appella, E.; Minucci, S.;
Pandolfi,

[143602]P. P.; Pelicci, P. G.: PML regulates p53 acetylation and pre-
mature

[143603]senescence induced by oncogenic Ras. Nature 406:
207-210, 2000.

[143604]25718.

[143605]Tong, J.-H.; Dong, S.; Geng, J.-P.; Huang, W.; Wang, Z.-Y.;
Sun,

[143606]G.-L.; Chen, S.-J.; Chen, Z.; Larsen, C.-J.; Berger, R.:
Molecular

[143607]rearrangements of the MYL gene in acute promyelocytic
leukemia (APL,

[143608]M3) define a breakpoint cluster region as well as some
molecular variants. Oncogene 7:

[143609]311-316, 1992.

[143610]25719.

[143611]Wang, Z. G.; Delva, L.; Gaboli, M.; Rivi, R.; Giorgio, M.;
Cordon-Cardo,

[143612]C.; Grosveld, F.; Pandolfi, P. P.: Role of PML in cell growth
and

[143613]the retinoic acid pathway. Science 279: 1547-1551, 1998.

[143614]25720.

[143615]Bunz, F.; Kobayashi, R.; Stillman, B.: cDNAs encoding the
large

[143616]subunit of human replication factor C. Proc. Nat. Acad.
Sci. 90:

[143617]11014-11018, 1993.

[143618]25721.

[143619]Lossie, A. C.; Haugen, B. R.; Wood, W. M.; Camper, S. A.;
Gordon,

[143620]D. F.: Chromosomal localization of the large subunit of
mouse replication

[143621]factor C in the mouse and human. Mammalian Genome 6:
58-59, 1995.

[143622]25722.

[143623]Luckow, B.; Bunz, F.; Stillman, B.; Lichter, P.; Schutz, G.:
Cloning,

[143624]expression, and chromosomal localization of the
140-kilodalton subunit

[143625]of replication factor C from mice and humans. Molec. Cell.
Biol. 14:

[143626]1626-1634, 1994.

[143627]25723.

[143628]Pennaneach, V.; Salles-Passador, I.; Munshi, A.; Brickner,
H.;

[143629]Regazzoni, K.; Dick, F.; Dyson, N.; Chen, T.-T.; Wang, J. Y.
J.; Fotedar,

[143630]R.; Fotedar, A.: The large subunit of replication factor C promotes

[143631]cell survival after DNA damage in an LxCxE motif- and Rb-dependent

[143632]manner. Molec. Cell 7: 715-727, 2001.

[143633]25724.

[143634]Uchiumi, F.; Ohta, T.; Tanuma, S.: Replication factor C recognizes

[143635]5-prime-phosphate ends of telomeres. Biochem. Biophys. Res. Commun. 229:

[143636]310-315, 1996.

[143637]25725.

[143638]Schmickel, R. D.: Contiguous gene syndromes: a component of recognizable

[143639]syndromes. J. Pediat. 109: 231-241, 1986.

[143640]25726.

[143641]Olson, E.; Srivastava, D.: Molecular pathways controlling heart

[143642]development. Science 272: 671-676, 1996.

[143643]25727.

[143644]Cunningham, J. M.; Vanin, E. F.; Tran, N.; Valentine, M.;
Jane,
[143645]S. M.: The human transcription factor CP2 (TFCP2), a com-
ponent of
[143646]the human gamma-globin stage selector protein, maps to
chromosome
[143647]region 12q13 and is within 250 kb of the NF-E2 gene. Ge-
nomics 30:
[143648]398-399, 1995.
[143649]25728.
[143650]Jane, S. M.; Nienhuis, A. W.; Cunningham, J. M.:
Hemoglobin switching
[143651]in man and chicken is mediated by a heteromeric complex
between the
[143652]ubiquitous transcription factor CP2 and a developmentally
specific
[143653]protein. EMBO J. 13: 197-105, 1995.
[143654]25729.
[143655]Lambert, J.-C.; Goumidi, L.; Wavrant-De Vrieze, F.;
Frigard, B.;
[143656]Harris, J. M.; Cummings, A.; Coates, J.; Pasquier, F.; Cotel,
D.;

- [143657] Gaillac, M.; St. Clair, D.; Mann, D. M. A.; Hardy, J.; London, C.
- [143658] L.; Amouyel, P.; Chartier-Harlin, M.-C.: The transcriptional factor
- [143659] LBP-1c/CP2/LSF gene on chromosome 12 is a genetic determinant of Alzheimer's
- [143660] disease. Hum. Molec. Genet. 9: 2275-2280, 2000.
- [143661] 25730.
- [143662] Swendeman, S. L.; Spielholz, C.; Jenkins, N. A.; Gilbert, D. J.;
- [143663] Copeland, N. G.; Sheffery, M.: Characterization of the genomic structure,
- [143664] chromosomal location, promoter, and developmental expression of the
- [143665] alpha-globin transcription factor CP2. J. Biol. Chem. 269: 11663-11671,
- [143666] 1994.
- [143667] 25731.
- [143668] Taylor, A. E.; Yip, A.; Brayne, C.; Easton, D.; Evans, J. G.; Xuereb,
- [143669] J.; Cairns, N.; Esiri, M. M.; Rubinsztein, D. C.: Genetic association

[143670] of an LBP-1c/CP2/LSF gene polymorphism with late onset
Alzheimer's

[143671] disease. J. Med. Genet. 38: 232-233, 2001.

[143672] 25732.

[143673] Bandara, L. R.; La Thangue, N. B.: Adenovirus E1a prevents
the

[143674] retinoblastoma gene product from complexing with a cel-
lular transcription

[143675] factor. Nature 351: 494-497, 1991.

[143676] 25733.

[143677] Chen, C.-R.; Kang, Y.; Siegel, P. M.; Massague, J.: E2F4/5
and

[143678] p107 as Smad cofactors linking the TGF-beta receptor to
c-myc repression. Cell 110:

[143679] 19-32, 2002.

[143680] 25734.

[143681] Girling, R.; Partridge, J. F.; Bandara, L. R.; Burden, N.;
Totty,

[143682] N. F.; Hsuan, J. J.; La Thangue, N. B.: A new component of
the transcription

[143683] factor DRTF1/E2F. Nature 362: 83-87, 1993. Note: Cor-

rection: Nature

[143684]365: 468 only, 1993.

[143685]25735.

[143686]Malas, S.; Duthie, S. M.; Mohri, F.; Lovell-Badge, R.;

Episkopou,

[143687]V.: Cloning and mapping of the human SOX1: a highly

conserved gene

[143688]expressed in the developing brain. Mammalian Genome 8:

866–868,

[143689]1997.

[143690]25736.

[143691]Zhang, Y.; Venkatraj, V. S.; Fischer, S. G.; Warburton, D.;

Chellappan,

[143692]S. P.: Genomic cloning and chromosomal assignment of

the E2F dimerization

[143693]partner TFDP gene family. Genomics 39: 95–98, 1997.

[143694]25737.

[143695] Moloney, D. J.; Panin, V. M.; Johnston, S. H.; Chen, J.; Shao,
[143696] L.; Wilson, R.; Wang, Y.; Stanley, P.; Irvine, K. D.; Halti-
wanger,
[143697] R. S.; Vogt, T. F.: Fringe is a glycosyltransferase that mod-
ifies
[143698] Notch. Nature 406: 369–375, 2000.
[143699] 25738.
[143700] Sestan, N.; Artavanis-Tsakonas, S.; Rakic, P.: Contact-
dependent
[143701] inhibition of cortical neurite growth mediated by Notch
signaling. Science 286:
[143702] 741–746, 1999.
[143703] 25739.
[143704] Tanigaki, K.; Nogaki, F.; Takahashi, J.; Tashiro, K.;
Kurooka,
[143705] H.; Honjo, T.: Notch1 and Notch3 instructively restrict
bFGF-responsive
[143706] multipotent neural progenitor cells to an astroglial fate.
Neuron 29:
[143707] 45–55, 2001.
[143708] 25740.

- [143709]Barton, D. E.; Foellmer, B. E.; Du, J.; Tamm, J.; Derynck, R.;
- [143710]Francke, U.: Chromosomal mapping of genes for trans-
forming growth
- [143711]factors beta-2 and beta-3 in man and mouse: dispersion
of TGF-beta
- [143712]gene family. Oncogene Res. 3: 323-331, 1988.
- [143713]25741.
- [143714]Dean, M.; Park, M.; Vande Woude, G. F.: Characterization
of the
- [143715]rearranged TPR-MET oncogene breakpoint. Molec. Cell.
Biol. 7: 921-924,
- [143716]1987.
- [143717]25742.
- [143718]Lemmink, H. H.; Mochizuki, T.; van den Heuvel, L. P. W. J.;
Schroder,
- [143719]C. H.; Barrientos, A.; Monnens, L. A. H.; van Oost, B. A.;
Brunner,
- [143720]H. G.; Reeders, S. T.; Smeets, H. J. M.: Mutations in the
type IV
- [143721]collagen alpha-3 (COL4A3) gene in autosomal recessive
Alport syndrome. Hum.

[143722]Molec. Genet. 3: 1269–1273, 1994.

[143723]25743.

[143724]Gibson, R. A.; Buchwald, M.; Roberts, R. G.; Mathew, C. G.:
Characterisation

[143725]of the exon structure of the Fanconi anaemia group C
gene by vectorette

[143726]PCR. Hum. Molec. Genet. 2: 35–38, 1993.

[143727]25744.

[143728]Gregersen, N.; Andresen, B. S.; Corydon, M. J.; Corydon, T.
J.;

[143729]Olsen, R. K. J.; Bolund, L.; Bross, P.: Mutation analysis in
mitochondrial

[143730]fatty acid oxidation defects: exemplified by acyl–CoA de-
hydrogenase

[143731]deficiencies, with special focus on genotype–phenotype
relationship. Hum.

[143732]Mutat. 18: 169–189, 2001.

[143733]25745.

[143734]Onkenhout, W.; Venizelos, V.; Scholte, H. R.; De Klerk, J. B.

[143735]C.; Poorthuis, B. J. H. M.: Intermediates of unsaturated
fatty acid

- [143736]oxidation are incorporated in triglycerides but not in phospholipids
- [143737]in tissues from patients with mitochondrial beta-oxidation defects. J.
- [143738]Inherit. Metab. Dis. 24: 337-344, 2001.
- [143739]25746.
- [143740]Lemmink, H. H.; Schroder, C. H.; Monnens, L. A. H.; Smeets, H.
- [143741]J. M.: The clinical spectrum of type IV collagen mutations. Hum.
- [143742]Mutat. 9: 477-499, 1997.
- [143743]25747.
- [143744]Goldmuntz, E.; Wang, Z.; Roe, B. A.; Budarf, M. L.: Cloning, genomic
- [143745]organization, and chromosomal localization of human citrate transport
- [143746]protein to the DiGeorge/velocardiofacial syndrome minimal critical
- [143747]region. Genomics 33: 271-276, 1996.
- [143748]25748.
- [143749]Heisterkamp, N.; Mulder, M. P.; Langeveld, A.; ten Hoeve,

J.; Wang,

[143750]Z.; Roe, B. A.; Groffen, J.: Localization of the human mitochondrial

[143751]citrate transporter protein gene to chromosome 22q11 in the DiGeorge

[143752]syndrome critical region. Genomics 29: 451–456, 1995.

[143753]25749.

[143754]Iacobazzi, V.; Lauria, G.; Palmieri, F.: Organization and sequence

[143755]of the human gene for the mitochondrial citrate transport protein. DNA

[143756]Seq. 7: 127–139, 1997.

[143757]25750.

[143758]Kaplan, R. S.; Mayor, J. A.; Wood, D. O.: The mitochondrial tricarboxylate

[143759]transport protein: cDNA cloning, primary structure, and comparison

[143760]with other mitochondrial transport proteins. J. Biol. Chem. 268:

[143761]13682–13690, 1993.

[143762]25751.

[143763]Stoffel, M.; Karayiorgou, M.; Espinosa, R., III; Le Beau, M.
M.

[143764]: The human mitochondrial citrate transporter gene
(SLC20A3) maps

[143765]to chromosome band 22q11 within a region implicated in
DiGeorge syndrome,

[143766]velo-cardio-facial syndrome and schizophrenia. Hum.
Genet. 98: 113-115,

[143767]1996.

[143768]25752.

[143769]Uberbacher, E. C.; Mural, R. J.: Locating protein-coding
regions

[143770]in human DNA sequences by a multiple sensor-neural
network approach. Proc.

[143771]Nat. Acad. Sci. 88: 11261-11265, 1991.

[143772]25753.

[143773]Gibson, R. A.; Hajianpour, A.; Buchwald, M.; Mathew, C.
G.: Screening

[143774]the FACC gene for mutations in patients with Fanconi
anaemia. (Abstract) Am.

[143775]J. Hum. Genet. 51 (suppl.): A216 only, 1992.

[143776]25754.

[143777]Vidal, R.; Frangione, B.; Rostagno, A.; Mead, S.; Revesz, T.;
Plant,

[143778]G.; Ghiso, J.: A stop-codon mutation in the BRI gene asso-
ciated with

[143779]familial British dementia. *Nature* 399: 776–781, 1999.

[143780]25755.

[143781]Joensuu, T.; Blanco, G.; Pakarinen, L.; Sistonen, P.; Kaari-
ainen,

[143782]H.; Brown, S.; de la Chapelle, A.; Sankila, E.-M.: Refined
mapping

[143783]of the Usher syndrome type III locus on chromosome 3,
exclusion of

[143784]candidate genes, and identification of the putative mouse
homologous

[143785]region. *Genomics* 38: 255–263, 1996.

[143786]25756.

[143787]Lamhonwah, A.-M.; Tein, I.: Carnitine uptake defect:
frameshift

[143788]mutations in the human plasmalemmal carnitine trans-
porter gene. *Biochem.*

[143789]Biophys. Res. Commun. 252: 396–401, 1998.

[143790]25757.

[143791]Matsuishi, T.; Hirata, K.; Terasawa, K.; Kato, H.; Yoshino, M.;

[143792]Ohtaki, E.; Hirose, F.; Nonaka, I.; Sugiyama, N.; Ohta, K.:
Successful

[143793]carnitine treatment in two siblings having lipid storage
myopathy

[143794]with hypertrophic cardiomyopathy. Neuropediatrics 16:
6–12, 1985.

[143795]25758.

[143796]Nezu, J.; Tamai, I.; Oku, A.; Ohashi, R.; Yabuuchi, H.;
Hashimoto,

[143797]N.; Nikaido, H.; Sai, Y.; Koizumi, A.; Shoji, Y.; Takada, G.;
Matsuishi,

[143798]T.; Yoshino, M.; Kato, H.; Ohura, T.; Tsujimoto, G.;
Hayakawa, J.;

[143799]Shimane, M.; Tsuji, A.: Primary systemic carnitine defi-
ciency is

[143800]caused by mutations in a gene encoding sodium ion-
dependent carnitine

[143801]transporter. Nature Genet. 21: 91–94, 1999.

[143802]25759.

[143803]Shoji, Y.; Koizumi, A.; Kayo, T.; Ohata, T.; Takahashi, T.;
Harada,

[143804]K.; Takada, G.: Evidence for linkage of human primary
systemic carnitine

[143805]deficiency with D5S436: a novel gene locus on chromo-
some 5q. Am.

[143806]J. Hum. Genet. 63: 101-108, 1998.

[143807]25760.

[143808]Tamai, I.; Ohashi, R.; Nezu, J.; Yabuuchi, H.; Oku, A.; Shi-
mane,

[143809]M.; Sai, Y.; Tsuji, A.: Molecular and functional identifica-
tion of

[143810]sodium ion-dependent, high affinity human carnitine
transporter OCTN2. J.

[143811]Biol. Chem. 273: 20378-20382, 1998.

[143812]25761.

[143813]Tein, I.; De Vivo, D. C.; Bierman, F.; Pulver, P.; De Meirleir,

[143814]L. J.; Cvitanovic-Sojat, L.; Pagon, R. A.; Bertini, E.; Dionisi-
Vici,

[143815]C.; Servidei, S.; Dimauro, S.: Impaired skin fibroblast car-

nitine

[143816]uptake in primary systemic carnitine deficiency manifested by childhood

[143817]carnitine-responsive cardiomyopathy. *Pediat. Res.* 28: 247–255, 1990.

[143818]25762.

[143819]Wang, Y.; Korman, S. H.; Ye, J.; Gargus, J. J.; Gutman, A.; Taroni,

[143820]F.; Garavaglia, B.; Longo, N.: Phenotype and genotype variation in

[143821]primary carnitine deficiency. *Genet. Med.* 3: 387–392, 2001.

[143822]25763.

[143823]Engelkamp, D.; Schafer, B. W.; Mattei, M. G.; Erne, P.; Heizmann,

[143824]C. W.: Six S100 genes are clustered on human chromosome 1q21: identification

[143825]of two genes coding for the two previously unreported calcium-binding

[143826]proteins S100D and S100E. *Proc. Nat. Acad. Sci.* 90: 6547–6551, 1993.

[143827]25764.

- [143828]Herrera, G. A.; Turbat-Herrera, E. A.; Lott, R. L.: S-100 protein
expression by primary and metastatic adenocarcinomas.
Am. J. Clin.
Path. 89: 168-176, 1988.
- [143831]25765.
- [143832]Isobe, T.; Tsugira, A.; Okuyama, T.: Amino acid sequence
of the
subunit structure of bovine brain S-100 protein (PAP 1-b).
J. Neurochem. 30:
921-923, 1978.
- [143835]25766.
- [143836]Moore, B. W.: A soluble protein characteristic of the ner-
vous
system. Biochem. Biophys. Res. Commun. 19: 739-744,
1965.
- [143838]25767.
- [143839]Morii, K.; Tanaka, R.; Takahashi, Y.; Minoshima, S.;
Fukuyama,
R.; Shimizu, N.; Kuwano, R.: Structure and chromosome
assignment

- [143841]of human S100 alpha and beta subunit genes. Biochem. Biophys. Res.
- [143842]Commun. 175: 185–191, 1991.
- [143843]25768.
- [143844]Most, P.; Bernotat, J.; Ehlermann, P.; Pleger, S. T.; Reppel, M.;
- [143845]Borries, M.; Niroomand, F.; Pieske, B.; Janssen, P. M. L.; Eschenhagen,
- [143846]T.; Karczewski, P.; Smith, G. L.; Koch, W. J.; Katus, H. A.; Remppis,
- [143847]A.: S100A1: a regulator of myocardial contractility. Proc. Nat.
- [143848]Acad. Sci. 98: 13889–13894, 2001.
- [143849]25769.
- [143850]Asada, Y.; Nadeau, J. H.: Fert is on mouse chromosome 11, not
- [143851]chromosome 17. Mammalian Genome 5: 830 only, 1994.
- [143852]25770.
- [143853]Hao, Q.–L.; Heisterkamp, N.; Groffen, J.: Isolation and sequence
- [143854]analysis of a novel human tyrosine kinase gene. Molec.

Cell. Biol. 9:

[143855]1587-1593, 1989.

[143856]25771.

[143857]Dryja, T. P.; Grondin, V. J.; Ringens, P.; Cotran, P.; Berson,

[143858]E. L.; Travis, G.: Isolation of human retinal cDNA fragments homologous

[143859]to the murine rds gene transcript. (Abstract) Invest. Ophthalmol. Vis.

[143860]Sci. 30 (suppl.): 43 only, 1989.

[143861]25772.

[143862]Farrar, G. J.; Jordan, S. A.; Kenna, P.; Humphries, M. M.; Kumar-Singh,

[143863]R.; McWilliam, P.; Allamand, V.; Sharp, E.; Humphries, P.: Autosomal

[143864]dominant retinitis pigmentosa: localization of a disease gene (RP6)

[143865]to the short arm of chromosome 6. Genomics 11: 870-874, 1991.

[143866]25773.

[143867]Farrar, G. J.; Kenna, P.; Jordan, S. A.; Kumar-Singh, R.; Humphries,

[143868]M. M.; Sharp, E. M.; Sheils, D. M.; Humphries, P.: A three-
base-pair

[143869]deletion in the peripherin-RDS gene in one form of retini-
tis pigmentosa. Nature 354:

[143870]478-480, 1991.

[143871]25774.

[143872]Feist, R. M.; White, M. F., Jr.; Skalka, H.; Stone, E. M.:
Choroidal

[143873]neovascularization in a patient with adult foveomacular
dystrophy

[143874]and a mutation in the retinal degeneration slow gene
(pro210-to-arg). Am.

[143875]J. Ophthalm. 118: 259-260, 1994.

[143876]25775.

[143877]Felbor, U.; Schilling, H.; Weber, B. H. F.: Adult vitelliform

[143878]macular dystrophy is frequently associated with mutations
in the peripherin/RDS

[143879]gene. Hum. Mutat. 10: 301-309, 1997.

[143880]25776.

[143881]Gass, J. D. M.: A clinicopathologic study of a peculiar
foveomacular

[143882]dystrophy. Trans. Am. Ophthal. Soc. 72: 139–155, 1974.

[143883]25777.

[143884]Jackson, K. E.; Mitchell, E. B.; Stone, E. M.; Ferrell, R. E.;

[143885]Gorin, M. B.: The identification of an exon–2 peripherin
mutation

[143886]in a family with heterogeneous manifestations of a butter-
fly pattern

[143887]macular dystrophy. (Abstract) Am. J. Hum. Genet. 53
(suppl.): 1177

[143888]only, 1993.

[143889]25778.

[143890]Jordan, S. A.; Farrar, G. J.; Kumar–Singh, R.; Kenna, P.;
Humphries,

[143891]M. M.; Allamand, V.; Sharp, E. M.; Humphries, P.: Autosomal dominant
mal dominant

[143892]retinitis pigmentosa (adRP; RP6): cosegregation of RP6
and the peripherin–RDS

[143893]locus in a late–onset family of Irish origin. Am. J. Hum.
Genet. 50:

[143894]634–639, 1992.

[143895]25779.

[143896]Billingsley, G. D.; Walter, M. A.; Hammond, G. L.; Cox, D.

W.:

[143897]Physical mapping of four serpin genes: alpha-

1-antitrypsin, alpha-1-antichymotrypsin,

[143898]corticosteroid-binding globulin, and protein C inhibitor,

within a

[143899]280-kb region on chromosome 14q32.1. Am. J. Hum.

Genet. 52: 343-353,

[143900]1993.

[143901]25780.

[143902]Clifton-Bligh, R. J.; Wentworth, J. M.; Heinz, P.; Crisp, M.

S.;

[143903]John, R.; Lazarus, J. H.; Ludgate, M.; Chatterjee, V. K.: Mu-

tation

[143904]of the gene encoding human TTF-2 associated with thy-

roid agenesis,

[143905]cleft palate and choanal atresia. Nature Genet. 19:

399-401, 1998.

[143906]25781.

[143907]Bird, T. D.; Koerker, R. M.; Leaird, B. J.; Vlcek, B. W.;

Thorning,

[143908]D. R.: Lipomembranous polycystic osteodysplasia (brain,

bone, and

[143909]fat disease): a genetic cause of presenile dementia. Neu-
rology 33:

[143910]81-86, 1983.

[143911]25782.

[143912]Paloneva, J.; Manninen, T.; Christman, G.; Hovanes, K.;
Mandelin,

[143913]J.; Adolfsson, R.; Bianchin, M.; Bird, T.; Miranda, R.;
Salmaggi,

[143914]A.; Tranebjaerg, L.; Konttinen, Y.; Peltonen, L.: Mutations
in two

[143915]genes encoding different subunits of a receptor signaling
complex

[143916]result in an identical disease phenotype. Am. J. Hum.
Genet. 71:

[143917]656-662, 2002.

[143918]25783.

[143919]Greco, A.; Ittmann, M.; Barletta, C.; Basilico, C.; Croce, C.
M.;

[143920]Cannizzaro, L. A.; Huebner, K.: Chromosomal localization
of human

[143921]genes required for G(1) progression in mammalian cells.

Genomics 4:

[143922]240–245, 1989.

[143923]25784.

[143924]Ittmann, M.; Greco, A.; Basilico, C.: Isolation of the human
gene

[143925]that complements a temperature-sensitive cell cycle mu-
tation in BHK

[143926]cells. Molec. Cell. Biol. 7: 3386–3393, 1987.

[143927]25785.

[143928]Bonyadi, M.; Cui, W.; Nagase, H.; Akhurst, R. J.: The TGF-
beta

[143929]type II receptor, Tgfbr2, maps to distal mouse chromo-
some 9. Genomics 33:

[143930]328–329, 1996.

[143931]25786.

[143932]Chen, R.-H.; Ebner, R.; Derynck, R.: Inactivation of the
type

[143933]II receptor reveals two receptor pathways for the diverse
TGF-beta

[143934]activities. Science 260: 1335–1338, 1993.

[143935]25787.

- [143936]Eshleman, J. R.; Markowitz, S. D.: Microsatellite instability
[143937]in inherited and sporadic neoplasms. *Curr. Opin. Oncol.* 7:
83–89,
[143938]1995.
[143939]25788.
- [143940]Hahm, K.-B.; Cho, K.; Lee, C.; Im, Y.-H.; Chang, J.; Choi,
S.-G.;
[143941]Sorensen, P. H. B.; Thiele, C. J.; Kim, S.-J.: Repression of
the
[143942]gene encoding the TGF- β type II receptor is a major
target of the
[143943]EWS-FLI1 oncoprotein. *Nature Genet.* 23: 222–227, 1999.
[143944]25789.
- [143945]Lin, H. Y.; Wang, X.-F.; Ng-Eaton, E.; Weinberg, R. A.;
Lodish,
[143946]H. F.: Expression cloning of the TGF- β type II receptor,
a functional
[143947]transmembrane serine/threonine kinase. *Cell* 68:
775–785, 1992.
[143948]25790.
- [143949]Lu, S.-L.; Kawabata, M.; Imamura, T.; Akiyama, Y.; No-

mizu, T.;

[143950]Miyazono, K.; Yuasa, Y.: HNPCC associated with germline mutation

[143951]in the TGF-beta type II receptor gene. (Letter) Nature Genet. 19:

[143952]17-18, 1998.

[143953]25791.

[143954]Markowitz, S.; Wang, J.; Myeroff, L.; Parsons, R.; Sun, L.; Lutterbaugh,

[143955]J.; Fan, R. S.; Zborowska, E.; Kinzler, K. W.; Vogelstein, B.; Brattain,

[143956]M.; Willson, J. K. V.: Inactivation of the type II TGF-beta receptor

[143957]in colon cancer cells with microsatellite instability. Science 268:

[143958]1336-1338, 1995.

[143959]25792.

[143960]Mathew, S.; Murty, V. V. V. S.; Cheifetz, S.; George, D.; Massague,

[143961]J.; Chaganti, R. S. K.: Transforming growth factor receptor gene

[143962]TGFBR2 maps to human chromosome band 3p22. Ge-

nomics 20: 114–115,

[143963]1994.

[143964]25793.

[143965]Myeroff, L. L.; Parsons, R.; Kim, S.-J.; Hedrick, L.; Cho, K.
R.;

[143966]Orth, K.; Mathis, M.; Kinzler, K. W.; Lutterbaugh, J.; Park,
K.; Bang,

[143967]Y.-J.; Lee, H. Y.; Park, J.-G.; Lynch, H. T.; Roberts, A. B.;
Vogelstein,

[143968]B.; Markowitz, S. D.: A transforming growth factor beta
receptor

[143969]type II gene mutation common in colon and gastric but
rare in endometrial

[143970]cancers with microsatellite instability. Cancer Res. 55:
5545–5547,

[143971]1995.

[143972]25794.

[143973]Parsons, R.; Myeroff, L. L.; Liu, B.; Willson, J. K. V.;
Markowitz,

[143974]S. D.; Kinzler, K. W.; Vogelstein, B.: Microsatellite instabil-
ity

[143975]and mutations of the transforming growth factor beta type

II receptor

[143976] gene in colorectal cancer. *Cancer Res.* 55: 5548–5550, 1995.

[143977] 25795.

[143978] Tanaka, S.; Mori, M.; Mafune, K.; Ohno, S.; Sugimachi, K.:

A

[143979] dominant negative mutation of transforming growth factor-beta receptor

[143980] type II gene in microsatellite stable oesophageal carcinoma. *Brit.*

[143981] *J. Cancer* 82: 1557–1560, 2000.

[143982] 25796.

[143983] Tannergard, P.; Liu, T.; Weger, A.; Nordenskjold, M.; Lindbloom,

[143984] A.: Tumorigenesis in colorectal tumors from patients with hereditary

[143985] non-polypoid colorectal cancer. *Hum. Genet.* 101: 51–55, 1997.

[143986] 25797.

[143987] Watanabe, T.; Wu, T.-T.; Catalano, P. J.; Ueki, T.; Satriano,

[143988] R.; Haller, D. G.; Benson, A. B., III; Hamilton, S. R.: *Molecu-*

lar

[143989]predictors of survival after adjuvant chemotherapy for
colon cancer. New

[143990]Eng. J. Med. 344: 1196–1206, 2001.

[143991]25798.

[143992]Bonnefont, J.-P.; Chretien, D.; Rustin, P.; Robinson, B.;
Vassault,

[143993]A.; Aupetit, J.; Charpentier, C.; Rabier, D.; Saudubray,
J.-M.; Munnich,

[143994]A.: Alpha-ketoglutarate dehydrogenase deficiency pre-
sented as congenital

[143995]lactic acidosis. J. Pediatr. 121: 255–258, 1992.

[143996]25799.

[143997]Guffon, N.; Lopez-Mediavilla, C.; Dumoulin, R.; Mousson,
B.; Godinot,

[143998]C.; Carrier, H.; Collombet, J. M.; Divry, P.; Mathieu, M.;
Guibaud,

[143999]P.: 2-Ketoglutarate dehydrogenase deficiency, a rare
cause of primary

[144000]hyperlactataemia: report of a new case. J. Inherit. Metab.
Dis. 16:

[144001]821–830, 1993.

[144002]25800.

[144003]Kohlschutter, A.; Behbehani, A.; Langenbeck, U.; Albani, M.; Heidemann,

[144004]P.; Hoffmann, G.; Kleineke, J.; Lehnert, W.; Wendel, U.: A familial

[144005]progressive neurodegenerative disease with 2-oxoglutaric aciduria. Europ.

[144006]J. Pediat. 138: 32-37, 1982.

[144007]25801.

[144008]Koike, K.: The gene encoding human 2-oxoglutarate dehydrogenase:

[144009]structural organization and mapping to chromosome 7p13-p14. Gene 159:

[144010]261-266, 1995.

[144011]25802.

[144012]Koike, K.; Urata, Y.; Goto, S.: Cloning and nucleotide sequence

[144013]of the cDNA encoding human 2-oxoglutarate dehydrogenase (lipoamide). Proc.

[144014]Nat. Acad. Sci. 89: 1963-1967, 1992.

[144015]25803.

[144016]Szabo, P.; Cai, X.; Ali, G.; Blass, J. P.: Localization of the
[144017]gene (OGDH) coding for the E1 α component of the alpha-
ketoglutarate
[144018]dehydrogenase complex to chromosome 7p13-p11.2.
Genomics 20: 324-326,
[144019]1994.
[144020]25804.
[144021]Kury, S.; Dreno, B.; Bezieau, S.; Giraudet, S.; Kharfi, M.;
Kamoun,
[144022]R.; Moisan, J.-P.: Identification of SLC39A4, a gene in-
volved in
[144023]acrodermatitis enteropathica. Nature Genet. 31: 239-240,
2002.
[144024]25805.
[144025]Rogers, E. E.; Eide, D. J.; Guerinot, M. L.: Altered selectivity
[144026]in an Arabidopsis metal transporter. Proc. Nat. Acad. Sci.
97: 12356-12360,
[144027]2000.
[144028]25806.
[144029]Wang, K.; Pugh, E. W.; Griffen, S.; Doheny, K. F.; Mostafa,
W.

- [144030]Z.; al-Aboosi, M. M.; el-Shanti, H.; Gitschier, J.: Homozygosity
- [144031]mapping places the acrodermatitis enteropathica gene on chromosomal
- [144032]region 8q24.3. Am. J. Hum. Genet. 68: 1055–1060, 2001.
- [144033]25807.
- [144034]Haberle, J.; Pauli, S.; Linnebank, M.; Kleijer, W. J.; Bakker,
- [144035]H. D.; Wanders, R. J. A.; Harms, E.; Koch, H. G.: Structure of the
- [144036]human argininosuccinate synthetase gene and an improved system for
- [144037]molecular diagnostics in patients with classical and mild citrullinemia. Hum.
- [144038]Genet. 110: 327–333, 2002.
- [144039]25808.
- [144040]Weve, H.: Ablatio falciformis congenita (retinal fold). Brit.
- [144041]J. Ophthal. 22: 456–470, 1938.
- [144042]25809.
- [144043]Barbosa, J.; Rich, S.; Dunsworth, T.; Swanson, J.: Linkage disequilibrium

[144044]between insulin-dependent diabetes and the Kidd blood group Jk(b)

[144045]allele. J. Clin. Endocr. Metab. 55: 193-195, 1982.

[144046]25810.

[144047]Harper, P. A.; Healy, P. J.; Dennis, J. A.; O'Brien, J. J.; Rayward,

[144048]D. H.: Citrullinaemia as a cause of neurological disease in neonatal

[144049]Friesian calves. Aust. Vet. J. 63: 378-379, 1986.

[144050]25811.

[144051]Harper, P. A. W.; Healy, P. J.; Dennis, J. A.: Animal model of

[144052]human disease: citrullinemia (argininosuccinate synthetase deficiency). Am.

[144053]J. Path. 135: 1213-1215, 1989.

[144054]25812.

[144055]Jackson, M. J.; Surh, L. C.; O'Brien, W. E.; Beaudet, A. L.:

[144056]Assignment of the structural gene for argininosuccinate synthetase

[144057]to proximal mouse chromosome 2. Genomics 6: 545-547, 1990.

[144058]25813.

[144059]Kobayashi, K.; Ichiki, H.; Saheki, T.; Tatsuno, M.;
Uchiyama,

[144060]C.; Nukada, O.; Yoda, T.: Structure of an abnormal messenger RNA

[144061]for argininosuccinate synthetase in citrullinemia. Hum.
Genet. 76:

[144062]27-32, 1987.

[144063]25814.

[144064]Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.;
Beaudet,

[144065]A. L.: Characterization of nine mutant alleles causing citrullinemia.

[144066](Abstract) Am. J. Hum. Genet. 45 (suppl.): A201 only,
1989.

[144067]25815.

[144068]Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.;
Beaudet,

[144069]A. L.: Heterogeneity of mutations in argininosuccinate
synthetase

[144070]causing human citrullinemia. J. Biol. Chem. 265:
11361-11367, 1990.

[144071]25816.

[144072]Kobayashi, K.; Kakinoki, H.; Fukushige, T.; Shaheen, N.;
Terazono,

[144073]H.; Saheki, T.: Nature and frequency of mutations in the
argininosuccinate

[144074]synthetase gene that cause classical citrullinemia. Hum.
Genet. 96:

[144075]454–463, 1995.

[144076]25817.

[144077]Kobayashi, K.; Rosenbloom, C.; Beaudet, A. L.; O'Brien, W.
E.

[144078]: Additional mutations in argininosuccinate synthetase
causing citrullinemia. Molec.

[144079]Biol. Med. 8: 95–100, 1991.

[144080]25818.

[144081]Kobayashi, K.; Saheki, T.; Imamura, Y.; Noda, T.; Inoue, I.;
Matuo,

[144082]S.; Hagihara, S.; Nomiya, H.; Jinno, Y.; Shimada, K.:
Messenger

[144083]RNA coding for argininosuccinate synthetase in citrulline-
mia. Am.

- [144084]J. Hum. Genet. 38: 667–680, 1986.
- [144085]25819.
- [144086]Kobayashi, K.; Shaheen, N.; Terazono, H.; Saheki, T.: Mu-
tations
[144087]in argininosuccinate synthetase mRNA of Japanese pa-
tients, causing
[144088]classical citrullinemia. Am. J. Hum. Genet. 55: 1103–1112,
1994.
- [144089]25820.
- [144090]Li, C.-M.; Chao, H.-K.; Liu, Y.-F.; Su, T.-S.: A nonsense
mutation
[144091]is responsible for the RNA-negative phenotype in human
citrullinaemia. Europ.
[144092]J. Hum. Genet. 9: 685–689, 2001.
- [144093]25821.
- [144094]Northrup, H.; Lathrop, M.; Lu, S.-Y.; Daiger, S. P.; Beaudet,
[144095]A. L.; O'Brien, W. E.: Multilocus linkage analysis with the
human
[144096]argininosuccinate synthetase gene. Genomics 5: 442–444,
1989.
- [144097]25822.

[144098] Sase, M.; Kobayashi, K.; Imamura, Y.; Saheki, T.; Nakano, K.;

[144099] Miura, S.; Mori, M.: Level of translatable messenger RNA coding for

[144100] argininosuccinate synthetase in the liver of the patients with quantitative-type

[144101] citrullinemia. Hum. Genet. 69: 130–134, 1985.

[144102] 25823.

[144103] Todd, S.; Naylor, S. L.: New chromosomal mapping assignments

[144104] for argininosuccinate synthetase pseudogene 1, interferon-beta-3 gene,

[144105] and the diazepam binding inhibitor gene. Somat. Cell Molec. Genet. 18:

[144106] 381–385, 1992.

[144107] 25824.

[144108] Hsieh, C.-L.; Swaroop, A.; Francke, U.: Chromosomal localization

[144109] and cDNA sequence of human RALB, a GTP binding protein. Somat. Cell

[144110] Molec. Genet. 16: 407–410, 1990.

[144111]25825.

[144112]Ahmad, F.; Goldstein, B. J.: Functional association between
the

[144113]insulin receptor and the transmembrane protein-tyrosine
phosphatase

[144114]LAR in intact cells. J. Biol. Chem. 272: 448-457, 1997.

[144115]25826.

[144116]Disteche, C. M.; Adler, D. A.; Tedder, T. F.; Saito, H.: Map-
ping

[144117]of the genes for LYAM1, a new lymphocyte adhesion
molecule, and for

[144118]LAR, a new receptor-linked protein tyrosine phosphatase,
to human

[144119]chromosome 1 (Abstract) Cytogenet. Cell Genet. 51: 990
only, 1989.

[144120]25827.

[144121]Harder, K. W.; Saw, J.; Miki, N.; Jirik, F.: Coexisting amplifi-
cations

[144122]of the chromosome 1p32 genes (PTPRF and MYCL1) en-
coding protein tyrosine

[144123]phosphatase LAR and L-myc in a small cell lung cancer
line. Genomics 27:

[144124]552-553, 1995.

[144125]25828.

[144126]Jirik, F. R.; Harder, K. W.; Melhado, I. G.; Anderson, L. L.;
Duncan,

[144127]A. M. V.: The gene for leukocyte antigen-related tyrosine
phosphatase

[144128](LAR) is localized to human chromosome 1p32, a region
frequently deleted

[144129]in tumors of neuroectodermal origin. Cytogenet. Cell
Genet. 61:

[144130]266-268, 1992.

[144131]25829.

[144132]Nam, H.-J.; Poy, F.; Krueger, N. X.; Saito, H.; Frederick, C.
A.

[144133]: Crystal structure of the tandem phosphatase domains of
RPTP LAR. Cell 97:

[144134]449-457, 1999.

[144135]25830.

[144136]O'Grady, P.; Krueger, N. X.; Streuli, M.; Saito, H.: Genomic
organization

[144137]of the human LAR protein tyrosine phosphatase gene and

alternative

[144138]splicing in the extracellular fibronectin type-III domains. J. Biol.

[144139]Chem. 269: 25193-25199, 1994.

[144140]25831.

[144141]Schaapveld, R. Q.; Schepens, J. T.; Robinson, G. W.; Attema, J.;

[144142]Oerlemans, F. T.; Fransen, J. A.; Streuli, M.; Wieringa, B.; Hennighausen,

[144143]L.; Hendriks, W. J.: Impaired mammary gland development and function

[144144]in mice lacking LAR receptor-like tyrosine phosphatase activity. Dev.

[144145]Biol. 188: 134-146, 1997.

[144146]25832.

[144147]Schaapveld, R. Q. J.; van den Maagdenberg, A. M. J. M.; Schepens,

[144148]J. T. G.; Olde Weghuis, D.; Geurts van Kessel, A.; Wieringa, B.; Hendriks,

[144149]W. J. A. J.: The mouse gene Ptp^{prf} encoding the leukocyte common antigen-related

[144150]molecule LAR: cloning, characterization, and chromoso-

mal localization. Genomics 27:

[144151]124–130, 1995.

[144152]25833.

[144153]Tsujioka, K.; Kawakami, N.; Uchino, Y.; Ichijo, T.; Furukawa,

[144154]T.; Saito, H.; Yamamoto, H.: Distinct functions of the two protein

[144155]tyrosine phosphatase domains of LAR (leukocyte common antigen-related)

[144156]on tyrosine dephosphorylation of insulin receptor. Molec. Endocr. 15:

[144157]271–280, 2001.

[144158]25834.

[144159]Ali, R. R.; Sarra, G.-M.; Stephens, C.; de Alwis, M.; Bainbridge,

[144160]J. W. B.; Munro, P. M.; Fauser, S.; Reichell, M. B.; Kinnon, C.; Hunt,

[144161]D. M.; Bhattacharya, S. S.; Thrasher, A. J.: Restoration of photoreceptor

[144162]ultrastructure and function in retinal degeneration slow mice by gene

[144163]therapy. Nature Genet. 25: 306–310, 2000.

[144164]25835.

[144165]Bascom, R. A.; Connell, G.; Garcia-Heras, J.; Collins, L.;
Ledbetter,

[144166]D.; Molday, R. S.; Kalnins, V.; McInnes, R. R.: Molecular
and ultrastructural

[144167]characterization of the products of the human retinopathy
candidate

[144168]genes ROM1 and RDS. (Abstract) Am. J. Hum. Genet. 47
(suppl.): A101

[144169]only, 1990.

[144170]25836.

[144171]Connell, G.; Bascom, R.; Molday, L.; Reid, D.; McInnes, R.
R.;

[144172]Molday, R. S.: Photoreceptor peripherin is the normal
product of

[144173]the gene responsible for retinal degeneration in the rds
mouse. Proc.

[144174]Nat. Acad. Sci. 88: 723-726, 1991.

[144175]25837.

[144176]Demant, P.; Ivanyi, D.; van Nie, R.: The map position of the
rds

[144177]gene on the 17th chromosome of the mouse. Tissue Anti-
gens 13: 53-55,

[144178]1979.

[144179]25838.

[144180]Inoue, C.; Shiga, K.; Takasawa, S.; Kitagawa, M.; Ya-
mamoto, H.;

[144181]Okamoto, H.: Evolutionary conservation of the insulinoma
gene rig

[144182]and its possible function. Proc. Nat. Acad. Sci. 84:
6659-6662,

[144183]1987.

[144184]25839.

[144185]Kitagawa, M.; Takasawa, S.; Kikuchi, N.; Itoh, T.; Teraoka,
H.;

[144186]Yamamoto, H.; Okamoto, H.: Rig encodes ribosomal pro-
tein S15: the

[144187]primary structure of mammalian ribosomal protein S15.
FEBS Lett. 283:

[144188]210-214, 1991.

[144189]25840.

[144190]Shiga, K.; Yamamoto, H.; Okamoto, H.: Isolation and char-

acterization

[144191] of the human homologue of rig and its pseudogenes: the functional

[144192] gene has features characteristic of housekeeping genes.

Proc. Nat.

[144193] Acad. Sci. 87: 3594–3598, 1990.

[144194] 25841.

[144195] Brody, L. C.; Abel, K. J.; Castilla, L. H.; Couch, F. J.; McKinley,

[144196] D. R.; Yin, G.-Y.; Ho, P. P.; Merajver, S.; Chandrasekharappa, S.

[144197] C.; Xu, J.; Cole, J. L.; Struwing, J. P.; Valdes, J. M.; Collins,

[144198] F. S.; Weber, B. L.: Construction of a transcription map surrounding

[144199] the BRCA1 locus of human chromosome 17. Genomics 25: 238–247, 1995.

[144200] 25842.

[144201] Hloch, P.; Schiedner, G.; Stahl, H.: Complete cDNA sequence of

[144202] the human p68 protein. Nucleic Acids Res. 18: 3045, 1990.

[144203] 25843.

[144204] Iggo, R.; Gough, A.; Xu, W.; Lane, D. P.; Spurr, N. K.: Chromosome

[144205] mapping of the human gene encoding the 68-kDa nuclear antigen (p68)

[144206] by using the polymerase chain reaction. Proc. Nat. Acad. Sci. 86:

[144207] 6211–6214, 1989.

[144208] 25844.

[144209] Iggo, R. D.; Xu, W.; Lane, D. P.; Gough, A.; Spurr, N. K.: Assignment

[144210] of P68 to chromosome 17 using PCR. (Abstract) Cytogenet. Cell Genet. 51:

[144211] 1017, 1989.

[144212] 25845.

[144213] Acker, J.; Mattei, M.-G.; Wintzerith, M.; Roeckel, N.; Depetris,

[144214] D.; Vigneron, M.; Keding, C.: Chromosomal localization of human

[144215] RNA polymerase II subunit genes. Genomics 20: 496–499, 1994.

[144216] 25846.

- [144217]Schulze, A.; Hansen, C.; Skakkebaek, N. E.; Brondum-Nielsen, K.;
- [144218]Ledbetter, D. H.; Tommerup, N.: Exclusion of SNRPN as a major determinant
- [144219]of Prader-Willi syndrome by a translocation breakpoint. Nature Genet. 12:
- [144220]452-454, 1996.
- [144221]25847.
- [144222]Schweizer, J.; Zynger, D.; Francke, U.: In vivo nuclease hypersensitivity
- [144223]studies reveal multiple sites of parental origin-dependent differential
- [144224]chromatin conformation in the 150 kb SNRPN transcription unit. Hum.
- [144225]Molec. Genet. 8: 555-566, 1999.
- [144226]25848.
- [144227]Shemer, R.; Hershko, A. Y.; Perk, J.; Mostoslavsky, R.; Tsuberi,
- [144228]B.; Cedar, H.; Buiting, K.; Razin, A.: The imprinting box of the
- [144229]Prader-Willi/Angelman syndrome domain. Nature Genet. 26: 440-443,

[144230]2000.

[144231]25849.

[144232]Sun, Y.; Nicholls, R. D.; Butler, M. G.; Saitoh, S.; Hainline,

[144233]B. E.; Palmer, C. G.: Breakage in the SNRPN locus in a balanced 46,XY,t(15;19)

[144234]Prader-Willi syndrome patient. Hum. Molec. Genet. 5: 517-524, 1996.

[144235]25850.

[144236]Sutcliffe, J. S.; Nakao, M.; Christian, S.; Orstavik, K. H.; Tommerup,

[144237]N.; Ledbetter, D. H.; Beaudet, A. L.: Deletions of a differentially

[144238]methylated CpG island at the SNRPN gene define a putative imprinting

[144239]control region. Nature Genet. 8: 52-58, 1994.

[144240]25851.

[144241]Wirth, J.; Back, E.; Huttenhofer, A.; Nothwang, H.-G.; Lich, C.;

[144242]Gross, S.; Menzel, C.; Schinzel, A.; Kioschis, P.; Tommerup, N.; Ropers,

[144243]H.-H.; Horsthemke, B.; Buiting, K.: A translocation break-

point cluster

[144244]disrupts the newly defined 3-prime end of the SNURF-SNRPN transcription

[144245]unit on chromosome 15. Hum. Molec. Genet. 10: 201-210, 2001.

[144246]25852.

[144247]Critcher, R.; Stitson, R. N. M.; Wade-Martins, R.; Easty, D. J.;

[144248]Farr, C. J.: Assignment of Sox4 to mouse chromosome 13 bands A3-A5

[144249]by fluorescence in situ hybridization; refinement of the human SOX4

[144250]location to 6p22.3 and of SOX20 to chromosome 17p12.3. Cytogenet.

[144251]Cell Genet. 81: 294-295, 1998.

[144252]25853.

[144253]Denny, P.; Swift, S.; Connor, F.; Ashworth, A.: An SRY-related

[144254]gene expressed during spermatogenesis in the mouse encodes a sequence-specific

[144255]DNA-binding protein. EMBO J. 11: 3705-3712, 1992.

[144256]25854.

- [144257]Farr, C. J.; Easty, D. J.; Ragoussis, J.; Collignon, J.; Lovell-Badge,
- [144258]R.; Goodfellow, P. N.: Characterization and mapping of the human
- [144259]SOX4 gene. Mammalian Genome 4: 577–584, 1993.
- [144260]25855.
- [144261]Hansen, G. M.; Skapura, D.; Justice, M. J.: Genetic profile of
- [144262]insertion mutations in mouse leukemias and lymphomas. Genome Res. 10:
- [144263]237–243, 2000.
- [144264]25856.
- [144265]Li, J.; et al; et al: Leukaemia disease genes: large-scale cloning
- [144266]and pathway predictions. Nature Genet. 23: 348–353, 1999.
- [144267]25857.
- [144268]Lund, A. H.; Turner, G.; Trubetskoy, A.; Verhoeven, E.; Wientjens,
- [144269]E.; Hulsman, D.; Russell, R.; DePinho, R. A.; Lenz, J.; van Lohuizen,

- [144270]M.: Genome-wide retroviral insertional tagging of genes involved
in cancer in Cdkn2a-deficient mice. Nature Genet. 32: 160–165, 2002.
- [144271]25858.
- [144273]Suzuki, T.; Shen, H.; Akagi, K.; Morse, H. C., III; Malley, J. D.; Naiman, D. Q.; Jenkins, N. A.; Copeland, N. G.: New genes involved
in cancer identified by retroviral tagging. Nature Genet. 32: 166–174, 2002. Note: Erratum: Nature Genet. 32: 331 only, 2002.
- [144276]25859.
- [144278]van de Wetering, M.; Oosterwegel, M.; van Norren, K.; Clevers, H.: Sox-4, an Sry-like HMG box protein, is a transcriptional activator
in lymphocytes. EMBO J. 12: 3847–3854, 1993.
- [144281]25860.
- [144282]Shoyab, M.; McDonald, V. L.; Bradley, J. G.; Todaro, G. J.: Amphiregulin:
a bifunctional growth-modulating glycoprotein produced

by the phorbol

[144284]12-myristate 13-acetate-treated human breast adenocarcinoma cell line

[144285]MCF-7. Proc. Nat. Acad. Sci. 85: 6528-6532, 1988.

[144286]25861.

[144287]Shoyab, M.; Plowman, G. D.; McDonald, V. L.; Bradley, J. G.; Todaro,

[144288]G. J.: Structure and function of human amphiregulin: a member of

[144289]the epidermal growth factor family. Science 243: 1074-1076, 1989.

[144290]25862.

[144291]Donahue, R. P.; Bias, W. B.; Renwick, J. H.; McKusick, V. A.:

[144292]Probable assignment of the Duffy blood group locus to chromosome 1

[144293]in man. Proc. Nat. Acad. Sci. 61: 949-955, 1968.

[144294]25863.

[144295]Jiang, Y.; Armstrong, D.; Albrecht, U.; Atkins, C. M.; Noebels,

[144296]J. L.; Eichele, G.; Sweatt, J. D.; Beaudet, A. L.: Mutation of the

[144297] Angelman ubiquitin ligase in mice causes increased cytoplasmic p53

[144298] and deficits of contextual learning and long-term potentiation. *Neuron* 21: 799–811, 1998.

[144300] 25864.

[144301] Kishino, T.; Lalande, M.; Wagstaff, J.: UBE3A/E6-AP mutations

[144302] cause Angelman syndrome. *Nature Genet.* 15: 70–73, 1997.

[144303] 25865.

[144304] Matsuura, T.; Sutcliffe, J. S.; Fang, P.; Galjaard, R.-J.; Jiang,

[144305] Y.; Benton, C. S.; Rommens, J. M.; Beaudet, A. L.: De novo truncating

[144306] mutations in E6-AP ubiquitin-protein ligase gene (UBE3A) in Angelman

[144307] syndrome. *Nature Genet.* 15: 74–77, 1997.

[144308] 25866.

[144309] Ono, H.; Kuno, Y.; Tanaka, H.; Yamashina, M.; Tsuyoshi, T.; Kondo,

[144310] N.; Orii, T.: A case of paroxysmal nocturnal hemoglobin-

uria without

[144311]deficiency of decay-accelerating factor on erythrocytes.

Blood 75:

[144312]1746–1747, 1990.

[144313]25867.

[144314]Petranka, J. G.; Fleenor, D. E.; Sykes, K.; Kaufman, R. E.;

Rosse,

[144315]W. F.: Structure of the CD59–encoding gene: further evidence of a

[144316]relationship to murine lymphocyte antigen Ly–6 protein.

Proc. Nat.

[144317]Acad. Sci. 89: 7876–7879, 1992.

[144318]25868.

[144319]Rosse, W. F.: Personal Communication. Durham, N. C.

6/3/1993.

[144320]25869.

[144321]Rosse, W. F.; Parker, C. J.: Paroxysmal nocturnal hemoglobinuria. Clin.

[144322]Haemat. 14: 105–125, 1985.

[144323]25870.

[144324]Rother, R. P.; Rollins, S. A.; Mennone, J.; Chodera, A.; Fi-

del,

[144325]S. A.; Bessler, M.; Hillmen, P.; Squinto, S. P.: Expression of recombinant

[144326]transmembrane CD59 in paroxysmal nocturnal hemoglobinuria B cells

[144327]confers resistance to human complement. Blood 84: 2604–2611, 1994.

[144328]25871.

[144329]Tone, M.; Walsh, L. A.; Waldmann, H.: Gene structure of human

[144330]CD59 and demonstration that discrete mRNAs are generated by alternative

[144331]polyadenylation. J. Molec. Biol. 227: 971–976, 1992.

[144332]25872.

[144333]Walsh, L. A.; Tone, M.; Thiru, S.; Waldmann, H.: The CD59 antigen--a

[144334]multifunctional molecule. Tissue Antigens 40: 213–220, 1992.

[144335]25873.

[144336]Yamashina, M.; Ueda, E.; Kinoshita, T.; Takami, T.; Ojima, A.;

[144337]Ono, H.; Tanaka, H.; Kondo, N.; Orii, T.; Okada, N.; Okada,

- H.; Inoue,
[144338]K.; Kitani, T.: Inherited complete deficiency of
20-kilodalton homologous
[144339]restriction factor (CD59) as a cause of paroxysmal noctur-
nal hemoglobinuria. New
[144340]Eng. J. Med. 323: 1184-1189, 1990.
[144341]25874.
- [144342]Milatovich, A.; Bolger, G.; Michaeli, T.; Francke, U.: Chro-
mosome
[144343]localizations of genes for five cAMP-specific phosphodi-
esterases in
[144344]man and mouse. Somat. Cell Molec. Genet. 20: 75-86,
1994.
[144345]25875.
- [144346]Loughney, K.; Martins, T. J.; Harris, E. A. S.; Sadhu, K.;
Hicks,
[144347]J. B.; Sonnenburg, W. K.; Beavo, J. A.; Ferguson, K.: Isola-
tion and
[144348]characterization of cDNAs corresponding to two human
calcium, calmodulin-regulated,
[144349]3-prime,5-prime-cyclic nucleotide phosphodiesterases. J.
Biol. Chem. 271:

[144350]796–806, 1996.

[144351]25876.

[144352]Wilson, D. E.; McKenna, L.: Assignment of the human gene
for phosphodiesterase

[144353]1A to chromosome 4. (Abstract) Am. J. Hum. Genet. 43:
A162 only,

[144354]1988.

[144355]25877.

[144356]Maruo, Y.; Nishizawa, K.; Sato, H.; Sawa, H.; Shimada, M.:
Prolonged

[144357]unconjugated hyperbilirubinemia associated with breast
milk and mutations

[144358]of the bilirubin uridine diphosphate–glu-
curonosyltransferase gene. Pediatrics 106:

[144359]e59, 2000. Note: Electronic Article.

[144360]25878.

[144361]Miles, J. S.; Moss, J. E.; Taylor, B. A.; Burchell, B.; Wolf,

[144362]C. R.: Mapping genes encoding drug–metabolizing en-
zymes in recombinant

[144363]inbred mice. Genomics 11: 309–316, 1991.

[144364]25879.

[144365]Moghrabi, N.; Clarke, D. J.; Boxer, M.; Burchell, B.: Identifi-
cation

[144366]of an A-to-G missense mutation in exon 2 of the UGT1
gene complex

[144367]that causes Crigler-Najjar syndrome type 2. Genomics 18:
171-173,

[144368]1993.

[144369]25880.

[144370]Moghrabi, N.; Clarke, D. J.; Burchell, B.; Boxer, M.: Coseg-
regation

[144371]of intragenic markers with a novel mutation that causes
Crigler-Najjar

[144372]syndrome type I: implication in carrier detection and pre-
natal diagnosis. Am.

[144373]J. Hum. Genet. 53: 722-729, 1993.

[144374]25881.

[144375]Moghrabi, N.; Sutherland, L.; Wooster, R.; Povey, S.; Boxer,
M.;

[144376]Burchell, B.: Chromosomal assignment of human phenol
and bilirubin

[144377]UDP-glucuronosyltransferase genes (UGT1A-subfamily).

Ann. Hum. Genet. 56:

[144378]81-91, 1992.

[144379]25882.

[144380]Mojarrabi, B.; Butler, R.; Mackenzie, P. I.: cDNA cloning
and

[144381]characterization of the human UDP glucuronosyltrans-
ferase, UGT1A3. Biochem.

[144382]Biophys. Res. Commun. 225: 785-790, 1996.

[144383]25883.

[144384]Monaghan, G.; McLellan, A.; McGeehan, A.; Li Volti, S.;
Mollica,

[144385]F.; Salemi, I.; Din, Z.; Cassidy, A.; Hume, R.; Burchell, B.:
Gilbert's

[144386]syndrome is a contributory factor in prolonged unconju-
gated hyperbilirubinemia

[144387]of the newborn. J. Pediat. 134: 441-446, 1999.

[144388]25884.

[144389]Nagai, F.; Homma, H.; Tanase, H.; Matsui, M.: Studies on
the

[144390]genetic linkage of bilirubin and androsterone UDP-

glucuronyltransferases

[144391]by cross-breeding of two mutant rat strains. *Biochem. J.*
252: 897-900,

[144392]1988.

[144393]25885.

[144394]Peters, W. H. M.; Jansen, P. L. M.; Nauta, H.: The molecular
[144395]weights of UDP-glucuronyltransferase determined with
radiation-inactivation

[144396]analysis: a molecular model of bilirubin UDP-
glucuronyltransferase. *J.*

[144397]*Biol. Chem.* 259: 11701-11705, 1984.

[144398]25886.

[144399]Ritter, J. K.; Chen, F.; Sheen, Y. Y.; Tran, H. M.; Kimura, S.;

[144400]Yeatman, M. T.; Owens, I. S.: A novel complex locus UGT1
encodes

[144401]human bilirubin, phenol, and other UDP-
glucuronosyltransferase isozymes

[144402]with identical carboxyl termini. *J. Biol. Chem.* 267:
3257-3261,

[144403]1992.

[144404]25887.

[144405]Ritter, J. K.; Crawford, J. M.; Owens, I. S.: Cloning of two

[144406]human liver bilirubin UDP-glucuronosyltransferase cDNAs
with expression

[144407]in COS-1 cells. J. Biol. Chem. 266: 1043-1047, 1991.

[144408]25888.

[144409]Ritter, J. K.; Yeatman, M. T.; Ferreira, P.; Owens, I. S.: Identification

[144410]of a genetic alteration in the code for bilirubin UDP-glucuronosyltransferase

[144411]in the UGT1 gene complex of a Crigler-Najjar type I patient. J. Clin.

[144412]Invest. 90: 150-155, 1992.

[144413]25889.

[144414]Ritter, J. K.; Yeatman, M. T.; Kaiser, C.; Gridelli, B.; Owens,

[144415]I. S.: A phenylalanine codon deletion at the UGT1 gene complex locus

[144416]of a Crigler-Najjar type I patient generates a pH-sensitive bilirubin

[144417]UDP-glucuronosyltransferase. J. Biol. Chem. 268: 23573-23579, 1993.

[144418]25890.

[144419]Rosatelli, M. C.; Meloni, A.; Faa, V.; Saba, L.; Crisponi, G.;
[144420]Clemente, M. G.; Meloni, G.; Piga, M. T.; Cao, A.: Molecular
analysis
[144421]of patients of Sardinian descent with Crigler–Najjar syn-
drome type
[144422]I. J. Med. Genet. 34: 122–125, 1997.
[144423]25891.
[144424]Sampietro, M.; Lupica, L.; Perrero, L.; Comino, A.; Martinez
di
[144425]Montemuros, F.; Cappellini, M. D.; Fiorelli, G.: The expres-
sion of
[144426]uridine diphosphate glucuronosyltransferase gene is a
major determinant
[144427]of bilirubin level in heterozygous beta–thalassaemia and
in glucose–6–phosphate
[144428]dehydrogenase deficiency. Brit. J. Haemat. 99: 437–439,
1997.
[144429]25892.
[144430]Sato, H.; Sakai, Y.; Koiwai, O.; Watanabe, T.: Mapping of
the
[144431]mouse bilirubin UDP–glucuronosyltransferase gene
(Gnt–1) to chromosome

[144432]1 by restriction fragment length variations. Biochem.

Genet. 30:

[144433]347–352, 1992.

[144434]25893.

[144435]Seppen, J.; Bosma, P. J.; Goldhoorn, B. G.; Bakker, C. T. M.;

[144436]Chowdhury, J. R.; Chowdhury, N. R.; Jansen, P. L. M.; Oude
Elferink,

[144437]R. P. J.: Discrimination between Crigler–Najjar type I and II
by

[144438]expression of mutant bilirubin uridine diphosphate–
glucuronosyltransferase. J.

[144439]Clin. Invest. 94: 2385–2391, 1994.

[144440]25894.

[144441]Strassburg, C. P.; Oldhafer, K.; Manns, M. P.; Tukey, R. H.:

[144442]Differential expression of the UGT1A locus in human liver,
biliary,

[144443]and gastric tissue: identification of UGT1A7 and UGT1A10
transcripts

[144444]in extrahepatic tissue. Molec. Pharm. 52: 212–220, 1997.

[144445]25895.

- [144446]Tukey, R. H.; Strassburg, C. P.: Human UDP-glucuronosyltransferases: metabolism, expression, and disease. Annu. Rev. Pharm. Toxicol. 40: 581-616, 2000.
- [144449]25896.
- [144450]Ueyama, H.; Koiwai, O.; Soeda, Y.; et al.: Analysis of the promoter of human bilirubin UDP-glucuronosyltransferase gene (UGT1*1) in relevance to Gilbert's syndrome. Hepatol. Res. 9: 152-163, 1997.
- [144453]25897.
- [144454]van Es, H. H.; Bout, A.; Liu, J.; Anderson, L.; Duncan, A. M.; Bosma, P.; Oude Elferink, R.; Jansen, P. L.; Roy Chowdhury, J.; Schurr, E.: Assignment of the human UDP glucuronosyltransferase gene (UGT1A1) to chromosome region 2q37. Cytogenet. Cell Genet. 63: 114-116, 1993.
- [144458]25898.
- [144459]Wooster, R.; Sutherland, L.; Ebner, T.; Clarke, D.; Da Cruz

e

[144460]Silva, O.; Burchell, B.: Cloning and stable expression of a
new member

[144461]of the human liver phenol/bilirubin:UDP-glucuronosyltransferase cDNA

[144462]family. Biochem. J. 278: 465-469, 1991.

[144463]25899.

[144464]Yamamoto, K.; Soeda, Y.; Kamisako, T.; Hosaka, H.;
Fukano, M.;

[144465]Sato, H.; Fujiyama, Y.; Adachi, Y.; Satoh, Y.; Bamba, T.:
Analysis

[144466]of bilirubin uridine 5-prime-diphosphate
(UDP)-glucuronosyltransferase

[144467]gene mutations in seven patients with Crigler-Najjar syndrome type

[144468]II. J. Hum. Genet. 43: 111-114, 1998.

[144469]25900.

[144470]James, P. F.; Grupp, I. L.; Grupp, G.; Woo, A. L.; Askew, G.
R.;

[144471]Croyle, M. L.; Walsh, R. A.; Lingrel, J. B.: Identification of a

[144472]specific role for the Na,K-ATPase alpha-2 isoform as a

regulator of

[144473]calcium in the heart. Molec. Cell 3: 555–563, 1999.

[144474]25901.

[144475]Kent, R. B.; Fallows, D. A.; Geissler, E.; Glaser, T.; Emanuel,

[144476]J. R.; Lalley, P. A.; Levenson, R.; Housman, D. E.: Genes

encoding

[144477]alpha and beta subunits of Na,K-ATPase are located on

three different

[144478]chromosomes in the mouse. Proc. Nat. Acad. Sci. 84:

5369–5373, 1987.

[144479]25902.

[144480]Yang-Feng, T. L.; Schneider, J. W.; Lindgren, V.; Shull, M.

M.;

[144481]Benz, E. J., Jr.; Lingrel, J. B.; Francke, U.: Chromosomal lo-

calization

[144482]of human Na⁺,K⁺-ATPase alpha- and beta-subunit

genes. Genomics 2:

[144483]128–138, 1988.

[144484]25903.

[144485]Kawakami, K.; Nojima, H.; Ohta, T.; Nagano, K.: Molecular

cloning

[144486]and sequence analysis of human Na,K-ATPase beta-subunit. Nucleic

[144487]Acids Res. 14: 2833-2844, 1986.

[144488]25904.

[144489]Lane, L. K.; Shull, M. M.; Whitmer, K. R.; Lingrel, J. B.:
Characterization

[144490]of two genes for the human Na,K-ATPase beta subunit.
Genomics 5:

[144491]445-453, 1989.

[144492]25905.

[144493]Gloor, S.; Antonicek, H.; Sweadner, K. J.; Pagliusi, S.;
Frank,

[144494]R.; Moos, M.; Schachner, M.: The adhesion molecule on
glia (AMOG)

[144495]is a homologue of the beta subunit of the Na,K-ATPase. J.
Cell Biol. 110:

[144496]165-174, 1990.

[144497]25906.

[144498]Hsieh, C.-L.; Cheng-Deutsch, A.; Gloor, S.; Schachner, M.;
Francke,

[144499]U.: Assignment of Amog (adhesion molecule on glia) gene

to mouse

[144500]chromosome 11 near Zfp-3 and Asgr-1,2 and to human chromosome 17. Somat.

[144501]Cell Molec. Genet. 16: 401-405, 1990.

[144502]25907.

[144503]Malo, D.; Schurr, E.; Levenson, R.; Gros, P.: Assignment of Na,K-ATPase

[144504]beta-(2)-subunit gene (Atpb-2) to mouse chromosome 11. Genomics 6:

[144505]697-699, 1990.

[144506]25908.

[144507]Martin-Vasallo, P.; Dackowski, P.; Emanuel, J. R.; Levenson, R.

[144508]: Identification of a putative isoform of the Na,K-ATPase beta subunit:

[144509]primary structure and tissue-specific expression. J. Biol. Chem. 164:

[144510]4613-4618, 1989.

[144511]25909.

[144512]Pagliusi, S.; Antonicek, H.; Gloor, S.; Frank, R.; Moos, M.; Schachner,

[144513]M.: Identification of a cDNA clone specific for the neural adhesion

[144514]molecule AMOG. J. Neurosci. Res. 22: 113–119, 1989.

[144515]25910.

[144516]Katzmarzyk, P. T.; Rankinen, T.; Perusse, L.; Deriaz, O.; Tremblay,

[144517]A.; Borecki, I.; Rao, D. C.; Bouchard, C.: Linkage and association

[144518]of the sodium potassium-adenosine triphosphatase alpha-2 and beta-1

[144519]genes with respiratory quotient and resting metabolic rate in the

[144520]Quebec Family Study. J. Clin. Endocr. Metab. 84: 2093–2097, 1999.

[144521]25911.

[144522]Rampazzo, C.; Gallinaro, L.; Milanesi, E.; Frigimelica, E.; Reichard,

[144523]P.; Bianchi, V.: A deoxyribonucleotidase in mitochondria: involvement

[144524]in regulation of dNTP pools and possible link to genetic disease. Proc.

[144525]Nat. Acad. Sci. 97: 8239–8244, 2000.

[144526]25912.

[144527]Akaba, K.; Kimura, T.; Sasaki, A.; Tanabe, S.; Wakabayashi, T.;

[144528]Hiroi, M.; Yasumura, S.; Maki, K.; Aikawa, S.; Hayasaka, K.:
Neonatal

[144529]hyperbilirubinemia and a common mutation of the bilirubin uridine

[144530]diphosphate-glucuronosyltransferase gene in Japanese. J. Hum. Genet. 44:

[144531]22-25, 1999.

[144532]25913.

[144533]Aono, S.; Yamada, Y.; Keino, H.; Sasaoka, Y.; Nakagawa, T.; Onishi,

[144534]S.; Mimura, S.; Koiwai, O.; Sato, H.: A new type of defect in the

[144535]gene for bilirubin uridine

5-prime-diphosphate-glucuronosyltransferase

[144536]in a patient with Crigler-Najjar syndrome type I. Pediat. Res. 35:

[144537]629-632, 1994.

[144538]25914.

- [144539]Beutler, E.; Gelbart, T.; Demina, A.: Racial variability in the
- [144540]UDP-glucuronosyltransferase 1 (UGT1A1) promoter: a
balanced polymorphism
- [144541]for regulation of bilirubin metabolism? Proc. Nat. Acad.
Sci. 95:
- [144542]8170–8174, 1998.
- [144543]25915.
- [144544]Bosma, P. J.; Goldhoorn, B.; Oude Elferink, R. P. J.; Sinaas-
appel,
- [144545]M.; Oostra, B. A.; Jansen, P. L. M.: A mutation in bilirubin
uridine
- [144546]5-prime-diphosphate-glucuronosyltransferase isoform 1
causing Crigler-Najjar
- [144547]syndrome type II. Gastroenterology 105: 216–220, 1993.
- [144548]25916.
- [144549]Akaba, K.; Kimura, T.; Sasaki, A.; Tanabe, S.; Ikegami, T.;
Hashimoto,
- [144550]M.; Umeda, H.; Yoshida, H.; Umetsu, K.; Chiba, H.; Yuasa,
I.; Hayasaka,
- [144551]K.: Neonatal hyperbilirubinemia and mutation of the
bilirubin uridine

[144552]diphosphate-glucuronosyltransferase gene: a common missense mutation

[144553]among Japanese, Koreans and Chinese. Biochem. Molec. Biol. Int. 46:

[144554]21-26, 1998.

[144555]25917.

[144556]Bosma, P. J.; Roy Chowdhury, J.; Bakker, C.; Gantla, S.; de Boer,

[144557]A.; Oostra, B. A.; Lindhout, D.; Tytgat, G. N. J.; Jansen, P. L. M.;

[144558]Oude Elferink, R. P. J.; Roy Chowdhury, N.: The genetic basis of

[144559]the reduced expression of bilirubin UDP-glucuronosyltransferase 1

[144560]in Gilbert's syndrome. New Eng. J. Med. 333: 1171-1175, 1995.

[144561]25918.

[144562]Bosma, P. J.; Roy Chowdhury, J.; Huang, T.-J.; Lahiri, P.; Oude

[144563]Elferink, R. P. J.; Van Es, H. H. G.; Lederstein, M.; Whittington,

[144564]P. F.; Jansen, P. L. M.; Roy Chowdhury, N.: Mechanisms of

inherited

[144565] deficiencies of multiple UDP-glucuronosyltransferase iso-
forms in two

[144566] patients with Crigler-Najjar syndrome, type I. FASEB J. 6:
2859-2863,

[144567] 1992.

[144568] 25919.

[144569] Thornton, M. A.; Poncz, M.; Korostishevsky, M.; Yakobson,
E.;

[144570] Usher, S.; Seligsohn, U.; Peretz, H.: The human platelet al-
pha-IIb

[144571] gene is not closely linked to its integrin partner beta-3.
Blood 94:

[144572] 2039-2047, 1999.

[144573] 25920.

[144574] von dem Borne, A. E. G.; Decary, F.: Nomenclature of
platelet

[144575] specific antigens. Brit. J. Haemat. 74: 239-240, 1990.

[144576] 25921.

[144577] Xiong, J.-P.; Stehle, T.; Diefenbach, B.; Zhang, R.; Dunker,
R.;

[144578]Scott, D. L.; Joachimiak, A.; Goodman, S. L.; Arnaout, M. A.: Crystal

[144579]structure of the extracellular segment of integrin alpha-V-beta-3. Science 294:

[144580]339-345, 2001.

[144581]25922.

[144582]Xiong, J.-P.; Stehle, T.; Zhang, R.; Joachimiak, A.; Frech, M.;

[144583]Goodman, S. L.; Arnaout, M. A.: Crystal structure of the extracellular

[144584]segment of integrin alpha-V-beta-3 in complex with an Arg-Gly-Asp

[144585]ligand. Science 296: 151-155, 2002.

[144586]25923.

[144587]Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly, J.;

[144588]Bowen-Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of the amphiregulin

[144589]and the platelet-growth factor receptor alpha genes to the proximal

[144590]long arm of chromosome 4. (Abstract) Cytogenet. Cell Genet. 51:

[144591]990, 1989.

[144592]25924.

[144593]Gronwald, R. G. K.; Adler, D. A.; Kelly, J. D.; Disteché, C.
M.;

[144594]Bowen-Pope, D. F.: The human PDGF receptor alpha-
subunit gene maps

[144595]to chromosome 4 in close proximity to c-kit. Hum. Genet.
85: 383-385,

[144596]1990.

[144597]25925.

[144598]Hol, F. A.; Geurds, M. P. A.; Chatkupt, S.; Shugart, Y. Y.;
Balling,

[144599]R.; Schrandt-Stumpel, C. T. R. M.; Johnson, W. G.; Hamel,
B. C. J.;

[144600]Mariman, E. C. M.: PAX genes and human neural tube de-
fects: an amino

[144601]acid substitution in PAX1 in a patient with spina bifida. J.
Med.

[144602]Genet. 8: 655-660, 1996.

[144603]25926.

[144604]Hsieh, C.-L.; Navankasattusas, S.; Escobedo, J. A.;

Williams, L.

[144605]T.; Francke, U.: Chromosomal localization of the gene for
AA-type

[144606]platelet-derived growth factor receptor (PDGFRA) in hu-
mans and mice. Cytogenet.

[144607]Cell Genet. 56: 160-163, 1991.

[144608]25927.

[144609]Ikuno, Y.; Kazlauskas, A.: TGF-beta-1-dependent con-
traction of

[144610]fibroblasts is mediated by th PDGF-alpha receptor. Invest.
Ophthalm.

[144611]Vis. Sci. 43: 41-46, 2002.

[144612]25928.

[144613]Joosten, P. H. L. J.; Hol, F. A.; van Beersum, S. E. C.; Peters,

[144614]H.; Hamel, B. C. J.; Afink, G. B.; van Zoelen, E. J. J.; Mari-
man,

[144615]E. C. M.: Altered regulation of platelet-derived growth fac-
tor receptor-alpha

[144616]gene-transcription in vitro by spina bifida-associated mu-
tant Pax1

[144617]proteins. Proc. Nat. Acad. Sci. 95: 14459-14463, 1998.

[144618]25929.

[144619]Sanke, T.; Bell, G. I.; Sample, C.; Rubenstein, A. H.; Steiner,

[144620]D. F.: An islet amyloid peptide is derived from an
89-amino acid

[144621]precursor by proteolytic processing. J. Biol. Chem. 263:
17243-17246,

[144622]1988.

[144623]25930.

[144624]Verchere, C. B.; D'Alessio, D. A.; Palmiter, R. D.; Weir, G.
C.;

[144625]Bonner-Weir, S.; Baskin, D. G.; Kahn, S. E.: Islet amyloid
formation

[144626]associated with hyperglycemia in transgenic mice with
pancreatic beta

[144627]cell expression of human islet amyloid polypeptide. Proc.
Nat. Acad.

[144628]Sci. 93: 3492-3496, 1996.

[144629]25931.

[144630]Westermarck, P.; Engstrom, U.; Johnson, K. H.; Westermarck,
G. T.;

[144631]Betsholtz, C.: Islet amyloid polypeptide: pinpointing amino

acid

[144632]residues linked to amyloid fibril formation. Proc. Nat.

Acad. Sci. 87:

[144633]5036–5040, 1990.

[144634]25932.

[144635]Hockaday, T. D. R.: Hypogonadism and life-long anosmia.

Postgrad.

[144636]Med. J. 42: 572–574, 1966.

[144637]25933.

[144638]Oliveira, L. M. B.; Seminara, S. B.; Beranova, M.; Hayes, F.

J.;

[144639]Valkenburgh, S. B.; Schipani, E.; Costa, E. M. F.; Latronico,

A. C.;

[144640]Crowley, W. F., Jr.; Vallejo, M.: The importance of autosomal genes

[144641]in Kallmann syndrome: genotype–phenotype correlations and neuroendocrine

[144642]characteristics. J. Clin. Endocr. Metab. 86: 1532–1538, 2001.

[144643]25934.

[144644]Bignon, C.; Roux–Dosseto, M.; Zeigler, M. E.; Mattei,

M.-G.; Lissitzky,

[144645]J.-C.; Wicha, M. S.; Martin, P.-M.: Genomic analysis of the
67-kDa

[144646]laminin receptor in normal and pathological tissues: circumstantial

[144647]evidence for retroposon features. Genomics 10: 481-485,
1991.

[144648]25935.

[144649]Gehlsen, K. R.; Dillner, L.; Engvall, E.; Ruoslahti, E.: The
human

[144650]laminin receptor is a member of the integrin family of cell
adhesion

[144651]receptors. Science 241: 1228-1229, 1988. Note: Erratum:
Science

[144652]245:342-343, 1989.

[144653]25936.

[144654]Jackers, P.; Minoletti, F.; Belotti, D.; Clausse, N.; Sozzi, G.;

[144655]Sobel, M. E.; Castronovo, V.: Isolation from a multigene
family of

[144656]the active human gene of the metastasis-associated multifunctional

[144657]protein 37LRP/p40 at chromosome 3p21.3. Oncogene 13:

495–503, 1996.

[144658]25937.

[144659]Lafreniere, R. G.; Brown, C. J.; Rider, S.; Chelly, J.; Taillon–
Miller,

[144660]P.; Chinault, A. C.; Monaco, A. P.; Willard, H. F.: 2.6 Mb
YAC contig

[144661]of the human X inactivation center region in Xq13: physical linkage

[144662]of the RPS4X, PHKA1, XIST and DXS128E genes. Hum.
Molec. Genet. 2:

[144663]1105–1115, 1993.

[144664]25938.

[144665]Montuori, N.; Muller, F.; De Riu, S.; Fenzi, G.; Sobel, M. E.;

[144666]Rossi, G.; Vitale, M.: Laminin receptors in differentiated
thyroid

[144667]tumors: restricted expression of the 67–kilodalton laminin
receptor

[144668]in follicular carcinoma cells. J. Clin. Endocr. Metab. 84:
2086–2092,

[144669]1999.

[144670]25939.

[144671]Sato, K.; Narumi, K.; Sakai, T.; Abe, T.; Kikuchi, T.; Matsushima,
[144672]K.; Sindoh, S.; Motomiya, M.: Cloning of 67-kDa laminin
receptor
[144673]cDNA and gene expression in normal and malignant cell
lines of the
[144674]human lung. Cancer Lett. 62: 199–203, 1992.
[144675]25940.
[144676]Tohgo, A.; Takasawa, S.; Munakata, H.; Yonekura, H.;
Hayashi, N.;
[144677]Okamoto, H.: Structural determination and characterization of a 40
[144678]kDa protein isolated from rat 40 S ribosomal subunit. FEBS
Lett. 340:
[144679]133–138, 1994.
[144680]25941.
[144681]Yow, H. K.; Wong, J. M.; Chen, H. S.; Lee, C. G.; Davis, S.;
Steele,
[144682]G. D., Jr.; Chen, L. B.: Increased mRNA expression of a
laminin-binding
[144683]protein in human colon carcinoma: complete sequence of
a full-length

- [144684]cDNA encoding the protein. Proc. Nat. Acad. Sci. 85:
6394–6398,
- [144685]1988. Note: Erratum: Proc. Nat. Acad. Sci. 86: 7032 only,
1989.
- [144686]25942.
- [144687]Oklu, R.; Hesketh, R.: The latent transforming growth fac-
tor beta
- [144688]binding protein (LTBP) family. Biochem. J. 352: 601–610,
2000.
- [144689]25943.
- [144690]Lloyd, D. J.; Trembath, R. C.; Shackleton, S.: A novel inter-
action
- [144691]between lamin A and SREBP1: implications for partial
lipodystrophy
- [144692]and other laminopathies. Hum. Molec. Genet. 11:
769–777, 2002.
- [144693]25944.
- [144694]Wydner, K. L.; McNeil, J. A.; Lin, F.; Worman, H. J.;
Lawrence,
- [144695]J. B.: Chromosomal assignment of human nuclear enve-
lope protein genes
- [144696]LMNA, LMNB1, and LBR by fluorescence in situ hybridiza-

tion. Genomics 32:

[144697]474–478, 1996.

[144698]25945.

[144699]Garbarz, M.; Devaux, I.; Bournier, O.; Grandchamp, B.;
Dhermy,

[144700]D.: Protein 4.1 Lille, a novel mutation in the downstream
initiation

[144701]codon of protein 4.1 gene associated with heterozygous
4,1(–) hereditary

[144702]elliptocytosis. Hum. Mutat. 5: 339–340, 1995.

[144703]25946.

[144704]Garbarz, M.; Dhermy, D.; Lecomte, M. C.; Feo, C.; Chave-
roche,

[144705]I.; Galand, C.; Bournier, O.; Bertrand, O.; Boivin, P.: A vari-
ant

[144706]of erythrocyte membrane skeletal protein band 4.1 asso-
ciated with

[144707]hereditary elliptocytosis. Blood 64: 1006–1015, 1984.

[144708]25947.

[144709]Geerdink, R. A.; Nijenhuis, L. E.; Huizinga, J.: Hereditary
elliptocytosis:

[144710]linkage data in man. Ann. Hum. Genet. 30: 363–378,
1967.

[144711]25948.

[144712]Jensson, O.; Jonasson, T.; Olafsson, O.: Hereditary ellipto-
cytosis

[144713]in Iceland. Brit. J. Haemat. 13: 844–854, 1967.

[144714]25949.

[144715]Kan, Y.-W.: Personal Communication. San Francisco, Calif.

[144716]2/28/1986.

[144717]25950.

[144718]Kuroda, S.; Takeuchi, T.; Nagamori, H.: Data on the linkage between

[144719]elliptocytosis and Rh blood type. Jpn. J. Hum. Genet. 5:
112-118,

[144720]1960.

[144721]25951.

[144722]Lambert, S.; Conboy, J.; Zail, S.: A molecular study of heterozygous

[144723]protein 4.1 deficiency in hereditary elliptocytosis. Blood
72: 1926-1929,

[144724]1988.

[144725]25952.

[144726]Lambert, S.; Zail, S.: Partial deficiency of protein 4.1 in
hereditary

[144727]elliptocytosis. Am. J. Hemat. 26: 263-272, 1987.

[144728]25953.

[144729]Lipton, E. L.: Elliptocytosis with hemolytic anemia: the effects
[144730]of splenectomy. Pediatrics 15: 67-82, 1955.
[144731]25954.

[144732]Lux, S. E.; Wolfe, L. C.: Inherited disorders of the red cell
[144733]membrane skeleton. Pediat. Clin. N. Am. 27: 463-486,
1980.
[144734]25955.

[144735]Marchesi, S. L.; Conboy, J.; Agre, P.; Letsinger, J. T.;
Marchesi,
[144736]V. T.; Speicher, D. W.; Mohandas, N.: Molecular analysis of
insertion/deletion
[144737]mutations in protein 4.1 in elliptocytosis. I. Biochemical
identification
[144738]of rearrangements in the spectrin/actin binding domain
and functional
[144739]characterizations. J. Clin. Invest. 86: 516-523, 1990.
[144740]25956.

[144741]McGuire, M.; Agre, P.: Three distinct variants of protein
4.1
[144742]in Caucasian hereditary elliptocytosis. (Abstract) Clin. Res.

35:

[144743]428A, 1987.

[144744]25957.

[144745]McGuire, M.; Smith, B. L.; Agre, P.: Distinct variants of erythrocyte

[144746]protein 4.1 inherited in linkage with elliptocytosis and Rh type in

[144747]three white families. Blood 72: 287–293, 1988.

[144748]25958.

[144749]Morle, L.; Garbarz, M.; Alloisio, N.; Girot, R.; Chaverroche, I.;

[144750]Boivin, P.; Delaunay, J.: The characterization of protein 4.1 Presles,

[144751]a shortened variant of RBC membrane protein 4.1. Blood 65: 1511–1517,

[144752]1985.

[144753]25959.

[144754]Morle, L.; Pothier, B.; Alloisio, N.; Ducluzeau, M.–T.; Marques,

[144755]S.; Olim, G.; Martins e Silva, J.; Feo, C.; Garbarz, M.; Chaverroche,

- [144756]I.; Boivin, P.; Delaunay, J.: Red cell membrane alteration involving
[144757]protein 4.1 and protein 3 in a case of recessively inherited
haemolytic
[144758]anemia. Europ. J. Haemat. 38: 447–455, 1987.
[144759]25960.
- [144760]Morton, N. E.: The detection and estimation of linkage between
[144761]the genes for elliptocytosis and the Rh blood type. Am. J.
Hum. Genet. 8:
[144762]80–96, 1956.
[144763]25961.
- [144764]Nielsen, J. A.; Strunk, K. W.: Homozygous hereditary elliptocytosis
[144765]as the cause of haemolytic anemia in infancy. Scand. J.
Haemat. 5:
[144766]486–496, 1968.
[144767]25962.
- [144768]Parra, M.; Gascard, P.; Walensky, L. D.; Snyder, S. H.; Mohandas,
[144769]N.; Conboy, J. G.: Cloning and characterization of 4.1G

(EPB41L2),

[144770]a new member of the skeletal protein 4.1 (EPB41) gene family. Genomics 49:

[144771]298–306, 1998.

[144772]25963.

[144773]Peters, J. C.; Rowland, M.; Israels, L. G.; Zipursky, A.: Erythrocyte

[144774]sodium transport in hereditary elliptocytosis. Canad. J. Physiol.

[144775]Pharm. 44: 817–827, 1966.

[144776]25964.

[144777]Roberts, J. A. F.: Genetic linkage in man, with particular reference

[144778]to the usefulness of very small bodies of data. Quart. J. Med. 14:

[144779]27–33, 1945.

[144780]25965.

[144781]Shi, Z.-T.; Afzal, V.; Coller, B.; Patel, D.; Chasis, J. A.; Parra,

[144782]M.; Lee, G.; Paszty, C.; Stevens, M.; Walensky, L.; Peters, L. L.;

[144783]Mohandas, N.; Rubin, E.; Conboy, J. G.: Protein

4.1R-deficient mice

[144784]are viable but have erythroid membrane skeleton abnormalities. J.

[144785]Clin. Invest. 103: 331-340, 1999.

[144786]25966.

[144787]Takakuwa, Y.; Tchernia, G.; Rossi, M.; Benabadji, M.; Mohandas,

[144788]N.: Restoration of normal membrane stability to unstable protein

[144789]4.1-deficient erythrocyte membranes by incorporation of purified protein

[144790]4.1. J. Clin. Invest. 78: 80-85, 1986.

[144791]25967.

[144792]Tang, C.-J. C.; Tang, T. K.: Rapid localization of membrane skeletal

[144793]protein 4.1 (EL1) to human chromosome 1p33-p34.2 by nonradioactive

[144794]in situ hybridization. Cytogenet. Cell Genet. 57: 119, 1991.

[144795]25968.

[144796]Tang, T. K.; Leto, T. L.; Correas, I.; Alonso, M. A.; Marchesi,

[144797]V. T.; Benz, E. J., Jr.: Selective expression of an erythroid-specific

[144798]isoform of protein 4.1. Proc. Nat. Acad. Sci. 85: 3713–3717, 1988.

[144799]25969.

[144800]Tchernia, G.; Mohandas, N.; Shohet, S. B.: Deficiency of skeletal

[144801]membrane protein band 4.1 in homozygous hereditary elliptocytosis:

[144802]implications for erythrocyte membrane stability. J. Clin. Invest. 68:

[144803]454–460, 1981.

[144804]25970.

[144805]Tomaselli, M. B.; John, K. M.; Lux, S. E.: Elliptical erythrocyte

[144806]membrane skeletons and heat-sensitive spectrin in hereditary elliptocytosis. Proc.

[144807]Nat. Acad. Sci. 78: 1911–1915, 1981.

[144808]25971.

- [144809] Brands, J. H. G. M.; Maassen, J. A.; Van Hemert, F. J.;
Amons,
[144810] R.; Moller, W.: The primary structure of the alpha subunit
of human
[144811] elongation factor 1: structural aspects of guanine-nu-
cleotide-binding
[144812] sites. Europ. J. Biochem. 155: 167-171, 1986.
[144813] 25972.
[144814] Ditzel, H. J.; Masaki, Y.; Nielsen, H.; Farnaes, L.; Burton, D.
[144815] R.: Cloning and expression of a novel human antibody-
-antigen pair
[144816] associated with Felty's syndrome. Proc. Nat. Acad. Sci. 97:
9234-9239,
[144817] 2000.
[144818] 25973.
[144819] Lund, A.; Knudsen, S. M.; Vissing, H.; Clark, B.; Tom-
merup, N.
[144820]: Assignment of human elongation factor 1-alpha genes:
EEF1A maps
[144821] to chromosome 6q14 and EEF1A2 to 20q13.3. Genomics
36: 359-361,

[144822]1996.

[144823]25974.

[144824]Opdenakker, G.; Cabeza-Arvelaiz, Y.; Fiten, P.; Dijkmans, R.; Van

[144825]Damme, J.; Volckaert, G.; Billiau, A.; Van Elsen, A.; Van der Schueren,

[144826]B.; Van den Berghe, H.; Cassiman, J.-J.: Human elongation factor

[144827]1-alpha: a polymorphic and conserved multigene family with multiple

[144828]chromosomal localizations. Hum. Genet. 75: 339-344, 1987.

[144829]25975.

[144830]Cohen, A. J.; Li, F. P.; Berg, S.; Marchetto, D. J.; Tsai, S.;

[144831]Jacobs, S. C.; Brown, R. S.: Hereditary renal-cell carcinoma associated

[144832]with chromosomal translocation. New Eng. J. Med. 301: 592-595, 1979.

[144833]25976.

[144834]Gemmell, R. M.; West, J. D.; Boldog, F.; Tanaka, N.; Robinson,

[144835]L. J.; Smith, D. I.; Li, F.; Drabkin, H. A.: The hereditary renal

[144836]cell carcinoma 3;8 translocation fuses FHIT to a patched-related gene,

[144837]TRC8. Proc. Nat. Acad. Sci. 95: 9572–9577, 1998.

[144838]25977.

[144839]Nakano, A.; Pulkkinen, L.; Murrell, D.; Rico, J.; Lucky, A. W.;

[144840]Garzon, M.; Stevens, C. A.; Robertson, S.; Pfendner, E.; Uitto, J.

[144841]: Epidermolysis bullosa with congenital pyloric atresia: novel mutations

[144842]in the beta-4 integrin gene (ITGB4) and genotype/phenotype correlations. Pediat.

[144843]Res. 49: 618–626, 2001.

[144844]25978.

[144845]Pulkkinen, L.; Bruckner-Tuderman, L.; August, C.; Uitto, J.: Compound

[144846]heterozygosity for missense mutation (L156P) and non-sense (R554X)

[144847]mutations in the beta-4 integrin gene (ITGB4) underlies mild, nonlethal

[144848]phenotype of epidermolysis bullosa with pyloric atresia.

Am. J. Path. 152:

[144849]935–941, 1998.

[144850]25979.

[144851]Pulkkinen, L.; Rouan, F.; Bruckner–Tuderman, L.; Wallerstein, R.;

[144852]Garzon, M.; Brown, T.; Smith, L.; Carter, W.; Uitto, J.: Novel
ITGB4

[144853]mutations in lethal and nonlethal variants of epidermolysis bullosa

[144854]with pyloric atresia: missense versus nonsense. Am. J.
Hum. Genet. 63:

[144855]1376–1387, 1998.

[144856]25980.

[144857]Shaw, L. M.; Rabinovitz, I., Wang, H. H.–F.; Toker, A., Mercurio,

[144858]A. M.: Activation of phosphoinositide 3–OH kinase by the
alpha–6/beta–4

[144859]integrin promotes carcinoma invasion. Cell 91: 949–960,
1997.

[144860]25981.

[144861]Suzuki, S.; Naitoh, Y.: Amino acid sequence of a novel integrin

[144862]beta-4 subunit and primary expression of the mRNA in epithelial cells. EMBO

[144863]J. 9: 757-763, 1990.

[144864]25982.

[144865]Vidal, F.; Aberdam, D.; Miquel, C.; Christiano, A. M.; Pulkkinen,

[144866]L.; Uitto, J.; Ortonne, J.-P.; Meneguzzi, G.: Integrin beta-4 mutations

[144867]associated with junctional epidermolysis bullosa with pyloric atresia. Nature

[144868]Genet. 10: 229-234, 1995.

[144869]25983.

[144870]Chany, C.; Vignal, M.; Couillin, P.; Van Cong, N.; Boue, J.; Boue,

[144871]A.: Chromosomal localization of human genes governing the interferon-induced

[144872]antiviral state. Proc. Nat. Acad. Sci. 72: 3129-3133, 1975.

[144873]25984.

[144874]Aplin, H. M.; Hirst, K. L.; Crosby, A. H.; Dixon, M. J.: Map-

ping

[144875]of the human dentin matrix acidic phosphoprotein gene
(DMP1) to the

[144876]dentinogenesis imperfecta type II critical region at chro-
mosome 4q21. Genomics 30:

[144877]347-349, 1995.

[144878]25985.

[144879]Fisher, L. W.; McBride, O. W.; Termine, J. D.; Young, M. F.:
Human

[144880]bone sialoprotein: deduced protein sequence and chro-
mosomal localization.

[144881]J. Biol. Chem. 265: 2347-2351, 1990.

[144882]25986.

[144883]Goold, R. D.; diSibio, G. L.; Xu, H.; Lang, D. B.; Dadgar, J.;

[144884]Magrane, G. G.; Dugaiczky, A.; Smith, K. A.; Cox, D. R.;
Masters,

[144885]S. B.; Myers, R. M.: The development of sequence-tagged
sites for

[144886]human chromosome 4. Hum. Molec. Genet. 2:
1271-1288, 1993.

[144887]25987.

[144888]Kerr, J. M.; Fisher, L. W.; Termine, J. D.; Wang, M. G.;
McBride,

[144889]O. W.; Young, M. F.: The human bone sialoprotein gene
(IBSP): genomic

[144890]localization and characterization. Genomics 17: 408–415,
1993.

[144891]25988.

[144892]Kim, R. H.; Shapiro, H. S.; Li, J. J.; Wrana, J. L.; Sodek, J.

[144893]: Characterization of the human bone sialoprotein (BSP)
gene and its

[144894]promoter sequence. Matrix Biol. 14: 31–40, 1994.

[144895]25989.

[144896]Young, M. F.; Ibaraki, K.; Kerr, J. M.; Lyu, M. S.; Kozak, C.
A.

[144897]: Murine bone sialoprotein (BSP): cDNA cloning, mRNA ex-
pression, and

[144898]genetic mapping. Mammalian Genome 5: 108–111, 1994.

[144899]25990.

[144900]Ho, A. S. Y.; Liu, Y.; Khan, T. A.; Hsu, D.-H.; Bazan, J. F.;
Moore,

[144901]K. W.: A receptor for interleukin 10 is related to interferon

receptors. Proc.

[144902]Nat. Acad. Sci. 90: 11267–11271, 1993.

[144903]25991.

[144904]Liu, Y.; Wei, S. H.–Y.; Ho, A. S.–Y.; de Waal Malefyt, R.;

Moore,

[144905]K. W.: Expression cloning and characterization of a human

IL-10 receptor. J.

[144906]Immun. 152: 1821–1829, 1994.

[144907]25992.

[144908]Barker, D. F.; Hostikka, S. L.; Zhou, J.; Chow, L. T.;

Oliphant,

[144909]A. R.; Gerken, S. C.; Gregory, M. C.; Skolnick, M. H.; Atkin,

C. L.;

[144910]Tryggvason, K.: Identification of mutations in the COL4A5
collagen

[144911]gene in Alport syndrome. Science 248: 1224–1227, 1990.

[144912]25993.

[144913]Draptchinskaia, N.; Gustavsson, P.; Andersson, B.; Petters-
son,

[144914]M.; Willig, T.–N.; Dianzani, I.; Ball, S.; Tchernia, G.; Klar, J.;

[144915]Matsson, H.; Tentler, D.; Mohandas, N.; Carlsson, B.; Dahl,

N.: The

[144916] gene encoding ribosomal protein S19 is mutated in Diamond-Blackfan

[144917] anaemia. *Nature Genet.* 21: 169–175, 1999.

[144918] 25994.

[144919] Tanzi, R. E.; St. George-Hyslop, P. H.; Gusella, J. F.: Molecular

[144920] genetics of Alzheimer disease amyloid. *J. Biol. Chem.* 266: 20579–20582,

[144921] 1991.

[144922] 25995.

[144923] Thinakaran, G.; Borchelt, D. R.; Lee, M. K.; Slunt, H. H.; Spitzer,

[144924] L.; Kim, G.; Ratovitsky, T.; Davenport, F.; Nordstedt, C.; Seeger,

[144925] M.; Hardy, J.; Levey, A. I.; Gandy, S. E.; Jenkins, N. A.; Copeland,

[144926] N. G.; Price, D. L.; Sisodia, S. S.: Endoproteolysis of presenilin

[144927] 1 and accumulation of processed derivatives in vivo. *Neuron* 17:

[144928] 181–190, 1996.

[144929]25996.

[144930]Struhl, G.; Adachi, A.: Requirements for Presenilin-dependent

[144931]cleavage of Notch and other transmembrane proteins.
Molec. Cell 6:

[144932]625–636, 2000.

[144933]25997.

[144934]Struhl, G.; Greenwald, I.: Presenilin is required for activity

[144935]and nuclear access of Notch in Drosophila. Nature 398:
522–525,

[144936]1999.

[144937]25998.

[144938]Takahashi, Y.; Koizumi, K.; Takagi, A.; Kitajima, S.; Inoue,
T.;

[144939]Koseki, H.; Saga, Y.: Mesp2 initiates somite segmentation
through

[144940]the Notch signalling pathway. Nature Genet. 25: 390–396,
2000.

[144941]25999.

[144942]Takashima, A.; Sato, M.; Mercken, M.; Tanaka, S.; Kondo,
S.; Honda,

- [144943]T.; Sato, K.; Murayama, M.; Noguchi, K.; Nakazato, Y.;
Takahashi,
- [144944]H.: Localization of Alzheimer-associated presenilin 1 in
transfected
- [144945]COS-7 cells. *Biochem. Biophys. Res. Commun.* 227:
423-426, 1996.
- [144946]26000.
- [144947]Trower, M. K.; Orton, S. M.; Purvis, I. J.; Sanseau, P.; Riley,
[144948]J.; Christodoulou, C.; Burt, D.; See, C. G.; Elgar, G.; Sher-
rington,
- [144949]R.; Rogaev, E. I.; St. George-Hyslop, P.; Brenner, S.; Dykes,
C. W.
- [144950]: Conservation of synteny between the genome of the
pufferfish (*Fugu*
- [144951]*rubripes*) and the region on human chromosome 14
(14q24.3) associated
- [144952]with familial Alzheimer disease (AD3 locus). *Proc. Nat.*
Acad. Sci. 93:
- [144953]1366-1369, 1996.
- [144954]26001.
- [144955]Tysoe, C.; Whittaker, J.; Xuereb, J.; Cairns, N. J.; Cruts, M.;

- [144956] Van Broeckhoven, C.; Wilcock, G.; Rubinsztein, D. C.: A
presenilin-1
[144957] truncating mutation is present in two cases with autopsy-
confirmed
[144958] early-onset Alzheimer disease. Am. J. Hum. Genet. 62:
70-76, 1998.
[144959] 26002.
- [144960] Van Broeckhoven, C.; Backhovens, H.; Cruts, M.; De Win-
ter, G.;
[144961] Bruylant, M.; Cras, P.; Martin, J.-J.: Mapping of a gene
predisposing
[144962] to early-onset Alzheimer's disease to chromosome
14q24.3. Nature
[144963] Genet. 2: 335-339, 1992.
[144964] 26003.
- [144965] Verkkoniemi, A.; Somer, M.; Rinne, J. O.; Myllykangas, L.;
Crook,
[144966] R.; Hardy, J.; Viitanen, M.; Kalimo, H.; Haltia, M.: Variant
Alzheimer's
[144967] disease with spastic paraparesis: clinical characterization.
Neurology 54:
[144968] 1103-1109, 2000.

[144969]26004.

[144970]Wasco, W.; Pettingell, W. P.; Jondro, P. D.; Schmidt, S. D.;
Gurubhagaratula,

[144971]S.; Rodes, L.; DiBlasi, T.; Romano, D. M.; Guenette, S. Y.;
Kovacs,

[144972]D. M.; Growdon, J. H.; Tanzi, R. E.: Familial Alzheimer's
chromosome

[144973]14 mutations. (Letter) Nature Med. 1: 848, 1995.

[144974]26005.

[144975]Wilson, C. A.; Doms, R. W.; Zheng, H.; Lee, V. M.-Y.: Pre-
senilins

[144976]are not required for A-beta-42 production in the early se-
cretory pathway. Nature

[144977]Neurosci. 5: 849-855, 2002.

[144978]26006.

[144979]Wittenburg, N.; Eimer, S.; Lakowski, B.; Rohrig, S.;
Rudolph,

[144980]C.; Baumeister, R.: Presenilin is required for proper mor-
phology

[144981]and function of neurons in C. elegans. Nature 406:
306-309, 2000.

[144982]26007.

[144983]Wolfe, M. S.; Xia, W.; Ostaszewski, B. L.; Diehl, T. S.; Kimberly,

[144984]W. T.; Selkoe, D. J.: Two transmembrane aspartates in presenilin-1

[144985]required for presenilin endoproteolysis and gamma-secretase activity. Nature 398:

[144986]513-517, 1999.

[144987]26008.

[144988]Xia, X.; Qian, S.; Soriano, S.; Wu, Y.; Fletcher, A. M.; Wang,

[144989]X.-J.; Koo, E. H.; Wu, X.; Zheng, H.: Loss of presenilin 1 is associated

[144990]with enhanced beta-catenin signaling and skin tumorigenesis. Proc.

[144991]Nat. Acad. Sci. 98: 10863-10868, 2001.

[144992]26009.

[144993]Ye, Y.; Lukinova, N.; Fortini, M. E.: Neurogenic phenotypes and

[144994]altered Notch processing in Drosophila presenilin mutants. Nature 398:

[144995]525-529, 1999.

[144996]26010.

[144997]Yu, H.; Saura, C. A.; Choi, S.-Y.; Sun, L. D.; Yang, X.; Handler,

[144998]M.; Kawarabayashi, T.; Younkin, L.; Fedeles, B.; Wilson, M. A.; Younkin,

[144999]S.; Kandel, E. R.; Kirkwood, A.; Shen, J.: APP processing and synaptic

[145000]plasticity in presenilin-1 conditional knockout mice. *Neuron* 31:

[145001]713-726, 2001.

[145002]26011.

[145003]Zhang, Z.; Hartmann, H.; Minh Do, V.; Abramowski, D.; Sturchler-Pierrat,

[145004]C.; Staufenbiel, M.; Sommer, B.; van de Weterling, M.; Clevers, H.;

[145005]Saftig, P.; De Strooper, B.; He, X.; Yanker, B. A.: Destabilization

[145006]of beta-catenin by mutations in presenilin-1 potentiates neuronal

[145007]apoptosis. *Nature* 395: 698-702, 1998.

[145008]26012.

[145009]Karrman, C.; Backman, B.; Dixon, M.; Holmgren, G.; Forsman, K.

[145010] Mapping of the locus for autosomal dominant amelogenesis imperfecta

[145011](AIH2) to a 4-Mb YAC contig on chromosome 4q11-q21. Genomics 39:

[145012]164-170, 1997.

[145013]26013.

[145014]Rajpar, M. H.; Harley, K.; Laing, C.; Davies, R. M.; Dixon, M.

[145015]J.: Mutation of the gene encoding the enamel-specific protein, enamelin,

[145016]causes autosomal-dominant amelogenesis imperfecta. Hum. Molec. Genet. 10:

[145017]1673-1677, 2001.

[145018]26014.

[145019]Bleyl, S.; Nelson, L.; Odelberg, S. J.; Ruttenberg, H. D.; Otterud,

[145020]B.; Leppert, M.; Ward, K.: A gene for familial total anomalous pulmonary

[145021]venous return maps to chromosome 4p13-q12. Am. J. Hum. Genet. 56:

[145022]408–415, 1995.

[145023]26015.

[145024]Blixt, A.; Mahlapuu, M.; Aitola, M.; Pelto-Huikko, M.; Enerback,

[145025]S.; Carlsson, P.: A forkhead gene, FoxE3, is essential for lens epithelial

[145026]proliferation and closure of the lens vesicle. Genes Dev. 14: 245–254,

[145027]2000.

[145028]26016.

[145029]Semina, E. V.; Brownell, I.; Mintz-Hittner, H. A.; Murray, J. C.;

[145030]Jamrich, M.: Mutations in the human forkhead transcription factor

[145031]FOXE3 associated with anterior segment ocular dysgenesis and cataracts. Hum.

[145032]Molec. Genet. 10: 231–236, 2001.

[145033]26017.

[145034]Siegelmann-Danieli, N.; Buetow, K. H.: Constitutional genetic

[145035]variation at the human aromatase gene (Cyp19) and

breast cancer risk. Brit.

[145036]J. Cancer 79: 456–463, 1999.

[145037]26018.

[145038]Simpson, E. R.; Michael, M. D.; Agarwal, V. R.; Hinshelwood, M.

[145039]M.; Bulun, S. E.; Zhao, Y.: Expression of the CYP19 (aromatase) gene:

[145040]an unusual case of alternative promoter usage. FASEB J. 11: 29–36,

[145041]1997.

[145042]26019.

[145043]Sparkes, R. S.; Mohandas, T.; Chen, S.; Besman, M. J.; Zollman,

[145044]S.; Shively, J. E.: Assignment of the aromatase gene to human chromosome

[145045]15q21. (Abstract) Cytogenet. Cell Genet. 46: 696–697, 1987.

[145046]26020.

[145047]Toda, K.; Merashima, M.; Kawamoto, T.; Sumimoto, H.; Yokoyama,

[145048]Y.; Kuribayashi, I.; Mitsuuchi, Y.; Maeda, T.; Yamamoto, Y.;

Sagara,

[145049]Y.; Ikeda, H.; Shizuta, Y.: Structural and functional characterization

[145050]of human aromatase P-450 gene. *Europ. J. Biochem.* 193: 559-565,

[145051]1990.

[145052]26021.

[145053]Wang, Z. J.; Jeffs, B.; Ito, M.; Achermann, J. C.; Yu, R. N.;

[145054]Hales, D. B.; Jameson, J. L.: Aromatase (Cyp19) expression is up-regulated

[145055]by targeted disruption of Dax1. *Proc. Nat. Acad. Sci.* 98: 7988-7993,

[145056]2001.

[145057]26022.

[145058]Whitlock, J. P., Jr.: The regulation of cytochrome P-450 gene

[145059]expression. *Annu. Rev. Pharm. Toxicol.* 26: 333-369, 1986.

[145060]26023.

[145061]Zhou, D.; Pompon, D.; Chen, S.: Structure-function studies of

[145062]human aromatase by site-directed mutagenesis: kinetic properties of

[145063]mutants pro308-to-phe, tyr361-to-phe, tyr361-to-leu, and phe406-to-arg. Proc.

[145064]Nat. Acad. Sci. 88: 410-414, 1991.

[145065]26024.

[145066]Luttrell, L. M.; Ferguson, S. S. G.; Daaka, Y.; Miller, W. E.;

[145067]Maudsley, S.; Della Rocca, G. J.; Lin, F.-T.; Kawakatsu, H.; Owada,

[145068]K.; Luttrell, D. K.; Caron, M. G.; Lefkowitz, R. J.: Beta-arrestin-dependent

[145069]formation of beta-2 adrenergic receptor-Src protein kinase complexes. Science 283:

[145070]655-661, 1999.

[145071]26025.

[145072]McDonald, P. H.; Chow, C.-W.; Miller, W. E.; Laporte, S. A.; Field,

[145073]M. E.; Lin, F.-T.; Davis, R. J.; Lefkowitz, R. J.: Beta-arrestin

[145074]2: a receptor-regulated MAPK scaffold for the activation of JNK3. Science 290:

[145075]1574-1577, 2000.

[145076]26026.

[145077]Ferns, G. A. A.; Stocks, J.; Ritchie, C.; Galton, D. J.: Genetic

[145078]polymorphisms of apolipoprotein C-III and insulin in survivors of

[145079]myocardial infarction. Lancet II: 300-303, 1985.

[145080]26027.

[145081]Holt, M.; Hogan, P. F.; Nurse, G. T.: The ovalocytosis polymorphism

[145082]on the western border of Papua New Guinea. Hum. Biol. 53: 23-34,

[145083]1981.

[145084]26028.

[145085]Hsu, L.; Morrison, M.: A new variant of the anion transport protein

[145086]in human erythrocytes. Biochemistry 24: 3086-3090, 1985.

[145087]26029.

[145088]Ideguchi, H.; Okubo, K.; Ishikawa, A.; Futata, Y.; Hamasaki, N.

[145089]: Band 3-Memphis is associated with a lower transport rate of phosphoenolpyruvate. Brit.

[145090]J. Haemat. 82: 122–125, 1992.

[145091]26030.

[145092]Iwase, S.; Ideguchi, H.; Takao, M.; Horiguchi–Yamada, J.;
Iwasaki,

[145093]M.; Takahara, S.; Sekikawa, T.; Mochizuki, S.; Yamada, H.:
Band 3

[145094]Tokyo: thr837–ala837 substitution in erythrocyte band 3
protein associated

[145095]with spherocytic hemolysis. Acta Haemat. 100: 200–203,
1998.

[145096]26031.

[145097]Jarolim, P.; Palek, J.; Rubin, H. L.; Prchal, J. T.; Korsgren,

[145098]C.; Cohen, C. M.: Band 3 Tuscaloosa: pro327–to–arg327
substitution

[145099]in the cytoplasmic domain of erythrocyte band 3 protein
associated

[145100]with spherocytic hemolytic anemia and partial deficiency
of protein

[145101]4.2. (Abstract) Blood 78 (suppl.): 252a, 1991.

[145102]26032.

[145103]Jarolim, P.; Rubin, H. L.; Brabec, V.; Chrobak, L.; Zolotarev,

[145104]A. S.; Alper, S. L.; Brugnara, C.; Wichterle, H.; Palek, J.:

Mutations

[145105]of conserved arginines in the membrane domain of erythroid band 3

[145106]lead to a decrease in membrane-associated band 3 and to the phenotype

[145107]of hereditary spherocytosis. Blood 85: 634–640, 1995.

[145108]26033.

[145109]Jarolim, P.; Rubin, H. L.; Liu, S.-C.; Cho, M. R.; Brabec, V.;

[145110]Derick, L. H.; Yi, S. J.; Saad, S. T. O.; Alper, S.; Brugnara, C.;

[145111]Golan, D. E.; Palek, J.: Duplication of 10 nucleotides in the erythroid

[145112]band 3 (AE1) gene in a kindred with hereditary spherocytosis and band

[145113]3 protein deficiency (band 3–Prague). J. Clin. Invest. 93: 121–130,

[145114]1994.

[145115]26034.

[145116]Inaba, M.; Yawata, A.; Koshino, I.; Sato, K.; Takeuchi, M.; Takakuwa,

- [145117]Y.; Manno, S.; Yawata, Y.; Kanzaki, A.; Sakai, J.; Ban, A.;
Ono, K.;
- [145118]Maede, Y.: Defective anion transport and marked spherocytosis with
- [145119]membrane instability caused by hereditary total deficiency
of red
- [145120]cell band 3 in cattle due to a nonsense mutation. J. Clin.
Invest. 97:
- [145121]1804–1817, 1996.
- [145122]26035.
- [145123]Inoue, T.; Kanzaki, A.; Kaku, M.; Yawata, A.; Takezono, M.;
Okamoto,
- [145124]N.; Wada, H.; Sugihara, T.; Yamada, O.; Katayama, Y.; Nagata, N.;
- [145125]Yawata, Y.: Homozygous missense mutation (band 3
Fukuoka: G130R):
- [145126]a mild form of hereditary spherocytosis with near-normal
band 3 content
- [145127]and minimal changes of membrane ultrastructure despite
moderate protein
- [145128]4.2 deficiency. Brit. J. Haemat. 102: 932–939, 1998.
- [145129]26036.

[145130]Jarolim, P.; Rubin, H. L.; Zakova, D.; Storry, J.; Reid, M. E.

[145131]: Characterization of seven low incidence blood group antigens carried

[145132]by erythrocyte band 3 protein. Blood 92: 4836–4843, 1998.

[145133]26037.

[145134]Jarolim, P.; Rubin, H. L.; Zhai, S.; Sahr, K. E.; Liu, S.-C.;

[145135]Mueller, T. J.; Palek, J.: Band 3 Memphis: a widespread polymorphism

[145136]with abnormal electrophoretic mobility of erythrocyte band 3 protein

[145137]caused by substitution AAG-to-GAG (lys-to-glu) in codon 56. Blood 80:

[145138]1592–1598, 1992.

[145139]26038.

[145140]Jay, D. G.: Role of band 3 in homeostasis and cell shape. Cell 86:

[145141]853–854, 1996.

[145142]26039.

[145143]Jenkins, P. B.; Abou-Alfa, G. K.; Dhermy, D.; Bursaux, E.; Feo,

[145144]C.; Scarpa, A. L.; Lux, S. E.; Garbarz, M.; Forget, B. G.; Gallagher,

[145145]P. G.: A nonsense mutation in the erythrocyte band 3 gene associated

[145146]with decreased mRNA accumulation in a kindred with dominant hereditary

[145147]spherocytosis. J. Clin. Invest. 97: 373–380, 1996.

[145148]26040.

[145149]Jones, G. L.; Edmundson, H. M.; Wesche, D.; Saul, A.: Human erythrocyte

[145150]band–3 has an altered N terminus in malaria-resistant Melanesian ovalocytosis. Biochim.

[145151]Biophys. Acta 1096: 33–40, 1990.

[145152]26041.

[145153]Kay, M. M. B.: Band 3 in aging and neurological disease. Ann.

[145154]N.Y. Acad. Sci. 621: 179–204, 1991.

[145155]26042.

[145156]Kay, M. M. B.; Bosman, G. J. C. G. M.; Lawrence, C.: Functional

[145157]topography of band 3: specific structural alteration linked

to functional

[145158]aberrations in human erythrocytes. Proc. Nat. Acad. Sci.

85: 492-496,

[145159]1988.

[145160]26043.

[145161]Kay, M. M. B.; Flowers, N.; Goodman, J.; Bosman, G.: Alter-
ation

[145162]in membrane protein band 3 associated with accelerated
erythrocyte

[145163]aging. Proc. Nat. Acad. Sci. 86: 5834-5838, 1989.

[145164]26044.

[145165]Kay, M. M. B.; Lawrence, C.; Bosman, G. J. C. G. M.: Molec-
ular

[145166]anatomy of an anemia. (Abstract) Clin. Res. 35: 599A,
1987.

[145167]26045.

[145168]Kay, M. M. B.; Marchalonis, J. J.; Hughes, J.; Watanabe, K.;
Schluter,

[145169]S. F.: Definition of a physiologic aging autoantigen by us-
ing synthetic

[145170]peptides of membrane protein band 3: localization of the

active antigenic

[145171]sites. Proc. Nat. Acad. Sci. 87: 5734–5738, 1990.

[145172]26046.

[145173]Kidson, C.; Lamont, G.; Saul, A.; Nurse, G. T.: Ovalocytic erythrocytes

[145174]from Melanesians are resistant to invasion by malaria parasites in

[145175]culture. Proc. Nat. Acad. Sci. 78: 5829–5832, 1981.

[145176]26047.

[145177]Langdon, R. G.; Holman, V. P.: Immunological evidence that band

[145178]3 is the major glucose transporter of the human erythrocyte membrane. Biochim.

[145179]Biophys. Acta 945: 23–32, 1988.

[145180]26048.

[145181]Lie-Injo, L. E.: Hereditary ovalocytosis and haemoglobin E-ovalocytosis

[145182]in Malayan aborigines. Nature 208: 1329, 1965.

[145183]26049.

[145184]Lie-Injo, L. E.; Fix, A.; Bolton, J. M.; Gilman, R. H.:

Haemoglobin

- [145185]E-hereditary elliptocytosis in Malayan aborigines. *Acta Haemat.* 47: 210-216, 1972.
- [145187]26050.
- [145188]Liu, S.-C.; Jarolim, P.; Rubin, H. L.; Palek, J.; Amato, D.; Hassan, K.; Zaik, M.; Sapak, P.: The homozygous state for the band 3 protein mutation in Southeast Asian ovalocytosis may be lethal. (Letter) *Blood* 84: 3590-3591, 1994.
- [145192]26051.
- [145193]Liu, S.-C.; Zhai, S.; Palek, J.; Golan, D. E.; Amato, D.; Hassan, K.; Nurse, G. T.; Babona, D.; Coetzer, T.; Jarolim, P.; Zaik, M.; Borwein, S.: Molecular defect of the band 3 protein in Southeast Asian ovalocytosis. *New Eng. J. Med.* 323: 1530-1538, 1990.
- [145197]26052.

[145198]Lux, S.; Bedrosian, C.; Shalev, O.; Morris, M.; Chasis, J.;
Davies,

[145199]K.; Savvides, P.; Telen, M.: Deficiency of band 3 in dominant hereditary

[145200]spherocytosis with normal spectrin content. (Abstract)
Clin. Res. 38:

[145201]300A, 1990.

[145202]26053.

[145203]Lux, S. E.; John, K. M.; Kopito, R. R.; Lodish, H. F.: Cloning

[145204]and characterization of band 3, the human erythrocyte
anion-exchange

[145205]protein (AE1). Proc. Nat. Acad. Sci. 86: 9089-9093, 1989.

[145206]26054.

[145207]Maillet, P.; Vallier, A.; Reinhart, W. H.; Wyss, E. J.; Ott, P.;

[145208]Texier, P.; Baklouti, F.; Tanner, M. J. A.; Delaunay, J.; Aloisio,

[145209]N.: Band 3 Chur: a variant associated with band
3-deficient hereditary

[145210]spherocytosis and substitution in a highly conserved position of transmembrane

[145211]segment 11. Brit. J. Haemat. 91: 804-810, 1995.

[145212]26055.

[145213]Carrington, M.; Kissner, T.; Gerrard, B.; Ivanov, S.; O'Brien,

[145214]S. J.; Dean, M.: Novel alleles of the chemokine–receptor
gene CCR5. Am.

[145215]J. Hum. Genet. 61: 1261–1267, 1997.

[145216]26056.

[145217]Chaganti, R. S. K.: Personal Communication. New York, N.
Y.

[145218]10/22/1993.

[145219]26057.

[145220]Chaudhuri, A.; Polyakova, J.; Zbrzezna, V .; Pogo, A. O.:
The

[145221]coding sequence of Duffy blood group gene in humans
and simians: restriction

[145222]fragment length polymorphism, antibody and malarial
parasite specificities,

[145223]and expression in nonerythroid tissues in Duffy–negative
individuals. Blood 85:

[145224]615–621, 1995.

[145225]26058.

[145226]Chaudhuri, A.; Polyakova, J.; Zbrzezna, V.; Williams, K.;
Gulati,

[145227]S.; Pogo, A. O.: Cloning of glycoprotein D cDNA, which
encodes the

[145228]major subunit of the Duffy blood group system and the
receptor for

[145229]the Plasmodium vivax malaria parasite. Proc. Nat. Acad.
Sci. 90:

[145230]10793–10797, 1993.

[145231]26059.

[145232]Cook, P. J. L.; Page, B. M.; Johnston, A. W.; Stanford, W. K.;

[145233]Gavin, J.: Four further families informative for 1q and the
Duffy

[145234]blood group. Cytogenet. Cell Genet. 22: 378–380, 1978.

[145235]26060.

[145236]Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Jacobs, P. A.;
Polani,

[145237]P. E.: Segregation of genetic markers in families with
chromosome

[145238]polymorphisms and structural rearrangements involving
chromosome no.

[145239]1. Ann. Hum. Genet. 37: 261-274, 1974.

[145240]26061.

[145241]Crawford, M. N.; Punnett, H. H.; Carpenter, G. G.: Deletion
of

[145242]the long arm of chromosome 16 and an unexpected Duffy
blood group

[145243]phenotype reveal a possible autosomal linkage. Nature
215: 1075-1076,

[145244]1967.

[145245]26062.

[145246]Gelpi, A. P.; King, M. C.: Association of Duffy blood groups

[145247]with the sickle cell trait. Hum. Genet. 32: 65-68, 1976.

[145248]26063.

[145249]Hadley, T. J.; David, P. H.; McGinniss, M. H.; Miller, L. H.:

[145250]Identification of an erythrocyte component carrying the
Duffy blood

[145251]group Fy-a antigen. Science 223: 597-599, 1984.

[145252]26064.

[145253]Hadley, T. J.; Peiper, S. C.: From malaria to chemokine re-

ceptor:

[145254]the emerging physiologic role of the Duffy blood group
antigen. Blood 89:

[145255]3077–3091, 1997.

[145256]26065.

[145257]Hamblin, M. T.; Di Rienzo, A.: Detection of the signature
of

[145258]natural selection in humans: evidence from the Duffy
blood group locus. Am.

[145259]J. Hum. Genet. 66: 1669–1679, 2000.

[145260]26066.

[145261]Hamblin, M. T.; Thompson, E. E.; Di Rienzo, A.: Complex
signatures

[145262]of natural selection at the Duffy blood group locus. Am. J.
Hum.

[145263]Genet. 70: 369–383, 2002.

[145264]26067.

[145265]Hessner, M. J.; Pircon, R. A.; Johnson, S. T.; Luhm, R. A.:
Prenatal

[145266]genotyping of the Duffy blood group system by allele-
specific polymerase

[145267]chain reaction. Prenatal Diag. 19: 41–45, 1999.

[145268]26068.

[145269]Horuk, R.; Chitnis, C. E.; Darbonne, W. C.; Colby, T. J.; Ry-
bicki,

[145270]A.; Hadley, T. J.; Miller, L. H.: A receptor for the malarial
parasite

[145271]Plasmodium vivax: the erythrocyte chemokine receptor.
Science 261:

[145272]1182–1184, 1993.

[145273]26069.

[145274]Howard, P. N.; Stoddard, G. R.; Goddard, M. W.; Seely, J.
R.:

[145275]Giemsa banding of chromosome 1qh+ and linkage analy-
sis. J. Med. Genet. 12:

[145276]44–48, 1975.

[145277]26070.

[145278]Iwamoto, S.; Li, J.; Sugimoto, N.; Okuda, H.; Kajii, E.: Char-
acterization

[145279]of the Duffy gene promoter: evidence for tissue-specific
abolishment

[145280]of expression in Fy(a–b–) of black individuals. Biochem.

Biophys.

[145281]Res. Commun. 222: 852–859, 1996.

[145282]26071.

[145283]Lautenberger, J. A.; Stephens, J. C.; O'Brien, S. J.; Smith, M.

[145284]W.: Significant admixture linkage disequilibrium across 30
cM around

[145285]the FY locus in African Americans. Am. J. Hum. Genet. 66:
969–978,

[145286]2000.

[145287]26072.

[145288]Lee, C. S. N.; Ying, K. L.; Bowen, P.: Position of the Duffy

[145289]locus on chromosome 1 in relation to breakpoints for
structural rearrangements. Am.

[145290]J. Hum. Genet. 26: 93–102, 1974.

[145291]26073.

[145292]Li, J.; Iwamoto, S.; Sugimoto, N.; Okuda, H.; Kajii, E.: Dinu-
cleotide

[145293]repeat in the 3-prime flanking region provides a clue to
the molecular

[145294]evolution of the Duffy gene. Hum. Genet. 99: 573–577,

1997.

[145295]26074.

[145296]Braun, A.; Kammerer, S.; Bohme, E.; Muller, B.; Roscher, A.
A.

[145297]: Identification of polymorphic sites of the human
bradykinin B(2)

[145298]receptor gene. Biochem. Biophys. Res. Commun. 211:
234-240, 1995.

[145299]26075.

[145300]Erdmann, J.; Hegemann, N.; Weidemann, A.; Kallisch, H.;
Hummel,

[145301]M.; Hetzer, R.; Fleck, E.; Regitz-Zagrosek, V.: Screening
the human

[145302]bradykinin B2 receptor gene in patients with cardiovascu-
lar diseases:

[145303]identification of a functional mutation in the promoter and
a new

[145304]coding variant (T21M). Am. J. Med. Genet. 80: 521-525,
1998.

[145305]26076.

[145306]Hess, J. F.; Borkowski, J. A.; Young, G. S.; Strader, C. D.;
Ransom,

- [145307]R. W.: Cloning and pharmacological characterization of a human bradykinin
- [145308](BK-2) receptor. Biochem. Biophys. Res. Commun. 184: 260-268, 1992.
- [145309]26077.
- [145310]Kammerer, S.; Braun, A.; Arnold, N.; Roscher, A. A.: The human
- [145311]bradykinin B(2) receptor gene: full length cDNA, genomic organization
- [145312]and identification of the regulatory region. Biochem. Biophys. Res.
- [145313]Commun. 211: 226-233, 1995.
- [145314]26078.
- [145315]Ma, J.; Wang, D.; Ward, D. C.; Chen, L.; Dessai, T.; Chao, J.;
- [145316]Chao, L.: Structure and chromosomal localization of the gene (BDKRB2)
- [145317]encoding human bradykinin B2 receptor. Genomics 23: 362-369, 1994.
- [145318]26079.
- [145319]Powell, S. J.; Slynn, G.; Thomas, C.; Hopkins, B.; Briggs, I.;
- [145320]Graham, A.: Human bradykinin B2 receptor: nucleotide

sequence analysis

[145321]and assignment to chromosome 14. Genomics 15:
435–438, 1993.

[145322]26080.

[145323]Taketo, M.; Yokoyama, S.; Rochelle, J.; Kimura, S.; Higashida,

[145324]H.; Taketo, M.; Seldin, M. F.: Mouse B2 bradykinin receptor
gene

[145325]maps to distal chromosome 12. Genomics 27: 222–223,
1995.

[145326]26081.

[145327]Lu, Q.; Sun, E. E.; Klein, R. S.; Flanagan, J. G.: Ephrin-B re-
verse

[145328]signaling is mediated by a novel PDZ–RGS protein and se-
lectively inhibits

[145329]G protein–coupled chemoattraction. Cell 105: 69–79,
2001.

[145330]26082.

[145331]Amara, S. G.; Arriza, J. L.; Leff, S. E.; Swanson, L. W.;
Evans,

[145332]R. M.; Rosenfeld, M. G.: Expression in brain of a messen-
ger RNA encoding

[145333]a novel neuropeptide homologous to calcitonin gene-related peptide.

[145334]Science 229: 1094–1097, 1985.

[145335]26083.

[145336]Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.; Geurts van

[145337]Kessel, A. H. M.; Baylin, S. B.; Nelkin, B. D.; Jansz, H. S.; Lips,

[145338]C. J. M.: The second human calcitonin/CGRP gene is located on chromosome

[145339]11. Hum. Genet. 70: 259–263, 1985.

[145340]26084.

[145341]Steenbergh, P. H.; Hoppener, J. W. M.; Zandberg, J.; Lips, C. J.

[145342]M.; Jansz, H. S.: A second human calcitonin/CGRP gene. FEBS Lett. 183:

[145343]403–407, 1985.

[145344]26085.

[145345]Arthur, J. S. C.; Elce, J. S.; Hegadorn, C.; Williams, K.; Greer,

[145346]P. A.: Disruption of the murine calpain small subunit gene,

Capn4:

[145347]calpain is essential for embryonic development but not for
cell growth

[145348]and division. Molec. Cell. Biol. 20: 4474–4481, 2000.

[145349]26086.

[145350]Ohno, S.; Emori, Y.; Suzuki, K.: Nucleotide sequence of a
cDNA

[145351]coding for the small subunit of human calcium-dependent
protease. Nucleic

[145352]Acids Res. 14: 5559 only, 1986.

[145353]26087.

[145354]Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Ohmi-
Imajoh,

[145355]S.; Suzuki, K.; Shimizu, Y.; Shimizu, N.: Four genes for the
calpain

[145356]family locate on four distinct human chromosomes.
(Abstract) Cytogenet.

[145357]Cell Genet. 51: 1054–1055, 1989.

[145358]26088.

[145359]Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Shimizu,
Y.;

[145360] Ohmi-Imajoh, S.; Shimizu, N.; Suzuki, K.: Four genes for the calpain

[145361] family locate on four distinct human chromosomes. Cytogenet. Cell

[145362] Genet. 53: 225-229, 1990.

[145363] 26089.

[145364] Sakihama, T.; Kakidani, H.; Zenita, K.; Yumoto, N.; Kikuchi, T.;

[145365] Sasaki, T.; Kannagi, R.; Nakanishi, S.; Ohmori, M.; Takio, K.; Titani,

[145366] K.; Murachi, T.: A putative Ca(2+)-binding protein: structure of

[145367] the light subunit of porcine calpain elucidated by molecular cloning

[145368] and protein sequence analysis. Proc. Nat. Acad. Sci. 82: 6075-6079,

[145369] 1985.

[145370] 26090.

[145371] Berchtold, M. W.; Egli, R.; Rhyner, J. A.; Hameister, H.; Strehler,

[145372] E. E.: Localization of the human bona fide calmodulin genes CALM1,

[145373]CALM2, and CALM3 to chromosomes 14q24–q31,
2p21.1–p21.3, and 19q13.2–q13.3. Genomics 16:

[145374]461–465, 1993.

[145375]26091.

[145376]Chin, D.; Winkler, K. E.; Means, A. R.: Characterization of
substrate

[145377]phosphorylation and use of calmodulin mutants to ad-
dress implications

[145378]from the enzyme crystal structure of calmodulin–depen-
dent protein

[145379]kinase I. J. Biol. Chem. 272: 31235–31240, 1997.

[145380]26092.

[145381]Drum, C. L.; Yan, S.–Z.; Bard, J.; Shen, Y.–Q.; Lu, D.; Soe-
laiman,

[145382]S.; Grabarek, Z.; Bohm, A.; Tang, W.–J.: Structural basis for
the

[145383]activation of anthrax adenylyl cyclase exotoxin by
calmodulin. Nature 415:

[145384]396–402, 2002.

[145385]26093.

[145386]Koller, M.; Schnyder, B.; Strehler, E. E.: Structural organi-

zation

[145387]of the human CaMIII calmodulin gene. Biochim. Biophys.

Acta 1087:

[145388]180–189, 1990.

[145389]26094.

[145390]Kretsinger, R. H.; Rudnick, S. E.; Weissman, L. J.: Crystal
structure

[145391]of calmodulin. J. Inorganic Biochem. 28: 289–302, 1986.

[145392]26095.

[145393]McPherson, J. D.; Hickie, R. A.; Wasmuth, J. J.; Meyskens,
F. L.;

[145394]Perham, R. N.; Strehler, E. E.; Graham, M. T.: Chromosomal
localization

[145395]of multiple genes encoding calmodulin. (Abstract) Cyto-
genet. Cell

[145396]Genet. 58: 1951 only, 1991.

[145397]26096.

[145398]Pegues, J. C.; Friedberg, F.: Multiple mRNAs encoding hu-
man calmodulin. Biochem.

[145399]Biophys. Res. Commun. 172: 1145–1149, 1990.

[145400]26097.

- [145401]Rhyner, J. A.; Ottiger, M.; Wicki, R.; Greenwood, T. M.;
Strehler,
[145402]E. E.: Structure of the human CALM1 calmodulin gene and
identification
[145403]of two CALM1-related pseudogenes CALM1P1 and
CALM1P2. *Europ. J. Biochem.* 225:
[145404]71–82, 1994.
[145405]26098.
[145406]Scambler, P. J.; McPherson, M. A.; Bates, G.; Bradbury, N.
A.;
[145407]Dormer, R. L.; Williamson, R.: Biochemical and genetic ex-
clusion
[145408]of calmodulin as the site of the basic defect in cystic fi-
brosis. *Hum.*
[145409]*Genet.* 76: 278–282, 1987.
[145410]26099.
[145411]Schumacher, M. A.; Rivard, A. F.; Bachinger, H. P.; Adel-
man, J.
[145412]P.: Structure of the gating domain of a $\text{Ca}(2+)$ -activated
 K^+ channel
[145413]complexed with $\text{Ca}(2+)/\text{calmodulin}$. *Nature* 410:
1120–1124, 2001.

[145414]26100.

[145415]Sen Gupta, B.; Detera-Wadleigh, S. D.; McBride, O. W.;
Friedberg,

[145416]F.: A calmodulin pseudogene on human chromosome 17.
Nucleic Acids

[145417]Res. 17: 2868 only, 1989.

[145418]26101.

[145419]Elbaz, A.; Vale-Santos, J.; Jurkat-Rott, K.; Lapie, P.;
Ophoff,

[145420]R. A.; Bady, B.; Links, T. P.; Piussan, C.; Vila, A.; Monnier,
N.;

[145421]Padberg, G. W.; Abe, K.; Feingold, N.; Guimaraes, J.;
Wintzen, A.

[145422]R.; van der Hoeven, J. H.; Saudubray, J. M.; Grunfeld, J. P.;
Lenoir,

[145423]G.; Nivet, H.; Echenne, B.; Frants, R. R.; Fardeau, M.;
Lehmann-Horn,

[145424]F.; Fontaine, B.: Hypokalemic periodic paralysis and the
dihydropyridine

[145425]receptor (CACNL1A3): genotype/phenotype correlations
for two predominant

[145426]mutations and evidence for the absence of a founder ef-

fect in 16 Caucasian

[145427]families. Am. J. Hum. Genet. 56: 374–380, 1995.

[145428]26102.

[145429]Gluecksohn-Waelsch, S.: Lethal genes and analysis of differentiation. Science 142:

[145430]1269–1276, 1963.

[145431]26103.

[145432]Gregg, R. G.; Couch, F.; Hogan, K.; Powers, P. A.: Assignment

[145433]of the human gene for the alpha-1 subunit of the skeletal muscle DHP-sensitive

[145434]Ca(2+) channel (CACNL1A3) to chromosome 1q31–q32. Genomics 15: 107–112,

[145435]1993.

[145436]26104.

[145437]Hogan, K.; Gregg, R. G.; Powers, P. A.: The structure of the

[145438]gene encoding the human skeletal muscle alpha-1 subunit of the dihydropyridine-sensitive

[145439]L-type calcium channel (CACNL1A3). Genomics 31: 392–394, 1996.

[145440]26105.

[145441]lles, D. E.; Segers, B.; Weghuis, D. O.; Suijkerbuijk, R.;
Mikala,
[145442]G.; Schwartz, A.; Wieringa, B.: Refined localization of the
alpha-1-subunit
[145443]of the skeletal muscle L-type voltage-dependent calcium
channel (CACNL1A3)
[145444]to human chromosome 1q32 by in situ hybridization. Ge-
nomics 19:
[145445]561-563, 1994.
[145446]26106.
[145447]Jurkat-Rott, K.; Lehmann-Horn, F.; Elbaz, A.; Heine, R.;
Gregg,
[145448]R. G.; Hogan, K.; Powers, P. A.; Lapie, P.; Vale-Santos, J. E.;
Weissenbach,
[145449]J.; Fontaine, B.: A calcium channel mutation causing hy-
pokalemic
[145450]periodic paralysis. Hum. Molec. Genet. 3: 1415-1419,
1994.
[145451]26107.
[145452]Pai, A. C.: Developmental genetics of a lethal mutation,
muscular
[145453]dysgenesis (mdg), in the mouse. I. Genetic analysis and

gross morphology. Dev.

[145454]Biol. 11: 82–92, 1965.

[145455]26108.

[145456]Sillen, A.; Sorensen, T.; Kantola, I.; Friis, M. L.; Gustavson,

[145457]K.–H.; Wadelius, C.: Identification of mutations in the
CACNL1A3

[145458]gene in 13 families of Scandinavian origin having hy-
pokalemic periodic

[145459]paralysis and evidence of a founder effect in Danish fami-
lies. Am.

[145460]J. Med. Genet. 69: 102–106, 1997.

[145461]26109.

[145462]Stewart, S. L.; Hogan, K.; Rosenberg, H.; Fletcher, J. E.:
Identification

[145463]of the arg1086his mutation in the alpha subunit of the
voltage-dependent

[145464]calcium channel (CACNA1S) in a North American family
with malignant

[145465]hyperthermia. Clin. Genet. 59: 178–184, 2001.

[145466]26110.

[145467]Tanabe, T.; Beam, K. G.; Powell, J. A.; Numa, S.: Restora-

tion

[145468]of excitation-contraction coupling and slow calcium current in dysgenic

[145469]muscle by dihydropyridine receptor complementary DNA.

Nature 336:

[145470]134-139, 1988.

[145471]26111.

[145472]Tanabe, T.; Takeshima, H.; Mikami, A.; Flockerzi, V.;

Takahashi,

[145473]H.; Kangawa, K.; Kojima, M.; Matsuo, H.; Hirose, T.; Numa,

S.: Primary

[145474]structure of the receptor for calcium channel blockers

from skeletal

[145475]muscle. Nature 328: 313-318, 1987.

[145476]26112.

[145477]Ambartsumian, N.; Tarabykina, S.; Grigorian, M.; Tulchin-

sky, E.;

[145478]Hulgaard, E.; Georgiev, G.; Lukanidin, E.: Characterization

of two

[145479]splice variants of metastasis-associated human mts1

gene. Gene 159:

[145480]125-130, 1995.

[145481]26113.

[145482]Jackson-Grusby, L. L.; Swiergiel, J.; Linzer, D. I.: A growth-related

[145483]mRNA in cultured mouse cells encodes a placental calcium binding protein.

[145484]Nucleic Acids Res. 15: 6677-6690, 1987.

[145485]26114.

[145486]Saris, C. J.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.; Noonan,

[145487]D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and tissue distribution

[145488]of the mRNA for bovine and murine p11, the S100-related light chain

[145489]of the protein-tyrosine kinase substrate p36 (calpactin I). J. Biol.

[145490]Chem. 262: 10663-10671, 1987.

[145491]26115.

[145492]Lefort, A.; Lecocq, R.; Libert, F.; Lamy, F.; Swillens, S.; Vassart,

[145493]G.; Dumont, J. E.: Cloning and sequencing of a calcium-binding protein

[145494]regulated by cyclic AMP in the thyroid. EMBO J. 8:
111–116, 1989.

[145495]26116.

[145496]Loughlin, J.; Irven, C.; Sykes, B.: Exclusion of the cartilage
[145497]link protein and the cartilage matrix protein genes as the
mutant

[145498]loci in several heritable chondrodysplasias. Hum. Genet.
94: 698–700,

[145499]1994.

[145500]26117.

[145501]Deak, F.; Piecha, D.; Bachrati, C.; Paulsson, M.; Kiss, I.: Pri-
mary

[145502]structure and expression of matrilin-2, the closest relative
of cartilage

[145503]matrix protein within the von Willebrand factor type A-like
module

[145504]superfamily. J. Biol. Chem. 272: 9268–9274, 1997.

[145505]26118.

[145506]Deak, F.; Wagener, R.; Kiss, I.; Paulsson, M.: The matrilins:

[145507]a novel family of oligomeric extracellular matrix proteins.
Matrix

[145508]Biol. 18: 55–64, 1999.

[145509]26119.

[145510]Hansson, A.-S.; Heinegard, D.; Holmdahl, R.: A new animal model

[145511]for relapsing polychondritis, induced by cartilage matrix protein

[145512](matrilin-1). J. Clin. Invest. 104: 589–598, 1999.

[145513]26120.

[145514]Jenkins, R. N.; Osborne–Lawrence, S. L.; Sinclair, A. K.; Eddy,

[145515]R. L., Jr.; Byers, M. G.; Shows, T. B.; Duby, A. D.: Structure and

[145516]chromosomal location of the human gene encoding cartilage matrix protein. J.

[145517]Biol. Chem. 265: 19624–19631, 1990.

[145518]26121.

[145519]Lang, B.; Rothenfusser, A.; Lanchbury, J. S.; Rauh, G.; Breedveld,

[145520]F. C.; Urlacher, A.; Albert, E. D.; Peter, H.-H.; Melchers, I.: Susceptibility

[145521]to relapsing polychondritis is associated with HLA–DR4.

Arthritis

[145522]Rheum. 36: 660–664, 1993.

[145523]26122.

[145524]Muratoglu, S.; Krysan, K.; Balazs, M.; Sheng, H.; Zakany, R.; Modis,

[145525]L.; Kiss, I.; Deak, F.: Primary structure of human matrilin-2, chromosome

[145526]location of the MATN2 gene and conservation of an AT-AC intron in

[145527]matrilin genes. Cytogenet. Cell Genet. 90: 323–327, 2000.

[145528]26123.

[145529]Wagener, R.; Kobbe, B.; Paulsson, M.: Primary structure of matrilin-3,

[145530]a new member of a family of extracellular matrix proteins related

[145531]to cartilage matrix protein (matrilin-1) and von Willebrand factor. FEBS

[145532]Lett. 413: 129–134, 1997.

[145533]26124.

[145534]Zeuner, M.; Straub, R. H.; Rauh, G.; Albert, E. D.; Scholmerich,

[145535]J.; Lang, B.: Relapsing polychondritis: clinical and immunogenetic

[145536]analysis of 62 patients. J. Rheum. 24: 96–101, 1997.

[145537]26125.

[145538]Boldyreff, B.; Klett, C.; Gottert, E.; Geurts van Kessel, A.; Hameister,

[145539]H.; Issinger, O.–G.: Assignment of casein kinase 2 alpha sequences

[145540]to two different human chromosomes. Hum. Genet. 89: 79–82, 1992.

[145541]26126.

[145542]Doray, B.; Ghosh, P.; Griffith, J.; Geuze, H. J.; Kornfeld, S.

[145543]: Cooperation of GGAs and AP–1 in packaging MPRs at the trans–Golgi

[145544]network. Science 297: 1700–1703, 2002.

[145545]26127.

[145546]Meisner, H.; Heller–Harrison, R.; Buxton, J.; Czech, M. P.: Molecular

[145547]cloning of the human casein kinase II alpha subunit. Biochemistry 28:

[145548]4072–4076, 1989.

[145549]26128.

[145550]Wirkner, U.; Voss, H.; Lichter, P.; Ansorge, W.; Pyerin, W.:
The

[145551]human gene (CSNK2A1) coding for the casein kinase II
subunit alpha

[145552]is located on chromosome 20 and contains tandemly ar-
ranged Alu repeats. Genomics 19:

[145553]257-265, 1994.

[145554]26129.

[145555]Wirkner, U.; Voss, H.; Lichter, P.; Weitz, S.; Ansorge, W.;
Pyerin,

[145556]W.: Human casein kinase II subunit alpha: sequence of a
processed

[145557](pseudo)gene and its localization on chromosome 11.
Biochim. Biophys.

[145558]Acta 1131: 220-222, 1992.

[145559]26130.

[145560]Yang-Feng, T. L.; Zheng, K.; Kopatz, I.; Naiman, T.;
Canaani, D.

[145561]: Mapping of the human casein kinase II catalytic subunit
genes: two

[145562]loci carrying the homologous sequences for the alpha subunit. Nucleic

[145563]Acids Res. 19: 7125–7129, 1991.

[145564]26131.

[145565]Lozeman, F. J.; Litchfield, D. W.; Piening, C.; Takio, K.; Walsh,

[145566]K. A.; Krebs, E. G.: Isolation and characterization of human cDNA

[145567]clones encoding the alpha and the alpha-prime subunits of casein kinase

[145568]II. Biochemistry 29: 8436–8447, 1990.

[145569]26132.

[145570]Xu, X.; Toselli, P. A.; Russell, L. D.; Seldin, D. C.: Globozoospermia

[145571]in mice lacking the casein kinase II alpha-prime catalytic subunit. Nature

[145572]Genet. 23: 118–121, 1999.

[145573]26133.

[145574]Wickenheisser, J. K.; Quinn, P. G.; Nelson, V. L.; Legro, R. S.;

[145575]Strauss, J. F., III; McAllister, J. M.: Differential activity of

[145576]the cytochrome P450 17-alpha-hydroxylase and steroido-
genic acute regulatory

[145577]protein gene promoters in normal and polycystic ovary
syndrome theca

[145578]cells. J. Clin. Endocr. Metab. 85: 2304-2311, 2000.

[145579]26134.

[145580]Geissler, E. N.; Liao, M.; Brook, J. D.; Martin, F. H.; Zsebo,

[145581]K. M.; Housman, D. E.; Galli, S. J.: Stem cell factor (SCF), a
novel

[145582]hematopoietic growth factor and ligand for c-kit tyrosine
kinase receptor,

[145583]maps on human chromosome 12 between 12q14.3 and
12qter. Somat. Cell

[145584]Molec. Genet. 17: 207-214, 1991.

[145585]26135.

[145586]Heissig, B.; Hattori, K.; Dias, S.; Friedrich, M.; Ferris, B.;

[145587]Hackett, N. R.; Crystal, R. G.; Besmer, P.; Lyden, D.; Moore,
M. A.

[145588]S.; Werb, Z.; Rafii, S.: Recruitment of stem and progenitor
cells

[145589]from the bone marrow niche requires MMP-9 mediated

release of Kit-ligand. Cell 109:

[145590]625-637, 2002.

[145591]26136.

[145592]Martin, F. H.; Suggs, S. V.; Langley, K. E.; Lu, H. S.; Ting, J.;

[145593]Okino, K. H.; Morris, C. F.; McNiece, I. K.; Jacobsen, F. W.;

Mendiaz,

[145594]E. A.; Birkett, N. C.; Smith, K. A.; and 15 others: Primary

structure

[145595]and functional expression of rat and human stem cell fac-

tor DNAs. Cell 63:

[145596]203-211, 1990.

[145597]26137.

[145598]Mathew, S.; Murty, V. V. V. S.; Hunziker, W.; Chaganti, R. S.

K.

[145599]: Subregional mapping of 13 single-copy genes on the

long arm of chromosome

[145600]12 by fluorescence in situ hybridization. Genomics 14:

775-779,

[145601]1992.

[145602]26138.

[145603]Abbott, C.; Piaggio, G.; Ammendola, R.; Solomon, E.;

Povey, S.;

[145604]Gounari, F.; De Simone, V.; Cortese, R.: Mapping of the
gene TCF2

[145605]for the transcription factor LFB3 to human chromosome
17 by polymerase

[145606]chain reaction. Genomics 8: 165–167, 1990.

[145607]26139.

[145608]Bach, I.; Mattei, M.–G.; Cereghini, S.; Yaniv, M.: Two mem-
bers

[145609]of an HNF1 homeoprotein family are expressed in human
liver. Nucleic

[145610]Acids Res. 19: 3553–3559, 1991.

[145611]26140.

[145612]Bingham, C.; Ellard, S.; Allen, L.; Bulman, M.; Shepherd,
M.; Frayling,

[145613]T.; Berry, P. J.; Clark, P. M.; Lindner, T.; Bell, G. I.; Ryffel,

[145614]G. U.; Nicholls, A. J.; Hattersley, A. T.: Abnormal nephron
development

[145615]associated with a frameshift mutation in the transcription
factor

[145616]hepatocyte nuclear factor–1–beta. Kidney Int. 57:

898–907, 2000.

[145617]26141.

[145618]Horikawa, Y.; Iwasaki, N.; Hara, M.; Furuta, H.; Hinokio, Y.;
Cockburn,

[145619]B. N.; Lindner, T.; Yamagata, K.; Ogata, M.; Tomonaga, O.;
Kuroki,

[145620]H.; Kasahara, T.; Iwamoto, Y.; Bell, G. I.: Mutation in hepa-
tocyte

[145621]nuclear factor-1-beta gene (TCF2) associated with MODY.
(Letter) Nature

[145622]Genet. 17: 384–385, 1997.

[145623]26142.

[145624]Menzel, R.; Kaisaki, P. J.; Rjasanowski, I.; Heinke, P.;
Kerner,

[145625]W.; Menzel, S.: A low renal threshold for glucose in dia-
betic patients

[145626]with a mutation in the hepatocyte nuclear factor-1-alpha
(HNF-1-alpha)

[145627]gene. Diabet. Med. 15: 816–820, 1998.

[145628]26143.

[145629]Nishigori, H.; Yamada, S.; Kohama, T.; Tomura, H.; Sho,

K.; Horikawa,
[145630]Y.; Bell, G. I.; Takeuchi, T.; Takeda, J.: Frameshift mutation, A263fsinsGG,
[145631]in the hepatocyte nuclear factor-1-beta gene associated
with diabetes
[145632]and renal dysfunction. Diabetes 47: 1354-1355, 1998.
[145633]26144.

[145634]Wild, W.; Pogge von Strandmann, E.; Nastos, A.; Senkel, S.;
Lingott-Frieg,
[145635]A.; Bulman, M.; Bingham, C.; Ellard, S.; Hattersley, A. T.;
Ryffel,
[145636]G. U.: The mutated human gene encoding hepatocyte nuclear factor
[145637]1-beta inhibits kidney formation in developing *Xenopus*
embryos. Proc.
[145638]Nat. Acad. Sci. 97: 4695-4700, 2000.
[145639]26145.

[145640]Kingsmore, S. F.; Watson, M. L.; Seldin, M. F.: Genetic
mapping
[145641]of the T lymphocyte-specific transcription factor 7 gene
on mouse
[145642]chromosome 11. Mammalian Genome 6: 378 only, 1995.

[145643]26146.

[145644]Roose, J.; Huls, G.; van Beest, M.; Moerer, P.; van der Horn, K.;

[145645]Goldschmeding, R.; Logtenberg, T.; Clevers, H.: Synergy between tumor

[145646]suppressor APC and the beta-catenin-Tcf4 target Tcf1. Science 285:

[145647]1923–1926, 1999.

[145648]26147.

[145649]van de Wetering, M.; Oosterwegel, M.; Holstege, F.; Dooyes, D.;

[145650]Suijkerbuijk, R.; Geurts van Kessel, A.; Clevers, H.: The human T

[145651]cell transcription factor-1 gene: structure, localization, and promoter

[145652]characterization. J. Biol. Chem. 267: 8530–8536, 1992.

[145653]26148.

[145654]van de Wetering, M.; Suijkerbuijk, R.; Geurts van Kessel, A.; Clevers,

[145655]H.: Assignment of the human T lymphocyte-specific transcription factor

[145656]TCF-1 to chromosome 5, band q31.1. (Abstract) Cyto-genet. Cell Genet. 58:

[145657]1906 only, 1991.

[145658]26149.

[145659]Verbeek, S.; Izon, D.; Hofhuis, F.; Robanus-Maandag, E.;
te Riele,

[145660]H.; van de Wetering, M.; Oosterwegel, M.; Wilson, A.; Mac-Donald, H.

[145661]R.; Clevers, H.: An HMG-box-containing T-cell factor re-quired for

[145662]thymocyte differentiation. Nature 374: 70-74, 1995.

[145663]26150.

[145664]Aplan, P. D.; Jones, C. A.; Chervinsky, D. S.; Zhao, X.;
Ellsworth,

[145665]M.; Wu, C.; McGuire, E. A.; Gross, K. W.: An scl gene prod-uct lacking

[145666]the transactivation domain induces bony abnormalities
and cooperates

[145667]with LMO1 to generate T-cell malignancies in transgenic
mice. EMBO

[145668]J. 16: 2408-2419, 1997.

[145669]26151.

[145670]Aplan, P. D.; Lombardi, D. P.; Ginsberg, A. M.; Cossman, J.;
Bertness,

[145671]V. L.; Kirsch, I. L.: Disruption of the human SCL locus by
'illegitimate'

[145672]V-(D)-J recombinase activity. Science 250: 1426-1429,
1990.

[145673]26152.

[145674]Aplan, P. D.; Lombardi, D. P.; Reaman, G. H.; Sather, H. N.;
Hammond,

[145675]G. D.; Kirsch, I. R.: Involvement of the putative
hematopoietic transcription

[145676]factor SCL in T-cell acute lymphoblastic leukemia. Blood
79: 1327-1333,

[145677]1992.

[145678]26153.

[145679]Brown, L.; Cheng, J.-T.; Chen, Q.; Siciliano, M. J.; Crist, W.;

[145680]Buchanan, G.; Baer, R.: Site-specific recombination of the
tal-1

[145681]gene is a common occurrence in human T-cell leukemia.
EMBO J. 9:

[145682]3343–3351, 1990.

[145683]26154.

[145684]Collazo–Garcia, N.; Scherer, P.; Aplan, P. D.: Cloning and characterization

[145685]of a murine SIL gene. *Genomics* 30: 506–513, 1995.

[145686]26155.

[145687]Izraeli, S.; Lowe, L. A.; Bertness, V. L.; Good, D. J.; Dorward,

[145688]D. W.; Kirsch, I. R.; Kuehn, M. R.: The SIL gene is required for

[145689]mouse embryonic axial development and left–right specification. *Nature* 399:

[145690]691–694, 1999.

[145691]26156.

[145692]Shiratori, A.; Okumura, K.; Nogami, M.; Taguchi, H.; Onozaki, T.;

[145693]Inoue, T.; Ando, T.; Shibata, T.; Izumi, M.; Miyazawa, H.; Hanaoka,

[145694]F.; Murakami, Y.; Eki, T.: Assignment of the 49–kDa (PRIM1) and 58–kDa

[145695](PRIM2A and PRIM2B) subunit genes of the human DNA

primase to chromosome

[145696]bands 1q44 and 6p11.1-p12. Genomics 28: 350-353,
1995. Note: Erratum:

[145697]Genomics 44: 251 only, 1997.

[145698]26157.

[145699]Roy, A. L.; Du, H.; Gregor, P. D.; Novina, C. D.; Martinez,
E.;

[145700]Roeder, R. G.: Cloning of an Inr- and E-box binding pro-
tein, TFII-I,

[145701]that interacts physically and functionally with USF1. EMBO
J. 16:

[145702]7091-7104, 1997.

[145703]26158.

[145704]Steingrimsson, E.; Sawadogo, M.; Gilbert, D. J.; Zervos, A.
S.;

[145705]Brent, R.; Blonar, M. A.; Fisher, D. E.; Copeland, N. G.;
Jenkins,

[145706]N. A.: Murine chromosomal location of five bHLH-Zip
transcription

[145707]factor genes. Genomics 28: 179-183, 1995.

[145708]26159.

[145709]Zhong, G.; Fan, P.; Ji, H.; Dong, F.; Huang, Y.: Identification
of a chlamydial protease-like activity factor responsible
for the
degradation of host transcription factors. J. Exp. Med.
193: 935-942,
2001.
26160.

[145714]Aasland, R.; Olsen, L. C.; Spurr, N. K.; Krokan, H. E.; Hel-
land,
D. E.: Chromosomal assignment of human uracil-DNA
glycosylase to
chromosome 12. Genomics 7: 139-141, 1990.
26161.

[145718]Dinner, A. R.; Blackburn, G. M.; Karplus, M.: Uracil-DNA
glycosylase
acts by substrate autocatalysis. Nature 413: 752-755,
2001.
26162.

[145721]Haug, T.; Skorpen, F.; Kvaloy, K.; Eftedal, I.; Lund, H.;
Krokan,

[145722]H. E.: Human uracil–DNA glycosylase gene: sequence organization,

[145723]methylation pattern, and mapping to chromosome 12q23–q24.1. Genomics 36:

[145724]408–416, 1996.

[145725]26163.

[145726]Haug, T.; Skorpen, F.; Lund, H.; Krokan, H. E.: Structure of the

[145727]gene for human uracil–DNA glycosylase and analysis of the promoter

[145728]function. FEBS Lett. 353: 180–184, 1994.

[145729]26164.

[145730]Meyer–Sieglar, K.; Mauro, D. J.; Seal, G.; Wurzer, J.; deRiel,

[145731]J. K.; Sirover, M. A.: A human nuclear uracil DNA glycosylase is

[145732]the 37–kDa subunit of glyceraldehyde–3–phosphate dehydrogenase. Proc.

[145733]Nat. Acad. Sci. 88: 8460–8464, 1991.

[145734]26165.

[145735]Nilsen, H.; Rosewell, I.; Robins, P.; Skjelbred, C. F.; Andersen,

[145736]S.; Slupphaug, G.; Daly, G.; Krokan, H. E.; Lindahl, T.; Barnes, D.

[145737]E.: Uracil–DNA glycosylase (UNG)–deficient mice reveal a primary

[145738]role of the enzyme during DNA replication. Molec. Cell 5: 1059–1065,

[145739]2000.

[145740]26166.

[145741]Olsen, L. C.; Aasland, R.; Wittwer, C. U.; Krokan, H. E.; Helland,

[145742]D. E.: Molecular cloning of human uracil–DNA glycosylase, a highly

[145743]conserved DNA repair enzyme. EMBO J. 8: 3121–3125, 1989.

[145744]26167.

[145745]Vollberg, T. M.; Siegler, K. M.; Cool, B. L.; Sirover, M. A.:

[145746]Isolation and characterization of the human uracil DNA

glycosylase

[145747]gene. Proc. Nat. Acad. Sci. 86: 8693–8697, 1989.

[145748]26168.

[145749]Hooper, M.; Hardy, K.; Handyside, A.; Hunter, S.; Monk,
M.: HPRT-deficient

[145750](Lesch–Nyhan) mouse embryos derived from germline col-
onization by

[145751]cultured cells. Nature 326: 292–295, 1987.

[145752]26169.

[145753]Kunicki, T. J.; Pidard, D.; Cazenave, J.–P.; Nurden, A. T.;
Caen,

[145754]J. P.: Inheritance of the human platelet alloantigen, PI(A1),
in

[145755]type I Glanzmann's thrombasthenia. J. Clin. Invest. 67:
717–724,

[145756]1981.

[145757]26170.

[145758]Letellier, S. J.; Hunter, J. B.; Aster, R. H.: Probable genetic

[145759]linkage between genes coding for platelet-specific anti-
gens of the

[145760]PI(A) and Bak systems. Am. J. Hemat. 29: 139–143, 1988.

[145761]26171.

[145762]Newman, P. J.; Seligsohn, U.; Lyman, S.; Collier, B. S.: The
molecular

[145763]genetic basis of Glanzmann thrombasthenia in the Iraqi-
Jewish and

[145764]Arab populations in Israel. Proc. Nat. Acad. Sci. 88:
3160-3164,

[145765]1991.

[145766]26172.

[145767]Newman, P. J.; Seligsohn, U.; Lyman, S.; Poncz, M.; Collier,
B.

[145768]S.: The molecular genetic basis of Glanzmann thrombas-
thenia in the

[145769]Iraqi-Jewish and Arab populations in Israel. (Abstract)
Clin. Res. 38:

[145770]467A only, 1990.

[145771]26173.

[145772]Haan, E. A.; Hull, Y. J.; White, S.; Cockington, R.; Charlton,

[145773]P.; Callen, D. F.: Tricho-rhino-phalangeal and branchio-
oto syndromes

[145774]in a family with an inherited rearrangement of chromo-

some 8q. Am.

[145775]J. Med. Genet. 32: 490–494, 1989.

[145776]26174.

[145777]Momeni, P.; Glockner, G.; Schmidt, O.; von Holtum, D.; Albrecht,

[145778]B.; Gillissen-Kaesbach, G.; Hennekam, R.; Meinecke, P.; Zabel, B.;

[145779]Rosenthal, A.; Horsthemke, B.; Ludecke, H.-J.: Mutations in a new

[145780]gene, encoding a zinc-finger protein, cause trichorhino-phalangeal

[145781]syndrome type I. Nature Genet. 24: 71–74, 2000.

[145782]26175.

[145783]Andersson, S.; Russell, D. W.: Structural and biochemical properties

[145784]of cloned and expressed human and rat steroid 5-alpha-reductases. Proc.

[145785]Nat. Acad. Sci. 87: 3640–3644, 1990.

[145786]26176.

[145787]Ellis, J. A.; Stebbing, M.; Harrap, S. B.: Genetic analysis of

[145788]male pattern baldness and the 5-alpha-reductase genes.

J. Invest.

[145789]Derm. 110: 849–853, 1998.

[145790]26177.

[145791]Harris, G.; Azzolina, B.; Baginsky, W.; Cimis, G.; Rasmusson, G.

[145792]H.; Tolman, R. L.; Raetz, C. R. H.; Ellsworth, K.: Identification

[145793]and selective inhibition of an isozyme of steroid
5-alpha-reductase

[145794]in human scalp. Proc. Nat. Acad. Sci. 89: 10787–10791,
1992.

[145795]26178.

[145796]Hsieh, C.-L.; Milatovich, A.; Russell, D.; Francke, U.: Chromosomal

[145797]mapping of human steroid 5 alpha-reductase gene
(SRD5A1) and pseudogene

[145798](SRD5AP1) in human and mouse. (Abstract) Cytogenet.
Cell Genet. 58:

[145799]1897 only, 1991.

[145800]26179.

[145801]Jenkins, E. P.; Andersson, S.; Imperato-McGinley, J.; Wil-

son, J.

[145802]D.; Russell, D. W.: Genetic and pharmacological evidence
for more

[145803]than one human steroid 5- α -reductase. J. Clin. Invest.
89: 293-300,

[145804]1992.

[145805]26180.

[145806]Jenkins, E. P.; Hsieh, C.-L.; Milatovich, A.; Normington, K.;
Berman,

[145807]D. M.; Francke, U.; Russell, D. W.: Characterization and
chromosomal

[145808]mapping of a human steroid 5- α -reductase gene and
pseudogene and

[145809]mapping of the mouse homologue. Genomics 11:
1102-1112, 1991.

[145810]26181.

[145811]Thigpen, A. E.; Silver, R. I.; Guileyardo, J. M.; Casey, M. L.;

[145812]McConnell, J. D.; Russell, D. W.: Tissue distribution and
ontogeny

[145813]of steroid 5- α -reductase isozyme expression. J. Clin.
Invest. 92:

[145814]903-910, 1993.

[145815]26182.

[145816]DeBose-Boyd, R. A.; Brown, M. S.; Li, W.-P.; Nohturfft, A.;
Goldstein,

[145817]J. L.; Espenshade, P. J.: Transport-dependent proteolysis
of SREBP:

[145818]relocation of Site-1 protease from Golgi to ER obviates the
need for

[145819]SREBP transport to Golgi. Cell 99: 703-712, 1999.

[145820]26183.

[145821]Hua, X.; Wu, J.; Goldstein, J. L.; Brown, M. S.; Hobbs, H. H.:

[145822]Structure of the human gene encoding sterol regulatory
element binding

[145823]protein-1 (SREBF1) and localization of SREBF1 and SREBF2
to chromosomes

[145824]17p11.2 and 22q13. Genomics 25: 667-673, 1995.

[145825]26184.

[145826]Osborne, T. F.: CREating a SCAP-less liver keeps SREBPs
pinned

[145827]in the ER membrane and prevents increased lipid synthe-
sis in response

[145828]to low cholesterol and high insulin. Genes Dev. 15:

1873–1878, 2001.

[145829]26185.

[145830]Shimano, H.; Shimomura, I.; Hammer, R. E.; Herz, J.; Goldstein,

[145831]J. L.; Brown, M. S.; Horton, J. D.: Elevated levels of SREBP-2 and

[145832]cholesterol synthesis in livers of mice homozygous for a targeted

[145833]disruption of the SREBP-1 gene. J. Clin. Invest. 100: 2115–2124,

[145834]1997.

[145835]26186.

[145836]Shimomura, I.; Hammer, R. E.; Richardson, J. A.; Ikemoto, S.; Bashmakov,

[145837]Y.; Goldstein, J. L.; Brown, M. S.: Insulin resistance and diabetes

[145838]mellitus in transgenic mice expressing nuclear SREBP-1c in adipose

[145839]tissue: model for congenital generalized lipodystrophy. Genes Dev. 12:

[145840]3182–3194, 1998.

[145841]26187.

[145842]Tobe, K.; Suzuki, R.; Aoyama, M.; Yamauchi, T.; Kamon, J.;
Kubota,
[145843]N.; Terauchi, Y.; Matsui, J.; Akanuma, Y.; Kimura, S.;
Tanaka, J.;
[145844]Abe, M.; Ohsumi, J.; Nagai, R.; Kadowaki, T.: Increased ex-
pression
[145845]of the sterol regulatory element-binding protein-1 gene
in insulin
[145846]receptor substrate-2 -/- mouse liver. J. Biol. Chem. 276:
38337-38340,
[145847]2001.
[145848]26188.
[145849]Wang, X.; Sato, R.; Brown, M. S.; Hua, X.; Goldstein, J. L.:
SREBP-1,
[145850]a membrane-bound transcription factor released by
sterol-regulated
[145851]proteolysis. Cell 77: 53-62, 1994.
[145852]26189.
[145853]Yokoyama, C.; Wang, X.; Briggs, M. R.; Admon, A.; Wu, J.;
Hua,
[145854]X.; Goldstein, J. L.; Brown, M. S.: SREBP-1, a basic-he-
lix-loop-helix-leucine

[145855]zipper protein that controls transcription of the low density lipoprotein

[145856]receptor gene. Cell 75: 187–197, 1993.

[145857]26190.

[145858]Achermann, J. C.; Ito, M.; Ito, M.; Hindmarsh, P. C.; Jameson,

[145859]J. L.: A mutation in the gene encoding steroidogenic factor-1 causes

[145860]XY sex reversal and adrenal failure in humans. (Letter) Nature Genet. 22:

[145861]125–126, 1999.

[145862]26191.

[145863]Biaison-Lauber, A.; Schoenle, E. J.: Apparently normal ovarian

[145864]differentiation in a prepubertal girl with transcriptionally inactive

[145865]steroidogenic factor 1 (NR5A1/SF-1) and adrenocortical insufficiency. Am.

[145866]J. Hum. Genet. 67: 1563–1568, 2000.

[145867]26192.

[145868]Bland, M. L.; Jamieson, C. A. M.; Akana, S. F.; Bornstein, S.

R.;

[145869]Eisenhofer, G.; Dallman, M. F.; Ingraham, H. A.: Haploinsufficiency

[145870]of steroidogenic factor-1 in mice disrupts adrenal development leading

[145871]to an impaired stress response. Proc. Nat. Acad. Sci. 97: 14488-14493,

[145872]2000.

[145873]26193.

[145874]Joosten, P. H. L. J.; Toepoel, M.; Mariman, E. C. M.; Van Zoelen,

[145875]E. J. J.: Promoter haplotype combinations of the platelet-derived

[145876]growth factor alpha-receptor gene predispose to human neural tube

[145877]defects. Nature Genet. 27: 215-217, 2001.

[145878]26194.

[145879]Mackenzie, P. I.; Owens, I. S.; Burchell, B.; Bock, K. W.; Bairoch,

[145880]A.; Belanger, A.; Fournel-Gigleux, S.; Green, M.; Hum, D. W.; Iyanagi,

[145881]T.; Lancet, D.; Louisot, P.; Magalou, J.; Roy Chowdhury, J.;

Ritter,

[145882]J. K.; Schachter, H.; Tephly, T. R.; Tipton, K. E.; Nebert, D. W.

[145883]: The UDP glycosyltransferase gene superfamily: recommended nomenclature

[145884]update based on evolutionary divergence. Pharmacogenetics 7: 255–269,

[145885]1997.

[145886]26195.

[145887]Corydon, M. J.; Vockley, J.; Rinaldo, P.; Rhead, W. J.; Kjeldsen,

[145888]M.; Winter, V.; Riggs, C.; Babovic-Vuksanovic, D.; Smeitink, J.; De

[145889]Jong, J.; Levy, H.; Sewell, A. C.; Roe, C.; Matern, D.; Dsouki, M.;

[145890]Gregersen, N.: Role of common gene variations in the molecular pathogenesis

[145891]of short-chain acyl-CoA dehydrogenase deficiency. Pediat. Res. 49:

[145892]18–23, 2001.

[145893]26196.

[145894]Colombo, R.: Age and origin of the PRNP E200K mutation

causing

[145895]familial Creutzfeldt–Jacob disease in Libyan Jews. (Letter)
Am. J.

[145896]Hum. Genet. 67: 528–531, 2000.

[145897]26197.

[145898]Cousens, J.; Smith, P. G.; Ward, H.; Everington, D.; Knight,
R.

[145899]S. G.; Zeidler, M.; Stewart, G.; Smith–Bathgate, E. A. B.;
Macleod,

[145900]M.–A.; Mackenzie, J.; Will, R. G.: Geographical distribution
of variant

[145901]Creutzfeldt–Jakob disease in Great Britain, 1994–2000.
Lancet 357:

[145902]1002–1007, 2001.

[145903]26198.

[145904]de Silva, R.; Ironside, J. W.; McCardle, L.; Esmonde, T.; Bell,

[145905]J.; Will, R.; Windl, O.; Dempster, M.; Estibeiro, P.; Lathe, R.:

[145906]Neuropathological phenotype and 'prion protein' genotype
correlation

[145907]in sporadic Creutzfeldt–Jakob disease. Neurosci. Lett. 179:
50–52,

[145908]1994.

[145909]26199.

[145910]Deslys, J.-P.; Jaegly, A.; d'Aignaux, J. H.; Mouthon, F.; de
Villemeur,

[145911]T. B.; Dormont, D.: Genotype at codon 129 and susceptibility to Creutzfeldt-Jakob

[145912]disease. Lancet 351: 1251 only, 1998.

[145913]26200.

[145914]Dlouhy, S. R.; Hsiao, K.; Farlow, M. F.; Foroud, T.; Con-
neally,

[145915]P. M.; Johnson, P.; Prusiner, S. B.; Hodes, M. E.; Ghetti, B.:
Linkage

[145916]of the Indiana kindred of Gerstmann-Straussler-Scheinker
disease to

[145917]the prion protein gene. Nature Genet. 1: 64-67, 1992.

[145918]26201.

[145919]Doh-ura, K.; Kitamoto, T.; Sakaki, Y.; Tateishi, J.: CJD dis-
crepancy.

[145920](Letter) Nature 353: 801-802, 1991.

[145921]26202.

[145922]Doh-ura, K.; Tateishi, J.; Sasaki, H.; Kitamoto, T.; Sakaki, Y.

[145923]Pro-to-leu change at position 102 of prion protein is the most common

[145924]but not the sole mutation related to Gerstmann-Straussler syndrome. Biochem.

[145925]Biophys. Res. Commun. 163: 974-979, 1989.

[145926]26203.

[145927]Duchen, L. W.; Poulter, M.; Harding, A. E.: Dementia associated

[145928]with a 216 base pair insertion in the prion protein gene: clinical

[145929]and neuropathological features. Brain 116: 555-567, 1993.

[145930]26204.

[145931]Erginel-Unaltuna, N.; Peoc'h, K.; Komurcu, E.; Acuner, T. T.;

[145932]Issever, H.; Laplanche, J.-L.: Distribution of the M129V polymorphism

[145933]of the prion protein gene in a Turkish population suggests a high

[145934]risk for Creutzfeldt-Jacob disease. Europ. J. Hum. Genet.

9: 965–968,

[145935]2001.

[145936]26205.

[145937]Farlow, M. R.; Yee, R. D.; Dlouhy, S. R.; Conneally, P. M.;
Azzarelli,

[145938]B.; Ghetti, B.: Gerstmann–Straussler–Scheinker disease. I.
Extending

[145939]the clinical spectrum. *Neurology* 39: 1446–1452, 1989.

[145940]26206.

[145941]Flechsig, E.; Shmerling, D.; Hegyi, I.; Raeber, A. J.; Fischer,

[145942]M.; Cozzio, A.; von Mering, C.; Aguzzi, A.; Weissmann, C.:
Prion

[145943]protein devoid of the octapeptide repeat region restores
susceptibility

[145944]to scrapie in PrP knockout mice. *Neuron* 27: 399–408,
2000.

[145945]26207.

[145946]Forloni, G.; Angeretti, N.; Chiesa, R.; Monzani, E.;
Salmona,

[145947]M.; Bugiani, O.: Neurotoxicity of a prion protein fragment.
(Letter) *Nature* 362:

[145948]543–546, 1993.

[145949]26208.

[145950]Gabizon, R.; Rosenmann, H.; Meiner, Z.; Kahana, I.; Kahana, E.;

[145951]Shugart, Y.; Ott, J.; Prusiner, S. B.: Mutation and polymorphism

[145952]of the prion protein gene in Libyan Jews with Creutzfeldt–Jakob disease

[145953](CJD). Am. J. Hum. Genet. 53: 828–835, 1993.

[145954]26209.

[145955]Gajdusek, D. C.: The transmissible amyloidoses: genetical control

[145956]of spontaneous generation of infectious amyloid proteins by nucleation

[145957]of configurational change in host precursors: kuru–CJD–GSS–scrapie–BSE. Europ.

[145958]J. Epidemiol. 7: 567–577, 1991.

[145959]26210.

[145960]Gambetti, P.; Petersen, R.; Monari, L.; Tabaton, M.; Autilio–Gambetti,

[145961]L.: Fatal familial insomnia and the widening spectrum of

prion diseases. Brit.

[145962]Med. Bull. 49: 980–994, 1993.

[145963]26211.

[145964]Ghetti, B.; Tagliavini, F.; Masters, C. L.; Beyreuther, K.; Gi-
accione,

[145965]G.; Verga, L.; Farlow, M. R.; Conneally, P. M.; Dlouhy, S. R.;
Azzarelli,

[145966]B.; Bugiani, O.: Gerstmann–Straussler–Scheinker disease.

II. Neurofibrillary

[145967]tangles and plaques with PrP-amyloid coexist in an af-
fected family. Neurology 39:

[145968]1453–1461, 1989.

[145969]26212.

[145970]Giaccone, G.; Verga, L.; Bugiani, O.; Frangione, B.; Serban,
D.;

[145971]Prusiner, S. B.; Farlow, M. R.; Ghetti, B.; Tagliavini, F.: Prion

[145972]protein preamyloid and amyloid deposits in Gerstmann–
Straussler–Scheinker

[145973]disease, Indiana kindred. Proc. Nat. Acad. Sci. 89:
9349–9353, 1992.

[145974]26213.

[145975]Goldfarb, L. G.; Brown, P.; Cervenakova, L.; Gajdusek, D.
C.:

[145976]Molecular genetic studies of Creutzfeldt-Jakob disease.
Molec. Neurobiol. 8:

[145977]89-97, 1994.

[145978]26214.

[145979]Goldfarb, L. G.; Brown, P.; Haltia, M.; Cathala, F.; McCom-
bie,

[145980]W. R.; Kovanen, J.; Cervenakova, L.; Goldin, L.; Nieto, A.;
Godec,

[145981]M. S.; Asher, D. M.; Gajdusek, D. C.: Creutzfeldt-Jakob
disease cosegregates

[145982]with the codon 178-asn PRNP mutation in families of Eu-
ropean origin. Ann.

[145983]Neurol. 31: 274-281, 1992.

[145984]26215.

[145985]Goldfarb, L. G.; Brown, P.; McCombie, W. R.; Goldgaber,
D.; Swergold,

[145986]G. D.; Wills, P. R.; Cervenakova, L.; Baron, H.; Gibbs, C. J.,
Jr.;

[145987]Gajdusek, D. C.: Transmissible familial Creutzfeldt-Jakob
disease

- [145988]associated with five, seven, and eight extra octapeptide coding repeats
- [145989]in the PRNP gene. Proc. Nat. Acad. Sci. 88: 10926–10930, 1991.
- [145990]26216.
- [145991]Goldfarb, L. G.; Brown, P.; Mitrova, E.; Cervenakova, L.; Goldin,
- [145992]L.; Korczyn, A. D.; Chapman, J.; Galvez, S.; Cartier, L.; Rubenstein,
- [145993]R.; Gajdusek, D. C.: Creutzfeldt–Jacob disease associated with the
- [145994]PRNP codon 200–lys mutation: an analysis of 45 families. Europ. J.
- [145995]Epidemiol. 7: 477–486, 1991.
- [145996]26217.
- [145997]Goldfarb, L. G.; Haltia, M.; Brown, P.; Nieto, A.; Kovanen, J.;
- [145998]McCombie, W. R.; Trapp, S.; Gajdusek, D. C.: New mutation in scrapie
- [145999]amyloid precursor gene (at codon 178) in Finnish Creutzfeldt–Jakob
- [146000]kindred. (Letter) Lancet 337: 425, 1991.

[146001]26218.

[146002]Goldfarb, L. G.; Mitrova, E.; Brown, P.; Toh, B. H.; Gaj-
dusek,

[146003]D. C.: Mutation in codon 200 of scrapie amyloid protein
gene in two

[146004]clusters of Creutzfeldt-Jakob disease in Slovakia. (Letter)
Lancet 336:

[146005]514-515, 1990.

[146006]26219.

[146007]Travis, G. H.; Sutcliffe, J. G.; Bok, D.: The retinal degenera-
tion

[146008]slow (rds) gene product is a photoreceptor disc mem-
brane-associated

[146009]glycoprotein. Neuron 6: 61-70, 1991.

[146010]26220.

[146011]Morris, C.; Heisterkamp, N.; Hao, Q. L.; Testa, J. R.; Grof-
fen,

[146012]J.: The human tyrosine kinase gene (FER) maps to chro-
mosome 5 and

[146013]is deleted in myeloid leukemias with a del(5q). Cytogenet.
Cell Genet. 53:

[146014]196–200, 1990.

[146015]26221.

[146016]Anderson, J.; Burns, H. D.; Enriquez–Harris, P.; Wilkie, A.
O.

[146017]M.; Heath, J. K.: Apert syndrome mutations in fibroblast
growth factor

[146018]receptor 2 exhibit increased affinity for FGF ligand. Hum.
Molec.

[146019]Genet. 7: 1475–1483, 1998.

[146020]26222.

[146021]Arman, E.; Haffner–Krausz, R.; Chen, Y.; Heath, J. K.;
Lonai, P.

[146022]: Targeted disruption of fibroblast growth factor (FGF) re-
ceptor 2

[146023]suggests a role for FGF signaling in pregastrulation mam-
malian development. Proc.

[146024]Nat. Acad. Sci. 95: 5082–5087, 1998.

[146025]26223.

[146026]Arman, E.; Haffner–Krausz, R.; Gorivodsky, M.; Lonai, P.:
Fgfr2

[146027]is required for limb outgrowth and lung–branching mor-

phogenesis. Proc.

[146028]Nat. Acad. Sci. 96: 11895–11899, 1999.

[146029]26224.

[146030]Chitayat, D.; Chun, K.: Reply to the letter to the editor by
Gripp

[146031]et al.--'not Antley–Bixler syndrome'. (Letter) Am. J. Med.
Genet. 83:

[146032]67–68, 1999.

[146033]26225.

[146034]Chun, K.; Siegel–Bartelt, J.; Chitayat, D.; Phillips, J.; Ray,

[146035]P. N.: FGFR2 mutation associated with clinical manifesta-
tions consistent

[146036]with Antley–Bixler syndrome. Am. J. Med. Genet. 77:
219–224, 1998.

[146037]26226.

[146038]Cohen, M. M., Jr.: Pfeiffer syndrome update, clinical sub-
types,

[146039]and guidelines for differential diagnosis. Am. J. Med.
Genet. 45:

[146040]300–307, 1993.

[146041]26227.

[146042]Cornejo-Roldan, L. R.; Roessler, E.; Muenke, M.: Analysis of the

[146043]mutational spectrum of the FGFR2 gene in Pfeiffer syndrome. Hum.

[146044]Genet. 104: 425-431, 1999.

[146045]26228.

[146046]den Dunnen, J. T.; Antonarakis, E.: Nomenclature for the description

[146047]of human sequence variations. Hum. Genet. 109: 121-124, 2001.

[146048]26229.

[146049]Deng, C.; Bedford, M.; Li, C.; Xu, X.; Yang, X.; Dunmore, J.;

[146050]Leder, P.: Fibroblast growth factor receptor-1 (FGFR-1) is essential

[146051]for normal neural tube and limb development. Dev. Biol. 185: 42-54,

[146052]1997.

[146053]26230.

[146054]Dionne, C. A.; Crumley, G.; Bellot, F.; Kaplow, J. M.; Searfoss,

[146055]G.; Ruta, M.; Burgess, W. H.; Jaye, M.; Schlessinger, J.:

Cloning

- [146056]and expression of two distinct high-affinity receptors cross-reacting
- [146057]with acidic and basic fibroblast growth factors. EMBO J. 9: 2685-2692,
- [146058]1990.
- [146059]26231.
- [146060]Dionne, C. A.; Modi, W. S.; Crumley, G.; O'Brien, S. J.; Schlessinger,
- [146061]J.; Jaye, M.: BEK, a receptor for multiple members of the fibroblast
- [146062]growth factor (FGF) family, maps to human chromosome 10q25.3-q26. Cytogenet.
- [146063]Cell Genet. 60: 34-36, 1992.
- [146064]26232.
- [146065]Gilbert, E.; Del Gatto, F.; Champion-Arnaud, P.; Gesnel, M.-C.;
- [146066]Breathnach, R.: Control of BEK and K-SAM splice sites in alternative
- [146067]splicing of the fibroblast growth factor receptor 2 pre-mRNA. Molec.
- [146068]Cell. Biol. 13: 5461-5468, 1993.

[146069]26233.

[146070]Gorlin, R. J.: Patient described by Chun et al. may not
present

[146071]Antley-Bixler syndrome. (Letter) Am. J. Med. Genet. 83: 64
only,

[146072]1999.

[146073]26234.

[146074]Gorry, M. C.; Preston, R. A.; White, G. J.; Zhang, Y.; Sing-
hal,

[146075]V. K.; Losken, H. W.; Parker, M. G.; Nwokoro, N. A.; Post, J.
C.;

[146076]Ehrlich, G. D.: Crouzon syndrome: mutations in two
spliceoforms of

[146077]FGFR2 and a common point mutation shared with Jack-
son-Weiss syndrome. Hum.

[146078]Molec. Genet. 4: 1387-1390, 1995.

[146079]26235.

[146080]Gripp, K. W.; Stolle, C. A.; McDonald-McGinn, D. M.;
Markowitz,

[146081]R. I.; Bartlett, S. P.; Katowitz, J. A.; Muenke, M.; Zackai, E.
H.

- [146082]: Phenotype of the fibroblast growth factor receptor 2
ser351-to-cys
[146083]mutation: Pfeiffer syndrome type III. Am. J. Med. Genet.
78: 356-360,
[146084]1998.
[146085]26236.
[146086]Gripp, K. W.; Zackai, E. H.; Cohen, M. M., Jr.: Not Antley-
Bixler
[146087]syndrome. (Letter) Am. J. Med. Genet. 83: 65-66, 1999.
[146088]26237.
[146089]Hajihosseini, M. K.; Wilson, S.; De Moerlooze, L.; Dickson,
C.
[146090]: A splicing switch and gain-of-function mutation in
Fgfr2-IIIc hemizygotes
[146091]causes Apert/Pfeiffer-syndrome-like phenotypes. Proc.
Nat. Acad.
[146092]Sci. 98: 3855-3860, 2001.
[146093]26238.
[146094]Hollway, G. E.; Suthers, G. K.; Haan, E. A.; Thompson, E.;
David,
[146095]D. J.; Gecz, J.; Mulley, J. C.: Mutation detection in FGFR2

craniosynostosis

[146096]syndromes. Hum. Genet. 99: 251–255, 1997.

[146097]26239.

[146098]Houssaint, E.; Blanquet, P. R.; Champion–Arnaud, P.; Gesnel, M.

[146099]C.; Torriglia, A.; Courtois, Y.; Breathnach, R.: Related fibroblast

[146100]growth factor receptor genes exist in the human genome. Proc. Nat.

[146101]Acad. Sci. 87: 8180–8184, 1990.

[146102]26240.

[146103]Ibrahimi, O. A.; Eliseenkova, A. V.; Plotnikov, A. N.; Yu, K.;

[146104]Ornitz, D. M.; Mohammadi, M.: Structural basis for fibroblast growth

[146105]factor receptor 2 activation in Apert syndrome. Proc. Nat. Acad.

[146106]Sci. 98: 7182–7187, 2001.

[146107]26241.

[146108]Jackson, R.: The lines of Blaschko: a review and reconsideration. Brit.

[146109]J. Derm. 95: 349–360, 1976.

[146110]26242.

[146111]Johnson, D.; Wall, S. A.; Mann, S.; Wilkie, A. O. M.: A novel
[146112]mutation, ala315 to ser, in FGFR2: a gene-environment
interaction
[146113]leading to craniosynostosis? Europ. J. Hum. Genet. 8:
571-577, 2000.

[146114]26243.

[146115]Williams, T. M.; Montoya, G.; Wu, Y.; Eddy, R. L.; Byers, M.
G.;

[146116]Shows, T. B.: The TCF8 gene encoding a zinc finger pro-
tein (Nil-2-a)
[146117]resides on human chromosome 10p11.2. Genomics 14:
194-196, 1992.

[146118]26244.

[146119]van Nie, R.; Ivanyi, D.; Demant, P.: A new H-2-linked mu-
tation,

[146120]rds, causing retinal degeneration in the mouse. Tissue
Antigens 12:

[146121]106-108, 1978.

[146122]26245.

[146123]Vine, A. K.; Schatz, H.: Adult-onset foveomacular pigment

epithelial

[146124] dystrophy. Am. J. Ophthal. 89: 680–691, 1980.

[146125] 26246.

[146126] Weleber, R. G.; Carr, R. E.; Murphey, W. H.; Sheffield, V. C.;

[146127] Stone, E. M.: Phenotypic variation including retinitis pigmentosa,

[146128] pattern dystrophy, and fundus flavimaculatus in a single family with

[146129] a deletion of codon 153 or 154 of the peripherin/RDS gene. Arch.

[146130] Ophthal. 111: 1531–1542, 1993.

[146131] 26247.

[146132] Wells, J.; Wroblewski, J.; Keen, J.; Inglehearn, C.; Jubb, C.;

[146133] Eckstein, A.; Jay, M.; Arden, G.; Bhattacharya, S.; Fitzke, F.; Bird,

[146134] A.: Mutations in the human retinal degeneration slow (RDS) gene can

[146135] cause either retinitis pigmentosa or macular dystrophy.

Nature Genet. 3:

[146136] 213–218, 1993.

[146137] 26248.

- [146138] Wroblewski, J. J.; Wells, J. A., III; Eckstein, A.; Fitzke, F.;
- [146139] Jubb, C.; Keen, J.; Inglehearn, C.; Bhattacharya, S.; Arden, G. B.;
- [146140] Jay, M.; Bird, A. C.: Macular dystrophy associated with mutations
- [146141] at codon 172 in the human retinal degeneration slow gene. *Ophthalmology* 101:
- [146142] 12–22, 1994.
- [146143] 26249.
- [146144] Eph Nomenclature Committee: Unified nomenclature for the Eph
- [146145] family receptors and their ligands, the ephrins. *Cell* 90: 403–404,
- [146146] 1997.
- [146147] 26250.
- [146148] Hattori, M.; Osterfield, M.; Flanagan, J. G.: Regulated cleavage
- [146149] of a contact-mediated axon repellent. *Science* 289: 1360–1365, 2000.
- [146150] 26251.
- [146151] Maru, Y.; Hirai, H.; Yoshida, M. C.; Takaku, F.: Evolution,

expression,

[146152]and chromosomal location of a novel receptor tyrosine kinase gene,

[146153]eph. Molec. Cell. Biol. 8: 3770–3776, 1988.

[146154]26252.

[146155]Yoshida, M. C.; Maru, H.; Hirai, H.; Takau, F.: Chromosomal location

[146156]of a novel receptor tyrosine kinase gene, EPH, on chromosome 7. (Abstract) Cytogenet.

[146157]Cell Genet. 51: 1113 only, 1989.

[146158]26253.

[146159]Boyd, A. W.; Ward, L. D.; Wicks, I. P.; Simpson, R. J.; Salvaris,

[146160]E.; Wilks, A.; Welch, K.; Loudovaris, M.; Rockman, S.; Busmanis, I.

[146161]: Isolation and characterization of a novel receptor-type protein

[146162]tyrosine kinase (hek) from a human pre-B cell line. J. Biol. Chem. 267:

[146163]3262–3267, 1992.

[146164]26254.

[146165]Fox, G. M.; Holst, P. L.; Chute, H. T.; Lindberg, R. A.;
Janssen,
[146166]A. M.; Basu, R.; Welcher, A. A.: cDNA cloning and tissue
distribution
[146167]of five human EPH-like receptor protein-tyrosine kinases.
Oncogene 10:
[146168]897-905, 1995.
[146169]26255.
[146170]Wicks, I. P.; Lapsys, N. M.; Baker, E.; Campbell, L. J.; Boyd,
[146171]A. W.; Sutherland, G. R.: Localization of a human receptor
tyrosine
[146172]kinase (ETK1) to chromosome region 3p11.2. Genomics
19: 38-41, 1994.
[146173]26256.
[146174]Wicks, I. P.; Wilkinson, D.; Salvaris, E.; Boyd, A. W.: Molec-
ular
[146175]cloning of HEK, the gene encoding a receptor tyrosine ki-
nase expressed
[146176]by human lymphoid tumor cell lines. Proc. Nat. Acad. Sci.
89: 1611-1615,
[146177]1992.

[146178]26257.

[146179]Agrawal, A.; Eastman, Q. M.; Schatz, D. G.: Transposition mediated

[146180]by RAG1 and RAG2 and its implications for the evolution of the immune

[146181]system. Nature 394: 744–751, 1998.

[146182]26258.

[146183]Blanquet, V.; Turleau, C.; Goossens, M.; Besmond, C.: Assignment

[146184]of the human recombination activating gene 1 (RAG1) to the 14q21.3–q22.2

[146185]region. Genomics 13: 488–489, 1992.

[146186]26259.

[146187]Fulop, G. M.; Phillips, R. A.: The scid mutation in mice causes

[146188]a general defect in DNA repair. Nature 347: 479–482, 1990.

[146189]26260.

[146190]Hikida, M.; Mori, M.; Takai, T.; Tomochika, K.; Hamatani, K.; Ohmori,

[146191]H.: Reexpression of RAG–1 and RAG–2 genes in activated

mature mouse

[146192]B cells. Science 274: 2091–2093, 1996.

[146193]26261.

[146194]Hiom, K.; Melek, M.; Gellert, M.: DNA transposition by the
RAG1

[146195]and RAG2 proteins: a possible source of oncogenic
translocations. Cell 94:

[146196]463–470, 1998.

[146197]26262.

[146198]Huye, L. E.; Purugganan, M. M.; Jiang, M.-M.; Roth, D. B.:
Mutational

[146199]analysis of all conserved basic amino acids in RAG–1 re-
veals catalytic,

[146200]step arrest, and joining-deficient mutants in the V(D)J re-
combinase. Molec.

[146201]Cell. Biol. 22: 3460–3473, 2002.

[146202]26263.

[146203]Ichihara, Y.; Hirai, M.; Kurosawa, Y.: Sequence and chro-
mosome

[146204]assignment to 11p13–p12 of human RAG genes. Immun.
Lett. 33: 277–284,

[146205]1992.

[146206]26264.

[146207]Janeway, C. A., Jr.: How the immune system works to protect the

[146208]host from infection: a personal view. Proc. Nat. Acad. Sci. 98:

[146209]7461–7468, 2001.

[146210]26265.

[146211]Kim, D. R.; Dai, Y.; Mundy, C. L.; Yang, W.; Oettinger, M. A.:

[146212]Mutations of acidic residues in RAG1 define the active site of the

[146213]V(D)J recombinase. Genes Dev. 13: 3070–3080, 1999.

[146214]26266.

[146215]Landree, M. A.; Wibbenmeyer, J. A.; Roth, D. B.: Mutational analysis

[146216]of RAG1 and RAG2 identifies three catalytic amino acids in RAG1 critical

[146217]for both cleavage steps of V(D)J recombination. Genes Dev. 13: 3059–3069,

[146218]1999.

[146219]26267.

[146220]McBlane, J. F.; van Gent, D. C.; Ramsden, D. A.; Romeo, C.;
Cuomo,

[146221]C. A.; Gellert, M.; Oettinger, M. A.: Cleavage at a V(D)J re-
combination

[146222]signal requires only RAG1 and RAG2 proteins and occurs
in two steps. Cell 83:

[146223]387–395, 1995.

[146224]26268.

[146225]Melek, M.; Gellert, M.: RAG1/2-mediated resolution of
transposition

[146226]intermediates: two pathways and possible consequences.
Cell 101:

[146227]625–633, 2000.

[146228]26269.

[146229]Mombaerts, P.; Iacomini, J.; Johnson, R. S.; Herrup, K.;
Tonegawa,

[146230]S.; Papaioannou, V. E.: RAG-1-deficient mice have no ma-
ture B and

[146231]T lymphocytes. Cell 68: 869–877, 1992.

[146232]26270.

[146233]Oettinger, M. A.: Personal Communication. Boston, Mass.
1/15/1993.

[146234]26271.

[146235]Sillekens, P. T. G.; Habets, W. J.; Beijer, R. P.; van Venrooij,

[146236]W. J.: cDNA cloning of the human U1 snRNA-associated A
protein: extensive

[146237]homology between U1 and U2 snRNP-specific proteins.
EMBO J. 6: 3841-3848,

[146238]1987.

[146239]26272.

[146240]Koyama, K.; Fukushima, Y.; Inazawa, J.; Tomotsune, D.;
Takahashi,

[146241]N.; Nakamura, Y.: The human homologue of the murine
Llg1h gene (LLGL)

[146242]maps within the Smith-Magenis syndrome region in
17p11.2. Cytogenet.

[146243]Cell Genet. 72: 78-82, 1996.

[146244]26273.

[146245]Forsberg, E.; Hirsch, E.; Frohlich, L.; Meyer, M.; Ekblom, P.;

[146246]Aszodi, A.; Werner, S.; Fassler, R.: Skin wounds and sev-
ered nerves

[146247] heal normally in mice lacking tenascin-C. Proc. Nat. Acad. Sci. 93:

[146248] 6594-6599, 1996.

[146249] 26274.

[146250] Gulcher, J. R.; Alexakos, M. J.; Le Beau, M. M.; Lemons, R. S.;

[146251] Stefansson, K.: Chromosomal localization of the human hexabrachion

[146252] (tenascin) gene and evidence for recent reduplication within the gene. Genomics 6:

[146253] 616-622, 1990.

[146254] 26275.

[146255] Gulcher, J. R.; Nies, D. E.; Alexakos, M. J.; Ravikant, N. A.;

[146256] Sturgill, M. E.; Marton, L. S.; Stefansson, K.: Structure of the

[146257] human hexabrachion (tenascin) gene. Proc. Nat. Acad. Sci. 88: 9438-9442,

[146258] 1991.

[146259] 26276.

[146260] Mitrovic, N.; Schachner, M.: Detection of tenascin-C in the nervous

[146261]system of the tenascin-C mutant mouse. J. Neurosci. Res.
42: 710-717,

[146262]1995.

[146263]26277.

[146264]Nies, D. E.; Hemesath, T. J.; Kim, J.-H.; Gulcher, J. R.; Ste-
fansson,

[146265]K.: The complete cDNA sequence of human hexabrachion
(tenascin):

[146266]a multidomain protein containing unique epidermal
growth factor repeats. J.

[146267]Biol. Chem. 266: 2818-2823, 1991.

[146268]26278.

[146269]Williams, T. M.; Moolten, D.; Burlein, J.; Romano, J.; Bhaer-
man,

[146270]R.; Godillot, A.; Mellon, M.; Rauscher, F. J., III; Kant, J. A.:

[146271]Identification of a zinc finger protein that inhibits IL-2
gene expression. Science 254:

[146272]1791-1794, 1991.

[146273]26279.

[146274]Ludecke, H.-J.; Schaper, J.; Meinecke, P.; Momeni, P.;
Gross, S.;

[146275] von Holtum, D.; Hirche, H.; Abramowicz, M. J.; Albrecht, B.; Apacik, C.; Christen, H.-J.; Claussen, U.; and 28 others: Genotypic and phenotypic spectrum in tricho-rhino-phalangeal syndrome types I and III. *Am. J. Hum. Genet.* 68: 81-91, 2001.

[146279] 26280.

[146280] Arya, R.; Lalloz, M. R. A.; Bellingham, A. J.; Layton, D. M.: Evidence for founder effect of the glu104-to-aspartic acid substitution and identification of new mutations in triosephosphate isomerase deficiency. *Hum. Mutat.* 10: 290-294, 1997.

[146284] 26281.

[146285] Asakawa, J.; Iida, S.: Origin of human triosephosphate isomerase isozymes: further evidence for the single structural locus hypothesis with Japanese variants. *Hum. Genet.* 71: 22-26, 1985.

[146288] 26282.

- [146289] McDonald, H.; Smailus, D.; Jenkins, H.; Adams, K.; Simpson, N.
- [146290] E.; Goodfellow, P. J.: Identification and characterization of a gene
- [146291] at D10S94 in the MEN2A region. Genomics 13: 344–348, 1992.
- [146292] 26283.
- [146293] Miwa, S.; Sato, T.; Murao, H.; Kozuru, M.; Ibayashi, H.: A new
- [146294] type of phosphofructokinase deficiency: hereditary non-spherocytic
- [146295] hemolytic anemia. Acta Haemat. Jpn. 35: 113–118, 1972.
- [146296] 26284.
- [146297] Vora, S.; DiMauro, S.; Spear, D.; Harker, D.; Danon, M. J.: Characterization
- [146298] of the enzymatic defect in late-onset muscle phosphofructokinase deficiency:
- [146299] new subtype of glycogen storage disease type VII. J. Clin. Invest. 80:
- [146300] 1479–1485, 1987.
- [146301] 26285.

- [146302]Bernardi, G.: Personal Communication. Paris, France
5/13/1992.
- [146303]26286.
- [146304]Boulard, M. R.; Bois, M.; Reviron, M.; Najean, Y.: Red-cell
phosphofructokinase
[146305]deficiency. New Eng. J. Med. 291: 978-979, 1974.
- [146306]26287.
- [146307]Chadefaux, B.; Rethore, M. O.; Allard, D.: Regional map-
ping of
[146308]liver type 6-phosphofructokinase isoenzyme on chromo-
some 21. Hum.
[146309]Genet. 68: 136-137, 1984.
- [146310]26288.
- [146311]Cox, D. R.; Kawashima, H.; Vora, S.; Epstein, C. J.: Regional
[146312]mapping of SOD-1, PRGS, and PFK-L on human chromo-
some 21. (Abstract) Cytogenet.
- [146313]Cell Genet. 37: 441-442, 1984.
- [146314]26289.
- [146315]Elson, A.; Levanon, D.; Brandeis, M.; Dafni, N.; Bernstein,
Y.;

- [146316]Danciger, E.; Groner, Y.: The structure of the human liver-type phosphofructokinase
- [146317]gene. Genomics 7: 47–56, 1990.
- [146318]26290.
- [146319]Etiemble, J.; Kahn, A.; Boivin, P.; Bernard, J. F.; Goude-
mand,
- [146320]M.: Hereditary hemolytic anemia with erythrocyte phosphofructokinase
- [146321]deficiency. Hum. Genet. 31: 83–91, 1976.
- [146322]26291.
- [146323]Etiemble, J.; Picat, C.; Simeon, J.; Blatrix, C.; Boivin, P.:
- [146324]Inherited erythrocyte phosphofructokinase deficiency:
molecular mechanism. Hum.
- [146325]Genet. 55: 383–390, 1980.
- [146326]26292.
- [146327]Gehrich, S. C.; Gekakis, N.; Sul, H. S.: Liver (B-type)
phosphofructokinase
- [146328]mRNA: cloning, structure, and expression. J. Biol. Chem.
263: 11755–11759,
- [146329]1988.

[146330]26293.

[146331]Kahn, A.; Etiemble, J.; Meienhofer, M. C.; Boivin, P.: Erythrocyte

[146332]phosphofructokinase deficiency associated with an unstable variant

[146333]of muscle phosphofructokinase. Clin. Chim. Acta 61: 415-419, 1975.

[146334]26294.

[146335]Levanon, D.; Danciger, E.; Dafni, N.; Groner, Y.: Genomic clones

[146336]of the human liver-type phosphofructokinase. Biochem. Biophys. Res.

[146337]Commun. 141: 374-380, 1986.

[146338]26295.

[146339]Pantelakis, S. N.; Karaklis, A. G.; Alexiou, D.; Vardas, E.; Valaes,

[146340]T.: Red cell enzymes in trisomy 21. Am. J. Hum. Genet. 22: 184-193,

[146341]1970.

[146342]26296.

[146343]Van Keuren, M.; Drabkin, H.; Hart, I.; Harker, D.; Patter-

son,

[146344]D.; Vora, S.: Regional assignment of human liver-type
6-phosphofructokinase

[146345]to chromosome 21q22.3 by using somatic cell hybrids
and a monoclonal

[146346]anti-L antibody. Hum. Genet. 74: 34-40, 1986.

[146347]26297.

[146348]Vora, S.; Davidson, M.; Seaman, C.; Miranda, A. F.; Noble,
N.

[146349]A.; Tanaka, K. R.; Frenkel, E. P.; DiMauro, S.: Heterogeneity
of

[146350]the molecular lesions in inherited phosphofructokinase
deficiency. J.

[146351]Clin. Invest. 72: 1995-2006, 1983.

[146352]26298.

[146353]Vora, S.; Durham, S.; de Martinville, B.; Francke, U.: As-
signment

[146354]of the genes for liver type phosphofructokinase (PFKL) to
chromosome

[146355]21 and for muscle type (PFKM) to region p32-q32 of
chromosome 1. (Abstract) Cytogenet.

[146356]Cell Genet. 32: 324 only, 1982.

[146357]26299.

[146358]Vora, S.; Francke, U.: Assignment of the human gene for
liver

[146359]type phosphofructokinase isozyme to chromosome 21 us-
ing somatic cell

[146360]hybrids. (Abstract) Pediat. Res. 15: 570 only, 1981.

[146361]26300.

[146362]Vora, S.; Francke, U.: Assignment of the human gene for
liver-type

[146363]6-phosphofructokinase isozyme (PFKL) to chromosome 21
by using somatic

[146364]cell hybrids and monoclonal anti-L antibody. Proc. Nat.
Acad. Sci. 78:

[146365]3738-3742, 1981.

[146366]26301.

[146367]Wang, D.; Fang, H.; Cantor, C. R.; Smith, C. L.: A contigu-
ous

[146368]NotI restriction map of band q22.3 of human chromosome
21. Proc.

[146369]Nat. Acad. Sci. 89: 3222-3226, 1992.

[146370]26302.

[146371]Vidal, R.; Revesz, T.; Rostagno, A.; Kim, E.; Holton, J. L.;
Bek,

[146372]T.; Bojsen-Moller, M.; Braendgaard, H.; Plant, G.; Ghiso, J.;
Frangione,

[146373]B.: A decamer duplication in the 3-prime region of the BRI
gene originates

[146374]an amyloid peptide that is associated with dementia in a
Danish kindred. Proc.

[146375]Nat. Acad. Sci. 97: 4920-4925, 2000.

[146376]26303.

[146377]Fathallah-Shaykh, H.; Wolf, S.; Wong, E.; Posner, J. B.:
Cloning

[146378]of a leucine-zipper protein recognized by the sera of pa-
tients with

[146379]antibody-associated paraneoplastic cerebellar degenera-
tion. Proc.

[146380]Nat. Acad. Sci. 88: 3451-3454, 1991.

[146381]26304.

[146382]Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.; Posner, J.
B.;

[146383]Siniscalco, M.: In situ mapping of the gene coding for a
leucine

[146384] zipper DNA binding protein (CDR 2) to the region between
two rare

[146385] fragile sites of autosome 16 (16p12–p13.1). (Abstract)
Cytogenet.

[146386] Cell Genet. 58: 1999–2000, 1991.

[146387] 26305.

[146388] Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.; Posner, J.
B.;

[146389] Siniscalco, M.: In situ mapping of the gene coding for a
leucine

[146390] zipper DNA binding protein (CDR62) to 16p12–16p13.1.
Genomics 13:

[146391] 1340–1342, 1992.

[146392] 26306.

[146393] Raimondi, E.; Rubboli, F.; Moralli, D.; Chini, B.; Fornasari,
D.;

[146394] Tarroni, P.; De Carli, L.; Clementi, F.: Chromosomal local-
ization

[146395] and physical linkage of the genes encoding the human al-
pha-3, alpha-5,

[146396] and beta-4 neuronal nicotinic receptor subunits. Ge-
nomics 12: 849–850,

[146397]1992.

[146398]26307.

[146399]Anand, R.; Lindstrom, J.: Nucleotide sequence of the human nicotinic

[146400]acetylcholine receptor beta-2 subunit gene. Nucleic Acids Res. 18:

[146401]4272, 1990.

[146402]26308.

[146403]De Fusco, M.; Becchetti, A.; Patrignani, A.; Annesi, G.; Gambardella,

[146404]A.; Quattrone, A.; Ballabio, A.; Wanke, E.; Casari, G.: The nicotinic

[146405]receptor beta-2 subunit is mutant in nocturnal frontal lobe epilepsy. Nature

[146406]Genet. 26: 275-276, 2000.

[146407]26309.

[146408]Lueders, K. K.; Elliott, R. W.; Marenholz, I.; Mischke, D.; DuPree,

[146409]M.; Hamer, D.: Genomic organization and mapping of the human and

[146410]mouse neuronal beta-1-nicotinic acetylcholine receptor

genes. Mammalian

[146411]Genome 10: 900–905, 1999.

[146412]26310.

[146413]Phillips, H. A.; Favre, I.; Kirkpatrick, M.; Zuberi, S. M.;
Goudie,

[146414]D.; Heron, S. E.; Scheffer, I. E.; Sutherland, G. R.; Berkovic,
S.

[146415]F.; Bertrand, D.; Mulley, J. C.: CHRN2 is the second
acetylcholine

[146416]receptor subunit associated with autosomal dominant
nocturnal frontal

[146417]lobe epilepsy. Am. J. Hum. Genet. 68: 225–231, 2001.

[146418]26311.

[146419]Picciotto, M. R.; Zoli, M.; Lena, C.; Bessis, A.; Lallemant,
Y.;

[146420]LeNovere, N.; Vincent, P.; Pich, E. M.; Brulet, P.; Changeux,
J.–P.

[146421]: Abnormal avoidance learning in mice lacking functional
high-affinity

[146422]nicotine receptor in the brain. Nature 374: 65–67, 1995.

[146423]26312.

[146424]Zoli, M.; Picciotto, M. R.; Ferrari, R.; Cocchi, D.; Changeux,
[146425]J.-P.: Increased neurodegeneration during ageing in mice
lacking
[146426]high-affinity nicotine receptors. EMBO J. 18: 1235-1244,
1999.
[146427]26313.

[146428]Tarroni, P.; Rubboli, F.; Chini, B.; Zwart, R.; Oortgiesen, M.;
[146429]Sher, E.; Clementi, F.: Neuronal-type nicotinic receptors in
human
[146430]neuroblastoma and small-cell lung carcinoma cell lines.
FEBS Lett. 312:
[146431]66-70, 1992.
[146432]26314.

[146433]Bonner, T. I.; Buckley, N. J.; Young, A. C.; Brann, M. R.:
Identification
[146434]of a family of muscarinic acetylcholine receptor genes.
Science 237:
[146435]527-532, 1987.
[146436]26315.

[146437]Eng, C. E. L.; Strom, C. M.: Analysis of three restriction
fragment

[146438]length polymorphisms in the human type II procollagen gene. Am. J.

[146439]Hum. Genet. 37: 719–732, 1985.

[146440]26316.

[146441]Eyre, D. R.; Weis, M. A.; Moskowitz, R. W.: Cartilage expression

[146442]of a type II collagen mutation in an inherited form of osteoarthritis

[146443]associated with a mild chondrodysplasia. J. Clin. Invest. 87: 357–361,

[146444]1991.

[146445]26317.

[146446]Francomano, C. A.; Liberfarb, R. M.; Hirose, T.; Maumenee, I.

[146447]H.; Streeten, E. A.; Meyers, D. A.; Pyeritz, R. E.: The Stickler

[146448]syndrome: evidence for close linkage to the structural gene for type

[146449]II collagen. Genomics 1: 293–296, 1987.

[146450]26318.

[146451]Francomano, C. A.; Maumenee, I.; Liberfarb, R.; Pyeritz, R. E.

[146452]: Cosegregation of Stickler syndrome and type II collagen gene alleles.

[146453](Abstract) Cytogenet. Cell Genet. 46: 615 only, 1987.

[146454]26319.

[146455]Garofalo, S.; Vuorio, E.; Metsaranta, M.; Rosati, R.; Toman, D.;

[146456]Vaughan, J.; Lozano, G.; Mayne, R.; Ellard, J.; Horton, W.; de Crombrughe,

[146457]B.: Reduced amounts of cartilage collagen fibrils and growth plate

[146458]anomalies in transgenic mice harboring a glycine-to-cysteine mutation

[146459]in the mouse type II procollagen alpha-1-chain gene. Proc. Nat. Acad.

[146460]Sci. 88: 9648-9652, 1991.

[146461]26320.

[146462]Gupta, S. K.; Leonard, B. C.; Damji, K. F.; Bulman, D. E.: A

[146463]frame shift mutation in a tissue-specific alternatively spliced exon

[146464]of collagen 2A1 in Wagner's vitreoretinal degeneration. Am. J. Ophthal. 133:

[146465]203–210, 2002.

[146466]26321.

[146467]Holderbaum, D.; Malemud, C. J.; Moskowitz, R. W.; Haqqi, T. M.

[146468]: Human cartilage from late stage familial osteoarthritis transcribes

[146469]type II collagen mRNA encoding a cysteine in position 519. Biochem.

[146470]Biophys. Res. Commun. 192: 1169–1174, 1993.

[146471]26322.

[146472]Horton, W. A.: Personal Communication. Houston, Texas 1987.

[146473]26323.

[146474]Horton, W. A.; Machado, M. A.; Ellard, J.; Campbell, D.; Bartley,

[146475]J.; Ramirez, F.; Vitale, E.; Lee, B.: Characterization of a type

[146476]II collagen gene (COL2A1) mutation identified in cultured chondrocytes

[146477]from human hypochondrogenesis. Proc. Nat. Acad. Sci. 89: 4583–4587,

[146478]1992.

[146479]26324.

[146480]Huerre-Jeanpierre, C.; Mattei, M.-G.; Weil, D.; Grzeschik, K.

[146481]H.; Chu, M.-L.; Sangiorgi, F. O.; Sobel, M. E.; Ramirez, F.; Junien,

[146482]C.: Further evidence for the dispersion of the human fibrillar collagen

[146483]genes. Am. J. Hum. Genet. 38: 26-37, 1986.

[146484]26325.

[146485]Huerre-Jeanpierre, C.; Mattei, M. G.; Weil, D.; Grzeschik, K.

[146486]H.; Sangiorgi, F. O.; Ramirez, F.; Junien, C.: The gene for human

[146487]alpha-1 type II collagen (COL2A1) maps to band 12q131. (Abstract) Cytogenet.

[146488]Cell Genet. 40: 657-658, 1985.

[146489]26326.

[146490]Kaitila, I.; Korokko, J.; Marttinen, E.; Ala-Kokko, L.: Phenotypic

[146491]expressions of a gly154-to-arg mutation in type II collagen in two

[146492]unrelated patients with spondyloepimetaphyseal dysplasia

(SEMD). Am.

[146493]J. Med. Genet. 63: 111-122, 1996.

[146494]26327.

[146495]Kaprio, J.; Kujala, U. M.; Peltonen, L.; Koskenvuo, M.: Genetic

[146496]liability to osteoarthritis may be greater in women than in men. (Letter) Brit.

[146497]Med. J. 313: 232 only, 1996.

[146498]26328.

[146499]Kelly, T. E.; Lichtenstein, J. R.; Dorst, J. P.: An unusual familial

[146500]spondyloepiphyseal dysplasia: 'spondyloperipheral dysplasia.'. Birth

[146501]Defects Orig. Art. Ser. XIII(3B): 149-165, 1977.

[146502]26329.

[146503]Knowlton, R. G.; Katzenstein, P.; Moskowitz, R. W.; Weaver, E.

[146504]J.; Jimenez, S. A.; Pathria, M. N.; Malesud, C. J.; Prockop, D. J.

[146505]: Genetic linkage of the type II procollagen gene to primary generalized

[146506]osteoarthritis with chondrodysplasia. (Abstract) Cyto-
genet. Cell

[146507]Genet. 51: 1024 only, 1989.

[146508]26330.

[146509]Knowlton, R. G.; Katzenstein, P. L.; Moskowitz, R. W.;
Weaver,

[146510]E. J.; Malemud, C. J.; Pathria, M. N.; Jimenez, S. A.;
Prockop, D.

[146511]J.: Genetic linkage of a polymorphism in the type II pro-
collagen

[146512]gene (COL2A1) to primary osteoarthritis associated with
mild chondrodysplasia. New

[146513]Eng. J. Med. 322: 526-530, 1990.

[146514]26331.

[146515]Korkko, J.; Cohn, D. H.; Ala-Kokko, L.; Krakow, D.;
Prockop, D.

[146516]J.: Widely distributed mutations in the COL2A1 gene pro-
duce achondrogenesis

[146517]type II/hypochondrogenesis. Am. J. Med. Genet. 92:
95-100, 2000.

[146518]26332.

[146519]Law, M. L.; Tung, L.; Morse, H. G.; Berger, R.; Jones, C.;
Cheah,

[146520]K. S. E.; Solomon, E.: The human type II collagen gene
(COL2A1) assigned

[146521]to 12q14.3. Ann. Hum. Genet. 50: 131–137, 1986.

[146522]26333.

[146523]Law, M. L.; Tung, L.; Morse, H. G.; Berger, R.; Jones, C.;
Cheah,

[146524]K. S. E.; Stoker, N. G.; Solomon, E.: Regional assignment of
the

[146525]human type II collagen gene (COL2A1) on chromosome
12. (Abstract) Cytogenet.

[146526]Cell Genet. 40: 678 only, 1985.

[146527]26334.

[146528]Learmonth, I. D.; Christy, G.; Beighton, P.: Namaqualand
hip

[146529]dysplasia: orthopaedic implications. Clin. Orthop. 218:
142–147,

[146530]1987.

[146531]26335.

[146532]Li, S. W.; Prockop, D. J.; Helminen, H.; Fassler, R.; Lapvete–

lainen,

[146533]T.; Kiraly, K.; Peltarri, A.; Arokoski, J.; Lui, H.; Arita, M.;

Khillan,

[146534]J. S.: Transgenic mice with targeted inactivation of the

Col2a1 gene

[146535]for collagen II develop a skeleton with membranous and

periosteal

[146536]bone but no endochondral bone. Genes Dev. 9:

2821–2830, 1995.

[146537]26336.

[146538]Lovell–Badge, R. H.; Bygrave, A.; Bradley, A.; Robertson, E.;

[146539]Tilly, R.; Cheah, K. S. E.: Tissue–specific expression of the

human

[146540]type II collagen gene in mice. Proc. Nat. Acad. Sci. 84:

2803–2807,

[146541]1987.

[146542]26337.

[146543]Schwenk, W. F.; Rizza, R. A.; Mandarino, L. J.; Gerich, J. E.;

[146544]Hayles, A. B.; Haymond, M. W.: Familial insulin resistance

and acanthosis

[146545]nigricans: presence of a postbinding defect. Diabetes 35:

33–37,

[146546]1986.

[146547]26338.

[146548]Harris, E. L.: Personal Communication. Dunedin, New Zealand

[146549]11/27/2001.

[146550]26339.

[146551]Rossaak, J. I.; van Rij, A. M.; Jones, G. T.; Harris, E. L.:

[146552]Association of the 4G/5G polymorphism in the promoter region of plasminogen

[146553]activator inhibitor–1 with abdominal aortic aneurysms. J. Vasc. Surg. 31:

[146554]1026–1032, 2000.

[146555]26340.

[146556]Steinberger, D.; Mulliken, J. B.; Muller, U.: Crouzon syndrome:

[146557]previously unrecognized deletion, duplication, and point mutation

[146558]within FGFR2 gene. Hum. Mutat. 8: 386–390, 1996.

[146559]26341.

[146560]Oettinger, M. A.; Schatz, D. G.; Gorka, C.; Baltimore, D.:
RAG-1

[146561]and RAG-2, adjacent genes that synergistically activate
V(D)J recombination. Science 248:

[146562]1517-1523, 1990.

[146563]26342.

[146564]Steinberger, D.; Reinhartz, T.; Unsold, R.; Muller, U.:
FGFR2

[146565]mutation in clinically nonclassifiable autosomal dominant
craniosynostosis

[146566]with pronounced phenotypic variation. Am. J. Med. Genet.
66: 81-86,

[146567]1996.

[146568]26343.

[146569]Steinberger, D.; Vriend, G.; Mulliken, J. B.; Muller, U.: The

[146570]mutations in FGFR2-associated craniosynostoses are clustered in five

[146571]structural elements of immunoglobulin-like domain III of
the receptor. Hum.

[146572]Genet. 102: 145-150, 1998. Note: Erratum: Hum. Genet.
102: 495 only,

[146573]1998.

[146574]26344.

[146575]Tartaglia, M.; Di Rocco, C.; Lajeunie, E.; Valeri, S.; Velardi,

[146576]F.; Battaglia, P. A.: Jackson–Weiss syndrome: identification
of two

[146577]novel FGFR2 missense mutations shared with Crouzon and
Pfeiffer craniosynostotic

[146578]disorders. Hum. Genet. 101: 47–50, 1997.

[146579]26345.

[146580]Tartaglia, M.; Valeri, S.; Velardi, F.; Di Rocco, C.; Battaglia,

[146581]P. A.: Trp290Cys mutation in exon IIIa of the fibroblast
growth factor

[146582]receptor 2 (FGFR2) gene is associated with Pfeiffer syn-
drome. Hum.

[146583]Genet. 99: 602–606, 1997.

[146584]26346.

[146585]Teebi, A. S.; Kennedy, S.; Chun, K.; Ray, P. N.: Severe and
mild

[146586]phenotypes in Pfeiffer syndrome with splice acceptor mu-
tations in

[146587]lexon IIIc of FGFR2. Am. J. Med. Genet. 107: 43–47, 2002.

[146588]26347.

[146589]Tsukuno, M.; Suzuki, H.; Eto, T.: Pfeiffer syndrome caused
by

[146590]haploinsufficient mutation of FGFR2. J. Craniofac. Genet.
Dev. Biol. 19:

[146591]183–188, 1999.

[146592]26348.

[146593]Van Dissel–Emiliani, F. M.; De Boer–Brouwer, M.; De Rooij,
D.

[146594]G.: Effect of fibroblast growth factor–2 on Sertoli cells and
gonocytes

[146595]in coculture during the perinatal period. Endocrinology
137: 647–654,

[146596]1996.

[146597]26349.

[146598]Vogt, A.: Dyskephalie (dysostosis craniofacialis, maladie
De

[146599]Crouzon 1912) und eine neuartige Kombination dieser
Krankheit mit

[146600]Syndaktylie der 4 Extremitaeten (Dyskephalodaktylie).

Klin. Mbl.

[146601]Augenheilk. 90: 441–454, 1933.

[146602]26350.

[146603]Wang, T.-J.; Huang, C.-B.; Tsai, F.-J.; Wu, J.-Y.; Lai, R.-B.;

[146604]Hsiao, M.: Mutation in the FGFR2 gene in a Taiwanese patient with

[146605]Beare–Stevenson cutis gyrata syndrome. Clin. Genet. 61: 218–221,

[146606]2002.

[146607]26351.

[146608]Wilkie, A. O. M.: Abnormal spliceform expression associated with

[146609]splice acceptor mutations in exon IIIc of FGFR2. (Letter) Am. J.

[146610]Med. Genet. 111: 105 only, 2002.

[146611]26352.

[146612]Wilkie, A. O. M.: Personal Communication. Oxford, U.K. 2/24/1996.

[146613]26353.

[146614]Wilkie, A. O. M.; Slaney, S. F.; Oldridge, M.; Poole, M. D.;

Ashworth,

[146615]G. J.; Hockley, A. D.; Hayward, R. D.; David, D. J.; Pulleyn, L. J.;

[146616]Rutland, P.; Malcolm, S.; Winter, R. M.; Reardon, W.: Apert syndrome

[146617]results from localized mutations of FGFR2 and is allelic with Crouzon

[146618]syndrome. Nature Genet. 9: 165–172, 1995.

[146619]26354.

[146620]Wong, L.-J. C.; Chen, T.-J.; Dai, P.; Bird, L.; Muenke, M.: Novel

[146621]SNP at the common primer site of exon IIIa of FGFR2 gene causes error

[146622]in molecular diagnosis of craniosynostosis syndrome. Am. J. Med.

[146623]Genet. 102: 282–285, 2001.

[146624]26355.

[146625]Xu, X.; Weinstein, M.; Li, C.; Naski, M.; Cohen, R. I.; Ornitz,

[146626]D. M.; Leder, P.; Deng, C.: Fibroblast growth factor receptor 2 (FGFR2)–mediated

[146627]reciprocal regulation loop between FGF8 and FGF10 is es–

essential for

[146628]limb induction. Development 125: 753–765, 1998.

[146629]26356.

[146630]Yu, K.; Herr, A. B.; Waksman, G.; Ornitz, D. M.: Loss of fibroblast

[146631]growth factor receptor 2 ligand-binding specificity in Apert syndrome. Proc.

[146632]Nat. Acad. Sci. 97: 14536–14541, 2000.

[146633]26357.

[146634]Chan, J.; Watt, V. M.: Eek and erk, new members of the eph subclass

[146635]of receptor protein-tyrosine kinases. Oncogene 6: 1057–1061, 1991.

[146636]26358.

[146637]Park, S.; Frisen, J.; Barbacid, M.: Aberrant axonal projections

[146638]in mice lacking EphA8 (Eek) tyrosine protein kinase receptors. EMBO

[146639]J. 16: 3106–3114, 1997.

[146640]26359.

[146641]Ganju, P.; Shigemoto, K.; Brennan, J.; Entwistle, A.; Reith,

A.

- [146642]D.: The Eck receptor tyrosine kinase is implicated in pattern formation during gastrulation, hindbrain segmentation and limb development. *Oncogene* 9: 1613–1624, 1994.
- [146645]26360.
- [146646]Lindberg, R. A.; Hunter, T.: cDNA cloning and characterization of eck, an epithelial cell receptor–tyrosine kinase in the eph/elk family of protein kinases. *Molec. Cell. Biol.* 10: 6316–6324, 1990.
- [146649]26361.
- [146650]Sulman, E. P.; Tang, X. X.; Allen, C.; Biegel, J. A.; Pleasure, D. E.; Brodeur, G. M.; Ikegaki, N.: ECK, a human EPH–related gene, maps to 1p36.1, a common region of alteration in human cancers. *Genomics* 40: 371–374, 1997.
- [146654]26362.

- [146655]Arpaia, E.; Shahar, M.; Dadi, H.; Cohen, A.; Roifman, C. M.:
Defective
- [146656]T cell receptor signaling and CD8(+) thymic selection in
humans lacking
- [146657]Zap-70 kinase. Cell 76: 947-958, 1994.
- [146658]26363.
- [146659]Chan, A. C.; Iwashima, M.; Turck, C. W.; Weiss, A.: ZAP-
70: a
- [146660]70 kd protein-tyrosine kinase that associates with the
TCR zeta chain. Cell 71:
- [146661]649-662, 1992.
- [146662]26364.
- [146663]Chan, A. C.; Kadlecsek, T. A.; Elder, M. E.; Filipovich, A. H.;
- [146664]Kuo, W.-L.; Iwashima, M.; Parslow, T. G.; Weiss, A.: ZAP-
70 deficiency
- [146665]in an autosomal recessive form of severe combined im-
munodeficiency. Science 264:
- [146666]1559-1601, 1994.
- [146667]26365.
- [146668]Colucci, F.; Schweighoffer, E.; Tomasello, E.; Turner, M.;
Ortaldo,

[146669]J. R.; Vivier, E.; Tybulewicz, V. L. J.; Di Santo, J. P.: Natural
 [146670]cytotoxicity uncoupled from the Syk and ZAP-70 intracel-
 lular kinases. Nature
 [146671]Immun. 3: 288-294, 2002.
 [146672]26366.
 [146673]Elder, M. E.; Lin, D.; Clever, J.; Chan, A. C.; Hope, T. J.;
 Weiss,
 [146674]A.; Parslow, T. G.: Human severe combined immunodefi-
 ciency due to
 [146675]a defect in ZAP-70, a T cell tyrosine kinase. Science 264:
 1596-1599,
 [146676]1994.
 [146677]26367.
 [146678]Ku, G.; Malissen, B.; Mattei, M.-G.: Chromosomal location
 of the
 [146679]Syk and ZAP-70 tyrosine kinase genes in mice and hu-
 mans. Immunogenetics 40:
 [146680]300-302, 1994.
 [146681]26368.
 [146682]Roifman, C. M.; Hummel, D.; Martinez-Valdez, H.;
 Thorner, P.; Doherty,

- [146683]P. J.; Pan, S.; Cohen, F.; Cohen, A.: Depletion of CD8(+) cells in
[146684]human thymic medulla results in selective immune deficiency. J. Exp.
[146685]Med. 170: 2177–2182, 1989.
[146686]26369.
- [146687]Saito, T.; Matsuda, Y.; Ito, H.; Fusaki, N.; Hori, T.; Yamamoto,
[146688]T.: Localization of Zap70, the gene for a T cell-specific protein
[146689]tyrosine kinase, to mouse and rat chromosomes by fluorescence in situ
[146690]hybridization and molecular genetic linkage analyses. Mammalian Genome 8:
[146691]45–46, 1997.
[146692]26370.
- [146693]Toyabe, S.-I.; Watanabe, A.; Harada, W.; Karasawa, T.; Uchiyama,
[146694]M.: Specific immunoglobulin E responses in ZAP-70-deficient patients
[146695]are mediated by Syk-dependent T-cell receptor signalling. Immunology 103:

[146696]164–171, 2001.

[146697]26371.

[146698]Oettinger, M. A.; Stanger, B.; Schatz, D. G.; Glaser, T.; Call,

[146699]K.; Housman, D.; Baltimore, D.: The recombination activating genes,

[146700]RAG1 and RAG2, are on chromosome 11p in humans and chromosome 2p in

[146701]mice. Immunogenetics 35: 97–101, 1992.

[146702]26372.

[146703]Roman, C. A. J.; Baltimore, D.: Genetic evidence that the RAG1

[146704]protein directly participates in V(D)J recombination through substrate

[146705]recognition. Proc. Nat. Acad. Sci. 93: 2333–2338, 1996.

[146706]26373.

[146707]Santagata, S.; Gomez, C. A.; Sobacchi, C.; Bozzi, F.; Abinun,

[146708]M.; Pasic, S.; Cortes, P.; Vezzoni, P.; Villa, A.: N-terminal RAG1

[146709]frameshift mutations in Omenn's syndrome: internal methionine usage

[146710]leads to partial V(D)J recombination activity and reveals a
fundamental

[146711]role in vivo for the N-terminal domains. Proc. Nat. Acad.
Sci. 97:

[146712]14572–14577, 2000.

[146713]26374.

[146714]Schatz, D. G.; Oettinger, M. A.; Baltimore, D.: The V(D)J re-
combination

[146715]activating gene, RAG-1. Cell 59: 1035–1048, 1989.

[146716]26375.

[146717]Schultz, H. Y.; Landree, M. A.; Qiu, J.; Kale, S. B.; Roth, D.

[146718]B.: Joining-deficient RAG1 mutants block V(D)J recombina-
tion in vivo

[146719]and hairpin opening in vitro. Molec. Cell 7: 65–75, 2001.

[146720]26376.

[146721]Schwarz, K.; Gauss, G. H.; Ludwig, L.; Pannicke, U.; Li, Z.;
Linder,

[146722]D.; Friedrich, W.; Seger, R. A.; Hansen-Hagge, T. E.;
Desiderio, S.;

[146723]Lieber, M. R.; Bartram, C. R.: RAG mutations in human B
cell-negative

[146724]SCID. Science 274: 97–99, 1996.

[146725]26377.

[146726]Sherrington, P. D.; Forster, A.; Seawright, A.; van Heyning,
gen,

[146727]V.; Rabbitts, T. H.: Human RAG2, like RAG1, is on chromosome 11 band

[146728]p13 and therefore not linked to ataxia telangiectasia complementation

[146729]groups. Genes Chromosomes Cancer 5: 404–406, 1992.

[146730]26378.

[146731]Villa, A.; Santagata, S.; Bozzi, F.; Giliani, S.; Frattini, A.;

[146732]Imberti, L.; Gatta, L. B.; Ochs, H. D.; Schwarz, K.; Notarangelo,

[146733]L. D.; Vezzoni, P.; Spanopoulou, E.: Partial V(D)J recombination

[146734]activity leads to Omenn syndrome. Cell 93: 885–896, 1998.

[146735]26379.

[146736]Wienholds, E.; Schulte-Merker, S.; Walderich, B.; Plasterk, R.

[146737]H. A.: Target-selected inactivation of the zebrafish rag1

gene. Science 297:

[146738]99–102, 2002.

[146739]26380.

[146740]Yu, W.; Misulovin, Z.; Suh, H.; Hardy, R. R.; Jankovic, M.;
Yannoutsos,

[146741]N.; Nussenzweig, M. C.: Coordinate regulation of RAG1
and RAG2 by

[146742]cell type–specific DNA elements 5–prime of RAG2. Science
285: 1080–1084,

[146743]1999.

[146744]26381.

[146745]Richard, G.; Brown, N.; Smith, L. E.; Terrinoni, A.; Melino,
G.;

[146746]MacKie, R. M.; Bale, S. J.; Uitto, J.: The spectrum of muta-
tions

[146747]in erythrokeratodermias--novel and de novo mutations in
GJB3. Hum.

[146748]Genet. 106: 321–329, 2000.

[146749]26382.

[146750]Levitt, R. C.; Olckers, A.; Meyers, S.; Fletcher, J. E.; Rosen-
berg,

[146751]H.; Isaacs, H.; Meyers, D. A.: Evidence for the localization
of a

[146752]malignant hyperthermia susceptibility locus (MHS2) to human chromosome

[146753]17q. Genomics 14: 562–566, 1992.

[146754]26383.

[146755]Olickers, A.; Meyers, D. A.; Meyers, S.; Taylor, E. W.; Fletcher,

[146756]J. E.; Rosenberg, H.; Isaacs, H.; Levitt, R. C.: Adult muscle sodium

[146757]channel alpha-subunit is a gene candidate for malignant hyperthermia

[146758]susceptibility. Genomics 14: 829–831, 1992.

[146759]26384.

[146760]Naim, H. Y.; Sterchi, E. E.; Lentze, M. J.: Structure, biosynthesis,

[146761]and glycosylation of human small intestinal maltase-glucoamylase. J.

[146762]Biol. Chem. 263: 19709–19717, 1988.

[146763]26385.

[146764]Nichols, B. L.; Eldering, J.; Avery, S.; Hahn, D.; Quaroni, A.;

[146765]Sterchi, E.: Human small intestinal maltase-glucoamylase

cDNA cloning:

[146766]homology to sucrase-isomaltase. J. Biol. Chem. 273:
3076-3081, 1998.

[146767]26386.

[146768]Schuler, G. D.; Boguski, M. S.; Stewart, E. A.; Stein, L. D.;
Gyapay,

[146769]G.; Rice, K.; White, R. E.; Rodriguez-Tome, P.; Aggarwal,
A.; Bajorek,

[146770]E.; Bentolila, S.; Birren, B. B.; Butler, A.; Castle, A. B.; Chi-
annilkulchai,

[146771]N.; Chu, A.; Clee, C.; Cowles, S.; Day, P. J. R.; Dibling, T.;
and

[146772]84 others: A gene map of the human genome. Science
274: 540-546,

[146773]1996.

[146774]26387.

[146775]Levine, A.; Cantoni, G. L.; Razin, A.: Inhibition of promoter

[146776]activity by methylation: possible involvement of protein
mediators. Proc.

[146777]Nat. Acad. Sci. 88: 6515-6518, 1991.

[146778]26388.

[146779]Nan, X.; Meehan, R. R.; Bird, A.: Dissection of the methyl-CpG

[146780]binding domain from the chromosomal protein MeCP2.
Nucleic Acids

[146781]Res. 21: 4886-4892, 1993.

[146782]26389.

[146783]Ohki, I.; Shimotake, N.; Fujita, N.; Jee, J.-G.; Ikegami, T.;
Nakao,

[146784]M.; Shirakawa, M.: Solution structure of the methyl-CpG
binding domain

[146785]of human MBD1 in complex with methylated DNA. Cell
105: 487-497,

[146786]2001.

[146787]26390.

[146788]Carrera, C. J.; Eddy, R. L.; Shows, T. B.; Carson, D. A.: As-
signment

[146789]of the gene for methylthioadenosine phosphorylase to hu-
man chromosome

[146790]9 by mouse-human somatic cell hybridization. Proc. Nat.
Acad. Sci. 81:

[146791]2665-2668, 1984.

[146792]26391.

[146793]Chilcote, R. R.; Brown, E.; Rowley, J. D.: Lymphoblastic leukemia

[146794]with lymphomatous features associated with abnormalities of the short

[146795]arm of chromosome 9. New Eng. J. Med. 313: 286-291, 1985.

[146796]26392.

[146797]Nobori, T.; Takabayashi, K.; Tran, P.; Orvis, L.; Batova, A.; Yu,

[146798]A. L.; Carson, D. A.: Genomic cloning of methylthioadenosine phosphorylase:

[146799]a purine metabolic enzyme deficient in multiple different cancers. Proc.

[146800]Nat. Acad. Sci. 93: 6203-6208, 1996.

[146801]26393.

[146802]Olopade, O. I.; Jenkins, R. B.; Ransom, D. T.; Malik, K.; Pomykala,

[146803]H.; Nobori, T.; Cowan, J. M.; Rowley, J. D.; Diaz, M. O.: Molecular

[146804]analysis of deletions of the short arm of chromosome 9 in human gliomas. Cancer

[146805]Res. 52: 2523–2529, 1992.

[146806]26394.

[146807]Olopade, O. I.; Pomykala, H. M.; Hagos, F.; Sveen, L. W.;
Espinosa,

[146808]R., III; Dreyling, M. H.; Gursky, S.; Stadler, W. M.; Le Beau,
M.

[146809]M.; Bohlander, S. K.: Construction of a 2.8–megabase
yeast artificial

[146810]chromosome contig and cloning of the human methylthioadenosine phosphorylase

[146811]gene from the tumor suppressor region on 9p21. Proc.
Nat. Acad. Sci. 92:

[146812]6489–6493, 1995.

[146813]26395.

[146814]Ragione, F. D.; Takabayashi, K.; Mastropietro, S.; Mercurio,
C.;

[146815]Oliva, A.; Russo, G. L.; Pietra, V. D.; Borriello, A.; Nobori,
T.;

[146816]Carson, D. A.; Zappia, V.: Purification and characterization
of recombinant

[146817]human 5–prime–methylthioadenosine phosphorylase:
definite identification

[146818]of coding cDNA. Biochem. Biophys. Res. Commun. 223:
514–519, 1996.

[146819]26396.

[146820]Williams–Ashman, H. G.; Seidenfeld, J.; Galletti, P.: Trends
in

[146821]the biochemical pharmacology of

5–prime–deoxy–5–prime–methylthioadenosine. Biochem.

[146822]Pharm. 31: 277–288, 1982.

[146823]26397.

[146824]Kniest, W.: Zur Abgrenzung der Dysostosis enchondralis
von der

[146825]Chondrodystrophie. Z. Kinderheilk. 43: 633–640, 1952.

[146826]26398.

[146827]Mauernee, I. H.; Traboulsi, E. I.: The ocular findings in
Kniest

[146828]dysplasia. Am. J. Ophthal. 100: 155–160, 1985.

[146829]26399.

[146830]Siggers, D. C.; Rimoin, D. L.; Dorst, J. P.; Doty, S. B.;
Williams,

[146831]B. R.; Hollister, D. W.; Silberberg, R.; Granley, R. E.; Kauf-
man,

[146832]R. L.; McKusick, V. A.: The Kniest syndrome. Birth Defects
Orig.

[146833]Art. Ser. 10(9): 193–208, 1974.

[146834]26400.

[146835]Spranger, J.; Winterpacht, A.; Zabel, B.: Kniest dysplasia:
Dr.

[146836]W. Kniest, his patient, the molecular defect. Am. J. Med.
Genet. 69:

[146837]79–84, 1997.

[146838]26401.

[146839]Wilkin, D. J.; Artz, A. S.; South, S.; Lachman, R. S.; Rimoin,

[146840]D. L.; Wilcox, W. R.; McKusick, V. A.; Stratakis, C. A.; Fran-
comano,

[146841]C. A.; Cohn, D. H.: Small deletions in the type II collagen
triple

[146842]helix produce Kniest dysplasia. Am. J. Med. Genet. 85:
105–112,

[146843]1999.

[146844]26402.

[146845]Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human
gene encoding

- [146846]methionyl-tRNA synthetase to chromosome 12. (Abstract)
Cytogenet.
- [146847]Cell Genet. 37: 437 only, 1984.
- [146848]26403.
- [146849]Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human
MARS gene,
[146850]encoding methionyl-tRNA synthetase, to chromosome
12 using human
[146851]x Chinese hamster cell hybrids. Somat. Cell Molec. Genet.
10: 225-234,
[146852]1984.
- [146853]26404.
- [146854]Grewal, P. K.; Holzfeind, P. J.; Bittner, R. E.; Hewitt, J. E.
[146855]: Mutant glycosyltransferase and altered glycosylation of
alpha-dystroglycan
[146856]in the myodystrophy mouse. Nature Genet. 28: 151-154,
2001.
- [146857]26405.
- [146858]Evans, D. G. R.; Huson, S. M.; Donnai, D.; Neary, W.; Blair,
V.;
- [146859]Newton, V.; Harris, R.: A clinical study of type 2 neurofi-

bromatosis. Quart.

[146860]J. Med. 84: 603–618, 1992.

[146861]26406.

[146862]Honda, M.; Arai, E.; Sawada, S.; Ohta, A.; Niimura, M.:

Neurofibromatosis

[146863]2 and neurilemmomatosis gene are identical. J. Invest.

Derm. 104:

[146864]74–77, 1995.

[146865]26407.

[146866]Parry, D. M.; Eldridge, R.; Kaiser–Kupfer, M. I.; Bouzas, E.

A.;

[146867]Pikus, A.; Patronas, N.: Neurofibromatosis 2 (NF2): clinical

characteristics

[146868]of 63 affected individuals and clinical evidence for hetero–

geneity. Am.

[146869]J. Med. Genet. 52: 450–461, 1994.

[146870]26408.

[146871]Eddy, R. L.; Kretschmer, P. J.; Fairhurst, J. L.; Shows, T. B.;

[146872]Bohlen, P.; O'Hara, B.; Kovesdi, I.: A human gene family of

neurite

[146873]outgrowth–promoting proteins: the gene for a heparin

binding neurite

[146874]outgrowth-promoting factor maps to 7q22-qter.

(Abstract) Cytogenet.

[146875]Cell Genet. 58: 1920 only, 1991.

[146876]26409.

[146877]Lai, S.; Czubayko, F.; Riegel, A. T.; Wellstein, A.: Structure

[146878]of the human heparin-binding growth factor gene
pleiotrophin. Biochem.

[146879]Biophys. Res. Commun. 187: 1113-1122, 1992.

[146880]26410.

[146881]Li, Y.-S.; Hoffman, R. M.; Le Beau, M. M.; Espinosa, R., III;
Jenkins,

[146882]N. A.; Gilbert, D. J.; Copeland, N. G.; Deuel, T. F.: Charac-
terization

[146883]of the human pleiotrophin gene: promoter region and
chromosomal localization. J.

[146884]Biol. Chem. 267: 26011-26016, 1992.

[146885]26411.

[146886]Li, Y.-S.; Milner, P. G.; Chauhan, A. K.; Watson, M. A.;
Hoffman,

[146887]R. M.; Kodner, C. M.; Milbrandt, J.; Deuel, T. F.: Cloning

and expression

[146888]of a developmentally regulated protein that induces mitogenic and

[146889]neurite outgrowth activity. Science 250: 1690–1694, 1990.

[146890]26412.

[146891]Milner, P. G.; Shah, D.; Veile, R.; Donis-Keller, H.; Kumar, B.

[146892]V.: Cloning, nucleotide sequence, and chromosome localization of

[146893]the human pleiotrophin gene. Biochemistry 31: 12023–12028, 1992.

[146894]26413.

[146895]Brightbill, H. D.; Libraty, D. H.; Krutzik, S. R.; Yang, R.-B.;

[146896]Bellsie, J. T.; Bieharski, J. R.; Maitland, M.; Norgard, M. V.; Plevy,

[146897]S. E.; Smale, S. T.; Brennan, P. J.; Bloom, B. R.; Godowski, P. J.;

[146898]Modlin, R. L.: Host defense mechanisms triggered by microbial lipoproteins

[146899]through toll-like receptors. Science 285: 732–736, 1999.

[146900]26414.

- [146901]Jankovic, D.; Kullberg, M. C.; Hieny, S.; Caspar, P.; Collazo,
[146902]C. M.; Sher, A.: In the absence of IL-12, CD4+ T cell re-
sponses to
[146903]intracellular pathogens fail to default to a Th2 pattern and
are host
[146904]protective in an IL-10-/- setting. Immunity 16: 429-439,
2002.
[146905]26415.
- [146906]Jouanguy, E.; Lamhamedi-Cherradi, S.; Altare, F.; Fon-
daneche,
[146907]M.-C.; Tuerlinckx, D.; Blanche, S.; Emile, J.-F.; Gaillard,
J.-L.;
[146908]Schreiber, R.; Levin, M.; Fischer, A.; Hivroz, C.; Casanova,
J.-L.
[146909]: Partial interferon-gamma receptor 1 deficiency in a child
with tuberculoid
[146910]bacillus Calmette-Guerin infection and a sibling with clinical tuberculosis. J.
[146911]Clin. Invest. 100: 2658-2664, 1997.
[146912]26416.
- [146913]Warrington, J. A.; Bengtsson, U.: High-resolution physical
mapping

[146914] of human 5q31-q33 using three methods: radiation hybrid mapping, interphase

[146915] fluorescence in situ hybridization, and pulsed-field gel electrophoresis. Genomics 24:

[146916] 395-398, 1994.

[146917] 26417.

[146918] Anderson, S. K.; Gallinger, S.; Roder, J.; Frey, J.; Young, H.

[146919] A.; Ortaldo, J. R.: A cyclophilin-related protein involved in the

[146920] function of natural killer cells. Proc. Nat. Acad. Sci. 90: 542-546,

[146921] 1993.

[146922] 26418.

[146923] Frey, J. L.; Bino, T.; Kantor, R. R. S.; Segal, D. M.; Giardina,

[146924] S. L.; Roder, J.; Anderson, S.; Ortaldo, J. R.: Mechanism of target

[146925] cell recognition by natural killer cells: characterization of a novel

[146926] triggering molecule restricted to CD3- large granular lymphocytes. J.

[146927] Exp. Med. 174: 1527-1536, 1991.

[146928]26419.

[146929]Rinfret, A.; Anderson, S. K.: IL-2 regulates the expression
of

[146930]the NK-TR gene via an alternate RNA splicing mechanism.
Molec. Immun. 30:

[146931]1307-1313, 1993.

[146932]26420.

[146933]Simons-Evelyn, M.; Young, H. A.; Anderson, S. K.: Charac-
terization

[146934]of the mouse Nktr gene and promoter. Genomics 40:
94-100, 1997.

[146935]26421.

[146936]Young, H. A.; Jenkins, N. A.; Copeland, N. G.; Simek, S.;
Lerman,

[146937]M. I.; Zbar, B.; Glenn, G.; Ortaldo, J. R.; Anderson, S. K.:
Localization

[146938]of a novel natural killer triggering receptor locus to hu-
man chromosome

[146939]3p23-p21 and mouse chromosome 9. Genomics 16:
548-549, 1993.

[146940]26422.

[146941]Scott, I. C.; Halila, R.; Jenkins, J. M.; Mehan, S.; Apostolou,
[146942]S.; Wingvist, R.; Callen, D. F.; Prockop, D. J.; Peltonen, L.;
Kadler,
[146943]K. E.: Molecular cloning, expression and chromosomal lo-
calization
[146944]of a human gene encoding a 33 kDa putative metallopep-
tidase (PRSM1). Gene 174:
[146945]135–143, 1996.
[146946]26423.
[146947]Baldwin, A. S., Jr.: The NF-kappa-B and I-kappa-B pro-
teins: new
[146948]discoveries and insights. Annu. Rev. Immun. 14: 649–683,
1996.
[146949]26424.
[146950]Chen, F.; Castranova, V.; Shi, X.; Demers, L. M.: New in-
sights
[146951]into the role of nuclear factor-kappa-B, a ubiquitous tran-
scription
[146952]factor in the initiation of diseases. Clin. Chem. 45: 7–17,
1999.
[146953]26425.

[146954]Liptay, S.; Schmid, R. M.; Perkins, N. D.; Meltzer, P.; Al-
therr,
[146955]M. R.; McPherson, J. D.; Wasmuth, J. J.; Nabel, G. J.: Related
subunits
[146956]of NF-kappa-B map to two distinct loci associated with
translocations
[146957]in leukemia, NFKB1 and NFKB2. Genomics 13: 287-292,
1992.
[146958]26426.

[146959]Mathew, S.; Murty, V. V. V. S.; Dalla-Favera, R.; Chaganti,
R.
[146960]S. K.: Chromosomal localization of genes encoding the
transcription
[146961]factors, c-rel, NF-kappa-Bp50, NF-kappa-Bp65, and lyt-
10 by fluorescence
[146962]in situ hybridization. Oncogene 8: 191-193, 1993.
[146963]26427.

[146964]Ozes, O. N.; Mayo, L. D.; Gustin, J. A.; Pfeffer, S. R.; Pfe-
fer,
[146965]L. M.; Donner, D. B.: NF-kappa-B activation by tumour
necrosis factor
[146966]requires the Akt serine-threonine kinase. Nature 401:

82–85, 1999.

[146967]26428.

[146968]Romashkova, J. A.; Makarov, S. S.: NF- κ B is a target
of

[146969]AKT in anti-apoptotic PDGF signalling. Nature 401:
86–90, 1999.

[146970]26429.

[146971]Zhong, H.; May, M. J.; Jimi, E.; Ghosh, S.: The phosphory-
lation

[146972]status of nuclear NF- κ B determines its association
with CBP/p300

[146973]or HDAC-1. Molec. Cell 9: 625–636, 2002.

[146974]26430.

[146975]Claudio, E.; Brown, K.; Park, S.; Wang, H.; Siebenlist, U.:
BAFF-induced

[146976]NEMO-independent processing of NF- κ B2 in matur-
ing B cells. Nature

[146977]Immun. 3: 958–965, 2002.

[146978]26431.

[146979]Neri, A.; Chang, C.-C.; Lombardi, L.; Salina, M.; Corradini,
P.;

- [146980]Maiolo, A. T.; Chaganti, R. S. K.; Dalla-Favera, R.: B cell lymphoma-associated
[146981]chromosomal translocation involves candidate oncogene
lyt-10, homologous
[146982]to NF-kappa-B p50. Cell 67: 1075-1087, 1991.
[146983]26432.
- [146984]Adams, B. S.; Leung, K.; Meltzer, P. S.; Lewis, K. A.; Wag-
ner-McPherson,
[146985]C.; Evans, G. A.; Nabel, G. J.: Localization of the gene en-
coding
[146986]R-kappa-B (NFRKB), a tissue-specific DNA binding pro-
tein, to chromosome
[146987]11q24-q25. Genomics 14: 270-274, 1992.
[146988]26433.
- [146989]Sanyal, S.; Sandstrom, D. J.; Hoeffler, C. A.; Ramaswami,
M.:
[146990]AP-1 function upstream of CREB to control synaptic plas-
ticity in Drosophila. Nature 416:
[146991]870-874, 2002.
[146992]26434.
- [146993]Arheden, K.; Mandahl, N.; Strombeck, B.; Isaksson, M.;

Mitelman,

[146994]F.: Chromosome localization of the human oncogene INT1 to 12q13 by

[146995]in situ hybridization. Cytogenet. Cell Genet. 47: 86–87, 1988.

[146996]26435.

[146997]Gavin, B. J.; McMahon, J. A.; McMahon, A. P.: Expression of multiple

[146998]novel Wnt–1/int–1–related genes during fetal and adult mouse development. Genes

[146999]Dev. 4: 2319–2332, 1990.

[147000]26436.

[147001]Kirikoshi, H.; Sekihara, H.; Katoh, M.: WNT10A and WNT6, clustered

[147002]in human chromosome 2q35 region with head–to–tail manner, are strongly

[147003]coexpressed in SW480 cells. Biochem. Biophys. Res. Commun. 283:

[147004]798–805, 2001.

[147005]26437.

[147006]McMahon, A. P.: The Wnt family of developmental regula–

tors. Trends

[147007]Genet. 8: 236–242, 1992.

[147008]26438.

[147009]Nusse, R.; Brown, A.; Papkoff, J.; Scambler, P.; Shackleford, G.;

[147010]McMahon, A.; Moon, R.; Varmus, H.: A new nomenclature for int-1 and

[147011]related genes: the Wnt gene family. (Letter) Cell 64: 231–232, 1991.

[147012]26439.

[147013]Nusse, R.; van't Veer, L.; Geurts van Kessel, A.; van Agthoven,

[147014]A.; Bootsma, D.; Varmus, H.: Chromosomal localization of a human

[147015]homologue of a putative mammary tumor oncogene. (Abstract) Cytogenet.

[147016]Cell Genet. 37: 556–557, 1984.

[147017]26440.

[147018]Pellegrino, J. E.; Lensch, M. W.; Muenke, M.; Chance, P. F.: Clinical

[147019]and molecular analysis in Joubert syndrome. Am. J. Med.

Genet. 72:

[147020]59–62, 1997.

[147021]26441.

[147022]Thomas, K. R.; Capecchi, M. R.: Targeted disruption of the
murine

[147023]int-1 proto-oncogene resulting in severe abnormalities in
midbrain

[147024]and cerebellar development. Nature 346: 847–850, 1990.

[147025]26442.

[147026]Andrews, J.; Erdjument, H.; Nicholson, D. C.: Hereditary
coproporphyrria:

[147027]incidence in a large English family. J. Med. Genet. 21:
341–349,

[147028]1984.

[147029]26443.

[147030]Barnes, H. D.; Whittaker, N.: Hereditary coproporphyrria
with acute

[147031]intermittent manifestations. Brit. Med. J. 2: 1102–1104,
1965.

[147032]26444.

[147033]Barohn, R. J.; Sanchez, J. A.; Anderson, K. E.: Acute pe-

ripheral

[147034]neuropathy due to hereditary coproporphyria. Muscle
Nerve 17: 793-799,

[147035]1994.

[147036]26445.

[147037]Berger, H.; Goldberg, A.: Hereditary coproporphyria. Brit.
Med.

[147038]J. 2: 85-88, 1955.

[147039]26446.

[147040]Brodie, M. J.; Thompson, G. G.; Moore, M. R.; Beattie, A.
D.; Goldberg,

[147041]A.: Hereditary coproporphyria: demonstration of the ab-
normalities

[147042]in haem biosynthesis in peripheral blood. Quart. J. Med.
46: 229-241,

[147043]1977.

[147044]26447.

[147045]Cacheux, V.; Martasek, P.; Fougerousse, F.; Delfau, M. H.;
Druart,

[147046]L.; Tachdjian, G.; Grandchamp, B.: Localization of the hu-
man coproporphyrinogen

[147047]oxidase gene to chromosome band 3q12. Hum. Genet.
94: 557–559, 1994.

[147048]26448.

[147049]Connon, J. J.; Turkington, V.: Hereditary coproporphyria.
Lancet II:

[147050]263–264, 1968.

[147051]26449.

[147052]Cripps, D. J.; Peters, H. A.: Stool porphyrins in acute inter-
mittent

[147053]and hereditary coproporphyria: adverse effects of tran-
quilizers. Arch.

[147054]Neurol. 23: 80–84, 1970.

[147055]26450.

[147056]Delfau-Larue, M.-H.; Martasek, P.; Grandchamp, B.: Co-
protophyrinogene

[147057](sic) oxidase: gene organization and description of a mu-
tation leading

[147058]to exon 6 skipping. Hum. Molec. Genet. 3: 1325–1330,
1994.

[147059]26451.

[147060]Doss, M.; von Tiepermann, R.; Kopp, W.: Harderoporphyryn

coproporphyria.

[147061](Letter) Lancet I: 292 only, 1984.

[147062]26452.

[147063]Elder, G. H.; Evans, J. O.; Thomas, N.; Cox, R.; Brodie, M.
J.;

[147064]Moore, M. R.; Goldberg, A.; Nicholson, D. C.: The primary
enzyme

[147065]defect in hereditary coproporphyria. Lancet II: 1217-1219,
1976.

[147066]26453.

[147067]Goldberg, A.; Rimington, C.; Lochhead, A. C.: Hereditary
coproporphyria. Lancet I:

[147068]632-636, 1967.

[147069]26454.

[147070]Grandchamp, B.; Phung, N.; Nordmann, Y.: Homozygous
case of hereditary

[147071]coproporphyria. (Letter) Lancet II: 1348-1349, 1977.

[147072]26455.

[147073]Grandchamp, B.; Weil, D.; Nordmann, Y.; Van Cong, N.; de
Verneuil,

[147074]H.; Foubert, C.; Gross, M.-S.: Assignment of the human

coproporphyrinogen

[147075]oxidase to chromosome 9. Hum. Genet. 64: 180–183,
1983.

[147076]26456.

[147077]Haeger–Aronsen, B.; Stathers, G.; Swahn, G.: Hereditary
coproporphyrinogen

[147078]study of a Swedish family. Ann. Intern. Med. 69: 221–227,
1968.

[147079]26457.

[147080]Hunter, J. A. A.; Khan, S. A.; Hope, E.; Beattie, A. D.; Bev-
eridge,

[147081]G. W.; Smith, A. W. M.; Goldberg, A.: Hereditary copropor-
phyria.

[147082]Photosensitivity, jaundice and neuropsychiatric manifesta-
tions associated

[147083]with pregnancy. Brit. J. Derm. 84: 301–310, 1971.

[147084]26458.

[147085]Kohno, H.; Furukawa, T.; Yoshinaga, T.; Tokunaga, R.;
Taketani,

[147086]S.: Coproporphyrinogen oxidase: purification, molecular
cloning,

[147087]and induction of mRNA during erythroid differentiation. J.

Biol.

[147088]Chem. 268: 21359–21363, 1993.

[147089]26459.

[147090]Lamoril, J.; Deybach, J.-C.; Puy, H.; Grandchamp, B.; Nordmann,

[147091]Y.: Three novel mutations in the coproporphyrinogen oxidase gene. Hum.

[147092]Mutat. 9: 78–80, 1997.

[147093]26460.

[147094]Lamoril, J.; Martasek, P.; Deybach, J.-C.; Da Silva, V.; Grandchamp,

[147095]B.; Nordmann, Y.: A molecular defect in coproporphyrinogen oxidase

[147096]gene causing harderoporphyria, a variant form of hereditary coproporphyria. Hum.

[147097]Molec. Genet. 4: 275–278, 1995.

[147098]26461.

[147099]Lamoril, J.; Puy, H.; Gouya, L.; Rosipal, R.; Da Silva, V.; Grandchamp,

[147100]B.; Foint, T.; Bader-Meunier, B.; Dommergues, J. P.; Deybach, J. C.;

- [147101]Nordmann, Y.: Neonatal hemolytic anemia due to inherited harderoporphyria:
[147102]clinical characteristics and molecular basis. Blood 91:
1453–1457,
[147103]1998.
[147104]26462.
- [147105]Lamoril, J.; Puy, H.; Whatley, S. D.; Martin, C.; Woolf, J. R.;
[147106]Da Silva, V.; Deybach, J.-C.; Elder, G. H.: Characterization
of mutations
[147107]in the CPO gene in British patients demonstrates absence
of genotype–phenotype
[147108]correlation and identifies relationship between hereditary
coproporphyria
[147109]and harderoporphyria. Am. J. Hum. Genet. 68:
1130–1138, 2001.
[147110]26463.
- [147111]Lomholt, J. C.; With, T. K.: Hereditary coproporphyria: a
family
[147112]with unusually few and mild symptoms. Acta Med. Scand.
186: 83–85,
[147113]1969.

[147114]26464.

[147115]Martasek, P.; Camadro, J. M.; Delfau-Larue, M.-H.; Dumas, J.-B.;

[147116]Montagne, J. J.; De Verneuil, H.; Labbe, P.; Grandchamp, B.: Molecular

[147117]cloning, sequencing, and functional expression of a cDNA encoding

[147118]human coproporphyrinogen oxidase. Proc. Nat. Acad. Sci. 91: 3024-3028,

[147119]1994.

[147120]26465.

[147121]Martasek, P.; Nordmann, Y.; Grandchamp, B.: Homozygous hereditary

[147122]coproporphyrinemia caused by an arginine to tryptophane substitution in

[147123]coproporphyrinogen oxidase and common intragenic polymorphisms. Hum.

[147124]Molec. Genet. 3: 477-480, 1994.

[147125]26466.

[147126]McIntyre, N.; Pearson, A. J. G.; Allan, D. J.; Craske, S.; West,

[147127]G. M. L.; Moore, M. R.; Beattie, A. D.; Paxton, J.; Goldberg, A.:

[147128]Hepatic delta-aminolaevulinic acid synthetase in an attack of hereditary

[147129]coproporphyrinemia and during remission. Lancet I: 560-564, 1971.

[147130]26467.

[147131]Nordmann, Y.; Grandchamp, B.; de Verneuil, H.; Phung, L.; Cartigny,

[147132]B.; Fontaine, G.: Harderoporphyria: a variant hereditary coproporphyrinemia. J.

[147133]Clin. Invest. 72: 1139-1149, 1983.

[147134]26468.

[147135]Roberts, D. T.; Brodie, M. J.; Moore, M. R.; Thompson, G. G.;

[147136]Goldberg, A.; MacSween, R. N. M.: Hereditary coproporphyrinemia presenting

[147137]with photosensitivity induced by the contraceptive pill. Brit. J.

[147138]Derm. 96: 549-554, 1977.

[147139]26469.

[147140]Dickinson, D. P.; Zhao, Y.; Thiesse, M.; Siciliano, M. J.: Direct

[147141]mapping of seven genes encoding human type 2 cystatins to a single

[147142]site located at 20p11.2. Genomics 24: 172–175, 1994.

[147143]26470.

[147144]Gopal Rao, V. V.; Schnittger, S.; Abrahamson, M.; Hansmann, I.

[147145]: Cystatin–C (CST3), the candidate gene for the hereditary cystatin–C

[147146]amyloid angiopathy (HCCAA) maps to or close to human chromosome 20p11.22.

[147147](Abstract) Cytogenet. Cell Genet. 58: 2029 only, 1991.

[147148]26471.

[147149]Saitoh, E.; Kim, H.–S.; Smithies, O.; Maeda, N.: Human cysteine–proteinase

[147150]inhibitors: nucleotide sequence analysis of three members of the cystatin

[147151]gene family. Gene 61: 329–338, 1987.

[147152]26472.

[147153]Saitoh, E.; Sabatini, L. M.; Eddy, R. L.; Shows, T. B.; Azen,

E.

[147154]A.; Isemura, S.; Sanada, K.: The human cystatin C gene (CST3) is

[147155]a member of the cystatin gene family which is localized on chromosome

[147156]20. Biochem. Biophys. Res. Commun. 162: 1324–1331, 1989.

[147157]26473.

[147158]Cruzen, M. E.; Bengtsson, U.; McMahon, J.; Wasmuth, J. J.; Arfin,

[147159]S. M.: Assignment of the cysteinyl-tRNA synthetase gene (CARS) to

[147160]11p15.5. Genomics 15: 692–693, 1993.

[147161]26474.

[147162]Takahashi, E.; Hori, T.; O'Connell, P.; Leppert, M.; White, R.

[147163]; R-banding and nonisotopic in situ hybridization: precise localization

[147164]of the human type II collagen gene (COL2A1). Hum. Genet. 86: 14–16,

[147165]1990.

[147166]26475.

[147167]Nebert, D. W.; Gonzalez, F. J.: P450 genes: structure, evolution,
[147168]and regulation. *Annu. Rev. Biochem.* 56: 945–993, 1987.
[147169]26476.

[147170]Paolini, M.; Cantelli–Forti, G.; Perocco, P.; Pedulli, G. F.;
[147171]Abdel–Rahman, S. Z.; Legator, M. S.: Co–carcinogenic effect of beta–carotene.
[147172](Letter) *Nature* 398: 760–761, 1999.
[147173]26477.

[147174]Espinoza, H.; Cox, C. J.; Semina, E. V.; Amendt, B. A.: A
molecular
[147175]basis for differential developmental anomalies in Axen–
feld–Rieger
[147176]syndrome. *Hum. Molec. Genet.* 11: 743–753, 2002.
[147177]26478.

[147178]Letinic, K.; Zoncu, R.; Rakic, P.: Origin of GABAergic neurons
[147179]in the human neocortex. *Nature* 417: 645–649, 2002.
[147180]26479.

[147181]McGuinness, T.; Porteus, M. H.; Smiga, S.; Bulfone, A.;

Kingsley,

[147182]C.; Qiu, M.; Liu, J. K.; Long, J. E.; Xu, D.; Rubenstein, J. L. R.

[147183]: Sequence, organization, and transcription of the *Dlx-1*
and the *Dlx-2*

[147184]locus. *Genomics* 35: 473–485, 1996.

[147185]26480.

[147186]Ozcelik, T.; Porteus, M. H.; Rubenstein, J. L. R.; Francke, U.

[147187]: *DLX2* (*TES1*), a homeobox gene of the Distal-less family,
assigned

[147188]to conserved regions on human and mouse chromosomes
2. *Genomics* 13:

[147189]1157–1161, 1992.

[147190]26481.

[147191]Porteus, M. H.; Brice, A. E. J.; Bulfone, A.; Usdin, T. B.; Ciaranello,

[147192]R. D.; Rubenstein, J. L. R.: Isolation and characterization of
a

[147193]library of cDNA clones that are preferentially expressed in
the embryonic

[147194]telencephalon. *Molec. Brain Res.* 12: 7–22, 1992.

[147195]26482.

- [147196] Qiu, M.; Bulfone, A.; Martinez, S.; Meneses, J. J.; Shimamura,
mura,
- [147197] K.; Pedersen, R. A.; Rubenstein, J. L.: Null mutation of Dlx-
2 results
- [147198] in abnormal morphogenesis of proximal first and second
branchial arch
- [147199] derivatives and abnormal differentiation in the forebrain.
Genes
- [147200] Dev. 9: 2523–2538, 1995.
- [147201] 26483.
- [147202] Salvador, J. M.; Hollander, M. C.; Nguyen, A. T.; Kopp, J. B.;
- [147203] Barisoni, L.; Moore, J. K.; Ashwell, J. D.; Fornace, A. J., Jr.:
- [147204] Mice lacking the p53–effector gene Gadd45a develop a
lupus–like syndrome. Immunity 16:
- [147205] 499–508, 2002.
- [147206] 26484.
- [147207] Takekawa, M.; Saito, H.: A family of stress–inducible
GADD45–like
- [147208] proteins mediate activation of the stress–responsive
MTK1/MEKK4 MAPKKK. Cell 95:
- [147209] 521–530, 1998.

[147210]26485.

[147211]Dhermy, D.; Garbarz, M.; Lecomte, M. C.; Feo, C.;
Bournier, O.;

[147212]Chaveroche, I.; Gautero, H.; Galand, C.; Boivin, P.: Heredi-
tary elliptocytosis:

[147213]clinical, morphological and biochemical studies of 38
cases. Nouv.

[147214]Rev. Franc. Hemat. 28: 129-140, 1986.

[147215]26486.

[147216]Fairbanks, G.; Steck, T. L.; Wallach, D. F. H.: Elec-
trophoretic

[147217]analysis of the major polypeptides of the human erythro-
cyte membrane. Biochemistry 10:

[147218]2606-2617, 1971.

[147219]26487.

[147220]Feo, C. J.; Fischer, S.; Piau, J. P.; Grange, M. J.; Tchernia,

[147221]G.: Premiere observation de l'absence d'une proteine de la
membrane

[147222]erythrocytaire (bande 4-1) dans un cas d'anemie ellipto-
cytaire familiale. Nouv.

[147223]Rev. Franc. Hemat. 22: 315-325, 1980.

[147224]26488.

[147225]Fishbein, W. N.; Armbrustmacher, V. W.; Griffin, J. L.;
Davis,

[147226]J. I.; Foster, W. D.: Levels of adenylate deaminase, adeny-
late kinase,

[147227]and creatine kinase in frozen human muscle biopsy speci-
mens relative

[147228]to type1/type2 fiber distribution: evidence for a carrier
state of

[147229]myoadenylate deaminase deficiency. Ann. Neurol. 15:
271-277, 1984.

[147230]26489.

[147231]Fishbein, W. N.; Griffin, J. L.; Nagarajan, K.; Winkert, J. W.;

[147232]Armbrustmacher, V. W.: Immunologic uniqueness of mus-
cle adenylate

[147233]deaminase (mAD) and genetic transmission of the defi-
ciency state.

[147234](Abstract) Clin. Res. 27: 274A only, 1979.

[147235]26490.

[147236]Genetta, T.; Morisaki, H.; Morisaki, T.; Holmes, E. W.: A
novel

[147237]bipartite intronic splicing enhancer promotes the inclusion
of a mini-exon

[147238]in the AMP deaminase 1 gene. J. Biol. Chem. 276:
25589-25597, 2001.

[147239]26491.

[147240]Kar, N. C.; Pearson, C. M.: Muscle adenylate deaminase
deficiency:

[147241]report of six new cases. Arch. Neurol. 38: 279-281, 1981.

[147242]26492.

[147243]Kelemen, J.; Bradley, W. G.; DiMauro, S.: Reply to J. B. Shu-
mate.

[147244](Letter) Neurology 33: 1534 only, 1983.

[147245]26493.

[147246]Kelemen, J.; Rice, D. R.; Bradley, W. G.; Munsat, T. L.; Di-
Mauro,

[147247]S.; Hogan, E. L.: Familial myoadenylate deaminase defi-
ciency and

[147248]exertional myalgia. Neurology 32: 857-863, 1982.

[147249]26494.

[147250]Lecky, B. R. F.: Failure of D-ribose in myoadenylate deam-
inase

[147251]deficiency. (Letter) Lancet I: 193 only, 1983.

[147252]26495.

[147253]Loh, E.; Rebbeck, T. R.; Mahoney, P. D.; DeNofrio, D.;
Swain,

[147254]J. L.; Holmes, E. W.: Common variant in AMPD1 gene pre-
dicts improved

[147255]clinical outcome in patients with heart failure. Circulation
99:

[147256]1422–1425, 1999.

[147257]26496.

[147258]Morisaki, H.; Higuchi, I.; Abe, M.; Osame, M.; Morisaki, T.:

[147259]First missense mutations (R388W and R425H) of AMPD1
accompanied with

[147260]myopathy found in a Japanese patient. Hum. Mutat. 16:
467–472, 2000.

[147261]26497.

[147262]Morisaki, H.; Morisaki, T.; Newby, L. K.; Holmes, E. W.: Al-
ternative

[147263]splicing: a mechanism for phenotypic rescue of a common
inherited

[147264]defect. J. Clin. Invest. 91: 2275–2280, 1993.

[147265]26498.

[147266]Morisaki, T.; Gross, M.; Morisaki, H.; Pongratz, D.; Zollner,

[147267]N.; Holmes, E. W.: Molecular basis of AMP deaminase deficiency in

[147268]skeletal muscle. Proc. Nat. Acad. Sci. 89: 6457-6461,
1992.

[147269]26499.

[147270]Morisaki, T.; Sabina, R. L.; Holmes, E. W.: Adenylate deaminase:

[147271]a multigene family in humans and rats. J. Biol. Chem. 265:
11482-11486,

[147272]1990.

[147273]26500.

[147274]Morton, C. C.; Eddy, R. L.; Shows, T. B.; Clark, P. R. H.;
Sabina,

[147275]R. L.; Holmes, E. W.: Human AMP deaminase-1 gene
(AMPD1) is mapped

[147276]to chromosome 1. (Abstract) Cytogenet. Cell Genet. 51:
1048-1049,

[147277]1989.

[147278]26501.

- [147279]Moseley, W. S.; Morisaki, T.; Sabina, R. L.; Holmes, E. W.; Seldin, M. F.: Ampd-2 maps to distal mouse chromosome 3 in linkage with Ampd-1. Genomics 6: 572-574, 1990.
- [147282]26502.
- [147283]Sabina, R. L.; Morisaki, T.; Clarke, P.; Eddy, R.; Shows, T. B.; Morton, C. C.; Holmes, E. W.: Characterization of the human and rat myoadenylate deaminase genes. J. Biol. Chem. 265: 9423-9433, 1990.
- [147286]26503.
- [147287]Sabina, R. L.; Ogasawara, N.; Holmes, E. W.: Expression of three stage-specific transcripts of AMP deaminase during myogenesis. Molec. Cell. Biol. 9: 2244-2246, 1989.
- [147290]26504.
- [147291]Sabina, R. L.; Swain, J. L.; Olanow, C. W.; Bradley, W. G.; Fishbein,

[147292]W. N.; DiMauro, S.; Holmes, E. W.: Myoadenylate deaminase deficiency:

[147293]functional and metabolic abnormalities associated with disruption

[147294]of the purine nucleotide cycle. J. Clin. Invest. 73: 720–730, 1984.

[147295]26505.

[147296]Sabina, R. L.; Swain, J. L.; Patten, B. M.; Ashizawa, T.; O'Brien,

[147297]W. E.; Holmes, E. W.: Disruption of the purine nucleotide cycle:

[147298]a potential explanation for muscle dysfunction in myoadenylate deaminase

[147299]deficiency. J. Clin. Invest. 66: 1419–1423, 1980.

[147300]26506.

[147301]Shumate, J. B.: Myoadenylate deaminase deficiency--a nonfamilial,

[147302]nondisease? (Letter) Neurology 33: 1533–1534, 1983.

[147303]26507.

[147304]Shumate, J. B.; Kaiser, K. K.; Carroll, J. E.; Brooke, M. H.:

[147305]Adenylate deaminase deficiency in a hypotonic infant. J.

Pediat. 96:

[147306]885–887, 1980.

[147307]26508.

[147308]Sinkeler, S. P. T.; Joosten, E. M. G.; Wevers, R. A.; Oei, T.

[147309]L.; Jacobs, A. E. M.; Veerkamp, J. H.; Hamel, B. C. J.:

Myoadenylate

[147310]deaminase deficiency: a clinical, genetic, and biochemical
study in

[147311]nine families. Muscle Nerve 11: 312–317, 1988.

[147312]26509.

[147313]Valen, P. A.; Nakayama, D. A.; Veum, J.; Sulaiman, A. R.;
Wortmann,

[147314]R. L.: Myoadenylate deaminase deficiency and forearm is-
chemic exercise

[147315]testing. Arthritis Rheum. 30: 661–668, 1987.

[147316]26510.

[147317]van den Berghe, G.; Hers, H. G.: Abnormal AMP deaminase
in primary

[147318]gout. (Letter) Lancet II: 1090 only, 1980.

[147319]26511.

[147320]Eddy, R. L.; Mahnke-Zizelman, D. K.; Bausch-Jurken, M.
T.; Sabina,

[147321]R. L.; Shows, T. B.: Distribution of the AMP deaminase
multigene

[147322]family within the human genome: assignment of the
AMPD2 to chromosome

[147323]1p21-p34 and AMPD3 to chromosome 11p13-pter.
(Abstract) Human Genome

[147324]Mapping Workshop 93 24 only, 1993.

[147325]26512.

[147326]Ogasawara, N.; Goto, H.; Yamada, Y.; Nishigaki, I.; Itoh, T.;
Hasegawa,

[147327]I.; Park, K. S.: Deficiency of AMP deaminase in erythro-
cytes. Hum.

[147328]Genet. 75: 15-18, 1987.

[147329]26513.

[147330]Sermusvitayawong, K.; Wang, X.; Nagabukuro, A.; Mat-
suda, Y.; Morisaki,

[147331]H.; Toyama, K.; Mukai, T.; Morisaki, T.: Genomic organi-
zation of

[147332]Ampd3, heart-type AMPD gene, located in mouse chro-
mosome 7. Mammalian

- [147333]Genome 8: 767–769, 1997.
- [147334]26514.
- [147335]Yamada, Y.; Goto, H.; Ogasawara, N.: Cloning and nucleotide sequence
[147336]of the cDNA encoding human erythrocyte-specific AMP deaminase. Biochim.
[147337]Biophys. Acta 1171: 125–128, 1992.
- [147338]26515.
- [147339]Yamada, Y.; Goto, H.; Ogasawara, N.: A point mutation responsible
[147340]for human erythrocyte AMP deaminase deficiency. Hum. Molec. Genet. 3:
[147341]331–334, 1994.
- [147342]26516.
- [147343]Allegrucci, C.; Liguori, L.; Mezzasoma, I.; Minelli, A.: A1 adenosine
[147344]receptor in human spermatozoa: its role in the fertilization process. Molec.
[147345]Genet. Metab. 71: 381–386, 2000.
- [147346]26517.

[147347]Bennhold, H.; Peters, H.; Roth, E.: Ueber einen Fall von
kompletter

[147348]Analbuminaemie ohne wesentliche klinische Krankheits-
zeichen. Verh.

[147349]Dtsch. Ges. Inn. Med. 60: 630–634, 1954.

[147350]26518.

[147351]Blumberg, B. S.; Martin, J. R.; Melartin, L.: Alloalbumine-
mia:

[147352]albumin Naskapi in Indians of the Ungava. J.A.M.A. 203:
180–185,

[147353]1968.

[147354]26519.

[147355]Boman, H.; Hermodson, M.; Hammond, C. A.; Motulsky, A.
G.: Analbuminemia

[147356]in an American Indian girl. Clin. Genet. 9: 513–526, 1976.

[147357]26520.

[147358]Brennan, S. O.: The molecular abnormality of albumin
Parklands:

[147359]365 asp-to-his. Biochim. Biophys. Acta 830: 320–324,
1985.

[147360]26521.

- [147361] Brennan, S. O.; Arai, K.; Madison, J.; Laurell, C.-B.; Gal-
liano,
[147362] M.; Watkins, S.; Peach, R.; Myles, T.; George, P.; Putnam, F.
W.:
[147363] Hypermutable CpG dinucleotides in the propeptide-en-
coding sequence
[147364] of the human albumin gene. Proc. Nat. Acad. Sci. 87:
3909-3913,
[147365] 1990.
[147366] 26522.
- [147367] Brennan, S. O.; Carrell, R. W.: A circulating variant of hu-
man
[147368] proalbumin. Nature 274: 908-909, 1978.
[147369] 26523.
- [147370] Brennan, S. O.; Fellowes, A. P.: Albumin Hawkes Bay; a low
level
[147371] variant caused by loss of a sulphhydryl group at position
177. Biochim.
[147372] Biophys. Acta 1182: 46-50, 1993.
[147373] 26524.
- [147374] Brennan, S. O.; Herbert, P.: Albumin Canterbury (313 lys-

to-asn):

[147375]a point mutation in the second domain of serum albumin.

Biochim.

[147376]Biophys. Acta 912: 191–197, 1987.

[147377]26525.

[147378]Brennan, S. O.; Myles, T.; Peach, R. J.; Donaldson, D.;

George,

[147379]P. M.: Albumin Redhill (–1 arg, ala320-to-thr): a glycoprotein variant

[147380]of human serum albumin whose precursor has an aberrant signal peptidase

[147381]cleavage site. Proc. Nat. Acad. Sci. 87: 26–30, 1990.

[147382]26526.

[147383]Campagnoli, M.; Rossi, A.; Palmqvist, L.; Flisberg, A.;

Niklasson,

[147384]A.; Minchotti, L.; Galliano, M.: A novel splicing mutation causes

[147385]an undescribed type of analbuminemia. Biochim. Biophys. Acta 1586:

[147386]43–49, 2002.

[147387]26527.

- [147388]Carlson, J.; Sakamoto, Y.; Laurell, C.-B.; Madison, J.;
Watkins,
[147389]S.; Putnam, F. W.: Alloalbuminemia in Sweden: structural
study and
[147390]phenotypic distribution of nine albumin variants. Proc.
Nat. Acad.
[147391]Sci. 89: 8225-8229, 1992.
[147392]26528.
- [147393]Cormode, E. J.; Lyster, D. M.; Israels, S.: Analbuminemia in
[147394]a neonate. J. Pediat. 86: 862-867, 1975.
[147395]26529.
- [147396]Dammacco, F.; Miglietta, A.; D'Addabbo, A.; Fratello, A.;
Moschetta,
[147397]R.; Bonomo, L.: Analbuminemia: report of a case and re-
view of the
[147398]literature. Vox Sang. 39: 153-161, 1980.
[147399]26530.
- [147400]Darlington, G.: Personal Communication. New Haven,
Conn. and
[147401]New York, N. Y. 9/17/1974.

[147402]26531.

[147403]Darlington, G. J.; Bernhard, H. P.; Ruddle, F. H.: Human
serum

[147404]albumin phenotype activation in mouse hepatoma-human
leukocyte cell

[147405]hybrids. Science 185: 859-862, 1974.

[147406]26532.

[147407]Dayhoff, M. O.: Serum albumin. Atlas of Protein Sequence
and

[147408]Structure. Washington, D. C.: National Biomedical Re-
search Foundation

[147409](pub.) 5: 1972. Pp. D316.

[147410]26533.

[147411]DeCosimo, D. R.; Fang, S.-L.; Braverman, L. E.: Prevalence
of

[147412]familial dysalbuminemic hyperthyroxinemia in Hispanics.
(Letter) Ann.

[147413]Intern. Med. 107: 780-781, 1987.

[147414]26534.

[147415]Dugaiczuk, A.: Personal Communication. Riverside, Calif.
4/29/1989.

[147416]26535.

[147417]Dugaiczky, A.; Law, S. W.; Dennison, O. E.: Nucleotide sequence

[147418]and the encoded amino acids of human serum albumin mRNA. Proc. Nat.

[147419]Acad. Sci. 79: 71–75, 1982.

[147420]26536.

[147421]Earle, D. P.; Hutt, M. P.; Schmid, K.; Gitlin, D.: Observations

[147422]on double albumin: a genetically transmitted serum protein anomaly. J.

[147423]Clin. Invest. 38: 1412–1420, 1959.

[147424]26537.

[147425]Efremov, G. D.; Braend, M.: Serum albumin: polymorphism in man. Science 146:

[147426]1679–1680, 1964.

[147427]26538.

[147428]Esumi, H.; Okui, M.; Sato, S.; Sugimura, T.; Nagase, S.: Absence

[147429]of albumin mRNA in the liver of analbuminemic rats. Proc. Nat. Acad.

[147430]Sci. 77: 3215–3219, 1980.

[147431]26539.

[147432]Esumi, H.; Takahashi, Y.; Sato, S.; Nagase, S.; Sugimura,
T.:

[147433]A seven–base–pair deletion in an intron of the albumin
gene of analbuminemic

[147434]rats. Proc. Nat. Acad. Sci. 80: 95–99, 1983.

[147435]26540.

[147436]Esumi, H.; Takahashi, Y.; Sekiya, T.; Sato, S.; Nagase, S.;
Sugimura,

[147437]T.: Presence of albumin mRNA precursors in nuclei of
albuminemic

[147438]rat liver lacking cytoplasmic albumin mRNA. Proc. Nat.
Acad. Sci. 79:

[147439]734–738, 1982.

[147440]26541.

[147441]Fine, J. M.; Abdo, Y.; Rochu, D.; Rousseaux, J.; Dautrevaux,
M.

[147442]: Identification of the human albumin variant 'Gainesville'
with proalbumin

[147443]'Christchurch'. Blood Transf. Immunohemat. 26: 341–346,

1983.

[147444]26542.

[147445]Fine, J. M.; Marneux, M.; Rochu, D.: Human albumin genetic variants:

[147446]an attempt at a classification of European allotypes. Am. J. Hum.

[147447]Genet. 40: 278–286, 1987.

[147448]26543.

[147449]Franklin, S. G.; Wolf, S. I.; Ozdemir, Y.; Yuregir, G. T.; Isbir,

[147450]T.; Blumberg, B. S.: Albumin Naskapi variant in North American Indians

[147451]and Eti Turks. Proc. Nat. Acad. Sci. 77: 5480–5482, 1980.

[147452]26544.

[147453]Franklin, S. G.; Wolf, S. I.; Zweidler, A.; Blumberg, B. S.:

[147454]Localization of the amino acid substitution site in a new variant

[147455]of human serum albumin, albumin Mexico–2. Proc. Nat. Acad. Sci. 77:

[147456]2505–2509, 1980.

[147457]26545.

[147458]Fraser, G. R.; Harris, H.; Robson, E. B.: A new genetically determined

[147459]plasma-protein in man. Lancet I: 1023-1024, 1959.

[147460]26546.

[147461]Galliano, M.; Minchiotti, L.; Ferri, G.; Iadarola, P.; Zapponi,

[147462]M. C.; Fine, J. M.: Structural characterization of the human albumin

[147463]variant 'Pollibauer'. Blood Transf. Immunohemat. 27: 597-602, 1984.

[147464]26547.

[147465]Galliano, M.; Minchiotti, L.; Iadarola, P.; Ferri, G.; Zapponi,

[147466]M. C.; Castellani, A. A.: The amino acid substitution in albumin

[147467]Roma: 321 glu-to-lys. FEBS Lett. 233: 100-104, 1988.

[147468]26548.

[147469]Galliano, M.; Minchiotti, L.; Iadarola, P.; Porta, F.; Stoppini,

[147470]M.; Zapponi, M. C.; Ferri, G.; Castellani, A. A.: Genetic variants

[147471]of human serum albumin. Prog. Med. Lab. 2: 475-477, 1988.

[147472]26549.

[147473]Galliano, M.; Minchiotti, L.; Iadarola, P.; Stoppini, M.; Giagnoni,

[147474]P.; Watkins, S.; Madison, J.; Putnam, F. W.: Protein and DNA sequence

[147475]analysis of a 'private' genetic variant: albumin Ortonovo (glu505-to-lys). Biochim.

[147476]Biophys. Acta 1225: 27-32, 1993.

[147477]26550.

[147478]Galliano, M.; Minchiotti, L.; Iadarola, P.; Zapponi, M. C.; Ferri,

[147479]G.; Castellani, A. A.: Structural characterization of a chain termination

[147480]mutant of human serum albumin. J. Biol. Chem. 261: 4283-4287, 1986.

[147481]26551.

[147482]Galliano, M.; Minchiotti, L.; Porta, F.; Rossi, A.; Ferri, G.;

[147483]Madison, J.; Watkins, S.; Putnam, F. W.: Mutations in genetic variants

[147484]of human serum albumin found in Italy. Proc. Nat. Acad. Sci. 87:

[147485]8721–8725, 1990.

[147486]26552.

[147487]Galliano, M.; Minchiotti, L.; Stoppini, M.; Tarnoky, A. L.: A

[147488]new proalbumin variant: albumin Jaffna (–1 arg-to-leu).

FEBS Lett. 255:

[147489]295–299, 1989.

[147490]